

K-STABILITY OF SPECIAL GUSHEL-MUKAI MANIFOLDS

YUCHEN LIU AND LINSHENG WANG

ABSTRACT. Gushel-Mukai manifolds are specific families of n -dimensional Fano manifolds of Picard rank 1 and index $n-2$ where $3 \leq n \leq 6$. A Gushel-Mukai n -fold is either ordinary, i.e. a hyperquadric section of a quintic Del Pezzo $(n+1)$ -fold, or special, i.e. it admits a double cover over the quintic Del Pezzo n -fold branched along an ordinary Gushel-Mukai $(n-1)$ -fold. In this paper, we prove that a general special Gushel-Mukai n -fold is K-stable for every $3 \leq n \leq 6$. Furthermore, we give a description of the first and last walls of the K-moduli of the pair (M, cQ) , where M is the quintic Del Pezzo fourfold (or fivefold) and Q is an ordinary Gushel-Mukai threefold (or fourfold). Besides, we compute δ -invariants of quintic Del Pezzo fourfolds and fivefolds which were shown to be K-unstable by K. Fujita, and show that they admit Kähler-Ricci solitons.

1. INTRODUCTION

The theory of K-stability, first introduced by Tian [Tia97] and later reformulated algebraically by Donaldson [Don02], is an algebraic stability theory that characterizes the existence of canonical metrics on Fano varieties. According to the Yau-Tian-Donaldson correspondence [CDS15, Tia15, Ber16, BBJ21, LTW22, Li22, LXZ22], a Fano variety is K-polystable if and only if it admits a Kähler-Einstein metric. Thus it is an important problem to check K-stability for explicit Fano manifolds. In dimension 2, K-stability of Del Pezzo surfaces was completely determined in [Tia90]. For Fano threefolds, this problem was largely solved, see e.g. [LX19, AZ23, ACC⁺23]. In higher dimensions, much progress has been made for Fano hypersurfaces, see e.g. [Che01, Fuj19, AZ22, AZ23, Liu22]. Despite these developments, it remains a challenge to determine K-stability for most Fano manifolds in higher dimensions.

In this paper, we study the K-stability of Gushel-Mukai manifolds which are specific families of n -dimensional Fano manifolds of Picard rank 1 and index $n-2$ where $3 \leq n \leq 6$. Let us recall relevant notions. Let X be an n -dimensional Fano manifold with Picard group generated by an ample divisor H . The *index* of X is the positive integer r such that $-K_X \sim rH$. The self-intersection number $d := (H^n)$ is called the *degree* of X . For $r \geq n$, Kobayashi and Ochiai proved in [KO73] that X is isomorphic to either \mathbb{P}^n (where $r = n+1$) or a smooth quadric hypersurface in \mathbb{P}^{n+1} (where $r = n$). Fano manifolds with index $r = n-1$, known as *Del Pezzo manifolds*, were classified by T. Fujita and Iskovskikh from 1977 to 1988; see for example [Fuj90]. Fano manifolds with index $r = n-2$ were studied by Gushel [Gus83] and Mukai [Muk89]. It was proved by Mukai that there are only 10 possibilities for the degree d . For the case $d = 10$, the corresponding Fano manifolds are called *Gushel-Mukai (GM) manifolds*. See [Deb20] for a comprehensive survey on Gushel-Mukai manifolds.

There are two types of GM manifolds: *ordinary GM n -folds* Q_n are quadratic sections of quintic Del Pezzo $(n + 1)$ -folds M_{n+1} (which is a smooth $(n + 1)$ -dimensional linear section of $\text{Gr}(2, 5)$); *special GM n -folds* X_n are double covers of quintic Del Pezzo n -folds M_n branched along an ordinary GM $(n - 1)$ -fold Q_{n-1} . For $3 \leq n \leq 6$, GM n -folds are Fano manifolds. It was proved by [AZ23, Theorem 5.1] that all the smooth GM threefolds are K-stable.

Our first main result proves the K-stability for a general member of special GM manifolds.

Theorem 1.1. *A general special GM manifold of dimension 4, 5, or 6 is K-stable.*

Since the parameter space of the special GM n -folds is an irreducible closed subvariety of the parameter space of the GM n -folds, which is also irreducible (see Section 2 for details), we have the following corollary by openness of K-stability [BL22, BLX22, LXZ22].

Corollary 1.2. *A general GM manifold of dimension 4, 5, or 6 is K-stable.*

Recall that a special GM n -fold X_n admits a double cover $X_n \rightarrow M_n$ branched along an ordinary GM $(n - 1)$ -fold Q_{n-1} (see Section 2). By [LZ22a, Theorem 1.2] and [Zhu21], we know that the K-polystability of X_n is equivalent to the K-polystability of the log Fano pair $(M_n, \frac{1}{2}Q_{n-1})$. Hence it's natural to study the K-stability of quintic Del Pezzo manifolds M_n at first.

The quintic Del Pezzo surfaces and threefolds are K-polystable, which were proved by [Tia90] and [CS09] respectively. However, the quintic Del Pezzo fourfolds and fivefolds are K-unstable. They are among the first examples of K-unstable Fano manifolds with Picard number one found by K. Fujita [Fuj17]. He found a 2-plane $S \subseteq M_4 \subseteq M_5$ (which is of “non-vertex type”, see Section 3.1 for detail) such that $\beta(E_S) < 0$, where E_S is the exceptional divisor of the blowup of M_4 or M_5 along S . We further show that

Theorem 1.3. *The delta invariant of M_4 is $\delta(M_4) = \frac{25}{27}$, which is minimized by E_S .*

However, the delta invariant of M_5 is not minimized by E_S (where $\frac{A_{M_5}(E_S)}{S(E_S)} = \frac{45}{46}$, see Section 4.1 for details). There exists a 3-space $W = \mathbb{P}^3 \subseteq M_5$ such that $\frac{A_{M_5}(E_W)}{S(E_W)} = \frac{15}{16} < \frac{45}{46}$, where E_W is the exceptional divisor of the blowup of M_5 along W . We show that

Theorem 1.4. *The delta invariant of M_5 is $\delta(M_5) = \frac{15}{16}$, which is minimized by E_W .*

Next, we study the wall-crossing problem of the K-moduli spaces of (M_n, cQ_{n-1}) for $n = 4$ or 5 . Recall that for any Fano manifold X and divisor $D \in |-rK_X|$, where $r > 0$ is a rational number such that rK_X is Cartier, the *K-semistable domain* of (X, D) is defined by

$$\text{Kss}(X, D) = \{c \in [0, \min\{1, r^{-1}\}] : (X, cD) \text{ is K-semistable}\}.$$

We determine the K-semistable domains of (M_4, Q_3) and (M_5, Q_4) . This is an essential step in the proof of Theorem 1.1.

Theorem 1.5. *For general $Q_3 \in |\mathcal{O}_{M_4}(2)|$ and $Q_4 \in |\mathcal{O}_{M_5}(2)|$, we have*

$$\text{Kss}(M_4, Q_3) = \left[\frac{1}{9}, \frac{7}{8} \right], \quad \text{Kss}(M_5, Q_4) = \left[\frac{1}{8}, \frac{4}{5} \right].$$

The last walls are obtained by computing beta invariants at ord_{Q_3} and ord_{Q_4} . However, the first walls are more mysterious. To overcome the difficulty, we construct special quadric sections $Q_{3,0}$ on M_4 and $Q_{4,0}$ on M_5 to proceed K-polystable pairs, see Section 6 for the definition of $Q_{3,0}$ and $Q_{4,0}$.

Theorem 1.6. *The log Fano pairs $(M_4, \frac{1}{9}Q_{3,0})$ and $(M_5, \frac{1}{8}Q_{4,0})$ are K-polystable.*

With a similar strategy, we may consider the weighted K-stability instead of the K-stability of log Fano pairs. For the soliton candidates ξ_0 on M_4 and η_0 on M_5 (see Section 7), we have

Theorem 1.7. *The Fano pairs (M_4, ξ_0) and (M_5, η_0) are weighted K-polystable.*

Hence by [HL23, BLXZ23], we get the existence of Kähler-Ricci solitons on quintic Del Pezzo manifolds, which answers a question asked by [Ino19, Del22].

Corollary 1.8. *Any quintic Del Pezzo fourfold or fivefold admits a Kähler-Ricci soliton.*

The paper is organized as follows. In Section 2, we recall the definition of the quintic Del Pezzo manifolds and the GM manifolds. Next, we give a detailed description of the quintic Del Pezzo fourfolds and fivefolds, and compute their delta invariants in Section 3 and 4 respectively. We prove the K-stability of general special GM fourfolds and fivefolds in Section 5. Finally, we study the wall-crossing problem of the K-moduli spaces of (M_4, cQ_3) and (M_5, cQ_4) in Section 6, and we show that M_4 and M_5 both admit Kähler-Ricci solitons in the last section.

Acknowledgments. We would like to thank Minghao Miao, Lu Qi, Fei Si, and Shengxuan Zhou for helpful discussions. The first author was partially supported by NSF CAREER Grant DMS-2237139 and the Alfred P. Sloan Foundation. The second author would like to thank Gang Tian for his support and guidance.

2. PRELIMINARIES

Throughout this paper, we work over the field of complex numbers \mathbb{C} .

We recall the definition of GM manifolds and quintic Del Pezzo manifolds in this section.

Let V_5 be a \mathbb{C} -vector spaces of dimension 5. Let $\text{Gr} = \text{Gr}(2, V_5) \subseteq \mathbb{P}(\wedge^2 V_5) =: \mathbb{P}$ be the Grassmannian of 2-dimensional subspaces in V_5 , and $\text{CGr} \subseteq \mathbb{P}(\mathbb{C} \oplus \wedge^2 V_5) =: \tilde{\mathbb{P}}$ be the projective cone over Gr with vertex $v = \mathbb{P}(\mathbb{C})$. We have the natural projection map $\pi : \tilde{\mathbb{P}} \setminus \{v\} \rightarrow \mathbb{P}$, which restricts to $\pi_{\text{Gr}} : \text{CGr} \setminus \{v\} \rightarrow \text{Gr}$. For $3 \leq n \leq 6$ we define

$$\begin{aligned} X_n &= \text{CGr} \cap H_{6-n} \cap Q, \\ M_n &= \text{Gr} \cap H'_{6-n}, \\ Q_{n-1} &= \text{Gr} \cap H'_{6-n} \cap Q', \end{aligned}$$

where $H_{6-n} \subseteq \tilde{\mathbb{P}}, H'_{6-n} \subseteq \mathbb{P}$ are linear subspaces of codimension $6 - n$, and $Q \subseteq \tilde{\mathbb{P}}, Q' \subseteq \mathbb{P}$ are quadric hypersurfaces. We assume that X_n, M_n and Q_{n-1} are smooth, hence $v \notin Q$.

It is not difficult to show that X_n is a Fano manifold of dimension n with $\text{Pic}(X) = \langle H \rangle$, where $H = \mathcal{O}_{\tilde{\mathbb{P}}}(1)|_{X_n}$. We also have $(-K_{X_n}) = (n-2)H$ and $H^n = 10$. On the other hand, we see that M_n is a Fano manifold with $-K_{M_n} = \langle H \rangle$, where $H = \mathcal{O}_{\mathbb{P}}(1)|_{M_n}$. And $(-K_{M_n}) = (n-1)H, H^n = 5$. We have the following classification result due to Mukai [Muk89] and T. Fujita [Fuj81] respectively.

Definition-Theorem 2.1 ([Muk89]). *Let X_n be a Fano manifold of dimension n with Picard number one, index $n - 2$, and degree 10. Then $n \in \{3, 4, 5, 6\}$ and there exists a codimension $(6 - n)$ subspace $H_{6-n} \subseteq \tilde{\mathbb{P}}$ and a quadric hypersurface $Q \subseteq \tilde{\mathbb{P}}$ such that*

$$X_n = \text{CGr} \cap H_{6-n} \cap Q.$$

The manifolds X_n are called Gushel-Mukai (GM) manifolds. There are two different types of X_n :

- (1) *If $v \in H_{6-n}$, then we have $H'_{6-n} = H_{6-n} \cap \mathbb{P} \subseteq \tilde{\mathbb{P}}$, and π induces a double cover $X_n \rightarrow M_n$ branched along some $Q_{n-1} \subseteq H'_{6-n}$. We say that X_n is a special Gushel-Mukai manifold;*
- (2) *If $v \notin H_{6-n}$, then $n \leq 5$ and π maps $H_{6-n} \subseteq \tilde{\mathbb{P}}$ isomorphically to $H'_{5-n} = \pi(H_{6-n}) \subseteq \mathbb{P}$, hence maps X_n isomorphically to $Q_n = \pi(X_n) \subseteq H'_{5-n}$. We say that X_n is an ordinary Gushel Mukai manifold.*

Note that if $n = 6$ then every Gushel-Mukai manifold is special.

Definition-Theorem 2.2 ([Fuj81]). *Let M_n be a Fano manifold of dimension n with Picard number one, index $n - 1$, and degree 5. Then M_n is a linear section of $\text{Gr}(2, V_5) \subseteq \mathbb{P}(\wedge^2 V_5)$. The manifolds M_n are called quintic Del Pezzo manifolds.*

- Remark 2.3.**
- (1) *If $n = 1$ or 2 , then the above definition extends verbatim to smooth GM curves and GM surfaces. By [DK18], a smooth GM curve is precisely a Clifford general smooth curve of genus 6, and a smooth GM surface is precisely a Brill-Noether general smooth polarized K3 surface of degree 10.*
 - (2) *By the discussions above we know that for any special GM n -fold X_n , there exists a double cover $\pi : X_n \rightarrow M_n$ branched along an ordinary GM $(n - 1)$ -fold Q_{n-1} . On the other hand, the parameter space of all the GM n -folds is irreducible and contains the parameter space of special GM n -folds as an irreducible subvariety.*

By [Fuj81], we know that the quintic Del Pezzo manifolds are unique.

Theorem 2.4 ([Fuj81]). *For any $2 \leq n \leq 6$, all the quintic del Pezzo n -folds are isomorphic to each other.*

The quintic Del Pezzo fourfolds and fivefolds are among the first examples of K-unstable Fano manifolds with Picard number one found by K. Fujita [Fuj17].

Theorem 2.5 ([Fuj17]). *The quintic Del Pezzo fourfolds and fivefolds are K-unstable.*

On the other hand, by [Tia90] and [CS09] we know that the quintic Del Pezzo surface M_2 and threefold M_3 are K-polystable. Since $M_6 = \text{Gr}(2, V_5)$ is a homogeneous space, it is K-polystable, see for example [ACC⁺23, Theorems 1.2.5 and 1.4.7].

The following result summarizes fundamental properties of the moduli spaces of smooth GM manifolds.

Theorem 2.6 ([DK20, Deb20]). *For any $3 \leq n \leq 6$, there exists a quasi-projective irreducible coarse moduli space \mathbf{M}_n^{GM} parameterizing smooth GM n -folds. Moreover, $\dim \mathbf{M}_n^{\text{GM}} = 25 - \frac{(6-n)(5-n)}{2}$. The moduli space \mathbf{M}_n^{GM} is the union of an open subvariety $\mathbf{M}_{n,\text{ord}}^{\text{GM}}$ and an irreducible closed subvariety $\mathbf{M}_{n,\text{spe}}^{\text{GM}}$, complement to each other, that parameterize ordinary and special GM n -folds respectively. Moreover, the following properties hold.*

- (1) $\mathbf{M}_{n,\text{ord}}^{\text{GM}} = \emptyset$ if and only if $n = 6$.
- (2) $\mathbf{M}_{n,\text{spe}}^{\text{GM}} \cong \mathbf{M}_{n-1,\text{ord}}^{\text{GM}}$ for $4 \leq n \leq 6$.
- (3) $\mathbf{M}_{3,\text{spe}}^{\text{GM}}$ is isomorphic to the moduli space $\mathbf{M}_{2,\text{ord,ss}}^{\text{GM}}$ parameterizing strongly smooth GM surfaces, see [DK18, Definition 3.15].

3. THE QUINTIC DEL PEZZO FOURFOLDS

We overview basic facts of the quintic Del Pezzo fourfolds in this section. We first compute the automorphism group by viewing the quintic Del Pezzo fourfolds as linear sections of $\text{Gr}(2, V_5)$ via the Plücker embedding. T.Fujita classified the quintic Del Pezzo manifolds by showing that they are birational to the projective space in an explicit way. This is the key ingredient in our computation of the delta minimizer of the Del Pezzo fourfolds.

3.1. Properties of quintic Del Pezzo fourfolds. Let $M = M_4$ be a quintic Del Pezzo fourfold, with the Plücker polarization $\mathcal{O}_M(1)$. We denote by $S \subseteq M$ a plane of *non-vertex type* (introduced by [Fuj81] with the property $c_2(N_{S/M}) = 2$, which will be formulated explicitly in Section 3.1). Let $\pi : \tilde{M} \rightarrow M$ be the blowup of M along S with exceptional divisor E_S . There is a unique divisor $D_9 \in |\mathcal{O}_M(1)|$ with $\text{ord}_{E_S}(D_9) = 2$, and let \tilde{D}_9 be the strict transform of D_9 on \tilde{M} . It was shown in [Fuj81, (10.9)] that there is a morphism $\tau : \tilde{M} \rightarrow \mathbb{P}^4$, which is the blowup of \mathbb{P}^4 along a twisted cubic curve C_3 in a hyperplane $H_9 \subseteq \mathbb{P}^4$ with exceptional divisor \tilde{D}_9 , and E_S is the strict transform of H_9 . Hence \tilde{D}_9 and E_S intersect transversally. Denote $F = \tilde{D}_9 \cap E_S$. Write the restrictions as $\pi_{E_S} : E_S \rightarrow S$ and $\tau_{E_S} : E_S \rightarrow H_9$. We will show that there is an $\text{Aut}(M)$ -invariant conic curve $C_2 \subseteq S$. Denote by $\tilde{T} = \pi^{-1}(C_2) \subseteq E_S$. Then $T = \tau_{E_S}(\tilde{T}) \subseteq H_9$ is the tangent developable of C_3 , which is a quartic surface with a node along C_3 .

We compute the intersection numbers of some divisors on E_S . Since $\tau_{E_S} : E_S \rightarrow H_9 \cong \mathbb{P}^3$ is the blowup of \mathbb{P}^3 along a twisted cubic C_3 with exceptional divisor F and the normal bundle $N_{C_3/H_9} \cong \mathcal{O}_{C_3}(5)^{\oplus 2}$, the restriction $\tau_F : F \rightarrow C_3$ is a trivial \mathbb{P}^1 -bundle with $F|_F = -h_F + 5l_F$, where $l_F = \tau_F^* \mathcal{O}_{C_3}(1)$ and h_F is the pull-back of $\mathcal{O}_{\mathbb{P}^1}(1)$ on a fiber. Note that $\mathcal{O}_{H_9}(1)|_{C_3} \sim \mathcal{O}_{C_3}(3)$,

and $(\tau_{E_S}^* \mathcal{O}_{H_9}(1))|_F \sim \tau_F^*(\mathcal{O}_{H_9}(1)|_{C_3})$. We omit $\tau_{E_S}^*$ and have

$$(1) \quad \mathcal{O}_{H_9}(1)^2 \cdot F = 0, \quad \mathcal{O}_{H_9}(1) \cdot F^2 = -3, \quad F^3 = -10.$$

We shall present a quintic Del Pezzo fourfold explicitly in the projective space by writing down the defining equations for the convenience of computing the automorphism group and to study the Sarkisov link structure presented above. Recall that $\text{Gr} = \text{Gr}(2, V_5)$ is embedded in $\mathbb{P}(\wedge^2 V_5)$ via Plücker embedding, and $M = M_4$ is a smooth intersection $\text{Gr} \cap H \cap H'$, where H, H' are hyperplanes in $\mathbb{P}(\wedge^2 V_5)$. Let $\{e_0, e_1, e_2, e_3, e_4\}$ be a basis of V_5 , and $\{e_j^*\} \subseteq V_5^*$ be the corresponding dual basis. Set $e_{ij} = e_i \wedge e_j, e_{ij}^* = e_i^* \wedge e_j^*$. Any point of $\wedge^2 V_5$ can be denoted by

$$X_0 e_{01} + X_1 e_{02} + X_2 e_{03} + X_3 e_{04} + X_4 e_{12} + X_5 e_{13} + X_6 e_{14} + X_7 e_{23} + X_8 e_{24} + X_9 e_{34},$$

where $X_i \in \mathbb{C}$, and $[X_0, \dots, X_9]$ forms a projective coordinate of $\mathbb{P}(\wedge^2 V_5)$. The defining equations of $G \subseteq \mathbb{P}(\wedge^2 V_5)$ are

$$\begin{cases} X_0 X_7 - X_1 X_5 + X_2 X_4 = 0, \\ X_0 X_8 - X_1 X_6 + X_3 X_4 = 0, \\ X_0 X_9 - X_2 X_6 + X_3 X_5 = 0, \\ X_1 X_9 - X_2 X_8 + X_3 X_7 = 0, \\ X_4 X_9 - X_5 X_8 + X_6 X_7 = 0. \end{cases}$$

Any hyperplane $H \subseteq \mathbb{P}(\wedge^2 V_5)$ corresponds to a linear functional $l \in \wedge^2 V_5^*$, which can be viewed as a skew-symmetric 2-form on V_5 . We define

$$l = 2(e_{03}^* + e_{14}^*), \quad l' = 2(e_{13}^* + e_{24}^*),$$

whose corresponding skew-symmetric matrices under the basis $\{e_i\}$ are

$$J = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \end{pmatrix}, \quad J' = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \end{pmatrix}.$$

The hyperplanes defined by l, l' are $H = \{X_2 + X_6 = 0\}$ and $H' = \{X_5 + X_8 = 0\}$. One may check that $M = G \cap H \cap H' \subseteq \mathbb{P}_{X_0, X_1, X_2, X_3, X_4, X_7, X_8, X_9}^7$ is smooth, hence is a quintic Del Pezzo fourfold. And the defining equation of $M \subseteq \mathbb{P}^7$ becomes

$$\begin{cases} X_0 X_7 + X_1 X_8 + X_2 X_4 = 0, \\ X_0 X_8 + X_1 X_2 + X_3 X_4 = 0, \\ X_0 X_9 + X_2^2 - X_3 X_8 = 0, \\ X_1 X_9 - X_2 X_8 + X_3 X_7 = 0, \\ X_4 X_9 + X_8^2 - X_2 X_7 = 0. \end{cases}$$

It was shown in [Fuj81, (10.9)] that there are two types of planes on M , that is, the *non-vertex type* planes S and the *vertex type* planes T . They are characterized by the second Chern classes of

their normal bundles in M :

$$c_2(N_{S/M}) = 2, \quad c_2(N_{T/M}) = 1.$$

In our explicit case, they are

$$\begin{aligned} S &= \{X_2 = X_3 = X_7 = X_8 = X_9 = 0\}, \\ R_\lambda &= \{\lambda^2 X_0 - \lambda X_1 + X_4 = X_2 + \lambda X_3 = X_7 + \lambda^3 X_3 = X_8 - \lambda^2 X_3 = X_9 = 0\}, \lambda \in \mathbb{C}, \\ R_\infty &= \{X_0 = X_2 = X_3 = X_8 = X_9 = 0\}. \end{aligned}$$

K.Fujita proved in [Fuj17] that $\beta(E_S) < 0$, hence M is a K-unstable.

With this explicit $M = M_4$, we could study the diagram $M \leftarrow \tilde{M} \rightarrow \mathbb{P}^4$ locally.

Let's work on the affine open subset $\{X_0 = 1\}$ of M , which is isomorphic to \mathbb{A}^4 with coordinate $(X_1, X_4, X_2, X_3) =: (u, v, x, y)$. Hence

$$\begin{cases} X_7 = u(ux + vy) - vx, \\ X_8 = -(ux + vy), \\ X_9 = -y(ux + vy) - x^2, \end{cases}$$

and $S = \{x = y = 0\}$. So $\tilde{M} = \text{Bl}_{(x,y)} M = \{\xi_0 y = \xi_1 x\} \subseteq M \times \mathbb{P}_{\xi_0, \xi_1}^1$. We also replace \tilde{M} by the affine open subset $\{\xi_1 = 1\}$ and denote by $s = \xi_0$, which is isomorphic to \mathbb{A}^4 with coordinate (u, v, s, y) . And the morphism π is just

$$\pi : \tilde{M} \rightarrow M, \quad (u, v, s, y) \mapsto (u, v, sy, y).$$

Hence the exceptional divisor of $\pi : \tilde{M} \rightarrow M$ is $E_S = \{y = 0\}$, and

$$\begin{cases} \pi^*(X_2) = sy, \\ \pi^*(X_3) = y, \\ \pi^*(X_7) = (u(us + v) - vs)y, \\ \pi^*(X_8) = -(us + v)y, \\ \pi^*(X_9) = -(s^2 + us + v)y^2. \end{cases}$$

The morphism $\tau : \tilde{M} \rightarrow \mathbb{P}^4$ is determined by $|\mathcal{O}_M(1) - E_S| = \pi^*\langle X_2, X_3, X_7, X_8, X_9 \rangle$. Locally,

$$\tau : (u, v, s, y) \mapsto [s, 1, u(us + v) - vs, -(us + v), -(s^2 + us + v)y].$$

The strict transform of $D_9 = \{X_9 = 0\}$ is $\tilde{D}_9 = \{s^2 + us + v = 0\}$, which is contracted by $\tau : \tilde{M} \rightarrow \mathbb{P}^4$ to the twisted cubic curve $C_3 : \mathbb{A}^1 \rightarrow \mathbb{P}^4, t \mapsto [1, t, t^2, t^3, 0]$. This is just the exceptional divisor of the blowup $\tau : \tilde{M} \rightarrow \mathbb{P}^4$ of this twisted cubic curve. We also see that $\tau(E_S) = H_9 \cong \mathbb{P}^3$ is the projective 3-space spanned by C_3 .

Consider the tangent developable T of C_3 in $H_9 = \mathbb{P}_{X,1,Z,Y}^3$

$$\{4Y(Y - X^2)^2 - 4X(Y - X^2)(Z - XY) + (Z - XY)^2 = 0\}.$$

This also defines the cone over the tangent developable of C_3 in \mathbb{P}^4 , whose strict transform to \tilde{M} is

$$\tilde{Q}_0 = \{4Y - 4X(-u) + (-u)^2 = 0\} = \{u^2 - 4v = 0\},$$

where $X = s, Y = -(us + v)$ and $Z = u(us + v) - vs$ (hence $Z - XY = (-u)(Y - X^2)$). Hence this is also the strict transform via π of the quadric hypersurface

$$(2) \quad Q_0 = \{X_1^2 - 4X_0X_4 = 0\} \subseteq M.$$

Remark 3.1. We will see that Q_0 is invariant under a maximal reductive subgroup G of $\text{Aut}(M)$. The log Fano pair (M, cQ_0) is essential in our study of the K-stability of special GM fourfolds.

We denote by $\tilde{T} = \tilde{Q}_0 \cap E_S$ the strict transform via τ of the tangent developable T of $C_3 \subseteq H_9$, and by $C_2 = Q_0 \cap S$ the corresponding conic curve in $S = \mathbb{P}_{X_0, X_1, X_4}^2$. Then $R = \pi^{-1}(C_2)$ and $\pi : R \rightarrow C_2$ is a trivial \mathbb{P}^1 -bundle. On the other hand, the restrictions of π and τ to E_S are

$$\begin{aligned} \pi_{E_S} : E_S &\rightarrow S, & (u, v, s) &\mapsto (u, v), \\ \tau_{E_S} : E_S &\rightarrow \mathbb{P}^3, & (u, v, s) &\mapsto [s, 1, u(us + v) - vs, -(us + v)]. \end{aligned}$$

We see that E_S is a \mathbb{P}^1 -bundle over S via π , and is the blowup of $H_9 \subseteq \mathbb{P}^3$ along the twisted cubic C_3 with exceptional divisor $F = \tilde{D}_9 \cap E_S$. And $\pi_F : F \rightarrow S$ is a double cover ramified along $C_2 \subseteq S$.

3.2. The automorphism group of a quintic Del Pezzo fourfold. We compute the automorphism group of $M = M_4$ in our explicit setting in this subsection, see [PVdV99]. We would like to mention that there is a detailed study of $\text{Aut}(M)$ in [PZ18].

Note that $M = \{p \in \text{Gr} : l(p) = l'(p) = 0\}$ and $\text{Aut}(\text{Gr}) = \text{PGL}_5(\mathbb{C})$. For any $P \in \text{GL}_5(\mathbb{C})$, $P(M) = M$ is equivalent that $P^*\langle l, l' \rangle = \langle l, l' \rangle$. Hence

$$\text{Aut}(M) = \{P \in \text{GL}_5(\mathbb{C}) : \langle P^T J P, P^T J' P \rangle = \langle J, J' \rangle\} / \mathbb{G}_m,$$

$$\text{aut}(M) = \{Q \in \mathfrak{sl}_5(\mathbb{C}) : \langle Q^T J + JQ, Q^T J' + J'Q \rangle = \langle J, J' \rangle\}.$$

For any $Q = (q_{ij}) \in \text{aut}(M)$, there exist $a, b, c, d \in \mathbb{C}$ such that

$$\begin{cases} Q^T J + JQ &= aJ + bJ', \\ Q^T J' + J'Q &= cJ + dJ'. \end{cases}$$

We get some linear equations in q_{ij} , by solving which we see that there exist $e, f, g, h \in \mathbb{C}$ such that

$$(3) \quad Q = \begin{pmatrix} -2d & 2b & 0 & e & f \\ c & -a - d & b & f & g \\ 0 & 2c & -2a & g & h \\ 0 & 0 & 0 & a + 2d & -c \\ 0 & 0 & 0 & -b & 2a + d \end{pmatrix}.$$

The Lie algebra $\text{aut}(M)$ is non-reductive. Let $G \subseteq \text{Aut}(M)$ be the reductive subgroup whose Lie algebra is generated by

$$(4) \quad \begin{pmatrix} -2d & 2b & 0 & 0 & 0 \\ c & -a-d & b & 0 & 0 \\ 0 & 2c & -2a & 0 & 0 \\ 0 & 0 & 0 & a+2d & -c \\ 0 & 0 & 0 & -b & 2a+d \end{pmatrix}.$$

Then the quadric divisor Q_0 of M defined by $\{X_1^2 - 4X_0X_4 = 0\}$ is G -invariant, and $G = \text{Aut}(M, Q_0)$. It's clear that $D_9 = \{X_9 = 0\} \cap M$ and $S \subseteq D_9$ are G -invariant. We denote by $D_{9,Q} = Q_0 \cap D_9$, then $C_2 = D_{9,Q} \cap S = Q_0 \cap S$ is the unique G -invariant curve in $S \cong \mathbb{P}^2$. The point $o_9 = [0, \dots, 0, 1] \in Q_0$ is the unique G -invariant point on Q_0 . By [PVdV99, Proposition 6.8], the $\text{Aut}(M)$ -orbit decomposition of M is

$$M = (M \setminus D_9) \sqcup (D_9 \setminus S) \sqcup (S \setminus C_2) \sqcup C_2.$$

Moreover, the G -orbit decomposition of M is

$$\begin{aligned} M = & \left(M \setminus (Q_0 \cup D_9) \right) \sqcup \left(Q_0 \setminus (D_{9,Q} \cup o_9) \right) \sqcup \left(D_9 \setminus (D_{9,Q} \cup S) \right) \sqcup \\ & (D_{9,Q} \setminus C_2) \sqcup (S \setminus C_2) \sqcup C_2 \sqcup o_9. \end{aligned}$$

3.3. The delta function on the co-weight lattice. We first present an Okounkov body of M_4 in this subsection, which will be used in the computation of the expected vanishing order (S -invariant). Consider the affine chart $U_9 = M \cap \{X_9 \neq 0\}$, then

$$\begin{cases} x_0 &= x_2^2 - x_3x_8, \\ x_1 &= x_2x_8 - x_3x_7, \\ x_4 &= x_8^2 - x_2x_7, \end{cases}$$

where $x_i = X_i/X_9 (i = 0, 1, 2, 3, 4, 7, 8, 9)$ and $U_9 \cong \mathbb{A}_{x_2, x_3, x_7, x_8}^4$. The coordinate hyperplanes induce a faithful valuation $\mathfrak{v} : K(M)^\times \rightarrow \mathbb{Z}^4$ as follows (with respect to the lexicographic ordering of \mathbb{Z}^4)

$$\begin{cases} \mathfrak{v}(x_2) = (1, 0, 0, 0), \\ \mathfrak{v}(x_3) = (0, 1, 0, 0), \\ \mathfrak{v}(x_7) = (0, 0, 1, 0), \\ \mathfrak{v}(x_8) = (0, 0, 0, 1), \end{cases} \quad \begin{cases} \mathfrak{v}(x_0) = (0, 1, 0, 1), \\ \mathfrak{v}(x_1) = (0, 1, 1, 0), \\ \mathfrak{v}(x_4) = (0, 0, 0, 2). \end{cases}$$

The convex polytope spanned by the above seven points in \mathbb{R}^4 (denoted by Δ) has volume $\frac{5}{24}$, which is equal to $\text{vol}(\mathcal{O}(1))/4!$. Hence Δ is an Okounkov body of the polarized variety $(M, \mathcal{O}_M(1))$.

We denote by $P_a(t) = \exp(-tQ_a)$ and $P_d(t) = \exp(-tQ_d) \in \text{Aut}(M)$ the one-parameter groups determined by

$$Q_a = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix}, \quad Q_d = \begin{pmatrix} -2 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then M admits a $\mathbb{T} = \mathbb{G}_m^2$ -action generated by P_a and P_d , whose actions on $\mathbb{P} \cong \mathbb{P}^7$ are

$$\begin{aligned} P_a(t) \cdot [X_i] &= [tX_0, t^2X_1, t^{-1}X_2, t^{-2}X_3, t^3X_4, tX_7, X_8, t^{-3}X_9], \\ P_d(t) \cdot [X_i] &= [t^3X_0, t^2X_1, X_2, tX_3, tX_4, t^{-2}X_7, t^{-1}X_8, t^{-3}X_9]. \end{aligned}$$

The weights of the \mathbb{T} -action on \mathbb{P}^7 are

$$\begin{aligned} \eta &= (1, 2, -1, -2, 3, 1, 0, -3), \\ \zeta &= (3, 2, 0, 1, 1, -2, -1, -3). \end{aligned}$$

Let $N_{\mathbb{Z}} = \text{Hom}(\mathbb{G}_m, \mathbb{T})$, $M_{\mathbb{Z}} = \text{Hom}(\mathbb{T}, \mathbb{G}_m)$ be the co-weight and weight lattice of the \mathbb{T} -action. Then N is generated by η and ζ . Any $(a, b) \in \mathbb{R}^2$ determines a one-parameter subgroup generated by $a\eta + b\zeta$. We denote by $\nu_{(a,b)}$ the valuation induced by $a\eta + b\zeta \in N_{\mathbb{R}}$. Hence the delta function $\delta(\nu) = A_M(\nu)/S_M(\nu)$ is defined on $N_{\mathbb{R}}$. The delta function is invariant up to scaling, that is, $\delta(\nu_{(ka, kb)}) = \delta(\nu_{(a,b)})$ for any $k \in \mathbb{R}_{>0}$.

One may shift the weight such that all the numbers in the weight are non-negative and at least one number is equal to zero. Explicitly, let $m_{(a,b)} = \min_{0 \leq i \leq 9} \text{wt}_{(a,b)}(i)$ and $i_{(a,b)}$ be the minimizer. We define the *shifted weight* by $\overline{\text{wt}}_{(a,b)} = \text{wt}_{(a,b)} - m_{(a,b)}\text{wt}_1$, where $\text{wt}_1 = (1, \dots, 1)$. Then the valuation $\nu_{(a,b)}$ on M is determined by

$$(5) \quad \nu_{(a,b)}(X_i) = \overline{\text{wt}}_{(a,b)}(i),$$

for any $0 \leq i \leq 9, i \neq 5, 6$. For example, consider the one-parameter subgroup defined by ζ , that is, $(a, b) = (0, 1)$. We have $m_{(0,1)} = -3$ and $i_{(0,1)} = 9$. Then the shifted weight is

$$\overline{\text{wt}}_{(0,1)} = (6, 5, 3, 4, 4, 1, 2, 0),$$

which defines a quasimonomial valuation $\nu_{(0,1)}$ on U_9 using formula (5).

Next, we compute the delta function on $N_{\mathbb{R}}$. For example, consider the valuation $\nu_{(0,1)} = \nu_{\zeta}$ computed above, which is a monomial valuation on U_9 with weight $(3, 4, 1, 2)$ with respect to x_2, x_3, x_7, x_8 , where $x_i = X_i/X_9 (i = 2, 3, 7, 8)$ is the coordinate chart of U_9 . Hence by [JM12, Proposition 5.1], the log discrepancy of $\nu_{(0,1)}$ is

$$A_M(\nu_{\zeta}) = 3 + 4 + 1 + 2 = 10.$$

One may show that the G -function on Δ of ν_{ζ} is

$$G_{\zeta}(w, x, y, z) = 3w + 4x + y + 2z.$$

Similarly, the G -function of $\nu_\eta = \nu_{(1,0)}$ is

$$G_\eta(w, x, y, z) = 2w + x + 4y + 3z.$$

Hence the expected vanishing order of ν_ζ with respect to $-K_M = \mathcal{O}_M(3)$ is

$$S(-K_M, \nu_\zeta) = 3S(\mathcal{O}_M(1), \nu_\zeta) = \frac{3}{\text{vol}(\Delta)} \int_\Delta G d\rho = \frac{48}{5}.$$

We see that the value of the delta function on ν_ζ is $\delta(\nu_\zeta) = \frac{A_M(\nu_\zeta)}{S(-K_M, \nu_\zeta)} = \frac{25}{24} > 1$.

On the other hand, we consider the valuation $\nu_{(0,-1)} = \nu_{-\zeta}$, whose values on X_i are

$$(0, 1, 3, 2, 2, 5, 4, 6).$$

Note that $\nu_{(0,1)}(X_i) + \nu_{-\zeta}(X_i) = 6$. So we have $G_\zeta + G_{-\zeta} = 6$ on Δ . Hence

$$S(\mathcal{O}_M(1), \nu_\zeta) + S(\mathcal{O}_M(1), \nu_{-\zeta}) = 6,$$

hence $S(-K_M, \nu_{-\zeta}) = \frac{42}{5}$. The valuation $\nu_{-\zeta}$ is supported on the affine chart $U_0 = M \cap \{X_0 = 1\} \cong \mathbb{A}^4$ (with coordinates x_1, x_2, x_3, x_4), and is monomial with weight $(1, 3, 2, 2)$. Hence $A_M(\nu_{-\zeta}) = 1 + 3 + 2 + 2 = 8$, and $\delta(\nu_{-\zeta}) = \frac{20}{21} < 1$. Symmetrically, we have $\delta(\nu_\eta) = \delta(\nu_{(1,0)}) = \frac{25}{24}$ and $\delta(\nu_{-\eta}) = \delta(\nu_{(-1,0)}) = \frac{20}{21}$.

For general $\nu_{(a,b)} = \nu_{a\eta+b\zeta}$, the weight is $(4a+6b, 5a+5b, 2a+3b, a+4b, 6a+4b, 4a+b, 3a+2b, 0)$.

If $a, b > 0$, then $\nu_{a\eta+b\zeta}$ has the same center as ν_ζ , and is monomial with weight $(2a+3b, a+4b, 4a+b, 3a+2b)$ with respect to x_2, x_3, x_7, x_8 . Hence $A_M(\nu_{a\eta+b\zeta}) = 10(a+b)$, and the G -function of $\nu_{a\eta+b\zeta}$ on Δ is $G_{a\eta+b\zeta} = aG_\eta + bG_\zeta$. So $S(\nu_{a\eta+b\zeta}) = aS(\nu_\eta) + bS(\nu_\zeta) = \frac{48}{5}(a+b)$, and $\delta(\nu_{a\eta+b\zeta}) = \frac{25}{24}$.

If $a \geq b, b < 0$, we simply assume $b = -1$. Then the minimizer of $\text{wt}_{(a,-1)} = a\eta - \zeta$ is

$$i_{(a,-1)} = \begin{cases} 9 & 4 \leq a, \\ 3 & \frac{2}{3} \leq a \leq 4, \\ 0 & -1 \leq a \leq \frac{2}{3}, \end{cases}$$

In each case, the valuation $\nu_{(a,-1)} = \nu_{a\eta-\zeta}$ is monomial on $U_i, i = i_{(a,-1)}$ with respect to the coordinate chart. More precisely, the weight of the monomial valuation and the coordinate charts are

$$\begin{cases} (2a-3, a-4, 4a-1, 3a-2), & (x_2, x_3, x_7, x_8), & 4 \leq a, \\ (3a-2, 4a-1, a+1, -a+4), & (x_0, x_1, x_2, x_9), & \frac{2}{3} \leq a \leq 4, \\ (a+1, -2a+3, -3a+2, 2a+2), & (x_1, x_2, x_3, x_4), & -1 \leq a \leq \frac{2}{3}. \end{cases}$$

It is not difficult to compute the G -function on Δ

$$G_{a\eta-\zeta} = \begin{cases} aG_\eta - G_\zeta & 4 \leq a, \\ aG_\eta - G_\zeta + (4-a) & \frac{2}{3} \leq a \leq 4, \\ aG_\eta - G_\zeta + (6-4a) & -1 \leq a \leq \frac{2}{3}. \end{cases}$$

Hence the log discrepancies and the S -invariants follow

$$A_M(\nu_{a\eta-\zeta}) = \begin{cases} 10(a-1), \\ 7a+2, \\ -2a+8, \end{cases} \quad S(-K_M; \nu_{a\eta-\zeta}) = \frac{3}{5} \cdot \begin{cases} 16(a-1) & 4 \leq a, \\ 11a+4 & \frac{2}{3} \leq a \leq 4, \\ -4a+14 & -1 \leq a \leq \frac{2}{3}. \end{cases}$$

So the delta function on $\{a \geq -1, b = -1\}$ is

$$\delta(\nu_{a\eta-\zeta}) = \frac{5}{3} \cdot \begin{cases} 5/8 & 4 \leq a, \\ (7a+2)/(11a+4) & \frac{2}{3} \leq a \leq 4, \\ (a-4)/(2a-7) & -1 \leq a \leq \frac{2}{3}. \end{cases}$$

Symmetrically, we have the delta function on $\{a = -1, b \geq -1\}$. We conclude that

$$\delta(\nu_{(a,b)}) = \delta(\nu_{a\eta+b\zeta}) = \frac{5}{3} \cdot \begin{cases} (4a+b)/(7a+2b) & a \leq b \leq -\frac{2}{3}a, \\ (2a-7b)/(4a-11b) & -\frac{2}{3}a \leq b \leq -4a, \\ 5/8 & a \geq -4b \text{ \& } b \geq -4a, \\ (7a-2b)/(11a-4b) & -\frac{2}{3}b \leq a \leq -4b, \\ (a+4b)/(2a+7b) & b \leq a \leq -\frac{2}{3}b. \end{cases}$$

Hence the delta function on $N_{\mathbb{R}}$ is minimized by $\nu_{(-1,-1)}$, whose weight is

$$5 \cdot (0, 0, 1, 1, 0, 1, 1, 2),$$

which is the \mathbf{K} -unstable center ord_{E_S} (up to rescaling by 5) of M_4 found by [Fuj17].

3.4. The delta minimizer of a quintic Del Pezzo fourfold. It was proved in [Fuj17] that $\beta(E_S) < 0$. We have shown that the delta function is minimized by E_S on $N_{\mathbb{R}}$. In this subsection, we will further show that E_S is a minimizer of $\delta(M)$.

By the celebrated theory of [LXZ22, Zhu21], there exists a $\text{Aut}(M)$ -invariant prime divisor E over M computing the delta invariant. Hence $\delta(M, -K_M) = \delta_{C_2}(M, -K_M)$ since $C_2 \subseteq M$ is the unique minimal $\text{Aut}(M)$ -orbit.

Theorem 3.2. *The delta invariant of M_4 is $\delta(M_4) = \frac{25}{27}$ and is minimized by E_S .*

We have shown that

$$\frac{A_M(E_S)}{S(-K_M; E_S)} = \delta(\nu_{(-1,-1)}) = \frac{25}{27}.$$

Note that $-K_M \cong \mathcal{O}_M(3)$. It suffices to show

$$(6) \quad \delta_{C_2}(M, \mathcal{O}_M(1)) \geq \frac{25}{9}.$$

We will prove the inequality using the following strengthened version of the Abban-Zhuang estimate [AZ22, Theorem 3.2], which is stated at the end of the proof of [Zhu21, Lemma 4.9]. See also [MW23, Theorem 4.6].

Let U be a quasi-projective variety and (X, Δ) be a klt pair with a projective morphism $f : X \rightarrow U$. Let F be a plt-type divisor over X with the associated plt-type blowup $\pi : Y \rightarrow X$. We set Δ_Y

to be the strict transform of Δ , and Δ_F be the different of Δ_Y on F . Let V_\bullet be a multi-graded linear series on X , and W_\bullet be the refinement of V_\bullet by F . For any closed subvariety $Z \subseteq U$, We define

$$(7) \quad \delta_Z(X, \Delta; V_\bullet) := \inf_v \frac{A_{X, \Delta}(v)}{S(V_\bullet; v)},$$

where the infimum runs over all the valuations v on X whose center on U contain Z .

Theorem 3.3. *Assume that $Z \subseteq f(F)$. Then*

$$(8) \quad \delta_Z(X, \Delta; V_\bullet) \geq \min \left\{ \frac{A_{X, \Delta}(F)}{S(V_\bullet; F)}, \delta_Z(F, \Delta_F; W_\bullet) \right\}.$$

Remark 3.4. It is clear that $\delta_Z(X, \Delta; V_\bullet) = \inf_{Z'} \delta_{Z'}(X, \Delta; V_\bullet)$ where the infimum runs over all the subvarieties Z' of X dominating Z . The above strengthened version of the Abban-Zhuang estimate says that there is no need to consider all those Z' appearing in the infimum but only the refinement center $C_X(F) \subseteq X$.

Proof of Theorem 3.2. We take refinement of $R_\bullet = R(M, \mathcal{O}(1))$ by E_S , F and C succeedingly, and get multi-graded linear series $W_{\bullet\bullet}^{E_S}, W_{\bullet\bullet}^F$ respectively. Recall that $\tau_{E_S} : E_S \rightarrow \mathbb{P}^3$ is the blowup of a twisted cubic curve C_3 with exceptional divisor F , and $\pi_{E_S} : E_S \rightarrow \mathbb{P}^2$ is a \mathbb{P}^1 -bundle with $\tilde{T} = \pi_{E_S}^{-1}(C_2)$, which is the strict transform of the tangent developable $T \subseteq \mathbb{P}^3$ of C_3 . The divisors F and \tilde{T} in E_S tangent along a curve C of order 2. They are the only G -invariant subvarieties of E_S .

By Theorem 3.3, we have

$$\delta_{C_2}(M; \mathcal{O}_M(1)) \geq \min \left\{ \frac{A_M(E_S)}{S(\mathcal{O}(1); E_S)}, \frac{A_{E_S}(F)}{S(W_{\bullet\bullet}^{E_S}; F)}, \frac{A_F(C)}{S(W_{\bullet\bullet}^F; C)} \right\}.$$

We already have $A_M(E_S) = 2$ and $S(\mathcal{O}(1); E_S) = 18/25$. Let's compute other invariants on the right hand side of the inequality.

We first compute the refinement $W_{\bullet\bullet}^{E_S}$. We make the assumption that $u \in \mathbb{Q}, m \in \mathbb{N}$ such that $mu \in \mathbb{N}$ in the following. Recall that $\tilde{D}_9 \subseteq \tilde{M}$ is the strict transform of $D_9 = \{X_9 = 0\} \subseteq M$ and $\pi^*D_9 = \tilde{D}_9 + 2E_S$. The nef and big divisor $\pi^*\mathcal{O}_M(1) - E_S$ determines the morphism $\tau : \tilde{M} \rightarrow \mathbb{P}^4$ contracting \tilde{D}_9 to a twisted cubic curve C_3 in the hyperplane $H_9 \subseteq \mathbb{P}^4$. We have the following decomposition

$$\begin{aligned} \pi^*\mathcal{O}_M(1) - uE_S &\sim_{\mathbb{Q}} \tilde{D}_9 + (2-u)E_S \\ &= (2-u)(\tilde{D}_9 + E_S) - (1-u)\tilde{D}_9 \sim_{\mathbb{Q}} (2-u)\tau^*\mathcal{O}_{\mathbb{P}^4}(1) - (1-u)\tilde{D}_9, \end{aligned}$$

for $0 \leq u \leq 2$. Hence $\pi^*\mathcal{O}_M(1) - uE_S$ is an ample \mathbb{Q} -divisor for $0 < u < 1$, and the fixed part of $\pi^*\mathcal{O}_M(1) - uE_S$ is $(u-1)\tilde{D}_9$ for $1 \leq u \leq 2$. Recall that $\tau_{E_S} : E_S \rightarrow H_9 = \mathbb{P}^3$ is the blowup of H_9 along a twisted cubic C_3 with exceptional divisor $F = \tilde{D}_9|_{E_S}$. The refinement is defined as the quotient $W_{m, mu}^{E_S} = \mathcal{F}_{E_S}^{mu} V_m / \mathcal{F}_{E_S}^{mu+1} V_m$, which is embedded in $H^0(E_S, m[\pi^*\mathcal{O}_M(1) - uE_S]|_{E_S})$. Using Kawamata-Viehweg vanishing theorem, we conclude that

$$W_{m, mu}^{E_S} = \begin{cases} H^0\left(E_S, m\left[(2-u)\tau_{E_S}^*\mathcal{O}_{H_9}(1) - (1-u)F\right]\right) & 0 \leq u \leq 1, \\ H^0\left(E_S, m(2-u)\tau_{E_S}^*\mathcal{O}_{H_9}(1) + m(u-1)F\right) & 1 \leq u \leq 2. \end{cases}$$

In the second row we mean that the subspace of

$$H^0\left(E_S, m[(2-u)\tau_{E_S}^* \mathcal{O}_{H_9}(1) + (u-1)F]\right),$$

consisting of sections with $\text{ord}_F \geq m(u-1)$. We will simply denote this graded linear series by

$$W_{(1,u)}^{E_S} = \begin{cases} H^0\left(E_S, \mathcal{O}_{H_9}(2-u) - (1-u)F\right) & 0 \leq u \leq 1, \\ H^0\left(E_S, \mathcal{O}_{H_9}(2-u)\right) + (u-1)F & 1 \leq u \leq 2. \end{cases}$$

We could compute $S(W_{\bullet\bullet}^{E_S}; F)$ now. If $0 \leq u \leq 1$, we have naturally

$$(9) \quad \mathcal{F}_F^v W_{(1,u)}^{E_S} = vF + H^0\left(E_S, \mathcal{O}_{H_9}(2-u) - (1-u+v)F\right),$$

which is non-vanishing for $0 \leq v \leq u/2$. If $1 \leq u \leq 2$ and $u-1 \leq v$, the filtration $\mathcal{F}_F^v W_{(1,u)}^{E_S}$ is also formulated as (9) which is non-vanishing for $v \leq u/2$. Finally if $1 \leq u \leq 2$ and $v \leq u-1$, the filtration is trivial:

$$(10) \quad \mathcal{F}_F^v W_{(1,u)}^{E_S} = (u-1)F + H^0\left(E_S, \mathcal{O}_{H_9}(2-u)\right).$$

In a word, there are two polyhedral regions

$$\begin{aligned} \Delta_1^F &= \{(u, v) : 0 \leq v \leq u/2, u-1 \leq v\}, \\ \Delta_2^F &= \{(u, v) : 0 \leq v \leq u-1, u \leq 2\}, \end{aligned}$$

in the (u, v) -plane such that the filtration is formulated as (9) when $(u, v) \in \Delta_1$ and as (10) when $(u, v) \in \Delta_2$. In the former case, we have

$$\begin{aligned} \text{vol}(\mathcal{F}_F^{(v)} W_{(1,u)}^{E_S}) &= \left((2-u)\tau_{E_S}^* \mathcal{O}_{H_9}(1) - (1-u+v)F\right)^3, \\ &= (2-u)^3 \cdot 1 - 3(2-u)^2(1-u+v) \cdot 0 \\ &\quad + 3(2-u)(1-u+v)^2 \cdot (-3) - (1-u+v)^3 \cdot (-10) \\ &= (2-u)^3 - 9(2-u)(1-u+v)^2 + 10(1-u+v)^3, \end{aligned}$$

where the second equality follows from (1). In the latter case

$$\text{vol}(\mathcal{F}_F^{(v)} W_{(1,u)}^{E_S}) = \mathcal{O}_{H_9}(2-u)^3 = (2-u)^3.$$

Recall $\text{vol}(W_{\bullet\bullet}^{E_S}) = \text{vol}(V_\bullet) = \text{vol}(\mathcal{O}_M(1)) = 5$. Hence we have

$$(11) \quad S(W_{\bullet\bullet}^{E_S}; F) = \frac{1}{5/4!} \int_{\Delta_1^F \cup \Delta_2^F} \frac{\text{vol}(\mathcal{F}_F^{(v)} W_{(1,u)}^{E_S})}{3!} dudv = \frac{1}{5} < \frac{9}{25}.$$

Next we take refinement of $W_{\bullet\bullet}^{E_S}$ by F , which is

$$W_{(1,u,v)}^F = H^0\left(F, (1-u+v)h_F + (1+2u-5v)l_F\right),$$

for $(u, v) \in \Delta_1^F = \{(u, v) : 0 \leq v \leq u/2, 1-u+v \geq 0\}$. Hence

$$\mathcal{F}_C^{(w)} W_{(1,u,v)}^F = H^0\left(F, (1-u+v-w)h_F + (1+2u-5v-w)l_F\right),$$

which is non-vanishing for $(u, v, w) \in \Delta^C$ where

$$\Delta^C = \{(u, v, w) : 0 \leq v \leq u/2, 0 \leq w \leq 1 - u + v, w \leq 1 + 2u - 5v\}.$$

Hence

$$(12) \quad S(W_{\bullet\bullet\bullet}^F; C) = \frac{1}{5/4!} \int_{\Delta^C} \frac{\text{vol}(\mathcal{F}_C^{(w)} W_{(1,u,v)}^F)}{2!} dudvdw = \frac{1}{5} < \frac{9}{25}.$$

So $\delta_{C_2}(M; \mathcal{O}_M(1)) \geq \min\{\frac{25}{9}, 5, 5\} = \frac{25}{9}$. We conclude that $\delta(M_4; \mathcal{O}(1)) = \frac{25}{9}$. \square

4. THE QUINTIC DEL PEZZO FIVEFOLDS

We find the delta minimizer of a quintic Del Pezzo fivefold in this section.

4.1. The automorphism group and delta invariant of a quintic Del Pezzo fivefold. Let $M = \text{Gr}(2, V_5) \cap \{X_5 + X_8 = 0\} \subseteq \mathbb{P}^9$, which is a smooth quintic Del Pezzo fivefold. We shall omit X_5 and view M as a subvariety of $\mathbb{P}_{X_0, X_1, X_2, X_3, X_4, X_6, X_7, X_8, X_9}^8$. With the same argument in the previous section, one can show that the endomorphism algebra $\text{aut}(M) \subseteq \mathfrak{sl}_5$ is generated by

$$\begin{pmatrix} -2\lambda & \lambda_1 & \lambda_2 & \lambda_3 & \lambda_4 \\ 0 & a & b & f & g \\ 0 & c & d & g & h \\ 0 & p & q & \lambda - a & -c \\ 0 & q & r & -b & \lambda - d \end{pmatrix}.$$

There are only two orbits on M via the $\text{Aut}(M)$ -action, that is, the closed orbit $W = \{X_4 = X_5 = X_6 = X_7 = X_8 = X_9 = 0\} \cong \mathbb{P}^3 \subseteq \mathbb{P}^9$ and the open orbit $M \setminus W$, see [PVdV99, Proposition 5.3]. Let E_W be the exceptional divisor of the blowup $\pi : \tilde{M} \rightarrow M$ of M along W . Then we have

$$\frac{A_M(E_W)}{S(-K_M; E_W)} = \frac{15}{16} < \frac{45}{46} = \frac{A_M(E_S)}{S(-K_M; E_S)},$$

where the second equality follows from [Fuj17]. For the first one, we state proof now. Using Schubert calculus, one can show that the Chern classes of $N_{W/M}$ are $c_1 = 0, c_2 = \sigma_1^2$, where $\sigma_1 = c_1(M)$. Let $h = c_1(\mathcal{O}_{E_W/W}(1))$. Then we have intersection numbers

$$(13) \quad \begin{pmatrix} \sigma_1^4 & \sigma_1^3 h & \sigma_1^2 h^2 & \sigma_1 h^3 & h^4 \\ 0 & 1 & 0 & -1 & 0 \end{pmatrix}.$$

Hence we have $\text{vol}_{\tilde{M}|E_W}(\mathcal{O}_M(1) - tE_W) = (\sigma_1 + th)^4 = 4t - 4t^3$, and

$$(14) \quad S(\mathcal{O}_M(1); E_W) = \frac{n}{V} \int_0^1 t \cdot (4t - 4t^3) \cdot dt = \frac{8}{15}.$$

We conclude by $-K_M = \mathcal{O}(4)$ and $A_M(E_W) = 2$. Furthermore, we have

Theorem 4.1. *The delta invariant of $M = M_5$ is $\delta(M_5) = \frac{15}{16}$ and is minimized by E_W .*

Proof. Let $W_{\bullet\bullet}^{E_W}$ be the refinement of $R_{\bullet} = R(M, \mathcal{O}(1))$ by E_W . By equivariant K-stability and Theorem 3.3 it suffices to show that

$$S(W_{\bullet\bullet}; Z) \leq \frac{4}{15},$$

for any prime divisor $Z \subseteq E_W$ dominating W . We firstly see that $\mathcal{O}_M(1) - tE_W$ is ample for $0 < t < 1$ and is not big for $t > 1$. Hence $W_{(1,t)}^{E_W} = \mathcal{O}_W(1) + \mathcal{O}_{E_W/W}(t)$ for $0 < t < 1$, which is always ample. So $\text{vol}(W_{(1,t)}) = (\sigma_1 + th)^4 = 4t - 4t^3$. Since $\pi : E_W \rightarrow W$ is a \mathbb{P}^1 -bundle and $Z \subseteq E_W$ dominates W , we have $Z \sim \mathcal{O}_W(a) + \mathcal{O}_{E_W/W}(b)$ for $1 \leq b \leq a$. Then we have

$$\begin{aligned} S(W_{\bullet\bullet}; Z) &= \frac{1}{5/5!} \left(\int_0^{\frac{b}{a}} \int_0^{\frac{t}{b}} \frac{\text{vol}(W_{(1,t)} - sZ)}{4!} ds dt, \right. \\ &\quad \left. + \int_{\frac{b}{a}}^1 \int_0^{\frac{1-t}{a-b}} \frac{\text{vol}(W_{(1,t)} - sZ)}{4!} ds dt \right) = \frac{1}{6a} \leq \frac{1}{6}. \end{aligned}$$

□

We are also interested in the pairs (M, Q) for $Q \in |\mathcal{O}_M(2)|$, in particular for those Q with large symmetries. They are closely related to GM varieties. For $M = M_4$, there is a quadric divisor Q_0 (defined by $X_1^2 - 4X_0X_4$) such that the subgroup $\text{Aut}(M_4, Q_0) \subseteq \text{Aut}(M_4)$ is reductive, see 4. For $M = M_5$, we consider the quadric divisor Q_0 defined by $\{X_0X_2 + X_1X_3 = 0\}$. Then the endomorphism algebra of the pair (M, Q_0) is generated by

$$\begin{pmatrix} -2\lambda & 0 & 0 & 0 & 0 \\ 0 & a & b & 0 & 0 \\ 0 & c & d & 0 & 0 \\ 0 & 0 & 0 & \lambda - a & -c \\ 0 & 0 & 0 & -b & \lambda - d \end{pmatrix}.$$

We use the same notions $Q_x, P_x(t) = \exp(-tQ_x)$ as Section 3.2. Note that M admits a $\mathbb{T} = \mathbb{G}_m^3$ -action generated by P_λ, P_a and P_d , whose actions on $\mathbb{P} \cong \mathbb{P}^8$ are given by

$$\begin{aligned} P_\lambda(t) \cdot [X_i] &= [t^2 X_0, t^2 X_1, t^1 X_2, t^2 X_3, X_4, t^{-1} X_6, t^{-1} X_7, t^{-1} X_8, t^{-2} X_9], \\ P_a(t) \cdot [X_i] &= [t^{-1} X_0, X_1, t X_2, X_3, t^{-1} X_4, t^{-1} X_6, t X_7, X_8, t X_9], \\ P_d(t) \cdot [X_i] &= [X_0, t^{-1} X_1, X_2, t X_3, t^{-1} X_4, t X_6, t^{-1} X_7, X_8, t X_9]. \end{aligned}$$

We denote by η, ς, ζ the weights of the one-parameter subgroups P_λ, P_a, P_d respectively, that is

$$\begin{aligned} \eta &= (2, 2, 1, 1, 0, -1, -1, -1, -2), \\ \varsigma &= (-1, 0, 1, 0, -1, -1, 1, 0, 1), \\ \zeta &= (0, -1, 0, 1, -1, 1, -1, 0, 1). \end{aligned}$$

Let $G = \text{Aut}(M, Q_0)$. Then the generic orbit of the G -action on M is open. Note that W is G -invariant and has G -orbits $Q_{0,W} \setminus (l_{01} \cup l_{23}), l_{01} = \{X_0 = X_1 = 0\}, l_{23} = \{X_2 = X_3 = 0\}$. We denote by $K = \{X_0 = X_1 = X_2 = X_3 = 0\} \subseteq M$, then K is isomorphic to the quadric

threefold $Q^3 = \{Y_4Y_9 + Y_8^2 + Y_6Y_7 = 0\} \subseteq \mathbb{P}_{Y_4, Y_6, Y_7, Y_8, Y_9}^4$. Note that $K \subseteq Q_0$ is G -invariant. The hyperplane sections $H_4^Q = \{Y_4 = 0\}$, $H_9^Q = \{Y_9 = 0\} \subseteq K$ are both cone over conic $q = H_4^Q \cap H_9^Q = \{Y_6Y_7 + Y_8^2 = 0\} \subseteq \mathbb{P}_{Y_6, Y_7, Y_8}^2$ with vertices o_4, o_9 respectively. We have the following orbit decomposition

$$\begin{aligned} M = & \left(M \setminus (Q_0 \cup W) \right) \sqcup \left(Q_0 \setminus (K \cup Q_{0,W}) \right) \sqcup \\ & \left(W \setminus Q_{0,W} \right) \sqcup \left(Q_{0,W} \setminus (l_{01} \cup l_{23}) \right) \sqcup l_{01} \sqcup l_{23} \sqcup \\ & \left(K \setminus (H_4^Q \cup H_9^Q) \right) \sqcup \left(H_4^Q \setminus (o_4 \cup q) \right) \sqcup \left(H_9^Q \setminus (o_9 \cup q) \right) \sqcup q \sqcup o_4 \sqcup o_9. \end{aligned}$$

4.2. Properties of quintic Del Pezzo fivefolds. We make some preparations for computing the delta invariant $\delta(M_5, cQ_0)$ for suitable c . Note that $\text{Aut}(M, Q_0)$ is generated by the one-parameter subgroups with weights $-\frac{1}{5}(2\eta + \varsigma + \zeta)$, $-\frac{1}{5}(\eta - 2\varsigma - 2\zeta)$ and an SL_2 . We consider the toric divisors corresponding to these two one-parameter subgroups, which are E_W and E_S respectively.

The blowup $\tilde{M} \rightarrow M$ naturally embeds in $M \times \mathbb{P}_{Y_4, Y_6, Y_7, Y_8, Y_9}^4$. Let $\tau : \tilde{M} \rightarrow \mathbb{P}^4$ be the natural projection, and we have $\tau^* \mathcal{O}_{\mathbb{P}^4}(1) \sim \pi^* \mathcal{O}_M(1) - E_W$. Hence we get the bi-homogeneous equations defining $E_W \subseteq W \times \mathbb{P}_{Y_4, Y_6, Y_7, Y_8, Y_9}^4$

$$\begin{cases} X_0Y_7 + X_1Y_8 + X_2Y_4 = 0, \\ X_0Y_8 - X_1Y_6 + X_3Y_4 = 0, \\ X_0Y_9 - X_2Y_6 - X_3Y_8 = 0, \\ X_1Y_9 - X_2Y_8 + X_3Y_7 = 0. \end{cases}$$

If we consider it as equations in X_0, X_1, X_2, X_3 , then the determinant of coefficients is $(Y_4Y_9 + Y_8^2 + Y_6Y_7)^2$. Hence the restriction $\tau_W : E_W \rightarrow \mathbb{P}^4$ is a \mathbb{P}^1 -bundle over $Q^3 = \{Y_4Y_9 + Y_8^2 + Y_6Y_7 = 0\} \subseteq \mathbb{P}^4$. One may also consider the blowup of M along K with exceptional divisor E_K . Then E_W and E_K are isomorphic over Q^3 . The inverse image $L_{23} = \pi^{-1}(l_{23})$ is defined by

$$\begin{cases} X_0Y_7 + X_1Y_8 = 0, \\ X_0Y_8 - X_1Y_6 = 0, \end{cases} \subseteq l_{23} \times \mathbb{P}_{Y_4, Y_6, Y_7, Y_8}^3,$$

which is contained in $\tilde{Q}_{0,W} = \pi^{-1}(Q_{0,W})$. Hence $L_{23} \cong \mathbb{F}_2$ and the image of $\tau_{L_{23}}$ is just $H_9^Q \cong \mathbb{P}(1, 1, 2)$. The line $l = \{Y_6 = Y_7 = Y_8 = 0\} \subseteq L_{23}$ is contracted by $\tau_{L_{23}}$ to the vertex o_9 , and it is a minimal G -orbit of E_W . Note that L_{23} is contained in the pull-back $H_9 = \tau^{-1}(H_9^Q) \subseteq W \times \mathbb{P}_{Y_4, Y_6, Y_7, Y_8}^3$, which is defined as

$$\begin{cases} X_0Y_7 + X_1Y_8 + X_2Y_4 = 0, \\ X_0Y_8 - X_1Y_6 + X_3Y_4 = 0, \\ -X_2Y_6 - X_3Y_8 = 0, \\ -X_2Y_8 + X_3Y_7 = 0. \end{cases}$$

Let $\tilde{E}_W \rightarrow E_W$ be the blowup of E_W along l with exceptional divisor $E_l \cong l \times \mathbb{P}^2$. Then the morphism $\tau_W : E_W \rightarrow Q^3$ lifts to $\tilde{\tau}_W : \tilde{E}_W \rightarrow \tilde{Q}^3$, where \tilde{Q}^3 is the blowup of Q^3 at o_9 . And the

restriction of $\tilde{\tau}_W$ to E_l is just the second projection. Let $\tilde{L}_{23}, \tilde{Q}_{0,W} \subseteq \tilde{E}_W$ be the strict transform of L_{23} and $Q_{0,W}$ respectively. Then the curve $C = \tilde{L}_{23} \cap E_l \subseteq E_l$ is defined as

$$\begin{cases} X_0\tilde{Y}_7 + X_1\tilde{Y}_8 = 0, \\ X_0\tilde{Y}_8 - X_1\tilde{Y}_6 = 0, \end{cases} \subseteq l \times \mathbb{P}_{\tilde{Y}_6, \tilde{Y}_7, \tilde{Y}_8}^2 = E_l,$$

which is contained in the surface $R = \tilde{Q}_{0,W} \cap E_l = \{Y_6Y_7 + Y_8^2 = 0\} \subseteq E_l$.

Remark 4.2. The valuation ord_{E_l} (over E_W) is just the restriction of ord_{E_S} (over M) to E_W .

Finally, let $\tilde{\tilde{E}}_W \rightarrow \tilde{E}_W$ be the blowup of \tilde{E}_W along \tilde{L}_{23} with exceptional divisor $E_{\tilde{L}_{23}}$. Then the strict transform $\tilde{\tilde{E}}_l$ of E_l is the blowup of E_l along C with exceptional divisor $E_C = \tilde{\tilde{E}}_l \cap E_{\tilde{L}_{23}}$. We shall denote by $\tilde{\tilde{R}} \subseteq \tilde{\tilde{E}}_l$ the strict transform of R and $\tilde{\tilde{C}} = \tilde{\tilde{R}} \cap E_C$.

4.3. Computing S -invariants over E_W . We will compute the delta invariant of (M, cQ_0) in the last section for suitable c . By G -equivariant K -stability it suffices to compute $\delta_{l_{23}}(M, cQ_0)$. We take refinement of $R_\bullet = R(M, \mathcal{O}(1))$ by E_W, E_l and E_C succeedingly, and get multi-graded linear series $W_{\bullet\bullet\bullet}^{E_W}, W_{\bullet\bullet\bullet}^{E_l}, W_{\bullet\bullet\bullet\bullet}^{E_C}$ respectively.

We denote by $L = \pi^*\mathcal{O}_W(1)$ and $H = \tau^*\mathcal{O}_{Q^3}(1)$, then $L + \mathcal{O}_{E_W/W}(1) = H$. Let $\alpha \subseteq \tilde{L}_{23} \subseteq \tilde{E}_W$ be the strict transform of a fiber of $L_{23} \rightarrow l_{23}$, and $\beta \subseteq \tilde{H}_9$ be a curve intersect l transversally at a point such that it is mapped to a ruling of $H_9^Q = \mathbb{P}(1, 1, 2)$ isomorphically via τ , and mapped to a line in W isomorphically via π . Then $E_l \cdot \alpha = E_l \cdot \beta = 1$ and

$$\begin{pmatrix} L \cdot \alpha & H \cdot \alpha & L \cdot \beta & H \cdot \beta \\ 0 & 1 & 1 & 1 \end{pmatrix}.$$

Recall that $W_{(1,t)}^{E_W} = L + \mathcal{O}_{E_W/W}(t) = (1-t)L + tH$. Then

$$\begin{aligned} \alpha \cdot (W_{(1,t)}^{E_W} - sE_l) &= t - s, \\ \beta \cdot (W_{(1,t)}^{E_W} - sE_l) &= 1 - s. \end{aligned}$$

Hence $W_{(1,t)}^{E_W} - sE_l$ is ample for $0 < s < t$. We denote by $\tilde{L} = L - E_l, \tilde{H} = H - E_l$, where \tilde{H} is semiample but \tilde{L} has base locus \tilde{L}_{23} . If $t < s < 1$, then $W_{(1,t)}^{E_W} - sE_l$ has asymptotic base locus \tilde{L}_{23} . Blowing it up we get $\tilde{\tilde{E}}_W$ and $\tilde{\tilde{L}} = \tilde{L} - E_{\tilde{L}_{23}}$ is semiample. Consequently, we have

$$W_{(1,t)}^{E_W} - sE_l = \begin{cases} H^0\left(\tilde{\tilde{E}}_W, (1-t)L + (t-s)H + s\tilde{H}\right) & 0 \leq s \leq t, \\ H^0\left(\tilde{\tilde{E}}_W, (s-t)\tilde{\tilde{L}} + (1-s)L + t\tilde{H}\right) + (s-t)E_{\tilde{L}_{23}} & t \leq s \leq 1, \\ H^0\left(\tilde{\tilde{E}}_W, (1-t)\tilde{\tilde{L}} + (1+t-s)\tilde{H}\right) + (s-t)E_{\tilde{L}_{23}} + (s-1)\tilde{H}_9 & 1 \leq s \leq 1+t. \end{cases}$$

By (13), we have intersection numbers

$$(15) \begin{pmatrix} L^4 & L^3H & L^2H^2 & LH^3 & H^4 \\ 0 & 1 & 2 & 2 & 0 \end{pmatrix}, \quad \begin{pmatrix} L\tilde{H}^3 & L^2\tilde{H}^2 & L^3\tilde{H} & L\tilde{L}^3 & L^2\tilde{L}^2 & L^3\tilde{L} \\ 1 & 2 & 1 & -1 & 0 & 0 \end{pmatrix}.$$

Note that $N_{\tilde{L}_{23}/\tilde{E}_W} \cong \tilde{L}^{\oplus 2}|_{\tilde{L}_{23}}$ and $[\tilde{L}_{23}] = [\tilde{L}]^2$ in the Chow ring, we have

$$(16) \begin{pmatrix} \tilde{L}^4 & \tilde{L}^3\tilde{H} & \tilde{L}^2\tilde{H}^2 & \tilde{L}\tilde{H}^3 & \tilde{H}^4 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} \tilde{L}^3L & \tilde{L}^2L^2 & \tilde{L}L^3 & \tilde{L}^2\tilde{H}L & \tilde{L}\tilde{H}^2L & \tilde{L}\tilde{H}L^2 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix}.$$

Hence

$$\text{vol}(W_{(1,t)}^{E_W} - sE_l) = \begin{cases} 4(1-t)((1-t)^2t + 3(1-t)t^2 + 2t^3 - s^3) & 0 \leq s \leq t, \\ 4(s-t)(3(1-s)^2t + 3(1-s)t^2 + t^3) + \\ 4(1-s)t((1-s)^2 + 3(1-s)t + t^2) & t \leq s \leq 1, \\ 4(1-t)(1+t-s)^3 & 1 \leq s \leq 1+t, \end{cases}$$

$$(17) \quad S(W_{\bullet\bullet}^{E_W}; E_l) = \frac{1}{5/5!} \int_0^1 \int_0^{1+t} \frac{\text{vol}(W_{(1,t)}^{E_W} - sE_l)}{4!} ds dt = \frac{23}{30}.$$

Next we compute $W_{\bullet\bullet}^{E_l}$ and $S(W_{\bullet\bullet}^{E_l}; E_C)$. Recall that $E_l \cong l \times \mathbb{P}^2$. We denote by $L_E = \text{pr}_1^* \mathcal{O}_l(1)$ and $H_E = \text{pr}_2^* \mathcal{O}_{\mathbb{P}^2}(1)$. Then $L|_{E_l} = L_E, H|_{E_l} = 0$ and $E_l|_{E_l} = -H_E$. Moreover, $\tilde{H}_9|_{E_l} = R$ is a trivial \mathbb{P}^1 bundle over the conic $q \subseteq \mathbb{P}^2$, and $\tilde{L}_{23}|_{E_l} = C \subseteq R$. The strict transform \tilde{E}_l of E_l on \tilde{E}_W is the blowup of E_l along C with exceptional divisor $E_C = E_{\tilde{L}_{23}}|_{\tilde{E}_l}$. We also have $\tilde{H}_9|_{\tilde{E}_l} = \tilde{R}$ and $\tilde{R} \cap E_C = \tilde{C}$. In a word, we have

$$(18) \quad \begin{pmatrix} L & H & E_l & \tilde{H}_9 & \tilde{L}_{23} \\ L_E & 0 & -H_E & R & C \end{pmatrix}, \quad \begin{pmatrix} E_{\tilde{L}_{23}} & \tilde{H}_9 \\ E_C & \tilde{R} \end{pmatrix},$$

where the first rows consist of divisors and subvarieties of \tilde{E}_W and \tilde{E}_W , the second rows consist of their restrictions to E_l or \tilde{E}_l . Since C is the complete intersection of two divisors in $|L_E + H_E|$, we denote by $\tilde{B}_E = L_E + H_E - E_C$. Hence

$$W_{(1,t,s)}^{E_l} = \begin{cases} H^0(E_l, (1-t)L_E + sH_E) & 0 \leq s < t, \\ H^0(\tilde{E}_l, (s-t)\tilde{B}_E + (1-s)L_E + tH_E) + (s-t)E_C & t \leq s \leq 1, \\ H^0(\tilde{E}_l, (1-t)\tilde{B}_E + (1+t-s)H_E) + (s-t)E_C + (s-1)\tilde{R} & 1 \leq s \leq 1+t. \end{cases}$$

A straightforward computation shows that

$$W_{(1,t,s)}^{E_l} - rE_C = \begin{cases} H^0(\tilde{E}_l, r\tilde{B}_E + (1-t-r)L_E + (s-r)H_E) & (r, s, t) \in \Delta_A, \\ H^0(\tilde{E}_l, (1-t)\tilde{B}_E + (1-t+s-2r)H_E) + (-1+t+r)\tilde{R} & (r, s, t) \in \Delta_B, \\ H^0(\tilde{E}_l, (s-t)\tilde{B}_E + (1-s)L_E + tH_E) + (-t+s-r)E_C & (r, s, t) \in \Delta_{C_1}, \\ H^0(\tilde{E}_l, (1-t)\tilde{B}_E + (1+t-s)H_E) \\ + (-t+s-r)E_C + (s-1)\tilde{R} & (r, s, t) \in \Delta_{C_2}, \end{cases}$$

where

$$\begin{aligned}
\Delta_A &= \{0 < t < 1, 0 < s < 1, ((s < t, r > 0) || (s > t, r > s - t)), \\
&\quad ((s + t < 1, r < s) || (s + t > 1, r < 1 - t))\}, \\
\Delta_B &= \{0 < t < 1, -1 + t + s - 2r > 0, ((0 < s < 1, s + t > 1, r > 1 - t) \\
&\quad || (1 < s < 1 + t, r > s - t))\}, \\
\Delta_{C_1} &= \{0 < t < 1, t < s < 1, 0 < r < s - t\}, \\
\Delta_{C_2} &= \{0 < t < 1, 1 < s < 1 + t, 0 < r < s - t\}.
\end{aligned}$$

Since $N_{C/E_l} \cong (L_E + H_E)^{\oplus 2}$, we have intersection numbers

$$(19) \quad \begin{pmatrix} E_C^3 & E_C^2 L_E & E_C^2 H_E \\ -6 & -1 & -2 \end{pmatrix}, \quad \begin{pmatrix} \tilde{B}_E L_E^2 & \tilde{B}_E L_E H_E & \tilde{B}_E H_E^2 & \tilde{B}_E^2 \\ 0 & 1 & 1 & 0 \end{pmatrix}.$$

Hence

$$\text{vol}(W_{(1,t,s)}^{E_l} - rE_C) = \begin{cases} 6r(1-t-r)(s-r) + 3(1-t)(s-r)^2 & (r, s, t) \in \Delta_A, \\ 3(1-t)(1-t+s-2r)^2 & (r, s, t) \in \Delta_B, \\ 6(s-t)(1-s)t + 3(1-t)t^2 & (r, s, t) \in \Delta_{C_1}, \\ 3(1-t)(1+t-s)^2 & (r, s, t) \in \Delta_{C_2}. \end{cases}$$

$$(20) \quad S(W_{\bullet\bullet\bullet}^{E_l}; E_C) = \frac{1}{5/5!} \int_{\Delta_A \cup \Delta_B \cup \Delta_{C_1} \cup \Delta_{C_2}} \frac{\text{vol}(W_{(1,t,s)}^{E_l} - rE_C)}{3!} dr ds dt = \frac{154}{405}.$$

Finally we compute $W_{\bullet\bullet\bullet}^{E_C}$ and $S(W_{\bullet\bullet\bullet}^{E_C}; \tilde{C})$. Recall that $E_C \cong C \times \mathbb{P}^1$ is a trivial bundle. We denote by $l_E = \text{pr}_1^* \mathcal{O}_C(1)$ and $h_E = \text{pr}_2^* \mathcal{O}_{\mathbb{P}^1}(1)$. Then $L_E|_{E_C} = l_E, H_E|_{E_C} = 2l_E, E_C|_{E_C} = -h_E + 3l_E$ and $\tilde{B}_E|_{E_C} = h_E$. We also have $\tilde{R}|_{E_C} = \tilde{C}$. In a word

$$(21) \quad \begin{pmatrix} L_E & H_E & E_C & \tilde{B}_E & \tilde{R} \\ l_E & 2l_E & -h_E + 3l_E & h_E & \tilde{C} \end{pmatrix}.$$

Hence

$$W_{(1,t,s,r)}^{E_C} - u\tilde{C} = \begin{cases} H^0(E_C, (r-u)h_E + (1-t+2s-3r-u)l_E) & (u, r, s, t) \in \Delta_{AA}, \\ H^0(\tilde{E}_l, (1-t)h_E + 2(1-t+s-2r)l_E) \\ + (-1+t+r-u)\tilde{C} & (u, r, s, t) \in \Delta_{BB}, \end{cases}$$

where

$$\begin{aligned}\Delta_{AA} &= \{0 < t < 1, 0 < s < 1, ((s < t, r > 0) \parallel (s > t, r > s - t)), \\ &\quad ((s + t < 1, r < s) \parallel (s + t > 1, r < 1 - t)), 1 - t + 2s - 2r > 0, 0 < u < r\} \\ &\cup \{0 < t < 1, -1 + t + s - 2r > 0, ((0 < s < 1, s + t > 1, r > 1 - t) \\ &\quad \parallel (1 < s < 1 + t, r > s - t)), 1 - t + 2s - 2r > 0, -1 + t + r < u < r\}, \\ \Delta_{BB} &= \{0 < t < 1, -1 + t + s - 2r > 0, ((0 < s < 1, s + t > 1, r > 1 - t) \\ &\quad \parallel (1 < s < 1 + t, r > s - t)), 1 - t + 2s - 2r > 0, 0 < u < -1 + t + r\}.\end{aligned}$$

Hence

$$(22) \quad \begin{aligned}\text{vol}(W_{(1,t,s,r)}^{EC} - u\tilde{C}) &= \begin{cases} 2(r-u)(1-t+2s-3r-u) & (u, r, s, t) \in \Delta_{AA}, \\ 4(1-t)(1-t+s-2r) & (u, r, s, t) \in \Delta_{BB}, \end{cases} \\ S(W_{\bullet\bullet\bullet\bullet}^{EC}; \tilde{C}) &= \frac{1}{5/5!} \int_{\Delta_{AA} \cup \Delta_{BB}} \frac{\text{vol}(W_{(1,t,s,r)}^{EC} - u\tilde{C})}{2!} du dr ds dt = \frac{221}{2430}.\end{aligned}$$

5. K-STABILITY OF GENERAL SPECIAL GUSHEL-MUKAI MANIFOLDS

Let X be a GM n -fold, and $M \subseteq \mathbb{P}^{n+3}$ be the quintic Del Pezzo n -fold. When X is special, there is a double cover $p : X \rightarrow M$ branched along a smooth quadric divisor Q which is an ordinary GM $(n-1)$ -fold. We have $p^*(K_M + \frac{1}{2}Q) = K_X$. By [LZ22a, Theorem 1.2], we see that X is K-semistable if and only if the log Fano pair $(M, \frac{1}{2}Q)$ is K-semistable.

We first recall some basic results about the projective cone over a log Fano pair. Let (V, Δ_V) be a $(n-1)$ -dimensional log Fano pair such that $L = -\frac{1}{r}(K_V + \Delta_V)$ is an ample Cartier divisor for some rational number $r \in (0, n]$. We define the projective cone over V with polarization L by

$$Y = \bar{\mathcal{C}}(V, L) := \text{Proj}\left(\bigoplus_{m \geq 0} \bigoplus_{0 \leq \lambda \leq m} H^0(V, (m-\lambda)L)s^\lambda\right),$$

which is the union of the affine cone $\mathcal{C} = \text{Spec}(\bigoplus_{m \geq 0} H^0(V, mL))$ and a divisor $V_\infty \cong V$ at infinity. Let Δ_Y be closure of $\Delta_V \times \mathbb{C}^*$ in Y . Then $-(K_Y + \Delta_Y)$ is Cartier and

$$-(K_Y + \Delta_Y) \sim_{\mathbb{Q}} (1+r)V_\infty.$$

By [LX20, Proposition 5.3] or Theorem 6.2, we have

Proposition 5.1. *If (V, Δ_V) is K-semistable, then $(Y, \Delta_Y + (1 - \frac{r}{n})V_\infty)$ is K-semistable.*

We will use the following result about degeneration to normal cone. See for example [LZ22b, Lemma 2.12] and [ZZ22, Lemma 5.3].

Proposition 5.2. *Let X be a Fano variety of dimension n , and V be a prime divisor on X such that $-K_X \sim_{\mathbb{Q}} (1+r)V$ with $0 < r \leq n$. Then (X, V) specially degenerates to (Y, V_∞) , where $Y = \bar{\mathcal{C}}(V, H)$ and $H = V|_V$. In particular, if V is K-semistable, then $(X, (1 - \frac{r}{n})V)$ is K-semistable.*

We are ready to state the main theorem.

Theorem 5.3. *General special GM-manifolds are K-stable.*

Proof. It was proved by [AZ22] that smooth GM threefolds are all K-stable. We need to prove that there exists a K-stable special GM n -fold for $n = 4, 5, 6$. Then by openness of K-stability we are done, since the automorphism groups of GM manifolds are discrete and the parameter space of special GM n -folds is a subvariety of the parameter space of GM n -folds.

We first prove that general GM n -folds are K-semistable for $n = 4, 5, 6$. It suffices to find smooth quadric divisor $Q_{n-1} \subseteq M_n$ such that $(M_n, \frac{1}{2}Q_{n-1})$ is K-semistable.

For $n = 4$, note that Q is a GM threefold, which is K-stable by [AZ23, Theorem C]. Since $-K_M \sim_{\mathbb{Q}} \frac{3}{2}Q$, we have $r = \frac{1}{2}$ and $1 - \frac{r}{n} = \frac{7}{8} =: c_{\max}$. By Proposition 5.2 we conclude that $(M, \frac{7}{8}Q)$ is K-semistable. On the other hand, we consider a smooth hyperplane section H of M . Then $H \cong M_3$ which is K-polystable by [ACC⁺23, Example 3.4.1]. With the same method, since $-K_M = 3H$, we have $r = 2, 1 - \frac{r}{n} = \frac{1}{2}$. Hence $(M, \frac{1}{2}H)$ is K-semistable. One can construct a family that deform $(X, \frac{1}{2}H)$ to $(X, \frac{1}{4}Q)$ for some smooth quadric divisor Q of M . By the openness of K-semistability, we conclude that there exists smooth quadric divisor Q_0 on M such that $(X, \frac{1}{4}Q_0)$ is K-semistable. Finally, by interpolation of K-stability ([ADL19, Proposition 2.13]), we see that (M, cQ_0) is K-semistable for any $\frac{1}{4} \leq c \leq \frac{7}{8}$. In particular, $(M, \frac{1}{2}Q_0)$ is K-semistable.

For $n = 5$, we consider the pair $(M = M_5, \frac{1}{2}(H + H'))$, where $M = M_5, H, H'$ are two distinct smooth hyperplane sections of M with smooth intersection $I = H \cap H'$. So H and H' are M_4 and $I \cong M_3$. By Proposition 5.2, this pair can specially degenerate to $(\overline{\mathcal{C}}(H, \mathcal{O}_H(1)), \frac{1}{2}(H_{\infty} + \mathcal{E}))$, where $\mathcal{E} = \overline{\mathcal{C}}(I, \mathcal{O}_I(1))$. Applying Proposition 5.1 to $(H, \frac{1}{2}I)$ and $r = \frac{5}{2}$, since $\mathcal{O}_H(1) \sim_{\mathbb{Q}} -\frac{1}{r}(K_H + \frac{1}{2}I)$, we see that $(\overline{\mathcal{C}}(H, \mathcal{O}_H(1)), \frac{1}{2}(H_{\infty} + \mathcal{E}))$ is K-semistable. Then by the openness of K-semistability, $(M, \frac{1}{2}(H + H'))$ is K-semistable. Similar to the proof of $n = 4$ case, one can deform $(M, \frac{1}{2}(H + H'))$ to some $(M, \frac{1}{2}Q)$ and we get the K-semistability of $(M, \frac{1}{2}Q_0)$ for some smooth quadric divisor Q_0 .

We also consider the degeneration induced by some $Q \subseteq M_5$, which is an ordinary GM fourfold. This will be useful in the proof of K-stability. In the $n = 4$ case, we already have a K-semistable special GM fourfold. By openness of K-semistability, there exists a K-semistable ordinary GM fourfold Q . Since $-K_M \sim 2Q$ we have $r = 1, c_{\max} = \frac{4}{5}$. Then by Proposition 5.2 we see that $(M, \frac{4}{5}Q)$ is K-semistable. By interpolation we have (M, cQ) is K-semistable for any $\frac{1}{2} \leq c \leq \frac{4}{5}$.

For $n = 6$, we already have a K-polystable base $M = M_6 = \text{Gr}(2, 5)$. The same argument in the previous paragraph shows that $(M, \frac{3}{4}Q)$ is K-semistable for some Q . By interpolation we see that (M, cQ) is K-semistable for any $0 \leq c \leq \frac{3}{4}$. In particular, $(M, \frac{1}{2}Q)$ is K-semistable.

Next, we prove that general GM n -folds are K-stable for $n = 4, 5, 6$. We prove by induction on n using Theorem 6.2 in the next section.

Since any ordinary GM threefold $Q_3 \subseteq M_4$ is K-stable, we see that (M_4, cQ_3) is K-stable for any $0 \ll c < \frac{7}{8}$ by Theorem 6.2. By interpolation, (M_4, cQ_3) is K-stable for any $\frac{1}{4} < c < \frac{7}{8}$ (this holds

for $\frac{1}{9} < c < \frac{7}{8}$, see Theorem 6.5). In particular $(M_4, \frac{1}{2}Q_3)$ is K-stable, and we get a K-stable special GM fourfold. Then the openness of K-stability gives a K-stable ordinary GM fourfold $Q_4 \subseteq M_5$. By Theorem 6.2 again, (M_5, cQ_4) is K-stable for all $0 \ll c < \frac{4}{5}$, hence for all $\frac{1}{8} < c < \frac{4}{5}$ by interpolation and Theorem 6.6. Also by the openness of K-stability, we get a K-stable ordinary GM fivefold $Q_5 \subseteq M_6$. With the same argument we conclude that (M_6, cQ_5) is K-stable for all $0 < c < \frac{3}{4}$. In particular, we get a K-stable special GM sixfold. \square

6. THE K-SEMISTABLE DOMAINS OF PAIRS (M, cQ)

We consider the wall-crossing problem of the K-moduli spaces of the log Fano pairs (M, cQ) in this section, where M is a quintic Del Pezzo fourfold or fivefold and Q is an ordinary GM threefold or fourfold respectively. Note that the general theory of K-moduli wall crossing in the proportional case was established in [ADL19, Zho23].

For $M = M_4$, one may compute delta for the divisorial valuations ord_{E_S} and ord_{Q_3} . A straightforward computation shows that the smoothness of Q_3 implies that $S \not\subseteq Q_3$. So we have

$$\begin{aligned} \frac{A_{M_4, cQ_3}(E_S)}{S(-K_{M_4} - cQ_3; E_S)} &= \frac{25}{9(3-2c)}, \\ \frac{A_{M_4, cQ_3}(Q_3)}{S(-K_{M_4} - cQ_3; Q_3)} &= \frac{10(1-c)}{3-2c}. \end{aligned}$$

Hence (M_4, cQ_3) is K-semistable only if $\frac{1}{9} \leq c \leq \frac{7}{8}$.

For $M = M_5$, similarly we have for ord_{E_W} and ord_{Q_4} ,

$$\begin{aligned} \frac{A_{M_5, cQ_4}(E_W)}{S(-K_{M_5} - cQ_4; E_W)} &= \frac{15}{8(2-c)}, \\ \frac{A_{M_5, cQ_4}(Q_4)}{S(-K_{M_5} - cQ_4; Q_4)} &= \frac{6(1-c)}{2-c}. \end{aligned}$$

Hence (M_5, cQ_4) is K-semistable only if $\frac{1}{8} \leq c \leq \frac{4}{5}$. Moreover, we have the following theorem.

Theorem 6.1. *For general $Q_3 \in |\mathcal{O}_{M_4}(2)|$ and $Q_4 \in |\mathcal{O}_{M_5}(2)|$, we have*

$$\text{Kss}(M_4, Q_3) = \left[\frac{1}{9}, \frac{7}{8} \right], \quad \text{Kss}(M_5, Q_4) = \left[\frac{1}{8}, \frac{4}{5} \right].$$

6.1. The last wall. We show that the pair (M, cQ) is K-stable for those c sufficiently close to the last wall by the following theorem, which is a minor strengthening of the cone construction.

Theorem 6.2. *Let (X, Δ) be a log Fano pair of dimension n , $V \subseteq X$ be a prime divisor not contained in $\text{Supp}(\Delta)$ such that $(X, \Delta + V)$ is lc at the generic point of V . We denote by Δ_V the different of Δ on V . Assume that $-K_X - \Delta \sim_{\mathbb{Q}} (1+r)V$ for some rational number $0 < r < n$. Then we have*

(a) *If (V, Δ_V) is K-semistable, then $(X, \Delta + (1 - \frac{r}{n})V)$ is K-semistable;*

(b) *If (V, Δ_V) is K-stable, then $(X, \Delta + cV)$ is K-stable for any $c \in \mathbb{Q}_{\geq 0}$ such that*

$$(23) \quad \max \left\{ (r+1)(1 - \delta(X, \Delta)), \left(1 - \frac{r}{n}\right) - \frac{r}{n}(n+1)(\delta(V, \Delta_V) - 1) \right\} < c < 1 - \frac{r}{n}.$$

Remark 6.3. The theorem says that if (V, Δ_V) is K-stable, then $(X, \Delta + cV)$ is K-stable for $c = 1 - \frac{r}{n} - \varepsilon$ for any $0 < \varepsilon \ll 1$.

Proof. For any $c \in \mathbb{Q}_{\geq 0}$ such that the pair $(X, \Delta + cV)$ is log Fano, we denote by V_\bullet the graded linear series of $-K_X - \Delta - cV$ and by W_\bullet^V the refinement of V_\bullet by V . Then for any $x \in V$, [AZ22, Theorem 3.2] implies that

$$(24) \quad \delta_x(X, \Delta + cV) \geq \min \left\{ \frac{(1-c)(n+1)}{r+1-c}, \delta_x(V, \Delta_V; W_\bullet^V) \right\}.$$

Since $V \sim_{\mathbb{Q}} \frac{1}{r+1}(-K_X - \Delta)$, the multi-graded linear series W_\bullet^V is almost complete in the sense of [AZ22, Definition 2.16]. It is clear that W_\bullet^V has no fixed part. Hence by [AZ22, (3.1)], we have

$$\delta_x(V, \Delta_V; W_\bullet^V) = \delta_x(V, \Delta_V; c_1(W_\bullet^V)) = \frac{r(n+1)}{n(r+1-c)} \delta_x(V, \Delta_V),$$

where

$$\begin{aligned} c_1(W_\bullet^V) &= \left((-K_X - \Delta - cV) - S(-K_X - \Delta - cV; V)V \right)|_V \\ &= \frac{n(r+1-c)}{r(n+1)}(-K_V - \Delta_V). \end{aligned}$$

Thus the right hand side of (24) is greater than 1 if

$$\left(1 - \frac{r}{n}\right) - \frac{r}{n}(n+1)(\delta(V, \Delta_V) - 1) < c < 1 - \frac{r}{n}.$$

For any $x \notin V$, we have directly

$$\delta_x(X, \Delta + cV; -K_X - \Delta - cV) = \frac{r+1}{r+1-c} \delta_x(X, \Delta; -K_X - \Delta),$$

which is greater than 1 if $c > (r+1)(1 - \delta(X, \Delta))$. \square

Remark 6.4. Apply the theorem to (M_4, cQ) . We have shown that $\delta(M_4) = \frac{25}{27}$. Hence $(r+1)(1 - \delta(M_4)) = \frac{1}{9} < \frac{7}{8}$. By [AZ23, Theorem 5.12], we know that the smooth GM threefold Q is K-stable. We conclude that (M_4, cQ) is K-stable for any $0 \ll c < \frac{7}{8}$.

With the same argument in the proof of Theorem 5.3, we get a K-stable ordinary GM fourfold $Q \subseteq M_5$. Apply the theorem to (M_5, cQ) . Since $\delta(M_5) = \frac{15}{16}$, we have $(r+1)(1 - \delta(M_5)) = \frac{1}{8} < \frac{4}{5}$. Hence (M_5, cQ) is K-stable for any $0 \ll c < \frac{4}{5}$.

6.2. The first wall. For $M = M_4$. If $c < \frac{1}{9}$, then (M, cQ) is destabilized by E_S . We consider the pair $(M, \frac{1}{9}Q)$ with $Q \cap S = C_2$. The product test configuration induced by $-\frac{1}{5}(\eta + \zeta)$ (see Section 3.2) degenerating Q to the cone $Q_0 = \{X_1^2 - 4X_0X_4 = 0\} \subseteq M$. We have

Theorem 6.5. *The pair $(M_4, \frac{1}{9}Q_0)$ is K-polystable.*

Proof. We first prove that the pair is K-semistable. Note that $G := \text{Aut}(M, Q_0) \cong \text{GL}_2$ has one dimensional center \mathbb{T} whose action is generated by $-\frac{1}{5}(\eta + \zeta)$. The curve C_2 is one of the two

minimal orbits of the G -action on M (the other is the closed point $[0, 0, 0, 0, 0, 0, 0, 1]$). It suffices to show that $\delta_{C_2}(M, \frac{1}{9}Q_0) \geq 1$. We follow the calculation in the proof of Theorem 3.2.

Recall that $\tau_{E_S} : E_S \rightarrow \mathbb{P}^3$ is the blowup of a twisted cubic curve C_3 with exceptional divisor F , and $\pi_{E_S} : E_S \rightarrow \mathbb{P}^2$ is a \mathbb{P}^1 -bundle with $\tilde{T} = \pi_{E_S}^{-1}(C_2)$, which is the strict transform of the tangent developable $T \subseteq \mathbb{P}^3$ of C_3 . The divisors F and \tilde{T} in E_S tangent along a curve C of order 2. They are the only G -invariant subvarieties of E_S . Also note that $\pi_*^{-1}(Q_0) \cap E_S = \tilde{T}$ intersecting transversally.

We take refinement of $R_\bullet = R(M, \mathcal{O}(1))$ by E_S, F and C succeedingly. By (11) and (12),

$$S(W_{\bullet\bullet}^{E_S}; F) = \frac{1}{5}, \quad S(W_{\bullet\bullet}^F; C) = \frac{1}{5}.$$

The restriction of $\frac{1}{9}\tilde{T}$ on F is $\frac{2}{9}C$. Then

$$\frac{A_{E_S, \frac{1}{9}\tilde{T}}(F)}{S(W_{\bullet\bullet}^{E_S}; F)} = 5, \quad \frac{A_{F, \frac{2}{9}C}(C)}{S(W_{\bullet\bullet}^F; C)} = \frac{35}{9}.$$

Note that $-K_M - \frac{1}{9}Q_0 = \frac{25}{9}\mathcal{O}(1)$. By Theorem 3.3, we have

$$\begin{aligned} \delta_{C_2}(M, \frac{1}{9}Q_0) &\geq \frac{9}{25} \min \left\{ \frac{A_{M, \frac{1}{9}Q_0}(E_S)}{S(\mathcal{O}(1); E_S)}, \frac{A_{E_S, \frac{1}{9}\tilde{T}}(F)}{S(W_{\bullet\bullet}^{E_S}; F)}, \frac{A_{F, \frac{2}{9}C}(C)}{S(W_{\bullet\bullet}^F; C)} \right\} \\ &= \min \left\{ 1, \frac{9}{5}, \frac{7}{5} \right\} = 1. \end{aligned}$$

Hence $(M, \frac{1}{9}Q_0)$ is \mathbf{K} -semistable.

Next, we prove that $(M, \frac{1}{9}Q_0)$ is \mathbf{K} -polystable. Let $\mathbb{T} \subseteq G$ be the one-parameter subgroup generated by $-\frac{1}{5}(\eta + \zeta)$, and N the co-weight lattice of the \mathbb{T} -action. Then E_S is a toric divisor of \mathbb{T} , and we have naturally a \mathbb{T} -equivariant surjective rational map $f : \tilde{M} \dashrightarrow E_S$, which makes E_S a quotient of \tilde{M} by the \mathbb{T} -action. Hence $\text{Val}_M^{\mathbb{T}, \circ} \cong \text{Val}_{E_S}^{\circ} \times N_{\mathbb{R}}$. If $(M, \frac{1}{9}Q_0)$ is not \mathbf{K} -polystable, then by [Zhu21, Corollary 4.11], it is not G -equivariantly \mathbf{K} -polystable. Hence by [LX14, Theorem 7] there exists a G -equivariant non-product type special TC $(\mathcal{X}, \Delta, \mathcal{L})$ of $(M, \frac{1}{9}Q_0)$ with $\text{Fut}(\mathcal{X}, \Delta, \mathcal{L}) = 0$. Then $v = \text{ord}_{\mathcal{X}_0}|_{\mathcal{X}_1}$ is a G -invariant valuation on M not of the form wt_{ξ} for any $\xi \in N_{\mathbb{R}}$ with $\beta_{M, \frac{1}{9}Q_0}(v) = 0$, which is equivalent to $A_{M, \frac{1}{9}Q_0}(v)/S(\mathcal{O}(1); v) = \frac{25}{9}$. In particular, v is \mathbb{T} -invariant. Hence $v = v_{0, \xi}$ for some G -invariant valuation v_0 on E_S and $\xi \in N_{\mathbb{R}}$. Note that v_0 is also a valuation on M via the surjective rational map $f : \tilde{M} \dashrightarrow E_S$. Hence $A_{M, \frac{1}{9}Q_0}(v_0) = A_{E_S, \frac{1}{9}\tilde{T}}(v_0)$. Note that $R_\bullet = R(M, \mathcal{O}(1))$ admits a weight decomposition $R_m = \bigoplus_{\alpha \in \mathbb{Z}} R_{m, \alpha}$ with respect to the \mathbb{T} -action. Since E_S is a toric divisor of \mathbb{T} , we have a canonical \mathbb{T} -equivariant isomorphism of \mathbb{N}^2 -graded linear series $R_{\bullet\bullet} \cong W_{\bullet\bullet}^{E_S}$. Hence the filtration induced by v_0 on $R_{\bullet\bullet}, W_{\bullet\bullet}^{E_S}$ are the same via this isomorphism. So $S(\mathcal{O}(1); v_0) = S(W_{\bullet\bullet}^{E_S}; v_0)$ and

$$\frac{25}{9} = \frac{A_{M, \frac{1}{9}Q_0}(v)}{S(\mathcal{O}(1); v)} \geq \frac{A_{M, \frac{1}{9}Q_0}(v_0)}{S(\mathcal{O}(1); v_0)} = \frac{A_{E_S, \frac{1}{9}\tilde{T}}(v_0)}{S(W_{\bullet\bullet}^{E_S}; v_0)} \geq \delta_C(E_S, \frac{1}{9}\tilde{T}; W_{\bullet\bullet}^{E_S}) \geq \frac{25}{9} \min \left\{ \frac{9}{5}, \frac{7}{5} \right\},$$

where the first inequality follows from the definition of reduced delta invariant [XZ20, Remark A.2] and the second inequality follows from the G -invariance of v_0 . We get a contradiction. Hence $(M, \frac{1}{9}Q_0)$ is \mathbf{K} -polystable. \square

For $M = M_5$. If $c < \frac{1}{8}$, then (M, cQ) is destabilized by E_W . Let $Q_0 = \{X_0X_2 + X_1X_3 = 0\} \subseteq M$. We consider those Q with $Q_W = Q_{0,W}$. Then the product test configuration of M induced by $-\frac{1}{5}(2\eta + \varsigma + \zeta)$ (see Section 4.1) degenerating Q to the cone Q_0 . We have

Theorem 6.6. *The pair $(M_5, \frac{1}{8}Q_0)$ is K-polystable.*

Proof. We take refinements of $R_\bullet = R(M, \mathcal{O}(1))$ by E_W , E_l and E_C succeedingly (see Section 4.2), and get multi-graded linear series $W_{\bullet\bullet}^{E_W}$, $W_{\bullet\bullet\bullet}^{E_l}$, $W_{\bullet\bullet\bullet\bullet}^{E_C}$ respectively. Note that $-K_M - \frac{1}{8}Q_0 \sim_{\mathbb{Q}} \frac{15}{4}\mathcal{O}(1)$. By Theorem 3.3, (14), (17), (20) and (22), we have

$$\begin{aligned} \delta_{l_{23}}(M, \frac{1}{8}Q_0) &\geq \frac{4}{15} \min \left\{ \frac{A_{M, \frac{1}{8}Q_0}(E_W)}{S(\mathcal{O}(1); E_W)}, \frac{A_{E_W, \frac{1}{8}Q_{0,W}}(E_l)}{S(W_{\bullet\bullet}^{E_W}; E_l)}, \frac{A_{E_l, \frac{1}{8}R}(E_C)}{S(W_{\bullet\bullet\bullet}^{E_l}; E_C)}, \frac{A_{E_C, \frac{1}{8}\tilde{C}}(\tilde{C})}{S(W_{\bullet\bullet\bullet\bullet}^{E_C}; \tilde{C})} \right\} \\ &= \min \left\{ 1, 1, \frac{405}{308}, \frac{567}{221} \right\} = 1. \end{aligned}$$

Hence $(M, \frac{1}{8}Q_0)$ is K-semistable.

We prove that $(M, \frac{1}{8}Q_0)$ is K-polystable with the same argument in the proof of Theorem 6.5. Let $G = \text{Aut}(M, Q_0)$ and $\mathbb{T}_1, \mathbb{T}_2 \subseteq G$ be the one-parameter subgroups generated by $-\frac{1}{5}(2\eta + \varsigma + \zeta)$ and $-\frac{1}{5}(\eta - 2\varsigma - 2\zeta)$ respectively, and N the co-weight lattice of the $\mathbb{T} = \mathbb{T}_1 \times \mathbb{T}_2$ -action. Then E_W is a toric divisor of \mathbb{T}_1 , and we have naturally a \mathbb{T}_1 -equivariant surjective rational map $f_1 : \tilde{M} \dashrightarrow E_W$, which makes E_W a quotient of \tilde{M} by the \mathbb{T}_1 -action. The one-parameter subgroup \mathbb{T}_2 acts on E_W faithfully and E_l is a toric divisor of such an action. Hence we have a \mathbb{T}_2 -equivariant surjective rational map $f_2 : \tilde{E}_W \dashrightarrow E_l$, which makes E_l a quotient of \tilde{E}_W by the \mathbb{T}_2 -action. Hence $\text{Val}_M^{\mathbb{T}, \circ} \cong \text{Val}_{E_l}^{\circ} \times N_{\mathbb{R}}$. Note that \mathbb{T} is the center of G and $G \cong \mathbb{T} \times \text{SL}_2$. The \mathbb{T} -action on E_l is trivial, hence the G -action on E_l descends to the SL_2 -action. And $C \subseteq E_l$ is the only minimal orbit of the SL_2 -action.

If $(M, \frac{1}{8}Q_0)$ is not K-polystable, then following the same argument in the previous proof, there exists a non-trivial G -invariant valuation v_0 on E_l and $\xi \in N_{\mathbb{R}}$ such that $A_{M, \frac{1}{8}Q_0}(v_{0, \xi})/S(\mathcal{O}(1); v_{0, \xi}) = \frac{15}{4}$. On the other hand, we have a canonical \mathbb{T} -equivariant isomorphism of \mathbb{N}^3 -graded linear series $R_{\bullet\bullet\bullet} \cong W_{\bullet\bullet\bullet}^{E_l}$, and the filtration induced by v_0 on $R_{\bullet\bullet\bullet}$, $W_{\bullet\bullet\bullet}^{E_l}$ are the same via this isomorphism. So

$$\frac{15}{4} = \frac{A_{M, \frac{1}{8}Q_0}(v_{0, \xi})}{S(\mathcal{O}(1); v_{0, \xi})} \geq \frac{A_{M, \frac{1}{8}Q_0}(v_0)}{S(\mathcal{O}(1); v_0)} = \frac{A_{E_l, \frac{1}{8}R}(v_0)}{S(W_{\bullet\bullet\bullet}^{E_l}; v_0)} \geq \delta_C(E_l, \frac{1}{8}R; W_{\bullet\bullet\bullet}^{E_l}) \geq \frac{15}{4} \min \left\{ \frac{405}{308}, \frac{567}{221} \right\},$$

which is a contradiction. Hence $(M, \frac{1}{8}Q_0)$ is K-polystable. \square

Proof of Theorem 6.1. It follows directly from Remark 6.4, Theorem 6.5 and 6.6. \square

Remark 6.7. The wall crossing for K-moduli spaces of (M_4, cQ_3) and (M_5, cQ_4) are closely related to K-moduli spaces of Gushel-Mukai threefolds and fourfolds. In fact, the K-moduli space of GM threefolds (resp. GM fourfolds) is isomorphic to the K-moduli space of $(M_4, \frac{7}{8}Q_3)$ (resp. $(M_5, \frac{4}{5}Q_4)$) using degeneration to the normal cone. Moreover, if we restrict to $c = \frac{1}{2}$, then the corresponding K-moduli spaces of pairs are isomorphic to K-moduli spaces of special GM fourfolds and fivefolds.

7. KÄHLER-RICCI SOLITONS ON QUINTIC DEL PEZZO FOURFOLDS AND FIVEFOLDS

In this section, we show that M_4 and M_5 both admit Kähler-Ricci solitons following the argument in [MW23], which answers a question asked by [Ino19, Del22]. By the work of [HL23, BLXZ23] we know that a Fano manifold X admits a Kähler-Ricci soliton if and only if (X, ξ_0) is weighted K-polystable for some holomorphic vector field ξ_0 on X (which is called a *soliton candidate*, see for example [MW23, Section 3.1]).

For Del Pezzo fourfolds M_4 , consider the \mathbb{G}_m -action induced by $-\frac{1}{5}(\eta + \zeta)$ (see Section 3.2) with toric divisor E_S . Since $A_{M_4}(E_S) = 2$ and $|\mathcal{O}(1) - uE_S|_{\mathbb{Q}} \neq \emptyset$ for $u \in [0, 2]$, the moment polytope of this \mathbb{G}_m -action on $-K_{M_4} = \mathcal{O}(3)$ is $\mathbf{P} = [-2, 4] \ni 3u - 2 =: u'$. Recall that the soliton candidate ξ_0 with respect to this \mathbb{G}_m -action is the minimizer of the \mathbf{H} -functional

$$\mathbf{H}(\xi) = \log\left(\int_{\mathbf{P}} e^{-u'\xi} \cdot \mathrm{DH}_{\mathbf{P}}(du')\right),$$

where $\xi \in \mathbb{N}(\mathbb{G}_m)_{\mathbb{R}} \cong \mathbb{R}$ and $\mathrm{DH}_{\mathbf{P}}(du') = 3^4 \cdot \mathrm{vol}(W_{(1,u)}^{E_S})du$. We can numerically solve

$$\xi_0 \approx 0.16838665311714196.$$

Theorem 7.1. *The pair (M_4, ξ_0) is weighted K-polystable.*

Proof. By definition of ξ_0 , we have

$$\int_0^2 (3u - 2)e^{-(3u-2)\xi_0} \cdot \mathrm{vol}(W_{(1,u)}^{E_S})du = 0.$$

Recall that the g -weighted volume is defined by

$$\begin{aligned} \frac{\mathrm{vol}^g(\mathcal{F}_{E_S}^{(u_1)} R_{\bullet})}{4!} &:= \int_{u \geq u_1} e^{-(3u-2)\xi_0} \frac{\mathrm{vol}(W_{(1,u)}^{E_S})}{3!} du, \\ \mathbf{v}^g &:= \frac{\mathrm{vol}^g(R_{\bullet})}{4!} = \int_0^2 e^{-(3u-2)\xi_0} \frac{\mathrm{vol}(W_{(1,u)})}{3!} du \\ &\approx 0.2055698662861948. \end{aligned}$$

Let $R_{\bullet} = R(M_4, \mathcal{O}(1))$. Using integration by part, we have

$$S^g(R_{\bullet}; E_S) := \frac{1}{\mathbf{v}^g} \int_0^2 \frac{\mathrm{vol}^g(\mathcal{F}_{E_S}^{(u)} R_{\bullet})}{4!} du = \frac{1}{\mathbf{v}^g} \int_0^2 u e^{-(3u-2)\xi_0} \frac{\mathrm{vol}(W_{(1,u)}^{E_S})}{3!} du = \frac{2}{3} = \frac{1}{3} A_{M_4}(E_S).$$

Moreover, we have

$$\begin{aligned} S^g(W_{\bullet\bullet}^{E_S}; F) &= \frac{1}{\mathbf{v}^g} \int_{\Delta_1^F \cup \Delta_2^F} e^{-(3u-2)\xi_0} \cdot \frac{\mathrm{vol}(\mathcal{F}_F^{(v)} W_{(1,u)}^{E_S})}{3!} dudv \approx 0.179638, \\ S^g(W_{\bullet\bullet\bullet}^F; C) &= \frac{1}{\mathbf{v}^g} \int_{\Delta^C} e^{-(3u-2)\xi_0} \cdot \frac{\mathrm{vol}(\mathcal{F}_C^{(w)} W_{(1,u,v)}^F)}{2!} dudv dw \approx 0.211933. \end{aligned}$$

We conclude that

$$\begin{aligned} \delta_{C_2}^g(M) &\geq \frac{1}{3} \min \left\{ \frac{A_{M_4}(E_S)}{S^g(R_\bullet; E_S)}, \frac{A_{E_S}(F)}{S^g(W_{\bullet\bullet}^{E_S}; F)}, \frac{A_F(C)}{S^g(W_{\bullet\bullet}^F; C)} \right\} \\ &= \min \left\{ 1, \frac{1/3}{0.179638}, \frac{1/3}{0.211933} \right\} = 1. \end{aligned}$$

By [MW23, Corollary 5.5], we see that (M_4, ξ_0) is weighted K-semistable. Then following the same argument of Theorem 6.5 (see also [MW23, MW24]), it is weighted K-polystable. \square

For Del Pezzo fivefolds M_5 , consider the \mathbb{G}_m -action induced by $-\frac{1}{5}(2\eta + \varsigma + \zeta)$ (see Section 4.1) with toric divisor E_W . Since $A_{M_5}(E_W) = 2$ and $|\mathcal{O}(1) - tE_W|_{\mathbb{Q}} \neq \emptyset$ for $t \in [0, 1]$, the moment polytope of this \mathbb{G}_m -action on $-K_{M_5} = \mathcal{O}(4)$ is $\mathbf{P} = [-2, 2] \ni 4t - 2 =: t'$. We also numerically solve the soliton candidate

$$\eta_0 \approx 0.1693945440748772.$$

Theorem 7.2. *The pair (M_5, η_0) is weighted K-polystable.*

Proof. With the same argument in the proof of Theorem 7.1, we have

$$\begin{aligned} \mathbf{v}^g &= \int_0^1 e^{-(4t-2)\eta_0} \frac{\text{vol}(W_{(1,t)}^{E_W})}{4!} dt \\ &\approx 0.04119805228615477, \\ S^g(R_\bullet; E_W) &= \frac{1}{4} A_{M_5}(E_W). \end{aligned}$$

Moreover, we have

$$\begin{aligned} S^g(W_{\bullet\bullet}^{E_W}; E_l) &= \frac{1}{\mathbf{v}^g} \int_0^1 e^{-(4t-2)\eta_0} \int_0^{1+t} \frac{\text{vol}(W_{(1,t)}^{E_W} - sE_l)}{4!} ds dt = \frac{3}{4}, \\ S^g(W_{\bullet\bullet\bullet}^{E_l}; E_C) &= \frac{1}{\mathbf{v}^g} \int_{\Delta_A \cup \Delta_B \cup \Delta_{C_1} \cup \Delta_{C_2}} e^{-(4t-2)\eta_0} \cdot \frac{\text{vol}(W_{(1,t,s)}^{E_l} - rE_C)}{3!} dr ds dt \approx 0.390484, \\ S^g(W_{\bullet\bullet\bullet}^{E_C}; \tilde{C}) &= \frac{1}{\mathbf{v}^g} \int_{\Delta_{AA} \cup \Delta_{BB}} e^{-(4t-2)\eta_0} \cdot \frac{\text{vol}(W_{(1,t,s,r)}^{E_C} - u\tilde{C})}{2!} du dr ds dt \approx 0.089469. \end{aligned}$$

We conclude that

$$\begin{aligned} \delta_{l_{23}}^g(M_5) &\geq \frac{1}{4} \min \left\{ \frac{A_M(E_W)}{S^g(R_\bullet; E_W)}, \frac{A_{E_W}(E_l)}{S^g(W_{\bullet\bullet}^{E_W}; E_l)}, \frac{A_{E_l}(E_C)}{S^g(W_{\bullet\bullet\bullet}^{E_l}; E_C)}, \frac{A_{E_C}(\tilde{C})}{S^g(W_{\bullet\bullet\bullet}^{E_C}; \tilde{C})} \right\} \\ &= \min \left\{ 1, 1, \frac{1/2}{0.390484}, \frac{1/4}{0.089469} \right\} = 1. \end{aligned}$$

Now the same argument in Theorem 6.6 and 7.1 shows that (M_5, η_0) is weighted K-polystable. \square

REFERENCES

- [ACC⁺23] Carolina Araujo, Ana-Maria Castravet, Ivan Cheltsov, Kento Fujita, Anne-Sophie Kaloghiros, Jesus Martinez-Garcia, Constantin Shramov, Hendrik Süß, and Nivedita Viswanathan. *The Calabi problem for Fano threefolds*, volume 485 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2023.
- [ADL19] Kenneth Ascher, Kristin DeVleming, and Yuchen Liu. Wall crossing for K-moduli spaces of plane curves. *Proc. Lond. Math. Soc.*, to appear, 2019. arXiv:1909.04576.
- [AZ22] Hamid Abban and Ziquan Zhuang. K-stability of Fano varieties via admissible flags. *Forum Math. Pi*, 10:Paper No. e15, 43, 2022.
- [AZ23] Hamid Abban and Ziquan Zhuang. Seshadri constants and K -stability of Fano manifolds. *Duke Math. J.*, 172(6):1109–1144, 2023.
- [BBJ21] Robert Berman, Sebastian Boucksom, and Mattias Jonsson. A variational approach to the Yau-Tian-Donaldson conjecture. *J. Amer. Math. Soc.*, 34(3):605–652, 2021.
- [Ber16] Robert J. Berman. K-polystability of \mathbb{Q} -Fano varieties admitting Kähler-Einstein metrics. *Invent. Math.*, 203(3):973–1025, 2016.
- [BL22] Harold Blum and Yuchen Liu. Openness of uniform K-stability in families of \mathbb{Q} -Fano varieties. *Ann. Sci. Éc. Norm. Supér. (4)*, 55(1):1–41, 2022.
- [BLX22] Harold Blum, Yuchen Liu, and Chenyang Xu. Openness of K-semistability for Fano varieties. *Duke Math. J.*, 171(13):2753–2797, 2022.
- [BLXZ23] Harold Blum, Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. The existence of the Kähler-Ricci soliton degeneration. *Forum Math. Pi*, 11:Paper No. e9, 28, 2023.
- [CDS15] Xiuxiong Chen, Simon Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities, II: Limits with cone angle less than 2π , III: Limits as cone angle approaches 2π and completion of the main proof. *J. Amer. Math. Soc.*, 28(1):183–197, 199–234, 235–278, 2015.
- [Che01] I. A. Cheltsov. Log canonical thresholds on hypersurfaces. *Mat. Sb.*, 192(8):155–172, 2001.
- [CS09] Ivan Cheltsov and Constantin Shramov. Extremal metrics on del Pezzo threefolds. *Tr. Mat. Inst. Steklova*, 264:37–51, 2009.
- [Deb20] Olivier Debarre. Gushel-Mukai varieties. 2020. arXiv:2001.03485.
- [Del22] Thibaut Delcroix. Examples of K-unstable Fano manifolds. *Ann. Inst. Fourier (Grenoble)*, 72(5):2079–2108, 2022.
- [DK18] Olivier Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: classification and birationalities. *Algebr. Geom.*, 5(1):15–76, 2018.
- [DK20] Olivier Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: moduli. *Internat. J. Math.*, 31(2):2050013, 59, 2020.
- [Don02] S. K. Donaldson. Scalar curvature and stability of toric varieties. *J. Differential Geom.*, 62(2):289–349, 2002.
- [Fuj81] Takao Fujita. On the structure of polarized manifolds with total deficiency one. II. *J. Math. Soc. Japan*, 33(3):415–434, 1981.
- [Fuj90] Takao Fujita. On singular del Pezzo varieties. In *Algebraic geometry (L'Aquila, 1988)*, volume 1417 of *Lecture Notes in Math.*, pages 117–128. Springer, Berlin, 1990.
- [Fuj17] Kento Fujita. Examples of K-unstable Fano manifolds with the Picard number 1. *Proc. Edinb. Math. Soc. (2)*, 60(4):881–891, 2017.
- [Fuj19] Kento Fujita. K-stability of Fano manifolds with not small alpha invariants. *J. Inst. Math. Jussieu*, 18(3):519–530, 2019.
- [Gus83] N.P. Gushel'. Fano varieties of genus 6. *Izv. Math.*, 21:445–459, 1983.

- [HL23] Jiyuan Han and Chi Li. On the Yau-Tian-Donaldson conjecture for generalized Kähler-Ricci soliton equations. *Comm. Pure Appl. Math.*, 76(9):1793–1867, 2023.
- [Ino19] Eiji Inoue. The moduli space of Fano manifolds with Kähler-Ricci solitons. *Adv. Math.*, 357:106841, 65, 2019.
- [JM12] Mattias Jonsson and Mircea Mustață. Valuations and asymptotic invariants for sequences of ideals. *Ann. Inst. Fourier (Grenoble)*, 62(6):2145–2209 (2013), 2012.
- [KO73] Shoshichi Kobayashi and Takushiro Ochiai. Characterizations of complex projective spaces and hyperquadrics. *J. Math. Kyoto Univ.*, 13:31–47, 1973.
- [Li22] Chi Li. G -uniform stability and Kähler-Einstein metrics on Fano varieties. *Invent. Math.*, 227(2):661–744, 2022.
- [Liu22] Yuchen Liu. K -stability of cubic fourfolds. *J. Reine Angew. Math.*, 786:55–77, 2022.
- [LTW22] Chi Li, Gang Tian, and Feng Wang. The uniform version of Yau-Tian-Donaldson conjecture for singular Fano varieties. *Peking Math. J.*, 5(2):383–426, 2022.
- [LX14] Chi Li and Chenyang Xu. Special test configuration and K -stability of Fano varieties. *Ann. of Math. (2)*, 180(1):197–232, 2014.
- [LX19] Yuchen Liu and Chenyang Xu. K -stability of cubic threefolds. *Duke Math. J.*, 168(11):2029–2073, 2019.
- [LX20] Chi Li and Chenyang Xu. Stability of valuations and Kollár components. *J. Eur. Math. Soc. (JEMS)*, 22(8):2573–2627, 2020.
- [LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to K -stability. *Ann. of Math. (2)*, 196(2):507–566, 2022.
- [LZ22a] Yuchen Liu and Ziwen Zhu. Equivariant K -stability under finite group action. *Internat. J. Math.*, 33(1):Paper No. 2250007, 21, 2022.
- [LZ22b] Yuchen Liu and Ziquan Zhuang. On the sharpness of Tian’s criterion for K -stability. *Nagoya Math. J.*, 245:41–73, 2022.
- [Muk89] Shigeru Mukai. Biregular classification of Fano 3-folds and Fano manifolds of coindex 3. *Proc. Nat. Acad. Sci. U.S.A.*, 86(9):3000–3002, 1989.
- [MW23] Minghao Miao and Linsheng Wang. Kähler-Ricci solitons on Fano threefolds with non-trivial moduli. 2023. arXiv:2309.14212.
- [MW24] Minghao Miao and Linsheng Wang. Optimal degenerations of K -unstable Fano threefolds. 2024. arXiv:2401.13999.
- [PVdV99] J. Piontowski and A. Van de Ven. The automorphism group of linear sections of the Grassmannians $G(1, N)$. *Doc. Math.*, 4:623–664, 1999.
- [PZ18] Yuri Prokhorov and Mikhail Zaidenberg. Fano-Mukai fourfolds of genus 10 as compactifications of \mathbb{C}^4 . *Eur. J. Math.*, 4(3):1197–1263, 2018.
- [Tia90] Gang Tian. On Calabi’s conjecture for complex surfaces with positive first Chern class. *Invent. Math.*, 101(1):101–172, 1990.
- [Tia97] Gang Tian. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.*, 130(1):1–37, 1997.
- [Tia15] Gang Tian. K -stability and Kähler-Einstein metrics. *Comm. Pure Appl. Math.*, 68(7):1085–1156, 2015.
- [XZ20] Chenyang Xu and Ziquan Zhuang. On positivity of the CM line bundle on K -moduli spaces. *Ann. of Math. (2)*, 192(3):1005–1068, 2020.
- [Zho23] Chuyu Zhou. On wall-crossing for K -stability. *Adv. Math.*, 413:Paper No. 108857, 26, 2023.
- [Zhu21] Ziquan Zhuang. Optimal destabilizing centers and equivariant K -stability. *Invent. Math.*, 226(1):195–223, 2021.
- [ZZ22] Kewei Zhang and Chuyu Zhou. Delta invariants of projective bundles and projective cones of Fano type. *Math. Z.*, 300(1):179–207, 2022.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208, USA

Email address: yuchenl@northwestern.edu

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING, 210008, CHINA

Email address: linsheng_wang@outlook.com