

TRANSVERSE GEOMETRIC FORMALITY

GEORGES HABIB, KEN RICHARDSON, AND ROBERT WOLAK

ABSTRACT. A Riemannian metric on a closed manifold is said to be geometrically formal if the wedge product of any two harmonic forms is harmonic; equivalently, the interior product of any two harmonic forms is harmonic. Given a Riemannian foliation on a closed manifold, we say that a bundle-like metric is transversely geometrically formal if the interior product of any two basic harmonic forms is basic harmonic. In this paper, we examine the geometric and topological consequences of this condition.

CONTENTS

1. Introduction	2
2. Preliminaries	3
3. Transverse formality	6
3.1. General results concerning transverse formality	6
3.2. Formality versus transverse formality	10
3.3. Transverse Lie Foliations	11
3.4. Transverse formality is not a transverse property	13
4. Transverse formality for Riemannian flows, K -contact and Sasaki manifolds	15
4.1. General results for Riemannian flows	15
4.2. Sasakian manifolds	18
4.3. Classification of three-dimensional Riemannian flows that are transversely formal	20
5. Examples of Riemannian foliations with transverse formality	20
5.1. Geometric formality and transverse formality	20
5.2. The Hopf fibration of \mathbb{S}^3	21
5.3. The Carrière example	22
5.4. Solvmanifold admitting transversely formal Riemannian foliations	22
6. Appendix	25
References	29

Date: May 17, 2024.

2010 *Mathematics Subject Classification.* 53C12; 53C25; 58A14.

Key words and phrases. Riemannian foliation, isometric flow, Laplacian, basic cohomology, geometric formality.

The authors acknowledge that the research cooperation was funded by the program Excellence Initiative – Research University at the Jagiellonian University in Krakow within the framework of the research group Reeb-Reinhart 2022. The second author acknowledges support of Alfried Krupp Wissenschaftskolleg.

1. INTRODUCTION

The notion of formality, as introduced in [42] and explored further in [14], played a very important role in the study of the topology of manifolds. In particular, the authors demonstrated that compact Kähler manifolds are formal, thus the formality is an obstruction to the existence of a Kähler structure. Moreover, there are examples of symplectic, non-Kähler solvmanifolds; cf. [46]. Formal solvmanifolds were investigated in depth in [28].

In [30] Kotschick proposed a finer notion of formality for Riemannian manifolds: a compact Riemannian manifold (M, g) is geometrically formal if the wedge product of any two harmonic forms is a harmonic form. Therefore the space of harmonic forms is an algebra with the wedge product, and according to the Hodge theorem it is a minimal model for the cohomology of M . Thus any geometrically formal Riemannian manifold (M, g) is formal. The investigation into the geometrical formality has been continued by Kotschick and some other researchers resulting in numerous interesting theorems (cf. [2],[20], [31]), among them a fine classification of geometrically formal Riemannian manifolds in low dimensions in [4]. In [36], the author gave a complete description of Kähler manifolds where all harmonic one-forms have constant length, among other results.

Riemannian foliated manifolds (technically, Riemannian manifolds with foliations endowed with bundle-like metrics) form a subclass of Riemannian manifolds that appears naturally in many geometrical and applied problems. The foliated aspect is best expressed by the study of the transverse Riemannian geometry of the foliation. Moreover, one can investigate the interplay between the transverse and global properties of the foliated manifold. One of the tools is the basic cohomology $H^*(M, \mathcal{F})$, which in the case of compact Riemannian foliated manifolds is finite dimensional, [17], [27], [38]. Additionally, the Hodge theory for the basic cohomology was developed as well as the Poincaré duality property, cf. [26].

Continuing this line of research we introduce the notion of transverse geometric formality for a compact Riemannian foliated manifold (M, g, \mathcal{F}) . We might like to say that (M, g, \mathcal{F}) is transversely geometrically formal if any wedge product of two basic harmonic forms is basic harmonic, but the presence of mean curvature causes issues with such a definition; since the basic Laplacian does not commute with the basic Hodge star-operator, under such a definition basic harmonic forms would not have constant norm, which is a key property of formal metrics. Instead, the appropriate definition of transverse geometric formality is that the interior product of any two basic harmonic forms is also basic harmonic. If the basic mean curvature is zero, then these two properties coincide, as they do for formal metrics on Riemannian manifolds.

In Section 2 we recall, for the reader's convenience, some definitions and properties of Riemannian foliated manifolds. The notion of transverse geometric formality is introduced in Section 3, in which we provide the definition and investigate the basic properties of general Riemannian foliated manifolds. We show an upper bound on the dimensions of basic cohomology satisfied by foliations admitting transversely formal metrics in Corollary 3.5. In Theorem 3.10, we consider a bundle-like metric on the foliation that satisfies transverse formality and has basic harmonic mean curvature. In this case, if the dimension of a cohomology group is maximal, the foliation is necessarily minimal.

We next investigate the relation between geometric formality and transverse geometric formality in Subsection 3.2. For example, in Proposition 3.15, we show that for a Riemannian foliated manifold with basic mean curvature and involutive normal bundle, geometric formality implies transverse formality. Next, we study transverse geometric formality of

transversely Lie foliations, a class of Riemannian foliations which, due to Molino's structure theorem, cf. [35], are the basic building blocks of any compact Riemannian foliated manifold. We prove various results in this section, including Proposition 3.23, where we show that if a bundle-like metric is transversely formal and has basic harmonic mean curvature with maximal $\dim H^1(M, \mathcal{F})$, then the foliation is a \mathbb{R}^q -Lie foliation. In Subsection 3.4, we demonstrate that geometric formality is not a transverse property in Häefliger's sense, and we show for example that taut foliations that are transversely formal must be minimal.

Section 4 is dedicated to the detailed study of one-dimensional Riemannian foliations, i.e. Riemannian flows, and their subclasses, namely isometric flows, K-contact manifolds, and Sasakian manifolds. Among other results, we prove in Theorem 4.10 that if a Riemannian flow is minimal and transversely geometrically formal and satisfies a Ricci curvature condition, some inequalities are satisfied between the basic and ordinary cohomology. In Theorem 4.12, on closed Sasakian manifold, geometric formality implies that the foliation defined by the Reeb vector field is transversely geometrically formal. We classify all one-dimensional transversely geometrically formal foliations on closed three-manifolds in Theorem 4.15.

In Section 5, we present some relevant examples. First, we show that S^3 foliated by the Hopf fibration is transversely geometrically formal. Then we investigate Carrière's example of a non-isometric flow on a 3-manifold and verify that it is transversely geometrically formal. Finally, we construct some Riemannian foliations on a solvmanifold and study their geometric formality. In the appendix we give an example of a nontaut Riemannian foliation of codimension 4, with noninvolutive normal bundle carrying a transversely formal bundle-like metric.

2. PRELIMINARIES

Let (M, g) be a closed Riemannian manifold of dimension m , endowed with a transversely oriented Riemannian foliation \mathcal{F} of codimension q and dimension $p = m - q$. The foliation \mathcal{F} is given by an integrable subbundle $T\mathcal{F}$ of TM of rank p . Let $Q = TM/T\mathcal{F} \cong (T\mathcal{F})^\perp$ be the normal bundle of the foliation. In this paper we will in fact let $Q = (T\mathcal{F})^\perp$ to simplify exposition. We assume the metric g to be bundle-like, meaning that the induced metric g_Q of g on Q satisfies the holonomy invariance property. This means

$$\mathcal{L}_X g_Q = 0,$$

for any $X \in \Gamma(T\mathcal{F})$, where \mathcal{L}_X denotes the Lie derivative in the direction of X . Such metrics have been studied for decades; see [39, 35, 44]. The existence of bundle-like metrics assures the existence of a transverse Levi-Civita connection ∇ on $\Gamma(Q)$, i.e. a torsion-free connection compatible with the metric g_Q extending the Bott partial connection; see [44, Ch. 3]. This connection can be extended in the usual way to act on $\Gamma(\wedge^* Q^*)$. For simplicity we will assume that our foliated manifolds are oriented and transversely oriented.

Notation 2.1. *We call the triple (M, \mathcal{F}, g) an oriented Riemannian foliated manifold (ORFM) if M is an oriented, closed Riemannian manifold with metric g , and \mathcal{F} is a transversely oriented Riemannian foliation for which g is bundle-like.*

Following [44] we define the mean curvature form κ (or its dual vector $\kappa^\#$) as

$$\kappa^\# = \sum_{j=1}^p \pi \left(\nabla_{f_j}^M f_j \right),$$

where $\pi : TM \rightarrow Q$ is the projection, $\{f_j\}_{j=1,\dots,p}$ is a local orthonormal frame of $T\mathcal{F}$, and ∇^M is the Levi-Civita connection on M .

The characteristic form $\chi_{\mathcal{F}}$ of the foliation \mathcal{F} is the leafwise volume form and is defined as follows:

$$\chi_{\mathcal{F}}(Y_1, \dots, Y_p) = \det(g(Y_i, f_j)_{i,j=1,\dots,p}),$$

where Y_1, \dots, Y_p are any vector fields on M .

The characteristic form $\chi_{\mathcal{F}}$ and the mean curvature form κ are related by the Rummmler's formula, [41]:

$$d\chi_{\mathcal{F}} = -\kappa \lrcorner \chi_{\mathcal{F}} + \varphi_0,$$

where φ_0 satisfies the property $\chi_{\mathcal{F}} \lrcorner \varphi_0 = 0$; here, \lrcorner denotes interior product of differential forms. The mean curvature one-form κ depends on the choice of the bundle-like metric. The normal bundle Q is involutive if and only if $\varphi_0 = 0$.

On any Riemannian foliated manifold (M, g, \mathcal{F}) , the set of all basic r -forms ($0 \leq r \leq q$) is

$$\Omega^r(M, \mathcal{F}) = \{\alpha \in \Omega^r(M) : X \lrcorner \alpha = X \lrcorner d\alpha = 0 \text{ for all vectors } X \in \Gamma(T\mathcal{F})\},$$

which is a subcomplex of the differential forms $\Omega^r(M)$ on M . The basic cohomology is then defined as being the cohomology associated to the exterior differential restricted to basic forms, that is

$$H^r(M, \mathcal{F}) = \frac{\ker(d : \Omega^r(M, \mathcal{F}) \rightarrow \Omega^{r+1}(M, \mathcal{F}))}{\text{im}(d : \Omega^{r-1}(M, \mathcal{F}) \rightarrow \Omega^r(M, \mathcal{F}))},$$

for $0 \leq r \leq q$. It is shown in [17], [27], [38] that the basic cohomology is always finite dimensional but does not necessarily satisfy Poincaré duality (see [12] for an example).

The restriction of the bundle-like metric to the normal bundle of the foliation of the oriented Riemannian manifold (M, g, \mathcal{F}) defines the basic Hodge star operator $\bar{*}$, cf. [44],

$$\bar{*} : \Omega^r(M, \mathcal{F}) \rightarrow \Omega^{q-r}(M, \mathcal{F}).$$

This operator is related to the usual Hodge $*$ -operator on (M, g) by the formula

$$\bar{*}\alpha = (-1)^{p(q-r)} * (\alpha \wedge \chi_{\mathcal{F}})$$

for any $\alpha \in \Omega^r(M, \mathcal{F})$. The pointwise inner product between basic forms is defined in the usual way as; if $\alpha, \beta \in \Omega^r(M, \mathcal{F})$ we have

$$\langle \alpha, \beta \rangle = \bar{*}(\alpha \wedge \bar{*}\beta).$$

In this case, the standard scalar product satisfies

$$\langle \alpha, \beta \rangle = \int_M \alpha \wedge \bar{*}\beta \wedge \chi_{\mathcal{F}}.$$

A Riemannian foliation on a compact manifold is said to be taut if there exists a Riemannian metric which makes all its leaves minimal submanifolds. Tautness is characterized by the nonvanishing of the top dimensional basic cohomology, i.e., $H^q(M, \mathcal{F}) \neq 0$ (see [33]).

Let $L^2(\Omega(M))$ and $L^2(\Omega(M, \mathcal{F}))$ denote the closures of $\Omega(M)$ and $\Omega(M, \mathcal{F})$ with respect to the L^2 inner product on forms $\langle \cdot, \cdot \rangle$, and let $P_b : L^2(\Omega(M)) \rightarrow L^2(\Omega(M, \mathcal{F}))$ denote the orthogonal projection (see [38]). We denote $\kappa_b = P_b \kappa$. By [1], κ_b is always closed, and it defines a cohomology class $[\kappa_b] \in H^1(M, \mathcal{F})$ that is independent of the choice of a bundle-like metric. By [15], there exists a bundle-like metric g' whose mean curvature is basic ($\kappa = \kappa_b$),

such that g and g' define the same metric on Q . Another property that characterises tautness of a Riemannian foliation is that $[\kappa_b] = 0$ (see [33]).

The formal adjoint δ_b of d in the complex $\Omega^*(M, \mathcal{F})$ with respect to the scalar product $\langle \cdot, \cdot \rangle$ is the operator

$$\delta_b = (-1)^{q(r+1)+1} \bar{*}(d - \kappa_b \wedge) \bar{*} : \Omega^r(M, \mathcal{F}) \rightarrow \Omega^{r-1}(M, \mathcal{F}).$$

The basic Laplacian is the operator on basic forms defined by

$$\Delta_b = \delta_b d + d \delta_b.$$

A basic form α is called **basic harmonic** if and only if $\Delta_b \alpha = 0$; equivalently, $d\alpha = 0$ and $\delta_b \alpha = 0$.

The transverse volume form ν is a basic q -form; it satisfies $\delta_b \nu = \kappa_b \lrcorner \nu = \bar{*} \kappa_b$. In particular, the transverse volume form ν is basic harmonic if and only if the basic component of mean curvature is zero. This is certainly the case if the foliation is minimal, but general bundle-like metrics on a taut Riemannian foliation do not necessarily satisfy this condition.

The basic Hodge theorem was proved in special cases by [17], [27] and in general by [38]. In particular, each basic cohomology class contains a unique basic harmonic form.

Let us introduce the twisted differential

$$d_{\kappa_b} := d - \kappa_b \wedge .$$

Since κ_b is closed, the twisted differential satisfies $d_{\kappa_b}^2 = 0$ and, therefore, defines a chain complex. We denote by $H_{\kappa_b}^*(M, \mathcal{F})$ the corresponding cohomology. The formal adjoint of d_{κ_b} is $\delta_{\kappa_b} := \delta_b - \kappa_b \lrcorner$. The two adjoint operators δ_b and δ_{κ_b} are related by

$$\bar{*} \delta_b = \delta_{\kappa_b} \bar{*}.$$

Hence, we introduce the twisted Laplacian operator as

$$\Delta_{\kappa_b} := d_{\kappa_b} \delta_{\kappa_b} + \delta_{\kappa_b} d_{\kappa_b}.$$

As for the basic cohomology, we can show that there is a basic Hodge theorem for the twisted Laplacian and that each cohomology class in $H_{\kappa_b}^*(M, \mathcal{F})$ contains a unique basic κ_b -**harmonic** form. Here, a basic form α is called κ_b -harmonic if $\Delta_{\kappa_b} \alpha = 0$; equivalently, $d\alpha = \kappa_b \wedge \alpha$ and $\delta_b \alpha = \kappa_b \lrcorner \alpha$.

For the basic cohomology of compact Riemannian foliated manifolds we have the following twisted duality theorem (cf. [44, Theorem 7.54], with modifications from [38] and [1]):

Theorem 2.2. *Let (M, \mathcal{F}, g) be an ORFM. The pairing $\alpha \otimes \beta \mapsto \int_M \alpha \wedge \beta \wedge \chi_{\mathcal{F}}$ induces a nondegenerate pairing*

$$H^r(M, \mathcal{F}) \otimes H_{\kappa_b}^{q-r}(M, \mathcal{F}) \rightarrow \mathbb{R}$$

of finite dimensional spaces.

It is not difficult to check that the transversal Hodge star $\bar{*}$ operator maps basic harmonic r -forms to basic κ_b -harmonic $(q - r)$ -forms, and vice-versa. This is a consequence of the fact that $\Delta_b \bar{*} = \bar{*} \Delta_{\kappa_b}$ and $\Delta_{\kappa_b} \bar{*} = \bar{*} \Delta_b$; see [44, proof of Theorem 7.54], with modifications from [1, Corollary 3.3 and proof] or [38, Proposition 2.2]. Therefore, by the basic Hodge theorem for Δ_b and Δ_{κ_b} , we have that

$$H^*(M, \mathcal{F}) \simeq H_{\kappa_b}^{q-*}(M, \mathcal{F}).$$

This also explains the reason why Poincaré duality does not hold in general for the basic cohomology.

Finally, we end with the following result that we will use in the sequel.

Lemma 2.3. *If (M, \mathcal{F}, g) has basic mean curvature, then every basic harmonic one-form is harmonic.*

Proof. Let $p = \dim \mathcal{F}$. If α is a basic harmonic one-form, by [38, Proposition 2.4] we have that $d\alpha = 0$ and

$$\begin{aligned} 0 &= \delta_b \alpha = (\delta + (\kappa_b - \kappa) \lrcorner + (-1)^p \varphi_{0 \lrcorner} \circ \chi_{\mathcal{F}} \wedge) \alpha \\ &= (\delta + 0 + (-1)^p \varphi_{0 \lrcorner} \circ \chi_{\mathcal{F}} \wedge) \alpha = \delta \alpha, \end{aligned}$$

since $\varphi_{0 \lrcorner}(\chi_{\mathcal{F}} \wedge \cdot)$ is always zero on basic one-forms. \square

3. TRANSVERSE FORMALITY

In this section, we introduce the notion of transverse geometric formality of an ORFM both for taut and nontaut Riemannian foliations. The main problem in the nontaut case is the fact that the basic Hodge star operator does not commute with the basic Laplacian. An equivalent definition using the interior product of forms avoids this awkwardness. We investigate the basic cohomology of a transversely formal ORFM and their influence on the topology of the foliated manifold. Particular attention is paid to transversely Lie foliations. Then the relation between geometric formality and transverse formality is studied. Finally, it is shown that transverse formality is not a transverse property.

Definition 3.1. *Let (M, \mathcal{F}, g) be an ORFM. We say that (M, \mathcal{F}, g) is **transversely formal** (or **transversely geometrically formal**) if the wedge product of any basic harmonic form and basic κ_b -harmonic form is basic κ_b -harmonic.*

Definition 3.2. *Let (M, \mathcal{F}, g) be an ORFM. We say that (M, \mathcal{F}, g) is **transversely r -formal** (or **transversely geometrically r -formal**) for $0 \leq r \leq q$ if the wedge product of any basic harmonic r -form and basic κ_b -harmonic $(q - r)$ -form is basic κ_b -harmonic.*

Remark 3.3. *Note that in [37], r -formality is the condition that the wedge product of any two harmonic r -forms is harmonic. In our foliation case, it turns out to be more convenient to use the definition above, in light of Lemma 3.4.*

We note that transverse formality and formality are distinct; an ORFM may satisfy one property but not the other. See Section 5, which contains several interesting examples of Riemannian foliations that satisfy transverse formality.

3.1. General results concerning transverse formality. In this subsection, we prove the fundamental properties of transversely formal metrics.

Lemma 3.4. *Suppose that (M, \mathcal{F}, g) is transversely r -formal. Then the pointwise inner product of any two basic harmonic r -forms is constant. In particular, basic harmonic r -forms have constant length. The same is true for basic κ_b -harmonic $(q - r)$ -forms.*

Proof. If the r -forms α, β are basic harmonic and ν is the transverse volume q -form, then the pointwise inner product satisfies

$$(\alpha, \beta) \nu = \alpha \wedge \bar{*} \beta,$$

and $\bar{*} \beta$ is basic κ_b -harmonic. Since $H_{\kappa_b}^q(M, \mathcal{F}) \cong H^0(M, \mathcal{F}) \cong \mathbb{R}$ is generated by the basic κ_b -harmonic form ν . Thus, (α, β) is constant (and also $(\bar{*} \alpha, \bar{*} \beta)$ is constant). \square

Corollary 3.5. *Let \mathcal{F} be a codimension q foliation of a compact manifold M that admits a transversely r -formal bundle-like metric g . Then the basic cohomology satisfies $\dim H^r(M, \mathcal{F}) \leq \binom{q}{r}$ for $0 \leq r \leq q$.*

Proof. From the lemma above, for the transversely formal metric, the number of linearly independent harmonic forms of degree r is at most the rank of $\wedge^r Q^* = \binom{q}{r}$. \square

Remark 3.6. *For ordinary n -dimensional closed manifolds, the existence of a geometrically formal metric implies that $\dim H^1(M) \neq n-1$ (see [30, Theorem 6]). However, for particular transversely formal foliations of codimension q , the dimension of $H^1(M, \mathcal{F})$ might be equal to $q-1$ (see Corollary 3.11); the proof does not generalize because of the interaction with the mean curvature. See for example Section 5.3 for an example of a codimension 2 foliation that has a transversely formal metric where $\dim H^1(M, \mathcal{F}) = 1$. If, however, the foliation is taut, we do have $\dim H^1(M, \mathcal{F}) \neq q-1$ (see Corollary 3.26).*

As equivalent definition of transverse formality is given in the next proposition:

Proposition 3.7. *The Riemannian foliated manifold (M, \mathcal{F}, g) is transversely formal if and only if the interior product of any two basic harmonic forms is basic harmonic. The same is true for two basic κ_b -harmonic forms.*

Proof. Suppose that (M, \mathcal{F}, g) is transversely formal. If α, β are two basic harmonic forms, then $\bar{*}\beta$ is basic κ_b -harmonic, so by transverse formality $\alpha \wedge \bar{*}\beta$ is basic κ_b -harmonic. Thus, $\alpha \lrcorner \beta = \pm \bar{*}(\alpha \wedge \bar{*}\beta)$ is basic harmonic. A similar proof works for two basic κ_b -harmonic forms.

Conversely, suppose that (M, \mathcal{F}, g) has the required property, and suppose that α is basic harmonic and β is basic κ_b -harmonic. Then $\bar{*}\beta$ is basic harmonic, and thus $\alpha \lrcorner \bar{*}\beta = \pm \bar{*}(\alpha \wedge \beta)$ is basic harmonic, so that $\alpha \wedge \beta$ is basic κ_b -harmonic. A similar proof works for the κ_b -harmonic case. \square

In the following, we will study the case where the foliation is transversely formal and that $\dim H^r(M, \mathcal{F}) = \binom{q}{r}$ for some r . We will see that this is a restriction on the geometry of the foliation. For this, we need several lemmas:

Lemma 3.8. *Suppose that (M, \mathcal{F}, g) is transversely formal, and κ is basic harmonic. If α is basic harmonic r -form, then*

$$\nabla_{\kappa^\#} \alpha + \sum_{i=1}^q \nabla_{e_i} \kappa \wedge (e_i \lrcorner \alpha) = 0,$$

where $\{e_i\}_{i=1, \dots, q}$ be a local orthonormal frame of $\Gamma(Q)$.

Proof. With the assumptions, we have that $d\alpha = 0$ and $\kappa^\# \lrcorner \alpha$ is basic harmonic by Proposition 3.8, thus it is also closed. Then

$$\mathcal{L}_{\kappa^\#} \alpha = d(\kappa^\# \lrcorner \alpha) + \kappa^\# \lrcorner (d\alpha) = 0.$$

Recall the Lie derivative along any vector field X on differential r -forms can be related to the Levi-Civita connection on M by the formula $\mathcal{L}_X = \nabla_X^M + S_X^{[r]}$ where $S_X^{[r]}$ is the canonical extension of the endomorphism $S = \nabla^M X$ to differential forms defined by

$$(S_X^{[r]} \omega)(X_1, \dots, X_r) = \sum_{j=1}^r \omega(X_1, \dots, \nabla_{X_j}^M X, \dots, X_r)$$

for any differential r -form ω and $X_1, \dots, X_r \in TM$. Equivalently, this extension is equal to

$$S_X^{[r]} = \sum_{l=1}^p f_l \wedge S(f_l) \lrcorner + \sum_{i=1}^q e_i \wedge S(e_i) \lrcorner,$$

where $\{f_l\}_{l=1, \dots, p}$ is a local orthonormal frame of $T\mathcal{F}$ and $\{e_i\}_{i=1, \dots, q}$ is a local orthonormal frame of Q . Hence for $X = \kappa^\#$, we get that $\nabla_{\kappa^\#}^M \alpha + S_{\kappa^\#}^{[r]} \alpha = 0$. Applying this last identity to sections in Q , we deduce the required identity, since the form α is basic and $S = \nabla^M \kappa$ is a symmetric endomorphism as κ is closed. \square

Lemma 3.9. *Suppose that the ORFM (M, \mathcal{F}, g) is transversely formal, and suppose that the mean curvature form κ is basic harmonic. Then for any two basic harmonic forms α, β , the pointwise inner product $(\nabla_{\kappa^\#} \alpha, \beta)$ is identically zero. The same fact is true for two basic κ -harmonic forms.*

Proof. Let α and β be two basic harmonic forms. From Lemma 3.8, we have that

$$\nabla_{\kappa^\#} \alpha + \sum_{i=1}^q \nabla_{e_i} \kappa \wedge e_i \lrcorner \alpha = 0.$$

We take the pointwise inner product with β to get

$$(\nabla_{\kappa^\#} \alpha, \beta) + \sum_{i=1}^q (e_i \lrcorner \alpha, \nabla_{e_i} \kappa \lrcorner \beta) = 0. \quad (3.1)$$

Switching the roles of α and β , we get

$$(\nabla_{\kappa^\#} \beta, \alpha) + \sum_{i=1}^q (e_i \lrcorner \beta, \nabla_{e_i} \kappa \lrcorner \alpha) = 0. \quad (3.2)$$

Note that $\sum_{i=1}^q (e_i \lrcorner \alpha, \nabla_{e_i} \kappa \lrcorner \beta) = \sum_{i=1}^q (e_i \lrcorner \beta, \nabla_{e_i} \kappa \lrcorner \alpha)$, because $\nabla \kappa$ is a symmetric two-tensor as κ is closed. Thus, subtracting (3.1) and (3.2), we get

$$(\nabla_{\kappa^\#} \alpha, \beta) = (\nabla_{\kappa^\#} \beta, \alpha).$$

But, since (α, β) is constant as the metric is transversely formal, we get that

$$0 = \kappa^\# (\alpha, \beta) = (\nabla_{\kappa^\#} \alpha, \beta) + (\alpha, \nabla_{\kappa^\#} \beta),$$

so that $(\nabla_{\kappa^\#} \alpha, \beta) = 0$ which is the statement of the lemma. The last part follows from the fact that $\bar{*}$ maps the space of basic κ -harmonic forms isometrically to the space of basic harmonic forms and that $\bar{*}$ is a transversal isometry. \square

Theorem 3.10. *Suppose that the ORFM (M, \mathcal{F}, g) is transversely formal and has basic harmonic mean curvature. If $\dim H^r(M, \mathcal{F}) = \binom{q}{r}$ for some r with $0 < r \leq q$, then the foliation is minimal.*

Proof. If $r = q$, $\dim H^r(M, \mathcal{F}) = 1$, so the foliation is taut. Since the mean curvature is basic harmonic, it must be zero. Now suppose $\dim H^r(M, \mathcal{F}) = \binom{q}{r}$ for some r with $0 < r < q$. By Lemma 3.9, $(\nabla_{\kappa^\#} \alpha, \beta) = 0$ for all basic harmonic r -forms α, β . Since $\dim H^r(M, \mathcal{F}) = \binom{q}{r} = \text{rank } \wedge^r Q^*$, then $\nabla_{\kappa^\#} \alpha = 0$. Here we are using the fact that the pointwise inner product of basic harmonic k -forms is constant, so that if a set of forms is linearly dependent at one point, then it is linearly dependent globally. Now Lemma 3.8

gives that $\sum_{i=1}^q \nabla_{e_i} \kappa \wedge e_i \lrcorner \alpha = 0$ for all basic harmonic r -forms α , which means in fact that $\sum_{i=1}^q \nabla_{e_i} \kappa \wedge e_i \lrcorner \alpha = 0$ for all $\alpha \in \wedge^r Q^*$. Since the extension $S_\kappa^{[r]}$ of the symmetric endomorphism $S = \nabla \kappa$ is identically zero, the endomorphism S must itself be zero which implies $\nabla \kappa = 0$. But then

$$0 = \delta_b \kappa = -e_i \lrcorner \nabla_{e_i} \kappa + |\kappa|^2 = |\kappa|^2$$

Then κ is identically zero. □

As a direct consequence, we get the following result

Corollary 3.11. *Suppose that the ORFM (M, \mathcal{F}, g) is transversely formal and has basic harmonic mean curvature. If the foliation is nontaut, then $1 \leq \dim H^1(M, \mathcal{F}) \leq q - 1$.*

Proof. Since the foliation is nontaut and the mean curvature is basic harmonic, we get that $\dim H^1(M, \mathcal{F}) \geq 1$. Now, Corollary 3.5 gives that $\dim H^1(M, \mathcal{F}) \leq q$. Since again the mean curvature cannot be zero, Theorem 3.10 implies that $\dim H^1(M, \mathcal{F}) < q$. Hence, we deduce the statement. □

Proposition 3.12. *Let (M, \mathcal{F}, g) be an ORFM with basic mean curvature such that there exists a basic harmonic one-form α that has unit length. Then there exists a codimension one minimal Riemannian foliation \mathcal{F}_α on M such that \mathcal{F} saturates the leaves of \mathcal{F}_α and the universal cover $(\widetilde{M}, \widetilde{\mathcal{F}}_\alpha, \widetilde{g})$ is a product codimension one Riemannian foliation.*

Proof. We are given that α is a basic harmonic one-form of constant length 1, so that it is in fact also harmonic (Lemma 2.3). Then $*\alpha$ is also a harmonic form of constant length, and we have $d\alpha = 0$ and $d(*\alpha) = 0$. Then α is the characteristic form of the foliation given by the flow of the vector field $\alpha^\#$, and $*\alpha$ is the characteristic form of the normal bundle. By Rummler's formula, the $d\alpha = 0$ implies the flow is minimal (geodesic) and the normal bundle is involutive and defines a Riemannian foliation \mathcal{F}_α of codimension one. By $d(*\alpha) = 0$, we see that the foliation \mathcal{F}_α is minimal. The form α generates the basic cohomology $H^*(M, \mathcal{F}_\alpha)$, and this cohomology injects into the cohomology of M . By [7, Corollary 3.3] and [8, Section 5], the metric and foliations lift to the universal cover $(\widetilde{M}, \widetilde{\mathcal{F}}, \widetilde{\mathcal{F}}_\alpha, \widetilde{g})$ so that the lifted $\widetilde{\mathcal{F}}_\alpha$ is a product foliation, and $T\widetilde{\mathcal{F}} \subseteq T\widetilde{\mathcal{F}}_\alpha$. □

Remark 3.13. *In the case of transverse formality, the proposition above produces k orthogonal codimension one minimal Riemannian foliations \mathcal{F}_{α_j} for an orthonormal set of basic harmonic one-forms $\{\alpha_j\}$, where $k = \dim H^1(M, \mathcal{F})$, such that the original foliation \mathcal{F} is the contained the intersection of all the \mathcal{F}_{α_j} .*

Proposition 3.14. *Let (M, \mathcal{F}, g) be a nontaut ORFM, such that the metric g is transversely formal and κ is basic harmonic (thus harmonic). Then $H = \kappa^\#$ generates a geodesic flow such that $\nabla_\bullet H$ is symmetric and the normal bundle to H is integrable and is the tangent bundle to a minimal Riemannian foliation that contains \mathcal{F} as a subfoliation.*

Proof. Since $|\kappa|^2$ is constant, $H = \kappa^\#$ has constant norm and generates a geodesic flow. Since $d\kappa = 0 = d(*\kappa)$, by Rummler's formula normal bundle to H is integrable and is the tangent bundle to a minimal Riemannian foliation. □

3.2. Formality versus transverse formality. The main results show that geometric formality of the Riemannian manifold coupled with the assumption that the foliation has basic mean curvature in some special cases implies transverse formality.

Proposition 3.15. *Suppose (M, \mathcal{F}, g) is an ORFM with basic mean curvature and integrable normal bundle. Assume that (M, g) is geometrically formal. Then (M, \mathcal{F}, g) is transversely formal. If moreover the foliation is nontaut, the pointwise inner product of a basic κ -harmonic form and a basic harmonic form is always zero.*

Proof. Since the mean curvature is basic (i.e. $\kappa = \kappa_b$) and the normal bundle is integrable, every basic harmonic form is also harmonic, because the basic codifferential is a restriction of the ordinary differential δ (see [38, Proposition 2.4]). Also conversely, any harmonic form which is basic is a basic harmonic form. Let α be a basic κ -harmonic r -form and let β be a basic harmonic s -form. Since (M, g) is geometrically formal, $\beta \lrcorner (\overline{*}\alpha)$ is harmonic. As β is basic, we have $\beta \lrcorner (\overline{*}\alpha) = \pm \overline{*}(\beta \wedge \alpha)$ and, thus, $\overline{*}(\alpha \wedge \beta)$ is harmonic. But this last form is also basic, hence it becomes basic harmonic, so that $\alpha \wedge \beta$ is a basic κ -harmonic. Thus, (M, \mathcal{F}, g) is transversely formal. In addition, if $r = s$, then $\overline{*}\alpha \wedge \beta = (\alpha, \beta)\nu$ is harmonic and thus has constant length. Then (α, β) is a constant. If the foliation is nontaut, then ν is not harmonic, since $\delta\nu = \delta_b\nu = \kappa \lrcorner \nu$, so we have $(\alpha, \beta) = 0$. \square

Remark 3.16. *Note that it is possible that a bundle-like metric on a Riemannian foliation is both transversely formal and geometrically formal, but where the normal bundle is not integrable. See, for example, the Hopf fibration in Section 5.2.*

Lemma 3.17. *If the ORFM (M, \mathcal{F}, g) is geometrically formal and has basic mean curvature, then the wedge product of any two basic harmonic 1-forms is basic harmonic.*

Proof. If α and β are basic harmonic 1-forms, then they are also harmonic, because δ_b is a restriction of δ on 1-forms (see [38, Proposition 2.4]). Then $\alpha \wedge \beta$ is harmonic, since M is formal. On the other hand, from [38, Proposition 2.4], we have

$$\delta_b(\alpha \wedge \beta) = \delta(\alpha \wedge \beta) + (-1)^{\dim \mathcal{F}} \varphi_{0 \lrcorner}(\chi_F \wedge \alpha \wedge \beta).$$

Since $\delta(\alpha \wedge \beta) = 0$ and $\varphi_{0 \lrcorner}(\chi_F \wedge \alpha \wedge \beta)$ is orthogonal to basic forms and is thus 0, we deduce $\delta_b(\alpha \wedge \beta) = 0$. Therefore $\alpha \wedge \beta$ is basic harmonic. \square

Theorem 3.18. *Let (M, \mathcal{F}, g) be an ORFM such that (M, g) is geometrically formal and \mathcal{F} has codimension two with basic mean curvature. Then (M, \mathcal{F}, g) is transversely formal. If moreover the foliation is nontaut, then $\dim H^1(M, \mathcal{F}) = 1$.*

Proof. Suppose α is a basic harmonic 1-form, and β is a basic κ -harmonic 1-form. Then $\overline{*}\beta$ is a basic harmonic 1-form. Thus, α and $\overline{*}\beta$ are harmonic forms. Since the metric is formal, we have $(\alpha, \overline{*}\beta) = (\alpha, \beta)$ is constant. But $\alpha \wedge \beta = \pm (\alpha, \overline{*}\beta)\nu$, so since $(\alpha, \overline{*}\beta)$ is constant and ν is κ -harmonic, $\alpha \wedge \beta$ is κ -harmonic. Hence M is transversely formal. By Corollary 3.5 we have $\dim H^1(M, \mathcal{F}) \leq 2$. Notice here $H^1(M, \mathcal{F})$ cannot be zero since the foliation is nontaut. We shall now prove that $\dim H^1(M, \mathcal{F}) = 2$ cannot occur. Assume it were the case and let us denote by $\{\alpha_1, \alpha_2\}$ a basis of $H^1(M, \mathcal{F})$; then by Lemma 3.17 one would get that $\alpha_1 \wedge \alpha_2$ is basic harmonic, hence belonging to $H^2(M, \mathcal{F}) = 0$, since the foliation is nontaut. Thus, $\alpha_1 = c\alpha_2$ for some $c \in \mathbb{R}$, a contradiction. Therefore $\dim H^1(M, \mathcal{F}) = 1$. \square

Remark 3.19. *From Corollary 3.11, one can see that for codimension 2 nontaut transversely formal foliations, we have that $\dim H^1(M, \mathcal{F}) = 1$. But this is true under the extra*

assumption that κ is basic harmonic, which is not the case in Theorem 3.18. However, we need the assumption that the metric should be geometrically formal.

3.3. Transverse Lie Foliations. Transverse foliations form a very particular class of Riemannian foliations which play an important role in the Molino structure theorem, cf. [35]. For this class of foliations we obtain some very strong results about transverse formality of foliations with dense leaves.

A foliation \mathcal{F} of codimension q on a manifold M is called a transverse Lie foliation if its normal bundle Q admits a global trivialization X_1, \dots, X_q by foliated vector fields which form a Lie algebra \mathfrak{g} . Let G be a simply connected Lie group whose Lie algebra is isomorphic to \mathfrak{g} . Let $x_0 \in M$. Then there exists a covering $p : \widehat{M} \rightarrow M$ of the manifold M , a surjective submersion $D : \widehat{M} \rightarrow G$ of connected fibres, and a homomorphism $h : \pi_1(M, x_0) \rightarrow G$ such that D is $\pi_1(M, x_0)$ -equivariant with respect to the natural action of $\pi_1(M, x_0)$ on \widehat{M} and the action of G by left translations. The fibres of D are the leaves of the lifted foliation $\widehat{\mathcal{F}}$. In this case, basic forms on (M, \mathcal{F}) correspond bijectively to $\Gamma = \text{im}(h)$ -invariant forms on G ; thus the complex of basic forms $A^*(M, \mathcal{F})$ can be identified with the complex of Γ -invariant forms $A^*(G)^\Gamma$. If the leaves of \mathcal{F} are dense in M , then the subgroup Γ is dense in G . Therefore, if a transverse Lie foliation has dense leaves, then basic forms are in one-to-one correspondence with left-invariant forms on G .

The normal part of a bundle-like metric g corresponds to a left-invariant Riemannian metric \hat{g} on G . Assume that our foliation is minimal, i.e. $\kappa = 0$. Then any transversely harmonic basic form is closed and coclosed. The corresponding left-invariant form on G is also closed and coclosed (and thus harmonic) for the corresponding left-invariant Riemannian metric on the Lie group G .

First, assume that the group G is compact and the Riemannian metric \hat{g} is bi-invariant. Then harmonic forms on G are just bi-invariant forms, [25, 34]. This fact permits us to formulate the following proposition.

Proposition 3.20. *Let \mathcal{F} be a transverse \mathfrak{g} -Lie foliation of a compact manifold M with dense leaves. If the corresponding simply connected Lie group is compact, then the foliation \mathcal{F} admits a bundle-like metric for which the foliation is transversely formal.*

Proof. Take a bi-invariant Riemannian metric \hat{g} on G . We can lift it to a bundle-like metric on (M, \mathcal{F}) . As G is compact and simply connected $H^q(M, \mathcal{F}) = H^q(G) \neq 0$. Thus the foliation is taut, and we can modify the metric \hat{g} along the leaves to a minimal Riemannian metric. Then the basic harmonic forms correspond to bi-invariant forms on G ; hence any wedge product of such forms is also basic harmonic. \square

Second, consider a transverse Lie foliation modelled on a nilpotent Lie group $G = N$. For such foliations we have the following result.

Theorem 3.21. *Let \mathcal{F} be a transverse \mathfrak{g} -Lie foliation of a compact manifold M with dense minimal leaves. If the Lie algebra \mathfrak{g} is nilpotent with rational structure constants and the foliation is transversely formal, then $N = \mathbb{R}^q$.*

Proof. The basic cohomology $H^*(M, \mathcal{F})$ is isomorphic to the cohomology $H^*(\mathfrak{n})$ of the Lie algebra $\mathfrak{n} = \text{Lie}(N)$. The module of basic harmonic forms is a minimal model for the cohomology $H^*(\mathfrak{n})$. The Lie group N admits a lattice Γ_0 , and the cohomology of the compact nilmanifold $M_0 = N/\Gamma_0$ is also isomorphic to $H^*(\mathfrak{n})$. Thus the compact nilmanifold M_0 is

formal. The Hasegawa result ensures the nilmanifold M_0 is a torus, cf. [23]. Thus the Lie group N must be abelian, hence $N = \mathbb{R}^q$. \square

Lastly, consider a transverse Lie foliation modelled on a solvable Lie group $G = S$. Hisashi Kasuya's results [28] on formality of solvmanifolds can be used to formulate the following result for Lie foliations.

Theorem 3.22. *Let \mathcal{F} be a taut transversely \mathfrak{g} -Lie foliation of a compact manifold M with dense minimal leaves modelled on a special solvable Lie group G , i.e. $G = \mathbb{R}^k \times_{\varphi} \mathbb{R}^s$ with a semisimple action φ . Moreover assume that G admits a lattice. Then there exists a bundle-like Riemannian metric such that the foliation is transversely formal for this bundle-like metric.*

Proof. Taking into account [28, Theorem 1.1], the proof is analogous to that of Theorem 3.21. \square

Next, we show that transverse formality and maximality of the rank of $H^1(M, \mathcal{F})$ implies that the manifold has the structure of an abelian Lie foliation.

Let (M, \mathcal{F}, g) be an ORFM. Assume that the foliation \mathcal{F} is minimal, i.e. $\kappa = 0$. Then the transverse volume form ν is a basic harmonic q -form. Assume that \mathcal{F} is transversely formal.

Let α and β be basic harmonic 1-forms, and let Y be any vector field tangent to \mathcal{F} . Then

$$\begin{aligned} 0 &= d\beta(\alpha^{\sharp}, Y) = \alpha^{\sharp}\beta(Y) - Y\beta(\alpha^{\sharp}) - \beta([\alpha^{\sharp}, Y]) \\ &= -Yg(\beta^{\sharp}, \alpha^{\sharp}) - \beta([\alpha^{\sharp}, Y]) = -\beta([\alpha^{\sharp}, Y]). \end{aligned} \quad (3.3)$$

Let α, β, γ be basic harmonic 1-forms. Then

$$\begin{aligned} 0 &= d\gamma(\alpha^{\sharp}, \beta^{\sharp}) \\ &= \alpha^{\sharp}\gamma(\beta^{\sharp}) - \beta^{\sharp}\gamma(\alpha^{\sharp}) - \gamma([\alpha^{\sharp}, \beta^{\sharp}]) \\ &= \alpha^{\sharp}g(\gamma^{\sharp}, \beta^{\sharp}) - \beta^{\sharp}g(\gamma^{\sharp}, \alpha^{\sharp}) - \gamma([\alpha^{\sharp}, \beta^{\sharp}]) = -\gamma([\alpha^{\sharp}, \beta^{\sharp}]), \end{aligned} \quad (3.4)$$

as $g(\gamma^{\sharp}, \beta^{\sharp})$ is a constant function, since $\gamma \wedge \bar{*}\beta = g(\gamma^{\sharp}, \beta^{\sharp})\nu$.

Recall that a codimension q foliation (M, \mathcal{F}) is called a \mathfrak{g} -Lie foliation if there exist foliated vector fields X_1, \dots, X_q whose projections to Q are linearly independent such that for any $1 \leq i, j \leq q$, $[X_i, X_j] = \sum_k c_{ij}^k X_k \text{ mod } \mathcal{F}$, where the c_{ij}^k are the structure constants of the Lie algebra \mathfrak{g} . See [18].

Proposition 3.23. *Let (M, \mathcal{F}, g) be an ORFM that is transversely formal with κ basic harmonic. Suppose $\dim H^1(M, \mathcal{F}) = q = \text{codim}(\mathcal{F})$, and let $\alpha_1, \dots, \alpha_q$ be an orthonormal basis of the space of basic harmonic 1-forms. Then the foliated vector fields $\alpha_1^{\sharp}, \dots, \alpha_q^{\sharp}$ commute mod \mathcal{F} . Therefore the foliation \mathcal{F} is \mathbb{R}^q -Lie foliation.*

Proof. By Theorem 3.10, $\kappa = 0$. Observe that $\cap_{k=1}^q \ker(\alpha_k) = T\mathcal{F}$. Since 3.3 holds for any leafwise vector field Y , each α_k^{\sharp} is a foliated vector field. By 3.4, since γ can be chosen to be any α_k , each $[\alpha_j^{\sharp}, \alpha_m^{\sharp}]$ must be a leafwise vector field and is thus 0 mod \mathcal{F} . \square

3.4. Transverse formality is not a transverse property. A property of Riemannian foliations (M, \mathcal{F}, g) is called a **transverse property (or transverse invariant)** if the property (or invariant) only depends on the metric induced on $TM/T\mathcal{F}$ (the *Riemannian foliation structure*) and not on the leafwise metric or the choice of orthogonal space $(T\mathcal{F})^\perp$. Examples of transverse properties and invariants are tautness, the basic Euler characteristic, eigenvalues of the twisted basic Laplacian (cf. [21]). The following demonstrates that transverse formality of (M, \mathcal{F}, g) depends on the entire metric g and is not simply dependent on the Riemannian foliation structure.

Proposition 3.24. *Let (M, \mathcal{F}, g) have basic harmonic mean curvature κ . Then for any other metric g' that restricts to the same metric on the normal bundle, the basic projection of the mean curvature form κ'_b is basic harmonic if and only if $\kappa'_b = \kappa$.*

Proof. Let δ_b the basic adjoint of d for the metric g ; then $\delta'_b = \delta_b + dh \lrcorner$ is the basic adjoint of d with respect to the metric g' if $\kappa'_b = \kappa + dh$; we have that h is a basic function. If κ'_b is basic g' -harmonic, then

$$\begin{aligned} 0 &= (d + \delta'_b) \kappa'_b = (d + \delta_b + dh \lrcorner) (\kappa + dh) \\ &= (d + \delta_b) dh + dh \lrcorner \kappa + dh \lrcorner dh \\ &= \Delta_b h + (dh, \kappa) + |dh|^2. \end{aligned}$$

Integrating over the manifold, we obtain (with $\langle \bullet, \bullet \rangle$ denoting the L^2 inner product)

$$\begin{aligned} 0 &= \langle \Delta_b h, 1 \rangle + \langle dh, \kappa \rangle + \langle dh, dh \rangle \\ &= \langle h, \Delta_b 1 \rangle + \langle h, \delta_b \kappa \rangle + \langle dh, dh \rangle \\ &= 0 + 0 + \|dh\|^2, \end{aligned}$$

which can happen only if h is constant and thus $\kappa'_b = \kappa$. □

The following theorem demonstrates that on taut foliations, only certain metrics can be transversely formal, and the choice depends on more than the transverse metric.

Theorem 3.25. *Let (M, \mathcal{F}, g) be a foliation of a closed manifold with a bundle-like metric that is transversely formal. If (M, \mathcal{F}) is taut, then the basic component κ_b of the mean curvature one-form must vanish.*

Proof. Suppose that (M, \mathcal{F}, g) is taut and transversely formal, and suppose that the basic component κ_b of mean curvature satisfies $\kappa_b = df$ for a basic function f . Then the transverse volume form ν has length 1, and $e^f \nu$ is basic harmonic, since $d(e^f \nu) = 0$ and

$$\begin{aligned} \delta_b(e^f \nu) &= (-\bar{*}d\bar{*} + df \lrcorner)(e^f \nu) = -\bar{*}d(e^f) + e^f df \lrcorner \nu \\ &= -e^f \bar{*}(df) + e^f df \lrcorner \nu = -e^f df \lrcorner \nu + e^f df \lrcorner \nu = 0. \end{aligned}$$

Also, $|e^f \nu| = e^f$, and by Lemma 3.4, e^f must be constant, so that $\kappa_b = 0$. □

Corollary 3.26. *Let (M, \mathcal{F}, g) be a taut, codimension q foliation of a closed manifold with a bundle-like metric that is transversely formal. Then the wedge product of any two basic harmonic forms is basic harmonic, and $\dim H^1(M, \mathcal{F}) \neq q - 1$.*

Proof. By the last theorem, the basic Laplacian Δ_b and the twisted basic Laplacian Δ_{κ_b} coincide as operators, since $\kappa_b = 0$. Then the wedge product of any two basic harmonic forms is basic harmonic, and $\bar{*}$ maps basic harmonic forms to basic harmonic forms. Then we see

that if $\alpha_1, \dots, \alpha_{q-1}$ are linearly independent basic harmonic one-forms, then $\overline{*}(\alpha_1 \wedge \dots \wedge \alpha_{q-1})$ is an additional basic harmonic one-form that is not a linear combination of the others. \square

The following results show that in general the transverse formality property depends on more than just the transverse metric.

Proposition 3.27. *Let (M, g, \mathcal{F}) be a nontaut Riemannian, transversely formal foliation that has basic harmonic mean curvature. We assume further that the leaves are not dense, and that the mean curvature has some nonzero component orthogonal to the leaf closures at some point. Then there exists a bundle-like metric g' that induces the same transverse Riemannian structure as g , such that (M, g', \mathcal{F}) is not transversely formal.*

Proof. With the given assumptions, let κ be the basic harmonic mean curvature one-form of (M, g, \mathcal{F}) , and let \overline{L} denote a leaf closure such that κ^\sharp is not contained in $T\overline{L}$ at some point x_0 of \overline{L} . Then, for $x \in M$, we let $r = r(x)$ denote the distance from x to \overline{L} , and we construct a basic function f as a function of r such that $f(x) > 0$ for $x \in \overline{L}$ and $f(x) = 0$ for $r(x) \geq \epsilon$ for a fixed, sufficiently small $\epsilon > 0$. We can then make these choices of ϵ and f such that $\kappa^\sharp(f) = (\kappa, df)$ is nonzero (and basic).

Next, let g_L and g_Q denote the restrictions of g to $T\mathcal{F}$ and to Q , respectively. We define the bundle-like metric $g' = e^{-f}g_L \oplus g_Q$. Now, let $\alpha = \kappa + dh$ be the basic harmonic representative of κ in the metric g' , for some basic function h . We note that $|\kappa|$ is constant with respect to both the metrics g and g' .

Suppose now that g' is also transversely formal, so that α has constant length (with respect to both g and g'). Then $|\alpha|^2 = |\kappa|^2 + |dh|^2 + 2(\kappa, dh)$ is constant, implying that $|dh|^2 + 2(\kappa, dh)$ is constant. At a critical point of h this constant is zero, so in fact $|dh|^2 = -2(\kappa, dh)$ on M . Integrating over (M, g) , we see that $\int_M |dh|^2 dv_g = -2 \int_M (\kappa, dh) dv_g = -2 \int_M (\delta_b \kappa, h) dv_g = 0$, so that h is constant, and $\alpha = \kappa$. Since $\delta'_b \alpha = 0$, and $\delta'_b = \delta_b + \frac{\dim \mathcal{F}}{2} df \lrcorner$, we have $0 = \delta'_b \alpha = \delta_b \kappa + \frac{\dim \mathcal{F}}{2} (\kappa, df) = \frac{\dim \mathcal{F}}{2} (\kappa, df) \neq 0$. This is a contradiction, so we have that (M, g', \mathcal{F}) is not transversely formal. \square

Remark 3.28. *Note that by examining the proof above, we can find bundle-like metrics arbitrarily close to transversely formal metrics that are not transversely formal and yet induce the same transverse metric. We see a specific example of this situation for the Carrière flow (M, \mathcal{F}, g) of Section 5.3, where multiplying the stated leafwise metric by a general positive basic function causes the transverse formality property to fail.*

The following corollary shows the existence of bundle-like metrics that are not transversely formal on all taut foliations that do not have dense leaves.

Proposition 3.29. *Let (M, g, \mathcal{F}) be a taut Riemannian, transversely formal foliation. We assume further that the leaves are not dense. Then there exists a bundle-like metric g' that induces the same transverse Riemannian structure as g , such that (M, g', \mathcal{F}) is not transversely formal.*

Proof. By Theorem 3.25, any taut, transversely formal foliation must have $\kappa_b = 0$. By multiplying the leafwise metric by any nonconstant basic function, we obtain a new bundle-like metric g' with $\kappa_b \neq 0$ and such that g and g' induce the same transverse Riemannian structure. By Theorem 3.25, (M, g', \mathcal{F}) is not transversely formal. \square

If (M, \mathcal{F}) is a foliation with dense leaves, then transverse formality is indeed a transverse property.

Proposition 3.30. *Let (M, \mathcal{F}, g) be a foliation with dense leaves with bundle-like metric that is transversely formal. Then any other metric g' that induces the same transverse metric on $TM/T\mathcal{F}$ as g is also transversely formal.*

Proof. Assume (M, \mathcal{F}, g) is as given, with g' the new metric. The mean curvature must change by $\kappa'_b = \kappa_b + dh$ for a basic function h , which is necessarily constant, so that $\kappa'_b = \kappa_b$. Note that the basic codifferential δ_b must be the same for both metrics. Thus, the basic Laplacian is the same operator on basic forms in both metrics, and thus the property of transverse formality is preserved. \square

4. TRANSVERSE FORMALITY FOR RIEMANNIAN FLOWS, K -CONTACT AND SASAKI MANIFOLDS

Riemannian flows, that is Riemannian foliations of 1-dimensional leaves, have been the object of in depth research for several decades due to numerous applications. The rich theory permits us to obtain interesting results on geometric transverse formality as well as to test some general conjectures.

Notation 4.1. *Let $(M^{q+1}, \mathcal{F}, g)$ be an ORFM. If $\dim \mathcal{F} = 1$ with the foliation be given by the integral curves of a unit vector field ξ , we say that $(M^{q+1}, \mathcal{F}, g, \xi)$ is a 1-ORFM. Throughout this section, we will also denote by $H_{a\kappa_b}^*(M, \mathcal{F})$ the basic cohomology of the flow associated with the twisted differential $d_{a\kappa_b} := d - a\kappa_b \wedge$ for $a \in \mathbb{R}$.*

4.1. General results for Riemannian flows. Recall that Rummler's formula reduces for Riemannian flows to

$$d\xi^b = -\kappa \wedge \xi^b + \varphi_0,$$

where, as before, $\varphi_0 = \sum_{j=1}^q e^j \wedge \nabla_{e_j} \xi^b \in \Gamma(\Lambda^2 Q^*)$ is the Euler form and $\kappa = (\nabla_\xi \xi)^b \in \Gamma(Q^*)$ is the mean curvature one-form. Here $\{e_j\}_{j=1, \dots, q}$ is an orthonormal frame of $\Gamma(Q)$ with corresponding coframe $\{e^j\}$. We have the following crucial lemma for Riemannian flows.

Lemma 4.2. ([22, Lemma 2.4]) *Let (M, \mathcal{F}, g, ξ) be a 1-ORFM. If the mean curvature κ of the flow is basic, then $d\varphi_0 = -\kappa \wedge \varphi_0$ and, thus, the Euler form φ_0 is a basic 2-form.*

Proof. Using Rummler's formula, we compute

$$d\varphi_0 = d(\kappa \wedge \xi^b) = d\kappa \wedge \xi^b - \kappa \wedge d\xi^b = -\kappa \wedge \varphi_0.$$

In the last equality, we used the fact that κ is a closed form, as it is basic. Hence to show that φ_0 is basic, we compute

$$\xi \lrcorner d\varphi_0 = -\xi \lrcorner (\kappa \wedge \varphi_0) = 0,$$

since φ_0 is a 2-form on Q . \square

Remark 4.3. *The metric may always be chosen so that κ is basic, by [15]. The conclusion of the Lemma above is in general false if the mean curvature is not basic.*

In the following, we compute the codifferential of the basic form φ_0 , in two different ways. We have

Proposition 4.4. *Let (M, \mathcal{F}, g, ξ) be a 1-ORFM. Assume that the mean curvature κ is basic and harmonic. Then $\delta_b \varphi_0 = (\Delta - |\varphi_0|^2 - |\kappa|^2) \xi^b$.*

Proof. Since φ_0 is a basic form and κ is basic, we know by [38, Proposition 2.4], that

$$\delta_b \varphi_0 = \delta \varphi_0 - \varphi_0 \lrcorner (\xi^b \wedge \varphi_0) = \delta d \xi^b + \delta(\kappa \wedge \xi^b) - |\varphi_0|^2 \xi^b.$$

As the flow is Riemannian, we have that $\delta \xi^b = 0$ and, thus, $\Delta \xi^b = \delta d \xi^b$. Now, an easy computation shows that

$$\delta(\kappa \wedge \xi^b) = (\delta \kappa) \xi^b + \nabla_{\xi}^M \kappa - \nabla_{\kappa^{\#}}^M \xi^b = [\xi, \kappa^{\#}]^b = -|\kappa|^2 \xi^b,$$

where we have used that $[\xi, \kappa^{\#}] = g(\nabla_{\xi}^M \kappa^{\#} - \nabla_{\kappa^{\#}}^M \xi, \xi) \xi = -|\kappa|^2 \xi$, since $\kappa^{\#}$ is basic and that $\delta \kappa = \delta_b \kappa = 0$, as κ is harmonic. Therefore, $\delta_b \varphi_0 = \Delta \xi^b - (|\kappa|^2 + |\varphi_0|^2) \xi^b$. \square

One can easily see from Lemma 4.2 and Proposition 4.4, that for a minimal Riemannian flow, the 2-form φ_0 is a basic harmonic form if and only if ξ is an eigenvector for the Laplacian Δ on M associated with the eigenvalue $|\varphi_0|^2$. In the following, we denote by ϕ the skew-symmetric endomorphism on $\Gamma(TM)$ defined by $g(\phi(\cdot), \cdot) = -g(\nabla^M \xi, \cdot)$. A simple calculation shows that $\varphi_0(\cdot, \cdot) = -2g(\phi(\cdot), \cdot)$.

Proposition 4.5. *(in [16, proof of Proposition 6.2 for minimal flows]) Let (M, \mathcal{F}, g, ξ) be a 1-ORFM, such that the mean curvature is basic. If $\text{Ric}(\xi) = \lambda \xi$ for some function λ , then $\delta_b \varphi_0 = -\kappa \lrcorner \varphi_0$.*

Proof. Fix a point of M and choose a foliated orthonormal frame $\{e_i\}_{i=1, \dots, q}$ of $\Gamma(Q)$ such that $\nabla e_i = 0$ at the point in question. Recall here that ∇ is the transversal Levi-Civita connection. For any $Y \in \Gamma(Q)$ parallel at the same point, we compute

$$\begin{aligned} 0 &= \text{Ric}(\xi, Y) = \sum_{i=1}^q R(e_i, Y, \xi, e_i) \\ &= \sum_{i=1}^q g(-\nabla_{e_i}^M \phi Y + \nabla_Y^M \phi e_i - \nabla_{[e_i, Y]}^M \xi, e_i) \\ &= \sum_{i=1}^q -g(\nabla_{e_i}^M \phi Y, e_i) - g(\phi e_i, \nabla_Y^M e_i) - g([e_i, Y], \xi) g(\kappa^{\#}, e_i), \end{aligned}$$

where we use the fact that $[e_i, Y]$ is leafwise since both e_i and Y are parallel at the point in question. Then, projecting ∇^M to Q , we get that

$$\begin{aligned} 0 &= -\sum_{i=1}^q g((\nabla_{e_i} \phi)(Y), e_i) - \sum_{i=1}^q g(\phi e_i, \nabla_Y e_i) - 2 \sum_{i=1}^q g(\phi(e_i), Y) g(\kappa^{\#}, e_i) \\ &= \frac{1}{2}(\delta_b \varphi_0)(Y) + \frac{1}{2} \varphi_0(\kappa^{\#}, Y). \end{aligned}$$

Therefore, we deduce that $\delta_b \varphi_0 = -\kappa \lrcorner \varphi_0$. \square

Combining Lemma 4.2 with Proposition 4.5, we deduce that for Riemannian flows with basic mean curvature if $\text{Ric}(\xi) = \lambda \xi$, the Euler form φ_0 is a basic $(-\kappa)$ -harmonic form. Thus φ_0 is an element of $H_{-\kappa}^2(M, \mathcal{F})$. Notice here that one can compute the function λ explicitly.

Proposition 4.6. *Let (M, \mathcal{F}, g, ξ) be a 1-ORFM, such that the mean curvature κ is basic. If $\text{Ric}(\xi) = \lambda \xi$ for some function λ , then $\lambda = -\delta_b \kappa + |\phi|^2$.*

Proof. We fix a point of M and choose a foliated orthonormal frame $\{e_i\}_{i=1,\dots,q}$ of $\Gamma(Q)$ such that $\nabla e_i = 0$ at that point. Then

$$\begin{aligned}
\lambda = \text{Ric}(\xi, \xi) &= \sum_{i=1}^q R(e_i, \xi, \xi, e_i) \\
&= \sum_{i=1}^q g(\nabla_{e_i}^M \nabla_{\xi}^M \xi - \nabla_{\xi}^M \nabla_{e_i}^M \xi - \nabla_{[e_i, \xi]}^M \xi, e_i) \\
&= \sum_{i=1}^q g(\nabla_{e_i}^M \kappa^{\#}, e_i) + g(\nabla_{\xi}^M \phi e_i, e_i) - g([e_i, \xi], \xi) g(\kappa^{\#}, e_i) \\
&= \sum_{i=1}^q g(\nabla_{e_i}^M \kappa^{\#}, e_i) - \sum_{i=1}^q g(\phi e_i, \nabla_{\xi}^M e_i) - |\kappa|^2 \\
&= -\delta_b \kappa + |\phi|^2.
\end{aligned}$$

Here, we used that $[e_i, \xi]$ is leafwise since e_i is foliated. Also, we used that $\nabla_{\xi}^M e_i = -\phi(e_i) + [e_i, \xi]$. \square

In particular, if κ is basic harmonic, then $\lambda = |\phi|^2$. We also notice that the condition $\text{Ric}(\xi) = \lambda \xi$ is not very restrictive since, for example, it is automatically satisfied for K -contact flows on a $(2m+1)$ -dimensional manifold, with $\lambda = 2m$ (see [6, Theorem 7.1]).

Lemma 4.7. *(In [16, Proposition 6.2] for minimal flows) Let (M, \mathcal{F}, g, ξ) be a 1-ORFM, such that the mean curvature κ is basic. If the Euler class $[\varphi_0] \in H_{-\kappa}^2(M, \mathcal{F})$ is nontrivial, then $H_{-\kappa}^1(M, \mathcal{F}) \simeq H_{-\kappa}^1(M)$, and $1 \leq \dim H_{-\kappa}^2(M, \mathcal{F}) \leq 1 + \dim H_{-\kappa}^2(M)$. When $\dim H_{-\kappa}^2(M, \mathcal{F}) = 1$, we get that $H_{-\kappa}^2(M) \hookrightarrow H^1(M, \mathcal{F})$.*

Proof. By the Gysin sequence [40, Proposition 2.30], we have

$$0 \rightarrow H_{-\kappa}^1(M, \mathcal{F}) \rightarrow H_{-\kappa}^1(M) \rightarrow H^0(M, \mathcal{F}) \xrightarrow{\wedge[\varphi_0]} H_{-\kappa}^2(M, \mathcal{F}) \rightarrow H_{-\kappa}^2(M) \rightarrow H^1(M, \mathcal{F}).$$

We have $H^0(M, \mathcal{F}) \cong \mathbb{R}$, generated by constants, and the map $\wedge[\varphi_0]$ is injective and thus with one-dimensional range. Then $H_{-\kappa}^1(M, \mathcal{F}) \simeq H_{-\kappa}^1(M)$, and the inequality follows immediately. When $\dim H_{-\kappa}^2(M, \mathcal{F}) = 1$, the last map in the sequence is clearly injective. \square

Remark 4.8. *One way to ensure that the Euler class is nonzero is to impose the condition $\text{Ric}(\xi) = \lambda \xi$ with $\lambda > 0$.*

Lemma 4.9. *Let (M, \mathcal{F}, g, ξ) be a 1-ORFM, such that the mean curvature κ is basic.*

(1) *If $H^r(M, \mathcal{F}) \cong 0$ for some integer r , then*

$$H_{-\kappa}^{r+2}(M, \mathcal{F}) \hookrightarrow H_{-\kappa}^{r+2}(M),$$

and $\dim H_{-\kappa}^{r+1}(M, \mathcal{F}) \geq \dim H_{-\kappa}^{r+1}(M)$. We also have that

$$\dim H_{-\kappa}^{r+2}(M) \leq \dim H_{-\kappa}^{r+2}(M, \mathcal{F}) + \dim H^{r+1}(M, \mathcal{F}).$$

(2) *If $H_{-\kappa}^{r+1}(M) \cong 0$ and $H_{-\kappa}^{r+2}(M) \cong 0$ for some integer r , then*

$$H^r(M, \mathcal{F}) \simeq H_{-\kappa}^{r+2}(M, \mathcal{F}).$$

Proof. (1) By the Gysin sequence in [40, Proposition 2.30],

$$H_{-\kappa}^{r+1}(M, \mathcal{F}) \rightarrow H_{-\kappa}^{r+1}(M) \rightarrow H^r(M, \mathcal{F}) \xrightarrow{\wedge[\varphi_0]} H_{-\kappa}^{r+2}(M, \mathcal{F}) \hookrightarrow H_{-\kappa}^{r+2}(M) \rightarrow H^{r+1}(M, \mathcal{F})$$

We have that the map $\wedge[\varphi_0]$ above is the zero map. Thus the following map is injective. The statement (2) follows from the same sequence. \square

Theorem 4.10. *Let (M, \mathcal{F}, g, ξ) be a 1-ORFM, such that the flow is minimal. Suppose that $\text{Ric}(\xi) = \lambda\xi$ for some function $\lambda > 0$. Assume that the flow is transversely formal and that $H^r(M, \mathcal{F}) = 0$ for some r . Then any basic harmonic $(r+2)$ -form is harmonic. If moreover $\dim H^1(M, \mathcal{F}) \geq 1$, then we have*

$$\dim H^{r+1}(M) \leq \dim H^{r+1}(M, \mathcal{F}) \leq \dim H^{r+2}(M).$$

Proof. Let α be a basic harmonic $(r+2)$ -form. Since the flow is minimal, the Euler form φ_0 is a basic harmonic 2-form, by Proposition 4.5. By the transverse formality assumption, the form $\varphi_0 \lrcorner \alpha$ is a basic harmonic r -form and is zero, since $H^r(M, \mathcal{F}) = 0$. Now,

$$\delta\alpha = \delta_b\alpha + \varphi_0 \lrcorner (\xi \wedge \alpha) = \delta_b\alpha = 0.$$

Hence α is harmonic. To prove the second part, we consider the map $H^{r+1}(M, \mathcal{F}) \rightarrow H^{r+2}(M); \alpha \mapsto \theta^i \wedge \alpha$ where $\{\theta^i\}$ is a basis of $H^1(M, \mathcal{F})$. This map is clearly injective, since $\theta^i \lrcorner \alpha$ is an element in $H^r(M, \mathcal{F}) = 0$. Hence $\dim H^{r+1}(M, \mathcal{F}) \leq \dim H^{r+2}(M)$. The other part of the inequality comes from the first part of Lemma 4.9. \square

4.2. Sasakian manifolds. We now consider Sasakian manifolds, which are special cases of isometric flows with very particular transverse geometric structure, namely a Kähler metric.

We will use the following setup. Let $(M^{2n+1}, \mathcal{F}, g, \xi)$ be a 1-ORFM that is a minimal flow, and we let

$$\begin{aligned} \eta &= \xi^\flat \\ d\eta &= \varphi_0. \end{aligned}$$

The endomorphism $\phi : TM \rightarrow TM$ and metric g are chosen so that

$$\begin{aligned} \varphi_0(X, Y) &= 2g(X, \phi Y) \\ \phi^2 &= -I + \eta \otimes \xi \\ g(\phi X, \phi Y) &= g(X, Y) - \eta(X)\eta(Y) \end{aligned}$$

for $X, Y \in \Gamma(TM)$. The normality condition for Sasakian manifolds is

$$(\nabla_X^M \phi)(Y) = g(X, Y)\xi - \eta(Y)X.$$

Note that this condition implies our formula for ϕ in the last section:

$$\phi(\bullet) = -\nabla_{\bullet}^M \xi.$$

Remark 4.11. *We are using notation consistent with that used in, for example, [19] and [11]. Other researchers use metrics that differ from ours by a constant, such as [6].*

Theorem 4.12. *Let (M, g) be a closed Sasakian manifold of dimension $2n+1$ with Reeb vector field ξ . Let \mathcal{F} be the corresponding Riemannian flow defined by ξ . If (M, g) is geometrically formal, then (M, \mathcal{F}, g) is transversely formal.*

Proof. Suppose (M, g) is geometrically formal. By [20, Theorem 1.2], M is a rational homology sphere, so that $H^r(M) \cong 0$ for $1 \leq r < n$. By [6, Theorem 7.1], K -contact flows satisfy the Ricci curvature condition in Lemma 4.5, and that includes Sasakian manifolds. Then, by Lemma 4.7, $H^1(M, \mathcal{F}) \simeq 0$, and $H^2(M, \mathcal{F}) \simeq \mathbb{R}$. Using Lemma 4.9, we deduce that $H^r(M, \mathcal{F}) \simeq 0$ for r odd and that $H^r(M, \mathcal{F}) \simeq \mathbb{R}$ for r even. Then M is transversely formal. \square

Remark 4.13. *The converse of the statement in Theorem 4.12 is in general false. From [11, Theorem 1.1], any quotient of a Heisenberg group $H(1, 2n)$ by a cocompact discrete subgroup Γ admits a canonical Sasakian structure. In fact, these are the only compact nilmanifolds that admit Sasakian structures and are metric fibrations over a flat torus. Hence, the first cohomology group $H^1(M)$ is isomorphic to $H^1(\mathbb{T}^{2n}) \simeq \mathbb{R}^{2n}$. Thus, by [20, Theorem 1.2], none of these manifolds are geometrically formal but they are transversely formal, since the basic harmonic forms are the pullbacks of the harmonic forms on that flat torus.*

In the following theorem, we show that transverse formality of closed Sasakian manifolds implies some formality properties of wedge products of harmonic forms of specific degrees.

Theorem 4.14. *Let (M, g) be a closed Sasakian manifold of dimension $2n + 1$ with Reeb vector field ξ . Let \mathcal{F} be the corresponding Riemannian flow defined by ξ . If (M, \mathcal{F}, g) is transversely formal, then*

- (1) *The wedge product of two harmonic forms respectively of degrees less than n and greater than $n + 1$ is harmonic.*
- (2) *Assume that $H^r(M, \mathcal{F}) = 0$ for some integer r . The wedge product of two harmonic forms respectively of degrees p and q with $p + q = r + 2$ and $p, q \leq n$ is harmonic.*
- (3) *Assume that $H^r(M, \mathcal{F}) = 0$ for some integer r . The wedge product of two harmonic forms respectively of degrees p and q with $p + q = 4n - r$ and $p, q \geq n + 1$ is harmonic.*

Proof. Let α and β be harmonic forms of degree $\leq n$. Then [43, Theorem 4.1] implies that α, β are basic forms, and

$$\varphi_0 \lrcorner \alpha = 0, \quad \varphi_0 \lrcorner \beta = 0.$$

Then by [38, Proposition 2.4],

$$\delta_b \alpha = \delta \alpha - \varphi_0 \lrcorner (\xi^\flat \wedge \alpha) = \delta \alpha = 0.$$

The same is true for β , so that α and β are basic harmonic. By transverse formality, $\alpha \wedge \beta$ is basic harmonic, and also $\alpha \lrcorner \beta$ is basic harmonic. Now, we have

$$\delta(\alpha \lrcorner \beta) = \delta_b(\alpha \lrcorner \beta) + \varphi_0 \lrcorner (\xi^\flat \wedge (\alpha \lrcorner \beta)) = \delta_b(\alpha \lrcorner \beta) = 0.$$

Thus, $\alpha \lrcorner \beta$ is harmonic. Next, suppose α is a harmonic form of degree $\leq n$, and suppose that β is a harmonic form of degree $\geq n + 1$. Then $*\beta$ has degree $\leq n$, then $\alpha \lrcorner *\beta$ is harmonic by the previous argument. But $\alpha \lrcorner *\beta = \pm *(\alpha \wedge \beta)$, so $\alpha \wedge \beta$ is harmonic. This proves the first part of the theorem. To prove the second part, for α and β of degrees less than n , the form $\alpha \wedge \beta$ is basic harmonic by the argument above. If the degree of $\alpha \wedge \beta$ is $r + 2$, then Proposition 4.10 yields the second part. The last statement follows from the second part by using the Hodge star operator. \square

4.3. Classification of three-dimensional Riemannian flows that are transversely formal. In this part, we are going to classify three-dimensional Riemannian flows that are transversely formal. For this, consider a 1-ORFM of three-manifold M . If the flow is nontaut, then by Carrière classification [12, Theorem III.A.1], it is foliated diffeomorphic to a hyperbolic torus. The standard metric on these manifolds is transversely formal (see Section 4.5). Hence, we are left with taut Riemannian flows. But Theorem 3.25 tells us that for such flows, transverse formality implies that the basic component of the mean curvature of the flow vanishes. Again by Carrière classification, the manifold M is diffeomorphic to one of the following:

- \mathbb{T}^3 and the flow is generated by a linear flow on \mathbb{T}^3 .
- $\mathbb{S}^1 \times \mathbb{S}^2$ and the flow is generated by an irrational rotation suspension.
- $L_{p,q}$ (Lens space) and the flow has 2 closed leaves and is generated by the flow on \mathbb{S}^3 as described in [24].
- Seifert fibration.

Let us treat each case separately. When M is \mathbb{T}^3 (resp. $\mathbb{S}^1 \times \mathbb{S}^2$), the formality of the metric in \mathbb{T}^3 (resp. $\mathbb{S}^1 \times \mathbb{S}^2$) gives transverse formality by Proposition 3.18. When M is the Lens space, the computations of the mean curvature done in [24] shows that such flow is minimal if and only if it is a Hopf fibration which is not possible because of the closedness of the orbits. We are left with the Seifert fibration. For this particular case, we will assume furthermore that $\text{Ric}(\xi) = \lambda\xi$ with $\lambda = |\phi|^2 > 0$. By Lemma 4.5, φ_0 is basic harmonic and transverse formality gives that φ_0 has constant norm, so that $|\varphi_0|^2 = 2|\phi|^2$ is a positive constant. Recall here $\phi = -\nabla^M \xi$ is the skew-symmetric endomorphism on $\Gamma(TM)$. In dimension three and since $\phi(\xi) = 0$, we can write ϕ as the transformation associated to the matrix $\begin{pmatrix} 0 & -b \\ b & 0 \end{pmatrix}$ with b constant. If we rescale the Riemannian metric, the manifold $(M, \mathcal{F}, b^2g, \frac{1}{b}\xi)$ is Sasakian, and the matrix of the new ϕ is $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, a complex structure. From [5, Theorem 8] and its restatement in [10, Section 7], either M is a quotient of a round 3-sphere, a Heisenberg 3-manifold, or $\widetilde{SL}(2, \mathbb{R})/\Gamma$ with Γ a discrete subgroup of the connected component of the isometry group of $\widetilde{SL}(2, \mathbb{R})$ in the natural metric. Observe that quotients of the round sphere and Heisenberg 3-manifolds are necessarily transversely formal, but quotients of the type $\widetilde{SL}(2, \mathbb{R})/\Gamma$ have $\dim H^1(M, \mathcal{F}) > 2$ and thus are not transversely formal.

Theorem 4.15. *Suppose that (M, \mathcal{F}, g, ξ) is a 1-ORFM on a 3-manifold. Assume that M is transversely formal. Then M is either \mathbb{T}^3 , $\mathbb{S}^1 \times \mathbb{S}^2$, the hyperbolic torus or a Seifert fibration. Assume moreover that $\text{Ric}(\xi) = \lambda\xi$ with $\lambda > 0$, then up to a rescaling of the metric and the flow, M is a quotient of \mathbb{S}^3 or is a Heisenberg 3-manifold.*

5. EXAMPLES OF RIEMANNIAN FOLIATIONS WITH TRANSVERSE FORMALITY

5.1. Geometric formality and transverse formality. The following examples illustrate that transverse formality and geometric formality need not coexist for metrics on foliations.

Example 5.1. *In this example, the manifold (M, g) is formal, but the Riemannian foliation (M, \mathcal{F}, g) is not transversely formal. In [32, Theorem 24], Kotschick and Terzic showed that the example introduced by Totaro in [45, Section 1], which is a biquotient of $\mathbb{S}^3 \times \mathbb{S}^3 \times \mathbb{S}^3$ with the standard metric, is not geometrically formal. This gives a Riemannian foliation*

by the fibers of this submersion, such that the total space is formal, but the foliation is not transversely formal.

We already mentioned in Remark 4.13 that the central foliation for some Heisenberg manifolds is transversely formal, but the metric on the manifold itself is not formal. In fact, from [23], the only formal nilmanifolds are tori. We give another simple example here.

Example 5.2. *In this example, the Riemannian foliation (M, \mathcal{F}, g) is transversely formal, but the manifold (M, g) is not formal for any bundle-like metric. Let H be a closed hyperbolic surface, which has first Betti number at least 4. Let $M = H \times \mathbb{S}^1$, which has first Betti number at least 5. Then the codimension one foliation of M with leaves of the form $H \times \{\theta\}$ is clearly Riemannian and transversely formal for the product metric. However, by [30, Theorem 6], the first Betti number of a geometrically formal 3-manifold must be 0, 1, or 3; thus the manifold M is not formal for any metric.*

5.2. The Hopf fibration of \mathbb{S}^3 . Consider the sphere $M = \mathbb{S}^3(r) \subseteq \mathbb{R}^4$ of radius r with metric induced from the standard Euclidean metric in coordinates (x, y, z, w) . We consider the one-dimensional foliation \mathcal{F} determined by the unit tangent $\xi = \frac{1}{r}(x\partial_y - y\partial_x + z\partial_w - w\partial_z)$, so that the characteristic (leafwise volume) form is $\xi^\flat = \frac{1}{r}(xdy - ydx + zdw - wdz)$, by which we really mean $\frac{1}{r}i^*(xdy - ydx + zdw - wdz)$, where $i : \mathbb{S}^3(r) \rightarrow \mathbb{R}^4$ is the inclusion. That means we are considering equivalence classes of forms, so that for example $x^2 + y^2 + z^2 + w^2 = r^2$ and $xdx + ydy + zdz + wdw = 0$. We consider $e^1 = \frac{1}{r}(xdz - zdx + wdy - ydw)$ and $e^2 = \frac{1}{r}(ydz - zdy + xdw - wdx)$, unit one-forms on $\mathbb{S}^3(r)$. We check that ξ^\flat, e^1, e^2 is an oriented orthonormal basis of T^*M at each point, and we choose the orientation so that $*\xi^\flat = e^1 \wedge e^2$. A computation shows that

$$(d\xi^\flat, e^1 \wedge e^2) = \frac{2}{r^3}(x^2 + y^2 + z^2 + w^2) = \frac{2}{r},$$

so that

$$d\xi^\flat = \varphi_0 = \frac{2}{r}e^1 \wedge e^2.$$

From Rummel's formula, we realize that this means that \mathcal{F} is minimal (i.e. geodesic) and that the normal bundle is not integrable.

$$\begin{aligned} \Delta\xi^\flat &= \delta d\xi^\flat = \frac{2}{r} * d * (e^1 \wedge e^2) = \frac{2}{r} * d\xi^\flat \\ &= \frac{4}{r^2} * (e^1 \wedge e^2) = \frac{4}{r^2} \xi^\flat, \end{aligned}$$

so that ξ^\flat (and thus $d\xi^\flat = \varphi_0 = \frac{2}{r}e^1 \wedge e^2$) is an eigenvalue of the Laplacian corresponding to eigenvalue $\frac{4}{r^2}$. We note also that as a basic form, φ_0 is closed, so that by using the transverse star operator and the fact that \mathcal{F} is minimal,

$$\delta_b \varphi_0 = -\bar{*}d\bar{*}(\varphi_0) = -\bar{*}d\bar{*}\left(\frac{2}{r}e^1 \wedge e^2\right) = -\frac{2}{r}\bar{*}d\xi^\flat = 0,$$

so that φ_0 is basic harmonic. We see that $|\varphi_0|^2 = \frac{4}{r^2}$, a constant. In fact, $\frac{r^2}{4}\varphi_0$ is the transverse volume form of (M, \mathcal{F}) .

We see that the basic cohomology $H^k(M, \mathcal{F})$ is 1 in degree 0 and 2 (generated by 1, φ_0 , respectively), and since $M/\mathcal{F} \cong \mathbb{S}^2$, we get that $H^1(M, \mathcal{F}) \cong 0$. Thus, we have that the metric on M is both formal and transversely formal, and our calculations are consistent with Proposition 3.18 and with Theorem 4.12, as $r = 1$ corresponds to the Sasaki metric.

5.3. The Carrière example. We will compute the various types of harmonic forms of the Carrière example from [12] in the 3-dimensional case. Let A be a matrix in $\mathrm{SL}_2(\mathbb{Z})$ of trace strictly greater than 2. We denote respectively by V_1 and V_2 the eigenvectors associated with the eigenvalues λ and $\frac{1}{\lambda}$ of A with $\lambda > 1$ irrational. Let the hyperbolic torus $M = \mathbb{T}_A^3$ be the quotient of $\mathbb{T}^2 \times \mathbb{R}$ by the equivalence relation which identifies (m, t) to $(A(m), t + 1)$. We choose the bundle-like metric (letting (x, s, t) denote the local coordinates in the V_2 direction, V_1 direction, and \mathbb{R} direction, respectively) as

$$g = \lambda^{-2t} dx^2 + \lambda^{2t} ds^2 + dt^2.$$

First, we notice that the mean curvature of the flow is $\kappa = \kappa_b = \log(\lambda) dt$, since $\chi_{\mathcal{F}} = \lambda^{-t} dx$ is the characteristic form and $d\chi_{\mathcal{F}} = -\log(\lambda) \lambda^{-t} dt \wedge dx = -\kappa \wedge \chi_{\mathcal{F}}$. Since $\varphi_0 = 0$, the normal bundle is involutive, and we can see this as the resulting foliation is given by the (y, t) coordinates. The transverse volume form is $\nu = \lambda^t ds \wedge dt = d\left(\frac{1}{\log \lambda} \lambda^t ds\right)$, and we see that $d\nu = 0 = (d - \kappa \wedge) \nu$. Since the flow is nontaut, this is consistent with the fact that $H^0(M, \mathcal{F}) \cong H_\kappa^2(M, \mathcal{F}) \cong \mathbb{R}$. The cohomology groups of M satisfy $H^j(M) \cong \mathbb{R}$, and the generating cohomology classes are generated by $1, dt, dx \wedge ds, dx \wedge ds \wedge dt$; all are clearly closed. We note that the codifferential δ satisfies $\delta = (-1)^k * d*$ on k -forms, so that we can check if the forms are harmonic by evaluating $d*$. We see that $d(*dt) = d(dx \wedge ds) = 0$, $d*(dx \wedge ds) = d(dt) = 0$, $d*(dx \wedge ds \wedge dt) = d(1) = 0$, so the generators mentioned are all harmonic. The generators are also constant length, and the wedge product of any two harmonic forms is harmonic, so that (M, g) is geometrically formal. Next, observe that the basic cohomology groups satisfy $H^0(M, \mathcal{F}) \cong H^1(M, \mathcal{F}) \cong \mathbb{R}$, with generators 1 and dt , both clearly closed. Since δ_b is a restriction of δ when the mean curvature is basic and the normal bundle is involutive, we see that both 1 and dt are basic harmonic forms. Also, $H_\kappa^1(M, \mathcal{F}) = H_\kappa^2(M, \mathcal{F}) \cong \mathbb{R}$, and they are generated by the basic κ_b -harmonic forms $\bar{*}dt = \lambda^t ds$ and $\bar{*}1 = \nu = \lambda^t ds \wedge dt$. We see that wedge product of basic harmonic and basic κ -harmonic forms is basic κ -harmonic, because the only nontrivial case is $(dt) \wedge (\lambda^t ds) = \nu$, so (M, \mathcal{F}, g) is both formal and transversely formal. This example demonstrates Proposition 3.18.

5.4. Solvmanifold admitting transversely formal Riemannian foliations. We consider the example given in [13] and in [3], and we construct three Riemannian foliations on this manifold, all of which are transversely formal. We consider the Lie group G_k of matrices

$$\begin{pmatrix} e^{kz} & 0 & 0 & x \\ 0 & e^{-kz} & 0 & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

all variables in \mathbb{R} , with k fixed, $e^k + e^{-k} = 2 \cosh(k) \in \mathbb{Z} \setminus \{2\}$. This forms a connected solvable Lie group with basis $\{dx - kxdz, dy + kydz, dz\}$ of right-invariant forms.

Let $\Gamma_k < G_k$ be a discrete subgroup such that $N_k = G_k / \Gamma_k$ is compact. One such example (adapted from an example in [9]) is defined using $\tau = e^k = \frac{3+\sqrt{5}}{2}$; $2 \cosh k = 3$, which satisfies $\tau^2 - 3\tau + 1 = 0$. We let

$$\Gamma_k = \left\{ \left(\begin{array}{cccc} \tau^s & 0 & 0 & n + m\tau \\ 0 & \tau^{-s} & 0 & n + m\tau^{-1} \\ 0 & 0 & 1 & s \\ 0 & 0 & 0 & 1 \end{array} \right) : s, n, m \in \mathbb{Z} \right\}$$

We check that if $A(s, n, m)$ is the matrix above, $A(s_1, n_1, m_1) A(s_2, n_2, m_2) = A(s_1 + s_2, a, b)$, where a and b are specific integers, found by rewriting τ^s as an integer linear combination of 1 and τ and verifying that τ^{-s} is the same linear combination of 1 and τ^{-1} . Thus Γ is indeed a discrete subgroup, and it can be checked that G_k/Γ_k is compact.

The basis $\{dx - kxdz, dy + kydz, dz\}$ descends to a global basis of one-forms $\{\alpha, \beta, \gamma\}$ over N_k . The group $H^1(N_k)$ is one dimensional with generator $[\gamma]$, and $H^2(N_k)$ is also one-dimensional with generator $[\alpha \wedge \beta]$.

Let $\lambda \in \mathbb{R}$ be a number such that $\lambda[\alpha \wedge \beta] \in H^2(N_k, \mathbb{Z})$. For nonzero $n \in \mathbb{Z}$, let $M_{n,k}$ be the total space of a principal \mathbb{S}^1 -bundle over N_k with Chern class $n\lambda[\alpha \wedge \beta]$; such a bundle exists by the result in [29]. For simplicity, we denote the pullbacks of forms on N_k to $M_{n,k}$ by the same notations.

Let η be a connection form with curvature

$$d\eta = n\lambda\alpha \wedge \beta.$$

The set $\{\alpha, \beta, \gamma, \eta\}$ forms a global basis of one-forms on the 4-dimensional manifold $M_{n,k}$. We consider the metric which makes this basis orthonormal.

Relations between these forms from [3]:

$$\begin{aligned} d\alpha &= -k\alpha \wedge \gamma \\ d\beta &= k\beta \wedge \gamma \\ d\gamma &= 0 \\ d\eta &= n\lambda\alpha \wedge \beta. \end{aligned} \tag{5.1}$$

We see $d(*\gamma) = d(\alpha \wedge \beta \wedge \eta) = d(\alpha \wedge \beta) \wedge \eta$, and $d(\alpha \wedge \beta) = 0$, so that $d(*\gamma) = 0$, so γ is the harmonic form that generates $H^1(M_{n,k}) \cong \mathbb{R}$, and thus $*\gamma = \alpha \wedge \beta \wedge \eta$ is the harmonic form that generates $H^3(M_{n,k}) \cong \mathbb{R}$. We check that $H^2(M_{n,k}) \cong 0$, because although $\alpha \wedge \beta$ is closed, $d(\frac{1}{n\lambda}\eta) = \alpha \wedge \beta$, and similarly all other closed two-forms are exact. Thus, we easily check that the wedge product of any two harmonic forms is harmonic, so that $M_{n,k}$ is geometrically formal.

Let $\omega = k\gamma$ and $\omega' = -k\gamma$. The forms $\alpha, \alpha \wedge \eta, \alpha \wedge \gamma, \alpha \wedge \gamma \wedge \eta$ are ω -harmonic,

$$d_\omega \left(\frac{1}{2k}\beta \right) = \beta \wedge \gamma,$$

so the latter is d_ω -exact. In fact, these four ω -harmonic forms generate all of the cohomology groups H_ω^* . We have $d_\omega(\alpha \wedge \eta) = 0 = \delta_\omega(\alpha \wedge \eta)$, and also $\beta, \beta \wedge \eta, \beta \wedge \gamma, \beta \wedge \gamma \wedge \eta$ are ω' -harmonic and generate $H_{\omega'}^*$, and

$$d_{\omega'} \left(-\frac{1}{2k}\alpha \right) = \alpha \wedge \gamma,$$

so the latter is $d_{\omega'}$ -exact.

The dual vectors to the orthonormal basis of one-forms are (in order) A, B, C, H . Then we have (for example $n\lambda = d\eta(A, B) = -\eta([A, B])$)

$$\begin{aligned}
[A, C] &= kA \\
[A, B] &= -n\lambda H \\
[B, C] &= -kB \\
[A, H] &= [B, H] = [C, H] = 0.
\end{aligned} \tag{5.2}$$

We note that a distribution L of $TM_{n,k}$ is the tangent bundle to a Riemannian foliation with bundle-like metric if and only if L is involutive and

$$\langle [X, V], Y \rangle + \langle X, [Y, V] \rangle + V \langle X, Y \rangle = 0 \tag{5.3}$$

for all $V \in \Gamma L$, $X, Y \in \Gamma L^\perp$ (see [44, (5.13), Theorem 5.17]). The only possible rank one distribution with these properties is $L = \text{span} \{H\}$. We check that $\langle [A, H], B \rangle + \langle A, [B, H] \rangle = 0$, $\langle [A, H], C \rangle + \langle A, [C, H] \rangle = 0$, $\langle [B, H], C \rangle + \langle B, [C, H] \rangle = 0$, etc., producing a Riemannian flow. For this flow, η is the characteristic form, and $d\eta = n\lambda\alpha \wedge \beta$, so the Euler form is $n\lambda\alpha \wedge \beta$, and the mean curvature is zero, so this is an isometric flow with normal bundle that is not involutive.

Now we compute $H^*(M_{n,k}, \mathcal{F})$. We have $H^0(M_{n,k}, \mathcal{F}) \cong \mathbb{R}$ (generated by constants), $H^3(M_{n,k}, \mathcal{F}) \cong \mathbb{R}$ (generated by $\alpha \wedge \beta \wedge \gamma$). From (5.1), the only closed one-forms mod exact forms are multiples of γ , and $\delta_b \gamma = 0$, so γ is basic harmonic. Thus $H^1(M_{n,k}, \mathcal{F}) \cong \mathbb{R}$. Then we also have that $H^2(M_{n,k}, \mathcal{F}) \cong \mathbb{R}$, generated by $\bar{*}\gamma = \alpha \wedge \beta$, which must also be basic harmonic. We see that the wedge product of any basic harmonic form (linear combination of $1, \gamma, \alpha \wedge \beta, \alpha \wedge \beta \wedge \gamma$) with any other is also basic harmonic. Thus, $(M_{n,k}, \mathcal{F})$ is transversely formal in this metric.

From (5.2) and (5.3), the only possible two-dimensional Riemannian foliations are spanned by $\{A, H\}$ and $\{B, H\}$. We consider the codimension two foliation with $T\mathcal{F}_2 = \text{span} \{B, H\}$. Then the characteristic form of this foliation is $\beta \wedge \eta$, and $d(\beta \wedge \eta) = -k\gamma \wedge (\beta \wedge \eta)$, so the mean curvature is $\kappa = \kappa_b = k\gamma$, which is not exact, and the normal bundle is involutive. The codimension two foliation is Riemannian with $H^0(M_{n,k}, \mathcal{F}_2) \cong \mathbb{R}$, $H^2(M_{n,k}, \mathcal{F}_2) \cong 0$, $H_\kappa^0(M_{n,k}, \mathcal{F}_2) \cong 0$, $H_\kappa^2(M_{n,k}, \mathcal{F}_2) \cong \mathbb{R}$, spanned by the basic κ -harmonic volume form $\nu = -\alpha \wedge \gamma = d(\frac{1}{k}\alpha)$. To check $H^1(M_{n,k}, \mathcal{F}_2)$, we see from (5.1) that multiples of γ are the only closed basic forms mod exact forms, and

$$\delta_b \gamma = -\bar{*}d\bar{*}\gamma + \kappa_b \lrcorner \gamma = -\bar{*}d(\alpha) + k = -\bar{*}(-k\alpha \wedge \gamma) + k = -k + k = 0,$$

so we see that $H^1(M_{n,k}, \mathcal{F}_2)$ is spanned by γ , which is basic harmonic (and thus κ is basic harmonic). Then we see that $H_\kappa^1(M_{n,k}, \mathcal{F}_2) \cong \mathbb{R}$, and this is spanned by the basic κ -harmonic form $\bar{*}\gamma = \alpha$. We see in this case that the wedge product of any basic harmonic form (combination of 1 and γ) with any basic κ -harmonic form (combination of α and $\gamma \wedge \alpha$) is also basic κ -harmonic. Therefore, we have transverse formality. We can see that if we define another codimension two Riemannian foliation by the span of $\{A, H\}$, the results would be similar.

The only codimension three Riemannian foliation has tangent bundle $T\mathcal{F}_3 = \ker \gamma$, which is spanned by $\{A, B, H\}$. Since $|\gamma| = 1$, the foliation is minimal, which we can also check by $d(\alpha \wedge \beta \wedge \eta) = 0$. The basic cohomology injects into ordinary cohomology, and we see that $H^1(M_{n,k}, \mathcal{F}_3)$ is generated by the basic harmonic form γ . We clearly have transverse formality in this case (always true for codimension one foliations).

6. APPENDIX

In this section, we construct a 7-dimensional solvmanifold with an interesting 3-dimensional nontaut Riemannian foliation with transversely formal metric and for which the normal bundle is not involutive. Let $G = \{M(x, y, t, a, b, s, z) : (x, y, t, a, b, s, z) \in \mathbb{R}^7\}$ denote the solvable Lie group of matrices defined below, which implicitly depends on the nonzero integers n_1, n_2 and nonzero real numbers k_1, k_2 that are chosen such that $e^{k_1} + e^{-k_1} = 2 \cosh(k_1) = K_1 \in \mathbb{Z}_{>3}$, $e^{k_2} + e^{-k_2} = 2 \cosh(k_2) = K_2 \in \mathbb{Z}_{>3}$. We let $\kappa_1 = e^{k_1}$, $\kappa_2 = e^{k_2}$, so that each κ_j satisfies $k^2 - K_j k + 1 = k^{-2} - K_j k^{-1} + 1 = 0$. We define the 9×9 matrix $M(x, y, t, a, b, s, z)$ by

$$\begin{aligned} M(x, y, t, a, b, s, z)_{11} &= e^{k_1 z} \\ M(x, y, t, a, b, s, z)_{33} &= e^{-k_1 z} \\ M(x, y, t, a, b, s, z)_{44} &= e^{k_2 z} \\ M(x, y, t, a, b, s, z)_{66} &= e^{-k_2 z} \\ M(x, y, t, a, b, s, z)_{22} &= M(x, y, t, a, b, s, z)_{55} = 1 \\ M(x, y, t, a, b, s, z)_{77} &= M(x, y, t, a, b, s, z)_{88} = 1 \\ M(x, y, t, a, b, s, z)_{21} &= -n_1 y e^{k_1 z} \\ M(x, y, t, a, b, s, z)_{45} &= -n_2 b e^{k_2 z} \\ M(x, y, t, a, b, s, z)_{\bullet 9} &= (x, t, y, a, s, b, z, 0, 1)^T, \end{aligned}$$

with all other entries zero.

A calculation shows that

$$\begin{aligned} &M(x_1, y_1, t_1, a_1, b_1, z_1, s_1) M(x_2, y_2, t_2, a_2, b_2, s_2, z_2) \\ &= M(x_1 + x_2 e^{k_1 z_1}, y_1 + y_2 e^{-k_1 z_1}, t_1 + t_2 - n_1 y_1 x_2 e^{k_1 z_1}, a_1 + a_2 e^{k_2 z_1}, \\ &\quad b_1 + b_2 e^{-k_2 z_1}, s_1 + s_2 - n_2 b_1 a_2 e^{k_2 z_1}, z_1 + z_2). \end{aligned}$$

From this we see that the left-invariant vector fields are such that for $g \in M$, $(L_g)_* \partial_x = X = e^{k_1 z} \partial_x - n_1 y e^{k_1 z} \partial_t$, $(L_g)_* \partial_y = Y = e^{-k_1 z} \partial_y$, $(L_g)_* \partial_t = T = \partial_t$, $(L_g)_* \partial_a = A = e^{k_2 z} \partial_a - n_2 b e^{k_2 z} \partial_s$, $(L_g)_* \partial_b = B = e^{-k_2 z} \partial_b$, $(L_g)_* \partial_s = S = \partial_s$, $(L_g)_* \partial_z = Z = \partial_z$.

The corresponding global dual basis of left-invariant one-forms is $\xi = e^{-k_1 z} dx$, $v = e^{k_1 z} dy$, $\tau = dt + n_1 y dx$, $\alpha = e^{-k_2 z} da$, $\beta = e^{k_2 z} db$, $\sigma = ds + n_2 b da$, $\zeta = dz$.

We have the relations

$$\begin{aligned} d\xi &= k_1 \xi \wedge \zeta, & d\alpha &= k_2 \alpha \wedge \zeta, \\ dv &= -k_1 v \wedge \zeta, & d\beta &= -k_2 \beta \wedge \zeta, \\ d\tau &= -n_1 dx \wedge dy = -n_1 \xi \wedge v, & d\sigma &= -n_2 da \wedge db = -n_2 \alpha \wedge \beta, \\ d\zeta &= 0. \end{aligned} \tag{6.1}$$

The corresponding bracket relations are:

$$\begin{aligned} [X, Z] &= k_1 X & [A, Z] &= k_2 A \\ [Y, Z] &= -k_1 Y & [B, Z] &= -k_2 B \\ [X, Y] &= -n_1 T & [A, B] &= -n_2 S \end{aligned}$$

with all other brackets zero. The differentials of two-forms are as follows:

$$\begin{aligned}
d(\xi \wedge v) &= 0, & d(\alpha \wedge \beta) &= 0, \\
d(v \wedge \tau) &= -k_1 v \wedge \zeta \wedge \tau, & d(\beta \wedge \sigma) &= -k_2 \beta \wedge \zeta \wedge \sigma, \\
d(\xi \wedge \tau) &= k_1 \xi \wedge \zeta \wedge \tau, & d(\alpha \wedge \sigma) &= k_2 \alpha \wedge \zeta \wedge \sigma, \\
d(\tau \wedge \zeta) &= -n_1 \xi \wedge v \wedge \zeta, & d(\sigma \wedge \zeta) &= -n_2 \alpha \wedge \beta \wedge \zeta, \\
d(\tau \wedge \alpha) &= -n_1 \xi \wedge v \wedge \alpha - k_2 \tau \wedge \alpha \wedge \zeta & d(\xi \wedge \sigma) &= k_1 \xi \wedge \zeta \wedge \sigma + n_2 \xi \wedge \alpha \wedge \beta \\
d(\tau \wedge \beta) &= -n_1 \xi \wedge v \wedge \beta + k_2 \tau \wedge \beta \wedge \zeta & d(v \wedge \sigma) &= -k_1 v \wedge \zeta \wedge \sigma + n_2 v \wedge \alpha \wedge \beta \\
d(\tau \wedge \sigma) &= -n_1 \xi \wedge v \wedge \sigma + n_2 \tau \wedge \alpha \wedge \beta & d(\xi \wedge \zeta) &= 0 \\
d(\xi \wedge \alpha) &= k_1 \xi \wedge \zeta \wedge \alpha - \xi \wedge k_2 \alpha \wedge \zeta = (-k_1 - k_2) \xi \wedge \alpha \wedge \zeta \\
d(\xi \wedge \beta) &= k_1 \xi \wedge \zeta \wedge \beta + \xi \wedge k_2 \beta \wedge \zeta = (-k_1 + k_2) \xi \wedge \beta \wedge \zeta \\
d(v \wedge \alpha) &= -k_1 v \wedge \zeta \wedge \alpha - v \wedge k_2 \alpha \wedge \zeta = (k_1 - k_2) v \wedge \alpha \wedge \zeta \\
d(v \wedge \beta) &= -k_1 v \wedge \zeta \wedge \beta + v \wedge k_2 \beta \wedge \zeta = (k_1 + k_2) v \wedge \beta \wedge \zeta \\
d(v \wedge \zeta) &= 0 \\
d(\alpha \wedge \zeta) &= 0 \\
d(\beta \wedge \zeta) &= 0.
\end{aligned} \tag{6.2}$$

Let Γ be a discrete subgroup of G such that $\Gamma \backslash G$ is compact. One such example is the set of matrices

$$\begin{pmatrix}
(\kappa_1)^z & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x + y\kappa_1 \\
A & 1 & 0 & 0 & 0 & 0 & 0 & 0 & t \\
0 & 0 & (\kappa_1)^{-z} & 0 & 0 & 0 & 0 & 0 & x + y(\kappa_1)^{-1} \\
0 & 0 & 0 & (\kappa_2)^z & 0 & 0 & 0 & 0 & a + b\kappa_2 \\
0 & 0 & 0 & B & 1 & 0 & 0 & 0 & s \\
0 & 0 & 0 & 0 & 0 & (\kappa_2)^{-z} & 0 & 0 & a + b(\kappa_2)^{-1} \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & z \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}$$

with $A = -n_1 (x(\kappa_1)^z + y(\kappa_1)^{z-1})$, $B = -n_2 (a + b(\kappa_2)^{-1})(\kappa_2)^z$, and such that $(x, y, t, a, b, s, z) \in \mathbb{Z}^7$.

Using the equations satisfied by κ_1, κ_2 , one may indeed verify that Γ is a subgroup. The left-invariant vector fields X, Y, T, A, B, S, Z and the left-invariant forms $\xi, v, \tau, \alpha, \beta, \sigma, \zeta$ descend to well defined one-forms on $\Gamma \backslash G$ that satisfy the same relations (6.1). We note that none of these forms are exact on the compact manifold, whereas all were exact on G . We choose the metric for which these vector fields form an orthonormal basis for the tangent space of $\Gamma \backslash G$.

We consider the three-dimensional distribution spanned by A, S , and T , which satisfies the Frobenius condition so that it is the tangent space to a foliation \mathcal{F} . We check (5.3) to determine if the foliation is Riemannian. Since all brackets of basis vectors with S and T are zero, we need only check

$$\begin{aligned}
\langle [Z, A], \bullet \rangle + \langle Z, [\bullet, A] \rangle &= 0 \\
\langle [B, A], \bullet \rangle + \langle B, [\bullet, A] \rangle &= 0
\end{aligned}$$

where \bullet is any choice of section of L^\perp . Thus, this foliation is Riemannian. The characteristic form of the foliation is $\chi_{\mathcal{F}} = \alpha \wedge \sigma \wedge \tau$, and we compute

$$\begin{aligned} d\chi_{\mathcal{F}} &= d(\alpha \wedge \sigma \wedge \tau) = d\alpha \wedge \sigma \wedge \tau - \alpha \wedge d\sigma \wedge \tau + \alpha \wedge \sigma \wedge d\tau \\ &= k_2\alpha \wedge \zeta \wedge \sigma \wedge \tau - 0 - n_1\alpha \wedge \sigma \wedge \xi \wedge v \\ &= -k_2\zeta \wedge \chi_{\mathcal{F}} - n_1\alpha \wedge \sigma \wedge \xi \wedge v, \end{aligned}$$

so that the mean curvature form is $k_2\zeta$ and the Euler form is $\varphi_0 = -n_1\alpha \wedge \sigma \wedge \xi \wedge v$, which is nonzero. Thus, this foliation is nontaut and also as nonintegrable normal bundle. Also note that $k_2\zeta$ is basic, and

$$\begin{aligned} \delta_b(\kappa_b) &= \delta_b(k_2\zeta) = -k_2\bar{*}d\bar{*}\zeta + \kappa_{b\lrcorner}k_2\zeta \\ &= -k_2\bar{*}d(-\xi \wedge v \wedge \beta) + k_2^2 \\ &= k_2\bar{*}(0 \wedge \beta + \xi \wedge v \wedge (-k_2\beta \wedge \zeta)) + k_2^2 \\ &= -k_2^2\bar{*}(\xi \wedge v \wedge \beta \wedge \zeta) + k_2^2 = -k_2^2 + k_2^2 = 0. \end{aligned}$$

Thus, $\kappa = k_2\zeta$ is basic harmonic.

We now compute some of the de Rham cohomology of $\Gamma \setminus G$ using (6.1) and (6.2). We see that $H^1(\Gamma \setminus G) \cong \mathbb{R}$, generated by $[\zeta]$. The group $H^2(\Gamma \setminus G) \cong 0$ (all two closed forms listed in (6.2) are exact). Then $H^5(\Gamma \setminus G) \cong H^2(\Gamma \setminus G) \cong 0$, and $H^6(\Gamma \setminus G) \cong \mathbb{R}$ is generated by $[\xi \wedge v \wedge \tau \wedge \alpha \wedge \beta \wedge \sigma]$. If this manifold is indeed formal, then we note that all harmonic 4-forms ω must be in the kernel of $\zeta \wedge$, because otherwise $\zeta \wedge$ would yield a nontrivial harmonic 5-form. But this means all harmonic 4-forms are of the form $\zeta \wedge$ (harmonic 3-form), which in turn means each harmonic 3-form is a constant linear combination of the wedge product of three of the forms $\xi, v, \tau, \alpha, \beta, \sigma$. We note that $\xi \wedge v \wedge \tau$ and $\alpha \wedge \beta \wedge \sigma$ are clearly closed and not exact, so that the rank of $H^3(\Gamma \setminus G)$ is at least 2.

Next, we consider the basic cohomology $H^*(\Gamma \setminus G, \mathcal{F})$, where the foliation has codimension four. The basic forms must be algebraic combinations of the forms ξ, v, β, ζ , because they span the conormal bundle Q^* . We only need to check if their differentials have no leafwise components. We see that $d\xi = k_1\xi \wedge \zeta$, $dv = -k_1v \wedge \zeta$, $d\beta = -k_2\beta \wedge \zeta$, and $d\zeta = 0$, so only ζ is basic. Since it is also closed and not exact, $H^1(\Gamma \setminus G, \mathcal{F})$ is generated by $[\zeta]$. We have seen above that ζ is basic-harmonic, since $\delta_b(\zeta) = 0$. Next, we have that locally $\bigwedge^2 Q^*$ is spanned by $\xi \wedge v, v \wedge \beta, v \wedge \zeta, \xi \wedge \beta, \beta \wedge \zeta, \xi \wedge \zeta$. Their differentials are respectively 0, $(k_1 + k_2)v \wedge \beta \wedge \zeta, 0, (-k_1 + k_2)\xi \wedge \beta \wedge \zeta, 0, 0$, so in fact all of these forms are basic. If $k_1 + k_2 \neq 0$, $-k_1 + k_2 \neq 0$, we see that $\xi \wedge v, v \wedge \zeta, \beta \wedge \zeta, \xi \wedge \zeta$ are closed, yet of those $v \wedge \zeta = d\left(-\frac{1}{k_1}v\right)$, $\beta \wedge \zeta = d\left(-\frac{1}{k_2}\beta\right)$, $\xi \wedge \zeta = d\left(\frac{1}{k_1}\xi\right)$; $\xi \wedge v$ is not the differential of any basic one-form, so that $H^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}$ with generator $\xi \wedge v$. We see also that

$$\begin{aligned} \delta_b(\xi \wedge v) &= -\bar{*}d\bar{*}(\xi \wedge v) + \kappa_{b\lrcorner}(\xi \wedge v) \\ &= -\bar{*}d(\beta \wedge \zeta) + 0 = 0, \end{aligned}$$

so that $\xi \wedge v$ is basic harmonic. If $k_1 = -k_2$, $v \wedge \beta$ is also closed and is not exact, so that $H^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}^2$, spanned by $\xi \wedge v$ and $v \wedge \beta$. In this case,

$$\begin{aligned} \delta_b(v \wedge \beta) &= -\bar{*}d\bar{*}(v \wedge \beta) + \kappa_{b\lrcorner}(v \wedge \beta) \\ &= -\bar{*}d(\xi \wedge \zeta) + 0 = 0, \end{aligned}$$

and also $v \wedge \beta$ is basic harmonic. If $k_1 = k_2$, $\xi \wedge \beta$ is also closed and is not exact, so that $H^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}^2$, spanned by $\xi \wedge v$ and $\xi \wedge \beta$. In this case,

$$\begin{aligned} \delta_b(\xi \wedge \beta) &= -\bar{*}d\bar{*}(\xi \wedge \beta) + \kappa_{b\lrcorner}(\xi \wedge \beta) \\ &= -\bar{*}d(-v \wedge \zeta) + 0 = 0, \end{aligned}$$

so also $\xi \wedge \beta$ is basic harmonic.

The three forms in $\bigwedge^3 Q^*$ are spanned locally by $v \wedge \beta \wedge \zeta$, $\xi \wedge \beta \wedge \zeta$, $\xi \wedge v \wedge \zeta$, $\xi \wedge v \wedge \beta$. We see that their differentials are, respectively, 0, 0, 0, $-k_2\xi \wedge v \wedge \beta \wedge \zeta$. We see that $v \wedge \beta \wedge \zeta = d\left(\frac{1}{k_1+k_2}v \wedge \beta\right)$ if $k_1 \neq -k_2$, $\xi \wedge \beta \wedge \zeta = d\left(\frac{1}{-k_1+k_2}\xi \wedge \beta\right)$ if $k_1 \neq k_2$, and finally $\xi \wedge v \wedge \zeta$ is not the differential of any basic form. So if $k_1 \neq -k_2$ and $k_1 \neq k_2$, $H^3(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}$, generated by $\xi \wedge v \wedge \zeta$. Note

$$\begin{aligned} \delta_b(\xi \wedge v \wedge \zeta) &= -\bar{*}d\bar{*}(\xi \wedge v \wedge \zeta) + \kappa_{b\lrcorner}(\xi \wedge v \wedge \zeta) \\ &= -\bar{*}d(-\beta) + k_2\xi \wedge v = -k_2\bar{*}(\beta \wedge \zeta) + k_2\xi \wedge v \\ &= -k_2\bar{*}(\beta \wedge \zeta) + k_2\xi \wedge v = -k_2\xi \wedge v + k_2\xi \wedge v = 0, \end{aligned}$$

so this form is basic harmonic. If $k_1 = k_2$, $\xi \wedge \beta \wedge \zeta$ generates another cohomology class, and

$$\begin{aligned} \delta_b(\xi \wedge \beta \wedge \zeta) &= -\bar{*}d\bar{*}(\xi \wedge \beta \wedge \zeta) + \kappa_{b\lrcorner}(\xi \wedge \beta \wedge \zeta) \\ &= -\bar{*}dv + k_2\xi \wedge \beta = -\bar{*}(-k_1v \wedge \zeta) + k_2\xi \wedge \beta \\ &= k_1(-\xi \wedge \beta) + k_2\xi \wedge \beta = 0, \end{aligned}$$

so that form is also basic harmonic. If $k_1 = -k_2$, $v \wedge \beta \wedge \zeta$ generates another cohomology class, and

$$\begin{aligned} \delta_b(v \wedge \beta \wedge \zeta) &= -\bar{*}d\bar{*}(v \wedge \beta \wedge \zeta) + \kappa_{b\lrcorner}(v \wedge \beta \wedge \zeta) \\ &= -\bar{*}d(-\xi) + k_2v \wedge \beta = -\bar{*}(-k_1\xi \wedge \zeta) + k_2v \wedge \beta \\ &= k_1v \wedge \beta + k_2v \wedge \beta = 0, \end{aligned}$$

and that form is also basic harmonic. In either of these last two cases, $H^3(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}^2$. Finally, $H^4(\Gamma \setminus G, \mathcal{F}) \cong 0$ since the foliation is nontaut. We now calculate the κ -harmonic forms, using the differential $d - \kappa\wedge$. We have $H_\kappa^0(\Gamma \setminus G, \mathcal{F}) \cong 0$ since κ is not basic exact. Among the one-forms, we see that

$$\begin{aligned} d\xi &= k_1\xi \wedge \zeta, dv = -k_1v \wedge \zeta, d\beta = -k_2\beta \wedge \zeta, d\zeta = 0 \\ \kappa \wedge \xi &= -k_2\xi \wedge \zeta, \kappa \wedge v = -k_2v \wedge \zeta, \kappa \wedge \beta = -k_2\beta \wedge \zeta, \kappa \wedge \zeta = 0. \end{aligned}$$

From the above, ζ is basic κ -exact, since $(d - \kappa\wedge)\left(-\frac{1}{k_2}\right) = \zeta$, and β is basic κ -closed and not basic κ -exact, and ξ is basic κ -closed if and only if $k_1 = -k_2$, and v is basic κ -closed if and only if $k_1 = k_2$. In all of these cases, these generating forms are basic κ -harmonic. We could also tell this since $\bar{*}\delta_b = \pm(d - \kappa_{b\lrcorner})\bar{*}$ and $\bar{*}d = \pm(\delta_b - \kappa_{b\lrcorner})\bar{*}$ on basic forms. In summary, $H_\kappa^1(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}$ unless $k_1 = \pm k_2$, in which case $H_\kappa^1(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}^2$. Similarly, we see that $H_\kappa^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}$, generated by the basic κ -harmonic form $\bar{*}(\xi \wedge v) = \beta \wedge \zeta$, if $k_1 \neq \pm k_2$. If $k_1 = k_2$, we have the additional basic κ -harmonic form $\bar{*}(-\xi \wedge \beta) = v \wedge \zeta$, and if $k_1 = -k_2$, we have $\bar{*}(v \wedge \beta) = \xi \wedge \zeta$, also basic κ -harmonic. Thus, $H_\kappa^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}$ unless $k_1 = \pm k_2$, in which case $H_\kappa^2(\Gamma \setminus G, \mathcal{F}) \cong \mathbb{R}^2$. Finally, $H_\kappa^3(\Gamma \setminus G, \mathcal{F})$ is generated by $-\bar{*}\zeta = \xi \wedge v \wedge \beta$, also basic κ -harmonic. We also observe that in all these cases, the wedge product of a basic harmonic form and a basic κ -harmonic form is basic κ -harmonic, verifying that $(\Gamma \setminus G, \mathcal{F})$ with the given metric is transversely formal.

REFERENCES

1. Jesús A. Álvarez López, *The basic component of the mean curvature of Riemannian foliations*, Ann. Global Anal. Geom. **10** (1992), no. 2, 179–194.
2. Victor Bangert and Mikhail Katz, *An optimal Loewner-type systolic inequality and harmonic one-forms of constant norm*, Comm. Anal. Geom. **12** (2004), no. 3, 703–732.
3. Augustin Banyaga, *Examples of non d_ω -exact locally conformal symplectic forms*, J. Geom. **87** (2007), no. 1-2, 1–13.
4. Christian Bär, *Geometrically formal 4-manifolds with nonnegative sectional curvature*, Comm. Anal. Geom. **23** (2015), no. 3, 479–497.
5. Florin Alexandru Belgun, *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), no. 1, 1–40.
6. David E. Blair, *Riemannian geometry of contact and symplectic manifolds*, second ed., Progress in Mathematics, vol. 203, Birkhäuser Boston, Ltd., Boston, MA, 2010.
7. Robert A. Blumenthal and James J. Hebda, *de Rham decomposition theorems for foliated manifolds*, Ann. Inst. Fourier (Grenoble) **33** (1983), no. 2, 183–198.
8. ———, *Ehresmann connections for foliations*, Indiana Univ. Math. J. **33** (1984), no. 4, 597–611.
9. Christoph Bock, *On low-dimensional solvmanifolds*, Asian J. Math. **20** (2016), no. 2, 199–262.
10. Charles P. Boyer, Krzysztof Galicki, and Paola Matzeu, *On eta-Einstein Sasakian geometry*, Comm. Math. Phys. **262** (2006), no. 1, 177–208.
11. Beniamino Cappelletti-Montano, Antonio De Nicola, Juan Carlos Marrero, and Ivan Yudin, *Sasakian nilmanifolds*, Int. Math. Res. Not. Ino. 15, 6648–6660.
12. Yves Carrière, *Flots riemanniens*, no. 116, 1984, Transversal structure of foliations (Toulouse, 1982), pp. 31–52.
13. Luis C. de Andrés, Luis A. Cordero, Marisa Fernández, and José J. Mencía, *Examples of four-dimensional compact locally conformal Kähler solvmanifolds*, Geom. Dedicata **29** (1989), no. 2, 227–232.
14. Pierre Deligne, Phillip Griffiths, John Morgan, and Dennis Sullivan, *Real homotopy theory of Kähler manifolds*, Invent. Math. **29** (1975), no. 3, 245–274.
15. Demetrio Domínguez, *Finiteness and tenseness theorems for Riemannian foliations*, Amer. J. Math. **120** (1998), no. 6, 1237–1276.
16. Fida El Chami and Georges Habib, *The Bochner formula for Riemannian flows*, Results Math. **77** (2022), no. 1, Paper No. 25, 24.
17. Aziz El Kacimi-Alaoui and Gilbert Hector, *Décomposition de Hodge basique pour un feuilletage riemannien*, Ann. Inst. Fourier (Grenoble) **36** (1986), no. 3, 207–227.
18. Edmond Fedida, *Sur les feuilletages de Lie*, C. R. Acad. Sci. Paris Sér. A-B **272** (1971), A999–A1001.
19. Tamehiro Fujitani, *Complex-valued differential forms on normal contact Riemannian manifolds*, Tohoku Math. J. (2) **18** (1966), 349–361.
20. Jean-François Grosjean and Paul-Andi Nagy, *On the cohomology algebra of some classes of geometrically formal manifolds*, Proc. Lond. Math. Soc. (3) **98** (2009), no. 3, 607–630.
21. Georges Habib and Ken Richardson, *Modified differentials and basic cohomology for Riemannian foliations*, J. Geom. Anal. **23** (2013), no. 3, 1314–1342.
22. ———, *Riemannian flows and adiabatic limits*, Internat. J. Math. **29** (2018), no. 2, 1850011, 21.
23. Keizo Hasegawa, *Minimal models of nilmanifolds*, Proc. Amer. Math. Soc. **106** (1989), no. 1, 65–71.
24. James J. Hebda, *An example relevant to curvature pinching theorems for Riemannian foliations*, Proc. Amer. Math. Soc. **114** (1992), no. 1, 195–199.
25. William V. D. Hodge, *The theory and applications of harmonic integrals*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1989, Reprint of the 1941 original, With a foreword by Michael Atiyah.
26. Franz W. Kammer and Philippe Tondeur, *Duality theorems for foliations*, no. 116, 1984, Transversal structure of foliations (Toulouse, 1982), pp. 108–116.
27. ———, *de Rham-Hodge theory for Riemannian foliations*, Math. Ann. **277** (1987), no. 3, 415–431.
28. Hisashi Kasuya, *Geometrical formality of solvmanifolds and solvable Lie type geometries*, Geometry of transformation groups and combinatorics, RIMS Kōkyūroku Bessatsu, vol. B39, Res. Inst. Math. Sci. (RIMS), Kyoto, 2013, pp. 21–33.

29. Shoshichi Kobayashi, *Principal fibre bundles with the 1-dimensional toroidal group*, Tohoku Math. J. (2) **8** (1956), 29–45.
30. Dieter Kotschick, *On products of harmonic forms*, Duke Math. J. **107** (2001), no. 3, 521–531.
31. ———, *Geometric formality and non-negative scalar curvature*, Pure Appl. Math. Q. **13** (2017), no. 3, 437–451. MR 3882204
32. Dieter Kotschick and Svjetlana Terzić, *Geometric formality of homogeneous spaces and of biquotients*, Pacific J. Math. **249** (2011), no. 1, 157–176.
33. Xosé Masa, *Duality and minimality in Riemannian foliations*, Comment. Math. Helv. **67** (1992), no. 1, 17–27.
34. Krzysztof Maurin, *The Riemann legacy*, Mathematics and its Applications, vol. 417, Kluwer Academic Publishers Group, Dordrecht, 1997, Riemannian ideas in mathematics and physics, Translated from the Polish by Jerzy Kowalski-Glikman.
35. Pierre Molino, *Riemannian foliations*, Progress in Mathematics, vol. 73, Birkhäuser Boston, Inc., Boston, MA, 1988, Translated from the French by Grant Cairns, With appendices by Cairns, Y. Carrière, É. Ghys, E. Salem and V. Sergiescu.
36. Paul-Andi Nagy, *On length and product of harmonic forms in Kähler geometry*, Math. Z. **254** (2006), no. 1, 199–218.
37. Liviu Ornea and Mihaela Pilca, *Remarks on the product of harmonic forms*, Pacific J. Math. **250** (2011), no. 2, 353–363.
38. Efton Park and Ken Richardson, *The basic Laplacian of a Riemannian foliation*, Amer. J. Math. **118** (1996), no. 6, 1249–1275.
39. Bruce L. Reinhart, *Foliated manifolds with bundle-like metrics*, Ann. of Math. (2) **69** (1959), 119–132.
40. José Ignacio Royo Prieto, *Estudio cohomológico de flujos riemannianos*, Ph.D. thesis, University of the Basque Country, 2004.
41. Hansklaus Rummeler, *Quelques notions simples en géométrie riemannienne et leurs applications aux feuilletages compacts*, Comment. Math. Helv. **54** (1979), no. 2, 224–239.
42. Dennis Sullivan, *Differential forms and the topology of manifolds*, Manifolds—Tokyo 1973 (Proc. Internat. Conf., Tokyo, 1973), Published for the Mathematical Society of Japan by University of Tokyo Press, Tokyo, 1975, pp. 37–49.
43. Shun-ichi Tachibana, *On a decomposition of C-harmonic forms in a compact Sasakian space*, Tohoku Math. J. (2) **19** (1967), 198–212.
44. Philippe Tondeur, *Geometry of foliations*, Monographs in Mathematics, vol. 90, Birkhäuser Verlag, Basel, 1997.
45. Burt Totaro, *Curvature, diameter, and quotient manifolds*, Math. Res. Lett. **10** (2003), no. 2-3, 191–203.
46. Aleksy Tralle and John Oprea, *Symplectic manifolds with no Kähler structure*, Lecture Notes in Mathematics, vol. 1661, Springer-Verlag, Berlin, 1997.

LEBANESE UNIVERSITY, FACULTY OF SCIENCES II, DEPARTMENT OF MATHEMATICS, P.O. BOX 90656
 FANAR-MATN, LEBANON, AND UNIVERSITÉ DE LORRAINE, CNRS, IECL, 54506 NANCY, FRANCE
Email address, G. Habib: g Habib@ul.edu.lb

DEPARTMENT OF MATHEMATICS, TEXAS CHRISTIAN UNIVERSITY, FORT WORTH, TEXAS 76129, USA
Email address, K. Richardson: k.richardson@tcu.edu

WYDZIAŁ MATEMATYKI I INFORMATYKI UJ, UL. PROF. STANISŁAWA ŁOJASIEWICZA 6, 30-348
 KRAKÓW, POLAND
Email address, R. Wolak: robert.wolak@uj.edu.pl