

# TOEPLITZ ALGEBRA AND SYMBOL MAP VIA BEREZIN TRANSFORM ON $H^2(\mathbb{D}^n)$

MO JAVED AND AMIT MAJI†

ABSTRACT. Let  $\mathcal{T}(L^\infty(\mathbb{T}))$  be the Toeplitz algebra, that is, the  $C^*$ -algebra generated by the set  $\{T_\varphi : \varphi \in L^\infty(\mathbb{T})\}$ . Douglas's theorem on *symbol map* states that there exists a  $C^*$ -algebra homomorphism from  $\mathcal{T}(L^\infty(\mathbb{T}))$  onto  $L^\infty(\mathbb{T})$  such that  $T_\varphi \mapsto \varphi$  and the kernel of the homomorphism coincides with commutator ideal in  $\mathcal{T}(L^\infty(\mathbb{T}))$ . In this paper, we use the *Berezin transform* to study results akin to Douglas's theorem for operators on the Hardy space  $H^2(\mathbb{D}^n)$  over the open unit polydisc  $\mathbb{D}^n$  for  $n \geq 1$ . We further obtain a class of bigger  $C^*$ -algebras than the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  for which the analog of *symbol map* still holds true.

## 1. INTRODUCTION

One of the major concerns of operator theory and function theory has generally been the study of operators associated with the spaces of holomorphic functions. We often come across linear operators induced by functions (usually called the symbol of the operator), such as Toeplitz operators, Hankel operators, and composition operators; on the other hand, we can investigate functions that are induced by an operator. A great illustration of the relationship between operators and functions is the *Berezin transform*.

The *Berezin transform*, which was first introduced in [3] as a tool in quantization, is named after F. A. Berezin. Usually, it involves an open subset  $\mathcal{G}$  of  $\mathbb{C}^n$  ( $n \geq 1$ ) and a reproducing kernel Hilbert space  $H(\mathcal{G})$  with a reproducing kernel, say,  $K_z$  for  $z \in \mathcal{G}$ . Then, for a bounded operator  $A$  on  $H(\mathcal{G})$ , its *Berezin transform*, denoted as  $\tilde{A}$ , is a complex-valued function on  $\mathcal{G}$  defined by

$$\tilde{A}(z) = \langle Ak_z, k_z \rangle_{H(\mathcal{G})} \quad (z \in \mathcal{G}),$$

where  $k_z = K_z / \|K_z\|_{H(\mathcal{G})}$  is the normalized reproducing kernel of  $H(\mathcal{G})$ . Thus every bounded linear operator  $A$  on  $H(\mathcal{G})$  induces a bounded function  $\tilde{A}$  on  $\mathcal{G}$ . Indeed,  $\|\tilde{A}\|_\infty \leq \|A\|$ . Therefore, the *Berezin transform* of an operator yields a crucial information about the operator. Beyond its original purpose, the *Berezin transform* has shown usefulness in a number of contexts, including the Hardy space, the Bergman space ([17], [1]), the Bargmann-Segal space [5]. It has also established important relationships with the Bloch space and functions of bounded mean oscillation, and the detailed investigation of these connections can be found in [18]. It has also found many applications in many areas of mathematical analysis and mathematical physics (see [4], [13], [19]).

---

2010 *Mathematics Subject Classification.* 47B35, 47A13, 32A65.

*Key words and phrases.* Hardy space, Toeplitz operator, Hankel operator, Toeplitz algebra, Symbol map, Berezin transform.

Toeplitz operators emerge to be one of the most distinguishable classes of concrete operators because of their immensely useful and widespread applications including operator theory, function theory, operator algebras, and associated disciplines. Furthermore, it is preferable to think of these operators as elements within operator algebras, like the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}))$ , that is, the  $C^*$ -algebra generated by the set  $\{T_\varphi : \varphi \in L^\infty(\mathbb{T})\}$  of Toeplitz operators on the Hardy space  $H^2(\mathbb{D})$  (see [9], [11]). The *symbol map*, which was initially discovered and utilized by Douglas, proves to be an important tool in the study of the Toeplitz algebra. The classical result of Douglas's *symbol map* [11, chapter 7] is as follows:

**Theorem 1.1.** *Let  $\mathcal{T}(L^\infty(\mathbb{T}))$  denote the  $C^*$ -algebra generated by the set  $\{T_\varphi : \varphi \in L^\infty(\mathbb{T})\}$ . Then there exists a  $C^*$ -homomorphism*

$$\sigma : \mathcal{T}(L^\infty(\mathbb{T})) \rightarrow L^\infty(\mathbb{T})$$

such that kernel of  $\sigma$  is the commutator ideal, that is, the ideal in  $\mathcal{T}(L^\infty(\mathbb{T}))$  generated by the commutators  $[A, B] = AB - BA$ ;  $A, B \in \mathcal{T}(L^\infty(\mathbb{T}))$ , and  $\sigma(T_\varphi) = \varphi$  for all  $\varphi \in L^\infty(\mathbb{T})$ .

The  $C^*$ -homomorphism  $\sigma$  is called the *symbol map*. However, Douglas's proof of Theorem 1.1 was mere existential and much abstract in nature until 1982, when Barría and Halmos in [2] presented a constructive proof of Theorem 1.1. More specifically, Barría and Halmos obtained the symbol of an operator  $A \in \mathcal{T}(L^\infty(\mathbb{T}))$  as the symbol of the Toeplitz operator in the SOT-limit of the sequence  $\{T_z^{*m} A T_z^m\}_{m=1}^\infty$ , where  $T_z$  denotes the unilateral shift on the Hardy space  $H^2(\mathbb{D})$ . However, due to the nuances of limits in the strong operator topology, it is still very difficult to find the symbol of any operator  $A$  in  $\mathcal{T}(L^\infty(\mathbb{T}))$ . The *Berezin transform* is a useful tool in this way to obtain the symbol of an operator  $A \in \mathcal{T}(L^\infty(\mathbb{T}))$  more concretely. Indeed, the *Berezin transform* of a Toeplitz operator on Hardy space reduces to the Poisson integral of the symbol of that Toeplitz operator (see Theorem 3.2 below). In particular, Engliš [12] developed a technique using the *Berezin transform* in which he obtained the symbol of an operator  $A \in \mathcal{T}(L^\infty(\mathbb{T}))$  as the radial boundary function of the *Berezin transform* of  $A$ . The key observation in his proof is that

$$T_\varphi K_\xi = \varphi(\xi) K_\xi,$$

for the densely defined Toeplitz operator  $T_\varphi$  with symbol  $\varphi \in H^2(\mathbb{D})^\perp$  and  $K_\xi$  is the reproducing kernel of  $H^2(\mathbb{D})$ . It uses the fact that  $\bar{\varphi} \in H^2(\mathbb{D})$  whenever  $\varphi \in H^2(\mathbb{D})^\perp$ . However, this technique fails in the case  $\mathbb{D}^n$  for  $n > 1$ , for an example: if we take  $\varphi(z_1, z_2) = \bar{z}_1 z_2$  then both  $\varphi$  and  $\bar{\varphi}$  are in  $H^2(\mathbb{D}^2)^\perp$ , and  $T_\varphi K_\xi \neq \varphi(\xi) K_\xi$  for  $\xi \in \mathbb{D}^2$ , where  $K_\xi(z) = \frac{1}{(1 - \bar{\xi}_1 z_1)(1 - \bar{\xi}_2 z_2)}$  is a reproducing kernel of  $H^2(\mathbb{D}^2)$ . Thus the natural question arises:

**Question:** Is the Douglas's symbol map valid in the setting of the open unit polydisc  $\mathbb{D}^n$  for  $n > 1$ ?

In this paper we present a comprehensive study of symbols of operators on  $H^2(\mathbb{D}^n)$ . The main contribution of this article is to address the above question with the help of the *Berezin transform*. More precisely, we obtain the following result (see Theorem 4.3 below):

**Theorem.** Let  $T \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ , the Toeplitz algebra of operators on  $H^2(\mathbb{D}^n)$  for  $n \geq 1$ . Then  $\tilde{T} \rightarrow \varphi$  radially for some  $\varphi \in L^\infty(\mathbb{T}^n)$ . Moreover, the map

$$\Sigma : \mathcal{T}(L^\infty(\mathbb{T}^n)) \rightarrow L^\infty(\mathbb{T}^n), \quad T \mapsto \varphi$$

is a  $C^*$ -algebra homomorphism with  $\ker(\Sigma) = \mathcal{I}$ , the *semi-commutator ideal* in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  and  $\Sigma(T_\varphi) = \varphi$  for any Toeplitz operator  $T_\varphi$ .

The rest of the paper is outlined as follows: Section 2 is devoted to establishing the necessary background, and along the way, we fix some notations and state some properties of Poisson integral and *Berezin transform*. In Section 3, we prove an important result concerning the radial limits, which serves as a key ingredient in the proof of the *symbol map*. Section 4 concerns with the main result, that is, the *symbol map* (see Theorem 4.3), and we depict through an example that Theorem 4.3 can not be extended to the whole of  $\mathfrak{B}(H^2(\mathbb{D}^n))$ , the  $C^*$ -algebra of all bounded operators on  $H^2(\mathbb{D}^n)$ . In section 5, we present a family of larger  $C^*$ -algebras than the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  for which the analog of symbol map still holds true. In section 6, we conclude with some open questions.

## 2. PRELIMINARIES

Let  $\mathbb{Z}_+$  denote the set of all non-negative integers,  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  be the open unit disc in the complex plane  $\mathbb{C}$ ,  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  the unit circle in  $\mathbb{C}$ , and  $\mathfrak{B}(\mathcal{H})$  denote the set of all bounded linear operators on a Hilbert space  $\mathcal{H}$ . For  $n \geq 1$ , the Cartesian product of  $n$  copies of  $\mathbb{D}$ , denoted by  $\mathbb{D}^n$ , is the unit polydisc in  $\mathbb{C}^n$  and the  $n$ -torus  $\mathbb{T}^n$  denotes the distinguished boundary of  $\mathbb{D}^n$ . Then  $H^2(\mathbb{D}^n)$  denotes as the Hilbert space of all analytic functions  $f$  on  $\mathbb{D}^n$  for which

$$\|f\|_{H^2(\mathbb{D}^n)} := \sup_{0 \leq r < 1} \left( \int_{\mathbb{T}^n} |f(r\zeta)|^2 d\mathbf{m}_n(\zeta) \right)^{1/2} < \infty,$$

where  $\mathbf{m}_n$  stands for the normalised Lebesgue measure on  $\mathbb{T}^n$ . We shall write bold faced  $\boldsymbol{\xi}$  and  $\boldsymbol{\zeta}$ , respectively as vectors  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$  and  $\boldsymbol{\zeta} = (\zeta_1, \dots, \zeta_n) \in \mathbb{T}^n$  for  $n > 1$ . For  $1 \leq p < \infty$ ,  $L^p(\mathbb{T}^n) := L^p(\mathbb{T}^n, d\mathbf{m}_n)$  is the Banach space of equivalence classes of Lebesgue complex measurable functions  $f$  such that  $|f|^p$  is integrable on  $\mathbb{T}^n$  with respect to  $\mathbf{m}_n$  and  $p = 2$  gives the Hilbert space  $L^2(\mathbb{T}^n)$  of square-integrable equivalence classes of functions with respect to  $\mathbf{m}_n$ . Also  $L^\infty(\mathbb{T}^n)$  denotes the commutative  $C^*$ -algebra of essentially bounded complex measurable functions on  $\mathbb{T}^n$  with respect to  $\mathbf{m}_n$ , and  $H^\infty(\mathbb{D}^n)$  stands for the space of bounded holomorphic functions on  $\mathbb{D}^n$ . Let  $\{e_{\mathbf{k}}\}_{\mathbf{k} \in \mathbb{Z}^n}$  denote the standard orthonormal basis for  $L^2(\mathbb{T}^n)$ , where  $e_{\mathbf{k}}(\boldsymbol{\zeta}) = \zeta_1^{k_1} \dots \zeta_n^{k_n}$  for  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$ . Consider the closed subspace  $H^2(\mathbb{T}^n)$  of  $L^2(\mathbb{T}^n)$  defined by

$$H^2(\mathbb{T}^n) := \{f \in L^2(\mathbb{T}^n) : \langle f, e_{\mathbf{k}} \rangle_{L^2(\mathbb{T}^n)} = 0 \text{ for all } \mathbf{k} \in \mathbb{Z}^n \setminus \mathbb{Z}_+^n\}.$$

We then identify the Hardy space  $H^2(\mathbb{D}^n)$  (via radial limits) with  $H^2(\mathbb{T}^n)$  by the unitary map

$$H^2(\mathbb{D}^n) \ni \tilde{f} = \sum_{\mathbf{k} \in \mathbb{Z}_+^n} a_{\mathbf{k}} \mathbf{z}^{\mathbf{k}} \mapsto \sum_{\mathbf{k} \in \mathbb{Z}_+^n} a_{\mathbf{k}} e_{\mathbf{k}} = f \in H^2(\mathbb{T}^n).$$

Now onwards, without any ambiguity, we shall interchangeably write  $H^2(\mathbb{D}^n)$  (respectively, its elements  $\tilde{f}$ ) in place of  $H^2(\mathbb{T}^n)$  (respectively, its elements  $f$ ) and vice-versa. Furthermore, for  $\tilde{f}, \tilde{g} \in H^2(\mathbb{D}^n)$  and  $f, g \in H^2(\mathbb{T}^n)$ , we have

$$\langle \tilde{f}, \tilde{g} \rangle_{H^2(\mathbb{D}^n)} = \langle f, g \rangle_{L^2(\mathbb{T}^n)}.$$

For  $\varphi \in L^\infty(\mathbb{T}^n)$ , the Toeplitz operator  $T_\varphi$  is a bounded operator on  $H^2(\mathbb{D}^n)$  given by

$$T_\varphi f = P_{H^2(\mathbb{D}^n)}(\varphi f) \quad (f \in H^2(\mathbb{D}^n)),$$

and the Hankel operator  $H_\varphi$  with symbol  $\varphi \in L^\infty(\mathbb{T}^n)$  is a bounded operator from  $H^2(\mathbb{D}^n)$  to  $H^2(\mathbb{D}^n)^\perp$  defined by

$$H_\varphi f = P_{H^2(\mathbb{D}^n)^\perp}(\varphi f) \quad (f \in H^2(\mathbb{D}^n)),$$

where  $P_{H^2(\mathbb{D}^n)}$  is the orthogonal projection of  $L^2(\mathbb{T}^n)$  onto  $H^2(\mathbb{D}^n)$  and  $P_{H^2(\mathbb{D}^n)^\perp} = I_{L^2(\mathbb{T}^n)} - P_{H^2(\mathbb{D}^n)}$ . Clearly, the adjoint operator  $T_\varphi^* = T_{\bar{\varphi}}$ , where  $\bar{\varphi}$  is the complex conjugate of  $\varphi$ . It is easy to check that  $[T_\varphi, T_\psi] = H_\varphi^* H_\psi$  for  $\varphi, \psi \in L^\infty(\mathbb{T}^n)$ .

We frequently use the identification of the Hardy space over the polydisc  $H^2(\mathbb{D}^n)$  with  $H^2(\mathbb{D}) \otimes \cdots \otimes H^2(\mathbb{D})$ , the  $n$ -fold ( $n > 1$ ) tensor product of Hilbert space  $H^2(\mathbb{D})$  via the canonical unitary map  $z_1^{m_1} \cdots z_n^{m_n} \mapsto z^{m_1} \otimes \cdots \otimes z^{m_n}$ , where  $m_i \in \mathbb{Z}_+$ . Thus, we shall often write  $(f_1 \otimes \cdots \otimes f_n) = f_1 \cdots f_n$ , where  $f_j \in H^2(\mathbb{D})$ . The Hardy space over the polydisc  $H^2(\mathbb{D}^n)$  is a reproducing kernel Hilbert space on  $\mathbb{D}^n$  for  $n \geq 1$ . For  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$ , the (Szegő) kernel  $K_{\boldsymbol{\xi}}$  of  $H^2(\mathbb{D}^n)$  is defined as

$$K_{\boldsymbol{\xi}}(\boldsymbol{z}) = \prod_{j=1}^n \frac{1}{(1 - \bar{\xi}_j z_j)}, \quad (\boldsymbol{z} \in \mathbb{D}^n)$$

and it reproduces the value of each  $f \in H^2(\mathbb{D}^n)$ , that means  $f(\boldsymbol{\xi}) = \langle f, K_{\boldsymbol{\xi}} \rangle$ . The normalised reproducing kernel is denoted by

$$k_{\boldsymbol{\xi}}(\boldsymbol{z}) = \frac{K_{\boldsymbol{\xi}}(\boldsymbol{z})}{\|K_{\boldsymbol{\xi}}\|} = \frac{\sqrt{(1 - |\xi_1|^2) \cdots (1 - |\xi_n|^2)}}{(1 - \bar{\xi}_1 z_1) \cdots (1 - \bar{\xi}_n z_n)}, \quad (\boldsymbol{z} \in \mathbb{D}^n).$$

In particular, the reproducing kernel  $K_\xi$  ( $\xi \in \mathbb{D}$ ) of the Hardy space over the disc  $H^2(\mathbb{D})$  is given by

$$K_\xi(z) = \frac{1}{1 - \bar{\xi}z} \quad (z \in \mathbb{D}),$$

and  $k_\xi(z) = \frac{\sqrt{1 - |\xi|^2}}{1 - \bar{\xi}z}$  is the normalised reproducing kernel of  $H^2(\mathbb{D})$ .

Let  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$ ,  $\boldsymbol{\zeta} = (\zeta_1, \dots, \zeta_n) \in \mathbb{T}^n$ ,  $\xi_j = r_j e^{i\theta_j}$  and  $\zeta_j = e^{it_j}$  where  $0 \leq r_j < 1$  and  $\theta_j, t_j \in [0, 2\pi)$  for all  $1 \leq j \leq n$ . The Poisson kernel  $P(\boldsymbol{\xi}, \boldsymbol{\zeta})$  for the unit polydisc  $\mathbb{D}^n$  is defined as

$$\begin{aligned} P(\boldsymbol{\xi}, \boldsymbol{\zeta}) &= \prod_{j=1}^n \frac{1 - |\xi_j|^2}{|1 - \xi_j \bar{\zeta}_j|^2} \\ &= P_{r_1}(\theta_1 - t_1) \cdots P_{r_n}(\theta_n - t_n), \end{aligned}$$

where  $P_r(\theta) = \frac{1-r^2}{1-2r\cos\theta+r^2}$  is the Poisson kernel for the unit disc  $\mathbb{D}$ . For a complex Borel measure  $\mu$  on  $\mathbb{T}^n$ , its Poisson integral, denoted by  $P[d\mu]$ , is the integral given by

$$P[d\mu](\boldsymbol{\xi}) = \int_{\mathbb{T}^n} P(\boldsymbol{\xi}, \boldsymbol{\zeta}) d\mu(\boldsymbol{\zeta}) \quad (\boldsymbol{\xi} \in \mathbb{D}^n).$$

For  $\varphi \in L^1(\mathbb{T}^n)$ , its harmonic extension  $\tilde{\varphi}$  into  $\mathbb{D}^n$  is defined by the Poisson integral formula

$$\tilde{\varphi}(\boldsymbol{\xi}) = P[\varphi d\mathbf{m}_n](\boldsymbol{\xi}) = \int_{\mathbb{T}^n} P(\boldsymbol{\xi}, \boldsymbol{\zeta}) \varphi(\boldsymbol{\zeta}) d\mathbf{m}_n(\boldsymbol{\zeta}) \quad (\boldsymbol{\xi} \in \mathbb{D}^n).$$

Let  $\Phi : \mathbb{D}^n \rightarrow \mathbb{C}$  and  $\varphi : \mathbb{T}^n \rightarrow \mathbb{C}$  be any two complex functions. Define  $\hat{\Phi}$  as

$$\hat{\Phi}(\boldsymbol{\zeta}) = \lim_{r \rightarrow 1^-} \Phi(r\boldsymbol{\zeta})$$

for all  $\boldsymbol{\zeta} \in \mathbb{T}^n$  at which the (radial) limit on the right exists. We say that ‘ $\Phi$  approaches  $\varphi$  radially’, denoted by ‘ $\Phi \rightarrow \varphi$  radially’ or ‘ $\Phi(\boldsymbol{\xi}) \rightarrow \varphi$  radially’ if

$$\hat{\Phi}(\boldsymbol{\zeta}) = \varphi(\boldsymbol{\zeta}) \quad \text{for a.e. } \boldsymbol{\zeta} \in \mathbb{T}^n.$$

**Theorem 2.1.** (See Theorem 2.3.1 in [16]) *Let  $\varphi \in L^1(\mathbb{T}^n)$ , and  $\nu$  be a measure on  $\mathbb{T}^n$  such that  $\nu$  is singular with respect to  $\mathbf{m}_n$ . If  $\Phi = P[\varphi d\mathbf{m}_n + d\nu]$ , then*

$$\Phi \rightarrow \varphi \quad \text{radially.}$$

*In particular, if  $\nu = 0$ , then  $\tilde{\varphi} \rightarrow \varphi$  radially.*

The Berezin transform of  $T \in \mathfrak{B}(H^2(\mathbb{D}^n))$  is a complex function on  $\mathbb{D}^n$  defined by

$$\tilde{T}(\boldsymbol{\xi}) = \langle Tk_{\boldsymbol{\xi}}, k_{\boldsymbol{\xi}} \rangle \quad \text{for all } \boldsymbol{\xi} \in \mathbb{D}^n.$$

Here  $k_{\boldsymbol{\xi}}$  is the normalised reproducing kernel of  $H^2(\mathbb{D}^n)$ .

The following notions are important and used throughout the paper:

- (1) For  $\varphi, \psi \in L^\infty(\mathbb{T}^n)$ , the *semicommutator* is defined as

$$[T_\varphi, T_\psi] = T_{\varphi\psi} - T_\varphi T_\psi.$$

- (2)  $\mathcal{T}(L^\infty(\mathbb{T}^n)) := C^*$ -algebra generated by the set  $\{T_\varphi : \varphi \in L^\infty(\mathbb{T}^n)\}$ , called the Toeplitz algebra of operators on  $H^2(\mathbb{D}^n)$ .

- (3)  $\mathcal{C} :=$  closed ideal in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  generated by the set  $\{[A, B] = AB - BA : A, B \in \mathcal{T}(L^\infty(\mathbb{T}^n))\}$ , called the *commutator ideal*.

- (4)  $\mathcal{S} :=$  closed ideal in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  generated by the set  $\{[T_\varphi, T_\psi] : \varphi, \psi \in L^\infty(\mathbb{T}^n)\}$ , called the *semicommutator ideal*.

### 3. ON RADIAL LIMITS

In this section we shall discuss about some key results on radial limits which will be used in the next section. Before moving into our main result, we will fix some notations first.

Suppose  $H^2(\mathbb{D})^1 := H^2(\mathbb{D})$  and  $H^2(\mathbb{D})^{-1} := H^2(\mathbb{D})^\perp$ . For  $\alpha = (\alpha_1, \dots, \alpha_n) \in \{-1, 1\}^n$  and  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$ , we write

$$H^2(\mathbb{D}^n)^\alpha := \bigvee \{e_{\mathbf{k}} : k_j < 0 \text{ if } \alpha_j = -1 \text{ and } k_j \geq 0 \text{ if } \alpha_j = 1 \text{ for } 1 \leq j \leq n\}.$$

It is easy to check that  $H^2(\mathbb{D}^n)^\alpha$  is a closed subspace of  $L^2(\mathbb{T}^n)$  for all  $\alpha \in \{-1, 1\}^n$ . We identify  $H^2(\mathbb{D}^n)^\alpha$  with the  $n$ -fold tensor product of Hilbert spaces  $H^2(\mathbb{D})^{\alpha_j}$  for  $1 \leq j \leq n$  by the unitary map  $e_{\mathbf{k}} \mapsto e_{k_1} \otimes \cdots \otimes e_{k_n}$  for all  $k_j < 0$  (provided  $\alpha_j = -1$ ) and  $k_j \geq 0$  (provided  $\alpha_j = 1$ ), where

$$(e_{k_1} \otimes \cdots \otimes e_{k_n})(\zeta) = \zeta_1^{k_1} \cdots \zeta_n^{k_n},$$

for all  $\zeta \in \mathbb{T}^n$ ,  $k_j < 0$  (provided  $\alpha_j = -1$ ) and  $k_j \geq 0$  (provided  $\alpha_j = 1$ ). We shall simply write  $\otimes_{j=1}^n f_j = \prod_{j=1}^n f_j$  for  $f_j \in H^2(\mathbb{D})^{\alpha_j}$ , and also write bold faced  $\mathbf{1} = (1, \dots, 1) \in \{-1, 1\}^n$ .

The following lemma is an easy consequence of the Poisson integral:

**Lemma 3.1.** *Let  $f, g \in L^2(\mathbb{T}^n)$ . Then*

$$\langle fk_{\xi}, gk_{\xi} \rangle_{L^2(\mathbb{T}^n)} = (\widetilde{f\bar{g}})(\xi), \quad (\xi \in \mathbb{D}^n).$$

Moreover,  $\langle fk_{\xi}, gk_{\xi} \rangle_{L^2(\mathbb{T}^n)} \rightarrow f\bar{g}$  radially.

*Proof.* Suppose  $f, g \in L^2(\mathbb{T}^n)$ . Now

$$\begin{aligned} \langle fk_{\xi}, gk_{\xi} \rangle_{L^2(\mathbb{T}^n)} &= \int_{\mathbb{T}^n} (fk_{\xi})(\zeta) (\overline{gk_{\xi}})(\zeta) d\mathbf{m}_n(\zeta) \\ (3.1) \qquad \qquad \qquad &= \int_{\mathbb{T}^n} (f\bar{g})(\zeta) |k_{\xi}(\zeta)|^2 d\mathbf{m}_n(\zeta). \end{aligned}$$

Note that

$$|k_{\xi}(\zeta)|^2 = \frac{(1 - |\xi_1|^2) \cdots (1 - |\xi_n|^2)}{|1 - \bar{\xi}_1 \zeta_1|^2 \cdots |1 - \bar{\xi}_n \zeta_n|^2} = P(\xi, \zeta),$$

where  $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$  and  $\zeta = (\zeta_1, \dots, \zeta_n) \in \mathbb{T}^n$ . Therefore, the equation (3.1) yields

$$\langle fk_{\xi}, gk_{\xi} \rangle_{L^2(\mathbb{T}^n)} = (\widetilde{f\bar{g}})(\xi).$$

Since  $f, g \in L^2(\mathbb{T}^n)$ ,  $f\bar{g} \in L^1(\mathbb{T}^n)$ . Hence, Theorem 2.1 infers that  $\langle fk_{\xi}, gk_{\xi} \rangle_{L^2(\mathbb{T}^n)} \rightarrow f\bar{g}$  radially.  $\blacksquare$

The following theorem establishes the fact that the Berezin transform of  $T_\varphi$  is the harmonic extension of  $\varphi$ :

**Theorem 3.2.** *For  $\varphi \in L^\infty(\mathbb{T}^n)$ ,  $\widetilde{T_\varphi}(\xi) = \widetilde{\varphi}(\xi)$  for all  $\xi \in \mathbb{D}^n$ . Furthermore,  $\widetilde{T_\varphi} \rightarrow \varphi$  radially.*

*Proof.* Let  $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$ . Then

$$\widetilde{T_\varphi}(\xi) = \langle T_\varphi k_{\xi}, k_{\xi} \rangle = \langle P_{H^2(\mathbb{D}^n)}(\varphi k_{\xi}), k_{\xi} \rangle = \langle \varphi k_{\xi}, k_{\xi} \rangle_{L^2(\mathbb{T}^n)} = \widetilde{\varphi}(\xi),$$

where the last equality follows from Lemma 3.1.

Now Theorem 2.1 gives  $\widetilde{T_\varphi} \rightarrow \varphi$  radially.  $\blacksquare$

**Lemma 3.3.** *If  $x \in H^2(\mathbb{D})^\perp$ , then  $P_{H^2(\mathbb{D})}(xk_{\xi}) = \widetilde{x}(\xi)k_{\xi}$  for all  $\xi \in \mathbb{D}$ .*

*Proof.* Let  $z \in \mathbb{D}$ . Then

$$\begin{aligned} \langle P_{H^2(\mathbb{D})}(xK_\xi), K_z \rangle_{H^2(\mathbb{D})} &= \langle xK_\xi, K_z \rangle_{L^2(\mathbb{T})} \\ &= \langle K_\xi, \bar{x}K_z \rangle_{H^2(\mathbb{D})}. \end{aligned}$$

Since  $x \in H^2(\mathbb{D})^\perp$ ,  $\bar{x} \in H^2(\mathbb{D})$ , and hence  $\bar{x}K_z \in H^2(\mathbb{D})$ . Thus, by reproducing property

$$\begin{aligned} \langle P_{H^2(\mathbb{D})}(xK_\xi), K_z \rangle_{H^2(\mathbb{D})} &= \overline{\tilde{x}(\xi)K_z(\xi)} \\ &= \tilde{x}(\xi) \langle K_\xi, K_z \rangle_{H^2(\mathbb{D})} \\ &= \langle \tilde{x}(\xi)K_\xi, K_z \rangle_{H^2(\mathbb{D})} \quad (\text{for all } z \in \mathbb{D}). \end{aligned}$$

Therefore,  $P_{H^2(\mathbb{D})}(xk_\xi) = \tilde{x}(\xi)k_\xi$ . ■

We now state the main result of this section.

**Theorem 3.4.** *Let  $f \in H^2(\mathbb{D}^n)^\alpha$  and  $g \in H^2(\mathbb{D}^n)^\beta$  for  $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_n) \in \{-1, 1\}^n$ . Then*

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow f\bar{g} \text{ radially,}$$

that is,

$$\langle P_{H^2(\mathbb{D}^n)}(fk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(gk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \rightarrow f(e^{i\theta_1}, \dots, e^{i\theta_n})\bar{g}(e^{i\theta_1}, \dots, e^{i\theta_n})$$

as  $r \rightarrow 1^-$  a.e.  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n$ .

*Proof.* Suppose that  $f \in H^2(\mathbb{D}^n)^\alpha$  and  $g \in H^2(\mathbb{D}^n)^\beta$  for  $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_n) \in \{-1, 1\}^n$ . If either  $\alpha = \mathbf{1}$  or  $\beta = \mathbf{1}$ , then

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle = \langle fk_\xi, gk_\xi \rangle,$$

and hence  $\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow f\bar{g}$  radially by using Lemma 3.1.

We now assume that  $\alpha \neq \mathbf{1}$  and  $\beta \neq \mathbf{1}$ . Firstly, consider

$$I = \{j : \alpha_j = -1\} \quad \text{and} \quad J = \{j : \beta_j = -1\}.$$

Let  $f = f_1 \otimes \dots \otimes f_n$  and  $g = g_1 \otimes \dots \otimes g_n$ , where  $f_j \in H^2(\mathbb{D})^{\alpha_j}$  and  $g_j \in H^2(\mathbb{D})^{\beta_j}$  for all  $1 \leq j \leq n$ . Then

$$\begin{aligned} \langle P_{H^2(\mathbb{D}^n)}(fK_\xi), K_z \rangle &= \langle fK_\xi, K_z \rangle_{L^2(\mathbb{T}^n)} \\ &= \langle (f_1 \otimes \dots \otimes f_n)(K_{\xi_1} \otimes \dots \otimes K_{\xi_n}), K_{z_1} \otimes \dots \otimes K_{z_n} \rangle_{L^2(\mathbb{T}^n)} \\ &= \langle f_1K_{\xi_1} \otimes \dots \otimes f_nK_{\xi_n}, K_{z_1} \otimes \dots \otimes K_{z_n} \rangle_{L^2(\mathbb{T}^n)} \\ &= \prod_{j=1}^n \langle f_jK_{\xi_j}, K_{z_j} \rangle_{L^2(\mathbb{T})} \\ &= \prod_{j=1}^n \langle P_{H^2(\mathbb{D})}(f_jK_{\xi_j}), K_{z_j} \rangle_{H^2(\mathbb{D})} \\ &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left( \prod_{j=1}^n \langle P_{H^2(\mathbb{D})}(f'_jK_{\xi_j}), K_{z_j} \rangle_{H^2(\mathbb{D})} \right), \end{aligned}$$

where  $f'_j = \begin{cases} f_j & \text{if } j \notin I; \\ 1 & \text{if } j \in I. \end{cases}$ , and the last equality follows from Lemma 3.3. Therefore,

$$\begin{aligned} \langle P_{H^2(\mathbb{D}^n)}(fK_\xi), K_z \rangle &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left( \prod_{j=1}^n \langle f'_j K_{\xi_j}, K_{z_j} \rangle_{H^2(\mathbb{D})} \right) \\ &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left\langle \bigotimes_{j=1}^n f'_j K_{\xi_j}, \bigotimes_{j=1}^n K_{z_j} \right\rangle \\ &= \left\langle \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left( \bigotimes_{j=1}^n f'_j K_{\xi_j} \right), K_z \right\rangle, \end{aligned}$$

for all  $z \in \mathbb{D}^n$ . Hence

$$P_{H^2(\mathbb{D}^n)}(fk_\xi) = \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left( \bigotimes_{j=1}^n f'_j k_{\xi_j} \right), \text{ where } f'_j = \begin{cases} f_j & \text{if } j \notin I; \\ 1 & \text{if } j \in I. \end{cases}$$

Similarly, we have

$$P_{H^2(\mathbb{D}^n)}(gk_\xi) = \left( \prod_{j \in J} \tilde{g}_j(\xi_j) \right) \left( \bigotimes_{j=1}^n g'_j k_{\xi_j} \right), \text{ where } g'_j = \begin{cases} g_j & \text{if } j \notin J; \\ 1 & \text{if } j \in J. \end{cases}$$

Now

$$\begin{aligned} \langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle &= \left\langle \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \left( \bigotimes_{j=1}^n f'_j k_{\xi_j} \right), \left( \prod_{j \in J} \tilde{g}_j(\xi_j) \right) \left( \bigotimes_{j=1}^n g'_j k_{\xi_j} \right) \right\rangle \\ &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \overline{\left( \prod_{j \in J} \tilde{g}_j(\xi_j) \right)} \left\langle \left( \bigotimes_{j=1}^n f'_j k_{\xi_j} \right), \left( \bigotimes_{j=1}^n g'_j k_{\xi_j} \right) \right\rangle \\ &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \overline{\left( \prod_{j \in J} \tilde{g}_j(\xi_j) \right)} \prod_{j=1}^n \langle f'_j k_{\xi_j}, g'_j k_{\xi_j} \rangle \\ &= \left( \prod_{j \in I} \tilde{f}_j(\xi_j) \right) \overline{\left( \prod_{j \in J} \tilde{g}_j(\xi_j) \right)} \prod_{j=1}^n \widetilde{(f'_j g'_j)}(\xi_j) \end{aligned}$$

(By Lemma 3.1)



$$\begin{aligned}
& \text{(By Theorem 2.1)} \quad \rightarrow \left( \prod_{j \in I} f_j \right) \overline{\left( \prod_{j \in J} g_j \right)} \left( \prod_{j=1}^n f'_j \overline{g'_j} \right) \\
& = \left( \prod_{j \in I} f_j \right) \overline{\left( \prod_{j \in J} g_j \right)} \left( \prod_{j=1}^n f'_j \right) \left( \prod_{j=1}^n \overline{g'_j} \right) \\
& = \left( \prod_{j \in I} f_j \right) \left( \prod_{j \in J} \overline{g_j} \right) \left( \prod_{j \in I^c} f_j \right) \left( \prod_{j \in J^c} \overline{g_j} \right) \\
& = \left( \prod_{j=1}^n f_j \right) \left( \prod_{j=1}^n \overline{g_j} \right) \\
& = f \overline{g}.
\end{aligned}$$

This proves that

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow f \overline{g} \text{ radially.}$$

Suppose that  $f$  is a finite sum of elements of the form  $f_1 \otimes \cdots \otimes f_n$ , where  $f_j \in H^2(\mathbb{D})^{\alpha_j}$ , and  $g$  is a finite sum of elements of the form  $g_1 \otimes \cdots \otimes g_n$ , where  $g_j \in H^2(\mathbb{D})^{\beta_j}$  for all  $1 \leq j \leq n$ . That is,

$$f = \sum_{p=1}^l (f_{1p} \otimes \cdots \otimes f_{np}) \quad \text{and} \quad g = \sum_{q=1}^{l'} (g_{1q} \otimes \cdots \otimes g_{nq}),$$

where  $f_{jp} \in H^2(\mathbb{D})^{\alpha_j}$  and  $g_{jq} \in H^2(\mathbb{D})^{\beta_j}$  for all  $1 \leq j \leq n$ . Therefore,

$$\begin{aligned}
& \langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \\
& = \left\langle P_{H^2(\mathbb{D}^n)} \left( \left( \sum_{p=1}^l (f_{1p} \otimes \cdots \otimes f_{np}) \right) k_\xi \right), P_{H^2(\mathbb{D}^n)} \left( \left( \sum_{q=1}^{l'} (g_{1q} \otimes \cdots \otimes g_{nq}) \right) k_\xi \right) \right\rangle \\
& = \left\langle \sum_{p=1}^l P_{H^2(\mathbb{D}^n)}((f_{1p} \otimes \cdots \otimes f_{np})k_\xi), \sum_{q=1}^{l'} P_{H^2(\mathbb{D}^n)}((g_{1q} \otimes \cdots \otimes g_{nq})k_\xi) \right\rangle \\
& = \sum_{p=1}^l \sum_{q=1}^{l'} \langle P_{H^2(\mathbb{D}^n)}((f_{1p} \otimes \cdots \otimes f_{np})k_\xi), P_{H^2(\mathbb{D}^n)}((g_{1q} \otimes \cdots \otimes g_{nq})k_\xi) \rangle.
\end{aligned}$$

From the above, we obtain that

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow \sum_{p=1}^l \sum_{q=1}^{l'} (f_{1p} \otimes \cdots \otimes f_{np}) \overline{(g_{1q} \otimes \cdots \otimes g_{nq})} = f \overline{g} \text{ radially.}$$

Finally, assume that  $f \in H^2(\mathbb{D})^{\alpha_1} \otimes \cdots \otimes H^2(\mathbb{D})^{\alpha_n}$  and  $g \in H^2(\mathbb{D})^{\beta_1} \otimes \cdots \otimes H^2(\mathbb{D})^{\beta_n}$ . Then we shall show that

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow f \overline{g} \text{ radially.}$$

Now there exist sequences  $(f^{(m)})_{m=1}^\infty$  and  $(g^{(m)})_{m=1}^\infty$  such that  $\|f^{(m)} - f\|_{L^2(\mathbb{T}^n)} \rightarrow 0$  and  $\|g^{(m)} - g\|_{L^2(\mathbb{T}^n)} \rightarrow 0$  as  $m \rightarrow \infty$ , where each  $f^{(m)}$  and  $g^{(m)}$  are the finite sums of elements of the form  $f_1 \otimes \cdots \otimes f_n$  and  $g_1 \otimes \cdots \otimes g_n$ , respectively, and  $f_j \in H^2(\mathbb{D})^{\alpha_j}$  and  $g_j \in H^2(\mathbb{D})^{\beta_j}$  for all  $1 \leq j \leq n$ . Since  $f^{(m)} \rightarrow f$  in  $L^2(\mathbb{T}^n)$ , there exists a subsequence  $(f^{(m_t)})_{t=1}^\infty$  of  $(f^{(m)})_{m=1}^\infty$  which converges to  $f$  pointwise a.e. on  $\mathbb{T}^n$ . Similarly, there exists a subsequence  $(g^{(m_t)})_{t=1}^\infty$  of  $(g^{(m)})_{m=1}^\infty$  such that  $(g^{(m_t)})_{t=1}^\infty$  converges to  $g$  pointwise a.e. on  $\mathbb{T}^n$ . Thus

$$f^{(m_t)} \overline{g^{(m_t)}} \rightarrow f \overline{g} \text{ pointwise a.e. } (e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n.$$

Suppose  $E_0$  is a measurable null subset of  $\mathbb{T}^n$  such that

$$f^{(m_t)}(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g^{(m_t)}}(e^{i\theta_1}, \dots, e^{i\theta_n}) \rightarrow f(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g}(e^{i\theta_1}, \dots, e^{i\theta_n})$$

as  $t \rightarrow \infty$  for all  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n \setminus E_0$ . We have also shown that for each  $m \geq 1$ ,

$$\langle P_{H^2(\mathbb{D}^n)}(f^{(m)} k_\xi), P_{H^2(\mathbb{D}^n)}(g^{(m)} k_\xi) \rangle \rightarrow f^{(m)} \overline{g^{(m)}} \text{ radially.}$$

That means there exists a measurable null subset  $E_m$  of  $\mathbb{T}^n$  such that

$$\langle P_{H^2(\mathbb{D}^n)}(f^{(m)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g^{(m)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \rightarrow f^{(m)}(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g^{(m)}}(e^{i\theta_1}, \dots, e^{i\theta_n})$$

as  $r \rightarrow 1^-$ , and  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n \setminus E_m$ . Let  $E = \bigcup_{m=0}^\infty E_m$ . Then  $E$  is clearly a measurable null subset of  $\mathbb{T}^n$ . Also  $f^{(m)} k_\xi \rightarrow f k_\xi$  and  $g^{(m)} k_\xi \rightarrow g k_\xi$  in  $L^2(\mathbb{T}^n)$  because  $k_\xi \in L^\infty(\mathbb{T}^n)$ . Hence

$$P_{H^2(\mathbb{D}^n)}(f^{(m)} k_\xi) \rightarrow P_{H^2(\mathbb{D}^n)}(f k_\xi) \text{ and } P_{H^2(\mathbb{D}^n)}(g^{(m)} k_\xi) \rightarrow P_{H^2(\mathbb{D}^n)}(g k_\xi) \text{ in } H^2(\mathbb{D}^n).$$

as  $m \rightarrow \infty$ . Let  $\varepsilon > 0$  be any number and fix  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n \setminus E$ . Then there exists  $N_1 \in \mathbb{N}$  such that for  $m \geq N_1$

$$|\langle P_{H^2(\mathbb{D}^n)}(f k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle -$$

$$\langle P_{H^2(\mathbb{D}^n)}(f^{(m)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g^{(m)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle| < \frac{\varepsilon}{3}.$$

Now choose  $M > N_1$  such that

$$\left| f^{(m_M)}(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g^{(m_M)}}(e^{i\theta_1}, \dots, e^{i\theta_n}) - f(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g}(e^{i\theta_1}, \dots, e^{i\theta_n}) \right| < \frac{\varepsilon}{3}.$$

Also there exists  $s \in (0, 1)$  such that

$$|\langle P_{H^2(\mathbb{D}^n)}(f^{(m_M)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g^{(m_M)} k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle -$$

$$f^{(m_M)}(e^{i\theta_1}, \dots, e^{i\theta_n}) \overline{g^{(m_M)}}(e^{i\theta_1}, \dots, e^{i\theta_n})| < \frac{\varepsilon}{3}$$

for all  $r \in (s, 1)$ . Note that

$$\begin{aligned}
& \left| \langle P_{H^2(\mathbb{D}^n)}(fk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(gk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle - f(e^{i\theta_1}, \dots, e^{i\theta_n})\overline{g}(e^{i\theta_1}, \dots, e^{i\theta_n}) \right| \\
& \leq \left| \langle P_{H^2(\mathbb{D}^n)}(fk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(gk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \right. \\
& \quad \left. - \langle P_{H^2(\mathbb{D}^n)}(f^{(m_M)}k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g^{(m_M)}k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \right| \\
& + \left| \langle P_{H^2(\mathbb{D}^n)}(f^{(m_M)}k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(g^{(m_M)}k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \right. \\
& \quad \left. - f^{(m_M)}(e^{i\theta_1}, \dots, e^{i\theta_n})\overline{g^{(m_M)}}(e^{i\theta_1}, \dots, e^{i\theta_n}) \right| \\
& + \left| f^{(m_M)}(e^{i\theta_1}, \dots, e^{i\theta_n})\overline{g^{(m_M)}}(e^{i\theta_1}, \dots, e^{i\theta_n}) - f(e^{i\theta_1}, \dots, e^{i\theta_n})\overline{g}(e^{i\theta_1}, \dots, e^{i\theta_n}) \right|.
\end{aligned}$$

Hence

$$\begin{aligned}
& \left| \langle P_{H^2(\mathbb{D}^n)}(fk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}), P_{H^2(\mathbb{D}^n)}(gk_{(re^{i\theta_1}, \dots, re^{i\theta_n})}) \rangle \right. \\
& \quad \left. - f(e^{i\theta_1}, \dots, e^{i\theta_n})\overline{g}(e^{i\theta_1}, \dots, e^{i\theta_n}) \right| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon
\end{aligned}$$

whenever  $r \in (s, 1)$ .

This completes the proof.  $\blacksquare$

We have an immediate consequence of the above result as follows.

**Corollary 3.5.** *If  $f, g \in L^2(\mathbb{T}^n)$ , then*

$$\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle \rightarrow f\overline{g} \text{ radially.}$$

*Proof.* Note that

$$L^2(\mathbb{T}^n) = \bigoplus_{\alpha \in \{-1, 1\}^n} H^2(\mathbb{D}^n)^\alpha.$$

Let  $f, g \in L^2(\mathbb{T}^n)$  such that

$$f = \bigoplus_{\alpha \in \{-1, 1\}^n} f_\alpha \quad \text{and} \quad g = \bigoplus_{\beta \in \{-1, 1\}^n} g_\beta,$$

where  $f_\alpha \in H^2(\mathbb{D}^n)^\alpha$  and  $g_\beta \in H^2(\mathbb{D}^n)^\beta$  for all  $\alpha, \beta \in \{-1, 1\}^n$ . Therefore,

$$\begin{aligned}
\langle P_{H^2(\mathbb{D}^n)}(fk_\xi), P_{H^2(\mathbb{D}^n)}(gk_\xi) \rangle &= \sum_{\alpha \in \{-1, 1\}^n} \sum_{\beta \in \{-1, 1\}^n} \langle P_{H^2(\mathbb{D}^n)}(f_\alpha k_\xi), P_{H^2(\mathbb{D}^n)}(g_\beta k_\xi) \rangle \\
&\rightarrow \sum_{\alpha \in \{-1, 1\}^n} \sum_{\beta \in \{-1, 1\}^n} f_\alpha \overline{g_\beta} = f\overline{g} \quad \text{radially,}
\end{aligned}$$

which follows from Theorem 3.4.  $\blacksquare$

#### 4. SYMBOL MAP FOR $H^2(\mathbb{D}^n)$

In this section we shall discuss about the *symbol map* on the Toeplitz algebra on  $H^2(\mathbb{D}^n)$ . Before that, we need some results in the sequel.

**Theorem 4.1.** *For  $\varphi \in L^\infty(\mathbb{T}^n)$ ,  $\|H_\varphi k_\xi\| \rightarrow 0$  radially. Moreover, if  $\varphi, \psi \in L^\infty(\mathbb{T}^n)$ , then  $\widetilde{[T_\varphi, T_\psi]} \rightarrow 0$  radially.*

*Proof.* Suppose  $\varphi \in L^\infty(\mathbb{T}^n)$ . Then by definition

$$\begin{aligned} H_\varphi k_\xi &= P_{H^2(\mathbb{D}^n)^\perp}(\varphi k_\xi) \\ &= P_{H^2(\mathbb{D}^n)^\perp}((P_{H^2(\mathbb{D}^n)}\varphi + P_{H^2(\mathbb{D}^n)^\perp}\varphi)k_\xi) \\ &= P_{H^2(\mathbb{D}^n)^\perp}((P_{H^2(\mathbb{D}^n)}\varphi)k_\xi) + P_{H^2(\mathbb{D}^n)^\perp}((P_{H^2(\mathbb{D}^n)^\perp}\varphi)k_\xi). \end{aligned}$$

Since  $k_\xi \in H^\infty(\mathbb{D}^n)$ , clearly  $(P_{H^2(\mathbb{D}^n)}\varphi)k_\xi \in H^2(\mathbb{D}^n)$ . Therefore,

$$H_\varphi k_\xi = P_{H^2(\mathbb{D}^n)^\perp}((P_{H^2(\mathbb{D}^n)^\perp}\varphi)k_\xi).$$

Set  $y = P_{H^2(\mathbb{D}^n)^\perp}\varphi$ . Then

$$H_\varphi k_\xi = P_{H^2(\mathbb{D}^n)^\perp}(yk_\xi).$$

Again

$$yk_\xi = P_{H^2(\mathbb{D}^n)}(yk_\xi) + P_{H^2(\mathbb{D}^n)^\perp}(yk_\xi).$$

Thus  $\|yk_\xi\|^2 = \|P_{H^2(\mathbb{D}^n)}(yk_\xi)\|^2 + \|P_{H^2(\mathbb{D}^n)^\perp}(yk_\xi)\|^2$ , and hence

$$\|H_\varphi k_\xi\|^2 = \|yk_\xi\|^2 - \|P_{H^2(\mathbb{D}^n)}(yk_\xi)\|^2.$$

Now Lemma 3.1 yields

$$\|yk_\xi\|^2 = \langle yk_\xi, yk_\xi \rangle = \widetilde{|y|^2}(\xi).$$

Therefore,

$$(4.1) \quad \|H_\varphi k_\xi\|^2 = \widetilde{|y|^2}(\xi) - \|P_{H^2(\mathbb{D}^n)}(yk_\xi)\|^2.$$

Now Corollary 3.5 yields

$$\|P_{H^2(\mathbb{D}^n)}(yk_\xi)\|^2 \rightarrow |y|^2 \quad \text{radially.}$$

Hence, from equation (4.1), we conclude that

$$\|H_\varphi k_\xi\| \rightarrow 0 \quad \text{radially.}$$

Note that

$$\left| \widetilde{[T_\varphi, T_\psi]}(\xi) \right| = |\langle H_\varphi^* H_\psi k_\xi, k_\xi \rangle| = |\langle H_\psi k_\xi, H_\varphi k_\xi \rangle| \leq \|H_\psi k_\xi\| \|H_\varphi k_\xi\| \rightarrow 0 \quad \text{radially.}$$

This finishes the proof of the second part. ■

**Theorem 4.2.** *Let  $S \in \mathcal{S}$ . Then  $\widetilde{S} \rightarrow 0$  radially.*

*Proof.* Note that the set of all finite sums of operators of the form  $A[T_\varphi, T_\psi]B$ , where  $A, B \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ , forms a dense subset of  $\mathcal{S}$ . Now the set of all finite sums of operators of the form  $T_{\varphi_1}T_{\varphi_2}\cdots T_{\varphi_n}$  is dense in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  and hence the span of the operators of the form

$$T_{\varphi_1}T_{\varphi_2}\cdots T_{\varphi_n}[T_\varphi, T_\psi]T_{\psi_1}T_{\psi_2}\cdots T_{\psi_m}$$

form a dense subset of  $\mathcal{S}$ . Therefore, it suffices to prove for  $S = T_{\varphi_1}T_{\varphi_2}\cdots T_{\varphi_n}[T_\varphi, T_\psi]T_{\psi_1}T_{\psi_2}\cdots T_{\psi_m}$ , where  $\varphi, \varphi_i, \psi, \psi_i \in L^\infty(\mathbb{T}^n)$ . Again for any symbol  $u, v, w \in L^\infty(\mathbb{T}^n)$ ,

$$\begin{aligned} T_w[T_u, T_v] &= T_w T_{uv} - T_w T_u T_v \\ &= (T_w T_{uv} - T_{wuv}) + (T_{wuv} - T_{wu}T_v) + (T_{wu}T_v - T_w T_u T_v) \\ &= -[T_w, T_{uv}] + [T_{wu}, T_v] + [T_w, T_u]T_v. \end{aligned}$$

Thus, it is enough to assume that  $S \in \mathcal{S}$  is of the form

$$S = [T_\varphi, T_\psi]A = H_{\overline{\varphi}}^* H_\psi A \quad (A \in \mathcal{T}(L^\infty(\mathbb{T}^n)), \varphi, \psi \in L^\infty(\mathbb{T}^n)).$$

Therefore,

$$\begin{aligned} |\tilde{S}(\boldsymbol{\xi})| &= |\langle H_{\overline{\varphi}}^* H_\psi A k_\xi, k_\xi \rangle| \\ &= |\langle H_\psi A k_\xi, H_{\overline{\varphi}} k_\xi \rangle| \\ &= \|H_\psi A k_\xi\| \|H_{\overline{\varphi}} k_\xi\| \\ &\leq \|H_\psi A\| \|H_{\overline{\varphi}} k_\xi\| \rightarrow 0 \quad \text{radially,} \end{aligned}$$

which follows from Theorem 4.1. ■

We are now in a position to state the main result of this section.

**Theorem 4.3.** *Let  $T \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ . Then  $\tilde{T} \rightarrow \varphi$  radially for some  $\varphi \in L^\infty(\mathbb{T}^n)$ . Moreover, the map*

$$\Sigma : \mathcal{T}(L^\infty(\mathbb{T}^n)) \rightarrow L^\infty(\mathbb{T}^n), \quad T \mapsto \varphi$$

*is a  $C^*$ -algebra homomorphism with  $\ker(\Sigma) = \mathcal{S}$  and  $\Sigma(T_\varphi) = \varphi$  for any Toeplitz operator  $T_\varphi$ .*

*Proof.* We shall prove by induction that

$$\text{for any } m \geq 2 \text{ if } \varphi_1, \dots, \varphi_m \in L^\infty(\mathbb{T}^n), \text{ then } T_{\varphi_1} \cdots T_{\varphi_m} - T_{\varphi_1 \cdots \varphi_m} \in \mathcal{S}.$$

Recall that for any two symbols  $\varphi_1, \varphi_2 \in L^\infty(\mathbb{T}^n)$ ,  $T_{\varphi_1} T_{\varphi_2} - T_{\varphi_1 \varphi_2} = -[T_{\varphi_1}, T_{\varphi_2}] \in \mathcal{S}$ .

Suppose now  $T_{\varphi_1} \cdots T_{\varphi_m} - T_{\varphi_1 \cdots \varphi_m} \in \mathcal{S}$  for some  $m > 2$  and  $\varphi_1, \dots, \varphi_m \in L^\infty(\mathbb{T}^n)$ .

Note that

$$\begin{aligned} T_{\varphi_1} \cdots T_{\varphi_m} T_{\varphi_{m+1}} - T_{\varphi_1 \cdots \varphi_m \varphi_{m+1}} &= (T_{\varphi_1} \cdots T_{\varphi_m} T_{\varphi_{m+1}} - T_{\varphi_1 \cdots \varphi_m} T_{\varphi_{m+1}}) \\ &\quad + (T_{\varphi_1 \cdots \varphi_m} T_{\varphi_{m+1}} - T_{(\varphi_1 \cdots \varphi_m) \varphi_{m+1}}) \\ &= (T_{\varphi_1} \cdots T_{\varphi_m} - T_{\varphi_1 \cdots \varphi_m}) T_{\varphi_{m+1}} \\ &\quad - [T_{\varphi_1 \cdots \varphi_m}, T_{\varphi_{m+1}}]. \end{aligned}$$

Since  $\mathcal{S}$  is a closed ideal in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ , we have

$$T_{\varphi_1} \cdots T_{\varphi_m} T_{\varphi_{m+1}} - T_{\varphi_1 \cdots \varphi_m \varphi_{m+1}} \in \mathcal{S}.$$

Now the collection of all finite sums of operators of the form  $T_{\varphi_1} \cdots T_{\varphi_m}$  is dense in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ .

Thus operators of the form

$$T = T_\varphi + S \quad (\varphi \in L^\infty(\mathbb{T}^n), S \in \mathcal{S})$$

is a dense subset of  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ . Consider

$$\mathcal{T}_0(L^\infty(\mathbb{T}^n)) = \{T_\varphi + S : \varphi \in L^\infty(\mathbb{T}^n), S \in \mathcal{S}\}.$$

Suppose  $T = T_\varphi + S$  for some  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ . Then, Theorem 3.2 and Theorem 4.2 yield  $\tilde{T} \rightarrow \varphi$  radially. Define a map  $\Sigma_0 : \mathcal{T}_0(L^\infty(\mathbb{T}^n)) \rightarrow L^\infty(\mathbb{T}^n)$  as

$$(4.2) \quad \Sigma_0(T) = \varphi \quad (T = T_\varphi + S \in \mathcal{T}_0(L^\infty(\mathbb{T}^n))).$$

We shall show that  $\Sigma_0$  is well-defined. Suppose  $T_{\varphi_1} + S_1 = T_{\varphi_2} + S_2$ . Then

$$T_{\varphi_1} - T_{\varphi_2} = S_2 - S_1 \implies T_{\varphi_1 - \varphi_2} = S_2 - S_1.$$

Now Theorem 4.2 gives  $\widetilde{T}_{\varphi_1 - \varphi_2} = \widetilde{S_2 - S_1} \rightarrow 0$  radially. Again Theorem 3.2 infers  $\varphi_1 - \varphi_2 = 0$  a.e. and hence  $\varphi_1 = \varphi_2$  a.e..

Suppose  $T_{\varphi_1} + S_1, T_{\varphi_2} + S_2 \in \mathcal{T}_0(L^\infty(\mathbb{T}^n))$  and  $\alpha \in \mathbb{C}$ . Then

$$\begin{aligned} \Sigma_0(\alpha(T_{\varphi_1} + S_1) + (T_{\varphi_2} + S_2)) &= \Sigma_0(T_{\alpha\varphi_1 + \varphi_2} + \alpha S_1 + S_2) \\ \text{(By (4.2))} \quad &= \alpha\varphi_1 + \varphi_2 \\ &= \alpha\Sigma_0(T_{\varphi_1} + S_1) + \Sigma_0(T_{\varphi_2} + S_2). \end{aligned}$$

Again if  $T \in \mathfrak{B}(H^2(\mathbb{D}^n))$ , then

$$\left| \widetilde{T}(\boldsymbol{\xi}) \right| = |\langle Tk_{\boldsymbol{\xi}}, k_{\boldsymbol{\xi}} \rangle| \leq \|T\| \quad (\boldsymbol{\xi} \in \mathbb{D}^n).$$

Suppose  $T = T_\varphi + S$  for  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ . Then

$$\left| \widetilde{T}(re^{i\theta_1}, \dots, re^{i\theta_n}) \right| \leq \|T\| \quad \text{for all } 0 \leq r < 1, (e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n.$$

Letting  $r \rightarrow 1^-$ , we obtain  $|\varphi(e^{i\theta_1}, \dots, e^{i\theta_n})| \leq \|T\|$  a.e.  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n$ . Hence

$$\|\Sigma_0(T)\|_{L^\infty(\mathbb{T}^n)} = \|\varphi\|_{L^\infty(\mathbb{T}^n)} \leq \|T\|.$$

Therefore,  $\Sigma_0$  is a linear and bounded map from  $\mathcal{T}_0(L^\infty(\mathbb{T}^n))$  to  $L^\infty(\mathbb{T}^n)$ .

We now extend  $\Sigma_0$  to  $\overline{\mathcal{T}_0(L^\infty(\mathbb{T}^n))} = \mathcal{T}(L^\infty(\mathbb{T}^n))$ , denoting its extension by  $\Sigma$ . Thus  $\Sigma : \mathcal{T}(L^\infty(\mathbb{T}^n)) \rightarrow L^\infty(\mathbb{T}^n)$  is a linear and bounded map. Suppose  $T \in \mathcal{T}(L^\infty(\mathbb{T}^n))$  and set  $\varphi = \Sigma(T)$ . Then there exists a sequence  $\{T_m\}_{m=1}^\infty$ , where each  $T_m = T_{\varphi_m} + S_m \in \mathcal{T}_0(L^\infty(\mathbb{T}^n))$ , such that  $\|T_m - T\| \rightarrow 0$  as  $m \rightarrow \infty$ . Therefore,

$$\|T_{\varphi_m} - T_\varphi\| = \|T_{\varphi_m - \varphi}\| = \|\varphi_m - \varphi\|_{L^\infty(\mathbb{T}^n)} = \|\Sigma(T_m) - \Sigma(T)\|_{L^\infty(\mathbb{T}^n)} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Thus

$$S_m = T_m - T_{\varphi_m} \rightarrow T - T_\varphi \quad \text{as } m \rightarrow \infty.$$

Let  $S = T - T_\varphi$ . Then  $S \in \mathcal{S}$  as  $\{S_m\}$  in  $\mathcal{S}$  and  $\mathcal{S}$  is closed, and hence if  $T \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ , then  $T = T_\varphi + S$ , where  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ . Thus

$$\mathcal{T}(L^\infty(\mathbb{T}^n)) = \{T_\varphi + S : \varphi \in L^\infty(\mathbb{T}^n), S \in \mathcal{S}\}.$$

Therefore, Theorem 3.2 and Theorem 4.2 give  $\widetilde{T} \rightarrow \varphi$  radially.

To show  $\Sigma$  is a  $C^*$ -algebra homomorphism, let  $T = T_\varphi + S$ , where  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ . Note that

$$(\widetilde{T_\varphi + S})^* = \widetilde{T_\varphi + S}^* = \widetilde{T_\varphi} + \widetilde{S}^* = \widetilde{T_\varphi} + \widetilde{\overline{S}} \rightarrow \overline{\varphi} \quad \text{radially.}$$

Therefore,  $\Sigma(T^*) = \overline{\Sigma(T)}$ . Suppose  $T_1 = T_{\varphi_1} + S_1$  and  $T_2 = T_{\varphi_2} + S_2$ , where  $\varphi_1, \varphi_2 \in L^\infty(\mathbb{T}^n)$  and  $S_1, S_2 \in \mathcal{S}$ . Then

$$\begin{aligned} T_1 T_2 &= T_{\varphi_1} T_{\varphi_2} + T_{\varphi_1} S_2 + S_1 T_{\varphi_2} + S_1 S_2 \\ &= T_{\varphi_1 \varphi_2} + (T_{\varphi_1} T_{\varphi_2} - T_{\varphi_1 \varphi_2}) + (T_{\varphi_1} S_2 + S_1 T_{\varphi_2} + S_1 S_2). \end{aligned}$$

Since  $\mathcal{S}$  is an ideal in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ , clearly  $(T_{\varphi_1}T_{\varphi_2} - T_{\varphi_1\varphi_2}) + (T_{\varphi_1}S_2 + S_1T_{\varphi_2} + S_1S_2) \in \mathcal{S}$ . Hence

$$\widetilde{T_1T_2} \rightarrow \varphi_1\varphi_2 \text{ radially.}$$

It follows that  $\Sigma(T_1T_2) = \Sigma(T_1)\Sigma(T_2)$  for any  $T_1, T_2 \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ .

Finally, it is clear from the definition that  $\ker(\Sigma) = \mathcal{S}$ , the semicommutator ideal  $\mathcal{S}$  in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ , and  $\Sigma(T_\varphi) = \varphi$  for all  $\varphi \in L^\infty(\mathbb{T}^n)$ . This finishes the proof.  $\blacksquare$

The next result readily follows from the above theorem.

**Corollary 4.4.** *For  $T \in \mathcal{T}(L^\infty(\mathbb{T}^n))$ ,  $[\widetilde{T^*}, T] \rightarrow 0$  radially.*

We shall now show that the symbol map can not be extended to the whole of  $\mathfrak{B}(H^2(\mathbb{D}^n))$  in the following example (see [12] for  $n = 1$  case).

**Example 4.5.** Consider the operator  $T$  on  $H^2(\mathbb{D})$ , defined by

$$Tz^k = z^{2k+1} \quad (k \in \mathbb{Z}_+),$$

where  $\{z^k\}_{k \in \mathbb{Z}_+}$  denotes the standard orthonormal basis for  $H^2(\mathbb{D})$ . Now

$$\langle Tz^k, Tz^l \rangle = \langle z^{2k+1}, z^{2l+1} \rangle = \delta_{k,l} = \langle z^k, z^l \rangle, \quad (k, l \in \mathbb{Z}_+).$$

Therefore,  $T$  is an isometry on  $H^2(\mathbb{D})$ . Thus,  $T^*T = I_{H^2(\mathbb{D})}$  and  $TT^* = P_{\mathcal{R}(T)}$ , the orthogonal projection onto the range  $\mathcal{R}(T) = \bigvee_{k=0}^{\infty} \{z^{2k+1}\}$ . Hence

$$(T^*T - TT^*)z^k = (I - P_{\mathcal{R}(T)})z^k = \begin{cases} 0 & \text{if } k \text{ is odd;} \\ z^k & \text{if } k \text{ is even.} \end{cases}$$

That means  $[\widetilde{T^*}, T]$  is an orthogonal projection onto  $\mathcal{M} = \bigvee \{z^{2k} : k \in \mathbb{Z}_+\}$ . Therefore, for  $\xi \in \mathbb{D}$

$$\begin{aligned} [\widetilde{T^*}, T](\xi) &= \|P_{\mathcal{M}}k_\xi\|^2 = (1 - |\xi|^2)\|P_{\mathcal{M}}K_\xi\|^2 \\ &= (1 - |\xi|^2) \left\| P_{\mathcal{M}} \left( \sum_{k=0}^{\infty} \bar{\xi}^k z^k \right) \right\|^2 \\ &= (1 - |\xi|^2) \left\| \sum_{k=0}^{\infty} \bar{\xi}^{2k} z^{2k} \right\|^2 \\ &= (1 - |\xi|^2) \sum_{k=0}^{\infty} |\xi|^{4k} \\ &= \frac{(1 - |\xi|^2)}{(1 - |\xi|^4)} = \frac{1}{1 + |\xi|^2} \rightarrow \frac{1}{2} \text{ radially.} \end{aligned}$$

Now we consider the operator  $S$  on  $H^2(\mathbb{D}^n)$  as

$$S = \overbrace{T \otimes \cdots \otimes T}^{n\text{-times}}.$$

Then

$$\begin{aligned}
S^*S - SS^* &= (T \otimes \cdots \otimes T)^*(T \otimes \cdots \otimes T) - (T \otimes \cdots \otimes T)(T \otimes \cdots \otimes T)^* \\
&= (T^*T \otimes T^*T \otimes \cdots \otimes T^*T) - (TT^* \otimes TT^* \otimes \cdots \otimes TT^*) \\
&= I_{H^2(\mathbb{D}^n)} - (TT^* \otimes TT^* \otimes \cdots \otimes TT^*).
\end{aligned}$$

Thus for  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_+^n$ ,

$$[S^*, S]z^{\mathbf{k}} = \begin{cases} z^{\mathbf{k}} & \text{if } k_j \text{ is even for some } j; \\ 0 & \text{if } k_j \text{ is odd for all } j. \end{cases}$$

Hence  $[S^*, S]$  is an orthogonal projection onto

$$\mathcal{N} = \bigvee \{z^{\mathbf{k}} : \mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_+^n, k_j \text{ is even for some } 1 \leq j \leq n\},$$

that is,  $[S^*, S] = P_{\mathcal{N}}$ .

Now for  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$ ,

$$\begin{aligned}
\widetilde{[S^*, S]}(\boldsymbol{\xi}) &= \|P_{\mathcal{N}}k_{\boldsymbol{\xi}}\|^2 = \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left\| P_{\mathcal{N}} \left( \sum_{\mathbf{k} \in \mathbb{Z}_+^n} \bar{\boldsymbol{\xi}}^{\mathbf{k}} z^{\mathbf{k}} \right) \right\|^2 \\
&= \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left\| \sum_{\substack{\mathbf{k} \in \mathbb{Z}_+^n \\ \text{some } k_j \text{ even}}} \bar{\boldsymbol{\xi}}^{\mathbf{k}} z^{\mathbf{k}} \right\|^2 \\
&= \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left( \sum_{\text{some } k_j \text{ even}} |\xi_1|^{2k_1} \cdots |\xi_n|^{2k_n} \right) \\
&\geq \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left( \sum_{\text{all } k_j \text{ even}} |\xi_1|^{2k_1} \cdots |\xi_n|^{2k_n} \right) \\
&= \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left( \sum_{k_1 \text{ even}} |\xi_1|^{2k_1} \right) \cdots \left( \sum_{k_n \text{ even}} |\xi_n|^{2k_n} \right) \\
&= \left( \prod_{j=1}^n (1 - |\xi_j|^2) \right) \left( \sum_{m=0}^{\infty} |\xi_1|^{4m} \right) \cdots \left( \sum_{m=0}^{\infty} |\xi_n|^{4m} \right) \\
&= \frac{\prod_{j=1}^n (1 - |\xi_j|^2)}{\prod_{j=1}^n (1 - |\xi_j|^4)} = \prod_{j=1}^n (1 + |\xi_j|^2)^{-1} \geq \frac{1}{2^n} > 0.
\end{aligned}$$

Hence, by corollary 4.4 we conclude that the symbol calculus can not be extended to any  $C^*$ -algebra that contains  $S$ .



## 5. EXTENSION OF TOEPLITZ ALGEBRA

In this section, we present a family of  $C^*$ -algebras containing the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  and the analogs of Theorem 4.3 still hold.

Let  $\psi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ . Consider

$$(5.1) \quad \mathcal{B}(\psi, S) = \{A \in \mathfrak{B}(H^2(\mathbb{D}^n)) : \|A(T_\psi + S)k_\xi\| \text{ and } \|A^*(T_\psi + S)k_\xi\| \rightarrow 0 \text{ radially}\}.$$

In particular, if  $\psi = 1$  a.e. on  $\mathbb{T}^n$  and  $S = \mathbf{0}$ , where  $\mathbf{0}$  denotes the zero operator on  $H^2(\mathbb{D}^n)$ , then we simply write  $\mathcal{B} = \mathcal{B}(1, \mathbf{0})$ .

**Lemma 5.1.**  $\mathcal{B}(\psi, \mathbf{0})$  is a  $C^*$ -algebra for all  $\psi \in L^\infty(\mathbb{T}^n)$ .

*Proof.* Let  $A_1, A_2 \in \mathcal{B}(\psi, \mathbf{0})$  and  $c \in \mathbb{C}$ . Then  $cA_1 + A_2, A_1A_2 \in \mathcal{B}(\psi, \mathbf{0})$  as

$$\begin{aligned} \|(cA_1 + A_2)T_\psi k_\xi\| &\leq |c|\|A_1T_\psi k_\xi\| + \|A_2T_\psi k_\xi\|, \\ \|(cA_1 + A_2)^*T_\psi k_\xi\| &\leq |c|\|A_1^*T_\psi k_\xi\| + \|A_2^*T_\psi k_\xi\|, \end{aligned}$$

$$\|A_1A_2T_\psi k_\xi\| \leq \|A_1\|\|A_2T_\psi k_\xi\| \quad \text{and} \quad \|(A_1A_2)^*T_\psi k_\xi\| \leq \|A_2^*\|\|A_1^*T_\psi k_\xi\|.$$

From the definition, it follows that if  $A \in \mathcal{B}(\psi, \mathbf{0})$ , then  $A^* \in \mathcal{B}(\psi, \mathbf{0})$ .

To see that  $\mathcal{B}(\psi, \mathbf{0})$  is closed, let  $(A_m)$  be a sequence in  $\mathcal{B}(\psi, \mathbf{0})$  such that  $\|A - A_m\| \rightarrow 0$  as  $m \rightarrow \infty$  for some  $A \in \mathfrak{B}(H^2(\mathbb{D}^n))$ . Now for any  $\varepsilon > 0$  choose  $N \in \mathbb{N}$  such that  $\|A - A_N\| < \frac{\varepsilon}{2(1+\|T_\psi\|)}$ . Therefore,

$$(5.2) \quad \begin{aligned} \|AT_\psi k_\xi\| &\leq \|(A - A_N)T_\psi k_\xi\| + \|A_NT_\psi k_\xi\| \\ &\leq \|A - A_N\|\|T_\psi\| + \|A_NT_\psi k_\xi\| \\ &< \varepsilon/2 + \|A_NT_\psi k_\xi\|. \end{aligned}$$

Since  $\|A_mT_\psi k_\xi\| \rightarrow 0$  radially for each  $m \in \mathbb{N}$ , there exists a measure zero subset  $E_m$  of  $\mathbb{T}^n$  such that

$$\|A_mT_\psi k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}\| \rightarrow 0 \quad \text{as } r \rightarrow 1^-,$$

for all  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n \setminus E_m$ . Consider  $E = \cup_{m=1}^\infty E_m$  of  $\mathbb{T}^n$ , and hence  $E$  is a measure zero subset. Thus for a fixed  $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in \mathbb{T}^n \setminus E$ , we choose  $0 < s < 1$  such that  $\|A_NT_\psi k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}\| < \varepsilon/2$  whenever  $s < r < 1$ . Hence the equation (5.2) implies that

$$s < r < 1 \implies \|AT_\psi k_{(re^{i\theta_1}, \dots, re^{i\theta_n})}\| < \varepsilon.$$

Therefore,  $\|AT_\psi k_\xi\| \rightarrow 0$  radially. Now  $\|A_m^* - A^*\| = \|A_m - A\| \rightarrow 0$  as  $m \rightarrow \infty$  and  $\|A_m^*T_\psi k_\xi\| \rightarrow 0$  radially for all  $m \in \mathbb{N}$ . It follows that  $\|A^*T_\psi k_\xi\| \rightarrow 0$  radially. Thus  $A \in \mathcal{B}(\psi, \mathbf{0})$  and hence  $\mathcal{B}(\psi, \mathbf{0})$  is closed.  $\blacksquare$

**Lemma 5.2.**  $\mathcal{C} \subseteq \mathcal{S} \subseteq \mathcal{B}$ .

*Proof.* Firstly, it is easy to see that  $\mathcal{C} \subseteq \mathcal{S}$ . Let  $\varphi, \psi \in L^\infty(\mathbb{T}^n)$ . Then

$$\begin{aligned} \|[T_\varphi, T_\psi]k_\xi\| &= \|H_\varphi^*H_\psi k_\xi\| \leq \|H_\varphi^*\|\|H_\psi k_\xi\|, \\ \text{and } \|[T_\varphi, T_\psi]^*k_\xi\| &= \|(H_\varphi^*H_\psi)^*k_\xi\| \leq \|H_\psi^*\|\|H_\varphi k_\xi\|. \end{aligned}$$

Now Theorem 4.1 infers that  $[T_\varphi, T_\psi] \in \mathcal{B}$ , and hence  $\mathcal{C} \subseteq \mathcal{S} \subseteq \mathcal{B}$ .  $\blacksquare$

**Lemma 5.3.**  $\mathcal{B}(\psi, S) = \mathcal{B}(\psi, \mathbf{0})$  for all  $\psi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ .

*Proof.* It follows easily from Lemma 5.2 that  $\mathcal{B}(\psi, \mathbf{0}) \subseteq \mathcal{B}(\psi, S)$ .

Suppose  $\psi \in L^\infty(\mathbb{T}^n)$  and  $S \in \mathcal{S}$ , and let  $A \in \mathcal{B}(\psi, S)$ . Then  $\|Sk_\xi\| \rightarrow 0$  radially, and we also have

$$\begin{aligned} \|AT_\psi k_\xi\| &\leq \|A(T_\psi + S)k_\xi\| + \|A\|\|Sk_\xi\| \rightarrow 0, \\ \|A^*T_\psi k_\xi\| &\leq \|A^*(T_\psi + S)k_\xi\| + \|A\|\|Sk_\xi\| \rightarrow 0. \end{aligned}$$

Therefore,  $A \in \mathcal{B}(\psi, \mathbf{0})$ . ■

**Theorem 5.4.** Let  $\varphi \in L^2(\mathbb{T}^n)$ . Then  $\|P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) - \tilde{\varphi}(\xi)k_\xi\| \rightarrow 0$  radially. In particular,  $\|T_\varphi k_\xi - \tilde{\varphi}(\xi)k_\xi\| \rightarrow 0$  radially for  $\varphi \in L^\infty(\mathbb{T}^n)$ .

*Proof.* First, assume that  $\varphi \in L^2(\mathbb{T}^n)$ . Then we have

$$\begin{aligned} &\|P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) - \tilde{\varphi}(\xi)k_\xi\|^2 \\ &= \|P_{H^2(\mathbb{D}^n)}(\varphi k_\xi)\|^2 + \|\tilde{\varphi}(\xi)k_\xi\|^2 - \langle P_{H^2(\mathbb{D}^n)}(\varphi k_\xi), \tilde{\varphi}(\xi)k_\xi \rangle - \langle \tilde{\varphi}(\xi)k_\xi, P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) \rangle \\ &= \|P_{H^2(\mathbb{D}^n)}(\varphi k_\xi)\|^2 + \|\tilde{\varphi}(\xi)k_\xi\|^2 - \overline{\tilde{\varphi}(\xi)} \langle P_{H^2(\mathbb{D}^n)}(\varphi k_\xi), k_\xi \rangle - \tilde{\varphi}(\xi) \langle k_\xi, P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) \rangle. \end{aligned}$$

Now Corollary 3.5 yields

$$\|P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) - \tilde{\varphi}(\xi)k_\xi\| \rightarrow 0 \quad \text{radially.}$$

Finally, if  $\varphi \in L^\infty(\mathbb{T}^n)$ , then  $P_{H^2(\mathbb{D}^n)}(\varphi k_\xi) = T_\varphi k_\xi$  and hence the required result follows. ■

**Proposition 5.5.** If  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $A \in \mathcal{B}(\psi, \mathbf{0})$ , then  $T_\varphi A, AT_\varphi \in \mathcal{B}(\psi, \mathbf{0})$ .

*Proof.* To show  $T_\varphi A$  and  $AT_\varphi$  are in  $\mathcal{B}(\psi, \mathbf{0})$  for  $A \in \mathcal{B}(\psi, \mathbf{0})$  and  $\varphi \in L^\infty(\mathbb{T}^n)$ , it suffices to check that  $\|AT_\varphi T_\psi k_\xi\| \rightarrow 0$  radially. Note that

$$(5.3) \quad \begin{aligned} \|AT_\varphi T_\psi k_\xi\| &\leq \|AT_\varphi T_\psi k_\xi - \tilde{\varphi}(\xi)AT_\psi k_\xi\| + \|\tilde{\varphi}(\xi)AT_\psi k_\xi\| \\ &\leq \|A\|\|T_\varphi T_\psi k_\xi - \tilde{\varphi}(\xi)T_\psi k_\xi\| + \|\varphi\|_\infty \|AT_\psi k_\xi\|. \end{aligned}$$

Also

$$\|T_\varphi T_\psi k_\xi - \tilde{\varphi}(\xi)T_\psi k_\xi\|^2 = (T_\psi^* T_\varphi^* T_\varphi T_\psi)(\xi) + |\tilde{\varphi}(\xi)|^2 \|T_\psi k_\xi\|^2 - 2\operatorname{Re} \left( \overline{\tilde{\varphi}(\xi)} (T_\psi^* T_\varphi T_\psi)(\xi) \right).$$

Applying Theorems 2.1, 4.3, and 3.2, we obtain

$$\|T_\varphi T_\psi k_\xi - \tilde{\varphi}(\xi)T_\psi k_\xi\|^2 \rightarrow |\varphi\psi|^2 + |\varphi\psi|^2 - 2|\varphi\psi|^2 = 0 \quad \text{radially.}$$

Hence  $\|AT_\varphi T_\psi k_\xi\| \rightarrow 0$  radially follows from (5.3). ■

An immediate consequence of Proposition 5.5 is the following.

**Lemma 5.6.** Let  $\psi_1$  and  $\psi_2$  be two inner functions on  $\mathbb{D}^n$  and let  $\psi = \overline{\psi_1}\psi_2$ . Then  $\mathcal{B}(\psi, \mathbf{0}) = \mathcal{B}$ .

*Proof.* Follows from Proposition 5.5. ■

Now we record the following properties of the collection  $\mathcal{B}$  defined in (5.1):

**Remark 5.7.** Suppose  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $T_\varphi \in \mathcal{B}(\psi, \mathbf{0})$ , where  $|\psi| > 0$  a.e. on  $\mathbb{T}^n$ . Then

$$|\widetilde{T_{\varphi\psi}}(\boldsymbol{\xi})| \leq \|T_{\varphi\psi}k_{\boldsymbol{\xi}}\| \leq \|[T_\varphi, T_\psi]k_{\boldsymbol{\xi}}\| + \|T_\varphi T_\psi k_{\boldsymbol{\xi}}\| \rightarrow 0 \quad \text{radially.}$$

Now Theorem 3.2 yields that  $\varphi\psi \equiv 0$  a.e. on  $\mathbb{T}^n$ . But  $|\psi| > 0$  a.e. on  $\mathbb{T}^n$ , hence  $\varphi \equiv 0$  a.e. on  $\mathbb{T}^n$ . Thus, any Toeplitz operator in  $\mathcal{B}(\psi, \mathbf{0})$  corresponds to the symbol 0.

For  $\psi \in L^\infty(\mathbb{T}^n)$  with  $|\psi| = 1$  a.e. on  $\mathbb{T}^n$ , consider

$$\mathcal{A}_\psi := \{T_\varphi + A : \varphi \in L^\infty(\mathbb{T}^n), A \in \mathcal{B}(\psi, \mathbf{0})\}.$$

**Theorem 5.8.** *The collection  $\mathcal{A}_\psi$  satisfies the following:*

- (1) *If  $T \in \mathcal{A}_\psi$ , then  $\widetilde{T} \rightarrow \varphi$  radially for some  $\varphi \in L^\infty(\mathbb{T}^n)$ .*
- (2)  *$\mathcal{A}_\psi$  is a  $C^*$ -algebra.*
- (3) *Define*

$$\Sigma_\psi : \mathcal{A}_\psi \rightarrow L^\infty(\mathbb{T}^n) \quad \text{by} \quad \Sigma_\psi(T) = \varphi,$$

where  $\varphi$  is as in (1). Then  $\Sigma_\psi$  induces a  $C^*$ -algebra homomorphism from  $\mathcal{A}_\psi$  onto  $L^\infty(\mathbb{T}^n)$  such that  $\Sigma_\psi(T_\varphi) = \varphi$  for all  $\varphi \in L^\infty(\mathbb{T}^n)$ .

*Proof.* (1) Let  $T \in \mathcal{A}_\psi$ . Then  $T = T_\varphi + A$  for some  $\varphi \in L^\infty(\mathbb{T}^n)$ ,  $A \in \mathcal{B}(\psi, \mathbf{0})$ . Now Theorem 3.2 implies that

$$\widetilde{T} = \widetilde{T}_\varphi + \widetilde{A} \rightarrow \varphi \quad \text{radially.}$$

Also

$$\begin{aligned} |\widetilde{A}(\boldsymbol{\xi})| &= |\langle Ak_{\boldsymbol{\xi}}, k_{\boldsymbol{\xi}} \rangle| \\ &= |\langle \psi Ak_{\boldsymbol{\xi}}, \psi k_{\boldsymbol{\xi}} \rangle| \\ &\leq |\langle P_{H^2(\mathbb{D}^n)}(\psi Ak_{\boldsymbol{\xi}}), P_{H^2(\mathbb{D}^n)}(\psi k_{\boldsymbol{\xi}}) \rangle| + |\langle P_{H^2(\mathbb{D}^n)^\perp}(\psi Ak_{\boldsymbol{\xi}}), P_{H^2(\mathbb{D}^n)^\perp}(\psi k_{\boldsymbol{\xi}}) \rangle| \\ &= |\langle T_\psi Ak_{\boldsymbol{\xi}}, T_\psi k_{\boldsymbol{\xi}} \rangle| + |\langle H_\psi Ak_{\boldsymbol{\xi}}, H_\psi k_{\boldsymbol{\xi}} \rangle| \\ &\leq \|A^* T_\psi^* T_\psi k_{\boldsymbol{\xi}}\| + \|H_\psi A\| \|H_\psi k_{\boldsymbol{\xi}}\|. \end{aligned}$$

Thus, Proposition 5.5 and Theorem 4.1 infer that  $\widetilde{A}(\boldsymbol{\xi}) \rightarrow 0$  radially.

- (2) It is immediate from Lemma 5.1 that  $\mathcal{A}_\psi$  is linear space and closed under adjoint. Let  $\varphi_1, \varphi_2 \in L^\infty(\mathbb{T}^n)$ , and  $A_1, A_2 \in \mathcal{B}(\psi, \mathbf{0})$ . Then

$$\begin{aligned} (T_{\varphi_1} + A_1)(T_{\varphi_2} + A_2) &= T_{\varphi_1}T_{\varphi_2} + T_{\varphi_1}A_2 + A_1T_{\varphi_2} + A_1A_2 \\ (5.4) \quad &= T_{\varphi_1\varphi_2} - ([T_{\varphi_1}, T_{\varphi_2}] - T_{\varphi_1}A_2 - A_1T_{\varphi_2} - A_1A_2). \end{aligned}$$

Using Lemma 5.1, Lemma 5.2, and Proposition 5.5, we have  $\mathcal{A}_\psi$  is closed under multiplication.

To show that  $\mathcal{A}_\psi$  is closed, we show  $\mathcal{A}_\psi = \overline{\mathcal{A}_\psi}$ . Consider

$$\eta_0 : \mathcal{A}_\psi \rightarrow L^\infty(\mathbb{T}^n) \quad \text{defined by} \quad \eta_0(T_\varphi + A) = \varphi.$$

It is easily seen that  $\eta_0$  is a well-defined bounded linear map from  $\mathcal{A}_\psi$  onto  $L^\infty(\mathbb{T}^n)$ . Suppose  $\eta$  is the bounded linear extension of  $\eta_0$  from  $\overline{\mathcal{A}_\psi}$  onto  $L^\infty(\mathbb{T}^n)$ . Assume that

$T \in \overline{\mathcal{A}_\psi}$ . Then there exists a sequence  $\{T_{\varphi_m} + A_m\}$  in  $\mathcal{A}_\psi$  such that  $\|(T_{\varphi_m} + A_m) - T\| \rightarrow 0$  as  $m \rightarrow \infty$ , and hence  $\varphi_m = \eta(T_{\varphi_m} + A_m) \rightarrow \eta(T)$  in  $L^\infty(\mathbb{T}^n)$ . Thus

$$\|T_{\varphi_m} - T_{\eta(T)}\| = \|\varphi_m - \eta(T)\|_\infty \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Therefore,  $A_m = (T_{\varphi_m} + A_m) - T_{\varphi_m} \rightarrow T - T_{\eta(T)}$  as  $m \rightarrow \infty$ . Since  $\mathcal{B}(\psi, \mathbf{0})$  is closed,  $T - T_{\eta(T)} \in \mathcal{B}(\psi, \mathbf{0})$ . Therefore,  $T = T_{\eta(T)} + (T - T_{\eta(T)}) \in \mathcal{A}_\psi$ .

- (3) It is easy to check that the mapping  $\Sigma_\psi$  is well-defined linear, preserves conjugation. Also (5.4) yields that the map is multiplicative too. Thus  $\Sigma_\psi$  is a  $C^*$ -algebra homomorphism from  $\mathcal{A}_\psi$  onto  $L^\infty(\mathbb{T}^n)$ , and also  $\Sigma_\psi(T_\varphi) = \varphi$  with  $\ker(\Sigma_\psi) = \mathcal{B}(\psi, \mathbf{0})$ . ■

It is obvious from Lemma 5.2 that  $\mathcal{T}(L^\infty(\mathbb{T}^n)) \subseteq \mathcal{A}_\psi$ . Following example shows that Theorem 5.8 is indeed an extension of Theorem 4.3.

**Example 5.9.** Let  $P = P_{\mathcal{M}}$  be the orthogonal projection of  $H^2(\mathbb{D})$  onto the closed subspace  $\mathcal{M}$ , where  $\mathcal{M} := \bigvee_{m=0}^\infty \{z^{2^m}\}$ . Then for  $m \geq 1$

$$\|(T_z P - P T_z) z^{2^m}\| = \|z^{2^{m+1}} - P z^{2^{m+1}}\| = 1.$$

But  $z^{2^m} \rightarrow 0$  weakly as  $m \rightarrow \infty$ , and hence  $T_z P - P T_z$  is not a compact operator. Therefore  $P \notin \mathcal{T}(L^\infty(\mathbb{T}^n))$  (see [2]). Also it is shown in [12, Example 8] that  $\|P k_{r e^{i\theta}}\| \rightarrow 0$  as  $r \rightarrow 1^-$  for a.e.  $e^{i\theta} \in \mathbb{T}$ . Consider an operator  $A$  on  $H^2(\mathbb{D}^n)$  defined as

$$A = \overbrace{P \otimes I \otimes \cdots \otimes I}^{n\text{-times}},$$

where  $I$  denotes the identity operator on  $H^2(\mathbb{D})$ . Now for  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{D}^n$ ,

$$\begin{aligned} \|A k_{\boldsymbol{\xi}}\| &= \|(P \otimes I \otimes \cdots \otimes I)(k_{\xi_1} \otimes k_{\xi_2} \otimes \cdots \otimes k_{\xi_n})\| \\ &= \|P k_{\xi_1} \otimes k_{\xi_2} \otimes \cdots \otimes k_{\xi_n}\| \\ &= \|P k_{\xi_1}\| \|k_{\xi_2}\| \cdots \|k_{\xi_n}\| \\ &= \|P k_{\xi_1}\|. \end{aligned}$$

It follows that  $\|A k_{\boldsymbol{\xi}}\| \rightarrow 0$  radially, and  $A \in \mathcal{A}_\psi$ , in fact  $A \in \mathcal{B}$ . Let  $T_{z_i}$  denote the multiplication operator on  $H^2(\mathbb{D}^n)$  by the  $i$ -th coordinate function  $z_i$  for  $1 \leq i \leq n$ , that is,  $(T_{z_i} f)(\boldsymbol{\xi}) = \xi_i f(\boldsymbol{\xi})$  for all  $\boldsymbol{\xi} \in \mathbb{D}^n$ ,  $f \in H^2(\mathbb{D}^n)$ . Consider

$$\mathbb{E} := \{T \in \mathfrak{B}(H^2(\mathbb{D}^n)) : T_{z_i}^* T T_{z_i} - T \text{ is compact for all } 1 \leq i \leq n\}.$$

Then  $\mathbb{E}$  is a norm-closed algebra. Now if  $T_\varphi$  is Toeplitz operator on  $H^2(\mathbb{D}^n)$ , then  $T_{z_i}^* T_\varphi T_{z_i} = T_\varphi$  for all  $1 \leq i \leq n$  (see [15]). Hence  $\mathbb{E}$  contains all Toeplitz operators on  $H^2(\mathbb{D}^n)$ . We show that  $A \notin \mathcal{T}(L^\infty(\mathbb{T}^n))$  by showing that  $A \notin \mathbb{E}$ . Now

$$T_{z_1}^* A T_{z_1} - A = (T_{z_1}^* P T_{z_1} - P) \otimes I \otimes \cdots \otimes I,$$

and  $T_{z_1}^* P T_{z_1} - P$  is not compact. Therefore,  $T_{z_1}^* A T_{z_1} - A$  is not compact (see [14]) and hence  $A \notin \mathbb{E}$ .

Denote

$$\mathcal{A} = \{T_\varphi + A : \varphi \in L^\infty(\mathbb{T}^n), A \in \mathcal{B}\}.$$

Note that if  $A \in \mathcal{B}$  then Proposition 5.5 gives that  $A \in \mathcal{B}(\psi, \mathbf{0})$  for each  $\psi \in L^\infty(\mathbb{T}^n)$ , and hence  $\mathcal{A} \subseteq \mathcal{A}_\psi$ . In the next result we prove that the class  $\mathcal{A}$  is maximal onto which the *symbol map* exists and  $T_\varphi \rightarrow \varphi$ , provided that the symbol of an operator is obtained as the radial limit of its *Berezin transform*.

**Proposition 5.10.** *Let  $\mathcal{A}$  be any closed  $C^*$ -subalgebra of  $\mathfrak{B}(H^2(\mathbb{D}^n))$  containing Toeplitz operators such that there exists a  $C^*$ -homomorphism  $\mathfrak{S}$  from  $\mathcal{A}$  onto  $L^\infty(\mathbb{T}^n)$  with  $\mathfrak{S}(T_\varphi) = \varphi$  for all  $\varphi \in L^\infty(\mathbb{T}^n)$ , and for each  $T \in \mathcal{A}$ ,  $\mathfrak{S}(T)$  is the radial limit of  $\tilde{T}$ . Then  $\mathcal{A} \subseteq \mathcal{A}$ . In particular,  $\mathcal{A}_\psi = \mathcal{A}$ , where  $\mathcal{A}_\psi$  is as in Theorem 5.8.*

*Proof.* Let  $T \in \mathcal{A}$ . Suppose  $\mathfrak{S}(T) = \varphi$  for some  $\varphi$  in  $L^\infty(\mathbb{T}^n)$ . Then  $T - T_\varphi \in \ker(\mathfrak{S})$  and hence

$$\mathcal{A} = \{T_\varphi + A : \varphi \in L^\infty(\mathbb{T}^n), A \in \ker(\mathfrak{S})\}.$$

Now if  $A \in \ker(\mathfrak{S})$ , then  $A^*A$  is so. Therefore,

$$\|Ak_\xi\|^2 = \langle Ak_\xi, Ak_\xi \rangle = \langle A^*Ak_\xi, k_\xi \rangle = \widetilde{(A^*A)}(\xi) \rightarrow 0 \text{ radially.}$$

In the similar fashion, it can be shown that  $\|A^*k_\xi\| \rightarrow 0$  radially. Thus  $\ker(\mathfrak{S}) \subseteq \mathcal{B}$ , and hence  $\mathcal{A} \subseteq \mathcal{A}$ .

Following Theorem 5.8, in particular, we have  $\mathcal{A}_\psi = \mathcal{A}$  for all  $\psi \in L^\infty(\mathbb{T}^n)$  with  $|\psi| = 1$  a.e. on  $\mathbb{T}^n$ . ■

We shall conclude this section with the following application of our *symbol map* result, which gives a partial answer in some sense to the question raised by Barriá and Halmos in [2]:

Which projections belong to the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}))$ ?

Using the *symbol map*, that is, Theorem 4.3, we shall show that if  $P$  is a projection in  $\mathcal{T}(L^\infty(\mathbb{T}))$  which is a finite rank perturbation of a Toeplitz operator, then the rank of  $P$  is finite or  $I_{H^2(\mathbb{D})} - P$  is finite rank, that is, co-finite. However, for  $n > 1$ , if a projection in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  is compact perturbation of a Toeplitz operator, then its rank is either finite or co-finite. More precisely, we have the following result:

**Theorem 5.11.** *Let  $P$  be an orthogonal projection in the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$ . For  $n = 1$ , if  $P = T_\varphi + F$ , where  $\varphi \in L^\infty(\mathbb{T})$  and  $F$  is of finite rank, then either  $P$  or  $I_{H^2(\mathbb{D})} - P$  is of finite rank. For  $n > 1$ , if  $P = T_\varphi + K$ , where  $\varphi \in L^\infty(\mathbb{T}^n)$  and  $K$  is compact, then either  $P$  or  $I_{H^2(\mathbb{D}^n)} - P$  is of finite rank.*

*Proof.* Suppose  $P$  is an orthogonal projection in the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  such that  $P$  is either finite or compact perturbation of a Toeplitz operator  $T_\varphi$  for  $n \geq 1$ . Then Theorem 4.3 yields  $\bar{\varphi} = \varphi$  and  $\varphi^2 = \varphi$  a.e. on  $\mathbb{T}^n$ . Hence  $\varphi = \chi_E$ , where  $E = \{\zeta \in \mathbb{T}^n : \varphi(\zeta) = 1\}$ , a measurable subset of  $\mathbb{T}^n$ .

For  $n = 1$ : If  $P = T_{\chi_E} + F$ , where  $\varphi \in L^\infty(\mathbb{T})$  and  $F$  is of finite rank, then  $P^2 = P$  gives

$$T_{\chi_E} - T_{\chi_E}^2 = F^2 - F + T_{\chi_E}F + FT_{\chi_E},$$

and hence  $T_{\chi_E}T_{\chi_E^c} = T_{\chi_E} - T_{\chi_E}^2$  is of finite rank. Therefore, either  $T_{\chi_E} = 0$  or  $T_{\chi_E^c} = 0$  by [7, Corollary 2.2]. Thus, either  $P$  or  $I_{H^2(\mathbb{D})} - P$  is of finite rank.

For  $n > 1$ : we shall get

$$T_{\chi_E} - T_{\chi_E}^2 = K^2 - K + T_{\chi_E}K + KT_{\chi_E},$$

where  $K$  is a compact operator. But  $T_{\chi_E}T_{\chi_E^c} = T_{\chi_E} - T_{\chi_E}^2 = H_{\chi_E}^* H_{\chi_E}$  which is compact, and hence  $H_{\chi_E}$  is compact. Now [6, Corollary 5], gives all compact Hankel operators are zero whenever  $n > 1$ . Thus  $T_{\chi_E}T_{\chi_E^c} = 0$  implies either  $T_{\chi_E} = 0$  or  $T_{\chi_E^c} = 0$  by Theorem 2.1 in [7]. Therefore,  $P = T_{\chi_E} + K$  gives either  $P$  or  $I_{H^2(\mathbb{D}^n)} - P$  is compact. Since both  $P$  and  $I_{H^2(\mathbb{D}^n)} - P$  are orthogonal projections and all compact orthogonal projections are of finite rank; therefore either  $P$  or  $I_{H^2(\mathbb{D}^n)} - P$  is of finite rank.

This completes the proof. ■

## 6. CONCLUDING REMARKS

The Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  provides an effective way to study the Toeplitz operators as elements of a  $C^*$ -algebra. More specifically, Theorem 4.3 shows that the study of the Toeplitz algebra  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  is largely reduces to that of the semi-commutator ideal  $\mathcal{S}$ . Let  $\theta$  be an inner function on  $\mathbb{D}^n$ , and  $\mathcal{K}_\theta = H^2(\mathbb{D}^n) \ominus \theta H^2(\mathbb{D}^n)$ . Suppose  $P_\theta$  denotes the orthogonal projection of  $H^2(\mathbb{D}^n)$  onto  $\mathcal{K}_\theta$ . Then

$$P_\theta = I_{H^2(\mathbb{D}^n)} - T_\theta T_{\bar{\theta}} = [T_\theta, T_{\bar{\theta}}] \in \mathcal{S}.$$

In fact, if  $\varphi = \theta_1 \bar{\eta}_1$  and  $\psi = \theta_2 \bar{\eta}_2$ , where  $\theta_i, \eta_i$  are inner functions, then

$$\begin{aligned} [T_\varphi, T_\psi] &= T_{\varphi\psi} - T_\varphi T_\psi \\ &= T_{\theta_1 \bar{\eta}_1 \theta_2 \bar{\eta}_2} - T_{\theta_1 \bar{\eta}_1} T_{\theta_2 \bar{\eta}_2} \\ &= T_{\bar{\eta}_1} T_{\bar{\eta}_2} T_{\theta_1} T_{\theta_2} - T_{\bar{\eta}_1} T_{\theta_1} T_{\bar{\eta}_2} T_{\theta_2} \\ &= T_{\bar{\eta}_1} T_{\bar{\eta}_2} T_{\theta_1} (I - T_{\eta_2} T_{\bar{\eta}_2}) T_{\theta_2} \\ &= T_{\bar{\eta}_1} T_{\bar{\eta}_2} T_{\theta_1} P_{\eta_2} T_{\theta_2}. \end{aligned}$$

In the case for  $n = 1$ , the semi-commutator ideal coincides with the commutator ideal in  $\mathcal{T}(L^\infty(\mathbb{T}))$  due to the availability of Douglas-Rudin theorem; again from the above calculation the semi-commutator ideal coincides with the closed ideal in  $\mathcal{T}(L^\infty(\mathbb{T}))$  generated by all orthogonal projections on the model space. Thus, Douglas-Rudin theorem (see [10]) plays an important role in this study. We now conclude this paper by raising some questions which naturally arise in our paper:

**Question 1.** Is it true that the class  $\{\bar{\varphi}\psi : \varphi, \psi \in H^\infty(\mathbb{T}^n)\}$  is norm-dense in  $L^\infty(\mathbb{T}^n)$  for  $n > 1$ ?

**Question 2.** Does the semi-commutator ideal coincide with the commutator ideal in  $\mathcal{T}(L^\infty(\mathbb{T}^n))$  for  $n > 1$ ?

It is noteworthy to point out that the affirmative answer to Question 1 gives an affirmative answer to Question 2, which we shall try to carry out in the future paper.

*Data availability:* Data sharing is not applicable to this article as no data sets were generated or analysed during the current study.

*Declarations*

*Conflict of interest:* The authors have no competing interests to declare.

## REFERENCES

- [1] S. Axler and D. Zheng, *The Berezin transform on the Toeplitz algebra*, *Studia Math.* **127** (1998), 113–136.
- [2] J. Barriá and P. R. Halmos, *Asymptotic Toeplitz operators*, *Trans. Amer. Math. Soc.* **273** (1982), no. 2, 621–630.
- [3] F. A. Berezin, *Covariant and contravariant symbols of operators*, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **36** (1972), 1134–1167.
- [4] F. A. Berezin, *Quantization of complex symmetric spaces*, (Russian) *Izv. Akad. Ser. Mat.* **39** 1975, no. 2, 363–402.
- [5] C. Berger and L. Coburn, *Toeplitz operators and quantum mechanics*, *J. Funct. Anal.* **68** (1986), 273–299.
- [6] M. Cotlar, C. Sadosky, *Abstract, weighted, and multidimensional Adamjan-Arov-Krein theorem, and the singular numbers of Sarason commutants*, *Integral Equations Operator Theory* **17** (1993), 169–201.
- [7] X. Ding, *Products of Toeplitz operators on the polydisk*, *Integral Equations Operator Theory* **45** (2003), 389–403.
- [8] A. Brown, P. Halmos, *Algebraic properties of Toeplitz operators*, *J. Reine Angew. Math.* **100** (1963), 89–102.
- [9] L. A. Coburn, *The  $C^*$ -algebra generated by an isometry*, *Bull. Amer. Math. Soc.* **73** (1967), 722–726.
- [10] R. G. Douglas, W. Rudin, *Approximation by inner functions*, *Pacific J. of Math.* **31** (1969), no. 2, 313–320.
- [11] R. G. Douglas, *Banach algebra techniques in operator theory*, Academic Press Vol. 179, 1972.
- [12] M. Engliš, *Toeplitz operators and the Berezin transform on  $H^2$* , *Linear Algebra Appl.* **232** (1995), 171–204.
- [13] M. Engliš, *Berezin quantization and reproducing kernels on complex domains*, *Trans. Amer. Math. Soc.* **348** 1996, no. 2, 411–479.
- [14] C. Kubrusly and J. Zanni, *A note on compactness of tensor products*, *Acta Math. Univ. Comenian. (N.S.)* **84** (2015), no. 1, 59–62.
- [15] A. Maji, J. Sarkar and S. Sarkar, *Toeplitz and asymptotic Toeplitz operators on  $H^2(\mathbb{D}^n)$* , *Bull. Sci. Math.* **146** (2018), 33–49.
- [16] W. Rudin, *Function theory in polydiscs*, W. A. Benjamin, Inc., New York, 1969.
- [17] K. Stroethoff, *Algebraic properties of Toeplitz operators on the Hardy space via the Berezin transform*, in: *Contemp. Math.*, **232**, Amer. Math. Soc., Providence, RI, 1999, pp. 313–319.
- [18] K. Zhu, *VMO, ESV, and Toeplitz operators on the Bergman space*, *Trans. Amer. Math. Soc.* **302** (1987), 617–646.
- [19] K. Zhu, *Analysis on Fock Spaces*, New York: Springer, 2012.

INDIAN INSTITUTE OF TECHNOLOGY ROORKEE, DEPARTMENT OF MATHEMATICS, ROORKEE-247 667,  
UTTARAKHAND, INDIA

*Email address:* javediitr07@gmail.com

INDIAN INSTITUTE OF TECHNOLOGY ROORKEE, DEPARTMENT OF MATHEMATICS, ROORKEE-247 667,  
UTTARAKHAND, INDIA

*Email address:* amit.maji@ma.iitr.ac.in, amit.iitm07@gmail.com, (†Corresponding author)