REVIEW MANUSCRIPT

Molecular techniques employed in CTG(Ser1) and CTG(Ala) D-xylose

metabolizing yeast clades for strain design and industrial applications

Ana Paula Wives^{1,2,†}, Isabelli Seiler de Medeiros Mendes^{1,2,†}, Sofia Turatti dos Santos^{1,2,†}, and Diego Bonatto^{1,2,*}

¹Laboratório de Biologia Molecular e Computacional, Centro de Biotecnologia da UFRGS, Departamento de Biologia Molecular e Biotecnologia, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

²Bioprocess and Biotechnology for Food Research Center (Biofood), Food Science and Technology Institute (ICTA), Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, Porto Alegre, RS, 91501-970, Brazil.

Short title: Molecular techniques for D-xylose fermenting yeasts.

*Corresponding author:

Diego Bonatto Centro de Biotecnologia da UFRGS - Laboratory 212 Departamento de Biologia Molecular e Biotecnologia Universidade Federal do Rio Grande do Sul – UFRGS Avenida Bento Gonçalves 9500 - Prédio 43421 Caixa Postal 15005 Porto Alegre – Rio Grande do Sul BRAZIL 91509-900 Phone: (+55 51) 3308-9410 E-mail: diego.bonatto@ufrgs.br Contract/grant sponsor: CNPq, FAPERGS ⁺ These authors contributed equally to the work.

Abstract

D-xylose is the second most abundant monosaccharide found in lignocellulose and is of biotechnological importance for producing second-generation ethanol and other high-value chemical compounds. The conversion of d-xylose to ethanol is promoted by microbial fermentation, mainly by bacteria, yeasts, or filamentous fungi. Among yeasts, species belonging to the CTG(Ser1) or CTG(Ala) clade display a remarkable ability to ferment D-xylose to ethanol and other compounds; however, these yeasts are not employed on an industrial scale owing to their poor fermentative performance compared to conventional yeasts, such as Saccharomyces cerevisiae, and also because of the lack of a molecular toolbox for the development of new strains tailored to fermentation stress tolerance and performance. Thus, the purpose of this review was to evaluate the major molecular tools (e.g., transformation markers and techniques, vectors, regulatory sequences, and gene editing techniques) available for the most studied yeasts of the CTG(Ser1) clade, such as Scheffersomyces, Spathaspora, Candida and Yamadazyma species, and the CTG(Ala) clade, representative *Pachysolen tannophilus*. Furthermore, we synthesized state-of-the-art molecular developments and perspectives to design D-xylose-fermenting yeast strains.

Keywords: D-xylose; molecular tools; CTG(Ser1) clade; CTG(Ala) clade; bioethanol.

Introduction

D-xylose is a major pentose found in lignocellulosic biomass and is composed of a xylan polysaccharide found in the hemicellulose fraction of lignocellulose [1,2]. It corresponds to 5–35% of the total carbohydrates found in lignocellulose biomass, depending on the plant source [1], and can be broadly classified into softwood (gymnosperms) and hardwood (angiosperms). In addition to hemicellulose, spent sulfite liquor (SSL), a byproduct of the cellulose acidic sulfite pulping process, contains 15–50% D-xylose and many other compounds, such as pentoses, hexoses, furfural and its derivatives, organic acids, and lignosulfonates [2,3]. The amount of D-xylose found in SSL depends on the wood type; softwood SSL contains a higher proportion of hexoses (e.g., glucose) than hardwood SSL, which is rich in D-xylose [3].

Considering its industrial importance, D-xylose has broad applications, ranging from biofuel production to value-added chemicals in food, health products, fuel additives, and drug manufacturing [4]. However, converting D-xylose to different chemicals by microbial fermentation is challenging, and only a few yeast species can naturally metabolize D-xylose in addition to genetically modified strains of the yeast *Saccharomyces cerevisiae* [5–7]. In some countries, softwood SSL has been used to produce second-generation bioethanol and additives for feedstock and civil construction [3]. However, untreated hardwood SSL is still not feasible for industrial applications because of the presence of high concentrations of inhibitory molecules for microbial cell growth and metabolism (e.g., acetic acid, furfural, and phenols) [5,8,9]. To solve this problem, different methods have been developed to reduce the formation of inhibitory molecules in hardwood SSL and other lignocellulose derivatives, including the addition of chemical compounds, such as inorganic hydroxides and reducing agents (e.g.,

sulfite), or the use of liquid-liquid and/or solid-liquid extraction procedures, thermal treatment, and biological agents (e.g., laccase and filamentous fungi) [8]. Among the methods employed, there is an increase in the fermentability of treated hardwood SSL and lignocellulose derivatives [8]. However, many of these methods require additional steps and/or treatment conditions that make their use economically unfeasible on a large industrial scale [3].

Therefore, an ideal yeast species that can tolerate inhibitory molecules present in SSL and lignocellulose derivatives with highly efficient D-xylose metabolism is desirable, and different strategies have been employed to obtain such yeast strains and species. An initial approach is to investigate yeast biodiversity in biomes, ecosystems, and ecological niches. In this approach, D-xylose-metabolizing yeast species have been isolated and characterized, such as Pachysolen tannophilus, which was initially present in Castanea vesca (sweet chestnut) and Acacia mearnsii (A. mollissima; black wattle) extracts used in leather tanning [10,11], and displayed high tolerance to the inhibitory molecules found in hardwood SSL [12]. Another characterized yeast genus isolated from the gut of the wood-boring beetle Odontotaenius disjunctus (Coleoptera: Passalidae) and Phrenapates bennetti (Coleoptera: Tenebrionidae) is Spathaspora sp., which can metabolize D-xylose under anaerobic conditions [13]. These yeast species are defined as "non-conventional yeasts" since they display genetic and phenotypic traits that differ from the "conventional yeasts," such as Saccharomyces cerevisiae [6], and constitute an unexplored source of genes and biochemical pathways with interest in industrial applications. On the other hand, the use of conventional yeasts for D-xylose fermentation has several advantages over non-conventional yeasts, especially in terms of metabolism, physiology, fermentative predictability, stress tolerance, and a large set of molecular tools available for genetic modifications and breeding selection of new strains [1,14,15]. In Saccharomyces cerevisiae, the enzymes of two major xylose assimilation pathways can be heterologously expressed: the oxidoreductase and isomerase pathways [14,16]. Non-conventional yeasts that naturally metabolize D-xylose employ the oxidoreductase pathway, which depends on the activity of two enzymes, xylose reductase (XR, EC 1.1.1.30) and xylitol dehydrogenase (XDH, EC 1.1.1.9), which convert D-xylose to xylulose via xylitol [14]. Bacteria (e.g., Escherichia coli) and chytrid fungi (e.g., Piromyces sp. and Orpinomyces sp.) mainly employ the xylose isomerase (XI, EC 5.3.1.5) pathway [14,17]. To date, the generation of recombinant S. cerevisiae strains able to fully convert D-xylose to ethanol by overexpression of XR-XDH or XI enzymes has only been partially achieved for a variety of reasons, including low enzyme activity, improper recombinant protein folding, metabolic imbalance (mainly due to the different requirements of XR-XDH enzymes for cofactors such as NAD(P)H/NAD(P)⁺), accumulation of xylitol as a by-product, restrained cell growth, and lack of proper D-xylose transporters, among others [14,15]. Considering the bottlenecks associated with the heterologous expression of oxidoreductase or isomerase pathways in S. cerevisiae, the use of non-conventional yeast species naturally adapted to D-xylose as a carbon source appears to be a logical pathway to explore in terms of industrial production of ethanol. Therefore, the use of different molecular tools to generate new non-conventional yeast strains is essential. This review focuses on the molecular tools available for the most studied non-conventional yeast species for the CTG(Ser1) or CTG(Ala) clade employed for second-generation bioethanol production from D-xylose, such as Scheffersomyces

stipitis, Spathaspora passalidarum, Candida intermedia, Yamadazyma tenuis, and Pachysolen tannophilus.

Scheffersomyces stipitis

The yeast genus Scheffersomyces includes some important species known to efficiently convert D-xylose to ethanol, including Scheffersomyces stipitis (initially classified Pichia stipitis), Scheffersomyces shehatae, Scheffersomyces as parashehatae, and Scheffersomyces illinoinensis [13,18]. These yeast species are found in symbiosis within the gut of the wood-ingesting beetles, Odontotaenius disjunctus (Coleoptera: Passalidae) and Verres sternbergianus (Coleoptera: Passalidae). Characteristically, the guts of wood-ingesting beetles are oxygen-limited and have high amounts of partially degraded wood, leading to the evolution of efficient biochemical pathways to process D-xylose and complex polysaccharides in Scheffersomyces stipitis as indicated by complete genome sequencing of this yeast in 2007 [18].

Scheffersomyces belongs to the fungal CTG(Ser1) clade, which comprises other biotechnologically relevant yeast species, such as *Spathaspora passalidarum*, and is characterized by a CTG codon that encodes serine instead of leucine [19]. The reassignment of the CTG codon in the fungal CTG(Ser1) clade imposes challenges in terms of coding sequence translation, particularly for the generation of functional transformation markers commonly used for non-conventional yeast selection, such as hygromycin B (*hygR*), nourseothricin (*natR*), geneticin (*kamR*), and phleomycin (*bleR*). The coding sequence for these transformation selection markers requires codon optimization to generate a functional protein in *Scheffersomyces* sp., which was performed by Matanović et al. for *S. stipitis* [20] (Table 1). In this sense, two antibiotic-resistance markers were optimized for *S. stipitis* selection, namely *kamR* and *bleR*, with *S. stipitis* partially resistant to phleomycin and geneticin [21,22] (Table 1). In addition, the *hygR* marker was codon-optimized and proven effective in *S. stipitis* and other D-xylose-fermenting yeasts [23]. Additionally, a series of auxotrophic *S. stipitis* strains were obtained for uracil (*ura3*), leucine (*leu2*), histidine (*his3-1*), and tryptophan (*trp5-10*) biosynthesis, allowing the selection of transformed cells with vectors harboring the respective prototrophic complementation markers (Table 1) in culture media lacking specific amino acids and nitrogen bases [22,24–26]. Unfortunately, the number of auxotrophic *S. stipitis* strains available is limited compared with that of conventional yeasts. However, it is expected that introducing gene editing or DNA recombination tools could increase the number of auxotrophic markers in *S. stipitis*. In this sense, adapting a *lox*P/Cre recombination system in *S. stipitis* allows for the recycling of antibiotic resistance markers [22], which can also be applied to generate new single or combined auxotrophic markers.

Regarding the development of vectors for *S. stipitis*, an early study described a vector containing a 2 μ m replicon from *S. cerevisiae* and a non-optimized *kamR* sequence (pUCKm8 vector; Table 2) that could replicate in *S. stipitis* [21]. However, the authors did not evaluate the vector stability. Matanović et al. and Yang et al. [20,26] described plasmids containing antibiotic resistance or prototrophic (*URA3*) markers with an autonomously replicating sequence (*ARS2*) isolated from *S. stipitis* (pRS54 and pJM2/6 vector series; Table 2). The authors showed that these vectors could be episomally maintained for at least 50 generations or stably integrated into the genome of *S. stipitis* when linearized [26] (Table 2). Another important achievement was the development of a series of stable centromeric vectors containing a centromeric

sequence derived from chromosome five (*CEN5*) in addition to an *ARS* (*ARS2*) [27]. *CEN5* increases the expression of heterologous sequences, including Cas9, which allows for better gene knockout efficiency [27,28]. Interestingly, *CEN5*, as well as other centromeric sequences from *S. stipitis*, appear to be part of the long terminal repeat (LTR) of Tps5 (a Ty5-type retrotransposon) that forms a cluster in the centromeres of *S. stipitis* [29]. Finally, integrative vectors using the *URA3* (YIPs1; Table 2) or rDNA 18S sequence from *Spathaspora passalidarum*, which contain different codon-adapted antibiotic-resistance markers (pRTH series; Table 2), were successfully employed for *S. stipitis* transformation with high stability, albeit with low efficiency [23,26].

Developing vectors for S. stipitis is essential for establishing a gene editing system using CRISPR-Cas9 [27,28]. However, as observed in almost all non-conventional yeasts [30], S. stipitis preferentially repairs DNA double-strand breaks induced by Cas9 by a non-homologous end-joining (NHEJ) DNA repair mechanism instead of homologous recombination (HR), generating mainly insertion-deletions (indels) in the DNA target molecule and hampering the use of gene editing techniques such as allele replacement [31]. A common way to circumvent the NHEJ pathway is to delete the coding sequences of the Ku complex, which is composed of Ku70 and Ku80 proteins responsible for directing the molecular components of the NHEJ pathway to the DSB lesion site [32]. The $ku70\Delta/ku80\Delta$ technique was employed for S. stipitis gene editing with a high HR rate and gene knockdown expression using dead Cas9 (dCas9) coupled with a repressor factor [31]. However, a significant drawback of the $ku70\Delta/ku80\Delta$ technique is diminished cell growth and telomere instability [30], which impair the industrial usability of $ku70\Delta/ku80\Delta$ strains for most non-conventional yeast species. Finally, the development of the so-called "bio-luminescent indicator nullified by Cas9-actuated recombination" (BLINCAR) allows the delivery and/or deletion of multiple genetic elements within *S. stipitis* genome [33] using replicative and/or integrative vectors combining the Ss*ARS2* replicon or long segments (\ge 2.0 kb) of amplified target genome sequences with different antibiotic resistance markers, like *hygR* (pWR vector series; Table 2).

Efficient transformation techniques are fundamental for the development of new strains of *Scheffersomyces* sp. and other D-xylose-fermenting yeasts. An early successful electroporation protocol for *S. stipitis* was designed by Ho et al. [21] using the pUCKm8 vector (Tables 2 and 3); however, transformation efficiency was not evaluated. Further, an electroporation efficiency of 6.0×10^2 to 8.6×10^3 colony-forming units (CFU) per ug of DNA was achieved using the pJM2/6 vector series (Tables 2 and 3) [26]. Cao et al. [31] improved electroporation efficiency by increasing the voltage to 2.5 kV, obtaining 1.0×10^3 CFU for 1 µg of a 10 kb plasmid. In addition to electroporation, *Scheffersomyces* employed a modified lithium acetate/polyethylene glycol (LiAc/PEG) transformation procedure, although the transformation efficiency was not indicated (Table 3) [33,34].

Considering the development of DNA expression devices (e.g., genes), the availability of promoters and terminators for *Scheffersomyces* sp. and other non-conventional yeast species is crucial for yeast strain design. Different numbers of constitutive and inducible promoters have been described in *S. stipitis* (Table 4). Strong constitutive promoters, such as *TEF1*p and *TDH3*p, and terminators from *XYL1* and *XYL2* were isolated from *S. stipitis* and employed to express the fumarate biosynthesis pathway in *Rhizopus oryzae* [35]. In addition, *TEF1* promoters from different species, such as *Blastobotrys adeninivorans* have been shown to function in *S. stipitis* and other

xylose-metabolizing yeasts, such as *Spathaspora passalidarum*, *Candida jeffriesii*, and *Candida amazonensis* [23]. In the same study, the authors showed that native promoter sequences from *S. stipitis XYL1.1*, *S. passalidarum ADH1*, and *S. passalidarum XYL1.1* could drive the expression of *EGFP* in their respective hosts, but not necessarily in different D-xylose-fermenting yeasts, pointing to genera- or species-specific transcriptional regulatory mechanisms [23]. Similarly, terminator sequences from D-xylose-fermenting yeasts are not necessarily interchangeable among different species; however, *CYC1*t from *S. passalidarum* promotes gene expression in *S. stipitis* [23]. Additionally, the derived promoter and terminator sequences from *TEF1* of *Eremothecium gossypii* have been proven to work on *S. stipitis* strains and other xylose-fermenting yeasts [20]. Finally, the inducible promoters *GAL1* and *XKS1* (Table 4) from *S. stipitis* could be alternatives for regulating the expression of specific genes in the presence of galactose and D-xylose, respectively [33].

Spathaspora passalidarum

The genus *Spathaspora* contains 22 described yeast species that use D-xylose from lignocellulosic residues to produce ethanol and/or xylitol with high efficiency under aerobic or anaerobic conditions, even in the presence of glucose and cellobiose [13,36]. The most characterized species of *Spathaspora* is *S. passalidarum*, isolated from the wood-boring beetle *O. disjunctus* (Coleoptera: Passalidae) [37], followed by *S. arborariae*, which was isolated from rotting wood in the Atlantic Rain Forest and Cerrado ecosystem in Brazil [38]. As they belong to the CTG(Ser1) clade [19], the molecular tools available for *Spathaspora* sp. are even more limited than those for *S. stipitis;* however, they are more or less interchangeable between both yeast genera, especially antibiotic-resistant transformation markers (Table 1) [23]. To date, no studies

have described the use of prototrophic and/or dominant markers for auxotrophic *S. passalidarum* strains (Table 1).

Different vectors optimized for *Spathaspora* are also scarce. An integrative vector containing the rDNA 18S sequence from *S. passalidarum* and codon-adapted *hygR* was used to express *EGFP* (pRTH vector series; Table 2) with a low transformation efficiency [23]. In addition, an epissomal vector containing a non-optimized codon *kamR* marker and a 2 µm replicon from *S. cerevisiae* (YEp352-km vector; Table 2) was shown to replicate in *S. passalidarum* [39].

Two lithium acetate transformation protocols were developed for *S. passalidarum* (Table 3). One protocol employed the LiAc/PEG/ssDNA method using one microgram of the linearized integrative vector series pRTH (Table 2), followed by incubation at 30 °C for 3 h in a rich (YPD) medium [23], and the second employed a LiAc/PEG/single-stranded DNA (ssDNA) optimized method for the YEp352-km vector (Table 2) [39]. Both protocols can be employed for *S. passalidarum* transformation, albeit with low efficiency (31–50 transformants/µg of pRTH vector for the first protocol and approximately 170 transformants/µg of YEp352-km vector under optimized conditions for the second protocol; Table 3) [39]. In addition, to date, no electroporation or trans-kingdom conjugation protocols have been described for *Spathaspora* species and/or strains (Table 3).

To our knowledge, no gene editing tools have been developed or adapted for *Spathaspora* species. Finally, promoters from *S. passalidarum ADH1, XYL1.1,* and *E. gossypii TEF1* can drive heterologous *EGFP* expression in this yeast (Table 4) [20,23]. Moreover, different terminator sequences, mainly derived from *S. passalidarum* genes (and the *S. cerevisiae CYC1* terminator), were proven to work on this yeast (Table 4).

Interestingly, the *S. passalidarum XYL1.1* terminator does not work on *S. stipitis* [23], indicating that not all DNA components are interchangeable between *S. passalidarum* and *S. stipitis*.

Candida intermedia and Yamadazyma tenuis

The genus *Candida* represents a large group that contains important yeast species, such as pathogenic *C. albicans, C. glabrata, C. parapsilosis, C. tropicalis,* and *C. krusei* [40]. Many *Candida* species can metabolize D-xylose, including *C. insectorum* and *C. jeffriesii* (both isolated from wood-ingesting insects), *C. ligosa* and *C. tropicalis* (isolated from rotting wood), and *C. famata* and *C. guilliermondii* (isolated from fruits) [41]. In addition to *Candida*, the *Yamadazyma* monophyletic clade was described as a sister group of *Candida* isolated from tree bark and is capable of metabolizing D-xylose [42,43]. Two species of *Candida* and *Yamadazyma* have been well studied for D-xylose fermentation to bioethanol, namely *C. intermedia* and *Y. tenuis* (formerly known as *Candida tenuis*) [44]. Similar to *Scheffesomyces* and *Spathaspora, C. intermedia* and *Y. tenuis* belong to the CTG(Ser1) clade [43,45]. Moreover, molecular tools widely available for *Candida* (mainly designed for *C. albicans*) can potentially be employed for *Yamadazyma* owing to their phylogenetic proximity to *Candida* clade.

C. intermedia has also been used for xylitol and single-cell proteins from D-xylose [46], has an efficient D-xylose transport system [47], and can metabolize glucose and D-xylose at high concentrations [48]. A few molecular tools have been developed for *C. intermedia*, including a split-marker deletion approach for gene editing based on vectors containing the *C. albicans* dominant marker nourseothricin acetyltransferase (*CaNAT1*) (Tables 1 and 2) flanked by 1 kb sequences with homology to the target sequence (template DNA) [45]. The authors employed electroporation for

vector transformation with low efficiency (Table 3). Using this approach, the authors improved the HR rates of target sequence editing to 70% [45]. However, we could not find any other reports indicating the use of precision gene-editing techniques for C. intermedia and Y. tenuis, such as Cas9. Despite this, the use of Cas9 technology for C. albicans and other Candida species is well established and can be adapted for C. intermedia and Y. tenuis [49]. Similarly, promoters, terminators, transformation markers, different transformation techniques, and replicative plasmids containing ARS or integrative plasmids that are available for C. albicans, C. tropicalis, C. glabatra, and C. maltosa could potentially be used for C. intermedia and Y. tenuis [50,51]. The bovine serum albumin (BSA)-responsive promoter (SAP2) of C. albicans has been shown to regulate the expression of a recombinant flippase (FLP) in C. intermedia in culture media containing BSA as a unique nitrogen source [45], reinforcing the idea that the molecular tools available for C. albicans can be directly employed for C. intermedia genetic manipulation. Finally, we did not find any molecular tools for Y. tenuis (Tables 1 to 4).

Pachysolen tannophilus

Pachysolen tannophilus is a D-xylose-fermenting model yeast belonging to the CTG(Ala) clade [52]. However, molecular tools for the genetic analysis and/or modification of *P. tannophilus* have been less developed than other D-xylose-fermenting yeasts, such as *S. stipitis*. Considering transformation markers, *P. tannophilus* auxotrophs for different amino acid requirements have been generated by UV mutagenesis [53–55], where *leu*⁻ strains are conventionally employed for transformation with plasmids containing the *LEU2* gene from *S. cerevisiae* (Table 1) because its coding sequence contains only the CTG codon and does not interfere with the activity of the

Leu2 protein [56]. However, similar to yeasts from the CTG(Ser1) clade, antibiotic-resistance transformation marker coding sequences should have their codons optimized for *P. tannophilus* (e.g., changing the CTG codon to TTG codon) because the codon CTG codifies to Ala instead of Leu in this yeast [56]. An early attempt to express the kamR transformation marker in P. tannophilus was unsuccessful, probably because of the non-optimized kamR coding sequence [57]. Furthermore, it was shown that even codon-optimized kamR and aureobasidin A resistance genes (AUR1-C) were unable to confer resistance to both geneticin and aureobasidin A in P. tannophilus under the control of native S. cerevisiae LEU2 and AUR1 promoters (Table 1) [56], pointing that S. cerevisiae promoters negatively affect the expression of these markers. Reinforcing this idea, a codon-optimized hygromycin B-resistance (hygR) coding sequence controlled by P. tannophilus native glyceraldehyde-3-phosphate dehydrogenase promoter and terminator has been shown to work in P. tannophilus [58]. Interestingly, S. cerevisiae hexokinase isoenzyme 2 (HXK2) was employed as a dominant transformation marker in hexose-metabolizing negative mutants of *P. tannophilus* (Table 1) [59].

Very few vectors are available for *P. tannophilus* (Table 2). Epissomal vectors containing the *S. cerevisiae* 2 µm replicon combined with the prototrophic *S. cerevisiae LEU2* marker (YEp13 or pACT2-Cas9c vectors; Table 2) or replicative vectors containing the *ARS* from *S. cerevisiae* combined with the dominant marker *HXK2* (YRp vector; Table 2) have been used for *P. tannophilus* transformation [51,56], but no data have been reported on plasmid stability in this yeast, and no integrative vectors have been reported to date for *P. tannophilus*. Additionally, the use of a centromeric sequence from *S. cerevisiae* (*CEN4*) and *ARS1* from the same yeast proved ineffective for plasmid replication and stability in *P. tannophilus* (Table 2), indicating that the design of

centromeric vectors depends on the characterization of centromeres and ARS in P. tannophilus [56].

Different transformation methods based on the use of lithium acetate (LiAc) alone [59] or combined with ssDNA and PEG have been employed for *P. tannophilus* (Table 3) [56]. The efficiency of LiAc transformation was reported to be 40-50 transformant/µg vector [59], whereas the combined LiAc-ssDNA-PEG technique achieved an efficiency of 3×10^4 transformants/µg DNA (Table 3) [56]. Interestingly, trans-kingdom conjugation between *P. tannophilus* and *Escherichia coli* has been applied with relative success to promote the transfer of vectors [53,56], with 2.99 × 10⁻² transconjugants per recipient at a donor-to-recipient ratio of 1000:1 (Table 3) [56]. To date, no electroporation technique has been reported in *P. tannophilus*.

Although this yeast has been known since 1957 [11], little has been done to prospect and/or characterize regulatory sequences (e.g., promoters and terminators) for heterologous gene expression or gene reengineering, except for the promoter and terminator of *P. tannophilus GAPDH* gene (Table 4), which has been proven to be efficient in driving the expression of *hygR* [58]. Moreover, little effort has been made to develop transformation markers, vectors, and gene-editing techniques for this yeast species. Considering that *P. tannophilus* belongs to a few groups of yeasts capable of tolerating the toxic components of hardwood SSL [12] and can convert glycerol to ethanol [60], the development of molecular tools could improve the biotechnological importance of this yeast for biofuel production.

Conclusion

The D-xylose-fermenting yeasts of the CTG(Ser1) and CTG(Ala) clades are an important source of genotypes and phenotypes useful for converting lignocellulosic

biomass to biofuels and other industrially important chemical compounds. However, many of these yeast species are still unsuitable for industrial applications because of the lack of knowledge about their physiology and molecular tools that enable researchers to develop new strains tailored for industrial applications compared to conventional yeasts (e.g., S. cerevisiae). A major barrier to the development of molecular tools for these yeasts is the reassignment of the CTG codon for Ser or Ala, which impairs the use of existing vectors, transformation markers, regulatory sequences, and precise gene editing technologies available for conventional yeasts. Moreover, highly efficient transformation techniques and characterization of centromeres and autonomously replicative sequences for yeasts of the CTG(Ser1) and CTG(Ala) clades for the development of stable vectors still lag behind those observed for conventional yeasts. Considering that synthetic biology technologies, such as gene synthesis, cloning, and genome annotation tools, are advancing rapidly, the development of new molecular tools is expected to increase, allowing the design of new strains adapted for industrial applications.

Funding

This work was supported by "CNPq/MCTI/CT-BIOTEC N° 30/2022 - Linha 2: Novas tecnologias em Biotecnologia" from Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq [grant number 440226/2022-8], by "Programa Pesquisador Gaúcho - PqG" from Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul - FAPERGS [Edital FAPERGS 07/2021 and grant number 21/2551-0001958-1], by "INOVA CLUSTERS TECNOLÓGICOS" from Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul - Estado do Rio Grande do Sul - FAPERGS [Edital FAPERGS 07/2021 and grant number 21/2551-0001958-1], by

22/2551-0000834-8] and by "Programa de Redes Inovadoras de Tecnologias Estratégicas do Rio Grande Do Sul – RITEs-RS" from Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul - FAPERGS [Edital FAPERGS 06/2021 and grant number 22/2551-0000397-4]. The sponsors had no role in the study design, collection, analysis, interpretation of data, writing of the report, or decision to submit the article for publication.

Authors contributions

Ana Paula Wives: Writing - Review & Editing, Visualization. Isabelli Seiler de Medeiros Mendes: Writing - Review & Editing, Visualization. Sofia Turatti dos Santos: Writing - Review & Editing, Visualization. Diego Bonatto: Conceptualization, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration, Funding acquisition.

Declaration of interest

Declarations of interest: none.

References

- [1] A. Ochoa-Chacón, A. Martinez, H.M. Poggi-Varaldo, L. Villa-Tanaca, A.C. Ramos-Valdivia, T. Ponce-Noyola, Xylose metabolism in bioethanol production: *Saccharomyces cerevisiae* vs non-*Saccharomyces* yeasts, BioEnergy Res. 15 (2022) 905–923. https://doi.org/10.1007/s12155-021-10340-x.
- [2] S.S. Helle, T. Lin, S.J.B. Duff, Optimization of spent sulfite liquor fermentation, Enzyme Microb. Technol. 42 (2008) 259–264.

https://doi.org/10.1016/j.enzmictec.2007.10.004.

- [3] S.R. Pereira, D.J. Portugal-Nunes, D.V. Evtuguin, L.S. Serafim, A.M.R.B. Xavier, Advances in ethanol production from hardwood spent sulphite liquors, Process Biochem. 48 (2013) 272–282. https://doi.org/10.1016/j.procbio.2012.12.004.
- [4] V. Narisetty, R. Cox, R. Bommareddy, D. Agrawal, E. Ahmad, K. Kumar Pant, A. Kumar Chandel, S. Kant Bhatia, D. Kumar, P. Binod, V. Kumar Gupta, V. Kumar, Valorisation of xylose to renewable fuels and chemicals, an essential step in augmenting the commercial viability of lignocellulosic biorefineries, Sustain. Energy Fuels 6 (2022) 29–65. https://doi.org/10.1039/D1SE00927C.
- [5] N. Bezuidenhout, D.C.S. Rorke, E. Van Rensburg, D. Diedericks, J.F. Görgens, Ethanol production from non-detoxified hardwood spent sulfite liquor in submerged fed-batch culture using advanced yeasts, Biomass Convers. Biorefinery (2023). https://doi.org/10.1007/s13399-023-03992-6.
- [6] J.C. Bergmann, D. Trichez, W.G. de Morais Junior, T.G.S. Ramos, T.F. Pacheco, C.V.G.C. Carneiro, V.M. Honorato, L.A. Serra, J.R.M. Almeida, Biotechnological Application of non-conventional yeasts for xylose valorization, in: A. Sibirny (Ed.), Non-Conv. Yeasts Basic Res. Appl., Springer International Publishing, Cham, 2019: pp. 23–74. https://doi.org/10.1007/978-3-030-21110-3_2.
- [7] J.C. Du Preez, B.A. Prior, A quantitative screening of some xylose-fermenting yeast isolates, Biotechnol. Lett. 7 (1985) 241–246. https://doi.org/10.1007/BF01042370.
- [8] L.J. Jönsson, B. Alriksson, N.-O. Nilvebrant, Bioconversion of lignocellulose: inhibitors and detoxification, Biotechnol. Biofuels 6 (2013) 16. https://doi.org/10.1186/1754-6834-6-16.
- [9] M.J. Taherzadeh, K. Karimi, Fermentation inhibitors in ethanol processes and

different strategies to reduce their effects, in: Biofuels, Elsevier, 2011: pp. 287–311. https://doi.org/10.1016/B978-0-12-385099-7.00012-7.

- [10] P.J. Slininger, P.L. Bolen, C.P. Kurtzman, *Pachysolen tannophilus:* Properties and process considerations for ethanol production from d-xylose, Enzyme Microb. Technol. 9 (1987) 5–15. https://doi.org/10.1016/0141-0229(87)90043-3.
- [11] J. Boidin, J. Adzet, Deux curieuses levuses isolees d'extraits tannants d'origine végétale: *Pachysolen (nov. gen.) tannophilus nov. sp.* et *P. pelliculatus nov. sp*, Bull. soc. mycol. France 73 (1957) 331–342.
- [12] E.M. Lohmeier-Vogel, C.R. Sopher, H. Lee, Intracellular acidification as a mechanism for the inhibition by acid hydrolysis-derived inhibitors of xylose fermentation by yeasts, J. Ind. Microbiol. Biotechnol. 20 (1998) 75–81. https://doi.org/10.1038/sj.jim.2900484.
- [13] R.M. Cadete, C.A. Rosa, The yeasts of the genus Spathaspora: potential candidates for second-generation biofuel production, Yeast 35 (2018) 191–199. https://doi.org/10.1002/yea.3279.
- [14] S. Kwak, Y.-S. Jin, Production of fuels and chemicals from xylose by engineered Saccharomyces cerevisiae: a review and perspective, Microb. Cell Factories 16 (2017) 82. https://doi.org/10.1186/s12934-017-0694-9.
- [15] B. de O. Vargas, J.R. Dos Santos, G.A.G. Pereira, F. da S.B. de Mello, An atlas of rational genetic engineering strategies for improved xylose metabolism in *Saccharomyces cerevisiae*, PeerJ 11 (2023) e16340. https://doi.org/10.7717/peerj.16340.
- [16] S.H. Bhosale, M.B. Rao, V.V. Deshpande, Molecular and industrial aspects of glucose isomerase., Microbiol. Rev. 60 (1996) 280–300.

- [17] H.R. Harhangi, A.S. Akhmanova, R. Emmens, C. van der Drift, W.T.A.M. de Laat, J.P. van Dijken, M.S.M. Jetten, J.T. Pronk, H.J.M. Op den Camp, Xylose metabolism in the anaerobic fungus *Piromyces* sp. strain E2 follows the bacterial pathway, Arch. Microbiol. 180 (2003) 134–141. https://doi.org/10.1007/s00203-003-0565-0.
- [18] T.W. Jeffries, I.V. Grigoriev, J. Grimwood, J.M. Laplaza, A. Aerts, A. Salamov, J. Schmutz, E. Lindquist, P. Dehal, H. Shapiro, Y.-S. Jin, V. Passoth, P.M. Richardson, Genome sequence of the lignocellulose-bioconverting and xylose-fermenting yeast Pichia stipitis, Nat. Biotechnol. 25 (2007) 319–326. https://doi.org/10.1038/nbt1290.
- [19] N. Papon, V. Courdavault, M. Clastre, Biotechnological potential of the fungal CTG clade species in the synthetic biology era, Trends Biotechnol. 32 (2014) 167–168. https://doi.org/10.1016/j.tibtech.2013.10.006.
- [20] A. Matanović, K. Arambašić, B. Žunar, A. Štafa, M. Svetec Miklenić, B. Šantek, I.-K. Svetec, Toolbox for genetic transformation of non-conventional *Saccharomycotina* yeasts: High efficiency transformation of yeasts belonging to the *Schwanniomyces* genus, J. Fungi 8 (2022) 531. https://doi.org/10.3390/jof8050531.
- [21] N.W.Y. Ho, D. Petros, X.X. Deng, Genetic transformation of xylose-fermenting yeast *Pichia stipitis*, Appl. Biochem. Biotechnol. 28 (1991) 369–375. https://doi.org/10.1007/BF02922616.
- [22] J.M. Laplaza, B.R. Torres, Y.-S. Jin, T.W. Jeffries, *Sh ble* and Cre adapted for functional genomics and metabolic engineering of *Pichia stipitis*, Enzyme Microb. Technol. 38 (2006) 741–747. https://doi.org/10.1016/j.enzmictec.2005.07.024.
- [23] H. Li, H. Fan, Y. Li, G.Y. Shi, Z.Y. Ding, Z.H. Gu, L. Zhang, Construction and application of multi-host integrative vector system for xylose-fermenting yeast, FEMS Yeast Res. 17 (2017) fox055. https://doi.org/10.1093/femsyr/fox055.

- [24] P. Lu, B.P. Davis, J. Hendrick, T.W. Jeffries, Cloning and disruption of the b-isopropylmalate dehydrogenase gene (*LEU2*) of *Pichia stipitis* with *URA3* and recovery of the double auxotroph, Appl. Microbiol. Biotechnol. 49 (1998) 141–146. https://doi.org/10.1007/s002530051150.
- [25] M. Piontek, J. Hagedorn, C.P. Hollenberg, G. Gellissen, A.W.M. Strasser, Two novel gene expression systems based on the yeasts *Schwanniomyces occidentalis* and *Pichia stipitis*, Appl. Microbiol. Biotechnol. 50 (1998) 331–338. https://doi.org/10.1007/s002530051300.
- [26] V.W. Yang, J.A. Marks, B.P. Davis, T.W. Jeffries, High-efficiency transformation of *Pichia stipitis* based on its *URA3* gene and a homologous autonomous replication sequence, *ARS2*, Appl. Environ. Microbiol. 60 (1994) 4245–4254. https://doi.org/10.1128/aem.60.12.4245-4254.1994.
- [27] M. Cao, A.S. Seetharam, A.J. Severin, Z. Shao, Rapid isolation of centromeres from *Scheffersomyces stipitis*, ACS Synth. Biol. 6 (2017) 2028–2034. https://doi.org/10.1021/acssynbio.7b00166.
- [28] M. Cao, M. Gao, C.L. Lopez-Garcia, Y. Wu, A.S. Seetharam, A.J. Severin, Z. Shao, Centromeric DNA facilitates non-conventional yeast genetic engineering, ACS Synth. Biol. 6 (2017) 1545–1553. https://doi.org/10.1021/acssynbio.7b00046.
- [29] A.Y. Coughlan, K.H. Wolfe, The reported point centromeres of Scheffersomyces stipitis are retrotransposon long terminal repeats, Yeast 36 (2019) 275–283. https://doi.org/10.1002/yea.3375.
- [30] M. Lu, S. Billerbeck, Improving homology-directed repair by small molecule agents for genetic engineering in unconventional yeast?—Learning from the engineering of mammalian systems, Microb. Biotechnol. 17 (2024) e14398.

https://doi.org/10.1111/1751-7915.14398.

- [31] M. Cao, M. Gao, D. Ploessl, C. Song, Z. Shao, CRISPR–mediated genome editing and gene repression in *Scheffersomyces stipitis*, Biotechnol. J. 13 (2018) 1700598. https://doi.org/10.1002/biot.201700598.
- [32] S. Zahid, M. Seif El Dahan, F. lehl, P. Fernandez-Varela, M.-H. Le Du, V. Ropars, J.B. Charbonnier, The multifaceted roles of Ku70/80, Int. J. Mol. Sci. 22 (2021) 4134. https://doi.org/10.3390/ijms22084134.
- [33] W.D. Reichard, S.E. Smith, J.B. Robertson, BLINCAR: a reusable bioluminescent and Cas9-based genetic toolset for repeatedly modifying wild-type *Scheffersomyces stipitis*, mSphere 8 (2023) e00224-23. https://doi.org/10.1128/msphere.00224-23.
- [34] K.O. Barros, M. Mader, D.J. Krause, J. Pangilinan, B. Andreopoulos, A. Lipzen, S.J. Mondo, I.V. Grigoriev, C.A. Rosa, T.K. Sato, C.T. Hittinger, Oxygenation influences xylose fermentation and gene expression in the yeast genera *Spathaspora* and *Scheffersomyces*, Biotechnol. Biofuels Bioprod. 17 (2024) 20. https://doi.org/10.1186/s13068-024-02467-8.
- [35] L. Wei, J. Liu, H. Qi, J. Wen, Engineering Scheffersomyces stipitis for fumaric acid production from xylose, Bioresour. Technol. 187 (2015) 246–254. https://doi.org/10.1016/j.biortech.2015.03.122.
- [36] T.M. Long, Y.-K. Su, J. Headman, A. Higbee, L.B. Willis, T.W. Jeffries, Cofermentation of glucose, xylose, and cellobiose by the beetle-associated yeast *Spathaspora passalidarum*, Appl. Environ. Microbiol. 78 (2012) 5492–5500. https://doi.org/10.1128/AEM.00374-12.
- [37] N.H. Nguyen, S.-O. Suh, C.J. Marshall, M. Blackwell, Morphological and ecological similarities: wood-boring beetles associated with novel xylose-fermenting

yeasts, *Spathaspora passalidarum gen. sp. nov.* and *Candida jeffriesii sp. nov.*, Mycol. Res. 110 (2006) 1232–1241. https://doi.org/10.1016/j.mycres.2006.07.002.

- [38] R.M. Cadete, R.O. Santos, M.A. Melo, A. Mouro, D.L. Gonçalves, B.U. Stambuk, F.C.O. Gomes, M.-A. Lachance, C.A. Rosa, *Spathaspora arborariae* sp. nov., a d-xylose-fermenting yeast species isolated from rotting wood in Brazil, FEMS Yeast Res. 9 (2009) 1338–1342. https://doi.org/10.1111/j.1567-1364.2009.00582.x.
- [39] Y. Chen, W. Huang, G. Fu, J. Guo, M. Liu, X. Guo, D.-G. Xiao, A genetic transformation protocol for the xylose-fermenting yeast *Spathaspora passalidarum*, Eng. Life Sci. 15 (2015) 550–555. https://doi.org/10.1002/elsc.201400232.
- S. Giri, A. Kindo, A review of *Candida* species causing blood stream infection,
 Indian J. Med. Microbiol. 30 (2012) 270–278.
 https://doi.org/10.4103/0255-0857.99484.
- [41] R. Kaewwichian, P. Khunnamwong, S. Am-In, S. Jindamorakot, M. Groenewald, S. Limtong, *Candida xylosifermentans* sp. nov., a d-xylose-fermenting yeast species isolated in Thailand, Int. J. Syst. Evol. Microbiol. 69 (2019) 2674–2680. https://doi.org/10.1099/ijsem.0.003505.
- [42] M.A.B. Haase, J. Kominek, Q.K. Langdon, C.P. Kurtzman, C.T. Hittinger, Genome sequence and physiological analysis of *Yamadazyma laniorum* f.a. sp. nov. and a reevaluation of the apocryphal xylose fermentation of its sister species, Candida tenuis, FEMS Yeast Res. 17 (2017) fox019. https://doi.org/10.1093/femsyr/fox019.
- [43] D. Libkind, N. Čadež, D.A. Opulente, Q.K. Langdon, C.A. Rosa, J.P. Sampaio, P. Gonçalves, C.T. Hittinger, M.A. Lachance, Towards yeast taxogenomics: lessons from novel species descriptions based on complete genome sequences, FEMS Yeast Res. 20 (2020) foaa042. https://doi.org/10.1093/femsyr/foaa042.

- [44] J. Ruchala, A.A. Sibirny, Pentose metabolism and conversion to biofuels and high-value chemicals in yeasts, FEMS Microbiol. Rev. 45 (2021) fuaa069. https://doi.org/10.1093/femsre/fuaa069.
- [45] K.V.R. Peri, F. Faria-Oliveira, A. Larsson, A. Plovie, N. Papon, C. Geijer, Split-marker-mediated genome editing improves homologous recombination frequency in the CTG clade yeast *Candida intermedia*, FEMS Yeast Res. 23 (2023) foad016. https://doi.org/10.1093/femsyr/foad016.
- [46] J. Wu, J. Hu, S. Zhao, M. He, G. Hu, X. Ge, N. Peng, Single-cell protein and xylitol production by a novel yeast strain *Candida intermedia* FL023 from lignocellulosic hydrolysates and xylose, Appl. Biochem. Biotechnol. 185 (2018) 163–178. https://doi.org/10.1007/s12010-017-2644-8.
- [47] M.J. Leandro, P. Gonçalves, I. Spencer-Martins, Two glucose/xylose transporter genes from the yeast *Candida intermedia:* first molecular characterization of a yeast xylose–H⁺ symporter, Biochem. J. 395 (2006) 543–549. https://doi.org/10.1042/BJ20051465.
- [48] M. Saito, H. Nagasaki, S. Watanabe, T. Fujimoto, Ethanol production from sugars in hydrolysates of cellulosic biomass resources with xylose-fermenting yeast *Candida intermedia* 4-6-4T2, J. Jpn. Pet. Inst. 60 (2017) 127–136. https://doi.org/10.1627/jpi.60.127.
- [49] L. Lombardi, J. Oliveira-Pacheco, G. Butler, Plasmid-based CRISPR-Cas9 gene editing in multiple *Candida* species, mSphere 4 (2019) 10.1128/msphere.00125-19. https://doi.org/10.1128/msphere.00125-19.
- [50] J. Berman, P.E. Sudbery, *Candida albicans:* A molecular revolution built on lessons from budding yeast, Nat. Rev. Genet. 3 (2002) 918–931.

https://doi.org/10.1038/nrg948.

- [51] T.-T. Wang, Y.-J. Choi, B.H. Lee, Transformation systems of non- Saccharomyces Yeasts, Crit. Rev. Biotechnol. 21 (2001) 177–218. https://doi.org/10.1080/20013891081719.
- [52] S. Mühlhausen, P. Findeisen, U. Plessmann, H. Urlaub, M. Kollmar, A novel nuclear genetic code alteration in yeasts and the evolution of codon reassignment in eukaryotes, Genome Res. 26 (2016) 945–955. https://doi.org/10.1101/gr.200931.115.
- [53] G.T. Hayman, P.L. Bolen, Movement of shuttle plasmids from *Escherichia coli* into yeasts other than *Saccharomyces cerevisiae* using trans-kingdom conjugation, Plasmid 30 (1993) 251–257. https://doi.org/10.1006/plas.1993.1056.
- [54] J.-K. Ahn, W.-G. An, J.-D. Lee, H.-K. Jun, Comparative studies on xylose utilization and fermentation by *Pachysolen tannophilus* fusants and auxotrophs, Korean J. Mycol. 19 (1991) 276–281.
- [55] N.K. Harner, P.K. Bajwa, P.A. Formusa, G.D. Austin, M.B. Habash, J.T. Trevors, C.-K. Chan, C.-Y. Ho, H. Lee, Determinants of tolerance to inhibitors in hardwood spent sulfite liquor in genome shuffled *Pachysolen tannophilus* strains, Antonie Van Leeuwenhoek 108 (2015) 811–834. https://doi.org/10.1007/s10482-015-0537-9.
- [56] G.-Y. Mei, P.K. Bajwa, M. Dashtban, C.-Y. Ho, H. Lee, Transfer of plasmid into the pentose-fermenting yeast *Pachysolen tannophilus*, J. Microbiol. Methods 148 (2018) 97–103. https://doi.org/10.1016/j.mimet.2018.03.013.
- [57] X. Liu, Conversion of the biodiesel by-product glycerol by the non-conventional yeast *Pachysolen tannophilus*, phd, Technical University of Denmark, 2012.
- [58] R. Riley, S. Haridas, K.H. Wolfe, M.R. Lopes, C.T. Hittinger, M. Göker, A.A.

Salamov, J.H. Wisecaver, T.M. Long, C.H. Calvey, A.L. Aerts, K.W. Barry, C. Choi, A. Clum, A.Y. Coughlan, S. Deshpande, A.P. Douglass, S.J. Hanson, H.-P. Klenk, K.M. LaButti, A. Lapidus, E.A. Lindquist, A.M. Lipzen, J.P. Meier-Kolthoff, R.A. Ohm, R.P. Otillar, J.L. Pangilinan, Y. Peng, A. Rokas, C.A. Rosa, C. Scheuner, A.A. Sibirny, J.C. Slot, J.B. Stielow, H. Sun, C.P. Kurtzman, M. Blackwell, I.V. Grigoriev, T.W. Jeffries, Comparative genomics of biotechnologically important yeasts, Proc. Natl. Acad. Sci. 113 (2016) 9882–9887. https://doi.org/10.1073/pnas.1603941113.

- [59] D.N. Wedlock, R.J. Thornton, Transformation of a glucose negative mutant of *Pachysolen tannophilus* with a plasmid carrying the cloned hexokinase PII gene from *Saccharomyces cerevisiae*, Biotechnol. Lett. 11 (1989) 601–604. https://doi.org/10.1007/BF01025265.
- [60] X. Liu, P.R. Jensen, M. Workman, Bioconversion of crude glycerol feedstocks into ethanol by *Pachysolen tannophilus*, Bioresour. Technol. 104 (2012) 579–586. https://doi.org/10.1016/j.biortech.2011.10.065.

Tables

Table 1. Transformation markers employed in CTG(Ser1) and CTG(Ala) clade yeasts.

Yeast species	Transformation markers		
	Prototrophic markers	Antibiotic-resistance markers	Other marker (dominant markers)
Scheffersomyces stipitis	URA3; LEU2; HIS3-1; TRP5-10 [22,24–26]	kamR ^{1,2} ; bleR ^{1,2} ; natR ¹ ; hygR ¹ [20–23]	Not reported
Spathaspora passalidarum	Not reported	kamR ^{1,2} ; bleR ^{1,2} ; natR ¹ ; hygR ¹ [23]	Not reported
Candida intermedia	Not reported	Candida albicans natR [45]	Not reported
Yamadazyma tenuis	Not reported	Not reported	Not reported
Pachysolen tannophilus	SceLEU2 [56]	 Non-optimized codon sequence of kamR do not confer resistance to geneticin [57] Codon-optimized <i>kamR</i> and <i>SceAUR1-C</i> were observed not to confer resistance to geneticin and aureobasidin A, respectively [56]. Codon-optimized <i>hygR</i> confers resistance to hygromycin B [58] 	SceHXK2 [59]

¹Requires codon optimization.

²Scheffersomyces stipitis is partially resistant to phleomycin and geneticin (up to 2,000

µg/mL for geneticin) [23,33]

Abbreviation: Saccharomyces cerevisiae (Sce)

Table 2. Different vector types employed for the transformation of D-xylose fermenting

yeasts. The table indicates the replicon/integration type (boldface) and transformation

markers. In addition, the names of vectors are shown inside parentheses.

Yeast species	Vector type			
	Epissomal	Replicative	Centromeric	Integrative
Scheffersomyces stipitis	1. S. cerevisiae 2 μ replicon; kamR (pUCKm8 vector) [21]	 SsARS2; antibiotic resistance markers¹ (pRS54 series and pWR series) [20,33] SsARS2; SsURA3 (pJM2/6 series) [26] 	1. SsCEN5; SsARS2; SsURA3 [28]	 SsURA3² (YIPs1) [26] Spa18S³ rDNA; hygR (pRTH series) [23] SsURA3⁴, SsEGC3⁴, SsGSC2⁴, and SsCRF1⁴; antibiotic resistance markers¹ (pWR series) [33]
Spathaspora passalidarum	 S. cerevisiae 2 μ replicon; kamR (YEp352-km vector) [39] 	Not reported	Not reported	1. Spa18S ³ rDNA; <i>hygR</i> (pRTH series) [23]
Candida intermedia	Not reported	Not reported	Not reported	1. CiADE2 ⁵ ; CiXYL1_2C ⁵ ; CiLAC9 ⁵ ; CiLADA ⁵ ; C. albicans natR ⁵ [45]
Yamadazyma tenuis	Not reported	Not reported	Not reported	Not reported
Pachysolen tannophilus	 S. cerevisiae 2 μ replicon; SceLEU2 (YEp13) [53] S. cerevisiae 2 μ replicon; 	1. S. cerevisiae ARS; Sce <i>HXK2</i> (YRp) [59]	1. S. cerevisiae CEN4-ARS1 unable to replicate in P. tannophilus (pAUR123 and derivatives)	Not reported

Yeast species	Vector type			
	<i>SceLEU2</i> (pACT2-CA S9c) [56]		[56]	

¹Transformation markers: *kamR*; *bleR*; *natR*; *hygR*.

²The ~ 4.1 kb *SsURA3* sequence was cut with different restriction endonucleases and employed for *S. stipitis* TJ26 (*ura3-1*) electroporation. Southern blot data showed that *SsURA3* genome integration was random due to non-homologous end-joining recombination (NHEJ).

³The ~1.55 kb *Spa*18S rDNA sequence was cut with *Stul*, and the linearized vector was employed for yeast transformation.

⁴A 2.7 kb genomic segment containing the *SsURA3* gene was employed for vector integration (pWR58 vector) after cutting with *Eco*RV. The 2 kb genomic sequences containing *SsEGC34, SsGSC24,* and *SsCRF14* were amplified by PCR, cloned into the pWR88 vector, and linearized with *Eco*RV.

⁵Two vectors were generated in this study. Vector p*CaNAT1* and its derivative with flippase (FLP) recombinase (p*FLP_CaNAT1*). Both vectors contained the *Candida albicans natR* transformation marker under the control of the *EgTEF1* promoter and terminator, while *CaSAP2* promoter induces FLP recombinase. A homology sequence of ~1.0 kb on both sides of the indicated target genes was used for genome integration. Abbreviations: *Blastobotrys adeninivorans* (*Ba*); *Eremothecium gossypii (Eg); Pachysolen tannophilus (Pt*); *Saccharomyces cerevisiae (Sc); Scheffersomyces stipitis (Ss), Spathaspora passalidarum (Sp)*.

Table 3. Transformation techniques described for D-xylose fermenting yeasts. When

available, the transformation efficiency is indicated inside parentheses.

Yeast species	Transformation technique		
	Chemical	Electroporation	Trans-kingdom conjugation
Scheffersomyces stipitis	Yes (LiAc/PEG/ssDNA) - no reported efficiency [33,34]	Yes (6.0 × 10 ² to 8.6 × 10 ³ CFU per ug of DNA) [21,26,31]	Not reported
Spathaspora passalidarum	Yes 1. LiAc/PEG/ssD NA: 3.1 to 5.0 × 10 ¹ CFU per ug of DNA [23] 2. LiAc/PEG/ssD NA:~ 1.7 × 10 ² CFU per ug of DNA [39]	Not reported	Not reported
Candida intermedia	Not reported	Yes (7.8 × 10 ¹ CFU per ug of DNA) [45]	Not reported
Yamadazyma tenuis	Not reported	Not reported	Not reported
Pachysolen tannophilus	Yes (LiAc): 4.0 - 5.0 × 10 ¹ CFU per ug of DNA [59] (LiAc/PEG/ssDNA): 3.0 × 10 ⁴ CFU per ug of DNA [56]	Not reported	Yes (2.99 × 10 ⁻² transconjugants per recipient at donor-to-recipient ratio of 1000:1) [53,56]

Abbreviations: colony-forming units (CFU); lithium acetate (LiAc); polyethylene glycol

(PEG); single-stranded DNA (ssDNA).

Table 4. Regulatory sequences (promoters and terminators) used for the expression of coding sequences in D-xylose fermenting yeasts. Inducible promoter sequences are shown in boldface.

Yeast species	Regulatory sequences		
	Promoters	Terminators	
Scheffersomyces stipitis	SsTEF1; SsTDH3; BaTEF1; SsXYL1.1; EgTEF1; SsCCW12; SsPGK1; SsHHF1; SsADH1; SsCYC1; SsGAL1¹; SsXKS1¹ [20,23,33,35]	SsXYL1; SsXYL2; SpCYC1;SsADH1; SsADH2; SsACT1 [20,23,33,35]	
Spathaspora passalidarum	EgTEF1; SpTEF1; SpADH1; SsXYL1.1 [23]	ScCYC1; SpXYL1.1; SpTEF1; SpADH1 [23]	
Candida intermedia	<i>EgTEF1; CaSAP2²</i> [45]	<i>EgTEF1</i> [45]	
Yamadazyma tenuis	Not reported	Not reported	
Pachysolen tannophilus	PtGAPDH [58]	PtGAPDH [58]	

¹The SsGAL1 and SsXKS1 promoters are induced by the presence of galactose or

xylose in culture media.

²The *CaSAP2* promoter is induced by bovine serum albumin (BSA) in culture media without additional nitrogen sources.

Abbreviations: Blastobotrys adeninivorans (Ba); Candida albicans (Ca); Eremothecium

gossypii (Eg); Pachysolen tannophilus (Pt); Saccharomyces cerevisiae (Sc);

Scheffersomyces stipitis (Ss), Spathaspora passalidarum (Sp).