

Elliptic Quantum Groups

Hitoshi Konno

*Department of Mathematics, Tokyo University of Marine Science and Technology,
Etchujima, Koto, Tokyo 135-8533, Japan
hkonno0@kaiyodai.ac.jp*

Abstract

We expose the elliptic quantum groups in the Drinfeld realization associated with both the affine Lie algebra \mathfrak{g} and the toroidal algebra \mathfrak{g}_{tor} . There the level-0 and level $\neq 0$ representations appear in a unified way so that one can define the vertex operators as intertwining operators of them. The vertex operators are key for many applications such as a derivation of the elliptic weight functions, integral solutions of the (elliptic) q -KZ equation and a formulation of algebraic analysis of the elliptic solvable lattice models. Identifying the elliptic weight functions with the elliptic stable envelopes we make a correspondence between the level-0 representation of the elliptic quantum group and the equivariant elliptic cohomology. We also emphasize a characterization of the elliptic quantum groups as q -deformations of the W -algebras.

Keywords : Elliptic quantum group, Affine Lie algebra, Toroidal algebra, deformed W -algebra, q -KZ equation, Hypergeometric integral, Quantum integrable system, Quiver variety, Elliptic cohomology, Stable envelope

Key points :

- The elliptic quantum groups $U_{q,p}(\mathfrak{g})$ and $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ are formulated in terms of the Drinfeld generators in a uniform way¹ and equipped with H -Hopf algebroid structure as a coalgebra structure.
- $U_{q,p}(\mathfrak{g})$ is characterized both as a p -deformation of the quantum affine algebra $U_q(\mathfrak{g})$ and as a q -deformation of the W -algebra of the coset type $\mathfrak{g} \oplus \mathfrak{g} \supset \mathfrak{g}_{diag}$. In the same way, $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ is characterized both as a p -deformation of the quantum toroidal algebra $U_{q,\kappa}(\mathfrak{g}_{tor})$ and as a (q, κ) -deformation of affine quiver W -algebra.
- The vertex operators defined as intertwiners of the tensor product of the level-0 and $\neq 0$ modules are key objects for many applications.

¹One may unify them as “quiver elliptic quantum groups”, where $U_{q,p}(\mathfrak{g})$ is a finite Dynkin quiver type, whereas $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ is an affine Dynkin quiver type.

- The Gelfand-Tsetlin basis for the level-0 representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ is constructed explicitly by using the elliptic weight function as the change of basis matrix from the standard basis.
- The elliptic weight function of type \mathfrak{sl}_N is identified with the elliptic stable envelope for the equivariant elliptic cohomology of the cotangent bundle to the partial flag variety $E_T(T^*fl_\lambda)$. This allows us to make a dictionary between $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ and $E_T(T^*fl_\lambda)$.

1 Introduction

Quantum groups are algebraic systems associated with solutions of the Yang-Baxter equation (YBE). They are in general associative algebras classified by finite dimensional simple Lie algebras $\bar{\mathfrak{g}}$, affine Lie algebras \mathfrak{g} or toroidal algebras \mathfrak{g}_{tor} , and equipped with certain coalgebra structure. Typical examples are Yangians $Y(\bar{\mathfrak{g}})$, Yangian doubles $\mathcal{DY}(\bar{\mathfrak{g}})$, quantum groups $U_q(\bar{\mathfrak{g}})$, affine quantum groups $U_q(\mathfrak{g})$ and quantum toroidal algebras $U_{q,\kappa}(\mathfrak{g}_{tor})$. See for example [11, 26, 60].

Similarly elliptic quantum groups (EQG) are quantum groups associated with elliptic solutions of the YBE classified by affine Lie algebras \mathfrak{g} or toroidal algebras \mathfrak{g}_{tor} . In elliptic setting, it is known at least for the cases classified by \mathfrak{g} that there are two types of YBEs, the vertex type and the face type (Fig.1 and Fig.2). Accordingly, there are two types of elliptic quantum groups, the vertex type and the face type [29]. Both of them are dynamical quantum groups, whose universal R matrices [29] satisfy the dynamical YBE (DYBE) [21]².




Figure 1: The vertex type Yang-Baxter equation

There are several formulations of the elliptic quantum groups. They can be classified by their generators and coalgebra structures. See Table 1 : $\mathcal{A}_{q,p}(\widehat{\mathfrak{sl}}_N)$ and $\mathcal{B}_{q,\lambda}(\mathfrak{g})$ in terms of the Chevalley generators, $U_{q,p}(\mathfrak{g})$ in terms of the Drinfeld generators and $E_{q,p}(\widehat{\mathfrak{g}}_N)$ in terms of the L operators. Only $\mathcal{A}_{q,p}(\widehat{\mathfrak{sl}}_N)$ is the vertex type. The others are the face type. Coalgebra structures

²Obviously the face type YBE is dynamical [21]. The vertex type is also dynamical, because the elliptic nome p is a dynamical parameter getting a shift by a central element q^{-2c} and yields the new elliptic nome $p^* = pq^{-2c}$ [29].




Figure 2: The face type Yang-Baxter equation

are the quasi-Hopf algebra structure for $\mathcal{A}_{q,p}(\widehat{\mathfrak{sl}}_N)$, $\mathcal{B}_{q,\lambda}(\mathfrak{g})$, and the Hopf algebroid structure for $E_{q,p}(\widehat{\mathfrak{gl}}_N)$, $U_{q,p}(\mathfrak{g})$, respectively. For expositions of these formulations with historical remarks and arguments for their consistency, the reader can consult [45].

	co-algebra structure	generators
$\mathcal{A}_{q,p}(\widehat{\mathfrak{sl}}_N)$ (vertex type)	quasi-Hopf algebra	Chevalley
$\mathcal{B}_{q,\lambda}(\mathfrak{g})$ (face type)		
$E_{q,p}(\mathfrak{g})$ (face type)	Hopf Algebroid	L -operator
$U_{q,p}(\mathfrak{g})$ (face type)	Hopf Algebroid	Drinfeld

Table 1: Three formulations of the elliptic quantum groups

In this article, we expose the elliptic quantum group $U_{q,p}(\mathfrak{g})$ and its toroidal version $U_{q,\kappa,p}(\mathfrak{g}_{tor})$. They are generated by the Drinfeld generators, i.e. analogues of the loop generators of the affine Lie algebra \mathfrak{g} [33]. After giving definitions of $U_{q,p}(\mathfrak{g})$ and some typical representations of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$, we introduce the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ and describe their applications : derivations of the elliptic weight functions and integral solutions of the elliptic q -KZ equation, construction of the level-0 action on the Gelfand-Tsetlin basis, algebraic analysis of the elliptic solvable lattice models. We then describe a geometric interpretation of the level-0 representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ in terms of the equivariant elliptic cohomology of the cotangent bundle to the partial flag variety $E_T(T^*fl_\lambda)$. We also emphasize a characterization of $U_{q,p}(\mathfrak{g})$ as a q -deformation of the W -algebra of the coset type. In the final section we briefly describe a formulation of the elliptic quantum toroidal algebras, $U_{q,\kappa,p}(\mathfrak{g}_{tor})$, $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ and expose their connections to the

Macdonald theory, affine quiver W -algebra and application to instanton calculus of the super symmetric gauge theories.

2 Elliptic Algebra $U_{q,p}(\mathfrak{g})$

Through this paper, let p, q be generic complex numbers satisfying $|p|, |q| < 1$. Let $\bar{\mathfrak{g}}$ be a simple Lie algebra over \mathbb{C} and $\mathfrak{g} = X_l^{(1)}$ the corresponding untwisted affine Lie algebra with the generalized Cartan matrix $A = (a_{ij})_{i,j \in I}$, $I = \{0\} \cup \bar{I}$, $\bar{I} = \{1, \dots, l\}$, $\text{rank} A = l$. We denote by $B = (b_{ij})_{i,j \in I}$, $b_{ij} = d_i a_{ij}$ the symmetrization of A and set $q_i = q^{d_i}$. We also use the notations for $n \in \mathbb{Z}$,

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \quad [n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}.$$

We fix a realization $(\mathfrak{h}, \Pi, \Pi^\vee)$ of A , i.e. \mathfrak{h} is a $l+2$ -dimensional \mathbb{C} -vector space, $\Pi = \{\alpha_0, \alpha_1, \dots, \alpha_l\} \subset \mathfrak{h}^*$ a set of simple roots, and $\Pi^\vee = \{h_0, h_1, \dots, h_l\} \subset \mathfrak{h}$ a set of simple coroots satisfying $\langle \alpha_j, h_i \rangle = a_{ij}$ ($i, j \in I$) for a canonical pairing $\langle \cdot, \cdot \rangle : \mathfrak{h}^* \times \mathfrak{h} \rightarrow \mathbb{C}$ [33]. Set also $\mathcal{Q} = \sum_{i \in I} \mathbb{Z} \alpha_i$. We take $\{h_1, \dots, h_l, c, d\}$ as the basis of \mathfrak{h} and $\{\bar{\Lambda}_1, \dots, \bar{\Lambda}_l, \Lambda_0, \delta\}$ the dual basis satisfying

$$\langle \delta, d \rangle = 1 = \langle \Lambda_0, c \rangle, \quad \langle \bar{\Lambda}_i, h_j \rangle = \delta_{i,j},$$

with the other pairings being 0.

Let H_P be a \mathbb{C} -vector space spanned by P_0, P_1, \dots, P_l and H_P^* be its dual space spanned by Q_0, Q_1, \dots, Q_l with a pairing $\langle Q_i, P_j \rangle = a_{ij}$. For $\alpha = \sum_{i \in I} c_i \alpha_i \in \mathfrak{h}^*$, we set $P_\alpha = \sum_{i \in I} c_i P_i$, $Q_\alpha = \sum_{i \in I} c_i Q_i$. In particular $P_{\alpha_i} = P_i$, $Q_{\alpha_i} = Q_i$. Define also an analogue of the root lattice $\mathcal{R}_Q = \sum_{i \in I} \mathbb{Z} Q_i$.

Let us consider $H := \mathfrak{h} \oplus H_P$ and $H^* := \mathfrak{h}^* \oplus H_P^*$ with a pairing $\langle \cdot, \cdot \rangle : H^* \times H \rightarrow \mathbb{C}$ by extending those on $\mathfrak{h}^* \times \mathfrak{h}$ and $H_P^* \times H_P$ with $\langle \mathfrak{h}^*, H_P \rangle = 0 = \langle H_P^*, \mathfrak{h} \rangle$. We denote by $\mathbb{F} = \mathcal{M}_{H^*}$ the field of meromorphic functions on H^* . We regard a meromorphic function $g(h, P)$ of $h \in \mathfrak{h}, P \in H_P$ as an element in \mathbb{F} by $g(h, P)(\mu) = g(\langle \mu, h \rangle, \langle \mu, P \rangle)$ for $\mu \in H^*$.

2.1 Definition

The elliptic algebra $U_{q,p}(\mathfrak{g})$ is a topological algebra over $\mathbb{F}[[p]]$ generated by the Drinfeld generators

$$\alpha_{i,m}, \quad e_{i,n}, \quad f_{i,n}, \quad K_i^\pm, \quad q^{\pm c/2}, \quad q^d \quad (i \in \bar{I}, \quad m \in \mathbb{Z} \setminus \{0\}, \quad n \in \mathbb{Z}).$$

In order to write down the defining relations, it is convenient to introduce their generating functions $e_i(z)$, $f_i(z)$ and $\phi_i^\pm(z)$ called the elliptic currents.

$$\begin{aligned} e_i(z) &= \sum_{n \in \mathbb{Z}} e_{i,n} z^{-n}, & f_i(z) &= \sum_{n \in \mathbb{Z}} f_{i,n} z^{-n}, \\ \phi_i^+(q^{-c/2}z) &= K_i^+ \exp\left(- (q_i - q_i^{-1}) \sum_{n>0} \frac{\alpha_{i,-n}}{1-p^n} z^n\right) \exp\left((q_i - q_i^{-1}) \sum_{n>0} \frac{p^n \alpha_{i,n}}{1-p^n} z^{-n}\right), \\ \phi_i^-(q^{c/2}z) &= K_i^- \exp\left(- (q_i - q_i^{-1}) \sum_{n>0} \frac{p^n \alpha_{i,-n}}{1-p^n} z^n\right) \exp\left((q_i - q_i^{-1}) \sum_{n>0} \frac{\alpha_{i,n}}{1-p^n} z^{-n}\right). \end{aligned}$$

We also set $p^* = pq^{-2c}$. The defining relations are given as follows. For $g(h, P) \in \mathbb{F}$,

$$\begin{aligned} q^{\pm c/2} &: \text{central}, \\ g(h, P)e_j(z) &= e_j(z)g(h + \langle \alpha_j, h \rangle, P - \langle Q_j, P \rangle), & g(h, P)f_j(z) &= f_j(z)g(h - \langle \alpha_j, h \rangle, P), \\ [g(h, P), \alpha_{i,m}] &= 0, & [g(h, P), q^d] &= 0, & g(h, P)K_j^\pm &= K_j^\pm g(h, P - \langle Q_j, P \rangle), \\ [q^d, \alpha_{j,m}] &= q^m \alpha_{j,m}, & [q^d, x_j^\pm(z)] &= x_j^\pm(q^{-1}z) \\ [K_i^\pm, K_j^\pm] &= [K_i^\pm, K_j^\mp] = 0 = [K_i^\pm, \alpha_{j,m}], \\ K_i^\pm e_j(z)(K_i^\pm)^{-1} &= q^{\mp \langle \alpha_j, h_i \rangle} e_j(z), & K_i^\pm f_j(z)(K_i^\pm)^{-1} &= q^{\pm \langle \alpha_j, h_i \rangle} f_j(z), \end{aligned}$$

$$\begin{aligned} [\alpha_{i,m}, \alpha_{j,n}] &= \delta_{m+n,0} \frac{[a_{ij}m]_i q^{cm} - q^{-cm}}{m} \frac{1-p^m}{q_j - q_j^{-1}} \frac{1-p^m}{1-p^*m} q^{-cm}, \\ [\alpha_{i,m}, e_j(z)] &= \frac{[a_{ij}m]_i}{m} \frac{1-p^m}{1-p^*m} q^{-cm} z^m e_j(z), \\ [\alpha_{i,m}, f_j(z)] &= -\frac{[a_{ij}m]_i}{m} z^m f_j(z), \\ (z - q^{bij}w)g_{ij}(w/z; p^*)e_i(z)e_j(w) &= (q^{bij}z - w)g_{ij}(z/w; p^*)e_j(w)e_i(z), \\ (z - q^{-bij}w)g_{ij}(w/z; p)^{-1}f_i(z)f_j(w) &= (q^{-bij}z - w)g_{ij}(z/w; p)^{-1}f_j(w)f_i(z), \\ [e_i(z), f_j(w)] &= \frac{\delta_{i,j}}{q_i - q_i^{-1}} \left(\delta(q^c w/z) \phi_i^-(q^{c/2}w) - \delta(q^{-c}w/z) \phi_i^+(q^{c/2}z) \right), \\ &+ \text{Serre relations.} \end{aligned}$$

Here we set

$$g_{ij}(z; s) = \exp\left(- \sum_{m>0} \frac{1}{m} \frac{q^{bijm} - q^{-bijm}}{1-s^m} (sz)^m\right) \in \mathbb{C}[[s]][[z]].$$

These relations are treated as formal Laurent series in the argument of the elliptic currents i.e. z, w etc.. All the coefficients in z, w etc. are well defined in the p -adic topology.

It is easy to find that in the limit $p \rightarrow 0$ the above relations except for those indicating the non-commutativity of \mathbb{F} and $e_i(z), f_i(z), K_j^\pm$ go to the defining relations of the quantum affine

algebra $U_q(\mathfrak{g})$ [15]. One also finds that the non-commutativity of \mathbb{F} can be realized as the one of P_j and Q_i by setting $[P_i, Q_j] = a_{i,j}$ ($i, j \in I$). Hence by using the group algebra $\mathbb{C}[\mathcal{R}_Q]$ of \mathcal{R}_Q , i.e. $e^{Q_\alpha}, e^{Q_\beta}, e^{Q_\alpha}e^{Q_\beta} = e^{Q_\alpha+Q_\beta}, e^0 = 1 \in \mathbb{C}[\mathcal{R}_Q]$, one obtains the following isomorphism [17].

For generic p, q ,

$$U_{q,p}(\mathfrak{g})/pU_{q,p}(\mathfrak{g}) \cong (U_q(\mathfrak{g}) \otimes \mathbb{F}[[p]]) \sharp \mathbb{C}[\mathcal{R}_Q].$$

Here the smash product \sharp expresses the non-commutativity between \mathbb{F} and $\mathbb{C}[\mathcal{R}_Q]$.

Remark. For representations, on which $q^{\pm c/2}$ take complex values e.g. $q^{\pm k/2}$ (see Sec.4), we treat p and $p^* = pq^{-2k}$ as generic complex numbers satisfying $|p| < 1$ and $|p^*| < 1$. Then one has

$$g_{ij}(z; s) = \frac{(sq^{b_{ij}}z; s)_\infty}{(sq^{-b_{ij}}z; s)_\infty}, \quad (z; s)_\infty = \prod_{n=0}^{\infty} (1 - zs^n)$$

for $|sq^{\pm b_{ij}}z| < 1$, $s = p, p^*$. Hence in the sense of analytic continuation, one can rewrite the relations of $e_i(z)$ and $e_j(w)$ and of $f_i(z)$ and $f_j(w)$ as

$$z\theta_{p^*}(q^{b_{ij}}w/z)e_i(z)e_j(w) = -w\theta_{p^*}(q^{b_{ij}}z/w)e_j(w)e_i(z), \quad (2.1)$$

$$z\theta_p(q^{-b_{ij}}w/z)f_i(z)f_j(w) = -w\theta_p(q^{-b_{ij}}z/w)f_j(w)f_i(z), \quad (2.2)$$

respectively. Here $\theta_p(z)$ denotes the odd theta function given by

$$\theta_p(z) = (z; p)_\infty (p/z; p)_\infty (p; p)_\infty.$$

The additive notations are also often used. Introduce $r, r^* = r - k \in \mathbb{R}_{>0}$ and set $p = q^{2r}, p^* = pq^{-2k} = q^{2r^*}$,

$$E_i(z) := e_i(z)z^{-\frac{P\alpha_i-1}{r^*}}, \quad F_i(z) := f_i(z)z^{\frac{(P+h)\alpha_i-1}{r}}.$$

Then from (2.1), (2.2) and the non-commutativity of h, P with $e_i(z), f_i(z)$ one obtains

$$F_i(z_1)F_j(z_2) = \frac{[u_1 - u_2 - a_{ij}/2]}{[u_1 - u_2 + a_{ij}/2]} F_j(z_2)F_i(z_1),$$

$$E_i(z_1)E_j(z_2) = \frac{[u_1 - u_2 + a_{ij}/2]^*}{[u_1 - u_2 - a_{ij}/2]^*} E_j(z_2)E_i(z_1).$$

Here we set $z_i = q^{2u_i}$ ($i = 1, 2$). The symbols $[u]$ and $[u]^*$ denote Jacobi's odd theta functions given by

$$[u] = q^{\frac{u^2}{r}-u}\theta_p(q^{2u}), \quad [u]^* = q^{\frac{u^2}{r^*}-u}\theta_{p^*}(q^{2u}).$$

They have the following quasi periodicity.

$$\begin{aligned} [u + r] &= -[u], & [u + r\tau] &= -e^{-\pi i\tau} e^{-2\pi iu/r} [u], \\ [u + r^*]^* &= -[u]^*, & [u + r^*\tau^*]^* &= -e^{-\pi i\tau^*} e^{-2\pi iu/r^*} [u]^*, \end{aligned}$$

where $p = e^{-2\pi i/\tau}$, $p^* = e^{-2\pi i/\tau^*}$.

2.2 Coalgebra structure

In order to formulate a coalgebra structure over $\mathbb{F} = \mathcal{M}_{H^*}$, which does not commute with the Drinfeld generators, one needs to extend the Hopf algebra to the H -Hopf algebroid. For the elliptic algebra $\mathcal{U} = U_{q,p}(\mathfrak{g})$, it is roughly sketched as follows. For expositions of the background concepts of H -Hopf algebroid, the reader can consult [16, 35, 41].

Let $P + h = \sum_{i \in I} c_i(P_i + h_i) \in H$, $c_i \in \mathbb{C}$. The algebra \mathcal{U} is a H -algebra by

$$\begin{aligned} \mathcal{U} &= \bigoplus_{\alpha, \beta \in \mathfrak{h}^*} \mathcal{U}_{\alpha, \beta} \\ (\mathcal{U})_{\alpha\beta} &= \left\{ x \in U \mid q^{P+h} x q^{-(P+h)} = q^{\langle \alpha, P+h \rangle} x, \quad q^P x q^{-P} = q^{\langle Q_\beta, P \rangle} x \quad \forall P + h, P \in H \right\} \end{aligned}$$

with the two moment maps $\mu_l, \mu_r : \mathbb{F} \rightarrow \mathcal{U}_{0,0}$ defined by

$$\mu_l(\widehat{f}) = f(h, P + h, p) \in \mathbb{F}[[p]], \quad \mu_r(\widehat{f}) = f(h, P, p^*) \in \mathbb{F}[[p^*]]$$

for $\widehat{f} = f(h, P, p^*) \in \mathbb{F}[[p^*]]$. Here $p^* = pq^{-2c}$ as before.

The tensor product $\mathcal{U} \widetilde{\otimes} \mathcal{U}$ is the \mathfrak{h}^* -bigraded vector space with

$$(\mathcal{U} \widetilde{\otimes} \mathcal{U})_{\alpha\beta} = \bigoplus_{\gamma \in \mathfrak{h}^*} (\mathcal{U}_{\alpha\gamma} \otimes_{\mathcal{M}_{H^*}} \mathcal{U}_{\gamma\beta}),$$

where $\otimes_{\mathcal{M}_{H^*}}$ denotes the ordinary tensor product modulo the following relation.

$$\mu_r(\widehat{f}) a \widetilde{\otimes} b = a \widetilde{\otimes} \mu_l(\widehat{f}) b \quad a, b \in \mathcal{U}.$$

Let us regard $T_\alpha = e^{-Q_\alpha} \in \mathbb{C}[\mathcal{R}_Q]$ as a shift operator

$$(T_\alpha \widehat{f}) = f(h, P + \langle Q_\alpha, P \rangle, p).$$

Then $\mathcal{D} = \{\widehat{f} e^{-Q_\alpha} \mid \widehat{f} \in \mathbb{F}, e^{-Q_\alpha} \in \mathbb{C}[\mathcal{R}_Q]\}$ becomes the H -algebra having the property

$$\mathcal{U} \cong \mathcal{U} \widetilde{\otimes} \mathcal{D} \cong \mathcal{D} \widetilde{\otimes} \mathcal{U}$$

by $a \cong a \widetilde{\otimes} T_{-\beta} \cong T_{-\alpha} \widetilde{\otimes} a$ for all $a \in \mathcal{U}_{\alpha\beta}$.

Let $\pi_V, V = \bigoplus_i \mathbb{C}v_i$ be the vector representation of $\bar{\mathfrak{g}}$ and consider the elliptic dynamical R -matrix $R^+(z, \Pi) \in \text{End}(V \otimes V)$ of type \mathfrak{g} [29, 39].

For example, the $\widehat{\mathfrak{sl}}_N$ type is given by

$$\begin{aligned} R^+(z, \Pi) &= \rho^+(z) \bar{R}(z, \Pi), \\ \bar{R}(z, \Pi) &= \sum_{j=1}^N E_{j,j} \otimes E_{j,j} + \sum_{1 \leq j_1 < j_2 \leq N} (b(u, (P+h)_{j_1, j_2}) E_{j_1, j_1} \otimes E_{j_2, j_2} + \bar{b}(u) E_{j_2, j_2} \otimes E_{j_1, j_1} \\ &\quad + c(u, (P+h)_{j_1, j_2}) E_{j_1, j_2} \otimes E_{j_2, j_1} + \bar{c}(u, (P+h)_{j_1, j_2}) E_{j_2, j_1} \otimes E_{j_1, j_2}), \end{aligned} \quad (2.3)$$

where $E_{i,j} v_k = \delta_{j,k} v_i$, $z = q^{2u}$, $\Pi_{j,k} = q^{2(P+h)_{j,k}}$, $(P+h)_{j,k} = \sum_{i=j}^{k-1} (P_i + h_i)$,

$$\begin{aligned} \rho^+(z) &= q^{-\frac{N-1}{N}} z^{\frac{N-1}{rN}} \frac{\Gamma(z; p, q^{2N}) \Gamma(q^{2N} z; p, q^{2N})}{\Gamma(q^2 z; p, q^{2N}) \Gamma(q^{2N-2} z; p, q^{2N})}, \\ b(u, s) &= \frac{[s+1][s-1][u]}{[s]^2[u+1]}, \quad \bar{b}(u) = \frac{[u]}{[u+1]}, \\ c(u, s) &= \frac{[1][s+u]}{[s][u+1]}, \quad \bar{c}(u, s) = \frac{[1][s-u]}{[s][u+1]}. \end{aligned}$$

The symbol $\Gamma(z; p, q^{2N})$ denotes the elliptic Gamma function defined by

$$\Gamma(z; p, q) = \frac{(pq/z; p, q)_\infty}{(z; p, q)_\infty}, \quad (z; p, q)_\infty = \prod_{m, n=0}^{\infty} (1 - zp^m q^n), \quad |p|, |q| < 1. \quad (2.4)$$

Let $L^+(z) = \sum_{i,j} E_{i,j} L_{i,j}^+(z) \in \text{End}(V) \otimes \tilde{\mathcal{U}}[[z, z^{-1}]]$ be the the L -operator satisfying the dynamical RLL -relation [29]

$$R^{+(12)}(z_1/z_2, \Pi) L^{+(1)}(z_1) L^{+(2)}(z_2) = L^{+(2)}(z_2) L^{+(1)}(z_1) R^{+*(12)}(z_1/z_2, \Pi^*).$$

Here $R^{+*}(z, \Pi^*)$ denotes the same elliptic R -matrix $R^+(z, \Pi^*)$ with $\Pi_{i,j}^* = q^{2P_{i,j}}$ except for replacing the elliptic nome p by p^* . Define two H -algebra homomorphisms, the counit $\varepsilon : \mathcal{U} \rightarrow \mathcal{D}$ and the comultiplication $\Delta : \mathcal{U} \rightarrow \mathcal{U} \otimes \mathcal{U}$, and the antihomomorphism $S : \mathcal{U} \rightarrow \mathcal{U}$ by

$$\begin{aligned} \varepsilon(L_{i,j}^+(z)) &= \delta_{i,j} T_{Q_{\varepsilon_i}} \quad (n \in \mathbb{Z}), \quad \varepsilon(e^Q) = e^Q, \\ \varepsilon(\mu_l(\hat{f})) &= \varepsilon(\mu_r(\hat{f})) = \hat{f} T_0, \\ \Delta(L_{i,j}^+(z)) &= \sum_k L_{i,k}^+(z) \tilde{\otimes} L_{k,j}^+(z), \\ \Delta(e^Q) &= e^Q \tilde{\otimes} e^Q, \\ \Delta(\mu_l(\hat{f})) &= \mu_l(\hat{f}) \tilde{\otimes} 1, \quad \Delta(\mu_r(\hat{f})) = 1 \tilde{\otimes} \mu_r(\hat{f}), \\ S(L_{i,j}^+(z)) &= (L^+(z)^{-1})_{ij}, \\ S(e^Q) &= e^{-Q}, \quad S(\mu_r(\hat{f})) = \mu_l(\hat{f}), \quad S(\mu_l(\hat{f})) = \mu_r(\hat{f}). \end{aligned}$$

Here $\alpha_i = \epsilon_i - \epsilon_{i+1}$ ($i \in \bar{I}$). Then the set $(U_{q,p}(\mathfrak{g}), H, \mathcal{M}_{H^*}, \mu_l, \mu_r, \Delta, \varepsilon, S)$ becomes a H -Hopf algebroid [41, 44–46].

An explicit realization of $L^+(z)$ in terms of the elliptic currents of \mathcal{U} was studied in [28, 36, 44, 46]. In general, the connection between $L^+(z)$ and the elliptic currents is given as follows. Let $\{h^1, \dots, h^l\}$ be the dual basis to $\{h_1, \dots, h_l\}$ of $\bar{\mathfrak{h}}$ and (π_V, V) be the vector representation of $\bar{\mathfrak{g}}$. Define [29]

$$\begin{aligned} L^-(z) &= (\text{Ad}(q^{-2\theta_V(P)}) \otimes \text{id})(q^{2T_V} L^+(zpq^{-c}), \\ \theta_V(P) &= -\frac{1}{2} \sum_j (\pi_V(h_j) \pi_V(h^j) + 2P_j \pi_V(h^j)), \\ T_V &= \sum_j \pi_V(h_j) \otimes h^j. \end{aligned}$$

One finds that $L^\pm(z)$ satisfy

$$R^{\pm(12)}(z_1/z_2, \Pi) L^{\pm(1)}(z_1) L^{\pm(2)}(z_2) = L^{\pm(2)}(z_2) L^{\pm(1)}(z_1) R^{\pm*(12)}(z_1/z_2, \Pi^*), \quad (2.5)$$

$$R^{+(12)}(q^c z_1/z_2, \Pi) L^{+(1)}(z_1) L^{-(2)}(z_2) = L^{-(2)}(z_2) L^{+(1)}(z_1) R^{+*(12)}(q^{-c} z_1/z_2, \Pi^*). \quad (2.6)$$

Here $R^\pm(z, \Pi)$, $R^{\pm*}(z, \Pi^*)$ and $L^+(z)$ are related to $R_{V_V}^\pm(z, \lambda+h)$, $R_{V_V}^\pm(z, \lambda)$ and $L_V^+(z, \lambda)$ in [29] by

$$\begin{aligned} R^\pm(z, \Pi) &= R_{V_V}^\pm(z, \lambda+h), & R^{\pm*}(z, \Pi^*) &= R_{V_V}^\pm(z, \lambda), \\ L^+(z) &= L_V^+(z, \lambda) e^{-\sum_i E_{i,i} Q_{\epsilon_i}} \end{aligned}$$

with $\lambda = (r-k+h^\vee)d + \sum_i (P_{\alpha'_i} + 1) \bar{h}^i$, $\lambda+h = (r+h^\vee)d + \sum_i (P_{\alpha'_i} + h_{\alpha'_i} + 1) \bar{h}^i$, $[P_{i,j}, Q_{\epsilon_k}] = \delta_{i,k} - \delta_{j,k}$ and α'_i being the simple root of the Langlands dual Lie algebra of $\bar{\mathfrak{g}}$ [39]. Then consider the following Gauss decomposition of $L^\pm(z)$.

$$L^\pm(z) = F^\pm(z) K^\pm(z) E^\pm(z).$$

Here $K^\pm(z)$ are diagonal matrices with entries $K_j^\pm(z)$, whereas $F^\pm(z)$ (resp. $E^\pm(z)$) are upper (resp. lower) triangular matrices with entries $F_{i,j}^\pm(z)$ (resp. $E_{j,i}^\pm(z)$) ($i < j$) and all diagonal entries 1. By combining the relations for these entries obtained from (2.5)-(2.6), one finds the following identification with the elliptic currents $E_j(z)$, $F_j(z)$ and $\phi_j^\pm(z)$.

$$\begin{aligned} E_j(zq^{j-c/2}) &= \mu^* \left(E_{j+1,j}^+(zq^{c/2}) - E_{j+1,j}^-(zq^{-c/2}) \right), \\ F_j(zq^{j-c/2}) &= \mu \left(F_{j,j+1}^+(zq^{-c/2}) - F_{j,j+1}^-(zq^{c/2}) \right), \\ \phi_j^\pm(zq^{-c/2}q^j) &= \kappa K_j^\pm(z) K_{j+1}^\pm(z)^{-1}. \end{aligned}$$

Here μ, μ^*, κ are constants satisfying

$$\mu\mu^* = -\kappa \frac{q}{q - q^{-1}} \frac{(p^*; p^*)_\infty}{\theta_{p^*}(q^2)}.$$

See for example [44, 45].

Note also that to formulate a H -Hopf algebroid structure one may also use the Drinfeld comultiplication for the elliptic currents [47, 48] instead of the standard comultiplication for $L^+(z)$ given in the above. The resultant H -Hopf algebroid is completely different coalgebra structure from the above. These two coalgebra structures have their own appropriate applications. In particular, they define different vertex operators as intertwining operators of $U_{q,p}(\mathfrak{g})$ -modules, even if the modules are the same. The standard one is appropriate to discuss the problems exposed in the subsequent sections Sec.5-10, whereas the Drinfeld comultiplication is appropriate to discuss problems related to the deformed W -algebras. See Sec.3 and Sec.11. One should also note that for the quantum toroidal algebras, whatever they are trigonometric or elliptic, only the coalgebra structures associated with the Drinfeld comultiplication are available at the moment, because of a lack of expressions of the (elliptic) R -matrices and the L -operators.

3 $U_{q,p}(\mathfrak{g})$ as Deformed W -algebra

The elliptic algebra $U_{q,p}(\mathfrak{g})$ has another characterization as a q -deformation of the W -algebra $\mathcal{W}_{\bar{\mathfrak{g}}}$ of the coset type $(\mathfrak{g})_{r-h\nu-k} \oplus (\mathfrak{g})_k \supset (\mathfrak{g}_{\text{diag}})_{r-h\nu}$ [9, 12, 27], or more precisely as an algebra of screening currents of the deformation of $\mathcal{W}_{\bar{\mathfrak{g}}}$. Here we assume \mathfrak{g} is an untwisted affine Lie algebra and realized as a central extension of the loop algebra [33]. The generating functions of \mathfrak{g} is called the currents. The symbol $(\mathfrak{g})_k$ means to consider \mathfrak{g} at the level k representation. A key for understanding this characterization of $U_{q,p}(\mathfrak{g})$ is the (generalized) Feigin-Fuchs (FF) construction of $\mathcal{W}_{\bar{\mathfrak{g}}}$ from \mathfrak{g} [12, 19]. There one starts from the level- k currents of \mathfrak{g} in the Lepowsky-Wilson (LW) realization [53] and the corresponding energy-momentum (EM) tensor of the Wess-Zumino-Witten (WZW) model, i.e. the generating function of the Virasoro algebra. The LW realization of \mathfrak{g} is a realization of the currents in terms of the $\text{rk } \mathfrak{g}$ number of free bosons and the level- k Z -algebra. The EM tensor is obtained by the Sugawara construction in terms of the currents. Then the FF construction deforms the EM tensor by introducing the so-called background charge term, which depends on a parameter r , and at the same time deforms the free boson part of the currents of \mathfrak{g} by r and the Z -algebra part by adding the dynamical parameters. The deformed EM tensor becomes the generating function of the Virasoro generators of the coset W -algebra $\mathcal{W}_{\bar{\mathfrak{g}}}$, whereas the deformed currents of \mathfrak{g} becomes the so-called screening currents, i.e. the primary field of conformal dimension one w.r.t. this new Virasoro generators. All the

generating functions of $\mathcal{W}\bar{\mathfrak{g}}$, the Virasoro as well as the other generators of higher conformal dimensions, are characterized as the intersection of the kernel of the screening operators i.e. certain contour integral of the screening currents, on $U(\mathfrak{g})$ [18]. In this sense, we refer the resultant r -deformation (+ modification by the dynamical parameters) of the affine Lie algebra \mathfrak{g} to the W -algebra $\mathcal{W}\bar{\mathfrak{g}}$. When one consider a q -deformation of it, \mathfrak{g} and its r -deformation $\mathcal{W}\bar{\mathfrak{g}}$ turn out to give $U_q(\mathfrak{g})$ and its p -deformation $U_{q,p}(\mathfrak{g})$ with $p = q^{2r}$, respectively.

For example, the currents of the level $k(\neq 0)$ affine Lie algebra $\widehat{\mathfrak{sl}}_2$ has the following LW realization.

$$e(z) = Z^+(z) : \exp \left\{ \sqrt{2k} \sum_{m \neq 0} \frac{a_m}{km} z^{-m} \right\} :, \quad f(z) = Z^-(z) : \exp \left\{ -\sqrt{2k} \sum_{m \neq 0} \frac{a_m}{km} z^{-m} \right\} :,$$

where $Z^\pm(z)$ are the generating functions of the level- k Z -algebra, and a_m ($m \in \mathbb{Z} \setminus \{0\}$) generate the Heisenberg subalgebra of $\widehat{\mathfrak{sl}}_2$ satisfying

$$[\sqrt{2k}a_m, \sqrt{2k}a_n] = 2mk\delta_{m+n,0}. \quad (3.1)$$

The generating function $\widetilde{\phi}(z) = \sum_{m \neq 0} \frac{a_m}{m} z^{-m}$ is called free boson. The Z -algebra is realized in terms of the \mathbb{Z}_k -parafermions $\Psi(z), \Psi^\dagger(z)$ and the zero mode operators α, h satisfying $[h, \alpha] = 2$ as follows.

$$Z^+(z) = \Psi(z)e^{\alpha}z^{h/k}, \quad Z^-(z) = \Psi^\dagger(z)e^{-\alpha}z^{-h/k}. \quad (3.2)$$

The corresponding 2d conformal field theory (CFT) is the $\widehat{\mathfrak{sl}}_2$ WZW model generated by the EM tensor

$$T(z) = \frac{1}{2}(\partial\widetilde{\phi}(z))^2 + (\text{zero-modes term}) + T_{PF}(z).$$

Here $T_{PF}(z)$ denotes the EM tensor of the \mathbb{Z}_k -parafermions. The central charge of the models is $c_{WZW} = \frac{3k}{k+2}$.

Then the FF construction is the following procedure. Deform $T(z)$ by adding the background charge term to $T(z)$

$$T(z) \mapsto T_{FF}(z) = T(z) + \sqrt{\frac{k}{2r(r-k)}} \partial^2 \widetilde{\phi}(z).$$

This makes $T_{FF}(z)$ the generating function of the coset Virasoro algebra $(\widehat{\mathfrak{sl}}_2)_{r-2-k} \oplus (\widehat{\mathfrak{sl}}_2)_k \supset (\widehat{\mathfrak{sl}}_2)_{r-2}$ whose central charge is $c_{Vir} = \frac{3k}{k+2} \left(1 - \frac{2(k+2)}{r(r-k)}\right)$. At the same time, deform the currents of $\widehat{\mathfrak{sl}}_2$ by scaling the coefficients of the free boson and by adding the dynamical parameter P and

its conjugate Q satisfying $[P, Q] = 2$ to the Z -algebra part :

$$e(z) \mapsto S^+(z) = Z^+(z)e^{-Q}z^{-\frac{P-1}{r-k}} : \exp \left\{ \sqrt{\frac{2kr}{r-k}} \sum_{m \neq 0} \frac{a_m}{km} z^{-m} \right\} :, \quad (3.3)$$

$$f(z) \mapsto S^-(z) = Z^-(z)z^{\frac{P+h-1}{r}} : \exp \left\{ -\sqrt{\frac{2k(r-k)}{r}} \sum_{m \neq 0} \frac{a_m}{km} z^{-m} \right\} :. \quad (3.4)$$

It then turns out $S^\pm(z)$ become the screening currents of the Virasoro algebra generated by $T_{FF}(z)$.

Now let us consider a q -deformation of these constructions. It is known that the quantum affine algebra $U_q(\mathfrak{g})$ has an analogue of the LW realization. See for example [17]. In our case, the level- k currents of $U_q(\widehat{\mathfrak{sl}}_2)$ are realized in terms of the Heisenberg generators \tilde{a}_m and the level- k q -deformed Z operators $Z_q^\pm(z)$. The former satisfy

$$[\tilde{a}_m, \tilde{a}_n] = \frac{[2m]_q [km]_q}{m} q^{-k|m|} \delta_{m+n,0}. \quad (3.5)$$

This is a q -deformation of (3.1), i.e. in the limit $q \rightarrow 1$ one recovers (3.1). The operators $Z_q^\pm(z)$ are realized in the same way as (3.2) by replacing $\Psi(z), \Psi^\dagger(z)$ with appropriate q -parafermions [17, 38]. The level- k currents of $U_q(\widehat{\mathfrak{sl}}_2)$ are hence realized as

$$e_q(z) = Z_q^+(z) : \exp \left\{ \sum_{m \neq 0} \frac{\tilde{a}_m}{[km]_q} z^{-m} \right\} :, \quad f_q(z) = Z_q^-(z) : \exp \left\{ -\sum_{m \neq 0} \frac{\tilde{a}_m}{[km]_q} z^{-m} \right\} :.$$

Then a q -analogue of the FF construction (3.3)-(3.4) is obtained by noting the following observations.

- In (3.3) and (3.4), the Z -algebra part remains the same as in $e(z)$ and $f(z)$ except for getting a modification by the dynamical parameters
- The commutation relation of the Heisenberg generators a_m with the scaled coefficients can be read as from $S^+(z)$,

$$[\sqrt{\frac{2kr}{r-k}} a_m, \sqrt{\frac{2kr}{r-k}} a_n] = 2mk \frac{r}{r-k} \delta_{m+n,0}, \quad (3.6)$$

and from $S^-(z)$,

$$[\sqrt{\frac{2k(r-k)}{r}} a_m, \sqrt{\frac{2k(r-k)}{r}} a_n] = 2mk \frac{r-k}{r} \delta_{m+n,0}. \quad (3.7)$$

The first point indicates that the q - Z -algebra operators $Z_q^\pm(z)$ in $e_q(z)$ and $f_q(z)$ should remain the same except for getting a modification by the dynamical parameters P and Q in a q -deformed

FF process. The second point suggests to use the Heisenberg generators α_m or α'_m satisfying the following p -deformed commutation relation from (3.5).

$$[\alpha_m, \alpha_n] = \frac{[2m]_q [km]_q}{m} \frac{1 - p^m}{1 - p^{*m}} q^{-km} \delta_{m+n,0}, \quad (3.8)$$

$$[\alpha'_m, \alpha'_n] = \frac{[2m]_q [km]_q}{m} \frac{1 - p^{*m}}{1 - p^m} q^{km} \delta_{m+n,0}. \quad (3.9)$$

Here we set $p = q^{2r}$, $p^* = pq^{-2k} = q^{2(r-k)}$, and α_m, α'_m are not independent

$$\alpha'_m = \frac{1 - p^{*m}}{1 - p^m} q^{km} \alpha_m.$$

These are q -deformations of (3.6) and (3.7). Note that (3.8) is nothing but the relations for the Heisenberg subalgebra of $U_{q,p}(\widehat{\mathfrak{sl}}_2)$ at level k given in Sec. 2.1. One hence obtains the following q -analogue of the FF construction.

$$\begin{aligned} e_q(z) &\mapsto S_q^+(z) = Z_q^+(z) e^{-Q} z^{-\frac{P-1}{r-k}} : \exp \left\{ \sum_{m \neq 0} \frac{\alpha_m}{[km]_q} z^{-m} \right\} :, \\ f_q(z) &\mapsto S_q^-(z) = Z_q^-(z) z^{\frac{P+h-1}{r}} : \exp \left\{ - \sum_{m \neq 0} \frac{\alpha'_m}{[km]_q} z^{-m} \right\} : . \end{aligned}$$

Then the currents $S_q^+(z)$ and $S_q^-(z)$ turn out to give a realization of the level- k elliptic currents $e(z)$ and $f(z)$ of $U_{q,p}(\widehat{\mathfrak{sl}}_2)$, respectively [17, 38]. Hence regarding $e(z)$ and $f(z)$ as screening currents, one can define a deformation of the W -algebra $\mathcal{W}\mathfrak{sl}_2$ of the coset type at level k as the kernel of the screening operators associated to either $e(z)$ or $f(z)$.

In general, in the FF process the Z -algebra structure of \mathfrak{g} remains the same as in the LW realization to realize the screening currents of $\mathcal{W}\bar{\mathfrak{g}}$ of the coset type $(\mathfrak{g})_{r-h^\vee-k} \oplus (\mathfrak{g})_k \supset (\mathfrak{g}_{\text{diag}})_{r-h^\vee}$ except for a modification by adding the dynamical parameters. Correspondingly, the q - Z -algebra structure of $U_q(\mathfrak{g})$ remains the same to realize the elliptic currents $e_i(z), f_i(z)$ of $U_{q,p}(\mathfrak{g})$ as a p -deformation of $U_q(\mathfrak{g})$ except for a modification by the dynamical parameters [17, 38]. Then by using $e_i(z)$ and $f_i(z)$ as screening currents, one defines a deformation of $\mathcal{W}\bar{\mathfrak{g}}$ as the intersection of the kernel of the screening operators associated with either $e_i(z)$'s or $f_i(z)$'s. We denote the

resultant deformed W -algebra by $\mathcal{W}_{p,p^*}\bar{\mathfrak{g}}$.

$$\begin{array}{ccc}
 & U_q(\mathfrak{g}) & \\
 q\text{-deform. } \nearrow & & \searrow p = q^{2r}\text{-deform. + dynamical param.} \\
 \mathfrak{g} & & U_{q,p}(\mathfrak{g}) \\
 r\text{-deform. + dynamical param. } \searrow & & \nearrow q\text{-deform.} \\
 \text{(the Feigin-Fuchs construction)} & & \\
 & \mathcal{W}_{\bar{\mathfrak{g}}} & \\
 & \parallel & \\
 & (\mathfrak{g})_{r-h^\vee-k} \oplus (\mathfrak{g})_k \supset (\mathfrak{g})_{r-h^\vee} &
 \end{array}$$

Figure 3 : Two characterizations of $U_{q,p}(\mathfrak{g})$

Note that here we have obtained two elliptic nomes p and p^* naturally in a q -deformation of the FF construction. The parameters p, p^* correspond to q, t in Frenkel-Reshetikhin's formulation of the deformed W -algebra $W_{q,t}(\bar{\mathfrak{g}})$ [24]³. It is remarkable that the appearance of two elliptic nomes is also consistent to the quasi-Hopf formulation of the face type EQG $\mathcal{B}_{q,\lambda}(\mathfrak{g})$ [29], where p is treated as a dynamical parameter and p^* is nothing but a dynamical shift of p by q^{-2c} .

The generating function of $\mathcal{W}_{p,p^*}\bar{\mathfrak{g}}$ can also be constructed explicitly in terms of the vertex operators of $U_{q,p}(\mathfrak{g})$ w.r.t. the Drinfeld comultiplication. The same construction is valid also in the elliptic quantum toroidal algebras, which give a realization of the deformed affine quiver W -algebras. See Sec.11 for the case $\mathfrak{gl}_{1,tor}$.

Moreover in Sec.5, we discuss the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ derived as intertwining operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ -modules w.r.t. the standard comultiplication. Constructing correlation functions of them, one finds that they play the role of the vertex operators of the corresponding deformed W -algebra $\mathcal{W}_{p,p^*}\widehat{\mathfrak{sl}}_N$. See Sec.7 and 8.

4 Representations

A representation of H -algebra is called a dynamical representation. We skip all formal descriptions of it and expose only the level-0 and the level-1 dynamical representations of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$.

³Frenkel-Reshetikhin's deformed W -algebras are the Hamiltonian reduction type, which are isomorphic to the coset type only for the simply laced $\bar{\mathfrak{g}}$. For example, for the case B_l the coset construction gives Fateev-Lukyanov's WB_l algebra [54], which is different from $W(B_l)$ obtained by the quantum Hamiltonian reduction of $B_l^{(1)}$. See for example [9].

Here one says that a $U_{q,p}(\mathfrak{g})$ -module has level k , if $q^{c/2}$ acts as the scalar $q^{k/2}$ on it. For detailed description of the dynamical representations, the reader can consult [16, 35, 41].

4.1 The evaluation representation

Let $\mathcal{V} = \bigoplus_{i=1}^N \mathbb{F}v_i \otimes 1$ and set $\mathcal{V}_z = \mathcal{V}[z, z^{-1}]$. The following gives the level-0 dynamical action of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on \mathcal{V}_z .

$$\begin{aligned} \pi_z(q^{c/2}) &= 1, \quad \pi_z(d) = -z \frac{d}{dz}, \\ \pi_z(\alpha_{j,m}) &= \frac{[m]_q}{m} (q^{j-N+1}z)^m (q^{-m}E_{j,j} - q^mE_{j+1,j+1}), \\ \pi_z(e_j(w)) &= \frac{(pq^2; p)_\infty}{(p; p)_\infty} E_{j,j+1} \delta(q^{j-N+1}z/w) e^{-Q_{\alpha_j}}, \\ \pi_z(f_j(w)) &= \frac{(pq^{-2}; p)_\infty}{(p; p)_\infty} E_{j+1,j} \delta(q^{j-N+1}z/w), \\ \pi_z(\phi_j^+(w)) &= q^{-\pi(h_j)} e^{-Q_{\alpha_j}} \frac{\theta_p(q^{-j+N-1+2\pi(h_j)\frac{w}{z}})}{\theta_p(q^{-j+N-1}w/z)}, \\ \pi_z(\phi_j^-(w)) &= q^{\pi(h_j)} e^{-Q_{\alpha_j}} \frac{\theta_p(q^{j-N+1-2\pi(h_j)\frac{z}{w}})}{\theta_p(q^{j-N+1}z/w)}, \quad j = 1, \dots, N-1. \end{aligned}$$

Here $\pi(h_j) = E_{j,j} - E_{j+1,j+1}$, and the element $e^Q \in \mathbb{C}[\mathcal{R}_Q]$ acts on $f(h, P)v \otimes 1$ by $e^Q \cdot (f(h, P)v \otimes 1) = f(h, P - \langle P, Q \rangle)v \otimes e^Q$. This is called the evaluation representation associated with the vector representation with the evaluation parameter z . In particular, one has

$$\pi_z(L_{i,j}^+(w))_{k,l} = R^+(w/z, \Pi^*)_{ik}^{jl}.$$

4.2 The level-1 highest weight representation

Let Λ_0 and $\Lambda_a = \Lambda_0 + \bar{\Lambda}_a$ ($a = 1, \dots, N-1$) be the fundamental weights of $\widehat{\mathfrak{sl}}_N$. For generic $\nu \in \mathfrak{h}^*$, set

$$\mathcal{V}(\Lambda_a + \nu, \nu) = \mathbb{F} \otimes_{\mathbb{C}} (\mathcal{F}_\alpha \otimes e^{\Lambda_a} \mathbb{C}[\mathcal{Q}]) \otimes e^{Q_\nu} \mathbb{C}[\mathcal{R}_Q],$$

where $\mathcal{F}_\alpha = \mathbb{C}[\alpha_{j,-m} \ (j = 1, \dots, N-1, m \in \mathbb{N}_{>0})]$ denotes the Fock module of the Heisenberg algebra. We here consider the group algebra $\mathbb{C}[\mathcal{Q}] \otimes \mathbb{C}[\mathcal{R}_Q]$ with the following central extension.

$$\begin{aligned} e^{\alpha_j} e^{\alpha_k} &= (-1)^{(\alpha_j, \alpha_k)} q^{\frac{1}{r}(\delta_{j,k+1} - \delta_{k,j-1})} e^{\alpha_k} e^{\alpha_j}, \\ e^{Q_{\alpha_j}} e^{Q_{\alpha_k}} &= q^{(\frac{1}{r} - \frac{1}{r^*})(\delta_{j,k+1} - \delta_{j,k-1})} e^{Q_{\alpha_k}} e^{Q_{\alpha_j}}, \\ e^{Q_{\alpha_j}} e^{\alpha_k} &= q^{\frac{1}{r}(\delta_{j,k+1} - \delta_{k,j-1})} e^{\alpha_k} e^{Q_{\alpha_j}}. \end{aligned}$$

Then the space $\mathcal{V}(\Lambda_a + \nu, \nu)$ is the level-1 irreducible highest weight $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ -module with the highest weight $(\Lambda_a + \nu, \nu)$ w.r.t. the set $(P + h, P)$ by the action

$$\begin{aligned} E_j(z) &= : \exp \left\{ - \sum_{n \neq 0} \frac{1}{[n]_q} \alpha_{j,n} z^{-n} \right\} : e^{\alpha_j} e^{-Q_{\alpha_j}} z^{h_{\alpha_j} + 1} (q^{N-j} z)^{-\frac{P_{\alpha_j} - 1}{r^*}}, \\ F_j(z) &= : \exp \left\{ \sum_{n \neq 0} \frac{1}{[n]_q} \alpha'_{j,n} z^{-n} \right\} : e^{-\alpha_j} z^{-h_{\alpha_j} + 1} (q^{N-j} z)^{\frac{(P+h)\alpha_j - 1}{r}}, \\ \phi_j^+(q^{-1/2} z) &= q^{-h_j} e^{-Q_{\alpha_j}} : \exp \left\{ (q - q^{-1}) \sum_{m \neq 0} \frac{\alpha_{j,m}}{1 - p^m} p^m z^{-m} \right\} :, \\ \phi_j^-(q^{1/2} z) &= q^{h_j} e^{-Q_{\alpha_j}} : \exp \left\{ (q - q^{-1}) \sum_{m \neq 0} \frac{\alpha_{j,m}}{1 - p^m} z^{-m} \right\} : \quad (1 \leq j \leq N - 1). \end{aligned}$$

The highest weight vector is given by $1 \otimes e^{\Lambda_a} \otimes e^{Q_\nu}$.

Note that one has the following natural decomposition.

$$\mathcal{V}(\Lambda_a + \nu, \nu) = \bigoplus_{\xi, \eta \in \mathcal{Q}} \mathcal{F}_{a,\nu}(\xi, \eta), \quad \mathcal{F}_{a,\nu}(\xi, \eta) = \mathbb{F} \otimes_{\mathbb{C}} (\mathcal{F}_\alpha \otimes e^{\Lambda_a + \xi}) \otimes e^{Q_{\nu + \eta}}.$$

It turns out that $\mathcal{F}_{a,\nu}(\xi, \eta)$ is isomorphic to the Verma module of the W -algebra $\mathcal{W}\mathfrak{sl}_N$ with the central charge $(N - 1) \left(1 - \frac{N(N+1)}{r(r-1)} \right)$ and the highest weight $\frac{1}{r(r-1)} |r(\nu + \eta + \rho) - (r - 1)(\Lambda_a + \nu + \xi + \eta + \rho)|^2$ [17].

5 Vertex Operators

The vertex operators defined for the infinite dimensional quantum groups such as $U_q(\mathfrak{g})$, $U_{q,p}(\mathfrak{g})$ and $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ have proven to be key objects in many applications [10, 23, 30, 37, 47]. Here we expose the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ [28, 36], which are used in a derivation of the elliptic weight functions (Sec.6), integral solutions of the elliptic q -KZ equation and the vertex function (Sec.7) and also in the algebraic analysis of the elliptic lattice models (Sec.8).

Let \mathcal{V}_z be as in Sec.4.1 and $\mathcal{V}(\Lambda_a + \nu, \nu)$ denote the irreducible level-1 highest weight $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ -module with the highest weight $(\Lambda_a + \nu, \nu)$ in Sec.4.2. Let $\Lambda_{a+N} = \Lambda_a$. The (type I) vertex operator $\Phi(z)$ is the intertwiner of the $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ -modules

$$\Phi(z) : \mathcal{V}(\Lambda_a + \nu, \nu) \rightarrow \mathcal{V}_z \otimes \mathcal{V}(\Lambda_{a-1} + \nu, \nu)$$

satisfying the intertwining relations

$$\Delta(x)\Phi(z) = \Phi(z)x \quad \forall x \in U_{q,p}(\widehat{\mathfrak{sl}}_N). \quad (5.1)$$

The components of the vertex operator are defined by

$$\Phi(zq^{-1})u = \sum_{\mu=1}^N v_{\mu} \tilde{\otimes} \Phi_{\mu}(z) u, \quad \forall u \in \mathcal{V}(\lambda, \nu).$$

One can solve the linear equation (5.1) by using the representations in the last section uniquely up to an overall constant factor. The result is summarized as follows.

$$\begin{aligned} \Phi_{\mu}(z) &= a_{\mu,N} \oint_{\mathbb{T}^{N-\mu}} \prod_{m=\mu}^{N-1} \frac{dt_m}{2\pi i t_m} \Phi_N(z) F_{N-1}(t_{N-1}) F_{N-2}(t_{N-2}) \cdots F_{\mu}(t_{\mu}) \varphi_{\mu}(z, t_{\mu}, \dots, t_{N-1}; \Pi), \\ \Phi_N(z) &= : \exp \left\{ \sum_{m \neq 0} (q^m - q^{-m}) \mathcal{E}'_m z^{-m} \right\} : e^{-\bar{\epsilon}_N (-z)^{-h_{\epsilon_N}} z^{\frac{1}{r}(P+h)_{\epsilon_N}}}, \end{aligned} \quad (5.2)$$

where we set $\bar{\epsilon}_i = \epsilon_i - \sum_{j=1}^N \epsilon_j / N$, $\Pi = \{\Pi_{\mu,m} \ (m = \mu + 1, \dots, N)\}$,

$$\mathbb{T}^{N-\mu} = \{t \in \mathbb{C}^{N-\mu} \mid |t_{\mu}| = \cdots = |t_{N-1}| = 1\},$$

and

$$\varphi_{\mu}(z, t_{\mu}, \dots, t_{N-1}; \Pi) = \prod_{m=\mu}^{N-1} \frac{[v_{m+1} - v_m + (P+h)_{\mu,m+1} - \frac{1}{2}][1]}{[v_{m+1} - v_m + \frac{1}{2}][(P+h)_{\mu,m+1}]}$$

with $z = q^{2u}$, $t_m = q^{2v_m}$, $v_N = u$. We assume $|p| < |z| < 1$. The symbol \mathcal{E}'_m^j denotes a linear combination of $\alpha'_{j,m}$ obtained by solving

$$\alpha'_{j,m} = [m]_q^2 (q - q^{-1}) (\mathcal{E}'_m^j - q^{-m} \mathcal{E}'_{m+1}^j), \quad \sum_{j=1}^N q^{(j-1)m} \mathcal{E}'_m^j = 0.$$

The most important property of $\Phi_{\mu}(z)$ is the following commutation relation.

$$\Phi_{\mu_2}(z_2) \Phi_{\mu_1}(z_1) = \sum_{\mu'_1, \mu'_2=1}^N R(z_1/z_2, \Pi)_{\mu'_1 \mu'_2}^{\mu_1 \mu_2} \Phi_{\mu'_1}(z_1) \Phi_{\mu'_2}(z_2), \quad (5.3)$$

where

$$R(z, \Pi) = \mu(z) \bar{R}(z, \Pi), \quad \mu(z) = z^{-\frac{r-1}{r} \frac{N-1}{N}} \frac{\Gamma(pz; p, q^{2N}) \Gamma(q^{2N} z; p, q^{2N})}{\Gamma(q^2 z; p, q^{2N}) \Gamma(pq^{2N-2} z; p, q^{2N})}.$$

For example, this yields the transition property of the elliptic weight function (Sec.6) as well as the R -matrix coefficients in the elliptic q -KZ connection (Sec.7).

6 Elliptic Weight Functions

The weight functions are objects in the theory of (q -) hypergeometric integrals. They play the role of a basis of the (twisted) de Rham cohomology [4, 56, 59, 70]. The elliptic version of the weight function was first introduced for the \mathfrak{sl}_2 type in [70]⁴. We here expose the elliptic weight functions of the \mathfrak{sl}_N type, which can be derived by considering a composition of the vertex operators given in the last section. In the next section, we show that they play the role in elliptic hypergeometric integral solutions of the elliptic q -KZ equation. Moreover, the elliptic weight functions are identified with the elliptic stable envelopes [2] associated with Nakajima quiver variety X . This makes clear a connection between representation theory of EQG and geometry of the elliptic cohomology (Sec.10).

Let $\Phi_\mu(z)$ be the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ in the last section and consider their composition

$$\phi_{\mu_1 \cdots \mu_n}(z_1, \cdots, z_n) := \Phi_{\mu_1}(z_1) \cdots \Phi_{\mu_n}(z_n) : \mathcal{F}_{a,\nu}(\xi, \eta) \rightarrow \mathcal{F}_{a',\nu}(\xi, \eta).$$

Here $a' \equiv a - n \pmod{N}$. Let $[1, n] = \{1, \cdots, n\}$ and define the index sets $I_l := \{i \in [1, n] \mid \mu_i = l\}$ ($l = 1, \cdots, N$). Set also $\lambda_l := |I_l|$, $\lambda := (\lambda_1, \cdots, \lambda_N)$. Then $I = (I_1, \cdots, I_N)$ is a partition of $[1, n]$, i.e.

$$I_1 \cup \cdots \cup I_N = [1, n], \quad I_k \cap I_l = \emptyset \quad (k \neq l).$$

Thus obtained partition I is often denoted by I_{μ_1, \cdots, μ_n} . For $\lambda = (\lambda_1, \cdots, \lambda_N) \in \mathbb{N}^N$, let \mathcal{I}_λ be the set of all partitions $I = (I_1, \cdots, I_N)$ satisfying $|I_l| = \lambda_l$ ($l = 1, \cdots, N$). Set also $\lambda^{(l)} := \lambda_1 + \cdots + \lambda_l$, $I^{(l)} := I_1 \cup \cdots \cup I_l$ and let $I^{(l)} =: \{i_1^{(l)} < \cdots < i_{\lambda^{(l)}}^{(l)}\}$. Note that \mathcal{I}_λ specifies the coordinate flags of the partial flag variety $fl_\lambda : 0 = V_0 \subset V_1 \subset \cdots \subset V_{N-1} \subset V_N = \mathbb{C}^n$ with $\dim V_l = \lambda^{(l)}$.

Now let us substitute the realization of $\Phi_\mu(z)$ (5.2) to $\phi_{\mu_1 \cdots \mu_n}(z_1, \cdots, z_n)$. In each $\Phi_\mu(z)$, we assign the integration variable $t_a^{(l)}$ to the elliptic current $F_l(*)$ ($l = 1, \cdots, N-1$) as its argument if $\Phi_\mu(z)$ is the $i_a^{(l)}$ -th vertex operator i.e. $z = z_{i_a^{(l)}}$. After rearranging the order of the elliptic currents F_l 's and Φ_N 's and taking their normal ordering as specified in $\widetilde{\Phi}(t, z)$ in the below, one obtains the following expression for $I = I_{\mu_1 \cdots \mu_n}$,

$$\phi_{\mu_1 \cdots \mu_n}(z_1, \cdots, z_n) = \oint_{\mathbb{T}^M} \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \frac{dt_a^{(l)}}{2\pi i t_a^{(l)}} \widetilde{\Phi}(t, z) \widetilde{W}_I(t, z, \Pi), \quad |p| < |z_1|, \cdots, |z_n| < 1, \quad (6.1)$$

⁴Strictly speaking, the elliptic weight functions in [70] were used as a pole subtraction matrix, in the terminology of [2], in the q -hypergeometric integral solution of the q -KZ equation. They specifies the cycles of the integral. On the other hand, there the role of a basis of the co-cycle was played by the trigonometric weight functions.

$$\begin{aligned} \tilde{\Phi}(t, z) = & \Phi_N(z_1) \cdots \Phi_N(z_n) :: F_{N-1}(t_1^{(N-1)}) \cdots F_{N-1}(t_{\lambda^{(N-1)}}^{(N-1)}) : \cdots : F_1(t_1^{(1)}) \cdots F_1(t_{\lambda^{(1)}}^{(1)}) : \\ & \times \prod_{1 \leq k < l \leq n} \langle \Phi_N(z_k) \Phi_N(z_l) \rangle^{Sym} \prod_{l=1}^{N-1} \prod_{1 \leq a < b \leq \lambda^{(l)}} \langle F_l(t_a^{(l)}) F_l(t_b^{(l)}) \rangle^{Sym}, \end{aligned}$$

$$\tilde{W}_I(t, z, \Pi) = \text{Sym}_{t^{(1)}} \cdots \text{Sym}_{t^{(N-1)}} \tilde{U}_I(t, z, \Pi),$$

$$\begin{aligned} \tilde{U}_I(t, z, \Pi) = & \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \left(\frac{[v_b^{(l+1)} - v_a^{(l)} + (P+h)_{\mu_s, l+1} - C_{\mu_s, l+1}(s)][1]}{[v_b^{(l+1)} - v_a^{(l)} + 1][(P+h)_{\mu_s, l+1} - C_{\mu_s, l+1}(s)]} \Big|_{i_s^{(N)} = i_b^{(l+1)} = i_a^{(l)}} \right. \\ & \times \left. \prod_{\substack{b=1 \\ i_b^{(l+1)} > i_a^{(l)}}}^{\lambda^{(l+1)}} \frac{[v_b^{(l+1)} - v_a^{(l)}]}{[v_b^{(l+1)} - v_a^{(l)} + 1]} \prod_{b=a+1}^{\lambda^{(l)}} \frac{[v_a^{(l)} - v_b^{(l)} - 1]}{[v_a^{(l)} - v_b^{(l)}]} \right), \end{aligned}$$

where we set $t_a^{(l)} = q^{2v_a^{(l)}}$, $t_s^{(N)} = z_s = q^{2u_s}$, $M = \sum_{l=1}^{N-1} \lambda^{(l)}$ and $C_{\mu_s, l+1}(s) = \sum_{j=s+1}^n \langle \bar{\epsilon}_{\mu_j}, h_{\mu_s, l+1} \rangle$. The symbol $\langle \rangle^{Sym}$ denotes the symmetric part of the operator product expansion coefficient $\langle \rangle$ defined by for $\mathcal{O} = \Phi_N, F_l$,

$$\mathcal{O}(w_a) \mathcal{O}(w_b) = \langle \mathcal{O}(w_a) \mathcal{O}(w_b) \rangle : \mathcal{O}(w_a) \mathcal{O}(w_b) : .$$

Hence $\tilde{\Phi}(t, z)$ is an operator valued symmetric function in $z = (z_1, \dots, z_n)$ as well as in $t^{(l)} = (t_1^{(l)}, \dots, t_{\lambda^{(l)}}^{(l)})$ for each l . The function $\tilde{W}_I(t, z, \Pi)$ is the elliptic weight function of type \mathfrak{sl}_N . It is obtained by collecting all non-symmetric part of $\langle \rangle$'s and factors arisen from the exchange among Φ_N 's and F_l 's and by making symmetrizations $\text{Sym}_{t^{(l)}}$ of all entries in $t^{(l)}$. Hence $\tilde{W}_I(t, z, \Pi)$ is a symmetric function in $v_a^{(l)}$ ($a = 1, \dots, \lambda^{(l)}$) for each l .

The weight function $\tilde{W}_I(t, z, \Pi)$ has several nice properties such as

- The triangular property: For $I, J \in \mathcal{I}_\lambda$,

$$\begin{aligned} (1) \quad & \tilde{W}_J(z_I, z, \Pi) = 0 \text{ unless } I \leq J. \\ (2) \quad & \tilde{W}_I(z_I, z, \Pi) = \prod_{1 \leq k < l \leq N} \prod_{a \in I_k} \prod_{\substack{b \in I_l \\ a < b}} \frac{[u_b - u_a]}{[u_b - u_a + 1]} \end{aligned}$$

Here \leq denotes the partial ordering defined by

$$I \leq J \Leftrightarrow i_a^{(l)} \leq j_a^{(l)} \quad \forall l, a.$$

for $I^{(l)} = \{i_1^{(l)} < \dots < i_{\lambda^{(l)}}^{(l)}\}$ and $J^{(l)} = \{j_1^{(l)} < \dots < j_{\lambda^{(l)}}^{(l)}\}$ ($l = 1, \dots, N$). We also denote by $t = z_I$ the specialization $t_a^{(l)} = z_{i_a^{(l)}}$ ($l = 1, \dots, N-1, a = 1, \dots, \lambda^{(l)}$) [66].

- The transition property:

$$\begin{aligned} & \tilde{W}_{I \dots \mu_{i+1} \mu_i \dots}(t, \dots, z_{i+1}, z_i, \dots, \Pi) \\ & = \sum_{\mu'_i, \mu'_{i+1}} \bar{R}(z_i/z_{i+1}, \Pi q^{-2 \sum_{j=i}^n \langle \bar{\epsilon}_{\mu_j}, h \rangle})_{\mu_i \mu_{i+1}}^{\mu'_i \mu'_{i+1}} \tilde{W}_{I \dots \mu'_i \mu'_{i+1} \dots}(t, \dots, z_i, z_{i+1}, \dots, \Pi) \end{aligned}$$

as well as the orthogonality, quasi-periodicity, the shuffle product formula compatible to the wheel conditions etc. [42]. These properties are used to identify $\widetilde{W}_I(t, z, \Pi)$ with the elliptic stable envelope for the cotangent bundle to the partial flag variety (Sec.10).

7 Integral Solution of the Elliptic q -KZ equation

Let us consider the n -point function $\phi_{\mu_1 \dots \mu_n}(z_1, \dots, z_n)$ with the zero weight condition $\sum_{i=1}^n \bar{\epsilon}_{\mu_i} = 0$, which is equivalent to consider $\lambda = (m^N)$ with $mN = n$. In this case, one can take a trace over the space $\mathcal{F}_{a,\nu}(\xi, \eta)$. It then turns out that the trace gives a solution of the elliptic q -KZ equation due to the cyclic property of the trace and the commutation relation of the vertex operators (5.3) [22, 42]. From (6.1), one gets the following elliptic hypergeometric integral.

$$\begin{aligned} \mathrm{tr}_{\mathcal{F}_{a,\nu}} \left(Q^{-d} \Phi_{\mu_1}(z_1) \cdots \Phi_{\mu_N}(z_N) \right) &= \oint_{\mathbb{T}^M} \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \frac{dt_a^{(l)}}{2\pi i t_a^{(l)}} e(t, \Pi) \Phi(t, z) \widetilde{W}_I(t, z, \Pi), \\ e(t, \Pi) &= \exp \left\{ \frac{\sum_{l=1}^{N-1} \log(\Pi_l / \Pi_{l+1}) \sum_a \log(t_a^{(l)})}{\log p} \right\}, \\ \Phi(t, z) &= \prod_{l=1}^{N-1} \left[\prod_{a=1}^{\lambda^{(l)}} \prod_{b=1}^{\lambda^{(l+1)}} \frac{\Gamma(t_a^{(l)} / t_b^{(l+1)}; p, Q)}{\Gamma(p^* t_a^{(l)} / t_b^{(l+1)}; p, Q)} \prod_{1 \leq a < b \leq \lambda^{(l)}} \frac{\Gamma(p^* t_a^{(l)} / t_b^{(l)}, p^* t_b^{(l)} / t_a^{(l)}; p, Q)}{\Gamma(t_a^{(l)} / t_b^{(l)}, t_b^{(l)} / t_a^{(l)}; p, Q)} \right]. \end{aligned}$$

Note that only the weight function $\widetilde{W}_I(t, z, \Pi)$ carries the tensor indices through I . We expect that $\widetilde{W}_I(t, z, \Pi), I \in \mathcal{I}_{(m^N)}$ form a basis of the twisted de Rham cohomology associated with $T_{Q, z_i} \Phi(t, z) / \Phi(t, z), i = 1, \dots, n$, where T_{Q, z_i} is the shift operator $T_{Q, z_i} f(\dots, z_i, \dots) = f(\dots, Qz_i, \dots)$. Furthermore, to specify the cycle of the integral one may insert to the integral an additional elliptic weight function $\widetilde{W}_J(t, z, \Pi), J \in \mathcal{I}_{(m^N)}$, whose elliptic nome is Q instead of p . Thus obtained elliptic hypergeometric integrals labeled by $I, J \in \mathcal{I}_{(m^N)}$ are conjectured to give fundamental solutions of the elliptic q -KZ equation. See [70] for the trigonometric \mathfrak{sl}_2 case.

In the trigonometric limit $Q \rightarrow 0$, the trace goes to the following vacuum expectation value.

$$\begin{aligned} \langle 0 | \Phi_{\mu_1}(z_1) \cdots \Phi_{\mu_N}(z_N) | 0 \rangle &= \oint_{\mathbb{T}^M} \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \frac{dt_a^{(l)}}{2\pi i t_a^{(l)}} e(t, \Pi) \Phi^{trig.}(t, z) \widetilde{W}_I(t, z, \Pi), \\ \Phi^{trig.}(t, z) &= \prod_{l=1}^{N-1} \left[\prod_{a=1}^{\lambda^{(l)}} \prod_{b=1}^{\lambda^{(l+1)}} \frac{(p^* t_a^{(l)} / t_b^{(l+1)}; p)_\infty}{(t_a^{(l)} / t_b^{(l+1)}; p)_\infty} \prod_{1 \leq a < b \leq \lambda^{(l)}} \frac{(t_a^{(l)} / t_b^{(l)}, t_b^{(l)} / t_a^{(l)}; p)_\infty}{(p^* t_a^{(l)} / t_b^{(l)}, p^* t_b^{(l)} / t_a^{(l)}; p)_\infty} \right]. \end{aligned}$$

This gives the integral formula for the vertex function for the equivariant K-theory $K_T(T^* fl_\lambda)$ with $T = (\mathbb{C}^\times)^n \times \mathbb{C}^\times$, which is the generating function of counting the quasi-maps from \mathbb{P}^1 to $T^* fl_\lambda$ [49]. Here the elliptic weight function \widetilde{W}_I plays the role of the pole subtraction matrix [2]. In particular, for the full flag variety case i.e. $\lambda^{(l)} = l$ ($l = 1, \dots, N-1$) with $n = N$, it gives

the integral formula for the Macdonald symmetric function [59], which is also known as the 3d vortex partition function of the $T[U(N)]$ theory, if one drops the weight function \widetilde{W}_I from the integrand. Indeed the first and the second blocks in $\Phi(t, z)$ are elliptic analogues of the Cauchy type reproducing kernel and the Macdonald weight function, respectively, by reading (p, p^*) as (q, t) in the Macdonald theory. Hence the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ play the role of the vertex operators of the deformed W -algebra $\mathcal{W}_{p,p^* \mathfrak{sl}_N}$ [3, 25].

8 Algebraic Analysis of the Elliptic Lattice Models

The algebraic analysis [30] is a scheme of mathematical formulation of the solvable 2d lattice models defined by the R -matrices. It formulates the model on the infinite lattice directly by using both finite and infinite dimensional representations of the corresponding affine quantum groups $U_q(\mathfrak{g})$ or elliptic quantum groups $U_{q,p}(\mathfrak{g})$. In particular, the infinite dimensional representations are identified with the spaces of states of the model specified by the ground state boundary condition, where the configurations at enough far sites from the center of the lattice is given by the ground state configurations. In addition, the two types intertwining operators, the type I and II vertex operators, of the quantum groups realize the local operators of the model such as spin operators acting on the infinite dimensional representations (type I) and the creation operators of the physical excitations (type II), respectively. Then the traces of compositions of the vertex operators give correlation functions as well as form factors of the models. Hence these are characterized as solutions of the (elliptic) q -KZ equations [22, 23, 30] as discussed in the last section for the type I case. In this sense, the algebraic analysis is an off-critical extension of CFT [8].

There are series of elliptic solvable lattice models defined by the elliptic solutions of the face type YBE, i.e. the elliptic dynamical R -matrices, associated with affine Lie algebras [5, 13, 32, 50, 51]. The critical behavior of such models is described by the W -algebra $\mathcal{W}_{\bar{\mathfrak{g}}}$ of the coset type $(\mathfrak{g})_{r-h^\vee-k} \oplus (\mathfrak{g})_k \supset (\mathfrak{g}_{\text{diag.}})_{r-h^\vee}$ [13, 14]. Here r is taken as a real positive parameter and called the restriction height. Accordingly the elliptic quantum group $U_{q,p}(\mathfrak{g})$ plays a central role in the algebraic analysis of these models as a deformation of the W -algebra $\mathcal{W}_{p,p^* \bar{\mathfrak{g}}}$. See Sec.3. The spaces of states with the ground state boundary condition are identified with the irreducible $\mathcal{W}_{p,p^* \bar{\mathfrak{g}}}$ -modules, whose direct sum are isomorphic to the $U_{q,p}(\mathfrak{g})$ -modules (Sec.4.2). The type I and II vertex operators of $U_{q,p}(\mathfrak{g})$ provide correlation functions and form factors of the model. See [22, 37, 55].

In the case $\mathfrak{g} = \widehat{\mathfrak{sl}}_N$, one also has the vertex type elliptic lattice models [6, 7]. These are

related to the face models by the so-called vertex-face correspondence [6, 31]. The algebraic analysis of the vertex models is carried out based on the one of the face models by applying the vertex-face correspondence [10, 37, 52].

9 Level-0 Action on the Gelfand-Tsetlin Basis

Let (π_z, \mathcal{V}_z) be the evaluation representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ in Sec.4.1. The action of the L -operator $L^+(1/w)$ on $\mathcal{V}_w \otimes \mathcal{V}_z$ is extended to the tensor space $\mathcal{V}_w \otimes \mathcal{V}_{z_1} \otimes \cdots \otimes \mathcal{V}_{z_n}$ by the comultiplication

$$\begin{aligned} & (\pi_{z_1} \otimes \cdots \otimes \pi_{z_n}) \Delta'^{(n-1)}(L^\pm(1/w)) \\ &= \bar{R}^{(0n)}(z_n/w, \Pi^* q^{2\sum_{j=1}^{n-1} h^{(j)}}) \bar{R}^{(0n-1)}(z_{n-1}/w, \Pi^* q^{2\sum_{j=1}^{n-2} h^{(j)}}) \cdots \bar{R}^{(01)}(z_1/w, \Pi^*). \end{aligned}$$

Here we take the opposite comultiplication Δ' following [43]. This is the action on the standard basis

$$v_I := v_{\mu_1} \otimes \cdots \otimes v_{\mu_n}, \quad I = I_{\mu_1 \cdots \mu_n} \in \mathcal{I}_\lambda, \quad \lambda \in \mathbb{N}^N, \quad |\lambda| = n.$$

Let us consider the following change of basis [43].

$$\xi_I = \sum_{J \in \mathcal{I}_\lambda} \widetilde{W}_J(z_I^{-1}, z^{-1}, \Pi q^{2\sum_{j=1}^n \langle \epsilon_{\mu_j}, h \rangle}) v_J. \quad (9.1)$$

The transition matrix is given by the specialized elliptic weight function so that it is lower triangular (Sec.6). It turns out that the new vectors ξ_I ($I \in \mathcal{I}_\lambda, \lambda \in \mathbb{N}^N, |\lambda| = n$) form the Gelfand-Tsetlin (GT) basis of $\mathcal{V}_{z_1} \otimes \cdots \otimes \mathcal{V}_{z_n}$, on which the commuting subalgebra generated by $\alpha_{j,m}$ ($j = 1, \dots, N-1, m \in \mathbb{Z} \setminus \{0\}$) is diagonalized simultaneously. In fact the level-0 action of the elliptic currents of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on ξ_I is given by

$$\begin{aligned} \phi_j^\pm(w) \xi_I &= \prod_{a \in I_j} \frac{[u_a - v + 1]}{[u_a - v]} \Big|_{\pm} \prod_{b \in I_{j+1}} \frac{[u_b - v - 1]}{[u_b - v]} \Big|_{\pm} e^{-Q_{\alpha_j}} \xi_I \\ E_j(w) \xi_I &= a^* \sum_{i \in I_{j+1}} \delta(z_i/w) \prod_{\substack{k \in I_{j+1} \\ \neq i}} \frac{[u_i - u_k + 1]}{[u_i - u_k]} e^{-Q_{\alpha_j}} \xi_{I^i} \\ F_j(w) \xi_I &= a \sum_{i \in I_j} \delta(z_i/w) \prod_{\substack{k \in I_j \\ \neq i}} \frac{[u_k - u_i + 1]}{[u_k - u_i]} \xi_{I^i} \end{aligned}$$

where $aa^* = -\frac{1}{q-q^{-1}} \frac{[0]'}{[1]}$, and $I = (I_1, \dots, I_N)$, $I^{i'} \in \mathcal{I}_{(\lambda_1, \dots, \lambda_j+1, \lambda_{j+1}-1, \dots, \lambda_N)}$, $I^i \in \mathcal{I}_{(\lambda_1, \dots, \lambda_j-1, \lambda_{j+1}+1, \dots, \lambda_N)}$ with

$$\begin{aligned} (I^{i'})_j &= I_j \cup \{i\}, & (I^{i'})_{j+1} &= I_{j+1} - \{i\}, & (I^{i'})_k &= I_k \quad (k \neq j, j+1), \\ (I^i)_j &= I_j - \{i\}, & (I^i)_{j+1} &= I_{j+1} \cup \{i\}, & (I^i)_k &= I_k \quad (k \neq j, j+1). \end{aligned}$$

Note that this action is given in terms of the partitions in completely a combinatorial way. In the next section, this is identified with a geometric action of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on the equivariant elliptic cohomology $\bigoplus_{\lambda \in \mathbb{N}^N, |\lambda|=n} E_T(T^*fl_\lambda)$ under the identification of the elliptic weight function with the elliptic stable envelope.

10 Geometric Interpretation

The notion of stable envelopes was initiated in [57, 65]. They form a good basis of equivariant cohomology and K-theory of Nakajima quiver varieties [61, 62] and has nice applications to enumerative geometry, geometric representation theory and quantum integrable systems. In particular, the transition matrices between stable envelopes defined for different chambers give the R -matrices satisfying the YBE so that they provide a new geometric formulation of quantum groups as well as quantum integrable systems associated with the quiver varieties. The elliptic version of stable envelopes was defined in [2] for the equivariant elliptic cohomology $E_T(X)$ of the Nakajima quiver varieties X . Remarkably they provide the elliptic dynamical R -matrix as their transition matrix, which coincides for example with (2.4) for the case $X = T^*fl_\lambda$. We here expose an identification of the elliptic stable envelopes for $E_T(T^*fl_\lambda)$ with the elliptic weight functions for $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ described in Sec.6. This makes more transparent the correspondence between geometry of quiver variety and representation of EQG. For the definition of the elliptic cohomology and the elliptic stable envelopes, the reader can consult [2].

The elliptic stable envelope $\text{Stab}_{\mathcal{E}}(F_I)$ for $E_T(T^*fl_\lambda)$ at the torus fixed point F_I labeled by the partition $I \in \mathcal{I}_\lambda$ can be constructed explicitly by the abelianization procedure [2, 43]. On the other hand, let us consider the elliptic weight function modified as

$$\mathcal{W}_I(t, z, \Pi) = \frac{H_\lambda(t, z) \widetilde{W}_I(t, z, \Pi)}{E_\lambda(t)},$$

$$H_\lambda(t, z) = \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \prod_{b=1}^{\lambda^{(l+1)}} \left[v_b^{(l+1)} - v_a^{(l)} + 1 \right], \quad E_\lambda(t) = \prod_{l=1}^{N-1} \prod_{a=1}^{\lambda^{(l)}} \prod_{b=1}^{\lambda^{(l)}} [v_b^{(l)} - v_a^{(l)} + 1].$$

This yields the expression

$$\begin{aligned}
\mathcal{W}_I(t, z, \Pi) &= \text{Sym}_{t^{(1)}} \cdots \text{Sym}_{t^{(N-1)}} U_I(t, z, \Pi), \\
U_I(t, z, \Pi) &= \prod_{l=1}^{N-1} \frac{\prod_{a=1}^{\lambda^{(l)}} u_I^{(l)}(t_a^{(l)}, t^{(l+1)}, \Pi_{\mu_{i_a^{(l)}, l+1} q}^{-2C_{\mu_{i_a^{(l)}, l+1}(i_a^{(l)})}})}{\prod_{1 \leq a < b \leq \lambda^{(l)}} [v_a^{(l)} - v_b^{(l)}][v_b^{(l)} - v_a^{(l)} - 1]}, \\
u_I^{(l)}(t_a^{(l)}, t^{(l+1)}, \Pi_{j,k}) &= \frac{\left[v_b^{(l+1)} - v_a^{(l)} + (P+h)_{j,k} \right]}{[(P+h)_{j,k}]} \Bigg|_{i_b^{(l+1)}=i_a^{(l)}} \\
&\quad \times \prod_{\substack{b=1 \\ i_b^{(l+1)} > i_a^{(l)}}}^{\lambda^{(l+1)}} \left[v_b^{(l+1)} - v_a^{(l)} \right] \prod_{\substack{b=1 \\ i_b^{(l+1)} < i_a^{(l)}}}^{\lambda^{(l+1)}} \left[v_b^{(l+1)} - v_a^{(l)} + 1 \right].
\end{aligned}$$

Comparing these expressions, one finds $\text{Stab}_{\mathfrak{C}}(F_I)$ and $\mathcal{W}_I(z, t, \Pi)$ are identical as

$$\text{Stab}_{\mathfrak{C}}(F_I) = \mathcal{W}_{\sigma_0(I)}(\tilde{t}, \sigma_0(z^{-1}), \Pi^{-1}). \quad (10.1)$$

Here for the longest element σ_0 in \mathfrak{S}_n and $I = I_{\mu_1 \cdots \mu_n}$, we define $\sigma_0(I) = I_{\mu_{\sigma_0(1)} \cdots \mu_{\sigma_0(n)}}$ and set $\sigma_0(I) = (\tilde{I}_1, \dots, \tilde{I}_N)$, $\tilde{I}^{(l)} = \tilde{I}_1 \cup \cdots \cup \tilde{I}_l$ and $\tilde{I}^{(l)} = \{\tilde{i}_1^{(l)} < \cdots < \tilde{i}_{\lambda^{(l)}}^{(l)}\}$. Then \tilde{t} denotes the set of all $t_a^{(l)}$ corresponding to $\tilde{i}_a^{(l)}$. The chamber \mathfrak{C} is taken as $|z_1| < \cdots < |z_n|$. In addition, the restriction $t = z_J$ defined in Sec.6 can be identified with the restriction of the stable class $\text{Stab}_{\mathfrak{C}}(F_I)$ to the fixed point F_J . One then finds

$$\text{Stab}_{\mathfrak{C}}(F_I)|_{F_J} = \mathcal{W}_{\sigma_0(I)}(z_J^{-1}, \sigma_0(z^{-1}), \Pi^{-1}). \quad (10.2)$$

Such coincidence as well as the integral formula for the vertex function obtained as correlation function of the vertex operators in Sec.7 lead us to the following dictionary from the level-0 representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ to the equivariant elliptic cohomology $\text{E}_T(T^*fI_\lambda)$.

evaluation parameters z_1, \dots, z_n	\leftrightarrow	equivariant parameters $\in (\mathbb{C}^\times)^n$
dynamical parameters Π_j	\leftrightarrow	Kähler parameters $\in \text{Pic}_T(X) \otimes_{\mathbb{Z}} E$
integration variables $t_a^{(l)}$ ($a = 1, \dots, \lambda^{(l)}$)	\leftrightarrow	Chern roots of the tautological vector bundle \mathcal{V}_l of rk $\lambda^{(l)}$ on the l -th vertex of the quiver diagram
the weight (μ_1, \dots, μ_n)	\leftrightarrow	the fixed point $F_{I_{\mu_1, \dots, \mu_n}}$

Furthermore, denoting the fixed point classes by $[I]$ one has the localization formula for the stable envelope.

$$\text{Stab}_{\mathfrak{C}}(F_K) = \sum_{I \in \mathcal{I}_\lambda} \frac{\text{Stab}_{\mathfrak{C}}(F_K)|_{F_I}}{N(z_I)} [I],$$

where $N(z_I)$ is a certain normalization. Under the identification (10.2), one can compare this with the change of basis relation from the GT basis ξ_I to the standard basis v_K , i.e. the reversed relation of (9.1) obtained by applying the orthogonality property of the elliptic weight functions. Then one finds that these two are identical under the correspondence [43]

$$\begin{aligned} \text{standard bases } v_I &\leftrightarrow \text{stable classes } \text{Stab}_{\mathcal{E}}(F_I) \\ \text{the Gelfand-Tsetlin bases } \xi_I &\leftrightarrow \text{fixed point classes } [I]. \end{aligned}$$

This allows us to interpret the level-0 action of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on the GT basis of $\mathcal{V}_{z_1} \otimes \cdots \otimes \mathcal{V}_{z_n}$ in Sec.9 as an action on the fixed point classes in $\bigoplus_{\lambda \in \mathbb{N}^N, |\lambda|=n} E_T(T^* fl_\lambda)$.

11 Elliptic Quantum Toroidal Algebras

A toroidal algebra \mathfrak{g}_{tor} associated with a complex simple Lie algebra $\bar{\mathfrak{g}}$ is a canonical two-dimensional central extension of the double loop Lie algebra $\mathbb{C}^\times \times \mathbb{C}^\times \rightarrow \bar{\mathfrak{g}}$. Its quantum version, i.e. quantum toroidal algebra, was introduced in [26]. We denote it by $U_{q,\kappa}(\mathfrak{g}_{tor})$. It is formulated as an extension of the quantum affine algebra $U_q(\mathfrak{g})$ in the Drinfeld realization [15] by introducing the new generators, i.e. the 0-th generators w.r.t. the index set I , and replacing the finite type Cartan matrix $(a_{i,j})$ $i, j \in \bar{I}$ appearing in the defining relation of $U_q(\mathfrak{g})$ with the affine type generalized Cartan matrix $(a_{i,j})$ $i, j \in I$. This procedure is called the quantum affinization. In particular, for the case $\bar{\mathfrak{g}} = \mathfrak{gl}_N$ one can introduce an extra parameter κ due to a cyclic property of the affine Dynkin diagram of type $A_{N-1}^{(1)}$.

In the same way, the elliptic quantum toroidal algebra $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ is formulated as the quantum affinization of the elliptic algebra $U_{q,p}(\mathfrak{g})$ by adding the 0-th generators $\alpha_{0,m}, e_{0,n}, f_{0,n}, K_0^\pm$ ($m \in \mathbb{Z} \setminus \{0\}, n \in \mathbb{Z}$) [48]. As a result, the element $\prod_{i \in I} (K_i^+)^{a_i^\vee}$ becomes a central element. Here a_i^\vee ($i \in I$) denote the colabel of the affine Dynkin diagram [33]. In addition, $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ possesses two subalgebras, both of which are isomorphic to the elliptic algebra $U_{q,p}(\mathfrak{g})$: the one generated by $\alpha_{i,l}, k_i^\pm, e_{i,n}, f_{i,n}, q^{\pm c/2}, q^d$ ($i \in \bar{I}, l \in \mathbb{Z} \setminus \{0\}, n \in \mathbb{Z}$) and the other by $e_{i,0}, f_{i,0}, K_i^\pm, q^d$ ($i \in I$). These are elliptic analogues of those in $U_{q,\kappa}(\mathfrak{g}_{tor})$ [26].

On the other hand, the \mathfrak{gl}_1 version of the quantum toroidal algebra $U_{q,t}(\mathfrak{gl}_{1,tor})$ was introduced in [58] as a q, t -deformation of the $W_{1+\infty}$ algebra. Remarkably, $U_{q,t}(\mathfrak{gl}_{1,tor})$ is isomorphic to the elliptic Hall algebra [68]. It is also remarkable that representations of $U_{q,t}(\mathfrak{gl}_{1,tor})$ have a deep connection to the Macdonald theory [20, 25, 58, 69]. In addition, representations of $U_{q,t}(\mathfrak{gl}_{1,tor})$ provide a relevant scheme of calculation of the instanton partition functions [63] as well as a study of the Alday-Gaiotto-Tachikawa (AGT) correspondence [1] for the 5d and 6d lifts of the 4d $\mathcal{N} = 2$ SUSY gauge theories, which are known as linear quiver gauge theories.

The elliptic quantum toroidal algebra $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ was formulated in [47]. It has several nice representations such as the level-(0,0) representation realized in terms of the elliptic Ruijsenaars difference operators [67] as well as an elliptic analogue of the q -Fock representation of $U_{q,t}(\mathfrak{gl}_{1,tor})$ which is expected to give a geometric representation on the equivariant elliptic cohomology of the Hilbert scheme of points on \mathbb{C}^2 , $\bigoplus_n E_T(\text{Hilb}_n(\mathbb{C}^2))$. These suggest a deep connection of $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ to an expected elliptic analogue of the theory of Macdonald symmetric function. It is also remarkable that $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ realizes a deformation of the Jordan quiver W -algebra $\mathcal{W}_{p,p^*}(\Gamma(\widehat{A}_0))$ introduced in [34] in the same way as the elliptic algebra $U_{q,p}(\mathfrak{g})$ realizes the deformed W -algebra $\mathcal{W}_{p,p^*}\bar{\mathfrak{g}}$ discussed in Sec.3. Namely, the level-(1, N) elliptic currents of $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ gives a realization of the screening currents and a composition of the vertex operators w.r.t. the Drinfeld comultiplication gives a realization of the generating functions of $\mathcal{W}_{p,p^*}(\Gamma(\widehat{A}_0))$. Moreover thus obtained realization turns out to provide a relevant scheme to the instanton calculus of the 5d and 6d lifts of the 4d $\mathcal{N} = 2^*$ SUSY gauge theories, i.e. the $\mathcal{N} = 2$ SUSY gauge theories coupled with the adjoint matter [63], known as the Jordan quiver gauge theories⁵. There one of the essential feature is that $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ at the level (1, N) possesses the four parameters q, t, p, p^* satisfying one constraint $p/p^* = t/q$, which play the role of the $SU(4)$ Ω -deformation parameters [64].

12 Summary

We have exposed the EQGs $U_{q,p}(\mathfrak{g})$, $U_{q,\kappa,p}(\mathfrak{g}_{tor})$ and $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ in the Drinfeld realization. The main features of them are summarized as follows.

- They allow to treat both the level-0 and level $\neq 0$ representations in a unified way. In particular the level $\neq 0$ representation yields the two elliptic nomes p, p^* , which are identified with q, t in the Macdonald theory and the deformed W -algebras. In the same way $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ possesses the $SU(4)$ Ω -deformation parameters q, t, p, p^* in the level (1, N) representation.
- The coalgebra structure of them allows to formulate the vertex operators as intertwining operators of modules, where the level-0 and level $\neq 0$ representations are mixed as tensor product.
- The vertex operators are important objects having many applications to formulations of mathematical and physical quantities such as elliptic weight functions, integral solutions

⁵It is also called the ADHM quiver gauge theories.

of the elliptic q -KZ equation, vertex functions, correlation functions and form factors in algebraic analysis of elliptic solvable lattice models etc..

- They realize a deformation of the W -algebras including the affine quiver types. In particular, the vertex operators of EQGs provide those of the deformed W -algebras.

We also have shown a correspondence between the level-0 representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ and the equivariant elliptic cohomology $E_T(T^*fI_\lambda)$ by deriving the elliptic weight functions, the integral solutions of the (elliptic) q -KZ equation as well as the level-0 action on the Gelfand-Tsetlin basis. In this process, the vertex operators of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ play an essential role. However, in the geometry side neither the non-zero level representations nor the vertex operators intertwining them have yet been understood. It should be outstanding to formulate the vertex operators of EQGs geometrically.

In addition, knowing that the formulations of elliptic quantum groups and equivariant elliptic cohomologies possess the equivariant and Kähler parameters in a unified way, one of their promising application seems to be the 3d-mirror symmetry for a pair of supersymmetric gauge theories with $\mathcal{N} = 4$ supersymmetry [2]. This provides an important example of the deeper mathematical problem of symplectic duality for conical symplectic resolutions. In such gauge theories, Nakajima quiver varieties are known to give moduli spaces of vacua called the Higgs branch. The 3d-mirror symmetry states that the Higgs branch of the dual theory conjecturally coincides with the Coulomb branch of the original one in such a way that the roles of the equivariant and the Kähler parameters of the dual theories are exchanged. To show the duality at the level of physical quantities such as vertex functions would be outstanding.

References

- [1] L.F. Alday, D. Gaiotto and Y. Tachikawa, Liouville Correlation Functions from Four Dimensional Gauge Theories, *Lett.Math.Phys.* **91** (2010), 167–197.
- [2] M. Aganagic and A. Okounkov, Elliptic Stable Envelopes, Preprint 2016, arXiv:1604.00423.
- [3] M. Aganagic and S.Shakirov, Gauge/vortex duality and AGT, *Math. Phys. Stud.* Springer, Cham (2016), 419–448.
- [4] K. Aomoto and M. Kita, Theory of Hypergeometric Functions, 1994, Maruzen Pub. (in Japanese); English Edition, 2011, Springer Monographs in Mathematics, Springer Japan.

- [5] G.E. Andrews and R.J. Baxter and P.J. Forrester. Eight vertex SOS model and generalized Rogers-Ramanujan-type identities. *J.Stat.Phys.*,35 (1984) 193–266.
- [6] R.J. Baxter, Exactly solved models in statistical mechanics, Academic Press, London (1982)
- [7] A.A. Belavin, Dynamical symmetry of integrable quantum systems. *Nucl. Phys.* **B180** [FS2] (1981), 189-200.
- [8] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, Infinite conformal symmetry in two-dimensional quantum field theory, *Nuclear Phys. B* **241** (1984), 333–380.
- [9] P. Bouwknegt and K. Schoutens, W Symmetry in Conformal Field Theory, *Phys.Rep.* **223** (1993), 183–276.
- [10] J.-S. Caux, H. Konno, M. Sorrell and R. Weston, Exact Form-factor Results for the Longitudinal Structure Factor of the Massless XXZ Model in Zero Field, *J. Stat. Mech.* (2012), P01007 (40pages).
- [11] V. Chari and A. Pressley, A Guide to Quantum Groups, 1994, Cambridge Univ. Press.
- [12] P. Christe and F. Ravanini, $G_N \otimes G_L / G_{N+L}$ Conformal Field Theories and Their Modular Invariant Partition Functions. *Int.J.Mod.Phys.* **A4**, 897-920 (1989)
- [13] E. Date, M. Jimbo, A. Kuniba, T. Miwa and M. Okado. Exactly solvable SOS models. *Nucl. Phys. B*, 290 [FS20]: 231–273, 1987
- [14] E. Date and M. Jimbo and M. Okado, Crystal Base and q -Vertex Operators, *Comm. Math. Phys.*, **155**, 1993, 47–69.
- [15] V.G. Drinfeld, A New Realization of Yangians and Quantized Affine Algebras. *Soviet Math. Dokl.* **36** (1988) 212-216.
- [16] P. Etingof and A. Varchenko, Exchange Dynamical Quantum Groups, *Comm.Math.Phys.* **205** (1999), 19–52.
- [17] R.M. Farghly, H. Konno, K. Oshima, Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{g}})$ and Quantum Z -algebras, *Alg. Rep. Theory* **18** (2014), 103-135.
- [18] B. Feigin and E. Frenkel, Affine Kac-Moody Algebras at the Critical Level and Gelfand-Dikii Algebras, *Int.J.Mod.Phys. A* **7**, Suppl.1A, Proceedings of the RIMS Project 1991, “Infinite Analysis” (1992), 197–215.

- [19] B. Feigin and D. Fuchs, Skew-symmetric invariant differential operators on the line and Verma modules over the Virasoro algebra, *Funktsional. Anal. i Prilozhen.* **16** (1982), 47–63. English translation: *Funct. Annal. Appl.* **16** (1982), 114–126; Verma modules over a Virasoro algebra. *Funktsional. Anal. i Prilozhen.* **17** (1983), 91–92. English translation: *Funct. Annal. Appl.* **17** (1983), 241–242.
- [20] B. Feigin and A. Tsybaliuk, Bethe Subalgebras of $U_q(\widehat{\mathfrak{gl}}_n)$ via Shuffle Algebras, *Selecta Math. (N.S.)* **22** (2016), no. 2, 979–1011.
- [21] G. Felder, Elliptic Quantum Groups, *Proc. ICMP Paris-1994* (1995), 211–218.
- [22] O. Foda, M. Jimbo, T. Miwa, K. Miki and A. Nakayashiki, Vertex Operators in Solvable Lattice Models, *J. Math. Phys.* **35** (1994) 13–46.
- [23] I. B. Frenkel and N. Reshetikhin, Quantum Affine Algebras and Holonomic Difference Equations, *Comm. Math. Phys.* **146** (1992) 1–60.
- [24] E. Frenkel and N. Reshetikhin, Deformation of W -Algebras Associated to Simple Lie Algebras, *Comm. Math. Phys.* **197** (1998) 1–32.
- [25] M. Fukuda, Y. Ohkubo and J. Shiraishi, Generalized Macdonald Functions on Fock Tensor Spaces and Duality Formula for Changing Preferred Direction, arXiv:1903.05905.
- [26] V. Ginzburg, M. Kapranov and E. Vasserot, Elliptic Algebras and Equivariant Elliptic Cohomology I, Preprint (1995), arXiv:q-alg/9505012.
- [27] P. Goddard and A. Kent and D. Olive. Virasoro algebras and coset space models *Phys. Lett. B*,152:88–,1985; Unitary representations of the Virasoro and super-Virasoro algebras *Comm. Math. Phys.*,103:105–119,1986.
- [28] M. Jimbo, H. Konno, S. Odake and J. Shiraishi, Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_2)$: Drinfeld Currents and Vertex Operators, *Comm. Math. Phys.* **199** (1999), 605–647
- [29] M. Jimbo, H. Konno, S. Odake and J. Shiraishi, Quasi-Hopf Twistors for Elliptic Quantum Groups, *Transformation Groups* **4** (1999), 303–327.
- [30] M. Jimbo and T. Miwa. *Algebraic Analysis of Solvable Lattice Models*. CBMS Regional Conference Series in Mathematics vol. 85, AMS, 1994.
- [31] M. Jimbo and T. Miwa and M. Okado, Solvable Lattice Models whose States are Dominant Integral Weights of $A_{n-1}^{(1)}$, *Lett. Math. Phys.* **14** (1987), 123–131.

- [32] M. Jimbo and T. Miwa and M. Okado, Solvable Lattice Models Related to the Vector Representation of Classical Simple Lie Algebras, *Comm. Math. Phys.* **116** (1988), 507–525.
- [33] V. G. Kac, Infinite Dimensional Lie algebras, *3rd. ed. Cambridge University Press*, 1990.
- [34] T. Kimura and V. Pestun, Quiver W -algebras *Lett.Math.Phys.* **108** 1351-1381 (2018)
- [35] E. Koelink and H. Rosengren, Harmonic Analysis on the $SU(2)$ Dynamical Quantum Group, *Acta.Appl.Math.*, **69** (2001), 163–220.
- [36] T. Kojima and H. Konno, The Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ and the Drinfeld Realization of the Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(\widehat{\mathfrak{sl}}_N)$, *Comm. Math. Phys.* **239** (2003), 405-447.
- [37] T. Kojima, H. Konno and R. Weston, The Vertex-Face Correspondence and Correlation Functions of the Fusion Eight-Vertex Models I: The General Formalism, *Nucl. Phys.* **B720** (2005), 348-398.
- [38] H. Konno, An Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_2)$ and the Fusion RSOS Models, *Comm. Math. Phys.* **195** (1998), 373–403.
- [39] H. Konno, Dynamical R Matrices of Elliptic Quantum Groups and Connection Matrices for the q -KZ Equations, *SIGMA*, **2** (2006), Paper 091, 25 pages.
- [40] H. Konno, Elliptic Quantum Group $U_{q,p}(\widehat{\mathfrak{sl}}_2)$ and Vertex Operators, *J.Phys.A***41** (2008) 194012.
- [41] H. Konno, Elliptic Quantum Group $U_{q,p}(\widehat{\mathfrak{sl}}_2)$, Hopf Algebroid Structure and Elliptic Hypergeometric Series, *J. Geom. Phys.* **59** (2009), 1485-1511.
- [42] H. Konno, Elliptic Weight Functions and Elliptic q -KZ Equation, *Journal of Integrable Systems* **2** (2017) 1-43. doi: 10.1093/integr/xyx011.
- [43] H. Konno, Elliptic Stable Envelopes and Finite-dimensional Representations of Elliptic Quantum Group, *Journal of Integrable Systems* **3** (2018) 1-43. doi: 10.1093/integr/xyy012.
- [44] H. Konno, Elliptic Quantum Groups $U_{q,p}(\widehat{\mathfrak{gl}}_N)$ and $E_{q,p}(\widehat{\mathfrak{gl}}_N)$, *Adv.Studies in Pure Math.* **76** (2018), 347-417.
- [45] H. Konno, Elliptic Quantum Groups: Representations and Related Geometry, Springer Briefs in Mathematical Physics (2020) Springer.

- [46] H. Konno and K. Oshima, Elliptic Quantum Group $U_{q,p}(B_N^{(1)})$ and Vertex Operators, *RIMS Kokyuroku Bessatsu* , **B62** (2017) 97-148.
- [47] H. Konno and K. Oshima, Elliptic Quantum Toroidal Algebra $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ and Affine Quiver Gauge Theories, *Lett.Math.Phys.* **113** (2023) 32, 64 pages.
- [48] H. Konno and K. Oshima, Elliptic Quantum Toroidal Algebras, Z -algebra Structure and Representations, preprint (2023), *Alg. Rep. Theory* (2024) <https://doi.org/10.1007/s10468-024-10251-3>.
- [49] P. Koroteev, P. Pushkar, A. Smirnov and A. Zeitlin, Quantum K-theory of Quiver Varieties and Many-body Systems, Preprint 2017, arXiv:1705.10419.
- [50] A. Kuniba, Exact Solution of Solid-on-solid Models for Twisted Affine Lie Algebras $A_{2n}^{(2)}$ and $A_{2n-1}^{(2)}$, *Nucl. Phys.*, **B355**, 1991, 801–821.
- [51] A. Kuniba and J. Suzuki, Exactly solvable $G_2^{(1)}$ solid-on-solid models. , *Phys. Lett.*, **A160**, 1991, 216–222.
- [52] M. Lashkevich and Y. Pugai. Free field construction for correlation functions of the eight-vertex model. *Nucl. Phys.* **B516** (1998), 623-651.
- [53] J. Lepowsky and R.L. Wilson, A New Family of Algebras Underlying the Rogers-Ramanujan Identities and Generalizations, *Proc. Natl. Acad. Sci. USA* **78** (1981) 7254-7258; The Structure of Standard Modules, I: Universal Algebras and the Roger-Ramanujan Identities, *Invent.Math.***77** (1984) 199–290.
- [54] S.L. Lukyanov and V.A. Fateev , Additional Symmetries and Exactly-Soluble Models in Two-Dimensional Conformal Field Theory, *Sov. Sci. Rev. A. Phys.***15** (1990), 1-117.
- [55] S. Lukyanov and Y. Pugai, Multi-point Local Height Probabilities in the Integrable RSOS Model, *Nucl.Phys.* **B473**(1996), 631-658.
- [56] A. Matsuo, Jackson Integrals of Jordan Pochhammer Type and Quantum Knizhnik Zamolodchikov Equations, *Comm. Math.Phys.* **151** (1993) 263–273; Quantum Algebra Structure of Certain Jackson Integrals, *Comm. Math.Phys.* **157** (1993) 479–498.
- [57] D. Maulik and A. Okounkov, Quantum Groups and Quantum Cohomology , Preprint 2012, arXiv:1211.1287,
- [58] K. Miki, A (q, γ) analog of the $W_{1+\infty}$ algebra. *J. Math. Phys.* **48**, 123520, 35pp. (2007)

- [59] K. Mimachi, A Solution to Quantum Knizhnik-Zamolodchikov Equations and Its Application to Eigenvalue Problems of the Macdonald Type, *Duke Math. J.* **85** (1996) 635–658.
- [60] A. Molev, Yangians and Classical Lie Algebras, *Mathematical Surveys and Monographs* **143** (2007) AMS.
- [61] H. Nakajima, Instantons on ALE Spaces, Quiver Varieties and Kac-Moody Algebras, *Duke Math. J.* **76** (1994) 365–416.
- [62] H. Nakajima, Quiver Varieties and Kac-Moody Algebras, *Duke Math. J.* **91** (1998) 515–560.
- [63] N. Nekrasov, Seiberg - Witten Prepotential From Instanton Counting. *Adv. Theor. Math. Phys.* **7** 831-864 (2004)
- [64] N. Nekrasov, BPS/CFT Correspondence: Non-perturbative Dyson-Schwinger Equations and qq -Characters. *J.High Energy Phys.* **03** 181 (2016)
- [65] A. Okounkov, Lectures on K-theoretic Computations in Enumerative Geometry. (2015), arXiv: 1512.07363.
- [66] R. Rimányi, V. Tarasov and A. Varchenko, Trigonometric Weight Functions as K -theoretic Stable Envelope Maps for the Cotangent Bundle of a Flag Variety, *J.Geom.Phys.*, **94** (2015) 81–119.
- [67] S.N.M. Ruijsenaars, First order analytic difference equations and integrable quantum systems, *J. Math. Phys.*, **38** (1997), 1069–1146.
- [68] O. Schiffmann, Drinfeld Realization of the Elliptic Hall Algebra. arXiv:1004.2575.
- [69] O. Schiffmann and E. Vasserot, The Elliptic Hall Algebra, Cherednik Hecke Algebras and Macdonald Polynomials. *Composito Math.* **147** 188-234 (2011)
- [70] V. Tarasov and A. Varchenko, Geometry of q -Hypergeometric Functions, Quantum Affine Algebras and Elliptic Quantum Groups, *Astérisque* **246** (1997) Société Mathématique de France.