

Generic behavior of differentially positive systems on a globally orderable Riemannian manifold

Lin Niu*

School of Mathematics and Physics
University of Science and Technology Beijing
Beijing, 100083, P. R. China

Yi Wang†

School of Mathematical Sciences
University of Science and Technology of China
Hefei, Anhui, 230026, P. R. China

Abstract

Differentially positive systems are the nonlinear systems whose linearization along trajectories preserves a cone field on a smooth Riemannian manifold. One of the embryonic forms for cone fields in reality is originated from the general relativity. By utilizing the Perron-Frobenius vector fields and the Γ -invariance of cone fields, we show that generic (i.e., “almost all” in the topological sense) orbits are convergent to certain single equilibrium. This solved a reduced version of Forni-Sepulchre’s conjecture in 2016 for globally orderable manifolds.

Keywords: Differential positivity, Causal order, Generic convergence, Homogeneous space, Cone field

AMS Subject Classification (2020): 37C20, 37C65, 53C30, 22F30

1 Introduction

Differential analysis provides a general framework for investigating a nonlinear dynamical system by analyzing the linearization of the system at every point in the state space. The motivation is that the local behavior of a system often has strong implications for the global nonlinear behavior. The current series of papers focus on the nonlinear dynamical system whose

*Supported by the National Natural Science Foundation of China No.12201034 and 12090012.

†Supported by the National Natural Science Foundation of China No.12331006.

linearization along trajectories preserves a cone field. Here, a cone field C_M assigns to each point x of a manifold M a closed convex cone $C_M(x)$ in the tangent space $T_x M$ of x .

One of the embryonic forms for cone fields in reality is originated from the general relativity, for which the time-orientable space-time (i.e., a connected 4-dimensional Hausdorff C^∞ -manifold endowed with a Lorentz metric) naturally generates at each point a Lorentzian cone in tangent space (see, e.g. [2, 12, 43] or Appendix A.1). The structure of a cone field in the time-orientable space-time turns out to be one of the crucial mathematical tools in the study of general relativity including causality theory, singularity theory and black holes, etc.. Among others, the well-known non-spacelike curve (also called causal curve) in these theories (c.f. Hawking and Ellis [12] and Penrose [43]) is actually one whose tangent vector at each point falls in the Lorentzian cone field. Moreover, the causally related points in a space-time should be joined by a non-spacelike curve.

From this point of view, a cone field on the manifold M naturally induces a “conal order relation” as follows: two points $x, y \in M$ are *conal ordered*, denoted by $x \leq_M y$, if there exists a conal curve on M beginning at x and ending at y . Here, a conal curve is a piecewise smooth curve whose tangent vector lies in the cone at every point along the curve wherever it is defined (see Definition 2.1 and Fig.1(a)). In particular, in the setting of space-times (see [11, 29, 48] and [16, 42, 54]), the causal curves are actually the conal curves; while, the causally related points are the ones that are conal ordered. Besides, conal order also has various applications in hyperbolic partial differential equations and harmonic analysis (see [5, 6]), as well as in the theory of Wiener-Hopf operators on symmetric spaces [18].

It deserves to point out that, unlike the standard order relation induced by a single closed convex cone in a topological vector space, the conal order relation “ \leq_M ” induced by a cone field on M is certainly reflexive and transitive, but not always antisymmetric. In fact, there are examples of M that contain the closed conal curves (e.g., the closed timelike curves in time-orientable space-times [12, Chapter 5]), which reveals that the anti-symmetry fails. Moreover, the relation “ \leq_M ” is not necessarily closed, i.e., the set $\{(x, y) \in M \times M : x \leq_M y\}$ is not a closed subset in $M \times M$ (see [26, p.299] or [35, p.470]). For instance, such set is not necessarily closed in Minkowski space (see [12, p.183] or [43, p.12]).

In many recent works [7–9, 33, 34], the nonlinear system whose linearization along trajectories preserves a convex cone field is also referred as differentially positive system, the flow of which naturally keeps the conal order “ \leq_M ”. To the best of our knowledge, Forni and Sepulchre [7–9] first studied the dynamics of differentially positive systems. Among others, they constructed on M a canonical defined vector field \mathcal{W} , called the Perron-Frobenius vector field, such that the vector $\mathcal{W}(x)$ lies in the interior of the cone $C_M(x)$ at each point $x \in M$. By appealing to the Perron-Frobenius vector field \mathcal{W} , a dichotomy characterization of limit sets for differentially positive systems was provided. Based on this, Forni and Sepulchre [9, p.353] further posed the following

Conjecture ([Forni and Sepulchre [9]). *For almost every $x \in M$, the ω -limit set $\omega(x)$ is given by either a fixed point, or a limit cycle, or fixed points and connecting arcs.*

This conjecture predicts what typically happens to orbits, as time increases to $+\infty$. The description of typical properties of orbits usually means those shared by “most” orbits.

In our present work, we will tackle this conjecture and made an attempt to establish the asymptotic behavior of “most” orbits for differentially positive systems. For this purpose, we first introduce the following assumption for M :

(H1) M is globally orderable equipped with a continuous solid cone field.

(H1) means that the conal order “ \leq_M ” on M is a partial order (see [16, 17, 26, 35] and Definition 2.2); and hence, it is actually anti-symmetric. This occurs naturally in various situations. One of the well-known examples is a homogeneous space of positive definite matrices with the affine-invariant cone field (see, e.g, [32, Section 3]). As a matter of fact, (H1) excludes the occurrence of the closed conal curves (see [26, Section 5]). In the setting of space-times, such closed conal (causal) curves would seem to generate paradoxes involving causality and are said to “violate causality” ([2, 12, 13, 29, 43]).

As a consequence, by (H1), the possibility of a limit cycle, or fixed points and connecting arcs (which are all closed conal curves) can be ruled out. In other words, Forni-Sepulchre’s Conjecture is naturally reduced to the following

Conjecture A. *For almost every $x \in M$, the ω -limit set $\omega(x)$ is a singleton.*

In the present paper, we will prove conjecture A under the following two reasonable assumptions:

(H2) The conal order “ \leq_M ” is quasi-closed.

(H3) Both the cone field C_M and the Riemannian metric on M are Γ -invariant.

More precisely, our main result is the following

Theorem A. *Assume that (H1)-(H3) hold. Then, for almost every $x \in M$, the ω -limit set $\omega(x)$ is a singleton.*

This theorem, in a detailed version, will be proved in Section 3 (see Theorem 3.1). It concludes that generic (i.e., “almost all” in the topological sense) orbits are convergent to a single equilibrium.

The quasi-closedness of “ \leq_M ” in (H2) means that $x \leq_M y$ whenever $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$ and $x_n \ll_M y_n$ for all integer $n \geq 0$ (see Definition 2.3). Here, we write $x \ll_M y$ if there exists a so-called *strictly* conal curve (whose tangent vector lies in the interior of cone at

every point along the curve) on M beginning at x and ending at y (see Definition 2.1). We point out that (H2) is motivated by the causal continuity of space-times in Hawking and Sachs [13], while the notation “ \ll_M ” is derived from the timelike curve in general relativity (see e.g., [43]). In fact, one can prove that the conal order “ \leq_M ” is indeed quasi-closed in many causally continuous space-times (see Theorem A.1 in the Appendix).

As for (H3), a cone field C_M is called Γ -invariant if there exists a linear invertible mapping $\Gamma(x_1, x_2) : T_{x_1}M \rightarrow T_{x_2}M$ for all $x_1, x_2 \in M$, such that $\Gamma(x_1, x_2)C_M(x_1) = C_M(x_2)$. While, the Riemannian metric $(\cdot, \cdot)_x$ on M is Γ -invariant if $(u, v)_{x_1} = (\Gamma(x_1, x_2)u, \Gamma(x_1, x_2)v)_{x_2}$ for all $x_1, x_2 \in M$ and $u, v \in T_{x_1}M$. Actually, Γ -invariance is motivated by the homogeneous structure of the manifolds. A well-known example admitting (H3) is a homogeneous space $M = G/H$ of a connected Lie group G assigned to each point $x \in M$ a closed convex cone in the tangent space T_xM such that the cone field is invariant under the action of G (see e.g., [17, 26, 35]); and a homogeneous Riemannian metric on such homogeneous space is naturally Γ -invariant (see e.g., [1, 14]). In the Appendix A.2, more detailed structures of the globally orderable homogeneous spaces are presented. We further mention here that the homogeneous cone field comes from the Lie theory. It is shown that cone fields arise as quotients these so-called Lie wedge fields (see [16, Lemma VI.1.5]). This provides a crucial link between the Lie theory of semigroups and that of cone fields on manifolds (see [16, 17, 19, 26, 35]). Very recently, the works in [31, 36–39] form an ongoing project aiming at the connections between causal structures on homogeneous spaces, algebraic quantum field theory, modular theory of operator algebras and unitary representations of Lie groups.

Needless to say, differential positivity in Forni and Sepulchre [9] with respect to a constant cone field on a flat space M is reduced to the classical monotonicity ([4, 15, 21–23, 30, 44–46, 49–53, 55, 56]) with respect to a closed convex cone. From this point of view, differentially positive systems can be regarded as a natural generalization of the so-called classical monotone systems to nonlinear manifolds. We pointed out that the order introduced by a closed convex cone is a closed partial order (see the detail in Appendix A.3). In such a flat space, Theorem A automatically implies the celebrated Hirsch’s Generic Convergence Theorem [21].

The paper is organized as follows. In Section 2, we introduce some notations and definitions and summarize some preliminary results. Theorem A (i.e., Theorem 3.1) with its proof will be given in Section 3. Several fundamental tools and critical lemmas, which turns out be useful in the proof of Theorem A (or Theorem 3.1), will be postponed in Section 4. Finally, in the Appendix, we will present the order structures on space-times, the globally orderable homogeneous spaces with homogeneous cone fields, as well as the differential positivity in the flat spaces.

2 Notations and preliminary results

Throughout this paper, M will be reserved as a smooth manifold of dimension n . The tangent bundle is denoted by TM and the tangent space at a point $x \in M$ by T_xM . M is endowed with a Riemannian metric tensor, represented by an inner product $(\cdot, \cdot)_x$ on the tangent space T_xM . We set $|v|_x := \sqrt{(v, v)_x}$ for any $v \in T_xM$ (sometimes we omit the subscripts x in $|\cdot|_x$) and the Riemannian metric endows the manifold with the Riemannian distance d . We assume that (M, d) is a complete metric space.

Let E be a n dimensional real linear space. A nonempty closed subset C of the linear space E is called a *closed convex cone* if $C + C \subset C$, $\alpha C \subset C$ for all $\alpha \geq 0$, and $C \cap (-C) = \{0\}$. A convex cone C is *solid* if its interior $\text{Int}C \neq \emptyset$. A convex cone C' in E is said to *surround* a cone C if $C \setminus \{0\} \subset \text{Int}C'$.

A *cone field* on a manifold M is a map $x \mapsto C_M(x)$, such that $C_M(x)$ is a closed convex cone in T_xM for each $x \in M$. A manifold equipped with a cone field is called a *conal manifold*. A cone field is *solid* if each convex cone $C_M(x)$ is solid.

A cone field C_M is called a Γ -*invariant* cone field if there exists a linear invertible mapping $\Gamma(x_1, x_2) : T_{x_1}M \rightarrow T_{x_2}M$ for each $x_1, x_2 \in M$, such that $\Gamma(x_1, x_2)C_M(x_1) = C_M(x_2)$. Moreover, Γ is continuous with respect to (x_1, x_2) and $\Gamma(x, x) = \text{Id}_x$ for all $x \in M$, where $\text{Id}_x : T_xM \rightarrow T_xM$ is the identity map. We say that the Riemannian metric is Γ -*invariant* under the linear invertible mapping Γ if $(u, v)_{x_1} = (\Gamma(x_1, x_2)u, \Gamma(x_1, x_2)v)_{x_2}$ for all $x_1, x_2 \in M$ and $u, v \in T_{x_1}M$. In this case, $|v|_{x_1} = |\Gamma(x_1, x_2)v|_{x_2}$ holds for all $x_1, x_2 \in M$ and $v \in T_{x_1}M$.

Pick a smooth chart $\Phi : U \rightarrow V$, where $U \subset M$ is an open set containing x and $V \subset \mathbb{R}^n$ is an open set. Let $C_M^\Phi(y) \subset \mathbb{R}^n$ denote the representation of the cone field in the chart, i.e., $d\Phi(y)(C_M(y)) = \{\Phi(y)\} \times C_M^\Phi(y)$. Following [26], a cone field $x \mapsto C_M(x)$ on M is *upper semicontinuous* at $x \in M$ if given a smooth chart $\Phi : U \rightarrow \mathbb{R}^n$ at x and a convex cone C' surrounding $C_M^\Phi(x)$, there exists a neighborhood W of x such that $C_M^\Phi(y) \subset C'$ for all $y \in W$. The cone field is *lower semicontinuous* at x if given any open set N such that $N \cap C_M^\Phi(x)$ is non-empty, there exists a neighborhood W of x such that $N \cap C_M^\Phi(y)$ is non-empty for all $y \in W$. A cone field is *continuous at x* if it is both lower and upper semicontinuous at x and *continuous* if it is continuous at every x .

Definition 2.1. A continuous piecewise smooth curve $t \mapsto \gamma(t)$ defined on $[t_0, t_1]$ into a conal manifold M is called a *conal curve* if $\gamma'(t) \in C_M(\gamma(t))$ whenever $t_0 \leq t < t_1$, in which the derivative is the right hand derivative at those finitely many points where the derivative is not continuous. (See Fig.1(a)). Moreover, γ is called a *strictly conal curve* if $\gamma'(t) \in \text{Int}C_M(\gamma(t))$ for $t_0 \leq t < t_1$.

For two points $x, y \in M$, we say x and y are ordered, denoted by $x \leq_M y$, if there exists a conal curve $\alpha : [t_0, t_1] \subset \mathbb{R} \rightarrow M$ such that $\alpha(t_0) = x$ and $\alpha(t_1) = y$. This relation is an *order* on M . In fact, it is always reflexive (i.e., $x \leq_M x$ for all $x \in M$) and transitive (i.e., $x \leq_M y$ and $y \leq_M z$ implies $x \leq_M z$). The order “ \leq_M ” is referred to as the *conal order*. We write $x \ll_M y$ if

there exists a so-called *strictly conal* curve γ with $\gamma(t_0) = x$ and $\gamma(t_1) = y$. Clearly, the relation “ \ll_M ” is always transitive. Let U, V be the subsets of M . We write $U \leq_M V$ (resp., $U \ll_M V$) if $x \leq_M y$ (resp., $x \ll_M y$) for any $x \in U$ and $y \in V$.

The following proposition implies that the relation “ \ll_M ” is *open*.

Proposition 2.1. *If $x \ll_M y$, then there exist neighborhoods U of x , V of y such that $U \ll_M V$.*

Proof. We only prove the existence of V , i.e., if $x \ll_M y$, there exists a neighborhood V of y such that $x \ll_M z$ for each $z \in V$. Since $x \ll_M y$, there is a strictly conal curve $\gamma_1 : [t_0, t_1] \subset \mathbb{R} \rightarrow M$ such that $\gamma_1(t_0) = x$ and $\gamma_1(t_1) = y$. Let $\gamma(t) = \gamma_1(t_1 - t)$, $t \in [0, t_1 - t_0]$. Then $\gamma(0) = y$, $\gamma(t_1 - t_0) = x$ and $\gamma'(0) \in \text{Int}\{-C_M(\gamma(0))\}$.

Let $\Phi : U \rightarrow \Phi(U) \subset \mathbb{R}^n$ be a chart with $\Phi(\gamma(0)) = 0$ such that $\Phi(U)$ is convex. We set $-\tilde{C}(\Phi(z)) = d\Phi(z)(-C_M(z))$ for $z \in U$ and $\alpha = \Phi \circ \gamma$. Then $\alpha'(0) \in \text{Int}\{-\tilde{C}(0)\}$. So, we can choose a compact convex neighborhood B of $\alpha'(0)$ in the interior of $-\tilde{C}(0)$ and set $W = \mathbb{R}^+ B := \{\lambda B : \lambda > 0\} \subset -\tilde{C}(0)$. Since the cone field is continuous, there is a neighborhood $V \subset M$ of $\gamma(0)$ such that $W \subset -\tilde{C}(z)$ for all $z \in U' = \Phi(V)$ and U' is convex.

On the other hand, $\alpha'(0) = \lim_{t \rightarrow 0^+} \frac{1}{t} \alpha(t) \in \text{Int}W$. Then there is a $s_0 > 0$ such that $\alpha((0, s_0]) \subset \text{Int}W$ and $\alpha([0, s_0]) \subset U'$. Let B' be an open convex neighborhood of $\alpha(s_0)$ in $W \cap U'$.

Let $B'' = \{\alpha(s_0) - u : u \in B'\} \cap U'$, then B'' is an open neighborhood of $\alpha(0)$. For each $v \in B''$, we set $\alpha_v(t) = \alpha(s_0) + t(v - \alpha(s_0))$, $t \in [0, 1]$. Then $\alpha_v(0) = \alpha(s_0)$, $\alpha_v(1) = v$ and $\alpha'_v(t) = v - \alpha(s_0) \in -B' \subset \tilde{C}(\alpha_v(t))$.

Thus, there is a point $p \in \gamma_1$ and an open neighborhood D of y such that $p \ll_M z$ for all $z \in D$. Since $x \ll_M p$, we obtain $x \ll_M z$ for all $z \in D$. \square

According to our standing assumption (H1) in the introduction, we now give the following crucial definitions:

Definition 2.2. A conal manifold is said to be *globally orderable* if the order “ \leq_M ” is a partial order which is locally order convex.

The conal order “ \leq_M ” is a *partial* order relation if it is additionally antisymmetric (i.e., $x \leq_M y$ and $y \leq_M x$ implies $x = y$). A partial order “ \leq ” is *locally order convex* if it has a basis of neighborhoods at each point that are order convex ($z, x \in U$ implies $\{y : z \leq y \leq z\} \subset U$). See [26, 35].

Remark 2.1. In the terminology of general relativity, each point p in a globally orderable manifold M is called a strong point (see Lawson [26, Section 5]), by which means that every neighborhood U of p contains a smaller neighborhood V of p such that every conal curve that begins in V and leaves U terminates outside of V (see e.g., [12, p.192], [43, p.28] or [2, p.59]).

Remark 2.2. The *globality of the conal order*, i.e., whether the conal order associated with a cone field can be extended to a partial order on the global manifold, is a central problem in [26].

The equivalence between globality of the conal order in a homogeneous manifold and globality of the Lie wedge in the Lie group has been shown in [26, Section 5] (see also [35, Theorem 1.6] and [17, Section 4.3]). Globality of a conal order occurs naturally in various situations; for example, the conal order induced by the affine-invariant cone field on the homogeneous space of positive definite matrices, as we mentioned above, is a partial order (see [32, Theorem 2]). Besides, if the manifold is globally orderable, one can exclude the occurrence of the closed conal curves (see [26, Section 5]).

Definition 2.3. The order “ \leq_M ” is *quasi-closed* if $x \leq_M y$ whenever $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$ and $x_n \ll_M y_n$ for all n .

Remark 2.3. In general, the conal order is not a closed order ($x \leq_M y$ whenever $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$ and $x_n \leq_M y_n$), see [26, p.299] and [35, p.470]. The quasi-closed order relationship here is inspired by the causal continuity of space-times studied by Hawking and Sachs in [13]. One may refer to Appendix A.1 for more details.

Now, we consider a system Σ generated by a smooth vector field f on M equipped with a continuous solid cone field C_M . The induced flow by system Σ is denoted by φ_t . We write $d\varphi_t(x)$ as the tangent map from $T_x M$ to $T_{\varphi_t(x)} M$. Let $x \in M$, the *positive semiorbit* (resp., *negative semiorbit*) of x , be denoted by $O^+(x) = \{\varphi_t(x) : t \geq 0\}$ (resp., $O^-(x) = \{\varphi_t(x) : t \leq 0\}$). The *full orbit* of x is denoted by $O(x) = O^+(x) \cup O^-(x)$. An *equilibrium* is a point x such that $O(x) = \{x\}$. Let E be the set of all the equilibria of φ_t . The ω -limit set $\omega(x)$ of x is defined by $\omega(x) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} \varphi_t(x)}$. A point $z \in \omega(x)$ if and only if there exists a sequence $\{t_i\}$, $t_i \rightarrow \infty$, such that $\varphi_{t_i}(x) \rightarrow z$ as $i \rightarrow \infty$. If $O^+(x)$ is precompact, then $\omega(x)$ is nonempty, compact, connected, and invariant (i.e., $\varphi_t(\omega(x)) = \omega(x)$ for any $t \in \mathbb{R}$). A point x is called a *convergent point* if $\omega(x)$ consists of a single point of E . The set of all convergent points is denoted by C .

Throughout the paper, we always assume that all orbits are forward complete, which means $\varphi_t(x)$ is well defined for all $t \geq 0$, and the orbit of x has compact closure for each $x \in M$.

Definition 2.4. The system Σ is said to be *differentially positive* (DP) with respect to C_M if

$$d\varphi_t(x)C_M(x) \subseteq C_M(\varphi_t(x)), \quad \forall x \in M, \quad \forall t \geq 0.$$

And the differentially positive system Σ is said to be *strongly differentially positive* (SDP) with respect to C_M if

$$d\varphi_t(x)\{C_M(x) \setminus \{0\}\} \subseteq \text{Int}C_M(\varphi_t(x)), \quad \forall x \in M, \quad \forall t > 0.$$

See Fig.1(b).


In the present paper, we focus on the strongly differentially positive system Σ on a Riemannian manifold M . We first give the following two useful preliminary results.

Proposition 2.2. *If $x \leq_M y$, then $\varphi_t(x) \leq_M \varphi_t(y)$ for $t \geq 0$. Moreover, $\varphi_t(x) \ll_M \varphi_t(y)$ for all $t > 0$.*

Proof. If $x \leq_M y$, there exists a conal curve $\gamma(s)$ such that $\gamma(0) = x$ and $\gamma(1) = y$. Since Σ is SDP and $\frac{d}{ds}\gamma(s) \in C_M(\gamma(s)) \setminus \{0\}$, then $\frac{d}{ds}\varphi_t(\gamma(s)) = d\varphi_t(\gamma(s))\frac{d}{ds}\gamma(s) \in \text{Int}C_M(\varphi_t(\gamma(s)))$ for $t > 0$. So, $\varphi_t(\gamma(s))$ is a strictly conal curve. \square

Proposition 2.3. *If $x \leq_M y$, then for $t_0 > 0$, there exist neighborhoods U of x , V of y such that $\varphi_t(U) \ll_M \varphi_t(V)$ for any $t \geq t_0$.*

Proof. Since $x \leq_M y$, $\varphi_{t_0}(x) \ll_M \varphi_{t_0}(y)$ for $t_0 > 0$. One can take neighborhoods \bar{U} of $\varphi_{t_0}(x)$ and \bar{V} of $\varphi_{t_0}(y)$ such that $\bar{U} \ll_M \bar{V}$ by Proposition 2.1. By the continuity of φ_{t_0} , there are neighborhoods U of x , V of y such that $\varphi_{t_0}(U) \subset \bar{U}$ and $\varphi_{t_0}(V) \subset \bar{V}$. \square



(a) A conal curve γ on manifold M .

(b) Strongly differentially positive system on manifold M

Fig 1: Conal curves and strongly differentially positive flows on manifold M .

As mentioned in the introduction, we hereafter impose the following hypotheses:

(H1) M is a globally orderable conal manifold equipped with a continuous solid cone field C_M .

(H2) The conal order “ \leq_M ” is quasi-closed.

(H3) Both the cone field C_M and the Riemannian metric on M are Γ -invariant.

Before ending this section, we give the following critical lemma, which turns out to be important for the proof of our main results in the forthcoming sections.

Lemma 2.1. *Assume that (H1)-(H2) hold. Then*

(a) *The ω -limit set cannot contain two points x and y with $x \leq_M y$;*

(b) *If $x \leq_M y$, then either $\omega(x) \ll_M \omega(y)$, or $\omega(x) = \omega(y) \subset E$.*

For the sake of the completeness, we will postpone its proof in Section 4.

3 Proof of Theorem A

In this section, we will focus on the generic behavior of the SDP flow φ_t under the hypotheses (H1)-(H3).

Theorem 3.1. (Generic Convergence) *Assume that (H1)-(H3) hold. If the cone field admits a C^∞ -section and φ_t satisfies the compactness condition (P), then $\text{Int}C$ is dense in M .*

Recall that C denotes the set of all convergent points. Here, we say that a cone field admits a C^∞ -section if there is a C^∞ vector field X such that $X(x) \in C_M(x)$ for all $x \in M$. We also formulate the *compactness condition (P)*: For each $x_0 \in M$, $\cup_{n \geq 0} \omega(x_n)$ has compact closure contained in M , where $\{x_n\} \rightarrow x_0$ with $x_n \leq_M x_{n+1} \leq_M x_0$ (or $x_0 \leq_M x_{n+1} \leq_M x_n$) for $n \geq 1$.

Remark 3.1. We point out that the homogeneous cone field on a homogeneous space naturally admits C^∞ -sections (see [26] or Proposition A.6 in the Appendix). While the compactness condition (P) is a relatively weak compactness requirement which is frequently satisfied (see e.g., [50, 51]).

The proof of Theorem 3.1 will be divided into several steps. To proceed it, we recall the so-called *Perron-Frobenius vector field*, which was first introduced by Forni and Sepulchre [9]. A continuous vector field v on an invariant compact set $\Omega \subset M$ is called a *Perron-Frobenius vector field* on Ω , if for each $x \in \Omega$,

- (i) $v(x) \in \text{Int}C_M(x)$ with $|v(x)|_x = 1$;
- (ii) $v(\varphi_t(x)) = \frac{d\varphi_t(x)v(x)}{|d\varphi_t(x)v(x)|}$ for $t \geq 0$;
- (iii) $\lim_{t \rightarrow \infty} d_{\varphi_t(x)}(d\varphi_t(x)u, v(\varphi_t(x))) = 0$ for all $u \in C_M(x) \setminus \{0\}$,

where $d_{\varphi_t(x)}(\cdot, \cdot)$ is the Hilbert Metric induced by $C_M(\varphi_t(x))$ (see [9, Section VI]).

Lemma 3.1. *Let $\omega(x) \subset E$ and v be the Perron-Frobenius vector field on $\omega(x)$. If $\omega(x)$ is not a singleton, then there exist $\tau > 0$ and $\rho(e) > 1$ such that*

$$d\varphi_\tau(e)v(e) = \rho(e)v(e), \quad \text{for } e \in \omega(x).$$

Moreover, there exists $\rho > 1$ such that $\rho(e) > \rho$ for all $e \in \omega(x)$.

Proof. Fix $\tau > 0$. Since $e \in \omega(x) \subset E$, then $\varphi_t(e) = \varphi_\tau(e) = e$ for all $t \geq 0$. So, $d\varphi_\tau(e)v(e) = \rho(e)v(e)$ for some $\rho(e) > 0$.

We then pick a smooth chart $\Phi : U \rightarrow W$, where $U \subset M$ is the coordinate neighborhood of e , W is an open set in \mathbb{R}^n . Using this coordinate map Φ , we write $\tilde{e} = \Phi(e)$, $\tilde{v}(\tilde{e}) = d\Phi(e)v(e)$ and $\tilde{C}(\tilde{e}) = d\Phi(e)C_M(e)$. Let $A = \Phi \circ \varphi_\tau \circ \Phi^{-1}$. Then $A(\tilde{e}) = \tilde{e}$ and $dA(\tilde{e}) = d\Phi(e) \circ d\varphi_\tau(e) \circ d\Phi^{-1}(\tilde{e}) : T_{\tilde{e}}\mathbb{R}^n \rightarrow T_{\tilde{e}}\mathbb{R}^n$. So, $dA(\tilde{e})\tilde{v}(\tilde{e}) = d\Phi(e) \circ d\varphi_\tau(e) \circ d\Phi^{-1}(\tilde{e})\tilde{v}(\tilde{e}) = d\Phi(e) \circ d\varphi_\tau(e)v(e) = d\Phi(e)\rho(e)v(e) = \rho(e)d\Phi(e)v(e) = \rho(e)\tilde{v}(\tilde{e})$. Since system is SDP and $\tilde{C}(\tilde{e}) = d\Phi(e)C_M(e)$, then

$dA(\tilde{e})(\tilde{C}(\tilde{e}) \setminus \{0\}) \subset \text{Int}\tilde{C}(\tilde{e})$. Moreover, $\tilde{v}(\tilde{e}) \in \text{Int}\tilde{C}(\tilde{e})$, since $v(e) \in \text{Int}C_M(e)$. By the Perron-Frobenius Theorem, we obtain that $\rho(e) = \sigma(dA(\tilde{e}))$, where $\sigma(dA(\tilde{e}))$ is the spectral radius of $dA(\tilde{e})$.

Since $\omega(x)$ is not a singleton, we obtain that there exists a sequence $e_n \in \omega(x) \cap U$ such that $e_n \neq e$ and $e_n \rightarrow e$. Let $\tilde{e}_n = \Phi(e_n)$. Then $\tilde{e} - \tilde{e}_n = A(\tilde{e}) - A(\tilde{e}_n) = dA(\tilde{e})(\tilde{e} - \tilde{e}_n) + o(|\tilde{e} - \tilde{e}_n|)$, where $o(|\tilde{e} - \tilde{e}_n|)/|\tilde{e} - \tilde{e}_n| \rightarrow 0$ as $n \rightarrow \infty$. Let $w_n = \tilde{e} - \tilde{e}_n/|\tilde{e} - \tilde{e}_n|$, then $w_n = dA(\tilde{e})w_n + r_n$, where $r_n \rightarrow 0$ as $n \rightarrow \infty$. If $w_n \rightarrow w$ as $n \rightarrow \infty$, we obtain that $w = dA(\tilde{e})w$. Hence, $\sigma(dA(\tilde{e})) \geq 1$.

If $\sigma(dA(\tilde{e})) = 1$, then $w \in \text{Int}\tilde{C}(\tilde{e})$. Hence, there exists a neighborhood \tilde{N} of w in the interior of $\tilde{C}(\tilde{e})$ such that there is a convex neighborhood \tilde{W} of \tilde{e} satisfies that for any $\tilde{z} \in \tilde{W}$, $\tilde{N} \subset \tilde{C}(\tilde{z})$. Thus, there is a $N > 0$ such that $\tilde{e}_n \in \tilde{W}$ and $w_n \in \tilde{N}$ for all $n \geq N$. Let $\tilde{\beta}(s) = s\tilde{e} + (1-s)\tilde{e}_n$ for $s \in [0, 1]$, then $\frac{d}{ds}\tilde{\beta}(s) = \tilde{e} - \tilde{e}_n \in \text{Int}\tilde{C}(\tilde{\beta}(s))$. Then, $\Phi^{-1}(\tilde{\beta}(s))$ is a conal curve in M connecting e_n and e . Thus, $e_n \leq_M e$. Since $e_n, e \in \omega(x)$, it is a contradiction to Lemma 2.1(a). Thus, we obtain that $\rho(e) > 1$. The conclusion of the lemma follows from the compactness of $\omega(x)$ and the continuity of the spectral radius (see e.g., [24, 28]). \square

Proposition 3.1. *If $x \leq_M y$, then either $\omega(x) \ll_M \omega(y)$, or $\omega(x) = \omega(y) = \{e\}$ for some $e \in E$.*

Proof. We just need to prove the case that $\omega(x) = \omega(y) = \{e\}$ for some $e \in E$ by Lemma 2.1(b). Suppose that $\omega(x) = \omega(y) = K \subset E$. Let $\tau > 0$ be the one in Lemma 3.1. Let $x_n = \varphi_{n\tau}(x)$, $y_n = \varphi_{n\tau}(y)$ for $n \geq 1$. Since $x \leq_M y$, there exists a conal curve $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = x$, $\gamma(1) = y$ and $\frac{d}{ds}\gamma(s) \in C_M(\gamma(s))$. Let $\gamma_n(s) = \varphi_{n\tau}(\gamma(s))$, then $\gamma_n(s)$ is a strictly conal curve connecting x_n and y_n such that $x_n \ll_M y_n$. Suppose that K contains more than a single element.

By passing to a subsequence if necessary, we assume that $x_n \rightarrow p$, $y_n \rightarrow q$ as $n \rightarrow \infty$, where $p, q \in K$. Thus, $p = q$. Otherwise, $p \neq q$ with $p \leq_M q$, which is a contradiction to Lemma 2.1(a). The length of γ_n is $L(\gamma_n) = \int_0^1 \left| \frac{d}{ds}\gamma_n(s) \right|_{\gamma_n(s)} ds$. We assert that $L(\gamma_n) \rightarrow 0$ as $n \rightarrow \infty$.

M is a globally orderable conal manifold, then every point in M is a strong point (see [26, Proposition 5.3]). We claim that for any open neighborhood U of p , there exists a $N > 0$ such that $\gamma_n \subset U$ for all $n \geq N$. In fact, suppose there exist a neighborhood U_1 of p and a subsequence $\{\gamma_{n_k}\}$ such that γ_{n_k} leaves U_1 for all k . Since p is a strong point, there exists $V_1 \subset U_1$ open containing p such that every conal curve that begins in V_1 and leaves U_1 terminates outside of V_1 (see [26, Lemma 5.2]). On the other hand, $\gamma_{n_k}(0) \rightarrow p$ and $\gamma_{n_k}(1) \rightarrow p$, then there exists a $k_0 > 0$ such that for $k \geq k_0$, $\gamma_{n_k}(0)$ and $\gamma_{n_k}(1)$ are in V_1 , which is a contradiction. Thus, we have the claim. Since M is locally compact, we can find an open neighborhood U of p such that \bar{U} is compact. By the previous claim, there is a N such that $\gamma_n \subset U$ for all $n \geq N$. Let $\Omega = \{z \in M : z \text{ is a limit point of a sequence } \{z_n\}, z_n \in \gamma_n\}$. Clearly, $p \in \Omega$. If there is a $q \in \Omega$ with $q \neq p$, then there exist neighborhoods W_1 of p and W_2 of q such that $W_1 \cap W_2 = \emptyset$ since M is a Hausdorff space. $q \in \Omega$ implies that there is a subsequence $\{z_{n_i}\}$ such that $z_{n_i} \rightarrow q$, where $z_{n_i} \in \gamma_{n_i}$. Then there is a $I_1 > 0$ such that for $i \geq I_1$,

$z_{n_i} \in W_2$ with $z_{n_i} \notin W_1$, which contradicts the previous claim for neighborhood W_1 of p . Hence, $\Omega = p$. If there exist a $\alpha > 0$ and a subsequence n_j such that $L(\gamma_{n_j}) \geq \alpha > 0$, then $\Omega_1 = \{z \in M : z \text{ is a limit point of a sequence } \{z_{n_j}\}, z_{n_j} \in \gamma_{n_j}\} = p$, which is a contradiction. So, $L(\gamma_n) \rightarrow 0$ as $n \rightarrow \infty$.

Since x_n is attracted to K , one can choose $e_n \in K$ such that $d(x_n, e_n) \rightarrow 0$ as $n \rightarrow \infty$. On the other hand, $L(\gamma_n) \rightarrow 0$ as $n \rightarrow \infty$. Hence, $\max_{0 \leq s \leq 1} d(\gamma_n(s), e_n) \rightarrow 0$ as $n \rightarrow \infty$. Then there exist a $\bar{N} > 0$ and a coordinate neighborhood U of p such that $\gamma_n, e_n \in U$ for all $n \geq \bar{N}$. Since there is a smooth coordinate chart $\Phi : U \rightarrow W \subset \mathbb{R}^n$, all notations in the following have coordinate representations. With the diffeomorphism Φ , we treat the following notations both in manifold M and \mathbb{R}^n .

For each $s \in [0, 1]$, we have

$$\begin{aligned}
\left| \frac{d}{ds} \gamma_{n+1}(s) \Big|_{\gamma_{n+1}(s)} \right| &= |d\varphi_\tau(\gamma_n(s)) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(\gamma_n(s))} \\
&= |\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(e_n)} \\
&\geq -|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) \frac{d}{ds} \gamma_n(s) - d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(e_n)} \\
&\quad + |d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(e_n)},
\end{aligned} \tag{3.1}$$

where Γ is defined as above.

Since $d(\gamma_n(s), e_n) \rightarrow 0$ as $n \rightarrow \infty$, and Γ is continuous with $\Gamma(x, x) = \text{Id}_x$ for any $x \in M$, then for any $\delta_1 > 0$, there is a $N_1 \geq \bar{N}$ such that for $n \geq N_1$, $\|\Gamma(\gamma_n(s), e_n) - \Gamma(e_n, e_n)\| \leq \delta_1$. Since φ_t is smooth, then for any $\delta_2 > 0$, there is a $N_2 \geq \bar{N}$ such that for $n \geq N_2$, $\|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) - \Gamma(\varphi_\tau(e_n), \varphi_\tau(e_n)) d\varphi_\tau(e_n)\| \leq \delta_2$. Let $\delta = \min\{\delta_1, \delta_2\}$ and $N = \max\{N_1, N_2\}$, then for $n \geq N$, we can obtain that $\|\Gamma(\gamma_n(s), e_n) - \Gamma(e_n, e_n)\| \leq \delta$ and $\|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) - \Gamma(\varphi_\tau(e_n), \varphi_\tau(e_n)) d\varphi_\tau(e_n)\| \leq \delta$. If there is a $L > 0$ such that $\|d\varphi_\tau(x)\| \leq L$ for all $x \in K$, then for $n \geq N$, we have that

$$\begin{aligned}
&|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) \frac{d}{ds} \gamma_n(s) - d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(e_n)} \\
&\leq \|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) - d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n)\| \cdot \left| \frac{d}{ds} \gamma_n(s) \Big|_{\gamma_n(s)} \right| \\
&\leq (\|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n)) d\varphi_\tau(\gamma_n(s)) - \Gamma(\varphi_\tau(e_n), \varphi_\tau(e_n)) d\varphi_\tau(e_n)\| \\
&\quad + \|d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n) - d\varphi_\tau(e_n) \Gamma(e_n, e_n)\|) \cdot \left| \frac{d}{ds} \gamma_n(s) \Big|_{\gamma_n(s)} \right| \\
&\leq (\delta + \delta \|d\varphi_\tau(e_n)\|) \cdot \left| \frac{d}{ds} \gamma_n(s) \Big|_{\gamma_n(s)} \right| \\
&\leq \delta(1 + L) \left| \frac{d}{ds} \gamma_n(s) \Big|_{\gamma_n(s)} \right|.
\end{aligned} \tag{3.2}$$

Next, we deal with the term $|d\varphi_\tau(e_n) \Gamma(\gamma_n(s), e_n) \frac{d}{ds} \gamma_n(s) \Big|_{\varphi_\tau(e_n)}$ in inequation (3.1). Due to Lemma 3.1, we have that $d\varphi_\tau(e_n)v(e_n) = \rho(e_n)v(e_n)$, where $\rho(e_n) > \rho > 1$ and v is the Perron-

Frobenius vector field. And we obtain that $|\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s)-v(\gamma_n(s))|_{\gamma_n(s)} \rightarrow 0$ as $n \rightarrow \infty$. Since Γ preserves the metric, $|\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s) - \Gamma(\gamma_n(s), e_n)v(\gamma_n(s))|_{e_n} \rightarrow 0$. Since $\Gamma(\cdot, \cdot)$ and $v(\cdot)$ are continuous, we have $|\Gamma(\gamma_n(s), e_n)v(\gamma_n(s)) - \Gamma(e_n, e_n)v(e_n)|_{e_n} \rightarrow 0$ as $d(\gamma_n(s), e_n) \rightarrow 0$. Hence, we obtain that $|\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s) - v(e_n)|_{e_n} \rightarrow 0$ as $n \rightarrow \infty$. Since $\|d\varphi_\tau(e_n)\|$ is bounded for $e_n \in K$, let $\epsilon_n = |d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s) - d\varphi_\tau(e_n)v(e_n)|_{\varphi_\tau(e_n)}$, then $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and we have that

$$\begin{aligned} & |d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s)|_{\varphi_\tau(e_n)} \\ &= |d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s) - d\varphi_\tau(e_n)v(e_n) + d\varphi_\tau(e_n)v(e_n)|_{\varphi_\tau(e_n)} \\ &\geq |d\varphi_\tau(e_n)v(e_n)|_{\varphi_\tau(e_n)} - |d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{1}{|\frac{d}{ds}\gamma_n(s)}\frac{d}{ds}\gamma_n(s) - d\varphi_\tau(e_n)v(e_n)|_{\varphi_\tau(e_n)} \\ &= \rho(e_n)|v(e_n)|_{e_n} - \epsilon_n \\ &> \rho - \epsilon_n, \end{aligned}$$

Thus,

$$|d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{d}{ds}\gamma_n(s)|_{\varphi_\tau(e_n)} > (\rho - \epsilon_n)|\frac{d}{ds}\gamma_n(s)|_{\gamma_n(s)}. \quad (3.3)$$

It follows from (3.1), (3.2) and (3.3) that

$$\begin{aligned} |\frac{d}{ds}\gamma_{n+1}(s)|_{\gamma_{n+1}(s)} &\geq -|\Gamma(\varphi_\tau(\gamma_n(s)), \varphi_\tau(e_n))d\varphi_\tau(\gamma_n(s))\frac{d}{ds}\gamma_n(s) - d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{d}{ds}\gamma_n(s)|_{\varphi_\tau(e_n)} \\ &\quad + |d\varphi_\tau(e_n)\Gamma(\gamma_n(s), e_n)\frac{d}{ds}\gamma_n(s)|_{\varphi_\tau(e_n)} \\ &\geq -\delta(1+L)|\frac{d}{ds}\gamma_n(s)|_{\gamma_n(s)} + (\rho - \epsilon_n)|\frac{d}{ds}\gamma_n(s)|_{\gamma_n(s)} \\ &= (\rho - \epsilon_n - \delta(1+L))|\frac{d}{ds}\gamma_n(s)|_{\gamma_n(s)}. \end{aligned}$$

Since $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and $\rho > 1$ by the Lemma 3.1, we can choose δ small enough such that there exist a $\tilde{N} > N$ and $l > 1$ satisfying for all $n \geq \tilde{N}$, $\rho - \epsilon_n - \delta(1+L) \geq l > 1$. Thus, we obtain that $|\frac{d}{ds}\gamma_{n+1}(s)|_{\gamma_{n+1}(s)} \geq l|\frac{d}{ds}\gamma_n(s)|_{\gamma_n(s)}$. So, $L(\gamma_{n+1}) \geq lL(\gamma_n)$, which is a contradiction to $L(\gamma_n) \rightarrow 0$ as $n \rightarrow \infty$. Thus, K is a singleton. \square

Lemma 3.2. *Suppose that the cone field on M admits a C^∞ -section V , then for each $x \in M$, there exist $\epsilon_x > 0$ and a conal curve $\gamma_x : (-\epsilon_x, \epsilon_x) \rightarrow M$ such that $\gamma_x(0) = x$ and $\frac{d}{ds}\gamma_x(s) = V(\gamma_x(s))$ for $s \in (-\epsilon_x, \epsilon_x)$.*

Proof. See [27, Proposition 9.2]. In fact, γ_x is the integral curve of V . \square

By the Lemma 3.2, we obtain that for any $x \in M$, there exists a conal curve passing through x . Thus, we say that x can be approximated from below (resp. above) in M (i.e., there exists a sequence $\{x_n\}$ in M such that $x_n \leq_M x_{n+1} \leq_M x$ (resp. $x \leq_M x_{n+1} \leq_M x_n$) for $n \geq 1$ and $x_n \rightarrow x$ as $n \rightarrow \infty$).

Lemma 3.3. *If $x_0 \in M$ satisfies that there exists a sequence $\{x_n\}$ such that $x_n \leq_M x_{n+1} \leq_M x_0$ for $n \geq 1$ and $x_n \rightarrow x_0$, then one of the following alternatives must occur :*

- (1) *There exists $p \in E$ such that $\omega(x_n) \ll_M \omega(x_{n+1}) \ll_M p = \omega(x_0)$ for all $n \geq 1$.*
- (2) *$\omega(x_n) = \omega(x_0) = p \in E$ for $n \geq 1$.*
- (3) *There exists $p \in E$ such that $\omega(x_n) = p \ll_M \omega(x_0)$ for all $n \geq 1$.*

Proof. Let $x_0 \in M$ have the property that it is approximated from below in M by a sequence $\{x_n\}$ such that $x_n \leq_M x_{n+1} \leq_M x_0$ for $n \geq 1$ and $x_n \rightarrow x_0$. By Proposition 3.1, either there exists a $N > 0$ such that $\omega(x_n) = \omega(x_m)$ for all $n, m \geq N$, or there is a subsequence $\{x_{n_i}\}$ such that $\omega(x_{n_i}) \ll_M \omega(x_{n_{i+1}})$ for all $i \geq 1$. Thus, we assume that either $\omega(x_n) = \omega(x_m)$ for all n, m , or $\omega(x_n) \ll_M \omega(x_{n+1})$, where $x_n \rightarrow x_0$.

Suppose that $\omega(x_n) \ll_M \omega(x_{n+1})$ for $n \geq 1$ and $x_n \rightarrow x_0$ as $n \rightarrow \infty$. We first obtain that $\omega(x_n) \ll_M \omega(x_0)$ for all $n \geq 1$. In fact, if there exists $n_0 > 0$ such that $\omega(x_{n_0}) = \omega(x_0)$, then $\omega(x_n) = \omega(x_0)$ for all $n \geq n_0$, which is a contradiction. Let $\Omega = \{y : y = \lim_{n \rightarrow \infty} y_n, y_n \in \omega(x_n)\}$. By the compactness condition (P), the set Ω belongs to a compact set $\overline{\bigcup_{n \geq 0} \omega(x_n)}$. Furthermore, Ω is nonempty. If $\{y_{n_k}\}$ and $\{y_{m_k}\}$ are two subsequences of $\{y_n\}$ such that $y_{n_k} \rightarrow p$ and $y_{m_k} \rightarrow q$ as $k \rightarrow \infty$. Since $y_n \in \omega(x_n)$, then for each k , there exists $l(k)$ such that $y_{n_k} \ll_M y_{m_{l(k)}}$. Thus, $p \leq_M q$. A similar argument shows that $q \leq_M p$, then $p = q$. So, $\lim_{n \rightarrow \infty} y_n = p$ and $p \in \Omega$. Thus, Ω is nonempty. Suppose that $u, w \in \Omega$ have the property that there exist two sequences $\{u_n\}, \{w_n\}$ with $u_n, w_n \in \omega(x_n)$ and $u_n \rightarrow u, w_n \rightarrow w$ as $n \rightarrow \infty$. Since $u_n \ll_M w_{n+1}$ (resp. $w_n \ll_M u_{n+1}$), we obtain that $u \leq_M w$ (resp. $w \leq_M u$). Then $u = w$. Thus, Ω is a singleton. On the other hand, since each $\omega(x_n)$ is invariant, we obtain that Ω is positively invariant. So, $\Omega = \{p\} \in E$. It is easy to see from the arbitrariness of y_n in $\omega(x_n)$ that $\lim_{n \rightarrow \infty} \text{dist}(\omega(x_n), p) = 0$. Since $\omega(x_n) \ll_M \omega(x_0)$, then $p \leq_M \omega(x_0)$. If $p \in \omega(x_0)$, then $\omega(x_0) = p$ by Lemma 2.1(a). Thus, x_0 is a convergent point and (1) holds. If $p \notin \omega(x_0)$, we will get a contradiction. Since $p \in E$ and $\omega(x_0)$ is invariant, we obtain that $p \ll_M \omega(x_0)$. For each $z \in \omega(x_0)$, there exist $t_z > 0$, a neighborhood U_z of p and a neighborhood V_z of z such that $\varphi_t(U_z) \ll_M \varphi_t(V_z)$ for all $t \geq t_z$. Since $\{V_z\}$ is an open cover of $\omega(x_0)$, we obtain that $\omega(x_0) \subset \bigcup_{i=1}^n V_{z_i} = V$, where $z_i \in \omega(x_0)$. Meanwhile, $U = \bigcap_{i=1}^n U_{z_i}$ is a neighborhood of p . Let $t_0 = \max_{1 \leq i \leq n} \{t_{z_i}\}$, then $\varphi_t(U) \ll_M \varphi_t(V)$ for $t \geq t_0$. On the other hand, there exists $t_1 > 0$ such that $\varphi_{t_1}(x_0) \in V$. Since $x_n \rightarrow x_0$, there is a $N_1 > 0$ such that $\varphi_{t_1}(x_{N_1}) \in V$. By $p \in E$, we have that $p \ll_M \varphi_t(x_{N_1})$ for $t > t_0 + t_1$. Thus, $p \leq_M \omega(x_{N_1})$. By the definition of Ω , $\omega(x_n) \ll_M \omega(x_{n+1}) \leq_M p$ for all $n \geq 1$. Thus, $\omega(x_{N_1}) = p$ and $\omega(x_n) = p$ for all $n > N_1$, which is a contradiction. Hence, we have proved the case (1).

Suppose that $\omega(x_n) = \omega(x_m)$ for all n, m . By Proposition 3.1, $\omega(x_n) = p \in E$ for all $n > 1$. Since $x_n \leq_M x_0$, then $\omega(x_n) = \omega(x_0)$ or $\omega(x_n) \ll_M \omega(x_0)$. So, (2) and (3) hold. \square

Remark 3.2. An analogous result holds if x_0 is approximated from above.

Now, we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. Suppose that $x_0 \in M \setminus \text{Int}C$, then we will prove that $x_0 \in \overline{\text{Int}C}$. In fact, there is a sequence $\{x_n\} \subset M \setminus C$ such that $x_n \rightarrow x_0$. Since the cone field admits a C^∞ -section, then all x_n can be approximated from below and above in M . Without loss of generality, we assume that for each x_n , there exists a sequence $\{z_m^n\}$ such that $z_m^n \leq_M z_{m+1}^n \leq_M x_n$ for $m \geq 1$ and $z_m^n \rightarrow x_n$ as $m \rightarrow \infty$. Since $\{x_n\} \subset M \setminus C$ for all n , then the case (3) of Lemma 3.3 holds for each sequence $\{z_m^n\}$.

We claim that $x_n \in \overline{\text{Int}C}$ for each n . If the claim holds, then $x_0 \in \overline{\text{Int}C}$ since $x_n \rightarrow x_0$ and $\overline{\text{Int}C}$ is a closed set. Thus, we obtain the theorem.

Now, it suffices to prove the claim. For each n , $z_m^n \rightarrow x_n$ as $m \rightarrow \infty$ and $\{x_n\} \subset M \setminus C$, then there exists a $p \in E$ such that $\omega(z_m^n) = p \ll_M \omega(x_n)$ for all $m \geq 1$. For $y \in \omega(x_n)$, there exist a neighborhood W_y of y and $t_y > 0$ such that $p \ll_M \varphi_t(W_y)$ for $t \geq t_y$. Since $\{W_y\}$ is an open cover of $\omega(x_n)$, we obtain that $\omega(x_n) \subset \bigcup_{i=1}^k W_{y_i} = W$, where $y_i \in \omega(x_n)$. Let $t_0 = \max_{1 \leq i \leq k} \{t_{y_i}\}$, then $p \ll_M \varphi_t(W)$ for $t \geq t_0$. On the other hand, there is a $t_1 > 0$ such that $\varphi_{t_1}(x_n) \in W$. Furthermore, one can choose a neighborhood U of x_n such that $\varphi_{t_1}(U) \subset W$. If $x \in U$ with $x \leq_M x_n$, then there exist a neighborhood V of x , $V \subset U$, a neighborhood O of x_n and $t_2 > 0$ such that $\varphi_t(V) \ll_M \varphi_t(O)$ for $t \geq t_2$. We can choose a $L > 0$ such that $z_L^n \in O$, then we have $\varphi_t(V) \ll_M \varphi_t(z_L^n)$ for $t \geq t_2$. Since $V \subset U$, $\varphi_{t_1}(U) \subset W$ and $p \ll_M \varphi_t(W)$ for $t \geq t_0$, we obtain that $p \ll_M \varphi_{t+t_1}(V)$ for $t \geq t_0$. Thus, $p \ll_M \varphi_t(V) \ll_M \varphi_t(z_L^n)$ for $t \geq t_0 + t_1 + t_2$. Since $\omega(z_m^n) = p$ for all $m \geq 1$, then $\omega(s) = p \in E$ for all $s \in V$. Hence, $x \in \text{Int}C$. Since the cone field admits a C^∞ -section, then there is a sequence $\{u_m^n\}$ in U such that $u_m^n \leq_M x_n$ and $u_m^n \rightarrow x_n$ as $m \rightarrow \infty$. By the previous proof, $u_m^n \in \text{Int}C$. Thus, $x_n \in \overline{\text{Int}C}$. And hence, we have proved the claim. \square

4 Proof of Lemma 2.1

In this section, we focus on proving the critical Lemma 2.1. We first need the following proposition.

Proposition 4.1. *If $x \leq_M \varphi_T(x)$ for some $T > 0$, then $\varphi_t(x) \rightarrow p \in E$ as $t \rightarrow \infty$.*

Proof. If $x \leq_M \varphi_T(x)$ and Σ is SDP, then for $t_0 > 0$, there exist neighborhoods U of x , V of $\varphi_{t_0}(x)$ such that $\varphi_{t_0}(U) \ll_M \varphi_{t_0}(V)$ by Proposition 2.3.

By the continuity, there exists $0 < \epsilon_0 < t_0$ such that $\varphi_{t_0}(x) \ll_M \varphi_{t_0+T+\epsilon}(x)$ for $\epsilon \in (-\epsilon_0, \epsilon_0)$. We first claim that $\omega(x)$ is a τ -periodic orbit for any $\tau \in (T - \epsilon_0, T + \epsilon_0)$. We just prove the case of $\tau = T$ and a similar argument for $\tau \in (T - \epsilon_0, T + \epsilon_0)$. If $x \leq_M \varphi_T(x)$, then $\varphi_{nT}(x) \ll_M \varphi_{(n+1)T}(x)$ for $n = 1, 2, \dots$. Thus, there exists $p \in M$ such that $\varphi_{nT}(x) \rightarrow p$ as $n \rightarrow \infty$. In fact, if there exist two sequences $\varphi_{n_k T}(x)$ and $\varphi_{n_l T}(x)$ satisfying $\varphi_{n_k T}(x) \rightarrow p$ and $\varphi_{n_l T}(x) \rightarrow q$, then for each k , there is a $l(k)$ such that $\varphi_{n_k T}(x) \ll_M \varphi_{n_{l(k)} T}(x)$. Thus,

$p \leq_M q$ by (H2). A similar argument shows that $q \leq_M p$. So, $p = q$ by (H1). Consider the orbit of p , $\varphi_{t+T}(p) = \varphi_{t+T}(\lim_{n \rightarrow \infty} \varphi_{nT}(x)) = \lim_{n \rightarrow \infty} \varphi_{(n+1)T+t}(x) = \varphi_t(p)$ for all $t \geq 0$. Thus, $O(p)$ is a T -periodic orbit. Suppose that there exist $q \in M$ and $\{t_j\}$ such that $t_j \rightarrow \infty$ and $\varphi_{t_j}(x) \rightarrow q$ as $j \rightarrow \infty$. For each j , $t_j = n_j T + s_j$, $n_j \in \{0, 1, 2, \dots\}$, $0 \leq s_j < T$, then $\varphi_{t_j}(x) = \varphi_{n_j T + s_j}(x) = \varphi_{s_j}(\varphi_{n_j T}(x))$. Thus, $\varphi_{t_j}(x) = \varphi_{s_j}(\varphi_{n_j T}(x)) \rightarrow \varphi_s(p)$ as $j \rightarrow \infty$ where $s \in [0, T]$ such that $s_j \rightarrow s$ as $j \rightarrow \infty$. So, $\omega(x) = O(p)$.

Since $\omega(x)$ is a τ -periodic orbit for all $\tau \in (T - \epsilon_0, T + \epsilon_0)$ and $\omega(x) = O(p)$, then $\varphi_{t+\tau}(p) = \varphi_t(p)$ for all $t \geq 0$ (i.e., $O(p)$ is τ -periodic). Let P be the set of all periods of $\varphi_t(p)$. It is easy to see that $(T - \epsilon_0, T + \epsilon_0) \subset P$ and $\varphi_{t+s}(p) = \varphi_t(\varphi_s(p)) = \varphi_t(\varphi_{s+T}(p)) = \varphi_t(p)$ for all $s \in [0, \epsilon_0)$ and $t \geq 0$. Thus, $[0, \epsilon_0) \subset P$. We next prove that $T + \epsilon_0 \in P$. In fact, let $\epsilon \in (0, \epsilon_0)$ and $t = \epsilon_0 - \epsilon \in (0, \epsilon_0)$, then $\varphi_{T+\epsilon_0}(p) = \varphi_{T+\epsilon+(\epsilon_0-\epsilon)}(p) = \varphi_{T+\epsilon}(\varphi_t(p)) = \varphi_{T+\epsilon}(p) = p$. Thus, $T + \epsilon_0 \in P$. Then $[0, \epsilon_0] \subset P$.

For each $t > 0$, $t = n\epsilon_0 + \tau$, where $n \in \{0, 1, 2, \dots\}$, and $0 \leq \tau < \epsilon_0$. $\varphi_t(p) = \varphi_{n\epsilon_0+\tau}(p) = p$. Thus, $p \in E$ and $\omega(x) = p$. \square

Now, we prove Lemma 2.1(a).

Proof of Lemma 2.1(a). If there exist two points $x, y \in \omega(z)$ with $x \leq_M y$, then we can find a neighborhood U of x , a neighborhood V of y for $t_0 > 0$ such that $\varphi_{t_0}(U) \ll_M \varphi_{t_0}(V)$ by Proposition 2.3. Since $x, y \in \omega(z)$, there exist $0 < t_1 < t_2$ such that $\varphi_{t_1}(z) \in U$ and $\varphi_{t_2}(z) \in V$. Then $\varphi_{t_0+t_1}(z) \ll_M \varphi_{t_0+t_2}(z) = \varphi_{t_2-t_1}(\varphi_{t_0+t_1}(z))$. By Proposition 4.1, $\varphi_t(z) \rightarrow p \in E$. So, $\omega(z) = p$, which is a contradiction. \square

In order to prove Lemma 2.1(b), we further need several propositions.

Proposition 4.2. *If $x \leq_M y$, $t_k \rightarrow \infty$, $\varphi_{t_k}(x) \rightarrow p$ and $\varphi_{t_k}(y) \rightarrow p$ as $k \rightarrow \infty$, then $p \in E$.*

Proof. Since Σ is SDP, there exist a neighborhood U of x , a neighborhood V of y for $t_0 > 0$ such that $\varphi_{t_0}(U) \ll_M \varphi_{t_0}(V)$ by Proposition 2.3. Let $\delta > 0$ be small such that $\{\varphi_l(x) : 0 \leq l \leq \delta\} \subset U$ and $\{\varphi_l(y) : 0 \leq l \leq \delta\} \subset V$. Then $\varphi_s(x) = \varphi_{t_0}(\varphi_{s-t_0}(x)) \ll_M \varphi_{t_0}(y)$ for any $t_0 \leq s \leq t_0 + \delta$. Thus, $\varphi_{t_k-t_0}(\varphi_s(x)) \ll_M \varphi_{t_k-t_0}(\varphi_{t_0}(y))$ for k large enough. Let $r = s - t_0$ and $k \rightarrow \infty$, we obtain $\varphi_r(p) \leq_M p$ for all $r \in [0, \delta]$. As a similar argument, we obtain $p \leq_M \varphi_r(p)$ for all $r \in [0, \delta]$. So, $p = \varphi_r(p)$ for all $r \in [0, \delta]$. For any $t > 0$, we write $t = n\delta + \tau$, where $\tau \in [0, \delta)$ and $n \in \{0, 1, 2, \dots\}$. So, $\varphi_t(p) = \varphi_{n\delta+\tau}(p) = p$. Thus, $p \in E$. \square

Proposition 4.3. *If $x \leq_M y$, then $\omega(x) \cap \omega(y) \subset E$.*

Proof. Let $p \in \omega(x) \cap \omega(y)$, there exists a sequence $\{t_k\}$ such that $t_k \rightarrow \infty$ and $\varphi_{t_k}(x) \rightarrow p$ as $k \rightarrow \infty$. By a subsequence, we assume that $\varphi_{t_k}(y) \rightarrow q$ as $k \rightarrow \infty$. Since $x \leq_M y$ and Σ is SDP, we obtain $p \leq_M q$ and $p, q \in \omega(y)$. The Lemma 2.1(a) means $p = q$ and Proposition 4.2 implies $p \in E$. \square

Lemma 4.1. *Let $x \leq_M y$, $p \in \omega(x)$, $q \in \omega(y)$ and $p \ll_M q$. If p (or q) is an equilibrium, then $\omega(x) \ll_M \omega(y)$.*

Proof. Assume that $p \in E$. Since $q \in \omega(y)$ and $p \ll_M q$, there exists a sequence $\{t_k\} \rightarrow \infty$ such that $\varphi_{t_k}(y) \rightarrow q$ and hence, there exists a $k(q) > 0$ such that $p \ll_M \varphi_{t_k(q)}(y)$. Thus, $p = \varphi_t(p) \ll_M \varphi_t(\varphi_{t_k(q)}(y))$ for all $t > 0$. Thus, $p \leq_M \omega(y)$. By Lemma 2.1(a), $p \notin \omega(y)$. Moreover, $p \ll_M \omega(y)$. In fact, since $\omega(y)$ is invariant, for any $z \in \omega(y)$, there exists $T > 0$ such that $\varphi_{-T}(z) \in \omega(y)$ and $p \leq_M \varphi_{-T}(z)$. Thus, $p \ll_M z$.

Since $p \in \omega(x)$, there exists a sequence $\{t_l\} \rightarrow \infty$ such that $\varphi_{t_l}(x) \rightarrow p$. Since $p \ll_M \omega(y)$, there exists a $l(p)$ such that $\varphi_{t_l(p)}(x) \ll_M \omega(y)$. Thus, $\varphi_t(\varphi_{t_l(p)}(x)) \ll_M \varphi_t(\omega(y))$ for all $t > 0$. Since $\omega(y)$ is compact and invariant, we obtain that $\omega(x) \leq_M \omega(y)$. By Lemma 2.1(a), $\omega(x) \cap \omega(y) = \emptyset$. Since $\omega(x), \omega(y)$ are compact and invariant, then $\omega(x) \ll_M \omega(y)$. In fact, let $a \in \omega(x)$ and $b \in \omega(y)$, there is a $T_1 > 0$ such that $\varphi_{-T_1}(a) \in \omega(x)$ and $\varphi_{-T_1}(b) \in \omega(y)$ with $\varphi_{-T_1}(a) \leq_M \varphi_{-T_1}(b)$, then $a \ll_M b$.

A similar argument is used if $q \in E$. □

Lemma 4.2. *Assume that $K \subset M$ is a compact set in which the flow φ_t is SDP. Then there exists $\delta > 0$ with the following property. Let ψ_t denote the flow of a C^1 vector field g such that $g \in U_\delta(f)$, where $U_\delta(f)$ is a δ -neighborhood of f in space of C^1 vector fields on M with the C^1 topology. Then there exists $t_0 > 0$ such that if K is positively invariant under the flow of ψ_t , then ψ_t is SDP for all $t \geq t_0$.*

Proof. We first assume that $t \in [t_0, 2t_0]$, where $t_0 > 0$ is fixed. Since Σ is SDP, we obtain that $d\varphi_t(x)v \in \text{Int}C_M(\varphi_t(x))$ for all $v \in C_M(x)$ with $|v|_x = 1$. With a coordinate chart, we treat the following notations both in manifold and \mathbb{R}^n . By the continuity of the cone field, there exist neighborhoods U of $\varphi_t(x)$ and V of $d\varphi_t(x)v$ such that $V \subset C_M(y)$ for all $y \in U$. So, there exists $\delta > 0$ such that for $g \in U_\delta(f)$, $\psi_t(x) \in U$ and $d\psi_t(x)v \in V$ (see e.g., [40, 41]). Thus, $d\psi_t(x)v \in \text{Int}C_M(\psi_t(x))$ for $t \in [t_0, 2t_0]$, $v \in C_M(x) \setminus \{0\}$.

For $t \geq 2t_0$, let us write $t = r + kt_0$, where $r \in [t_0, 2t_0)$ and $k \in \{1, 2, \dots\}$. Define $x_j = \psi_{jt_0}(x)$, $j = 1, 2, \dots, k$. It is clear that $x_j \in K$ if K is positively invariant. Then $d\psi_t(x)v = d\psi_r(x_k)d\psi_{t_0}(x_{k-1}) \cdots d\psi_{t_0}(x)v$. By the preceding proof, $d\psi_t(x)v \in \text{Int}C_M(\psi_t(x))$ for $t > 2t_0$.

Thus, we have proved that ψ_t is SDP for all $t \geq t_0$. □

Lemma 4.3. *Let $x \leq_M y$, $p \in \omega(x)$, $q \in \omega(y)$ and $p \ll_M q$. If p (or q) belongs to a periodic orbit, then $\omega(x) \ll_M \omega(y)$.*

Proof. Assume that $p \in \gamma$, γ is a periodic orbit and q is not an equilibrium (the other case is similar).

If $\gamma \cap \omega(y) \neq \emptyset$, then $p \in \omega(y)$. Thus, $p \in E$ by Proposition 4.3, which is a contradiction. Thus, $\omega(y) \subset M \setminus \gamma$. Since $q \in \omega(y)$, then the orbit closure of q is in the compact set $\omega(y) \subset M \setminus \gamma$.

By the Closing Lemma (see e.g., [20, 47]), there is a C^1 vector field g whose flow ψ_t has a closed orbit β_g passing through q . Moreover, g can be chosen to C^1 approximate f as closely as

desired and to coincides with f outside a given neighborhood U of the orbit closure of q with respect to f such that $U \cap \gamma = \emptyset$. Thus, ψ_t is eventually SDP by Lemma 4.2 and γ is also a closed orbit of ψ_t . In the following, we write γ_f (resp. γ_g) as the orbit generated by f (resp. g). So, $\gamma = \gamma_f = \gamma_g$.

For the system generated by vector field g , γ_g and β_g are two periodic orbits and $p \in \gamma_g$, $q \in \beta_g$ with $p \ll_M q$. Then $\gamma_g \ll_M q$.

In fact, if $p \ll_M q$, then there is a $\epsilon_0 > 0$ such that $\psi_\epsilon(p) \ll_M q$ for all $\epsilon \in [0, \epsilon_0]$. Let $T_1, T_2 > 0$ be the periods of γ_g and β_g and $a = \frac{T_2}{T_1}$. If a is a rational number, let $a = \frac{n_1}{n_2}$, $n_1, n_2 \in \{1, 2, \dots\}$, then $n_1 T_1 = n_2 T_2$. Define $\epsilon_1 = \sup\{\epsilon > 0 : \psi_\tau(p) \ll_M q \text{ for all } \tau \in [0, \epsilon]\}$. Suppose that $\epsilon_1 < \infty$. Then $\psi_{\epsilon_1}(p) \leq_M q$. Since $\psi_{\epsilon_1}(p) = \psi_{n_1 T_1}(\psi_{\epsilon_1}(p)) = \psi_{n_2 T_2}(\psi_{\epsilon_1}(p)) \ll_M \psi_{n_2 T_2}(q) = q$, one has $\psi_{\epsilon_1}(p) \ll_M q$. Thus, there exists $\delta > 0$ such that $\psi_{\epsilon_1 + \epsilon}(p) \ll_M q$ for all $\epsilon \in [0, \delta]$, which is a contradiction. Thus, $\psi_\epsilon(p) \ll_M q$ for all $\epsilon \geq 0$. So, $\gamma_g \ll_M q$. If a is an irrational number and $T_2 = a T_1$, the set $W = \{\psi_{n T_2}(\psi_\epsilon(p)) : n = 1, 2, \dots\}$ is dense in γ_g for any fixed $\epsilon \in [0, \epsilon_0]$. For any $z \in \gamma_g$, there exists a sequence $\{z_i\} \subset W$ such that $z_i \ll_M q$ and $z_i \rightarrow z$ as $i \rightarrow \infty$. Thus, $\gamma_g \leq_M q$. So, $\gamma_g = \psi_{T_2}(\gamma_g) \ll_M q$.

For system Σ , we obtain that $\gamma \ll_M q$. Since $q \in \omega(y)$, then there is a sequence $\{t_k\} \rightarrow \infty$ such that $\varphi_{t_k}(y) \rightarrow q$ as $k \rightarrow \infty$. Then there exists $k(q) > 0$ such that $\gamma \ll_M \varphi_{t_{k(q)}}(y)$. Thus, we obtain that $\gamma \ll_M \varphi_t(\varphi_{t_{k(q)}}(y))$ for all $t > 0$. So, $\gamma \leq_M \omega(y)$. On the other hand, $\gamma \cap \omega(y) = \emptyset$ and $\gamma, \omega(y)$ are invariant, then $\gamma \ll_M \omega(y)$. Since $\gamma \subset \omega(x)$, in a similar way we can obtain that $\omega(x) \ll_M \omega(y)$. \square

Proposition 4.4. *Let $x \leq_M y$, $p \in \omega(x)$, $q \in \omega(y)$ and $p \ll_M q$. Then $\omega(x) \ll_M \omega(y)$.*

Proof. If $p \in E$, then $\omega(x) \ll_M \omega(y)$ by Lemma 4.1. In the following, we assume that p is not an equilibrium. Thus, $p \notin \omega(y)$.

By the Closing Lemma (see e.g., [20, 47]), there exists $T > 0$ such that for any $\epsilon > 0$, we can choose a vector field g such that $g = f$ outside the ϵ -neighborhood N_ϵ of the set $\{\varphi_t(p) : |t| \leq T\}$. Moreover, ψ_t with respect to the vector field g has a closed orbit γ_g passing through p and ψ_t is eventually SDP. For small ϵ , $\omega(y) \subset M \setminus N_\epsilon$. It is easy to see that there exists a y_0 such that $p \leq_M y_0$ and $\omega_g(y_0) = \omega(y)$, where $\omega_g(y_0) = \{z \in M : \text{there exists a sequence } t_k \rightarrow \infty \text{ such that } \psi_{t_k}(y_0) \rightarrow z\}$. In fact, since $p \ll_M q$, there is a neighborhood U_q of q such that $U_q \cap N_\epsilon = \emptyset$ and $p \ll_M U_q$. Then there is a $t_q > 0$ such that $\varphi_{t_q}(y) \in U_q$. Let $y_0 = \varphi_{t_q}(y)$, then $p \ll_M y_0$ and $\omega_g(y_0) = \omega(y)$ since $g = f$ outside N_ϵ with $\omega(y) \subset M \setminus N_\epsilon$. Since $p \in \gamma_g$, $q \in \omega_g(y_0)$ and $p \ll_M q$, then $\gamma_g = \omega_g(p) \ll_M \omega_g(y_0)$ by Lemma 4.3. Thus, $p \ll_M \omega(y)$. Since $p \in \omega(x)$, there exists a sequence $\{t_k\} \rightarrow \infty$ such that $\varphi_{t_k}(x) \rightarrow p$. Then there exists $k(p) > 0$ such that $\varphi_{t_{k(p)}}(x) \ll_M \omega(y)$. Thus, we obtain that $\varphi_t(\varphi_{t_{k(p)}}(x)) \ll_M \varphi_t(\omega(y))$ for all $t > 0$. So, $\omega(x) \leq_M \omega(y)$. Since $\omega(x)$ and $\omega(y)$ are nonordering invariant sets and Σ is SDP, then $\omega(x) \cap \omega(y) = \emptyset$ and $\omega(x) \ll_M \omega(y)$. \square

Now, we are ready to prove Lemma 2.1(b).

Proof of Lemma 2.1(b). If $p \in \omega(x)$, then there exists a sequence $t_k \rightarrow \infty$ such that $\varphi_{t_k}(x) \rightarrow p$. By passing to a subsequence if necessary, we assume that $\varphi_{t_k}(y) \rightarrow q \in \omega(y)$. Since Σ is SDP and $x \leq_M y$, we obtain that $p \leq_M q$. If $p \neq q$, then $\varphi_t(p) \ll_M \varphi_t(q)$ for any $t > 0$. Thus, $\omega(x) \cap \omega(y) = \emptyset$ and $\omega(x) \ll_M \omega(y)$ by Proposition 4.4.

If $p = q$, then $p = q \in E$ by Proposition 4.2. If $\omega(x) = \omega(y)$, then $\omega(x) = \omega(y) \subset E$ by Proposition 4.3. If $\omega(x) \neq \omega(y)$, then there exists $p_1 \in \omega(x) \setminus \omega(y)$ such that there is a sequence $t_l \rightarrow \infty$ with $\varphi_{t_l}(x) \rightarrow p_1$. Therefore, there is $q_1 \in \omega(y)$ such that $\varphi_{t_l}(y) \rightarrow q_1$ by a subsequence if necessary. Thus, $p_1 \neq q_1$ and $p_1 \leq_M q_1$. So, $\varphi_t(p_1) \ll_M \varphi_t(q_1)$ for any $t > 0$ and hence, we obtain that $\omega(x) \cap \omega(y) = \emptyset$ and $\omega(x) \ll_M \omega(y)$ by Proposition 4.4, which is a contradiction. \square

A Appendix

A.1 Order structures on space-times

A *Lorentz metric* g for a smooth manifold M of dimension four is a smooth nondegenerate symmetric tensor field of type $(0,2)$ on M such that for each $p \in M$, by suitable choice of the basis, $g|_p$ has the matrix $\text{diag}(+1, +1, +1, -1)$. A *space-time* (M, g) is a connected C^∞ Hausdorff manifold M of dimension four with a Lorentz metric g .

Remark A.1. Although the arguments refer to 4-dimensional space-times, the results can be extended to a space-time of $n(\geq 2)$ -dimensions. See [2, 12, 43].

Let (M, g) be a space-time, a vector $v \in T_p M$ is said to be *timelike*, *null*, *spacelike* according to whether $g(v, v)$ is negative, zero, or positive, respectively. The non-spacelike (i.e., timelike and null) vectors in $T_p M$ form two so-called *Lorentzian cones* C and $-C$ (see e.g., [12]). Furthermore, the timelike vectors form the interior of the Lorentzian cones. See Fig.A.1.




Fig A.1: The timelike vectors, null vectors and spacelike vectors in $T_p M$.

A space-time (M, g) is said to be *time-orientable* if M admits a continuous Lorentzian cone field C_M , which is generated by Lorentz metric g .

In the remained of this subsection, we assume that space-time (M, g) is time-orientable. Thus, M is a conal manifold with respect to a continuous Lorentzian cone field C_M . In such situation, the non-spacelike vectors, which belong to cone field C_M , are called future directed.

A *non-spacelike curve* is a continuous piecewise smooth curve whose tangent vector is future directed non-spacelike. A *timelike curve* is a continuous piecewise smooth curve whose tangent vector is future directed timelike. Thus, in a time-orientable space-time, a non-spacelike (resp., timelike) curve is a conal (resp., strictly conal) curve and vice versa. The order “ \leq_M ” (resp., “ \ll_M ”) on M is well-defined by the non-spacelike (resp., timelike) curves.

A non-spacelike curve is also known as *causal curve*. In general relativity, each point of manifold M corresponds to an event. And a signal can be sent from p to q if there is a (future directed) causal curve from p to q . Thus, closed causal curves generate paradoxes involving causality (i.e., violate causality). As a result, we assume that the space-time (M, g) is *causal*, i.e., (M, g) contain no closed non-spacelike (conal) curves (see e.g., [2, 12, 13, 43]).

For a given point $p \in M$, the *chronological future* $I^+(p)$, *chronological past* $I^-(p)$, *causal future* $J^+(p)$, and *causal past* $J^-(p)$ of p are defined as follows:

- $I^+(p) = \{q \in M : p \ll_M q\}$; $I^-(p) = \{q \in M : q \ll_M p\}$;
- $J^+(p) = \{q \in M : p \leq_M q\}$; $J^-(p) = \{q \in M : q \leq_M p\}$.

Remark A.2. In some articles, the set $J^+(p)$ would be called *forward set* or *reachable set* from p and be written as $\uparrow p$; the set $J^-(p)$ would be called *backward set* or *controllable set* from p and be written as $\downarrow p$.

Remark A.3. For any $p \in M$, $I^\pm(p)$ are open by Proposition 2.1.

I^+ is said to be *inner continuous* at $p \in M$ if each compact set $K \subset I^+(p)$, there exists a neighborhood $U(p)$ of p such that $K \subset I^+(q)$ for each $q \in U(p)$. I^+ is said to be *outer continuous* at $p \in M$ if each compact set $K \subset M \setminus \overline{I^+(p)}$, there exists a neighborhood $U(p)$ of p such that $K \subset M \setminus \overline{I^+(q)}$ for each $q \in U(p)$. The *inner* and *outer continuity* of I^- are defined dually. See [13].

Proposition A.1. For any $p \in M$, $I^\pm(p)$ are inner continuous.

Proof. Suppose $K \subset I^-(p)$ is compact. For any $x \in I^-(p)$, i.e., $x \ll_M p$, there is a strictly conal curve γ such that $\gamma(0) = x$ and $\gamma(1) = p$. Let $w = \gamma(\frac{1}{2})$, then $x \ll_M w \ll_M p$, i.e., $w \in I^-(p)$ and $x \in I^-(w)$. Since $I^-(w)$ is open, then $I^-(w)$ is an open neighborhood of x . Thus, $\{I^-(w) : w \in I^-(p)\}$ is an open covering of K . Since K is a compact set, we choose w_1, w_2, \dots, w_n to determine a finite subcovering. On the other hand, $w_i \in I^-(p)$ implies that $p \in I^+(w_i)$, $i = 1, 2, \dots, n$. So, $U = \bigcap_{i=1}^n I^+(w_i)$ is an open neighborhood of p . For any

$u \in U$, $u \in I^+(w_i)$, $i = 1, 2, \dots, n$. Then $w_i \in I^-(u)$. For any $y \in I^-(w_i)$, since $y \ll_M w_i$ and $w_i \ll_M u$, then $y \ll_M u$, i.e., $y \in I^-(u)$. So, $I^-(w_i) \subset I^-(u)$ for $i = 1, 2, \dots, n$. Thus, $K \subset \bigcup_{i=1}^n I^-(w_i) \subset I^-(u)$. So, we have proved that $I^-(p)$ is inner continuous.

A similar argument is used for $I^+(p)$. □

Proposition A.2. (1) $x \ll_M y$, $y \leq_M z$ implies $x \ll_M z$,

(2) $x \leq_M y$, $y \ll_M z$ implies $x \ll_M z$.

Proof. See [12, p183] and [43, Proposition 2.18]. □

Remark A.4. In [3, 10], Proposition A.2 is called *push-up lemma*. The push-up lemma fails in the case where the spacetime metric (i.e., Lorentz metric) is continuous, see [3, Example 1.11] or [10].

Proposition A.3. $I^\pm(p) \subset J^\pm(p) \subset \overline{I^\pm(p)} = \{q \in M : I^\pm(q) \subset I^\pm(p)\}$.

Proof. It is clear that $I^+(p) \subset J^+(p)$. See [43, Proposition 3.3] for $\overline{I^+(p)} = \{q \in M : I^+(q) \subset I^+(p)\}$ and [43, Proposition 3.9] for $J^+(p) \subset \overline{I^+(p)}$. □

Theorem A.1. For any $p \in M$, if $J^\pm(p)$ are closed and $I^\pm(p)$ are outer continuous, then the conal order " \leq_M " is quasi-closed.

Proof. If $x_n \ll_M y_n$ for all n and $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$, we just need to prove that $y \in J^+(x) = \overline{I^+(x)}$ by Proposition A.3. Suppose that $y \notin \overline{I^+(x)}$, i.e., $y \in M \setminus \overline{I^+(x)}$. Then there exists a compact set K containing y such that $K \subset M \setminus \overline{I^+(x)}$. Since $y_n \rightarrow y$ as $n \rightarrow \infty$, there is a $N_1 > 0$ such that $y_n \in K$ for all $n > N_1$. On the other hand, I^+ is outer continuous at x , then there exists a neighborhood U of x such that for any $z \in U$, $K \subset M \setminus \overline{I^+(z)}$. Since $x_n \rightarrow x$ as $n \rightarrow \infty$, there is a $N_2 > 0$ such that $x_n \in U$ and $K \subset M \setminus \overline{I^+(x_n)}$ for all $n > N_2$. Let $N = \max\{N_1, N_2\}$. If $n > N$, then $x_n \in U$ and $y_n \in K$ with $y_n \in I^+(x_n)$, which is a contradiction. Thus, $y \in J^+(x) = \overline{I^+(x)}$. □

Proposition A.4. The following conditions are equivalent.

(A) For all p and q in M , $I^+(p) \subset I^+(q)$ if and only if $I^-(q) \subset I^-(p)$;

(B) For all p and q in M , $p \in \overline{J^+(q)}$ if and only if $q \in \overline{J^-(p)}$.

Proof. The conclusion can be immediately obtained with the Proposition A.3. See also [13, Proposition 1.3]. □

Proposition A.5. If any one of the equivalent conditions in Proposition A.4 holds, then $I^\pm(p)$ are outer continuous for any $p \in M$.

Proof. We first assert that for $v \in M \setminus \overline{I^-(p)}$, there exists $w \in I^+(p)$ such that $v \in M \setminus \overline{I^-(w)}$. Before proving this assertion, we will show how it implies this Proposition. Let $K \subset M \setminus \overline{I^-(p)}$ be compact. This assertion implies that $\{M \setminus \overline{I^-(w)} : w \in I^+(p)\}$ is an open covering of $K \subset M \setminus \overline{I^-(p)}$. Choose a finite subcovering $\{M \setminus \overline{I^-(w_i)} : w_i \in I^+(p), i = 1, 2, \dots, n\}$ determined by w_1, w_2, \dots, w_n and $U = \bigcap_{i=1}^n I^-(w_i)$. Then U is a neighborhood of p such that for any $u \in U$, $K \subset M \setminus \overline{I^-(u)}$. In fact, since $w_i \in I^+(p)$, then $p \in I^-(w_i)$, $i = 1, 2, \dots, n$. So, $p \in \bigcap_{i=1}^n I^-(w_i) = U$. For any $u \in U$, $u \in I^-(w_i)$ for $i = 1, 2, \dots, n$. Then $I^-(u) \subset I^-(w_i)$ for $i = 1, 2, \dots, n$. Thus, $M \setminus \overline{I^-(w_i)} \subset M \setminus \overline{I^-(u)}$ for $i = 1, 2, \dots, n$. So, $K \subset M \setminus \overline{I^-(u)}$. Then $I^-(p)$ is outer continuous.

It remains to prove the assertion. Suppose that $v \in \overline{I^-(w)}$ for any $w \in I^+(p)$, then we will get a contradiction. Since $v \in \overline{I^-(w)}$, then $I^-(v) \subset I^-(w)$ by Proposition A.3. Since condition (A) of Proposition A.4 holds, then $I^+(w) \subset I^+(v)$ for all $w \in I^+(p)$. On the other hand, for any $y \in I^+(p)$, i.e., $p \ll_M y$, there is a strictly conal curve $\gamma : [0, 1] \mapsto M$ such that $\gamma(0) = p$ and $\gamma(1) = y$. Let $x = \gamma(\frac{1}{2})$, then $p \ll_M x \ll_M y$. Thus, $y \in I^+(x)$ and $x \in I^+(p)$. Since $I^+(w) \subset I^+(v)$ for all $w \in I^+(p)$, then $I^+(x) \subset I^+(v)$. So, $y \in I^+(v)$ for any $y \in I^+(p)$. Then $I^+(p) \subset I^+(v)$. By Proposition A.3, $p \in \overline{I^+(v)}$. Thus, $v \in \overline{I^-(p)}$ by Proposition A.4, which is a contradiction with $v \in M \setminus \overline{I^-(p)}$. \square

Remark A.5. If for all p and q in M , either $I^+(p) = I^+(q)$ or $I^-(p) = I^-(q)$ implies $p = q$, then that I^\pm are outer continuous is equivalent to any one of the conditions in Proposition A.4. See [13, Theorem 2.1].

Lemma A.1. *If $J^\pm(p)$ are closed for all $p \in M$, then $I^\pm(p)$ are outer continuous.*

Proof. If $J^\pm(p)$ are closed for all $p \in M$, then condition (B) of Proposition A.4 holds. So, $I^\pm(p)$ are outer continuous by Proposition A.5. \square

Corollary A.1. *If $J^\pm(p)$ are closed for all $p \in M$, then the conal order “ \leq_M ” is quasi-closed.*

Proof. If $J^\pm(p)$ are closed for all $p \in M$, then $I^\pm(p)$ are outer continuous by Lemma A.1. Thus, the conal order “ \leq_M ” is quasi-closed by Theorem A.1. \square

A.2 Globally orderable homogeneous spaces

Let G be a connected Lie group and M be a smooth manifold. A *left action* of Lie group G on manifold M is a smooth map $\theta : G \times M \rightarrow M$ satisfying $\theta(e, x) = x$ and $\theta(g_1 g_2, x) = \theta(g_1, \theta(g_2, x))$ for all $g_1, g_2 \in G$, $x \in M$, where e is the identity element in G . We write $g \cdot x$ or gx for $\theta(g, x)$. The action is said to be *transitive* if for every pair of points $x, y \in M$, there exists $g \in G$ such that $g \cdot x = y$. For each $x \in M$, the *isotropy group* of x , denoted by G_x , is the set $G_x = \{g \in G : g \cdot x = x\}$.

A smooth manifold endowed with a transitive smooth action by a Lie group G is called a *homogeneous G -space* (or a *homogeneous space*).

Let G be a Lie group and $H \subset G$ be a closed subgroup. The *left coset space* $G/H = \{gH : g \in G\}$ is a smooth manifold of dimension $(\dim G - \dim H)$, and the left action of G on G/H is given by $g_1 \cdot (g_2H) = (g_1g_2)H$. Hence, G/H is a homogeneous space (see [27, Theorem 21.17]).

Let G be a Lie group and M be a homogeneous G -space. If p is any point of M , then the isotropy group G_p is a closed subgroup of G , and the $F : G/G_p \rightarrow M$ defined by $F(gG_p) = g \cdot p$ is an equivariant diffeomorphism (see [27, Theorem 21.18]). Because of this equivariant diffeomorphism, we can define a homogeneous space to be a coset space of the form G/H , where G is a Lie group and H is a closed subgroup of G .

Let G be a Lie group and fix $a \in G$. Define the *left translation* map $L_a : G \rightarrow G$ by $L_a(g) = ag$. The left translation map is diffeomorphism since it is smooth with smooth inverse. The inverse of L_a is clearly the map $L_{a^{-1}}$. The diffeomorphism L_g induces a vector space isomorphism $dL_g|_e : \mathfrak{g} = T_eG \rightarrow T_gG$, where \mathfrak{g} is the Lie algebra of G and e is the identity element in G .

Let $M = G/H$ be a homogeneous space and the natural projection $\pi : G \rightarrow G/H$, $\pi(g) = gH$ be a submersion. For each $a \in G$, define the *left translation* $\tau_a : G/H \rightarrow G/H$ by $\tau_a(gH) = agH$. Then the left translations τ_g are related to the left translations L_g on the Lie group G by $\pi \circ L_g = \tau_g \circ \pi$ for each $g \in G$. Let \mathfrak{h} be the Lie algebra of H and $o = \pi(e) = eH$. The differential $d\pi|_e : T_eG \rightarrow T_oM$ is a vector space homomorphism with $\ker d\pi|_e = \mathfrak{h}$, we obtain that $\mathfrak{g}/\mathfrak{h} \cong T_oM$, where $\mathfrak{g}/\mathfrak{h}$ is the set of cosets $X + \mathfrak{h} = \{X + Y : Y \in \mathfrak{h}\}$ for $X \in \mathfrak{g}$ (see [16, P488] or [1, P71]).

A *wedge* W is a closed and convex subset of a vector space that is invariant by scaling with real positive numbers (see e.g., [16] and [25]). Thus, a convex cone is a wedge in a vector space. A *wedge field* W_M on a manifold M assigns to each point $x \in M$ a wedge $W_M(x)$ in the tangent space T_xM .

Let $\Phi : G \times M \rightarrow M$ be any left group action on M such that each of the maps $\tau_g : M \rightarrow M$ defined by $\tau_g(x) = \Phi(g, x) = g \cdot x$ forms a diffeomorphism of M . Then a wedge field W_M is said to be *G -invariant* or *homogeneous* if $d\tau_g|_x(W_M(x)) = W_M(g \cdot x)$ for all $g \in G$ and $x \in M$.

Lemma A.2. *Let H be a closed subgroup of a Lie group G and W a wedge in \mathfrak{g} such that (i) $W \cap -W = \mathfrak{h}$, and (ii) $\text{Ad}(H)(W) = W$, where Ad is the adjoint representation of G . Define W_G and C_M by*

$$W_G(g) = dL_g|_e W, \quad C_M(x) = d\tau_g|_o C,$$

where $M = G/H$, e is the identity element in G , $o = eH$ is the base point in M , $x = gH$, and C is the convex cone in T_oM obtained as the projection of W onto $\mathfrak{g}/\mathfrak{h}$. Then, W_G is an invariant wedge field on G and C_M is a well-defined homogeneous or G -invariant cone field on M . Moreover, for each $g \in G$,

$$d\pi|_g(W_G) = C_M(\pi(g)),$$

where $\pi : G \rightarrow M$ is the canonical projection $\pi(g) = gH$.

Proof. See [16, Lemma VI.1.5]. □

Proposition A.6. *The homogeneous cone field on a homogeneous space is continuous and admits C^∞ sections.*

Proof. See [26, Proposition 4.6]. □

Theorem A.2. *Let C_M be a homogeneous cone field on $M = G/H$ as described in Lemma A.2. If $S = \overline{\langle \exp W \rangle H} \subset G$, then $S = \pi^{-1}(\overline{\{x \in M : o \leq_M x\}})$ and G/H is globally orderable with respect to C_M if and only if $W = \mathbf{L}(S)$, where $\mathbf{L}(S) = \{Z \in \mathfrak{g} : \exp(\mathbb{R}^+ Z) \subset S\}$.*

Proof. This Theorem is derived from [35, Theorem 1.6] or [17, Theorem 4.21]. Where $\exp : \mathfrak{g} \rightarrow G$ is the Lie group exponential map, $\exp(tZ)$ with $Z \in \mathfrak{g}, t \in \mathbb{R}$ is the one-parameter subgroup on G (see e.g., [27]) and \leq_M is the order generated by cone field C_M . □

Let $M = G/H$ be a homogeneous space with base-point $o = eH$. A Riemannian metric $(\cdot, \cdot)_p$, $p \in M$, is said to be *G-invariant* or *homogeneous*, if it satisfies $(X, Y)_o = (d\tau_g(X), d\tau_g(Y))_{g \cdot o}$ for each $g \in G$ and $X, Y \in T_o M$.

Proposition A.7. *Let G be a Lie group, H a closed subgroup, then the space G/H is complete in any G -invariant metric.*

Proof. See [14, p148]. □

A homogeneous space $M = G/H$ is called *reductive* if there exists a subspace \mathfrak{m} of \mathfrak{g} such that $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ and $\text{Ad}(h)\mathfrak{m} \subset \mathfrak{m}$ for all $h \in H$. Hence, $\mathfrak{m} \cong T_o M$ (see [1]). The next Proposition gives a simple description of G -invariant Riemannian metrics on a homogeneous space.

Proposition A.8. *Let $M = G/H$ be a reductive homogeneous space. Then there is a one-to-one correspondence between G -invariant Riemannian metrics on $M = G/H$ and $\text{Ad}^{G/H}$ -invariant inner products $\langle \cdot, \cdot \rangle$ on $\mathfrak{m} \cong T_o M$; that is, $\langle X, Y \rangle = \langle \text{Ad}^{G/H}(h)X, \text{Ad}^{G/H}(h)Y \rangle$ for all $X, Y \in \mathfrak{m}$, $h \in H$.*

Proof. See [1, Proposition 5.1]. The homomorphism $\text{Ad}^{G/H} : H \rightarrow GL(T_o M)$ such that $\text{Ad}^{G/H}(h) = d\tau_h|_o$ is the isotropy representation of the homogeneous space $M = G/H$. □

A.3 Differential positivity in flat spaces

Let M be the n -dimensional Euclidean space \mathbb{R}^n and C is a closed convex cone of M . There is a partial order “ \leq ” on M generated by cone C ($x \leq y$ if and only if $y - x \in C$). It should be pointed out that the order introduced by a closed convex cone is a closed partial order.

We consider the cone field on M defined by $C_M(x) = \{x\} \times C \subset TM = \mathbb{R}^n \times \mathbb{R}^n$. Such cone field is said to be a *contact cone field*. Thus, we can define the order “ \leq_M ” on M with respect to cone field C_M ($x \leq_M y$ if and only if there exists a conal curve $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = x$, $\gamma(1) = y$ and $\frac{d}{ds}\gamma(s) \in C_M(\gamma(s)) = C$).

It is easy to see that the constant cone field on Euclidean space \mathbb{R}^n satisfies the smoothness conditions and invariance condition in the previous sections.

The next proposition implies that the order “ \leq_M ” generated by constant cone field C_M agrees with the partial order “ \leq ” on M generated by cone C (see [26, Proposition 1.10]).

Proposition A.9. *Let M be a n -dimensional Euclidean space and C is a convex cone in M such that C forms a constant cone field C_M . Then for $x, y \in M$, $x \leq y$ if and only if $x \leq_M y$.*

Proof. If $x \leq y$, i.e., $y - x \in C$, then we choose a curve $\alpha(s) = sy + (1 - s)x$, where $s \in [0, 1]$. Thus, $\alpha(0) = x$, $\alpha(1) = y$, and $\frac{d}{ds}\alpha(s) = y - x \in C$. So, $\alpha(s)$ is a conal curve and $x \leq_M y$.

If $x \leq_M y$, then there is a conal curve $\alpha : [0, 1] \rightarrow M$ such that $\alpha(0) = x$, $\alpha(1) = y$ and $\frac{d}{ds}\alpha(s) \in C$. For any $\lambda \in C^* \setminus \{0\}$, where C^* is the dual cone of C , $\lambda(\frac{d}{ds}\alpha(s)) \geq 0$. On the other hand, $\lambda(y - x) = \lambda(\alpha(1) - \alpha(0)) = \lambda(\alpha(1)) - \lambda(\alpha(0))$. Since $\frac{d}{ds}\lambda(\alpha(s)) = \lambda(\frac{d}{ds}\alpha(s)) \geq 0$, then $\lambda(\alpha(1)) \geq \lambda(\alpha(0))$. So, $\lambda(y - x) \geq 0$. Thus, we obtain that $y - x \in C$, i.e., $x \leq y$. \square

And a similar result can be obtained for the order “ \ll ” with “ \ll_M ”, where $x \ll y$ if and only if $y - x \in \text{Int}C$.

Let $\frac{dx}{ds} = f(x)$ be a dynamical system in \mathbb{R}^n with the flow φ_t . The system is said to be *monotone* with respect to partial order “ \leq ” if $\varphi_t(x) \leq \varphi_t(y)$ whenever $x \leq y$ and $t \geq 0$ and *strongly monotone* if $\varphi_t(x) \ll \varphi_t(y)$ whenever $x \leq y$, $x \neq y$ and $t > 0$ (see [22] and [50]).

The following Proposition shows that in \mathbb{R}^n , a monotone system is differentially positive (see [9, Theorem 1]).

Proposition A.10. *Let M be the n -dimensional Euclidean space and C is a convex cone in M such that C forms a constant cone field C_M . Then a system is monotone with respect to partial order “ \leq ” generated by cone C if and only if this system is differentially positive with respect to cone field C_M .*

Proof. By Proposition A.9, order “ \leq ” and order “ \leq_M ” are equivalent.

Suppose that the system is differentially positive. For $x \leq y$, there exists a conal curve $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = x$, $\gamma(1) = y$ and $\frac{d}{ds}\gamma(s) \in C$. Then, we obtain that $\varphi_t(\gamma(\cdot))$ is also a conal curve for each $t \geq 0$. In fact, $\frac{d}{ds}\varphi_t(\gamma(s)) = d\varphi_t(\gamma(s))\frac{d}{ds}\gamma(s)$. Since φ_t is differentially positive and $\frac{d}{ds}\gamma(s) \in C$, then $\frac{d}{ds}\varphi_t(\gamma(s)) \in C$. So, $\varphi_t(x) \leq \varphi_t(y)$ for $t \geq 0$. And hence, φ_t is monotone with respect to “ \leq ”.

If the system is monotone. For $x \in M$ and $v \in C$, then there is a conal curve $\gamma(s)$, $s \in I$, such that $\gamma(0) = x$, $\frac{d}{ds}\gamma(s)|_{s=0} = v$ and $x \leq \gamma(s)$ with $s \geq 0$. Since φ_t is monotone, then for each $t \geq 0$, $\varphi_t(x) \leq \varphi_t(\gamma(s))$, i.e., $\varphi_t(\gamma(s)) - \varphi_t(\gamma(0)) \in C$. Thus, $\frac{d}{ds}\varphi_t(\gamma(s))|_{s=0} = \lim_{\Delta s \rightarrow 0^+} \frac{\varphi_t(\gamma(\Delta s)) - \varphi_t(\gamma(0))}{\Delta s} \in C$. So, we obtain that $d\varphi_t(x)v = \frac{d}{ds}\varphi_t(\gamma(s))|_{s=0} \in C$. And hence, the system is differentially positive. \square

References

- [1] A. Arvanitoyeorgos, An introduction to Lie groups and the geometry of homogeneous spaces, vol 22. American Mathematical Society, Providence, 2003.
- [2] J. K. Beem and P. E. Ehrlich, Global Lorentzian Geometry, Monographs and Textbooks in Pure and Applied Mathematics, 67. Marcel Dekker, Inc., New York, 1981.
- [3] P. T. Chruściel and J. D. E. Grant, On Lorentzian causality with continuous metrics, *Class. Quantum Grav.* **29**(2012), 145001.
- [4] I. Chueshov, Monotone Random Systems Theory and Applications, Lecture Notes in Mathematics, vol. 1779, Springer-Verlag, Berlin, 2002.
- [5] J. Faraut, Algèbres de Volterra et transformation de Laplace sphérique sur certains espaces symétriques ordonnés, *Symp. Math.* **29**(1987), 183-196.
- [6] J. Faraut, Espaces symétriques ordonnés et algèbres de Volterra, *J. Math. Soc. Japan*, **43**(1991), 133-147.
- [7] F. Forni and R. Sepulchre, Differential analysis of nonlinear systems: Revisiting the pendulum example, in 53rd IEEE Conference of Decision and Control, Los Angeles, IEEE, Piscataway, NJ, 2014, 3848-3859.
- [8] F. Forni, Differential positivity on compact sets, in 2015 54th IEEE Conference on Decision and Control (CDC), IEEE, Piscataway, NJ, 2015, 6355-6360.
- [9] F. Forni and R. Sepulchre, Differentially positive systems, *IEEE Trans. Automat. Control*, **61**(2016), 346-359.
- [10] L. García-Heveling, Causality theory of spacetimes with continuous Lorentzian metrics revisited, *Class. Quantum Grav.* **38**(2021), 145028.
- [11] A. K. Guts and A. V. Levichev, On the Foundations of Relativity Theory, *Sov. Math. Dokl.* **30**(1984), 253-257.
- [12] S. H. Hawking and G. F. R. Ellis, The large scale structure of space-time, Cambridge University Press, Cambridge, 1973.
- [13] S. H. Hawking and R. K. Sachs, Causally continuous space-times, *Commun. Math. Phys.* **35**(1974), 287-296.
- [14] S. Helgason, Differential Geometry, Lie Groups, and Symmetric Spaces, Amer. Math. Soc., 2001.

- [15] P. Hess and P. Poláčik, Boundedness of prime periods of stable cycles and convergence to fixed points in discrete monotone dynamical systems, *SIAM J. Math. Anal.* **24**(1993), 1312-1330.
- [16] J. Hilgert, K. H. Hofmann and J. Lawson, Lie groups, convex cones, and semigroups, Oxford University Press, Oxford, 1989.
- [17] J. Hilgert and K.-H. Neeb, Lie semigroups and their applications, *Lecture Notes in Math.* **1552**, Springer, 1993.
- [18] J. Hilgert and K.-H. Neeb, Wiener-Hopf operators on ordered homogeneous spaces, I, *J. Funct. Anal.* **132**(1995), 86-118.
- [19] J. Hilgert and G. Ólafsson, Causal Symmetric Spaces. *Geometry and Harmonic Analysis, Perspectives in Mathematics*, vol. 18, Academic Press, 1997.
- [20] M. W. Hirsch, Systems of differential equations which are competitive or cooperative II: convergence almost everywhere, *SIAM J. Math. Anal.* **16**(1985), 423-439.
- [21] M. W. Hirsch, Stability and convergence in strongly monotone dynamical systems, *J. Reine Angew. Math.* **383**(1988), 1-53.
- [22] M. W. Hirsch and H. L. Smith, Monotone dynamical systems, *Handbook of Differential Equations: Ordinary Differential Equations*, vol. 2, Elsevier, Amsterdam 2005.
- [23] J. Jiang and X. Zhao, Convergence in monotone and uniformly stable skew-product semiflows with applications, *J. Reine Angew. Math.* **589**(2005), 21-55.
- [24] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1976.
- [25] M. A. Krasnosel'skij, Je. A. Lifshits and A. V. Sobolev, *Positive linear systems, the method of positive operators*, Heldermann Verlag, Berlin, 1989.
- [26] J. D. Lawson, Ordered manifolds, invariant cone fields, and semigroups, *Forum Math.* **1**(1989), 273-308.
- [27] J. M. Lee, *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*, Springer, 2 edition, 2013.
- [28] B. Lemmens and R. D. Nussbaum, *Nonlinear Perron-Frobenius Theory*, Cambridge, Cambridge Univ. Press, 2012.
- [29] A. Levichev, Sufficient conditions for the nonexistence of closed causal curves in homogeneous space-times, *Izv. Phys.* **10**(1985), 118-119.

- [30] J. Mierczyński and W. Shen, Spectral Theory for Random and Nonautonomous Parabolic Equations and Applications, Chapman Hall/CRC Monogr. Surv. Pure Appl. Math., vol. 139, CRC Press, Boca Raton, FL, 2008.
- [31] V. Morinelli and K.-H. Neeb, Covariant homogeneous nets of standard subspaces, Commun. Math. Phys. **386**(2021), 305-385.
- [32] C. Mostajeran and R. Sepulchre, Ordering positive definite matrices, Inf. Geom. **1**(2018), 287-313.
- [33] C. Mostajeran and R. Sepulchre, Monotonicity on homogeneous spaces, Math. Control Signals Systems, **30**(2018), 1-25.
- [34] C. Mostajeran and R. Sepulchre, Positivity, monotonicity, and consensus on Lie groups, SIAM J. Control Optim. **56**(2018), 2436-2461.
- [35] K.-H. Neeb, Conal orders on homogeneous spaces, Inventiones mathematicae **104**(1991), 467-496.
- [36] K.-H. Neeb and G. Ólafsson, Nets of standard subspaces on Lie groups, Advances in Math. **384**(2021), 107715.
- [37] K.-H. Neeb and G. Ólafsson, Wedge domains in compactly causal symmetric spaces, Int. Math. Res. Not. IMRN **12**(2023), 10209-10312.
- [38] K.-H. Neeb and G. Ólafsson, Wedge domains in non-compactly causal symmetric spaces, Geometriae Dedicata **217**(2023) 30.
- [39] K.-H. Neeb, B. Ørsted and G. Ólafsson, Standard subspaces of Hilbert spaces of holomorphic functions on tube domains, Commun. Math. Phys. **386**(2021), 1437-1487.
- [40] L. Niu, Eventually competitive systems generated by perturbations, Electronic Journal of Differential Equations **121**(2019) 1-12.
- [41] L. Niu and X. Xie, Generic Poincaré-Bendixson theorem for singularly perturbed monotone systems with respect to cones of rank-2, Proc. Amer. Math. Soc. **151** (2023) 4199-4212.
- [42] G. I. Olshanskii, Convex cones in symmetric Lie algebras, Lie semigroups and invariant causal (order) structures on pseudo-Riemann symmetric spaces, Soviet Math. Dokl., Amer. Math. Soc. translation **26**(1982) 97-101.
- [43] R. Penrose, Techniques of differential topology in relativity, Regional Conference Series in Applied Math. vol. 7, SIAM, Philadelphia, 1972.
- [44] P. Poláčik, Convergence in smooth strongly monotone flows defined by semilinear parabolic equations, J. Differ. Equ. **79**(1989) 89-110.

- [45] P. Poláčik, Parabolic equations: asymptotic behavior and dynamics on invariant manifolds, in: Handbook on Dynamical systems, vol. 2, Elsevier, Amsterdam, 2002, 835-883.
- [46] P. Poláčik and I. Tereščák, Convergence to cycles as a typical asymptotic behavior in smooth strongly monotone discrete-time dynamical systems, Arch. Ration. Mech. Anal. **116**(1992) 339-360.
- [47] C. Pugh, An improved Closing Lemma and general density theorem, Amer. J. Math. **89**(1967) 1010-1021.
- [48] I. E. Segal, Mathematical Cosmology and Extragalactic Astronomy, Academic Press, New York, 1976.
- [49] W. Shen and Y. Yi, Almost automorphic and almost periodic dynamics in skew-product semiflows, Mem. Am. Math. Soc. **136**(647) 1998.
- [50] H. L. Smith, Monotone Dynamical Systems, an introduction to the theory of competitive and cooperative systems, Math. Surveys and Monographs, 41, Amer. Math. Soc., Providence, Rhode Island 1995.
- [51] H. L. Smith, Monotone dynamical systems: Reflections on new advances and applications, Discrete Contin. Dyn. Syst. **37**(2017), 485-504.
- [52] H. L. Smith and H. R. Thieme, Convergence for strongly ordered preserving semiflows, SIAM J. Math. Anal. **22**(1991), 1081-1101.
- [53] I. Tereščák, Dynamics of C^1 smooth strongly monotone discrete-time dynamical systems, preprint, Comenius University, Bratislava, 1994.
- [54] E. B. Vinberg, Invariant cones and orderings in Lie groups, Funct. Anal. Appl. **14**(1980), 1-13.
- [55] Y. Wang and J. Yao, Dynamics alternatives and generic convergence for C^1 -smooth strongly monotone discrete dynamical systems, J. Differ. Equ. **269**(2020), 9804-9818.
- [56] Y. Wang and J. Yao, Sharpened dynamics alternative and its C^1 -robustness for strongly monotone discrete dynamical systems, J. Funct. Anal. **283**(2022), 109538.