Rate Optimality and Phase Transition for User-Level Local Differential Privacy

Alexander Kent, Thomas B. Berrett and Yi Yu

Department of Statistics, University of Warwick

May 21, 2024

Abstract

Most of the literature on differential privacy considers the *item-level* case where each user has a single observation, but a growing field of interest is that of *user-level* privacy where each of the n users holds T observations and wishes to maintain the privacy of their entire collection.

In this paper, we derive a general minimax lower bound, which shows that, for locally private user-level estimation problems, the risk cannot, in general, be made to vanish for a fixed number of users even when each user holds an arbitrarily large number of observations. We then derive matching, up to logarithmic factors, lower and upper bounds for univariate and multidimensional mean estimation, sparse mean estimation and non-parametric density estimation. In particular, with other model parameters held fixed, we observe phase transition phenomena in the minimax rates as T the number of observations each user holds varies.

In the case of (non-sparse) mean estimation and density estimation, we see that, for T below a phase transition boundary, the rate is the same as having nT users in the item-level setting. Different behaviour is however observed in the case of s-sparse d-dimensional mean estimation, wherein consistent estimation is impossible when d exceeds the number of observations in the itemlevel setting, but is possible in the user-level setting when $T \gtrsim s \log(d)$, up to logarithmic factors. This may be of independent interest for applications as an example of a high-dimensional problem that is feasible under local privacy constraints.

Keywords: User-level differential privacy; Local differential privacy; Minimax optimality.

1 Introduction

Advances in data science and machine learning have demonstrated the immense statistical utility of large data sets when used as training data for a range of problems. Growing user concerns and regulatory demands (e.g. Zaeem and Barber, 2020), however, have raised issues of protecting the privacy of individuals whose data are collected and analysed. The desirable outcome of protecting an individual's privacy in a quantifiable manner whilst maintaining the statistical utility of the data has led to the development of formal frameworks for the notion of privacy, with *differential privacy* (Dwork et al., 2006) emerging as the "gold standard". These ideas have since been developed and implemented by technology companies such as Google (Erlingsson et al., 2014) and Meta (Yousefpour et al., 2021), and public bodies such as the United States Census Bureau (Bureau, 2023).

First formalised in what is called the *central model* of differential privacy, where there is a trusted data aggregator who has access to the original data before producing a private output statistic, there is also the *local model*, where data are privatised before being sent to a data aggregator. This is a more stringent requirement but results in greater privacy protection. The local model, whilst at the time not yet formalised under the differential privacy framework, is one of the older instances of privacy, dating back to the randomised response mechanism of Warner (1965), designed to avoid the issue of evasive responses in surveys where an individual lies when answering a potentially incriminating question.

Originating from computer science and cryptography, differential privacy has more recently attracted interest from the statistics community. A rich field of questions can be generated by aiming to introduce optimal private protocols for familiar statistical problems and thus to quantify the cost of privacy; see, for example, work on mean estimation (e.g. Duchi et al., 2018), density estimation (e.g. Butucea et al., 2020; Sart, 2023), hypothesis testing (e.g. Berrett and Butucea, 2020; Lam-Weil et al., 2022) and change point

analysis (e.g. Berrett and Yu, 2021). There is also a growing appreciation of the connections between differential privacy and robust statistics, both in the central (Liu et al., 2022; Hopkins et al., 2023) and local models (Li et al., 2023; Chhor and Sentenac, 2023), and techniques from the robust statistics literature can often be leveraged to construct private estimators. Questions of optimality are usually answered by deriving information-theoretic lower bounds, for which there are now general theoretical tools developed for both the central (e.g. Wasserman and Zhou, 2010; Bun et al., 2014; Cai et al., 2021, 2023) and local models (e.g. Duchi et al., 2018; Acharya et al., 2023a).

Whilst traditionally both the central and local models of differential privacy assume each user possesses a single observation, this is not always the case in practice. When a user possesses multiple observations, two frameworks are possible, the event-level setting and the stronger user-level setting. In the event-level setting, each user is subject to a level of privacy which protects against an inference attempt for any one item of the sample. In the user-level setting, each user's data must be sufficiently privatised to protect against an inference attempt on the entire sample at once. Naively applying existing techniques used in the event-level setting to the user-level setting under the local model either leads to a reduction in the privacy guarantee proportional to the number of data points a user holds, or results in estimators with greater error introduced for privatisation than the benefits provided by the additional data. As such, new methods are in demand to protect the user's entire collection of data, whilst simultaneously leveraging the additional data to provide improved performance over using only a single observation per user.

The user-level setup has been explored in the central model, where each user provides their entire collection, before privatisation, to the data aggregator (e.g. Liu et al., 2020; Levy et al., 2021; Ghazi et al., 2023). There are comparatively fewer studies on the local model of user-level privacy, particularly in a statistical context. The local model of user-level privacy has, so far, been considered for a few problems such as mean estimation (Girgis et al., 2022; Bassily and Sun, 2023), stochastic convex optimisation (Bassily and Sun, 2023), empirical risk minimisation (Girgis et al., 2022) and discrete density estimation (Acharya et al., 2023b). To the best of our knowledge, however, minimax optimality of existing estimators in most of these problems is unproven, even in univariate mean estimation, and the question of what happens as the number of observations a single user holds grows to infinity has not been addressed. We, in this paper, establish the minimax rates in a range of canonical statistical problems and discover interesting phase transition phenomena in these rates.

Lastly, we remark on connections between user-level local differential privacy and other decentralised learning problems. One can view local privacy as having each user's data pass through a channel that injects noise to privatise the data. Statistical inference under local information constraints (see e.g. Cai and Wei, 2024, and the references therein), for instance, can be viewed as having each user's data pass through a channel which compresses their data into a specified number of binary digits before being passed to the data aggregator, and there exist connections in how minimax lower bounds can be obtained in both these frameworks (e.g. Acharya et al., 2023a). Another similar topic is that of federated learning, introduced in McMahan et al. (2017) (see e.g. Kairouz et al. 2021 for a more general overview), where a central server aims to fit a model using the data held by a collection of clients, without the central server having access to all the data, due to privacy requirements and storage constraints, among other reasons.

1.1 General setup

We first introduce the framework of differential privacy (DP). Given data $\{X^{(i)}\}_{i=1}^n \subset \mathcal{Y}$, we consider a conditional distribution $Q: \sigma(\mathcal{Z}) \times \mathcal{Y}^n \to [0, 1]$ induced via a random mapping from \mathcal{Y}^n to a (potentially different) space \mathcal{Z} , where $\sigma(\mathcal{Z})$ denotes a σ -algebra on \mathcal{Z} . Writing $X^{(k:l)} = (X_k, X_{k+1}, \ldots, X_l)$ and fixing $\alpha \geq 0$, the mechanism Q is said to be α -DP if

$$\sup_{S \in \sigma(\mathcal{Z})} \frac{Q(Z \in S \mid X^{(1:n)} = x^{(1:n)})}{Q(Z \in S \mid X^{(1:n)} = x^{\prime(1:n)})} \le e^{\alpha}, \quad \forall x^{(1:n)}, x^{\prime(1:n)} \in \mathcal{Y}^n \quad \text{with} \quad \sum_{i=1}^n \mathbb{1}\{x^{(i)} \neq x^{\prime(i)}\} \le 1.$$

The value of α controls the strength of the privacy constraint, with smaller values imposing more restrictive conditions. In this work, we consider the case $0 < \alpha \leq 1$, known as the high-privacy regime. We exclude the degenerate case of $\alpha = 0$ in which no inference is possible.

For the more stringent local differential privacy (LDP) condition, we consider a family of conditional distributions $\{Q_i\}_{i=1}^n$ where $Q_1 : \sigma(\mathcal{Z}) \times \mathcal{Y} \to [0, 1]$ and $Q_i : \sigma(\mathcal{Z}) \times \mathcal{Y} \times \mathcal{Z}^{i-1} \to [0, 1]$ for $i \in \{2, \ldots, n\}$. Here the *i*-th individual privatises their data $X^{(i)}$ according to Q_i to produce a private observation $Z^{(i)}$. For $\alpha \geq 0$, we say that such a collection of conditional distributions $\{Q_i\}_{i=1}^n$ satisfies α -LDP if, for all

 $i \in \{1, 2, \dots, n\}$

$$\sup_{S \in \sigma(\mathcal{Z})} \frac{Q_i(Z^{(i)} \in S \mid X^{(i)} = x^{(i)}, Z^{(1:i-1)} = z^{(1:i-1)})}{Q_i(Z^{(i)} \in S \mid X^{(i)} = x^{\prime(i)}, Z^{(1:i-1)} = z^{(1:i-1)})} \le e^{\alpha}, \quad \forall x^{(i)}, x^{\prime(i)} \in \mathcal{Y} \text{ and } \forall z^{(1:i-1)} \in \mathcal{Z}^{i-1}.$$
(1)

Such a collection $\{Q_i\}_{i=1}^n$ is called *sequentially interactive*. In the case that, for each $i \in \{1, 2, ..., n\}$, $Z^{(i)}$ only depends on $X^{(i)}$, we say that the collection is instead *non-interactive*, and the conditional distributions take the simpler form $Q_i(Z^{(i)} \in S \mid X^{(i)} = x^{(i)})$.

We suppose that each user holds $T \ge 1$ independent and identically distributed (i.i.d.) data points, denoted by $X_{1:T}^{(i)}$ and taking values in $\mathcal{X} = \mathcal{Y}^T$. Most of the existing differential privacy literature assumes T = 1 whereas in this work we will consider general T. Let \mathcal{Z} denote the output space, which may depend on T. We say that, for $\alpha \ge 0$, a collection of conditional distributions $\{Q_i\}_{i=1}^n$ constitutes an α -LDP mechanism at the user-level if, for all $i \in \{1, \ldots, n\}$, the following inequalities hold

$$\sup_{S \in \sigma(\mathcal{Z})} \frac{Q_i(Z^{(i)} \in S \mid X_{1:T}^{(i)} = x_{1:T}^{(i)}, Z^{(1:i-1)} = z^{(1:i-1)})}{Q_i(Z^{(i)} \in S \mid X_{1:T}^{'(i)} = x_{1:T}^{'(i)}, Z^{(1:i-1)} = z^{(1:i-1)})} \le e^{\alpha}, \quad \forall x_{1:T}^{(i)}, x_{1:T}^{'(i)} \in \mathcal{X}, \quad z^{(1:i-1)} \in \mathcal{Z}^{i-1}.$$
(2)

If we relax this requirement so that the inequalities only need to holds when $x_t^{(i)} = x_t^{\prime(i)}$ for all except at most one $t \in \{1, \ldots, T\}$, then we obtain the definition of LDP at the *event-level*. Note, however, that this relaxation would not prevent privacy leakage due to repeated measures of the same individual's data. Writing $Q = \{Q_i\}_{i=1}^n$ when there is no ambiguity, we refer to any conditional distribution Q satisfying (2) as a user-level α -LDP mechanism, and denote the set of all such mechanisms by Q_{α} . As we focus on this model of privacy in this work, we refer to user-level local differential privacy as *user-level privacy* for brevity, specifying the distinction between the local and central models only when relevant.

As for the minimax framework, denote by \mathcal{P} the family of distributions that an observation $X_t^{(i)}$ is generated from. To quantify the best possible performance of estimators based on the privatised data generated by mechanisms Q satisfying (2), we consider the user-level α -LDP minimax risk, that is,

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) = \inf_{Q \in \mathcal{Q}_{\alpha}} \inf_{\hat{\theta}} \sup_{P \in \mathcal{P}} \mathbb{E}_{P,Q} \left\{ \Phi \circ \rho\left(\hat{\theta}, \theta\left(P\right)\right) \right\},\tag{3}$$

where

- the quantity of interest to be estimated is $\theta(P) \in \theta(\mathcal{P})$, denoting a functional on the space \mathcal{P} ;
- the function ρ is a metric on the space $\theta(\mathcal{P})$ and $\Phi : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ is a non-decreasing function with $\Phi(0) = 0$;
- the outermost infimum is taken over all user-level α -LDP privacy mechanisms generating the privatised data $Z^{(i)}$;
- the inner infimum is taken over all measurable functions $\hat{\theta} = \hat{\theta}(Z^{(1)}, \dots, Z^{(n)})$ of the privatised data generated by the privacy mechanism Q, conditional on the raw data generated from $P \in \mathcal{P}$; and
- the risk is the expectation with respect to both the distribution P of the data and the privacy mechanism Q.

We note that the usual α -LDP risk where each user has a single item, as considered in, for example, Duchi et al. (2018), can be recovered as a special case of (3) by considering $\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$. In an abuse of notation, denote the minimax risk without privacy by $\mathcal{R}_{n,1,\infty}(\theta(\mathcal{P}), \Phi \circ \rho)$, with $\alpha = \infty$.

Remark 1. As each observation a user holds is distributed according to a distribution P, we see that user-level privacy is the natural extension of (1) where we view each user's single data point as from the product distribution $P' = P^{\otimes T}$ and where the quantity being estimated $\theta(P') = \theta(P)$ regardless of T.

Being a stronger notion of privacy, user-level privacy provides a stronger guarantee than event-level privacy at the cost of reduced statistical utility of the privatised output. What is not so clear however, is whether the user-level framework with n users each possessing an independent sample of size T is preferable in terms of statistical utility to the traditional LDP framework having nT many users each with one observation. On the one hand, under user-level privacy a greater amount of noise may be

needed to privatise the user's sample, suggesting the user-level setup is at a disadvantage. On the other hand, it is possible that under the user-level setup, a user can discern useful information locally from their collection of items before submitting a privatised value, potentially providing an advantage. To distinguish between these two setups, we refer to the case where each user holds a single data point as the item-level case, so we will be interested in comparing the performance of item-level α -LDP with nT users against user-level α -LDP with n users, each with a sample of size T. Note, however, that with the same α , this results in a stricter privacy protection for the user-level. This is because in the user-level setting, every T observations share α privacy budget, while every one observation enjoys α privacy budget in the item-level setting. Lacking a widely-accepted strategy for a fairer comparison, we stick to this comparison while acknowledging the stricter privacy protection imposed in user-level privacy.

1.2 Contributions

In this paper, we aim to investigate the aforementioned question. The list of contributions is summarised below.

- In Section 2, we discuss the case where the number of items a user holds diverges whilst the number of users is kept fixed. As a user would be able to estimate any functional of the distribution arbitrarily well given an infinite and independent sample, we consider the minimax risk of estimating a quantity where each user knows the quantity exactly, but cannot report this value without randomising it to satisfy α -LDP. We develop general upper and lower bounds on the minimax risk for any estimation problem. The lower bound in particular has relevance beyond this thought experiment as it also applies to the user-level setting that we consider in specific problems, and will be used to characterise when phase transitions occur in the fundamental limits.
- In Section 3, for $d \in \mathbb{N}$, we consider *d*-dimensional mean estimation for distributions supported on unit ℓ_2 - and ℓ_{∞} -balls. Mean estimation procedures and proofs of their minimax optimality in the item-level case have been previously developed in Duchi et al. (2018). The problem has also been considered in the central user-level setting in Levy et al. (2021), and in the local setting, without showing minimax optimality, for estimators for distributions supported on the ℓ_{∞} -ball with some further assumptions in Girgis et al. (2022) and Bassily and Sun (2023). We build upon this prior work by only imposing a restriction on the support of data without further assumptions, considering both ℓ_2 - and ℓ_{∞} -balls, and deriving minimax rates. In particular, we show that a phase transition occurs when T is larger than a quantity exponential in $n\alpha^2$ and further increases in T no longer improve estimation.
- In Section 4, we consider sparse mean estimation. Unlike the mean estimation problems in Section 3 where, for small T - up until the phase transition, the optimal minimax rate is the same (up to logarithmic factors) as having nT users in the item-level setup, in Section 4 we see a prominent difference between minimax risks of the user- and item-level setups. For comparison, without privacy restrictions, assuming that the mean has at most one non-zero entry enables consistent estimators as long as the dimension is not exponential in the number of users. Duchi et al. (2018), however, shows that under item-level LDP, no consistent estimator exists if $n\alpha^2 = O(d)$. We will show that in the user-level setting, consistent estimators exist even if d is polynomial in $n\alpha^2$ as long as $\log(d) = O(T)$. This shows that there exist estimation problems that are significantly easier in the user-level setting compared to the item-level setting.
- In Section 5, we consider non-parametric density estimation. The estimation of discrete densities under user-level local privacy has been considered in Acharya et al. (2023b) where an estimator is developed and its minimax optimality established in a regime where the value of T is not too large. We consider the case of continuous densities lying in a function family defined through a Sobolev ellipsoid condition on the coefficients of its basis expansion. As well as constructing an estimator and proving its near-minimax optimality in all regimes of T, we will see that unlike the two previously considered parametric problems where the phase transition boundary is exponential in $n\alpha^2$, the phase transition boundary for this non-parametric problem is polynomial in $n\alpha^2$.

1.3 Notation and organisation

For $n \in \mathbb{N}$, we write $[n] = \{1, \ldots, n\}$. For a collection of vectors $\{x^{(i)}\}_{i \in [n]} \subset \mathbb{R}^d$, denote $x^{(1:n)} = \{x^{(1)}, \ldots, x^{(n)}\}$. For $a, b \in \mathbb{R}$, let $a \wedge b = \min(a, b)$ and $a \vee b = \max(a, b)$. For non-negative real sequences

 $\{a_n\}_{n\in\mathbb{N}}$ and $\{b_n\}_{n\in\mathbb{N}}$, $a_n \lesssim b_n$ denotes the existence of a constant C > 0 such that $\limsup_{n\to\infty} a_n/b_n \leq C$, $a_n \gtrsim b_n$ denotes that $b_n \lesssim a_n$ and $a_n \asymp b_n$ denotes that $a_n \lesssim b_n \lesssim a_n$. We say a random variable X is σ -sub-Gaussian if $\mathbb{P}(|X| > \varepsilon) \leq 2e^{-\varepsilon^2/(2\sigma^2)}$ for any $\varepsilon \geq 0$. For a distribution P, denote the n-fold product distribution arising from n i.i.d. observations as $P^{\otimes n}$. For two distributions P, Q, let $D_{\mathrm{KL}}(P||Q)$ and $D_{\mathrm{TV}}(P,Q)$ denote the Kullback–Leibler divergence and total variation distance between P and Q respectively. Given a convex set $\mathcal{C} \subset \mathbb{R}^d$, denote the projection of $x \in \mathbb{R}^d$ onto \mathcal{C} as $\Pi_{\mathcal{C}}(x)$. We define the Laplace distribution with scale $\lambda > 0$ to be the distribution. For a vector $v \in \mathbb{R}^d$ and $j \in [d]$ denote the j-th co-ordinate of v as v_j ; write $\|v\|_0 = \sum_{i=1}^d \mathbb{1}\{v_i \neq 0\}$. For two vectors $v, v' \in \mathbb{R}^d$, denote the Hamming distance between v and v' by $H(v,v') = \|v - v'\|_0$. For any set S, let |S| be its cardinality. For a given radius r > 0 and dimension $d \in \mathbb{N}$, we write $\mathbb{B}_2(r) = \{x \in \mathbb{R}^d : \|x\|_2 \leq r\}$ and $\mathbb{B}_{\infty}(r) = \{x \in \mathbb{R}^d : \|x\|_{\infty} \leq r\}$. Throughout this work, for simplicity we omit the taking the floor and/or ceiling of quantities for which an integer value is expected.

In the rest of this paper, we consider the case of diverging T in Section 2, and finite T in Sections 3, 4 and 5, on d-dimensional mean, s-sparse d-dimensional mean and density estimation, respectively. The paper is concluded with more discussion in Section 6, with all technical details in the Appendices.

2 Infinite observations per user: $T = \infty$

In this section, we will consider the limiting behaviour as T diverges to infinity. Whilst without any privacy constraints the error typically vanishes as $T \to \infty$, with privacy constraints, we will show that this is no longer the case. To understand what happens as T diverges, we consider the case where each user knows exactly the functional to estimate. Given a family of distributions \mathcal{P} , we consider the family

$$\mathcal{P}^{\infty} = \{ \delta_{\theta} : \theta \in \theta(\mathcal{P}) \}, \text{ where } \theta(\mathcal{P}) = \{ \theta(P) : P \in \mathcal{P} \},$$
(4)

and δ_{θ} is the point mass at θ . The sample space is therefore $\theta(\mathcal{P})$. For any $\theta \in \theta(\mathcal{P})$, the corresponding functional to estimate for $\delta_{\theta} \in \mathcal{P}^{\infty}$ is $\theta(\delta_{\theta}) = \theta$.

Our motivation is that if each user has an infinite sample size, then they would be able to estimate the functional perfectly. The problem is therefore equivalent to the case where the data distribution is a point mass on the value of the functional. With some abuse of notation and the user-level α -LDP minimax risk defined in (3), we define

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) = \mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}^{\infty}), \Phi \circ \rho), \tag{5}$$

and refer to this estimation problem as the *infinite-T* problem giving rise to the infinite-*T* minimax rate. Of particular interest is whether $\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ is of the same rate as $\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ as $T \to \infty$, and how the user-level rate $\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ compares to $\mathcal{R}_{nT,1,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ – the item-level rate with an equivalent number of observations.

We first demonstrate in Example 1 that it is wrong to take the lower bounds from the item-level case and to assume that $\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ and $\mathcal{R}_{nT,1,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho)$ are of the same order. In fact, as our results in Section 4 show, it is generally not even true that $\mathcal{R}_{n,T,\alpha}$ is lower bounded by a constant multiple of $\mathcal{R}_{nT,1,\alpha}$, so these two risks are generally incomparable.

Example 1. Estimating the mean of a distribution from the family $\mathcal{P} = \{P : \mathbb{E}_P(X) \in [-1,1]\}$, we have that the user-level LDP minimax risk with respect to the squared error loss is lower bounded as

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), (\cdot)^2) \gtrsim \min\left\{1, \frac{1}{nT\alpha^2}\right\}.$$
 (6)

When T = 1, (6) coincides with the item-level lower bound on $\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}), (\cdot)^2)$ (Corollary 1 in Duchi et al., 2018). We will show in Section 3 that, for a certain range of T, (6) is tight up to logarithmic factors, but not for the whole range of T. Indeed, we will also show that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), (\cdot)^2) \gtrsim e^{-12n\alpha^2}.$$
 (7)

This second lower bound, independent of T, demonstrates that for a fixed n, even as $T \to \infty$, the risk is bounded away from 0. The proofs of (6) and (7) are provided in Appendix A.

In the rest of this section, we will show that this non-vanishing phenomenon is not specific to the estimation problem in Example 1, but general for user-level LDP problems.

2.1 General minimax rates

We now introduce general upper and lower bounds for infinite-T LDP estimation problems. The lower bound developed here will also apply to the finite-T case.

We also construct a general method, providing tight upper bounds for the infinite-T case for a range of illustrative problems.

Theorem 1 (General infinite-T rates). Given a family of distributions \mathcal{P} and $\delta > 0$, let $N(\delta)$ be the δ -covering number of the metric space (Θ, ρ) with $\Theta = \theta(\mathcal{P})$ and suppose that $N(2\delta) > 1$. For $\alpha \in [0, 1]$, the LDP minimax risk satisfies that

$$\frac{\Phi(\delta)}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{\log(N(2\delta))} \right\} \le \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \le \Phi(\delta) + \Phi(\operatorname{diam}(\Theta))N(\delta)e^{-n\alpha^2/20}, \tag{8}$$

where diam(Θ) = sup_{$\theta, \theta' \in \Theta$} $\rho(\theta, \theta')$.

The proof of Theorem 1 is provided in Appendix A. The upper bound is due to a non-interactive procedure to be presented and the lower bound follows from an application of Fano's inequality along with a bound on the private Kullback–Leibler divergence (Duchi et al., 2018, Corollary 3). Of importance is the fact that the proof of the lower bound uses no properties of the family of distributions other than the metric entropy of the space $\theta(\mathcal{P})$. As the parameter space $\theta(\mathcal{P})$ remains the same across the item-level, user-level and infinite-T setups, the lower bound in Theorem 1 is applicable across all three, leading immediately to the following result.

Corollary 2. With the same conditions and notation as Theorem 1, we have that, for all $T \ge 1$, the user-level private minimax risk is lower bounded as

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \ge \frac{\Phi(\delta)}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{\log(N(2\delta))} \right\}.$$
(9)

Remark 2. As the lower bound in (9) is independent of T, an important consequence of Corollary 2 is that for any user-level LDP estimation problem it is impossible for the minimax risk to vanish as T diverges, provided that there exists $\delta > 0$ such that $12n\alpha^2 + \log(2) < \log(N(2\delta))$.

Letting δ_{LB} and δ_{UB} satisfy

$$N(2\delta_{\rm LB}) \ge \exp(\left[24n\alpha^2 + 2\log(2)\right])$$
 and $\Phi(\delta_{\rm UB}) \ge \Phi\{\operatorname{diam}(\Theta)\}N(\delta_{\rm UB})e^{-n\alpha^2/20}$,

we have that

$$\Phi(\delta_{\rm LB}) \lesssim \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \lesssim \Phi(\delta_{\rm UB}).$$

We now construct a general estimator which attains the upper bound in (8). The estimator involves constructing a δ -covering of the ρ -metric space of interest, and uses a private voting procedure to select the ball believed to be closest to the unknown parameter. We note that this voting-based strategy is commonly used in the privacy literature, sometimes under the name of unary encoding. See e.g. Wang et al. (2017) for an overview of this technique, and e.g. Girgis et al. (2022) for an application in user-level privacy. The procedure is formalised as follows.

Step 1. (Initialisation.) For $\delta > 0$, let $N(\delta)$ be the δ -covering number of the metric space (Θ, ρ) and $\mathcal{B} = \{B_j, j \in [N(\delta)]\}$ be a δ -cover, with θ_j being the centre of B_j for $j \in [N(\delta)]$. Define

$$B'_1 = B_1, \quad B'_j = B_j \setminus \bigcup_{k \in [j-1]} B_k, \quad j > 1.$$
 (10)

Step 2. (Voting.) For each user $i \in [n]$ with observation $X^{(i)} = \theta$, let $V^{(i)} = (\mathbb{1}\{X^{(i)} \in B'_j\})_{j \in [N(\delta)]}$.

Let $\{U_{i,j}\}_{i\in[n], j\in[N(\delta)]}$ be i.i.d. from Unif[0,1] and

$$\widetilde{V}_{j}^{(i)} = \begin{cases} V_{j}^{(i)}, & U_{i,j} \le e^{\alpha/2}/(1+e^{\alpha/2}), \\ 1-V_{j}^{(i)}, & \text{otherwise}, \end{cases} \quad i \in [n], \ j \in [N(\delta)].$$
(11)

Step 3. (Consolidation.) Set

$$j^* = \min \underset{j \in [N(\delta)]}{\operatorname{arg\,max}} \left(\sum_{i=1}^n \widetilde{V}_j^{(i)} \right) \quad \text{and} \quad \hat{\theta} = \theta_{j^*}$$

Note that the privatisation in (11) is non-interactive and satisfies α -LDP by Lemma 16.

With this algorithm in hand, we focus on the form of the upper bound in (8), consisting of two terms. The term $\Phi(\delta)$ is the error that arises when the correct ball is chosen. In this case, the error is at most $\Phi(\delta)$ by the definition of the constructed covering of the space. The other term is the error incurred when a different ball is chosen, in which case the error varies depending on the distance between the parameter and the centre of the chosen ball. This error is therefore upper bounded by the worst case distance $\Phi(\text{diam}(\Theta))$. This adverse outcome occurs when, after privatising the votes, the number of votes for the correct ball is less than that of some other ball, and the probability of this adverse outcome is upper bounded by $N(\delta)e^{-n\alpha^2/20}$. The exponential term is due to the use of Hoeffding's inequality when considering the privatised votes.

There is a trade-off between taking δ smaller to obtain a finer covering of Θ , which would result in a smaller $\Phi(\delta)$, and the difficulty in identifying the ball the users are voting for after privatisation as $N(\delta)$ increases. Loosely speaking, as δ decreases, there are more candidates to vote for, and so more opportunities for an incorrect ball to obtain more votes after privatisation by random chance. When obtaining upper bounds, we set δ to balance these two terms to obtain a sharp upper bound.

2.2 Application of Theorem 1

In this subsection we apply Theorem 1 in a range of problems. For suitably chosen values of δ we obtain minimax rates up to constants and, in the case of exponential bounds, constants in the exponent, the proofs of which are contained in Appendix A.

2.2.1 Mean estimation

Consider the mean estimation problems for the classes of distributions

$$\mathcal{P}_d = \left\{ P : \operatorname{supp}(P) \subseteq \mathbb{B}_{\infty}(1) \subset \mathbb{R}^d \right\} \quad \text{and} \quad \mathcal{P}'_d = \left\{ P : \operatorname{supp}(P) \subseteq \mathbb{B}_2(1) \subset \mathbb{R}^d \right\}, \tag{12}$$

with the mean of a distribution P denoted as $\theta(P) = \mathbb{E}_P(X)$.

Proposition 3. (i) Assuming that $n\alpha^2 > 60d \log(6d)$, it holds that

$$e^{-Cn\alpha^2/d} \lesssim \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \lesssim e^{-cn\alpha^2/d},$$
(13)

where \mathcal{P}_d is denoted in (12), $\mathcal{R}_{n,\infty,\alpha}$ is the infinite-T minimax risk rate defined in (5) and C > c > 0 are absolute constants.

(ii) Assuming that $n\alpha^2 > 60d$, it holds that

$$e^{-C'n\alpha^2/d} \lesssim \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}'_d), \|\cdot\|_2^2) \lesssim e^{-c'n\alpha^2/d},\tag{14}$$

where \mathcal{P}'_d is denoted in (12), $\mathcal{R}_{n,\infty,\alpha}$ is the infinite-T minimax risk rate defined in (5) and C' > c' > 0 are absolute constants.

The primary component of the proof of Proposition 3 is to upper and lower bound the covering numbers of the spaces $\theta(\mathcal{P}_d)$ and $\theta(\mathcal{P}'_d)$, before applying the general results Theorem 1.

The item-level counterpart of Proposition 3 is shown in Duchi et al. (2018), where the minimax rates are

$$\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \asymp d^2/(n\alpha^2) \quad \text{and} \quad \mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}_d'), \|\cdot\|_2^2) \asymp d/(n\alpha^2).$$
(15)

Compared to these polynomial rates, Proposition 3 exhibits exponentially decaying rates. In order to achieve vanishing minimax rates, in the ℓ_2 -ball scenarios (\mathcal{P}'_d) , the same dependence on d is observed in both the infinite-T and item-level cases, i.e. $d/(n\alpha^2) \to 0$. On the other hand, such dependence is substantially weakened in the infinite-T case when considering the ℓ_{∞} -ball scenario (\mathcal{P}_d) . While in both setups the ℓ_{∞} -ball scenario suffers from a worse dependence on the dimension d compared to the ℓ_2 -ball one, the infinite-T setup only requires $n\alpha^2 \gtrsim d \log(d)$, as opposed to $n\alpha^2 \gtrsim d^2$ for the item-level rate.

2.2.2 Sparse mean estimation

Consider the high-dimensional sparse mean estimation problem where the family of distributions is

$$\mathcal{P}_{d,s} = \left\{ P : \operatorname{supp}(P) \subseteq \mathbb{B}_{\infty}(1) \subset \mathbb{R}^{d}, \ \|\mathbb{E}_{P}(X)\|_{0} \leq s \right\},\tag{16}$$

with the mean of distribution P denoted as $\theta(P) = \mathbb{E}_P(X)$.

Proposition 4. Assuming that $n\alpha^2 > 120s \log(6d)$, it holds that

$$e^{-Cn\alpha^2/s} \lesssim \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_2^2) \lesssim e^{-cn\alpha^2/s},\tag{17}$$

where $\mathcal{P}_{d,s}$ is denoted in (16), $\mathcal{R}_{n,\infty,\alpha}$ is the infinite-T minimax risk rate defined in (5) and C > c > 0 are absolute constants.

Proposition 4 is shown by obtaining suitable bounds on the covering number of the metric space $(\theta(\mathcal{P}_{d,s}), \|\cdot\|_2)$, where $\|\cdot\|_2$ is the vector ℓ_2 -norm, and by applying Theorem 1. The upper bound on the covering number of $\theta(\mathcal{P}_{d,s})$ comes from considering a union of s-dimensional subspaces which cover the space $\theta(\mathcal{P}_{d,r})$. For the lower bound we explicitly construct a packing of $\theta(\mathcal{P}_{d,s})$, described in the proof of Lemma 18 in Appendix D. This is an interesting byproduct on its own and gives a lower bound on the covering number.

Without privacy constraints, one can show that $\mathcal{R}_{n,1,\infty}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_2^2) \approx s \log(ed/s)/n$ (e.g. Han, 2016, Theorem 8.2.6), showing that the dimension d can grow exponentially large in n. On the other hand, it is shown in Theorem 3 in Acharya et al. (2023a) that for $s > 4 \log(d)$, $\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_2^2) \approx sd/(n\alpha^2)$

We see that, item-level LDP constraints are disastrous for sparse mean estimation, where the dimension cannot be linear in $n\alpha^2$ for consistent estimation, let alone exponential. On the other hand, we see that in the infinite-*T* case, the rates in Proposition 4 and (13) match when replacing *d* by *s*. This shows that in the infinite-*T* case, the fundamental limits of the *s*-dimensional mean and *s*-sparse *d*-dimensional mean problems are the same, up to constants in exponential. This hints at potential benefits of user-level LDP for finite-*T* in the context of sparse mean estimation to be explored further in Section 4.

2.2.3 Non-parametric density estimation

For the non-parametric density estimation problem we consider the *elliptical Sobolev space* (e.g. equation (1.91) in Tsybakov, 2009) defined below. Define the trigonometric basis

$$\varphi_1(x) = 1, \quad \varphi_{2j}(x) = \sqrt{2}\cos(2\pi jx) \quad \text{and} \quad \varphi_{2j+1}(x) = \sqrt{2}\sin(2\pi jx), \quad j \in \mathbb{N},$$
 (18)

Given a smoothness parameter $\beta \in \mathbb{N}$ and a radius r > 0, the Sobolev class of functions of smoothness β and radius r is given by

$$\mathcal{S}_{\beta,r} = \left\{ f \in L^2([0,1]) : f = \sum_{j=1}^{\infty} \theta_j \varphi_j, \sum_{j=1}^{\infty} j^{2\beta} \theta_j^2 \le r^2 \right\}.$$
(19)

This leads to the definition of the sub-class of Sobolev densities

$$\mathcal{F}_{\beta,r} = \left\{ f \in \mathcal{S}_{\beta,r} : f > 0, \int f(x) \, \mathrm{d}x = 1 \right\}.$$
(20)

We denote by P_f the distribution induced by the density f and define the class of all such distributions

$$\mathcal{P}_{\beta,r} = \{ P_f : f \in \mathcal{F}_{\beta,r} \}, \qquad (21)$$

where the functional to estimate is $\theta(P_f) = f$ and the error is measured by the squared $L^2([0,1])$ norm denoted by $\|\cdot\|_2^2$ where

$$||f||_{2}^{2} = \int_{\mathcal{X}} \{f(x)\}^{2} \, \mathrm{d}x.$$
(22)

Proposition 5. There exists a constant $C'_{\beta} > 0$ depending only on β such that, whenever $n\alpha^2 \ge C'_{\beta}$, it holds that

$$\frac{c_{\beta}}{(n\alpha^2)^{2\beta}} \lesssim \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{\beta,1}), \|\cdot\|_2^2) \lesssim \frac{C_{\beta}}{(n\alpha^2)^{2\beta}},\tag{23}$$

where $\mathcal{P}_{\beta,1}$ is denoted in (21), $\mathcal{R}_{n,\infty,\alpha}$ is the infinite-T minimax risk rate defined in (5) and $C_{\beta} > c_{\beta} > 0$ are constants depending only on β .

¹For s = 1, Duchi et al. (2018) shows that the non-interactive minimax risk, as opposed to the interactive risks considered throughout this work and in Acharya et al. (2023a), is of order $d \log(2d)/(n\alpha^2)$.

The upper and lower bounds of Proposition 5 are derived by obtaining suitable upper and lower bounds on the covering number of the space $\mathcal{F}_{\beta,1}$ and by applying Theorem 1. For the upper bound on the complexity of $\mathcal{F}_{\beta,1}$, it suffices to use the standard upper bound on the covering number of $\mathcal{S}_{\beta,r}$ (e.g. Wainwright, 2019, Example 5.12); for the lower bound, we explicitly construct a packing of $\mathcal{F}_{\beta,1}$, described in the proof of Lemma 17 in Appendix D, which is again a byproduct of its own interest.

Compared to Proposition 3, we see that the minimax risk is polynomial in $n\alpha^2$ rather than exponential. This occurs due to the fact that the covering number of the $\mathcal{F}_{\beta,1}$ is exponential in $1/\delta$, rather than polynomial as is the case for the spaces arising from the classes in (12) and (16). This also shows a significant difference between parametric and non-parametric rates in user-level LDP settings.

2.3 Discussion

In this section we have discussed the minimax rates for the infinite-T case, where the data are drawn from distributions in \mathcal{P}^{∞} , defined in (4). Comparisons of the minimax rates corresponding to \mathcal{P}^{∞} and those to \mathcal{P} , in the three problems we discussed in this section are collected below, where we suppress multiplicative constants and constants in exponents.

	d -dim. mean (ℓ_2 -ball)	Sparse mean	Density (Sobolev β -smooth)
Distribution class \mathcal{P}	$d/(n\alpha^2)$	$sd/(n\alpha^2)$	$(n\alpha^2)^{-2\beta/(2\beta+2)}$
Distribution class \mathcal{P}^∞	$e^{-n\alpha^2/d}$	$e^{-n\alpha^2/s}$	$(n\alpha^2)^{-2\beta}$

Table 1: Comparison of infinite-T user-level rates, and item-level mean estimation and density rates derived in Duchi et al. (2018), and sparse mean estimation rate derived in Acharya et al. (2023a). Multiplicative constants and constants in exponents are suppressed for illustration convenience.

We note the significant difference between the two cases, for example in the user-level setting with the exponential rate in mean estimation, or the lack of dependence on d in the sparse rate except through the required assumption $n\alpha^2 \gtrsim 120 \log(6d)$, which is itself no stronger (up to constants) than what is assumed in the non-private analogue. We also note that the rate is not always exponential, as seen in the non-parametric density estimation, where a polynomial rate is still observed, but with a different exponent from that in the item-level case.

With these results in hand for the infinite-T case we will next consider the finite-T user-level LDP case. By virtue of Corollary 2, we have that all the lower bounds developed thus far also hold for the minimax risk for finite-T. We note the parallels to Levy et al. (2021), where it was shown that for the problems considered therein, the minimax risk for the central user-level framework does not vanish even as $T \to \infty$, answering a conjecture of Amin et al. (2019). We thus resolve the local model analogue of this conjecture where one asks what these T-independent lower bounds will be in the local model.

Further, intuition suggests that we might expect that as $T \to \infty$ with other quantities held fixed, the minimax risk for user-level LDP will demonstrate similar behaviour in the limit as in the infinite-Tcases above as each user can estimate the functional increasingly well. We will shortly see that this is indeed the case, with the user-level rates demonstrating a phase transition once T is sufficiently large, matching the rates we obtained in this section.

3 Mean estimation

Recalling the families \mathcal{P}_d and \mathcal{P}'_d defined in (12), we estimate the mean of a distribution P so that $\theta(P) = \mathbb{E}_P(X)$ and measure the error by the squared ℓ_2 -loss.

Theorem 6. (i) Assuming that $n\alpha^2 > \widetilde{C}d\log(ed)$, where $\widetilde{C} > 0$ is an absolute constant, we have that

$$\frac{d^2}{nT\alpha^2} \vee e^{-Cn\alpha^2/d} \lesssim \mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \lesssim \frac{d^2\log(nT\alpha^2/d)}{nT\alpha^2} + e^{-cn\alpha^2/d},\tag{24}$$

where \mathcal{P}_d is denoted in (12) for the ℓ_{∞} case, $\mathcal{R}_{n,T,\alpha}$ is the user-level minimax risk rate defined in (3) and C > c > 0 are absolute constants.

(ii) Assuming that $n\alpha^2 > \widetilde{C}' d\log(ed)$, where $\widetilde{C}' > 0$ is an absolute constant, we have that

$$\frac{d}{nT\alpha^2} \vee e^{-C'n\alpha^2/d} \lesssim \mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}'_d), \|\cdot\|_2^2) \lesssim \min\left\{1, \frac{d\{\log(nT\alpha^2)\}^2}{nT\alpha^2} + e^{-c'n\alpha^2/d}\right\}$$

where \mathcal{P}'_d is denoted in (12) for the ℓ_2 case, $\mathcal{R}_{n,T,\alpha}$ is the user-level minimax risk rate defined in (3) and C' > c' > 0 are absolute constants.

In both cases, we see that lower bounds are the maxima of two parts, the polynomial term analogous to the standard item-level rate (Duchi et al., 2018, Corollary 4), and the exponential term inherited from Theorem 1.

The estimation methods in deriving upper bounds are similar to that in Girgis et al. (2022) and Bassily and Sun (2023), and repeat standard user-level estimation techniques of first localising to a small region where the mean is believed to lie, and then obtaining a refined estimator on the chosen region.

The exponential-decay minimax rate once T is sufficiently large is a phenomenon new to the literature, and confirms the intuition of Section 2 that the infinite-T rate, for this problem given by Proposition 3, should coincide with the finite-T rate for large T.

In the rest of this section, we present the estimation methods for the univariate case in Section 3.1, the multivariate case with ℓ_{∞} -ball in Section 3.2 and the multivariate case with ℓ_2 -ball in Section 3.3, with discussion in Section 3.4. The upper bounds follow from the analysis, in Appendix B, of the constructed estimators, with the lower bounds proven in Appendix C.

3.1 Univariate procedure

The estimation procedure developed for the univariate case will be used as a building block for both the ℓ_2 - and ℓ_{∞} -ball cases of the multivariate problem. The intuition for the univariate case is as follows. As each user has an i.i.d. sample of size T, each user is able to estimate the mean within O(1/T) mean squared error accuracy. For $\alpha \in (0, 1)$, if we were to naively privatise this procedure via, e.g. the Laplacian mechanism, to ensure α -LDP the added noise would need to have $O(1/\alpha^2)$ variance, dominating O(1/T).

To avoid this problem, the procedure splits the users into two halves. Without loss of generality, assume that the number of the users n is even. We partition [-1, 1] into non-overlapping sub-intervals of width 2δ , with δ to be specified. The first half of users calculate their own sample means, identify which sub-intervals contain their sample means and publicise indicators. The central statistician then produces an interval of width $O(\delta)$ that will contain the mean with high probability.

The second half calculate their own sample means, project the sample means onto the interval output from the first half and apply the Laplacian mechanism. As the width of the initially estimated interval is $O(\delta)$ rather than O(1), it suffices to perturb with noise with variance of order $O(\delta^2/\alpha^2)$ rather than $O(1/\alpha^2)$. By choosing $\delta = O(1/\sqrt{T})$, up to logarithmic factors, we are able to take advantage of the T observations each user holds.

We note that for large values of T we do not make use of all T observations, but instead cap the value of T at T^* , a value exponential in $n\alpha^2$, making users discard excess data. This step is taken to prevent the value of δ becoming too small, which would result in the total number of sub-intervals becoming too large, and hence impossible to reliably select the correct interval from the private votes. Further, discarding excess observations does not reduce the utility of our final estimator. To be specific, using only T^* observations in calculating sample means in (26) and (30) does not reduce the accuracy of the final estimator as if all T observations were used. This is because the estimation error rate is dominated by either δ or the added Laplacian noise required for privacy. As a byproduct, this also results in a lower storage and computational cost.

The procedure is formalised as follows.

Step 1. (Initialisation.) Let $T^* = T \wedge e^{n\alpha^2/K} \ge 1$ with K > 0 an absolute constant and let

$$\delta = \left\{ 2\log(nT^*\alpha^2)/T^* \right\}^{1/2}.$$
(25)

Let $\{I_j, j \in [N(\delta)]\}$ be a partition of [-1, 1], with $I_j = [-1 + 2(j-1)\delta, -1 + 2j\delta), j \in [N(\delta) - 1]$ and $I_{N(\delta)} = [-1 + 2\{N(\delta) - 1\}\delta, -1 + 2N(\delta)\delta].$

Step 2. (Localisation.) Calculate

$$\hat{\theta}^{(i)} = \frac{1}{T^*} \sum_{t=1}^{T^*} X_t^{(i)} \text{ and } V_j^{(i)} = \mathbb{1}\{\hat{\theta}^{(i)} \in I_{j-1} \cup I_j \cup I_{j+1}\}, \quad i \in [n/2], \, j \in [N(\delta)],$$
(26)

with $I_0 = I_{N(\delta)+1} = \emptyset$. Let $U_{i,j}$, $i \in [n/2], j \in [N(\delta)]$, be i.i.d. from Unif[0, 1] and set

$$\widetilde{V}_{j}^{(i)} = \begin{cases} V_{j}^{(i)}, & U_{i,j} \le e^{\alpha/6}/(1+e^{\alpha/6}), \\ 1-V_{j}^{(i)}, & \text{otherwise}, \end{cases} \quad i \in [n/2], \ j \in [N(\delta)].$$

$$(27)$$

Choose

$$j^* = \min \underset{j \in [N(\delta)]}{\operatorname{arg\,max}} \sum_{i=1}^{n/2} \widetilde{V}_j^{(i)}.$$
(28)

Let L, U denote the lower and upper end points of I_{j^*} respectively,

$$\tilde{L} = L - 6\delta, \quad \tilde{U} = U + 6\delta \quad \text{and} \quad \tilde{I}_{j^*} = [\tilde{L}, \tilde{U}].$$
(29)

Step 3. (Refinement.) Let

$$\hat{\theta}^{(i)} = \Pi_{\tilde{I}_{j^*}} \left(\frac{1}{T^*} \sum_{t=1}^{T^*} X_t^{(i)} \right) + \frac{14\delta}{\alpha} \ell_i, \quad i \in [n] \setminus [n/2],$$
(30)

where the ℓ_i 's are i.i.d. standard Laplace random variables. The final estimator is defined as

$$\hat{\theta} = \frac{1}{n/2} \sum_{i=n/2+1}^{n} \hat{\theta}^{(i)}.$$

The privatisation in (27) satisfies α -LDP by Lemma 16, as does the estimator (30) due to being an instance of the Laplacian mechanism.

With this estimator for the univariate case, we now consider the multivariate generalisations, where the univariate procedure is applied co-ordinatewise. For the ℓ_{∞} -ball this can be done to the orginal data, but for the ℓ_2 -ball we first apply a transformation.

3.2 Multivariate procedure for ℓ_{∞} -ball

For the multivariate procedure for the ℓ_{∞} -ball, we split the *n* users into *d* many equally-sized folds, each of which is assigned to estimate a co-ordinate using the univariate procedure detailed in Section 3.1. Since we assume that $n\alpha^2 \gtrsim d\log(ed)$, we may assume without loss of generality that *n* is a multiple of 2*d*.

The mean squared error of the final estimator and the upper bound in (24) have a factor of d^2 . One d is from the fact that each co-ordinate is estimated by n/d users, and the other d is introduced by the sum of d-many terms when summing the error for each co-ordinate.

The procedure is formalised as follows.

Step 1. (Initialisation.) For $j \in [d]$, let

$$N_{j,1} = \left\{ (j-1)\frac{n}{2d} + 1, \dots, j\frac{n}{2d} \right\}, N_{j,2} = \left\{ (d+j-1)\frac{n}{2d} + 1, \dots, (d+j)\frac{n}{2d} \right\}, N_j = N_{j,1} \cup N_{j,2}.$$
(31)

Let $T^* = T \wedge e^{n\alpha^2/(Kd)} \ge 1$ with K > 0 an absolute constant and let

$$\delta = \left\{ 2 \log(nT^* \alpha^2 / d) / T^* \right\}^{1/2}.$$

Let $\{I_j, j \in [N(\delta)]\}$ be a partition of [-1, 1], with $I_j = [-1 + 2(j-1)\delta, -1 + 2j\delta), j \in [N(\delta) - 1]$, and $I_{N(\delta)} = [-1 + 2(N(\delta) - 1)\delta, -1 + 2N(\delta)\delta]$.

Step 2. (Estimation.) For each $j \in [d]$, perform the following.

Calculate

$$\hat{\theta}^{(i)} = \frac{1}{T^*} \sum_{t=1}^{T^*} X_t^{(i)} \text{ and } V_k^{(i)} = \mathbb{1}\{\hat{\theta}_j^{(i)} \in I_{k-1} \cup I_k \cup I_{k+1}\}, \quad i \in N_{j,1}, \, k \in [N(\delta)],$$

with $I_0 = I_{N(\delta)+1} = \emptyset$. Let $U_{i,k}$, $i \in N_{j,1}$, $k \in [N(\delta)]$, be i.i.d. from Unif[0, 1] and set

$$\widetilde{V}_{k}^{(i)} = \begin{cases} V_{k}^{(i)}, & U_{i,k} \le e^{\alpha/6}/(1+e^{\alpha/6}), \\ 1-V_{k}^{(i)}, & \text{otherwise,} \end{cases} \quad i \in N_{j,1}, \, k \in [N(\delta)]. \end{cases}$$

Choose

$$k_j^* = \min \underset{k \in [N(\delta)]}{\arg \max} \sum_{i \in N_{j,1}} \widetilde{V}_k^{(i)}.$$

Let L_j, U_j denote the lower and upper end points of $I_{k_j^*}$ respectively,

$$\tilde{L}_j = L_j - 6\delta, \quad \tilde{U}_j = U_j + 6\delta \text{ and } \tilde{I}_{k_j^*} = [\tilde{L}_j, \tilde{U}_j].$$

Step 3. (Refinement.) For each $j \in [d]$, perform the following:

Let

$$\hat{\theta}^{(i)} = \Pi_{\tilde{I}_{k_{j}^{*}}} \left(\frac{1}{T^{*}} \sum_{t=1}^{T^{*}} X_{t,j}^{(i)} \right) + \frac{14\delta}{\alpha} \ell_{i}, \quad i \in N_{j,2}$$

where the ℓ_i are i.i.d. standard Laplace random variables. Let the refined estimator for the *j*-th co-ordinate be

$$\hat{\theta}_j = \frac{1}{n/(2d)} \sum_{i \in N_{j,2}} \hat{\theta}^{(i)} \in \mathbb{R}.$$

Step 4. (Consolidation.) The final estimator is defined as

$$\hat{\theta} = \sum_{j=1}^{d} e_j \hat{\theta}_j \in \mathbb{R}^d, \tag{32}$$

where e_j is the *j*-th standard basis vector of \mathbb{R}^d .

3.3 Multivariate procedure for ℓ_2 -ball

For the ℓ_{∞} -ball, splitting the sample and applying the univariate estimator to each co-ordinate attains the optimal rate. This procedure however does not attain the optimal rate for estimation on the ℓ_2 -ball. As seen in the lower bound in Theorem 6, for estimation on the ℓ_2 -ball, the dependence on the dimension is linear rather than quadratic, but applying the same sample splitting method with no modification yields a quadratic rate. We instead carry out an encoding step which, loosely speaking, improves the concentration of data by a factor of $O(1/\sqrt{d})$, resulting in a linear rate in d. Such encoding has been implemented in the context of differential privacy, including Kashin's representation (see e.g. Lyubarskii and Vershynin 2010 for an introduction and e.g. Chen et al. 2020 for an application to differential privacy) and random rotations. The latter method has been used for private mean estimation in both central (e.g. Levy et al., 2021) and local (e.g. Girgis et al., 2022) frameworks.

Without loss of generality, we assume that the dimension is a power of 2, as we can append auxiliary values of 0 to all the data. This dyadic design does not affect the final rate up to constants.

We first define the Hadamard matrices, denoting H_k for k a power of 2, as the family of matrices which can be constructed inductively from

$$H_1 = (1)$$
 and $H_k = \begin{pmatrix} H_{k-1} & H_{k-1} \\ H_{k-1} & -H_{k-1} \end{pmatrix}$, $k > 1$.

We then denote by W_k a diagonal matrix with diagonal entries being i.i.d. Rademacher random variables. Construct the rotation matrix

$$R_k = H_k W_k / \sqrt{k}. \tag{33}$$

Rotating data using the rotation matrix R_k , with high probability, ensures that the resulting data concentrate better in ℓ_{∞} -norm. This is made precise in the following result, derived from the proof of Levy et al. (2021, Lemma 2).

Lemma 7 (Levy et al., 2021, Lemma 2). Assume that the dimension $d \in \mathbb{N}$ is a power of 2 and fix any $j \in [d]$. For the rotation matrix R_d in (33), vectors $x, x_0 \in \mathbb{R}^d$ and $\gamma \in (0, 1)$, we have, with probability at least $1 - \gamma$, that

$$|(R_d x)_j - (R_d x_0)_j| \le \frac{10||x - x_0||_2 \{\log(1/\gamma)\}^{1/2}}{d^{1/2}}.$$

Applying this random transformation to the data before using the univariate estimation procedure to each co-ordinate yields the following algorithm.

Step 1. (Initialisation.) For $j \in [d]$, let

$$N_{j,1} = \left\{ (j-1)\frac{n}{2d} + 1, \dots, j\frac{n}{2d} \right\}, N_{j,2} = \left\{ (d+j-1)\frac{n}{2d} + 1, \dots, (d+j)\frac{n}{2d} \right\}, N_j = N_{j,1} \cup N_{j,2}.$$

Let $T^* = T \wedge e^{-n\alpha^2/(Kd)} \ge 1$ with K > 0 an absolute constant, R_d be a random rotation matrix defined in (33), and

$$\delta = \left[\frac{C\{\log(nT^*\alpha^2)\}^2}{dT^*}\right]^{1/2},\tag{34}$$

for an absolute constant C > 0.

Let $\{I_j, j \in [N(\delta)]\}$ be a partition of [-1, 1], with $I_j = [-1 + 2(j-1)\delta, -1 + 2j\delta), j \in [N(\delta) - 1]$ and $I_{N(\delta)} = [-1 + 2(N(\delta) - 1)\delta, -1 + 2N(\delta)\delta].$

Step 2. (Estimation.) For each $j \in [d]$, perform the following.

Calculate

$$\hat{\theta}^{(i)} = \frac{1}{T^*} \sum_{t=1}^{T^*} X_t^{(i)} \text{ and } V_k^{(i)} = \mathbb{1}\{ (R_d \hat{\theta}^{(i)})_j \in I_{k-1} \cup I_k \cup I_{k+1} \}, \quad i \in N_{j,1}, \, k \in [N(\delta)], \quad (35)$$

with $I_0 = I_{N(\delta)+1} = \emptyset$ and R_d in (33). Let $U_{i,k}$, $i \in N_{j,1}$, $k \in [N(\delta)]$, be i.i.d. from Unif[0, 1] and set

$$\widetilde{V}_{k}^{(i)} = \begin{cases} V_{k}^{(i)}, & U_{i,k} \le e^{\alpha/6}/(1+e^{\alpha/6}), \\ 1-V_{k}^{(i)}, & \text{otherwise}, \end{cases} \quad i \in N_{j,1}, \, k \in [N(\delta)]. \end{cases}$$

Choose

$$k_j^* = \min \argmax_{k \in [N(\delta)]} \sum_{i \in N_{j,1}} \widetilde{V}_k^{(i)}.$$

Let L_j, U_j denote the lower and upper end points of $I_{k_i^*}$ respectively,

$$\tilde{L}_j = L_j - 6\delta, \quad \tilde{U}_j = U_j + 6\delta \text{ and } \tilde{I}_{k_j^*} = [\tilde{L}_j, \tilde{U}_j]$$

Step 3. (Refinement.) For each $j \in [d]$, perform the following:

Let

$$\tilde{\theta}^{(i)} = \Pi_{\tilde{I}_{k_{j}^{*}}} \left\{ \frac{1}{T^{*}} \sum_{t=1}^{T^{*}} \left(R_{d} X_{t}^{(i)} \right)_{j} \right\} + \frac{14\delta}{\alpha} \ell_{i}, \quad i \in N_{j,2}.$$

where the ℓ_i are i.i.d. standard Laplace random variables. Let the refined estimator for the *j*-th co-ordinate be

$$\tilde{\theta}_j = \frac{1}{n/(2d)} \sum_{i \in N_{j,2}} \tilde{\theta}^{(i)} \in \mathbb{R}.$$
(36)

Step 4. (Consolidation.) The final estimator is defined as

$$\hat{\theta} = R_d^{-1} \sum_{j=1}^d e_j \tilde{\theta}_j \in \mathbb{R}^d,$$
(37)

where e_j is the *j*-th standard basis vector of \mathbb{R}^d .

In addition to the random rotation technique that is used explicitly in the above algorithm, we also use the fact that the ℓ_2 -norm of an average of random variables that are supported on an ℓ_2 -ball concentrates better than only using the fact that the random variables are sub-Gaussian due to having bounded support. This is because, due to being supported on an ℓ_2 -ball, the ℓ_2 -norms of the random variables are themselves bounded random variables, and hence satisfy the sub-Gaussian tail bound; see Jin et al. (2019) for a detailed discussion.

3.4 Discussion

In this section we derived the user-level LDP minimax rates for mean estimation problems. One interesting observation is that the user-level rate for n users each with sample size T is equivalent to the item-level rate with nT users, until T is sufficiently large that the infinite-T rate from Proposition 3 dominates. Similar limiting behaviour is observed in the central user-level privacy setting in Levy et al. (2021) where the minimax risk of mean estimation (on both the ℓ_2 - and ℓ_{∞} -balls) is lower bounded independently of T.

We consider the following table which compares the minimax rates for a range of differential privacy settings, and the non-private case as a baseline, all with a total sample size of nT. We note that despite using the strictest notion of privacy of all the settings, our rates still compare favourably, matching those of item-level LDP with an equivalent number of samples when T is smaller than some quantity exponential in $n\alpha^2$, and with an exponentially decaying rate otherwise.

Setting	Minimax rate	Reference
Non-Private	d/(nT)	e.g. Wu (2016)
Local Item-Level	$d/(nT\alpha^2)$	Duchi et al. (2018)
Local User-Level $(T$ -small)	$d/(nT\alpha^2)$	Theorem 6
Local User-Level $(T-large)$	$e^{-n\alpha^2/d}$	Theorem 6
Central Item-level	$d/(nT) + d/(n^2T^2\alpha^2)$	e.g. Levy et al. (2021)
Central User-level $(T$ -small)	$d/(nT) + d/(n^2T\alpha^2)$	Levy et al. (2021)

Table 2: Comparison of minimax optimal rates for *d*-dimensional mean estimation of distributions supported on the unit ℓ_2 -ball with total sample size of nT in all. Dependence on multiplicative constants, constants in exponents, and logarithmic factors are suppressed for brevity.

One notable difference between the central and local settings however, taking estimation on the ℓ_2 ball as an example, is the linear rather than quadratic factor of 1/T in the second term for the central user-level rate in Table 2. In particular, this shows that under central privacy constraints it is not the case that the user-level rate with n users and T independent observations is the same as the item-level rate with nT users. Hence, a key difference between the behaviour of central and local user-level privacy for mean estimation is that in the local case, as long as T is not too large, there is no difference in the minimax error rate between the item-level problems with an equivalent total sample size, whereas in the central case the user-level problem is strictly harder for all values of T.

Comparing our results to existing results on mean estimation under user-level local differential privacy in Girgis et al. (2022) and Bassily and Sun (2023), these previous works only consider the case of the ℓ_{∞} -ball, that is, the class \mathcal{P}_d , and impose further sub-Gaussianity assumptions beyond that which is already implied by being supported on an ℓ_{∞} -ball. On the other hand, in Theorem 6, we make no such assumptions. This leads to a different optimal rate for the family \mathcal{P}_d , where a quadratic rather than linear dependence on the dimension is observed. We further consider the case of distributions supported on ℓ_2 -balls in the family \mathcal{P}'_d . To the best of our knowledge, this is the first consideration of distributions compactly supported on ℓ_2 -balls under user-level LDP constraints as opposed to ℓ_{∞} -balls, and the resulting different dependence on the dimension d. We consider the full range of possible values of T whereas these prior works only consider the case that T is not larger than a quantity exponential in $n\alpha^2/d$.

Lastly, regarding the assumptions on the support of the distributions in Theorem 6, we remark that it would be sufficient to assume that the data are sub-Gaussian rather than bounded. This is because the localisation step only requires the parameter space to be compact, though some minor adjustments would be needed for data lying outside of candidate sub-intervals, and the compact support is used only through its implication that the data are sub-Gaussian.

4 Sparse mean estimation

In this section, we consider the problem of user-level LDP sparse mean estimation. Recalling the family $\mathcal{P}_{d,s}$ defined in (16), we estimate the mean of a distribution $P \in \mathcal{P}_{d,s}$ so that $\theta(P) = \mathbb{E}_P(X)$ and measure the error by the squared ℓ_2 -loss. **Theorem 8.** For s such that $16 \log(ed)/3 \le s \le d$, assuming that $n\alpha^2 > \widetilde{C}s \log(ed)$, where $\widetilde{C} > 0$ is an absolute constant, we then have that

$$\left[\frac{s}{T} \wedge s \left\{ \left(1 + \frac{d}{n\alpha^2}\right)^{1/T} - 1 \right\} \right] \vee e^{-\frac{Cn\alpha^2}{s}} \lesssim \mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_2^2) \\ \lesssim \min\left\{s, \frac{s \log(dnT\alpha^2)}{T} + e^{-cn\alpha^2/s}, \frac{sd\{\log(dnT\alpha^2)\}^2}{nT\alpha^2} + e^{-cn\alpha^2/d}\right\}, \quad (38)$$

where $\mathcal{P}_{d,s}$ is the family of distributions as in (16), $\mathcal{R}_{n,T,\alpha}$ is the user-level minimax risk rate defined in (3) and C > c > 0 are absolute constants.

Remark 3. The estimation procedures constructed to derive the upper bounds hold for any $s \in [d]$. The assumption that $s \geq 16 \log(ed)/3$ is only required for the lower bound. For the regime that $s < 16 \log(ed)/3$ or the case that $16 \log(ed)/3 > d$, more refined analysis might be required and we will leave this for future work. We also remark that in $s \log(dnT\alpha^2)/T$, the term d is unavoidable based on the current proof. The term $sd\{\log(dnT\alpha^2)\}^2/(nT\alpha^2)$, however, can be sharpened to $sd\{\log(nT\alpha^2)\}^2/(nT\alpha^2)$, which in fact is the upper bound we derived in the proof. We keep the current presentation to facilitate comparison between the two terms.

In the lower bound, there are two terms. The second is the infinite-T rate inherited from Theorem 1, and the first by an application of a strengthened form of Assouad's lemma. These lower bounds are proven in Appendix C and the upper bounds follow from the analysis, in Appendix B, of two constructed estimators to be introduced shortly.

We first inspect the complex first term of the lower bound and consider its form in two different regimes under the condition $T \gtrsim \log\{d/(n\alpha^2)\}$, though we briefly note that in the case that T = 1 we obtain the same item-level lower bound to that in Acharya et al. (2023a).

Regime 1. Consider $n\alpha^2 \lesssim d^{\gamma}$ for some $0 < \gamma \leq 1$. This is the usual regime of interest for sparse problems, where the effective sample size is smaller than the ambient dimension. Focusing on the $\gamma < 1$ case for simplicity, we derive the form of the lower bound, noting that the same rate can be obtained when $\gamma = 1$ with a similar calculation.

When $0 < \gamma < 1$, we have

$$\left(1+\frac{d}{n\alpha^2}\right)^{1/T} - 1 \asymp \frac{\log\{1+d/(n\alpha^2)\}}{T} \gtrsim \frac{\log(1+d^{1-\gamma})}{T} \asymp \frac{\log(d)}{T},$$

where the first equality up to constants holds as the expression is bounded due to the assumption $T \gtrsim \log\{d/(n\alpha^2)\}$. By the condition that $s \gtrsim \log(d)$, the lower bound takes the form $s/T \vee e^{-Cn\alpha^2/s}$.

For the upper bound, we note that the second term in the minimum is smaller than the third in this regime, giving an upper bound of $\min\{s, s \log(dnT\alpha^2)/T + e^{-cn\alpha^2/s}\}$. Consider the mild condition that $s \log(dnT\alpha^2)/T + e^{-cn\alpha^2/s} \leq s$. We obtain an upper bound of the order $s \log(dnT\alpha^2)/T + e^{-cn\alpha^2/s}$, matching the lower bound and leading to a minimax rate of order, up to logarithmic factors,

$$s/T \vee e^{-Cn\alpha^2/s}$$

Regime 2. Consider $n\alpha^2 \geq \widetilde{C}d\log(dnT\alpha^2)$ where \widetilde{C} is as in Theorem 8. We first note in this regime that

$$e^{-cn\alpha^2/s} \le e^{-cn\alpha^2/d} \le e^{-c\widetilde{C}\log(dnT\alpha^2)} \le \frac{1}{dnT\alpha^2},$$

for \tilde{C} sufficiently large. Hence, we have that the polynomial term dominates in the lower bound, and the polynomial terms dominate in both the second and third terms in the minimum in the upper bound.

We then note that

$$\left(1 + \frac{d}{n\alpha^2}\right)^{1/T} - 1 \asymp \frac{d}{nT\alpha^2},$$

where the equality up to constants holds as $d/(n\alpha^2)$ is bounded in this regime. Hence, the lower bound takes the form $sd/(nT\alpha^2)$.

For the upper bound, in this regime we have that $s/T \ge \tilde{C}sd\log(dnT\alpha^2)/(nT\alpha^2)$ and so the third term in the minimum is less than the second term. This gives an upper bound of min $\{s, sd\{\log(dnT\alpha^2)\}^2/(nT\alpha^2)\}$. Consider the mild condition that $sd\{\log(dnT\alpha^2)\}^2/(nT\alpha^2) \le s$. We obtain an upper bound of the order $sd\{\log(dnT\alpha^2)\}^2/(nT\alpha^2)$, matching the lower bound and leading to a minimax rate of order, up to logarithmic factors,

$$sd/(nT\alpha^2)$$

The upper bound in Theorem 8 is obtained by a two-component procedure detailed in Section 4.1. We note that the procedure requires the knowledge of the true sparsity level s, the consequences of the misspecification will be discussed in Section 4.2.

4.1 Estimation procedures

Depending on the value of T, we employ one of two estimation procedures. The first is used when T is sufficiently large so that each user can locally identify the *s*-many non-zero entries of the mean to reduce the problem to an *s*-dimensional mean estimation problem. When T is small, we threshold the estimator obtained by naively applying the ℓ_{∞} -ball estimator to improve its performance under the sparsity assumption.

In particular, the first procedure is conducted when $n\alpha^2 < \tilde{C}d\log(dnT\alpha^2)$ for \tilde{C} as in Theorem 8, and the second procedure otherwise. It is easy to verify that with this choice we pick the estimator with the smaller error of the two where the first and second procedures have errors corresponding to the second and third terms respectively in the minimum in the upper bound in (38).

4.1.1 Variable selection-based procedure

For the first procedure, deployed when $n\alpha^2 \leq \tilde{C}d\log(dnT\alpha^2)$, the first half of users carry out variable selection locally with their own data to identify the *s*-many non-zero entries of the mean. Then, using a hashing-based voting method inspired by Zhou et al. (2022), the users vote for the co-ordinates they believe containing the non-zero entries. The other half of users then conduct *s*-dimensional mean estimation on the resulting ℓ_{∞} -ball.

The procedure is formalised as follows.

Step 1. (Initialisation.) Let $T^* = T \wedge e^{n\alpha^2/(Ks)} \geq 1$ with K > 0 an absolute constant and let

$$\varepsilon = \left\{ \frac{2\log(dnT^*\alpha^2)}{T^*} \right\}^{1/2}.$$
(39)

Step 2. (Localisation.) Let $\hat{\omega}^{(i)}$ be the *d*-dimensional binary vector with entries

$$\hat{\omega}_{j}^{(i)} = \mathbb{1}\left\{ \left| \frac{1}{T^{*}} \sum_{t=1}^{T^{*}} X_{t,j}^{(i)} \right| \ge \varepsilon \right\}, \quad i \in [n/2], \ j \in [d].$$

$$(40)$$

Let

$$Z_i = \Pi_{[-\eta,\eta]} \left(\sum_{j=1}^d r_{i,j} \hat{\omega}_j^{(i)} \right) + \frac{2\eta}{\alpha} \ell_i, \quad i \in [n/2]$$

where $r_{i,j}$ for $i \in [n/2]$, $j \in [d]$ are i.i.d. Rademacher random variables, ℓ_i for $i \in [n/2]$ are i.i.d. standard Laplace random variables and $\eta = (30s)^{1/2}$.

Let

$$\mathcal{I} = \left\{ j \in [d] : \frac{2}{n} \sum_{i=1}^{n/2} r_{i,j} Z_i \ge 1/2 \right\}.$$
(41)

Step 3. (Estimation.) If $|\mathcal{I}| \leq s$, let $\mathcal{I}' = \mathcal{I}$, and if $|\mathcal{I}| > s$ let \mathcal{I}' be the first s elements of \mathcal{I} , so that $\mathcal{I}' = \{\iota_1, \iota_2, \ldots, \iota_{s'}\}$ where $1 \leq \iota_1 < \iota_2 < \ldots < \iota_{s'}$ and $s' \leq s$. Let

$$Y_t^{(i)} = \left(X_{t,\iota_1}^{(i)}, \dots, X_{t,\iota_{s'}}^{(i)}\right), \quad i \in [n] \setminus [n/2], \ t \in [T].$$

Apply the ℓ_{∞} -ball estimation procedure, i.e. Steps 1-4 in Section 3.2, on data $\{Y_t^{(i)}\}_{i \in [n], t \in [T]}$ to obtain estimator $\hat{\theta} \in \mathbb{R}^{s'}$. The final estimator $\hat{\theta}$ in \mathbb{R}^d is defined via its co-ordinates

$$\hat{\theta}_j = \begin{cases} \bar{\theta}_j, & j \in \mathcal{I}_{s'}, \\ 0, & \text{otherwise,} \end{cases} \quad j \in [d].$$

$$\tag{42}$$

The strength of this variable selection method comes from the effectiveness of the hashing-based voting method to identify the non-zero entries of the mean vector. To identify these non-zero entries, one could naively use the voting procedure in Section 3, which involves either (1) splitting users to estimate each co-ordinate separately or (2) privatising a *d*-dimensional binary vector with *s*-many non-zero entries (c.f. Lemma 16). These two strategies result in factors *d* and s^2 in the final rate, respectively. Instead, the hashing-based method compresses the votes of the users to a one-dimensional quantity without splitting users, by exploiting Rademacher random variables. The votes for a specific co-ordinate can then be extracted from this univariate quantity and the final rate is merely inflated by a factor of $\log(d)$.

4.1.2 Thresholding-based procedure

For the second procedure, conducted when $n\alpha^2 > \tilde{C}d\log(nT\alpha^2)$, we apply the ℓ_{∞} -ball procedure of Section 3.2 with an additional thresholding step applied to the estimator thus obtained. This reduces the d^2 factor in the polynomial term of the error for the ℓ_{∞} -ball procedure in (24) to a factor of sd.

The algorithm is formalised as follows.

Step 1. (Initialisation.) For $j \in [d]$, let

$$N_{j,1} = \left\{ (j-1)\frac{n}{2d} + 1, \dots, j\frac{n}{2d} \right\}, N_{j,2} = \left\{ (d+j-1)\frac{n}{2d} + 1, \dots, (d+j)\frac{n}{2d} \right\}, N_j = N_{j,1} \cup N_{j,2}.$$

Let $T^* = T \wedge e^{n\alpha^2/(Kd)} \ge 1$ with K > 0 an absolute constant and let

$$\varepsilon = C'\delta \left\{ \frac{d\log(nT^*\alpha^2)}{n\alpha^2} \right\}^{1/2},\tag{43}$$

where C' > 0 is an absolute constant and

$$\delta = \left\{ 4 \log(nT^* \alpha^2) / T^* \right\}^{1/2}.$$
 (44)

Let $\{I_j, j \in [N(\delta)]\}$ be a partition of [-1, 1], with $I_j = [-1 + 2(j-1)\delta, -1 + 2j\delta), j \in [N(\delta) - 1]$ and $I_{N(\delta)} = [-1 + 2(N(\delta) - 1)\delta, -1 + 2N(\delta)\delta]$.

Step 2. (Estimation.) For each $j \in [d]$, perform the following.

Calculate

$$\tilde{\theta}^{(i)} = \frac{1}{T^*} \sum_{t=1}^{T^*} X_t^{(i)} \text{ and } V_k^{(i)} = \mathbb{1}\{\tilde{\theta}_j^{(i)} \in I_{k-1} \cup I_k \cup I_{k+1}\}, \quad i \in N_{j,1}, \, k \in [N(\delta)], \tag{45}$$

with $I_0 = I_{N(\delta)+1} = \emptyset$. Let $U_{i,k}$, $i \in N_{j,1}$, $k \in [N(\delta)]$, be i.i.d. from Unif[0, 1] and set

$$\widetilde{V}_{k}^{(i)} = \begin{cases} V_{k}^{(i)}, & U_{i,k} \le e^{\alpha/6}/(1+e^{\alpha/6}), \\ 1-V_{k}^{(i)}, & \text{otherwise}, \end{cases} \quad i \in N_{j,1}, \, k \in [N(\delta)]. \end{cases}$$

Choose

$$k_j^* = \min \argmax_{k \in [N(\delta)]} \sum_{i \in N_{j,1}} \widetilde{V}_k^{(i)}.$$

Let L_j, U_j denote the lower and upper end points of $I_{k_i^*}$ respectively,

$$\tilde{L}_j = L_j - 6\delta, \quad \tilde{U}_j = U_j + 6\delta \quad \text{and} \quad \tilde{I}_{k_j^*} = [\tilde{L}_j, \tilde{U}_j].$$
(46)

Step 3. (Refinement.) For each $j \in [d]$, perform the following:

Let

$$\tilde{\theta}^{(i)} = \Pi_{\tilde{I}_{k_j^*}} \left(\frac{1}{T^*} \sum_{t=1}^{T^*} X_{t,j}^{(i)} \right) + \frac{14\delta}{\alpha} \ell_i, \quad i \in N_{j,2}$$

where the ℓ_i are i.i.d. standard Laplace random variables. Let the refined estimator for the *j*-th co-ordinate be

$$\tilde{\theta}_j = \frac{1}{n/(2d)} \sum_{i \in N_{j,2}} \tilde{\theta}^{(i)} \in \mathbb{R}.$$

Step 4. (Consolidation.) The interim estimator is defined as

$$\tilde{\theta} = \sum_{j=1}^d e_j \hat{\theta}_j \in \mathbb{R}^d,$$

where e_j is the *j*-th standard basis vector of \mathbb{R}^d .

Step 5. (Truncation.) The thresholded interim estimator $\tilde{\theta}_{\text{thresh}} \in \mathbb{R}^d$ is defined with co-ordinates

$$\tilde{\theta}^{\text{thresh}} = \tilde{\theta}_j \mathbb{1}\{|\tilde{\theta}_j| > 3\varepsilon\}, \quad j \in [d].$$
(47)

Denote a set of the *s*-many largest entries of $\tilde{\theta}^{\text{thresh}}$ by $\mathcal{I} = (\iota_1, \ldots, \iota_s)$ where $|\tilde{\theta}^{\text{thresh}}_{\iota_j}| \ge |\tilde{\theta}^{\text{thresh}}_{\iota_{j-1}}|$ for $1 < j \le s$. The final estimator $\hat{\theta} \in \mathbb{R}^d$ is defined by taking the largest *s* entries of $\tilde{\theta}^{\text{thresh}}$ via

$$\hat{\theta}_j = \begin{cases} \tilde{\theta}_j^{\text{thresh}} & \text{if } j \in \mathcal{I}, \\ 0 & \text{otherwise.} \end{cases}$$
(48)

4.2 Discussion

This section demonstrates, through sparse mean estimation, an important example where in some regimes the user-level framework provides a decidedly superior minimax rate over the item-level framework with an equivalent number of total data points, in contrast to the findings of Section 3. In particular, even as $d/(n\alpha^2) \to \infty$ we saw that, as long as $\log(d/n\alpha^2)/T \to 0$, the rate is improved to $s \log(d)/T$ up to further logarithmic factors compared to the item-level rate with nT observations of $sd/(nT\alpha^2)$ (Acharya et al., 2023a). The enabling factor for this is that with more than $\log\{d/(n\alpha^2)\}$ -many observations a user can, with high probability, successfully identify the relevant non-zero co-ordinates locally, without the need for any communication with the data aggregator. This suggests that similar phenomena of regimes with superior user-level performance may exist in other problems where there is similar potential for localisation.

The following table includes comparisons of the minimax rates for a range of differential privacy settings, and the non-private case as a baseline, all with an effective sample size of nT. We note, similarly to Section 3, our rates still compare favourably, in fact out-performing local item-level in some regimes, despite using the strictest notion of privacy.

Setting	Minimax rate	Reference
Non-Private	$s\log(ed/s)/(nT)$	e.g. Han (2016)
Local Item-Level	$sd/(nT\alpha^2)$	Acharya et al. (2023a)
Local User-Level $(n, T$ -small)	s/T	Theorem 8
Local User-Level $(n$ -large, T -small)	$sd/(nT\alpha^2)$	Theorem 8
Local User-Level $(T$ -large)	$e^{-n\alpha^2/s}$	Theorem 8
Central Item-level ²	$s \log(d) / (nT\alpha) + \{s \log(d)\}^2 / (n^2 T^2 \alpha^2)$	Cai et al. (2021)

Table 3: Comparison of minimax optimal rates for *d*-dimensional *s*-sparse mean estimation of distributions supported on the unit ℓ_{∞} -ball, all with effective sample size of nT. Dependence on multiplicative constants, constants in exponents, and logarithmic factors are suppressed for brevity.

²In the absence of existing results for central item-level privacy for estimation of sparse means on the ℓ_{∞} -ball, we compare to results obtained under Gaussian distribution assumptions.

Figure 1 below provides an illustration of the difference in performance between the user-level setup, and that of item-level with an equivalent number of samples, depending on the relationship between the values of $n\alpha^2/d$ and T. We note again that this is a harsh comparison for the user-level framework as the α privacy budget has to simultaneously protect across all T observations a user holds, as opposed to the item-level setting where each observation enjoys the entire α budget. In Figure 1a, we see that for estimation on an ℓ_2 -ball, user-level is never better than item-level, with equivalent performance as long as T is not too large. In contrast, in Figure 1b, the behaviour where user-level outperforms item-level in certain regimes is apparent.



Figure 1: Comparison of performance of user-level LDP and nT-item level LDP estimator performance. Constants, constants in exponents, and logarithmic factors are omitted for ease of comparison.

Mis-specification of *s*. Recalling that both of procedures introduced in Section 4.1 require the true sparsity level *s* as an input, we remark on the consequences of mis-specification. If underestimated, i.e. the input $s_1 < s$, then each method suffers an additional bias of at most $s - s_1 \leq s$. In the selection-based procedure, this arises due to failing to select a non-zero co-ordinate. For the thresholding-based procedure, this arises at the final step when only the largest $s_1 < s$ co-ordinates of the estimator are kept. This gives an upper bound on the risk of the estimator of

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_{2}^{2}\right] \lesssim s - s_{1} + \min\left\{s, \frac{s_{1}\log(dnT\alpha^{2})}{T} + e^{-cn\alpha^{2}/s_{1}}, \frac{s_{1}d\{\log(dnT\alpha^{2})\}^{2}}{dnT\alpha^{2}} + e^{-cn\alpha^{2}/d}\right\}.$$

If overestimated, i.e. $s_2 > s$, then no extra bias will be introduced but larger variance will be due to estimating more co-ordinates. The risk of both estimation procedures see s replaced with s_2 accordingly, giving the upper bound

$$\mathbb{E}\left[\|\hat{\theta}-\theta\|_2^2\right] \lesssim \min\left\{s, \frac{s_2\log(dnT\alpha^2)}{T} + e^{-cn\alpha^2/s_2}, \frac{s_2d\{\log(dnT\alpha^2)\}^2}{dnT\alpha^2} + e^{-cn\alpha^2/d}\right\}.$$

As such, we see that in practice, it is generally safer to choose a relatively large s when the truth is unknown.

Choice of the estimation procedure. In Section 4.1 we recommended to use the first procedure – the variable selection-based method when $n\alpha^2 \leq \tilde{C}d\log(dnT\alpha^2)$, and the second procedure – the thresholding-based method otherwise. We now discuss the estimation error upper bound if an estimator is chosen against recommendation.

As we mentioned at the beginning of Section 4.1, the error control of the first procedure corresponds to the second term in the upper bound in (38), and the second procedure corresponds to the third term. When the choice is made against our recommendation, i.e. chosen when $n\alpha^2 > \tilde{C}d\log(dnT\alpha^2)$, the error control of the first procedure still holds. Whereas for the second procedure, the error control will change. To be specific, when $n\alpha^2 \approx d$, the corresponding error control will always be bounded below from a positive constant. This is due to the term $e^{-n\alpha^2/(Kd)}$ arising from an application of the *d*-dimensional ℓ_{∞} -ball estimator of Section 3. From this, we see that if $n\alpha^2 \simeq d$, the error contains a constant term bounding the error away from zero.

Previous work with the same terminology. Zhou et al. (2022) share the term of user-level privacy for sparse mean estimation with us, but the setups considered differ. In Zhou et al. (2022), it is assumed that each user holds a single, possibly high-dimensional, vector. The item-level privacy in Zhou et al. (2022) is to protect against a perturbation of a single entry of this vector, whereas user-level privacy protects against any two different vectors the user may hold. In our setup, we assume that each user holds multiple, possibly high-dimensional, vectors. The item-level considered in our paper corresponds to protecting against the perturbation of one observation from a user's collection, regardless of how many entries are perturbed, whilst user-level corresponds to the entire collection, perturbing both every vector and every co-ordinate of any vector. Further, in Zhou et al. (2022) sparsity is defined to mean that at most *s*-many entries of a user's vector will be non-zero, whereas in our work, the sparsity assumption is applied to the mean vector, whilst all the entries of a user's data may be non-zero.

We also note that whilst we implement the same hashing method as Zhou et al. (2022), our analysis of its performance is different. In terms of the inference goal, Zhou et al. (2022) aim at deriving a high-probability ℓ_{∞} -upper bound on the estimation error, whilst we conduct variable selection based on binary vectors. In terms of the technical details, the truncation intervals in our paper have width $O(s^{1/2})$, compared to $O(s^{1/2} \log(n/\beta))$ in Zhou et al. (2022). This is achieved through a tighter analysis and different inference goals, and allows us to avoid further logarithmic factors in the error controls.

5 Non-parametric density estimation

In this section, we consider the user-level LDP non-parametric density estimation problem. Recalling the Sobolev class of densities $\mathcal{F}_{\beta,r}$ defined in (20), for a density $f \in \mathcal{F}_{\beta,r}$, we denote by P_f the induced distribution the data are drawn from. Our aim is to estimate the density $\theta(P_f) = f$ and we measure error through the L_2 -norm as defined in (22). We have the following result.

Theorem 9. Assuming that $n\alpha^2 \geq \widetilde{C}_{\beta}$, where $\widetilde{C}_{\beta} > 0$ is an absolute constant depending only on β , we then have that

$$c_{\beta}\left\{(nT\alpha^{2})^{-\frac{2\beta}{2\beta+2}} \vee (n\alpha^{2})^{-2\beta}\right\} \lesssim \mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}_{\beta,1}), \|\cdot\|_{2}^{2}) \lesssim C_{\beta}\left\{(nT\alpha^{2})^{-\frac{2\beta}{2\beta+2}}\log(nT\alpha^{2}) + (n\alpha^{2})^{-2\beta}\left\{\log(n\alpha^{2})\right\}^{2\beta+3}\right\},$$

where $\mathcal{P}_{\beta,1}$ is the family of distributions as in (21), $\mathcal{R}_{n,T,\alpha}$ is the user-level minimax risk rate defined in (3), and $C_{\beta} > c_{\beta} > 0$ are absolute constants depending only on β .

We see that the lower bound in Theorem 9 consists of two parts, the first analogous to the standard item-level rate (Duchi et al., 2018, Corollary 4), and the second inherited from Theorem 1. Unlike in the mean estimation problems we considered, the rate in the large T regime is polynomial rather than exponential, appearing to be a non-parametric rate. Nevertheless, this again demonstrates the intuition of Section 2 wherein we see that the rate for sufficiently large T coincides with the infinite-T rate in Proposition 5. The lower bounds is proven in Appendix C. An estimation procedure achieving the upper bound is presented below, the performance of which is analysed in Appendix B.

5.1 Estimation procedure

The trigonometric basis (18) allows us to translate a density estimation problem into mean estimation, where we estimate a truncation of the sequence $\{\theta_j\}_{j\geq 1}$ of basis coefficients given in (19). In particular, we use the fact that for any $f \in \mathcal{F}_{\beta,r}$ and $j \in \mathbb{N}$, we have $\int f(x)\varphi_j(x) dx = \theta_j$ by the orthonormality of the trigonometric basis. Hence, if the data $\{X_i\}_{i\in[n]}$ are i.i.d. from the distribution P_f , then the sample mean of $\{\varphi_j(X_i)\}_{i\in[n]}$ is a consistent estimator of θ_j . Importantly, for all $j \in \mathbb{N}$, we have that $\sup_x |\varphi_j(x)| \leq \sqrt{2}$, allowing us (after rescaling) to invoke the results of Section 3 for estimation on a bounded ℓ_{∞} -ball.

The procedure is formalised as follows.

Step 1. (Initialisation.) Let

$$M = \begin{cases} (nT\alpha^2)^{1/(2\beta+2)}, & T \le (n\alpha^2)^{2\beta+1} / \{C\log(n\alpha^2)\}^{2\beta+2}, \\ n\alpha^2 / \{C\log(n\alpha^2)\}, & \text{otherwise.} \end{cases}$$
(49)

where C > 0 is some sufficiently large constant depending only on β .

Let

$$Y_t^{(i)} = (\varphi_j(X_t^{(i)}), j \in [M])^\top \in [0, \sqrt{2}]^M, \quad i \in [n], t \in [T].$$

Step 2. (Estimation.) Apply the ℓ_{∞} -ball estimation procedure, i.e. Steps 1-4 in Section 3.2, on data $\{Y_t^{(i)}\}_{i \in [n], t \in [T]}$ to obtain estimator $\hat{\theta} \in \mathbb{R}^M$. The final estimator is defined as

$$\hat{f} = \sum_{j=1}^{M} \hat{\theta}_j \varphi_j.$$
(50)

In a standard non-parametric projection estimation procedure, one estimates the density by obtaining estimates of some finite number of coefficients through calculating the sample mean of the basis functions applied to data (e.g. Tsybakov, 2009, Chapter 1). The item-level LDP non-parametric estimation procedure in Duchi et al. (2018) is a private analogue, where the mean estimation step is carried out using a suitable (item-level) LDP mean estimation technique. We see that in the user-level setting, our estimator \hat{f} in (50) operates in a similar fashion. The only adjustment is to use a user-level LDP mean estimation technique.

As for the cut-off level M, the larger it is, the less approximation bias is introduced but the greater the variance associated with a higher dimensional mean estimation problem is. As a sketch, one can see that for T sufficiently large, the error introduced by the mean estimation component is $O\{\exp(n\alpha^2/M)\}$ and the error from truncation is $O(1/M^{2\beta})$. Balancing these two terms yields the choice of M and consequently the upper bound in Theorem 9.

5.2 Discussion

In this section, we saw that the user-level rate for n users with sample size T is equivalent up to the item-level rate with nT users, until T is sufficiently large so that the infinite-T rate dominates, as was the case with the mean estimation problems of Section 3 and unlike sparse mean estimation as in Section 4.

The comparisons of the minimax rate for a range of differential privacy settings, and the non-private case as a baseline, all with a total sample size of nT, are presented in Table 4. We start by noting that, the central item-level rate consists of the sum of two terms, the non-private rate $(nT)^{-2\beta/(2\beta+1)}$ and the cost for privacy $(nT\alpha)^{-2\beta/(\beta+1)}$. As is usual, provided α is not too small, the privacy cost is dominated by the non-private rate.

On the other hand, for the local item-level setting, we see that the privacy cost dominates the nonprivate rate. This behaviour is again seen in the local user-level setting when T is small. Further, the rate for T sufficiently large is polynomial, albeit smaller than the small-T case, owing to the larger metric entropy of the function space and the result of Corollary 2.

Setting	Minimax rate	Reference
Non-Private	$(nT\alpha^2)^{-2\beta/(2\beta+1)}$	e.g. Tsybakov (2009)
Local Item-Level	$(nT\alpha^2)^{-2\beta/(2\beta+2)}$	Duchi et al. (2018)
Local User-Level $(T$ -small)	$(nT\alpha^2)^{-2\beta/(2\beta+2)}$	Theorem 9
Local User-Level $(T-large)$	$(n\alpha^2)^{-2\beta}$	Theorem 9
Central Item-Level	$(nT)^{-2\beta/(2\beta+1)} + (nT\alpha)^{-2\beta/(\beta+1)}$	Cai et al. (2021)

Table 4: Comparison of minimax optimal rates for non-parametric density estimation for β -Sobolev smooth densities supported on [0, 1] with effective sample size of nT in all. Dependence on multiplicative constants, constants in exponents, and logarithmic factors are suppressed for brevity.

Regarding existing results on user-level privacy, Acharya et al. (2023b) show that the minimax risk of discrete density estimation (under the high privacy regime $\alpha \leq 1$) for a distribution with N outcomes is $O\{N/(nT\alpha^2)\}$. However, this holds only for the regime $n\alpha^2 \leq CN \log(T)$ for some constant C > 0, and what happens outside of this regime is not considered. In light of our results on mean estimation, this regime suggests a phase transition may occur once T is exponential in $n\alpha^2/N$, akin to d-dimensional mean estimation on the ℓ_2 -ball. This is perhaps to be expected as discrete density estimation is similarly a parametric problem. For a sketch for the lower bound, suppose that $\delta > 0$ is sufficiently small and is such that $1/\delta$ is an integer. One can construct a 2δ -separated packing with respect to the L_2 -norm for densities (or equivalently, the ℓ_2 -norm for the vector of probabilities of outcomes) of the space of discrete densities on N outcomes, indexed by the probability vector $p \in [0, 1]^N$ denoting the probability of each outcome, as

$$\mathcal{V} = \{ p \in \{0, \delta, \dots, (1/\delta - 1)\delta, 1\}^N : p_1 + \dots + p_N = 1 \}$$

= $\{ \delta x : x \in \{0, 1, \dots, 1/\delta\}^N, \, \delta x_1 + \dots + \delta x_N = 1 \}.$

In particular, by a standard stars-and-bars counting method (e.g. Feller, 1968, Chapter 3), it can be shown that

$$|\mathcal{V}| = \binom{1/\delta - 1}{N - 1} \ge \left(\frac{1/\delta - 1}{N - 1}\right)^{N - 1}.$$

Assuming that $n\alpha^2 \ge C(N-1)\log(e(N-1))$ for some sufficiently large absolute constant C > 0 and setting

$$\delta = \left\{ 1 + (N-1) \exp\left(\frac{24n\alpha^2 + 2\log(2)}{N-1}\right) \right\}^{-1},\$$

we have that

$$\begin{aligned} \mathcal{R}_{n,\infty,\alpha} \Big(\theta(\mathcal{P}), \|\cdot\|_2^2 \Big) \gtrsim \left\{ 1 + (N-1) \exp\left(\frac{24n\alpha^2 + 2\log(2)}{N-1}\right) \right\}^{-2} \\ \gtrsim \frac{1}{(N-1)^2} \exp\left(\frac{-24n\alpha^2 - 2\log(2)}{N-1}\right) \gtrsim e^{-cn\alpha^2/(N-1)} \end{aligned}$$

where c > 0 is an absolute constant, the first inequality is by an application of Theorem 1, and the last two both use the assumption $n\alpha^2 \ge C(N-1)\log(e(N-1))$.

Hence, we see that the infinite-T minimax lower bound for discrete density estimation is exponential, matching that for mean estimation rather than the polynomial rate for non-parametric density estimation.

6 Conclusions

In this paper, we considered a range of canonical statistical problems under the user-level local differential privacy framework. We provided, up to poly-logarithmic factors and constants in exponents when exponential rates occur, matching upper and lower bounds on the minimax rates in all regimes of Tfor non-sparse mean estimation and non-parametric density estimation, and similarly for sparse mean estimation up to a logarithmic gap in the regime. We further characterise phase transition phenomena as the number of samples held by each user varies.

	d -dim. mean (ℓ_2 -ball)	s-sparse mean	Density
Small- T rate	$d/(nT\alpha^2)$	$\min\{s/T, sd/(nT\alpha^2)\}$	$(nT\alpha^2)^{-2\beta/(2\beta+2)}$
Large- T rate	$e^{-n\alpha^2/d}$	$e^{-n\alpha^2/s}$	$(n\alpha^2)^{-2\beta}$
Phase transition $(T = \cdot)$	$O(e^{n\alpha^2/d})$	$\begin{cases} O(e^{n\alpha^2/s}), & d/(n\alpha^2) \gtrsim 1\\ O(e^{n\alpha^2/d}), & d/(n\alpha^2) \lesssim 1 \end{cases}$	$O\left((n\alpha^2)^{2\beta+1}\right)$

Table 5: Comparison of user-level rates and phase transitions. Dependence on multiplicative constants, constants in exponents, and logarithmic factors are suppressed for brevity.

Minimax optimal rates for user-level LDP problems, in particular proofs of lower bounds, have not been studied systematically before. The only existing instance of such work (Acharya et al., 2023b) proves matching upper and lower bounds, but restricts to a certain range of T, not considering the behaviour as T diverges. We on the other hand consider all regimes of T. This provides insight on the fundamental limits of these problems in a way that appears to be intrinsically linked to the metric entropy of the parameter space.

First observed in Acharya et al. (2023b), the user-level rate matches the item-level rate with an equivalent number of users in the "small-T" regime in discrete distribution estimation problems. It was not known whether this was a special case or more general. We see the same behaviour in mean and non-parametric density estimation problems, but different behaviour for sparse mean estimation problems, bucking this trend and showing it is not universal behaviour for user-level LDP estimation.

It is also particularly striking as the high-dimensional sparse mean estimation problem is intractable under item-level privacy, but becomes feasible under the user-level framework. This itself may have useful applications in practice, but could also motivate further exploration to identify other statistical problems where item-level privacy is pessimistic, especially where there is an indication of the ability for users to localise onto relevant factors of the data locally.

Considering the method used to derive the upper bound in Theorem 1, whilst the method only works in the infinite-T case, it suggests a general method that might be applied to the finite-T case. For example, as can be seen from the univariate estimation method in Section 3, one proceeds by covering the parameter space with intervals (resp. balls) of a certain width (resp. radius), using a private voting procedure to select a candidate, and then localising on this candidate to extract the benefits of each user having multiple samples. However, difficulties arise in the finite-T case when trying to vote on a candidate as the error in the estimate of the functional can result in the votes being split over neighbouring candidates, with the number of neighbouring candidates increasing as the dimension increases. If some technique to circumvent these issues can be developed, it would provide a general procedure for finite-T that would provide an upper bound that intrinsically depends on the metric entropy of the space. Combined with the general lower bound of Theorem 8, this would show that the fundamental limits of user-level LDP estimation are entirely dependent on, and can be deduced from, the metric entropy of the parameter space.

Acknowledgements

The second author was supported by Engineering and Physical Sciences Research Council (EPSRC) New Investigator Award EP/W016117/1. The third author was partially supported by the Philip Leverhulme Prize.

References

- Jayadev Acharya, Clément L Canonne, Ziteng Sun, and Himanshu Tyagi. Unified lower bounds for interactive high-dimensional estimation under information constraints. *Advances in Neural Information Processing Systems*, 36:51133–51165, 2023a.
- Jayadev Acharya, Yuhan Liu, and Ziteng Sun. Discrete distribution estimation under user-level local differential privacy. In International Conference on Artificial Intelligence and Statistics, pages 8561– 8585. PMLR, 2023b.
- Kareem Amin, Alex Kulesza, Andres Munoz, and Sergei Vassilvtiskii. Bounding user contributions: A bias-variance trade-off in differential privacy. In *International Conference on Machine Learning*, pages 263–271. PMLR, 2019.
- Raef Bassily and Ziteng Sun. User-level private stochastic convex optimization with optimal rates. In *International Conference on Machine Learning*, pages 1838–1851. PMLR, 2023.
- Thomas B Berrett and Cristina Butucea. Locally private non-asymptotic testing of discrete distributions is faster using interactive mechanisms. *Advances in Neural Information Processing Systems*, 33:3164– 3173, 2020.
- Thomas B Berrett and Yi Yu. Locally private online change point detection. Advances in Neural Information Processing Systems, 34:3425–3437, 2021.
- Mark Bun, Jonathan Ullman, and Salil Vadhan. Fingerprinting codes and the price of approximate differential privacy. *Proceedings of the forty-sixth annual ACM symposium on Theory of computing*, pages 1–10, 2014.
- United States Census Bureau. Statistical Safeguards, 2023. URL https://www.census.gov/about/policies/privacy/statistical_safeguards.html. Accessed: 25-3-2024.
- Cristina Butucea, Amandine Dubois, Martin Kroll, and Adrien Saumard. Local differential privacy: Elbow effect in optimal density estimation and adaptation over Besov ellipsoids. *Bernoulli*, 26(3):1727 – 1764, 2020.

- T Tony Cai and Hongji Wei. Distributed gaussian mean estimation under communication constraints: Optimal rates and communication-efficient algorithms. *Journal of Machine Learning Research*, 25(37): 1–63, 2024.
- T Tony Cai, Yichen Wang, and Linjun Zhang. The cost of privacy: Optimal rates of convergence for parameter estimation with differential privacy. *The Annals of Statistics*, 49(5):2825–2850, 2021.
- T Tony Cai, Yichen Wang, and Linjun Zhang. Score attack: A lower bound technique for optimal differentially private learning. arXiv preprint arXiv:2303.07152, 2023.
- Wei-Ning Chen, Peter Kairouz, and Ayfer Ozgur. Breaking the communication-privacy-accuracy trilemma. Advances in Neural Information Processing Systems, 33:3312–3324, 2020.
- Julien Chhor and Flore Sentenac. Robust estimation of discrete distributions under local differential privacy. In International Conference on Algorithmic Learning Theory, pages 411–446. PMLR, 2023.
- John C Duchi, Michael I Jordan, and Martin J Wainwright. Minimax optimal procedures for locally private estimation. *Journal of the American Statistical Association*, 113(521):182–201, 2018.
- Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. In Shai Halevi and Tal Rabin, editors, *Theory of Cryptography*, pages 265–284, Berlin, Heidelberg, 2006. Springer Berlin Heidelberg.
- Ulfar Erlingsson, Vasyl Pihur, and Aleksandra Korolova. Rappor: Randomized aggregatable privacypreserving ordinal response. In Proceedings of the 2014 ACM SIGSAC conference on computer and communications security, pages 1054–1067, 2014.
- William Feller. An Introduction to Probability Theory and Its Applications. Wiley, New York, NY, 3rd edition, 1968. ISBN 9780471257080.
- Badih Ghazi, Pritish Kamath, Ravi Kumar, Pasin Manurangsi, Raghu Meka, and Chiyuan Zhang. On user-level private convex optimization. In *International Conference on Machine Learning*, volume 202, pages 11283–11299. PMLR, 2023.
- Antonious M Girgis, Deepesh Data, and Suhas Diggavi. Distributed user-level private mean estimation. In 2022 IEEE International Symposium on Information Theory (ISIT), pages 2196–2201. IEEE, 2022.
- Fang Han. Lecture 8: Minimax Lower Bounds: LeCam, Fano, and Assouad, 2016. URL https://sites.stat.washington.edu/people/fanghan/teaching/STAT583/minimax.pdf. Accessed: 25-3-2024.
- Samuel B Hopkins, Gautam Kamath, Mahbod Majid, and Shyam Narayanan. Robustness implies privacy in statistical estimation. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, pages 497–506, 2023.
- Chi Jin, Praneeth Netrapalli, Rong Ge, Sham M Kakade, and Michael I Jordan. A short note on concentration inequalities for random vectors with subgaussian norm. *arXiv preprint arXiv:1902.03736*, 2019.
- Peter Kairouz, H Brendan McMahan, Brendan Avent, Aurélien Bellet, Mehdi Bennis, Arjun Nitin Bhagoji, Kallista Bonawitz, Zachary Charles, Graham Cormode, Rachel Cummings, et al. Advances and open problems in federated learning. *Foundations and trends in machine learning*, 14(1–2): 1–210, 2021.
- Joseph Lam-Weil, Béatrice Laurent, and Jean-Michel Loubes. Minimax optimal goodness-of-fit testing for densities and multinomials under a local differential privacy constraint. *Bernoulli*, 28(1):579–600, 2022.
- Daniel Levy, Ziteng Sun, Kareem Amin, Satyen Kale, Alex Kulesza, Mehryar Mohri, and Ananda Theertha Suresh. Learning with user-level privacy. Advances in Neural Information Processing Systems, 34:12466–12479, 2021.
- Mengchu Li, Thomas B Berrett, and Yi Yu. On robustness and local differential privacy. *The Annals of Statistics*, 51(2):717–737, 2023.

- Xiyang Liu, Weihao Kong, and Sewoong Oh. Differential privacy and robust statistics in high dimensions. In *Conference on Learning Theory*, pages 1167–1246. PMLR, 2022.
- Yuhan Liu, Ananda Theertha Suresh, Felix Xinnan X Yu, Sanjiv Kumar, and Michael Riley. Learning discrete distributions: user vs item-level privacy. Advances in Neural Information Processing Systems, 33:20965–20976, 2020.
- Yurii Lyubarskii and Roman Vershynin. Uncertainty principles and vector quantization. IEEE Transactions on Information Theory, 56(7):3491–3501, 2010.
- Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-efficient learning of deep networks from decentralized data. In Artificial intelligence and statistics, pages 1273–1282. PMLR, 2017.
- Mathieu Sart. Density estimation under local differential privacy and hellinger loss. *Bernoulli*, 29(3): 2318–2341, 2023.
- Alexandre B Tsybakov. Introduction to Nonparametric Estimation. Springer New York, New York, NY, 2009.
- Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science.* Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2018.
- Martin J Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019.
- Tianhao Wang, Jeremiah Blocki, Ninghui Li, and Somesh Jha. Locally differentially private protocols for frequency estimation. In 26th USENIX Security Symposium (USENIX Security 17), pages 729–745, 2017.
- Stanley L Warner. Randomized response: A survey technique for eliminating evasive answer bias. *Journal* of the American Statistical Association, 60(309):63–69, 1965.
- Larry Wasserman and Shuheng Zhou. A statistical framework for differential privacy. Journal of the American Statistical Association, 105(489):375–389, 2010.
- Yihong Wu. Lecture 18: Density estimation and structured estimation, 2016. URL http://www.stat.yale.edu/~yw562/teaching/598/lec18.pdf. Accessed: 23-4-2024.
- Ashkan Yousefpour, Igor Shilov, Alexandre Sablayrolles, Davide Testuggine, Karthik Prasad, Mani Malek, John Nguyen, Sayan Ghosh, Akash Bharadwaj, Jessica Zhao, et al. Opacus: User-friendly differential privacy library in pytorch. arXiv preprint arXiv:2109.12298, 2021.
- Bin Yu. Assouad, Fano, and Le Cam. In Festschrift for Lucien Le Cam, pages 423–435. Springer New York, New York, NY, 1997.
- Razieh Nokhbeh Zaeem and K Suzanne Barber. The effect of the gdpr on privacy policies: Recent progress and future promise. ACM Transactions on Management Information Systems (TMIS), 12 (1):1–20, 2020.
- Mingxun Zhou, Tianhao Wang, TH Hubert Chan, Giulia Fanti, and Elaine Shi. Locally differentially private sparse vector aggregation. In 2022 IEEE Symposium on Security and Privacy (SP), pages 422–439. IEEE, 2022.

Appendices

The appendices are organised as follows: In Appendix A, we collect the results relevant to Section 2. For the user-level rates of Section 3, Section 4 and Section 5, we separate the proofs of the upper bounds and lower bounds into Appendix B and Appendix C respectively. Lastly, in Appendix D we collect auxiliary technical details.

A Proofs of results in Section 2

Proof of Example 1. In this proof, we derive the two results (6) and (7) separately.

Sample size T-dependent lower bound (6). We construct two distributions P_0 and P_1 where

$$P_0(X=0) = P_0(X=1) = \frac{1}{2}$$
 and $P_1(X=0) = 1 - P_0(X=1) = \frac{1}{2} + 2\delta$,

where $\delta = \min\{1/5, (192nT\alpha^2)^{-1/2}\}$. Noting that $|\mathbb{E}_{P_0}(X) - \mathbb{E}_{P_1}(X)| = 2\delta$, it follows from a private version of Le Cam's lemma (Duchi et al., 2018, Proposition 1) that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}),(\cdot)^2) \geq \frac{\delta^2}{2} \left\{ 1 - \sqrt{3n\alpha^2 D_{\mathrm{KL}}(P_0^{\otimes T} \| P_1^{\otimes T})} \right\} = \frac{\delta^2}{2} \left\{ 1 - \sqrt{3nT\alpha^2 D_{\mathrm{KL}}(P_0 \| P_1)} \right\}$$
$$\geq \frac{\delta^2}{2} \left(1 - \sqrt{48nT\alpha^2\delta^2} \right) \gtrsim \min\left\{ 1, \frac{1}{nT\alpha^2} \right\},$$

where the first inequality is due to the fact that $(e^{\alpha}-1)^2 \leq 3\alpha^2$ for $\alpha \in (0,1]$; the identity follows from the assumption that for each user the *T* samples are independent and the additivity of the Kullback–Leibler divergence for product distributions; the second inequality holds by noticing that for $\delta \in (0, 1/5)$,

$$D_{\mathrm{KL}}(P_0 \| P_1) = -\frac{1}{2} \log (1 + 4\delta) - \frac{1}{2} \log (1 - 4\delta) \le 16\delta^2;$$

and the final inequality holds by the choice of δ . This concludes the proof of the inequality (6).

Sample size T-independent lower bound (7). We consider two distributions P_0 and P_1 where

$$P_0(X=0) = P_1(X=1) = 1.$$

For any user-level α -LDP privacy mechanism Q, let

$$M_j(\cdot) = \int Q(\cdot|x) \,\mathrm{d}P_j^{\otimes T}(x), \quad j \in \{0,1\}.$$

It then follows that

$$\begin{aligned} \mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}),(\cdot)^2) \geq &\frac{1}{2} \left\{ 1 - D_{\mathrm{TV}}(M_0^n,M_1^n) \right\} \geq \frac{1}{4} e^{-D_{\mathrm{KL}}(M_0^n,M_1^n)} \\ \geq &\frac{1}{4} e^{-12n\alpha^2 D_{\mathrm{TV}}(P_0^{\otimes T},P_1^{\otimes T})} \geq \frac{1}{4} e^{-12n\alpha^2}, \end{aligned}$$

where the first inequality follows from Le Cam's lemma (e.g. Yu, 1997, Lemma 1); the second is from the Bretagnolle–Huber inequality (e.g. Tsybakov, 2009, Equation (2.25)); the third is from Corollary 3 in Duchi et al. (2018); and the last is due to the fact that the total variation distance is trivially upper bounded by 1. Hence, inequality (7) is proved which concludes the proof.

Proof of Theorem 1. We prove the lower and upper bounds separately, and the claim follows consequently.

The lower bound.

This proof is an application of Fano's lemma (e.g. Yu, 1997, Lemma 3). For $\delta > 0$, let $\{P_v : v \in \mathcal{V} \subset \Theta\} \subset \mathcal{P}$ satisfy the following: (i) For any $v, v' \in \mathcal{V}$ with $v \neq v'$, it holds that $\rho(\theta(P_v), \theta(P_{v'})) \geq 2\delta$. (ii) With $M(2\delta)$ being the 2δ -packing number in the metric space $(\Theta, \rho), |\mathcal{V}| = M(2\delta)$. Note that $M(2\delta) \leq N(\delta) < \infty$ (e.g. Vershynin, 2018, Lemma 4.2.8). It then holds that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \geq \inf_{\substack{Q \in \mathcal{Q}_{\alpha} \\ \hat{\theta} \ v \in \mathcal{V}}} \inf_{\substack{v \in \mathcal{V} \\ v \in \mathcal{V}}} \mathbb{E}_{P_{v},Q} \left[\Phi \circ \rho(\hat{\theta}, \theta(P_{v})) \right] \\
\geq \inf_{\substack{Q \in \mathcal{Q}_{\alpha} \\ Q \in \mathcal{Q}_{\alpha}}} \frac{\Phi(\delta)}{2} \left\{ 1 - \frac{K_{n} + \log(2)}{\log(|\mathcal{V}|)} \right\} \geq \frac{\Phi(\delta)}{2} \inf_{\substack{Q \in \mathcal{Q}_{\alpha} \\ Q \in \mathcal{Q}_{\alpha}}} \left\{ 1 - \frac{K_{n} + \log(2)}{\log(N(2\delta))} \right\},$$
(51)

where the second inequality follows from Fano's inequality (e.g. Yu, 1997), where

$$K_n = \max_{v,v' \in \mathcal{V}} D_{\mathrm{KL}}(M_v^n \| M_{v'}^n)$$

and the final inequality follows from the fact that $M(2\delta) \ge N(2\delta)$ (e.g. Vershynin, 2018, Lemma 4.2.8). We further note that

$$\sup_{Q \in \mathcal{Q}_{\alpha}} \max_{v,v' \in \mathcal{V}} D_{\mathrm{KL}}(M_{v}^{n} \| M_{v'}^{n}) \le 12n\alpha^{2} D_{\mathrm{TV}}^{2}(P_{v}, P_{v'}) \le 12n\alpha^{2},$$
(52)

where the second inequality follows from Corollary 3 in Duchi et al. (2018) and the fact that $(e^{\alpha} - 1)^2 \leq 3\alpha^2$ for $\alpha \in (0, 1]$; and the final inequality follows from the fact that the total variation distance is trivially upper bounded by 1. Combining (51) and (52) leads to

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \geq \frac{\Phi(\delta)}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{\log(N(2\delta))} \right\},\,$$

as claimed.

The upper bound.

Step 1: Preparation. Recall the non-overlapping covering set $\widetilde{\mathcal{B}}$ constructed in (10) with $|\widetilde{\mathcal{B}}| = N_{\widetilde{\mathcal{B}}} \leq N(\delta)$. For $P \in \mathcal{P}$, denote by $l \in [N_{\widetilde{\mathcal{B}}}]$ the fixed but unknown value such that $\theta(P) \in B'_l \in \widetilde{\mathcal{B}}$. For $\alpha > 0$, write $\pi_{\alpha} = e^{\alpha}/(1+e^{\alpha})$. It follows from the construction that $V_j^{(i)} = \mathbb{1}\{j = l\}$, for $i \in [n]$ and $j \in [N_{\widetilde{\mathcal{B}}}]$. By the construction (11), it holds, for $Z_j = \sum_{i=1}^n \widetilde{V}_j^i$ the total privatised votes for the *j*-th ball, that

$$Z_j \sim \operatorname{Bin}(n, \pi_{\alpha/2}) \mathbb{1}\{j = l\} + \operatorname{Bin}(n, 1 - \pi_{\alpha/2}) \mathbb{1}\{j \neq l\}, \quad j \in [N_{\widetilde{\mathcal{B}}}]$$

In addition, $\{Z_j\}_{j \in [N_{\widetilde{\mathcal{B}}}]}$ is a collection of mutually independent random variables. Define the event $A = \bigcap_{j \neq l} \{Z_l > Z_j\}$. The rest of the proof is concerned with controlling the risks on the event A and its complement A^c .

Step 2: Probability Control. It holds that

$$\mathbb{P}(A^c) \le \sum_{j \in [N_{\widetilde{\mathcal{B}}}] \setminus \{l\}} \mathbb{P}(Z_l \le Z_j) \le N(\delta) \mathbb{P}(Z_l \le n - Z') = N(\delta) \mathbb{P}(Z \le n),$$
(53)

where $Z' \sim \operatorname{Bin}(n, \pi_{\alpha/2})$ is independent of $\{Z_j\}_{j \in [N_{\tilde{g}}]}$, and $Z = Z_l + Z'$. By construction, Z_l and Z' are i.i.d. and $Z \sim \operatorname{Bin}(2n, \pi_{\alpha/2})$. Therefore Z has the same distribution as $\sum_{i=1}^{2n} W_i$, where $\{W_i\}_{i=1}^{2n}$ is a collection of independent $\operatorname{Ber}(\pi_{\alpha/2})$ random variables. It then follows from Hoeffding's inequality (e.g. Wainwright, 2019, Proposition 2.5) that

$$\mathbb{P}(Z \le n) = \mathbb{P}\left(\sum_{i=1}^{2n} W_i \le n\right) \le \exp\left(-4n(\pi_{\alpha/2} - 1/2)^2\right) \le e^{-n\alpha^2/20},\tag{54}$$

where the last inequality follows from the fact that $\pi_{\alpha/2} = e^{\alpha/2}/(1+e^{\alpha/2})$ and that $\alpha \in (0,1]$. Combining (53) and (54), it follows that

$$\mathbb{P}(A^c) \le N(\delta)e^{-n\alpha^2/20}.$$

Step 3: Conclusion. On the event A, due to the construction, we have that $\hat{\theta}, \theta(P) \in B_l$ and hence that $\rho(\hat{\theta}, \theta(P)) \leq \delta$. On the event A^c , we can only guarantee that $\hat{\theta}, \theta(P) \in \Theta$ and hence that $\rho(\hat{\theta}, \theta(P)) \leq \text{diam}(\Theta)$. From this and the bound on $\mathbb{P}(A^c)$ we obtain that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \le \Phi(\delta)\mathbb{P}(A) + \Phi(\operatorname{diam}(\Theta))\mathbb{P}(A^c) \le \Phi(\delta) + \Phi(\operatorname{diam}(\Theta))N(\delta)e^{-n\alpha^2/20},$$

nired.

as required.

Proof of Proposition 3. We prove the ℓ_{∞} - and ℓ_2 -cases separately. In each case, we first upper and lower bound the covering numbers of the spaces and then apply Theorem 1.

The ℓ_{∞} -ball case. For $\delta \in (0, 1)$, to upper bound the covering number $N(\delta)$ of $\mathbb{B}_{\infty}(1)$, we have that

$$N(\delta) \leq \frac{\operatorname{Vol}((2/\delta)\mathbb{B}_{\infty}(1) + \mathbb{B}_{2}(1))}{\operatorname{Vol}(\mathbb{B}_{2}(1))} \leq \frac{\operatorname{Vol}((2/\delta + 1)\mathbb{B}_{\infty}(1))}{\operatorname{Vol}(\mathbb{B}_{2}(1))} \leq \left(\frac{3}{\delta}\right)^{d} \frac{\operatorname{Vol}(\mathbb{B}_{\infty}(1))}{\operatorname{Vol}(\mathbb{B}_{2}(1))} = \left(\frac{3}{\delta}\right)^{d} \frac{2^{d}\Gamma(1 + d/2)}{\pi^{d/2}} \leq \left(\frac{18d}{\pi\delta^{2}}\right)^{d/2},$$
(55)

where the first inequality follows from Lemma 5.7(b) in Wainwright (2019), by setting \mathbb{B} , \mathbb{B}' , $\|\cdot\|$ and $\|\cdot\|'$ therein as $\mathbb{B}_{\infty}(1)$, $\mathbb{B}_{2}(1)$, $\|\cdot\|_{\infty}$ and $\|\cdot\|_{2}$; the second from the fact that $\mathbb{B}_{2}(1) \subseteq \mathbb{B}_{\infty}(1)$; the third from the fact that $\delta \in (0, 1)$; and the last is due to $\Gamma(1 + x) \leq x^{x}$ for all x > 0.

For the lower bound on $N(\delta)$, we have that, for any $\delta \in (0, 1)$,

$$N(\delta) \ge \left(\frac{1}{\delta}\right)^d \frac{\operatorname{Vol}(\mathbb{B}_{\infty}(1))}{\operatorname{Vol}(\mathbb{B}_2(1))} = \left(\frac{1}{\delta}\right)^d \frac{2^d \Gamma(1+d/2)}{\pi^{d/2}} \ge \left(\frac{2d}{3\pi\delta^2}\right)^{d/2},\tag{56}$$

where the first inequality is from Lemma 5.7(a) in Wainwright (2019) and the final inequality from $\Gamma(1+x) \ge (x/3)^x$ using Stirling's approximation. Combining (55) and (56), we have that

$$\left(\frac{d}{5\delta^2}\right)^{d/2} \le N(\delta) \le \left(\frac{6d}{\delta^2}\right)^{d/2}.$$
(57)

We now apply Theorem 1 to obtain upper and lower bounds on the minimax rate. To obtain the lower bound, we let $\delta = (d/20)^{1/2} e^{-24n\alpha^2/d - \log(4)/d} < 1$, where the inequality holds due to the assumption that $n\alpha^2 > 60d \log(6d)$. An application of the general lower bound in Theorem 1 gives

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \ge \frac{\delta^2}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{\log(N(2\delta))} \right\} \ge \frac{\delta^2}{2} \left\{ 1 - \frac{24n\alpha^2 + \log(4)}{d\log(d/(20\delta^2))} \right\}$$
$$= \frac{d}{40} e^{-48n\alpha^2/d - \log(16)/d} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{d\{24n\alpha^2/d + \log(4)/d\}} \right\}$$
$$= \frac{d}{80} e^{-48n\alpha^2/d - \log(16)/d} \gtrsim e^{-Cn\alpha^2/d}, \tag{58}$$

where C > 0 is an absolute constant; the first inequality is from the first inequality in (8); the second from the first inequality in (57); the first equality follows from the choice of δ ; and the last inequality holds as $n\alpha^2 \gtrsim 1$ which follows from the assumption $n\alpha^2 > 60d \log(6d)$. To obtain an upper bound, we set $\delta = (6d)^{1/2}e^{-n\alpha^2/(40d)} < 1$ where the inequality holds due to the

To obtain an upper bound, we set $\delta = (6d)^{1/2} e^{-n\alpha^2/(40d)} < 1$ where the inequality holds due to the assumption $n\alpha^2 > 60d \log(6d)$. An application of the general upper bound in Theorem 1 gives that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{d,r}), \|\cdot\|_2^2) \le \delta^2 + \{\operatorname{diam}(\Theta)\}^2 N(\delta) e^{-n\alpha^2/20} \le \delta^2 + 4d(6d/\delta^2)^{d/2} e^{-n\alpha^2/20} = 6de^{-n\alpha^2/(20d)} + 4de^{-n\alpha^2[1/20-1/40]} \le e^{-cn\alpha^2/d},$$
(59)

where c > 0 is an absolute constant; the first inequality follows from the upper bound in (8); the second from the second inequality in (57) and the fact that diam(Θ) = $2d^{1/2}$; the equality holds by the choice of δ ; and the final inequality comes from the assumption that $n\alpha^2 > 60d \log(6d)$ which allows the factor of d to be absorbed into the exponential term. By combining (58) and (59), we conclude the proof of (13).

The ℓ_2 -ball case. It follows from Example 5.8 of Wainwright (2019) that, for any $\delta \in (0, 1)$, the δ -covering number of $\mathbb{B}_2(1)$ satisfies

$$(1/\delta)^d \le N(\delta) \le (3/\delta)^d. \tag{60}$$

To obtain the lower bound on the minimax risk, we let $\delta = 2^{-1}e^{-24n\alpha^2/d - \log(4)/d} < 1$. An application of the general lower bound in Theorem 1 gives that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \ge \frac{\delta^2}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{\log(N(2\delta))} \right\} \ge \frac{\delta^2}{2} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{d\log(1/(2\delta))} \right\}$$
$$= \frac{1}{8} e^{-48n\alpha^2/d - \log(16)/d} \left\{ 1 - \frac{12n\alpha^2 + \log(2)}{d\{24n\alpha^2/d + \log(4)/d\}} \right\}$$
$$= \frac{1}{16} e^{-48n\alpha^2/d - \log(16)/d} \gtrsim e^{-C'n\alpha^2/d}, \tag{61}$$

where C' > 0 is an absolute constant; the first inequality is from (8); the second from the first inequality in (60); the first equality follows from the choice of δ ; and the last inequality holds as $n\alpha^2 \gtrsim 1$ which follows from the assumption $n\alpha^2 > 60d$.

To obtain the upper bound, we let $\delta = 3e^{-n\alpha^2/(30d)} < 1$, where the inequality holds due to the assumption that $n\alpha^2 > 60d$. An application of the general upper bound in Theorem 1 gives that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{d,1}), \|\cdot\|_2^2) \le \delta^2 + \{\operatorname{diam}(\Theta)\}^2 N(\delta) e^{-n\alpha^2/20} \le \delta^2 + 4(3/\delta)^d e^{-n\alpha^2/20}$$

$$=9e^{-n\alpha^2/(15d)} + 4e^{-n\alpha^2(1/20 - 1/30)} \lesssim e^{-c'n\alpha^2/d},$$
(62)

where c' > 0 is an absolute constant; the first inequality follows the upper bound in (8); the second from (60) and the fact that diam(Θ) = 2; and the equality holds by the choice of δ . By combining (61) and (62), we conclude the proof of (14).

Proof of Proposition 4. For $\delta > 0$, we first obtain the required bounds on the δ -covering number $N(\delta)$ of the metric space $(\Theta, \|\cdot\|_2)$, where $\Theta = \theta(\mathcal{P}_{d,s})$ and $\|\cdot\|_2$ is the vector ℓ_2 -norm.

For an upper bound on $N(\delta)$, we note that there are $\binom{d}{s}$ -many choices for the non-zero elements of the parameter vector, and for each choice, the covering number of the resulting space can be bounded by (57) with s in place of d therein. Hence, for any $\delta \in (0, 1)$, we have that

$$N(\delta) \le {\binom{d}{s}} \left(\frac{6s}{\delta^2}\right)^{\frac{s}{2}} \le \left(\frac{ed}{s}\right)^s \left(\frac{6s}{\delta^2}\right)^{\frac{s}{2}},\tag{63}$$

where the second inequality comes from the bound $\binom{d}{s} \leq (ed/s)^s$ for $s \in [d]$ due to Stirling's formula.

For a lower bound on $N(\delta)$, by Lemma 18 and for any $\delta \in (0, 1/4)$, it holds that

$$N(\delta) \ge \left(\frac{d}{s}\right)^s \left\{\frac{s(1/\delta - 2)^2}{20}\right\}^{\frac{s}{2}}.$$
(64)

To obtain the lower bound on the risk, we let

$$\delta = \left(\frac{2(20s)^{1/2}}{d}e^{24n\alpha^2/s + \log(4)/s} + 4\right)^{-1} < \frac{1}{8}$$

where the inequality holds for $d \ge 2$ due to the assumption $n\alpha^2 > 120s \log(6d)$. An application of the general lower bound in Theorem 1 gives that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_{2}^{2}) \geq \frac{\delta^{2}}{2} \left\{ 1 - \frac{12n\alpha^{2} + \log(2)}{\log(N(2\delta))} \right\} \geq \frac{\delta^{2}}{2} \left(1 - \frac{24n\alpha^{2} + \log(4)}{s\log[\{d^{2}/(20s)\}\{1/(2\delta) - 2\}^{2}]} \right)$$
$$= \left(\frac{2\sqrt{20s}}{d} e^{24n\alpha^{2}/s + 2\log(2)/s} + 4 \right)^{-2} \left(1 - \frac{1}{2} \right) \gtrsim \frac{d^{2}}{s} e^{-Cn\alpha^{2}/s}, \tag{65}$$

where C > 0 is an absolute constant; the first inequality is from (8); the second from the bound (64); the equality follows from the choice of δ ; and the last inequality holds as $(s^{1/2}/d)e^{24n\alpha^2/s+\log(4)/s} \gtrsim 1$ due to the assumption $n\alpha^2 \geq 120s \log(6d)$.

To obtain the upper bound, we let $\delta = 6^{1/2} e^{ds^{-1/2}} e^{-n\alpha^2/(40s)} < 1$, where the inequality holds due to the assumption $n\alpha^2 \ge 120s \log(6d)$. An application of the general upper bound in Theorem 1 shows that

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{P}_{d,s}), \|\cdot\|_{2}^{2}) \leq \delta^{2} + \{\operatorname{diam}(\Theta)\}^{2} N(\delta) e^{-n\alpha^{2}/20} \leq \delta^{2} + 4s \left(\frac{ed}{s}\right)^{s} \left(\frac{6s}{\delta^{2}}\right)^{\frac{s}{2}} e^{-n\alpha^{2}/20} \\ = 6e^{2} \frac{d^{2}}{s} e^{-n\alpha^{2}/(20s)} + 4s e^{-n\alpha^{2}[1/20-1/40]} \lesssim e^{-cn\alpha^{2}/s}, \tag{66}$$

where c > 0 is an absolute constant; the first inequality follows from the upper bound in (8); the second from (63) and the fact that diam(Θ) = $2s^{1/2}$; the equality holds by the choice of δ ; and the final inequality comes from the assumption that $n\alpha^2 > 120s \log(6d)$. By combining (65) and (66), we conclude the proof of (17).

Proof of Proposition 5. For $\delta > 0$, we first obtain the required bounds on the δ -covering number $N(\delta)$ of the metric space $(\Theta, \|\cdot\|_2)$, where $\Theta = \theta(\mathcal{P}_{\beta,1})$ and $\|\cdot\|_2$ is the $L^2([0,1])$ norm. We have that

$$\exp\left(c_{\beta}(1/\delta)^{1/\beta}\right) \le N(\delta) \le \exp\left(C_{\beta}(1/\delta)^{1/\beta}\right),\tag{67}$$

where the upper bound holds by noting $\mathcal{F}_{\beta,1} \subset \mathcal{S}_{\beta,1}$ and by an upper bound on the covering number of $\mathcal{S}_{\beta,1}$ (e.g. Wainwright, 2019, Example 5.12), and the lower bound follows from Lemma 17, for δ sufficiently small.

To obtain the lower bound in (23), we set

$$\delta = \frac{1}{2} \left(\frac{c_{\beta}}{c_{\beta}' \{24n\alpha^2 + 2\log(2)\}} \right)^{\beta}$$

where $c'_{\beta} > 0$ is a sufficiently large absolute constant, so that δ is sufficiently small therefore the lower bound in (67) holds. Applying the lower bound in Theorem 1 leads to

$$\begin{aligned} \mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{F}_{\beta,1}), \|\cdot\|_{2}^{2}) &\geq \frac{\delta^{2}}{2} \left\{ 1 - \frac{12n\alpha^{2} + \log(2)}{\log(N(2\delta))} \right\} \geq \frac{\delta^{2}}{2} \left\{ 1 - \frac{12n\alpha^{2} + \log(2)}{c_{\beta}\{1/(2\delta)\}^{1/\beta}} \right\} \\ &= \frac{1}{8} \left(\frac{c_{\beta}}{c_{\beta}'\{24n\alpha^{2} + 2\log(2)\}} \right)^{2\beta} \left(1 - \frac{1}{2c_{\beta}'} \right) \gtrsim \left(\frac{c_{\beta}}{c_{\beta}'n\alpha^{2}} \right)^{2\beta} \end{aligned}$$

where the first inequality is from the lower bound in (8), the second from the first inequality in (67), the first equality follows from the choice of δ , and the last holds under the assumption that $n\alpha^2 \gtrsim 1$. Setting $c''_{\beta} = (c_{\beta}/c'_{\beta})^{2\beta}$ shows that the lower bound can be written as

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{F}_{\beta,1}), \|\cdot\|_2^2) \gtrsim \frac{c_{\beta}''}{(n\alpha^2)^{2\beta}},\tag{68}$$

as required.

To obtain the upper bound in (23), we let $\delta = \{40C_{\beta}/(n\alpha^2)\}^{\beta}$. When $n\alpha^2 \ge C'_{\beta}$, where $C'_{\beta} > 0$ is a sufficiently large constant, we have δ sufficiently small so that the upper bound in (67) holds. Applying the upper bound in Theorem 1 leads to

$$\mathcal{R}_{n,\infty,\alpha}(\theta(\mathcal{F}_{\beta,1}), \|\cdot\|_{2}^{2}) \leq \delta^{2} + \{\operatorname{diam}(\Theta)\}^{2} N(\delta) e^{-n\alpha^{2}/20} \leq \delta^{2} + 2 \exp\left(C_{\beta}(1/\delta)^{1/\beta} - n\alpha^{2}/20\right)$$
$$= \{40C_{\beta}/(n\alpha^{2})\}^{2\beta} + 2e^{-n\alpha^{2}/40} \lesssim \{C_{\beta}''/(n\alpha^{2})\}^{2\beta}, \tag{69}$$

where $C_{\beta}^{\prime\prime} > 0$ is some constant depending on β , the first inequality follows from the upper bound in (8), the second from the second inequality in (67) and the fact that diam(Θ) $\leq 2^{1/2}$, the equality holds by the choice of δ and simplifying the exponential term; and the final inequality comes from noting that for any value of β the exponential term is dominated by the polynomial. By combining (68) and (69), we conclude the proof of (23).

B Proof of user-level upper bounds

B.1 Proof of Theorem 6 upper bound (univariate)

Proof of Theorem 6 (d = 1).

Step 1: initial estimator interval $[\tilde{L}, \tilde{U}]$. Recall that n is assumed to be even. We denote by $l \in [N(\delta)]$ the fixed but unknown index such that the truth $\theta \in I_l$. Further, for each $i \in [n]$ and $t \in [T]$, the random variable $X_t^{(i)} \in [-1, 1]$. Hence, recalling the definition of $\hat{\theta}^{(i)}$ for $i \in [n/2]$ in (26), we see that

$$\mathbb{P}\left(\left|\hat{\theta}^{(i)} - \theta\right| \ge \delta\right) = \mathbb{P}\left\{\left|\frac{1}{T^*}\sum_{t=1}^T X_t^{(i)}\right| \ge \delta\right\} \le 2e^{-T^*\delta^2/2} = \frac{2}{nT^*\alpha^2} \le \frac{1}{4}$$
(70)

where the first inequality is due to Hoeffding's inequality for bounded random variables (e.g. Wainwright, 2019, Proposition 2.5); the second equality by the definition of δ from (25); and the last inequality by the assumption that $n\alpha^2 \geq \tilde{C}$ for some sufficiently large \tilde{C} and $T^* \geq 1$. Consequently, we can bound the following probabilities

$$\mathbb{P}(V_l^{(i)} = 0) \le \mathbb{P}\left(|\hat{\theta}^{(i)} - \theta| \ge \delta\right) \le \frac{1}{4}$$

and

$$\mathbb{P}(V_j^{(i)} = 1) \le \mathbb{P}\left(|\hat{\theta}^{(i)} - \theta| \ge \delta\right) \le \frac{1}{4}, \quad j \in \{k \in [N(\delta)] : |l - k| > 2\},\$$

following from (70). For any x > 0, write $\pi_x = e^x/(1+e^x)$. Then, for $k \in [N(\delta)]$, we have that

$$p_k := \mathbb{P}(\widetilde{V}_k^{(i)} = 1) = (2\pi_{\alpha/6} - 1)\mathbb{P}(V_k^{(i)} = 1) + (1 - \pi_{\alpha/6})$$

Hence, $p_l \ge (2\pi_{\alpha/6} + 1)/4$ and $p_j \le (3 - 2\pi_{\alpha/6})/4$ for $j \in \{k \in [N(\delta)] : |l - k| > 2\}$.

Letting $Z_j = \sum_{i=1}^{n/2} \widetilde{V}_j^{(i)}$ denote the total privatised votes for the *j*-th sub-interval, we consider the event

$$A = \bigcap_{j \in [N(\delta)]: |l-j| > 2} \{ Z_l > Z_j \}.$$
 (71)

We note that

$$Z_l - Z_j = \sum_{i=1}^{n/2} (\widetilde{V}_l^{(i)} - \widetilde{V}_j^{(i)}),$$

where for each $i \in [n/2]$, it holds that $\widetilde{V}_l^{(i)} - \widetilde{V}_j^{(i)} \in \{-1, 0, 1\}$ and that $\mathbb{E}[\widetilde{V}_l^{(i)} - \widetilde{V}_j^{(i)}] = p_l - p_j$. Then

$$\mathbb{P}(Z_l \le Z_j) = \mathbb{P}\left(\sum_{i=1}^{n/2} \left(\widetilde{V}_l^{(i)} - \widetilde{V}_j^{(i)}\right) \le 0\right) = \mathbb{P}\left(\sum_{i=1}^{n/2} \left(\widetilde{V}_l^{(i)} - \widetilde{V}_j^{(i)} - p_l + p_j\right) \le -n/2(p_l - p_j)\right)$$
$$\le e^{-n(p_l - p_j)^2/4} \le e^{-n(1/2 - \pi_{\alpha/6})^2/4} \le e^{-2n\alpha^2/K}.$$
(72)

for K > 0 an absolute constant, where the first inequality comes from Hoeffding's inequality and the third by the fact that $\alpha^2/K \leq (1/2 - \pi_{\alpha/6})^2$ holds for $\alpha \in (0, 1]$ for K sufficiently large.

Hence, we have

$$\mathbb{P}(A^c) \le \sum_{j:|l-j|>2} \mathbb{P}(Z_l \le Z_j) \le N(\delta) e^{-2n\alpha^2/K} \le \frac{3}{\delta \wedge 1} e^{-2n\alpha^2/K},\tag{73}$$

where the first inequality is by the union bound; the second by (72); and the last by bounding the covering number using (60). We note that for the covering number bound we require $\delta < 1$, so to account for when this fails to hold we take the minimum $\delta \wedge 1$ which suffices for an upper bound.

On the event A, we have that j^* , as defined in (28), satisfies $j^* \in \{l-2, l-1, l, l+1, l+2\}$ almost surely. As the distance between the parameter $\theta \in I_l$ and the closest endpoint of either I_{l-2} or I_{l+2} is at most 4δ , expanding the endpoints of the chosen interval I_{j^*} by 6δ as in (29) ensures that $\min\{\theta - \tilde{L}, \tilde{U} - \theta\} \ge 2\delta$, equivalently that $\theta \in [\tilde{L} + 2\delta, \tilde{U} - 2\delta]$, regardless of the value of j^* .

Step 2: Final estimator. It remains to consider the refined estimator using the remaining users $i \in [n] \setminus [n/2]$. Recalling the private estimator $\hat{\theta}^{(i)}$ as defined in (30) and writing $\overline{X^{(i)}} = (T^*)^{-1} \sum_{t=1}^{T^*} X_t^{(i)}$ for $i \in [n]$, we have for any $i \in [n] \setminus [n/2]$ that

$$\mathbb{E}\left[\left|\hat{\theta}-\theta\right|^{2}\right] = \mathbb{E}\left[\mathbb{E}\left\{\left|\hat{\theta}-\theta\right|^{2}\mid j^{*}\right\}\right] = \mathbb{E}\left[\left\{\mathbb{E}(\hat{\theta}|j^{*})-\theta\right\}^{2} + \operatorname{Var}(\hat{\theta}|j^{*})\right] \\
= \mathbb{E}\left[\left\{\mathbb{E}(\hat{\theta}^{(i)}-\overline{X^{(i)}}|j^{*})\right\}^{2} + \frac{2}{n}\operatorname{Var}(\hat{\theta}^{(i)}|j^{*})\right] \\
= \mathbb{E}\left[\left\{\mathbb{E}\left(\left(\hat{\theta}^{(i)}-\overline{X^{(i)}}\right)\mathbb{1}\{A\}\mid j^{*}\right) + \mathbb{E}\left(\left(\hat{\theta}^{(i)}-\overline{X^{(i)}}\right)\mathbb{1}\{A^{c}\}\mid j^{*}\right)\right\}^{2} + \frac{2}{n}\operatorname{Var}(\hat{\theta}^{(i)}|j^{*})\right] \\
\leq 2\mathbb{E}\left[\left\{\mathbb{E}\left(\left(\hat{\theta}^{(i)}-\overline{X^{(i)}}\right)\mathbb{1}\{A\}\mid j^{*}\right)\right\}^{2} + \left\{\mathbb{E}\left(\left(\hat{\theta}^{(i)}-\overline{X^{(i)}}\right)\mathbb{1}\{A^{c}\}\mid j^{*}\right)\right\}^{2} + \frac{1}{n}\operatorname{Var}(\hat{\theta}^{(i)}|j^{*})\right] \\
= 2\mathbb{E}\{(I) + (II) + (III)\},$$
(74)

where the first equality is by the tower law of expectation; the second is by the bias-variance decomposition; the third by the definition of $\hat{\theta}$, the fact that $\overline{X^{(i)}}$ is an unbiased estimator of θ and the fact that $\overline{X^{(i)}}$ is independent of j^* ; and the inequality follows from the Cauchy–Schwarz inequality.

Term (*III*). The variance term can be upper bounded as

$$\operatorname{Var}(\hat{\theta}^{(i)}|j^*) = \operatorname{Var}\left(\Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}}) \mid j^*\right) + \operatorname{Var}\left(\frac{14\delta}{\alpha}\ell_i\right) \le 49\delta^2 + \frac{392\delta^2}{\alpha^2} \le \frac{441\delta^2}{\alpha^2},\tag{75}$$

where the variance of the truncated sample mean is bounded as it is itself a bounded random variable taking values on an interval of width 14δ and the last inequality uses the fact that $\alpha \in (0, 1]$.

Term (I). We have, following a similar argument as in Duchi et al. (2018, Appendix B.1), that

$$\mathbb{E}\left[\left|\overline{X^{(i)}} - \Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}})\right| \mathbbm{1}\{A\} \mid j^*\right]$$

$$= \mathbb{E}\left[\left|\overline{X^{(i)}} - \tilde{U}\right| \mathbb{1}\left\{\overline{X^{(i)}} > \tilde{U}\right\} \mathbb{1}\left\{A\right\} \left|j^*\right] + \mathbb{E}\left[\left|\overline{X^{(i)}} - \tilde{L}\right| \mathbb{1}\left\{\overline{X^{(i)}} < \tilde{L}\right\} \mathbb{1}\left\{A\right\} \left|j^*\right]\right]$$

$$\leq 2\mathbb{P}\left(\left\{\overline{X^{(i)}} > \tilde{U}\right\} \cap A \left|j^*\right) + 2\mathbb{P}\left(\left\{\overline{X^{(i)}} < \tilde{L}\right\} \cap A \left|j^*\right),$$
(76)

where the inequality uses the result that $\overline{X^{(i)}} - \tilde{U} \leq 2$ which follows from the facts $\overline{X^{(i)}} \in [-1, 1]$ and $\tilde{U} \ge -1 + 2\delta + 6\delta = 8\delta - 1 \ge -1$ and similarly the result that $\overline{X^{(i)}} - \tilde{L} \ge -2$ which uses the fact that $\tilde{L} < 1.$

The first term in (76) can be bounded as

$$\mathbb{P}\left(\left\{\overline{X^{(i)}} > \tilde{U}\right\} \cap A \mid j^*\right) = \mathbb{P}\left(\left\{\overline{X^{(i)}} - \theta > \tilde{U} - \theta\right\} \cap A \mid j^*\right) \\
\leq \mathbb{P}\left(\left\{\overline{X^{(i)}} - \theta > 2\delta\right\} \cap A\right) \\
\leq \mathbb{P}\left(\overline{X^{(i)}} - \theta > 2\delta\right) \le e^{-T^*\delta^2/2},$$
(77)

where the first inequality is due to the facts that $\overline{X^{(i)}}$ is independent of j^* and that, on the event A, $\tilde{U} - \theta > 2\delta$; and the last inequality follows from Hoeffding's inequality. A similar argument gives the same inequality for the other term where we note that $\theta - \tilde{L} > 2\delta$ on A. From this, we have that

$$\left\{ \mathbb{E}\left[\left(\hat{\theta}^{(i)} - \overline{X^{(i)}}\right) \mathbb{1}\{A\} \mid j^* \right] \right\}^2 = \left\{ \mathbb{E}\left[\left(\Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}}) - \overline{X^{(i)}}\right) \mathbb{1}\{A\} \mid j^* \right] \right\}^2 \le 16e^{-T^*\delta^2}, \tag{78}$$

where in the equality we use the fact that the Laplacian noise variables ℓ_i are independent of all other randomness in the estimator with mean zero, and the inequality comes from combining (76) and (77).

Term (II). We have that

$$\mathbb{E}\left[\left\{\mathbb{E}\left(\left(\hat{\theta}^{(i)} - \overline{X^{(i)}}\right)\mathbb{1}\left\{A^{c}\right\} \mid j^{*}\right)\right\}^{2}\right] = \mathbb{E}\left[\left\{\mathbb{E}\left(\left\{\Pi_{\tilde{I}_{j^{*}}}(\overline{X^{(i)}}) - \overline{X^{(i)}}\right\}\mathbb{1}\left\{A^{c}\right\} \mid j^{*}\right)\right\}^{2}\right] \\ \leq 4\mathbb{E}\left[\mathbb{P}(A^{c}|j^{*})^{2}\right] \leq 4\mathbb{P}(A^{c}) \leq \frac{12}{\delta \wedge 1}e^{-2n\alpha^{2}/K}$$
(79)

where the equality follows from the fact that the Laplacian noise variables ℓ_i are independent of all other randomness in the estimator and are mean zero; the first inequality comes from the fact that both $X^{(i)}$ and $\Pi_{\tilde{I}_{*}}(\overline{X^{(i)}})$ take values in [-1,1]; and the last by (73).

We combine (74), (75), (78) and (79) to see that

$$\begin{split} \mathbb{E}\left[|\hat{\theta} - \theta|^{2}\right] &\leq 32e^{-T^{*}\delta^{2}} + \frac{24}{\delta \wedge 1}e^{-2n\alpha^{2}/K} + \frac{882\delta^{2}}{n\alpha^{2}} \\ &\lesssim \frac{1}{nT^{*}\alpha^{2}} + \left(\frac{(T^{*})^{1/2}}{\{\log(nT^{*}\alpha^{2})\}^{1/2}} \vee 1\right)e^{-2n\alpha^{2}/K} + \frac{\log(nT^{*}\alpha^{2})}{nT^{*}\alpha^{2}} \\ &\lesssim \frac{\log(nT^{*}\alpha^{2})}{nT^{*}\alpha^{2}} + \frac{(T^{*})^{1/2}}{\{\log(nT^{*}\alpha^{2})\}^{1/2}}e^{-2n\alpha^{2}/K} + e^{-2n\alpha^{2}/K}, \\ &\lesssim \frac{\log(nT^{*}\alpha^{2})}{nT^{*}\alpha^{2}}, \end{split}$$

where the second inequality comes from substituting in the value of δ as in (25); the third by bounding the maximum term by the sum of the two terms; and the fourth using the fact that $e^{-2n\alpha^2/K} \leq (T^*)^{-3/2}e^{-(n/2)\alpha^2/K} \lesssim 1/\{(T^*)^{3/2}n\alpha^2\}$ and controlling the log term in the denominator using the assumption that $n\alpha^2 > \tilde{C}$ for some sufficiently large \tilde{C} . In the case $T \le e^{n\alpha^2/K}$, we have $T^* = T$ which immediately gives the upper bound in (24). In the

case $T > e^{n\alpha^2/K}$, we have $T^* = e^{n\alpha^2/K}$, which gives

$$\frac{\log(nT^*\alpha^2)}{nT^*\alpha^2} = \left(\frac{\log(n\alpha^2)}{n\alpha^2} + \frac{\log(T^*)}{n\alpha^2}\right)\frac{1}{T^*} \lesssim e^{-n\alpha^2/K},$$

where the inequality comes from the assumption $n\alpha^2 \geq \widetilde{C}$, and from substituting in the value $e^{n\alpha^2/K}$ for T^* . Putting together the pieces, we have

$$\mathbb{E}\left[|\hat{\theta} - \theta|^2\right] \lesssim \begin{cases} \frac{\log(nT\alpha^2)}{nT\alpha^2} & \text{when } T \le e^{n\alpha^2/K}, \\ e^{-n\alpha^2/K} & \text{otherwise.} \end{cases}$$
(80)

Lastly, bounding the error by the sum of the errors in the two cases for T completes the proof.

B.2 Proof of Theorem 6 upper bound (multivariate)

Proof of Theorem 6 (ℓ_{∞} -ball case). The estimation procedure in Section 3.2 is equivalent to splitting the sample and applying the univariate procedure, defined in Section 3.1, to each co-ordinate in turn. For $j \in [d]$, the collection of users indexed by the set N_j in (31), uses the *j*-th co-ordinate of their data to estimate the *j*-th co-ordinate of the mean. We note that the assumption that $n\alpha^2 > \tilde{C}d\log(ed)$ ensures that the required assumption for the univariate sub-problems is satisfied.

The estimator for the *j*-th co-ordinate $\hat{\theta}_j$ has error bounded as

$$\mathbb{E}\left[|\hat{\theta}_j - \theta_j|^2\right] \lesssim \begin{cases} \frac{d \log(nT\alpha^2/d)}{nT\alpha^2}, & T \le e^{n\alpha^2/(Kd)}, \\ e^{-n\alpha^2/(Kd)}, & \text{otherwise}, \end{cases}$$
(81)

which follows from the univariate error in (80), noting the factors of d arising from the number of users in each collection being such that $|N_j| = n/d$ for $j \in [d]$. For the final estimator $\hat{\theta}$ as defined in (32), the error can be bounded as

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_2^2\right] = \sum_{j=1}^d \mathbb{E}\left[|\hat{\theta}_j - \theta_j|_2^2\right] \lesssim \begin{cases} \frac{d^2 \log(nT\alpha^2/d)}{nT\alpha^2}, & T \le e^{n\alpha^2/(Kd)}, \\ de^{-n\alpha^2/(Kd)}, & \text{otherwise}, \end{cases}$$
(82)

where the equality comes from the decomposition of multivariate mean squared error and the inequality by summing up d-many copies of (81). Lastly, we have that

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_2^2\right] \lesssim \frac{d^2 \log(nT\alpha^2/d)}{nT\alpha^2} + e^{-cn\alpha^2/(Kd)},\tag{83}$$

which comes from bounding the error by the sum of the errors in the two cases for T and using the assumption that $n\alpha^2 > \tilde{C}d\log(ed)$ for some sufficiently large \tilde{C} to absorb the factor of d into the exponential term. This completes the proof.

Proof of Theorem 6 (ℓ_2 -ball case). We recall that we assume without loss of generality that the dimension d is a power of 2, noting that we can always append zero values to the data until this holds, with this procedure not affecting the final minimax rate except up to constants. We then note that we can always take the origin of the ℓ_2 -ball as a trivial estimator. We upper bound the error of this trivial estimator by the radius of the ball. As a consequence, the upper bound on the risk consists of the minimum between the constant 1, and the error of the estimator we proposed in Section 3.3. In the rest of the proof, we focus on our proposed estimator.

The proof for the ℓ_2 -ball case also involves considering each co-ordinate separately and applying a univariate estimation procedure to each co-ordinate. However, due to the application of the random rotation of Lemma 7, the steps are not identical to the ℓ_{∞} case.

Step 1: random rotations. We are to show a user's estimator deviates from the truth by δ with small probability.

As $X_t^{(i)} \in \mathbb{B}_2(1)$ for all $i \in [n]$ and $t \in [T]$, we have that $||X_t^{(i)} - \theta||_2 \in [0, 2]$. Hence, we have by Jin et al. (2019, Lemma 1) that for some absolute constant $c_1 > 0$ and for all $\varepsilon > 0$,

$$\mathbb{P}(\|X_t^{(i)} - \theta\|_2 \ge \varepsilon) \le 2e^{-\varepsilon^2/(2c_1)}$$

Recalling the definition of $\hat{\theta}^{(i)}$ in (35), for each $i \in [n/2]$ we have by Lemma 15 that there exists a universal constant $c_2 > 0$ such that, for any $\gamma > 0$,

$$\|\hat{\theta}^{(i)} - \theta\|_2 \le c_2 \left\{ \frac{\log(2d/\gamma)}{T^*} \right\}^{1/2}$$
(84)

holds with probability at least $1 - \gamma$. We define the event

$$A_{i} = \left\{ \|\hat{\theta}^{(i)} - \theta\|_{2} \le c_{2} \left\{ \frac{\log(nT^{*}\alpha^{2})}{T^{*}} \right\}^{1/2} \right\}$$

from which we have that

$$\mathbb{P}(A_i^c) \le \frac{2d}{nT^*\alpha^2} \tag{85}$$

due to (84) with $\gamma = 2d/(nT^*\alpha^2)$ therein.

We thus have

$$\begin{split} \mathbb{P}\left(|(R\hat{\theta}^{(i)})_j - (R\theta)_j| \ge \delta\right) &\le \mathbb{P}\left(\left\{|(R\hat{\theta}^{(i)})_j - (R\theta)_j| \ge \delta\right\} \cap A_i\right) + \mathbb{P}(A_i^c) \\ &\le \mathbb{P}\left(\left\{|(R\hat{\theta}^{(i)})_j - (R\theta)_j| \ge \left(\frac{C}{dT^*}\right)^{1/2} \log(nT^*\alpha^2)\right\} \cap A_i\right) + \frac{2d}{nT^*\alpha^2} \\ &\le \mathbb{P}\left(\left\{|(R\hat{\theta}^{(i)})_j - (R\theta)_j| \ge \|\hat{\theta}^{(i)} - \theta\|_2 \left(\frac{C\log(nT^*\alpha^2)}{c_2^2d}\right)^{1/2}\right\} \cap A_i\right) + \frac{2d}{nT^*\alpha^2} \\ &\le \frac{1}{nT^*\alpha^2} + \frac{2d}{nT^*\alpha^2} \le \frac{1}{4} \end{split}$$

where the second inequality uses (85) and substitutes in the value of δ ; the third uses the definition of the event A_i ; the penultimate inequality follows from applying Lemma 7 with $\gamma = 1/(nT^*\alpha^2)$ therein, provided $C \geq 100c_2^2$; and the final using the assumption that $n\alpha^2 \geq \tilde{C}d\log(ed)$ for \tilde{C} sufficiently large and that $T \geq 1$.

Step 2: co-ordinate-wise treatment. We now focus on a single fixed co-ordinate $j \in [d]$ and follow the same steps as in the proof of the univariate case with some modifications to account for the random rotation. We first recall that for $j \in [d]$ and $k \in [N(\delta)]$, we let $Z_{j,k} = \sum_{i \in N_{j,1}} \widetilde{V}_k^{(i)}$ denote the total number of privatised votes from the *j*-th group of users for the *k*-th sub-interval and denote by $l_j \in [N(\delta)]$ the fixed but unknown index such that $(R_d \theta)_j \in I_{l_j}$. We consider the event

$$B_j = \bigcap_{k:|l_j - k| > 2} \{ Z_{j,l_j} > Z_{j,k} \},\tag{86}$$

which is analogous to the event (71) from the univariate case. On the event B_j , whichever interval is selected is sufficiently close to $R_d\theta_j$ so that $R_d\theta_j \in \tilde{I}_{k_j^*}$ and further, due to inflating the end points of the intervals, $\min\{(R_d\theta)_j - \tilde{L}_j, \tilde{U}_j - (R_d\theta)_j\} > 2\delta$. Bounding the probability of the complement by the same analysis as that in (72) and (73), we have

$$\mathbb{P}(B_j^c) \le \frac{3}{\delta \wedge 1} e^{-2n\alpha^2/(Kd)} \tag{87}$$

where K is as appears in (72) and, as in (73), we require $\delta < 1$, so to account for when this fails to hold we take the minimum $\delta \wedge 1$ which suffices for an upper bound. Writing $\overline{X^{(i)}} = (T^*)^{-1} \sum_{t=1}^{T^*} X_t^{(i)}$ for $i \in [n]$, we define the final events

$$D_{i,j} = \left\{ |(R_d \overline{X^{(i)}})_j - (R_d \theta)_j| \le \frac{10 ||\overline{X^{(i)}} - \theta||_2 \{\log(nT^* \alpha^2)\}^{1/2}}{d^{1/2}} \right\}.$$

By applying Lemma 7 with $\gamma = 1/(nT^*\alpha^2)$ therein we see that we have

$$\mathbb{P}(D_{i,j}^c) < 1/(nT^*\alpha^2). \tag{88}$$

Recalling the final estimator (37), we have that

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_{2}^{2}\right] = \mathbb{E}\left[\|R_{d}\hat{\theta} - R_{d}\theta\|_{2}^{2}\right] = \sum_{j=1}^{d} \mathbb{E}\left[|\tilde{\theta}_{j} - (R_{d}\theta)_{j}|^{2}\right]$$
(89)

where the first equality holds as the rotation R_d is an orthogonal transformation. Then, focusing on a single co-ordinate, we have for all $j \in [d]$ and any $i \in N_{j,2}$ that

$$\begin{split} \mathbb{E}\left[\left|\tilde{\theta}_{j}-(R_{d}\theta)_{j}\right|^{2}\right] \\ &\leq 2\mathbb{E}\left[\mathbb{E}\left\{\left\{\tilde{\theta}^{(i)}-(R_{d}\overline{X^{(i)}})_{j}\right\}\mathbb{1}\left\{B_{j}\right\} \mid k_{j}^{*}, R_{d}\right\}^{2}+\mathbb{E}\left\{\left\{\tilde{\theta}^{(i)}-(R_{d}\overline{X^{(i)}})_{j}\right\}\mathbb{1}\left\{(B_{j})^{c}\right\} \mid k_{j}^{*}, R_{d}\right\}^{2} \\ &\quad +\frac{d}{n}\mathrm{Var}(\tilde{\theta}^{(i)}|k_{j}^{*}, R_{d})\right] \end{split}$$

$$= 2\mathbb{E}\{(I) + (II) + (III)\},\tag{90}$$

where the inequality is by a similar decomposition to that in (74).

Step 2.1: term (III). The variance term can easily be bounded as

$$\operatorname{Var}(\tilde{\theta}^{(i)}|k_j^*, R_d) = \operatorname{Var}\left(\Pi_{\tilde{I}_{k_j^*}}\{(R_d \overline{X^{(i)}})_j\} \mid k_j^*, R_d\right) + \operatorname{Var}\left(\frac{14\delta}{\alpha}\ell_i\right) \le 49\delta^2 + \frac{392\delta^2}{\alpha^2} \le \frac{441\delta^2}{\alpha^2}, \quad (91)$$

where the variance of the truncated sample mean is bounded as it is itself a bounded random variable taking values on an interval of width 14δ and the last inequality uses the fact that $\alpha \in (0, 1]$.

Step 2.2: term (I). Following the same arguments as in (76) and (77), we have that

$$\mathbb{E}\left[\left| (R_{d}\overline{X^{(i)}})_{j} - \Pi_{\tilde{I}_{k_{j}^{*}}}\{(R_{d}\overline{X^{(i)}})_{j}\} \middle| \mathbb{1}\{B_{j}\} \middle| k_{j}^{*}, R_{d} \right] \\
\leq 2\mathbb{P}\left(\left\{ (R_{d}\overline{X^{(i)}})_{j} > \tilde{U}_{j} \right\} \cap B_{j} \middle| k_{j}^{*}, R_{d} \right) + 2\mathbb{P}\left(\left\{ (R_{d}\overline{X^{(i)}})_{j} < \tilde{L}_{j} \right\} \cap B_{j} \middle| k_{j}^{*}, R_{d} \right) \\
\leq 2\mathbb{P}\left(\left\{ \left| (R_{d}\overline{X^{(i)}})_{j} - (R_{d}\theta)_{j} \middle| > 2\delta \right\} \cap B_{j} \middle| k_{j}^{*}, R_{d} \right),$$
(92)

where the second inequality is due to the fact that on the event B_j , $\tilde{U}_j - (R_d\theta)_j > 2\delta$ and $(R_d\theta)_j - \tilde{L}_j > 2\delta$. We upper bound the contribution of (92) in the event $D_{i,j}$, that

$$\mathbb{E}\left[\mathbb{P}\left(\left\{\left|(R_{d}\overline{X^{(i)}})_{j}-(R_{d}\theta)_{j}\right|>2\delta\right\}\cap B_{j}\mid k_{j}^{*},R_{d}\right)^{2}\right]\leq\mathbb{P}\left(\left\{\left|(R_{d}\overline{X^{(i)}})_{j}-(R_{d}\theta)_{j}\right|>2\delta\right\}\cap B_{j}\right)\\
\leq\mathbb{P}\left(\left\{\left|(R_{d}\overline{X^{(i)}})_{j}-(R_{d}\theta)_{j}\right|>2\delta\right\}\cap B_{j}\cap D_{i,j}\right)+\mathbb{P}(D_{i,j}^{c})\\
\leq\mathbb{P}\left(\left\{\left\|\overline{X^{(i)}}-\theta\right\|_{2}\geq\frac{d^{1/2}\delta}{5\{\log(nT^{*}\alpha^{2})\}^{1/2}}\right\}\cap B_{j}\cap D_{i,j}\right)+\mathbb{P}(D_{i,j}^{c})\\
\leq2d\exp\left(-\frac{T^{*}d\delta^{2}}{25c_{3}\log(nT^{*}\alpha^{2})}\right)+\frac{1}{nT^{*}\alpha^{2}},$$
(93)

where the third inequality holds on the event $D_{i,j}$ and the fourth inequality follows from applying Lemma 15, with some absolute constant $c_3 > 0$, for the first term and by (88) for the second term.

We simplify the exponential term by noting that

$$\exp\left(-\frac{T^*d\delta^2}{25c_3\log(nT^*\alpha^2)}\right) = \exp\left(-\frac{C\{\log(nT^*\alpha^2)\}^2}{25c_3\log(nT^*\alpha^2)}\right) \le \left(\frac{1}{nT^*\alpha^2}\right)^2,\tag{94}$$

where in the last line we take $C > 50c_3$.

Step 2.3: term (II). We have by the same argument as (79) that

$$\mathbb{E}\left(\left[\mathbb{E}\left\{\left\{\tilde{\theta}^{(i)} - (R_d \overline{X^{(i)}})_j\right\}\mathbb{1}\left\{B_j^c\right\} \mid k_j^*, R_d\right\}\right]^2\right) \le 4\mathbb{P}(B_j^c).$$
(95)

We hence have that

$$\mathbb{E}[|(\tilde{\theta}_j - (R_d\theta)_j)|^2] \le 2d \exp\left(-\frac{T^* d\delta^2}{25c_3 \log(nT^*\alpha^2)}\right) + \frac{1}{nT^*\alpha^2} + 8\mathbb{P}(B_j^c) + \frac{882d\delta^2}{n\alpha^2}$$
$$\lesssim \frac{d}{(nT^*\alpha^2)^2} + \frac{1}{nT^*\alpha^2} + \left(\frac{3}{\delta} \lor 1\right) e^{-2n\alpha^2/(Kd)} + \frac{d\delta^2}{n\alpha^2}$$
$$\lesssim \frac{1}{nT^*\alpha^2} + \left(\frac{3}{\delta} \lor 1\right) e^{-2n\alpha^2/(Kd)} + \frac{d\delta^2}{n\alpha^2},\tag{96}$$

where the first inequality is due to (90), (91), (92), (93) and (95); the second is from (94), (95) and (87); and the final using the assumption that $n\alpha^2 > \tilde{C}' d\log(ed)$ for some sufficiently large \tilde{C}' .

Step 3: completing the proof. Summing up the error for each co-ordinate gives

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_2^2\right] = \sum_{j=1}^d \mathbb{E}\left[|\tilde{\theta}_j - (R_d\theta)_j|^2\right]$$

$$\begin{split} &\lesssim \frac{d}{nT^*\alpha^2} + d\left(\frac{1}{\delta} + 1\right) e^{-2n\alpha^2/(Kd)} + \frac{d^2\delta^2}{n\alpha^2} \\ &\lesssim \frac{d^{3/2}(T^*)^{1/2}}{\log(nT^*\alpha^2)} e^{-2n\alpha^2/(Kd)} + de^{-2n\alpha^2/(Kd)} + \frac{d\{\log(nT^*\alpha^2)\}^2}{nT^*\alpha^2} \\ &\lesssim \frac{d^{3/2}}{T^*} e^{-n\alpha^2/(2Kd)} + \frac{d\{\log(nT^*\alpha^2)\}^2}{nT^*\alpha^2} \\ &\lesssim \frac{1}{T^*} e^{-C'n\alpha^2/d} + \frac{d\{\log(nT^*\alpha^2)\}^2}{nT^*\alpha^2} \\ &\lesssim \frac{d\{\log(nT^*\alpha^2)\}^2}{nT^*\alpha^2}, \end{split}$$

where C' > 0 is some absolute constant; the equality is by (89); the first inequality comes from summing up *d*-many copies of (96) and bounding the maximum term by the sum; the second inequality comes from substituting in the value of δ as in (34); the third using the fact that $e^{-2n\alpha^2/(Kd)} \leq (T^*)^{-3/2}e^{-n\alpha^2/(2Kd)}$ and controlling the log term in the denominator using the assumption that $n\alpha^2 > \tilde{C}' d\log(ed)$ for some sufficiently large \tilde{C}' ; and the penultimate by using the fact that $n\alpha^2 > \tilde{C}' d\log(ed)$ for some sufficiently large \tilde{C}' to absorb the factor of $d^{3/2}$ into the exponential term.

large \tilde{C}' to absorb the factor of $d^{3/2}$ into the exponential term. Thus, when $T \leq e^{n\alpha^2/(Kd)}$, we have that $T^* = T$ immediately giving the result for this case. When $T > e^{n\alpha^2/(Kd)}$, we have that $T^* = e^{n\alpha^2/(Kd)}$ giving

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_{2}^{2}\right] \leq \frac{d\{\log(nT^{*}\alpha^{2})\}^{2}}{n\alpha^{2}}e^{-n\alpha^{2}/(Kd)}$$
$$\lesssim \frac{d}{n\alpha^{2}}\left(\log(n\alpha^{2}) + \frac{n\alpha^{2}}{d}\right)^{2}e^{-n\alpha^{2}/(Kd)} \lesssim e^{-C''n\alpha^{2}/d}$$

where C'' > 0 is an absolute constant. Hence, we have that the error of the estimator $\hat{\theta}$ is bounded as

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|_2^2\right] \lesssim \begin{cases} \frac{d\{\log(nT\alpha^2)\}^2}{nT\alpha^2}, & T \le e^{n\alpha^2/(Kd)}\\ e^{-C''n\alpha^2/d}, & \text{otherwise.} \end{cases}$$

Lastly, bounding the error by the sum of the two cases for the different values of T completes the proof.

B.3 Proof of Theorem 8 upper bound

Proof of Theorem 8 (The first procedure). We define the following sets of indices

$$S_1 = \{ j \in [d] : |\theta_j| > 2\varepsilon \}, \quad S_2 = \{ j \in [d] : 0 < |\theta_j| \le 2\varepsilon \} \text{ and } S_0 = \{ j \in [d] : \theta_j = 0 \}.$$

For $j \in [d]$, denote

$$I_j = (n/2)^{-1} \sum_{i=1}^{n/2} r_{i,j} Z_i$$
(97)

which is such that $\mathcal{I} = \{j \in [d] : I_j \ge 1/2\}$. We then consider the following event

$$A = \left(\bigcap_{j \in S_1} \{I_j \ge 1/2\}\right) \cap \left(\bigcap_{j \in S_0} \{I_j < 1/2\}\right).$$

$$(98)$$

On this event, those co-ordinates in S_1 are correctly identified and those in S_0 are correctly rejected. We then consider the error of the estimator on the event A and its complement.

The construction of the estimator $\hat{\theta}$ in (42) ensures that $\hat{\theta}_j = 0$ for $j \notin \mathcal{I}'$, from which we obtain

$$\|\hat{\theta} - \theta\|_2^2 = \sum_{j \notin \mathcal{I}'} \theta_j^2 + \sum_{j \in \mathcal{I}'} (\tilde{\theta}_j - \theta_j)^2.$$
(99)

We consider the first summand in (99). On A, we have that $S_1 \subseteq \mathcal{I}$, and we also have $S_0 \cap \mathcal{I} = \emptyset$ and hence $|\mathcal{I}| \leq s$, giving $\mathcal{I}' = \mathcal{I}$. We have no guarantees on the membership of co-ordinates of S_2 in \mathcal{I} , and so the error is at most $|S_2|(2\varepsilon)^2 \leq 4s\varepsilon^2$. On the complement event A^c , the error can be bounded by the worst case error of 4s, giving

$$\sum_{j \notin \mathcal{I}'} \theta_j^2 = \sum_{j \notin \mathcal{I}'} \theta_j^2 \mathbb{1}\{A\} + \sum_{j \notin \mathcal{I}'} \theta_j^2 \mathbb{1}\{A^c\} \le 4s\varepsilon^2 + 4s\mathbb{1}\{A^c\} \quad \text{a.s.}$$
(100)

As the second summand in (99) corresponds to the ℓ_{∞} -ball estimator of Section 3.2, the mean squared error incurred in this term is bounded by (82) where the dimension of the problem is $s' \leq s$. We now show that the error of the s'-dimensional problem can be bounded by the error of the s-dimensional problem up to constants.

Indeed, we first note that $s^2 \log(nT\alpha^2/s)/(nT\alpha^2)$ is an increasing function of s due to the assumption $n\alpha^2 > \widetilde{C}s \log(ed)$. Further, with c as in (83), $e^{-cn\alpha^2/s}$ as a function of s is also increasing.

It remains to consider the case where $e^{cn\alpha^2/s} < T < e^{cn\alpha^2/s'}$ where the exponential rate of the s-dimensional problem must be compared to the polynomial rate of the s'-dimensional problem. We see that

$$\frac{(s')^2 \log(nT\alpha^2/s')}{nT\alpha^2} \le \left(\frac{s^2 \log(n\alpha^2) + n\alpha^2/s'}{n\alpha^2}\right) e^{-cn\alpha^2/s} \lesssim s^2 e^{-cn\alpha^2/s} \lesssim e^{-c'n\alpha^2/s}$$

where c' > 0 is an absolute constant; the first inequality is by the fact that $e^{-cn\alpha^2/s} < T < e^{-cn\alpha^2/s'}$ in the considered regime; and the last inequality by the fact that $n\alpha^2 > \tilde{C}s \log(ed)$ for some sufficiently large \tilde{C} .

Hence, the error of the s-dimensional ℓ_{∞} -ball estimator is indeed an increasing function of s up to constants, giving

$$\mathbb{E}\left[\sum_{j\in\mathcal{I}'} (\tilde{\theta}_j - \theta_j)^2\right] \lesssim \begin{cases} \frac{s^2 \log(nT\alpha^2/s)}{nT\alpha^2}, & T \le e^{n\alpha^2/(Ks)}, \\ e^{-c'n\alpha^2/s}, & \text{otherwise.} \end{cases}$$
(101)

We then combine (99), (100), Lemma 14, and (101) to see that

$$\mathbb{E}\left(\|\hat{\theta} - \theta\|_2^2\right) \lesssim \frac{s \log(dnT^*\alpha^2)}{T^*} + \begin{cases} \frac{s^2 \log(nT\alpha^2/s)}{nT\alpha^2}, & T \le e^{n\alpha^2/(Ks)}, \\ e^{-c'n\alpha^2/s}, & \text{otherwise}, \end{cases}$$

where we substitute in the value of ε as in (39). When $T \leq e^{n\alpha^2/(Ks)}$, we have that $T^* = T$, and hence,

$$\mathbb{E}\left(\|\hat{\theta} - \theta\|_2^2\right) \lesssim \frac{s\log(dnT\alpha^2)}{T} + \frac{s^2\log(nT\alpha^2/s)}{nT\alpha^2} \lesssim \frac{s\log(dnT\alpha^2)}{T},$$

where the second inequality follows from the assumption that $n\alpha^2 > \widetilde{C}s \log(ed)$ for some suitably large constant \widetilde{C} . When instead $T > e^{n\alpha^2/(Ks)}$, we have that $T^* = e^{n\alpha^2/(Ks)}$, giving

$$\mathbb{E}\left(\|\hat{\theta}-\theta\|_{2}^{2}\right) \lesssim \{s\log(dn\alpha^{2})+n\alpha^{2}\}e^{-n\alpha^{2}/(Ks)}+e^{-c'n\alpha^{2}/s} \lesssim e^{-c''n\alpha^{2}/s},$$

where c'' > 0 is some absolute constant, the first inequality comes from the value of T^* , and the second by the assumption that $n\alpha^2 > \tilde{C}s\log(d)$ for some suitably large constant $\tilde{C} > 0$ which allows the multiplicative prefactors to be absorbed into their respective exponential factor. Hence, we have

$$\mathbb{E}\left(\|\hat{\theta} - \theta\|_{2}^{2}\right) \lesssim \begin{cases} s \log(dnT\alpha^{2})/T, & T \leq e^{n\alpha^{2}/(Ks)}, \\ e^{-c''n\alpha^{2}/s}, & \text{otherwise.} \end{cases}$$

Lastly, bounding the error by the sum of the two cases for the different values of T completes the proof.

Proof of Theorem 8 (Second Procedure). We recall that this second procedure is only used when $n\alpha^2 > \tilde{C}d\log(dnT\alpha^2)$, which is henceforth assumed throughout this proof. We proceed by showing that the entries of the estimator $\tilde{\theta}$ obtained from the ℓ_{∞} -ball procedure of Section 3.2 are suitably close to the true values in θ with high probability so that that the thresholding procedure in (47) correctly shrinks to zero the co-ordinates of $\hat{\theta}$ which correspond to zeros of θ .

It suffices to consider a single fixed co-ordinate $j \in [d]$ as the ℓ_{∞} -ball estimation procedure is independent across co-ordinates.

For $i \in [n]$ we write $\overline{X^{(i)}} = (T^*)^{-1} \sum_{t=1}^{T^*} X_{t,j}^{(i)}$ and $Z_{j,k} = \sum_{i \in N_{j,1}} \widetilde{V}_k^{(i)}$ for the total number of privatised votes for the k-th sub-interval. We denote by $l_j \in [N(\delta)]$ the fixed but unknown index such that $\theta_j \in I_{l_j}$ and define the event

$$A_j = \bigcap_{k:|l_j - k| > 2} \{ Z_{j,l_j} > Z_{j,k} \},\$$

which is analogous to the event (71) considered in the analysis of the univariate mean estimator. Bounding the probability of the complement by the same analysis as that in (72) and (73), we have

$$\mathbb{P}(A_j^c) \le \frac{3}{\delta \wedge 1} e^{-2n\alpha^2/(Kd)} \tag{102}$$

where K is as appears in (72) and, as in (73), we require $\delta < 1$, so to account for when this fails to hold we take the minimum $\delta \wedge 1$ which suffices for an upper bound.

We denote by I_{j^*} the chosen interval for refinement and denote the lower and upper endpoints of this interval by L_j and U_j respectively. Recalling the inflated intervals \tilde{L}_j and \tilde{U}_j defined in (46), we now proceed to show that for a fixed user $i \in N_{j,2}$, on the event A_j the truncated sample mean is close to the sample mean with high probability. Indeed, we first note that

$$\mathbb{E}\left[\left|\Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}}) - \overline{X^{(i)}}\right| \mathbb{1}\{A_j\} \mid j^*\right] \le 4e^{-T^*\delta^2/2} \quad \text{a.s.}$$
(103)

which follows from the same argument as (76) and (77), noting the different value of δ in this case.

Hence, we have

$$\mathbb{P}\left(\left\{\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,2}}\left(\Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}})-\overline{X^{(i)}}\right)\right| > \varepsilon\right\} \cap A_j \left|j^*\right) \\
\leq \frac{1}{\varepsilon}\mathbb{E}\left[\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,2}}\left(\Pi_{\tilde{I}_{j^*}}(\overline{X^{(i)}})-\overline{X^{(i)}}\right)\right| \mathbb{1}\{A_j\} \left|j^*\right] \le \frac{4e^{-T^*\delta/2}}{\varepsilon} \\
\leq \frac{4}{C}\left(\frac{nT^*\alpha^2}{2d\{\log(nT^*\alpha^2)\}^2}\right)^{1/2}e^{-T^*\delta^2/2} \lesssim \frac{1}{nT^*\alpha^2} \quad \text{a.s.}$$
(104)

where the first inequality is by Markov's inequality; the second by (103); the last two inequalities by the values of ε and δ as in (43) and (44) respectively.

We also note the definition of a sub-Exponential random variable (e.g. Wainwright, 2019, Definition 2.7) where it is easy to verify that the Laplacian random variables are sub-Exponential with parameters (2, 2). Hence, by e.g. Wainwright (2019, Equation 2.18) we have that

$$\mathbb{P}\left(\left\{\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,1}}\left(\frac{14\delta}{\alpha}\ell_i\right)\right| > \varepsilon\right\} \cap A_j\right) \le 2\begin{cases} e^{-\frac{n}{16d}\{\varepsilon\alpha/(14\delta)\}^2}, & \text{for } 0 \le \varepsilon\alpha/(14\delta) \le 2\\ e^{-\frac{n}{8d}\{\varepsilon\alpha/(14\delta)\}}, & \text{for } \varepsilon\alpha/(14\delta) > 2. \end{cases}$$
(105)

We note that

$$\frac{\varepsilon\alpha}{14\delta} = \frac{C'}{14} \left(\frac{d\log(nT^*\alpha^2)}{n\alpha^2}\right)^{1/2} \le \frac{C'}{14\widetilde{C}} \le 2$$

for \tilde{C} taken sufficiently large using the fact $n\alpha^2 > \tilde{C}d\log(dnT\alpha^2)$. Hence we have the sub-Gaussian regime in (105), giving

$$\mathbb{P}\left(\left\{\left|\tilde{\theta}_{j}-\theta_{j}\right|>3\varepsilon\right\}\cap A_{j}\right)\leq \mathbb{P}\left(\left\{\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,1}}\left(\Pi_{\tilde{I}_{j^{*}}}(\overline{X^{(i)}})-\overline{X^{(i)}}\right)\right|>\varepsilon\right\}\cap A_{j}\right) \\ +\mathbb{P}\left(\left\{\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,1}}\left(\overline{X^{(i)}}-\theta_{j}\right)\right|>\varepsilon\right\}\cap A_{j}\right)+\mathbb{P}\left(\left\{\left|\frac{1}{n/(2d)}\sum_{i\in N_{j,1}}\frac{14\delta}{\alpha}\ell_{i}\right|>\varepsilon\right\}\cap A_{j}\right) \\ \lesssim \frac{1}{nT^{*}\alpha^{2}}+2e^{-\frac{nT^{*}\varepsilon^{2}}{4d}}+2e^{-\frac{n}{16d}\left(\frac{\varepsilon\alpha}{14\delta}\right)^{2}}=\frac{1}{nT^{*}\alpha^{2}}+2e^{-C'^{2}\left\{\log(nT^{*}\alpha^{2})\right\}^{2}/\alpha^{2}}+2e^{-C'^{2}\log(nT^{*}\alpha^{2})/(3136\alpha^{2})}$$

$$\lesssim \frac{1}{nT^*\alpha^2},\tag{106}$$

where in the second inequality the first probability term is bounded as in (104), the second term by Hoeffding's inequality, and the third term by the sum of the two regimes in (105). The equality comes from substituting the value of ε and δ as in (43) and (44) respectively, and the final inequality is from taking C' sufficiently large.

For a co-ordinate $j \in [d]$, define the event

$$B_j = \left\{ \left| \tilde{\theta}_j - \theta_j \right| \le 3\varepsilon \right\},\,$$

and let $B = \bigcap_{j \in [d]} B_j$, whence we have

$$\begin{split} \mathbb{P}(B^c) &\leq \sum_{j=1}^d \left\{ \mathbb{P}(A_j^c) + \mathbb{P}(B_j^c \cap A_j) \right\} \\ &\lesssim d\left(\frac{1}{\delta} + 1\right) e^{-2n\alpha^2/(Kd)} + d\mathbb{P}\left(\left\{ \left| \tilde{\theta}_j - \theta_j \right| > 3\varepsilon \right\} \cap A_j \right) \\ &\lesssim \left(\frac{d(T^*)^{1/2}}{\{\log(nT\alpha^2)\}^{1/2}} + d \right) e^{-2n\alpha^2/(Kd)} + \frac{d}{nT^*\alpha^2} \\ &\lesssim \frac{1}{T^*} e^{-cn\alpha^2/d} + \frac{d}{nT^*\alpha^2} \lesssim \frac{d}{nT^*\alpha^2}, \end{split}$$

where c > 0 is an absolute constant; the second inequality uses (102); the third by (106) and the value of δ as in (44); and the penultimate by controlling the exponential term as $e^{-2n\alpha^2/(Kd)} \leq (T^*)^{-3/2}e^{-n\alpha^2/(2Kd)}$ and absorbing the prefactor of d using the fact that $n\alpha^2 \geq \tilde{C}d\log(dnT\alpha^2)$ for \tilde{C} sufficiently large.

We now analyse the error of the estimator on the event B, and control its error off this event. We define the sets

$$S_1 = \{ j \in [d] : |\theta_j| > 6\varepsilon \}, \qquad S_2 = \{ j \in [d] : |\theta_j| \le 6\varepsilon \}, \qquad S_0 = \{ j \in [d] : \theta_j = 0 \}.$$

We note by the construction of the final estimator (48), the worst case error is 4s as at most s many co-ordinates are non-zero, and we see that

$$\mathbb{E}\left[\|\hat{\theta}-\theta\|_{2}^{2}\right] \leq 4s\mathbb{P}(B^{c}) + \mathbb{E}(\|\hat{\theta}-\theta\|_{2}^{2}\mathbb{1}\{B\})$$

$$\lesssim \frac{sd}{nT^{*}\alpha^{2}} + \sum_{j\in S_{1}\cup S_{2}}\mathbb{E}(|\hat{\theta}-\theta|^{2}\mathbb{1}\{B\}) + \sum_{j\in S_{0}}\mathbb{E}(|\hat{\theta}-\theta|^{2}\mathbb{1}\{B\})$$

$$\lesssim \frac{sd}{nT^{*}\alpha^{2}} + |S_{1}\cup S_{2}|\varepsilon^{2} \lesssim \frac{sd}{nT^{*}\alpha^{2}} + s\varepsilon^{2} \lesssim \frac{sd\{\log(nT^{*}\alpha^{2})\}^{2}}{nT^{*}\alpha^{2}},$$

where in the first inequality we bound the error on the event C^c by the worst case of 4s, and the third inequality uses the fact that on the event B, (i) co-ordinates in S_0 are correctly thresholded to 0 in the estimator $\hat{\theta}$, contributing no error, and (ii) co-ordinates in $S_1 \cup S_2$ contribute error at most $O(\varepsilon^2)$. The final inequality then follows from the value of ε as in (43).

In the case $T \leq e^{n\alpha^2/(Kd)}$, we have $T^* = T$ which immediately gives the desired rate. In the case $T > e^{n\alpha^2/(Kd)}$, we have $T^* = e^{n\alpha^2/(Kd)}$, which gives

$$\mathbb{E}\left[\|\hat{\theta} - \theta\|^2\right] \lesssim \frac{sd\{\log(nT^*\alpha^2)\}^2}{nT^*\alpha^2} = \frac{sd}{nT^*\alpha^2}\left\{\log(n\alpha^2) + \log(T^*)\right\}^2,$$
$$= \frac{sd}{n\alpha^2}\left(\log(n\alpha^2) + \frac{n\alpha^2}{Kd}\right)^2 e^{-n\alpha^2/(Kd)} \lesssim e^{-c'n\alpha^2/d},$$

where c' > 0 is an absolute constant; the second inequality comes from substituting in the value $e^{n\alpha^2/(Kd)}$ for T^* ; and the last inequality by the fact that $n\alpha^2 \ge \tilde{C}d\log(dnT\alpha^2)$ for some sufficiently large \tilde{C} . Trivially upper bounding the polynomial rate by inserting an extra factor of d into the log term to match that of the previously constructed estimator completes the proof.

B.4 Proof of Theorem 9 upper bound

Proof of Theorem 9.

Step 1: reducing from non-parametric density estimation to mean estimation. Due to the definition of \hat{f} in (50), it holds that

$$\mathbb{E}\left[\|\hat{f} - f\|_{2}^{2}\right] = \mathbb{E}\left[\int\{\hat{f}(x) - f(x)\}^{2} dx\right] = \mathbb{E}\left[\sum_{j=1}^{M} (\hat{\theta}_{j} - \theta_{j})^{2}\right] + \sum_{j=M+1}^{\infty} \theta_{j}^{2} \le \mathbb{E}\left[\|\hat{\theta} - \theta\|_{2}^{2}\right] + \frac{1}{M^{2\beta}}, (107)$$

where the second equality is due to the orthonormality of the basis (18) and the inequality follows from the definition of the Sobolev ellipsoid (19). As $\sup_j \sup_x |\varphi_j(x)| \leq \sqrt{2}$ for all $j \geq 1$, we have that $\theta \in \mathbb{B}_{\infty}(\sqrt{2}) \subset \mathbb{R}^M$. We recall the required assumption for the ℓ_{∞} -ball estimator in Theorem 6, that is, $n\alpha^2 > \widetilde{C}M \log(eM)$ where $\widetilde{C} > 0$ is the same constant as in Theorem 6. This is verified below.

Case 1. When $T \leq (n\alpha^2)^{2\beta+1}/\{C\log(n\alpha^2)\}^{2\beta+2}$, we have that $M = (nT\alpha^2)^{1/(2\beta+2)}$ and therefore $M \leq n\alpha^2/\{C\log(n\alpha^2)\}$. It then follows that

$$\widetilde{C}M\log(eM) \leq \frac{\widetilde{C}n\alpha^2}{C\log(n\alpha^2)}\log\left\{\frac{en\alpha^2}{C\log(n\alpha^2)}\right\} \leq \frac{\widetilde{C}n\alpha^2}{C} \leq n\alpha^2,$$

due to the assumption that $n\alpha^2 \geq \widetilde{C}_{\beta}$ for some sufficiently large \widetilde{C}_{β} and choosing $C \geq \widetilde{C}$.

Case 2. When $T > (n\alpha^2)^{2\beta+1}/\{C\log(n\alpha^2)\}^{2\beta+2}$, we have that $M = n\alpha^2/\{C\log(n\alpha^2)\}$ and therefore

$$\widetilde{C}M\log(eM) = \frac{\widetilde{C}n\alpha^2}{C\log(n\alpha^2)}\log\left\{\frac{en\alpha^2}{C\log(n\alpha^2)}\right\} \le \frac{\widetilde{C}n\alpha^2}{C} \le n\alpha^2,$$

due to the assumption that $n\alpha^2 \geq \widetilde{C}_{\beta}$ for some sufficiently large \widetilde{C}_{β} and choosing $C \geq \widetilde{C}$.

Step 2: applying Theorem 6. We proceed with two cases of M separately.

Step 2.1: case $T \leq (n\alpha^2)^{2\beta+1}/{C\log(n\alpha^2)}^{2\beta+2}$. For sufficiently large $C \geq K(2\beta+1)$ where K is as in (82), we have that

$$T \le (n\alpha^2)^{2\beta+1} \le \exp\left\{\frac{C}{K}\log(n\alpha^2)\right\} \le \exp\left\{\frac{1}{K}(n\alpha^2)^{\frac{2\beta+1}{2\beta+2}}T^{-1/(2\beta+2)}\right\} \le \exp\left(\frac{n\alpha^2}{KM}\right).$$

Hence, combining (82) and (107) we see that

$$\mathbb{E}\left[\|\hat{f} - f\|_{2}^{2}\right] \lesssim \frac{M^{2} \log(nT\alpha^{2}/M)}{nT\alpha^{2}} + \frac{1}{M^{2\beta}}, \\ = \frac{\log\{nT\alpha^{2}/(nT\alpha^{2})^{1/(2\beta+2)}\}}{(nT\alpha^{2})^{2\beta/(2\beta+2)}} + \frac{1}{(nT\alpha^{2})^{2\beta/(2\beta+2)}}, \\ \lesssim C_{\beta} \log(nT\alpha^{2})(nT\alpha^{2})^{-\frac{2\beta}{2\beta+2}},$$
(108)

where $C_{\beta} > 0$ is some constant depending on β and the equality comes from the value of M.

Step 2.2: case $T > (n\alpha^2)^{2\beta+1}/{C\log(n\alpha^2)}^{2\beta+2}$. Consider the two possible regimes for the mean estimation rate (82) separately.

When $T > \exp(n\alpha^2/(KM))$, the exponential rate occurs. Combining (82) and (107), we see that

$$\mathbb{E}\left[\|\hat{f} - f\|_{2}^{2}\right] \lesssim e^{-cn\alpha^{2}/M} + \frac{1}{M^{2\beta}} \\ = e^{-cC\log(n\alpha^{2})} + \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}} \\ \leq \frac{1}{(n\alpha^{2})^{2\beta}} + \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}} \lesssim \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}},$$
(109)

where c > 0 is an absolute constant; the equality from the value of M; and the second inequality by taking C sufficiently large so the $cC \ge 2\beta$.

When $T \leq \exp(n\alpha^2/(KM))$, the polynomial rate in (82) occurs. Note that

$$T \le e^{n\alpha^2/(KM)} = (n\alpha^2)^{C/K},$$
(110)

where the equality comes from the value of M. Combining (82) and (107), we see that

$$\mathbb{E}\left[\|\hat{f} - f\|_{2}^{2}\right] \lesssim \frac{M^{2} \log(nT\alpha^{2}/M)}{nT\alpha^{2}} + \frac{1}{M^{2\beta}} \\ \approx \frac{(n\alpha^{2})^{2}}{\{\log(n\alpha^{2})\}^{2}} \frac{\log(T\log(n\alpha^{2}))}{nT\alpha^{2}} + \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}} \\ \lesssim \frac{n\alpha^{2}}{T} \log(T) + \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}} \\ \lesssim \frac{n\alpha^{2}}{(n\alpha^{2})^{2\beta+1}} \{\log(n\alpha^{2})\}^{2\beta+3} + \frac{\{C\log(n\alpha^{2})\}^{2\beta}}{(n\alpha^{2})^{2\beta}} \lesssim \frac{\{\log(n\alpha^{2})\}^{2\beta+3}}{(n\alpha^{2})^{2\beta}}, \quad (111)$$

where the equality comes from the value of M and the second inequality comes from (110) and the fact that $T > (n\alpha^2)^{2\beta+1}/\{C\log(n\alpha^2)\}^{2\beta+2}\}$. Combining (109) and (111) shows that the rate is $O\{(n\alpha^2)^{-2\beta}\}$ up to constant and poly-logarithmic factors in both cases. Combining also with (108) completes the proof.

C Proof of user-level lower bounds

C.1 Preliminary results

To prove the lower bounds throughout this section we will use Assouad's method (see, e.g. Yu, 1997), in tandem with the results of Acharya et al. (2023a). While Assouad's method reduces the problem to bounding average pairwise TV distances between privatised versions of carefully constructed distributions, these latter results allow us to bound these TV distances uniformly over all suitable privacy mechanisms. For completeness, we will state a version of Assouad's method and the required results of Acharya et al. (2023a) which will be used. We first introduce them in the item-level case before generalising to the user-level case.

For some $k \in \mathbb{N}$, denote by $\mathcal{V} = \{-1, 1\}^k$ the hypercube and let $\{P_v : v \in \mathcal{V}\}$ be a family of distributions indexed by this hypercube. For each $v \in \mathcal{V}$, we write $\theta_v = \theta(P_v)$. The collection $\{P_v : v \in \mathcal{V}\}$ is 2 ϱ -Hamming separated with respect to the loss $\Phi \circ \rho$, if for all $v, v' \in \mathcal{V}$ we have that

$$\Phi\left(\frac{1}{2}\rho(\theta_v,\theta_{v'})\right) \ge 2\varrho \sum_{j=1}^k \mathbb{1}\{v_j \neq v'_j\}.$$
(112)

Note the factor of 1/2 inside $\Phi(\cdot)$; as is common in some formulations of Assouad's lemma, when we have ρ as the ℓ_p norm and $\Phi(\cdot) = (\cdot)^p$, we have a factor of 2^{-p} .

For each $j \in [k]$, we define the mixture distributions

$$P_{+j}^{n} = \frac{1}{2^{k-1}} \sum_{v \in \mathcal{V}: v_{j} = 1} P_{v}^{\otimes n} \quad \text{and} \quad P_{-j}^{n} = \frac{1}{2^{k-1}} \sum_{v \in \mathcal{V}: v_{j} = -1} P_{v}^{\otimes n}.$$
 (113)

Lemma 10 (Assouad's Method). Assuming that (112) holds, we have that

$$\mathcal{R}_{n,1,\infty}(\theta(\mathcal{P}), \Phi \circ \rho) \ge \varrho \sum_{j=1}^{k} \{1 - D_{\mathrm{TV}}(P_{+j}^{n}, P_{-j}^{n})\},\$$

where $\mathcal{R}_{n,1,\infty}$ is the non-private item-level minimax risk defined in Section 1.1.

Assouad's method is often presented in slightly differing forms and the above is a similar statement to those in Yu (1997) and Tsybakov (2009, Lemma 2.12).

For the distribution P_v and any α -LDP mechanism Q, we denote the distribution of the private outcomes for n users by $M_v^n(S) = \int Q(S \mid x^{(1:n)}) dP_v^{\otimes n}(x^{(1:n)})$ and the analogous private mixtures, for $j \in [k]$,

$$M_{+j}^{n}(S) = \frac{1}{2^{k-1}} \sum_{v \in \mathcal{V}: v_{j}=1} M_{v}^{n}(S) = \int Q(S \mid x^{(1:n)}) \,\mathrm{d}P_{+j}^{n}(x^{(1:n)}) \tag{114}$$

and

$$M_{-j}^{n}(S) = \frac{1}{2^{k-1}} \sum_{v \in \mathcal{V}: v_{j} = -1} M_{v}^{n}(S) = \int Q(S \mid x^{(1:n)}) \,\mathrm{d}P_{-j}^{n}(x^{(1:n)}) \tag{115}$$

Applying Lemma 10 we then obtain that

$$\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \ge \varrho \inf_{Q \in \mathcal{Q}_{\alpha}} \sum_{j=1}^{k} \{1 - \|M_{+j}^n - M_{-j}^n\|_{\mathrm{TV}}\},\$$

where $\mathcal{R}_{n,1,\alpha}$ is the item-level private risk defined in Section 1.1, and it remains to bound these TV distances uniformly over $Q \in \mathcal{Q}_{\alpha}$.

To this end, we now introduce the framework of Acharya et al. (2023a). For the considered family of distributions $\{P_v : v \in \mathcal{V}\}$ as in the earlier paragraphs, we require two assumptions to hold. For a vector $u \in \{-1, 1\}^k$ and $i \in [k]$, we define $u^{\oplus i}$ to be the vector with co-ordinates,

$$(u^{\oplus i})_j = \begin{cases} u_j, & j \neq i, \\ -u_j, & \text{otherwise} \end{cases}$$

Assumption 1. For every $v \in \mathcal{V}$ and $i \in [k]$, assume that $P_{v^{\oplus i}} \ll P_v$ and there exist measurable functions $\phi_{v,i} : \mathcal{Z} \to \mathbb{R}$ such that

$$\frac{\mathrm{d}P_{v^{\oplus i}}}{\mathrm{d}P_v} = 1 + \phi_{v,i}$$

It follows from Assumption 1 that $\mathbb{E}_{P_v}(\phi_{v,i}) = 0$ for all $i \in [k]$.

Assumption 2. Assume that there exists $\eta \ge 0$ such that for all $v \in V$ and $i, j \in [k]$ with $i \ne j$, it holds that

$$\mathbb{E}_{P_v}(\phi_{v,i}\phi_{v,j}) = 0 \quad and \quad \mathbb{E}_{P_v}(\phi_{v,i}^2) \le \eta^2.$$

For $i, j \in [k]$ and $v \in \mathcal{V}$, letting $\varphi_{v,i} = 1 + \phi_{v,i}$ the requirements of Assumption 2 become

$$\mathbb{E}_{P_v}(\varphi_{v,i}\varphi_{v,j}) = 1 \quad \text{and} \quad \mathbb{E}_{P_v}(\varphi_{v,i}^2) \le 1 + \eta^2.$$

The following result then follows immediately from Acharya et al. (2023a, Corollary 1) and the fact that $e^{\alpha} - 1 \leq 3\alpha$ for $\alpha \in (0, 1]$.

Lemma 11 (Acharya et al., 2023a). When Assumptions 1 and 2 hold, we have for any $\alpha \in (0, 1]$ and any $Q \in \mathcal{Q}_{\alpha}$ that

$$\frac{1}{k} \sum_{j=1}^{k} D_{\mathrm{TV}}(M_{+j}^{n}, M_{-j}^{n}) \le \sqrt{\frac{21}{k} n \alpha^{2} \eta^{2}},$$

where $\{M_{+j}, M_{-j}\}_{j \in [k]}$ are the private marginals defined in (114) and (115).

Combining the above result with Lemma 10, we obtain

$$\mathcal{R}_{n,1,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \ge \varrho k \left(1 - \sqrt{\frac{21}{k} n \alpha^2 \eta^2} \right).$$
(116)

We now consider the user-level analogue. With \mathcal{V} and the family $\{P_v : v \in \mathcal{V}\}$ as before, suppose that each user draws *T*-many i.i.d. observations from the distribution P_v such that each user's collection of data is distributed as $P_v^{\otimes T}$. Then, for each $j \in [k]$, we define the mixture distributions

$$P_{+j}^{n,T} = \frac{1}{2^{d-1}} \sum_{v \in \mathcal{V}: v_j = 1} (P_v^{\otimes T})^{\otimes n} \text{ and } P_{-j}^{n,T} = \frac{1}{2^{d-1}} \sum_{v \in \mathcal{V}: v_j = -1} (P_v^{\otimes T})^{\otimes n}.$$

Given any user-level α -LDP mechanism Q, we denote the private marginal for n users each with a sample of size T as

$$M_v^{n,T}(S) = \int Q(S \mid x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}) \,\mathrm{d}(P_v^{\otimes T})^{\otimes n}(x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}).$$

Consider the mixtures

$$M_{+j}^{n,T}(S) = \frac{1}{2^{d-1}} \sum_{v \in \mathcal{V}: v_j = 1} M_v^{n,T}(S) = \int Q(S \mid x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}) \,\mathrm{d}P_{\pm j}^{n,T}(x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)})$$

and

$$M_{-j}^{n,T}(S) = \frac{1}{2^{d-1}} \sum_{v \in \mathcal{V}: v_j = -1} M_v^{n,T}(S) = \int Q(S \mid x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}) \, \mathrm{d}P_{\pm j}^{n,T}(x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}).$$

Following the exact arguments above, we have that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \ge \varrho \sum_{j=1}^{k} \{1 - D_{\mathrm{TV}}(M_{+j}^{n,T}, M_{-j}^{n,T})\}.$$

Applying the item-level results above to the distributions $\tilde{P}_v = P_v^{\otimes T}$, we see that, after verifying Assumptions 1 and 2 for the distributions \tilde{P}_v in place of P_v , we will see that Lemma 11 holds in the user-level case where the dependence on T is absorbed into the quantity η^2 . In what follows, for each estimation problem of interest we will construct a family of distributions which obey the separation condition (112) and verify Assumptions 1 and 2.

C.2 Proof of Theorem 6 lower bound

Proof of Theorem 6.

Step 1: constructing a separated family. We consider the hypercube $\mathcal{V} = \{-1, 1\}^d$. For $v \in \mathcal{V}$, let P_v and P'_v be the distributions where, for $X \sim P_v$ and $X' \sim P'_v$, we have that the *j*-th, $j \in [d]$, co-ordinates are, independently of the other co-ordinates, distributed as

$$X_{j} = \begin{cases} 1, & \text{with probability } (1 + \delta v_{j})/2, \\ -1, & \text{with probability } (1 - \delta v_{j})/2, \end{cases}$$
(117)

and

$$X'_{j} = \begin{cases} d^{-1/2}, & \text{with probability } (1 + \delta v_{j})/2, \\ -d^{-1/2}, & \text{with probability } (1 - \delta v_{j})/2. \end{cases}$$
(118)

By construction we have that $P_v \in \mathcal{P}_d$ and $P'_v \in \mathcal{P}'_d$ as defined in (12), for all $v \in \mathcal{V}$. Note that $\theta(P_v) = \mathbb{E}_{P_v}(X) = \delta v$ and $\theta(P'_v) = \mathbb{E}_{P'_v}(X') = \delta v/d^{1/2}$. For the separation condition (112), it then holds that

$$\frac{1}{4} \|\theta(P_v) - \theta(P_{v'})\|_2^2 \ge \delta^2 \sum_{j=1}^d \mathbb{1}\{v_j \neq v'_j\} \quad \text{and} \quad \frac{1}{4} \|\theta(P'_v) - \theta(P'_{v'})\|_2^2 \ge \frac{\delta^2}{d} \sum_{j=1}^d \mathbb{1}\{v_j \neq v'_j\},$$

which shows that the distributions are δ^2 -separated and δ^2/d -separated for the ℓ_{∞} - and ℓ_2 -ball constructions respectively.

Step 2: Verifying Assumptions 1 and 2. We claim that, for both of the families of distributions obtained by taking the T-fold product of the distributions defined in (117) and (118), we have that Assumptions 1 and 2 hold with

$$\eta^{2} = \left(\frac{1+3\delta^{2}}{1-\delta^{2}}\right)^{T} - 1.$$
(119)

In this step, we show this holds for the construction (117). The same result for the construction (118) follows by replacing all instances of $\mathbb{1}\{x_{t,j} = \pm 1\}$ in what follows with $\mathbb{1}\{x_{t,j} = \pm d^{-1/2}\}$ and P_v by P'_v , with the calculations being otherwise identical.

We see that, for $i \in [d]$,

$$\frac{\mathrm{d}P_{v^{\oplus i}}^{\otimes T}}{\mathrm{d}P_{v}^{\otimes T}}(x_{1:T}) = \frac{\prod_{t=1}^{T} \prod_{j=1}^{d} \left(\frac{1+\delta v_{j}^{\oplus i}}{2}\right)^{\mathbb{I}\left\{x_{t,j}=1\right\}} \left(\frac{1-\delta v_{j}^{\oplus i}}{2}\right)^{\mathbb{I}\left\{x_{t,j}=-1\right\}}}{\prod_{t=1}^{T} \prod_{j=1}^{d} \left(\frac{1+\delta v_{j}}{2}\right)^{\mathbb{I}\left\{x_{t,j}=1\right\}} \left(\frac{1-\delta v_{j}}{2}\right)^{\mathbb{I}\left\{x_{t,j}=-1\right\}}}$$

$$=\prod_{t=1}^{T} \left(\frac{1-\delta v_i}{1+\delta v_i}\right)^{\mathbb{I}\{x_{t,i}=1\}} \left(\frac{1+\delta v_i}{1-\delta v_i}\right)^{\mathbb{I}\{x_{t,i}=-1\}} =:\varphi_{v,i}(x_{1:T})$$

This verifies Assumption 1. As for Assumption 2, we see further that have for $i \neq j$ that

$$\begin{split} & \mathbb{E}_{P_{v}}(\varphi_{v,i}\varphi_{v,j}) \\ &= \prod_{t=1}^{T} \mathbb{E}_{P_{v}} \left[\left(\frac{1-\delta v_{i}}{1+\delta v_{i}} \right)^{1\{X_{t,i}=1\}} \left(\frac{1+\delta v_{i}}{1-\delta v_{i}} \right)^{1\{X_{t,i}=-1\}} \left(\frac{1-\delta v_{j}}{1+\delta v_{j}} \right)^{1\{X_{t,j}=1\}} \left(\frac{1+\delta v_{j}}{1-\delta v_{j}} \right)^{1\{X_{1,j}=-1\}} \right] \\ &= \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1-\delta v_{i}}{1+\delta v_{i}} \right)^{1\{X_{1,i}=1\}} \left(\frac{1+\delta v_{i}}{1-\delta v_{i}} \right)^{1\{X_{1,i}=-1\}} \left(\frac{1-\delta v_{j}}{1+\delta v_{j}} \right)^{1\{X_{1,j}=1\}} \left(\frac{1+\delta v_{j}}{1-\delta v_{j}} \right)^{1\{X_{1,j}=-1\}} \right] \right\}^{T} \\ &= \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1+\delta v_{i}}{1-\delta v_{i}} \right) \left(\frac{1-\delta v_{i}}{1+\delta v_{i}} \right)^{21\{X_{1,i}=1\}} \left(\frac{1+\delta v_{j}}{1-\delta v_{j}} \right) \left(\frac{1-\delta v_{j}}{1+\delta v_{j}} \right)^{21\{X_{1,i}=1\}} \right] \right\}^{T} \\ &= \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1+\delta v_{i}}{1-\delta v_{i}} \right) \left(\frac{1-\delta v_{i}}{1+\delta v_{i}} \right)^{21\{X_{1,i}=1\}} \right] \right\}^{T} \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1+\delta v_{j}}{1-\delta v_{j}} \right) \left(\frac{1-\delta v_{j}}{1+\delta v_{j}} \right)^{21\{X_{1,j}=1\}} \right] \right\}^{T} , \end{split}$$

where in the second equality we use the fact that the $X_{t,i}$ are i.i.d. across $t \in [T]$, the third by the fact that $\mathbb{1}\{X_{t,i}=1\}=1-\mathbb{1}\{X_{t,i}=-1\}$, and the last by the independence of the co-ordinates of X_1 . Focusing on one of the expectation factors, we have that

$$\mathbb{E}_{P_v}\left[\left(\frac{1+\delta v_i}{1-\delta v_i}\right)\left(\frac{1-\delta v_i}{1+\delta v_i}\right)^{21\{X_{1,i}=1\}}\right] = \left(\frac{1+\delta v_i}{1-\delta v_i}\right)\mathbb{E}_{P_v}\left[\left(\frac{1-\delta v_i}{1+\delta v_i}\right)^{21\{X_{1,i}=1\}}\right]$$
$$= \left(\frac{1+\delta v_i}{1-\delta v_i}\right)\left[\left(\frac{1-\delta v_i}{1+\delta v_i}\right)^2\frac{(1+\delta v_i)}{2} + \frac{(1-\delta v_i)}{2}\right] = 1.$$

Hence, we have that $\mathbb{E}_{P_v}(\varphi_{v,i}\varphi_{v,j}) = 1$ for all $i \neq j$, as required.

To specify η^2 in Assumption 2, we consider that

$$\begin{split} \mathbb{E}_{P_{v}}(\varphi_{v,i}^{2}) &= \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1 - \delta v_{i}}{1 + \delta v_{i}} \right)^{21\{X_{1,i}=1\}} \left(\frac{1 + \delta v_{i}}{1 - \delta v_{i}} \right)^{21\{X_{1,i}=-1\}} \right] \right\}^{T} \\ &= \left\{ \mathbb{E}_{P_{v}} \left[\left(\frac{1 + \delta v_{i}}{1 - \delta v_{i}} \right)^{2} \left(\frac{1 - \delta v_{i}}{1 + \delta v_{i}} \right)^{41\{X_{1,i}=1\}} \right] \right\}^{T} \\ &= \left[\left(\frac{1 + \delta v_{i}}{1 - \delta v_{i}} \right)^{2} \left\{ \left(\frac{1 - \delta v_{i}}{1 + \delta v_{i}} \right)^{4} \frac{(1 + \delta v_{i})}{2} + \frac{(1 - \delta v_{i})}{2} \right\} \right]^{T} = \left(\frac{1 + 3\delta^{2}}{1 - \delta^{2}} \right)^{T} \end{split}$$

where the first equality comes from setting j = i in (120). Hence, we have that (119) holds.

Step 3: obtaining lower bounds. We set $\delta = \{cd/(nT\alpha^2)\}^{1/2}$ for some absolute constant c > 0. Hence, using our assumption that $n\alpha^2 \gtrsim d\log(ed)$, when c is sufficiently small we have that

$$\eta^{2} = \left(1 + \frac{4\delta^{2}}{1 - \delta^{2}}\right)^{T} - 1 \le \left(1 + \frac{8cd}{nT\alpha^{2}}\right)^{T} - 1 \le \frac{16cd}{n\alpha^{2}},\tag{121}$$

where the first and second inequalities follow once we choose c > 0 small enough that $\delta^2 \leq 1/2$ and $8cd/(n\alpha^2) < 1$, respectively. As Assumptions 1 and 2 hold, we have that for the ℓ_{∞} -ball case that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}_d), \|\cdot\|_2^2) \ge \frac{\delta^2 d}{2} \left(1 - \sqrt{\frac{21}{d}n\alpha^2\eta^2}\right) \ge \frac{cd^2}{2nT\alpha^2} \left(1 - \sqrt{336c}\right) \gtrsim \frac{d^2}{nT\alpha^2}.$$

where the first inequality is from (116) and the fact that the distributions are δ^2 -separated; the second from (121) and the value of δ ; and the last as c > 0 is sufficiently small. Combining with the infinite-T lower bound in (13), for which the required assumptions are satisfied, completes the proof for this case.

For the ℓ_2 -ball case, we similarly have that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}'_d), \|\cdot\|_2^2) \ge \frac{\delta^2}{2} \left(1 - \sqrt{\frac{21}{d}n\alpha^2\eta^2}\right) \ge \frac{cd}{2nT\alpha^2} \left(1 - \sqrt{336c}\right) \gtrsim \frac{d}{nT\alpha^2}$$

and combining with the infinite-T lower bound in (14), for which the required assumptions are satisfied, completes the proof.

C.3 Proof of Theorem 8 lower bound

As the distributions we are considering have s-sparse means, a standard Assouad's method construction where, for example, -1 corresponds to a zero entry for the mean and +1 corresponds to a non-zero entry, will fail as any $v \in \mathcal{V}$ with $\sum_{j=1}^{k} \mathbb{1}\{v_j = 1\} > s$ will not lie in the family we are considering. We instead consider a variant of Assouad's lemma, similar to that considered in Acharya et al. (2023a), that allows us to consider a family of distributions indexed by the hypercube even when not all such distributions are a member of the family our estimation problem considers.

Firstly, we require a strengthening of the separation condition (112) which holds with equality. We say the collection $\{P_v : v \in \mathcal{V}\}$ is 2 ρ -Hamming separated in equality with respect to the loss $\Phi \circ \rho$ if for all $v, v' \in \mathcal{V}$,

$$\Phi\left(\frac{1}{2}\rho(\theta_v,\theta_{v'})\right) = 2\varrho \sum_{j=1}^k \mathbb{1}\{v_j \neq v'_j\}.$$
(122)

We then have the following lemma and corollary, the proofs of which are contained in Appendix D.

Lemma 12. For $k \in \mathbb{N}$, let \mathcal{P} be a family of distributions and consider the hypercube $\mathcal{V} = \{-1, 1\}^k$, the elements of which index a family $\{P_v : v \in \mathcal{V}\} \subseteq \mathcal{P}$ satisfying the separation condition (122). Let $\mathcal{P}^* \subseteq \mathcal{P}$ be some subset of the family of distributions and V be a random variable on the set $\mathcal{V} = \{-1, 1\}^k$. Writing $\mathcal{V}^* = \{v \in \mathcal{V} : P_v \in \mathcal{P}^*\}$, we have that

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \ge \mathbb{E}_V \left[\mathbb{E}_{P_V} \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \right] - 2k \varrho \mathbb{P}(V \notin \mathcal{V}^*).$$

Corollary 13. Under the same conditions in Lemma 12, suppose that the random vector V consists of *i.i.d.* random variables as co-ordinates such that, for each $j \in [k]$, we have that $\mathbb{P}(V_j = 1) = \tau = 1 - \mathbb{P}(V_j = -1)$ with $\tau \leq 1/2$. Suppose also that for some constant c > 0, we have that $\mathbb{P}(V \notin \mathcal{V}^*) \leq c\tau$, then

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \ge 2\varrho \tau \left[\sum_{j=1}^k \{ 1 - D_{\mathrm{TV}}(P_{+j}, P_{-j}) \} - ck \right],$$

where P_{+j} and P_{-j} are the mixture distributions as defined in (113).

With these results, we now prove the lower bound for sparse mean estimation.

Proof of Theorem 8.

Step 1: constructing a separated family. We consider the hypercube $\mathcal{V} = \{-1, 1\}^d$. For $v \in \mathcal{V}$, let P_v be the distribution where, for $X \sim P_v$, we have that the *j*-th co-ordinate is, independently of the other co-ordinates, distributed as

$$X_{j} = \begin{cases} 1, & \text{with probability } \frac{1}{2} + \frac{\delta}{4}(v_{j}+1), \\ -1, & \text{with probability } \frac{1}{2} - \frac{\delta}{4}(v_{j}+1), \end{cases} \quad j \in [d].$$
(123)

Note that $\theta(P_v) = \mathbb{E}_{P_v}(X) = (\delta/2)(v+1)$, and hence the separation condition in (122) holds as

$$\frac{1}{4} \|\theta(P_v) - \theta(P_{v'})\|_2^2 = \frac{\delta^2}{4} \sum_{j=1}^d \mathbb{1}\{v_j \neq v'_j\}.$$

Step 2: verifying Assumptions 1 and 2. We claim that for the family of distributions obtained by taking the *T*-fold product of the distributions defined in (123), when $\delta \leq 2^{-1/2}$, Assumptions 1 and 2 hold with $\eta^2 = (1 + 2\delta^2)^T - 1$.

Step 2.1: Assumption 1. For $i \in [d]$, denote $\#_{+i}(x_{1:T}) = \sum_{t=1}^{T} \mathbb{1}\{x_{t,i} = 1\}$. We have that

$$\begin{split} \frac{\mathrm{d}P_{v^{\oplus i}}^{\otimes T}}{\mathrm{d}P_{v}^{\otimes T}}(x_{1:T}) &= \frac{\prod_{t=1}^{T} \prod_{j=1}^{d} \left(\frac{1}{2} + \frac{\delta}{4}(1+v_{j}^{\oplus i})\right)^{\mathbb{I}\left\{x_{t,j}=1\right\}} \left(\frac{1}{2} - \frac{\delta}{4}(1+v_{j}^{\oplus i})\right)^{\mathbb{I}\left\{x_{t,j}=-1\right\}}}{\prod_{t=1}^{T} \prod_{j=1}^{d} \left(\frac{1}{2} + \frac{\delta}{4}(1+v_{j})\right)^{\mathbb{I}\left\{x_{t,j}=1\right\}} \left(\frac{1}{2} - \frac{\delta}{4}(1+v_{j})\right)^{\mathbb{I}\left\{x_{t,j}=-1\right\}}}{\left(\frac{1}{2} - \frac{\delta}{4}(1+v_{j})\right)^{\mathbb{I}\left\{x_{t,i}=-1\right\}}} \\ &= \prod_{t=1}^{T} \left(\frac{1 + \frac{\delta}{2}(1-v_{i})}{1 + \frac{\delta}{2}(1+v_{i})}\right)^{\mathbb{I}\left\{x_{t,i}=1\right\}} \left(\frac{1 - \frac{\delta}{2}(1-v_{i})}{1 - \frac{\delta}{2}(1+v_{i})}\right)^{\mathbb{I}\left\{x_{t,i}=-1\right\}} \\ &= \left(\frac{1 - \frac{\delta}{2}(1-v_{i})}{1 - \frac{\delta}{2}(1+v_{i})}\right)^{T} \left(\frac{1 + \frac{\delta}{2}(1-v_{i})}{1 + \frac{\delta}{2}(1+v_{i})} \cdot \frac{1 - \frac{\delta}{2}(1+v_{i})}{1 - \frac{\delta}{2}(1-v_{i})}\right)^{\#_{+i}(x_{1:T})} = \varphi_{v,j}(x_{1:T}). \end{split}$$

We therefore verify Assumption 1.

Step 2.1: Assumption 2. Note that for $i, j \in [d], i \neq j$,

$$\begin{split} & \mathbb{E}_{P_{v}}\left(\varphi_{v,i}\varphi_{v,j}\right) \\ = & \mathbb{E}_{P_{v}}\left[\left(\frac{1-\frac{\delta}{2}(1-v_{i})}{1-\frac{\delta}{2}(1+v_{i})}\right)^{T}\left(\frac{1+\frac{\delta}{2}(1-v_{i})}{1+\frac{\delta}{2}(1+v_{i})} \cdot \frac{1-\frac{\delta}{2}(1+v_{i})}{1-\frac{\delta}{2}(1-v_{i})}\right)^{\#_{+i}(x_{1:T})}\right] \\ & \times \left(\frac{1-\frac{\delta}{2}(1-v_{j})}{1-\frac{\delta}{2}(1+v_{j})}\right)^{T}\left(\frac{1+\frac{\delta}{2}(1-v_{j})}{1+\frac{\delta}{2}(1+v_{j})} \cdot \frac{1-\frac{\delta}{2}(1+v_{j})}{1-\frac{\delta}{2}(1-v_{j})}\right)^{\#_{+j}(x_{1:T})}\right] \\ & = \left(\frac{1-\frac{\delta}{2}(1-v_{i})}{1-\frac{\delta}{2}(1+v_{i})} \cdot \frac{1-\frac{\delta}{2}(1-v_{j})}{1-\frac{\delta}{2}(1+v_{j})}\right)^{T} \mathbb{E}_{P_{v}}\left[\left(\frac{1+\frac{\delta}{2}(1-v_{i})}{1+\frac{\delta}{2}(1+v_{i})} \cdot \frac{1-\frac{\delta}{2}(1+v_{i})}{1-\frac{\delta}{2}(1-v_{i})}\right)^{\#_{+i}(x_{1:T})}\right] \\ & \times \mathbb{E}_{P_{v}}\left[\left(\frac{1+\frac{\delta}{2}(1-v_{j})}{1+\frac{\delta}{2}(1+v_{j})} \cdot \frac{1-\frac{\delta}{2}(1+v_{j})}{1-\frac{\delta}{2}(1-v_{j})}\right)^{\#_{+j}(x_{1:T})}\right]. \end{split}$$

In addition it holds that

$$\begin{split} \mathbb{E}_{P_{v}} \left[\left(\frac{1 + \frac{\delta}{2}(1 - v_{i})}{1 + \frac{\delta}{2}(1 + v_{i})} \cdot \frac{1 - \frac{\delta}{2}(1 + v_{i})}{1 - \frac{\delta}{2}(1 - v_{i})} \right)^{\#_{+i}(x_{1:T})} \right] \\ &= \prod_{t=1}^{T} \mathbb{E}_{P_{v}} \left[\left(\frac{1 + \frac{\delta}{2}(1 - v_{i})}{1 + \frac{\delta}{2}(1 + v_{i})} \cdot \frac{1 - \frac{\delta}{2}(1 + v_{i})}{1 - \frac{\delta}{2}(1 - v_{i})} \right) \mathbb{1}\{X_{t,i} = 1\} + \mathbb{1}\{X_{t,i} = -1\} \right] \\ &= \left[\left(\frac{1 + \frac{\delta}{2}(1 - v_{i})}{1 + \frac{\delta}{2}(1 + v_{i})} \cdot \frac{1 - \frac{\delta}{2}(1 + v_{i})}{1 - \frac{\delta}{2}(1 - v_{i})} \right) \left(\frac{1}{2} + \frac{\delta}{4}(1 + v_{i}) \right) + \left(\frac{1}{2} - \frac{\delta}{4}(1 + v_{i}) \right) \right]^{T} \\ &= \left(\frac{1 - \frac{\delta}{2}(1 + v_{i})}{1 - \frac{\delta}{2}(1 - v_{i})} \right)^{T} . \end{split}$$

We then have that $\mathbb{E}_{P_v}(\varphi_{v,i}\varphi_{v,j}) = 1$.

We now consider

$$\mathbb{E}_{P_{v}}(\varphi_{v,i}^{2}) = \left(\frac{1 - \frac{\delta}{2}(1 - v_{i})}{1 - \frac{\delta}{2}(1 + v_{i})}\right)^{2T} \mathbb{E}_{P_{v}}\left[\left(\frac{1 + \frac{\delta}{2}(1 - v_{i})}{1 + \frac{\delta}{2}(1 + v_{i})} \cdot \frac{1 - \frac{\delta}{2}(1 + v_{i})}{1 - \frac{\delta}{2}(1 - v_{i})}\right)^{2\#_{+i}(x_{1:T})}\right].$$

It holds that

$$\mathbb{E}_{P_{v}}\left[\left(\frac{1+\frac{\delta}{2}(1-v_{i})}{1+\frac{\delta}{2}(1+v_{i})}\cdot\frac{1-\frac{\delta}{2}(1+v_{i})}{1-\frac{\delta}{2}(1-v_{i})}\right)^{2\#_{+i}(x_{1:T})}\right]$$

$$= \left[\left(\frac{1 + \frac{\delta}{2}(1 - v_i)}{1 + \frac{\delta}{2}(1 + v_i)} \cdot \frac{1 - \frac{\delta}{2}(1 + v_i)}{1 - \frac{\delta}{2}(1 - v_i)} \right)^2 \left(\frac{1}{2} + \frac{\delta}{4}(1 + v_i) \right) + \left(\frac{1}{2} - \frac{\delta}{4}(1 + v_i) \right) \right]^T$$
$$= \left(\frac{(1 + \frac{\delta}{2}(1 + v))(1 + \frac{\delta^2}{4}(3v^2 - 2v - 1))}{(1 - \frac{\delta}{2}(1 - v))^2(1 + \frac{\delta}{2}(1 + v))} \right)^T$$

which combined with the fact that $v_i^2 \equiv 1$ for all $i \in [d]$, yields that

$$\mathbb{E}_{P_v}(\varphi_{v,i}^2) = \left(\frac{1 + \frac{\delta^2}{2}(1-v)}{1 - \frac{\delta^2}{2}(1+v)}\right)^T = \begin{cases} \left(\frac{1}{1-\delta^2}\right)^T, & v = 1\\ (1+\delta^2)^T, & v = -1. \end{cases}$$

As an upper bound suffices, we have that Assumption 2 holds with $\eta^2 = (1 + 2\delta^2)^T - 1$ when $\delta^2 \le 1/2$.

Step 3: obtaining lower bounds. Letting \mathcal{P} denote the *T*-fold product distributions induced by the construction (123), we consider the subset $\mathcal{P}^* = \{P \in \mathcal{P} : ||\mathbb{E}_P(X)||_0 \leq s\}$ and denote $\mathcal{V}^* = \{v \in \mathcal{V} : ||v||_0 \leq s\}$. We have that $v \in \mathcal{V}^*$ if and only if $P_v \in \mathcal{P}^*$. Lastly, for a fixed user-level α -LDP mechanism Q, denote the private marginals satisfying the sparsity condition

$$\mathcal{M}^* = \left\{ M(\cdot) = \int Q(\cdot | x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}) \, \mathrm{d}P(x_{1:T}^{(1)}, \dots, x_{1:T}^{(n)}) : P \in \mathcal{P}^* \right\}.$$

Let $V \in \{0,1\}^d$ be the random variable such that its co-ordinates are mutually independent with $\mathbb{P}(V_j = 1) = \tau = 1 - \mathbb{P}(V_j = -1)$ where $\tau = s/(2d) \leq 1/2$, and denote $V_+ = \sum_{j=1}^d \mathbb{1}\{V_j = 1\}$. In particular, we have that $V_+ \sim \operatorname{Bin}(d, \tau)$. By an application of Bernstein's inequality (e.g. Wainwright, 2019, Proposition 2.14), we have that

$$\mathbb{P}(V_{+} \ge 2d\tau) \le \exp\left(-\frac{(d\tau)^{2}}{2\left\{\sum_{j=1}^{d} \mathbb{P}(V_{j}=1) + d\tau/3\right\}}\right) \le e^{-3d\tau/8}$$
$$= e^{-3s/16} \le \frac{1}{d} \le \frac{3\tau}{8\log(ed)},$$

where the equality is by the value of τ ; the penultimate inequality by the requirement $16 \log(ed)/3 \leq s$; and the final inequality by the fact that $16 \log(ed)/3 \leq s$ and the value of τ .

Thus, we have that $\mathbb{P}(V \notin \mathcal{V}^*) \leq \frac{3\pi}{8}$ under the assumption $16 \log(ed)/3 \leq s \leq d$. Using Lemma 11 and Corollary 13 with the private marginals \mathcal{M}^* , we obtain

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \ge 2d\delta^2 \tau \left[1 - \sqrt{\frac{21n\alpha^2\eta^2}{d}} - \frac{3}{8}\right].$$

We now set

$$\delta = \left\{ \frac{1}{2} \left(1 + \frac{d}{84n\alpha^2} \right)^{\frac{1}{T}} - \frac{1}{2} \right\}^{1/2} \wedge \left(\frac{1}{2T} \right)^{1/2},$$
(124)

noting the requirement $\delta^2 \leq 1/2$ is satisfied. When $\delta = \left\{\frac{1}{2}\left(1 + \frac{d}{84n\alpha^2}\right)^{\frac{1}{T}} - \frac{1}{2}\right\}^{1/2}$, we have that $\eta^2 = d/(84n\alpha^2)$, giving that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \gtrsim s \left[\left(1 + \frac{d}{84n\alpha^2} \right)^{\frac{1}{T}} - 1 \right].$$

When $\delta = \{1/(2T)\}^{1/2}$, we have that $\eta^2 = (1 + 1/T)^{1/T} - 1 \leq 1$. In this case, the second term in (124) is smaller than the first, which upon rearrangement gives

$$\frac{21n\alpha^2}{d} \geq \frac{1}{4\{(1+1/T)^T-1\}} \geq \frac{1}{4}$$

Hence, we have

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \gtrsim \frac{s}{T} \left[1 - \sqrt{\frac{21n\alpha^2}{d}} - \frac{3}{8} \right] \gtrsim \frac{s}{T}.$$

These two choices of the value of δ together give a lower bound of

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}), \Phi \circ \rho) \gtrsim \frac{s}{T} \wedge s \left\{ \left(1 + \frac{d}{84n\alpha^2} \right)^{1/T} - 1 \right\}.$$

Lastly, combining with the infinite-T lower bound in (17), for which the required assumptions are satisfied, completes the proof.

C.4 Proof of Theorem 9 lower bound

We first define a Hölder space of functions denoted by $\mathcal{H}_{\beta,r}$, for $\beta \in \mathbb{N}$ and r > 0. For any $f[0,1] \to \mathbb{R}$, $f \in \mathcal{H}_{\beta,r}$, if it is $(\beta - 1)$ -times differentiable and satisfies that

$$|f^{(\beta-1)}(x) - f^{(\beta-1)}(y)| \le r|x-y|$$

Proof of Theorem 9.

Step 1: constructing a separated family. Consider the bump function

$$g(x) = \begin{cases} e^{-1/\{x(1/2-x)\}}, & x \in [0, 1/2], \\ -e^{-1/\{(x-1/2)(1-x)\}}, & x \in (1/2, 1], \\ 0, & \text{otherwise.} \end{cases}$$

and define the function g_{β} that $g_{\beta}(x) = c_{\beta}g(x)$ with a sufficiently small absolute constant $c_{\beta} > 0$ depending only on β , so that

$$\sup_{x \in [0,1]} \max\{|g_{\beta}^{(\beta)}(x)|, |g_{\beta}(x)|\} \le 1.$$

We remark that the choice of c_{β} exists, since $g(\cdot)$ is infinitely differentiable with bounded derivatives.

For $k \in \mathbb{N}$ to be specified and for each $j \in [k]$, we let

$$g_{\beta,j}(x) = \frac{r\pi^{\beta}}{2k^{\beta}}g_{\beta}\left(k\left(x-\frac{j-1}{k}\right)\right), \quad x \in \mathbb{R}.$$

Note that $g_{\beta,j}(x) \neq 0$ if and only if $x \in [(j-1)/k, j/k]$, and for $x, y \in [0, 1]$,

$$|g_{\beta,j}^{(\beta-1)}(x) - g_{\beta,j}^{(\beta-1)}(y)| = \frac{r\pi^{\beta}}{2k} \left| g_{\beta}^{(\beta-1)}(kx - (j-1)) - g_{\beta}^{(\beta-1)}(ky - (j-1)) \right| \le \frac{r\pi^{\beta}}{2} |x - y|$$

where the equality follows from the definition of $g_{\beta,j}^{(\beta-1)}(x)$, and the inequality via (i) an application of the mean value theorem, (ii) the fact that the derivative is bounded as $|g_{\beta}^{(\beta)}(x)| \leq 1$ for all x and (*iii*) $k \geq 1$. It then holds that $g_{\beta,j}^{(\beta-1)} \in \mathcal{H}_{\beta,r\pi^{\beta}/2}, j \in [k]$. We then define the family of densities indexed by the hypercube $\mathcal{V} = \{-1, 1\}^k$ by

$$\left\{f_v = 1 + \sum_{j=1}^k v_j g_{\beta,j} : v \in \mathcal{V}\right\} \subseteq \mathcal{F}_{\beta,r},\tag{125}$$

where the inclusion in the Sobolev space of densities $\mathcal{F}_{\beta,r}$ follows from

- the fact that each f_v is a sum of functions in $\mathcal{H}_{\beta,r\pi^\beta/2}$ with disjoint support and is thus itself is in $\mathcal{H}_{\beta,r\pi^\beta}$ (see e.g. Tsybakov, 2009, Section 2.6.1);
- the fact that (for integer β) $\mathcal{H}_{\beta,r\pi^{\beta}} \subseteq S_{\beta,r}$ by Definition 1.11 and Proposition 1.14 in Tsybakov (2009); and
- the fact that $g_{\beta,j}$ is small enough of a perturbation that $f_v \ge 0$, and that, since $\int_0^1 g_\beta(x) dx = 0$, we also have that $\int f_v(x) dx = 1$.

We also have that

$$\|f_v - f_{v'}\|_2^2 = 2\sum_{j=1}^k \mathbb{1}\{v_j \neq v'_j\} \int_{\frac{j-1}{k}}^{\frac{j}{k}} \{g_{\beta,j}(x)\}^2 \,\mathrm{d}x \ge \frac{c'_\beta r^2 \pi^{2\beta}}{2k^{2\beta+1}} \sum_{j=1}^k \mathbb{1}\{v_j \neq v'_j\},\tag{126}$$

where the inequality comes from integrating by substitution and the fact that $\int \{g_{\beta}(x)\}^2 dx > c'_{\beta}$ for some constant $c'_{\beta} > 0$ depending on β . The family of distributions is $c'_{\beta}r^2\pi^{2\beta}/(4k^{2\beta+1})$ -separated under squared- L_2 loss.

Step 2: Verifying Assumptions 1 and 2. We claim that, for the family of distributions obtained by taking the T-fold product of the distributions defined via the densities in (125), Assumptions 1 and 2 hold with

$$\eta^2 = \left(1 + \frac{2r^2\pi^{2\beta}}{k^{2\beta+1}}\right)^T - 1.$$

Step 2.1: Assumption 1. For any $v, v' \in \mathcal{V}$ and a fixed $x \in [0, 1]$, we have by the construction in (125) that $f_v(x) = 1 + v_j g_{\beta,l}(x)$, $f_{v'}(x) = 1 + v'_j g_{\beta,l}(x)$ for some $l \in [k]$. Further, for any $j \in [k]$

$$\sup_{x} |g_{\beta,j}(x)| \le \frac{r\pi^{\beta}}{2k^{\beta}} \sup_{x} |g_{\beta}(x)| \le \frac{r\pi^{\beta}c_{\beta}}{2k^{\beta}} \sup_{x} |g(x)| \le \frac{1}{2},$$
(127)

where the first two inequalities are from the definitions of the g_{β} and g, and the final inequality comes from taking $c_{\beta} > 0$ small enough. We then see that, for $i \in [k]$,

$$\frac{\mathrm{d}P_{v^{\oplus i}}^{\otimes T}}{\mathrm{d}P_{v}^{\otimes T}}(x_{1:T}) = \frac{\prod_{t=1}^{T} f_{v^{\oplus i}}(x_{t})}{\prod_{t=1}^{T} f_{v}(x_{t})} = \frac{\prod_{t=1}^{T} [f_{v}(x_{t}) - 2v_{i}g_{\beta,i}(x_{t})]}{\prod_{t=1}^{T} f_{v}(x_{t})}$$
$$= \prod_{t=1}^{T} \left[1 - \frac{2v_{i}g_{\beta,i}(x_{t})}{1 + v_{i}g_{\beta,i}(x_{t})} \right] =: \varphi_{v,i}(x_{1:T}),$$

where we use (127) to ensure that the denominators are bounded away from zero. We therefore verify Assumption 1.

Step 2.1: Assumption 2. Note that for $i, j \in [k], i \neq j$,

$$\begin{split} \mathbb{E}_{P_{v}}(\varphi_{v,i}\varphi_{v,j}) &= \prod_{t=1}^{T} \mathbb{E}_{P_{v}} \left[\left(1 - \frac{2v_{i}g_{\beta,i}(X_{t})}{1 + v_{i}g_{\beta,i}(X_{t})} \right) \left(1 - \frac{2v_{j}g_{\beta,j}(X_{t})}{1 + v_{j}g_{\beta,j}(X_{t})} \right) \right] \\ &= \left\{ \mathbb{E}_{P_{v}} \left[\left(1 - \frac{2v_{i}g_{\beta,i}(X_{1})}{1 + v_{i}g_{\beta,i}(X_{1})} \right) \left(1 - \frac{2v_{j}g_{\beta,j}(X_{1})}{1 + v_{j}g_{\beta,j}(X_{1})} \right) \right] \right\}^{T} \\ &= \left\{ \mathbb{E}_{P_{v}} \left[1 - \frac{2v_{i}g_{\beta,i}(X_{1})}{1 + v_{i}g_{\beta,i}(X_{1})} - \frac{2v_{j}g_{\beta,j}(X_{1})}{1 + v_{j}g_{\beta,j}(X_{1})} \right] \right\}^{T} \\ &= \left[1 - \int_{0}^{1} \left(\frac{2v_{i}g_{\beta,i}(x)}{1 + v_{i}g_{\beta,i}(x)} + \frac{2v_{j}g_{\beta,j}(x)}{1 + v_{j}g_{\beta,j}(x)} \right) f_{v}(x) \, \mathrm{d}x \right]^{T} \\ &= \left[1 - 2v_{i} \int_{0}^{1} g_{\beta,i}(x) \, \mathrm{d}x - 2v_{j} \int_{0}^{1} g_{\beta,j}(x) \, \mathrm{d}x \right]^{T} = 1, \end{split}$$

where the second equality uses the fact that the X_t are i.i.d. across $t \in [T]$; the third and the penultimate are due to the fact that $g_{\beta,i}$ and $g_{\beta,j}$ have disjoint support, $i \neq j$; and the last is by the fact that $\int g_{\beta,l}(x) dx = 0$ for all $l \in [k]$. Hence, we have that $\mathbb{E}_{P_v}(\varphi_{v,i}\varphi_{v,j}) = 1$ for all $i \neq j$.

We then consider that

$$\begin{split} \mathbb{E}_{P_{v}}(\varphi_{v,i}^{2}) &= \prod_{t=1}^{T} \mathbb{E}_{P_{v}} \left[\left(1 - \frac{2v_{i}g_{\beta,i}(X_{t})}{1 + v_{i}g_{\beta,i}(X_{t})} \right)^{2} \right] = \left\{ \mathbb{E}_{P_{v}} \left[\left(1 - \frac{2v_{i}g_{\beta,i}(X_{1})}{1 + v_{i}g_{\beta,i}(X_{1})} \right)^{2} \right] \right\}^{T} \\ &= \left\{ \mathbb{E}_{P_{v}} \left[1 - \frac{4v_{i}g_{\beta,i}(X_{1})}{1 + v_{i}g_{\beta,i}(X_{1})} + \left(\frac{2v_{i}g_{\beta,i}(X_{1})}{1 + v_{i}g_{\beta,i}(X_{1})} \right)^{2} \right] \right\}^{T}. \end{split}$$

We have that

$$\mathbb{E}_{P_v}\left[\frac{4v_ig_{\beta,i}(X_1)}{1+v_ig_{\beta,i}(X_1)}\right] = 0$$

due to the fact that $g_{\beta,i}$ and $g_{\beta,j}$ have disjoint support, $i \neq j$. It also holds that

$$\mathbb{E}_{P_{v}}\left[\frac{4g_{\beta,i}(X_{1})^{2}}{\{1+v_{i}g_{\beta,i}(X_{1})\}^{2}}\right] = \int_{0}^{1} \frac{4\{g_{\beta,i}(x)\}^{2}}{\{1+v_{i}g_{\beta,i}(x)\}^{2}} f_{v}(x) \,\mathrm{d}x = \int_{\frac{i-1}{k}}^{\frac{i}{k}} \frac{4\{g_{\beta,i}(x)\}^{2}}{1+v_{i}g_{\beta,i}(x)} \,\mathrm{d}x, \\ \leq \frac{2r^{2}\pi^{2\beta}}{k^{2\beta+1}} \int_{0}^{1} \{g_{\beta}(y)\}^{2} \,\mathrm{d}y \leq \frac{2r^{2}\pi^{2\beta}}{k^{2\beta+1}},$$

where in the first inequality use the substitution y = kx - (i-1) and use (127) to bound the denominator away from 0 and in the last inequality we use the fact that $|g_{\beta}(x)| \leq 1$ for all $x \in [0, 1]$. Hence, we have the upper bound

$$\mathbb{E}_{P_v}(\varphi_{v,i}^2) \le \left[1 + \frac{2r^2 \pi^{2\beta}}{k^{2\beta+1}}\right]^T,$$

which yields $\eta^2 = [1 + 2r^2 \pi^{2\beta} / k^{2\beta+1}]^T - 1$. We have now verified Assumption 2.

Step 3: obtaining lower bounds. We set

$$k = (168\pi^{2\beta})^{1/(2\beta+1)} (nT\alpha^2)^{\frac{1}{2\beta+2}}.$$

We then have, setting r = 1, that

$$\eta^{2} = \left(1 + \frac{2\pi^{2\beta}}{k^{2\beta+1}}\right)^{T} - 1 \le \frac{4\pi^{2\beta}T}{k^{2\beta+1}},\tag{128}$$

where the inequality holds under the condition

$$\frac{2\pi^{2\beta}}{k^{2\beta+1}} \le \frac{1}{T}, \quad \text{or, equivalently,} \quad T \le 84(n\alpha^2)^{2\beta+1}.$$
(129)

As Assumptions 1 and 2 hold, we have, for T satisfying (129), that

$$\mathcal{R}_{n,T,\alpha}(\theta(\mathcal{P}'_d), \|\cdot\|_2^2) \gtrsim \frac{c'_{\beta} r^2 \pi^{2\beta}}{4k^{2\beta}} \left(1 - \sqrt{\frac{21}{k} n \alpha^2 \eta^2} \right) \gtrsim \frac{c''_{\beta}}{(nT\alpha^2)^{2\beta/(2\beta+2)}} \left(1 - \sqrt{\frac{84\pi^{2\beta}}{(168\pi^{2\beta})^{2\beta+2/(2\beta+1)}}} \right),$$

$$\gtrsim c''_{\beta} (nT\alpha^2)^{-2\beta/(2\beta+2)},$$

where c''_{β} and c'''_{β} are constants depending only on β ; the first inequality is from (116) and the fact that the distributions are $c'_{\beta}r^2\pi^{2\beta}/(4k^{2\beta+1})$ -separated; the second from (128) and the value of k; and the last as C_{β} can be taken sufficiently large.

We observe that this restriction on the range of values of T is not problematic as when $T > (n\alpha^2)^{2\beta+1}$, we have $(n\alpha^2)^{-2\beta} > (nT\alpha^2)^{-2\beta/(2\beta+2)}$ and so the infinite-T lower bound in (23) (for which the required assumptions are satisfied) dominates in this case, and taking the maximum of the two lower bounds completes the proof.

D Auxiliary technical details

Proof of Lemma 12. For an estimator $\hat{\theta}$, we define \hat{v} to be any element of the set $\arg\min_{v\in\mathcal{V}}\rho(\theta(P_v),\hat{\theta})$. For any $v\in\mathcal{V}$, letting $\theta_v=\theta(P_v)$, it holds that

$$2\rho(\hat{\theta}, \theta_v) \ge \rho(\hat{\theta}, \theta_v) + \rho(\hat{\theta}, \theta_{\hat{v}}) \ge \rho(\theta_v, \theta_{\hat{v}}), \tag{130}$$

where the first inequality is by the definition of \hat{v} and the second by the triangle inequality for the metric ρ .

For $\mathcal{V} = \{-1, 1\}^k$ the hypercube which parameterises a family of distributions in \mathcal{P} , and for the subfamily of distributions $\mathcal{P}^* \subseteq \mathcal{P}$ under consideration, recall that $\mathcal{V}^* = \{v \in \mathcal{V} : P_v \in \mathcal{P}^*\}$ denotes the subset of \mathcal{V} which consists of v corresponding to distributions in \mathcal{P}^* .

Firstly, by (130) and fact that $\Phi(\cdot)$ is increasing, we see that

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \ge \sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \left(\frac{1}{2} \rho(\theta_{\hat{v}}, \theta(P)) \right) \right].$$
(131)

Let V be any random variable on the set \mathcal{V} and let V^* be the random variable arising from conditioning V on the event $\{V \in \mathcal{V}^*\}$. Then, as any average is less than or equal to a supremum, we have

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \left(\frac{1}{2} \rho(\theta_{\hat{v}}, \theta(P)) \right) \right] \ge \mathbb{E}_{V^*} \left[\mathbb{E}_{P_{V^*}} \left[\Phi \left(\frac{1}{2} \rho(\theta_{\hat{v}}, \theta(P_{V^*})) \right) \right] \right].$$

Writing $\theta(P_V) = \theta_V$ and using the law of total expectation, we also have

$$\mathbb{E}_{V}\left[\mathbb{E}_{P_{V}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V}))\right)\right]\right] = \mathbb{E}_{V}\left[\mathbb{E}_{P_{V}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V}))\right)\right] \mid V \in \mathcal{V}^{*}\right]\mathbb{P}(V \in \mathcal{V}^{*}) + \mathbb{E}_{V}\left[\mathbb{E}_{P_{V}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V}))\right)\right] \mid V \notin \mathcal{V}^{*}\right]\mathbb{P}(V \notin \mathcal{V}^{*}).$$

Trivially bounding the first probability term by 1 and bounding the risk in the second line using the worst case under the separation condition (122), we have

$$\mathbb{E}_{V}\left[\mathbb{E}_{P_{V}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V}))\right)\right]\right] \leq \mathbb{E}_{V}\left[\mathbb{E}_{P_{V}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V}))\right)\right] \mid V \in \mathcal{V}^{*}\right] + 2k\varrho\mathbb{P}(V \notin \mathcal{V}^{*}),\\ = \mathbb{E}_{V^{*}}\left[\mathbb{E}_{P_{V^{*}}}\left[\Phi\left(\frac{1}{2}\rho(\theta_{\hat{v}},\theta(P_{V^{*}}))\right)\right]\right] + 2k\varrho\mathbb{P}(V \notin \mathcal{V}^{*}).$$

Hence, we have that

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \ge \mathbb{E}_V \left[\mathbb{E}_{P_V} \left[\Phi \circ \rho(\hat{\theta}, \theta(P_V)) \right] \right] - 2k \varrho \mathbb{P}(V \notin \mathcal{V}^*),$$
(132)

which completes the proof.

Proof of Corollary 13. Given the random variable V, we denote for $j \in [k]$ the mixture distributions

$$P_{+j} = \sum_{v \in \mathcal{V}: v_j = 1} \mathbb{P}(V = v | V_j = 1) P_v \text{ and } P_{-j} = \sum_{v \in \mathcal{V}: v_j = -1} \mathbb{P}(V = v | V_j = -1) P_v.$$

We then have that

$$\begin{split} \mathbb{E}_{V} \left[\mathbb{E}_{P_{V}} \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \right] &\geq \mathbb{E}_{V} \left\{ \mathbb{E}_{P_{V}} \left[2\varrho \sum_{j=1}^{k} \mathbb{1}\{\hat{v}_{j} \neq V_{j}\} \right] \right\} \\ &= 2\varrho \sum_{j=1}^{k} \mathbb{E}_{V} \left[\mathbb{E}_{P_{V}} \left[\mathbb{1}\{\hat{v}_{j} \neq 1\} \mid V_{j} = 1\right] \right] \mathbb{P}(V_{j} = 1) \\ &+ 2\varrho \sum_{j=1}^{k} \mathbb{E}_{V} \left[\mathbb{E}_{P_{V}} \left[\mathbb{1}\{\hat{v}_{j} \neq -1\} \mid V_{j} = -1\right] \right] \mathbb{P}(V_{j} = -1) \\ &\geq 2\varrho \sum_{j=1}^{k} \left\{ P_{+j} \left(\hat{v}_{j} \neq 1 \right) \tau + P_{-j} \left(\hat{v}_{j} \neq -1 \right) \left(1 - \tau \right) \right\} \\ &\geq 2\varrho \tau \sum_{j=1}^{k} \left\{ P_{+j} \left(\hat{v}_{j} \neq 1 \right) + P_{-j} \left(\hat{v}_{j} \neq -1 \right) \right\} \\ &\geq 2\varrho \tau \sum_{j=1}^{k} \left\{ 1 - D_{\mathrm{TV}}(P_{+j}, P_{-j}) \right\}, \end{split}$$

where the first inequality is by (122) and (131), and in the penultimate we used the fact that $\tau \leq 1/2$ implies $1 - \tau \geq \tau$. Note that when $\tau = 1/2$, we recover the usual proof of Assouad's method.

Plugging in the above, and the assuming that $\mathbb{P}(V \notin \mathcal{V}^*) \leq c\tau$, into (132), we obtain that

$$\sup_{P \in \mathcal{P}^*} \mathbb{E}_P \left[\Phi \circ \rho(\hat{\theta}, \theta(P)) \right] \ge 2\varrho \tau \left[\sum_{j=1}^k \left\{ 1 - D_{\mathrm{TV}}(P_{+j}, P_{-j}) \right\} - ck \right].$$

Lemma 14. For the event A defined in (98), it holds that $\mathbb{P}(A^c) \leq 1/T^*$. Proof of Lemma 14. Recall the definition of $\hat{\omega}_j^{(i)}$ in (40). We denote

$$U_{i} = \sum_{j=1}^{d} r_{i,j} \hat{\omega}_{j}^{(i)}, \qquad \bar{U}_{i} = \Pi_{[-\eta,\eta]}(U_{i}),$$

noting that with this notation, we have that I_j as in (97) satisfies $I_j = (n/2)^{-1} \sum_{i=1}^{n/2} r_{i,j} \{ \bar{U}_i + (2\eta/\alpha)\ell_i \}$, and that $\bar{U}_i + (2\eta/\alpha)\ell_i$ is an α -LDP view of U_i . We also recall that the Rademacher random variables $\{r_{i,j}\}_{i \in [n/2], j \in [d]}$ are mutually independent and independent of the collection $\{\hat{\omega}_j^{(i)}\}_{i \in [n/2], j \in [d]}$. We now focus on I_j in the two cases where $j \in S_0$ and $j \in S_1$.

Case 1. We start by considering the expectation of $r_{i,j}\overline{U}_i$ for $j \in S_0$. Define the event $B_{i,j} = \{\hat{\omega}_j^{(i)} = 0\}$ where by the construction (40), we have that

$$\mathbb{P}(B_{i,j}^{c}) = \mathbb{P}(\hat{\omega}_{j}^{(i)} = 1) = \mathbb{P}\left(\left|\frac{1}{T^{*}}\sum_{t=1}^{T^{*}} X_{t,j}^{(i)}\right| \ge \varepsilon\right) \le 2e^{-\frac{T^{*}\varepsilon^{2}}{2}} \le \frac{2}{dnT^{*}\alpha^{2}} \le 1/5,$$
(133)

where the first inequality is by Hoeffding's inequality; the second by the definition of ε as in (39); and the last by the assumption $n\alpha^2 \geq \tilde{C}s \log(ed)$ for some sufficiently large \tilde{C} .

We also have that

$$\mathbb{E}[r_{i,j}\bar{U}_i] = \mathbb{E}[r_{i,j}\bar{U}_i\mathbb{1}(B^c_{i,j})], \qquad (134)$$

as on $B_{i,j}$ it holds that $r_{i,j}$ and \overline{U}_i are independent and so the expectation of their product is zero. Hence, we have that

$$\mathbb{E}[r_{i,j}\bar{U}_{i}\mathbb{1}(B_{i,j}^{c})] = \mathbb{E}[r_{i,j}(\bar{U}_{i} - U_{i})\mathbb{1}(B_{i,j}^{c})] + \mathbb{E}[r_{i,j}U_{i}\mathbb{1}(B_{i,j}^{c})]$$

$$= \mathbb{E}[r_{i,j}(\bar{U}_{i} - U_{i})\mathbb{1}(B_{i,j}^{c})] + \mathbb{E}\left[\left(1 + \sum_{k \neq j} (r_{i,j}r_{i,k}\hat{\omega}_{k}^{(i)})\right)\mathbb{1}(B_{i,j}^{c})\right]$$

$$= \mathbb{E}[r_{i,j}(\bar{U}_{i} - U_{i})\mathbb{1}(B_{i,j}^{c})] + \mathbb{P}(B_{i,j}^{c}), \qquad (135)$$

where the final line comes from the fact that $\mathbb{E}(r_{i,j}r_{i,k}) = 0$ for $j \neq k$. Then, writing $\tilde{U} = U_1 - r_{1,j}$, we have that

$$\mathbb{E}[r_{1,j}(U_1 - U_1)\mathbb{1}(B_{1,j}^c)] = \mathbb{E}[r_{1,j}(\eta - U_1)\mathbb{1}(\{U_1 \ge \eta\} \cap B_{1,j}^c)] + \mathbb{E}[r_{1,j}(-\eta - U_1)\mathbb{1}(\{U_1 \le -\eta\} \cap B_{1,j}^c)] = 2\mathbb{E}[r_{1,j}(\eta - U_1)\mathbb{1}(\{U_1 \ge \eta\} \cap B_{1,j}^c)] = \mathbb{E}[(\eta - 1 - \tilde{U})\mathbb{1}(\{\tilde{U} \ge \eta - 1\} \cap B_{1,j}^c)] - \mathbb{E}[(\eta + 1 - \tilde{U})\mathbb{1}(\{\tilde{U} \ge \eta + 1\} \cap B_{1,j}^c)] = -2\mathbb{P}(\{\tilde{U} \ge \eta + 1\} \cap B_{1,j}^c) + \mathbb{E}[(\eta - 1 - \tilde{U})\mathbb{1}(\{\eta - 1 \le \tilde{U} < \eta + 1\} \cap B_{1,j}^c)] \le 0.$$
(136)

Combining (133), (134), (135) and (136) we obtain

$$\mathbb{E}[r_{i,j}\bar{U}_i] \le \mathbb{P}(B_{i,j}^c) \le 1/5.$$
(137)

We also note for any fixed $j \in [d]$ that $r_{i,j}\overline{U}_i \in [-\eta,\eta]$ for $i \in [n/2]$ and that $\{r_{i,j}\overline{U}_i\}_{i\in[n/2]}$ are mutually independent. Hence, by Hoeffding's inequality, we have for x > 0 that

$$\mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}r_{i,j}\bar{U}_i - \mathbb{E}[r_{i,j}\bar{U}_i] \ge x\right) \le e^{-nx^2/(4\eta^2)}.$$
(138)

This gives that

$$\mathbb{P}(I_{j} \ge 1/2) = \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2} r_{i,j}(\bar{U}_{i} + (2\eta/\alpha)\ell_{i}) \ge 1/2\right) \\
= \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2} r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] + (2\eta/\alpha)\ell_{i} \ge 1/2 - \mathbb{E}[r_{1,j}\bar{U}_{1}]\right) \\
\leq \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2} r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] + (2\eta/\alpha)\ell_{i} \ge 3/10\right) \\
\leq \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2} r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] \ge 3/20\right) + \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2} (2\eta/\alpha)\ell_{i} \ge 3/20\right) \\
\leq \exp\left(-\frac{9n}{1600\eta^{2}}\right) + \exp\left(-\frac{9n\alpha^{2}}{25600\eta^{2}}\right) \le 2\exp\left(-\frac{9n\alpha^{2}}{25600\eta^{2}}\right),$$
(139)

where the second equality uses the independence and symmetry of $r_{i,j}$ and ℓ_i for any fixed *i* and *j*; the second equality uses (137); and in the third inequality the first term is via (138), and the second term by e.g. Wainwright (2019, Equation 2.18) where it is easy to verify that the Laplacian random variables are sub-Exponential with parameters (2, 2), (e.g. Wainwright, 2019, Definition 2.7).

Hence, by (139) and the union bound, we have that

$$\mathbb{P}\left(\bigcup_{j\in S_0} \{I_j \ge 1/2\}\right) \le 2d \exp\left(-\frac{9n\alpha^2}{25600\eta^2}\right),\tag{140}$$

where we use the fact that $|S_0| \leq d$.

Case 2. We start by considering the expectation of $r_{i,j}\overline{U}_i$ for $j \in S_1$. Define the event $D_{i,j} = \{\hat{\omega}_j^{(i)} = 1\}$ where by the construction (40), we have that

$$\mathbb{P}(D_{i,j}^c) = \mathbb{P}(\hat{\omega}_j^{(i)} = 0) = \mathbb{P}\left(\left|\frac{1}{T^*} \sum_{t=1}^T X_{t,j}^{(i)}\right| < \varepsilon\right)$$
$$\leq \mathbb{P}\left(\left|\frac{1}{T^*} \sum_{t=1}^T (X_{t,j}^{(i)} - \theta_j)\right| \ge \varepsilon\right) \le 2e^{-\frac{T^*\varepsilon^2}{2}} \le \frac{2}{nT^*\alpha^2} \le 1/10, \quad (141)$$

where the first inequality uses the fact that for $j \in S_1$, $|\theta_j| > 2\varepsilon$; the second inequality is by Hoeffding's inequality; the third by the definition of ε as in (39); and the last by the assumption $n\alpha^2 \geq \tilde{C}s \log(ed)$ for some sufficiently large \tilde{C} . We then have

$$\mathbb{E}[r_{i,j}\bar{U}_i] = \mathbb{E}[r_{i,j}\bar{U}_i\mathbb{1}(D_{i,j})]$$
(142)

as on $D_{i,j}^c$ it holds that $r_{i,j}$ and \bar{U}_i are independent and so the expectation of their product is zero. Hence, we have that

$$\mathbb{E}[r_{i,j}\bar{U}_i\mathbb{1}(D_{i,j})] = \mathbb{E}[r_{i,j}(\bar{U}_i - U_i)\mathbb{1}(D_{i,j})] + \mathbb{P}(D_{i,j})$$
(143)

by the same argument as (135). Then, writing $\tilde{U} = U_1 - r_{1,j}$, we have that

$$\begin{split} \mathbb{E}[r_{1,j}(\bar{U}_1 - U_1)\mathbb{1}(D_{1,j})] \\ &= -2\mathbb{P}(\{\tilde{U} \ge \eta + 1\} \cap D_{1,j}) + \mathbb{E}[(\eta - 1 - \tilde{U})\mathbb{1}(\{\eta - 1 \le \tilde{U} < \eta + 1\} \cap D_{1,j})] \\ &\ge -2\mathbb{P}(\{\tilde{U} \ge \eta + 1\} \cap D_{1,j}) - 2\mathbb{P}(\{\eta - 1 \le \tilde{U} < \eta + 1\} \cap D_{1,j}) \end{split}$$

$$= -2\mathbb{P}(\{\tilde{U} \ge \eta - 1\} \cap D_{1,j}) \ge -\frac{2}{(\eta - 1)^2} \mathbb{E}\left[\left(\sum_{k \ne j} r_{1,k}\hat{\omega}_k^{(1)}\right)^2\right]$$
$$\ge -\frac{2}{(\eta - 1)^2} \left(\sum_{k \in S_0} \mathbb{P}(\hat{\omega}_k^{(1)} = 1) + \sum_{k \in S_1 \cup S_2} \mathbb{P}(\hat{\omega}_k^{(1)} = 1)\right)$$
$$\ge -\frac{2}{(\eta - 1)^2} \left(\frac{2}{nT\alpha^2} + s\right) =: \zeta > -1/10, \tag{144}$$

where first equality is by the same calculations leading to (136); the second inequality is by Markov's inequality; the fourth inequality is by the same argument as (133) and the bounds $|S_0| \leq d$ and $|S_1 \cup S_2| \leq s$; and the final inequality is by the value of η and the assumption $n\alpha^2 \geq \tilde{C}s \log(ed)$ for some sufficiently large \tilde{C} . Combining (141), (142) (143) and the value ζ in (144), we obtain

$$\mathbb{E}[r_{i,j}\bar{U}_i\mathbb{1}(D_{i,j})] \ge \mathbb{P}(D_{i,j}) + \zeta > 1 - 1/10 - 1/10 \ge 4/5.$$
(145)

We then calculate

$$\mathbb{P}(I_{j} < 1/2) = \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}r_{i,j}(\bar{U}_{i} + (2\eta/\alpha)\ell_{i}) < 1/2\right) \\
= \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] + (2\eta/\alpha)\ell_{i} < 1/2 - \mathbb{E}[r_{1,j}\bar{U}_{1}]\right) \\
\leq \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] + (2\eta/\alpha)\ell_{i} < -3/10\right) \\
\leq \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}r_{i,j}\bar{U}_{i} - \mathbb{E}[r_{i,j}\bar{U}_{i}] < -3/20\right) + \mathbb{P}\left(\frac{1}{n/2}\sum_{i=1}^{n/2}(2\eta/\alpha)\ell_{i} < -3/20\right) \\
\leq \exp\left(-\frac{9n}{1600\eta^{2}}\right) + \exp\left(-\frac{9n\alpha^{2}}{25600\eta^{2}}\right) \leq 2\exp\left(-\frac{9n\alpha^{2}}{25600\eta^{2}}\right), \quad (146)$$

where the second equality uses the independence and symmetry of $r_{i,j}$ and ℓ_i for fixed *i* and *j*; the first inequality uses (145); and in the third inequality the first term is via (138) noting Hoeffding's inequality still holds for the negative $-r_{i,j}\bar{U}_i$, and the second term by e.g. Wainwright (2019, Equation 2.18) where it is easy to verify that the Laplacian random variables are sub-Exponential with parameters (2, 2), (e.g. Wainwright, 2019, Definition 2.7).

Hence, by (146) and the union bound, we have that

$$\mathbb{P}\left(\bigcup_{j\in S_1} \left\{I_j < 1/2\right\}\right) \le 2s \exp\left(-\frac{9n\alpha^2}{25600\eta^2}\right),\tag{147}$$

where we use the fact that $|S_1| \leq s$ by the sparsity assumption.

Combining (140) and (147) gives

$$\mathbb{P}(A^c) = \mathbb{P}\left(\left\{\bigcup_{j\in S_0} \{I_j \ge 1/2\}\right\} \cup \left\{\bigcup_{j\in S_1} \{I_j < 1/2\}\right\}\right)$$
$$\leq 4d \exp\left(-\frac{9n\alpha^2}{25600\eta^2}\right) = 4d \exp\left(-\frac{cn\alpha^2}{s}\right),$$

for c > 0 an absolute constant. Hence, we have that

$$\mathbb{P}(A^c) \le 4d \exp\left(-\frac{cn\alpha^2}{s}\right) \le e^{-n\alpha^2/(Ks)} = 1/T^*,$$

where the second inequality comes from the fact that $n\alpha^2 \ge \tilde{C}s \log(ed)$ for a sufficiently large constant \tilde{C} and taking K sufficiently large, and the equality comes from the value of T^* . Hence, the proof is completed.

Lemma 15 (Corollary 7 in Jin et al. 2019). Let X_i for $i \in [n]$ be a collection of i.i.d. mean-zero random variables taking values in \mathbb{R}^d such that, for some $\sigma^2 > 0$ and for all $i \in [n]$ and any $\varepsilon > 0$,

$$\mathbb{P}(\|X_i\|_2 \ge \varepsilon) \le 2e^{-\varepsilon^2/(2\sigma^2)}.$$

Then, writing $\overline{X} = n^{-1} \sum_{i=1}^{n} X_i$, there exists an absolute constant c > 0 such that for any $\gamma \in (0, 1)$,

$$\|\overline{X}\|_2 \le c \left\{ \frac{\sigma^2 \log(2d/\gamma)}{n} \right\}^{1/2}$$

holds with probability at least $1 - \gamma$. We equivalently have that

$$\mathbb{P}(\|\overline{X}\|_2 \ge \varepsilon) \le 2de^{-n\varepsilon^2/(c'\sigma^2)}.$$

where c' > 0 is some absolute constant.

Lemma 16. For $d \in \mathbb{N}$, let $X \in \{0,1\}^d$ be a binary vector with at most $m \in \mathbb{N}$ non-zero entries, i.e. $||X||_0 \leq m$. For any x > 0, write $\pi_x := e^x/(1 + e^x)$ and let $\{U_j\}_{j \in [d]}$ be a set of i.i.d. Unif(0,1)independent of X. The privatised vector $Z \in \{0,1\}^d$ with the co-ordinates

$$Z_j = \begin{cases} X_j, & U_j \le \pi_{\alpha/(2m)}, \\ 1 - X_j, & otherwise, \end{cases} \quad j \in [d],$$
(148)

is an α -LDP view of the vector X.

Proof of Lemma 16. Let $v \in \{0,1\}^d$ be an arbitrary binary vector. For $x, y \in \{0,1\}^d$ satisfying that $||x||_0, ||y||_0 \le m$, the privatised vector given by the mechanism (148) satisfies

$$\mathbb{P}(Z=v\mid X=x) = \prod_{j=1}^{d} \pi_{\alpha/(2m)}^{\mathbb{I}\{v_j=x_j\}} (1-\pi_{\alpha/(2m)})^{\mathbb{I}\{v_j\neq x_j\}} = \pi_{\alpha/(2m)}^{\#\{j:v_j=x_j\}} (1-\pi_{\alpha/(2m)})^{\#\{j:v_j\neq x_j\}} \\ = \left(\frac{\pi_{\alpha/(2m)}}{1-\pi_{\alpha/(2m)}}\right)^{n_x} (1-\pi_{\alpha/(2m)})^d = e^{\alpha n_x/(2m)} (1-\pi_{\alpha/(2m)})^d.$$

We then have that

$$\frac{\mathbb{P}(Z=v\mid X=x)}{\mathbb{P}(Z=v\mid X=y)} = e^{(n_x - n_y)\alpha/(2m)}.$$
(149)

Since x and y are non-zero on at most m co-ordinates each, they can differ from each other on at most 2m co-ordinates which implies $|n_x - n_y| \leq 2m$. Recalling that for a set A, $\sigma(A)$ denotes the sigma-algebra generated by A, we have that (149) holds for arbitrary v and so

$$\sup_{V \in \sigma(\{0,1\}^d)} \frac{\mathbb{P}(Z \in V \mid X = x)}{\mathbb{P}(Z \in V \mid X = y)} \le e^{\alpha}.$$

which shows the α -LDP condition as in (1) is satisfied.

Lemma 17. Given $\beta \in \mathbb{N}$ and r > 0, for $\delta > 0$ sufficiently small, the covering number $N(\delta)$ of the metric space $(\mathcal{F}_{\beta,r}, \|\cdot\|_2)$ with $\mathcal{F}_{\beta,r}$ defined (20), is lower bounded as,

$$N(\delta) \ge \exp\left(c_{\beta}(r/\delta)^{1/\beta}\right),$$

for some constant $c_{\beta} > 0$ depending only on β .

Proof of Lemma 17. To construct an explicit packing, we consider the construction in Appendix C.4 to obtain the family $\{f_v\}_{v\in\mathcal{V}}$ defined in (125) which is indexed by $\mathcal{V} = \{\pm 1\}^k$ for some $k \in \mathbb{N}$.

For $v, v' \in \mathcal{V}$, it follows from (126) that

$$\|f_v - f_{v'}\|_2^2 \ge \frac{c'_{\beta} r^2 \pi^{2\beta}}{2k^{2\beta+1}} \sum_{j=1}^k \mathbb{1}\{v_j \neq v'_j\},\$$

where $c'_{\beta} > 0$ is some constant depending on β .

By the Varshamov–Gilbert Lemma (e.g. Tsybakov, 2009, Lemma 2.9), there exists a subset $\mathcal{V}' \subseteq \mathcal{V}$ such that $|\mathcal{V}'| \geq 2^{k/8}$ and $\operatorname{Hamm}(u, u') \geq k/8$ for $u, u' \in \mathcal{V}'$, where $\operatorname{Hamm}(\cdot, \cdot)$ denotes the Hamming distance. For $u, u' \in \mathcal{V}'$, we therefore have that

$$\|f_u - f_{u'}\|_2^2 \ge \frac{c'_\beta r^2 \pi^{2\beta}}{16k^{2\beta}},$$

and hence we have $2^{k/8}$ -many distributions that are $\delta = (c'_{\beta})^{1/2} r \pi^{\beta} / (16k^{\beta})$ -separated.

Rearranging to get k in terms of δ , we have

$$2^{k/8} = \exp\left(\frac{\log(2)}{8} \left(\frac{(c_{\beta}')^{1/2} r \pi^{\beta}}{16\delta}\right)^{1/\beta}\right) \ge e^{c_{\beta}''(r/\delta)^{1/\beta}},$$

for some absolute constant $c''_{\beta} > 0$. This provides a lower bound on the δ -packing number whence a lower bound on the $\delta/2$ -covering number can be obtained, completing the proof.

Lemma 18. Given r > 0 and $d \in \mathbb{N}$, for any integer $s \in [d]$ and $0 < \delta < r/4$, the covering number $N(\delta)$ of the metric space $(\{\theta \in [-r, r]^d : \|\theta\|_0 \le s\}, \|\cdot\|_2)$ is lower bounded as,

$$N(\delta) \ge \left\{\frac{d}{s \wedge (d-s)}\right\}^{s \wedge (d-s)} \left\{\frac{s(r-2\delta)^2}{20\delta^2}\right\}^{\frac{s}{2}}.$$

Proof of Lemma 18. Note that when s = d, we can use the lower bound in (57) which is larger than, and thus implies, the lower bound (64) with s set to d therein. We hence assume s < d in what follows.

Writing $\Theta = \{\theta \in [-r, r]^d : \|\theta\|_0 \leq s\}$, we proceed by considering the packing number of the space $\Theta' = \{\theta \in [-r, r]^d : \|\theta\|_0 = s\}$, noting that since $\Theta' \subseteq \Theta$, this will give a lower bound on the packing number of Θ , and hence its covering number. We now decompose Θ' into a union of $\binom{d}{s}$ -many *s*-dimensional subspaces of $[-r, r]^d$. For a given indexing vector $\iota \in \{0, 1\}^d$, we denote the subspace

$$A_{\iota} = \{ x \in [-r, r]^d : x_j = 0 \text{ for } j \in [d] \text{ such that } \iota_j = 0 \}$$

We thus have that

$$\Theta' = \bigcup_{\iota \in \{0,1\}^d : \, \|\iota\|_0 = s} A_\iota$$

For any $\iota \in \{0, 1\}^d$, defining

$$B_{\iota} = \{ x \in A_{\iota} : |x_j| > \delta \text{ for } j \in [d] \text{ such that } \iota_j = 1 \} \subseteq A_{\iota},$$

we construct

$$\Theta'' = \bigcup_{\iota \in \{0,1\}^d : \|\iota\|_0 = s} B_\iota$$

Denoting the δ -packing numbers of Θ and Θ'' as $M(\delta)$ and $M''(\delta)$ respectively, we have the key observation that $M''(\delta)$ is precisely equal to the sum of the δ -packing numbers of the B_{ι} . This is because that, for two distinct index vectors $\iota^{(1)}$ and $\iota^{(2)}$, two points $x \in B_{\iota^{(1)}}$ and $y \in B_{\iota^{(2)}}$ necessarily satisfy $||x - y||_2 > 2^{1/2}\delta$.

For each $\iota \in \{0, 1\}^d$ such that $\|\iota\|_0 = s$, define the set of *s*-sparse sign vectors as $S_{\iota} = \{\varsigma \in \{-1, 0, 1\}^d : \varsigma_j = 0 \text{ for } j \in [d] \text{ such that } \iota_j = 0\}$. For $\iota \in \{0, 1\}^d$ and $\varsigma \in S_{\iota}$, we consider

$$C_{\iota,\varsigma} = \{ x \in A_{\iota} : \varsigma_j x_j > \delta \text{ for } j \in [d] \text{ such that } \iota_j = 1 \} \subseteq B_{\iota},$$

which leads to that

$$B_{\iota} = \bigcup_{\varsigma \in S_{\iota}} C_{\iota,\varsigma}.$$

For any $\iota \in \{0,1\}^d$, $\varsigma^{(1)}, \varsigma^{(2)} \in S_{\iota}$ with $\varsigma^{(1)} \neq \varsigma^{(2)}$, $x \in C_{\iota,\varsigma^{(1)}}$ and $y \in C_{\iota,\varsigma^{(2)}}$, we have that $||x-y||_2 > 2\delta$. Hence, the δ -packing number of B_{ι} is exactly equal to the sum of the δ -packing numbers of the $C_{\iota,\varsigma}$.

It remains to lower bound the packing number of $C_{\iota,\varsigma}$ for arbitrary ι and ς . By symmetry, we can assume without loss of generality that $\iota = \varsigma = \{1, \ldots, 1, 0, \ldots, 0\}$, that is, both the index vector ι and the sign vector ς have the first s entries as 1 and the rest as 0. It then holds that $C_{\iota,\varsigma} = \{x \in [-r, r]^d : (x_1, \ldots, x_s)^\top \in [\delta, r]^s, x_j = 0, s < j \le d\}$. Restricted to the first s co-ordinates, we see that $C_{\iota,\varsigma}$ is a shifted s-dimensional ℓ_{∞} -ball of radius $(r - \delta)/2$ and hence we can bound the packing number of $C_{\iota,\varsigma}$ by treating it as an s-dimensional ℓ_{∞} -ball and considering its covering number.

We can lower bound $N_{\mathbb{B}_{\infty}(r)}(\delta)$, the δ -covering number of an s-dimensional ℓ_{∞} -ball of radius r with respect to the metric $\|\cdot\|_2$, by a standard volumetric argument. Letting $\{\theta^{(j)}, j \in [N_{\mathbb{B}_{\infty}(r)}(\delta)]\}$ denote a cover for $\mathbb{B}_{\infty}(r)$, we have that

$$\mathbb{B}_{\infty}(r) \subseteq \bigcup_{j=1}^{N_{\mathbb{B}_{\infty}(r)}(\delta)} \{\theta^{(j)} + \mathbb{B}_{2}(\delta)\},\$$

which implies that

$$\operatorname{Vol}\left(\mathbb{B}_{\infty}(r)\right) \leq \operatorname{Vol}\left(\bigcup_{j=1}^{N_{\mathbb{B}_{\infty}(r)}(\delta)} \left\{\theta^{(j)} + \mathbb{B}_{2}(\delta)\right\}\right) \leq N_{\mathbb{B}_{\infty}(r)}(\delta)\operatorname{Vol}(\mathbb{B}_{2}(\delta)),$$

giving that

$$N_{\mathbb{B}_{\infty}(r)}(\delta) \ge \frac{\operatorname{Vol}(\mathbb{B}_{\infty}(r))}{\operatorname{Vol}(\mathbb{B}_{2}(\delta))} = \left(\frac{r}{\delta}\right)^{s} \frac{2^{s} \Gamma(1+s/2)}{\pi^{s/2}} \ge \left(\frac{dr^{2}}{5\delta^{2}}\right)^{\frac{s}{2}}.$$
(150)

Writing $M_{C_{\iota,\varsigma}}(\delta)$ and $N_{C_{\iota,\varsigma}}(\delta)$ for the δ -packing and δ -covering numbers of $C_{\iota,\varsigma}$ respectively, this gives the bound

$$M_{C_{\iota,\varsigma}}(\delta) \ge N_{\mathbb{B}_{\infty}\{(r-\delta)/2\}}(\delta) \ge \left\{\frac{s(r-\delta)^2}{20\delta^2}\right\}^{\frac{s}{2}},\tag{151}$$

the first inequality follows from the fact that the packing number of $C_{\iota,\varsigma}$ is lower bounded by its covering number and that $C_{\iota,\varsigma}$ can be viewed as an s-dimensional ℓ_{∞} -ball of radius $(r - \delta)/2$, and the second inequality by (150).

Finally, writing $M_{B_{\iota}}(\delta)$ for the packing number of B_{ι} , we have

$$M''(\delta) = \sum_{\iota \in \{0,1\}^d: \, \|\iota\|_0 = s} M_{B_\iota}(\delta) = \sum_{\iota \in \{0,1\}^d: \, \|\iota\|_0 = s} \sum_{\varsigma \in S_\iota} M_{C_{\iota,\varsigma}}(\delta) \ge {d \choose s} 2^s \left\{ \frac{s(r-\delta)^2}{20\delta^2} \right\}^{\frac{1}{2}},$$

where in the inequality we use the lower bound (151) and the fact that $|\{\iota : ||\iota||_0 = s\}| = {d \choose s}$ and $|I_{\iota}| = 2^s$. This gives a lower bound on $N(\delta)$ of

$$N(\delta) \ge M(2\delta) \ge M''(2\delta) \ge \left\{\frac{d}{s \wedge (d-s)}\right\}^{s \wedge (d-s)} \left\{\frac{s(r-2\delta)^2}{20\delta^2}\right\}^{\frac{s}{2}},$$

where in the first inequality we use the fact that $N(\delta) \ge M(2\delta)$, in the second the fact that $\Theta'' \subseteq \Theta$, and the bound $\binom{d}{s} \ge (d/s)^s$ for $1 \le s \le d/2$ for the final inequality.