

Convergence of the area functional on spaces with lower Ricci bounds and applications

Alessandro Cucinotta

May 21, 2024

Abstract

The goal of this note is to prove convergence results w.r.t. the area functional on metric measure spaces with a lower Ricci curvature bound. The paper can be divided in two parts.

In the first part, we show that the heat flow provides good approximation properties for the area functional on $\text{RCD}(K, \infty)$ spaces, implying that in this setting the area formula for functions of bounded variation holds and that the area functional coincides with its relaxation. Moreover, we obtain partial regularity and uniqueness results for functions whose epigraphs are perimeter minimizing.

In the second part of the paper, we consider Ricci limit spaces (and also finite dimensional $\text{RCD}(K, N)$ spaces for some results) and we show that, thanks to the previously obtained properties, minimizers of the area functional can be approximated with minimizers along the converging sequence of manifolds. As a first application, we show that minimizers of the area functional on non-collapsed Ricci limit spaces are locally Lipschitz and satisfy a-priori gradient estimates. Secondly, we obtain a Bernstein-type result for area minimizers with sublinear growth at infinity.

Contents

1 Preliminaries	5
1.1 Properties of RCD spaces	6
1.2 Minimal sets	8
1.3 Minimal graphs on manifolds	9
2 Heat flow approximation of the area and applications in $\text{RCD}(K, \infty)$ spaces	10
3 Approximation of minimal graphs in $\text{RCD}(K, N)$	18
4 Applications to non-collapsed Ricci limits	20

Introduction

In recent years there has been a growing interest in the study of metric (measure) spaces with a notion of lower bound on the Ricci curvature. Initially, this was motivated by the interest in Ricci limits, i.e. spaces arising as Gromov-Hausdorff limits of manifolds with a uniform lower Ricci curvature bound. Later on, intrinsic formulations of such curvature bound were found, leading to the theory of RCD spaces.

An $\text{RCD}(K, N)$ space is a metric measure space where $K \in \mathbb{R}$ plays the role of a lower bound on the Ricci curvature and $N \in [1, +\infty]$ plays the role of an upper bound on the dimension. This class includes measured Gromov-Hausdorff limits of smooth manifolds of fixed dimension with uniform Ricci curvature lower bounds and finite dimensional Alexandrov spaces with sectional curvature

bounded from below. Moreover, the $\text{RCD}(K, N)$ class is closed with respect to measured Gromov-Hausdorff convergence. For this reason, in the RCD setting, the study of objects that retain their properties along converging sequences of spaces is particularly relevant. This is one of the motivations that led to an extensive study of sets of finite perimeter and perimeter minimizing sets in the non-smooth context of $\text{RCD}(K, N)$ spaces.

In the note, we study some approximation properties of the area functional on RCD spaces and Ricci limits, extending a series of results that were previously obtained for sets of finite perimeter in this context. We then derive some new applications. The first central result concerns the possibility of approximating functions via the heat flow on $\text{RCD}(K, \infty)$ spaces in a way that involves the area. This is the content of Theorem 1. Given a metric measure space (X, d, \mathbf{m}) , an open set $\Omega \subset X$ and $u \in \text{BV}(\Omega)$, we denote by $|Du|$ its total variation measure, by $|\nabla u|$ the density of the absolutely continuous part of $|Du|$ w.r.t. \mathbf{m} and by $|D^s u|$ the singular part of $|Du|$. For every measurable set $E \subset \Omega$, we then define the area of u in E as

$$A(u, E) := \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E).$$

We also define $\text{Epi}(u) := \{(x, t) \in \Omega \times \mathbb{R} : t \leq u(x)\}$ and we denote by $L^0(\Omega)$ the set of measurable functions on Ω . Given two sets $A, B \subset X$, we denote by $P(A, B)$ the perimeter of A in B . Finally, we denote the heat flow on an $\text{RCD}(K, \infty)$ space by $P_t(\cdot)$.

Theorem 1. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space and let $u \in L^\infty(X) \cap \text{BV}(X)$. Then, for every bounded open set $E \subset X$ such that $|D^s u|(\partial E) = 0$, it holds*

$$\lim_{t \rightarrow 0} A(P_t(u), E) = A(u, E).$$

As a first application of Theorem 1 we obtain that the area formula holds on $\text{RCD}(K, \infty)$ spaces, i.e. the area of a function of bounded variation coincides with the perimeter of its epigraph. This is the content of Theorem 2. Before stating the theorem, we recall that related results were proved for PI spaces in [11], where it was conjectured that the area formula for functions of bounded variation should hold in that setting. In particular, [11, Theorem 5.1] implies that the area formula holds for Sobolev functions on PI spaces where weak and relaxed gradients coincide (and finite dimensional $\text{RCD}(K, N)$ spaces fit in this framework). Such formula was then proved for BV functions on finite dimensional $\text{RCD}(K, N)$ spaces in [12], while in the next theorem we extend the result to the infinite dimensional framework of $\text{RCD}(K, \infty)$ spaces.

Theorem 2. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subset X$ be open and bounded and let $u \in L^1(\Omega)$. Then $P(\text{Epi}(u), \Omega \times \mathbb{R}) < +\infty$ if and only if $u \in \text{BV}(\Omega)$. In this case, for every Borel set $E \subset \Omega$, it holds*

$$P(\text{Epi}(u), E \times \mathbb{R}) = A(u, E).$$

Moreover, if $u \in L^0(B_{2r}(x))$ and $P(\text{Epi}(u), B_{2r}(x) \times \mathbb{R}) < +\infty$, then $u \in \text{BV}(B_r(x))$ and the previous equation holds for every Borel set $E \subset B_r(x)$.

As anticipated, the previous result complements the one proved for doubling spaces supporting a Poincaré inequality in [11], where it was shown that if (X, d, \mathbf{m}) is such a space and $u \in \text{BV}(\Omega)$, then

$$P(\text{Epi}(u), \Omega \times \mathbb{R}) \geq A(u, \Omega).$$

Note that, although $\text{RCD}(K, \infty)$ spaces are not necessarily doubling, the proof of the inequality given in [11] works with very slight changes also in our setting. On the other hand, the proof that we give of the reverse inequality is a consequence of Theorem 1 and truly relies on the curvature assumption.

A second consequence of this convergence result is that on $\text{RCD}(K, \infty)$ spaces the relaxed area functional coincides with the area functional itself. This is the content of Theorem 3.

Let $\Omega \subset X$ be an open bounded set and let $u \in \text{BV}(\Omega)$. For every open set $E \subset \Omega$ we denote by $\text{Lip}(E)$ the set of Lipschitz functions on E and given $f \in \text{Lip}(E)$ we denote by $\text{lip}(f)$ its local Lipschitz constant. The relaxed area functional is defined by

$$\tilde{A}(u, E) := \inf \left\{ \liminf_{n \rightarrow +\infty} \int_E \sqrt{1 + \text{lip}(f_n)^2} \, d\mathbf{m} \mid (f_n)_n \subset L^1(E) \cap \text{Lip}(E), \|f_n - u\|_{L^1(E)} \rightarrow 0 \right\}.$$

Theorem 3. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subset X$ be open and bounded and let $u \in \text{BV}(\Omega)$. Then, for every open set $E \subset \Omega$, it holds*

$$\tilde{A}(u, E) = A(u, E).$$

The previous theorem extends the analogous result proved in [11] for Sobolev functions. Note that in general metric measure spaces (even if they are doubling and support a Poincaré inequality) the relaxed area functional does not coincide with the area functional itself, as shown in [30]. Theorem 3, in particular, allows to apply the mild regularity results for minimizers of the relaxed area functional in metric measure spaces established in [29] to the area functional itself in the $\text{RCD}(K, N)$ setting. The last result of the first part of the note concerns a uniqueness result for perimeter minimizers (which follows through the identification between area and perimeter) and it is given in Theorem 4. In what follows, when we refer to an $\text{RCD}(K, N)$ space, we implicitly mean that $N < +\infty$.

Theorem 4. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subsetneq X$ be an open set and let $u \in L^\infty(\Omega) \cap W^{1,1}(\Omega)$ be a function whose epigraph is perimeter minimizing. If $v \in L^\infty(\Omega)$ has epigraph which is perimeter minimizing and $\{u \neq v\} \subset\subset \Omega$, then $u = v$.*

Corollary. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, N)$ space, let $\Omega \subsetneq X$ be an open set and let $u \in W^{1,1}(\Omega)$ be a function whose epigraph is perimeter minimizing. If $v \in L^1(\Omega)$ has epigraph which is perimeter minimizing and $\{u \neq v\} \subset\subset \Omega$, then $u = v$.*

In the second part of the note, we apply the previously obtained theorems to prove more results in the finite dimensional setting of $\text{RCD}(K, N)$ spaces and Ricci limit spaces. We work with functions that minimize the area (see Definition 2.6), but we could have worked with functions whose epigraphs are perimeter minimizing since the two notions are equivalent (see Proposition 2.9).

A central result of this part of the note is Theorem 5, which shows that minimizers on a non-collapsed Ricci limit space can be approximated by minimizers along the approximating sequence of manifolds. Although technical in nature, Theorem 5 is at the core of all the subsequent results of the paper.

When considering pointed measured Gromov-Hausdorff convergence of spaces, we adopt the extrinsic viewpoint, meaning that we implicitly assume all spaces to be embedded in a common metric space (Z, d_Z) where the convergence is realized. In particular, balls in this setting are to be intended w.r.t. the distance in Z (where all spaces are embedded), while we will specify from time to time which measure is to be considered. Given a metric space (X, d) , we denote by H^n the n -dimensional Hausdorff measure induced by d . When we say that a metric space (M, d) is a Riemannian manifold we mean that the distance d is the Riemannian distance induced by the metric, which does not appear in our notation. We refer to the beginning of Section 3 for more details on the notation used.

Theorem 5. *Let $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let (M_k^n, d_k, x_k) be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $H_k^n(B_1(x_k)) \geq v > 0$ converging in pointed Gromov-Hausdorff sense to (X, d, x) . Let $r > 0$ be such that $B_r(x) \subsetneq X$ and let $u \in \text{BV}(B_r(x), H^n)$ be an area minimizer.*

For every $s \in (0, r)$, there exists a sequence of area minimizers $u_k \in \text{BV}(B_s(x), H_k^n)$ such that $u_k \mathbf{1}_{B_s(x)}$ converges in L^1 to $u \mathbf{1}_{B_s(x)}$ and $A(u_k, B_s(x), H_k^n) \rightarrow A(u, B_s(x), H^n)$.

Theorem 5 is useful because it allows to pass information from minimizers of the area on smooth manifolds to minimizers on the limit space. As a first instance of this idea, we obtain Theorem 6, which extends to Ricci limit spaces the a-priori gradient estimates for minimizers of the area functional on Riemannian manifolds found in [22].

Theorem 6. *Let $r > 0$, $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $\mathbf{H}_k^n(B_r(x_k)) \geq vr^n$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . If $u \in \text{BV}(B_r(x), \mathbf{H}^n)$ is an area minimizer, then there exists $c(K, n, v) > 0$ such that, setting*

$$c_r := c e^{cr^{-1}(u(x) - \inf_{y \in B_r(x)} u(y))},$$

it holds

$$\sup_{z \in B_{r/(15c_r)}(x)} \text{lip}(u)(z) \leq c_r.$$

A consequence of the previous result is the regularity of minimizers of the area functional on non-collapsed Ricci limit spaces. Observe that minimizers of the area on generic metric measure spaces can be far from being Lipschitz, as shown in [29].

Theorem 7. *Let $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $\mathbf{H}_k^n(B_1(x_k)) \geq v > 0$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . If $\Omega \subset X$ is open and $u \in \text{BV}(\Omega, \mathbf{H}^n)$ is an area minimizer, then $u \in \text{Lip}_{loc}(\Omega)$.*

Finally, Theorems 5 and 6 can be combined, under the assumption that the space in question has maximal volume growth, to obtain a stronger gradient estimate and a Bernstein-type result for minimizers with sublinear growth at infinity. This is the content of Theorems 8 and 9.

Theorem 8. *Let $n \in \mathbb{N}$ be fixed and let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq 0$ converging in pointed Gromov-Hausdorff sense to a space (X, \mathbf{d}, x) such that*

$$\lim_{r \rightarrow +\infty} \frac{\mathbf{H}^n(B_r(x))}{r^n} \geq v > 0.$$

If $u \in \text{BV}_{loc}(X, \mathbf{H}^n)$ is an area minimizer, then there exists $c(n, v) > 0$ such that, for every $r > 0$, it holds

$$\sup_{y \in B_{r/2}(x)} \text{lip}(u)(y) \leq c \left(1 + r^{-1} \sup_{y \in B_r(x)} |u(y) - u(x)| \right)^n.$$

Theorem 9. *Let $n \in \mathbb{N}$ be fixed and let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq 0$ converging in pointed Gromov-Hausdorff sense to a space (X, \mathbf{d}, x) such that*

$$\lim_{r \rightarrow +\infty} \frac{\mathbf{H}^n(B_r(x))}{r^n} \geq v > 0.$$

If $u \in \text{BV}_{loc}(X, \mathbf{H}^n)$ is an area minimizer such that

$$\lim_{y \rightarrow +\infty} \frac{\max\{-u(y), 0\}}{\mathbf{d}(x, y)} = 0,$$

then u is constant.

The proofs of the previous two theorems rely on the ones of the analogous results on Riemannian manifolds proved in [22]. Moreover, Theorem 9 is part of a series of recent results proving Bernstein-type theorems on manifolds with a lower Ricci curvature bound (see [23, 21, 15, 16, 20, 22]) and on metric measure spaces with with an analogous synthetic bound (see [18, 17]).

Acknowledgements. I would like to thank Prof. Andrea Mondino for his guidance.

1 Preliminaries

We will work on metric measure spaces (X, d, \mathbf{m}) , where (X, d) is a separable complete metric space and \mathbf{m} is a non-negative Borel measure on X which is finite on bounded sets. Given an open set $\Omega \subset X$ we denote by $\text{Lip}(\Omega)$, $\text{Lip}_c(\Omega)$ and $\text{Lip}_b(\Omega)$ respectively the set of Lipschitz functions, Lipschitz functions with compact support and bounded Lipschitz functions on Ω . Similarly, we denote by $C(\Omega)$ the set of continuous functions on Ω and by $L^0(\Omega)$ the one of measurable functions. If $u \in L^0(\Omega)$ we also define $\text{Epi}(u) := \{(x, t) \in \Omega \times \mathbb{R} : t \leq u(x)\}$. If $f \in \text{Lip}(\Omega)$ and $x \in \Omega$ we define

$$\text{lip}(f)(x) := \limsup_{y \rightarrow x} \frac{|f(x) - f(y)|}{d(x, y)} \quad \text{and} \quad L(f) := \sup_{x, y \in \Omega} \frac{|f(x) - f(y)|}{d(x, y)}.$$

We say that Ω' is well contained in Ω , and we denote it by $\Omega' \subset\subset \Omega$, if Ω' is bounded and $d(\Omega', \partial\Omega) > 0$. When we consider the spaces $L^1_{loc}(\Omega)$ and $\text{Lip}_{loc}(\Omega)$ we implicitly mean that we are localizing w.r.t. well contained subsets of Ω ; note that these are not necessarily precompact in Ω (although they will be compact if X is a finite dimensional $\text{RCD}(K, N)$ space). We now introduce Sobolev spaces in the setting of metric measure spaces, the main references being [14], [7], [8] and [25].

Definition 1.1. Let (X, d, \mathbf{m}) be a metric measure space and let $\Omega \subset X$ be an open set. A function $f \in L^2(\Omega)$ is said to be in the Sobolev space $W^{1,2}(\Omega)$ if there exists a sequence $\{f_i\}_{i \in \mathbb{N}} \subset \text{Lip}(\Omega)$ converging to f in $L^2(\Omega)$ such that

$$\limsup_{i \rightarrow +\infty} \int_{\Omega} \text{lip}(f_i)^2 d\mathbf{m} < +\infty.$$

For every $f \in W^{1,2}(\Omega)$ one can define a function $|\nabla f| \in L^2(\Omega)$ such that for every open set $A \subset \Omega$ we have

$$\int_A |\nabla f|^2 d\mathbf{m} = \inf \left\{ \liminf_{n \rightarrow +\infty} \int_A \text{lip}(f_n)^2 d\mathbf{m} \mid (f_n)_n \subset L^2(A) \cap \text{Lip}(A), \|f_n - f\|_{L^2(A)} \rightarrow 0 \right\}.$$

The quantity in the previous expression will be called Cheeger energy and will be denoted by $\text{Ch}(f)$ while $|\nabla f|$ will be called relaxed gradient. We define $\|f\|_{W^{1,2}(\Omega)} := \|f\|_{L^2(\Omega)} + \text{Ch}(f)$. One can check that with this norm the space $W^{1,2}(\Omega)$ is Banach.

Remark 1.2. Given a metric measure space (X, d, \mathbf{m}) , an open set $\Omega \subset X$ and a function $f \in W^{1,2}(\Omega)$, there exists a sequence $f_n \in \text{Lip}(\Omega)$ such that $\text{lip}_a(f_n) \rightarrow |\nabla f|$ in $L^2(\Omega)$, where

$$\text{lip}_a(f)(x) := \lim_{r \rightarrow 0} \sup_{y, z \in B_r(x)} \frac{|f(y) - f(z)|}{d(y, z)}.$$

This fact follows since in the definition of Sobolev spaces one could replace local Lipschitz constants lip with asymptotic Lipschitz constants lip_a and still obtain the same space (see [3] and [7, Remark 5.12]).

We now introduce functions of bounded variation following [35] (see also [4]).

Definition 1.3. Let (X, d, \mathbf{m}) be a metric measure space and let $\Omega \subset X$ be an open set. A function $f \in L^1(\Omega)$ is said to be of bounded variation if there exists a sequence of $\{f_i\}_{i \in \mathbb{N}} \subset \text{Lip}(\Omega)$ converging to f in $L^1(\Omega)$ such that

$$\limsup_{i \rightarrow +\infty} \int_{\Omega} \text{lip}(f_i) d\mathbf{m} < +\infty.$$

The space of such functions is denoted by $\text{BV}(\Omega)$.

For every $f \in \text{BV}(\Omega)$ and any open set $A \subset \Omega$ we define

$$|Df|(A) = \inf \left\{ \liminf_{n \rightarrow +\infty} \int_A \text{lip}(f_n) d\mathbf{m} \mid (f_n)_n \subset L^1(A) \cap \text{Lip}(A), \|f_n - f\|_{L^1(A)} \rightarrow 0 \right\}.$$

One can check that the quantity in the previous expression is the restriction to the open subsets of Ω of a finite measure. We denote by $|\nabla f|$ the density of $|Df|$ with respect to \mathbf{m} and by $|D^s f|$ the singular part of $|Du|$. We define $\|f\|_{\mathbf{BV}(\Omega)} := \|f\|_{L^1(\Omega)} + |Df|(\Omega)$. One can check that with this norm the space $\mathbf{BV}(\Omega)$ is Banach. The function f belongs to $W^{1,1}(\Omega)$ if $f \in \mathbf{BV}(\Omega)$ and $|Df| \ll \mathbf{m}$.

Definition 1.4. Let (X, d, \mathbf{m}) be a metric measure space and let $\Omega \subset X$ be open. We define the perimeter of $E \subset \Omega$ in Ω by

$$\mathbf{P}(E, \Omega) := \inf \left\{ \liminf_{n \rightarrow +\infty} \int_{\Omega} \text{lip}(f_n) d\mathbf{m} \mid (f_n)_n \subset L^1_{loc}(\Omega) \cap \text{Lip}(\Omega), \|f_n - 1_E\|_{L^1_{loc}(\Omega)} \rightarrow 0 \right\}.$$

We recall that $\mathbf{P}(E, \cdot)$ is the restriction to open sets of a Borel measure and that if $E \subset \Omega$ has finite measure, then it has finite perimeter if and only if $1_E \in \mathbf{BV}(\Omega)$ and in this case $|D1_E| = \mathbf{P}(E, \cdot)$.

Definition 1.5. We say that a metric measure space (X, d, \mathbf{m}) is infinitesimally Hilbertian if the space $W^{1,2}(X)$ is a Hilbert space.

We will consider pointed Gromov-Hausdorff (pGH) and pointed measured Gromov-Hausdorff (pmGH) convergence of metric measure spaces, and we refer to [27] for the relevant background. We recall that in the case of a sequence of uniformly locally doubling metric measure spaces $(X_i, d_i, \mathbf{m}_i, x_i)$ (as in the case of $\text{RCD}(K, N)$ spaces), pointed measured Gromov-Hausdorff convergence to (X, d, \mathbf{m}, x) can be equivalently characterized asking for the existence of a proper metric space (Z, d_z) such that all the metric spaces (X_i, d_i) are isometrically embedded into (Z, d_z) , $x_i \rightarrow x$ and $\mathbf{m}_i \rightarrow \mathbf{m}$ weakly in duality with continuous boundedly supported functions in Z . In this case we say that the convergence is realized in the space Z (see [27]).

1.1 Properties of RCD spaces

We now recall some properties of $\text{RCD}(K, \infty)$ spaces; these were introduced in [8] (see also [6, 25]) coupling the Curvature Dimension condition $\text{CD}(K, \infty)$, previously pioneered in [38, 39] and independently in [34], with the infinitesimal Hilbertianity assumption. The next proposition is an isoperimetric inequality that follows from the corresponding Poincaré inequality, which is proved in [37].

Proposition 1.6. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space and let $E \subset B_{2r}(x)$ be a set of finite perimeter. Then we have the following isoperimetric inequality:*

$$\min\{\mathbf{m}(E \cap B_r(x)), \mathbf{m}(B_r(x) \setminus E)\} \leq 4re^{K|r^2} \mathbf{P}(E, B_{2r}(x)).$$

Given $f : X \times \mathbb{R} \rightarrow \mathbb{R}$ and $(x, t) \in X \times \mathbb{R}$ we denote by f^t and f^x respectively the restriction of f to $X \times \{t\}$ and to $\{x\} \times \mathbb{R}$. The next proposition concerns a tensorization property of infinitesimally Hilbertian spaces and can be found in [11] and [8]. We denote by d_x and \mathbf{m}_x respectively the product distance and the product measure in the space $X \times \mathbb{R}$.

Proposition 1.7. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space and let $f \in \text{Lip}(X \times \mathbb{R})$. Then we have*

$$|\nabla f|^2(x, t) = |\nabla f^x|^2(t) + |\nabla f^t|^2(x) \quad \text{for } \mathbf{m}_x\text{-a.e. } (x, t) \in X \times \mathbb{R}.$$

We now recall some properties of the heat flow in the RCD setting, referring to [6, 8] for the proofs of these results. Given an $\text{RCD}(K, \infty)$ space (X, d, \mathbf{m}) , the heat flow $P_t : L^2(X) \rightarrow L^2(X)$ is the $L^2(X)$ -gradient flow of the Cheeger energy Ch . Let $\mathcal{P}_2(X)$ denote the space of probability measures with finite second moment. A key result in the RCD theory is that the dual heat semigroup $\bar{P}_t : \mathcal{P}_2(X) \rightarrow \mathcal{P}_2(X)$ defined by

$$\int_X d\bar{P}_t \mu := \int_X P_t f d\mu \quad \forall \mu \in \mathcal{P}_2(X), \quad \forall f \in \text{Lip}_b(X),$$

is K -contractive with respect to the Wasserstein distance W_2 and for $t > 0$ maps probability measures into absolutely continuous probability measures. This allows to define a stochastically complete heat

kernel $p_t : \mathsf{X} \times \mathsf{X} \rightarrow [0, +\infty)$, so that the definition of $P_t(f)$ can be then extended to L^∞ functions by setting

$$P_t(f)(x) := \int_{\mathsf{X}} f(y)p_t(x, y) \, d\mathbf{m}(y).$$

The heat flow has good approximation properties, in particular if $f \in W^{1,2}(\mathsf{X})$, then $P_t(f) \rightarrow f$ in $W^{1,2}(\mathsf{X})$; while if $f \in L^\infty(\mathsf{X})$, then $P_t f \in \text{Lip}(\mathsf{X})$ for every $t > 0$.

In [26] it was proved that if $f \in \text{BV}(\mathsf{X})$ then

$$|DP_t(f)| \leq e^{-Kt} \bar{P}_t(|Df|).$$

Combining the previous contraction estimate with a lower semicontinuity argument we obtain that if $\Omega \subset \mathsf{X}$ is open and such that $|Df|(\partial\Omega) = 0$, then

$$\lim_{t \rightarrow 0} |DP_t(f)|(\Omega) = |Df|(\Omega).$$

We next recall two basic lemmas on measures in metric measure spaces. Note in particular that these hold without the assumption that the space is proper. Given an \mathbb{R}^m -valued measure μ on X we denote by $|\mu|$ its variation. The first part of the next lemma follows from [5, Proposition 1.23], while the second part follows repeating the argument of [5, Proposition 1.47] using the density of continuous functions in $L^1(\mathsf{X})$ when $(\mathsf{X}, d, \mathbf{m})$ is $\text{RCD}(K, \infty)$.

Lemma 1.8. *Let $(\mathsf{X}, d, \mathbf{m})$ be an $\text{RCD}(K, \infty)$ space and let $f \in L^1(\mathsf{X})^m$. Consider the \mathbb{R}^m -valued measure $\mu := f\mathbf{m}$, then for every Borel set $B \subset \mathsf{X}$*

$$|\mu|(B) = \int_B |f| \, d\mathbf{m},$$

and for every open set $A \subset \mathsf{X}$

$$|\mu|(A) = \sup_{\substack{\{a_i\}_{i=1}^m \subset C(A) \\ \sum a_i^2 \leq 1}} \sum_{i=1}^m \int_A a_i f_i \, d\mathbf{m}.$$

The next lemma follows by standard measure theoretic arguments.

Lemma 1.9. *Let $(\mathsf{X}, d, \mathbf{m})$ be an $\text{RCD}(K, \infty)$ space and let μ, ν be positive Radon measures on X . If $\mu(A) \geq \nu(A)$ for every open set $A \subset \mathsf{X}$, then $\mu(E) \geq \nu(E)$ for every Borel set.*

We now recall some properties of $\text{RCD}(K, N)$ spaces. The finite dimensional $\text{RCD}(K, N)$ condition is obtained coupling the finite dimensional Curvature Dimension condition $\text{CD}(K, N)$ with the infinitesimal Hilbertianity assumption and was formalised in [25]. For a thorough introduction to the topic we refer to the survey [1] and the references therein. Let us also mention that in the literature one can find also the (a-priori weaker) $\text{RCD}^*(K, N)$. It was proved in [24, 10], that $\text{RCD}^*(K, N)$ is equivalent to the dimensional Bochner inequality. Moreover, [13] (see also [33]) proved that $\text{RCD}^*(K, N)$ and $\text{RCD}(K, N)$ coincide. We now recall some results on $\text{RCD}(K, N)$ spaces that we will need later in the note.

The $\text{RCD}(K, N)$ condition implies that the measure is locally doubling (see [38]) and, as mentioned before, the validity of a Poincaré inequality. In particular if f is a locally Lipschitz function on an $\text{RCD}(K, N)$ space, its relaxed gradient coincides with its local Lipschitz constant $\text{lip}(f)$ (see [31, Theorem 12.5.1] after [14]).

The next theorem is taken from [19, Theorem 1.2]. Given a metric space (X, d) , we denote by H^n the n -dimensional Hausdorff measure relative to d .

Theorem 1.10. *Let $(\mathsf{X}_i, d_i, \mathsf{H}_i^n, x_i)$ be a sequence of $\text{RCD}(K, n)$ spaces such that (X_i, d_i, x_i) converges in pGH sense to a metric space (X, d, x) . Then precisely one of the following holds:*

1.

$$\limsup_{i \rightarrow +\infty} \mathbf{H}_i^n(B_1(x_i)) > 0.$$

In this case the limsup is a limit and the pGH convergence can be improved to pmGH convergence to (X, d, \mathbf{H}^n, x) .

2.

$$\lim_{i \rightarrow +\infty} \mathbf{H}_i^n(B_1(x_i)) = 0.$$

In this case the Hausdorff dimension of X is bounded above by $n - 1$.

We now recall the definition of L^1 -convergence of functions along a sequence of spaces converging in pmGH sense. As anticipated, we use the so-called extrinsic viewpoint and, when considering spaces $X_i \rightarrow X$ in pmGH sense, we work in the space (Z, d_Z) which realizes the convergence. In particular, whenever we consider a ball $B_r(x)$ in this context, unless otherwise specified, we mean the ball of radius r centered in x in the space Z . Similarly, every function $u_i : X_i \rightarrow \mathbb{R}$ will be considered, unless otherwise specified, as a function on Z (extending it to zero outside of the embedding of X_i in Z). To avoid confusion we will specify from time to time which measure is to be considered when working with objects defined on Z .

Definition 1.11. Let (X_i, d_i, m_i, x_i) be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to (Y, d, m, y) . We say that the functions $f_i \in L^1(X_i, m_i)$ converge to $f \in L^1(X, m)$ in L^1 -sense if

$$\sigma \circ f_i m_i \rightarrow \sigma \circ f m \quad \text{and} \quad \int_{X_i} |f_i| dm_i \rightarrow \int_X |f| dm,$$

where $\sigma(z) := \text{sign} \sqrt{|z|}$ and weak convergence is intended w.r.t. boundedly supported functions in (Z, d_Z) , which is the space realizing the convergence. We say that $f_i \in L^1_{loc}(X_i, m_i)$ converge in L^1_{loc} -sense to $f \in L^1_{loc}(X, m)$ if $f_i 1_{B_r(x_i)} \rightarrow f 1_{B_r(x)}$ in L^1 -sense for every $r > 0$.

The next proposition is taken from [2, Proposition 3.3].

Proposition 1.12. Let (X_i, d_i, m_i, x_i) be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to (Y, d, m, y) . Let $f_i \in \text{BV}(X_i, m_i)$ with $\text{supp}(f_i) \subset B_r(x) \subset Z$, where (Z, d_Z) is the space realizing the convergence, be functions with

$$\sup_{i \in \mathbb{N}} \left(\|f_i\|_{\text{BV}(X_i, m_i)} + \|f_i\|_{L^\infty(X_i, m_i)} \right) < +\infty.$$

Then there exists a (non relabeled) subsequence and $f \in \text{BV}(X, m)$ with $\text{supp} f \subset \bar{B}_r(x)$ such that $f_i \rightarrow f$ in L^1 .

The next proposition is taken from [2, Proposition 3.6].

Proposition 1.13. Let (X_i, d_i, m_i, x_i) be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to (X, d, m, x) . If $u \in \text{BV}(X, m)$, and $u_i \in \text{BV}(X_i, m_i)$ is a sequence such that $u_i \rightarrow u$ in L^1 and $\sup_i \|u_i\|_{L^\infty(X_i, m_i)} < +\infty$, then for every open set $A \subset Z$, where (Z, d_Z) is the metric space realising the convergence, we have

$$|Du|(A) \leq \liminf_i |Du_i|(A).$$

1.2 Minimal sets

We now turn our attention to minimal sets. We only recall the properties that will be used later in the note.

Definition 1.14. Let (X, d, m) be a metric measure space and let $\Omega \subset X$ be an open set. Let $E \subset \Omega$ be a set of locally finite perimeter. We say that E is perimeter minimizing in Ω if for every $x \in \Omega$, $r > 0$ and $F \subset \Omega$ such that $F \Delta E \subset \subset B_r(x) \cap \Omega$ we have that $P(E, B_r(x) \cap \Omega) \leq P(F, B_r(x) \cap \Omega)$. If we say that E is perimeter minimizing we implicitly mean that $\Omega = X$.

The next theorem comes from [32, Theorem 4.2 and Lemma 5.1].

Theorem 1.15. *Let (X, d, m) be an $\text{RCD}(K, N)$ space and let $\Omega \subset X$ be an open set. There exist $C, \gamma_0 > 0$ depending only on K and N such that the following hold. If $E \subset X$ is a set minimizing the perimeter in $\Omega \subset X$, then, up to modifying E on an m -negligible set, for any $x \in \partial E$ and $r > 0$ such that $B_{2r}(x) \subset \Omega$ we have*

$$\frac{m(E \cap B_r(x))}{m(B_r(x))} > \gamma_0, \quad \frac{m(B_r(x) \setminus E)}{m(B_r(x))} > \gamma_0$$

and

$$\frac{m(B_r(x))}{Cr} \leq P(E, B_r(x)) \leq \frac{Cm(B_r(x))}{r}.$$

The next proposition is taken from [36, Theorem 2.43].

Proposition 1.16. *Let (X_i, d_i, m_i, x_i) be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to (Y, d, m, y) . Let $E_i \subset X_i$ be a sequence of Borel sets converging in L^1_{loc} -sense to $E \subset Y$. Assume that each E_i is perimeter minimizing in $B_{r_i}(x_i)$ and that $r_i \uparrow +\infty$. Then E is perimeter minimizing and in the metric space realizing the convergence we have that $\partial E_i \rightarrow \partial E$ in Kuratowski sense.*

We recall that given a space (X, d, m) we denote by d_X and m_X respectively the product distance and the product measure on $X \times \mathbb{R}$. Moreover, given a function $u \in \text{Lip}_{loc}(X)$ we will use the following notation for balls on the graph of u (these are actually intersections of balls in the product with the graph of u):

$$B_r^g(x, t) := B_r^X(x, t) \cap \text{graph}(u) \subset X \times \mathbb{R}.$$

Similarly, given a point $x \in X$, we will denote by \bar{x} its projection on the graph of u , i.e. $\bar{x} := (x, u(x))$. Whenever we evaluate a function $f : X \rightarrow \mathbb{R}$ on the graph of u we implicitly mean that we are referring to $f \circ \pi$, where $\pi : \text{graph}(u) \rightarrow X$ is the standard projection.

The next result follows from [17, Theorem 5.8].

Theorem 1.17. *There exists a constant $C > 0$ depending only on N such that if (X, d, m) is an $\text{RCD}(0, N)$ space and $u \in \text{Lip}_{loc}(B_{3R}(p))$ is a positive function whose epigraph minimizes the perimeter in $B_{3R}(p) \times \mathbb{R}$, setting $\bar{p} := (p, u(p))$, we have*

$$\sup_{B_{R/2}^g(\bar{p})} u \leq C \inf_{B_{R/2}^g(\bar{p})} u.$$

1.3 Minimal graphs on manifolds

In this section we recall some properties of minimal graphs on Riemannian manifolds. Given a manifold (M, g) and a solution of the minimal surface equation $u \in C^\infty(\Omega)$ on an open subset $\Omega \subset M$, we denote by Δ_g the laplacian on the graph of u .

In the next proposition and more in general in the note we will consider the laplacian Δ_g of a function f defined on $\Omega \subset M$. When doing so, we implicitly mean that we are applying the laplacian to the function $f \circ \pi$, where $\pi : \text{graph}(u) \rightarrow \Omega$ is the standard projection. The next proposition is the so-called Jacobi equation for minimal graphs and it follows by taking the second variation of the area of a minimal graph.

Proposition 1.18. *Let (M, g) be a Riemannian manifold, let $\Omega \subset M$ be open and let $u \in C^\infty(\Omega)$ be a solution of the minimal surface equation. Then, setting $v_u := \sqrt{1 + |\nabla u|^2}$ and denoting by A the second fundamental form of $\text{graph}(u)$ in $\Omega \times \mathbb{R}$, we have*

$$\Delta_g v_u^{-1} = -\frac{|A|^2}{v_u} - \frac{\text{Ric}(\nabla u, \nabla u)}{v_u^{3/2}},$$

or, equivalently,

$$\Delta_g \log v_u = |A|^2 + \frac{\text{Ric}(\nabla u, \nabla u)}{v_u^2} + |\nabla \log v_u|^2.$$

The next proposition is taken from [22, Lemma 5.7].

Proposition 1.19. *Let (M^n, g) be a manifold with $\text{Ric}_M \geq K$ and $H^n(B_1(x)) \geq v > 0$. Let $u \in C^\infty(B_{2r}(x))$ be a solution of the minimal surface equation.*

- *If $\phi \in C^\infty(B_r^g(\bar{x}))$ is a non-negative sub-harmonic function, then*

$$\sup_{B_{r/2}^g(\bar{x})} \phi \leq C(K, n, v) \int_{B_r^g(\bar{x})} \phi dH^n.$$

- *If $f \in C^\infty(B_{2r}^g(\bar{x}))$ is a non-negative super-harmonic function, then*

$$\inf_{B_r^g(\bar{x})} f \geq C'(K, n, v) \int_{B_r^g(\bar{x})} f dH^n.$$

The next theorem is the a-priori gradient estimate for minimal graphs and it is taken from [22, Theorem 6.1]. Although stated in a slightly weaker form, inspecting the proof of [22, Theorem 6.1], one can check that this is actually the version of the gradient estimate that is proved in the aforementioned reference.

Theorem 1.20. *Let $r > 0$ and let (M^n, g) be a complete non-compact Riemannian manifold with*

$$\text{Ric}_M \geq -(n-1)k^2r^{-2}$$

on $B_r(p) \subset M$ for some $k \geq 0$. Suppose that $H^n(B_r(p)) \geq vr^n$ for some constant $v > 0$ and let $u \in C^\infty(B_r(p))$ be a solution of the minimal surface equation. Then

$$\sup_{x \in B_{r/6}(\bar{p})} |\nabla u(x)| \leq ce^{cr^{-1}(u(p) - \inf_{x \in B_r(p)} u(x))}$$

where $c > 0$ depends only on k, n and v .

2 Heat flow approximation of the area and applications in $\text{RCD}(K, \infty)$ spaces

In this section, we prove Theorems 1 to 4. We implicitly assume that we are working on a fixed $\text{RCD}(K, \infty)$ space (X, d, \mathbf{m}) and that $\Omega \subset X$ is a fixed bounded open set. We denote by d_\times and \mathbf{m}_\times respectively the product distance and the product measure on $X \times \mathbb{R}$. The next two propositions contain the inequality in the area formula that was proved in [11] for PI spaces. Similar proofs work in our setting and we repeat them (with the few due modifications) for the sake of completeness as they will be needed to prove Theorems 1 and 2.

Definition 2.1. Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subset X$ be an open set and let $u \in \text{BV}(\Omega)$. For every measurable $E \subset \Omega$ we define the area of u on E as

$$A(u, E) := \int_E \sqrt{1 + |\nabla u|^2} d\mathbf{m} + |D^s u|(E).$$

Proposition 2.2. *If $u \in \text{BV}(\Omega)$, then for every Borel set $E \subset \Omega$, we have*

$$P(\text{Epi}(u), E \times \mathbb{R}) \geq A(u, E).$$

Proof. We will consider the measure $\mu := P(\text{Epi}(u), \cdot \times \mathbb{R})$ on the Borel subsets of Ω and we will show separately that

$$\mu \geq \sqrt{1 + |\nabla u|^2} \mathbf{m} \quad \text{and} \quad \mu \geq |D^s u|. \quad (1)$$

Since $\mathbf{m} \perp |D^s u|$ this will then imply that $\mu \geq \sqrt{1 + |\nabla u|^2} \mathbf{m} + |D^s u|$.

Thanks to Lemma 1.9, to obtain (1), it is sufficient to show that for every open set $E \subset \Omega$ we have

$$\mu(E) \geq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} \quad \text{and} \quad \mu(E) \geq |D^s u|(E).$$

We now repeat an argument from [11] to obtain that, for every open set $E \subset X$ and $a, b \in C(E)$ such that $a^2 + b^2 \leq 1$, it holds

$$\mu(E) \geq \int_E a \, d\mathbf{m} + \int_E b \, d|Du|. \quad (2)$$

We report the argument for the sake of completeness. Fix $N \in \mathbb{N}$ and let $g_n \in \text{Lip}(E \times (-N, N))$ be convergent to $1_{\text{Epi}(u)}$ in $L^1(E \times (-N, N))$. Modulo passing to a subsequence, we may assume that $g_n^t \rightarrow 1_{\text{Epi}(u)}^t$ in $L^1(E)$ for λ^1 -a.e. $t \in (-N, N)$ and that $g_n^x \rightarrow 1_{\text{Epi}(u)}^x$ in $L^1((-N, N))$ for \mathbf{m} -a.e. $x \in E$. For every $a, b \in C(E)$ with $a^2 + b^2 \leq 1$, applying Proposition 1.7, we get

$$\begin{aligned} \liminf_{n \rightarrow +\infty} \int_{E \times (-N, N)} |\nabla g_n| \, d\mathbf{m}_\times &\geq \liminf_{n \rightarrow +\infty} \int_{E \times (-N, N)} a(x) |\nabla g_n^x|(t) + b(x) |\nabla g_n^t|(x) \, d\mathbf{m}_\times(x, t) \\ &\geq \int_E \liminf_{n \rightarrow +\infty} \int_{(-N, N)} a(x) |\nabla g_n^x|(t) \, dt \, d\mathbf{m}(x) + \int_{(-N, N)} \liminf_{n \rightarrow +\infty} \int_X b(x) |\nabla g_n^t|(x) \, d\mathbf{m}(x) \, dt \\ &\geq \int_E a 1_{\{u \in (-N, N)\}} \, d\mathbf{m} + \int_{(-N, N)} \int_E b \, d|D1_{\{u > t\}}| \, dt. \end{aligned}$$

Since g_n is arbitrary, we get

$$\mathbf{P}(\text{Epi}(u), E \times (-N, N)) \geq \int_E a 1_{\{u \in (-N, N)\}} \, d\mathbf{m} + \int_{(-N, N)} \int_E b \, d|D1_{\{u > t\}}| \, dt,$$

so that letting $N \uparrow +\infty$ and using the coarea formula we deduce (2).

Equation (2) then implies that

$$\mu(E) \geq \int_E a + b |\nabla u| \, d\mathbf{m}.$$

Taking the supremum over all couples a, b of the aforementioned type and using Lemma 1.8, we get

$$\mu(E) \geq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m}.$$

Similarly, taking $a \equiv 0$ and $b \equiv 1$ in (2), we obtain

$$\mu(E) \geq |Du|(E) \geq |D^s u|(E),$$

concluding the proof. □

Proposition 2.3. *If $u \in \text{Lip}(\Omega)$, then, for every Borel set $E \subset \Omega$, it holds*

$$\mathbf{P}(\text{Epi}(u), E \times \mathbb{R}) = \mathbf{A}(u, E).$$

Proof. Thanks to the previous proposition, we only need to show that

$$\mathbf{P}(\text{Epi}(u), E \times \mathbb{R}) \leq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m}.$$

Thanks to Lemma 1.9, we only need to verify the previous inequality under the assumption that E is open. Let $\chi_\epsilon \in C^\infty(\mathbb{R})$ be monotonically convergent to $1_{(0, +\infty)}$, with values in $[0, 1]$ and $\int_{\mathbb{R}} \chi'_\epsilon \, dt \leq 1$. Then the functions

$$g_\epsilon(x, t) := \chi_\epsilon(u(x) - t)$$

are Lipschitz in $\Omega \times \mathbb{R}$ for every $\epsilon > 0$ and converge in $L^1_{loc}(\Omega \times \mathbb{R})$ to $\text{Epi}(u)$. Moreover Proposition 1.7 gives

$$|\nabla g_\epsilon|(x, t) = \chi'_\epsilon(u(x) - t) \sqrt{1 + |\nabla u|^2(x)} \quad \text{for } \mathbf{m}_X\text{-a.e. } (x, t) \in X \times \mathbb{R}.$$

Integrating both sides and using Fubini's Theorem, we get

$$\int_{E \times \mathbb{R}} |\nabla g_\epsilon| d\mathbf{m}_X \leq \int_E \sqrt{1 + |\nabla u|^2} dm,$$

which implies

$$\mathbf{P}(\text{Epi}(u), E \times \mathbb{R}) \leq \int_E \sqrt{1 + |\nabla u|^2} dm.$$

□

We now prove Theorem 1 from the Introduction.

Theorem 1. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space and let $u \in L^\infty(X) \cap \text{BV}(X)$. Then, for every bounded open set $E \subset X$ such that $|D^s u|(\partial E) = 0$, it holds*

$$\lim_{t \rightarrow 0} \mathbf{A}(P_t(u), E) = \mathbf{A}(u, E).$$

Proof. For every $t > 0$ set $u_t := P_t(u)$ and we note that this function is Lipschitz. We first show that

$$\int_E \sqrt{1 + |\nabla u|^2} dm + |D^s u|(E) \leq \liminf_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla u_t|^2} dm.$$

Using Propositions 2.2 and 2.3 and the lower semicontinuity of perimeters we get

$$\begin{aligned} \int_E \sqrt{1 + |\nabla u|^2} dm + |D^s u|(E) &\leq \mathbf{P}(\text{Epi}(u), E \times \mathbb{R}) \\ &\leq \liminf_{t \rightarrow 0} \mathbf{P}(\text{Epi}(u_t), E \times \mathbb{R}) = \liminf_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla u_t|^2} dm, \end{aligned}$$

as desired.

We now show that

$$\int_E \sqrt{1 + |\nabla u|^2} dm + |D^s u|(E) \geq \limsup_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla u_t|^2} dm.$$

Note that

$$\int_E \sqrt{1 + |\nabla u_t|^2} dm = \sup_{\substack{(a,b) \in C(E) \times C(E) \\ a^2 + b^2 \leq 1}} \int_E a + |\nabla u_t| b dm$$

and for every such pair (a, b) we have

$$\begin{aligned} \int_E a + |\nabla u_t| b dm &\leq \int_E a dm + \int_E b e^{-Kt} d\bar{P}_t(|Du|) \\ &\leq \int_E a dm + \int_E b e^{-Kt} P_t(|\nabla u|) dm + e^{-Kt} \bar{P}_t(|D^s u|)(E). \end{aligned}$$

Hence, passing to the supremum with respect to the pairs (a, b) , we get

$$\int_E \sqrt{1 + |\nabla u_t|^2} dm \leq \int_E \sqrt{1 + e^{-2Kt} P_t(|\nabla u|)^2} dm + e^{-Kt} \bar{P}_t(|D^s u|)(E).$$

Moreover, by definition of \bar{P}_t , we have that $\bar{P}_t(|D^s u|)$ converges weakly to $|D^s u|$ preserving the mass for every t , and since $|D^s u|(\partial E) = 0$ this implies that $\bar{P}_t(|D^s u|)(E) \rightarrow |D^s u|(E)$. In particular, passing to the limit as $t \rightarrow 0$ in the previous inequality, we get

$$\limsup_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla u_t|^2} dm \leq \int_E \sqrt{1 + |\nabla u|^2} dm + |D^s u|(E).$$

□

The next two results are needed to prove Theorem 2 from the Introduction.

Theorem 2.4. *Let $\Omega \subset \mathbb{X}$ be an open bounded set and let $u \in \text{BV}(\Omega)$. For every Borel set $E \subset \Omega$ it holds*

$$\text{P}(\text{Epi}(u), E \times \mathbb{R}) = \text{A}(u, E).$$

Proof. The inequality

$$\text{P}(\text{Epi}(u), E \times \mathbb{R}) \geq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E)$$

is true by Proposition 2.2 so we only show the reverse one.

Suppose first that $u \in \text{L}^\infty(\Omega) \cap \text{BV}(\Omega)$. It is clear that it is sufficient to consider the case when E is well contained in Ω and such that $|Du|(\partial E) = 0$ and then argue by approximation. So we assume that E has these properties and, modulo multiplying u by a cut off function which is equal to 1 in a neighborhood of \bar{E} , we can also suppose that $u \in \text{L}^\infty(\Omega) \cap \text{BV}(\Omega)$ with $\text{supp}(u) \subset\subset \Omega$. In particular, when we extend u to be 0 outside of Ω , we obtain a function in $u \in \text{L}^\infty(\mathbb{X}) \cap \text{BV}(\mathbb{X})$. When we refer to u in the next lines we implicitly mean its extension to \mathbb{X} . By lower semicontinuity of perimeters and the previous proposition we then get

$$\begin{aligned} \text{P}(\text{Epi}(u), E \times \mathbb{R}) &\leq \liminf_{t \rightarrow 0} \text{P}(\text{Epi}(P_t(u)), E \times \mathbb{R}) \\ &= \liminf_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla P_t(u)|^2} \, d\mathbf{m} = \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E), \end{aligned}$$

concluding the proof when $u \in \text{L}^\infty(\Omega) \cap \text{BV}(\Omega)$.

If $u \in \text{BV}(\Omega)$ is not bounded, we consider for every $k > 0$ the function $u^k := -k \vee u \wedge k$ and we exploit the lower semicontinuity of perimeters to obtain that for every open set $E \subset \Omega$ we have

$$\begin{aligned} \text{P}(\text{Epi}(u), E \times \mathbb{R}) &\leq \liminf_{k \rightarrow +\infty} \text{P}(\text{Epi}(u^k), E \times \mathbb{R}) \\ &= \liminf_{k \rightarrow +\infty} \int_E \sqrt{1 + |\nabla u^k|^2} \, d\mathbf{m} + |D^s u^k|(E). \end{aligned}$$

Moreover, since $|Du^k| \leq |Du|$ as measures, the last term is controlled by

$$\int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E),$$

concluding the proof. □

Proposition 2.5. *If $u : B_{2r}(x) \rightarrow \mathbb{R}$ is a measurable function whose epigraph has finite perimeter, then $u \in \text{BV}(B_r(x))$. Similarly if Ω is bounded and $u \in \text{L}^1(\Omega)$ has epigraph of finite perimeter, then $u \in \text{BV}(\Omega)$.*

Proof. Assume first that Ω is bounded and that $u \in \text{L}^1(\Omega)$ is a function whose epigraph has finite perimeter. Fix $N \in \mathbb{N}$ and let $g_n \in \text{Lip}(\Omega \times (-N, N))$ be convergent to $\mathbf{1}_{\text{Epi}(u)}$ in $\text{L}^1(\Omega \times (-N, N))$. Modulo passing to a subsequence, we may assume that $g_n^t \rightarrow \mathbf{1}_{\text{Epi}(u)}^t$ in $\text{L}^1(\Omega)$ for λ^1 -a.e. $t \in (-N, N)$. Then

$$\liminf_{n \rightarrow +\infty} \int_{\Omega \times (-N, N)} |\nabla g_n| \, d\mathbf{m}_\times \geq \liminf_{n \rightarrow +\infty} \int_{\Omega \times (-N, N)} |\nabla g_n^t| \, d\mathbf{m}_\times \geq \int_{-N}^N \text{P}(\{u > t\}, \Omega) dt.$$

Since g_n is arbitrary, letting $N \uparrow +\infty$, we deduce

$$\int_{\mathbb{R}} \text{P}(\{u > t\}, \Omega) dt < +\infty. \quad (3)$$

Since $u \in \text{L}^1(\Omega)$, this is sufficient to conclude that $u \in \text{BV}(\Omega)$.

If $u : B_{2r}(x) \rightarrow \mathbb{R}$ is measurable we need to prove that $u \in \mathbf{L}^1(B_r(x))$, and then we get that $u \in \mathbf{BV}(B_r(x))$ by the previous part of the proof. To this aim we write

$$\begin{aligned} \int_{B_r(x)} |u| \, d\mathbf{m} &= \int_0^\infty \mathbf{m}(\{|u| \geq t\} \cap B_r(x)) \, dt \\ &\leq T\mathbf{m}(B_r(x)) + \int_T^\infty \mathbf{m}(\{|u| \geq t\} \cap B_r(x)) \, dt. \end{aligned}$$

If $T \in \mathbb{N}$ is large enough, for every $t \geq T$ the isoperimetric inequality gives

$$\mathbf{m}(\{|u| \geq t\} \cap B_r(x)) \leq c(K, r)\mathbf{P}(\{|u| \geq t\}, B_{2r}(x)).$$

In particular, combining this with (3), we get

$$\int_{B_r(x)} |u| \, d\mathbf{m} \leq T\mathbf{m}(B_r(x)) + c(K, r) \int_T^\infty \mathbf{P}(\{|u| \geq t\}, B_{2r}(x)) \, dt < +\infty,$$

concluding the proof. \square

Combining Theorem 2.4 and the previous proposition we immediately get Theorem 2, which we recall below.

Theorem 2. *Let (X, d, \mathbf{m}) be an $\mathbf{RCD}(K, \infty)$ space, let $\Omega \subset X$ be open and bounded and let $u \in \mathbf{L}^1(\Omega)$. Then $\mathbf{P}(\mathbf{Epi}(u), \Omega \times \mathbb{R}) < +\infty$ if and only if $u \in \mathbf{BV}(\Omega)$. In this case, for every Borel set $E \subset \Omega$, it holds*

$$\mathbf{P}(\mathbf{Epi}(u), E \times \mathbb{R}) = \mathbf{A}(u, E).$$

Moreover, if $u \in \mathbf{L}^0(B_{2r}(x))$ and $\mathbf{P}(\mathbf{Epi}(u), B_{2r}(x) \times \mathbb{R}) < +\infty$, then $u \in \mathbf{BV}(B_r(x))$ and the previous equation holds for every Borel set $E \subset B_r(x)$.

We now turn our attention to Theorem 3, which will be an easy consequence of the approximation property of Theorem 1. As anticipated in the Introduction, given $u \in \mathbf{BV}(\Omega)$ and an open set $E \subset \Omega$, we define the relaxed area functional as

$$\tilde{\mathbf{A}}(u, E) := \inf \left\{ \liminf_{n \rightarrow +\infty} \int_E \sqrt{1 + \mathbf{lip}(f_n)^2} \, d\mathbf{m} \mid (f_n)_n \subset \mathbf{L}^1(E) \cap \mathbf{Lip}(E), \|f_n - u\|_{\mathbf{L}^1(E)} \rightarrow 0 \right\}.$$

In [11] it was proved that for every $f \in \mathbf{W}^{1,1}(\Omega)$, we have

$$\tilde{\mathbf{A}}(f, E) = \int_E \sqrt{1 + |\nabla f|^2} \, d\mathbf{m},$$

and in Theorem 3 we extend this to functions in $\mathbf{BV}(\Omega)$.

Theorem 3. *Let (X, d, \mathbf{m}) be an $\mathbf{RCD}(K, \infty)$ space, let $\Omega \subset X$ be open and bounded and let $u \in \mathbf{BV}(\Omega)$. Then, for every open set $E \subset \Omega$, it holds*

$$\tilde{\mathbf{A}}(u, E) = \mathbf{A}(u, E).$$

Proof. We first show that for every open set $E \subset \Omega$ we have

$$\tilde{\mathbf{A}}(u, E) \geq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E).$$

To this aim fix an open set E and let $\{u_n\}_{n \in \mathbb{N}} \subset \mathbf{Lip}(E)$ be such that

$$\int_E \sqrt{1 + |\nabla u_n|^2} \, d\mathbf{m} \rightarrow \tilde{\mathbf{A}}(u, E)$$

and $u_n \rightarrow u$ in $L^1(E)$. By the lower semicontinuity of perimeters, we get

$$A(u, E) = \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E) \leq \liminf_{n \rightarrow +\infty} \int_E \sqrt{1 + |\nabla u_n|^2} \, d\mathbf{m} = \tilde{A}(u, E).$$

We now show that

$$\tilde{A}(u, E) \leq A(u, E)$$

for every open set $E \subset \Omega$. To this aim it is sufficient to consider open sets $E \subset \subset \Omega$ such that $|D^s u|(\partial E) = 0$ and then argue by approximation. Let $u^k := -k \vee u \wedge k$ and note that for every $\epsilon > 0$ we have that if k is sufficiently large $\tilde{A}(u)(E) \leq \tilde{A}(u^k)(E) + \epsilon$. Using the lower semicontinuity of \tilde{A} , the fact that \tilde{A} is the area functional on Lipschitz functions and Theorem 1, we then obtain that

$$\begin{aligned} \tilde{A}(u^k, E) &\leq \liminf_{t \rightarrow 0} \tilde{A}(P_t(u^k))(E) = \liminf_{t \rightarrow 0} \int_E \sqrt{1 + |\nabla P_t(u^k)|^2} \, d\mathbf{m} \\ &= \int_E \sqrt{1 + |\nabla u^k|^2} \, d\mathbf{m} + |D^s u^k|(E). \end{aligned}$$

In particular we get that, if k is large enough, we have

$$\tilde{A}(u, E) \leq \int_E \sqrt{1 + |\nabla u^k|^2} \, d\mathbf{m} + |D^s u^k|(E) + \epsilon,$$

so that $\tilde{A}(u, E) \leq \int_E \sqrt{1 + |\nabla u|^2} \, d\mathbf{m} + |D^s u|(E) + \epsilon$. Since $\epsilon > 0$ is arbitrarily small, we conclude. \square

We conclude the section with some basic properties of minimizers of the area functional in $\text{RCD}(K, \infty)$ spaces that are needed to prove Theorem 4 from the Introduction.

Definition 2.6. Let (X, d, \mathbf{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subset X$ be an open set and let $u \in \text{BV}_{loc}(\Omega)$. We say that u is an area minimizer in $\Omega' \subset \subset \Omega$ if for every $f \in \text{BV}(\Omega')$ such that $\{f \neq u\} \subset \subset \Omega'$ we have

$$A(u, \Omega') \leq A(f, \Omega').$$

If we say that $u \in \text{BV}_{loc}(\Omega)$ is an area minimizer, we mean that the previous condition is satisfied for every $\Omega' \subset \subset \Omega$.

The following proposition follows from [29, Theorem 4.2] keeping in mind that in our setting the area functional coincides with the relaxed area functional thanks to Theorem 3. Since the setting of [29] is the one of PI spaces (and the proof of [29, Theorem 4.2] relies crucially on the doubling assumption), the next result holds in the finite dimensional setting of $\text{RCD}(K, N)$ spaces.

Proposition 2.7. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, N)$ space, let $\Omega \subset X$ be an open set and let $u \in \text{BV}_{loc}(\Omega)$ be an area minimizer, then $u \in L^\infty_{loc}(\Omega)$.*

A similar result holds for functions whose epigraph is perimeter minimizing adapting an argument that can be found in [28]. We report such argument with the due adaptations.

Proposition 2.8. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, N)$ space, let $\Omega \subset X$ be open and let $u \in L^1(\Omega)$ be a function whose epigraph minimizes the perimeter. Then $u \in L^\infty_{loc}(\Omega)$.*

Proof. Let $x \in \Omega$ and suppose for simplicity that $B_3(x) \subset \subset \Omega$. We will show that u is uniformly bounded in its Lebesgue points in $B_1(x)$, which will then imply the claim. To this aim let $y \in B_1(x)$ be a Lebesgue point for u and assume without loss of generality that $u(y) > 0$. Denoting by $\text{Epi}(u)$ the closed representative of the epigraph, we have that $(y, u(y)) \in \partial \text{Epi}(u)$, so that the density estimates from Theorem 1.15 imply that

$$\mathbf{m}_\times(B_1^\times(y, u(y)) \cap \text{Epi}(u)) \geq c(K, N).$$

In particular if $T \in \mathbb{N}$ and $u(y) > 2T$ we have

$$\begin{aligned} \mathbf{m}_\times(\text{Epi}(u) \cap B_1(y) \times [0, 2T]) &\geq \sum_{i=1}^T \mathbf{m}_\times(\text{Epi}(u) \cap B_1^\times(y, 2i)) \\ &\geq T \mathbf{m}_\times(B_1^\times(y, u(y)) \cap \text{Epi}(u)) \geq Tc(K, N). \end{aligned}$$

This implies that

$$\int_{B_2(x)} u \vee 0 \, d\mathbf{m} \geq \int_{B_1(y)} u \vee 0 \, d\mathbf{m} \geq Tc(K, N).$$

Putting everything together, if $y \in B_1(x)$ is a Lebesgue point of u and $u(y) > 2T$, then $T \leq C(x, u, K, N) < +\infty$. This shows that for every such y we have $u(y) \leq 2C(x, u, K, N) + 1$, proving the claim and the statement. \square

The proof of the next proposition follows by adapting arguments from [17, Section 3.2] and we report it for the sake of completeness.

Proposition 2.9. *Given an $\text{RCD}(K, \infty)$ space (X, d, \mathbf{m}) , an open set $\Omega \subset X$ and a function $u \in L^\infty(\Omega) \cap \text{BV}(\Omega)$, then u is a minimizer of the area functional in Ω if and only if its epigraph is a perimeter minimizer in $\Omega \times \mathbb{R}$.*

Proof. If the epigraph of u is perimeter minimizing, then for every $f \in \text{BV}(\Omega) \cap L^\infty(\Omega)$ such that $\{f \neq u\} \subset\subset \Omega$ we have $\text{Epi}(u) \Delta \text{Epi}(f) \subset\subset \Omega \times \mathbb{R}$, so that applying Theorem 2 we obtain

$$A(u, \Omega) \leq A(f, \Omega).$$

Similarly, if $f \in \text{BV}(\Omega)$ is such that $\{f \neq u\} \subset\subset \Omega$, then considering $f^c := -c \vee f \wedge c$ with $c := \|u\|_{L^\infty} + 1$ we get by the previous case that

$$A(u, \Omega) \leq A(f^c, \Omega) \leq A(f, \Omega).$$

We now prove the reverse implication, i.e. that if $u \in L^\infty(\Omega) \cap \text{BV}(\Omega)$ is a minimizer of the area functional in Ω , then its epigraph is a perimeter minimizer in $\Omega \times \mathbb{R}$.

To this aim, consider a set $E \subset \Omega \times \mathbb{R}$ such that $\text{Epi}(u) \Delta E \subset\subset \Omega \times \mathbb{R}$. Modulo translating vertically, we may suppose that there exists $c > 1$ such that u takes values in $(1, c - 1)$ and $\text{Epi}(u) \Delta E \subset\subset \Omega \times (1, c - 1)$. We then define $w(E) : \Omega \rightarrow \mathbb{R}$ by

$$w(E)(x) := \int_0^c 1_E(x, s) \, ds$$

and we claim that

$$w(E) \in \text{BV}(\Omega) \quad \text{and} \quad A(w(E), \Omega) \leq P(E, \Omega \times \mathbb{R}). \quad (4)$$

If the claim holds, noting that by construction $w(E)$ is a competitor for u , we get

$$P(\text{Epi}(u), \Omega \times \mathbb{R}) = A(u, \Omega) \leq A(w(E), \Omega) \leq P(E, \Omega \times \mathbb{R}).$$

To prove the claim, we consider a sequence $f_n \in \text{Lip}(\Omega \times (0, c))$ converging in $L^1(\Omega \times (0, c))$ to 1_E and such that

$$\lim_{n \rightarrow +\infty} |Df_n|(\Omega \times (0, c)) = P(E, \Omega \times \mathbb{R}).$$

Modulo truncating, we can assume that $f_n \equiv 1$ on $\Omega \times \{0\}$ and $f_n \equiv 0$ on $\Omega \times \{c\}$. We then define $w(f_n) : \Omega \rightarrow \mathbb{R}$ as

$$w(f_n)(x) := \int_0^c f_n(x, s) \, ds.$$

One can then check with arguments that were previously used in the note that $w(f_n) \in \text{Lip}(\Omega)$ and that

$$A(w(f_n), \Omega) \leq |Df_n|(\Omega \times (0, c)).$$

Similarly, one can check that $w(f_n) \rightarrow w(E)$ in $L^1(\Omega)$ as $n \rightarrow +\infty$. This implies that $w(E) \in \text{BV}(\Omega)$ as claimed. Concerning the area of $w(E)$, we get

$$A(w(E), \Omega) \leq \liminf A(w(f_n), \Omega) \leq \liminf |Df_n|(\Omega \times (0, c)) = P(E, \Omega \times \mathbb{R}),$$

proving the claim (4). \square

If the space in question is finite dimensional, the previous result holds without the assumption that u is bounded. This is the content of Proposition 2.10.

Proposition 2.10. *Let (X, d, \mathfrak{m}) be an $\text{RCD}(K, N)$ space, let $\Omega \subset X$ be open and let $u \in \text{BV}(\Omega)$, then u is a minimizer of the area functional in Ω if and only if its epigraph is a perimeter minimizer in $\Omega \times \mathbb{R}$.*

Proof. If u minimizes the area, then, by Proposition 2.7, u is locally bounded in Ω and we can conclude by Proposition 2.9.

Conversely, if the epigraph of u minimizes the perimeter, then u is locally bounded by Proposition 2.8. Consider a competitor $f \in \text{BV}(\Omega)$ for u and a subset $\Omega' \subset\subset \Omega$ such that $\{f \neq u\} \subset\subset \Omega'$. We define $f^c := -c \vee f \wedge c$ for $c := \|u\|_{L^\infty(\Omega')} + 1$, so that

$$A(u, \Omega') = P(\text{Epi}(u), \Omega' \times \mathbb{R}) \leq P(f^c, \Omega' \times \mathbb{R}) = A(f^c, \Omega') \leq A(f, \Omega'),$$

which implies

$$A(u, \Omega) \leq A(f, \Omega).$$

\square

We now prove Theorem 4 and its corollary.

Theorem 4. *Let (X, d, \mathfrak{m}) be an $\text{RCD}(K, \infty)$ space, let $\Omega \subsetneq X$ be an open set and let $u \in L^\infty(\Omega) \cap W^{1,1}(\Omega)$ be a function whose epigraph is perimeter minimizing. If $v \in L^\infty(\Omega)$ has epigraph which is perimeter minimizing and $\{u \neq v\} \subset\subset \Omega$, then $u = v$.*

Proof. By Theorem 2, $v \in \text{BV}(\Omega)$ and by Proposition 2.9, both u and v are area minimizers in Ω .

For every $t \in (0, 1)$ we consider the convex combination $tu + (1-t)v$. We have

$$A(tu + (1-t)v, \Omega) \leq tA(u, \Omega) + (1-t)A(v, \Omega) = A(u, \Omega).$$

Since $tu + (1-t)v$ is a competitor for u , the previous inequality implies

$$A(tu + (1-t)v, \Omega) = A(u, \Omega). \quad (5)$$

The function $a \mapsto \sqrt{1+a^2}$ is strictly convex on the positive real line. Hence, (5) implies that, for every $t \in (0, 1)$, it holds

$$|\nabla(tu + (1-t)v)| = |\nabla u| \quad \mathfrak{m}\text{-almost everywhere in } \Omega.$$

Taking the squares in the previous identity and evaluating at $t = 0$ and $t = 1/2$, we deduce

$$\begin{cases} \nabla u \cdot \nabla v = |\nabla u|^2 \\ |\nabla v|^2 = |\nabla u|^2, \end{cases}$$

which in turn implies

$$|\nabla(u-v)| = 0 \quad \mathfrak{m}\text{-almost everywhere in } \Omega.$$

Hence, the function $\psi := u-v$ satisfies $|D\psi| = |D^s\psi|$ and has support well contained in Ω . Moreover, the function $t \mapsto A(u+t\psi, \Omega)$ is constant in $(0, 1)$. Hence

$$0 = \frac{d}{dt}A(u+t\psi, \Omega) = \frac{d}{dt}|D^s(u+t\psi)|(\Omega) = \frac{d}{dt}|D^s(t\psi)|(\Omega) = |D^s\psi|(\Omega) = |D\psi|(\Omega).$$

In particular, ψ is constant. Since the support of ψ is well contained in Ω and $\Omega \subsetneq X$, it holds $\psi \equiv 0$. \square

Corollary. *Let (X, d, m) be an $\text{RCD}(K, N)$ space, let $\Omega \subsetneq X$ be an open set and let $u \in W^{1,1}(\Omega)$ be a function whose epigraph is perimeter minimizing. If $v \in L^1(\Omega)$ has epigraph which is perimeter minimizing and $\{u \neq v\} \subset \subset \Omega$, then $u = v$.*

Proof. By Lemma 2.8 both u and v are locally bounded. The statement follows by Theorem 4. \square

3 Approximation of minimal graphs in $\text{RCD}(K, N)$

The goal of this section is to prove Theorem 3.3, which is a crucial step towards Theorem 5.

We consider spaces (X_i, d_i, m_i, x_i) converging to a space (X, d, m, x) in pmGH sense and functions $u_i \in \text{BV}(X_i)$ converging in L^1 to $u \in \text{BV}(X)$. As anticipated, we use the so-called extrinsic viewpoint and we work in the space (Z, d_Z) which realizes the convergence. In particular, whenever we consider a ball $B_r(x)$ in this context, unless otherwise specified, we mean the ball of radius r centered in x in the space Z (observe that the intersection of this ball with the embedding of X in Z would give the corresponding ball in X). Similarly, unless otherwise stated, each function u_i will be considered as a function on Z (extending it to zero outside of the embedding of X_i in Z), so that it will make sense to consider products of the form $u_i 1_{B_r(x)} : Z \rightarrow \mathbb{R}$.

Given $f : Z \rightarrow \mathbb{R}$, we will write $A(f, B_r(x), m_i)$ meaning the area of f on $B_r(x) \subset Z$ w.r.t. to the pushforward of the measure m_i in Z and we will write $f \in L^\infty(B_r(x), m_i)$ (and the same notation will be used for Lebesgue and Sobolev spaces) to specify that the measure considered is again the pushforward of the measure m_i in Z .

Finally we will denote by $P_t^{X_i}$ the heat flow on the space X_i , noting that this coincides with the heat flow w.r.t. (the pushforward of) m_i on Z .

Theorem 3.1. *Let (X_i, d_i, m_i, x_i) be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to (X, d, m, x) .*

1. *If $u \in \text{BV}(X, m)$, and $u_i \in \text{BV}(X_i, m_i)$ is a sequence such that $u_i \rightarrow u$ in L^1 , then for every open set $A \subset Z$ we have*

$$A(u, A, m) \leq \liminf_i A(u_i, A, m_i).$$

2. *If $u \in \text{BV}(B_r(x), m)$, then for every $s \in (0, r)$ such that $|D^s u|(\partial B_s(x)) = 0$ there exists a sequence of Lipschitz functions $u_i \in L^1(B_s(x), m_i)$ such that $u_i 1_{B_s(x)}$ converges in L^1 -sense to $u 1_{B_s(x)} \in L^1(X, m)$ and $A(u_i, B_s(x), m_i) \rightarrow A(u, B_s(x), m)$.*

Proof. The first assertion follows by combining the localized lower semicontinuity of total variations given in Proposition 1.13 and the identification between area and perimeter given in Theorem 2.

We now prove the second assertion following the proof in [9, Theorem 8.1].

Fix $s \in (0, r)$ and observe that, modulo multiplying with a cut off function, we can assume that u has support well contained in $B_r(x)$. Also, modulo truncating (and using a diagonal argument), we can also assume that $u \in L^\infty(B_r(x), m)$ (as the areas of the truncated functions converge to the area of the initial function). In what follows, when we refer to u , we implicitly mean its extension to zero on the whole Z .

Since $|D^s u|(\partial B_s(x)) = 0$, Theorem 1 guarantees that

$$A(P_t^X(u), B_s(x), m) \rightarrow A(u, B_s(x), m) \quad \text{as } t \rightarrow 0.$$

Hence, if we manage to approximate in area the functions $P_t^X(u)$, we can then conclude with a diagonal argument. Each function $P_t^X(u)$ belongs to $W^{1,2}(Z, m)$, so that by Remark 1.2 we can consider a sequence $f_k \in \text{Lip}_c(Z)$ converging in $L^2(Z, m)$ to $P_t^X(u)$ and such that $\text{lip}_a(f_k) \rightarrow |\nabla P_t^X(u)|$ in $L^2(Z, m)$. We then fix k , and as i varies we get

$$\begin{aligned} A(f_k, B_s(x), m_i) &\leq \int_{B_s(x)} \sqrt{1 + \text{lip}_a^2 f_k} \, dm_i \\ &\leq \int_Z 1_{B_s(x)} \sqrt{1 + \text{lip}_a^2 f_k} \, dm_i. \end{aligned}$$

Taking a limit as i goes to infinity and taking into account that the last integrand is upper semi-continuous and that $\mathbf{m}(\partial B_s(x)) = 0$, we then obtain

$$\limsup_{i \rightarrow +\infty} \mathbf{A}(f_k, B_s(x), \mathbf{m}_i) \leq \int_Z 1_{\bar{B}_s(x)} \sqrt{1 + \text{lip}_a^2 f_k} \, d\mathbf{m} = \int_{B_s(x)} \sqrt{1 + \text{lip}_a^2 f_k} \, d\mathbf{m}.$$

Reindexing the sequence $\{f_k\}_{k \in \mathbb{N}}$ (so that each function is repeated a sufficient number of times) we then get

$$\limsup_{k \rightarrow +\infty} \mathbf{A}(f_k, B_s(x), \mathbf{m}_k) \leq \int_{B_s(x)} \sqrt{1 + \text{lip}^2 P_t(u)} \, d\mathbf{m} = \mathbf{A}(P_t(u), B_s(x), \mathbf{m}).$$

which is the desired approximation. \square

The same proof of the previous theorem can be repeated in the global setting to obtain the Γ -convergence of the area functional. In particular, replacing Proposition 1.13 with the results of [9], we deduce the aforementioned convergence in the case $N = \infty$.

Theorem 3.2. *Let $(X_i, \mathbf{d}_i, \mathbf{m}_i, x_i)$ be a sequence of $\text{RCD}(K, \infty)$ spaces converging in pmGH sense to $(X, \mathbf{d}, \mathbf{m}, x)$.*

1. *If $u \in \text{BV}(X, \mathbf{m})$, and $u_i \in \text{BV}(X_i, \mathbf{m}_i)$ is a sequence such that $u_i \rightarrow u$ in L^1 -sense, then*

$$\mathbf{A}(u, X, \mathbf{m}) \leq \liminf_i \mathbf{A}(u_i, X_i, \mathbf{m}_i).$$

2. *If $u \in \text{BV}(X, \mathbf{m})$, then there exists a sequence of functions $u_i \in \text{BV}(X_i, \mathbf{m}_i)$ converging in area to u .*

We are now in position to prove the main result of the section.

Theorem 3.3. *Let $(X_i, \mathbf{d}_i, \mathbf{m}_i, x_i)$ be a sequence of $\text{RCD}(K, N)$ spaces converging in pmGH sense to $(X, \mathbf{d}, \mathbf{m}, x)$. Let $r > 0$ be such that $B_r(x) \subsetneq X$ and let $u \in \text{BV}(B_r(x), \mathbf{m})$ be an area minimizer. For every $s \in (0, r)$, there exists $0 < \delta < r - s$ such that the following happens.*

There exists area minimizers $\tilde{u}_i \in \text{BV}(B_s(x), \mathbf{m}_i)$ and $\tilde{u} \in \text{BV}(B_{s+\delta}(x), \mathbf{m})$ such that $\tilde{u}_i 1_{B_s(x)}$ converges in L^1 to $\tilde{u} 1_{B_s(x)}$ and $\{\tilde{u} \neq u\} \subset\subset B_{s+\delta}(x)$.

Moreover, if $|D^s \tilde{u}|(\partial B_s(x)) = 0$, it holds $\mathbf{A}(\tilde{u}_i, B_s(x), \mathbf{m}_i) \rightarrow \mathbf{A}(\tilde{u}, B_s(x), \mathbf{m})$.

Proof. Let $0 < \epsilon < (r - s)/2$ be such that $|D^s u|(\partial B_{s+2\epsilon}(x)) = 0$. We set $\delta := 2\epsilon$. Consider the sequence $u_i \in \text{BV}(B_{s+2\epsilon}(x), \mathbf{m}_i) \cap L^\infty(B_{s+2\epsilon}(x), \mathbf{m}_i)$ converging in area to u given by Theorem 3.1. Modulo truncating, we can also assume that $\|u_i\|_{L^\infty(B_{s+2\epsilon}(x), \mathbf{m}_i)} \leq \|u\|_{L^\infty(B_{s+2\epsilon}(x), \mathbf{m})}$.

Then, we consider for every $i \in \mathbb{N}$ the function $\tilde{u}_i \in \text{BV}(B_{s+2\epsilon}(x), \mathbf{m}_i)$ such that $u_i = \tilde{u}_i$ on $B_{s+2\epsilon}(x) \setminus B_{s+\epsilon}(x)$ and

$$\mathbf{A}(\tilde{u}_i, B_{s+2\epsilon}(x), \mathbf{m}_i) \leq \mathbf{A}(f, B_{s+2\epsilon}(x), \mathbf{m}_i)$$

for every $f \in \text{BV}(B_{s+2\epsilon}(x), \mathbf{m}_i)$ such that $\{f \neq \tilde{u}_i\} \subset B_{s+\epsilon}(x)$. Existence of the functions \tilde{u}_i follows by the direct method of calculus of variations. We also note that the condition $u_i = \tilde{u}_i$ on $B_{s+2\epsilon}(x) \setminus B_{s+\epsilon}(x)$ is nonempty by the assumption that $B_r(x) \neq X$. Observe moreover that each \tilde{u}_i satisfies $\|\tilde{u}_i\|_{L^\infty(B_{s+2\epsilon}(x), \mathbf{m}_i)} \leq \|u_i\|_{L^\infty(B_{s+2\epsilon}(x), \mathbf{m}_i)}$ since, if this is not the case, we can truncate and obtain better competitors.

Consider then the function $\tilde{u} \in \text{BV}(B_{s+2\epsilon}(x), \mathbf{m})$ which is the L^1 -limit in $B_{s+2\epsilon}(x)$ of a (non relabeled) subsequence of the functions \tilde{u}_i (this exists thanks to Proposition 1.12 and a cutoff argument). We now claim that \tilde{u} is an area minimizer in $B_{s+2\epsilon}(x)$ w.r.t. \mathbf{m} .

To this aim, it is sufficient to show that

$$\mathbf{A}(\tilde{u}, B_{s+2\epsilon}(x), \mathbf{m}) = \mathbf{A}(u, B_{s+2\epsilon}(x), \mathbf{m}). \quad (6)$$

Observe that by construction $\{u \neq \tilde{u}\} \subset\subset B_{s+2\epsilon}(x)$ and for every i we have $\mathbf{A}(\tilde{u}_i, B_{s+2\epsilon}(x)) \leq \mathbf{A}(u_i, B_{s+2\epsilon}(x))$. Hence, since u is an area minimizer, we have that

$$\begin{aligned} \mathbf{A}(\tilde{u}, B_{s+2\epsilon}(x), \mathbf{m}) &\geq \mathbf{A}(u, B_{s+2\epsilon}(x), \mathbf{m}) = \lim_{i \rightarrow +\infty} \mathbf{A}(u_i, B_{s+2\epsilon}(x), \mathbf{m}_i) \\ &\geq \limsup_{i \rightarrow +\infty} \mathbf{A}(\tilde{u}_i, B_{s+2\epsilon}(x), \mathbf{m}_i) \geq \mathbf{A}(\tilde{u}, B_{s+2\epsilon}(x), \mathbf{m}), \end{aligned} \quad (7)$$

so that

$$A(\tilde{u}, B_{s+2\epsilon}(x), \mathbf{m}) = A(u, B_{s+2\epsilon}(x), \mathbf{m}).$$

This concludes the proof of the statement.

Suppose now that $|D^s \tilde{u}|(\partial B_s(x)) = 0$. By (7) we know that

$$A(\tilde{u}_i, B_{s+2\epsilon}(x), \mathbf{m}_i) \rightarrow A(\tilde{u}, B_{s+2\epsilon}(x), \mathbf{m}).$$

Convergence in area then follows by a standard lower semicontinuity argument combining the fact that $A(\tilde{u}, \partial B_s(x), \mathbf{m}) = 0$ and point 1 of Theorem 3.1. \square

In the next section, when using the previous theorem, we will occasionally say that the functions u_i converge in L^1 -sense to u in $B_s(x)$, meaning that $u_i \mathbf{1}_{B_s(x)} \rightarrow u \mathbf{1}_{B_s(x)}$ in L^1 -sense.

4 Applications to non-collapsed Ricci limits

We recall that, given a metric measure space (X, d, \mathbf{m}) and a function $u \in \text{Lip}_{loc}(X)$, we will use the following notation for balls on the graph of u :

$$B_r^g(x, t) := B_r^\times(x, t) \cap \text{graph}(u) \subset X \times \mathbb{R}.$$

Similarly, given a point $x \in X$, we will denote by \bar{x} its projection on the graph of u , i.e. $\bar{x} := (x, u(x))$.

In this section, we will often work in the setting of Riemannian manifolds. Whenever we say that we consider a Riemannian manifold (M^n, d) , we implicitly mean that we have a Riemannian metric g on M and that d is the Riemannian distance. Given a metric space (X, d) , we denote by H^n the n -dimensional Hausdorff measure induced by d .

In what follows, we will use the fact that minimizers of the area functional on smooth manifolds are smooth.

We now prove three basic lemmas.

Lemma 4.1. *Let $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let (M_k^n, d_k, x_k) be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ converging in pGH sense to (X, d, x) . For every $r > 0$, if k is large enough, then $H_k^n(B_r(x_k)) \geq H^n(B_r(x))/2$.*

Proof. If $H^n(B_r(x)) = 0$ the statement follows trivially. If instead $H^n(B_r(x)) > 0$, then the Hausdorff dimension of (X, d) is at least n , so that Theorem 1.10 gives that the sequence (M_k^n, d_k, H_k^n, x_k) converges in pmGH sense to (X^n, d, H^n, x) . The statement then follows by lower semicontinuity of measures w.r.t. weak convergence. \square

In the next lemma, we consider the behavior of the local Lipschitz constant of an area minimizing function $u \in \text{BV}(\Omega)$. Here, we implicitly mean that we are considering a precise representative for u , i.e. the function whose value in x is given by

$$\limsup_{s \rightarrow 0} \int_{B_s(x)} u \, d\mathbf{m}.$$

This will be implicitly used when we exploit the fact that $\text{graph}(u) \subset \partial \text{Epi}(u)$, where $\text{Epi}(u)$ denotes the closed representative of the epigraph of u .

Lemma 4.2. *Let $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let (M_i^n, d_i, H_i^n, x_i) be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_i^n} \geq K$ converging in pmGH sense to (X, d, \mathbf{m}, x) . If $u \in \text{BV}(B_r(x), \mathbf{m})$ is an area minimizer and there is a sequence of area minimizers $u_i \in \text{BV}(B_R(x_i), H_i^n)$ with $R > r$ converging in L_1 to u on $B_r(x)$ with*

$$\sup_{B_R(x_i)} \text{lip}(u_i) \leq c,$$

then

$$\sup_{B_r(x)} \text{lip}(u) \leq c.$$

Proof. We fix $\delta > 0$ and we claim that, for i large enough, we have

$$\sup_{B_r(x)} \text{lip}(u) \leq c + \delta.$$

To this end, suppose by contradiction that the statement is false. Therefore, there exists $y \in B_r(x) \cap X$ and a sequence $y_k \rightarrow y$ contained in $B_r(x) \cap X$ such that

$$\frac{|u(y) - u(y_k)|}{d(y, y_k)} > c.$$

Since the boundaries of the epigraphs of u_i converge in Kuratowski sense in $B_r(x) \times \mathbb{R}$ to $\partial \text{Epi}(u)$, and since $\text{graph}(u) \subset \partial \text{Epi}(u)$, we have that for every k there exists a sequence $y_i^k \in B_r(x) \cap M_i$ such that $y_i^k \rightarrow y_k$ in Z and $u_i(y_i^k) \rightarrow u(y_k)$. Similarly, there exists $y_i \in B_r(x) \cap M_i$ such that $y_i \rightarrow y$ in Z and $u_i(y_i) \rightarrow u(y)$. This implies that for every $k > 0$ fixed, for i large enough, we have

$$\frac{|u_i(y_i) - u_i(y_i^k)|}{d_i(y_i, y_i^k)} > c.$$

In particular, if k and i are sufficiently large, we will have that the geodesic in M_i connecting y_i and y_i^k is contained in $B_R(x_i)$. Integrating along this geodesic we have

$$|u_i(y_i) - u_i(y_i^k)| \leq cd_i(y_i, y_i^k),$$

a contradiction. □

Lemma 4.3. *Let (M, g) be a Riemannian manifold and let $u \in C^\infty(B_r(x))$ be a smooth function. If $0 < t \leq r$, $c > 1$ and*

$$\sup_{B_t^g(\bar{x})} \text{lip}(u) \leq c,$$

then

$$\pi(\bar{B}_t^g(\bar{x})) \supset B_{t/(2c)}(x).$$

Proof. Let $y \in B_{t/(2c)}(x)$ and let γ be a length minimizing geodesic connecting x and y . If u restricted to this geodesic takes value in $B_{t/(2c)}(x) \times (u(x) - t/2, u(x) + t/2)$, then there is nothing left to prove. So suppose by contradiction that this is not true. Without loss of generality this implies that there exists $\gamma(t_0)$ such that $u(\gamma(t_0)) = u(x) + t/2$, and we can also suppose that for every $s \in (0, t_0)$ we have $u(\gamma(s)) \in (u(x) - t/2, u(x) + t/2)$. In this case we have that

$$|u(x) - u(\gamma(t_0))| \leq cd(\gamma(t_0), x) < t/2,$$

a contradiction. □

We now prove a first regularity result for area minimizers in non-collapsed Ricci limit spaces.

Proposition 4.4. *Let $r > 0$, $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let (M_k^n, d_k, x_k) be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $H_k^n(B_r(x_k)) \geq vr^n$ converging in pointed Gromov-Hausdorff sense to (X, d, x) . Let $r/2 < s < r$, let $u_k \in \text{BV}(B_s(x), H_k^n)$ and $u \in \text{BV}(B_r(x), H^n)$ be area minimizers such that $u_k 1_{B_s(x)} \rightarrow u 1_{B_s(x)}$ in L^1 sense. There exists $c(K, n, v) > 0$ such that, setting*

$$c_r := ce^{cr^{-1}(u(x) - \inf_{y \in B_r(x)} u(y))},$$

it holds

$$\sup_{z \in B_{r/(15c_r)}(x)} \text{lip}(u)(z) \leq c_r.$$

Proof. Fix $r/2 < t < s$. Each function u_k is defined on $B_t(x_k)$ for k large enough and it is smooth on this ball. Moreover, thanks to Theorem 1.20, we have

$$\sup_{B_{t/6}^g(\bar{x}_k)} |\nabla u_k|(z) \leq ce^{ct^{-1}(u_k(x_k) - \inf_{y \in B_t(x_k)} u_k(y))}. \quad (8)$$

Observe now that the Kuratowski convergence of the graphs of u_k to the boundary of the epigraph of u in $B_s(x)$ implies that for every $\delta > 0$, if k is large enough, we have

$$\inf_{z \in B_t(x_k)} u_k(z) \geq \inf_{z \in B_s(x)} u(z) - \delta$$

and

$$u_k(x_k) \leq \sup_{y \in B_\delta(x)} u(y) + \delta.$$

Hence, for every $\delta > 0$ and for k large enough, the inequality (8) becomes

$$\sup_{B_{t/6}^g(\bar{x}_k)} |\nabla u_k|(z) \leq ce^{ct^{-1}(\sup_{y \in B_\delta(x)} u(y) - \inf_{y \in B_s(x)} u(y) + 2\delta)}. \quad (9)$$

We now set $c_\delta^t := ce^{ct^{-1}(\sup_{y \in B_\delta(x)} u(y) - \inf_{y \in B_s(x)} u(y) + 2\delta)}$ and we note that, by Lemma 4.3, it holds

$$\sup_{B_{t/(14c_\delta^t)}(x_k)} |\nabla u_k|(z) \leq c_\delta^t,$$

which, together with Lemma 4.2, implies that

$$\sup_{z \in B_{t/(15c_\delta^t)}(x)} \text{lip}(u)(z) \leq c_\delta^t. \quad (10)$$

This shows in particular that u is continuous in x , so that when we let $\delta \rightarrow 0$ we have that

$$\sup_{y \in B_\delta(x)} u(y) \rightarrow u(x).$$

Hence, taking the limit as $\delta \rightarrow 0$ in (10) and setting $c_0^t := ce^{ct^{-1}(u(x) - \inf_{y \in B_s(x)} u(y))}$, we get

$$\sup_{z \in B_{t/(15c_0^t)}(x)} \text{lip}(u)(z) \leq c_0^t \leq c_r := ce^{2cr^{-1}(u(x) - \inf_{y \in B_r(x)} u(y))}.$$

Now, taking the limit as $t \rightarrow r$ and using the same notation, we get

$$\sup_{z \in B_{r/(15c_r)}(x)} \text{lip}(u)(z) \leq c_r,$$

as claimed. \square

Corollary 4.5. *Let $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $\mathbf{H}_k^n(B_1(x_k)) \geq v > 0$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . Let $u_k \in \text{BV}(B_r(x), \mathbf{H}_k^n)$ and $u \in \text{BV}(B_r(x), \mathbf{H}^n)$ be area minimizers such that $u_k 1_{B_r(x)} \rightarrow u 1_{B_r(x)}$ in L^1 sense. Then $u \in \text{Lip}_{loc}(B_r(x))$.*

Proof. For every $y \in B_r(x)$ and every sequence $y_k \in M_k$ converging to y , we can find $\epsilon > 0$ and $v' > 0$ such that $\mathbf{H}_k^n(B_\epsilon(y_k)) \geq v' > 0$ for every k large enough (this follows from the doubling property of each of these manifolds and the uniform lower bound on the volume of the unit ball). Hence, the previous proposition guarantees that $u \in \text{Lip}(B_\delta(y))$ for $\delta > 0$ sufficiently small. By the arbitrariness of $y \in B_r(x)$ we conclude. \square

Theorem 5. *Let $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $H_k^n(B_1(x_k)) \geq v > 0$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . Let $r > 0$ be such that $B_r(x) \subsetneq X$ and let $u \in \text{BV}(B_r(x), \mathbb{H}^n)$ be an area minimizer.*

For every $s \in (0, r)$, there exists a sequence of area minimizers $u_k \in \text{BV}(B_s(x), \mathbb{H}_k^n)$ such that $u_k 1_{B_s(x)}$ converges in L_1 to $u 1_{B_s(x)}$ and $A(u_k, B_s(x), \mathbb{H}_k^n) \rightarrow A(u, B_s(x), \mathbb{H}^n)$.

Proof. By Theorem 3.3, there exists $0 < \delta < r - s$ and area minimizers $u_k \in \text{BV}(B_s(x), \mathbb{H}_k^n)$ and $\tilde{u} \in \text{BV}(B_{s+\delta}(x), \mathbb{H}^n)$, such that $u_k 1_{B_s(x)}$ converges in L_1 to $\tilde{u} 1_{B_s(x)}$ and $\{\tilde{u} \neq u\} \subset\subset B_{s+\delta}(x)$.

By the previous corollary, $\tilde{u} \in \text{Lip}_{loc}(B_{s+\delta}(x))$. By the corollary of Theorem 4, this implies $\tilde{u} = u$, so that $u_k 1_{B_s(x)}$ converges in L_1 to $u 1_{B_s(x)}$.

Moreover, since $|D^s \tilde{u}| \equiv 0$, Theorem 3.3 implies that

$$A(u_k, B_s(x), \mathbb{H}_k^n) \rightarrow A(\tilde{u}, B_s(x), \mathbb{H}^n) = A(u, B_s(x), \mathbb{H}^n).$$

This concludes the proof. \square

Theorem 6. *Let $r > 0$, $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $H_k^n(B_r(x_k)) \geq vr^n$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . If $u \in \text{BV}(B_r(x), \mathbb{H}^n)$ is an area minimizer, then there exists $c(K, n, v) > 0$ such that, setting*

$$c_r := ce^{cr^{-1}(u(x) - \inf_{y \in B_r(x)} u(y))},$$

it holds

$$\sup_{z \in B_{r/(15c_r)}(x)} \text{lip}(u)(z) \leq c_r.$$

Proof. The statement follows combining Theorem 5 and Proposition 4.4. \square

Theorem 7. *Let $v > 0$, $n \in \mathbb{N}$ and $K \in \mathbb{R}$ be fixed. Let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq K$ and $H_k^n(B_1(x_k)) \geq v > 0$ converging in pointed Gromov-Hausdorff sense to (X, \mathbf{d}, x) . If $\Omega \subset X$ is open and $u \in \text{BV}(\Omega, \mathbb{H}^n)$ is an area minimizer, then $u \in \text{Lip}_{loc}(\Omega)$.*

Proof. For every $y \in \Omega$ and every sequence $y_k \in M_k$ converging to y , we can find $\epsilon > 0$ such that $H_k^n(B_\epsilon(y_k)) \geq v > 0$ for every k large enough (this follows from the doubling property of each of these manifolds and the uniform lower bound on the volume of the unit ball). The statement then follows combining Theorem 5 and Corollary 4.5. \square

The next lemma, which will be used to prove Theorem 8, follows by repeating the argument of [22, Theorem 6.2], and for this reason some details in the proof are omitted.

Lemma 4.6. *Let $r_0 > 0$, $v > 0$, $n \in \mathbb{N}$. Let (M^n, \mathbf{d}, x) be a manifold with $\text{Ric}_{M^n} \geq 0$ and $H^n(B_{r_0}(x)) \geq vr_0^n$. If $u \in C^\infty(B_{2r_0}(x))$ is an area minimizer, then given $r > 0$ such that*

$$r^2 = r_0^2 - \left(\sup_{y \in B_r(x)} |u(y) - u(x)| \right)^2,$$

we have

$$\sup_{y \in B_r(x)} |\nabla u|(y) \leq c \left(1 + r^{-1} \sup_{y \in B_r(x)} |u(y) - u(x)| \right)^n,$$

where c depends only on n and v .

Proof. Thanks to Propositions 1.18 and 1.19, there exists a constant $c(n, v) > 0$ such that

$$\frac{1}{\mathbf{H}^n(B_{r_0}^g(\bar{x}))} \int_{B_{r_0}^g(\bar{x})} \frac{1}{\sqrt{1 + |\nabla u|^2}} d\mathbf{H}^n \leq c \inf_{y \in \pi(B_{r_0}^g(\bar{x}))} \frac{1}{\sqrt{1 + |\nabla u|^2}}(y),$$

where \mathbf{H}^n is the Hausdorff measure in $\mathbf{M} \times \mathbb{R}$ and we are using the previously defined convention for functions defined on \mathbf{M} evaluated on the graph of u .

Hence, possibly changing c at every step, we get

$$\sup_{y \in \pi(B_{r_0}^g(\bar{x}))} \sqrt{1 + |\nabla u|^2}(y) \leq c \frac{\mathbf{H}^n(B_{r_0}^g(\bar{x}))}{\mathbf{H}^n(\pi(B_{r_0}^g(\bar{x})))},$$

which, together with the fact that u is an area minimizer, implies that

$$\sup_{y \in \pi(B_{r_0}^g(\bar{x}))} \sqrt{1 + |\nabla u|^2}(y) \leq \frac{cr_0^n}{\mathbf{H}^n(\pi(B_{r_0}^g(\bar{x})))}.$$

By our hypothesis on $r > 0$ we then get that $B_r(x) \subset \pi(B_{r_0}^g(\bar{x}))$, while Bishop Gromov's inequality and the fact that $\mathbf{H}^n(B_{r_0}(x)) \geq vr_0^n$ imply that $\mathbf{H}^n(B_r(x)) \geq vr^n$. Putting these facts together we get, changing again $c > 0$,

$$\sup_{y \in \pi(B_{r_0}^g(\bar{x}))} \sqrt{1 + |\nabla u|^2}(y) \leq c \frac{r_0^n}{r^n} \leq c \left(1 + r^{-2} \sup_{y \in B_r(x)} |u(y) - u(x)|^2 \right)^{n/2},$$

which then implies the statement. \square

Theorem 8. Let $n \in \mathbb{N}$ be fixed and let $(\mathbf{M}_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{\mathbf{M}_k^n} \geq 0$ converging in pointed Gromov-Hausdorff sense to a space $(\mathbf{X}, \mathbf{d}, x)$ such that

$$\lim_{r \rightarrow +\infty} \frac{\mathbf{H}^n(B_r(x))}{r^n} \geq v > 0.$$

If $u \in \text{BV}_{loc}(\mathbf{X}, \mathbf{H}^n)$ is an area minimizer, then there exists $c(n, v) > 0$ such that, for every $r > 0$, it holds

$$\sup_{y \in B_{r/2}(x)} \text{lip}(u)(y) \leq c \left(1 + r^{-1} \sup_{y \in B_r(x)} |u(y) - u(x)| \right)^n.$$

Proof. Fix $0 < R_1 < R_2$ and consider the sequence of area minimizers $u_k \in \text{BV}(B_{R_2}(x), \mathbf{H}_k^n)$ converging in L^1 to u on $B_{R_2}(x)$ given by Theorem 5. Each function u_k is defined on $B_{R_1}(x_k)$ for k large enough and is smooth on this ball.

Consider then $r > 0$ such that

$$r^2 = R_1^2 - \left(\sup_{y \in B_r(x)} |u(y) - u(x)| \right)^2.$$

Since u is locally Lipschitz by Theorem 7, the Kuratowski convergence of the epigraphs of u_k to the epigraph of u implies that

$$\sup_{y \in B_r(x_k)} |u_k(y) - u_k(x_k)| \rightarrow \sup_{y \in B_r(x)} |u(y) - u(x)|.$$

In particular, setting $r_k > 0$ to be such that

$$r_k^2 = R_1^2 - \left(\sup_{y \in B_{r_k}(x_k)} |u_k(y) - u_k(x_k)| \right)^2,$$

we have that $r_k \rightarrow r$ as $k \rightarrow \infty$.

Moreover, combining Bishop Gromov's inequality on X with Lemma 4.1, we obtain that, for every k large enough, we have

$$\mathbf{H}_k^n(B_r(x_k)) \geq \frac{v}{2} r^n,$$

so that, by Lemma 4.6, we obtain

$$\sup_{y \in B_{r_k}(x)} |\nabla u_k|(y) \leq c(n, v) \left(1 + r_k^{-1} \sup_{y \in B_{r_k}(x_k)} |u_k(y) - u_k(x_k)| \right)^n.$$

Moreover, using Lemma 4.2 and the fact that $r_i \rightarrow r$, we get

$$\sup_{y \in B_{r/2}(x)} |\nabla u|(y) \leq c \left(1 + r^{-1} \sup_{y \in B_r(x)} |u(y) - u(x)| \right)^n. \quad (11)$$

Finally, the function $R_1 \mapsto r$ is increasing (by definition), continuous (since u is continuous), and tends to $+\infty$ at infinity (since u is locally bounded), so that (11) holds for every $r > 0$ by the arbitrariness of R_1 . \square

The next lemma will be used to prove Theorem 9 from the Introduction. The proof is adapted from the one in [23, beginning of Theorem 3.6].

Lemma 4.7. *Let (X, d, \mathbf{m}, x) be an $\text{RCD}(0, N)$ space. If $u \in \text{Lip}(X)$ is a function whose epigraph minimizes the perimeter and such that*

$$\lim_{y \rightarrow +\infty} \frac{\max\{-u(y), 0\}}{d(x, y)} = 0, \quad (12)$$

then

$$\lim_{y \rightarrow +\infty} \frac{|u(y)|}{d(x, y)} = 0.$$

Proof. Let $L := \mathbf{L}(u) \vee 1$, where $\mathbf{L}(u)$ is the Lipschitz constant of u . The Harnack inequality on the graph of u (Theorem 1.17) gives that, for every $r > 0$, it holds

$$\sup_{y \in B_r^g(\bar{x})} u(y) - \inf_{z \in B_r^g(\bar{x})} u(z) \leq C(u(x) - \inf_{z \in B_r^g(\bar{x})} u(z)),$$

so that

$$\sup_{y \in B_r^g(\bar{x})} u(y) \leq Cu(x) - (C - 1) \inf_{z \in B_r^g(\bar{x})} u(z).$$

Using that u is Lipschitz, we get

$$\sup_{y \in B_{r/(2L)}(x)} u(y) \leq Cu(x) - (C - 1) \inf_{z \in B_r(x)} u(z).$$

Moreover, from (12), for every $\delta > 0$ there exists $C_\delta > 0$ such that $u(y) \geq -C_\delta - \delta d(x, y)$ for every $y \in X$. Hence,

$$\sup_{y \in B_{r/(2L)}(x)} u(y) \leq Cu(x) + (C - 1)(C_\delta + \delta r).$$

This implies that

$$\limsup_{y \rightarrow +\infty} \frac{|u(y)|}{d(x, y)} \leq 2CL\delta,$$

which implies the statement by the arbitrariness of $\delta > 0$. \square

The proof of the next result, i.e. Theorem 9, is adapted from the one in [23, Theorem 3.6].

Theorem 9. Let $n \in \mathbb{N}$ be fixed and let $(M_k^n, \mathbf{d}_k, x_k)$ be a sequence of pointed Riemannian manifolds with $\text{Ric}_{M_k^n} \geq 0$ converging in pointed Gromov-Hausdorff sense to a space (X, \mathbf{d}, x) such that

$$\lim_{r \rightarrow +\infty} \frac{\mathbf{H}^n(B_r(x))}{r^n} \geq v > 0.$$

If $u \in \text{BV}_{loc}(X, \mathbf{H}^n)$ is an area minimizer such that

$$\lim_{y \rightarrow +\infty} \frac{\max\{-u(y), 0\}}{\mathbf{d}(x, y)} = 0,$$

then u is constant.

Proof. By Lemma 4.1 and Theorem 6 we know that $u \in \text{Lip}(X)$. Hence the previous lemma gives that

$$\lim_{y \rightarrow +\infty} \frac{|u(y)|}{\mathbf{d}(x, y)} = 0.$$

We claim that $|\nabla u| = 0$ in $B_1(x)$. A rescaling argument then gives $|\nabla u| = 0$ in $B_r(x)$ for every $r > 0$, yielding the statement.

To this aim, we fix $\epsilon > 0$ and we choose $R > 1$ large enough so that, for every $y \in X$ with $\mathbf{d}(x, y) > R$, it holds

$$|u(y)| \leq \epsilon \mathbf{d}(x, y).$$

Consider $R_1 > R$ to be fixed later and the minimizers $u_k \in C^\infty(B_{5R_1}(x) \cap M_k)$ converging in L^1 to u on $B_{5R_1}(x)$. For every k large enough, since the epigraphs of u_k converge in Kuratowski sense to the epigraph of u in $B_{5R_1}(x) \times \mathbb{R}$, we have that

$$|u_k(y)| \leq 2\epsilon \mathbf{d}_k(x, y) \tag{13}$$

for every $y \in B_{R_1}(x_k)$ such that $\mathbf{d}_k(x_k, y) > 2R$.

So far we estimated the growth of the functions u_k exploiting the growth condition on u ; now we estimate their Lipschitz constants. For every $s > 0$ let c_s be the quantity defined in Theorem 6 relative to u and let c_s^k denote the same quantity relative to each u_k . Note that by our assumption on the growth of u , we have that there exists $c > 1$ such that $c_s \leq c$ for every s .

Since the epigraphs of the functions u_k converge in Kuratowski sense to the one of u , we have that $c_{4R_1}^k \rightarrow c_{4R_1}$, so that by the a-priori gradient estimate Theorem 6, for k large enough,

$$\sup_{z \in B_{3R_1/(16c)}(x_k)} |\nabla u_k|(z) \leq 2c. \tag{14}$$

To shorten the notation we set $R_2 := 3R_1/(16c)$. A posteriori, since (14) holds for every R_1 large enough, we see that we could have chosen $R_1 > 0$ (large enough) so that

$$B_{2R}(x_k) \subset \pi(B_{R_2}^g(\bar{x}_k)). \tag{15}$$

We now denote by \mathbf{H}_k^n and ∇_\times respectively the Hausdorff measure and the gradient in $M_k \times \mathbb{R}$ w.r.t. the product distance, and by ∇_g the gradient on the graph of u_k . We claim that

$$\int_{B_{R_2}^g(\bar{x}_k)} |\nabla_g u_k|^2 d\mathbf{H}_k^n \leq 4\epsilon^2 \mathbf{H}_k^n(B_{2R_2}^g(\bar{x}_k)). \tag{16}$$

To prove the claim, we consider $\eta \in \text{Lip}(B_{2R_2}^\times(\bar{x}_k))$ with $\eta = 0$ on $\partial B_{2R_2}^\times(\bar{x}_k)$, $\eta = 1$ on $B_{R_2}^\times(\bar{x}_k)$ and $\nabla_\times \eta \leq 1/R_2$. Using the fact that u_k is harmonic on its graph, we then get

$$\int_{B_{R_2}^g(\bar{x}_k)} |\nabla_g u_k|^2 d\mathbf{H}_k^n \leq \int_{B_{2R_2}^g(\bar{x}_k)} |\nabla_g u_k|^2 \eta d\mathbf{H}_k^n \leq 4 \int_{B_{2R_2}^g(\bar{x}_k)} u_k^2 |\nabla_g \eta|^2 d\mathbf{H}_k^n.$$

Moreover, combining (13), (15) and the fact that $|\nabla_g \eta| \leq |\nabla_\times \eta|$, we deduce that

$$4 \int_{B_{2R_2}^g(\bar{x}_k)} u_k^2 |\nabla_g \eta|^2 d\mathbf{H}_k^n \leq 4\epsilon^2 \mathbf{H}_k^n(B_{2R_2}^g(\bar{x}_k)),$$

from which our claim (16) follows.

Finally, using the mean value property for sub-harmonic function on the graph of each u_k (i.e. Proposition 1.19) we get, thanks to (14) and (16), that

$$\begin{aligned} \sup_{B_2(x_k)} |\nabla u_k|^2 &\leq \sup_{B_{R_2}^g(x_k)} |\nabla u_k|^2 \leq \int_{B_{R_2}^g(x_k)} |\nabla u_k|^2 d\mathbf{H}_k^n \\ &\leq (1 + 4c^2) \int_{B_{R_2}^g(x_k)} |\nabla_g u_k|^2 d\mathbf{H}_k^n \leq 4\epsilon^2(1 + 4c^2) \frac{\mathbf{H}_k^n(B_{2R_2}^g(\bar{x}_k))}{\mathbf{H}_k^n(B_{R_2}^g(\bar{x}_k))} \leq C(u, n)\epsilon^2. \end{aligned}$$

In particular, by Lemma 4.2, we get

$$\sup_{B_1(x)} |\nabla u|^2 \leq C\epsilon^2.$$

Since $\epsilon > 0$ was arbitrary we conclude. \square

References

- [1] Luigi Ambrosio. “Calculus, heat flow and curvature-dimension bounds in metric measure spaces”. In: *Proceedings of the International Congress of Mathematicians, Rio de Janeiro 2018. Vol. I. Plenary lectures*. World Sci. Publ., Hackensack, NJ, 2018, pp. 301–340.
- [2] Luigi Ambrosio, Elia Brué, and Daniele Semola. “Rigidity of the 1-Bakry-Émery inequality and sets of finite perimeter in RCD spaces”. In: *Geom. Funct. Anal.* 29.4 (2019), pp. 949–1001. ISSN: 1016-443X,1420-8970. DOI: [10.1007/s00039-019-00504-5](https://doi.org/10.1007/s00039-019-00504-5). URL: <https://doi.org/10.1007/s00039-019-00504-5>.
- [3] Luigi Ambrosio, Maria Colombo, and Simone Di Marino. “Sobolev spaces in metric measure spaces: reflexivity and lower semicontinuity of slope”. In: *Variational methods for evolving objects*. Vol. 67. Adv. Stud. Pure Math. Math. Soc. Japan, [Tokyo], 2015, pp. 1–58. ISBN: 978-4-86497-028-0. DOI: [10.2969/aspm/06710001](https://doi.org/10.2969/aspm/06710001). URL: <https://doi.org/10.2969/aspm/06710001>.
- [4] Luigi Ambrosio and Simone Di Marino. “Equivalent definitions of BV space and of total variation on metric measure spaces”. In: *J. Funct. Anal.* 266.7 (2014), pp. 4150–4188. ISSN: 0022-1236,1096-0783. DOI: [10.1016/j.jfa.2014.02.002](https://doi.org/10.1016/j.jfa.2014.02.002). URL: <https://doi.org/10.1016/j.jfa.2014.02.002>.
- [5] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000, pp. xviii+434. ISBN: 0-19-850245-1.
- [6] Luigi Ambrosio, Nicola Gigli, Andrea Mondino, and Tapio Rajala. “Riemannian Ricci curvature lower bounds in metric measure spaces with σ -finite measure”. In: *Trans. Amer. Math. Soc.* 367.7 (2015), pp. 4661–4701. ISSN: 0002-9947,1088-6850. DOI: [10.1090/S0002-9947-2015-06111-X](https://doi.org/10.1090/S0002-9947-2015-06111-X). URL: <https://doi.org/10.1090/S0002-9947-2015-06111-X>.
- [7] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. “Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below”. In: *Invent. Math.* 195.2 (2014), pp. 289–391. ISSN: 0020-9910,1432-1297. DOI: [10.1007/s00222-013-0456-1](https://doi.org/10.1007/s00222-013-0456-1). URL: <https://doi.org/10.1007/s00222-013-0456-1>.

- [8] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. “Metric measure spaces with Riemannian Ricci curvature bounded from below”. In: *Duke Math. J.* 163.7 (2014), pp. 1405–1490. ISSN: 0012-7094,1547-7398. DOI: [10.1215/00127094-2681605](https://doi.org/10.1215/00127094-2681605). URL: <https://doi.org/10.1215/00127094-2681605>.
- [9] Luigi Ambrosio and Shouhei Honda. “New stability results for sequences of metric measure spaces with uniform Ricci bounds from below”. In: *Measure theory in non-smooth spaces*. Partial Differ. Equ. Meas. Theory. De Gruyter Open, Warsaw, 2017, pp. 1–51.
- [10] Luigi Ambrosio, Andrea Mondino, and Giuseppe Savaré. “Nonlinear Diffusion Equations and Curvature Conditions in Metric Measure Spaces”. In: *Memoirs of the American Mathematical Society* 262 (Sept. 2015). DOI: [10.1090/memo/1270](https://doi.org/10.1090/memo/1270).
- [11] Luigi Ambrosio, Andrea Pinamonti, and Gareth Speight. “Tensorization of Cheeger energies, the space $H^{1,1}$ and the area formula for graphs”. In: *Adv. Math.* 281 (2015), pp. 1145–1177. ISSN: 0001-8708,1090-2082. DOI: [10.1016/j.aim.2015.06.004](https://doi.org/10.1016/j.aim.2015.06.004). URL: <https://doi.org/10.1016/j.aim.2015.06.004>.
- [12] Gioacchino Antonelli, Camillo Brena, and Enrico Pasqualetto. “Subgraphs of BV functions on RCD spaces”. In: *Ann. Global Anal. Geom.* 65.2 (2024), Paper No. 14, 26. ISSN: 0232-704X,1572-9060. DOI: [10.1007/s10455-024-09945-0](https://doi.org/10.1007/s10455-024-09945-0). URL: <https://doi.org/10.1007/s10455-024-09945-0>.
- [13] Fabio Cavalletti and Emanuel Milman. “The globalization theorem for the curvature-dimension condition”. In: *Invent. Math.* 226.1 (2021), pp. 1–137. ISSN: 0020-9910,1432-1297. DOI: [10.1007/s00222-021-01040-6](https://doi.org/10.1007/s00222-021-01040-6). URL: <https://doi.org/10.1007/s00222-021-01040-6>.
- [14] J. Cheeger. “Differentiability of Lipschitz functions on metric measure spaces”. In: *Geom. Funct. Anal.* 9.3 (1999), pp. 428–517. ISSN: 1016-443X,1420-8970. DOI: [10.1007/s000390050094](https://doi.org/10.1007/s000390050094). URL: <https://doi.org/10.1007/s000390050094>.
- [15] Giulio Colombo, Marco Magliaro, Luciano Mari, and Marco Rigoli. “Bernstein and half-space properties for minimal graphs under Ricci lower bounds”. In: *Int. Math. Res. Not. IMRN* 23 (2022), pp. 18256–18290. ISSN: 1073-7928,1687-0247. DOI: [10.1093/imrn/rnab342](https://doi.org/10.1093/imrn/rnab342). URL: <https://doi.org/10.1093/imrn/rnab342>.
- [16] Giulio Colombo, Luciano Mari, and Marco Rigoli. “On minimal graphs of sublinear growth over manifolds with non-negative Ricci curvature”. 2023. arXiv: [2310.15620](https://arxiv.org/abs/2310.15620) [math.DG].
- [17] Alessandro Cucinotta. “Minimal Surface Equation and Bernstein Property on RCD spaces”. 2024. arXiv: [2403.02406](https://arxiv.org/abs/2403.02406) [math.DG].
- [18] Alessandro Cucinotta and Andrea Mondino. “Half Space Property in RCD(0,N) spaces”. 2024. arXiv: [2402.12230](https://arxiv.org/abs/2402.12230) [math.DG].
- [19] Guido De Philippis and Nicola Gigli. “Non-collapsed spaces with Ricci curvature bounded from below”. In: *J. Éc. polytech. Math.* 5 (2018), pp. 613–650. ISSN: 2429-7100,2270-518X. DOI: [10.5802/jep.80](https://doi.org/10.5802/jep.80). URL: <https://doi.org/10.5802/jep.80>.
- [20] Qi Ding. “Capacity for minimal graphs over manifolds and the half-space property”. 2023. arXiv: [2306.15137](https://arxiv.org/abs/2306.15137) [math.DG].
- [21] Qi Ding. “Liouville-type theorems for minimal graphs over manifolds”. In: *Anal. PDE* 14.6 (2021), pp. 1925–1949. ISSN: 2157-5045,1948-206X. DOI: [10.2140/apde.2021.14.1925](https://doi.org/10.2140/apde.2021.14.1925). URL: <https://doi.org/10.2140/apde.2021.14.1925>.
- [22] Qi Ding. “Poincaré inequality on minimal graphs over manifolds and applications”. 2023. arXiv: [2111.04458](https://arxiv.org/abs/2111.04458) [math.DG].
- [23] Qi Ding, Jürgen Jost, and Yuanlong Xin. “Minimal graphic functions on manifolds of non-negative Ricci curvature”. In: *Comm. Pure Appl. Math.* 69.2 (2016), pp. 323–371. ISSN: 0010-3640,1097-0312. DOI: [10.1002/cpa.21566](https://doi.org/10.1002/cpa.21566). URL: <https://doi.org/10.1002/cpa.21566>.
- [24] Matthias Erbar, Kazumasa Kuwada, and Karl-Theodor Sturm. “On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces”. In: *Inventiones mathematicae* 201 (Dec. 2014). DOI: [10.1007/s00222-014-0563-7](https://doi.org/10.1007/s00222-014-0563-7).

- [25] Nicola Gigli. “On the differential structure of metric measure spaces and applications”. In: *Mem. Amer. Math. Soc.* 236.1113 (2015), pp. vi+91. ISSN: 0065-9266,1947-6221. DOI: [10.1090/memo/1113](https://doi.org/10.1090/memo/1113). URL: <https://doi.org/10.1090/memo/1113>.
- [26] Nicola Gigli and Bang-Xian Han. “Independence on p of weak upper gradients on RCD spaces”. In: *J. Funct. Anal.* 271.1 (2016), pp. 1–11. ISSN: 0022-1236,1096-0783. DOI: [10.1016/j.jfa.2016.04.014](https://doi.org/10.1016/j.jfa.2016.04.014). URL: <https://doi.org/10.1016/j.jfa.2016.04.014>.
- [27] Nicola Gigli, Andrea Mondino, and Giuseppe Savaré. “Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows”. In: *Proceedings of the London Mathematical Society* 111 (Nov. 2013). DOI: [10.1112/plms/pdv047](https://doi.org/10.1112/plms/pdv047).
- [28] Enrico Giusti. *Minimal surfaces and functions of bounded variation*. Vol. 80. Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984, pp. xii+240. ISBN: 0-8176-3153-4. DOI: [10.1007/978-1-4684-9486-0](https://doi.org/10.1007/978-1-4684-9486-0). URL: <https://doi.org/10.1007/978-1-4684-9486-0>.
- [29] Heikki Hakkarainen, Juha Kinnunen, and Panu Lahti. “Regularity of minimizers of the area functional in metric spaces”. In: *Adv. Calc. Var.* 8.1 (2015), pp. 55–68. ISSN: 1864-8258,1864-8266. DOI: [10.1515/acv-2013-0022](https://doi.org/10.1515/acv-2013-0022). URL: <https://doi.org/10.1515/acv-2013-0022>.
- [30] Heikki Hakkarainen, Juha Kinnunen, Panu Lahti, and Pekka Lehtelä. “Relaxation and integral representation for functionals of linear growth on metric measure spaces”. In: *Anal. Geom. Metr. Spaces* 4.1 (2016), pp. 288–313. ISSN: 2299-3274. DOI: [10.1515/agms-2016-0013](https://doi.org/10.1515/agms-2016-0013). URL: <https://doi.org/10.1515/agms-2016-0013>.
- [31] Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, and Jeremy T. Tyson. *Sobolev spaces on metric measure spaces: An approach based on upper gradients*. Vol. 27. New Mathematical Monographs. Cambridge University Press, Cambridge, 2015, pp. xii+434. ISBN: 978-1-107-09234-1. DOI: [10.1017/CB09781316135914](https://doi.org/10.1017/CB09781316135914). URL: <https://doi.org/10.1017/CB09781316135914>.
- [32] Juha Kinnunen, Riikka Korte, Andrew Lorent, and Nageswari Shanmugalingam. “Regularity of sets with quasiminimal boundary surfaces in metric spaces”. In: *J. Geom. Anal.* 23.4 (2013), pp. 1607–1640. ISSN: 1050-6926,1559-002X. DOI: [10.1007/s12220-012-9299-z](https://doi.org/10.1007/s12220-012-9299-z). URL: <https://doi.org/10.1007/s12220-012-9299-z>.
- [33] Zhenhao Li. “The globalization theorem for $CD(K, N)$ on locally finite spaces”. In: *Ann. Mat. Pura Appl. (4)* 203.1 (2024), pp. 49–70. ISSN: 0373-3114,1618-1891. DOI: [10.1007/s10231-023-01352-9](https://doi.org/10.1007/s10231-023-01352-9). URL: <https://doi.org/10.1007/s10231-023-01352-9>.
- [34] John Lott and Cédric Villani. “Ricci curvature for metric-measure spaces via optimal transport”. In: *Ann. of Math. (2)* 169.3 (2009), pp. 903–991. ISSN: 0003-486X,1939-8980. DOI: [10.4007/annals.2009.169.903](https://doi.org/10.4007/annals.2009.169.903). URL: <https://doi.org/10.4007/annals.2009.169.903>.
- [35] Michele Miranda Jr. “Functions of bounded variation on “good” metric spaces”. In: *J. Math. Pures Appl. (9)* 82.8 (2003), pp. 975–1004. ISSN: 0021-7824. DOI: [10.1016/S0021-7824\(03\)00036-9](https://doi.org/10.1016/S0021-7824(03)00036-9). URL: [https://doi.org/10.1016/S0021-7824\(03\)00036-9](https://doi.org/10.1016/S0021-7824(03)00036-9).
- [36] Andrea Mondino and Daniele Semola. “Weak Laplacian bounds and minimal boundaries in non-smooth spaces with Ricci curvature lower bounds”. Accepted in: *Memoirs of the American Mathematical Society*, Preprint arXiv: 2107.12344.
- [37] Tapio Rajala. “Local Poincaré inequalities from stable curvature conditions on metric spaces”. In: *Calc. Var. Partial Differential Equations* 44.3-4 (2012), pp. 477–494. ISSN: 0944-2669,1432-0835. DOI: [10.1007/s00526-011-0442-7](https://doi.org/10.1007/s00526-011-0442-7). URL: <https://doi.org/10.1007/s00526-011-0442-7>.
- [38] Karl-Theodor Sturm. “On the geometry of metric measure spaces. I”. In: *Acta Math.* 196.1 (2006), pp. 65–131. ISSN: 0001-5962,1871-2509. DOI: [10.1007/s11511-006-0002-8](https://doi.org/10.1007/s11511-006-0002-8). URL: <https://doi.org/10.1007/s11511-006-0002-8>.
- [39] Karl-Theodor Sturm. “On the geometry of metric measure spaces. II”. In: *Acta Math.* 196.1 (2006), pp. 133–177. ISSN: 0001-5962,1871-2509. DOI: [10.1007/s11511-006-0003-7](https://doi.org/10.1007/s11511-006-0003-7). URL: <https://doi.org/10.1007/s11511-006-0003-7>.