QUANTUM-CLASSICAL MOTION OF CHARGED PARTICLES INTERACTING WITH SCALAR FIELDS

SHAHNAZ FARHAT

ABSTRACT. The goal of this article is to investigate the dynamics of semi-relativistic or nonrelativistic charged particles in interaction with a scalar meson field. Our main contribution is the derivation of the classical dynamics of a particle-field system as an effective equation of the quantum microscopic Nelson model, in the classical limit where the value of the Planck constant approaches zero ($\hbar \rightarrow 0$). Thus, we prove the validity of Bohr's correspondence principle, that is to establish the transition from quantum to classical dynamics. We use a Wigner measure approach to study such transition. Then, as a consequence of this interplay between classical and quantum dynamics, we establish the global well-posedness of the classical particle-field interacting system, despite the low regularity of the related vector field, which prevents the use of a fixed point argument.

KEYWORDS. Particle-field equation; Nelson model; Semi-classical analysis; Probabilistic representations; Measure theoritical techniques; Wigner measures; Density matrices; Liouville equations.

Contents

1. Introduction	2
1.1. General framework	3
1.2. Assumptions and main results	6
2. The classical system	7
2.1. The interaction representation	7
2.2. Properties of the particle-field equation	9
3. The quantum system	12
3.1. Quantum estimates	12
3.2. Self-adjointness of Nelson Hamiltonian	15
3.3. The dynamical equation	16
4. Existence of Wigner measure	21
4.1. Propagation of assumptions	21
4.2. Existence of unique Wigner measure	29
5. Derivation of the characteristic equations	31
5.1. Convergence	32
5.2. The characteristic equation	34
5.3. The Liouville equation	35
6. Proof of the main result	36
Appendix A. Prokhorov theorem	39
Appendix B. Useful results	39
Appendix C. Technical results about convergence	40
References	42

Date: May 22, 2024.

1. INTRODUCTION

Classical and quantum mechanics may initially appear to be fundamentally different, as classical mechanics deals with the trajectories of particles while quantum mechanics focuses on wave functions evolution. Furthermore, quantum mechanics are successful in describing microscopic objects, whereas macroscopic systems are better described by classical theories like classical mechanics and classical electrodynamics. The point at which quantum and classical physics are in accordance is known as the correspondence limit, or the classical limit. The Correspondence Principle has been introduced to quantum theory in 1920 by Niels Bohr. Such principle emphasizes the importance of understanding the relationship between the two theories and how they converge in specific scaling limits. Bohr proposed that as the quantum numbers increase, the system behaves more classically and the predictions of quantum mechanics align with those of classical mechanics. In the mathematical physics literature, the Bohr's principle is discussed in different frameworks (quantum mechanics, many-body theory, quantum field theory); and rigorously proved using mainly the Hepp's method [21]. However, this method applies only to a specific selection of density matrices, namely coherent states. In this article, we explore this principle by studying the convergence from quantum to classical dynamics in a particle-field interaction model as the value of the Planck constant, denoted by \hbar , approaches zero (see also [16, 20]).

On the other hand, the interaction between matter and fields has been a subject of great interest in recent decades. Here, the focus is on exploring the dynamics of charged particles and a scalar meson field interacting according to the Yukawa theory. Recall that the Yukawa theory models the strong nuclear force as an interaction between nucleons (non-relativistic or semi-relativistic particles) and mesons (fields). It is known that, despite the ability of classical mechanics in resolving many physical problems, there are still some phenomena that can not be explained by classical laws alone. Here, the low regularity of the vector field associated to the interacting system makes it difficult to construct global solutions using standard arguments. To overcome this issue, we use the quantum-classical transition of the Yukawa theory. Then, by employing transition, it becomes possible to construct global solutions for the classical interacting system. Another recent study [6] has also explored this type of convergence for non-relativistic particles interacting with the electro-magnetic field, considering the Pauli-Fierz Hamiltonian which converges to the Newton-Maxwell equation.

From a classical standpoint, in our case the dynamics are governed by a particle-field equation (1.1), also known as Newton-Klein-Gordon equation, which is a nonlinear system of coupled PDE-ODEs. Previous studies have examined this type of equation, as demonstrated in the articles [24, 25, 26]. These works focus on analyzing the long-term behavior of the solutions to a particle-field equation. Specifically, the authors of these articles introduce a form factor within the interacting system to ensure that the Hamiltonian remains bounded from below, and they assume that this form factor is compactly supported. In our investigation, we adopt a more general framework by imposing less restrictions on this form factor.

From a quantum standpoint, the time evolution is generated by the so-called Nelson Hamiltonian (1.5). The Nelson model was first introduced by Edward Nelson in [27, 28] to describe the interaction between particles (nucleons) and meson field (strong nuclear force). The Nelson model has been widely studied by many researchers, and a selection of relevant articles includes [1, 2, 4, 12, 13, 14, 15, 17, 18, 19, 22, 23, 30].

Our objectives are then:

- Proving the validity of Bohr's correspondence principle. More precisely, we want to establish a relationship between quantum and classical dynamics by showing that Nelson model reduces to the classical particle-field equation in the classical limit $\hbar \to 0$;
- Establishing the global well-posedness of a particle-field equation under weak assumptions on the form factor χ (see (1.6)) and on the potential V (see (1.7)).

The quantum dynamics have a well-defined global behavior. Our method involves transferring certain quantum regularization effects to the classical dynamics. This leads to the derivation of the classical dynamics of the particle-field system as an effective equation of a quantum microscopic dynamical system in the limit $\hbar \to 0$.

To achieve this scenario, we investigate the transition using Wigner measures approach in infinite dimensional bosonic quantum field theory. In recent years, this Wigner measure method have been used in many-body theory [10] and in quantum field theory [5] with an a priori knowledge of global well-posedness (GWP) for effective equations. Whereas in this work, our strategy furnishes global well-posedness and convergence at the same time. Usually, this convergence is non-trivial, and there is no prior guarantee of obtaining unique limits. However, we overcome this difficulty by relying on our assumptions.

The main results are the classical limit (Theorem 1.1) and the global well-posedness of a particle field equation (Theorem 1.3). To prove these outcomes, our strategy is summarized in the steps below:

- We first extract the quantum dynamical system using the family of density matrices $(\rho_{\hbar})_{\hbar}$ satisfying (1.8) and (1.9);
- Then, after proving the propagation (uniformly in any compact time interval) of the two uniform estimates (1.8) and (1.9), we take the limit to obtain the classical dynamics on the inverse Fourier transform of the Wigner measure. This results in a specific classical equation which is equivalent to a statistical Liouville equation, thanks to the regularities associated with the Wigner measure and vector field in this context;
- We employ then measure-theoretical methods [8, 9, 6] which provides us with the almost sure existence of global solutions. This requires us to prove the uniqueness of the solutions to a particle-field equation by using classical tools;
- Finally, we extend the existence result to all initial data. It is important to note, however, that the associated flow is Borel measurable with respect to initial data and may not be continuous.

1.1. General framework. This section provides a concrete mathematical description of the previous introduction. From a classical perspective, the dynamics are governed by a particlefield equation, as detailed in Paragraph 1.1.1. From a quantum perspective, the dynamics are governed by the Nelson Hamiltonian, which is explained in Paragraph 1.1.2.

1.1.1. The particle-field equation. Consider n fixed number of classical particles in the configuration space \mathbf{R}^d with $d \in \mathbf{N}^*$, interacting with field. Let M_j be the mass of the jth particle. The dynamic of the particles is characterized by their momenta $p_j \in \mathbf{R}^d$ and their positions $q_i \in \mathbf{R}^d$. Whereas, the field is described by $\alpha : \mathbf{R}^d \to \mathbf{C}$. Let $p = (p_1, \cdots, p_n), q = (q_1, \cdots, q_n)$ and $f_i : \mathbf{R}^d \longrightarrow \mathbf{R}$, the Hamiltonian of the particle-field system is

$$H(p,q,\alpha) = \sum_{j=1}^{n} f_j(p_j) + V(q_1,\cdots,q_n) + \int_{\mathbf{R}^d} \overline{\alpha(k)} \,\omega(k) \,\alpha(k) \,dk$$
$$+ \sum_{j=1}^{n} \int_{\mathbf{R}^d} \frac{\chi(k)}{\sqrt{\omega(k)}} \left[\alpha(k)e^{2\pi i k \cdot q_j} + \overline{\alpha(k)}e^{-2\pi i k \cdot q_j}\right] dk.$$

We consider two cases:

- Choosing $f_j(p_j) = \sqrt{p_j^2 + M_j^2}$ corresponds to the semi-relativistic case. Choosing $f_j(p_j) = p_j^2/2M_j$ corresponds to the non-relativistic case.

The parameter ω represents the dispersion relation defined by $\omega(k) = \sqrt{k^2 + m_f^2} \ge m_f > 0$, where m_f is the mass of the meson field. The function $V: \mathbf{R}^{dn} \to \mathbf{R}$ represents the external potential and $\chi : \mathbf{R}^d \to \mathbf{R}$ is the form factor. The equation of motion for the particle-field system is given by

$$\begin{cases} \partial_t p_j = -\frac{\partial H}{\partial q_j} = -\nabla_{q_j} V(q) - \int_{\mathbf{R}^d} 2\pi i k \frac{\chi(k)}{\sqrt{\omega(k)}} \left[\alpha(k) e^{2\pi i k \cdot q_j} - \overline{\alpha(k)} e^{-2\pi i k \cdot q_j} \right] dk; \\ \partial_t q_j = \frac{\partial H}{\partial p_j} = \nabla f_j(p_j); \\ i \partial_t \alpha = \frac{\partial H}{\partial \overline{\alpha}} = \omega(k) \ \alpha(k) + \sum_{j=1}^n \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot q_j}. \end{cases}$$
(1.1)

To clarify, the interaction term between particles and the scalar field has a specific form which is: linear in the field (for both semi and non-relativistic case); and in the momentum (only in the non-relativistic case). The solution $u = (p, q, \alpha)$ to (1.1) belongs to the following classical space

 $X^{\sigma} := \mathbf{R}^{dn} \times \mathbf{R}^{dn} \times \mathcal{G}^{\sigma}.$

where \mathcal{G}^{σ} with $\sigma \geq 0$ is the weighted L^2 lebesgue space endowed with the following norm

$$\|\alpha\|_{\mathcal{G}^{\sigma}}^{2} := \langle \alpha, \omega(\cdot)^{2\sigma} | \alpha \rangle_{L^{2}} = \int_{\mathbf{R}^{d}} \omega(k)^{2\sigma} |\alpha(k)|^{2} dk = \|\omega^{\sigma} | \alpha \|_{L^{2}}^{2}.$$

We have then for $u = (p, q, \alpha) \in X^{\sigma}$ the following norm

$$||u||_{X^{\sigma}}^{2} := \sum_{j=1}^{n} (|q_{j}|^{2} + |p_{j}|^{2}) + ||\alpha||_{\mathcal{G}^{\sigma}}^{2}.$$

The form factor serves as a way to term the interaction between particles and the field, by smoothing out the Hamiltonian and ensuring that it is bounded from below under certain assumptions. The magnitude of the coupling between the particles and the field is controlled by the form factor. We consider the energy space where the Hamiltonian is well-defined, namely $X^{1/2}$, but our main results are stated in the spaces X^{σ} with $\sigma \in [\frac{1}{2}, 1]$.

1.1.2. The Nelson model. The particle-field equation can be formally quantized to obtain the Nelson model. The Hilbert space of the quantized particle-field system is

$$\mathcal{H} := L^2(\mathbf{R}_x^{dn}, \mathbf{C}) \otimes \Gamma_s(L^2(\mathbf{R}_k^d, \mathbf{C})),$$

where $\Gamma_s(L^2(\mathbf{R}_k^d, \mathbf{C}))$ is the symmetric Fock space which could be identified with

$$\Gamma_s(L^2(\mathbf{R}^d_k, \mathbf{C})) := \bigoplus_{m=0}^{+\infty} L^2(\mathbf{R}^d, \mathbf{C})^{\bigotimes_s m} \simeq \bigoplus_{m=0}^{+\infty} L^2_s(\mathbf{R}^{dm}, \mathbf{C}).$$

We denote by $\mathcal{F}^m := L^2_s(\mathbf{R}^{dm}, \mathbf{C})$ the symmetric L^2 space over \mathbf{R}^{dm} . Let

$$X_n = (x_1, \cdots, x_n), \quad dX_n = dx_1 \cdots dx_n,$$

$$K_m = (k_1, \cdots, k_m), \quad dK_m = dk_1 \cdots dk_m.$$
(1.2)

Then, the Hilbert space \mathcal{H} is endowed with the following norm for all $\psi = \{\psi^m\}_{m\geq 0}$

$$\|\psi\|_{\mathcal{H}} := \left[\sum_{m=0}^{\infty} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} |\psi^{m}(X_{n}, K_{m})|^{2} dX_{n} dK_{m}\right]^{1/2}$$

Let \hat{p}_j and \hat{q}_j be the quantized momentum and position operators such that for all $j \in \{1, \dots, n\}$

$$\hat{p}_j = -i\hbar \nabla_{x_j}, \qquad \hat{q}_j = x_j,$$

The \hbar scaled creation-annihilation operators for the field are defined on Γ_s for any $f \in L^2(\mathbf{R}^d, \mathbf{C})$ as

$$\hat{a}_{\hbar}(f) = \int_{\mathbf{R}^d} \overline{f(k)} \, \hat{a}_{\hbar}(k) \, dk, \qquad \hat{a}^*_{\hbar}(f) = \int_{\mathbf{R}^d} f(k) \, \hat{a}^*_{\hbar}(k) \, dk \,,$$

where $\hat{a}_{\hbar}(k)$ and $\hat{a}_{\hbar}^{*}(k)$ are the creation-annihilation operator-valued distributions defined as follows

$$[\hat{a}_{\hbar}(k) \ \psi]^{m}(k_{1}, \cdots, k_{m}) = \sqrt{\hbar(m+1)} \ \psi^{m+1}(k, k_{1}, \cdots, k_{m});$$
$$[\hat{a}_{\hbar}^{*}(k) \ \psi]^{m}(k_{1}, \cdots, k_{m}) = \frac{\sqrt{\hbar}}{\sqrt{m}} \sum_{j=1}^{m} \delta(k-k_{j}) \ \psi^{m-1}(k_{1}, \cdots, \hat{k}_{j}, \cdots, k_{m})$$

In our case, we will work with the generalized \hbar scaled creation-annihilation operators. The two operators $\hat{a}^{\sharp}_{\hbar}(G) : \mathcal{H} \to \mathcal{H}$ are defined for

$$\begin{array}{rcl} G \,:\, L^2(\mathbf{R}^{dn}_x, \mathbf{C}) &\longrightarrow & L^2(\mathbf{R}^{dn}_x, \mathbf{C}) \otimes L^2(\mathbf{R}^{d}_k, \mathbf{C}) \\ \psi &\longmapsto & G \ \psi. \end{array}$$

with

$$(G \ \psi)(X_n, k) = \sum_{j=1}^n \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{-2\pi i k \cdot \hat{q}_j} \ \psi(X_n).$$

In general, we have

$$[\hat{a}_{\hbar}(G) \ \psi(X_n)]^m(K_m) = \sqrt{\hbar(m+1)} \sum_{j=1}^n \int_{\mathbf{R}^d} \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{2\pi i k \cdot \hat{q}_j} \ \psi^{m+1}(X_n; K_m, k) \ dk \ ; \tag{1.3}$$

$$[\hat{a}_{\hbar}^{*}(G) \ \psi(X_{n})]^{m}(K_{m}) = \frac{\sqrt{\hbar}}{\sqrt{m}} \sum_{j=1}^{m} \sum_{\ell=1}^{n} \frac{\chi(k_{j})}{\sqrt{\omega(k_{j})}} \ e^{-2\pi i k_{j} \cdot \hat{q}_{\ell}} \ \psi^{m-1}(X_{n}; k_{1}, \cdots, \hat{k}_{j}, \cdots, k_{m}).$$
(1.4)

Introduce the second quantization $d\Gamma(A) : \mathcal{H} \to \mathcal{H}$ for the self-adjoint operator A with $d\Gamma(A) \psi = \{[d\Gamma(A) \ \psi]^m\}_{m>0}$ and where

$$[d\Gamma(A) \ \psi]^m = \hbar \sum_{j=1}^m \psi \otimes \cdots \otimes \underbrace{A\psi}_{j^{th} \text{ position}} \otimes \cdots \otimes \psi.$$

The \hbar scaled number operator $\hat{N}_{\hbar} = d\Gamma(\text{Id})$ and the number operator \hat{N} are defined as follows

$$[\hat{N}_{\hbar} \psi]^m = \hbar \ m \ \psi^m, \qquad [\hat{N} \ \psi]^m = m \ \psi^m.$$

The free field Hamiltonian $d\Gamma(\omega): \mathcal{H} \to \mathcal{H}$ is defined as follows

$$[d\Gamma(\omega) \ \psi(X_n)]^m = \hbar \sum_{j=1}^m \omega(k_j) \ \psi^m(X_n; K_m).$$

Formally, one can express this as:

$$d\Gamma(\omega) = \int_{\mathbf{R}^d} \hat{a}^*_{\hbar}(k) \ \omega(k) \ \hat{a}_{\hbar}(k) \ dk.$$

The non-interacting Hamiltonian is defined as follows

$$\hat{H}_0 := \hat{H}_{01} + \hat{H}_{02}$$

where we have introduced the two terms \hat{H}_{01} and \hat{H}_{02} as follows

$$\hat{H}_{01} = \sum_{j=1}^{n} f_j(\hat{p}_j), \qquad \hat{H}_{02} = d\Gamma(\omega).$$

The interaction Hamiltonian $\hat{H}_1 : \mathcal{H} \to \mathcal{H}$ is defined in terms of \hat{a}_{\hbar} , \hat{a}^*_{\hbar} as in (1.3)-(1.4) as follows

$$\hat{H}_1 = \hat{a}_\hbar(G) + \hat{a}^*_\hbar(G)$$

The Nelson-Hamiltonian takes then the following form

$$\hat{H}_{\hbar} \equiv \hat{H} = \hat{H}_0 + V(\hat{q}) + \hat{H}_1.$$
(1.5)

The inclusion of a form factor χ in the interaction term of the particle-field equation ensures the well-definedness of the corresponding quantum dynamics and leads to a self-adjoint Nelson Hamiltonian. It has been demonstrated that, under certain mild assumptions on χ and the potential V, the unbounded operator \hat{H}_{\hbar} is indeed self-adjoint (as discussed in [5] and references therein). In the following, we aim to identify the minimal conditions on χ and V that enable further analysis.

1.2. Assumptions and main results. We have to impose the following assumptions on the external potential $V : \mathbf{R}^{dn} \mapsto \mathbf{R}$ and the form factor $\chi : \mathbf{R}^d \mapsto \mathbf{R}$ with $\sigma \ge 0$:

$$V \in \mathcal{C}_b^2(\mathbf{R}^{dn}; \mathbf{R}), \tag{1.6}$$

$$\omega(\cdot)^{\frac{3}{2}-\sigma}\chi(\cdot) \in L^2(\mathbf{R}^d;\mathbf{R}).$$
(1.7)

Note that the following identities hold true:

- If $\omega(\cdot)^{\frac{3}{2}-\sigma}\chi(\cdot) \in L^2(\mathbf{R}^d;\mathbf{R})$ then $\chi(\cdot) \in L^2(\mathbf{R}^d;\mathbf{R})$;
- If $\chi(\cdot) \in L^2(\mathbf{R}^d; \mathbf{R})$ then for any $\gamma > 0$, we have $\omega(\cdot)^{-\gamma}\chi(\cdot) \in L^2(\mathbf{R}^d; \mathbf{R})$.

Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices on \mathcal{H} of the particle-field quantum system. The main assumptions on the family of states $(\varrho_{\hbar})_{\hbar \in (0,1)}$ are:

$$\exists C_0 > 0, \ \forall \hbar \in (0,1), \quad \operatorname{Tr}[\varrho_\hbar \ d\Gamma(\omega^{2\sigma})] \le C_0, \tag{1.8}$$

$$\exists C_1 > 0, \ \forall \hbar \in (0,1), \quad \operatorname{Tr}[\varrho_\hbar \ (\hat{q}^2 + \hat{p}^2)] \le C_1.$$
 (1.9)

Remark that the following identities hold true:

- If $\operatorname{Tr}[\varrho_{\hbar} d\Gamma(\omega^{2\sigma})] \leq c_0$, then $\operatorname{Tr}[\varrho_{\hbar} d\Gamma(\omega)] \leq c'_0$ for some $c_0, c'_0 \in \mathbf{R}^*_+$;
- If $\operatorname{Tr}[\varrho_{\hbar} d\Gamma(\omega)] \leq c_1$, then $\operatorname{Tr}[\varrho_{\hbar} \hat{N}_{\hbar}] \leq c'_1$ for some $c_1, c'_1 \in \mathbf{R}^*_+$;
- If $\operatorname{Tr}[\varrho_{\hbar}(\hat{q}^2 + \hat{p}^2)] \leq c_2$, then $\operatorname{Tr}[\varrho_{\hbar}(\hat{H}_0 + 1)] \leq c'_2$ for some $c_2, c'_2 \in \mathbf{R}^*_+$.

The first result presented in this section concerns the flow of the particle-field equation.

Theorem 1.1 (Global well-posedness of the particle-field equation). Let $\sigma \in [\frac{1}{2}, 1]$. Assume (1.6) and (1.7) hold. Then for any initial condition $u_0 \in X^{\sigma}$ there exists a unique global strong solution $u(\cdot) \in \mathcal{C}(\mathbf{R}, X^{\sigma}) \cap \mathcal{C}^1(\mathbf{R}, X^{\sigma-1})$ of the particle-field equation (1.1). Moreover, the global flow map $u_0 \to \Phi_t(u_0) = u(t)$ associated to the particle-field equation (1.1) is Borel measurable.

The above global flow is not constructed from a fixed point argument, whereas it is constructed by means of statistical arguments. More precisely, we use measure theoritical techniques to construct this flow. And thus, it is only Borel measurble and not necessarily continuous. Denote by $\mathcal{P}(X^0)$ the set of all Borel probability measure over the space X^0 .

Definition 1.2 (Wigner measures). A Borel probability measure $\mu \in \mathcal{P}(X^0)$ is a Wigner measure of a family of density matrices $(\varrho_{\hbar})_{\hbar \in (0,1)}$ on the Hilbert space \mathcal{H} if and only if there exists a countable subset $\mathcal{A} \subset (0,1)$ with $0 \in \overline{\mathcal{A}}$ such that for any $\xi = (p_0, q_0, \alpha_0) \in X^0$:

$$\lim_{\hbar \to 0, \hbar \in \mathcal{A}} \operatorname{Tr} \left[\mathcal{W}(2\pi q_0, -2\pi p_0, \sqrt{2\pi\alpha_0}) \varrho_h \right] = \int_{X^0} e^{2\pi i \Re e \langle \xi, u \rangle_{X^0}} d\mu(u).$$

The next result concerns the classical limit which relies on the construction of a Wigner measure in the context of infinite-dimensional bosonic quantum field theory. This allows us to establish convergence from the quantum to the classical dynamics. Denote by

$$\mathcal{M}(\varrho_{\hbar}, \hbar \in \mathcal{A})$$

the set of all Wigner measure associated to the density matrices $(\varrho_{\hbar})_{\hbar \in \mathcal{A}}$.

Theorem 1.3 (Validity of Bohr's correspondence principle). Let $\sigma \in [\frac{1}{2}, 1]$ and assume (1.6) and (1.7) hold true. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices on \mathcal{H} satisfying (1.8) and (1.9). Let $(\hbar_n)_{n \in \mathbb{N}} \subset (0,1)$ such that $\hbar_n \longrightarrow 0$ and assume that $\mathcal{M}(\varrho_{\hbar_n}, n \in \mathbb{N}) = {\mu_0}$. Then for all times $t \in \mathbf{R}$, there exists a subsequence $(\hbar_{\ell})_{\ell \in \mathbf{N}}$ and a family of Borel probability measure $(\mu_t)_{t \in \mathbf{R}}$ such that

$$\mathcal{M}(e^{-i\frac{t}{\hbar_{\ell}}\hat{H}}\varrho_{\hbar_{\ell}}\ e^{i\frac{t}{\hbar_{\ell}}\hat{H}},\ \ell\in\mathbf{N})=\{\mu_t\},$$

where $\mu_t \in \mathcal{P}(X^0)$ satisfying

(i) μ_t is concentrated on X^{σ} i.e. $\mu_t(X^{\sigma}) = 1$;

(ii) $\mu_t = (\Phi_t)_{\sharp} \mu_0$, where $u_0 \mapsto \Phi_t(u_0) = u(t)$ is the Borel measurable global flow of the particle-field equation (1.1).

The result above indicates that when ρ_{\hbar} are density matrices on \mathcal{H} that approach the Wigner probability measure μ_0 as \hbar approaches zero, the evolved density matrices $\rho_{\hbar}(t)$ will converge to $\mu_t = (\Phi_t) \sharp \mu_0$ for all times t. Here, Φ_t is the flow that solves (1.1).

To demonstrate the aforementioned results, we adopt the following approach: Firstly, we employ classical techniques to establish the uniqueness property of the particle-field solutions. Subsequently, we establish crucial uniform propagation estimates on the quantum dynamics. Then, we present a probabilistic representation of measure-valued solutions for the Liouville's equation (see [8, 9]). This representation is used to construct a generalized global flow for a particle-field equation. As a conclusion, we establish by means of Wigner measures the global well-posedness for the particle-field equation and the Bohr's correspondence principle for the Nelson model.

2. The classical system

This section is dedicated to examining various classical properties of the particle-field equation. Firstly, in Subsection 2.1, we introduce the particle-field equation as a semi-linear partial differential equation and establish its interaction representation. In Subsection 2.2, we prove the uniqueness of the particle-field equation using this representation.

2.1. The interaction representation. The particle-field equation (1.1) takes the following form

$$\begin{cases} \frac{du(t)}{dt} = w(u(t)) = \mathcal{L}(u(t)) + \mathcal{N}(u(t)), \\ u(0) = u_0 \in X^{\sigma}, \end{cases}$$
(PFE)

where $t \to u(t) = (p(t), q(t), \alpha(t))$ is a solution, $\mathcal{L}(u) = (0, 0, -i\omega\alpha)$ is a linear operator such that $\mathcal{L}: X^{\sigma} \longrightarrow X^{\sigma-1}$ and \mathcal{N} is the nonlinearity given by

$$(\mathcal{N}(u))_{p_j} := -\nabla_{q_j} V(q) - \nabla_{q_j} I_j(q, \alpha),$$

$$(\mathcal{N}(u))_{q_j} := \nabla f_j(p_j),$$

$$(\mathcal{N}(u))_{\alpha}(k) := -i \sum_{j=1}^n \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot q_j},$$

(2.1)

where we have introduced $I_j: \mathbf{R}^{dn} \times L^2(\mathbf{R}^d, \mathbf{C}) \to \mathbf{R}$

$$I_j(q,\alpha) := \int_{\mathbf{R}^d} \frac{\chi(k)}{\sqrt{\omega(k)}} \left[\alpha(k) e^{2\pi i k \cdot q_j} + \overline{\alpha(k)} e^{-2\pi i k \cdot q_j} \right] dk$$
(2.2)

with

$$\nabla_{q_j} I_j(q,\alpha) = \int_{\mathbf{R}^d} 2\pi i k \; \frac{\chi(k)}{\sqrt{\omega(k)}} \left[\alpha(k) e^{2\pi i k \cdot q_j} - \overline{\alpha(k)} e^{-2\pi i k \cdot q_j} \right] dk. \tag{2.3}$$

We consider now the particle-field equation as a non-autonomous initial value problem over the Hilbert space X^{σ} with

$$\begin{cases} \frac{du(t)}{dt} = v(t, u(t)), \\ u(0) = u_0 \in X^{\sigma}. \end{cases}$$
(IVP)

The non-autonomous vector field v is defined in terms of the non-linearity $\mathcal{N} : X^{\sigma} \longrightarrow X^{\sigma}$ of the particle-field equation as well as the free field flow $\Phi_t^f : X^{\sigma} \longrightarrow X^{\sigma}$ as follows:

$$v(t,u) = \Phi^f_{-t} \circ \mathcal{N} \circ \Phi^f_t(u), \qquad (2.4)$$

where we have introduced the free field flow Φ_t^f as follows

$$\Phi_t^f(p,q,\alpha) = (p,q,e^{-it\omega(k)}\alpha).$$
(2.5)

Lemma 2.1 (Explicit expression for the vector field v). The vector field $v : \mathbf{R} \times X^{\sigma} \to X^{\sigma}$ takes the following explicit form:

$$(v(t,u))_{p_j} = \left(\mathcal{N} \circ \Phi_t^f(u)\right)_{p_j},$$

$$(v(t,u))_{q_j} = \left(\mathcal{N} \circ \Phi_t^f(u)\right)_{q_j},$$

$$(v(t,u))_{\alpha}(k) = e^{it\omega(k)} \left(\mathcal{N} \circ \Phi_t^f(u)\right)_{\alpha}(k),$$

$$(2.6)$$

where $v(t, u) = {}^{t} ((v(t, u))_{p_1}, \cdots, (v(t, u))_{p_n}, (v(t, u))_{q_1}, \cdots, (v(t, u))_{q_n}, v(t, u)_{\alpha}).$

Proof. The result follows from direct computations of v using the relation (2.4).

Proposition 2.2 (Equivalence between (PFE) and (IVP)). Assume (1.6) and (1.7) are satisfied. Let I be a bounded open interval containing the origin. Then, the statements below are equivalent:

- (1) $u(\cdot) \in \mathcal{C}^1(I, X^{\sigma})$ is a strong solution of (IVP);
- (2) $u(\cdot) \in \mathcal{C}(I, X^{\sigma})$ solves the following Duhamel formula

$$u(t) = u_0 + \int_0^t v(s, u(u)) \, ds, \quad \forall t \in I.$$

(3) The curve $t \mapsto \Phi_t^f(u(t)) \in \mathcal{C}(I, X^{\sigma}) \cap \mathcal{C}^1(I, X^{\sigma-1})$ is a strong solution to the particle-field equation (1.1).

Proof. The first two assertions can be proved easily since v is continuous vector field (by Lemma 2.5). Let us now prove the equivalence between (1) and (3). Suppose that $u(t) = (p(t), q(t), \alpha(t))$ is a solution to (IVP). Require to prove that

$$\tilde{u}(t) = (\tilde{p}(t), \tilde{q}(t), \tilde{\alpha}(t)) = \Phi_t^f(u(t)) = (p(t), q(t), e^{-it\omega(k)}\alpha(t)),$$

is a solution to (PFE). The first term

$$\partial_t \tilde{p}_j = \partial_t p_j = (v(t, u))_{p_j} = \left(\mathcal{N} \circ \Phi_t^f(u)\right)_{p_j} = (w(\tilde{u}))_{p_j}$$

The second term

$$\partial_t \tilde{q}_j = \partial_t q_j = \nabla f_j(p_j) = (\mathbf{w}(\tilde{u}))_{q_j}.$$

The third term

$$\partial_t \tilde{\alpha} = -i\omega(k) \ e^{-it\omega(k)} \ \alpha(k) + e^{-it\omega(k)} \partial_t \alpha$$

= $-i\omega(k) \ \tilde{\alpha}(k) + e^{-it\omega(k)} \ e^{it\omega(k)} \ \left(\mathcal{N} \circ \Phi^f_t(u)\right)_{\alpha}(k)$
= $-i\omega(k) \ \tilde{\alpha}(k) + (\mathcal{N}(\tilde{u}))_{\alpha}(k)$
= $(\mathbf{w}(\tilde{u}))_{\alpha}(k).$

We conclude that \tilde{u} is a solution to (PFE). Similarly, we can prove the reverse sense.

Let I be an open interval containing the origin. We are interested in strong solution to the particle-field equation (PFE) such that

$$u(\cdot) \in \mathcal{C}(I, X^{\sigma}) \cap \mathcal{C}^{1}(I, X^{\sigma-1}),$$

8

and (PFE) is satisfied for all $t \in I$. In particular, from the second assertion of Proposition 2.2, these solutions satisfy the following Duhamel formula for all $t \in I$

$$u(t) = \Phi_t^f(u(0)) + \int_0^t \Phi_{t-s}^f \circ \mathcal{N}(u(s)) \, ds, \qquad (2.7)$$

where $\Phi_t^f(\cdot)$ is the free field flow defined above in (2.5).

2.2. Properties of the particle-field equation. In this section, we establish various properties related to the particle-field equation and its time interaction representation. Of most significance is the recovery of the uniqueness property of solutions to the particle-field equation (PFE) on the energy space X^{σ} . Our approach starts with deriving estimates for $\nabla_{q_i} I_j(\cdot)$.

Lemma 2.3 (Estimates for $\nabla_{q_i} I_j$). We have the following two estimates.

(i) Assume $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Then, for all $(q, \alpha) \in \mathbf{R}^{dn} \times L^2(\mathbf{R}^d, \mathbf{C})$, for all $j \in \{1, \dots, n\}$, we have the following estimate

$$|\nabla_{q_j} I_j(q, \alpha)| \le 4\pi \|\omega^{1/2} \chi\|_{L^2} \|\alpha\|_{L^2}.$$
(2.8)

(ii) Assume (1.7) is satisfied. Then, for all $j \in \{1, \dots, n\}$, for all $q_1, q_2 \in \mathbf{R}^{dn}$ with $q_1 = (q_{1j})_j$ and $q_2 = (q_{2j})_j$, for all $\alpha_1, \alpha_2 \in \mathcal{G}^{\sigma}$, we have

$$\begin{aligned} |\nabla_{q_j} I_j(q_1, \alpha_1) - \nabla_{q_j} I_j(q_2, \alpha_2)| \\ &\leq 4\pi \|\omega^{1/2} \chi\|_{L^2} \|\alpha_1 - \alpha_2\|_{L^2} + 8\sqrt{2}\pi^2 \|\omega^{\frac{3}{2} - \sigma} \chi\|_{L^2} |q_{1j} - q_{2j}| \|\alpha_2\|_{\mathcal{G}^{\sigma}}. \end{aligned}$$

Proof. For (i), by Cauchy-Schwatrz inequality, we have $\forall (q, \alpha) \in \mathbf{R}^{dn} \times L^2(\mathbf{R}^d, \mathbf{C})$

$$\begin{aligned} |\nabla_{q_j} I_j(q, \alpha)| &= \Big| \int_{\mathbf{R}^d} \frac{\chi(k)}{\sqrt{\omega(k)}} 2\pi i k \Big[\alpha(k) e^{2\pi i k \cdot q_j} - \overline{\alpha(k)} e^{-2\pi i k \cdot q_j} \Big] \, dk \\ &\leq 4\pi \int_{\mathbf{R}^d} \Big| \frac{\chi(k)}{\sqrt{\omega(k)}} \, \omega(k) \Big| \, |\alpha(k)| \, dk \\ &\leq 4\pi \, \|\omega^{1/2} \chi\|_{L^2} \, \|\alpha\|_{L^2}. \end{aligned}$$

For (ii), by Cauchy-Schwartz inequality and using the estimate $|e^{iy} - 1| \leq \sqrt{2} |y|$, we have $\forall (q_1, \alpha_1), (q_2, \alpha_2) \in \mathbf{R}^{dn} \times \mathcal{G}^{\sigma}, \forall j \in \{1, \dots, n\}$ the following estimates

$$\begin{split} |\nabla_{q_{j}}I_{j}(q_{1},\alpha_{1}) - \nabla_{q_{j}}I_{j}(q_{2},\alpha_{2})| \\ &= \Big| \int_{\mathbf{R}^{d}} \frac{\chi(k)}{\sqrt{\omega(k)}} 2\pi ik \Big[\alpha_{1}(k)e^{2\pi ik \cdot q_{1j}} - \overline{\alpha_{1}(k)}e^{-2\pi ik \cdot q_{1j}} - \alpha_{2}(k)e^{2\pi ik \cdot q_{2j}} + \overline{\alpha_{2}(k)}e^{-2\pi ik \cdot q_{2j}} \Big] dk \Big| \\ &= \Big| \int_{\mathbf{R}^{d}} \frac{\chi(k)}{\sqrt{\omega(k)}} 2\pi ik \Big[(\alpha_{1}(k) - \alpha_{2}(k))e^{2\pi ik \cdot q_{1j}} + \alpha_{2}(k) \Big(e^{2\pi ik \cdot q_{1j}} - e^{2\pi ik \cdot q_{2j}} \Big) \\ &+ (\overline{\alpha_{2}(k)} - \overline{\alpha_{1}(k)})e^{-2\pi ik \cdot q_{1j}} + \overline{\alpha_{2}(k)} \Big(e^{-2\pi ik \cdot q_{2j}} - e^{-2\pi ik \cdot q_{1j}} \Big) \Big] \\ &\leq 4\pi \int_{\mathbf{R}^{d}} \Big| \frac{\chi(k)}{\sqrt{\omega(k)}} \omega(k) \Big| \Big[|\alpha_{1}(k) - \alpha_{2}(k)| + |\alpha_{2}(k) \left(e^{2\pi ik \cdot (q_{1j} - q_{2j})} - 1 \right) | \Big] \\ &\leq 4\pi \int_{\mathbf{R}^{d}} \Big| \sqrt{\omega(k)} \chi(k) \Big| \Big[|\alpha_{1}(k) - \alpha_{2}(k)| + |\alpha_{2}(k) \sqrt{2} \ 2\pi k \cdot (q_{1j} - q_{2j}) | \Big] \\ &\leq 4\pi \int_{\mathbf{R}^{d}} \Big| \sqrt{\omega(k)} \chi(k) \Big| \Big[|\alpha_{1}(k) - \alpha_{2}(k)| + 2\sqrt{2} \ \pi |\alpha_{2}(k)| \ \omega(k) \ |q_{1j} - q_{2j}| \Big] \\ &\leq 4\pi \int_{\mathbf{R}^{d}} \| \omega^{1/2} \chi \|_{L^{2}} \| \alpha_{1} - \alpha_{2} \|_{L^{2}} + 8\sqrt{2}\pi^{2} \| \omega^{\frac{3}{2} - \sigma} \chi \|_{L^{2}} \ |q_{1j} - q_{2j}| \ \|\alpha_{2} \|_{\mathcal{G}^{\sigma}}. \end{split}$$

The vector field \mathcal{N} , which characterizes the nonlinearity of the particle-field equation, possesses the following properties.

Proposition 2.4 (Continuity and boundedness of \mathcal{N}). Assume (1.6) and (1.7) are satisfied. Then, the nonlinearity $\mathcal{N}: X^{\sigma} \to X^{\sigma}$ is a continuous, and bounded on bounded sets, vector field.

Proof. Let us prove first that $\mathcal{N}: X^{\sigma} \to X^{\sigma}$ is bounded on bounded sets. Let $u \in X^{\sigma}$ be a bounded such that $||u||_{X^{\sigma}} \leq c_0$, for some $c_0 > 0$. Require to prove $||\mathcal{N}(u)||_{X^{\sigma}}^2 \leq c_1$ for some $c_1 > 0$. We have first with some $c_2 > 0$

$$\begin{split} \left| \left(\mathcal{N}(u) \right)_{p_j} \right| &= \left| -\nabla_{q_j} V(q) - \nabla_{q_j} I_j(q, \alpha) \right| \\ &\leq \left| \nabla_{q_j} V(q) \right| + \left| \nabla_{q_j} I_j(q, \alpha) \right| \\ &\leq \sum_{Lemma \ 2.3-(i)} \left\| \nabla_{q_j} V \right\|_{L^{\infty}} + 4\pi \left\| \omega^{1/2} \chi \right\|_{L^2} \left\| \alpha \right\|_{L^2} \\ &\leq \sup_{j=1}^n \left\| \nabla_{q_j} V \right\|_{L^{\infty}} + \frac{4\pi}{m_f^{\sigma}} \left\| \omega^{1/2} \chi \right\|_{L^2} \left\| \alpha \right\|_{\mathcal{G}^{\sigma}} \\ &\leq \sup_{j=1}^n \left\| \nabla_{q_j} V \right\|_{L^{\infty}} + \frac{4\pi}{m_f^{\sigma}} \left\| \omega^{1/2} \chi \right\|_{L^2} c_0 := c_2. \end{split}$$

We also have with some c_3 , $c_4 > 0$

$$\left| \left(\mathcal{N}(u) \right)_{q_j} \right| = \left| \nabla f_j(p_j) \right| \le c_3 \ |p_j| \le c_3 \ c_0 := c_4.$$

Finally, we have with some $c_5 > 0$

$$\begin{split} \| \left(\mathcal{N}(u) \right)_{\alpha} \|_{\mathcal{G}^{\sigma}}^{2} &= \| \omega^{\sigma} \left(\mathcal{N}(u) \right)_{\alpha} \|_{L^{2}}^{2} \\ &= \int_{\mathbf{R}^{d}} \Big| -i \sum_{j=1}^{n} \omega^{\sigma}(k) \; \frac{\chi(k)}{\sqrt{\omega(k)}} \; e^{-2\pi i k \cdot q_{j}} \Big|^{2} \; dk \\ &\lesssim n^{2} \; \| \omega^{\sigma - \frac{1}{2}} \chi \|_{L^{2}}^{2} := c_{5}, \end{split}$$

where $\|\omega^{\sigma-\frac{1}{2}}\chi\|_{L^2}$ is finite since $\sigma-\frac{1}{2}<\frac{3}{2}-\sigma$ for $\sigma\in[\frac{1}{2},1]$. This implies that there exists $c_1 > 0$ such that

$$\|\mathcal{N}(u)\|_{X^{\sigma}}^2 \le n \ (c_2^2 + c_4^2) + c_5 := c_1$$

It remains to prove the continuity of the nonlinear term $\mathcal{N}: X^{\sigma} \to X^{\sigma}$. Suppose that

$$u_{\ell} = (p_{\ell}, q_{\ell}, \alpha_{\ell}) \xrightarrow[\ell \to +\infty]{} u = (p, q, \alpha), \quad \text{ in } X^{\sigma} \text{ i.e. } \|u_{\ell} - u\|_{X^{\sigma}} \xrightarrow[\ell \to +\infty]{} 0$$

Require to prove

$$\mathcal{N}(u_{\ell}) \xrightarrow[\ell \to +\infty]{} \mathcal{N}(u) \text{ in } X^{\sigma} \text{ i.e. } \|\mathcal{N}(u_{\ell}) - \mathcal{N}(u)\|_{X^{\sigma}} \xrightarrow[\ell \to +\infty]{} 0.$$

Indeed, we have

$$\|\mathcal{N}(u_{\ell}) - \mathcal{N}(u)\|_{X^{\sigma}}^{2} = \sum_{j=1}^{n} \left[\left| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{p_{j}} \right|^{2} + \left| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{q_{j}} \right|^{2} \right] + \| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{\alpha} \|_{\mathcal{G}^{\sigma}}^{2}.$$
¹⁰

By Lemma 2.3-(ii), we can assert that

$$\begin{split} \left| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{p_{j}} \right| \\ &\leq \left| \nabla_{q_{j}} V(q_{\ell}) - \nabla_{q_{j}} V(q) \right| + \frac{4\pi}{m_{f}^{\sigma}} \| \omega^{1/2} \chi \|_{L^{2}} \| \alpha_{\ell} - \alpha \|_{\mathcal{G}^{\sigma}} + 8\sqrt{2}\pi^{2} \| \omega^{\frac{3}{2} - \sigma} \chi \|_{L^{2}} |q_{\ell j} - q_{j}| \| \alpha \|_{\mathcal{G}^{\sigma}} \\ &\leq \sum_{j'=1}^{n} \| \nabla_{q_{j'}} \nabla_{q_{j}} V \|_{L^{\infty}} |q_{\ell j'} - q_{j'}| + \frac{4\pi}{m_{f}^{\sigma}} \| \omega^{1/2} \chi \|_{L^{2}} \| \alpha_{\ell} - \alpha \|_{\mathcal{G}^{\sigma}} + 8\sqrt{2}\pi^{2} \| \omega^{\frac{3}{2} - \sigma} \chi \|_{L^{2}} |q_{\ell j} - q_{j}| \| \alpha \|_{\mathcal{G}^{\sigma}} \\ &\leq \left[n \sup_{j', j=1}^{n} \| \nabla_{q_{j'}} \nabla_{q_{j}} V \|_{L^{\infty}} + \frac{4\pi}{m_{f}^{\sigma}} \| \omega^{1/2} \chi \|_{L^{2}} + 8\sqrt{2}\pi^{2} \| \omega^{\frac{3}{2} - \sigma} \chi \|_{L^{2}} \| \alpha \|_{\mathcal{G}^{\sigma}} \right] \| u_{\ell} - u \|_{X^{\sigma}} \xrightarrow{0}{\rightarrow} 0. \end{split}$$

$$\left| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{q_j} \right| \le c \ |p_{\ell j} - p_j| \le c \ \|u_{\ell} - u\|_{X^{\sigma}} \xrightarrow[\ell \to +\infty]{} 0.$$

For the last term we have

$$\begin{split} \| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{\alpha} \|_{\mathcal{G}^{\sigma}}^{2} &= \int_{\mathbf{R}^{d}} \left| -i \sum_{j=1}^{n} \omega^{\sigma}(k) \; \frac{\chi(k)}{\sqrt{\omega(k)}} \left[e^{-2\pi i k \cdot q_{\ell j}} - e^{-2\pi i k \cdot q_{j}} \right] \right|^{2} \; dk \\ &= \int_{\mathbf{R}^{d}} \left| -i \sum_{j=1}^{n} \omega^{\sigma - \frac{1}{2}}(k) \; \chi(k) \left[e^{-2\pi i k \cdot q_{\ell j}} - e^{-2\pi i k \cdot q_{j}} \right] \right|^{2} \; dk, \end{split}$$

where we have

•
$$\int_{\mathbf{R}^{d}} \left| -i \sum_{j=1}^{n} \omega^{\sigma - \frac{1}{2}}(k) \chi(k) \left[e^{-2\pi i k \cdot q_{\ell j}} - e^{-2\pi i k \cdot q_{j}} \right] \right|^{2} dk \leq 4n^{2} \|\omega^{\sigma - \frac{1}{2}} \chi\|_{L^{2}}^{2} < +\infty;$$

• $\left| e^{-2\pi i k \cdot q_{\ell j}} - e^{-2\pi i k \cdot q_{j}} \right| \xrightarrow[\ell \to +\infty]{} 0.$

Hence by Lebesgue dominated convergence theorem, we get

$$\| \left(\mathcal{N}(u_{\ell}) - \mathcal{N}(u) \right)_{\alpha} \|_{\mathcal{G}^{\sigma}}^{2} \xrightarrow[\ell \to +\infty]{} 0$$

And thus,

$$\|\mathcal{N}(u_{\ell}) - \mathcal{N}(u)\|_{X^{\sigma}}^{2} \xrightarrow[\ell \to +\infty]{} 0.$$

The above theorem implies the following results on the vector field v.

Lemma 2.5 (Continuity and boundedness properties of the vector field v). Assume (1.6) and (1.7) are satisfied. Then, the vector field $v : \mathbf{R} \times X^{\sigma} \longrightarrow X^{\sigma}$ is continuous and bounded on bounded subsets of $\mathbf{R} \times X^{\sigma}$.

Proof. This is a consequence of the continuity and boundedness properties of the nonlinear term $\mathcal{N}: X^{\sigma} \to X^{\sigma}$ in Proposition 2.4.

As a consequence of the above properties, we have the following uniqueness property.

Proposition 2.6 (Uniqueness property). Assume (1.6) and (1.7) are satisfied. Let I be an open interval containing the origin and let $u_1, u_2 \in C(I, X^{\sigma})$ be two strong solutions of the particle-field equation (PFE) such that $u_1(0) = u_2(0)$. Then $u_1(t) = u_2(t)$ for all $t \in I$.

Proof. Note first that using Duhamel formula (2.7) as well as $u_1(0) = u_2(0)$, we have for all $t \ge 0$

$$\|u_1(t) - u_2(t)\|_{X^0} \le \int_0^t \|\mathcal{N}(u_1(s)) - \mathcal{N}(u_2(s))\|_{X^0} \, ds.$$

We claim that for all $s \in [0, t]$, there exists C > 0 such that

$$\|\mathcal{N}(u_1(s)) - \mathcal{N}(u_2(s))\|_{X^0} \le C \|u_1(s) - u_2(s)\|_{X^0}.$$

Indeed, by using the Mean Value Theorem for multivariate vector-valued function $\nabla_{q_j} V$, the first component yields to

$$\begin{split} \left| \left(\mathcal{N}(u_{1}(s)) - \mathcal{N}(u_{2}(s)) \right)_{p_{j}} \right| \\ &\leq \left| \nabla_{q_{j}} V(q_{1}(s)) - \nabla_{q_{j}} V(q_{2}(s)) \right| + \left| \nabla_{q_{j}} I_{j}(q_{1}(s), \alpha_{1}(s)) - \nabla_{q_{j}} I_{j}(q_{2}(s), \alpha_{2}(s)) \right| \\ &\lesssim \left\| \nabla_{q} \nabla_{q_{j}} V \right\|_{L^{\infty}} \left| q_{1}(s) - q_{2}(s) \right| + \left\| \alpha_{1}(s) - \alpha_{2}(s) \right\|_{L^{2}} + \left\| \alpha_{2}(s) \right\|_{\mathcal{G}^{\sigma}} \left| q_{1}(s) - q_{2}(s) \right| \\ &\lesssim \left\| u_{1}(s) - u_{2}(s) \right\|_{X^{0}}, \end{split}$$

where we have used for some bounded interval $J \subset I$

$$\|\alpha_2(s)\|_{\mathcal{G}^{\sigma}} \leq \sup_{s \in J} \|u_2(s)\|_{X^{\sigma}} < +\infty.$$

The second component yields to

$$\left(\mathcal{N}(u_1(s)) - \mathcal{N}(u_2(s))\right)_{q_j} \lesssim |p_1(s) - p_2(s)| \lesssim ||u_1(s) - u_2(s)||_{X^0}.$$

The third component yields to

$$\| \left(\mathcal{N}(u_1(s)) - \mathcal{N}(u_2(s)) \right)_{\alpha} \|_{L^2} \lesssim |q_1(s) - q_2(s)| \lesssim \| u_1(s) - u_2(s) \|_{X^0}.$$

Therefore, we get by combining the above three components the following estimate

$$\begin{aligned} \|u_1(t) - u_2(t)\|_{X^0} &\leq \int_0^t \|\mathcal{N}(u_1(s)) - \mathcal{N}(u_2(s))\|_{X^0} \, ds \\ &\leq C \, \int_0^t \|u_1(s) - u_2(s)\|_{X^0} \, ds. \end{aligned}$$

Then, by Gronwall's Lemma, we get $||u_1(t) - u_2(t)||_{X^0} = 0$. Thus, we get the desired result. \Box

Remark 2.7 (Local well-posendness in $X^{1/2}$). Under assumptions (1.6) and (1.7), for all initial data $u_0 \in X^{1/2}$, one can prove the existence of a unique local solution $u(\cdot) \in \mathcal{C}([0,T]; X^{1/2})$ to (1.1), where $T \in \mathbf{R}^+_*$ is small enough. This can be proved by means of standard fixed point argument.

Remark 2.8 (Global well-posendness in X^{σ}). In line with the above remark, one could also prove the global well-posedness in X^{σ} by using Granwall arguments and the conservation of Hamiltonian.

3. The quantum system

In Subsection 3.1, we prove some quantum estimates, which we then apply in Subsection 3.2 to establish the self-adjointness of the Nelson Hamiltonian using the Kato-Rellich theorem. Lastly, in Subsection 3.3, we discuss the dynamical equation for the quantum system.

3.1. Quantum estimates. Our initial focus here is on providing the reader with estimates that are necessary to establish the self-adjointness of the Nelson Hamiltonian. Denote by $\mathcal{L}(\mathcal{H})$ the set of all bounded operator and by $\mathcal{L}^{1}(\mathcal{H})$ the set of trace-class operators on \mathcal{H} .

Lemma 3.1 (Creation-Annihilation estimates). Let

$$F \in \mathcal{L}(L^2(\mathbf{R}^{dn}, dX_n), L^2(\mathbf{R}^{dn}, dX_n) \otimes L^2(\mathbf{R}^d, dk)).$$

(i) For every $\psi \in D(\hat{N}_{\hbar}^{1/2})$, we have

$$\|\hat{a}_{\hbar}(F)\psi\|_{\mathcal{H}} \le \|\hat{N}_{\hbar}^{1/2}\psi\|_{\mathcal{H}} \|F\|_{\mathcal{L}(L^{2},L^{2}\otimes L^{2})};$$
(3.1)

$$\|\hat{a}_{\hbar}^{*}(F)\psi\|_{\mathcal{H}} \leq \|(\hat{N}_{\hbar}+1)^{1/2}\psi\|_{\mathcal{H}} \|F\|_{\mathcal{L}(L^{2},L^{2}\otimes L^{2})}.$$
(3.2)

(ii) For all $\psi \in D((\hat{H}_{02})^{1/2})$, we have

$$\|\hat{a}_{\hbar}(F)\psi\|_{\mathcal{H}} \leq \left\|\frac{F}{\sqrt{\omega}}\right\|_{\mathcal{L}} \|(\hat{H}_{02}+1)^{1/2}\psi\|_{\mathcal{H}};$$
(3.3)

$$\|\hat{a}_{\hbar}^{*}(F)\psi\|_{\mathcal{H}}^{2} \leq \left\|\frac{F}{\sqrt{\omega}}\right\|_{\mathcal{L}}^{2} \|(\hat{H}_{02}+1)^{1/2}\psi\|_{\mathcal{H}}^{2} + \hbar \|F\|_{\mathcal{L}}^{2} \|\psi\|^{2}.$$
(3.4)

Proof. Let K_m and X_n as indicated in (1.2). For (i)-(3.1), we have with $\|\cdot\|_{\mathcal{L}} \equiv \|\cdot\|_{\mathcal{L}(L^2, L^2 \otimes L^2)}$

$$\begin{split} \|\hat{a}_{\hbar}(F)\psi\|_{\mathcal{H}}^{2} &= \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \left[\hat{a}_{\hbar}(F)\psi \right]^{m}(X_{n},K_{m}) \right|^{2} dX_{n} dK_{m} \\ &= \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \int_{\mathbf{R}^{d}} \sqrt{\hbar(m+1)} \ \overline{F(k)} \ \psi^{m+1}(X_{n},K_{m},k) dk \right|^{2} dX_{n} \ dK_{m} \\ &\leq _{c-s} \|F\|_{\mathcal{L}}^{2} \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left[\int_{\mathbf{R}^{d}} \left| \sqrt{\hbar(m+1)} \ \psi^{m+1}(X_{n},K_{m},k) \right|^{2} dk \right] dX_{n} \ dK_{m} \\ &\leq \|F\|_{\mathcal{L}}^{2} \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{d(m+1)}} \hbar(m+1) \ \left| \psi^{m+1}(X_{n},K_{m+1}) \right|^{2} dX_{n} \ dK_{m+1} \\ &\leq \|F\|_{\mathcal{L}}^{2} \|\hat{N}_{\hbar}^{1/2}\psi\|_{\mathcal{H}}^{2}. \end{split}$$

For (i)-(3.2), we have

$$\begin{aligned} \|\hat{a}_{\hbar}^{*}(F)\psi\|_{\mathcal{H}}^{2} &= \langle \psi, \ \hat{a}_{\hbar}(F)\hat{a}_{\hbar}^{*}(F)\psi \rangle \\ &= \langle \psi, \ \left[[\hat{a}_{\hbar}(F), \hat{a}_{\hbar}^{*}(F)] + \hat{a}_{\hbar}^{*}(F)\hat{a}_{\hbar}(F) \right]\psi \rangle \\ &\leq \hbar \ \|F\|_{\mathcal{L}}^{2} \ \|\psi\|^{2} + \|\hat{a}_{\hbar}(F)\psi\|^{2} \\ &\lesssim \|F\|_{\mathcal{L}}^{2} \ \|(\hat{N}_{\hbar} + 1)^{1/2} \ \psi\|^{2}. \end{aligned}$$

For (ii)-(3.3), we have

$$\begin{split} \|\hat{a}_{\hbar}(F) \psi\|_{\mathcal{H}}^{2} &= \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \left[\hat{a}_{\hbar}(F) \psi \right]^{m}(X_{n}, K_{m}) \right|^{2} dK_{m} \, dX_{n} \\ &= \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \int_{\mathbf{R}^{d}} \sqrt{\hbar(m+1)} \, \overline{F(k)} \, \psi^{m+1}(X_{n}, K_{m}, k) dk \right|^{2} dK_{m} \, dX_{n} \\ &= \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left[\int_{\mathbf{R}^{d}} \left| \sqrt{\hbar(m+1)} \, \frac{\overline{F(k)}}{\sqrt{\omega(k)}} \, \sqrt{\omega(k)} \, \psi^{m+1}(X_{n}, K_{m}, k) \right| dk \right]^{2} dK_{m} \, dX_{n} \\ &\leq \left\| \frac{F}{\sqrt{\omega}} \right\|_{\mathcal{L}}^{2} \, \sum_{m \ge 0} \int_{\mathbf{R}^{dm}} \left[\int_{\mathbf{R}^{d}} \left| \sqrt{\hbar(m+1)} \, \sqrt{\omega(k)} \, \psi^{m+1}(X_{n}, K_{m}, k) \right|^{2} dk \right] dk \, dK_{m} \, dX_{n} \\ &\leq \left\| \frac{F}{\sqrt{\omega}} \right\|_{\mathcal{L}}^{2} \, \sum_{m \ge 0} \int_{\mathbf{R}^{dm}} \int_{\mathbf{R}^{d}} \hbar(m+1) \, \omega(k) \, \left| \psi^{m+1}(X_{n}, K_{m}, k) \right|^{2} dk \, dK_{m} \, dX_{n} \\ &\leq \left\| \frac{F}{\sqrt{\omega}} \right\|_{\mathcal{L}}^{2} \, \left\| (\hat{H}_{02})^{1/2} \, \psi \right\|_{\mathcal{H}}^{2}. \end{split}$$

Similar discussion as for (i)-(3.2) works perfectly to prove (ii)-(3.4).

Lemma 3.2 (Field and number estimates). For all $\psi \in D(\hat{H}_{02})$, we have

(1) $\|\hat{H}_{02} \psi\|_{\mathcal{H}} \ge m_f \|\hat{N}_{\hbar} \psi\|_{\mathcal{H}};$ (2) $\|(\hat{N}_{\hbar} + 1)^{1/2}\psi\|_{\mathcal{H}} \le \varepsilon \|\hat{N}_{\hbar}\psi\|_{\mathcal{H}} + b(\varepsilon) \|\psi\|_{\mathcal{H}}$ for some $\varepsilon < 1$ and $b(\varepsilon) < +\infty$. *Proof.* For (1), we have

$$\begin{split} \|\hat{H}_{02} \psi\|_{\mathcal{H}}^2 &= \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \left[\hat{H}_{02} \psi \right]^m (X_n, k_1, \cdots, k_m) \right|^2 dX_n \ dk_1 \cdots dk_m \\ &= \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \hbar \sum_{l=1}^m \underbrace{\psi(k_l)}_{\ge m_f} \psi^m (X_n, k_1, \cdots, k_m) \right|^2 dX_n \ dk_1 \cdots dk_m \\ &\ge \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \hbar \ m \ m_f \ \psi^m (X_n, k_1, \cdots, k_m) \right|^2 \ dX_n \ dk_1 \cdots dk_m \\ &\ge m_f^2 \sum_{m \ge 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \hbar^2 \ m^2 \ \left| \ \psi^m (X_n, k_1, \cdots, k_m) \right|^2 dX_n \ dk_1 \cdots dk_m \\ &\ge m_f^2 \ \|\hat{N}_{\hbar} \psi\|_{\mathcal{H}}^2. \end{split}$$

For (2), we have

$$\begin{aligned} \|(\hat{N}_{\hbar}+1)^{1/2}\psi\|_{\mathcal{H}}^{2} &= \langle\psi, (\hat{N}_{\hbar}+1)\psi\rangle = \langle\psi, \hat{N}_{\hbar}\psi\rangle + \|\psi\|_{\mathcal{H}}^{2} \\ &\leq \|\psi\|_{\mathcal{H}} \|\hat{N}_{\hbar}\psi\|_{\mathcal{H}} + \|\psi\|_{\mathcal{H}}^{2} \\ &\leq \frac{\tilde{\varepsilon}}{2} \|\hat{N}_{\hbar}\psi\|_{\mathcal{H}}^{2} + (1+\frac{1}{2\tilde{\varepsilon}}) \|\psi\|_{\mathcal{H}}^{2}. \end{aligned}$$

This implies that, by choosing appropriate $\tilde{\varepsilon}$, there exists $\varepsilon < 1$ and $b(\varepsilon) < +\infty$ such that

$$\|(\hat{N}_{\hbar}+1)^{1/2}\psi\|_{\mathcal{H}} \leq \varepsilon \|\hat{N}_{\hbar}\psi\|_{\mathcal{H}} + b(\varepsilon) \|\psi\|_{\mathcal{H}}.$$

Below, we give an important inequality between \hat{H} and \hat{H}_0 which is useful for the coming discussions. Suppose $\hat{H} + a > 0$ and $\hat{H}_0 + b > 0$ with some $a, b \in \mathbf{R}$.

Lemma 3.3 (Equivalence between \hat{H} and \hat{H}_0). Assume (1.6) and $\omega^{-\frac{1}{2}}\chi \in L^2(\mathbf{R}^d, dk)$ are satisfied. Then there exists c, C > 0 independent of \hbar such that for all $\hbar \in (0, 1)$ and for all $\psi \in D(\hat{H}_0)$

$$c \langle \psi, (\hat{H}+a) \psi \rangle \leq \langle \psi, (\hat{H}_0+b) \psi \rangle \leq C \langle \psi, (\hat{H}+a) \psi \rangle.$$
(3.5)

In particular, for all $\psi \in D((\hat{H}_0)^{1/2})$, we have

$$c \| (\hat{H} + a)^{1/2} \psi \| \le \| (\hat{H}_0 + b)^{1/2} \psi \| \le C \| (\hat{H} + a)^{1/2} \psi \|.$$
(3.6)

Proof. Note that first we have the following estimates on V below

$$-\|V\|_{L^{\infty}} \le V \le \|V\|_{L^{\infty}} \Rightarrow \|V\|_{L^{\infty}} + V \ge 0.$$

We have

$$\begin{aligned} \langle \psi, \ (\hat{H}+a) \ \psi \rangle &= \langle \psi, \ \hat{H}_0 \ \psi \rangle + \langle \psi, V \ \psi \rangle + \langle \psi, \ \hat{H}_1 \ \psi \rangle + \langle \psi, a \ \psi \rangle \\ &\leq \langle \psi, \ \hat{H}_0 \ \psi \rangle + \|V\|_{L^{\infty}} \langle \psi, \ \psi \rangle + \|\psi\| \ \|\hat{H}_1 \ \psi\| + \langle \psi, a \ \psi \rangle \\ &\leq \langle \psi, \ \hat{H}_0 \ \psi \rangle + \|V\|_{L^{\infty}} \langle \psi, \ \psi \rangle + 2\|\omega^{-1/2} \ \chi\|_{L^2} \ \|\psi\| \ \|(\hat{H}_0+1)^{1/2} \ \psi\| + \langle \psi, a \ \psi \rangle \\ &\leq \langle \psi, \ \hat{H}_0 \ \psi \rangle + \|V\|_{L^{\infty}} \langle \psi, \ \psi \rangle + \|\omega^{-1/2} \ \chi\|_{L^2}^2 \ \|\psi\|^2 + \|(\hat{H}_0+1)^{1/2} \ \psi\|^2 + \langle \psi, a \ \psi \rangle \\ &\leq \langle \psi, \ \left(2\hat{H}_0 + \|V\|_{L^{\infty}} + \|\omega^{-1/2} \ \chi\|_{L^2}^2 + 1 + a\right) \ \psi \rangle \\ &\leq c \ \langle \psi, \ (\hat{H}_0 + b) \ \psi \rangle, \end{aligned}$$

where $c \in \mathbf{R}^+_*$ depends on $\|V\|_{L^{\infty}}$, $\|\omega^{-1/2} \chi\|_{L^2_{1,4}}^2$ and independent on \hbar .

On the reverse side, we have

$$\begin{aligned} \langle \psi, \ \hat{H}_0 \ \psi \rangle &= \langle \psi, \ \hat{H}_{01} \ \psi \rangle + \langle \psi, \hat{H}_{02} \ \psi \rangle \\ &\leq \langle \psi, \ (\hat{H}_{01} + V + \|V\|_{L^{\infty}}) \ \psi \rangle + \langle \psi, \hat{H}_{02} \ \psi \rangle. \end{aligned}$$

Then by using Lemma 3.1, we can also assert that for $\varepsilon > 0$, there exists $c_{\varepsilon} > 0$ (depends on the norm $\|\omega^{-1} \chi\|_{L^2}$) such that

$$|\langle \psi, \hat{H}_1 \psi \rangle| \le \|\omega^{-1}\chi\|_{L^2} \left[\frac{1}{\varepsilon} \langle \psi, \psi \rangle + \varepsilon \langle \psi, (\hat{H}_{02} + 1) \psi \rangle\right].$$

This means as quadratic form there exists a constant $c_{\varepsilon} \in \mathbf{R}^*_+$ which depends on ε such that

$$\hat{H}_{02} + \hat{H}_1 = (1 - \varepsilon) \ \hat{H}_{02} + \varepsilon \hat{H}_{02} + \hat{H}_1$$
$$\geq (1 - \varepsilon) \ \hat{H}_{02} - c_{\varepsilon} 1$$

This implies

$$\hat{H}_{02} \le \frac{1}{1-\varepsilon} \left[\hat{H}_{02} + \hat{H}_1 + c_\varepsilon \right].$$

We conclude that there exists C > 0 such that

$$\langle \psi, (\hat{H}_0 + b) \psi \rangle \leq \langle \psi, (\hat{H}_{01} + V + ||V||_{L^{\infty}}) \psi \rangle + \langle \psi, \frac{1}{1 - \varepsilon} [\hat{H}_{02} + \hat{H}_1 + c_{\varepsilon}] \psi \rangle$$

$$\leq C \langle \psi, (\hat{H} + a) \psi \rangle.$$

3.2. Self-adjointness of Nelson Hamiltonian. We prove here the self-adjointness of the Nelson Hamiltonian using the estimates provided in the previous section.

Proposition 3.4 (Self-adjointness of the Nelson Hamiltonian). Assume (1.6) and $\omega^{-\frac{1}{2}}\chi \in L^2(\mathbf{R}^d, dk)$. Then, the operator $\hat{H} : \mathcal{H} \to \mathcal{H}$ is self-adjoint operator on $D(\hat{H}_0) = D(\hat{H})$.

Proof. We have first with some $C_{\tilde{\varepsilon}} \in \mathbf{R}^*_+$

$$\begin{split} \|\hat{H}_{1}\psi\|_{\mathcal{H}} &= \|\hat{a}_{\hbar}(G)\psi + \hat{a}_{\hbar}^{*}(G)\psi\|_{\mathcal{H}} \\ &\leq \|\hat{a}_{\hbar}(G)\psi\|_{\mathcal{H}} + \|\hat{a}_{\hbar}^{*}(G)\psi\|_{\mathcal{H}} \\ &\leq 2\|G\| \|(\hat{N}_{\hbar}+1)^{1/2}\psi\|_{\mathcal{H}} \\ &\leq 2n \left\|\frac{\chi}{\sqrt{\omega}}\right\|_{L^{2}} \left[\tilde{\varepsilon}\|\hat{N}_{\hbar}\psi\|_{\mathcal{H}} + b(\tilde{\varepsilon})\|\psi\|_{\mathcal{H}}\right] \\ &\leq \frac{2n}{m_{f}} \left\|\frac{\chi}{\sqrt{\omega}}\right\|_{L^{2}} \tilde{\varepsilon} \|\hat{H}_{02} \ \psi\|_{\mathcal{H}} + C_{\tilde{\varepsilon}} \|\psi\|_{\mathcal{H}} \end{split}$$

where we have used the estimates in Lemmas 3.1 and 3.2. Choose $\tilde{\varepsilon}$ small enough such that $2n \left\| \chi/\sqrt{\omega} \right\|_{L^2} \tilde{\varepsilon} < m_f$. We conclude that there exists $\varepsilon < 1$ and $C_{\varepsilon} \in \mathbf{R}^*_+$ such that

$$\|\hat{H}_1\psi\|_{\mathcal{H}} \le \varepsilon \|\hat{H}_{02}\psi\|_{\mathcal{H}} + C_{\varepsilon} \|\psi\|_{\mathcal{H}}.$$

The operator \hat{H}_{02} is self-adjoint operator and \hat{H}_1 is symmetric operator. Thus by Kato-Rellich theorem, $\hat{H}_{02} + \hat{H}_1$ is self-adjoint on $D(\hat{H}_{02})$. Remark also that by (1.6), we have

$$\|V\psi\|_{L^2} \le \varepsilon \prime \|\hat{H}_{01}\psi\|_{L^2} + b \|\psi\|_{L^2}.$$

Then, again by Kato-Rellich theorem, $\hat{H}_{01} + V$ is self-adjoint on $D(\hat{H}_{01}) \subseteq D(V)$. We also have as a consequence of Kato-Rellich theorem that $\hat{H}_{01} + V \ge -c1$. This means $\hat{H}_{01} + V + c1 \ge 0$. This gives

$$\begin{aligned} \|\hat{H}_{02} \ \psi\|_{\mathcal{H}}^2 &= \langle \psi, \hat{H}_{02}^2 \ \psi \rangle \\ &\leq \langle \psi, \left(\hat{H}_{02} + \hat{H}_{01} + V + c1\right)^2 \psi \rangle \\ &\leq \| \left(\hat{H}_{02} + \hat{H}_{01} + V + c1\right) \psi \|_{\mathcal{H}}^2. \end{aligned}$$

We conclude

$$\|\hat{H}_1\psi\|_{\mathcal{H}} \le \varepsilon \|(\hat{H}_{02} + \hat{H}_{01} + V + c1)\psi\|_{\mathcal{H}} + C(\varepsilon) \|\psi\|_{\mathcal{H}}.$$

Remark that $\hat{H}_{01} + V + c1$ commutes with \hat{H}_{02} . This means $\hat{H}_{01} + V + c1 + \hat{H}_{02}$ is self adjoint in $D(\hat{H}_{01} + V + c1 + \hat{H}_{02}) = D(\hat{H}_{01} + \hat{H}_{02})$. By Kato-Rellich theorem, \hat{H} is self adjoint on $D(\hat{H}_0) = D(\hat{H}_{01} + \hat{H}_{02}) = D(\hat{H}_{01}) \cap D(\hat{H}_{02}).$

3.3. The dynamical equation. The primary objective of this section is to determine the dynamical equation of the quantum system. This equation should converge, as \hbar approaches zero, to a classical dynamical equation that involves the inverse Fourier transform of a specific Wigner measure. To achieve this, in Paragraph 3.3.1, we derive the Duhamel formula for the quantum system. Then, in Paragraph 3.3.2, we expand the commutator within this Duhamel formula.

3.3.1. Duhamel formula. We begin by introducing the Weyl Heisenberg operator, which acts on the entire interacting Hilbert space $\mathcal{H} = L^2(\mathbf{R}^{dn}, \mathbf{C}) \otimes \Gamma_s(\mathcal{G}^0)$, as the following map

$$\xi = (z, \alpha) \in X^0 \equiv \mathbf{C}^{dn} \oplus \mathcal{G}^0 \longmapsto \mathcal{W}(\xi) \equiv \mathcal{W}(z, \alpha) := W_1(z) \otimes W_2(\alpha)$$
(3.7)

where we have introduced with $\Im m\langle z, z' \rangle = q \cdot p' - p \cdot q', \ \forall (p,q), \ (p',q') \in \mathbf{R}^{dn} \times \mathbf{R}^{dn}$:

- the Weyl operator on the particle variable which is defined, for all $(p,q) \in \mathbf{R}^{dn} \times \mathbf{R}^{dn}$ and for $z = q + ip \in \mathbf{C}^{dn}$, as follows:

$$W_1(z) = e^{i \Im m \langle \hat{q} + i\hat{p}, z \rangle} = e^{i(p \cdot \hat{q} - q \cdot \hat{p})};$$
(3.8)

- the Weyl operator on the Fock space $\Gamma_s(L^2(\mathbf{R}^d, \mathbf{C}))$ which is defined for any $\alpha \in$ $L^2(\mathbf{R}^d, \mathbf{C})$ as follows:

$$W_2(\alpha) = e^{\frac{i}{\sqrt{2}}(\hat{a}_{\hbar}(\alpha) + \hat{a}_{\hbar}^*(\alpha))}.$$
(3.9)

The above operators satisfy the following commutation relations

$$W_1(z)W_1(z') = e^{-i\frac{\hbar}{2}\Im m \langle z, z' \rangle} W_1(z+z'), \quad \forall z, z' \in \mathbf{C}^{dn},$$
(3.10)

$$W_2(\alpha)W_2(\beta) = e^{-i\frac{\hbar}{2}\Im m \langle \alpha,\beta \rangle_{L^2}} W_2(\alpha+\beta), \quad \forall \alpha,\beta \in \mathcal{G}^0.$$
(3.11)

Below, we mention several crucial estimates that are necessary to establish a Duhamel formula for the evolved states of a quantum system. The prove of the following identities requires the estimates derived in Lemma 3.1, we refer the reader to [6] for more details on the proof.

Lemma 3.5 (Weyl Heisenberg estimates). There exists a constant C > 0 such that for any $\hbar \in (0,1)$

(i) for any
$$\alpha \in L^2(\mathbf{R}^d, \mathbf{C})$$
 and any $\psi \in D(\hat{N}_{\hbar})$
 $\|(\hat{N}_{\hbar})^{1/2} W_2(\alpha) \psi\|_{\Gamma_s} \leq C \|(\hat{N}_{\hbar} + 1)^{1/2} \psi\|_{\Gamma_s};$

(ii) for any $\alpha \in \mathcal{G}^{1/2}$ and any $\psi \in D(\hat{H}_0)$

$$|(\hat{H}_0)^{1/2} W_2(\alpha) \psi||_{\Gamma_s} \le C ||(\hat{H}_0 + 1)^{1/2} \psi||_{\Gamma_s};$$

(iii) for any $z \in \mathbf{C}^{dn}$ and any $\psi \in D((\hat{p}^2 + \hat{q}^2)^{1/2})$

$$\|(\hat{p}^2 + \hat{q}^2)^{1/2} W_1(z) \psi\|_{L^2(\mathbf{R}^{dn})} \le C \|(\hat{p}^2 + \hat{q}^2 + 1)^{1/2} \psi\|_{L^2(\mathbf{R}^{dn})}.$$

The matter here is to understand the propagation of the density matrices ρ_{\hbar} on the Hilbert space \mathcal{H} . To this end, we define

$$\varrho_{\hbar}(t) = e^{-i\frac{t}{\hbar}\hat{H}}\varrho_{\hbar} \ e^{i\frac{t}{\hbar}\hat{H}} \quad \text{and} \quad \tilde{\varrho}_{\hbar}(t) = e^{i\frac{t}{\hbar}\hat{H}_{02}}\varrho_{\hbar}(t) \ e^{-i\frac{t}{\hbar}\hat{H}_{02}}. \tag{3.12}$$

In order to prove the main results Theorems 1.1 and 1.3, it is necessary to identify the Wigner measures of the evolved state $\rho_{\hbar}(t)$. However, the complexity inherited from the interaction between particles and field makes direct identification unfeasible. Instead, we use the interaction representation $\tilde{\rho}_{\hbar}(t)$, which helps us overcome several nonlinearities that could lead to imprecise formulas. Furthermore, recovering the Wigner measures of $\rho_{\hbar}(t)$ from those of $\tilde{\rho}_{\hbar}(t)$ is not difficult. To this end, we start below derivation of the quantum dynamical system.

Proposition 3.6. Assume that (1.6) and $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). Then for all $\xi \in X^{1/2}$, for all $\hbar \in (0,1)$ and for all $t, t_0 \in \mathbf{R}$, we have

$$\operatorname{Tr}\left[\mathcal{W}(\xi)\tilde{\varrho}_{\hbar}(t)\right] = \operatorname{Tr}\left[\mathcal{W}(\xi)\tilde{\varrho}_{\hbar}(t_{0})\right] - \frac{i}{\hbar}\int_{t_{0}}^{t}\operatorname{Tr}\left(\left[\mathcal{W}(\xi),\hat{H}_{I}(s)\right]\tilde{\varrho}_{\hbar}(s)\right)ds,\tag{3.13}$$

where

$$\hat{H}_{I}(s) := e^{i\frac{s}{\hbar}\hat{H}_{02}}(\hat{H} - \hat{H}_{02}) \ e^{-i\frac{s}{\hbar}\hat{H}_{02}}.$$
(3.14)

Proof. By Duhamel's formula, we have

$$\operatorname{Tr}\left[\mathcal{W}(\xi)\tilde{\varrho}_{\hbar}(t)\right] = \operatorname{Tr}\left[\mathcal{W}(\xi)\tilde{\varrho}_{\hbar}(t_{0})\right] + \int_{t_{0}}^{t} \frac{d}{ds}\operatorname{Tr}\left[\mathcal{W}(\xi)\ \tilde{\varrho}_{\hbar}(s)\right]ds$$

We have also

$$\frac{d}{dt} \operatorname{Tr} \left[\mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar}(t) \right] = \lim_{s \to t} \frac{\operatorname{Tr} \left[\mathcal{W}(\xi) \ \left(\tilde{\varrho}_{\hbar}(t) - \tilde{\varrho}_{\hbar}(s) \right) \right]}{t - s}$$

Let $S = (\hat{H}_0 + 1)^{1/2}$. We start by

$$\begin{aligned} &\operatorname{Tr} \left[\mathcal{W}(\xi) \left(\tilde{\varrho}_{\hbar}(t) - \tilde{\varrho}_{\hbar}(s) \right) \right] \\ &= \operatorname{Tr} \left[\mathcal{W}(\xi) \left(e^{i\frac{t}{\hbar}\hat{H}_{02}} e^{-i\frac{t}{\hbar}\hat{H}} \varrho_{\hbar} e^{i\frac{t}{\hbar}\hat{H}} e^{-i\frac{t}{\hbar}\hat{H}_{02}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}} \varrho_{\hbar} e^{i\frac{s}{\hbar}\hat{H}} e^{-i\frac{s}{\hbar}\hat{H}_{02}} \right) \right] \\ &= \operatorname{Tr} \left[\mathcal{W}(\xi) \left(e^{i\frac{t}{\hbar}\hat{H}_{02}} e^{-i\frac{t}{\hbar}\hat{H}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}} \right) \varrho_{\hbar} e^{i\frac{t}{\hbar}\hat{H}} e^{-i\frac{t}{\hbar}\hat{H}_{02}} \right] \\ &+ \operatorname{Tr} \left[\mathcal{W}(\xi) e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}} \tilde{\varrho}_{\hbar} \left(e^{i\frac{t}{\hbar}\hat{H}} e^{-i\frac{t}{\hbar}\hat{H}_{02}} - e^{i\frac{s}{\hbar}\hat{H}} e^{-i\frac{s}{\hbar}\hat{H}_{02}} \right) \right] \\ &= \operatorname{Tr} \left[S^{-1} \ \mathcal{W}(\xi) \ S \ S^{-1} \left(e^{i\frac{t}{\hbar}\hat{H}_{02}} e^{-i\frac{t}{\hbar}\hat{H}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}} \right) \varrho_{\hbar} \ S \ S^{-1} \ e^{-i\frac{t}{\hbar}\hat{H}_{02}} \ S \right] \\ &+ \operatorname{Tr} \left[\mathcal{W}(\xi) \ e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}} \ \tilde{\varrho}_{\hbar} \ S \ S^{-1} \left(e^{i\frac{t}{\hbar}\hat{H}_{02}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} \right) \right] \end{aligned}$$

Remark that each step makes sense. Indeed, we have that

 $\mathcal{W}(\xi), \ e^{\frac{it}{\hbar}\hat{H}_{02}}, \ e^{\frac{it}{\hbar}\hat{H}}, \ S^{-1} \ e^{i\frac{t}{\hbar}\hat{H}} \ S, \ S^{-1} \ e^{-i\frac{t}{\hbar}\hat{H}_{02}} \ S \in \mathcal{L}(\mathcal{H}), \quad \varrho_{\hbar}, \ \varrho_{\hbar}(\hat{H}_{0}+1) \in \mathcal{L}^{1}(\mathcal{H}).$

$$\lim_{s \to t} S^{-1} \frac{\left(e^{i\frac{t}{\hbar}\hat{H}} e^{-i\frac{t}{\hbar}\hat{H}_{02}} - e^{i\frac{s}{\hbar}\hat{H}} e^{-i\frac{s}{\hbar}\hat{H}_{02}}\right)}{t - s} = \frac{i}{\hbar} S^{-1} e^{i\frac{t}{\hbar}\hat{H}} \left(\hat{H} - \hat{H}_{02}\right) e^{-i\frac{t}{\hbar}\hat{H}_{02}};$$
$$\lim_{s \to t} S^{-1} \frac{\left(e^{i\frac{t}{\hbar}\hat{H}_{02}} e^{-i\frac{t}{\hbar}\hat{H}} - e^{i\frac{s}{\hbar}\hat{H}_{02}} e^{-i\frac{s}{\hbar}\hat{H}}\right)}{t - s} = -\frac{i}{\hbar} S^{-1} e^{i\frac{t}{\hbar}\hat{H}_{02}} \left(\hat{H} - \hat{H}_{02}\right) e^{-i\frac{t}{\hbar}\hat{H}}.$$

Plugging these limits in the Duhamel's formula, we get the desired result.

3.3.2. The commutator expansion. The aim of this subsection is to expand the commutator $[\mathcal{W}(\xi), \hat{H}_I(s)]$ in the above Duhamel formula (3.13) in terms of the parameter $\hbar \in (0, 1)$.

Lemma 3.7 (Time evolved equation of $\hat{H}_I(s)$). For any $s \in \mathbf{R}$, the time evolved interaction term $\hat{H}_I(s)$ takes the following form

$$\hat{H}_{I}(s) = \sum_{j=1}^{n} f_{j}(\hat{p}_{j}) + V(\hat{q}) + \sum_{j=1}^{n} \hat{a}_{\hbar}(g_{j}(s)) + \hat{a}_{\hbar}^{*}(g_{j}(s)), \qquad (3.15)$$

where we have introduced

$$g_j(s) \equiv g_j(s)(\hat{q}) := \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot \hat{q}_j + is\omega(k)}.$$
(3.16)

Proof. We have

$$\hat{H} - \hat{H}_{02} = \sum_{j=1}^{n} f_j(\hat{p}_j) + V(\hat{q}) + \sum_{j=1}^{n} \hat{a}_\hbar(g_j) + \hat{a}^*_\hbar(g_j),$$

where the function g_j is given by

$$g_j \equiv g_j(\hat{q}) := \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{-2\pi i k \cdot \hat{q}_j}.$$
(3.17)

Then, we have with $\hat{q} = (\hat{q}_1, \cdots, \hat{q}_n)$

$$\hat{H}_{I}(s) = e^{i\frac{s}{\hbar}\hat{H}_{02}} \Big(\sum_{j=1}^{n} f_{j}(\hat{p}_{j}) + V(\hat{q}) + \sum_{j=1}^{n} \hat{a}_{\hbar}(g_{j}) + \hat{a}_{\hbar}^{*}(g_{j})\Big) e^{-i\frac{s}{\hbar}\hat{H}_{02}}.$$

It is sufficient then to look at the following identity

$$e^{i\frac{s}{\hbar}\hat{H}_{02}} \hat{a}^{\sharp}(g_j) e^{-i\frac{s}{\hbar}\hat{H}_{02}} = \hat{a}^{\sharp}(g_j(s)).$$

Now, since the Weyl operator $\mathcal{W}(\xi)$ is a unitary operator, we have

$$\frac{1}{\hbar} \left[\mathcal{W}(\xi), \hat{H}_I(s) \right] = \frac{1}{\hbar} \left(\mathcal{W}(\xi) \hat{H}_I(s) \mathcal{W}(\xi)^* - \hat{H}_I(s) \right) \mathcal{W}(\xi).$$
(3.18)

Lemma 3.8 (Expression for the commutators). For any $s \in \mathbf{R}$ and $\xi = (p_0, q_0, \alpha_0) \in X^{1/2}$, the following holds true with $q_0 = (q_{01}, \dots, q_{0n})$ and $p_0 = (p_{01}, \dots, p_{0n})$

$$\mathcal{W}(\xi)\hat{H}_{I}(s)\mathcal{W}(\xi)^{*} = \sum_{j=1}^{n} f_{j}(\hat{p}_{j} - \hbar p_{0j}) + V(\hat{q} - \hbar q_{0}) + \sum_{j=1}^{n} \hat{a}_{\hbar}(\tilde{g}_{j}(s)) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j}(s)) + \frac{i\hbar}{\sqrt{2}} \Big(\langle \alpha_{0}, \tilde{g}_{j}(s) \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} - \langle \tilde{g}_{j}(s), \alpha_{0} \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} \Big),$$

where we have introduced

$$\tilde{g}_j(s) := \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot (\hat{q}_j - \hbar q_{0j}) + is\omega(k)} = e^{2\pi i k \cdot q_{0j}\hbar} g_j(s).$$
(3.19)

Proof. Let $\hat{q}_j = (\hat{q}_j^{\nu})_{\nu=1,\cdots d}$ and $\hat{p}_j = (\hat{p}_j^{\nu})_{\nu=1,\cdots d}$. The results follow from the following identities

$$W_1(z_0) \ \hat{q}_j^{\nu} \ W_1(z_0)^* = \hat{q}_j^{\nu} - \hbar q_{0j}^{\nu}, \tag{3.20}$$

$$W_1(z_0) \ \hat{p}_j^{\nu} \ W_1(z_0)^* = \hat{p}_j^{\nu} - \hbar p_{0j}^{\nu}, \tag{3.21}$$

$$W_2(\alpha_0) \ \hat{a}_{\hbar}^*(f) \ W_2(\alpha)^* = \hat{a}_{\hbar}^*(f) + \frac{i\hbar}{\sqrt{2}} \langle \alpha_0, f \rangle_{L^2}, \tag{3.22}$$

$$W_2(\alpha_0) \ \hat{a}_{\hbar}(f) \ W_2(\alpha)^* = \hat{a}_{\hbar}(f) - \frac{i\hbar}{\sqrt{2}} \langle f, \alpha_0 \rangle_{L^2}.$$
(3.23)

We start proving the first identity (3.20). Recall from (3.10) that we have

$$W_1(z_0) = e^{i(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})}, \quad W_1(z_0)^* = e^{-i(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})}$$

Define

$$K(t) := e^{it(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})} \hat{q}_i^{\nu} e^{-it(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})}$$

Since \hat{q} and \hat{p} are self adjoint operators, we claim using Taylor expansions that

$$K(t) = K(0) + tK'(0).$$
(3.24)

Indeed, we have, using the commutation relation $[\hat{q}_{j}^{\nu}, \hat{p}_{j}^{\nu}] = i\hbar$, that

$$K'(0) = \frac{d}{dt}K(t)|_{t=0} = e^{it(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})} i[(p_0 \cdot \hat{q} - q_0 \cdot \hat{p}), \hat{q}_j^{\nu}] e^{-it(p_0 \cdot \hat{q} - q_0 \cdot \hat{p})}|_{t=0} = -\hbar q_{0j}^{\nu}.$$

This implies that $K^r(0) = 0$, for all $r \ge 2$. Take t = 1 in (3.24) and since $K(0) = \hat{q}_j^{\nu}$, we get (3.20). Similarly, we can prove the identity (3.21). Also the two identities (3.22) and (3.23) can be proved by simiar way using the commutation relations on the Fock space. In particular, the identity (3.20) gives

$$W_1(z_0) \ g_j(s) \ W_1(z_0)^* = \tilde{g}_j(s).$$

Lemma 3.9 (The expansion of the commutator). For any $s \in \mathbf{R}$ and $\xi = (p_0, q_0, \alpha_0) \in X^{1/2}$, we have the following expansion of the commutator in terms of the semiclassical parameter $\hbar \in (0, 1)$

$$\frac{1}{\hbar} \left[\mathcal{W}(\xi), \hat{H}_I(s) \right] = \left(\mathcal{B}_0(s, \hbar, \xi) + \hbar \mathcal{B}_1(s, \hbar, \xi) \right) \mathcal{W}(\xi).$$
(3.25)

The two terms B_0 and B_1 are identified as follows

$$B_{0}(s,\hbar,\xi) := -\sum_{j=1}^{n} \nabla f_{j}(\hat{p}_{j}) \cdot p_{0j} - \nabla V(\hat{q}) \cdot q_{0}$$

$$+ \sum_{j=1}^{n} \hat{a}_{\hbar} \Big(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar} \Big) + \hat{a}_{\hbar}^{*} \Big(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar} \Big)$$

$$+ \sum_{j=1}^{n} \frac{i}{\sqrt{2}} \Big(\langle \alpha_{0}, \tilde{g}_{j}(s) \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} - \langle \tilde{g}_{j}(s), \alpha_{0} \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} \Big),$$

$$B_{1}(s,\hbar,\xi) := \Theta_{1}(\hbar,\xi) + \Theta_{2}(\hbar,\xi), \qquad (3.27)$$

where Θ_1 and Θ_2 are identified below in the proof. Moreover, we have also the following estimates

$$\|(\hat{H}_{0}+1)^{-1/2}B_{0}(s,\hbar,\xi)(\hat{H}_{0}+1)^{-1/2}\|_{\mathcal{L}(\mathcal{H})} \lesssim \left(\|\chi\|_{L^{2}}+\left\|\sqrt{\omega}\,\chi\right\|_{L^{2}}\right)\,\|\xi\|_{X^{0}},\tag{3.28}$$

$$\|(\hat{H}_0+1)^{-1/2}\mathcal{B}_1(s,\hbar,\xi)(\hat{H}_0+1)^{-1/2}\|_{\mathcal{L}(\mathcal{H})} \lesssim \|\xi\|_{X^0}^2.$$
(3.29)

Proof. Exploiting Lemma 3.7 and Lemma 3.8 inside (3.18), the commutator expansion becomes

$$\frac{1}{\hbar} \left[\mathcal{W}(\xi), \hat{H}_{I}(s) \right] = \frac{1}{\hbar} \left[\sum_{j=1}^{n} \left(f_{j}(\hat{p}_{j} - \hbar p_{0j}) - f_{j}(\hat{p}_{j}) \right) + V(\hat{q} - \hbar q_{0}) - V(\hat{q}) + \sum_{j=1}^{n} \left(\hat{a}_{\hbar}(\tilde{g}_{j}(s) - g_{j}(s)) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j}(s) - g_{j}(s)) \right) + \frac{i\hbar}{\sqrt{2}} \sum_{j=1}^{n} \left(\langle \alpha_{0}, \tilde{g}_{j}(s) \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} - \langle \tilde{g}_{j}(s), \alpha_{0} \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} \right) \right] \mathcal{W}(\xi).$$
(3.30)

We start first by expanding the first line and then proving some estimates for the remaining terms. Let $X \in \mathbf{R}^d$ and $Y = (Y_1, \dots, Y_n) \in \mathbf{R}^{dn}$. We apply Taylor series to the two functions

$$t \longrightarrow A(t) := f_j(X - t \hbar p_{0j}),$$

$$t \longrightarrow B(t) := V(Y - t \hbar q_0).$$

We get

$$A(t) = A(0) + tA'(0) + \int_0^t A''(s)(t-s) \ ds,$$

and

$$B(t) = B(0) + tB'(0) + \int_0^t B''(s)(t-s) \, ds.$$

Let t = 1 in the above formulas and since \hat{p} and \hat{q} are self adjoint operators, we get

$$f_j(\hat{p}_j - \hbar p_{0j}) = f_j(\hat{p}_j) - \hbar \nabla f_j(\hat{p}_j) \cdot p_{0j} + \hbar^2 \underbrace{\int_0^1 p_{0j}^T H_{f_j}(\hat{p}_j - \hbar p_{0j}s) p_{0j} (1 - s) ds}_{:=\Theta_1(\hbar,\xi)}, \quad (3.31)$$

$$V(\hat{q} - \hbar q_0) = V(\hat{q}) - \hbar \nabla V(\hat{q}) \cdot q_0 + \hbar^2 \underbrace{\int_0^1 q_0^T H_V(\hat{q} - \hbar q_0 s) q_0 (1 - s) ds}_{:=\Theta_2(\hbar,\xi)},$$
(3.32)

where the notation \cdot^{T} represents the transpose. Moreover, the two terms H_{f_j} and H_V are respectively the Hessian matrices related to f_j and V. This implies

$$f_j(\hat{p}_j - \hbar p_{0j}) - f_j(\hat{p}_j) = -\hbar \nabla f_j(\hat{p}_j) \cdot p_{0j} + \hbar^2 \Theta_1(\hbar, \xi), \qquad (3.33)$$

$$V(\hat{q} - \hbar q_0) - V(\hat{q}) = -\hbar \nabla V(\hat{q}) \cdot q_0 + \hbar^2 \Theta_2(\hbar, \xi).$$
(3.34)

And thus, using (3.33)-(3.34), the commutator is expanded as indicated in (3.25). Now, to obtain the two estimates (3.28) and (3.29), we need first to prove that the function

$$F_j(\hbar, s) := (\tilde{g}_j(s) - g_j(s))/\hbar : L^2(\mathbf{R}^d, dx_j) \longrightarrow L^2(\mathbf{R}^d, dx_j) \otimes L^2(\mathbf{R}^d, dk)$$

is bounded uniformly in $\hbar \in (0, 1)$. Indeed, we have for all $\psi \in L^2(\mathbf{R}^d, dx_j)$

$$\begin{split} \|F_{j}(\hbar,s) \psi\|_{L^{2}_{x_{j}}\otimes L^{2}_{k}}^{2} &= \int_{\mathbf{R}^{d}} \int_{\mathbf{R}^{d}} \left| \left(F_{j}(\hbar,s) \psi\right)(x_{j},k) \right|^{2} dx_{j} dk \\ &= \int_{\mathbf{R}^{d}} \int_{\mathbf{R}^{d}} \left| \left[g_{j}(s) \left(\frac{e^{2\pi i k \cdot q_{0j}\hbar} - 1}{\hbar}\right)\psi\right](x_{j},k) \right|^{2} dx_{j} dk \\ &= \int_{\mathbf{R}^{d}} \int_{\mathbf{R}^{d}} \left|\frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot \hat{q}_{j} + is\omega(k)} \left(\frac{e^{2\pi i k \cdot q_{0j}\hbar} - 1}{\hbar}\right)\psi(x_{j}) \right|^{2} dx_{j} dk. \end{split}$$

Now, with the aid of Fubini and the estimate $|e^{iy} - 1| \le \sqrt{2}|y|$, we find that

$$\begin{aligned} \|F_{j}(\hbar,s) \ \psi\|_{L^{2}_{x_{j}}\otimes L^{2}_{k}}^{2} &\leq 8\pi^{2} \ \|\xi\|_{X^{0}}^{2} \ \int_{\mathbf{R}^{d}} \int_{\mathbf{R}^{d}} \left|\sqrt{\omega(k)}\chi(k)\right|^{2} \ \left|\psi(x_{j})\right|^{2} \ dx_{j} \ dk \\ &= 8\pi^{2} \ \|\chi\|_{\mathcal{G}^{1/2}}^{2} \ \|\xi\|_{X^{0}}^{2} \ \|\psi\|_{L^{2}_{x_{j}}}^{2}. \end{aligned}$$

We get finally, with some C > 0, that

$$\|F_j(\hbar, s)\|_{\mathcal{L}(L^2_{x_j}, L^2_{x_j} \otimes L^2_k)} \le C \|\chi\|_{\mathcal{G}^{1/2}} \|\xi\|_{X^0}.$$
(3.35)

Now, using the estimates in Lemma 3.1 on the creation-annihilation operators together with the above estimate for $F_j(\hbar, s)$, we can easily prove (3.28). It is also not hard to see that (3.29) hold true as a consequence of the fact that the Hessian matrices of f_j and V are bounded.

Our focus is on taking the classical limit $\hbar \to 0$. To accomplish this, it is crucial to establish a uniform bound on the expansion derived in Lemma 3.9, particularly for the remainder term. Let $S = (\hat{H}_0 + 1)^{1/2}$, we have

$$\operatorname{Tr}\left(\frac{1}{\hbar} \left[\mathcal{W}(\xi), \hat{H}_{I}(s)\right] \tilde{\varrho}_{\hbar}(s)\right) = \operatorname{Tr}\left[\underbrace{S^{-1} \operatorname{B}_{0}(s, \hbar, \xi) S^{-1}}_{\in \mathcal{L}(\mathcal{H})} \underbrace{S \operatorname{\mathcal{W}}(\xi) S^{-1}}_{\in \mathcal{L}(\mathcal{H})} \underbrace{S \operatorname{\tilde{\varrho}}_{\hbar}(s) S}_{\in \mathcal{L}^{1}(\mathcal{H})}\right] + \hbar \operatorname{Tr}\left[\underbrace{S^{-1}\operatorname{B}_{1}(\hbar, s, \xi) S^{-1}}_{\in \mathcal{L}(\mathcal{H})} \underbrace{S \operatorname{\mathcal{W}}(\xi) S^{-1}}_{\in \mathcal{L}(\mathcal{H})} \underbrace{S \operatorname{\tilde{\varrho}}_{\hbar}(s) S}_{\in \mathcal{L}^{1}(\mathcal{H})}\right]$$
(3.36)

- \triangleleft Lemma 3.9 assures that the first term in each of the above two lines in (3.36) is bounded.
- \triangleleft The Weyl-Heisenberg operator estimates presented in Lemma 3.5 guarantee that the bound of the second term in the above two lines in (3.36) holds.
- ⊲ The bound of the last term in each of the above two lines in (3.36) follows from Assumption (1.8) and (1.9) in conjunction with the equivalent relation between \hat{H} and \hat{H}_0 outlined in Lemma 3.3.

Our next step is to take the limit in the Duhamel formula (3.13) as \hbar approaches zero. Using the above arguments, we can disregard the remainder term when passing to the limit $\hbar \to 0$ in the Duhamel formula (3.13). We achieve this in the next section by extracting a subsequence.

4. EXISTENCE OF WIGNER MEASURE

According to Definition 1.2, the Wigner measures of $\tilde{\varrho}_{\hbar}(t)$ is obtained by taking limits of the following map:

$$\xi \to \operatorname{Tr} \left[\mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar}(t) \right]. \tag{4.1}$$

Thus, the first task is to verify that the Wigner measure associated to the above map is unique for all times. It is worth noting that, given our assumptions on the initial states $(\varrho_{\hbar})_{\hbar \in (0,1)}$, the associated set of Wigner measures

$$\mathcal{M}(\varrho_{\hbar}, \ \hbar \in (0,1))$$

is non-empty. To ensure that the sets of Wigner measures

$$\mathcal{M}(\varrho_{\hbar}(t), \ \hbar \in (0,1)) \text{ and } \mathcal{M}(\tilde{\varrho}_{\hbar}(t), \ \hbar \in (0,1))$$

are also non-empty, it is crucial to demonstrate that assumptions (1.8) and (1.9) can be uniformly propagated in time by both families of states $(\rho_{\hbar}(t))_{\hbar \in (0,1)}$ and $(\tilde{\rho}_{\hbar}(t))_{\hbar \in (0,1)}$. This is established in Subsection 4.1. Subsequently, in Subsection 4.2, we prove that the map (4.1) has a unique limit that holds for all times in compact interval.

4.1. **Propagation of assumptions.** In order to establish the existence of a unique Wigner measure that holds for all times, we demonstrate that if an initial state ρ_{\hbar} is localized uniformly in \hbar , then it will remain localized uniformly with respect to the semiclassical parameter $\hbar \in (0, 1)$ for all times in compact interval. We prove this result separately for particle operators in Paragraph 4.1.1 and for field operators in Paragraph 4.1.2. Finally, in Paragraph 4.1.3, we establish that both families of states $(\rho_{\hbar}(t))_{\hbar \in (0,1)}$ and $(\tilde{\rho}_{\hbar}(t))_{\hbar \in (0,1)}$ uniformly satisfy (1.8) and (1.9) for all times.

4.1.1. Position and Momentum operator estimates. In this part, we prove some uniform estimates (in \hbar) related to the two operators \hat{p}^2 and \hat{q}^2 .

Lemma 4.1 (Position operator's estimate). Assume that (1.6) and $\omega^{-1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Then, there exists constants $C_1, C_2 > 0$ such that for all $\psi \in D(\hat{H}_0^{1/2}) \cap D(\hat{q})$, all $t \in \mathbf{R}$ and all $\hbar \in (0, 1)$:

$$\langle e^{-i\frac{t}{\hbar}\hat{H}}\psi, \ \hat{q}^2 \ e^{-i\frac{t}{\hbar}\hat{H}}\psi \rangle \le C_1 \langle \psi, \ (\hat{H}_0 + \hat{q}^2 + 1) \ \psi \rangle \ e^{C_2|t|}.$$
 (4.2)

Proof. Let $\Theta_1(t) := \langle e^{-i\frac{t}{\hbar}\hat{H}}\psi, \ \hat{q}^2 e^{-i\frac{t}{\hbar}\hat{H}}\psi \rangle$. We have

$$\Theta_1(t) = \Theta_1(0) + \int_0^t \dot{\Theta}_1(s) \ ds$$

Then Stone's Theorem implies that

$$\dot{\Theta}_1(t) = \frac{1}{\hbar} \left\langle e^{-i\frac{t}{\hbar}\hat{H}}\psi, \ i[\hat{H}, \hat{q}^2] \ e^{-i\frac{t}{\hbar}\hat{H}}\psi \right\rangle.$$

Now, using some commutation relations, we get

$$i[\hat{H}, \hat{q}^2] = i \sum_{j=1}^n [f_j(\hat{p}_j), \hat{q}_j^2] = \hbar \sum_{j=1}^n \left[\nabla f_j(\hat{p}_j) \cdot \hat{q}_j + \hat{q}_j \cdot \nabla f_j(\hat{p}_j) \right].$$

Define $\psi(t) := e^{-i\frac{t}{\hbar}\hat{H}}\psi$. Since \hat{q}_j and $\nabla f_j(\hat{p}_j)$ are self-adjoint operators, we have the following estimates

$$\langle \psi(t), \ \nabla f_{j}(\hat{p}_{j}) \cdot \hat{q}_{j} \ \psi(t) \rangle \leq \| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \| \ \| \hat{q}_{j} \ \psi(t) \| \leq \frac{1}{2} \left[\| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \|^{2} + \| \hat{q}_{j} \ \psi(t) \|^{2} \right]$$

$$= \frac{1}{2} \left[\| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \|^{2} + \Theta_{1}(t) \right],$$

$$\langle \psi(t), \ \hat{q}_{j} \cdot \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \rangle \leq \| \hat{q}_{j} \ \psi(t) \| \ \| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \|^{2} + \| \hat{q}_{j} \ \psi(t) \|^{2} \right]$$

$$= \frac{1}{2} \left[\| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \|^{2} + \| \hat{q}_{j} \ \psi(t) \|^{2} \right]$$

$$= \frac{1}{2} \left[\| \nabla f_{j}(\hat{p}_{j}) \ \psi(t) \|^{2} + \Theta_{1}(t) \right],$$

$$(4.4)$$

where we have used the identity $2a \cdot b \leq a^2 + b^2$. At this stage, we have to consider separately the two cases: the semi-relativistic and the non-relativistic case since the function ∇f_j is bounded in the first case and not in the second one.

For semi-relativistic case:

Note that $\nabla f_j(\hat{p}_j)$ is a bounded operator. This implies that for some $c_1 > 0$, we have

$$\|\nabla f_j(\hat{p}_j) \psi(t)\|^2 \le c_1 \|\psi\|^2.$$

This gives

$$\Theta_1(t) \le \Theta_1(0) + c_1 \langle \psi, \psi \rangle \ t + \int_0^t \Theta_1(s) \ ds$$

Now using Gronwall's lemma and the estimate $te^t \leq e^{ct}$ for some c > 0, we find with some $C_1, C_2 > 0$

$$\Theta_1(t) \le \left[\Theta_1(0) + c_1 \langle \psi, \psi \rangle \ t\right] \ e^t \le C_1 \langle \psi, \ (\hat{q}^2 + 1) \ \psi \rangle \ e^{C_2|t|} \le C_1 \langle \psi, \ (\hat{H}_0 + \hat{q}^2 + 1) \ \psi \rangle \ e^{C_2|t|}.$$

For non-relativistic case:

We have $\nabla f_j(\hat{p}_j) = \hat{p}_j/M_j$. This implies (4.3) and (4.4) become

$$\begin{split} \langle \psi(t), \ \nabla f_j(\hat{p}_j) \cdot \hat{q}_j \ \psi(t) \rangle &\leq \frac{1}{2} \Big[\langle \psi(t), \ \frac{\hat{p}_j^2}{M_j^2} \ \psi(t) \rangle + \ \Theta_1(t) \Big] \\ &\leq \frac{1}{M_j} \ \langle \psi(t), \ \hat{H}_0 \ \psi(t) \rangle + \ \frac{1}{2} \Theta_1(t) \\ \langle \psi(t), \ \hat{q}_j \cdot \nabla f_j(\hat{p}_j) \ \psi(t) \rangle &\leq \frac{1}{2} \Big[\langle \psi(t), \ \frac{\hat{p}_j^2}{M_j^2} \ \psi(t) \rangle + \ \Theta_1(t) \Big] \\ &\leq \frac{1}{M_j} \ \langle \psi(t), \ \hat{H}_0 \ \psi(t) \rangle + \ \frac{1}{2} \Theta_1(t). \end{split}$$

By Lemma 3.3, we have

 $\langle e^{-i\frac{t}{\hbar}\hat{H}}\psi,\hat{H}_0 \ e^{-i\frac{t}{\hbar}\hat{H}}\psi\rangle \lesssim \langle e^{-i\frac{t}{\hbar}\hat{H}}\psi,(\hat{H}+1) \ e^{-i\frac{t}{\hbar}\hat{H}}\psi\rangle = \langle \psi,(\hat{H}+1) \ \psi\rangle \lesssim \langle \psi,(\hat{H}_0+1) \ \psi\rangle.$ This leads with some $c_2 \in \mathbf{R}^*_+$ to the following inequality

$$\Theta_1(t) \le \Theta_1(0) + c_2 \langle \psi, (\hat{H}_0 + 1) \psi \rangle \ t + \int_0^t \Theta_1(s) \ ds$$

Now using Gronwall's Lemma and the estimates $te^{ct} \leq e^{c't}$, we find with some C_1 , $C_2 > 0$

$$\begin{aligned} \Theta_1(t) &\leq \left[\Theta_1(0) + c_2 \langle \psi, (\hat{H}_0 + 1) | \psi \rangle | t \right] e^{\int_0^t 1 | ds} \leq C_1 \langle \psi, (\hat{H}_0 + \hat{q}^2 + 1) \psi \rangle | e^{C_2 |t|} \\ &\leq C_1 \langle \psi, (\hat{H}_0 + \hat{q}^2 + 1) \psi \rangle | e^{C_2 |t|}. \end{aligned}$$

Now, we give some uniform estimates for the momentum operator just in the semi-relativistic case: $f_j(\hat{p}_j) = \sqrt{\hat{p}_j^2 + M_j^2}$.

Lemma 4.2 (Momentum operator's estimate). Assume that (1.6) and $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Then, there exists constants $C_1, C_2 > 0$ such that for all $\psi \in D(\hat{H}_0^{1/2}) \cap D(\hat{p})$, all $t \in \mathbf{R}$ and all $\hbar \in (0, 1)$:

$$\langle e^{-i\frac{t}{\hbar}\hat{H}}\psi, \ \hat{p}^2 \ e^{-i\frac{t}{\hbar}\hat{H}}\psi\rangle \le C_1\langle\psi, \ (\hat{H}_0 + \hat{p}^2 + 1) \ \psi\rangle \ e^{C_2|t|}.$$
 (4.5)

Proof. Define

$$\Theta_2(t) := \langle \psi(t), \hat{p}^2 \psi(t) \rangle, \qquad \psi(t) := e^{-i\frac{t}{\hbar}\hat{H}} \psi.$$

We have that the map $t \to \Theta_2(t)$ is differentiable with

$$\dot{\Theta}_2(t) = \frac{i}{\hbar} \langle \psi(t), [\hat{H}, \hat{p}^2] \psi(t) \rangle$$

Then, Duhamel formula implies that

 \hat{a}^{\sharp}_{\hbar}

$$\Theta_2(t) = \Theta_2(0) + \int_0^t \dot{\Theta}_2(s) \ ds.$$

Let us compute first the explicit expression for the function $\dot{\Theta}_2(t)$. To do that, we need first to deal with the commutator $[\hat{H}, \hat{p}^2]$. Indeed, we have

$$\begin{bmatrix} \hat{H}, \hat{p}^2 \end{bmatrix} = \begin{bmatrix} d\Gamma(\omega) + \sum_{j=1}^n \sqrt{\hat{p}_j^2 + M_j^2} + V(\hat{q}) + \hat{H}_1, \hat{p}^2 \end{bmatrix}$$
$$= \begin{bmatrix} V(\hat{q}), \hat{p} \end{bmatrix} \hat{p} + \hat{p} \begin{bmatrix} V(\hat{q}), \hat{p} \end{bmatrix} + \begin{bmatrix} \hat{H}_1, \hat{p} \end{bmatrix} \hat{p} + \hat{p} \begin{bmatrix} \hat{H}_1, \hat{p} \end{bmatrix}$$

Recall that

$$(G) = \sum_{j=1}^{n} \hat{a}_{\hbar}^{\sharp}(g_{j}(\hat{q})), \qquad g_{j}(\hat{q})(k) := \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot \hat{q}_{j}}.$$

We can then assert that

$$\hat{a}_{\hbar}(G) = \sum_{j=1}^{n} \int_{\mathbf{R}^{d}} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{2\pi i k \cdot \hat{q}_{j}} \hat{a}_{\hbar}(k) dk =: \sum_{j=1}^{n} B_{j}(\hat{q}_{j}),$$
$$\hat{a}_{\hbar}^{*}(G) = \sum_{j=1}^{n} \int_{\mathbf{R}^{d}} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot \hat{q}_{j}} \hat{a}_{\hbar}^{*}(k) dk =: \sum_{j=1}^{n} B_{j}^{*}(\hat{q}_{j}),$$

where $q_j \to B_j^{\sharp}(q_j)$ is analytic function. We know that for any analytic function F

$$[F(\hat{q}_j), \hat{p}_j] = i\hbar \; \frac{\partial F(\hat{q}_j)}{\partial q_j},$$

This gives

$$\begin{split} & [V(\hat{q}), \hat{p}] \ \hat{p} = \sum_{j=1}^{n} i\hbar \ \nabla_{q_{j}} V(\hat{q}) \cdot \hat{p}_{j}, \\ & \hat{p} \ [V(\hat{q}), \hat{p}] = \sum_{j=1}^{n} i\hbar \ \hat{p}_{j} \cdot \nabla_{q_{j}} V(\hat{q}), \\ & [\hat{a}_{\hbar}(G), \hat{p}] \ \hat{p} = \left[\sum_{j=1}^{n} B_{j}(\hat{q}_{j}), \hat{p}\right] \ \hat{p} = \sum_{j=1}^{n} i\hbar \ \frac{\partial B_{j}(\hat{q}_{j})}{\partial q_{j}} \cdot \hat{p}_{j} = \sum_{j=1}^{n} i\hbar \ \hat{a}_{\hbar}(\tilde{g}_{j}) \cdot \hat{p}_{j}, \\ & \hat{p} \ [\hat{a}_{\hbar}(G), \hat{p}] = \sum_{j=1}^{n} i\hbar \ \hat{p}_{j} \cdot \hat{a}_{\hbar}(\tilde{g}_{j}), \\ & [\hat{a}_{\hbar}^{*}(G), \hat{p}] \ \hat{p} = \left[\sum_{j=1}^{n} B_{j}^{*}(\hat{q}_{j}), \hat{p}\right] \ \hat{p} = \sum_{j=1}^{n} i\hbar \ \frac{\partial B_{j}^{*}(\hat{q}_{j})}{\partial q_{j}} \cdot \hat{p}_{j} = \sum_{j=1}^{n} i\hbar \ \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \cdot \hat{p}_{j}, \\ & \hat{p} \ [\hat{a}_{\hbar}^{*}(G), \hat{p}] = \sum_{j=1}^{n} i\hbar \ \hat{p}_{j} \cdot \hat{a}_{\hbar}^{*}(\tilde{g}_{j}), \end{split}$$

where we have introduced the term \tilde{g}_j as follows

$$\tilde{g}_j := -2\pi i k \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{-2\pi i k \cdot \hat{q}_j}.$$

This implies that

$$\left[\hat{H}, \ \hat{p}^{2}\right] = i\hbar \ \sum_{j=1}^{n} \left[\hat{p}_{j} \cdot \left(\nabla_{q_{j}}V(\hat{q}) + \hat{a}_{\hbar}(\tilde{g}_{j}) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j})\right) + \left(\nabla_{q_{j}}V(\hat{q}) + \hat{a}_{\hbar}(\tilde{g}_{j}) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j})\right) \cdot \hat{p}_{j}\right].$$

We conclude that

$$\dot{\Theta}_{2}(t) = -\sum_{j=1}^{n} \left[\langle \psi(t), \ \hat{p}_{j} \cdot \left(\nabla_{q_{j}} V(\hat{q}) + \hat{a}_{\hbar}(\tilde{g}_{j}) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \right) \ \psi(t) \rangle \right. \\ \left. + \left\langle \psi(t), \ \left(\nabla_{q_{j}} V(\hat{q}) + \hat{a}_{\hbar}(\tilde{g}_{j}) + \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \right) \cdot \hat{p}_{j} \ \psi(t) \rangle \right].$$

We estimate now each part. Indeed, we have

$$\langle \psi(t), \ \hat{p}_{j} \cdot \nabla_{q_{j}} V(\hat{q}) \ \psi(t) \rangle = \langle \hat{p}_{j} \ \psi(t), \ \nabla_{q_{j}} V(\hat{q}) \ \psi(t) \rangle$$

$$\leq \| \hat{p}_{j} \ \psi(t) \| \ \| \nabla_{q_{j}} V(\hat{q}) \ \psi(t) \|$$

$$\leq \| \hat{p}_{j} \ \psi(t) \| \ \| \nabla_{q_{j}} V \|_{L^{\infty}} \ \| \psi \|$$

$$\leq \frac{1}{2} \ \| \nabla_{q_{j}} V \|_{L^{\infty}} \Big[\| \hat{p}_{j} \ \psi(t) \|^{2} + \| \psi \|^{2} \Big].$$

Similarly, we have

$$\langle \psi(t), \ \nabla_{q_j} V(\hat{q}) \cdot \hat{p}_j \ \psi(t) \rangle \leq \frac{1}{2} \ \| \nabla_{q_j} V \|_{L^{\infty}} \Big[\| \hat{p}_j \ \psi(t) \|^2 + \| \psi \|^2 \Big].$$

We have also using Lemmas 3.1 and 3.3

$$\begin{aligned} \langle \psi(t), \ \hat{a}_{\hbar}(\tilde{g}_{j}) \cdot \hat{p}_{j} \ \psi(t) \rangle &= \langle \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \ \psi(t), \ \hat{p}_{j} \ \psi(t) \rangle \\ &\leq \| \hat{p}_{j} \ \psi(t) \| \ \| \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \ \psi(t) \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H}_{02} + 1)^{1/2} \ \psi(t) \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H}_{0} + 1)^{1/2} \ \psi(t) \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H} + a)^{1/2} \ \psi(t) \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H} + a)^{1/2} \ \psi \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H}_{0} + 1)^{1/2} \ \psi \| \\ &\lesssim \left[\| \chi \|_{L^{2}} + \| \sqrt{\omega} \ \chi \|_{L^{2}} \right] \ \| \hat{p}_{j} \ \psi(t) \| \ \| (\hat{H}_{0} + 1)^{1/2} \ \psi \|^{2} \right]. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \langle \psi(t), \ \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \cdot \hat{p}_{j} \ \psi(t) \rangle &\lesssim \|\chi\|_{L^{2}} \left[\|\hat{p}_{j} \ \psi(t)\|^{2} + \|(\hat{H}_{0}+1)^{1/2} \ \psi\|^{2} \right], \\ \langle \psi(t), \ \hat{p}_{j} \cdot \hat{a}_{\hbar}(\tilde{g}_{j}) \ \psi(t) \rangle &\lesssim \|\chi\|_{L^{2}} \left[\|\hat{p}_{j} \ \psi(t)\|^{2} + \|(\hat{H}_{0}+1)^{1/2} \ \psi\|^{2} \right], \\ \langle \psi(t), \ \hat{p}_{j} \cdot \hat{a}_{\hbar}^{*}(\tilde{g}_{j}) \ \psi(t) \rangle &\lesssim \left[\|\chi\|_{L^{2}} + \|\sqrt{\omega} \ \chi\|_{L^{2}} \right] \left[\|\hat{p}_{j} \ \psi(t)\|^{2} + \|(\hat{H}_{0}+1)^{1/2} \ \psi\|^{2} \right]. \end{aligned}$$

We get at the end that there exists some $C \in \mathbf{R}^*_+$ depending on the quantities $\|\chi\|_{L^2}$, $\|\sqrt{\omega} \chi\|_{L^2}$ and $\|\nabla_{q_j}V\|_{L^{\infty}}$ such that

$$\dot{\Theta}_2(t) \le C \Big[\langle \psi, (\hat{H}_0 + 1) \psi \rangle + \Theta_2(t) \Big].$$

We find then

$$\Theta_2(t) \le \Theta_2(0) + C \langle \psi, (\hat{H}_0 + 1) \psi \rangle t + \int_0^t \Theta_2(s) ds.$$

This implies using Gronwall's Lemma that there exists $C_1, C_2 > 0$ depend on the quantities $\|\chi\|_{L^2}, \|\sqrt{\omega} \chi\|_{L^2}$ and $\|\nabla_{q_j}V\|_{L^{\infty}}$ such that

$$\Theta_2(t) \le C_1 \langle \psi, (\hat{H}_0 + \hat{p}^2 + 1) \psi \rangle e^{C_2|t|}$$

And thus the result follows.

4.1.2. Field operator's estimates. Below, we give some estimates for the field operator $d\Gamma(\omega^{2\sigma})$. Let Γ_{fin} be a dense subspace in the Fock space. Let $\psi \in D(\hat{H}_0)$ and define

$$\Theta_3(t) := \langle \psi(t), \ d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})} \ \psi(t) \rangle, \qquad \psi(t) := e^{-i\frac{t}{\hbar}\hat{H}} \ \psi.$$

Note that $d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})}$ is a bounded and positive approximation of $d\Gamma(\omega^{2\sigma})$ that strongly converges monotonically to it. The quantity $\Theta_3(t)$ is well-defined for each $t \in \mathbf{R}$ and $\delta > 0$. In addition, the map $t \to \Theta_3(t)$ is differentiable with

$$\dot{\Theta}_3(t) = \frac{i}{\hbar} \langle \psi(t), [\hat{H}, d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})}] \psi(t) \rangle.$$

Lemma 4.3. Assume (1.6) and $\omega^{\sigma-\frac{1}{2}}\chi \in L^2(\mathbf{R}^d, dk)$. For $\sigma \in [1/2, 1]$, there exists C > 0 such that for all $\delta > 0$, for all $\hbar \in (0, 1)$ and for all ϕ , $\psi \in \mathcal{C}_0^{\infty}(\mathbf{R}^{dn}) \otimes \Gamma_{fin}$:

$$\left|\frac{i}{\hbar}\langle\phi, \ [\hat{H}, d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}] \ \psi\rangle\right| \le C \ \|\omega^{\sigma-\frac{1}{2}}\chi\|_{L^{2}}\Big[\|\phi\| \ \|d\Gamma(\omega^{2\sigma})^{\frac{1}{2}}\psi\| + \|\psi\| \ \|d\Gamma(\omega^{2\sigma})^{\frac{1}{2}}\phi\|\Big].$$

Proof. Let us deal first with the term

$$\begin{split} \left[d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{H} \right] &= \left[d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{H}_1 \right] \\ &= \left[d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{a}_{\hbar}(G) + \hat{a}_{\hbar}^*(G) \right] \\ &= d\Gamma(\omega^{2\sigma}) \ \left[e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{a}_{\hbar}(G) \right] + \left[d\Gamma(\omega^{2\sigma}), \hat{a}_{\hbar}(G) \right] \ e^{-\delta d\Gamma(\omega^{2\sigma})} \\ &+ d\Gamma(\omega^{2\sigma}) \ \left[e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{a}_{\hbar}^*(G) \right] + \left[d\Gamma(\omega^{2\sigma}), \hat{a}_{\hbar}^*(G) \right] \ e^{-\delta d\Gamma(\omega^{2\sigma})}. \end{split}$$

We also have

- (i) $[d\Gamma(\omega^{2\sigma}), \hat{a}_{\hbar}(G)] = -\hbar \hat{a}_{\hbar}(\omega^{2\sigma}G),$
- (ii) $[d\Gamma(\omega^{2\sigma}), \hat{a}^*_{\hbar}(G)] = \hbar \ \hat{a}^*_{\hbar}(\omega^{2\sigma}G),$

(iii)
$$[e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{a}_{\hbar}(G)] = e^{-\delta d\Gamma(\omega^{2\sigma})} \hat{a}_{\hbar}(\beta G), \quad \beta = 1 - e^{-\delta \hbar \omega^{2\sigma}},$$

(iv)
$$[e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{a}_{\hbar}^{*}(G)] = \hat{a}_{\hbar}^{*}(-\beta \ G) \ e^{-\delta d\Gamma(\omega^{2\sigma})}, \quad \beta = 1 - e^{-\delta \ \hbar \ \omega^{2\sigma}}.$$

Using (i)-(ii)-(iii) and (iv), we get

$$\left[d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}, \hat{H}\right] = \hbar \Big(B_1 + B_2 + B_3\Big),$$

where we have introduced the three terms B_1 , B_2 and B_3 as follows

$$B_{1} := \left[\hat{a}_{\hbar}^{*}(\omega^{2\sigma}G) - \hat{a}_{\hbar}(\omega^{2\sigma}G)\right] e^{-\delta d\Gamma(\omega^{2\sigma})},$$

$$B_{2} := d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})} \hat{a}_{\hbar}\left(\frac{\beta G}{\hbar}\right),$$

$$B_{3} := d\Gamma(\omega^{2\sigma}) \hat{a}_{\hbar}^{*}\left(\frac{-\beta G}{\hbar}\right) e^{-\delta d\Gamma(\omega^{2\sigma})}.$$

We get then

$$\frac{i}{\hbar} \langle \phi, \left[\hat{H}, d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})} \right] \psi \rangle = -i \ \langle \phi, \ (B_1 + B_2 + B_3) \ \psi \rangle \\ = \underbrace{-i \langle \phi, \ B_1 \ \psi \rangle}_{(a)} + \underbrace{-i \langle \phi, \ B_2 \ \psi \rangle}_{(b)} + \underbrace{-i \langle \phi, \ B_3 \ \psi \rangle}_{(c)}.$$

For (b), we have

$$\begin{aligned} |\langle \phi, B_2 \psi \rangle| &= \left| \langle \phi, d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})} \hat{a}_{\hbar} \left(\frac{\beta G}{\hbar} \right) \psi \rangle \right| \\ &= \left| \langle \phi, \delta d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})} \hat{a}_{\hbar} \left(\frac{\beta G}{\delta \hbar} \right) \psi \rangle \right| \\ &\leq \|\phi\| \|\delta d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})} \hat{a}_{\hbar} \left(\frac{\beta G}{\delta \hbar} \right) \psi \| \\ &\leq \frac{1}{e} \|\phi\| \|\hat{a}_{\hbar} \left(\frac{\beta G}{\delta \hbar} \right) \psi\|, \end{aligned}$$

where in the last line we have used the fact that $\sup_{\delta>0} \|\delta d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})}\| \leq 1/e$. Remark also that we have with K_m and X_n as in (1.2)

$$= \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left| \int_{\mathbf{R}^{d}} \sum_{j=1}^{n} \omega^{\sigma-\frac{1}{2}}(k) \chi(k) \sqrt{\hbar(m+1)} \omega^{\sigma}(k) e^{2\pi i k \cdot \hat{q}_{j}} \psi^{m+1}(X_{n}, K_{m}, k) dk \right|^{2} dK_{m} dX_{m}$$

$$\leq n^{2} \|\omega^{\sigma-\frac{1}{2}} \chi\|_{2}^{2} \sum_{m\geq 0} \int_{\mathbf{R}^{dn}} \int_{\mathbf{R}^{dm}} \left[\int_{\mathbf{R}^{d}} \hbar(m+1) \omega^{2\sigma}(k) \left| \psi^{m+1}(X_{n}, K_{m}, k) \right|^{2} dk \right] dK_{m} dX_{n}$$

$$\leq n^{2} \|\omega^{\sigma-\frac{1}{2}} \chi\|_{2}^{2} \|d\Gamma(\omega^{2\sigma})^{1/2}\psi\|_{\mathcal{H}}^{2}.$$

Using the above estimates, we find that

$$|\langle \phi, B_2 \psi \rangle| \le \frac{n}{e} \|\omega^{\sigma - \frac{1}{2}} \chi\|_2 \|\phi\| \|d\Gamma(\omega^{2\sigma})^{1/2} \psi\|_{\mathcal{H}}.$$

For (c), remark first that we have

$$[d\Gamma(\omega^{2\sigma}), \ \hat{a}_{\hbar}^* \left(\frac{-\beta \ G}{\hbar}\right)] = \hbar \ \hat{a}_{\hbar}^* \left(\frac{-\omega^{2\sigma} \ \beta \ G}{\hbar}\right),$$

and

$$[d\Gamma(\omega^{2\sigma}), \ \hat{a}_{\hbar}^{*}\left(\frac{-\beta \ G}{\hbar}\right)] = d\Gamma(\omega^{2\sigma}) \ \hat{a}_{\hbar}^{*}\left(\frac{-\beta \ G}{\hbar}\right) - \hat{a}_{\hbar}^{*}\left(\frac{-\beta \ G}{\hbar}\right) \ d\Gamma(\omega^{2\sigma})$$

This implies that

$$d\Gamma(\omega^{2\sigma}) \ \hat{a}_{\hbar}^{*}\left(\frac{-\beta \ G}{\hbar}\right) = \hat{a}_{\hbar}^{*}\left(-\omega^{2\sigma} \ \beta \ G\right) + \hat{a}_{\hbar}^{*}\left(\frac{-\beta \ G}{\hbar}\right) \ d\Gamma(\omega^{2\sigma}).$$

Then, we have

$$\begin{aligned} |\langle \phi, B_3 | \psi \rangle| &\leq |\langle \phi, \hat{a}_{\hbar}^* \big(-\omega^{2\sigma} |\beta| G \big) |e^{-\delta d\Gamma(\omega^{2\sigma})} |\psi \rangle| + |\langle \phi, |\hat{a}_{\hbar}^* \big(\frac{-\beta|G|}{\hbar} \big) |d\Gamma(\omega^{2\sigma}) |e^{-\delta d\Gamma(\omega^{2\sigma})} |\psi \rangle| \\ &\lesssim ||\omega^{\sigma-\frac{1}{2}} |\chi||_2 ||\psi|| ||d\Gamma(\omega^{2\sigma})^{1/2} \phi||_{\mathcal{H}}, \end{aligned}$$

~ ~

where in the last line, we have used the same tricks as before as well as the fact that $|\beta| \leq 2$ and $e^{-\delta \ d\Gamma(\omega^{2\sigma})} \in \mathcal{L}(\mathcal{H})$ with $||e^{-\delta \ d\Gamma(\omega^{2\sigma})}|| \leq 1$. Similarly for (a), we can have by same techniques that

$$\begin{aligned} |\langle \phi, \hat{a}_{\hbar}^{*}(\omega^{2\sigma}G) \ e^{-\delta d\Gamma(\omega^{2\sigma})} \ \psi \rangle| &= |\langle \hat{a}_{\hbar}(\omega^{2\sigma}G) \ \phi, \ e^{-\delta d\Gamma(\omega^{2\sigma})} \ \psi \rangle| \\ &\lesssim \ \|\omega^{\sigma-\frac{1}{2}} \ \chi\|_{2} \ \|\psi\| \ \|d\Gamma(\omega^{2\sigma})^{1/2}\phi\|_{\mathcal{H}}. \end{aligned}$$

For the other term in (a), note that

$$[e^{-\delta d\Gamma(\omega^{2\sigma})}, \ \hat{a}_{\hbar}(\omega^{2\sigma}G)] = e^{-\delta d\Gamma(\omega^{2\sigma})} \ \hat{a}_{\hbar}(\beta \ \omega^{2\sigma}G).$$

This implies that

$$\hat{a}_{\hbar}(\omega^{2\sigma}G) \ e^{-\delta d\Gamma(\omega^{2\sigma})} = e^{-\delta d\Gamma(\omega^{2\sigma})} \ \hat{a}_{\hbar}((1-\beta) \ \omega^{2\sigma}G).$$

Then, using the above equality, we get

$$|\langle \phi, \hat{a}_{\hbar}(\omega^{2\sigma}G) \ e^{-\delta d\Gamma(\omega^{2\sigma})} \ \psi \rangle| \lesssim \|\omega^{\sigma-\frac{1}{2}} \ \chi\|_2 \|\phi\| \|d\Gamma(\omega^{2\sigma})^{1/2}\psi\|_{\mathcal{H}}.$$

We conclude that

$$|\langle \phi, B_1 \psi \rangle| \lesssim \|\omega^{\sigma - \frac{1}{2}} \chi\|_2 \left[\|\psi\| \| d\Gamma(\omega^{2\sigma})^{1/2} \phi\|_{\mathcal{H}} + \|\phi\| \| d\Gamma(\omega^{2\sigma})^{1/2} \psi\|_{\mathcal{H}} \right]$$

And thus, the final result follows.

Lemma 4.4. There exists C_1 , $C_2 \in \mathbf{R}^*_+$ such that

$$\Theta_3(t) \le C_1 \langle \psi, (d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})} + 1) \psi \rangle e^{C_2|t|}$$

Proof. Use the previous Lemma 4.3 with $\phi = \psi = \psi(t)$, we get that

$$\begin{split} \dot{\Theta}_{3}(t) &= \frac{\imath}{\hbar} \langle \psi(t), [\hat{H}, \ d\Gamma(\omega^{2\sigma}) \ e^{-\delta d\Gamma(\omega^{2\sigma})}] \ \psi(t) \rangle \\ &\lesssim \|\omega^{\sigma - \frac{1}{2}} \ \chi\|_{2} \left[\|\psi\| \ \|d\Gamma(\omega^{2\sigma})^{1/2} \psi(t)\|_{\mathcal{H}} \right] \\ &\lesssim \|\omega^{\sigma - \frac{1}{2}} \ \chi\|_{2} \left[\|\psi\| \ \|[d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})}]^{1/2} \psi(t)\|_{\mathcal{H}} \right] \\ &\lesssim \|\omega^{\sigma - \frac{1}{2}} \ \chi\|_{2} \left[\|\psi\|^{2} + \|[d\Gamma(\omega^{2\sigma}) e^{-\delta d\Gamma(\omega^{2\sigma})}]^{1/2} \psi(t)\|_{\mathcal{H}}^{2} \right] \\ &\lesssim c \langle \psi, \ \psi \rangle + c \ \Theta_{3}(t). \end{split}$$

And thus the result follows by applying the Gronwall's Lemma.

Lemma 4.5 (Field estimate). Assume that (1.6) and $\omega^{\sigma-\frac{1}{2}}\chi \in L^2(\mathbf{R}^d, dk)$. Then, there exists constants $C_1, C_2 > 0$ such that for all $\psi \in D(\hat{H}_0^{1/2}) \cap D(d\Gamma(\omega^{2\sigma})^{1/2})$, all $t \in \mathbf{R}$ and all $\hbar \in (0, 1)$:

$$\langle e^{-i\frac{t}{\hbar}\hat{H}}\psi, \ d\Gamma(\omega^{2\sigma}) \ e^{-i\frac{t}{\hbar}\hat{H}}\psi\rangle \leq C_1\langle\psi, \ (d\Gamma(\omega^{2\sigma})+1) \ \psi\rangle \ e^{C_2|t|}.$$
 (4.6)

Proof. It is a consequence of the previous Lemma 4.4. Indeed, the approximation map $e^{-\delta d\Gamma(\omega^{2\sigma})} d\Gamma(\omega^{2\sigma})$ converges strongly to $d\Gamma(\omega^{2\sigma})$. This leads to

$$\begin{split} \| \left(d\Gamma(\omega^{2\sigma}) \right)^{1/2} \psi(t) \|^2 &= \lim_{\delta \to 0} \| \left(e^{-\delta d\Gamma(\omega^{2\sigma})} \ d\Gamma(\omega^{2\sigma}) \right)^{1/2} \psi(t) \|^2 \\ &\lesssim \lim_{\delta \to 0} C_1 \ \| (e^{-\delta d\Gamma(\omega^{2\sigma})} \ d\Gamma(\omega^{2\sigma}) + 1)^{1/2} \ \psi \|^2 \ e^{C_2|t|} \\ &= C_1 \ \| (d\Gamma(\omega^{2\sigma}) + 1)^{1/2} \ \psi \|^2 \ e^{C_2|t|}. \end{split}$$

And thus, we achieve the desired result.

4.1.3. Propagation of estimates uniformly for all times. As a consequence of the previous estimates, the uniform bound on the initial states $(\varrho_{\hbar})_{\hbar \in (0,1)}$ propagates in time.

Lemma 4.6 (Propagation of the assumptions (1.8) and (1.9) in time). Assume (1.6) and $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). Then, the family of states $(\varrho_h(t))_{\hbar \in (0,1)}$ and $(\tilde{\varrho}_h(t))_{\hbar \in (0,1)}$ satisfy the same assumptions (1.8) and (1.9) uniformly for any $t \in \mathbf{R}$ in arbitrary compact interval.

Proof. Before we begin the proof, remark that by spectral decomposition, we have

$$\varrho_{\hbar} = \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ |e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m)\rangle \langle e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m)|,$$

where $\lambda_{\hbar}(m)$ are the eigenvalues and $\psi_{\hbar}(m)$ are their related eigenfunctions. Let J be a compact interval. Then for all $t \in J$:

• We have with some $c \in \mathbf{R}^*_+$ the following uniform estimate

$$\operatorname{Tr}\left[\tilde{\varrho}_{\hbar}(t) \ \hat{p}^{2}\right] = \operatorname{Tr}\left[e^{i\frac{t}{\hbar}\hat{H}_{02}} \ \varrho_{\hbar}(t) \ e^{-i\frac{t}{\hbar}\hat{H}_{02}} \ \hat{p}^{2}\right]$$

$$= \operatorname{Tr}\left[\varrho_{\hbar}(t) \ \hat{p}^{2}\right]$$

$$= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \|\hat{p} \ e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m)\|^{2}$$

$$= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m), \ \hat{p}^{2} \ e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m)\rangle$$

$$\leq C_{1} \ \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle \psi_{\hbar}(m), \ (\hat{H}_{0} + \hat{p}^{2} + 1) \ \psi_{\hbar}(m)\rangle \ e^{C_{2}|t|} \leq c,$$

where we have used Lemma 4.2 as well as assumptions (1.8) and (1.9).

• We have with some $c' \in \mathbf{R}^*_+$ the following estimate

$$\begin{aligned} \operatorname{Tr} \left[\tilde{\varrho}_{\hbar}(t) \ \hat{q}^{2} \right] &= \operatorname{Tr} \left[e^{i \frac{t}{\hbar} \hat{H}_{02}} \ \varrho_{\hbar}(t) \ e^{-i \frac{t}{\hbar} \hat{H}_{02}} \ \hat{q}^{2} \right] \\ &= \operatorname{Tr} \left[\varrho_{\hbar}(t) \ \hat{q}^{2} \right] \\ &= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \| \hat{q} \ e^{-i \frac{t}{\hbar} \hat{H}} \psi_{\hbar}(m) \|^{2} \\ &= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle e^{-i \frac{t}{\hbar} \hat{H}} \psi_{\hbar}(m), \ \hat{q}^{2} \ e^{-i \frac{t}{\hbar} \hat{H}} \psi_{\hbar}(m) \rangle \\ &\leq C_{1} \ \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle \psi_{\hbar}(m), \ (\hat{H}_{0} + \hat{q}^{2} + 1) \ \psi_{\hbar}(m) \rangle \ e^{C_{2}|t|} \leq c', \end{aligned}$$

where we have used Lemma 4.1 as well as assumptions (1.8) and (1.9). • We have for some c'' the following uniform estimate

$$\operatorname{Tr}\left[\tilde{\varrho}_{\hbar}(t) \ d\Gamma(\omega^{2\sigma})\right] = \operatorname{Tr}\left[e^{i\frac{t}{\hbar}\hat{H}_{02}} \ \varrho_{\hbar}(t) \ e^{-i\frac{t}{\hbar}\hat{H}_{02}} \ d\Gamma(\omega^{2\sigma})\right]$$
$$= \operatorname{Tr}\left[\varrho_{\hbar}(t) \ d\Gamma(\omega^{2\sigma})\right]$$
$$= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \|d\Gamma(\omega^{2\sigma})^{1/2} \ e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m)\|^{2}$$
$$= \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle e^{-i\frac{t}{\hbar}\hat{H}}\psi_{\hbar}(m), \ d\Gamma(\omega^{2\sigma}) \ e^{-i\frac{t}{\hbar}H}\psi_{\hbar}(m)\rangle$$
$$\leq C_{1} \ \sum_{m \in \mathbf{N}} \lambda_{\hbar}(m) \ \langle \psi_{\hbar}(m), \ (d\Gamma(\omega^{2\sigma}) + 1) \ \psi_{\hbar}(m)\rangle \ e^{C_{2}|t|} \leq c''.$$

where we have used Lemma 4.5 as well as assumptions (1.8).

4.2. Existence of unique Wigner measure. In this section, we prove that for any family of states $(\varrho_h)_{h \in (0,1)}$ which satisfies (1.8) and (1.9) and for any sequence $\hbar_n \to 0$, we can extract a subsequence $\hbar_{n_\ell} \to 0$ such that the set of Wigner measure

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N})$$

is singleton. The main results are stated below:

Proposition 4.7 (Existence of unique Wigner measure $\tilde{\mu}_t$ for all times). Assume that (1.6) and (1.7) hold true. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). For any sequence $(\hbar_n)_{n \in \mathbb{N}}$ in (0,1) such that $\hbar_n \to 0$, there exists a subsequence $(\hbar_{n_\ell})_{\ell \in \mathbb{N}}$ and a family of probability measures $(\tilde{\mu}_t)_{t \in \mathbb{R}}$ such that for all $t \in \mathbb{R}$,

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N}) = \{\tilde{\mu}_t\}.$$

Moreover, for every compact time interval J there exists a constant C > 0 such that for all times $t \in J$,

$$\int_{X^0} \|u\|_{X^{\sigma}}^2 d\tilde{\mu}_t(u) < C.$$
(4.7)

Proof. We prove the above proposition in two steps. Step 1 is dedicated to the extraction of a unique Wigner measure at fixed times. Step 2 generalizes for all times. To establish Step 1, it is necessary to recall the following result from [10, Theorem 6.2].

Proposition 4.8 (The set of Wigner measure is not empty). Let $(\varrho_h)_{h\in(0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). Then for all sequences $(\hbar_n)_{n\in\mathbb{N}}$ with $\lim_{n\to\infty} h_n = 0$, there exists a subsequence $(\hbar_{n_\ell})_{\ell\in\mathbb{N}}$ with $\lim_{\ell\to\infty} h_{n_\ell} = 0$ such that

$$\mathcal{M}(\varrho_{\hbar_{n_{\ell}}}, \ell \in \mathbf{N}) = \{\mu\}.$$

Moreover, we have

$$\int_{X^0} \|u\|_{X^0}^2 d\mu(u) < +\infty, \qquad \int_{X^0} \|u\|_{X^\sigma}^2 d\mu(u) < +\infty.$$
(4.8)

Step 1: Extraction of a unique Wigner measure at fixed times.

Let $s \in \mathbf{R}$ be a fixed time. Let $\hbar_n \to 0$. Then, by Proposition 4.8, there exists a subsequence $n \to \infty$ $(\hbar_{\ell})_{\ell \in \mathbf{N}} \equiv (\hbar_{n_{\ell}})_{\ell \in \mathbf{N}}$ such that $\hbar_{\ell} \to 0$ and a probability measure $\tilde{\mu}_s \in \mathcal{P}(X^0)$ such that $\ell \to \infty$

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{n_{\ell}}}(s), \ell \in \mathbf{N}) = \{\tilde{\mu}_s\}.$$

Moreover, we have the following integrability formula

$$\int_{X^0} \|u\|_{X^{\sigma}}^2 d\tilde{\mu}_s(u) = \int_{X^0} (p^2 + q^2 + \|\alpha\|_{\mathcal{G}^{\sigma}}^2) d\tilde{\mu}_s(p, q, \alpha) < +\infty.$$
(4.9)

The above integrability formula is a consequence of the following implications proved in [11, Lemma 3.12] for some C > 0:

- (i) If $\operatorname{Tr}[\varrho_{\hbar} \ \hat{N}_{\hbar}] \leq C \Longrightarrow \forall \mu \in \mathcal{M}(\varrho_{\hbar}; \hbar \in (0, 1)), \ \int_{X^{0}} \|\alpha\|_{\mathcal{G}^{0}}^{2} \ d\mu \leq C;$ (ii) If $\operatorname{Tr}[\varrho_{\hbar} \ d\Gamma(\omega^{2\sigma})] \leq C \Longrightarrow \forall \mu \in \mathcal{M}(\varrho_{\hbar}; \hbar \in (0, 1)), \ \int_{X^{0}} \|\alpha\|_{\mathcal{G}^{\sigma}}^{2} \ d\mu \leq C;$ (iii) If $\operatorname{Tr}[\varrho_{\hbar} \ (\hat{q}^{2} + \hat{p}^{2})] \leq C \Longrightarrow \forall \mu \in \mathcal{M}(\varrho_{\hbar}; \hbar \in (0, 1)), \ \int_{X^{0}} (q^{2} + p^{2}) \ d\mu \leq C.$

Now, by the help of uniform estimate in Lemma 4.6, we have the two family of states $(\varrho_h(t))_{h \in (0,1)}$ and $(\tilde{\varrho}_h(t))_{h\in(0,1)}$ satisfy uniformly the bounds of (i)-(ii)-(iii), one obtains then that (4.9) holds true as a consequence of (4.7) in Proposition 4.8.

Step 2: Generalization for all times.

Claim first that we have for all times $t \in \mathbf{R}$

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\ell}}(t), \ell \in \mathbf{N}) = \{\tilde{\mu}_t\}.$$
(4.10)

Let us now prove the integrability formula in Proposition 4.7. Recall that our density metrices $(\varrho_{\hbar})_{\hbar \in (0,1)}$ satisfies the assumptions (1.8) and (1.9). And, by Lemma 4.6, the family of states $(\tilde{\varrho}_{\hbar}(t))_{\hbar \in (0,1)}$ satisfies the same assumptions uniformly in any compact time interval J. Then, using (4.8), for all $t \in J$, we have (4.7). we come back now to prove the claim (4.10). Let $(t_i)_{i \in \mathbb{N}}$ be a countable dense set in \mathbf{R} . We have by Step 1 that

• for t_1 , for $\hbar_n \xrightarrow[n \to \infty]{} 0$, there exists a subsequence $\hbar_{\ell} \xrightarrow[\ell \to \infty]{} 0$ such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\ell}}(t_1), \ell \in \mathbf{N}) = \{\tilde{\mu}_{t_1}\}$$

• for t_2 , for $(\hbar_\ell)_{\ell \in \mathbf{N}}$, there exists a subsequence $(\hbar_{\phi_2(\ell)})_{\ell \in \mathbf{N}} \subset (\hbar_\ell)_{\ell \in \mathbf{N}}$ such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\phi_2(\ell)}}(t_2), \ell \in \mathbf{N}) = \{\tilde{\mu}_{t_2}\}.$$

• for t_3 , for $(\hbar_{\phi_2(\ell)})_{\ell \in \mathbf{N}}$, there exists a subsequence $(\hbar_{\phi_3(\ell)})_{\ell \in \mathbf{N}} \subset (\hbar_{\phi_2(\ell)})_{\ell \in \mathbf{N}}$ such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\phi_3(\ell)}}(t_3), \ell \in \mathbf{N}) = \{\tilde{\mu}_{t_3}\}.$$

• And so on, for t_j , for $(\hbar_{\phi_{i-1}(\ell)})_{\ell \in \mathbf{N}}$, there exists a subsequence $(\hbar_{\phi_i(\ell)})_{\ell \in \mathbf{N}} \subset (\hbar_{\phi_{i-1}(\ell)})_{\ell \in \mathbf{N}}$ such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\phi_j}(\ell)}(t_j), \ell \in \mathbf{N}) = \{\tilde{\mu}_{t_j}\}.$$

By diagonal arguments, we extract the subsequence $(\hbar_{\phi_{\ell}(\ell)})_{\ell}$ denoted by $(\hbar_{\ell})_{\ell}$ for simplicity such that for all $j \in \mathbf{N}$, we have

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{\ell}}(t_j), \ell \in \mathbf{N}) = \{\tilde{\mu}_{t_j}\}.$$

The above formula implies that for all $\xi = (p_0, q_0, \alpha_0) \in X^0$ and $\tilde{\xi} = (-2\pi q_0, 2\pi p_0, \sqrt{2\pi\alpha_0})$

$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \; \tilde{\varrho}_{\hbar_{\ell}}(t_j) \right] = \int_{X^0} e^{2\pi i \Re e \langle \xi, u \rangle_{X^0}} \; d\tilde{\mu}_{t_j}(u) \tag{4.11}$$

We have

$$\int_{X^0} \|u\|_{X^{\sigma}}^2 d\tilde{\mu}_{t_j}(u) < +\infty.$$

The above formula implies that the set of Wigner measure $\{\tilde{\mu}_{t_j}\}_{j\in\mathbb{N}}$ is tight in $\mathcal{P}(X^0)$. This implies that according to Prokhorov's theorem in Lemma A.1, for all $\xi \in X^0$, there exists a subsequence still denoted by t_j and a probability measure $\tilde{\mu}_t \in \mathcal{P}(X^0)$ such that $\tilde{\mu}_{t_j}$ converges weakly narrowly to $\tilde{\mu}_t$. This gives since the function $e^{2\pi i \Re e \langle \xi, u \rangle_{X^0}}$ is bounded that

$$\int_{X^0} e^{2\pi i \Re e \langle \xi, u \rangle_{X^0}} d\tilde{\mu}_{t_j}(u) \xrightarrow[t_j \to t]{} \int_{X^0} e^{2\pi i \Re e \langle \xi, u \rangle_{X^0}} d\tilde{\mu}_t(u)$$

Now, we need to prove that

$$\int_{X^0} e^{2\pi i \Re e\langle \xi, u \rangle_{X^0}} d\tilde{\mu}_{t_j}(u) \underset{t_j \to t}{\longrightarrow} \lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \ \tilde{\varrho}_{\hbar_\ell}(t) \right]$$
(4.12)

We start by

$$\left| \int_{X^{0}} e^{2\pi i \Re e \langle \xi, u \rangle_{X^{0}}} d\tilde{\mu}_{t_{j}}(u) - \lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \; \tilde{\varrho}_{\hbar_{\ell}}(t) \right] \right|$$

$$\leq \left| \int_{X^{0}} e^{2\pi i \Re e \langle \xi, u \rangle_{X^{0}}} d\tilde{\mu}_{t_{j}}(u) - \lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \; \tilde{\varrho}_{\hbar_{\ell}}(t_{j}) \right] \right| \qquad (1)$$

$$+ \left| \lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \; \tilde{\varrho}_{\hbar_{\ell}}(t_{j}) \right] - \lim_{\ell \to \infty} \operatorname{Tr} \left[\mathcal{W}(\tilde{\xi}) \; \tilde{\varrho}_{\hbar_{\ell}}(t) \right] \right| \qquad (2)$$

The quantity (1) is zero by (4.11). The quantity (2) is zero by using the following estimates:

(i) For $\xi \in X^0$, for all $t, t_0 \in J$ where J is compact interval, we have

$$\left| \operatorname{Tr} \Big[\mathcal{W}(\xi) \left(\tilde{\varrho}_{\hbar}(t) - \tilde{\varrho}_{\hbar}(t_0) \right) \Big] \right| \lesssim |t - t_0| \|\xi\|_{X^0} \Big[\|\xi\|_{X^0} + \|\chi\|_{L^2} + \|\chi\|_{\mathcal{G}^0} \Big];$$

(ii) For all $\xi_1, \xi_2 \in X^0$, for all $t \in J$ where J is compact interval, we have

$$\left| \operatorname{Tr} \left[\left(\mathcal{W}(\xi_1) - \mathcal{W}(\xi_2) \right) \, \tilde{\varrho}_{\hbar}(t) \right] \right| \lesssim \|\xi_1 - \xi_2\|_{X^0} \Big[\|\xi_1\|_{X^0} + \|\xi_2\|_{X^0} + 1 \Big]$$

For (i), we exploit (3.13), we have with $S = (\hat{H}_0 + 1)^{1/2}$

$$\left| \operatorname{Tr} \left[\mathcal{W}(\xi) \left(\tilde{\varrho}_{\hbar}(t) - \tilde{\varrho}_{\hbar}(t_0) \right) \right] \right| \leq |t - t_0| \left\| (\mathbf{B}_0 + \hbar \mathbf{B}_1) S^{-1} \right\|_{\mathcal{L}(\mathcal{H})} \left\| S \mathcal{W}(\xi) S^{-1} \right\|_{\mathcal{L}(\mathcal{H})} \left\| S \tilde{\varrho}_{\hbar}(t) \right\|_{\mathcal{L}^1(\mathcal{H})}.$$

Now using Lemma 3.5, the two estimates (3.28) and (3.29) and the two assumptions (1.8) and (1.9), we get the desired result.

For (ii), we have

$$\left| \operatorname{Tr} \left[\left(\mathcal{W}(\xi_1) - \mathcal{W}(\xi_2) \right) \, \tilde{\varrho}_{\hbar}(t) \right] \right| \leq \left\| \left(\mathcal{W}(\xi_1) - \mathcal{W}(\xi_2) \right) (\hat{N}_{\hbar} + 1)^{-1/2} \right\|_{\mathcal{L}(\mathcal{H})} \underbrace{ \left\| \left(\hat{N}_{\hbar} + 1 \right)^{1/2} \tilde{\varrho}_{\hbar}(t) \right\|_{\mathcal{L}^1(\mathcal{H})}}_{<\infty \text{ by (1.8)}} \right|_{\mathcal{L}^1(\mathcal{H})}$$

And thus, using following the same computations as in [10, Lemma 3.1], the result follows. \Box

5. DERIVATION OF THE CHARACTERISTIC EQUATIONS

Subsection 5.1 focuses on investigating the convergence of the quantum dynamics towards the evolution of the particle-field equation. In Subsection 5.2, we derive the characteristic equation that the Wigner measure satisfies. Finally, in Subsection 5.3, we demonstrate that this characteristic equation is equivalent to a Liouville equation.

5.1. Convergence. In this section, we take the classical limit $\hbar_{n_{\ell}} \to 0$ as $\ell \to \infty$ in the Duhamel formula (3.13) to derive the characteristics equation satisfied by the Wigner measure $(\tilde{\mu}_t)_{t \in \mathbf{R}}$.

Lemma 5.1 (Convergence). Assume (1.6) and $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). Then for all $\xi = (z_0, \alpha_0) \in X^0$ and all $t, t_0 \in \mathbf{R}$, the Duhamel formula (3.13) converges to the following characteristics equation

$$\int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_t(u) = \int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_{t_0}(u) - i \int_{t_0}^t \int_{X^0} b(s,\xi) e^{Q(\xi,u)} d\tilde{\mu}_s(u) ds$$
(5.1)

with

 $Q(\xi, u) = i\Im m \langle z, z_0 \rangle + \sqrt{2}i\Re e \langle \alpha_0, \alpha \rangle, \qquad \xi = (z_0, \alpha_0), \quad u = (z, \alpha), \tag{5.2}$

and where we have introduced

$$b(s,\xi) := -\sum_{j=1}^{n} \nabla f_j(p_j) \cdot p_{0j} - \sum_{j=1}^{n} \nabla q_j V(q) \cdot q_{0j} + \sum_{j=1}^{n} \left(\langle \alpha, b_j^0(s) \rangle_{L^2} + \langle b_j^0(s), \alpha \rangle_{L^2} \right) + \frac{i}{\sqrt{2}} \left(\langle \alpha_0, g_j(s) \rangle_{L^2} - \langle g_j(s), \alpha_0 \rangle_{L^2} \right).$$

The function $b_j^0(s)$ is such that $b_j^0(s) \equiv b_j^0(s)(p_j,q_j)$ is defined as follows

$$b_j^0(s) := 2\pi i k \cdot q_{0j} \ \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{-2\pi i k \cdot q_j + is\omega(k)}$$
(5.3)

Proof. From the Definition 1.2 of Wigner measure, we have

$$\lim_{\ell \to +\infty} \operatorname{Tr}[\mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar n_{\ell}}(s)] = \int_{X^0} e^{Q(\xi, u)} \, d\tilde{\mu}_s(u)$$

We plug (3.25) in the Duhamel's formula (3.13), we get

$$\int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_t(u) = \int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_{t_0}(u) - i \int_{t_0}^t \lim_{\ell \to \infty} \operatorname{Tr}[B_0(s,\hbar_{n_\ell},\xi) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] \, ds.$$

We have to prove then

$$\lim_{\ell \to +\infty} \operatorname{Tr}[\mathcal{B}_0(s, \hbar_{n_\ell}, \xi) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] = \int_{X^0} b(s, \xi) \ e^{Q(\xi, u)} \ d\tilde{\mu}_s(u).$$

We start with

$$\begin{aligned} \operatorname{Tr}[\operatorname{B}_{0}(s,\hbar_{n_{\ell}},\xi) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \\ &= -\sum_{j=1}^{n} \operatorname{Tr}[\nabla f_{j}(\hat{p}_{j}) \cdot p_{0j} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] - \sum_{j=1}^{n} \operatorname{Tr}[\nabla_{q_{j}} V(\hat{q}) \cdot q_{0j} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \\ &+ \sum_{j=1}^{n} \operatorname{Tr}[\hat{a}_{\hbar_{n_{\ell}}}\left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}}\right) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] + \operatorname{Tr}[\hat{a}^{*}_{\hbar_{n_{\ell}}}\left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}}\right) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \\ &+ \sum_{j=1}^{n} \frac{i}{\sqrt{2}} \Big(\operatorname{Tr}[\langle \alpha_{0}, \tilde{g}_{j}(s) \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] - \operatorname{Tr}[\langle \tilde{g}_{j}(s), \alpha_{0} \rangle_{L^{2}(\mathbf{R}^{d}, \mathbf{C})} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \Big). \end{aligned}$$

Let us start with the first two terms. We have

$$\lim_{\ell \to \infty} \operatorname{Tr}[\nabla f_j(\hat{p}_j) \cdot p_{0j} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] = \int_{X^0} e^{Q(\xi, u)} \ \nabla f_j(p_j) \cdot p_{0j} \ d\tilde{\mu}_s(u), \tag{5.4}$$

$$\lim_{\ell \to \infty} \operatorname{Tr}[\nabla_{q_j} V(\hat{q}) \cdot q_{0j} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] = \int_{X^0} e^{Q(\xi, u)} \ \nabla_{q_j} V(q) \cdot q_{0j} \ d\tilde{\mu}_s(u), \tag{5.5}$$

where we have used in the above two lines the convergent results in [6, Lemma B.1] since $\langle p_j \rangle^{-1} \nabla f_j(p_j) \cdot p_{0j} \in L^{\infty}$ and $\langle q \rangle^{-1} \nabla q_j V(q) \cdot q_{0j} \in L^{\infty}$. Let us deal now with the second line. The goal is to prove the following limit:

$$\lim_{\ell} \operatorname{Tr} \left[\hat{a}_{\hbar_{n_{\ell}}}^{*} \left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}} \right) \mathcal{W}(\xi) \; \tilde{\varrho}_{\hbar_{n_{\ell}}}(s) \right] = \int_{X^{0}} e^{Q(\xi, u)} \langle \alpha, b_{j}^{0}(s) \rangle_{L^{2}} d\tilde{\mu}_{s}(u).$$

We start then with

$$\begin{aligned} \left| \operatorname{Tr} \left[\hat{a}_{\hbar_{n_{\ell}}}^{*} \left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}} \right) \mathcal{W}(\xi) \; \tilde{\varrho}_{\hbar_{n_{\ell}}}(s) \right] - \int_{X^{0}} e^{Q(\xi, u)} \langle \alpha, b_{j}^{0}(s) \rangle_{L^{2}} d\tilde{\mu}_{s}(u) \right| \\ \leq \underbrace{\left| \operatorname{Tr} \left[\hat{a}_{\hbar_{n_{\ell}}}^{*} \left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}} - \tilde{b}_{j}^{0}(s) \right) \mathcal{W}(\xi) \; \tilde{\varrho}_{\hbar_{n_{\ell}}}(s) \right] \right|_{(1)} \\ + \underbrace{\left| \operatorname{Tr} \left[\hat{a}_{\hbar_{n_{\ell}}}^{*} \left(\tilde{b}_{j}^{0}(s) \right) \mathcal{W}(\xi) \; \tilde{\varrho}_{\hbar_{n_{\ell}}}(s) \right] - \int_{X^{0}} e^{Q(\xi, u)} \langle \alpha, b_{j}^{0}(s) \rangle_{L^{2}} d\tilde{\mu}_{s}(u) \right|, \end{aligned}$$

where

$$\tilde{b}_j^0(s) := 2\pi i k \cdot q_{0j} \ \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{-2\pi i k \cdot \hat{q}_j + is\omega(k)} \ .$$
(5.6)

For (1), let $S = (\hat{H}_0 + 1)^{1/2}$, we have

$$\begin{aligned} \left| \operatorname{Tr} \left[\hat{a}_{\hbar_{n_{\ell}}}^{*} \left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}} - \tilde{b}_{j}^{0}(s) \right) \mathcal{W}(\xi) \; \tilde{\varrho}_{\hbar_{n_{\ell}}}(s) \right] \right| \\ &\leq \underbrace{\left\| \frac{1}{\sqrt{\omega}} \left[\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}} - \tilde{b}_{j}^{0}(s) \right] \right\|_{\mathcal{L}(L^{2}, L^{2} \otimes L^{2})}}_{\stackrel{\longrightarrow}{\ell \to \infty}} \| S \; \mathcal{W}(\xi) \; S^{-1} \|_{\mathcal{L}(\mathcal{H})} \; \left\| S \; \tilde{\varrho}_{\hbar}(s) \right\|_{\mathcal{L}^{1}(\mathcal{H})}. \end{aligned}$$

The above convergence follows from dominated convergence theorem and our assumptions.

For (2), according to the expression (5.6), we have

$$\tilde{b}_j^0(s) = e^{-2\pi i k \cdot \hat{q}_j} \varphi_j(k),$$

for some $\varphi_j \in L^2(\mathbf{R}^d)$. Hence, applying Lemma C.2 in the appendix, we conclude that (2) converges to zero as $\ell \to \infty$. Similar discussions lead to

$$\lim_{\ell \to \infty} \operatorname{Tr}\left[\hat{a}_{\hbar_{n_{\ell}}}\left(\frac{\tilde{g}_{j}(s) - g_{j}(s)}{\hbar_{n_{\ell}}}\right) \,\mathcal{W}(\xi) \,\,\tilde{\varrho}_{\hbar_{n_{\ell}}}(s)\right] = \int_{X^{0}} e^{Q(\xi, u)} \,\,\langle b_{j}^{0}(s), \alpha \rangle_{L^{2}} \,d\tilde{\mu}_{s}(u)$$

We deal now with the last line

$$\begin{aligned} \operatorname{Tr}[\langle \alpha_{0}, \tilde{g}_{j}(s) \rangle_{L^{2}} \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{k}}}(s)] \\ &= \int_{\mathbf{R}^{d}} \overline{\alpha_{0}(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{is\omega(k)} e^{2\pi i k \cdot q_{0j} \hbar_{n_{\ell}}} \operatorname{Tr}[e^{-2\pi i k \cdot \hat{q}_{j}} \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \ dk \\ &= \int_{\mathbf{R}^{d}} \overline{\alpha_{0}(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{is\omega(k)} e^{2\pi i k \cdot q_{0j} \hbar_{n_{\ell}}} \operatorname{Tr}[W_{1}(-2\pi k, 0) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \ dk \\ &= \int_{\mathbf{R}^{d}} \overline{\alpha_{0}(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{is\omega(k)} e^{2\pi i k \cdot q_{0j} \hbar_{n_{\ell}}} \operatorname{Tr}[\mathcal{W}(-2\pi k, 0, 0) \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \ dk \\ &= \int_{\mathbf{R}^{d}} \overline{\alpha_{0}(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{is\omega(k)} e^{2\pi i k \cdot q_{0j} \hbar_{n_{\ell}}} \operatorname{Tr}[e^{-\frac{i\hbar_{n_{\ell}}}{2}\Im m \langle i2\pi k, q_{0} + ip_{0} \rangle} \ \mathcal{W}(p_{0} - 2\pi k e_{j}, q_{0}, \alpha_{0}) \ \tilde{\varrho}_{\hbar_{n_{\ell}}}(s)] \ dk. \end{aligned}$$

We conclude

$$\begin{split} &\lim_{\ell \to +\infty} \operatorname{Tr}[\langle \alpha_0, \tilde{g}_j(s) \rangle_{L^2} \ \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] \\ &= \int_{\mathbf{R}^d} \overline{\alpha_0(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{is\omega(k)} \lim_{\ell \to +\infty} \operatorname{Tr}[\mathcal{W}(p_0 - 2\pi k, q_0, \alpha_0) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] \ dk \\ &= \int_{\mathbf{R}^d} \overline{\alpha_0(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} \ e^{is\omega(k)} \ \int_{X^0} e^{-2\pi i k \cdot q_j} e^{Q(\xi, u)} \ d\tilde{\mu}_s(u) dk \end{split}$$

By Fubini, we get

$$\lim_{\ell \to +\infty} \operatorname{Tr}[\langle \alpha_0, \tilde{g}_j(s) \rangle_{L^2} \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)]$$

= $\int_{X^0} e^{Q(\xi, u)} \int_{\mathbf{R}^d} \overline{\alpha_0(k)} \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot q_j + i s \omega(k)} \ dk \ d\tilde{\mu}_s(u)$
= $\int_{X^0} e^{Q(\xi, u)} \ \langle \alpha_0, g_j(s) \rangle_{L^2} \ d\tilde{\mu}_s(u).$

Similar discussions also work to prove

$$\lim_{\ell \to +\infty} \operatorname{Tr}[\langle \tilde{g}_j(s), \alpha_0 \rangle_{L^2} \mathcal{W}(\xi) \ \tilde{\varrho}_{\hbar_{n_\ell}}(s)] = \int_{X^0} \langle g_j(s), \alpha_0 \rangle_{L^2} \ e^{Q(\xi, u)} \, d\tilde{\mu}_s(u).$$

5.2. The characteristic equation. Below, we derive the final form of the time-evolution equation satisfied by the Wigner measure $\tilde{\mu}_t$.

Corollary 5.2 (Characteristic equation). Assume (1.6) and $\omega^{1/2}\chi \in L^2(\mathbf{R}^d, dk)$. Then, the characteristic equation (5.1) can be further reduced to the following form

$$\int_{X^0} e^{2i\pi \Re e \langle y, u \rangle_{X^\sigma}} d\tilde{\mu}_t(u) = \int_{X^0} e^{2i\pi \Re e \langle y, u \rangle_{X^\sigma}} d\tilde{\mu}_{t_0}(u) + 2\pi i \int_{t_0}^t \int_{X^0} e^{2i\pi \Re e \langle y, u \rangle_{X^\sigma}} \Re e \langle v(s, u), y \rangle_{X^\sigma} d\tilde{\mu}_s(u) ds,$$
(5.7)

for all $t, t_0 \in \mathbf{R}$ and $y \in X^{\sigma}$.

Proof. Define

$$\tilde{\xi} := \left(\frac{z_0}{2i\pi}, \frac{\alpha_0}{\sqrt{2}\pi}\right) \in X^0, \quad with \quad \xi = (z_0, \alpha_0) \in X^0.$$

We claim that

$$b(s,\xi) = -2\pi \Re e \langle v(s,u), \tilde{\xi} \rangle_{X^0}.$$

Indeed, we first remark that

$$-2\pi \Re e \langle v(s,u), \tilde{\xi} \rangle_{X^0} = \underbrace{-2\pi \Re e \langle (v(s,u))_z, \frac{z_0}{2i\pi} \rangle}_{(1)} \underbrace{-2\pi \Re e \langle (v(s,u))_\alpha, \frac{\alpha_0}{\sqrt{2\pi}} \rangle_{L^2}}_{(2)}.$$

For (1), we have

$$-2\pi \Re e \langle (v(s,u))_{z}, \frac{z_{0}}{2i\pi} \rangle = -\Im m \langle (v(s,u))_{z}, z_{0} \rangle$$

$$= -\sum_{j=1}^{n} \left((v(s,u))_{q_{j}} \cdot p_{0j} - (v(s,u))_{p_{j}} \cdot q_{0j} \right)$$

$$= -\sum_{j=1}^{n} \nabla f_{j}(p_{j}) \cdot p_{0j} - \sum_{j=1}^{n} \nabla q_{j} V(q) \cdot q_{0j} + \sum_{j=1}^{n} \left(\langle \alpha, b_{j}^{0}(s) \rangle_{L^{2}} + \langle b_{j}^{0}(s), \alpha \rangle_{L^{2}} \right),$$

where recall that v(s, u) is as in (2.6) and $b_j^0(s)$ is as in (5.3).

For (2), we have

$$-2\pi \Re e \langle (v(s,u))_{\alpha}, \frac{\alpha_0}{\sqrt{2\pi}} \rangle_{L^2} = -\sqrt{2} \Re e \langle (v(s,u))_{\alpha}, \alpha_0 \rangle_{L^2}$$
$$= -\sqrt{2} \Re e \langle -i \sum_{j=1}^n \frac{\chi(k)}{\sqrt{\omega(k)}} e^{-2\pi i k \cdot q_j + i s \omega(k)}, \alpha_0 \rangle_{L^2}$$
$$= \sqrt{2} \sum_{j=1}^n \Im m \langle g_j(s), \alpha_0 \rangle_{L^2}.$$

where $g_j(s)$ is as in (3.16). On the other hand, we have

$$\frac{i}{\sqrt{2}} \Big(\langle \alpha_0, g_j(s) \rangle_{L^2(\mathbf{R}^d, \mathbf{C})} - \langle g_j(s), \alpha_0 \rangle_{L^2(\mathbf{R}^d, \mathbf{C})} \Big) \\ = -\sqrt{2} \Im m \langle \alpha_0, g_j(s) \rangle_{L^2(\mathbf{R}^d, \mathbf{C})} = \sqrt{2} \Im m \langle g_j(s), \alpha_0 \rangle_{L^2(\mathbf{R}^d, \mathbf{C})}$$

And, thus combining the above arguments, we prove the claimed results. The Characteristic equation (5.1) becomes then

$$\int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_t(u) = \int_{X^0} e^{Q(\xi,u)} d\tilde{\mu}_{t_0}(u) + 2\pi i \int_{t_0}^t \int_{X^0} e^{Q(\xi,u)} \Re e^{\langle v(s,u), \tilde{\xi} \rangle_{X^0}} d\tilde{\mu}_s(u) ds.$$
(5.8)

We have, with $Q(\xi, u)$ as in (5.2), that

$$Q(\xi, u) = 2\pi i \Re e \langle \tilde{\xi}, u \rangle_{X^0}.$$
(5.9)

We have also for all $y = (p, q, \alpha) \in X^{2\sigma}$ and all $\tilde{\xi} = (p, q, \omega^{2\sigma}\alpha) \in X^0$ that

$$\begin{aligned} &\Re e\langle y, u\rangle_{X^{\sigma}} = \Re e\langle \xi, u\rangle_{X^{0}}, \\ &\Re e\langle v(s, u), y\rangle_{X^{\sigma}} = \Re e\langle v(s, u), \tilde{\xi}\rangle_{X^{0}} \end{aligned}$$
(5.10)

By this way, plugging (5.9)-(5.10) in (5.8) gives that (5.7) is valid for all $y \in X^{2\sigma}$. The latter could be extended to all $y \in X^{\sigma}$ by dominated convergence theorem and the bound (5.12). \Box

5.3. The Liouville equation. In this part, we relate the characteristic equation (5.7) satisfied by the set of Wigner measures $(\tilde{\mu}_t)_{t \in \mathbf{R}}$ to a special Liouville equation. To do that, we need to have some integrability condition of the vector field v of (IVP) with respect to this Wigner measure and some regularities of the latter measure.

Lemma 5.3 (Integrability of the vector field v). Assume (1.6) and (1.7) hold true. Then, there exists a constant C > 0 such that for all $u = (p, q, \alpha) \in X^{\sigma}$,

$$\|v(t,u)\|_{X^{\sigma}} \le C(\|u\|_{X^{0}}^{2}+1).$$
(5.11)

Moreover, for any bounded open interval I,

$$\int_{I} \int_{X^{\sigma}} \|v(t,u)\|_{X^{\sigma}} d\tilde{\mu}_t(u) dt < +\infty.$$

$$(5.12)$$

Proof. The non-autonomous vector field v is defined in terms of the nonlinearity \mathcal{N} as indicated in (2.6). Then it is not hard to see by looking at the proof of Proposition 2.4 that

• in the semi-relativistic case, since the function $\nabla f_j(\hat{p}_j)$ is bounded, we get

$$\|v(t,u)\|_{X^{\sigma}} \le C(\|\alpha\|_{L^{2}}^{2}+1).$$
(5.13)

• in the non-relativistic case, we get

$$\|v(t,u)\|_{X^{\sigma}} \le C(\|\alpha\|_{L^{2}}^{2} + |p|^{2} + 1).$$
(5.14)

Thus, both inequalities (5.13) and (5.14) lead to (5.11). Now, the integrability condition (5.12) is a consequence of (4.7) in Proposition 4.7.

We establish now some regularity of the Wigner measures $(\tilde{\mu}_t)_{t\in\mathbf{R}}$ with respect to time.

Lemma 5.4 (Regular properties of the Wigner Measure $\tilde{\mu}_t$). The Wigner measures $(\tilde{\mu}_t)_{t \in \mathbf{R}}$ extracted in Proposition 4.7 satisfy

- (i) $\tilde{\mu}_t$ concentrates on X^{σ} i.e. $\tilde{\mu}_t(X^{\sigma}) = 1$;
- (ii) $\mathbf{R} \ni t \mapsto \tilde{\mu}_t \in \mathcal{P}(X^{\sigma})$ is weakly narrowly continuous.

Proof. For the first assertion (i), we have from Proposition 4.7 that

$$\int_{X^0} \|u\|_{X^{\sigma}}^2 \, d\tilde{\mu}_t(u) < C$$

And, from the Markov's inequality, we have

$$\tilde{\mu}_t(\{u \in X^0 : \|u\|_{X^{\sigma}} \ge \varepsilon\}) \le \frac{1}{\varepsilon^2} \ \tilde{\mu}_t(\|u\|_{X^{\sigma}}).$$

Let $\varepsilon \to \infty$, we get

$$\tilde{\mu}_t(\{u \in X^0; \ u \notin X^\sigma\}) = 0.$$

Hence, we get that the measure $\tilde{\mu}_t$ is concentrated in X^{σ} . The second assertion (ii) is proved in a similar fashion as in [6, Lemma 5.5] using Prokhorov's Theorem.

In the coming discussions, for more details, we refer the reader to Appendix A in [6]. Let I be an open bounded interval. Define the space of smooth cylindrical functions on $I \times X^{\sigma}$, denoted by $C_{0,cul}^{\infty}(I \times X^{\sigma})$, as follows

$$\begin{aligned} \mathcal{C}^{\infty}_{0,cyl}(I \times X^{\sigma}) &:= \Big\{ \phi : I \times X^{\sigma} \to \mathbf{R}; \ \phi(t,u) = \psi(t,\pi(u)), \ \psi \in \mathcal{C}^{\infty}_{0}(I \times \mathbf{R}^{d'}), \\ \pi : X^{\sigma} \to \mathbf{R}^{d'}, \ d' \in \mathbf{N} \Big\}, \end{aligned}$$

where $\pi: X^{\sigma} \to \mathbf{R}^{d'}$ is a projection of the form $\pi: u \to \pi(u) = (\Re e \langle u, e_1 \rangle_{X^{\sigma}}, \cdots, \Re e \langle u, e_{d'} \rangle_{X^{\sigma}})$, with $(e_1, \cdots, e_{d'})$ is an arbitrary orthonormal family of X^{σ} .

Proposition 5.5. The family of Wigner measures $(\tilde{\mu}_t)_{t \in \mathbf{R}}$ defined in Proposition 4.7 is a weakly narrowly continuous solution to the following Liouville equation

$$\int_{I} \int_{X^{\sigma}} \{\partial_{t} \phi(t, u) + \Re e \langle v(t, u), \nabla \phi(t, u) \rangle_{X^{\sigma}} \} d\tilde{\mu}_{t}(u) \ dt = 0,$$
(LE)

for any bounded open interval I containing the origin with $\phi \in C_{0,cul}^{\infty}(I \times X^{\sigma})$.

Proof. It is a direct consequence of Lemma B.1 by selecting $H \equiv X^{\sigma}$ which is a Hilbert space. More precisely, all the prerequists of Lemma B.1 are satisfied. Indeed, we have

- from Corollary 5.2 that the set of Wigner measures $\{\tilde{\mu}_t\}_{t\in I}$ solves the characteristic equation (5.7);
- from Lemma 5.4, we have checked that $\tilde{\mu}_t \in \mathcal{P}(X^{\sigma})$ is a weakly narrowly continuous;
- from Lemma 5.3, we have checked the integrability condition of v with respect to $\tilde{\mu}_t$.

And thus the result follows.

6. Proof of the main result

In order to prove the main Theorems 1.1 and 1.3, we must establish some identities. It is important to note that the statement of Theorem 1.1 is not related directly to the quantum dynamics and does not require any restrictions on it. Therefore, our plan is to ensure that the assumptions (1.8) and (1.9) are applied to a specific class of density matrices, namely the coherent states. To achieve this, we must first define the coherent states for the particle and field components separately, and then generalize to the entire interacting space since we are dealing with an interaction between particles and field. Let $u_0 = (z_0, \alpha_0) \in X^0$ and consider the family of coherent states

$$\mathcal{C}_{\hbar}(u_0) = \left| W_1(\frac{\sqrt{2}}{i\hbar} z_0) \psi \otimes W_2(\frac{\sqrt{2}}{i\hbar} \alpha_0) \Omega \right\rangle \left\langle W_1(\frac{\sqrt{2}}{i\hbar} z_0) \psi \otimes W_2(\frac{\sqrt{2}}{i\hbar} \alpha_0) \Omega \right|$$

where we have introduced

- → the coherent vector: $W_1(\frac{\sqrt{2}}{i\hbar}z_0)\psi$, centered on $z_0 \in \mathbf{C}^{dn}$ where $\psi(x) = (\pi\hbar)^{-dn/4} e^{-x^2/2\hbar} \in L^2(\mathbf{R}^{dn}, dx)$ is the normalized gaussian function on the particles related to the particle space $L^2(\mathbf{R}^{dn}, \mathbf{C})$.
- \rightarrow the coherent vector: $W_2(\frac{\sqrt{2}}{i\hbar}\alpha_0)\Omega$ in the Fock space, for $\alpha \in \mathcal{G}^0$ and Ω is the vacuum vector on the fock space.

It bears noting that these family of coherent states gives rise to a family of density matrices satisfying the assumptions (1.8) and (1.9).

Lemma 6.1 (The family of coherent states). The family of coherent states $(C_{\hbar}(u_0))_{\hbar \in (0,1)}$. satisfies

$$\mathcal{M}(\mathcal{C}_{\hbar}(u_0), \hbar \in (0, 1)) = \{\delta_{u_0}\},\$$

where δ_{u_0} is the Dirac measure centered on u_0 . Moreover, if $u_0 = (z_0, \alpha_0) \in X^{\sigma}$, then $(\mathcal{C}_{\hbar}(u_0))_{\hbar \in (0,1)}$ satisfies (1.8) and (1.9).

Proof. We have

$$\operatorname{Tr}(\mathcal{C}_{\hbar}(u_{0}) \ d\Gamma(\omega^{2\sigma})) = \|\alpha_{0}\|_{\mathcal{G}^{\sigma}}^{2}$$
$$\operatorname{Tr}(\mathcal{C}_{\hbar}(u_{0}) \ \hat{p}^{2}) = \langle \psi, \hat{p}^{2}\psi \rangle - 2p_{0}^{2}$$
$$\operatorname{Tr}(\mathcal{C}_{\hbar}(u_{0}) \ \hat{q}^{2}) = \langle \psi, \hat{q}^{2}\psi \rangle - 2q_{0}^{2}.$$

Below, we give useful lemma which relates the Wigner measure $\tilde{\mu}_t$ to μ_t in terms of the free field flow Φ_t^f .

Lemma 6.2 (Relations between the sets of Wigner measure). Let $(\varrho_{\hbar})_{h \in (0,1)}$ be a family of density matrices satisfying (1.8) and (1.9). Define

$$\tilde{\varrho}_{\hbar}(t) := e^{i\frac{t}{\hbar}d\Gamma(\omega)} \varrho_{\hbar} e^{-i\frac{t}{\hbar}d\Gamma(\omega)}$$

Then, we can assert that

- (1) the family of states $(\tilde{\varrho}_{\hbar}(t))_{\hbar \in (0,1)}$ satisfies (1.8) and (1.9);
- (2) for all sequences $(\hbar_n)_{n \in \mathbf{N}}$ with $\hbar_n \to 0$, there exists a subsequence \hbar_{n_ℓ} with $\hbar_{n_\ell} \to 0$ such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N}) = \{(\Phi_{-t}^{J})_{\sharp}\mu; \ \mu \in \mathcal{M}(\varrho_{\hbar_{n_{\ell}}}, \ell \in \mathbf{N})\},$$

where Φ^f_t is the free field flow as in (2.5).

Proof. The first assertion is a consequence of Lemma 4.6. Let $\mu \in \mathcal{M}(\varrho_{\hbar_{\ell}}, \ell \in \mathbf{N})$ and $\tilde{\mu}_t \in \mathcal{M}(\tilde{\varrho}_{\hbar_{\ell}}(t), \ell \in \mathbf{N})$. On one hand, we have

$$\lim_{\ell} \operatorname{Tr} \left[\tilde{\varrho}_{\hbar_{\ell}}(t) \ \mathcal{W}(\xi) \right] = \lim_{\ell} \operatorname{Tr} \left[\varrho_{\hbar_{\ell}} \ e^{-i\frac{t}{\hbar}d\Gamma(\omega)} \mathcal{W}(\xi) \ e^{i\frac{t}{\hbar}d\Gamma(\omega)} \right] = \lim_{\ell} \operatorname{Tr} \left[\varrho_{\hbar_{\ell}} \ \mathcal{W}(\Phi_{t}^{f}(\xi)) \right]$$
$$= \int_{X^{0}} e^{Q(\Phi_{t}^{f}(\xi), u)} \ d\mu(u) = \int_{X^{0}} e^{Q(\xi, \Phi_{-t}^{f}(u))} \ d\mu(u) = \int_{X^{0}} e^{Q(\xi, u)} \ d(\Phi_{-t}^{f})_{\sharp} \mu(u).$$

On the other hand, we have

$$\lim_{\ell} \operatorname{Tr} \left[\tilde{\varrho}_{\hbar_{\ell}}(t) \ \mathcal{W}(\xi) \right] = \int_{X^0} e^{Q(\xi, u)} \ d\tilde{\mu}_t(u).$$

We conclude then that

$$\tilde{\mu}_t = (\Phi^f_{-t})_{\sharp} \mu.$$

Below, we start the proof Theorems 1.1 and 1.3.

Proof of Theorem 1.1. Let $u_0 \in X^{\sigma}$ and defines the density matrices for all $\hbar \in (0, 1)$ as follows

$$\varrho_{\hbar} := \mathcal{C}_{\hbar}(u_0).$$

Then since $u_0 \in X^{\sigma}$, we can assert by Lemma 6.1 that the family of density matrices $(\varrho_{\hbar})_{\hbar \in (0,1)}$ satisfies (1.8) and (1.9). Thus, with this choice of density matrices and using the arguments of Proposition 4.7, we can assert that for each sequence $(\hbar_n)_{n \in \mathbb{N}}$ with $\hbar_n \to 0$, there exists a subsequence $(\hbar_{n_\ell})_{\ell \in \mathbb{N}}$ with $\hbar_{n_\ell} \to 0$ and a family of Borel probability measure $\{\tilde{\mu}_t\}_{t \in \mathbb{R}}$ in X^0 subsequence $(\hbar_{n_\ell})_{\ell \in \mathbb{N}}$ with $\hbar_{n_\ell} \to 0$ and a family of Borel probability measure $\{\tilde{\mu}_t\}_{t \in \mathbb{R}}$ in X^0

such that

$$\mathcal{M}(e^{i\frac{t}{\hbar_{n_{\ell}}}d\Gamma(\omega)} e^{-i\frac{t}{\hbar_{n_{\ell}}}\hat{H}} \mathcal{C}_{\hbar_{n_{\ell}}}(u_0) e^{i\frac{t}{\hbar_{n_{\ell}}}\hat{H}} e^{-i\frac{t}{\hbar_{n_{\ell}}}d\Gamma(\omega)}, \ell \in \mathbf{N}) = \{\tilde{\mu}_t\}.$$

Now, on one hand, we do have from Proposition 5.5 that $\{\tilde{\mu}_t\}_{t\in\mathbf{R}}$ is weakly narrowly continuous solution to the Liouville equation (LE); from the other hand, from Lemma 5.3, we can assert that all the prerequists to apply Theorem B.2 are in our hand. To recover the proof of Theorem 1.1, we follow the steps below.

⊲ We apply Theorem B.2 with the measure $\tilde{\mu}_t$ obtained above, we get the global well posedness of the initial value problem (IVP) $\tilde{\mu}_0$ -almost all initial data in X^σ as well as the existence of a generalized Borel measurable global flow $\tilde{\Phi}_t$ as follows

$$\begin{array}{ccccc} \tilde{\Phi}_t : \mathfrak{G} & \longrightarrow & X^{\sigma} \\ & u_0 & \longmapsto & u(t), \end{array}$$

where \mathfrak{G} is the ensemble of initial data obtained from Theorem B.2.

- \triangleleft Let $u_0 \in X^{\sigma}$. From Lemma 6.1, we have $\tilde{\mu}_0(\mathfrak{G}) = \delta_{u_0}(\mathfrak{G}) = 1$. This implies $u_0 \in \mathfrak{G}$;
- \triangleleft Use the equivalence between the solution to (IVP) and (1.1), we can show the existence and uniqueness of the solution of (1.1) with a generalized global flow

$$\Phi_t(u_0) = \Phi_t^f \circ \tilde{\Phi}_t(u_0)$$

where Φ_t^f is the free flow and $\tilde{\Phi}_t$ is the generalized flow of (IVP);

Proof of Theorem 1.3. We have here to prove the validity of Bohr's correspondence principle. Assume $(\varrho_{\hbar})_{\hbar}$ is a family of density matrices satisfying the Assumptions (1.8) and (1.9). Then, using Proposition 4.7, we can assert that for each sequence $(\hbar_n)_{n\in\mathbb{N}}$ with $\hbar_n \to 0$, there exists a subsequence $(\hbar_{n_{\ell}})_{\ell\in\mathbb{N}}$ with $\hbar_{n_{\ell}} \to 0$ and a family of Borel probability measures $\{\tilde{\mu}_t\}_{t\in\mathbb{R}}$ in X^0 such that

$$\mathcal{M}(\tilde{\varrho}_{\hbar_{n\ell}}(t), \ell \in \mathbf{N}) = \{\tilde{\mu}_t\}.$$

By Lemma 6.2, we have

$$\mathcal{M}(\varrho_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N}) = \{ (\Phi_t^f)_{\sharp} \tilde{\mu}_t; \ \tilde{\mu}_t \in \mathcal{M}(\tilde{\varrho}_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N}) \}.$$

This implies that

$$\mathcal{M}(\varrho_{\hbar_{n_{\ell}}}(t), \ell \in \mathbf{N}) = \{\mu_t\} = \{(\Phi^f_t)_{\sharp} \tilde{\mu}_t\}$$

From (ii) in Porbabilistic representation, we can assert that for any bounded Borel functions $\psi: X^{\sigma} \to \mathbf{R}$

$$\int_{X^{\sigma}} \psi(u) \ d\tilde{\mu}_t(u) = \int_{\mathcal{F}_I} \psi(e_t(u_0, u(\cdot))) \ d\eta(u_0, u(\cdot)).$$

Since, we have the generalized global flow Φ_t to (IVP), we get

$$e_t(u_0, u(\cdot)) = \tilde{\Phi}_t(e_0(u_0, u(\cdot))) = \tilde{\Phi}_t(u_0)$$

This gives

$$\int_{X^{\sigma}} \psi(u) \ d\tilde{\mu}_t(u) = \int_{X^{\sigma} \times \mathcal{C}(\overline{I}, X^{\sigma})} \psi \circ \tilde{\Phi}_t(e_0(u_0, u(\cdot))) \ d\eta(u_0, u(\cdot)) = \int_{X^{\sigma}} \psi \circ \tilde{\Phi}_t(u) \ d\tilde{\mu}_0(u).$$

We conclude that $\tilde{\mu}_t = (\Phi_t)_{\sharp} \tilde{\mu}_0$. This implies that:

$$\mu_t = (\Phi_t^f)_{\sharp} \tilde{\mu}_t = (\Phi_t^f \circ \tilde{\Phi}_t)_{\sharp} \tilde{\mu}_0 = (\Phi_t)_{\sharp} \tilde{\mu}_0 = (\Phi_t)_{\sharp} \mu_0$$

and where we have used $\tilde{\mu}_0 = \mu_0$ as a consequence of

$$\tilde{\varrho}_{\hbar}(0) = \varrho_{\hbar}(0) = \varrho_{\hbar}$$

APPENDIX A. PROKHOROV THEOREM

Let X be separable metric space. The proof of the following result is proved in [3, Theorem 5.1.3].

Theorem A.1 (Prokhorov Theorem). If a set $\mathcal{K} \subset \mathcal{P}(X)$ is tight i.e.

 $\forall \varepsilon > 0, \exists K_{\varepsilon} \text{ compact in } X \text{ such that } \mu(X \setminus K_{\varepsilon}) \leq \varepsilon, \forall \mu \in \mathcal{K},$

then \mathcal{K} is relatively compact in $\mathcal{P}(X)$.

APPENDIX B. USEFUL RESULTS

The following results relate the Liouville equations and the Characteristic equations satisfied by a family of Wigner measures. For more details, we refer the reader to [29, Proposition 4.2]. Let H be a Hilbert space.

Lemma B.1 (Equivalence). Let $v : \mathbf{R} \times \mathbf{H} \to \mathbf{H}$ be a continuous vector field such that it is bounded on bounded sets. Let $I \ni t \to \mu_t$ a weakly narrowly continuous curve in $\mathcal{P}(\mathbf{H})$ such that we have the following integrability condition

$$\int_I \int_{\mathcal{H}} \|v(t,u)\|_{\mathcal{H}} \ d\mu_t(u) \ dt < +\infty.$$

Then, the following statements are equivalent:

- (i) $\{\mu_t\}_{t\in I}$ is a solution of Liouville equation (LE);
- (ii) $\{\mu_t\}_{t\in I}$ solves the characteristic equation (5.7) for all $t \in I$ and for all $y \in H$.

The subsequent outcomes illustrate how to build the global solution to the (IVP) utilizing measure-theoretical approaches and certain probabilistic representations of the measure-valued solutions for the Liouville equation. Additional information on the topic can be found in [7] or in the Appendices of [6].

Theorem B.2 (Global flow of the initial value problem). Let $v : \mathbf{R} \times X^{\sigma} \to X^{\sigma}$ be a continuous vector field bounded on bounded sets. Assume

- $\exists t \in \mathbf{R} \to \tilde{\mu}_t \in \mathcal{P}(X^{\sigma})$ a weakly narrowly continuous solution to (LE) satisfying the integrability condition (5.12) on I;
- There is at most one solution of the initial value problem (IVP) over any bounded open interval I containing the origin.

Then for $\tilde{\mu}_0$ -almost all initial conditions u_0 in X^{σ} , there exists a unique global strong solution to (IVP). In addition, the set

 $\mathfrak{G} := \{ u_0 \in X^{\sigma} : \exists u(\cdot) \text{ a global strong solution of (IVP)} \\ with the initial condition u_0 \},\$

is Borel subset of X^{σ} with $\tilde{\mu}_0(\mathfrak{G}) = 1$ and for any time $t \in \mathbf{R}$ the map $u_0 \in \mathfrak{G} \to \tilde{\Phi}_t(u_0) = u(t)$ is Borel measurable.

Proposition B.3 (Superposition principle). There exists $\eta \in \mathcal{P}(X^{\sigma} \times \mathcal{C}(\overline{I}, X^{\sigma}))$ satisfying: (i) $\eta(\mathcal{F}_I) = 1$ where

$$\mathcal{F}_I := \left\{ (u_0, u(\cdot)) \in X^{\sigma} \times \mathcal{C}(\overline{I}, X^{\sigma}) : u(\cdot) \text{ satisfies (IVP) on } I \text{ with } u_0 \right\}$$

(ii) $\tilde{\mu}_t = (e_t)_{\sharp} \eta$, $\forall t \in I$, where the map

$$e_t: (u_0, u(\cdot)) \in X^{\sigma} \times \mathcal{C}(\overline{I}, X^{\sigma}) \to u(t) \in X^{\sigma}$$

is the evaluation map.

Appendix C. Technical results about convergence

Finally, we prove two technical lemmas which are useful for the study of the quantum-classical convergence in Subsection 5.1. We denote by \mathcal{F} the Fourier transform on \mathbf{R}^d .

Lemma C.1. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices on the Hilbert space \mathcal{H} satisfying (1.8)-(1.9) for $\sigma = \frac{1}{2}$. Assume that for some sequence $(\hbar_{\ell})_{\ell \in \mathbb{N}} \subset (0,1)$, $\hbar_{\ell} \to 0$, there exists a (unique) Borel measure $\mu \in \mathcal{P}(X^0)$ such that

$$\mathcal{M}(\varrho_{\hbar_{\ell}}, \ell \in \mathbf{N}) = \{\mu\}.$$

Then for any $\varphi \in L^2(\mathbf{R}^d)$, $\beta \in \mathcal{F}(L^1(\mathbf{R}^d))$ and $\xi \in X^0$, $j = 1, \cdots, n$,

$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\beta(\hat{q}_j) \, \hat{a}_{\hbar_{\ell}}(\varphi) \, \mathcal{W}(\xi) \, \varrho_{\hbar_{\ell}} \right] = \int_{X^0} \beta(q_j) \langle \varphi, \alpha \rangle_{L^2(\mathbf{R}^d)} \, e^{Q(\xi, u)} \, d\mu(u) \,,$$
$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\beta(\hat{q}_j) \, \hat{a}_{\hbar_{\ell}}^*(\varphi) \, \mathcal{W}(\xi) \, \varrho_{\hbar_{\ell}} \right] = \int_{X^0} \beta(q_j) \langle \alpha, \varphi \rangle_{L^2(\mathbf{R}^d)} \, e^{Q(\xi, u)} \, d\mu(u) \,,$$

with $u = (p,q,\alpha) \in X^0$ and $Q(\cdot, \cdot)$ is the phase given in (5.2).

Proof. The two limits are similar. By linear combinations one can use instead the fields operators

$$\hat{\phi}_{\hbar}(\varphi) = rac{\hat{a}_{\hbar}^{*}(\varphi) + \hat{a}_{\hbar}(\varphi)}{\sqrt{2}}$$
 and $\hat{\pi}_{\hbar}(\varphi) = rac{i\hat{a}_{\hbar}^{*}(\varphi) - i\hat{a}_{\hbar}(\varphi)}{\sqrt{2}}$.

So, it is enough to show

$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\beta(\hat{q}_j) \, \hat{\phi}_{\hbar_\ell}(\varphi) \, \mathcal{W}(\xi) \, \varrho_{\hbar_\ell} \right] = \sqrt{2} \int_{X^0} \beta(q_j) \Re e^{\langle \alpha, \varphi \rangle_{L^2(\mathbf{R}^d)}} \, e^{Q(\xi, u)} \, d\mu(u) \,. \tag{C.1}$$

Our goal is to prove (C.1). Since $\beta = \mathcal{F}(g)$ for some $g \in L^1(\mathbf{R}^d)$, one can write

$$\operatorname{Tr}\left[\beta(\hat{q}_{j})\,\hat{\phi}_{\hbar_{\ell}}(\varphi)\,\mathcal{W}(\xi)\,\varrho_{\hbar_{\ell}}\right] = \int_{\mathbf{R}^{d}}g(y)\,\operatorname{Tr}\left[e^{-2\pi i y\cdot\hat{q}_{j}}\,\hat{\phi}_{\hbar_{\ell}}(\varphi)\,\mathcal{W}(\xi)\,\varrho_{\hbar_{\ell}}\right]dy.\tag{C.2}$$

Furthermore, dominated convergence applies to the right hand side of (C.2) thanks to the assumptions (1.8)-(1.9) and the estimates in Lemma 3.1. Thus, the limit (C.1) reduces to

$$\lim_{\ell \to \infty} \operatorname{Tr} \left[e^{-2\pi i y \cdot \hat{q}_j} \, \hat{\phi}_{\hbar_\ell}(\varphi) \, \mathcal{W}(\xi) \, \varrho_{\hbar_\ell} \right] = \sqrt{2} \int_{X^0} e^{-2\pi i y \cdot q_j} \, \Re e^{\langle \alpha, \varphi \rangle_{L^2(\mathbf{R}^d)}} \, e^{Q(\xi, u)} \, d\mu(u) \,. \tag{C.3}$$

Now, applying [6, Lemma B.2], we obtain (C.3) for all $y \in \mathbf{R}^d$ since

$$e^{-2\pi i y \cdot \hat{q}_j} \mathcal{W}(\xi) = W_1(-2\pi y, 0) \, \mathcal{W}(\xi) = \mathcal{W}(-2\pi y, 0, 0) \, \mathcal{W}(\xi) = e^{i\hbar_\ell \pi y \cdot p_0} \mathcal{W}(\tilde{\xi})$$

with $\tilde{\xi} = (-2\pi y, 0, 0) + \xi$ and $\xi = (p_0, q_0, \alpha)$. Recall that the Weyl-Heisenberg operator $W_1(\cdot)$ is given in (3.10) while $\mathcal{W}(\cdot)$ is defined by (3.9)-(3.7).

Lemma C.2. Let $(\varrho_{\hbar})_{\hbar \in (0,1)}$ be a family of density matrices on the Hilbert space \mathcal{H} satisfying (1.8)-(1.9) for $\sigma = \frac{1}{2}$. Assume that for some sequence $(\hbar_{\ell})_{\ell \in \mathbf{N}} \subset (0,1), \ \hbar_{\ell} \to 0$, there exists a (unique) Borel measure $\mu \in \mathcal{P}(X^0)$ such that

$$\mathcal{M}(\varrho_{\hbar_{\ell}}, \ell \in \mathbf{N}) = \{\mu\}.$$

Then for any $\varphi \in L^2(\mathbf{R}_k^d), \xi \in X^0$ and $j = 1, \cdots, n$,

$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\hat{a}_{\hbar_{\ell}} (e^{-2\pi i \, k \cdot \hat{q}_{j}} \varphi) \, \mathcal{W}(\xi) \, \varrho_{\hbar_{\ell}} \right] = \int_{X^{0}} \langle e^{-2\pi i \, k \cdot q_{j}} \varphi, \alpha \rangle_{L^{2}(\mathbf{R}_{k}^{d})} \, e^{Q(\xi, u)} \, d\mu(u) \,,$$
$$\lim_{\ell \to \infty} \operatorname{Tr} \left[\hat{a}_{\hbar_{\ell}}^{*} (e^{-2\pi i \, k \cdot \hat{q}_{j}} \varphi)) \, \mathcal{W}(\xi) \, \varrho_{\hbar_{\ell}} \right] = \int_{X^{0}} \langle \alpha, e^{-2\pi i \, k \cdot q_{j}} \varphi \rangle_{L^{2}(\mathbf{R}_{k}^{d})} \, e^{Q(\xi, u)} \, d\mu(u) \,,$$

with $u = (p, q, \alpha) \in X^0$ and $Q(\cdot, \cdot)$ is the phase given in (5.2).

Proof. According to Definition 1.2 of Wigner measures and [10, Theorem 6.2 and Proposition [6.4], we deduce that

$$\mathcal{M}(\mathcal{W}(\xi)\varrho_{\hbar_{\ell}}, \ell \in \mathbf{N}) = \{e^{Q(\xi, \cdot)}\mu\}.$$

Here, we have used the extension of the notion of Wigner measures to trace-class operators which are not necessary non-negative nor trace normalized (see [10, Proposition 6.4]). Let $\{e_m\}_{m\in\mathbb{N}}$ be O.N.B of the Hilbert space $L^2(\mathbf{R}_k^d)$. The two limits are similar (almost conjugate) and it is enough to explain the argument for the second one. We denote $\rho_{\hbar\ell}(\xi) := \mathcal{W}(\xi) \rho_{\hbar\ell}$ and $b(\hat{q}_i) = e^{-2\pi i k \cdot \hat{q}_j} \varphi$. We have

$$\begin{aligned} \left| \operatorname{Tr} \left[\hat{a}_{\hbar_{\ell}}^{*}(b(\hat{q}_{j})) \ \varrho_{\hbar_{\ell}}(\xi) \right] - \int_{X^{0}} e^{Q(\xi,u)} \ \langle \alpha, b(q_{j}) \rangle_{L^{2}} \ d\mu(u) \right| \\ &\leq \underbrace{\left| \operatorname{Tr} \left[\left(\hat{a}_{\hbar_{\ell}}^{*}(b(\hat{q}_{j})) - \sum_{m=1}^{R} \hat{a}_{\hbar_{\ell}}^{*}(e_{m}) \mathcal{F}[\varphi \bar{e}_{m}](\hat{q}_{j}) \right] \right]_{(1)} \\ &+ \sum_{m=1}^{R} \underbrace{\left| \operatorname{Tr} \left[\hat{a}_{\hbar_{\ell}}^{*}(e_{m}) \mathcal{F}[\varphi \bar{e}_{m}](\hat{q}_{j}) \right] \right]_{(1)} \\ &+ \underbrace{\left| \int_{X^{0}} e^{Q(\xi,u)} \left(\sum_{m=1}^{R} \langle \alpha, e_{m} \rangle_{L^{2}} \mathcal{F}[\varphi \bar{e}_{m}](q_{j}) - \langle \alpha, b(q_{j}) \rangle_{L^{2}} \right) d\mu(u) \right| }_{(2)} , \end{aligned}$$

Using estimates as in Lemma 3.1 and assumptions (1.8)-(1.9), one proves

$$(1)^{2} \lesssim \left\| \langle x \rangle^{-1} \left(e^{-2\pi i k \cdot x} \varphi - \sum_{m=1}^{R} e_{m} \langle e_{m}, e^{-2\pi i k \cdot x} \varphi \rangle_{L^{2}(\mathbf{R}_{k}^{d})} \right) \right\|_{L^{\infty}(\mathbf{R}_{x}^{d}, L^{2}(\mathbf{R}_{k}^{d}))}^{2}$$
$$\lesssim \sup_{x \in \mathbf{R}^{d}} \sum_{m=R+1}^{\infty} \langle x \rangle^{-1} \left| \langle e_{m}, e^{-2\pi i k \cdot x} \varphi \rangle_{L^{2}(\mathbf{R}_{k}^{d})} \right|^{2}.$$

So, thanks to a further localization argument in the variable x combined to Dini's theorem, one concludes that (1) converges to zero uniformly in $\hbar \in (0,1)$ as $R \to \infty$. Similarly, using the pointwise convergence for any $x \in \mathbf{R}^d$,

$$\sum_{m=1}^{R} \langle \alpha, e_m \rangle_{L^2} \mathcal{F}[\varphi \bar{e}_m](x) - \langle \alpha, e^{-2\pi i k \cdot x} \varphi \rangle_{L^2} \xrightarrow[R \to \infty]{41} 0,$$

and dominated convergence, one concludes that (3) converges to zero as $R \to \infty$. Now, applying Lemma C.1 with $\beta := \mathcal{F}[\varphi \bar{e}_m] \in \mathcal{F}(L^1(\mathbf{R}^d))$, we obtain that (2) converges also to zero for any fixed $R \in \mathbf{N}$ as $\hbar_{\ell} \to 0$. Hence, using an $\varepsilon/3$ -argument we prove the claimed statement.

References

- A. Abdesselam and D. Hasler. Analyticity of the ground state energy for massless nelson models. Communications in Mathematical Physics, 310(2):511–536, jan 2012.
- [2] S. Albeverio. Scattering theory in a model of quantum fields. I. Journal of Mathematical Physics, 14(12):1800– 1816, December 1973.
- [3] L. Ambrosio, N. Gigli, and G. Savare. Gradient flows in metric spaces and in the space of probability measures. 01 2005.
- [4] Z. Ammari. Asymptotic completeness for a renormalized nonrelativistic hamiltonian in quantum field theory: The nelson model. *Mathematical Physics, Analysis and Geometry*, 3:217–285, 2000.
- [5] Z. Ammari and M. Falconi. Bohr's correspondence principle for the renormalized nelson model. SIAM Journal on Mathematical Analysis, 49(6):5031–5095, 2017.
- [6] Z. Ammari, M. Falconi, and F. Hiroshima. Towards a derivation of classical electrodynamics of charges and fields from qed. 2022.
- [7] Z. Ammari, S. Farhat, and V. Sohinger. Almost sure existence of global solutions for general initial value problems, 2023.
- [8] Z. Ammari and Q. Liard. On uniqueness of measure-valued solutions to Liouville's equation of Hamiltonian PDEs. Discrete Contin. Dyn. Syst., 38(2):723-748, 2018.
- [9] Z. Ammari, Q. Liard, and C. Rouffort. On well-posedness and uniqueness for general hierarchy equations of gross-pitaevskii and hartree type. Archive for Rational Mechanics and Analysis, 238, 11 2020.
- [10] Z. Ammari and F. Nier. Mean field limit for bosons and infinite dimensional phase-space analysis. Annales Henri Poincaré, 2008.
- [11] Z. Ammari and F. Nier. Mean field propagation of infinite dimensional wigner measures with a singular two-body interaction potential, 2014.
- [12] A. Arai. Ground State of the Massless Nelson Model Without Infrared Cutoff in a Non-Fock Representation. *Reviews in Mathematical Physics*, 13(9):1075–1094, January 2001.
- [13] V. Bach, J. Fröhlich, and I. M. Sigal. Spectral Analysis for Systems of Atoms and Molecules Coupled to the Quantized Radiation Field. *Communications in Mathematical Physics*, 207(2):249–290, January 1999.
- [14] T. Chen, C. Hainzl, N. Pavlović, and R. Seiringer. Unconditional uniqueness for the cubic Gross-Pitaevskii hierarchy via quantum de Finetti. Comm. Pure Appl. Math., 68(10):1845–1884, 2015.
- [15] T. Chen and N. Pavlović. Higher order energy conservation and global well-posedness of solutions for Gross-Pitaevskii hierarchies. Comm. Partial Differential Equations, 39(9):1597–1634, 2014.
- [16] M. Falconi. Classical limit of the nelson model with cutoff. Journal of Mathematical Physics, 54:012303– 012303, 2012.
- [17] J. Froehlich, M. Griesemer, and Benjamin Schlein. Asymptotic electromagnetic fields in models of quantummechanical matter interacting with the quantized radiation field. Advances in Mathematics, 164:349–398, 2000.
- [18] C. Gerard. On the Scattering Theory of Massless Nelson Models. Reviews in Mathematical Physics, 14(11):1165–1280, January 2002.
- [19] C. Gérard, F. Hiroshima, A. Panati, and A. Suzuki. Infrared problem for the nelson model on static spacetimes. *Communications in Mathematical Physics*, 308(2):543–566, sep 2011.
- [20] J. Ginibre, F. Nironi, and G. Velo. Partially classical limit of the nelson model. Annales Henri Poincare, 7:21–43, 01 2006.
- [21] K. Hepp. The classical limit for quantum mechanical correlation functions. Comm. Math. Phys., 35:265–277, 1974.
- [22] F. Hiroshima. Diamagnetic inequalities for systems of nonrelativistic particles with a quantized field. Reviews in Mathematical Physics - RMP, 8:185–203, 02 1996.
- [23] R. Høegh-Krohn. Asymptotic Fields in Some Models of Quantum Field Theory. II. Journal of Mathematical Physics, 10(4):639–643, April 1969.
- [24] A. Komech, H. Kunze, and M. Spohn. Effective Dynamics for a Mechanical Particle Coupled to a Wave Field. Communications in Mathematical Physics, 203(1):1–19, January 1999.
- [25] A. Komech and H. Spohn. Soliton-like asymptotics for a classical particle interacting with a scalar wave field. Nonlinear Analysis: Theory, Methods and Applications, 33(1):13–24, 1998.
- [26] A. Komech, H. Spohn, and M. Kunze. Long-time asymptotics for a classical particle interacting with a scalar wave field. *Communications in Partial Differential Equations*, 22(1-2):157–181, 1997.
- [27] E. Nelson. Interaction of Nonrelativistic Particles with a Quantized Scalar Field. J. Math. Phys., 5(9):1190– 1197, 1964.

- [28] E. Nelson. Schrödinger Particles Interacting with a Quantized Scalar Field. 1969.
- [29] C. Rouffort. On the general principle of the mean-field approximation for many-boson dynamics, 2018.
- [30] S. Teufel. Effective n-body dynamics for the massless nelson model and adiabatic decoupling without spectral gap. Annales Henri Poincaré, 3(5):939–965, oct 2002.