Large deviation for Gibbs probabilities at zero temperature and invariant idempotent probabilities for iterated function systems

Jairo K. Mengue *1 and Elismar R. Oliveira $^{\dagger 2}$

^{1,2}Universidade Federal do Rio Grande do Sul

May 22, 2024

Abstract

We consider two compact metric spaces J and X and a uniform contractible iterated function system $\{\phi_j : X \to X \mid j \in J\}$. For a Lipschitz continuous function A on $J \times X$ and for each $\beta > 0$ we consider the Gibbs probability $\rho_{\beta A}$. Our goal is to study a large deviation principle for such family of probabilities as $\beta \to +\infty$ and its connections with idempotent probabilities. In the non-place dependent case $(A(j, x) = A_j, \forall x \in X)$ we will prove that $(\rho_{\beta A})$ satisfy a LDP and -I (where I is the deviation function) is the density of the unique invariant idempotent probability for a mpIFS associated to A. In the place dependent case, we prove that, if $(\rho_{\beta A})$ satisfy a LDP, then -Iis the density of an invariant idempotent probability. Such idempotent probabilities were recently characterized through the Mañé potential and Aubry set, therefore we will obtain an identical characterization for -I.

Key words and phrases: iterated function systems, large deviation, idempotent measures, Maslov measures

2020 Mathematics Subject Classification: Primary 37A30, 37A50, 28A33, 46E27, 60B10,60F10; Secondary 15A80, 37C30.

1 Introduction

We consider two compact metric spaces (X, d_X) and (J, d_J) and denote by $C(X, \mathbb{R})$ and $C(J \times X, \mathbb{R})$, the set of continuous functions from X to \mathbb{R} and $J \times X$ to \mathbb{R} , respectively.

^{*}E-mail: jairo.mengue@ufrgs.br

[†]E-mail: elismar.oliveira@ufrgs.br

Definition 1.1. Let (X, d_X) and (J, d_J) be compact metric spaces. A uniformly contractible iterated function system $(X, (\phi_j)_{j \in J})$ is a family of maps $\{\phi_j : X \to X \mid j \in J\}$ satisfying: there exists $0 < \gamma < 1$ such that

$$d_X(\phi_{j_1}(x_1), \phi_{j_2}(x_2)) \le \gamma \cdot [d_J(j_1, j_2) + d_X(x_1, x_2)], \ \forall j_1, j_2 \in J, \forall x_1, x_2 \in X.$$
(1)

We also consider the continuous map $\phi: J \times X \to X$, defined by $\phi(j, x) = \phi_j(x)$.

By one hand, in [27] it is described a thermodynamic formalism for uniformly contractible IFS $(X, (\phi_j)_{j \in J})$. On the other hand, in [28] it is characterized the invariant idempotent probabilities for the so called max-plus IFS. In the present work we exhibit a connection between these two settings. Our objective is to study a large deviation principle (LDP) for Gibbs probabilities in thermodynamic formalism for IFS, when the temperature goes to zero, as an application of results in [28]. As we will see, the *Mañé potential* and *Aubry set* will play an important role in the description of the deviation function.

From now on we always consider the Borel sigma algebra in each metric space. Let $B: J \times X \to (0, +\infty)$ be a continuous function and ν be a fixed Borel probability on J satisfying $\operatorname{supp}(\nu) = J$. We define the transfer operator $\mathcal{L}_{\phi,B,\nu}: C(X,\mathbb{R}) \to C(X,\mathbb{R})$ by

$$\mathcal{L}_{\phi,B,\nu}(f)(x) = \int_J B(j,x) \cdot f(\phi_j(x)) \, d\nu(j),$$

for any $f \in C(X, \mathbb{R})$.

For each $\beta > 0$ (which is interpreted as the inverse of temperature in thermodynamic formalism) and a Lipschitz continuous function $A: J \times X \to \mathbb{R}$ we also consider the operator $\mathcal{L}_{\phi,e^{\beta A},\nu}$. It admits a unique pair $(\lambda_{\beta A}, h_{\beta A})$ where $\lambda_{\beta A}$ is a positive number and $h_{\beta A}: X \to \mathbb{R}$ is a positive and continuous function, such that $\sup_{x \in X} h_{\beta A}(x) = 1$ and $\mathcal{L}_{\phi,e^{\beta A},\nu}(h_{\beta A}) = \lambda_{\beta A} \cdot h_{\beta A}$. The function $q^{\beta}: J \times X \to \mathbb{R}$, defined by $q^{\beta}(j,x) := \frac{e^{\beta A(j,x)} \cdot h_{\beta A}(\phi_j(x))}{\lambda_{\beta A} \cdot h_{\beta A}(x)}$, satisfies $\mathcal{L}_{\phi,q^{\beta},\nu}(1) = 1$ and the dual operator of $\mathcal{L}_{\phi,q^{\beta},\nu}$ has a unique invariant probability $\rho_{\beta A}$ on X, that is,

$$\int_X f(x)d\rho_{\beta A}(x) = \int_X \int_J q^\beta(j,x) \cdot f(\phi_j(x)) \, d\nu(j) \, d\rho_{\beta A}(x), \ \forall f \in C(X,\mathbb{R}).$$
(2)

We call $\rho_{\beta A}$ a *Gibbs probability* on X. Our objective is to consider a large deviation principle for $(\rho_{\beta A})$ as $\beta \to +\infty$ and to present its connections with invariant idempotent probabilities.

The Ruelle (transfer) operator was introduced by D. Ruelle [31] in the context of equilibrium states - Gibbs measures - of infinite one-dimensional lattice gas. It was extensively studied for IFS in the last decades and a remarkable result was Fan's Theorem (see [13], Theorem 1.1) which shows the existence of a positive eigenfunction and a Gibbs probability for a contractive IFS with Dini continuous place dependent positive weights. Several extensions of this result appeared studying the Ruelle operator, its rate of convergence and the existence of eigenfunctions and eigenmeasures. In [18], for example, it is studied the conditions for the existence of such eigenfunctions together with the Perron-Frobenius property for non-expansive finite IFSs. In [33] it is studied the speed of convergence of the Ruelle operator for a finite IFS, where the weights are positive Lipschitz continuous and the system satisfies an average contractive condition. In [34], it is studied the existence of eigenfunctions and eigenmeasures of the Ruelle operator for a finite weakly contractive IFS.

A key generalization was given by Stenflo [32], where random iterations are used to represent the iterations of a so called IFS with probabilities, for an arbitrary measurable set of maps. Their approach is slightly different from the previous works on IFS with probabilities. Its main goal was to establish, when the transfer operator is Feller, that there exists a unique attracting invariant measure for the Markov chain generated by the IFS. It is worth to mention that this approach was previously introduced by [5], for an average-contractive system, to show the existence of invariant measures. In [5], it is assumed that J is finite and the maps are Lipschitz continuous, with place dependent and Dini continuous weights.

One can also introduce a thermodynamic formalism based on holonomic measures according to [23], [8], [27] and [7], where several results are obtained for fairly general IFSs.

There is a special class of IFS where the thermodynamic formalism is well known: if $J := \{1, ..., d\}$ is a finite set, $X := J^{\mathbb{N}}$ and $\phi_j(x) = \phi_j(x_1, x_2, x_3, ...) = (j, x_1, x_2, x_3, ...)$ where $x = (x_1, x_2, x_3, ...) \in X$, then $(\phi_j)_{j \in J}$ are the inverse branches of the shift map σ acting in $J^{\mathbb{N}}$. We call this model by full shift for finite alphabet. In this case, the thermodynamic formalism for zero temperature is also widely studied as well as its connection with ergodic optimization and max-plus algebra [2]. Considering this case, results concerning large deviations, when the temperature goes to zero, can be founded in [3] and [26].

In the setting of full shift for finite alphabet it is proved in [3] (more generally for sub-shifts of finite type) that $(\rho_{\beta A})$ satisfies a LDP, since A has a unique maximizing measure. In this case the deviation function I evaluated at any point $x \in X$, is given as by $I(x) = R(x) + R(\sigma(x)) + R(\sigma^2(x)) + \ldots$, where the function R is also presented. This is problematic for a general IFS, because we have at hands just the maps ϕ_j which in the full shift correspond to the inverse branches of σ .

In [26] the main idea is to use Prop. 3.2 of [30]. We believe that such strategy can not be replicated from the full shift for finite alphabet to a general IFS. In [26] it is also proved that the deviation function presented in [3] fails without the hypothesis of uniqueness of maximizing measure and another characterization of I is presented. The characterization of the deviation function for IFS in the present work is similar to one which appears in [26].

In order to study LDP for Gibbs probabilities in a uniformly contractible IFS we will apply recent results from [28]. We will consider the function $q = \lim_{\beta \to \infty} q^{\beta}$, when it exists. In the non-place dependent case $(A(j, x) = A_j, \forall x \in X)$ is easy to show that q exists. In this case, it is proved in [28] that there exists a unique invariant idempotent probability for the mpIFS (X, ϕ, q) and it is presented its density λ . In this case we will prove that $(\rho_{\beta A})$ satisfy a LDP with deviation function $I = -\lambda$. In the place dependent case, the situation is more complicated. As far as we know it is not proved a LDP or the existence of the limit q, even in the full shift for finite alphabet. In the present work we prove that, if $(\rho_{\beta A})$ satisfy a LDP and the limit q there exists, then $I = -\lambda$ where λ is the density of an invariant idempotent probability for the mpIFS (X, ϕ, q) . As such idempotent probabilities were recently characterized through the Mañé potential and Aubry set in [28], we will obtain a characterization for I using the Mañé potential and Aubry set too.

We would like to point out that despite the fact that the theory of IFS with compact space of maps is fairly old (see [19], [25], among others), only few works are known for such more general setting. Furthermore, as far as we know, the LDP theory for IFS, at zero temperature, is absent in the literature. Although, the thermodynamic formalism and multifractal analysis of Gibbs probabilities has shown notable advances in the last few years with many generalizations (for instance, [14], [15], [29] and [7], among others) thus it seems to be of paramount importance to find further connections and developments in this areas.

We propose, initially, in Section 2, to present a discussion concerning thermodynamic formalism for IFS, following mainly [27]. In the sequence, we extend to the setting of IFS some well known results, in thermodynamic formalism, for the zero temperature case. Right away, in Section 3 we present the main definitions and results contained in [28], concerning invariant idempotent probabilities for max-plus IFS. After the introduction of these elements we then present their connection by considering large deviation principles in Section 4.

2 Thermodynamic Formalism for IFS

In this section we describe the setting presented mainly in [27], concerning the thermodynamic formalism for IFS. The characterization of the entropy and its relation with the Kullback-Leibler divergence is presented in [21].

For each $\beta > 0$ and a Lipschitz continuous function $A: J \times X \to \mathbb{R}$ we consider the operator

$$\mathcal{L}_{\phi,e^{\beta A},\nu}(f)(x) = \int_{J} e^{\beta A(j,x)} \cdot f(\phi_j(x)) \, d\nu(j),$$

where ν is a fixed probability on J with $\operatorname{supp}(\nu) = J$.

Proposition 2.1. Suppose that $A : J \times X \to \mathbb{R}$ has Lipschitz constant Lip(A). Then $\mathcal{L}_{\phi,e^{\beta A},\nu}$ admits a unique pair $(\lambda_{\beta A}, h_{\beta A})$ where $\lambda_{\beta A}$ is a positive number and $h_{\beta A} : X \to \mathbb{R}$ is a positive and continuous function, such that $\sup_{x \in X} h_{\beta A}(x) = 1$ and $\mathcal{L}_{\phi,e^{\beta A},\nu}(h_{\beta A}) = \lambda_{\beta A} \cdot h_{\beta A}$. Furthermore, $h_{\beta A}$ is Lipschitz continuous and Lip $(h_{\beta A}) \leq e^{\beta \cdot \frac{\text{Lip}(A)}{1-\gamma}}$, where γ satisfies (1).

The proof of uniqueness can be found in [27], but details concerning the Lipschitz constant of $h_{\beta A}$ are missing. Let us then present such details following ideas of [6] and [4].

Proof. First, note that we can drop β in the computations. For each 0 < s < 1, we define the operator $T_s : C(X, \mathbb{R}) \to C(X, \mathbb{R})$ by

$$T_s(u)(x) = \log \int_J e^{A(j,x) + su(\phi_j(x))} d\nu(j).$$

We have $|T_s(u) - T_s(v)|_{\infty} \leq s|u - v|_{\infty}$, meaning that, T is a uniform contraction map on $C(X, \mathbb{R})$ with respect to the supremum norm $(|g|_{\infty} = \sup_{x \in X} |g(x)|)$. Let $u_s \in C(X, \mathbb{R})$ be the unique fixed point for T_s .

We claim that for any $s \in (0,1)$, the function u_s is Lipschitz continuous, with $\operatorname{Lip}(u_s) \leq \frac{\operatorname{Lip}(A)}{1-\gamma}$. Indeed, for any $x_1, x_2 \in X$ we have

$$u_s(x_1) - u_s(x_2) \le \max_j \{A(j, x_1) - A(j, x_2) + s[u_s(\phi_j(x_1)) - u_s(\phi_j(x_2))]\}.$$

By iterating this inequality we get

$$u_{s}(x_{1}) - u_{s}(x_{2}) \leq \max_{(j_{0}, j_{1}, \dots)} \left(\sum_{n=0}^{\infty} s^{n} [A(j_{n}, \phi_{j_{n-1}} \circ \dots \circ \phi_{j_{0}}(x_{1})) - A(j_{n}, \phi_{j_{n-1}} \circ \dots \circ \phi_{j_{0}}(x_{2}))] \right)$$
$$\leq \sum_{n=0}^{\infty} s^{n} Lip(A) \gamma^{n} d(x_{1}, x_{2}) = \frac{\operatorname{Lip}(A)}{1 - s\gamma} d(x_{1}, x_{2}) \leq \frac{\operatorname{Lip}(A)}{1 - \gamma} d(x_{1}, x_{2}).$$

Particularly $-\frac{\operatorname{Lip}(A)}{1-\gamma}\operatorname{diam}(X) \leq u_s - \max u_s \leq 0$ and then the family $\{u_s^* =$ $u_s - \max u_s \}_{0 \le s \le 1}$ is equicontinuous and uniformly bounded. Note also that, as $T_s(u_s) = u_s$, we obtain

$$-|A| + s \min u_s \le u_s(x) \le |A| + s \max u_s, \, \forall x \in X.$$

Consequently, $-|A| \leq (1-s) \min u_s \leq (1-s) \max u_s \leq |A|$, for any 0 < s < 1. Then, there exists a subsequence $s_n \to 1$ and a number $k \in \mathbb{R}$ such that $[(1 - k) + k] \in \mathbb{R}$ $(s_n) \max u_{s_n}] \rightarrow k$. By applying Arzela-Ascoli theorem we get a subsequence, also denoted by s_n , such that $\{u_{s_n}^*\}_{n\geq 1}$ converges uniformly to a function $u \in C(X, \mathbb{R})$. As $\operatorname{Lip}(u_s^*) = \operatorname{Lip}(u_s) \leq \frac{\operatorname{Lip}(A)}{1-\gamma}$ we get $\operatorname{Lip}(u) \leq \frac{\operatorname{Lip}(A)}{1-\gamma}$. In order to finish the proof, we will show that $h = e^u$ is an eigenfunction for $\mathcal{L}_{\phi,e^A,\nu}$ with associated eigenvalue e^k . Indeed, for any $s \in (0, 1)$ we have

$$e^{u_s^*(x)} = e^{u_s(x) - \max u_s} = e^{-(1-s)\max u_s + u_s(x) - s\max u_s}$$

= $e^{-(1-s)\max u_s} \int_J e^{A(j,x) + su_s(\phi_j(x)) - s\max u_s} d\nu(j).$

Taking the limit when $s \to 1$ we conclude that u satisfies

$$e^{u(x)} = e^{-k} \int_{J} e^{A(j,x) + u(\phi_j(x))} d\nu(j) = e^{-k} \mathcal{L}_{\phi,e^A,\nu}(e^u)(x).$$

Consider the function $q^{\beta}: J \times X \to \mathbb{R}$ which is defined by $q^{\beta}(j, x) := \frac{e^{\beta A(j,x)} \cdot h_{\beta A}(\phi_j(x))}{\lambda_{\beta A} \cdot h_{\beta A}(x)}$ and for each $j \in J$ consider also the function $q_j^{\beta} : X \to \mathbb{R}$, defined by $q_j^{\beta}(x) = q^{\beta}(j, x)$.

Replacing $e^{\beta A}$ by the function q^{β} we get the operator $\mathcal{L}_{\phi,q^{\beta},\nu}$ which satisfies

$$\mathcal{L}_{\phi,q^{\beta},\nu}(1) = 1,$$

that is, it has the eigenvalue $\lambda = 1$ associated to the eigenfunction h = 1. We say that this operator $\mathcal{L}_{\phi,q^{\beta},\nu}$ is normalized.

The dual of the normalized operator $\mathcal{L}_{\phi,q^{\beta},\nu}$, which we will denote by $\mathcal{M}_{\phi,q^{\beta},\nu}$, acting in Borel probabilities on X is given by

$$\mathcal{M}_{\phi,q^{\beta},\nu}(\mu)(f) = \int_{X} \mathcal{L}_{\phi,q^{\beta},\nu}(f)(x) \, d\mu(x) = \int_{X} \int_{J} q^{\beta}(j,x) \cdot f(\phi_{j}(x)) \, d\nu(j) \, d\mu(x),$$

for any $f \in C(X, \mathbb{R})$. There exists a unique probability $\rho_{\beta A}$ (Gibbs probability) which is invariant for $\mathcal{M}_{\phi,q^{\beta},\nu}$, that is, it satisfies (2).

We say that a probability π on $J \times X$ is holonomic if

$$\int g(x) d\pi(j, x) = \int g(\phi_j(x)) d\pi(j, x), \, \forall g \in C(X, \mathbb{R})$$

Let us denote by \mathcal{H} the set of holonomic probabilities on $J \times X$.

We have a natural holonomic probability $\pi_{\beta A} \in \mathcal{H}$ associated to βA , defined by

$$\int g(j,x) \, d\pi_{\beta A}(j,x) := \int_X \int_J q^\beta(j,x) \cdot g(j,x) \, d\nu(j) \, d\rho_{\beta A}(x), \ \forall g \in C(J \times X, \mathbb{R}).$$

This probability $\pi_{\beta A}$ is an equilibrium probability to βA in the following sense:

$$P_{\nu}(\beta A) := \sup_{\pi \in \mathcal{H}} \int \beta A \, d\pi + H_{\nu}(\pi) = \int \beta A \, d\pi_{\beta A} + H_{\nu}(\pi_{\beta A}), \tag{3}$$

where $H_{\nu}(\pi)$ is the relative entropy of π with respect to ν . More precisely, if the x-marginal of π is a probability ρ on X then

$$H_{\nu}(\pi) = \begin{cases} -\int \log(\operatorname{Jac}) d\pi & \text{if } d\pi = \operatorname{Jac}(j, x) d\nu(j) d\rho(x) \\ \\ -\infty & \text{if } \pi \text{ is not absolutely continuous} \\ & \text{with respect to } \nu \times \rho \end{cases}$$

The above defined entropy satisfies $H_{\nu}(\pi) = -D_{KL}(\pi | \nu \times \rho)$, where D_{KL} is the Kullback-Leibler divergence (see [21] for additional details).

We say that a measurable function Jac : $J \times X \to [0, +\infty)$ is a ν -Jacobian, if $\int_J \operatorname{Jac}(j, x) d\nu(j) = 1, \forall x \in X$. The functions 1 and q^{β} are ν -Jacobians. In [21] it is proved that

$$H_{\nu}(\pi) = -\sup\left\{\int \log(\operatorname{Jac}) \, d\pi \,|\, \operatorname{Jac} \text{ is a Lipschitz } \nu \operatorname{-Jacobian}\right\}.$$

Consequently

$$H_{\nu}(\pi) \le \int \log(1) \, d\pi = 0.$$

Furthermore,

$$H_{\nu}(\pi_{\beta A}) = -\int \log(q^{\beta}) \, d\pi_{\beta A}$$

This definition of entropy does not depend of the IFS, while the operator \mathcal{L} depends of ϕ , as can be observed from its definition. An interesting fact is that $P_{\nu}(\beta A) = \log(\lambda_{\beta A})$ (see [27]).

2.1 Zero temperature limits and ergodic optimization

This section follows ideas present in [22] and [2] which were adapted to the present setting. It extends to IFS some results concerning zero temperature limits. The zero temperature case corresponds to an analysis of the limits, as $\beta \to +\infty$, of the concepts introduced in above subsection (for instance: $\rho_{\beta A}$, $h_{\beta A}$, $\pi_{\beta A}$, $P_{\nu}(\beta A)$).

The following result generalizes one from [22] and will be useful in the present work.

Lemma 2.2. Let Y, Z be compact metric spaces. Let $W_{\beta} : Y \times Z \to \mathbb{R}$ be a family of measurable functions converging uniformly to a continuous function $W : Y \times Z \to \mathbb{R}$, as $\beta \to +\infty$, and let μ be a finite measure on Y with $\operatorname{supp}(\mu) = Y$. Then

$$\frac{1}{\beta} \log \int_{Y} e^{\beta W_{\beta}(y,z)} d\mu(y) \to \sup_{y \in Y} W(y,z)$$

uniformly on Z, as $\beta \to +\infty$. The same is true if we replace β by a sequence β_i which converges to $+\infty$.

Proof. As W is continuous and Y is compact we have that $\tilde{W}(z) := \sup_{y \in Y} W(y, z)$ defines a continuous function too. Fix an $\epsilon > 0$. As W, \tilde{W} are also uniformly continuous, there exists $\delta > 0$ such that

$$d(y_1, y_2) + d(z_1, z_2) < \delta \Rightarrow |W(y_1, z_1) - W(y_2, z_2)| < \epsilon$$

and

$$d(z_1, z_2) < \delta \Rightarrow |\tilde{W}(z_1) - \tilde{W}(z_2)| < \epsilon.$$

As $W_{\beta} \to W$ uniformly we can suppose also there exists $\beta_0 > 0$ such that

$$d(y_1, y_2) + d(z_1, z_2) < \delta \Rightarrow |W_\beta(y_1, z_1) - W_\beta(y_2, z_2)| < 2\epsilon, \text{ for } \beta > \beta_0.$$

Let $Z_0 \subseteq Z$ be a finite set such that $\bigcup_{z \in Z_0} B(z, \delta) = Z$. For each $z \in Z$, let $y_z \in Y$ be such that $W(y_z, z) > \tilde{W}(z) - \epsilon$. Then, for any $z \in Z$ and $y \in B(y_z, \delta)$, we have $W(y, z) > \tilde{W}(z) - 2\epsilon$. As $W_\beta \to W$ uniformly, there exists $\beta_1 > \beta_0$ such that $W_\beta(y, z) > \tilde{W}(z) - 3\epsilon$, for any $z \in Z$, $y \in B(y_z, \delta)$ and $\beta > \beta_1$.

As Z_0 is a finite set and $\operatorname{supp}(\mu) = Y$, there exists $\beta_2 > \beta_1$ such that

$$\frac{1}{\beta} \log \left(\mu \left(B(y_{z_0}, \delta) \right) \right) > -\epsilon, \, \forall z_0 \in Z_0, \, \beta > \beta_2.$$

Given any $z \in Z$, there exists $z_0 \in Z_0$ satisfying $d(z, z_0) < \delta$. If $\beta > \beta_2$ we have that

$$\frac{1}{\beta}\log\int_{Y}e^{\beta W_{\beta}(y,z)}d\mu(y) > \frac{1}{\beta}\log\int_{Y}e^{\beta(W_{\beta}(y,z_{0})-2\epsilon)}d\mu(y) \ge \frac{1}{\beta}\log\int_{B(y_{z_{0}},\delta)}e^{\beta(W_{\beta}(y,z_{0})-2\epsilon)}d\mu(y)$$
$$> \frac{1}{\beta}\log\left(\mu\left(B(y_{z_{0}},\delta)\right)e^{\beta(\tilde{W}(z_{0})-5\epsilon)}\right) > \tilde{W}(z_{0}) - 6\epsilon > \tilde{W}(z) - 7\epsilon.$$

On the other hand, there exists $\beta_4 > \beta_3$ such that, $W_{\beta}(y, z) < \tilde{W}(z) + \epsilon$, for any $\beta > \beta_4$ and $y \in Y$. Then, for $\beta > \beta_4$ we have

$$\frac{1}{\beta} \log \int_{Y} e^{\beta W_{\beta}(y,z)} d\mu(y) < \tilde{W}(z) + \epsilon.$$

Remember that $\log(\lambda_{\beta A}) = P_{\nu}(\beta A)$.

Definition 2.3. Let us define the number $m(A) := \sup_{\pi \in \mathcal{H}} \int A \, d\pi$.

A continuous function $V: X \to \mathbb{R}$ is a calibrated subaction of A if

$$\sup_{j \in J} [A(j, x) + V(\phi_j(x)) - V(x) - m(A)] = 0, \, \forall x \in X$$

A holonomic probability $\pi \in \mathcal{H}$ is maximizing with respect to A if $\int A d\pi = m(A)$.

Proposition 2.4. Consider a Lipschitz continuous function $A : J \times X \to \mathbb{R}$. Under the above framework, we have:

1.
$$\lim_{\beta \to +\infty} \frac{1}{\beta} \log(\lambda_{\beta A}) = \lim_{\beta \to +\infty} \frac{P_{\nu}(\beta A)}{\beta} = m(A).$$

- 2. Suppose that $\frac{1}{\beta_n} \log(h_{\beta_n A}) \to V$ uniformly, as $\beta_n \to +\infty$. Then V is a Lipschitz calibrated subaction of A.
- 3. Suppose that $\pi_{\beta_n A} \to \pi_{\infty}$ in the weak^{*} topology, as $\beta_n \to +\infty$. Then the probability π_{∞} is holonomic and maximizing to A.

Proof. By definition of $h_{\beta A}$ and $\lambda_{\beta A}$ we have

$$\lambda_{\beta A} = \int_{J} e^{\beta A(j,x) + \log(h_{\beta A}(\phi_j(x))) - \log(h_{\beta A}(x))} \, d\nu(j), \ \forall x \in X.$$
(4)

If $h_{\beta A}$ attains its minimum and maximum in x_1 and x_2 , respectively, we get:

$$\lambda_{\beta A} = \int_{J} e^{\beta A(j,x_2) + \log(h_{\beta A}(\phi_j(x_2))) - \log(h_{\beta A}(x_2))} \, d\nu(j) \le \int_{J} e^{\beta A(j,x_2)} \, d\nu(j) \le e^{\beta \cdot \max_{j,x} A(j,x)}$$

and similarly

$$\lambda_{\beta A} \ge \int_J e^{\beta A(j,x_1)} \, d\nu(j) \ge e^{\beta \cdot \min_{j,x} A(j,x)}.$$

Then

$$\min_{j,x} A(j,x) \le \frac{1}{\beta} \log(\lambda_{\beta A}) \le \max_{j,x} A(j,x).$$

Let k be an accumulation point of $\frac{1}{\beta}\log(\lambda_{\beta A})$ as $\beta \to +\infty$ and suppose that $\frac{1}{\beta_n}\log(\lambda_{\beta_n A}) \to k$. By Proposition 2.1 the family $\frac{1}{\beta}\log(h_{\beta A})$ is equicontinuous and uniformly bounded. By applying Arzela-Ascoli theorem, we get a subsequence of (β_n) , also denoted by (β_n) , and a Lipschitz continuous function V such that $\frac{1}{\beta_n}\log(h_{\beta_n A}) \to V$ uniformly. By taking $\lim_{\beta_n \to +\infty} \frac{1}{\beta_n}\log$ in both sides of (4) and applying Lemma 2.2 we obtain

$$k = \sup_{j} [A(j, x) + V(\phi_j(x)) - V(x)], \ \forall x \in X.$$

We claim that k = m(A). Indeed, by one hand, as $\pi_{\beta A}$ is holonomic, we have

$$\frac{1}{\beta}\log(\lambda_{\beta A}) = \frac{P_{\nu}(\beta A)}{\beta} = \frac{\int \beta A \, d\pi_{\beta A} + H_{\nu}(\pi_{\beta A})}{\beta} \stackrel{H_{\nu} \le 0}{\le} \frac{\int \beta A \, d\pi_{\beta A}}{\beta} = \int A \, d\pi_{\beta A} \le m(A)$$

Therefore $k \leq m(A)$. On the other hand, if π is any holonomic probability, as $k \geq A(j, x) + V(\phi_j(x)) - V(x)$ we get $k \geq \int A \, d\pi$. Therefore, $k \geq m(A)$.

We conclude that k = m(A) for any possible convergent sequence and then

$$\lim_{\beta \to +\infty} \frac{1}{\beta} \log(\lambda_{\beta A}) = m(A)$$

Clearly, we also get

$$m(A) = \sup_{j} [A(j, x) + V(\phi_j(x)) - V(x)], \ \forall x \in X,$$

(and then V is a calibrated subaction) for any uniform limit of $\frac{1}{\beta} \log(h_{\beta A})$, as $\beta \to \infty$.

Finally, suppose that $\pi_{\beta_n A} \to \pi_{\infty}$ in the weak^{*} topology, as $\beta_n \to +\infty$. Then π_{∞} is holonomic and $\int A \, d\pi_{\infty} \leq m(A)$. On the other hand,

$$m(A) = \lim_{\beta_n \to +\infty} \frac{P_{\nu}(\beta_n A)}{\beta} \stackrel{H_{\nu} \le 0}{\le} \lim_{\beta_n \to +\infty} \int A \, d\pi_{\beta_n A} = \int A \, d\pi_{\infty}.$$

Given $x \in X$, $n \in \mathbb{N}$ and a finite sequence $\omega = (j_1, j_2, ..., j_n) \in J^n$ we denote

$$\phi_{\omega}(x) = \phi_{(j_1,\dots,j_n)}(x) := \phi_{j_1} \circ \dots \circ \phi_{j_n}(x)$$

and

$$Sum(A, \omega, x) := A(j_1, \phi_{(j_2, \dots, j_n)}(x)) + A(j_2, \phi_{(j_3, \dots, j_n)}(x)) + \dots + A(j_n, x) - n \cdot m(A).$$

Given $x, y \in X$ and $\varepsilon > 0$ we define

$$S_{A,\varepsilon}(x,y) = \sup_{n \in \mathbb{N}} \left[\sup_{\omega \in J^n \mid d(x,\phi_{\omega}(y)) < \varepsilon} \operatorname{Sum}(A,\omega,y) \right],$$
(5)

which can be $-\infty$ if the set $\{\omega \in J^n | d(x, \phi_\omega(y)) < \varepsilon\}$ is empty for any *n*. Clearly $0 < \epsilon < \epsilon' \Rightarrow S_{q,\varepsilon} \leq S_{q,\varepsilon'}$, so we can define the *Mañé potential* $S_A : X \times X \to \mathbb{R} \cup \{-\infty\}$, by

$$S_A(x,y) = \lim_{\varepsilon \to 0} S_{A,\varepsilon}(x,y).$$
(6)

The $Aubry \ set$ is defined as

$$\Omega_A = \{ x \in X | S_A(x, x) = 0 \}.$$
(7)

Let V be a calibrated subaction to A and define $q(j, x) := A(j, x) + V(\phi_j(x)) - V(x) - m(A)$. Then, by definition of calibrated subaction, we have $\sup_{j \in J} q(j, x) = 0$, $\forall x \in X$. For any holonomic probability π on $J \times X$ we have $\int q \, d\pi = \int A \, d\pi - m(A)$. Then m(q) = 0 and a holonomic probability π is maximizing to A if and only if it is maximizing to q.

Theorem 2.5. Let $A: J \times X \to \mathbb{R}$ be a Lipschitz function and V be any continuous calibrated subaction to A. Let us define the function $q: J \times X \to \mathbb{R}$ by $q(j, x) := A(j, x) + V(\phi_j(x)) - V(x) - m(A)$. Then $S_q(x, y) = S_A(x, y) + V(x) - V(y)$ and $\Omega_q = \Omega_A$, which is a non empty set. Furthermore,

$$V(x) = \sup_{z \in \Omega_A} [S_A(z, x) + V(z)], \ \forall x \in X.$$
(8)

Proof. We have m(q) = 0, then, for $\omega = (j_1, ..., j_n)$,

Sum
$$(q, \omega, y) := q(j_1, \phi_{(j_2, \dots, j_n)}(y)) + q(j_2, \phi_{(j_3, \dots, j_n)}(y)) + \dots + q(j_n, y).$$

As $q(j,x) = A(j,x) + V(\phi_j(x)) - V(x) - m(A)$ we have

$$\operatorname{Sum}(q,\omega,y) = \operatorname{Sum}(A,\omega,y) + V(\phi_{\omega}(y)) - V(y).$$

Then

$$S_q(x,y) = \lim_{\varepsilon \to 0} \sup_{n \in \mathbb{N}} \left[\sup_{\omega \in J^n \mid d(x,\phi_\omega(y)) < \varepsilon} (\operatorname{Sum}(A,\omega,y) + V(\phi_\omega(y)) - V(y)) \right]$$
$$= S_A(x,y) + V(x) - V(y).$$

The Aubry sets satisfy $\Omega_A = \{x \in X \mid S_A(x, x) = 0\}$ and $\Omega_q = \{x \in X \mid S_q(x, x) = 0\}$. As $S_A(x, x) = S_q(x, x)$ we obtain $\Omega_A = \Omega_q$.

Let us denote also $q_j(x) = q(j, x)$. We still need to prove that $\Omega_A \neq \emptyset$ and $\sup_{z \in \Omega_A} S_q(z, x) = 0, \forall x \in X$. As V is a calibrated subaction we have $\sup_{j \in J} q(j, x) = 0, \forall x \in X$ and so $S_q \leq 0$. From now on we fix $x \in X$ and prove that there exists $\tilde{x} \in \Omega_A$ such that $S_q(\tilde{x}, x) = 0$.

Let (j_n) be a sequence of points of J satisfying $q_{j_1}(x) = 0$ and $q_{j_{n+1}}(\phi_{j_n} \circ \dots \circ \phi_{j_1}(x)) = 0$ for all $n \ge 1$. Let $x_n := \phi_{(j_n,\dots,j_1)}(x) = \phi_{j_n} \circ \dots \circ \phi_{j_1}(x)$.

As X is compact there exists a subsequence n_i and a point \tilde{x} such that $x_{n_i} \to \tilde{x}$. It follows from the definition of $S_q(\cdot, \cdot)$ that $S_q(\tilde{x}, x) \ge \lim_{n_i \to +\infty} \operatorname{Sum}(q, (j_{n_i}, ..., j_1), x) = 0$ and then $S_q(\tilde{x}, x) = 0$. In what follows we will prove that $\tilde{x} \in \Omega$.

Let $\omega_{k,l} := (j_k, j_{k-1}, ..., j_l)$ for $k \ge l$. By construction of (j_n) we have $q_{j_{n+1}}(x_n) = 0$ and

$$\operatorname{Sum}(q,\omega_{k,n+1},x_n) = 0, \ \forall k > n.$$
(9)

Given $\varepsilon > 0$ there exist $n, m \in \mathbb{N}$ such that n < m and $x_n, x_m \in B(\tilde{x}, \varepsilon/2)$. Observe that $x_m = \phi_{\omega_{m,n+1}}(x_n)$ and as ϕ satisfies (1) we get

$$d(\tilde{x}, \phi_{\omega_{m,n+1}}(\tilde{x})) \leq d(\tilde{x}, x_m) + d(x_m, \phi_{\omega_{m,n+1}}(\tilde{x}))$$
$$= d(\tilde{x}, x_m) + d(\phi_{\omega_{m,n+1}}(x_n), \phi_{\omega_{m,n+1}}(\tilde{x})) < \varepsilon.$$

It follows that

$$S_{q,\varepsilon}(\tilde{x},\tilde{x}) \ge \operatorname{Sum}(q,\omega_{m,n+1},\tilde{x}).$$

As A is Lipschitz continuous and ϕ satisfies (1), there exists C > 0 such that

$$\operatorname{Sum}(q,\omega_{m,n+1},\tilde{x}) = \operatorname{Sum}(A,\omega_{m,n+1},\tilde{x}) + V(\phi_{\omega_{m,n+1}}(\tilde{x})) - V(\tilde{x})$$
$$\geq \operatorname{Sum}(A,\omega_{m,n+1},x_n) - \frac{C}{1-\gamma}\frac{\varepsilon}{2} + V(\phi_{\omega_{m,n+1}}(\tilde{x})) - V(\tilde{x})$$
$$\operatorname{m}(q,\omega_{m,n+1},x_n) - \frac{C}{1-\gamma}\frac{\varepsilon}{2} + [V(\phi_{\omega_{m,n+1}}(\tilde{x})) - V(\phi_{\omega_{m,n+1}}(x_n))] - [V(\tilde{x}) - V(x_n)]$$

 $= \operatorname{Sum}(q, \omega_{m,n+1}, x_n) - \frac{C}{1-\gamma} \frac{c}{2} + [V(\phi_{\omega_{m,n+1}}(\tilde{x})) - V(\phi_{\omega_{m,n+1}}(x_n))] - [V(\tilde{x}) - V(x_n)].$

Therefore, applying also equation (9) and using that V is continuous, we have $S_q(\tilde{x}, \tilde{x}) = \lim_{\varepsilon \to 0} S_{q,\varepsilon}(\tilde{x}, \tilde{x}) \ge 0$. As we always have $S_q \le 0$, we complete the proof.

3 On invariant idempotent probabilities

In this section we recall definitions and results containing in [28]. The theory of idempotent probabilities was introduced by Maslov (also called Maslov measures) to model problems of optimization as integrals with respect to a max-plus structure, see [16], [17], [11] and [20] for classical results on Maslov measures, also [24], [35], [36], [10] and [9] for the existence of a unique invariant idempotent probability for non-place dependent IFSs.

Let us consider $\mathbb{R}_{max} := \mathbb{R} \cup \{-\infty\}$ endowed with the operations

- 1. \oplus : $\mathbb{R}_{\max} \times \mathbb{R}_{\max} \to \mathbb{R}_{\max}$, where $a \oplus b := \max(a, b)$ assuming $a \oplus -\infty := a$.
- 2. $\odot : \mathbb{R}_{\max} \times \mathbb{R}_{\max} \to \mathbb{R}_{\max}$, where $a \odot b := a + b$ assuming $a \odot -\infty := -\infty$.

We consider the following conventions and notations: for any function $f: A \to \mathbb{R}_{\max}$,

1. $\bigoplus_{a \in A} := \sup_{a \in A};$

2.
$$\bigoplus_{a \in A} f(a) = -\infty$$
, if $A = \emptyset$;

- 3. $\bigoplus_{a \in A} f(a) = -\infty$, if $f(a) = -\infty$, $\forall a \in A$.
- 4. $\bigoplus_{a \in A} f(a) = +\infty$, if $A \neq \emptyset$ and $\{f(a) \mid a \in A\}$ is not bounded above.

Definition 3.1. A functional $m : C(X, \mathbb{R}) \to \mathbb{R}$ is an idempotent probability on X if

- 1. $m(a \odot f) = a \odot m(f), \forall a \in \mathbb{R} \text{ and } \forall f \in C(X, \mathbb{R});$
- 2. $m(f \oplus g) = m(f) \oplus m(g), \forall f, g \in C(X, \mathbb{R});$
- 3. m(0) = 0.

We denote by I(X) the set of idempotent probabilities on X.

The idempotent probabilities are closely related with upper semi-continuous functions (u.s.c.). We denote by $U(X, \mathbb{R}_{\max})$ the set of u.s.c. functions $\lambda : X \to \mathbb{R}_{\max}$ satisfying $\{x \in X | \lambda(x) \neq -\infty\} \neq \emptyset$. The next result is well known in the literature (mainly from [16]) and a proof for the present setting can be founded in [28].

Theorem 3.2. A functional $\mu : C(X, \mathbb{R}) \to \mathbb{R}$ is an idempotent probability if and only if there exists $\lambda \in U(X, \mathbb{R}_{\max})$ satisfying

$$\mu(\psi) = \bigoplus_{x \in X} \lambda(x) \odot \psi(x), \quad \forall \, \psi \in C(X, \mathbb{R}) \quad and \quad \bigoplus_{x \in X} \lambda(x) = 0.$$

If $\mu \in I(X)$ there is a unique such function λ in $U(X, \mathbb{R}_{\max})$, which is called its density.

Definition 3.3. Let (X, d_X) and (J, d_J) be compact metric spaces. A max-plus IFS (mpIFS for short) $S = (X, (\phi_j)_{j \in J}, (q_j)_{j \in J})$ is a uniformly contractive IFS $(X, (\phi_j)_{j \in J})$ endowed with a normalized family of weights $(q_j)_{j \in J}$, which is a family

of functions $\{q_j : X \to \mathbb{R} \mid j \in J\}$ satisfying: i. there exists a constant C > 0 such that

$$|q_j(x_1) - q_j(x_2)| \le C \cdot d_X(x_1, x_2), \ \forall j \in J, \ \forall x_1, x_2 \in X;$$
(10)

ii.

$$\bigoplus_{j \in J} q_j(x) = 0, \ \forall x \in X;$$
(11)

iii. the function $q: J \times X \to \mathbb{R}$, defined by $q(j, x) = q_j(x)$, is continuous.

We also use the compact notation $\mathcal{S} = (X, \phi, q)$ to denote a mpIFS (assuming that the set of indices J is known). Considering the setting of above section, if $A: J \times X \to \mathbb{R}$ is Lipschitz continuous and $V: X \to \mathbb{R}$ is a Lipschitz calibrated subaction of A, then defining $q(j, x) = A(j, x) + V(\phi_j(x)) - V(x) - m(A)$ we get a normalized family of weights. This is the case of $q(j, x) = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(q^\beta(j, x))$, where $q^\beta(j, x) = \frac{e^{\beta A(j, x)} \cdot h_{\beta A}(\phi_j(x))}{h_{\beta A}(x) \cdot \lambda_{\beta A}}$, when such limit exists.

Definition 3.4. To each mpIFS $S = (X, \phi, q)$ we assign the following operators: 1. $\mathcal{L}^{0}_{\phi,q} : C(X, \mathbb{R}) \to C(X, \mathbb{R}), \text{ defined by}$

$$\mathcal{L}^{0}_{\phi,q}(f)(x) := \bigoplus_{j \in J} q_j(x) \odot f(\phi_j(x)).$$
(12)

2. $\mathcal{M}^0_{\phi,q}: I(X) \to I(X), \text{ defined by}$

$$\mathcal{M}^{0}_{\phi,q}(\mu) := \bigoplus_{j \in J} \mu(q_j \odot (f \circ \phi_j)).$$
(13)

3. $L^0_{\phi,q}: U(X, \mathbb{R}_{\max}) \to U(X, \mathbb{R}_{\max}), \text{ defined by}$

$$L^0_{\phi,q}(\lambda)(x) := \bigoplus_{(j,y)\in\phi^{-1}(x)} q_j(y) \odot \lambda(y).$$
(14)

The operators $\mathcal{L}^{0}_{\phi,q}$ and $\mathcal{M}^{0}_{\phi,q}$ play the hole of the operators $\mathcal{L}_{\phi,q^{\beta},\nu}$ and $\mathcal{M}_{\phi,q^{\beta},\nu}$, which were introduced in section 2, when considered the zero temperature case and the scale $\frac{1}{\beta}$ log (see also Lemma 2.2).

Next theorem establishes the relation between the three operators in Definition 3.4.

Theorem 3.5. Given a function $\lambda \in U(X, \mathbb{R}_{\max})$ satisfying $\bigoplus_x \lambda(x) = 0$ and the associated idempotent probability $\mu = \bigoplus_{x \in X} \lambda(x) \odot \delta_x \in I(X)$ we have that $\mathcal{M}^0_{\phi,q}(\mu) = \bigoplus_{x \in X} L_{\phi,q}(\lambda)(x) \odot \delta_x$, that is, $\mathcal{M}^0_{\phi,q}(\mu)$ has density $L_{\phi,q}(\lambda)$ where λ is the density of μ . Furthermore

$$\mathcal{M}^0_{\phi,q}(\mu)(f) = \mu(\mathcal{L}^0_{\phi,q}(f)),$$

for any $f \in C(X, \mathbb{R})$, that is, $\mathcal{M}^{0}_{\phi,q}$ is the max-plus dual of $\mathcal{L}^{0}_{\phi,q}$.

Definition 3.6. An idempotent probability $\mu \in I(X)$ with density $\lambda \in U(X, \mathbb{R}_{\max})$ is called invariant (with respect to the mpIFS) if it satisfies any of the following equivalent conditions:

1. $\mathcal{M}^{0}_{\phi,q}(\mu) = \mu;$ 2. $L_{\phi,q}(\lambda) = \lambda;$ 3. $\mu(\mathcal{L}^{0}_{\phi,q}(f)) = \mu(f), \text{ for any } f \in C(X, \mathbb{R}).$

In [28] the idempotent invariant probabilities were characterized using the Mañé potential and the Aubry set. Let us present these results below.

Theorem 3.7. Consider any function $\overline{\lambda} : X \to [-\infty, 0]$ satisfying $\bigoplus_{x \in X} \overline{\lambda}(x) = 0$ and $\overline{\lambda}(x) = 0$

$$\bar{\lambda}(x) = \bigoplus_{z \in \Omega_q} [S_q(x, z) \odot \bar{\lambda}(z)].$$

Then, $\overline{\mu}(f) := \bigoplus_{x \in X} (\overline{\lambda}(x) \odot f(x))$ is an idempotent probability which is invariant for the operator $\mathcal{M}^0_{\phi,q}$. Reciprocally, if $\mu \in I(X)$ is invariant for $\mathcal{M}^0_{\phi,q}$ and $\lambda \in U(X, \mathbb{R}_{\max})$ is its (unique u.s.c.) density, then λ satisfies the equation.

$$\lambda(x) = \bigoplus_{z \in \Omega_q} [S_q(x, z) \odot \lambda(z)].$$
(15)

We remark that we have a symmetric correspondence between the equation which characterizes the calibrated subactions in Theorem 2.5 and the equation which characterizes densities of invariant idempotent probabilities in Theorem 3.7. The main differences are the use of S_A instead S_q and the order of variables.

In [28] it is presented an example proving that a max-plus IFS can have infinitely many possible idempotent invariant probabilities. On the other hand, if $\phi_j(x) = \phi_j$ does not depend of x, the idempotent invariant probability is unique. This will play an important hole in Section 4 and we present this result below.

Given any $\omega = (j_1, j_2, ...) \in J^{\mathbb{N}}$, there exists a unique point, denoted by $\pi(\omega) \in X$ satisfying

$$\pi(\omega) = \lim_{n \to +\infty} \phi_{j_1} \circ \dots \circ \phi_{j_n}(x)$$

for any point x of X.

Theorem 3.8. Given a mpIFS, $S = (X, \phi, q)$, if $q_j(x) = q_j$ does not depend on x, then there exists a unique invariant idempotent probability for $\mathcal{M}^0_{\phi,q}$. Let $\lambda \in U(X, \mathbb{R}_{\max})$ be its density. Then for any $z \in \Omega$ we have

$$\lambda(x) = S_q(x, z) = \bigoplus_{(j_1, j_2, j_3, \dots) \in \pi^{-1}(x)} [q_{j_1} + q_{j_2} + q_{j_3} + \dots].$$

4 Zero temperature limits and Large Deviations

In [1] it is presented a necessary and sufficient condition for a family of probabilities to satisfy a large deviation principle through the notion of idempotent measures introduced by Maslov. In the present section we will prove that the idempotent **Definition 4.1.** Let $(\mu_{\beta})_{\beta>0}$ be a family of probabilities on X. We will say that (μ_{β}) satisfy a large deviation principle (LDP) if there exists a lower semi-continuous (l.s.c.) rate function $I: X \to [0, +\infty]$ such that 1. for each closed set $C \subset X$

$$\limsup_{\beta \to +\infty} \frac{1}{\beta} \log \mu_{\beta}(C) \le -\inf_{\omega \in C} I(\omega)$$

2. for each open set $U \subset X$

$$\liminf_{\beta \to +\infty} \frac{1}{\beta} \log \mu_{\beta}(U) \ge -\inf_{\omega \in U} I(\omega).$$

Remark: As X is open and closed and $\mu_{\beta}(X) = 1$, we obtain $\inf_{w \in X} I(w) = 0$. Furthermore, we assume $\log(0) = -\infty$, $\lim_{\beta \to +\infty} \frac{1}{\beta} \log(0) = -\infty$ and $-\inf_{x \in \emptyset} I(x) = -\infty$. Next two results are very useful in the present work.

Lemma 4.2. [Varadhan's Lemma] Let $(\mu_{\beta})_{\beta>0}$ be a family of probabilities on the compact metric space X which satisfies a LDP with rate function $I: X \to [0, +\infty]$. Then, for any function $f \in C(X, \mathbb{R})$ we have

$$\lim_{\beta \to \infty} \frac{1}{\beta} \log(\rho_{\beta A}(e^{\beta f})) = \sup_{x \in X} [f(x) - I(x)].$$

Proof. See Theorem 4.3.1 in [12].

Lemma 4.3. [Bric's Inverse Varadhan's Lemma] Let $(\mu_{\beta})_{\beta>0}$ be a family of probabilities on the compact metric space X. Suppose that for any function $f \in C(X, \mathbb{R})$ there exists the limit

$$\Gamma(f) := \lim_{\beta \to \infty} \frac{1}{\beta} \log(\rho_{\beta A}(e^{\beta f})).$$

Then (μ_{β}) satisfies a LDP. Furthermore, denoting by $I: X \to [0, +\infty]$ its deviation function, we have

$$\Gamma(f) = \sup_{x \in X} [f(x) - I(x)], \, \forall f \in C(X, \mathbb{R}).$$

Proof. See Theorem 4.4.2 in [12].

4.1 Irreducible Aubry set and a large deviation principle

In this subsection we investigate the uniqueness of idempotent invariant probabilities through the Aubry set and proves a LDP.

Definition 4.4. We will say that Ω_A is irreducible if $S_A(x, y) = 0$ for any $x, y \in \Omega_A$.

As an example, in [28] it is proved that Ω_q is irreducible if $q_j(x) = q_j$ does not depend of x.

Corollary 4.5. If Ω_A is irreducible then any calibrated subaction is constant on Ω_A and any two calibrated subactions are equal, up to the addition of a constant.

Proof. For any $x \in \Omega_A$ and any calibrated subaction V, applying Theorem 2.5, we have

$$V(x) = \sup_{z \in \Omega_A} S_A(z, x) + V(z) = \sup_{z \in \Omega_A} V(z).$$

This proves that V is constant in Ω_A . Let us call $V_{\Omega} := V(z), \forall z \in \Omega_A$. Then, for any $x \in X$ and any calibrated subaction V, we have

$$V(x) = \left(\sup_{z \in \Omega_A} S_A(z, x)\right) + V_{\Omega}.$$

In this way if V and U are any two calibrated subactions, we have $V(x) - U(x) = V_{\Omega} - U_{\Omega}, \forall x \in X$.

Corollary 4.6. If Ω_A is irreducible then there exists the uniform limit

$$V = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(h_{\beta A})$$

and consequently the uniform limit $q = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(q^{\beta})$, which satisfies $q(j, x) = A(j, x) + V(\phi_j(x)) - V(x) - m(A)$.

Proof. By construction of $h_{\beta A}$ we have $\sup_{x \in X} h_{\beta A}(x) = 1$ and the family $\frac{1}{\beta} \log(h_{\beta A})$ is equicontinuous (see Proposition 2.1). Let V be the unique calibrated subaction satisfying $\sup_{x \in X} V(x) = 0$ (any two of they differ just by a constant). By applying Arzela-Ascoli theorem we get $\frac{1}{\beta} \log(h_{\beta A}) \to V$ uniformly.

Proposition 4.7. It suppose that Ω_A is irreducible. Let $q = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(q^\beta) = A(j,x) + V(\phi_j(x)) - V(x) - m(A)$. There exists a unique invariant idempotent probability for the mpIFS (X, ϕ, q) . Its density λ satisfies $\lambda(x) = S_q(x, z), \forall z \in \Omega_A$.

Proof. From Theorem 2.5 we have $\Omega := \Omega_A = \Omega_q$. From Theorem 3.7, if ρ is an invariant idempotent probability with density λ we have

$$\lambda(x) = \sup_{z \in \Omega} S_q(x, z) + \lambda(z), \, \forall x \in X.$$

If $x \in \Omega$ then we obtain $\lambda(x) = \sup_{z \in \Omega} \lambda(z) =: \lambda_{\Omega}$ is a constant. If $x \in X$ we have then

$$\lambda(x) = \lambda_{\Omega} + \sup_{z \in \Omega} S_q(x, z).$$

As $\sup_{x \in X} \lambda(x) = 0$ and $\sup_{x \in X} \sup_{z \in \Omega} S_q(x, z) = 0$ we get $\lambda_{\Omega} = 0$ and then

$$\lambda(x) = \sup_{z \in \Omega} S_q(x, z)$$

In [28] it is proved that $S_q(x, z) \ge S_q(x, y) + S_q(y, z), \forall x, y, z \in X$. Then, if $z_1 \in \Omega$, we have $\lambda(x) = S_q(x, z_1)$, because

$$\lambda(x) \ge S_q(x, z_1) \ge \sup_{z \in \Omega} [S_q(x, z) + S_q(z, z_1)] = \sup_{z \in \Omega} S_q(x, z) = \lambda(x).$$

This proves that the invariant idempotent probability for the mpIFS (X, ϕ, q) is unique and its density is $\lambda(x) = S_q(x, z_1), \forall z_1 \in \Omega$.

Theorem 4.8. Suppose that Ω_A is irreducible. Let

$$q = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(q^{\beta}) = A(j, x) + V(\phi_j(x)) - V(x) - m(A).$$

Then $(\rho_{\beta A})$ satisfy a LDP with rate function

$$I(x) = -S_q(x, z) = -S_A(x, z) - V(x) + V(z), \, \forall z \in \Omega_A.$$

Furthermore,

$$\rho(f) := \sup_{x \in X} [f(x) - I(x)],$$

defines an idempotent probability with density -I, which is the unique invariant to the mpIFS (X, ϕ, q) . For any continuous function $f \in C(X, \mathbb{R})$ we have

$$\lim_{\beta \to \infty} \frac{1}{\beta} \log(\rho_{\beta A}(e^{\beta f})) = \rho(f).$$

Before we present the proof of Theorem 4.8, we need two lemmas.

Lemma 4.9. Let $g: J \times X \to \mathbb{R}$ be a continuous function and for each $\beta > 0$, let $g_{\beta}: J \times X \to \mathbb{R}$ be a continuous function. Suppose that $g_{\beta} \to g$ uniformly and that $\frac{1}{\beta} \log(\int_X e^{\beta \cdot g(j,x)} d\rho_{\beta A}(x))$ converges to a function $\Gamma^*(j)$ uniformly on J. Then

$$\frac{1}{\beta} \log(\int_X e^{\beta \cdot g_\beta(j,x)} \, d\rho_{\beta A}(x)) \to \Gamma^*(j)$$

uniformly on J. The same is true if we replace β by an increasing sequence β_i which converges to $+\infty$.

Proof. For any $\epsilon > 0$ there exists β_0 such that $g - \epsilon < g_\beta < g + \epsilon$, $\beta > \beta_0$ and

$$\Gamma^*(j) - \epsilon < \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g(j,x)} d\rho_{\beta A}(x) \right) < \Gamma^*(j) + \epsilon, \ \forall \beta > \beta_0.$$

Then we have, for $\beta > \beta_0$,

$$\Gamma^*(j) - 2\epsilon < \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g(j,x)} d\rho_{\beta A}(x) \right) - \epsilon = \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g(j,x) - \epsilon} d\rho_{\beta A}(x) \right)$$
$$\leq \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g_{\beta}(j,x)} d\rho_{\beta A}(x) \right) < \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g(j,x) + \epsilon} d\rho_{\beta A}(x) \right)$$
$$= \frac{1}{\beta} \log \left(\int_X e^{\beta \cdot g(j,x)} d\rho_{\beta A}(x) \right) + \epsilon < \Gamma^*(j) + 2\epsilon.$$

Lemma 4.10. Suppose for any $f: X \to \mathbb{R}$ there exists the limit

$$\Gamma(f) := \lim_{\beta \to +\infty} \frac{1}{\beta} \log \left(\int_X e^{\beta f(x)} d\rho_{\beta A}(x) \right).$$

Let $q: J \times X \to \mathbb{R}$ be a continuous function and consider the function $\Gamma^*: J \to \mathbb{R}$, defined by $\Gamma^*(j) = \Gamma(q_j + f \circ \phi_j)$, where $q_j(x) = q(j, x)$. Then $\frac{1}{\beta} \log(\int_X (e^{\beta(q+f \circ \phi)}) d\rho_{\beta A})$ converges uniformly to Γ^* on J. The same is true if we replace β by an increasing sequence β_i which converges to $+\infty$.

Proof. Fix any $\epsilon > 0$. As q and f are continuous, ϕ satisfies (1) and J is compact, there is $\delta > 0$ such that $d(j,l) < \delta \Rightarrow |q_j(x) - q_l(x) + f(\phi_j(x)) - f(\phi_l(x))| < \epsilon, \forall x \in X$. As J is compact and the balls $B(j,\delta), j \in J$, form an open covering of J we can obtain a finite set $J_0 \subset J$ such that the balls $B(j,\delta), j \in J_0$, form an open covering of J too. As this set J_0 is finite, there is $n_0 \in \mathbb{N}$ such that

$$\Gamma(q_j + f \circ \phi_j) - \epsilon < \frac{1}{\beta} \log(\int_X e^{\beta(q_j + f \circ \phi_j)} d\rho_{\beta A}) < \Gamma(q_j + f \circ \phi_j) + \epsilon, \ \forall j \in J_0, \forall \beta > n_0.$$

For any $l \in J$ there is $j \in J_0$ such that $d(l, j) < \delta$, then, $q_j + f \circ \phi_j - \epsilon < q_l + f \circ \phi_l < q_j + f \circ \phi_j + \epsilon$. From definition of Γ we immediately get $\Gamma(f + a) = \Gamma(f) + a$ for any $a \in \mathbb{R}$ and $[f \leq g \Rightarrow \Gamma(f) \leq \Gamma(g)]$. Then, for $\beta > n_0$,

$$\Gamma(q_{l} + f \circ \phi_{l}) - 3\epsilon \leq \Gamma(q_{j} + f \circ \phi_{j} + \epsilon) - 3\epsilon = \Gamma(q_{j} + f \circ \phi_{j}) - 2\epsilon$$

$$< \frac{1}{\beta} \log \left(\int_{X} e^{\beta(q_{j} + f \circ \phi_{j})} d\rho_{\beta A} \right) - \epsilon = \frac{1}{\beta} \log(\int_{X} e^{\beta(q_{j} + f \circ \phi_{j} - \epsilon)} d\rho_{\beta A})$$

$$\leq \frac{1}{\beta} \log \left(\int_{X} e^{\beta(q_{l} + f \circ \phi_{l})} d\rho_{\beta A} \right) < \frac{1}{\beta} \log \left(\int_{X} e^{\beta(q_{j} + f \circ \phi_{j} + \epsilon)} d\rho_{\beta A} \right)$$

$$< \Gamma(q_{j} + f \circ \phi_{j}) + 2\epsilon = \Gamma(q_{j} + f \circ \phi_{j} - \epsilon) + 3\epsilon \leq \Gamma(q_{l} + f \circ \phi_{l}) + 3\epsilon.$$

Proof of Theorem 4.8. From Proposition 4.7 there exists a unique invariant idempotent probability ρ for the mpIFS (X, ϕ, q) . Fix $\tilde{f} \in C(X, \mathbb{R})$ and suppose by contradiction there exists an increasing sequence of positive numbers (β_i) such that

$$\lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i \tilde{f}})) \neq \rho(\tilde{f}).$$

As $C(X, \mathbb{R})$ is separable, there exists a enumerable set F of continuous functions which is dense in $C(X, \mathbb{R})$ with the uniform norm. By applying a Cantor's diagonal argument we can construct a subsequence (also denoted by (β_i)) such that the limit $\lim_{\beta_i \to +\infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i f}))$ there exists for any $f \in F$. We denote by

$$\Gamma(f) := \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i f}))$$

for any $f \in F$.

Claim 1. The limit $\Gamma(g) := \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i g}))$ there exists for any $g \in C(X, \mathbb{R})$.

Proof of Claim 1. Given $\epsilon > 0$ there exists $f \in F$ such that $f(x) - \epsilon \leq g(x) \leq f(x) + \epsilon$ for any $x \in X$. Therefore

$$\Gamma(f) - \epsilon = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i (f-\epsilon)})) \le \liminf_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i g}))$$
$$\le \limsup_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i g})) \le \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i (f+\epsilon)})) \le \Gamma(f) + \epsilon.$$

Then

$$\limsup_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i g})) - \liminf_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i g})) \le 2\epsilon.$$

Making $\epsilon \to 0$ we get the existence of the limit $\Gamma(g)$. This finish the proof of claim.

Claim 2. The map $\Gamma: C(X, \mathbb{R}) \to \mathbb{R}$, given by $\Gamma(f) = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i f}))$ defines an idempotent probability.

Proof of Claim 2.

- 1. $\Gamma(0) = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(1)) = 0;$
- 2. $\Gamma(\lambda \odot \psi) = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i \lambda} \cdot e^{\beta_i \psi})) = \lambda \odot \Gamma(\psi)$, for all $\lambda \in \mathbb{R}$ and $\psi \in C(X, \mathbb{R})$;
- 3. $\Gamma(\varphi \oplus \psi) = \Gamma(\varphi) \oplus \Gamma(\psi)$, for all $\varphi, \psi \in C(X, \mathbb{R})$. To check this last equation just consider the sequence of inequalities:

$$\Gamma(\varphi \oplus \psi) = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i(\varphi \oplus \psi)})) \ge \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i \varphi})) = \Gamma(\varphi).$$

Similarly we get such inequality for ψ and so

$$\Gamma(\varphi \oplus \psi) \ge \Gamma(\varphi) \oplus \Gamma(\psi).$$

On the other hand

$$\Gamma(\varphi \oplus \psi) = \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i(\varphi \oplus \psi)}))$$
$$\leq \lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\rho_{\beta_i A}(e^{\beta_i \varphi}) + \rho_{\beta_i A}(e^{\beta_i \psi})) = \Gamma(\varphi) \oplus \Gamma(\psi).$$

Claim 3. The idempotent probability Γ is invariant for the operator $\mathcal{M}^{0}_{\phi,q}$. Proof of Claim 3. Replace f by $e^{\beta f}$ in equation (2) and then take $\lim_{\beta_i \to \infty} \frac{1}{\beta_i} \log(\cdot)$ in both sides of this equation. In this way we get

$$\frac{1}{\beta_i} \log\left(\int_X e^{\beta_i f} d\rho_{\beta_i A}(x)\right) = \frac{1}{\beta_i} \log\left(\int_X \int_J q_j^{\beta_i}(x) \cdot \left(e^{\beta_i f \circ \phi_j(x)}\right) d\nu(j) d\rho_{\beta_i A}(x)\right)$$

Applying Fubini's theorem we have

$$\frac{1}{\beta_i} \log \left(\int_X e^{\beta_i f} d\rho_{\beta_i A}(x) \right) = \frac{1}{\beta_i} \log \left(\int_J \int_X q_j^{\beta_i}(x) \cdot (e^{\beta_i f \circ \phi_j(x)}) d\rho_{\beta_i A}(x) d\nu(j) \right)$$
$$= \frac{1}{\beta_i} \log \left(\int_J \int_X e^{\beta_i [\frac{1}{\beta_i} \log \left(q_j^{\beta_i}(x) \cdot (e^{\beta_i f \circ \phi_j(x)}) \right)]} d\rho_{\beta_i A}(x) d\nu(j) \right).$$

As $\frac{1}{\beta_i} \log(q_j^{\beta_i}(x) \cdot (e^{\beta_i f \circ \phi_j(x)})) \to q(j,x) + f \circ \phi_j(x)$ uniformly, applying Lemma 4.9 and Lemma 4.10 we get that $\frac{1}{\beta_i} \log \int_X e^{\beta_i [\frac{1}{\beta_i} \log(q_j^{\beta_i}(x) \cdot (e^{\beta_i f \circ \phi_j(x)}))]} d\rho_{\beta_i A}(x)$ converges to $\Gamma(q_j + f \circ \phi_j)$ uniformly on J. Then, applying Lemma 2.2, we get

$$\frac{1}{\beta_i} \log\left(\int_J \int_X e^{\beta_i [\frac{1}{\beta_i} \log(q_j^{\beta_i}(x) \cdot (e^{\beta_i f \circ \phi_j(x)}))]} d\rho_{\beta_i A}(x) d\nu(j)\right) \to \sup_{j \in J} [\Gamma(q_j + f \circ \phi_j)].$$

Therefore,

$$\Gamma(f) = \sup_{j} [\Gamma(q_j + f \circ \phi_j)] \ \forall f \in C(X, \mathbb{R}).$$

This shows that Γ is invariant for the operator $\mathcal{M}^{0}_{\phi,q}$ acting on idempotent probabilities. This proves Claim 3.

From Proposition 4.7, there exists a unique invariant idempotent probability for $\mathcal{M}^{0}_{\phi,q}$ which we denote by ρ . Then $\Gamma = \rho$ and the initial hypothesis of this proof is false. We conclude that for any continuous function $f, \rho(f) = \lim_{\beta \to \infty} \frac{1}{\beta} \log(\rho_{\beta A}(e^{\beta f})).$ Let us call by λ the density of ρ . Then $\lambda \in U(X, \mathbb{R}_{\max})$ and from Proposition 4.7 it satisfies

$$\lambda(x) = S_q(x, z), \ \forall z \in \Omega_A.$$

We still need to prove that the Gibbs measures $\rho_{\beta A}$ satisfies a LDP with rate function $I(x) = -\lambda(x)$. But this is just a consequence of Theorem 3.2 and Lemma 4.3.

4.2Example: the non-place dependent case

In this subsection we suppose X and J are compact metric spaces. We consider a uniformly contractible IFS $\mathcal{S} = (X, (\phi_j)_{j \in J})$ and also fixed numbers $A_j, j \in J$, such that the function $A: J \to \mathbb{R}$, given by $A(j) = A_j$ is Lipschitz continuous.

For each $\beta > 0$, the operator $\mathcal{L}_{\phi,e^{\beta A},\nu}$ has the eigenfunction $h_{\beta A} = 1$ associated to the eigenvalue $\lambda_{\beta A} = \int_J e^{\beta A_j} d\nu(j)$. Then, we consider the numbers given by $q_j^{\beta} = \frac{e^{\beta A_j}}{\int_J e^{\beta A_i} d\nu(i)}, \ j \in J.$ Observe that $\int_J q_j^{\beta} d\nu(j) = 1.$ We will call $(q_j^{\beta})_{j \in J}$ a **Gibbs** **density** on J for temperature $\frac{1}{\beta}$. There exists a unique Borel probability $\rho_{\beta A}$ on X (Gibbs probability) which is invariant for the dual operator $\mathcal{M}_{\phi,q^{\beta},\nu}$. It satisfies

$$\rho_{\beta A}(f) = \int_{J} q_{j}^{\beta} \cdot \rho_{\beta A}(f \circ \phi_{j}) \, d\nu(j) \,\,\forall f \in C(X, \mathbb{R}).$$
(16)

As $q_j^\beta = \frac{e^{\beta A_j}}{\int_J e^{\beta A_i} d\nu(i)}$ we obtain from Lemma 2.2

$$q_j := \lim_{\beta \to \infty} \frac{1}{\beta} \log(q_j^\beta) = A_j - \max_{i \in J} A_i.$$

Then $q_j \leq 0$ and $\bigoplus_{j \in J} q_j = 0$. As A is also continuous, we conclude that $\mathcal{S} = (X, (\phi_j)_{j \in J}, (q_j)_{j \in J})$ is a max-plus IFS.

From Theorem 3.8 there exists a unique invariant idempotent probability ρ for $\mathcal{M}^{0}_{\phi,q}$. This probability satisfies

$$\rho(f) = \bigoplus_{j \in J} q_j \odot \rho(f \circ \phi_j) \ \forall f \in C(X, \mathbb{R})$$
(17)

and its density $\lambda \in U(X, \mathbb{R}_{\max})$ satisfies

$$\lambda(x) = \bigoplus_{(j_1, j_2, j_3, \dots) \in \pi^{-1}(x)} [q_{j_1} + q_{j_2} + q_{j_3} + \dots].$$

Theorem 4.11. Consider the above framework. Then the family of Gibbs probabilities ($\rho_{\beta A}$) satisfy a LDP and its rate function is given by

$$I(x) = -\bigoplus_{(j_1, j_2, j_3, \dots) \in \pi^{-1}(x)} [q_{j_1} + q_{j_2} + q_{j_3} + \dots].$$

Proof. From Theorem 3.8, if $x, z \in \Omega_A$ we have $S_q(x, z) = S_q(x, x) = 0$. Then Ω_A is irreducible and therefore the proof is consequence of Theorem 4.8.

4.3 The place dependent case

In this section we will prove the following result.

Theorem 4.12. Suppose that the family of invariant probabilities $(\rho_{\beta A})_{\beta>0}$ satisfy a LDP and denote by I its deviation function. Suppose that there exists the uniform limit $V = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(h_{\beta A})$. Then, the uniform limit $q = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(q^{\beta})$ there exists and it satisfies $q(j, x) = A(j, x) + V(\phi_j(x)) - V(x) - m(A)$, which defines a normalized family of weights. Furthermore, -I is the density of an idempotent invariant probability ρ for the mpIFS $S = (X, \phi, q)$, that is

$$-I(x) = \sup_{z \in \Omega} [S_q(x, z) - I(z)], \ \forall x \in X.$$
(18)

Proof. Suppose that the family of invariant probabilities $(\rho_{\beta A})_{\beta>0}$ satisfy a LDP and denote by I its deviation function. Then I is l.s.c and $\inf_{x\in X} I(x) = 0$. By applying Lemma 4.2 we get that, for any $f \in C(X, \mathbb{R})$, there exists the limit $\Gamma(f) := \lim_{\beta \to +\infty} \frac{1}{\beta} \log(\rho_{\beta A}(e^{\beta f}))$ and it satisfies

$$\Gamma(f) = \sup_{x \in X} [f(x) - I(x)].$$

Therefore, Γ defines an idempotent probability with density -I (observe that -I is u.s.c and $\sup_{x \in X} -I(x) = 0$).

Suppose there exists the uniform limit $V = \lim_{\beta \to +\infty} \frac{1}{\beta} \log(h_{\beta A})$. Then V is a calibrated subaction to A. We have, by definition of q^{β} ,

$$\frac{1}{\beta}\log(q^{\beta}(j,x)) = A(j,x) + \frac{1}{\beta}\log(h_{\beta A}(\phi_j(x))) - \frac{1}{\beta}\log(h_{\beta A}(x)) - \frac{1}{\beta}\log(\lambda_{\beta A}).$$

Then $\frac{1}{\beta} \log(q^{\beta}(j, x)) \to q(j, x)$ uniformly, where $q(j, x) = A(j, x) + V(\phi_j(x)) - V(x) - m(A)$. As V is a calibrated subaction, the function q defines a normalized family of weights. Let us consider the mpIFS $\mathcal{S} = (X, \phi, q)$.

Claim: Γ defines an idempotent probability which is invariant for the operator $\mathcal{M}^{0}_{\phi,q}$. *Proof of Claim:* It follows from the same reasoning as Claim 3 in the proof of Theorem 4.8.

Finally, as -I is the density of an invariant idempotent probability for the mpIFS $S = (X, \phi, q)$, we obtain, from Theorem 3.7 that

$$-I(x) = \sup_{z \in \Omega} [S_q(x, z) - I(z)].$$

Is a remarkable fact that equations (18) and (8) represents a kind of duality between calibrated subactions (associated to eigenfunction in zero temperature) and deviation functions (associated to Gibbs probabilities in zero temperature). On the other hand LDP provides a connection between Gibbs probabilities in thermodynamic formalism and invariant idempotent probabilities in mpIFS.

References

- AKIAN, M. Densities of idempotent measures and large deviations. Trans. Am. Math. Soc. 351, 11 (1999), 4515–4543.
- [2] BARAVIERA, A., LEPLAIDEUR, R., AND LOPES, A. Ergodic optimization, zero temperature limits and the max-plus algebra. Paper from the 29th Brazilian mathematics colloquium – 29° Colóquio Brasileiro de Matemática, Rio de Janeiro, Brazil, July 22 – August 2, 2013. Publ. Mat. IMPA. Rio de Janeiro: Instituto Nacional de Matemática Pura e Aplicada (IMPA), 2013.
- [3] BARAVIERA, A., LOPES, A. O., AND THIEULLEN, P. A large deviation principle for the equilibrium states of Hölder potentials: the zero temperature case. *Stoch. Dyn.* 6, 1 (2006), 77–96.

- [4] BARAVIERA, A. T., CIOLETTI, L. M., LOPES, A. O., MOHR, J., AND SOUZA, R. R. On the general one-dimensional XY model: positive and zero temperature, selection and non-selection. *Rev. Math. Phys.* 23, 10 (2011), 1063– 1113.
- [5] BARNSLEY, M., DEMKO, S., ELTON, J., AND GERONIMO, J. Invariant measures for markov processes arising from iterated function systems with placedependent probabilities. Ann. Inst. H. Poincar 'e Probab. Statist 24, 3 (1988), 367–394.
- [6] BOUSCH, T. La condition de walters. Ann. Sci. École Norm. Sup. (4) 34, 2 (2001), 287–311.
- [7] BRASIL, J. E., OLIVEIRA, E. R., AND SOUZA, R. R. Thermodynamic formalism for general iterated function systems with measures. *Qual. Theory Dyn. Syst. 22*, 1 (2023), 26. Id/No 19.
- [8] CIOLETTI, L., AND OLIVEIRA, E. R. Thermodynamic formalism for iterated function systems with weights, 2017.
- [9] DA CUNHA, R. D., OLIVEIRA, E. R., AND STROBIN, F. Existence of invariant idempotent measures by contractivity of idempotent Markov operators. J. Fixed Point Theory Appl. 25, 1 (2023), 11. Id/No 8.
- [10] DA CUNHA, R. D., OLIVEIRA, E. R., AND STROBIN, F. Fuzzy-set approach to invariant idempotent measures. *Fuzzy Sets and Systems* 457 (2023), 46–65. Non-additive measures and integrals (196p.).
- [11] DEL MORAL, P., AND DOISY, M. Maslov idempotent probability calculus. I. *Theory Probab. Appl.* 43, 4 (1998), 562–576 (1998) and teor. veroyatn. primen. 43, no. 4, 735–751.
- [12] DEMBO, A., AND ZEITOUNI, O. Large deviations techniques and applications., 2nd ed., corrected 2nd printing ed., vol. 38 of Stoch. Model. Appl. Probab. Berlin: Springer, 2010.
- [13] FAN, A., AND LAU, K.-S. Iterated function system and Ruelle operator. J. Math. Anal. Appl. 231, 2 (1999), 319–344.
- [14] HANUS, P., MAULDIN, R. D., AND URBAŃSKI, M. Thermodynamic formalism and multifractal analysis of conformal infinite iterated function systems. *Acta Math. Hungar. 96*, 1-2 (2002), 27–98.
- [15] JORDAN, T., AND POLLICOTT, M. Multifractal analysis and the variance of Gibbs measures. Journal of the London Mathematical Society 76, 1 (2007), 57–72.
- [16] KOLOKOL'TSOV, V. N., AND MASLOV, V. P. Idempotent analysis as a tool of control theory and optimal synthesis. I. Funct. Anal. Appl. 23, 1 (1989), 1–11.

- [17] KOLOKOLTSOV, V. N., AND MASLOV, V. P. Idempotent analysis and its applications. Appendix by Pierre Del Moral. Transl. from the Russian by V. E. Nazaikinskij, vol. 401 of Math. Appl., Dordr. Dordrecht: Kluwer Academic Publishers, 1997.
- [18] LAU, K.-S., AND YE, Y.-L. Ruelle operator with nonexpansive IFS. Stud. Math. 148, 2 (2001), 143–169.
- [19] LEWELLEN, G. B. Self-similarity. Rocky Mt. J. Math. 23, 3 (1993), 1023–1040.
- [20] LITVINOV, G. L., AND MASLOV, V. P. Idempotent mathematics: A correspondence principle and its applications to computing. *Russ. Math. Surv.* 51, 6 (1996), 1210–1211.
- [21] LOPES, A. O., AND MENGUE, J. K. On information gain, Kullback-Leibler divergence, entropy production and the involution kernel. *Discrete Contin. Dyn. Syst.* 42, 7 (2022), 3593–3627.
- [22] LOPES, A. O., MENGUE, J. K., MOHR, J., AND SOUZA, R. R. Entropy and variational principle for one-dimensional lattice systems with a general *a priori* probability: positive and zero temperature. *Ergodic Theory Dyn. Syst.* 35, 6 (2015), 1925–1961.
- [23] LOPES, A. O., AND OLIVEIRA, E. R. Entropy and variational principles for holonomic probabilities of IFS. *Discrete Contin. Dyn. Syst.* 23, 3 (2009), 937–955.
- [24] MAZURENKO, N., AND ZARICHNYI, M. Invariant idempotent measures. Carpathian Math. Publ. 10, 1 (2018), 172–178.
- [25] MENDIVIL, F. A generalization of ifs with probabilities to infinitely many maps. The Rocky Mountain journal of mathematics 28, 3 (1998), 1043–1051.
- [26] MENGUE, J. K. Large deviations for equilibrium measures and selection of subaction. Bulletin of the Brazilian Mathematical Society, New Series 49, 1 (2018), 17–42.
- [27] MENGUE, J. K., AND OLIVEIRA, E. R. Duality results for iterated function systems with a general family of branches. *Stoch. Dyn.* 17, 3 (2017), 1750021, 23.
- [28] MENGUE, J. K., AND OLIVEIRA, E. R. Invariant measures for placedependent idempotent iterated function systems. *Journal of Fixed Point Theory* and Applications 26, 2 (2024), 19.
- [29] MIHAILESCU, E. Thermodynamic formalism for invariant measures in iterated function systems with overlaps. Communications in Contemporary Mathematics 24, 06 (2022), 2150041.
- [30] PARRY, W., AND POLLICOTT, M. Zeta functions and the periodic orbit structure of hyperbolic dynamics. Astérisque 187, 187-188 (1990), 268.

- [31] RUELLE, D. Statistical mechanics of a one-dimensional lattice gas. Comm. Math. Phys. 9 (1968), 267–278.
- [32] STENFLO, Ö. Uniqueness of Invariant Measures for Place-Dependent Random Iterations of Functions, vol. 132. Springer, New York, NY, 2002, pp. 13–32.
- [33] YE, Y.-L. Convergence speeds of iterations of ruelle operator with weakly contractive ifs. Journal of Mathematical Analysis and Applications 279, 1 (2003), 151–167.
- [34] YE, Y.-L. Ruelle operator with weakly contractive iterated function systems. Ergodic Theory and Dynamical Systems 33, 4 (2013), 1265–1290.
- [35] ZAITOV, A. A. On a metric of the space of idempotent probability measures. Applied General Topology 21, 1 (2020), 35–51.
- [36] ZARICHNYI, M. M. Spaces and maps of idempotent measures. Izvestiya: Mathematics 74, 3 (2010), 481.