

On Hamiltonian projective billiards on boundaries of products of convex bodies

Alexey Glutsyuk^{*†‡}

May 24, 2024

*To Dmitry Valerievich Treschev on the occasion of his 60-th birthday.
To Sergei Vladimirovich Bolotin on the occasion of his 70-th birthday.*

Abstract

Let $K \subset \mathbb{R}_q^n$, $T \subset \mathbb{R}_p^n$ be two bounded strictly convex bodies (open subsets) with C^6 -smooth boundaries. We consider the product $\overline{K} \times \overline{T} \subset \mathbb{R}_{p,q}^{2n}$ equipped with the standard symplectic form $\omega = \sum_{j=1}^n dq_j \wedge dp_j$. The (K, T) -billiard orbits are continuous curves in the boundary $\partial(K \times T)$ whose intersections with the open dense subset $(K \times \partial T) \cup (\partial K \times T)$ are tangent to the characteristic line field given by kernels of the restrictions of the symplectic form ω to the tangent spaces to the boundary. For every $(q, p) \in K \times \partial T$ the characteristic line in $T_{(q,p)}\mathbb{R}^{2n}$ is directed by the vector $(\vec{n}(p), 0)$, where $\vec{n}(p)$ is the exterior normal to $T_p\partial T$, and similar statement holds for $(q, p) \in \partial K \times T$. The projection of each (K, T) -billiard orbit to K is an orbit of the so-called T -billiard in K . In the case, when T is centrally-symmetric, this is the billiard in \mathbb{R}_q^n equipped with Minkowski Finsler structure "dual to T ", with Finsler reflection law introduced in a joint paper by S.Tabachnikov and E.Gutkin in 2002. Studying (K, T) -billiard orbits is closely related to C.Viterbo's Symplectic Isoperimetric Conjecture and the famous Mahler Conjecture in convex geometry. We study the special case, when the T -billiard reflection law is the projective law introduced by S.Tabachnikov, i.e., given by projective involutions of the projectivized tangent spaces $T_q\mathbb{R}^n$, $q \in \partial K$. We show that this happens, if and only if T is an ellipsoid, or equivalently, if all the T -billiards are simultaneously affine equivalent to Euclidean billiards. As an application, we deduce analogous results for Finsler billiards.

^{*}CNRS, UMR 5669 (UMPA, ENS de Lyon), France. E-mail: aglutsyu@ens-lyon.fr

[†]HSE University, Moscow, Russia

[‡]Kharkevich Institute for Information Transmission Problems (IITP, RAS), Moscow

Contents

1 Finsler and (K, T)-billiards with projective reflection law. Introduction and main results	
1.1 Main results. The T -billiards with projective reflection law	5
1.2 Applications to Finsler billiards	8
2 Proof of Theorem 1.12 in two dimensions	12
3 Case of higher dimensions	16
3.1 Proof for planar sections	16
3.2 Proof of Theorem 1.12 in higher dimensions	24
4 Acknowledgements	25

1 Finsler and (K, T) -billiards with projective reflection law. Introduction and main results

Let $K \subset \mathbb{R}^n_{q_1, \dots, q_n}$ and $T \subset \mathbb{R}^n_{p_1, \dots, p_n}$ be two bounded strictly convex open subsets with C^6 -smooth boundaries. We consider their product $K \times T$ in the standard symplectic space \mathbb{R}^{2n} equipped with the symplectic form $\omega := \sum_{j=1}^n dq_j \wedge dp_j$. The open and dense subset $(K \times \partial T) \cup (\partial K \times T) \subset \partial(K \times T)$ carries the characteristic line field: the field of kernels of the restriction to $\partial(K \times T)$ of the symplectic form ω . It is well-known that the characteristic line field is directed by the vectors $(\vec{n}(p), 0) \in T_{q,p}\mathbb{R}^2$ for $(q, p) \in K \times \partial T$, respectively, $(0, -\vec{n}(q))$ for $(q, p) \in \partial K \times T$, where $\vec{n}(q)$, $\vec{n}(p)$ are the exterior normal vector fields on ∂K and ∂T respectively: see [6, 7] and the next paragraph. The above vectors induce the canonical orientation of the characteristic line field that can be also defined as follows. Let $H(p, q)$ be a non-negative Hamiltonian function on \mathbb{R}^{2n} such that $K \times T = \{H < 1\}$. We consider that H satisfies the following additional condition:

- for every fixed $p \in T$ the restriction to $\overline{K} \times \{p\}$ of the function H has non-zero gradient at the points of the boundary $\partial K \times \{p\}$, and the same holds for K interchanged with T .

The corresponding Hamiltonian vector field is tangent to the boundary $\partial(K \times T)$ and directs the above characteristic line field. Its restriction to $K \times \partial T$ coincides with the field $(\frac{\partial H}{\partial p}, 0)$, which is clearly co-directed with the field $(\vec{n}(p), 0)$. Similarly, its restriction to $\partial K \times T$ coincides with the field $(0, -\frac{\partial H}{\partial q})$, which is co-directed with the field $(0, -\vec{n}(q))$.

Definition 1.1 A continuous oriented curve in $\partial(K \times T)$ whose intersection

with the union $(K \times \partial T) \cup (\partial K \times T)$ is tangent and co-directed with the above characteristic line field is called an *orbit of the (K, T) -billiard*.

It is well-known, see [6, 7], that the dynamics of the above (K, T) -billiard (or equivalently, of the above Hamiltonian flow) is indeed "billiard-like". Namely, the Hamiltonian flow moves each point $(q, p_0) \in K \times \partial T$ so that q moves with a velocity parallel and co-directed with the exterior normal vector $\vec{n}(p_0)$ to ∂T at p_0 . Let L_0 denote the movement line of the point q , oriented by the velocity. As q reaches the boundary ∂K , say, at some point q_1 , the point p starts to move with a velocity parallel and co-directed with the interior normal vector to ∂K at q_1 , and q_1 remains fixed. This latter movement continues until p hits the boundary ∂T , say, at some point p_1 . Afterwards $q = q_1$ starts to move with a velocity parallel and co-directed to the exterior normal to ∂T at p_1 , etc. Let L_1 denote the oriented line of the latter movement of the point q . The line L_0 is directed out of K at q_1 , and L_1 is directed inside K at q_1 . See Figure 1. The *reflection map*

$$R_{K,q_1} : L_0 \mapsto L_1$$

is a diffeomorphism from the space of oriented lines through q_1 directed outside K at q_1 to the space of oriented lines through q_1 directed inside K at q_1 . The family of thus defined reflection maps R_{K,q_1} parametrized by the bouncing point q_1 will be briefly denoted R_K . For every $q_1 \in \partial K$ the map R_{K,q_1} is the restriction to the outward directed lines of a diffeomorphic involution $S^{n-1} \rightarrow S^{n-1}$ of the space of oriented lines through q_1 (identified with the sphere S^{n-1}) that fixes the lines lying in the tangent plane $T_{q_1} \partial K$. The projection $\pi_q : \mathbb{R}_q^n \times \mathbb{R}_p^n \rightarrow \mathbb{R}_q^n$ sends each (K, T) -billiard orbit to an orbit of the so-called T -billiard in K , see [6, 7, 10] and the next definition.

Definition 1.2 Let $K, T \subset \mathbb{R}^n$ be bounded strictly convex bodies. The T -billiard in K is the dynamical system on the space of oriented lines L intersecting K defined as follows. For every L take its last point q of intersection with the boundary ∂K in the sense of orientation. The *image of the line L under the T -billiard map* is its image under the reflection map $R_{K,q}$.

Remark 1.3 In the case, when the body T is centrally-symmetric, the T -billiard coincides with the Minkowski Finsler billiard introduced in a joint paper by E.Gutkin and S.Tabachnikov [10] where the Finsler unit ball is the dual to the body T . See the corresponding material in Subsection 1.2.

The next proposition is a well-known fact, see [6, p. 2011], which follows from definition.




Figure 1: The (K, T) -billiard orbits.

Proposition 1.4 *Let T be a ball. Then the T -billiard is the usual Euclidean billiard. Namely, for every $q_1 \in \partial K$ the reflection R_{K, q_1} is the standard reflection in the billiard in K at q_1 according to the usual reflection law: the angle of incidence is equal to the angle of reflection.*

Studying (K, T) -billiard orbits is closely related to C.Viterbo's Symplectic Isoperimetric Conjecture, see [14] and [6, p. 2006, conjecture 1.2], which states that for every convex body Σ in the standard symplectic space \mathbb{R}^{2n} and for every symplectic capacity the ratio of capacities of the body Σ and the unit ball is no greater than the ratio of their volumes taken in power $\frac{1}{n}$. Namely, it was shown in [11], see also [7], that the Hofer-Zehnder capacity of every convex bounded domain in \mathbb{R}^{2n} is equal to the minimal action of closed orbit of characteristic line field on its boundary. In particular, this holds for closed (K, T) -billiard orbits. As was shown in a joint paper by S.Artstein-Avidan, R.Karasev and Y.Ostrover, see [6, theorem 1.6], the Viterbo's Conjecture implies the famous Mahler Conjecture, which states that the product of every convex body $K \subset \mathbb{R}^n$ with its dual has volume no less than $\frac{4^n}{n!}$.

In [13] Sergei Tabachnikov introduced the projective billiards, which are common generalization of billiards on all the space forms of constant curvature, see the next definition.

The goal of the present paper is to describe those convex bodies T for which the corresponding T -billiard is a projective billiard. Our main result shows that this holds only in the case, when T is an ellipsoid (or more generally, a quadric, if T is not necessarily bounded).

The main results are stated in Subsection 1.1. As an application, in Subsection 1.2 we state and deduce similar results on Finsler billiards.

1.1 Main results. The T -billiards with projective reflection law

Definition 1.5 see [13]. A *projective billiard* is a smooth hypersurface $C \subset \mathbb{R}^n$ equipped with a transversal line field \mathcal{N} . For every $Q \in C$ the *projective billiard reflection involution* at Q acts on the space of oriented lines through Q as the affine involution $\mathbb{R}^n \rightarrow \mathbb{R}^n$ that fixes the points of the tangent line to C at Q , preserves the line $\mathcal{N}(Q)$ and acts on $\mathcal{N}(Q)$ as central symmetry with respect to the point¹ Q . In the case, when C is a strictly convex closed hypersurface, the *projective billiard map* acts on the *phase cylinder*: the space of oriented lines intersecting C . It sends an oriented line to its image under the above reflection involution at its last point of intersection with C in the sense of orientation. See Figure 2.




Figure 2: The projective billiard reflection.

Example 1.6 , see [13]. A usual Euclidean planar billiard is a projective billiard with transversal line field being the normal line field. Each space form of constant curvature, i.e., \mathbb{R}^n , S^n , \mathbb{H}^n , is realized as a hypersurface $\Sigma \subset \mathbb{R}_{x_1, \dots, x_{n+1}}^{n+1}$ equipped with an appropriate quadratic form Ψ , the Riemannian metric on the hypersurface being induced by the quadratic form:

$$\mathbb{R}^n = \{x_{n+1} = 1\}, \quad \Psi := \sum_{j=1}^n x_j^2;$$

¹In other words, two (non-oriented) lines a , b through Q are permuted by reflection at Q , if and only if they lie in one two-dimensional subspace (with origin at Q) and the quadruple of lines $T_Q C$, $\mathcal{N}(Q)$, a , b is harmonic: there exists a projective involution of the projectivization $\mathbb{R}P^1$ of the above subspace that fixes $T_Q C$, $\mathcal{N}(Q)$ and permutes a , b .

$$S^n = \left\{ \sum_{j=1}^{n+1} x_j^2 = 1 \right\}, \quad \Psi = \sum_{j=1}^{n+1} x_j^2;$$

$$\mathbb{H}^n = \left\{ \sum_{j=1}^n x_j^2 - x_{n+1}^2 = -1 \right\}, \quad \Psi = \sum_{j=1}^n x_j^2 - x_{n+1}^2.$$

The tautological projection $\pi : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}\mathbb{P}^n$ sends $\Sigma_+ := \Sigma \cap \{x_{n+1} > 0\}$ to the affine chart $\mathbb{R}^n = \{x_{n+1} = 1\} \subset \mathbb{R}\mathbb{P}^n_{[x_1:\dots:x_{n+1}]}$. Each Riemannian billiard on Σ bounded by a hypersurface $\alpha \subset \Sigma$ is projected to a projective projective billiard on the hypersurface $C = \pi(\alpha)$ with the transversal line field \mathcal{N} being the image of the normal line field to α by the differential $d\pi$. The projection π sends each orbit of the Riemannian billiard on α to an orbit of the corresponding projective billiard.

Definition 1.7 A projective billiard on a hypersurface $C \subset \mathbb{R}^n$ is said to be *affinely Euclidean*, if there exists an affine transformation $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ transforming it into a usual billiard, with usual reflection of lines from the image $F(C)$: this holds if and only if the image of the transversal line field \mathcal{N} on C under the differential dF is the normal line field to $F(C)$.

Proposition 1.8 1) Let $T \subset \mathbb{R}^n$ be an ellipsoid, and let $K \subset \mathbb{R}^n$ be a convex bounded open subset with a smooth boundary. Then the corresponding T -billiard reflection transformation R_K is a reflection defined by a projective billiard structure on ∂K .

2) The latter projective billiard structure is affinely Euclidean.

Proof Let us choose orthogonal coordinates p_1, \dots, p_n in which T is an ellipse $\sum_{j=1}^n \frac{p_j^2}{b_j^2} = 1$. Let us complete p_j by affine coordinates q_1, \dots, q_n on K so that $(q_1, \dots, q_n, p_1, \dots, p_n)$ are symplectic coordinates. The variable change

$$\tilde{p}_j = \frac{p_j}{b_j}, \quad \tilde{q}_j = b_j q_j \tag{1.1}$$

is symplectic and transforms $K \times T$ to the product of the rescaled image \tilde{K} of the set K and the unit ball in the \tilde{p} -coordinates. It conjugates the Hamiltonian flows with Hamiltonian function H in the coordinates (p, q) and (\tilde{p}, \tilde{q}) . In the coordinates (\tilde{q}, \tilde{p}) the points \tilde{q} move along billiard orbits in \tilde{K} , by Proposition 1.4. Therefore, the above rescaling sends the movements of the points $q \in K$ with reflections I_K to billiard orbits in \tilde{K} . Thus, the rescaling in the q -coordinates conjugates the reflection I_K at each point $q \in \partial K$ with the standard billiard reflection $I_{\tilde{K}}$ at \tilde{q} . This implies that the

reflection I_K is a projective billiard reflection: it is defined by the transversal line field \mathcal{N} on ∂K that is the pullback of the normal field on \tilde{K} under the rescaling (1.1) of the q -coordinates. The corollary is proved. \square

Corollary 1.9 *Statement 1) of Proposition 1.8 is valid in the case when T is bounded by an arbitrary (maybe unbounded) quadric.*

Corollary 1.9 follows from Statement 1) of Proposition 1.8 by analytic extension of the projectivity of reflection map $R_{K,q}$ as ∂T varies analytically in a family of complex quadrics.

We prove the next theorem, which is a converse statement.

Theorem 1.10 *Let in a (K, T) -billiard the reflection I_K be a projective billiard reflection. Let in addition the boundary ∂T be C^6 -smooth, closed and have positive definite second fundamental quadratic form everywhere. Then ∂T is an ellipsoid.*

Theorem 1.10 has an equivalent reformulation stated below. To state it, consider an arbitrary convex bounded domain T in \mathbb{R}^n whose boundary is C^6 -smooth and has positive definite second fundamental quadratic form. Let us identify ∂T with the unit sphere S^{n-1} via the Gauss map sending a point $p \in \partial T$ to the exterior normal vector to T at p . For a given class \mathcal{L} of parallel lines consider the involution $\mathcal{R}_{\mathcal{L}} : S^{n-1} \rightarrow S^{n-1}$ acting as follows. Take an arbitrary line $\Lambda \in \mathcal{L}$ intersecting ∂T , let A and B be their two intersection points (confluenting to one point in the case of tangency). Let $\vec{n}(A)$, $\vec{n}(B)$ be respectively the corresponding exterior normal vectors to ∂T . By definition,

$$\mathcal{R}_{\mathcal{L}}(\vec{n}(A)) = \vec{n}(B). \quad (1.2)$$

The map $\mathcal{R}_{\mathcal{L}}$ fixes vectors orthogonal to Λ ; their subset is identified with the unit sphere in $T_q \partial K$. We will also treat $\mathcal{R}_{\mathcal{L}}$ as a map acting on the space of oriented lines passing through the point $q \in \partial K$ where the normal vector $\vec{n}(q)$ to ∂K is parallel to Λ : the vectors $\vec{n}(A)$, $\vec{n}(B)$ are identified with the lines through q that are parallel and co-directed with them. We use the following

Proposition 1.11 *Fix an arbitrary point $q \in \partial K$. Let \mathcal{L} be the class of lines parallel to the normal line to ∂K at q . Then the T -billiard reflection map $R_{K,q}$ coincides with the corresponding map $\mathcal{R}_{\mathcal{L}}$ considered as a map acting on oriented lines passing through q and restricted to the subset of lines oriented "out of K " at q .*

The proposition follows from the discussion at the beginning of the section.

Theorem 1.12 *Let $\alpha \subset \mathbb{R}^n$ be a germ of hypersurface with positive definite second fundamental form. Consider the tautological projection $S^{n-1} \rightarrow \mathbb{RP}^{n-1}$, which is the double covering sending each unit vector to its projectivization: the point representing its ambient line. Let for every class \mathcal{L} of parallel lines (that contains at least one line intersecting α twice) the involution $\mathcal{R}_{\mathcal{L}}$ given by (1.2) (with $\partial T = \alpha$) be the lifting of a projective involution $\mathbb{RP}^{n-1} \rightarrow \mathbb{RP}^{n-1}$. Then α is a quadric.*

Theorem 1.10 follows from Theorem 1.12 and Proposition 1.11.

Remark 1.13 Maxim Arnold and Sergei Tabachnikov proved a result closely related to Theorem 1.12, see [5, theorem 3]: *if a smooth strictly convex closed hypersurface S admits a normal vector field N such that for every $x, y \in S$ one has $\langle N(x), x - y \rangle = -\langle N(y), x - y \rangle$, then S is an ellipsoid (ellipse in two dimensions).*

1.2 Applications to Finsler billiards

Let us recall the following well-known material from Finsler geometry, see, e.g., [1, 2, 3, 4, 8, 10]. A *Finsler manifold* is a manifold M equipped with a *Finsler structure*: a family of smooth strictly convex closed hypersurfaces $I_x \subset T_x M$ centrally-symmetric with respect to zero; they are called *indicatrices*. The *Finsler norm* on $T_x M$ is the norm for which the indicatrix I_x is the unit sphere. For every vector $v \in T_x M$ its Finsler norm is denoted by $L(x, v)$. A *Finsler geodesic* is a curve $\gamma : [t_0, t_1] \rightarrow M$ minimizing the functional

$$|\gamma| := \int_{t_0}^{t_1} L(\gamma(t), \dot{\gamma}(t)) dt$$

among curves with fixed endpoints.

Example 1.14 Recall (see, e.g., [10]) that a Finsler structure on \mathbb{R}^n is called *Minkowski*, if it is translation invariant. A Finsler structure on a domain $U \subset \mathbb{R}^n$ is *projective*, (or *projectively flat*), if all its geodesics are lines. It is well-known that Minkowski Finsler structures are projectively flat, see [10, p. 280], and so are the Riemannian metrics of spaces forms of constant curvature in appropriate chart. Conversely, a simply connected domain in \mathbb{R}^n equipped with a projectively flat Riemannian metric is always isometric (up to rescaling the metric by a constant factor) to a domain in a

space form of constant curvature (Beltrami Theorem, see, e.g., [12, section VI.2]).

E.Gutkin and S.Tabachnikov introduced the following *Finsler reflection law* in their joint paper [10]. Let $S \subset M$ be a hypersurface (for simplicity, strictly convex). Let $x \in S$, and let $a, b \in M$ be points such that there exist a geodesic arc from a to x and a geodesic arc from x to b . Let u, v denote their directing unit tangent vectors at x . We suppose that u and v lie on different sides from the tangent hyperplane $T_x S$. We say that the *Finsler reflection with respect to the hyperplane $T_x S$ sends u to v* , if x is a critical point of the function $g(x) := d(a, x) + d(x, b)$: the sum of lengths of the above geodesic arcs with fixed endpoints a and b . The following results were proved in [10, pp. 281, 282],

1) Thus defined reflection transformation depends only on x and $T_x S$ and is independent on a, b, S .

2) The reflection is a well-defined diffeomorphic involution $I_x \rightarrow I_x$ of the indicatrix that can be defined geometrically in one of the following equivalent ways.

a) Consider the *Legendre transform* $D : I_x \rightarrow T_x^* M \setminus \{0\}$, which sends each vector $v \in I_x$ to the covector $w \in T_x^* M$ for which $w(v) = 1$ and the kernel $\ker w \subset T_x M$ is parallel to the tangent hyperplane $T_v I_x \subset T_x M$. The latter pair of conditions is equivalent to the pair of conditions $w(v) = 1$ and $|w(u)| \leq 1$ for every $u \in I_x$. The hypersurface $J_x := D(I_x) \subset T_x^* M$ is a closed convex centrally-symmetric hypersurface, which is called the *figuratix* or the *dual to the indicatrix I_x* . The equivalent Finsler reflection law states that $u \in I_x$ is reflected to $v \in I_x$ with respect to a codimension one subspace $H = T_x S \subset T_x M$, if the linear functional $D(u) - D(v) \in T_x^* M$ vanishes on H . Recall that the *Legendre transformation is involutive*: the inverse $D^{-1} : J_x \rightarrow I_x$ coincides with the Legendre transformation defined by the hypersurface J_x instead of I_x .

b) The vector $u \in I_x$ is reflected to $v \in I_x$, if the hyperplanes tangent to I_x at u and at v respectively and the reflecting hyperplane H are concurrent: that is, either they are parallel, or they intersect themselves by a codimension two affine subspace in $T_x M$. See Figure 3.

Let us now consider that our Finsler manifold M is a domain in \mathbb{R}^n equipped with some Finsler structure. For every $x \in M$ we identify the cotangent space $T_x^* M$ with the corresponding tangent space $T_x M$ via the standard Euclidean scalar product. Then the figuratrix J_x is identified with a centrally-symmetric strictly convex closed hypersurface in $T_x M$, and the Legendre transform D sends each vector $u \in I_x$ to the vector $D(u) \in T_x M$




Figure 3: Finsler reflection law.

orthogonal to $T_x I_x$ and such that $\langle D(u), u \rangle = 1$. In other terms, the Legendre transform $D : I_x \rightarrow J_x$ is the Gauss map multiplied by a positive function.

Given a strictly convex closed surface $S \subset M$, we can define the *Finsler billiard* on S acting in the space of oriented Finsler geodesics intersecting S . Namely, take an oriented geodesic ℓ intersecting S and its last intersection point q with S in the sense of orientation. Its reflection image is the geodesic ℓ^* issued from q , directed inside the convex domain bounded by S .

The next proposition is well-known and follows immediately from definition, involutivity and the above Gauss map interpretation of the Legendre transform $D : I_x \rightarrow J_x$.

Proposition 1.15 (see, e.g., [6, p. 2011]). *Let $T \subset \mathbb{R}^n$ be a strictly convex bounded centrally symmetric convex body with smooth boundary. Then for every strictly convex bounded body $K \subset \mathbb{R}^n$ with smooth boundary the T -billiard in K (with reflection law given in Definition 1.2) is the Finsler billiard in the Minkowski Finsler space, where the indicatrix I_0 is dual to the boundary ∂T . Namely, $\partial T = J_0 = D(I_0)$ (after translation of the body T so that its center of symmetry becomes the origin).*

Theorem 1.10 together with Proposition 1.15 imply the following corollaries.

Corollary 1.16 *Let in a Minkowski Finsler space \mathbb{R}^n the Finsler reflection law be projective. Then the Finsler unit sphere is an ellipsoid. In particular, there exists an affine transformation $\mathbb{R}^n \rightarrow \mathbb{R}^n$ sending the Minkowski*

Finsler structure to the standard Euclidean metric and thus, sending orbits of each Finsler billiard to orbits of an Euclidean billiard with the standard reflection law.

Proof Let $J = J_0 = D(I_0)$ denote the figuratrix: the closed hypersurface dual to the unit sphere of the Finsler structure. Let T denote the convex body bounded by J . For every bounded strictly convex body $K \subset \mathbb{R}^n$ the (K, T) -billiard is projected to the T -billiard. The T -billiard reflection law is the Minkowski Finsler reflection law (Proposition 1.15). If it is projective, then ∂T is an ellipsoid (Theorem 1.10): $\partial T = E := \{ \langle Ax, x \rangle = 1 \}$, where A is a symmetric positive definite matrix. But the dual to an ellipsoid E is the ellipsoid $E^* := \{ \langle A^{-1}x, x \rangle = 1 \}$. Indeed, for every $x \in E$ the vector Ax is normal to $T_x E$, being half-gradient of the quadratic form $\langle Ax, x \rangle$. Its scalar product with x is equal to $\langle Ax, x \rangle = 1$, since $x \in E$. Hence, $Ax = D(x)$. On the other hand, $\langle A^{-1}(Ax), Ax \rangle = \langle x, Ax \rangle = 1$, since the matrix A is symmetric. Therefore, $Ax \in E^*$. Finally, the hypersurface $D(\partial T)$ is an ellipsoid. It coincides with the indicatrix I_0 , by involutivity of the Legendre transform. Corollary 1.16 is proved. \square

Corollary 1.17 *Let $U \subset \mathbb{R}^n$ be a simply connected domain equipped with a Finsler structure. Let the corresponding Finsler reflection law be projective. Then the Finsler structure is Riemannian, i.e., the corresponding unit spheres are ellipsoids.*

Proof At each point $q \in U$ for every codimension one vector subspace $H \subset T_q \mathbb{R}^n$ the involution defining the Finsler reflection law with respect to H is the same, as in the Minkowski Finsler space \mathbb{R}^n with the indicatrix I_0 being translated image of the indicatrix I_q . Therefore, if the reflection law in $T_q \mathbb{R}^n$ is projective, then this is also the case in the Minkowski space. Hence, the corresponding indicatrix I_0 is an ellipsoid (Corollary 1.16). This proves Corollary 1.17. \square

Corollary 1.18 *Let $U \subset \mathbb{R}^n$ be a simply connected domain equipped with a projectively flat Finsler structure. Let the corresponding Finsler reflection law be projective. Then (up to rescaling the metric by constant factor) U is isometric to a domain in a n -dimensional space form of constant curvature: either Euclidean space \mathbb{R}^n , or the unit sphere S^n , or the hyperbolic space \mathbb{H}^n .*

Proof The Finsler structure is Riemannian (Corollary 1.17). The simply connected Riemannian manifold U is projectively flat, by assumption.

Therefore, it is isometric (up to rescaling the metric by constant factor) to a domain in a space form of constant curvature (Beltrami Theorem, see [12, section VI.2]). Corollary 1.18 is proved. \square

2 Proof of Theorem 1.12 in two dimensions

In this section we consider that $n = 2$; thus, α is a planar curve.

Fix a point $p \in \alpha$. Let $\Gamma = \Gamma_p$ be the osculating conic for the curve α at p . This is the unique conic tangent to α with contact of order greater than 4, that is, having the same 4-jet, as α at p .

Recall that a point p of a curve α is called *sextatic*, if its osculating conic at p has contact of order greater than five with α , see [9, p.12].

The proof of Theorem 1.12 is based on the following well-known theorem.

Theorem 2.1 *If all the points of a connected planar curve α are sextatic, then α is a conic (or a connected component of a hyperbola).*

Proof The sextatic points are exactly those points where the affine curvature of the curve has zero derivative [9, p. 12, formula (3.33)]. Therefore, α has constant affine curvature, and hence, is a conic, see [9, pp. 9-10]. \square

Therefore, for the proof of Theorem 1.12 for $n = 2$ it suffices to prove the following lemma.

Lemma 2.2 *Fix an arbitrary point $p \in \alpha$. Let L_p denote the line tangent to α at p . Let $\mathcal{R}_{\mathcal{L}}$ be the involution (1.2) defined by $\partial T = \alpha$ and the class \mathcal{L} of lines parallel to L_p . Let $\mathcal{R}_{\mathcal{L}}$ be the lifting of a projective involution of \mathbb{RP}^1 . Then p is a sextatic point of the curve α .*

Proof We prove the lemma by contradiction. Suppose the contrary: at some point $p \in \alpha$ the conic $\Gamma = \Gamma_p$ and the curve α are locally graphs of functions that differ at 5-th Taylor terms. For both of them consider the corresponding involutions $\mathcal{R}_{\mathcal{L}}$,

$$f := \mathcal{R}_{\alpha, \mathcal{L}}, \quad g := \mathcal{R}_{\Gamma, \mathcal{L}},$$

defined by the class \mathcal{L} of lines parallel to L_p and by the curves α and Γ respectively. These are germs of one-dimensional C^5 -smooth involutions at their common fixed point O corresponding to the normal vector $\vec{n}(p)$. We prove the following proposition, which is the main part of the proof of the lemma.

Proposition 2.3 *In the above assumptions the involutions f and g have the same 2-jet at O (and even the same 3-jet), but $f \not\equiv g$.*

The involutions f and g are both liftings of projective involutions. This follows from assumptions for the involution f , which is defined by the curve α , and by Corollary 1.9 for the involution g , which is defined by the conic Γ . This together with the next proposition on a general pair of projective involutions will bring us to contradiction with Proposition 2.3.

Proposition 2.4 *Two arbitrary projective involutions $\mathbb{RP}^1 \rightarrow \mathbb{RP}^1$ with a common fixed point either coincide, or have different 2-jets at the fixed point.*

Proof Let f, g denote the involutions in question, and let O denote their common fixed point. The map f has yet another fixed point A . In an affine chart \mathbb{R}_t on $\mathbb{RP}^1 \setminus \{A\}$ centered at O one has

$$f(t) = -t, \quad g(t) = -\frac{t}{1+ct} = -t + ct^2 + O(t^3), \quad \text{as } x \rightarrow 0.$$

One has $c \neq 0$, if and only if $f \neq g$. In this case the germs f and g obviously have different 2-jets. The proposition is proved. \square

Proof of Proposition 2.3. The involutions f and g are defined to permute the exterior unit normal vectors to α and Γ respectively at points of intersection of a line ℓ parallel to L_p and the curve in question. The exterior normal vector $\vec{n}(y)$ to the underlying curve at a point y will be identified with the oriented tangent line L_y to the curve at y . Its orienting vector is obtained from the normal vector $\vec{n}(y)$ by rotation by $\frac{\pi}{2}$. Both $\vec{n}(y)$ and L_y will be identified with the slope of the line L_y : the tangent of its azimuth with respect to the line L_p . Thus, we can and will consider that the involutions f and g act on the slopes of the oriented tangent lines to the underlying curves.

Let us take positively oriented orthogonal affine coordinates (x, y) centered at $O = p$ so that the x -axis coincides with the oriented tangent line $L_p = L_O$. Then the curves α and Γ both lie in the upper half-plane (by choice of orientation). We can and will consider them as graphs of functions:

$$\alpha = \{y = h_\alpha(x)\}, \quad \Gamma = \{y = h_\Gamma(x)\}, \quad h_\alpha(x) \simeq h_\Gamma(x) \simeq \frac{1}{2}x^2, \quad \text{as } x \rightarrow 0.$$

Indeed, the latter asymptotics a priori holds with $\frac{1}{2}x^2$ replaced by νx^2 with $\nu > 0$. One can achieve that $\nu = \frac{1}{2}$ by applying a homothety, which does

not change our projectivity and jet (in)equality assumptions. One has

$$h_\alpha(x) = h_\Gamma(x) + cx^5 + o(x^5), \quad c \neq 0, \quad (2.1)$$

since the 5-jets of the curves α and Γ are supposed to be distinct, while their 4-jets are equal. See Figure 4. For every x the slope of the curve $\beta = \alpha, \Gamma$




Figure 4: The curve α and the conic Γ as graphs of functions near $p = O$ that differ in 5-th Taylor term.

at the point $(x, h_\beta(x))$ is equal to $h'_\beta(x)$, and one has

$$h'_\beta(x) \simeq x, \quad h'_\alpha(x) - h'_\Gamma(x) = 5cx^4 + o(x^4), \quad (2.2)$$

by (2.1). For every x small enough let

$$\zeta = \zeta(x) \text{ denote the point for which } h_\alpha(\zeta) = h_\Gamma(x), \text{ and } \zeta \simeq x, \quad (2.3)$$

$$\hat{x} = \hat{x}(x) \text{ denote the point distinct from } x \text{ where } h_\Gamma(\hat{x}) = h_\Gamma(x), \quad (2.4)$$

$$\tilde{\zeta} = \tilde{\zeta}(x) \text{ denote the point distinct from } \zeta \text{ where } h_\alpha(\tilde{\zeta}) = h_\alpha(\zeta), \quad (2.5)$$

see Figure 4.

Claim 1. *One has*

$$\zeta \simeq x, \quad \hat{x} \simeq -x, \quad \tilde{\zeta} \simeq -\zeta, \quad h'_\Gamma(x) \simeq x, \quad h'_\alpha(x) \simeq x, \quad (2.6)$$

$$\zeta = x - cx^4 + O(x^5), \quad \tilde{\zeta} = \hat{x} - cx^4 + O(x^5), \quad (2.7)$$

$$h'_\alpha(\zeta) = h'_\Gamma(x) + 4cx^4(1 + o(1)), \quad h'_\alpha(\tilde{\zeta}) = h'_\Gamma(\hat{x}) + 4cx^4(1 + o(1)). \quad (2.8)$$

Proof Formulas (2.6) follow from definition, since $h_\beta(x) \simeq \frac{1}{2}x^2$ for $\beta = \alpha, \Gamma$. One has $h_\Gamma(x) = h_\alpha(\zeta) = h_\Gamma(\zeta) + c\zeta^5 + o(\zeta^5)$, $\zeta \simeq x$. Hence,

$$h_\Gamma(\zeta) - h_\Gamma(x) = -c\zeta^5(1 + o(1)).$$

The difference in the latter left-hand side is asymptotic to $h'_\Gamma(x)(\zeta - x) \simeq \zeta(\zeta - x)$. This implies the first asymptotic equality in (2.7). Applying the same argument to \hat{x} and $\tilde{\zeta}$ yields the second equality in (2.7). Let us prove the first equality in (2.8); the proof of the second one is analogous. One has

$$\begin{aligned} h'_\alpha(\zeta) - h'_\Gamma(x) &= (h'_\alpha(\zeta) - h'_\Gamma(\zeta)) + (h'_\Gamma(\zeta) - h'_\Gamma(x)) \\ &= 5c\zeta^4 + o(\zeta^4) + h''_\Gamma(x)(\zeta - x) + O((\zeta - x)^2). \end{aligned}$$

Substituting the asymptotics $\zeta \simeq x$, $h''_\Gamma(x) \simeq 1$, $\zeta - x \simeq -cx^4$, to the latter right-hand side yields (2.8) and finishes the proof of the claim. \square

Recall that the involutions f and g act on the slopes of tangent lines so that

$$g(h'_\Gamma(x)) = h'_\Gamma(\hat{x}), \quad f(h'_\alpha(\zeta)) = h'_\alpha(\tilde{\zeta}), \quad (2.9)$$

$$h'_\Gamma(x) \simeq h'_\alpha(x) \simeq h'_\alpha(\zeta) \simeq x, \quad (2.10)$$

$$h'_\Gamma(\hat{x}) \simeq h'_\alpha(\hat{x}) \simeq h'_\alpha(\tilde{\zeta}) \simeq \hat{x} \simeq -x. \quad (2.11)$$

Claim 2. *One has*

$$f(t) = g(t) + 8ct^4(1 + o(1)). \quad (2.12)$$

Proof Clearly one has $f(t) \simeq g(t) \simeq -t$, $f'(t) \simeq g'(t) \simeq -1$. This together with (2.9)-(2.11) and (2.8) implies that

$$\begin{aligned} f(h'_\Gamma(x)) &= f(h'_\alpha(\zeta)) - f'(h'_\Gamma(x))(h'_\alpha(\zeta) - h'_\Gamma(x)) + O((h'_\alpha(\zeta) - h'_\Gamma(x))^2) \\ &= h'_\alpha(\tilde{\zeta}) + 4cx^4(1 + o(1)) = h'_\Gamma(\hat{x}) + (h'_\alpha(\tilde{\zeta}) - h'_\Gamma(\hat{x})) + 4cx^4(1 + o(1)) \\ &= g(h'_\Gamma(x)) + 8cx^4(1 + o(1)). \end{aligned}$$

This together with the asymptotics $h'_\Gamma(x) \simeq x$ proves (2.12). \square

Claim 2 immediately implies the statements of Proposition 2.3. \square

Thus, f and g are distinct projective involutions that have the same 2-jet at a common fixed point. This contradicts Proposition 2.4. The contradiction thus obtained finishes the proof of Lemma 2.2. \square

Lemma 2.2 together with Theorem 2.1 imply that the curve α is a conic. This proves Theorem 1.12.

3 Case of higher dimensions

Here we prove Theorem 1.12 in higher dimensions. First in Subsection 3.1 we prove that each planar section of the hypersurface α is a conic. Then in Subsection 3.2 we deduce that α is a quadric.

3.1 Proof for planar sections

Here we prove the following theorem.

Theorem 3.1 *Consider a C^6 -smooth germ α of hypersurface in \mathbb{R}^n at a point O with positive definite second fundamental quadratic form. Fix a two-dimensional plane Π intersecting α transversally. Set $\beta = \Pi \cap \alpha$. Let $L \subset \Pi$ be a line tangent to β , and let \mathcal{L} denote the class of lines parallel to L . Consider the corresponding germ of involution $f = \mathcal{R}_{\mathcal{L}} = \mathcal{R}_{\alpha, \mathcal{L}}$ acting on a domain in the unit sphere $S = S^{n-1}$, see (1.2): its germ at its fixed vector $\vec{n}(O) = \vec{n}_{\alpha}(O) = \vec{n}_{\Gamma}(O)$. Let for every line L tangent to β the corresponding mapping f be the lifting to S of a projective transformation. Then the intersection curve β is a conic.*

Corollary 3.2 *In the conditions of Theorem 1.12 each planar section of the hypersurface α is a conic.*

For the proof of Theorem 3.1 it suffices to prove the following lemma.

Lemma 3.3 *In the conditions of Theorem 3.1 let L be a line tangent to β , and let O be their tangency point. Let the map f corresponding to the given line L be the lifting to S of a projective transformation. Then O is a sextatic point of the planar curve $\beta \subset \Pi$.*

Lemma 3.3 together with Theorem 2.1 imply Theorem 3.1.

First let us sketch the proof of Lemma 3.3 given below. Let $\gamma \subset \Pi$ be the osculating conic to β at O . We have to show that it has contact greater than five with β . To do this, in the next proposition we construct an auxiliary quadric Γ containing γ that is tangent to α at O with contact of order at least three so that the normal vector fields of the hypersurfaces Γ and α along the curves γ and β respectively differ by a quantity of at least cubic order. We deal with them as germs of parametrized curves $\vec{n}_{\gamma}(x_1)$, $\vec{n}_{\beta}(x_1)$ in the unit sphere (identified with $\mathbb{R}P^{n-1}$ near $\vec{n}(O) := \vec{n}_{\gamma}(O) = \vec{n}_{\beta}(O)$ via the tautological projection). The involution $g = \mathcal{R}_{\Gamma, \mathcal{L}}$ is the lifting of a projective transformation, as is f . Using the above asymptotics we will

show (Proposition 3.6) that both projective involutions coincide: $f = g$. The projective involution f fixes points of the hyperplane H that is the projectivization of the $(n - 1)$ -dimensional subspace orthogonal to the line L . And it fixes exactly one more point $P \in \mathbb{R}\mathbb{P}^{n-1} \setminus H$. One has $\vec{n}(O) \in H$.

We prove Lemma 3.3 by contradiction. Suppose the contrary: the curves β and γ have contact of order exactly five. Then in orthogonal coordinates (x_1, x_n) on Π centered at O , with the x_1 -axis being the common tangent line to β and γ at O , the curves β and γ are graphs of functions

$$\beta = \{x_n = \nu_\alpha(x_1)\}, \quad \gamma = \{x_n = \nu_\Gamma(x_1)\}, \quad \nu_\alpha(x_1) - \nu_\Gamma(x_1) \simeq cx_1^5, \quad c \neq 0.$$

We identify O with the point in $\mathbb{R}\mathbb{P}^{n-1}$ represented by the vector $\vec{n}_\beta(O)$. We introduce appropriate affine coordinates (z, y) centered at $\vec{n}(O)$ on an affine chart in $\mathbb{R}\mathbb{P}^{n-1}$, $z \in \mathbb{R}$, $y = (y_1, \dots, y_{n-2}) \in \mathbb{R}^{n-2}$,

$$z(\vec{n}_\beta(x_1)) \simeq z(\vec{n}_\gamma(x_1)) \simeq x_1, \quad \text{as } x_1 \rightarrow 0,$$

and show that they satisfy the following statements:

- a) $H = \{z = 0\}$, and the projection $(y, z) \mapsto y$ coincides with the projection $\mathbb{R}\mathbb{P}^{n-1} \setminus \{P\} \rightarrow H$ from the fixed point P ;
- b) the projections of the curves $\vec{n}_\beta(x_1)$ and $\vec{n}_\gamma(x_1)$ to the z -coordinate differ by a quantity $5cx_1^4(1 + o(1))$, as $x_1 \rightarrow 0$; here $c \neq 0$.

The projective involution $f = g$ preserves each z -fiber $\{y\} \times \mathbb{R}\mathbb{P}_z^1$ and acts along it by a map with asymptotics $z \mapsto -z + O(z^2)$, as $z \rightarrow 0$, where the $O(z^2)$ depends analytically on y . For each parameter value x_1 we consider the corresponding parameter values $\zeta, \hat{x}_1, \tilde{\zeta}$ defined for the osculating planar curves γ and β , as in (2.3)–(2.5), see Figure 4. We show that

- the points $\vec{n}_\beta(\zeta)$ and $\vec{n}_\gamma(x_1)$ are $O(x_1^3)$ -close, and hence, lie in z -fibers with $O(x_1^3)$ -closed projections to the y -coordinates;
- their z -coordinates are $O(x_1)$ and differ by a quantity $4cx_1^4(1 + o(1))$;
- the same holds with ζ, x_1 replaced by the points $\tilde{\zeta}, \hat{x}_1$ respectively:

$$z(\vec{n}_\beta(\tilde{\zeta})) - z(\vec{n}_\gamma(\hat{x}_1)) = 4cx_1^4(1 + o(1)). \quad (3.1)$$

Therefore, the f -images of the points $\vec{n}_\beta(\zeta), \vec{n}_\gamma(x_1)$ have z -coordinates that differ by $-4cx_1^4(1 + o(1))$ plus a quantity $O(x_1^3) \times O(x_1^2) = O(x_1^5)$. Thus,

$$z(f(\vec{n}_\beta(\zeta))) - z(f(\vec{n}_\gamma(x_1))) = -4cx_1^4(1 + o(1)). \quad (3.2)$$

But $f(\vec{n}_\beta(\zeta)) = \vec{n}_\beta(\tilde{\zeta})$, $f(\vec{n}_\gamma(x_1)) = g(\vec{n}_\gamma(x_1)) = \vec{n}_\gamma(\hat{x}_1)$. Thus (3.2) contradicts to (3.1). The contradiction thus obtained will prove Lemma 3.3.

Proposition 3.4 *Given a point $O \in \mathbb{R}^n$, a germ α of C^6 -smooth hypersurface in \mathbb{R}^n at O and a two-dimensional plane Π through O transversal to α at O . Let L denote the intersection of the plane Π with the hyperplane tangent to α at O . Set $\beta := \Pi \cap \alpha$. Let the planar curve β have positive curvature at O (which is always the case if α has positive second fundamental form at O). Let $\gamma \subset \Pi$ denote the osculating conic at O for the curve β . There exists a unique quadric $\Gamma \subset \mathbb{R}^n$ containing the osculating conic γ that satisfies the following statements:*

(i) *The quadric Γ is tangent to α at O with contact of at least cubic order, i.e., has the same tangent hyperplane and the same second fundamental quadratic form at O , as α ;*

(ii) *Consider the osculating curves β and γ parametrized by the same parameter t , $t(O) = 0$, so that $\beta(t) - \gamma(t) = O(t^4)$. The smooth unit normal vector fields on α and Γ with the same orientation at O , being restricted to the curves β and γ and treated as vector functions of t , have the same 2-jet at 0.*

*The quadric Γ is called the **osculating quadric along the curve β** for the hypersurface α . It is unique in the class of complex quadrics.*

Addendum to Proposition 3.4. *The above quadric Γ exists and is unique under a weaker assumption, with γ being just some conic tangent to β with contact of order at least four (not necessarily at least five).*

Proof We prove the addendum (a stronger statement). Thus, everywhere below we will assume that γ is a conic tangent to β with contact of order at least 4. Condition (ii) is parametrization-independent: replacing $\gamma(t)$ by $\gamma(\tau(t)) = \gamma(t) + O(t^4)$ changes the unit normal vector to Γ at $\gamma(t)$ by a quantity of order at least three. The system of conditions (i) and (ii) is invariant under projective transformations. Indeed, invariance of condition (i) is obvious. Condition (ii) means that at points $\beta(t)$ and $\gamma(\tau)$ that are close to each other of order $O(t^4)$ the corresponding tangent hyperplanes to α and Γ respectively are $O(t^3)$ -close to each other as elements of Grassmannian manifold (which is in our case the dual projective space). The latter condition is obviously projective-invariant. Applying projective transformation of $\mathbb{R}P^n \supset \mathbb{R}^n$ we can and will consider that O is the origin in the affine chart \mathbb{R}^n , the tangent hyperplane to α at O is the coordinate hyperplane $\{x_n = 0\}$, the plane Π is the (x_1, x_n) -coordinate plane, and the conic γ is the parabola $\{x_n = x_1^2\}$. Set

$$\hat{x} := (x_2, \dots, x_{n-1}).$$

Then each quadric Γ tangent to the hyperplane $\{x_n = 0\}$ at O and containing γ is given by a quadratic equation of the type

$$x_1^2 + x_1 \sum_{j=2}^{n-1} c_j x_j + \langle A\hat{x}, \hat{x} \rangle - x_n \left(1 + \sum_{j=2}^{n-1} d_j x_j\right) = 0. \quad (3.3)$$

Or equivalently,

$$\begin{aligned} \Gamma &= \{h_\Gamma(x_1, \hat{x}) - x_n = 0\}, \quad h_\Gamma(x_1, \hat{x}) = \frac{x_1^2 + x_1 \sum_{j=2}^{n-1} c_j x_j + \langle A\hat{x}, \hat{x} \rangle}{1 + \sum_{j=2}^{n-1} d_j x_j} \\ &= (x_1^2 + x_1 \sum_{j=2}^{n-1} c_j x_j + \langle A\hat{x}, \hat{x} \rangle) \left(1 - \sum_{j=2}^{n-1} d_j x_j\right) + (\text{terms of order } \geq 4). \end{aligned} \quad (3.4)$$

Here A is a symmetric $(n-2) \times (n-2)$ -matrix. Each germ of C^6 -smooth hypersurface α tangent to Γ at O and having the same second quadratic form at O , as Γ , is locally a graph

$$\alpha = \{h_\alpha(x_1, \hat{x}) - x_n = 0\}, \quad (3.5)$$

where h_α is a C^6 -smooth function of the type

$$h_\alpha(x_1, \hat{x}) = x_1^2 + x_1 \sum_{j=2}^{n-1} c_j x_j + \langle A\hat{x}, \hat{x} \rangle + (\text{terms of order } \geq 3). \quad (3.6)$$

The intersection $\beta = \alpha \cap \Pi$ is a curve tangent to γ with contact of order at least 4, if and only if the restriction to the x_1 -axis of the higher order terms in (3.6) has trivial 3-jet. This means that h_α does not contain Taylor term x_1^3 . Let s_j denote the Taylor coefficients of the function h_α at $x_1^2 x_j$, $j = 2, \dots, n-1$. Finally,

$$\begin{aligned} h_\alpha(x_1, \hat{x}) - h_\Gamma(x_1, \hat{x}) &= x_1^2 \sum_{j=2}^{n-1} (s_j + d_j) x_j + x_1 O(\|\hat{x}\|^2) + O(\|\hat{x}\|^3) \\ &+ (\text{terms of order } \geq 4 \text{ in } (x_1, \hat{x})). \end{aligned} \quad (3.7)$$

Claim 3. *Condition (ii) holds if and only if $d_j = -s_j$.*

Proof The normal fields to the hypersurfaces $\xi = \alpha, \Gamma$ directed "down" are equal to

$$\vec{n}_\xi = \frac{\left(\frac{\partial h_\xi}{\partial x_1}, \dots, \frac{\partial h_\xi}{\partial x_{n-1}}, -1\right)}{\sqrt{1 + \sum_{j=1}^{n-1} \left(\frac{\partial h_\xi}{\partial x_j}\right)^2}}. \quad (3.8)$$

We have to compare them along the intersections of the hypersurfaces with the (x_1, x_n) -plane, which means comparing the restrictions of the vector functions (3.8) with $\xi = \alpha, \gamma$ to the x_1 -axis. The only terms in (3.7) that can contribute to linear or quadratic terms in x_1 in the partial derivatives $\frac{\partial h_\alpha}{\partial x_j}$ are

$$x_1^2 \sum_{j=2}^{n-1} (s_j + d_j)x_j, \quad (3.9)$$

since the contributions of the terms $x_1 O(\|\hat{x}\|^2) + O(\|\hat{x}\|^3)$ to the above partial derivatives vanish identically along the x_1 -axis, as does the coordinate vector function \hat{x} , and terms of order at least 4 contribute terms of order at least three. The contributions of terms (3.9) to the above partial derivatives are quadratic, and the derivatives in question are small of order at least one. Therefore, their contributions to the denominators in (3.8) for $\xi = \alpha$ are of order at least three. Finally, the statement saying that the vector functions $\vec{n}_\alpha, \vec{n}_\Gamma$ have the same 2-jet along the x_1 -axis is equivalent to the same statement for the numerators (the gradient fields) in (3.8). But the restriction to the x_1 -axis of the difference of the numerators in (3.8) corresponding to $\xi = \alpha, \Gamma$ is equal (up to terms of order at least three) to the contribution of (3.9): that is, to $(0, (s_2 + d_2)x_1^2, \dots, (s_{n-1} + d_{n-1})x_1^2, 0)$. The latter contribution has trivial second jet, if and only if it vanishes, i.e., $d_j = -s_j$ for all $j = 2, \dots, n-1$. The claim is proved. \square

Thus, all the terms in (3.3) are uniquely determined by conditions (i) and (ii). This proves Proposition 3.4 and its addendum. \square

Consider now new, affine orthogonal coordinates (x_1, \dots, x_n) centered at O such that the x_1 -axis is the common tangent line L to the curves γ and β , and the x_n -axis lies in the plane Π . In what follows we consider that the curves $\gamma, \beta \subset \Pi$ have unit curvature at O : one can achieve this by rescaling the coordinates (x_1, \dots, x_n) by homothety. The hypersurfaces $\xi = \Gamma, \alpha$ are graphs $\{x_n = h_\xi(x_1, \dots, x_{n-1})\}$ of functions $h_{\Gamma, \alpha}$. Their intersections with Π , that is, the curves γ and β , are graphs $\{x_n = \nu_\xi(x_1) := h_\xi(x_1, 0, \dots, 0)\}$. For every x_1 let $\hat{x}_1, \zeta, \tilde{\zeta}$ be the same, as in (2.3)–(2.5):

$$\zeta = \zeta(x_1) \text{ is the point for which } \nu_\alpha(\zeta) = \nu_\Gamma(x_1), \text{ and } \zeta \simeq x_1,$$

$$\hat{x}_1 = \hat{x}_1(x_1) \text{ is the point distinct from } x_1 \text{ where } \nu_\Gamma(\hat{x}_1) = \nu_\Gamma(x_1),$$

$$\tilde{\zeta} = \tilde{\zeta}(x_1) \text{ is the point distinct from } \zeta \text{ where } \nu_\alpha(\tilde{\zeta}) = \nu_\alpha(\zeta),$$

see Figure 4 (with h replaced by ν). We consider that the curves γ and β are parametrized by the parameter x_1 as graphs of functions $\nu_\Gamma(x_1)$ and $\nu_\alpha(x_1)$

respectively. Recall that we assume that the curves γ and β have tangency of contact of order exactly 5. Therefore,

$$\nu_\alpha(x_1) - \nu_\Gamma(x_1) = cx_1^5(1 + o(1)), \quad \text{as } x_1 \rightarrow 0; \quad c \neq 0. \quad (3.10)$$

Recall that we denote $f = \mathcal{R}_{\alpha, \mathcal{L}}$ and $g = \mathcal{R}_{\Gamma, \mathcal{L}}$. Here \mathcal{L} is the class of lines parallel to the line L tangent to the curves β, γ at O .

Proposition 3.5 *1) Consider the restrictions $\vec{n}_\gamma(x_1), \vec{n}_\beta(x_1)$ to the curves γ and β of the exterior unit normal vector fields to Γ and α respectively as functions of x_1 . One has*

$$g(\vec{n}_\gamma(x_1)) = \vec{n}_\gamma(\hat{x}_1), \quad f(\vec{n}_\beta(\zeta)) = \vec{n}_\beta(\tilde{\zeta}), \quad (3.11)$$

$$\vec{n}_\gamma(x_1) - \vec{n}_\beta(\zeta) = O(x_1^3), \quad \vec{n}_\gamma(\hat{x}_1) - \vec{n}_\beta(\tilde{\zeta}) = O(x_1^3). \quad (3.12)$$

2) Consider the orthogonal projections to Π of the vector fields $\vec{n}_\gamma, \vec{n}_\beta$. Let $\psi_\gamma(x_1)$ and $\psi_\beta(x_1)$ denote the arguments of the projections (their azimuths with respect to the x_1 -axis). One has

$$\psi_\beta(\zeta) - \psi_\gamma(x_1) = 4cx_1^4(1 + o(1)), \quad (3.13)$$

$$\psi_\beta(\tilde{\zeta}) - \psi_\gamma(\hat{x}_1) = 4cx_1^4(1 + o(1)), \quad (3.14)$$

Proof Formulas (3.11) follow from definition. One has

$$\vec{n}_\beta(x_1) - \vec{n}_\gamma(x_1) = O(x_1^3). \quad (3.15)$$

This follows from the fact that the restrictions to γ, β of the normal vector fields of the hypersurfaces Γ and α have the same 2-jet at O , which in its turn follows from Statement (ii) of Proposition 3.4. One has $\zeta - x_1 = O(x_1^4)$, by (2.7). This together with (3.15) implies the first formula in (3.12), and the proof of the second formula is analogous. Each argument $\psi_{\beta, \gamma}$ in (3.13), (3.14) is equal to $-\frac{\pi}{2}$ plus the azimuth of the oriented tangent line to the curve in question at the corresponding point. The tangents of the latter azimuths are the derivatives in (2.8). Formulas (3.13) and (3.14) follow from formulas (2.8), the two last statements and the fact that $\tan x$ has unit derivative at the origin. Proposition 3.5 is proved. \square

Proposition 3.6 *The involutions $f = \mathcal{R}_{\alpha, \mathcal{L}}$ and $g = \mathcal{R}_{\Gamma, \mathcal{L}}$ coincide.*

Proof One has

$$g(\vec{n}_\gamma(x_1)) - f(\vec{n}_\gamma(x_1)) = O(x_1^3), \quad \text{as } x_1 \rightarrow 0. \quad (3.16)$$

This follows from the fact that the points $\vec{n}_\gamma(x_1)$, $\vec{n}_\beta(\zeta)$ are $O(x_1^3)$ -close, as are their g - and f - images respectively, see (3.12) and (3.11), and smoothness of the maps f and g . On the other hand, we prove the following claim.

Claim 4. *If $f \neq g$, then $g(\vec{n}_\gamma(x_1)) - f(\vec{n}_\gamma(x_1))$ is not $O(x_1^3)$, as $x_1 \rightarrow 0$.*

Proof The projective involutions f and g have common hyperplane H of fixed points, which is the projectivization of the codimension one vector subspace $\{x_1 = 0\}$ orthogonal to L . Each involution $v = f, g$ has yet another fixed point $P_v \notin H$, fixes each line through P_v and acts there as a projective involution fixing P_v and its intersection point with H . Inequality $f \neq g$ is equivalent to the inequality $P_f \neq P_g$.

Let us identify O with the point in \mathbb{RP}^{n-1} represented by the vector $\vec{n}_\beta(O) = \vec{n}_\gamma(O)$ and introduce affine coordinates (z, y) , $y = (y_1, \dots, y_{n-2})$, centered at O on an affine chart in \mathbb{RP}^{n-1} so that $H = \{z = 0\}$ and along the curves $\vec{n}_\gamma(x_1)$ and $\vec{n}_\beta(x_1)$ one has $z \simeq x_1$, as $x_1 \rightarrow 0$. One has

$$z \circ g(\vec{n}_\gamma(x_1)) \simeq z \circ f(\vec{n}_\gamma(x_1)) \simeq -z(\vec{n}_\gamma(x_1)) \simeq -x_1, \quad \text{as } x_1 \rightarrow 0, \quad (3.17)$$

since g and f are involutions fixing points in H and γ is transversal to H . For every x_1 small enough the images $g(\vec{n}_\gamma(x_1))$, $f(\vec{n}_\gamma(x_1))$ lie in two lines $\Lambda_g(x_1)$ and $\Lambda_f(x_1)$ passing through $\vec{n}_\gamma(x_1)$ and P_g (respectively, $\vec{n}_\gamma(x_1)$ and P_f). The difference between the z -coordinates of each image and of its source point is asymptotic to $-2x_1$, by (3.17).

Case 1): $\Lambda_f(0) \neq \Lambda_g(0)$. Then the projections of the images $g(\vec{n}_\gamma(x_1))$, $f(\vec{n}_\gamma(x_1))$ to the hyperplane H along the z -axis differ by a quantity of order $c''x_1(1 + o(1))$, $c'' \neq 0$. Hence, the images do not differ by $O(x_1^3)$.

Case 2): $\Lambda_f(0) = \Lambda_g(0) = \Lambda$. We choose the above coordinates (z, y) so that in addition, the line Λ is the z -axis. Both projective involutions f and g fix the line Λ , and in the z -coordinate on Λ one has

$$f(z) - g(z) = az^2(1 + o(1)), \quad \text{as } z \rightarrow 0; \quad a \neq 0, \quad (3.18)$$

by Proposition 2.4. On the other hand, the functions $\Phi_1 := z \circ f$ and $\Phi_2 := z \circ g$ restricted to a small neighborhood of the point O equipped with the coordinates $z = (z, y)$ have difference with the asymptotics

$$\Phi_2(z) - \Phi_1(z) = \chi(y)z^2 + o(z^2), \quad \text{as } z \rightarrow 0, \quad (3.19)$$

where $\chi(y)$ is an analytic function. One has $\chi(0) = a \neq 0$, by (3.18). This implies that along the curve $\vec{n}_\gamma(x_1)$ the functions Φ_1 and Φ_2 differ by

a quantity $ax_1^2(1 + o(1)) \neq O(x_1^3)$. Therefore, the images $f(\vec{n}_\gamma(x_1))$ and $g(\vec{n}_\gamma(x_1))$ do not differ by a quantity $O(x_1^3)$. This proves Claim 4. \square

Claim 4 together with (3.16) imply that $f = g$. Proposition 3.6 is proved. \square

Let c be the constant factor in the right-hand side in (3.13). Let P be the fixed point of the involution $f = g$ that does not lie in H .

Claim 5. *Choosing appropriate coordinates (z, y) as at the beginning of the proof of Claim 4 one can achieve that the projection $(z, y) \mapsto y$ is the projection $\mathbb{R}\mathbb{P}^{n-1} \setminus \{P\} \rightarrow H$ from the point P and the following asymptotic equalities hold:*

$$z(\vec{n}_\beta(\zeta)) - z(\vec{n}_\gamma(x_1)) = 4cx_1^4(1 + o(1)), \quad (3.20)$$

$$z(\vec{n}_\beta(\tilde{\zeta})) - z(\vec{n}_\gamma(\hat{x}_1)) = 4cx_1^4(1 + o(1)) \text{ as } x_1 \rightarrow 0. \quad (3.21)$$

Proof Fix some affine coordinates $y = (y_1, \dots, y_{n-1})$ on H centered at O . We extend the coordinate vector function y to all of $\mathbb{R}\mathbb{P}^n \setminus \{P\}$ as the projection $\mathbb{R}\mathbb{P}^n \setminus \{P\} \rightarrow H_{y_1, \dots, y_{n-1}}$ from the point P . Let (x_1, \dots, x_n) be the orthogonal coordinates on \mathbb{R}^n introduced above. For every vector $\vec{n} \in \mathbb{R}^n$ set

$$z(\vec{n}) = -\frac{x_1(\vec{n})}{x_n(\vec{n})}.$$

The map $\vec{n} \mapsto z(\vec{n})$ is a projection $\mathbb{R}\mathbb{P}^{n-1} \setminus V \rightarrow \mathbb{R}\mathbb{P}_{[x_1: x_n]}^1$ from the codimension two projective subspace $V = \{x_1 = x_n = 0\} \subset H \setminus \{O\}$. (Recall that the latter O is identified with $\vec{n}_\gamma(O) = \vec{n}_\beta(O)$. One has $\vec{n}_\beta(O) \notin V$, since the vector $\vec{n}_\beta(O)$ is not orthogonal to the (x_1, x_n) -coordinate plane Π : it is orthogonal to the hyperplane $T_O\alpha = T_O\Gamma$ transversal to Π .) One has $z|_H \equiv 0$, by construction, since H is the projectivization of the $(n-1)$ -dimensional subspace $\{x_1 = 0\}$ orthogonal to the line L : the x_1 -axis in the orthogonal coordinates x_1, \dots, x_n . Afterwards formulas (3.20) and (3.21) follow from formulas (3.13) and (3.14): the z -coordinates in (3.20) and (3.21) are equal to the tangents of the arguments in (3.13) and (3.14) with added $\frac{\pi}{2}$, and $\tan \phi \simeq \phi$, as $\phi \rightarrow 0$. The claim is proved. \square

Proof of Lemma 3.3. One has $f = g$ (Proposition 3.6), hence,

$$\vec{n}_\gamma(\hat{x}_1) = f(\vec{n}_\gamma(x_1)), \quad \vec{n}_\beta(\tilde{\zeta}) = f(\vec{n}_\beta(\zeta)), \quad (3.22)$$

by definition.

Claim 6. *Let (z, y) be the coordinates as in Claim 5. Then*

$$z(f(\vec{n}_\beta(\zeta))) - z(f(\vec{n}_\gamma(x_1))) = -(z(\vec{n}_\beta(\zeta)) - z(\vec{n}_\gamma(x_1))) + O(x_1^5). \quad (3.23)$$

Proof Each pair "image-preimage" in (3.22) lies in one and the same z -fiber, since each z -fiber is f -invariant (it contains the fixed point P of the involution f). The y -coordinates (projections to H) of the latter z -fibers are $O(x_1^3)$ -close to each other: $y(\vec{n}_\gamma(x_1)) - y(\vec{n}_\beta(\zeta)) = O(x_1^3)$, by (3.12). On the other hand, the restrictions to the z -fibers of the involution f (written in the coordinate z) are of the form $f(z) = -z + \sum_{j=2}^5 b_j(y)z^j + o(z^5)$, as $z \rightarrow 0$, where $b_j(y)$ are analytic functions on a neighborhood of the point O . This and the above statements together imply that the restrictions of the projective involution f to the z -fibers containing $\vec{n}_\gamma(x_1)$ and $\vec{n}_\beta(\zeta)$ differ by a quantity $O(z^2x_1^3)$, as z and x_1 tend to zero. Let us introduce the auxiliary point $q(x_1)$ that lies in the same z -fiber, as $\vec{n}_\gamma(x_1)$, but has the same z -coordinate, as the other preimage $\vec{n}_\beta(\zeta)$. One has

$$z \circ f(\vec{n}_\beta(\zeta)) - z \circ f(q(x_1)) = O(x_1^5), \quad (3.24)$$

by the previous statement, since $z(\vec{n}_\gamma(x_1)) \simeq z(\vec{n}_\beta(x_1)) = O(x_1)$, and hence, $O(z^2x_1^3) = O(x_1^5)$. On the other hand,

$$z(q(x_1)) - z(\vec{n}_\gamma(x_1)) = z(\vec{n}_\beta(\zeta)) - z(\vec{n}_\gamma(x_1)) = 4cx_1^4(1 + o(1)),$$

by (3.20). Thus, the points $q(x_1)$ and $\vec{n}_\gamma(x_1)$ lie in the same z -fiber and have $O(x_1^4)$ -close z -coordinates that are $O(x_1)$. Therefore, the difference of the z -coordinates of their f -images is equal to minus the difference of their own z -coordinates times a quantity $1 + O(x_1)$, and $O(x_1)$ times $O(x_1^4)$ yields $O(x_1^5)$. This implies (3.23) with $\vec{n}_\beta(\zeta)$ replaced by $q(x_1)$. Together with (3.24), this implies (3.23). Claim 6 is proved. \square

The right-hand side in (3.23) is $-4cx_1^4(1 + o(1))$, by (3.20). But its left-hand side is equal to the left-hand side in (3.21), and hence, is $4cx_1^4(1 + o(1))$, by (3.21). The contradiction thus obtained proves Lemma 3.3. \square

Theorem 3.1 is proved, since it follows from Lemma 3.3.

3.2 Proof of Theorem 1.12 in higher dimensions

Theorem 3.7 [5, theorem 4] *Let a hypersurface in \mathbb{R}^n be such that each its planar section is a conic. Then it is a quadric.*

Theorem 3.1 implies that each planar section of the hypersurface α is a conic. This together with Theorem 3.7 proves Theorem 1.12.

4 Acknowledgements

I wish to thank Anastasiia Sharipova and Sergei Tabachnikov for introducing me to the area of Hamiltonian, Finsler and projective billiards and for helpful discussions.

References

- [1] Abate M.; Patrizio G. *Finsler Metrics – A Global Approach*. - Lecture Notes in Mathematics, Volume **1591**, Springer, Berlin, 1994.
- [2] Alvarez Paiva J.C. *Some problems on Finsler geometry*. - Handbook of Differential Geometry (editors: Franki J.E. Dillen, Leopold C.A. Verstraelen). North-Holland, Volume 2, 2006, Chapter 1, 1–33.
- [3] Alvarez Paiva J.C.; Durán C.E. *An introduction to Finsler geometry*. - Publicaciones de la Escuela Venezolana de Matemáticas, Caracas, Venezuela, 1998.
- [4] Arnold V. *Mathematical Methods of Classical Mechanics*. - Springer, Berlin, 1989.
- [5] Arnold, M.; Tabachnikov, S. *Remarks on Joachimsthal Integral and Poritsky Property*. - Arnold Math. J. **7** (2021), 483–491.
- [6] Artstein-Avidan, S.; Karasev, R.; Ostrover, Y. *From symplectic measurements to the Mahler Conjecture*. - Duke Math. J. **163** (2014), No. 11, 2003–2022.
- [7] Artstein-Avidan, S.; Ostrover, Y. *Bounds for Minkowski billiard trajectories in convex bodies*. - Int. Math. Res. Not. **2014** (2014), Issue 1, 165–193.
- [8] Bao D.; Chern S.-S.; Shen Z. *An Introduction to Riemann–Finsler Geometry*. - Springer, Berlin, 2000.
- [9] Buchin, S. *Affine Differential Geometry*. - Gordon & Breach, New York, 1983.
- [10] Gutkin, E.; Tabachnikov, S. *Billiards in Finsler and Minkowski geometries*. - J. Geom. Phys. **40** (2002), 277–301.
- [11] Hofer, H.; Zehnder, E. *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser, Basel, 1994.

- [12] Schouten, J. A. *Ricci-calculus. An introduction to tensor analysis and its geometrical applications.* - Die Grundlehren der mathematischen Wissenschaften. Vol. **10** (Second edition of 1923 original ed.). Berlin-Göttingen-Heidelberg, Springer-Verlag, 1954.
- [13] Tabachnikov, S. *Introducing projective billiards.* Ergod. Th. Dynam. Sys. **17** (1997), 957–976.
- [14] Viterbo, C. *Metric and isoperimetric problems in symplectic geometry.* - J. Amer. Math. Soc. **13** (2000), 411–431.