ON THE NUMBER OF GENERATORS OF GROUPS ACTING ARC-TRANSITIVELY ON GRAPHS

MARCO BARBIERI AND PABLO SPIGA

ABSTRACT. Given a finite connected graph Γ and a group G acting transitively on the vertices of Γ , we prove that the number of vertices of Γ and the cardinality of Gare bounded above by a function depending only on the cardinality of Γ and on the exponent of G. We also prove that the number of generators of a group G acting transitively on the arcs of a finite graph Γ cannot be bounded by a function of the valency alone.

1. INTRODUCTION

Let Γ be a locally finite connected graph, let G be a group acting transitively on the arcs of Γ . Suppose that, for any vertex α of Γ , the order of the vertex-stabilizer G_{α} is finite. In this note, an *arc-transitive graph* is a pair (Γ, G) with the properties just described. The *local action of* (Γ, G) is the transitive action that a vertex-stabilizer G_{α} induces on the neighbourhood $\Gamma(\alpha)$ of the fixed point α . In particular, the *valency of* Γ coincides the degree of the local action of (Γ, G) .

We say that a local action is graph-restrictive if, for any arc-transitive graphs (Γ, G) admitting such local action, then the order of the vertex-stabilizers is bounded from above by a constant depending only on the action. With this notation, the famous Weiss Conjecture (posed in [8]) states that primitive groups are graph-restrictive. We refer to [6] for an extensive study of this notion.

Let (Γ, G) be an arc-transitive graph of valency d whose local action is graph-restrictive. One can easily check that, by a connectedness argument (see the proof of Thereom 5), the group H generated by those elements d of G sending a prescribed vertex α to all its neighbours in turn is transitive. Therefore, by Frattini's Argument, $G = G_{\alpha}H$. In particular, $|G_{\alpha}| + d$ elements are sufficient to generate G. Hence the minimal number of generators for G, denoted by $\mathbf{d}(G)$ can be bounded with a function of the local action alone.

More surprisingly, for every arc-transitive graph (Γ, G) of valency at most 4, the minimal number of generators of G is bounded by a constant regardless of the local action. The result is trivial for $d \in \{1, 2\}$. For $d \in \{3, 4\}$, some deeper concepts enter the picture. For every arc-transitive graph (Γ, G) , there is a universal cover of the form $(\mathcal{T}_d, G_\alpha *_{G_{\alpha\beta}} G_{\{\alpha,\beta\}})$, where α and β are two adjacent vertices of Γ , adn \mathcal{T}_d is the infinite tree of valency d. For every amalgamated product appearing in the universal cover for valency 3 and 4, an explicit presentation has been produced: see [1, 2, 4]. A direct inspection of these presentations proves that the minimal number of generators for G is at most 10.

One could dare to conjecture that there exists a function $\mathbf{f} : \mathbb{N} \to \mathbb{N}$ that takes the valency of the graph Γ as input, and returns an upper bound for $\mathbf{d}(G)$. The main contribution of this note is proving that such a function cannot exist.

Theorem 1. There exists no function $\mathbf{f} : \mathbb{N} \to \mathbb{N}$ such that, for every arc-transitive graph (Γ, G) of valency d,

 $\mathbf{d}(G) \leq \mathbf{f}(d) \,.$

Remark 2. To prove the veracity of Theorem 1, we will exhibit an infinite family \mathcal{F} of pairs (Γ_h, G_h) such that the valency of the graphs is a constant (at least 8), while

²⁰²⁰ Mathematics Subject Classification. 20B25, 05C25.

Key words and phrases. Number of generators, arc-transitive, valency.

The authors are members of the GNSAGA INdAM and PRIN "Group theory and its applications" research group and kindly acknowledge their support.

 $\mathbf{d}(G_h)$ grows linearly with the exponent of the group. We would like to remark that the graphs Γ_h carry an oustanding similarity with those build in [3, 5, 7] to prove that, for some imprimitive local actions of degree 6, the order of the vertex-stabilizers grows exponentially.

We also observe that, in our construction, G is not the automorphism group of Γ . This prompts the following question.

Problem 3. Is there a function $\mathbf{f} : \mathbb{N} \to \mathbb{N}$ such that, if Γ is a connected arc-transitive graph of valency d, then

$$\mathbf{d}(\operatorname{Aut}(\Gamma)) \leq \mathbf{f}(d)$$
?

Moreover, for our current and limited knowledge of arc-transitive graphs (Γ, G) , having $\mathbf{d}(G)$ bounded appears to be quite common. Therefore, we ask the following.

Problem 4. Which assumption on the arc-transitive graph (Γ, G) are necessary to bound $\mathbf{d}(G)$ with a function of the valency (or the local action)?

To conclude, we give a bound on the order of the group G appearing in an arctransitive graph (Γ, G) depending on the valency d and the exponent of G. Our proof does not rely on the fact that the local action is transitive, thus the result holds for the larger class of vertex-transitive graphs. In this note, a *vertex-transitive graph* is a pair (Γ, G) where Γ is a finite connected graph and G is a subgroup of $\operatorname{Aut}(\Gamma)$ whose action on the vertex-set of Γ is transitive.

We underline that the exponent of G is not a local feature of the graph. For instance, if Γ is a cycle of odd length, then the local action is isomorphic to C_2 , while the exponent of Aut(Γ) is twice the length of the cycle. Finally, we point out that the function **g** appearing in Theorem 5 is not explicitly determined, but it depends on the solution of the Burnside Restricted Problem. Moreover, even for the few cases in which we can make it explicit, its growth is at least factorial in both parameters.

Theorem 5. Let (Γ, G) be a vertex-transitive graph. Suppose that the valency of Γ is d, and that the exponent of G is e. Then there exists a function $\mathbf{g} : \mathbb{N} \times \mathbb{N} \to \mathbb{N}$ such that

$$|G| \le \mathbf{g}(d, e)$$

2. Proof of Theorem 1

Let h be a positive integer, and let p be a prime. Set

$$H = C_{p^{h}} \times C_{p^{h}} = \langle a, b \mid a^{p^{h}} = b^{p^{h}} = [a, b] = 1 \rangle,$$

and consider the group algebra $\mathbb{F}_p[H]$. Define recursively the following chain of *H*-modules

$$\gamma_0 := \mathbb{F}_p[H], \quad \text{and, for any } i \ge 1,$$

$$\gamma_i := [\gamma_{i-1}, H] = \langle v - vh \mid v \in \gamma_{i-1}, h \in H \rangle_{\mathbb{F}_p}.$$

Consider the $(p^h \times p^h)$ -square unipotent matrix

$$M = \begin{pmatrix} 1 & 1 & 0 & \dots & 0 & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 \end{pmatrix},$$

and denote the identity matrix of size p^h by *id*. Observe that M is an element of order p^h in $\operatorname{GL}_{p^h}(p)$.

The natural basis for the group algebra $\mathbb{F}_p[H]$ is

$$(a^i b^j \mid i, j \in \{0, 1, \dots, p^h - 1\})$$
.

We have that

$$\mathcal{B} = \left(e_{xy} = (a-1)^x (b-1)^y \mid x, y \in \{0, 1, \dots, p^h - 1\}\right)$$

is also a basis. For convenience, we set $e_{xp^h} = e_{p^h y} = 0$, for every $x, y \in \{0, 1, \dots, p^h - 1\}$. Observe that

$$e_{xy} \cdot a = (a-1)^x (b-1)^y \cdot a$$

= $(a-1)^x (1+a-1)(b-1)^y$
= $(a-1)^x (b-1)^y + (a-1)^{x+1} (b-1)^y$
= $e_{xy} + e_{(x+1)y}$,

and

$$e_{xy} \cdot b = (a-1)^x (b-1)^y \cdot b$$

= $(a-1)^x (b-1)^y (1+b-1)$
= $(a-1)^x (b-1)^y + (a-1)^x (b-1)^{y+1}$
= $e_{xy} + e_{x(y+1)}$.

Therefore, in the basis \mathcal{B} , the action of a and b on $F_p[H]$ is given by the matrices by the matrices $M \otimes id$ and $id \otimes M$ respectively.

With a direct computation, we get that

$$\begin{split} \gamma_i &= \langle e_{xy} \mid x+y \geq i \rangle_{\mathbb{F}_p}, \quad \text{and} \\ \gamma_i / \gamma_{i+1} &= \langle e_{xy} + \gamma_{i+1} \mid x+y = i \rangle_{\mathbb{F}_p}. \end{split}$$

Recall that, for any *H*-module *V*, we denote by $\mathbf{d}_H(V)$ the minimal number of generators of *V* as an *H*-module. Since, by construction, γ_i/γ_{i+1} is a trivial section of $\mathbb{F}_p[H]$, we have that

$$\mathbf{d}_{H}(\gamma_{i}/\gamma_{i+1}) = \dim_{\mathbb{F}_{p}}(\gamma_{i}/\gamma_{i+1}) = \min\{i+1, 2p^{h} - i - 1\}.$$

Therefore, $\mathbf{d}_H(\gamma_{p^h-1}) = p^h$. In particular,

$$\mathbf{d}\left(\gamma_{p^{h}-1}\rtimes H\right)=p^{h}+2.$$

The group H acts regularly on the vertex-set of the 4-valent Cayley graph defined by

$$\Delta := \operatorname{Cay}(H, \{a, a^{-1}, b, b^{-1}\}).$$

Recall that, for any two graphs Γ , Δ , the wreath product of Δ by Γ , denoted by Γ wr Δ , is the graph having vertex-set $V\Gamma \times V\Delta$, where (δ_1, γ_1) and (δ_2, γ_2) are adjacent if either $\delta_1 = \delta_2$ and $\{\gamma_1, \gamma_2\}$ is an edge of Γ , or $\{\delta_1, \delta_2\}$ is an edge of Δ . We can finally define Γ_h as the wreath product of the empty graph on p vertices, $p\mathbf{K}_1$, by the Cayley graph Δ , that is, $\Gamma_h := p\mathbf{K}_1 \text{ wr } \Delta$. Note that, unless p = 2 and h = 1, Γ_h has valency 4p.

Observe that $\gamma_{p^h-1} \rtimes H$ embeds into $C_p \operatorname{wr} H$, which is a subgroup of the automorphism group of Γ_h . Moreover, as γ_{p^h-1} is not trivial, $\gamma_{p^h-1} \rtimes H$ must be transitive on the vertices of Γ_h . On the other hand, since $\gamma_{p^h-1} \rtimes H$ preserves the lifting of the labels $\{a, a^{-1}, b, b^{-1}\}$ from the Cayley graph Δ , this action is not arc-transitive. In particular, the local action of $(\Gamma_h, \gamma_{p^h-1} \rtimes H)$ is intransitive with 4 distinct orbits.

We need to extend H with some outer automorphisms to achieve arc-transitivity. Consider the automorphism of H defined on the generators by

 $\varphi: a \mapsto b, b \mapsto a, \text{ and } \psi: a \mapsto a^{-1}, b \mapsto b^{-1}.$

Observe that both are involution, and that $\langle \varphi, \psi \rangle$ is isomorphic to the Klein group. Extend the multiplication on $\mathbb{F}_p[H]$ by putting, for every $\varepsilon, \delta \in \mathbb{Z}_2$,

$$\left(\sum_{h\in H}\lambda_h h\right)\cdot\left(\varphi^{\varepsilon}\psi^{\delta}\right)=\sum_{h\in H}\lambda_h h^{\varphi^{\varepsilon}\psi^{\delta}}$$

With this operation, $\gamma_0 = \mathbb{F}_p[H]$ is an $\mathbb{F}_p[H \rtimes \langle \varphi, \psi \rangle]$ -module. Furthermore, for all indices i, since γ_i is a characteristic subgroup of $\gamma_0 \rtimes H$, then γ_i is an $\mathbb{F}_p[H \rtimes \langle \varphi, \psi \rangle]$ -submodule. In particular, the group $G_h := \gamma_{p^h-1} \rtimes (H \rtimes \langle \varphi, \psi \rangle)$ is a subgroup of $\operatorname{Aut}(\Gamma_h)$. Finally, as the index of H in $H \rtimes \langle \varphi, \psi \rangle$ is 4, we have that

$$\mathbf{d}_{H\rtimes\langle\varphi,\psi\rangle}(\gamma_{p^h-1}) \geq \frac{p^h}{4} \,.$$

Hence, we obtain that

$$\mathbf{d}(G_h) \ge \frac{p^h}{4} + 3.$$

Let us go back to the Cayley graph Δ . Observe that $\langle \varphi, \psi \rangle$ is transitive on the connection set of Δ . This implies that $H \rtimes \langle \varphi, \psi \rangle$ is an arc-transitive subgroup of Aut(Δ). Therefore, the local action of (Γ_h, G_h) transitively permutes the 4 orbits defined by $(\Gamma_h, \gamma_{p^h-1} \rtimes H)$. Hence, the action of G_h on the graph Γ_h is arc-transitive.

Therefore, for a fixed prime p, the family

$$\mathcal{F}_p = \{ (\Gamma_h, G_h) \mid h \ge 2 \}$$

contains only graphs of valency 4p, meanwhile

$$\lim_{h \to +\infty} \mathbf{d}(G_h) \ge \lim_{h \to +\infty} \frac{p^h}{4} + 3 = +\infty.$$

This concludes the proof of Theorem 1.

3. Proof of Theorem 5

Let (Γ, G) be a vertex-transitive graph, and suppose that d is the valency of Γ . Choose a vertex α of Γ , and, for any of its neighbours β , let $g_{\beta} \in G$ be an element such that

$$\alpha^{g_\beta} = \beta$$

(This element exists by the transitivity of G on the vertex-set of Γ .) Define the subgroup

$$H := \langle g_{\beta} \mid \beta \text{ is a neighbour of } \alpha \rangle \leq G.$$

By a routine connectedness argument, H is transitive on the vertex-set of Γ . Hence, by Frattini's Argument, $G = G_{\alpha}H$.

Let $\mathbf{B}(d, e)$ be the function solving the Burnside Restricted Problem for a finite group with d generators and exponent e. The existence of this function for all the choices of dand e was proved by Zel'manov in [9, 10]. Then, $|H| \leq B(d, e)$.

We now need to bound the order of G_{α} . Note that G_{α} can be embedded into $\operatorname{Sym}(V\Gamma - \{\alpha\})$, hence, as $|V\Gamma| \leq |H|$,

$$|G_{\alpha}| \le (\mathbf{B}(d, e) - 1)!$$

Therefore, we obtain that

$$|G| = |H||G_{\alpha}| \leq \mathbf{B}(d, e)(\mathbf{B}(d, e) - 1)!.$$

References

- D. Ž. Djoković. A class of finite group-amalgams. Proceedings of the American Mathematical Society, 80:22–26, 1980. 1
- [2] D. Ž. Djoković and G. L. Miller. Regular groups of automorphisms of cubic graphs. Journal of Combinatorial Theory, Series B, 29(2):195–230, 1980. 1
- [3] A. Hujdurović, P. Potočnik, and G. Verret. Three local actions in 6-valent arc-transitive graphs. Journal of Graph Theory, 99(2):207–216, 2022.
- [4] P. Potočnik. A list of 4-valent 2-arc-transitive graphs and finite faithful amalgams of index (4, 2). European Journal of Combinatorics, 30(5):1323–1336, 2009.
- [5] P. Potočnik, P. Spiga, and G. Verret. On the order of arc-stabilizers in arc-transitive graphs with prescribed local group. *Transactions of the American Mathematical Society*, 366(7). 2
- [6] P. Potočnik, P. Spiga, and G. Verret. On graph-restrictive permutation groups. Journal of Combinatorial Theory, Series B, 102(3):820–831, 2012.
- [7] P. Potočnik, P. Spiga, and G. Verret. On the nullspace of arc-transitive graphs over finite fields. Journal of Algebraic Combinatorics, 36:389–401, 2012. 2
- [8] R. Weiss. s-transitive graphs. Colloquia Mathematica Societatis János Bolyai, 25:827–847, 1978. 1
- [9] E. I. Zel'manov. Solution of the restricted Burnside problem for groups of odd exponent. Mathematics of the USSR Izvestiya, 36:41–60, 1991.
- [10] E. I. Zel'manov. A solution of the restricted Burnside problem for 2-groups. Mathematics of the USSR – Sbornik, 72(2):543–565, 1992. 3

Dipartimento di Matematica "Felice Casorati", University of Pavia, Via Ferrata 5, 27100 Pavia, Italy

Email address: marco.barbieri07@universitadipavia.it

Dipartimento di Matematica e Applicazioni , University of Milano-Bicocca, Via Cozzi 55, 20125 Milano, Italy

Email address: pablo.spiga@unimib.it