# The effect of regularization in high dimensional Cox regression

**Emanuele Massa** 

Physics of Machine Learning and Complex Systems, Radboud Universiteit, Nijmegen, 6525 AJ, The Netherlands emanuele.massa@donders.ru.nl

#### Abstract

We investigate analytically the behaviour of the penalized maximum partial likelihood estimator (PMPLE). Our results are derived for a generic separable regularization, but we focus on the elastic net. This penalization is routinely adopted for survival analysis in the high dimensional regime, where the Maximum Partial Likelihood estimator (no regularization) might not even exist. Previous theoretical results require that the number s of non-zero association coefficients is  $O(n^{\alpha})$ , with  $\alpha \in (0, 1)$  and n the sample size. Here we accurately characterize the behaviour of the PMPLE when s is proportional to n via the solution of a system of six non-linear equations that can be easily obtained by fixed point iteration. These equations are derived by means of the replica method and under the assumption that the covariates  $\mathbf{X} \in \mathbb{R}^p$  follow a multivariate Gaussian law with covariance  $I_p/p$ . The solution of the previous equations allows us to investigate the dependency of various metrics of interest and hence their dependency on the ratio  $\zeta = p/n$ , the fraction of true active components  $\nu = s/p$ , and the regularization strength. We validate our results by extensive numerical simulations.

# 1 Introduction

When the number of features is vastly larger than the number of observations, the estimator obtained by maximization of the Cox partial likelihood is not well defined. Hence the necessity of adding a regularization term in order to "fit" the model, i.e. define a well posed optimization problem. The elastic net regularization [27] is widely adopted in this context when the goal of the analysis is to obtain a more "parsimonious" model. By shrinking some of the coefficient exactly to zero, elastic net is capable of performing variable selection, even when p > n and the Lasso [26] might not be stable [28]. Because of this appealing characteristic, the use of lasso and elastic net has been investigated in the statistical literature for Generalized Linear Models [10, 3], but also for the Cox model [1, 15]. In particular, [1] shows that, with the Lasso regularization, the Penalized Maximum Partial Likelihood Estimator (PMPLE) is equal to a (biased) oracle estimator which knows the underlying true model, with overwhelming probability as the sample size n diverges. Besides some mild assumptions on the data generating process, their results require that the number of true active component s (the number of covariates that are effectively associated with the survival time) does not grow linearly with the number of observations n, i.e.  $s = O(n^{\alpha})$  for  $\alpha \in (0, 1)$ .

Here we characterize the behaviour of the PMPLE in the proportional regime, where the number of observation n, the number of covariates p and, importantly, the number of active covariates s, diverge proportionally, i.e.  $p = \zeta n$ ,  $s = \nu p$ . This is achieved by means of the replica method from statistical physics [9, 4, 17], which has already been applied in the recent past to study the behaviour of sparse linear regression [13, 20]. The replica method leads to six coupled non-linear equations whose solution grants access to virtually any metric of interest, e.g. the mean squared error for

the estimator of the association and of the survival function, but also the fraction of false positive and negatives in support recovery (as originally noted in [22] for logistic regression). A sufficient "working" assumption for the applicability of the replica method, which we shall also adopt here, is that the covariates  $\mathbf{X} \in \mathbb{R}^p$  are  $\mathbf{X} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p/p)$  [5, 13]. This requirement might be, in principle, relaxed to the less stringent requirement of i.i.d. covariates with finite variance of order O(1/p). It is relatively straightforward to extend our results to this scenario, hence we opt for the simpler setting. In any case we notice that the condition of Gaussian covariates is required in other approaches to high dimensional statistics in the proportional asymptotic regime, e.g. the Convex Gaussian Min Max Theorem (CGMT) [25, 24] or Approximate Message Passing (AMP)[7]. Both CGMT and AMP have been used to investigate the behaviour of sparse linear regression ([25, 19]), sparse logit regression [22] and regularized M-estimation for generalized linear models [24, 17]. The assumption of un-correlated covariates is needed to ensure that the non-linear equations can be neatly expressed as expectations over a small number of univariate random variables and depend solely on the ratios  $\zeta = p/n$  and  $\nu = s/p$ , rather then on p or n. In presence of correlations and for a non-rotationally invariant regularization, this seems to be not directly possible (even when the regularizer is serapable as in our case). The system of non-linear equations would, in that case, depend implicitly on a (non-separable) minimization over p variables, in the limit  $p \to \infty$ . Hence we opt for the assumption of uncorrelated covariates, knowing that this represents an ideal assumption. In any case we will see soon that the phenomenology of this model is already sufficiently rich to give interesting insights on high dimensional Cox regression with elastic net penalization.

#### 1.1 Our contribution

- We accurately characterize the asymptotic performance of the estimator obtained by optimizing the Cox partial likelihood with elastic net regularization in the proportional regime where the number of covariates included in the model p is  $p = \zeta n$ , and the number of active covariates s is  $s = \nu p$ , with n the sample size.
- We provide a sharp characterization of metrics of interest in inference such as the overlap of the inferred associations with the true ones, the mean squared error of the estimator of the associations and of the survival function, but also the fraction of false positive and negative related to the support recovery task.
- We provide a sharp characterization of metrics of interest in prediction such as the concordance-index, which measures the ability of the model to separate yet unseen high risk individuals from low risk ones.

Our investigation reveals that

- when p < n regularization always greatly improve the estimates of the Cox model, in terms
  of their Mean Squared Error;</li>
- when p > n and the number of active components is  $s = \nu p$  and  $\nu = O(1)$ , using an elastic net regularization which is more "lasso-like" does not lead to a significant improvement over the null model, in term of concordance index over a test set. A more "ridge-like" regularization achieves a slightly better prediction performance, at the cost of being less intepretable;

# 2 Setting and notation

Focusing on the case of right censored data, time to event data comprise observations reporting: i) the event indicator  $\Delta$ , equal to 1 if the subject experienced the event and 0 otherwise, ii) the observed time to event T if  $\Delta = 1$ , or the censoring time else, and iii) the covariate vector  $\mathbf{X} = (X_1, \ldots, X_p) \in \mathbb{R}^p$ , i.e. the list of characteristics of the subject. The most widely adopted model to deal with time to event data is the Cox semi-parametric proportional hazards model [14, 11, 6]. If censoring is non-informative, then the likelihood density of the event  $\Delta$  in the time interval [t, t + dt) given covariates  $\mathbf{X}$  and under the proportional hazards assumption reads

$$p(\Delta, t | \mathbf{X}' \boldsymbol{\beta}) = f(t | \mathbf{X}' \boldsymbol{\beta})^{\Delta} S_C(t)^{\Delta} \times S(t | \mathbf{X}' \boldsymbol{\beta})^{1-\Delta} f_c(t)^{1-\Delta} .$$
(1)

Above,  $f_c$  and  $S_C(x) = \int_x^\infty f_c(x) dx$  are , respectively,the density and survival functions for the censoring risk, while

$$S(t|\mathbf{X}'\boldsymbol{\beta}) := \exp\{-\Lambda(t)\exp(\mathbf{X}'\boldsymbol{\beta})\},\tag{2}$$

$$f(t|\mathbf{X}'\boldsymbol{\beta}) := -\frac{\mathrm{d}}{\mathrm{d}t}S(t|\mathbf{X}'\boldsymbol{\beta}) = \lambda(t)\exp\{\mathbf{X}'\boldsymbol{\beta} - \Lambda(t)\exp(\mathbf{X}'\boldsymbol{\beta})\},\tag{3}$$

are the survival (2) and density (3) functions for the primary risk (the risk under investigation). The latter' are parametrized by: i) the association parameters  $\beta \in \mathbb{R}^p$ , ii) the cumulative hazard function  $\Lambda$ . The log-likelihood (per data point) for a data-set of i.i.d censored observations  $\{(\Delta_i, T_i, \mathbf{X}_i)\}_{i=1}^n$ , reads

$$\mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\lambda}) = \frac{1}{n} \sum_{i=1}^n \left\{ \Delta_i \left( \log \boldsymbol{\lambda}(T_i) + \mathbf{X}'_i \boldsymbol{\beta} \right) - \boldsymbol{\Lambda}(T_i) \mathrm{e}^{\mathbf{X}'_i \boldsymbol{\beta}} \right\}$$
(4)

By first maximazing with respect to  $\lambda$  one obtains the so-called Breslow estimator [2, 8]:

$$\hat{\lambda}_n(t,\boldsymbol{\beta}) = \sum_{k=1}^n \frac{\Delta_k \delta(t-t_k)}{\sum_{j=1}^n \Theta(t_j-t_k) \mathrm{e}^{\mathbf{X}'_j \boldsymbol{\beta}}} , \qquad (5)$$

where  $\Theta(x)$  denotes the Heaviside step-function. Substituting in (4) and disregarding terms which do not depend on  $\beta$  it is possible to obtain the logarithm of the Cox partial likelihood per data point

$$\mathcal{PL}_{n}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{n} \left\{ \Delta_{i} \mathbf{X}_{i}^{\prime} \boldsymbol{\beta} - \Delta_{i} \log \left( \frac{1}{n} \sum_{j=1}^{n} \Theta(T_{j} - T_{i}) \mathrm{e}^{\mathbf{X}_{j}^{\prime} \boldsymbol{\beta}} \right) \right\}.$$
(6)

Optimizing the Cox partial likelihood one obtains an estimator  $\hat{\beta}_n$  and, consequently, an estimator for the cumulative hazard function of the primary risk via the Nelson-Aalen estimator

$$\hat{\Lambda}_n(t) = \sum_{i=1}^n \frac{\Delta_i \Theta(t - T_i)}{\sum_{j=1}^n \Theta(T_j - T_i) \mathrm{e}^{\mathbf{X}'_j \hat{\boldsymbol{\beta}}_n}} \tag{7}$$

which is obtained via integration of (5) with respect to t.

In the following we consider applying an elastic net regularization

$$r(\boldsymbol{\beta}) := \alpha \|\boldsymbol{\beta}\| + \frac{1}{2}\eta \|\boldsymbol{\beta}\|^2 \tag{8}$$

to the Cox partial likelihood, i.e. we are interested in the properties of the estimator that maximizes

$$\mathcal{PL}_{n}^{\alpha,\eta}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{n} \left\{ \Delta_{i} \mathbf{X}_{i}^{\prime} \boldsymbol{\beta} - \Delta_{i} \log \left( \frac{1}{n} \sum_{j=1}^{n} \Theta(T_{j} - T_{i}) \mathrm{e}^{\mathbf{X}_{j}^{\prime} \boldsymbol{\beta}} \right) \right\} - \frac{1}{n} r(\boldsymbol{\beta}) .$$
(9)

# **3** Typical behaviour of the penalized (partial) likelihood estimator

In the statistical physics approach to optimization, the optimal value of the objective function (9) is equivalent to minus the zero temperature free energy density of a fictitious physical system, i.e.

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \mathcal{P} \mathcal{L}_n(\hat{\boldsymbol{\beta}}_n) \Big] = \lim_{n \to \infty} \lim_{\gamma \to \infty} \frac{1}{n\gamma} \mathbb{E}_{\mathcal{D}} \Big[ \log \int e^{-\gamma \mathcal{H}_n(\boldsymbol{\beta}|\mathcal{D})} d\boldsymbol{\beta} \Big], \tag{10}$$

with Hamiltonian

$$\mathcal{H}_n(\boldsymbol{\beta}|\mathcal{D}) = \sum_{i=1}^n \Delta_i \Big[ \log \Big( \frac{1}{n} \sum_{j=1}^n \Theta(T_j - T_i) \mathrm{e}^{\mathbf{X}_i' \boldsymbol{\beta}} \Big) - \mathbf{X}_i' \boldsymbol{\beta} \Big] + r(\boldsymbol{\beta})$$
(11)

where we indicated with  $\mathcal{D}$  the data-set, i.e. the set of tuples  $\{\Delta_j, T_j, \mathbf{X}_j\}_{j=1}^n$ . The  $\beta$ s are the degrees of freedom of the system and the data-set  $\mathcal{D} = \{\Delta_j, T_j, \mathbf{X}_j\}_{j=1}^n$  plays the role of the quenched disorder.

We compute the right hand side of (10) via the replica method under the following assumptions over the data generating process: the event indicator and event time are generated as

$$\Delta = \Theta(C - Y), \quad T = \min\{Y, C\}$$
(12)

where  $C \sim f_C$  is the (latent) random variable censoring time and

$$Y|\mathbf{X} \sim f_0(y|\mathbf{X}'\boldsymbol{\beta}_0), \quad \mathbf{X} \sim \mathcal{N}(\mathbf{0}, \frac{1}{p}I_p)$$
 (13)

is the (latent) censoring time. This implies that the true (unknown) generative model is

$$f(\Delta, t | \mathbf{X}' \boldsymbol{\beta}_0) = f_0(t | \mathbf{X}' \boldsymbol{\beta}_0)^{\Delta} S_C(t)^{\Delta} \times S_0(t | \mathbf{X}' \boldsymbol{\beta}_0)^{1-\Delta} f_c(t)^{1-\Delta} .$$
(14)

Notice that we do not require that the model is correctly specified, i.e. we do not assume that the data are generated from a proportional hazard model. The scaling of the covariates is a standard assumption in virtually any previous study of high dimensional regression. The full replica derivation is included in the appendix B as it shares a significant overlap with the derivation of [18] and gives

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \mathcal{P} \mathcal{L}_n(\hat{\boldsymbol{\beta}}_n) \Big] = \underset{w, v, \tau, \hat{w}, \hat{v}, \hat{\tau}}{\operatorname{extr}} \mathcal{P} \mathcal{L}(w, v, \tau, \hat{w}, \hat{v}, \hat{\tau})$$
(15)

with

$$\mathcal{PL}(w, v, \tau, \hat{w}, \hat{v}, \hat{\tau}) = \frac{\zeta}{2\hat{\tau}} \Big( (w - \hat{w})^2 + v^2 + \hat{v}^2 (1 - \tau/\hat{\tau}) \Big) + \zeta \mathbb{E}_{Z,\beta_0} \Big[ \mathcal{M}_{\mathbf{r}(.)} \Big( \hat{w} \frac{\beta_0}{\theta_0} + \hat{v}Z, \hat{\tau} \Big) \Big] - \mathbb{E}_{\Delta,T,Z_0,Q} \Big[ \mathcal{M}_{g(.,\Lambda(T),\Delta)} \Big( wZ_0 + vQ, \tau \Big) \Big] + \text{const} \quad (16)$$

Above:

i)  $\mathcal{M}_{b(.)}$  is the Moureau envelope of a convex function  $b : \mathbb{R}^d \to \mathbb{R}$  [21], which is defined as

$$\mathcal{M}_{b(.)}(\mathbf{x},\alpha) = \min_{\mathbf{z}\in\mathbb{R}^d} \Big\{ \frac{1}{2\alpha} \|\mathbf{z} - \mathbf{x}\|^2 + b(\mathbf{z}) \Big\},\tag{17}$$

and we defined

$$g(x, y, z) := \exp(x)y - zx ; \qquad (18)$$

ii) the random variable  $\beta_0$  follows the law

$$\mathcal{P}_{\beta_0}(x) := \lim_{p \to \infty} \frac{1}{p} \sum_{\mu=1}^p \delta(x - \mathbf{e}'_{\mu} \boldsymbol{\beta}_0); \tag{19}$$

iii)  $Z_0, Q \sim \mathcal{N}(0, 1), Z_0 \perp Q$ , and

$$\Delta, T|Z_0 \sim f(\Delta, t|\theta_0 Z_0), \text{ with } \theta_0 := \|\beta_0\|/\sqrt{p}, \tag{20}$$

where iv) the function  $\Lambda$  is defined at  $w, v, \tau$  as the self consistent solution of

$$\Lambda(t) = \mathbb{E}_{\Delta,T,Z_0,Q}\left[\frac{\Delta\Theta(t-T)}{\mathcal{S}(T)}\right],\tag{21}$$

$$\mathcal{S}(t) = \mathbb{E}_{\Delta,T,Z_0,Q} \Big[ \Theta(T-t) \mathrm{e}^{wZ_0 + vQ + \tau\Delta - W_0 \left(\tau \mathrm{e}^{\tau\Delta + wZ_0 + vQ} \Lambda(T)\right)} \Big], \tag{22}$$

where  $W_0(.)$  is (the real branch of ) the Lambert W-function, which solves  $W_0(x) \exp W_0(x) = x$ , x > -1/e. The point of extremum in (15) satisfies the replica symmetric (RS) equations

$$w = \mathbb{E}_{Z,\beta_0} \left[ \beta_0 \varphi \right] / \theta_0 \tag{23}$$

$$\hat{v}\frac{\tau}{\hat{\tau}} = \mathbb{E}_{Z,\beta_0}\left[Z\varphi\right] \tag{24}$$

$$(w^2 + v^2) = \mathbb{E}_{Z,\beta_0} \left[ \varphi^2 \right]$$
(25)

$$\hat{w} = w - \frac{\hat{\tau}}{\zeta \tau} \left( w - \mathbb{E}_{\Delta, T, Z_0, Q} \left[ Z_0 \xi \right] \right)$$
(26)

$$v(1 - \zeta \tau / \hat{\tau}) = \mathbb{E}_{\Delta, T, Z_0, Q} \Big[ Q \xi \Big]$$
(27)

$$\zeta \hat{v}^2 = \frac{\hat{\tau}^2}{\tau^2} \mathbb{E}_{\Delta, T, Z_0, Q} \Big[ (\xi - wZ_0 - vQ)^2 \Big]$$
(28)

where

$$\xi := \operatorname{prox}_{g(.,\Lambda(T),\Delta)}(wZ_0 + vQ,\tau) = wZ_0 + vQ + \tau\Delta - W_0\Big(\tau e^{\tau\Delta + wZ_0 + vQ}\Lambda(T)\Big)$$
(29)

$$\varphi := \operatorname{prox}_{r(.)}(\hat{w}\beta_0/\theta_0 + \hat{v}Z, \alpha\hat{\tau}) = \frac{1}{1+\eta\hat{\tau}}\operatorname{st}(\hat{w}\beta_0 + \hat{v}Z, \alpha\hat{\tau})$$
(30)

with  $\operatorname{st}(x,\omega) := \Theta(x - |\omega|) (x - |\omega|\operatorname{sgn}(x))$  the soft thresholding operator.

#### 3.1 Interpretation

We show in appendix E that

$$\mathcal{P}(\Delta, t, h) := \lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \frac{1}{n} \sum_{i=1}^{n} \delta(t - T_{i}) \delta_{\Delta, \Delta_{i}} \delta(x - \mathbf{X}_{i}' \hat{\boldsymbol{\beta}}) \Big]$$
(31)

$$\mathcal{P}_{\varphi}(x) := \lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \left[ \frac{1}{p} \sum_{k=1}^{p} \delta(x - \mathbf{e}'_{k} \hat{\boldsymbol{\beta}}) \right].$$
(32)

admit the following expressions in the replica symmetric ansatze

$$\mathcal{P}_{\xi}(\Delta, t, h) = \mathbb{E}_{\Delta', T', Z'_{0}, Q'} \left[ \delta(t - T') \delta_{\Delta, \Delta'} \delta(h - \xi_{\star}) \right]$$
(33)

$$\mathcal{P}_{\varphi}(x) = \mathbb{E}_{\beta_0, Z} \Big[ \delta \big( x - \varphi_{\star} \big) \Big]$$
(34)

where  $\xi_*, \varphi_*$  are as in (29,30), computed at the fixed point of the RS equations, i.e. at  $w_*, v_*, \tau_*, \hat{w}_*, \hat{v}_*, \hat{\tau}_*$  and  $\Lambda_*(.)$  is the function solving (21, 22) evaluated at the fixed point. Hence we might regard  $\xi$  (29) and  $\varphi$  (30), as "probabilistic equivalents" for the entries of: i) the linear predictors  $\mathbf{X}'_i \hat{\boldsymbol{\beta}}$  and ii) the estimator  $\mathbf{e}'_{\mu} \hat{\boldsymbol{\beta}}$ . What we mean by probabilistic equivalent is that

$$\lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \frac{1}{n} \sum_{i=1}^{n} \ell_1(\mathbf{X}'_i \hat{\boldsymbol{\beta}}_n, \hat{\boldsymbol{\Lambda}}_n(T_i), \boldsymbol{\Delta}_i, T_i) \Big] = \sum_{\boldsymbol{\Delta} \in \{1,0\}} \int \mathcal{P}_{\boldsymbol{\xi}}(\boldsymbol{\Delta}, t, h) \ell_1(h, \boldsymbol{\Lambda}_{\star}(t), \boldsymbol{\Delta}, t) \mathrm{d}t \mathrm{d}h,$$
(35)

$$\lim_{p \to \infty} \mathbb{E}_{\mathcal{D}} \left[ \frac{1}{p} \sum_{\mu=1}^{p} \ell_2(\mathbf{e}'_{\mu} \hat{\boldsymbol{\beta}}_n) \right] = \int \mathcal{P}_{\varphi}(x) \ell_2(x) \, \mathrm{d}x \tag{36}$$

for any "reasonable" functions  $\ell_1, \ell_2$ , where  $\mathcal{D}$  is a short hand notation for the data-set  $\{(\Delta_1, T_1, \mathbf{X}_1), \dots, (\Delta_n, T_n, \mathbf{X}_n)\}$ . In turn (34) provides the following interpretation of the values  $w_{\star}, v_{\star}$ 

$$\hat{w}_n := \frac{\beta'_0 \beta_n}{\sqrt{\beta'_0 \beta_0}} \underset{n \to \infty}{\approx} w_\star, \quad \hat{v}_n^2 := \hat{\beta}'_n \hat{\beta}_n - w_n^2 \underset{n \to \infty}{\approx} v_\star^2.$$
(37)

Similarly (33) implies that the Replica Symmetric functional order parameter  $\Lambda(.)$  is equivalent to the Nelson-Aalen estimator computed over a large (ideally infinite) data-set, i.e.

$$\hat{\Lambda}_n(.) \underset{n \to \infty}{\approx} \Lambda_\star(.) . \tag{38}$$

In summary one can compute virtually any metric of interest over the training set via (34, 33). This is useful to understand how regularization impacts inference and prediction in the proportional regime. In particular what is the interplay of regularization with the sparsity of the underlying  $\beta_0$ . This will be the topic of the next sections.

#### 4 Simulation protocol

In sections (5, 6) we compare the solution of the RS equations (21, 22, 23, 24, 25, 26, 27, 28) with the corresponding finite sample size metrics computed from simulated data. Here we detail on the simulation protocol. For comparison purposes, it is handy to reparametrize the penalization via  $\lambda$  and  $\rho$  as follows

$$\alpha = \lambda \rho, \ \eta = \lambda (1 - \rho), \quad \text{with } \rho \in (0, 1] .$$
(39)

In all the simulations the true associations are generated as follows

$$\boldsymbol{\beta}_{0} = \left(\theta_{0} \sqrt{p} \, \boldsymbol{U}_{s}, \boldsymbol{0}_{p-s}\right), \quad \boldsymbol{U}_{s} \sim \operatorname{Unif}(\mathbb{S}_{s-1}) \,. \tag{40}$$

When  $p,s \rightarrow \infty$  we have that

$$\mathcal{P}_{\beta_0}(x) = \nu \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{1}{2}x^2/\sigma^2} + (1-\nu)\delta(x), \quad \nu = s/p \in [0,1], \quad \sigma = \theta_0/\sqrt{\nu}.$$
(41)

where  $\mathcal{P}_{\beta_0}$  is the limiting empirical distribution of the entries of  $\beta_0$ , defined in (19). The sparsity of the signal is controlled by  $\nu$  which is the fraction of "active" covariates that is actually correlated with the outcome. Under this assumption, the replica symmetric equations can be simplified further, to a somewhat simpler form

$$w = 2\frac{1}{1+\tilde{\eta}\hat{\tau}}\hat{w}\Phi(\chi_1)$$
(42)

$$\tau = 2\frac{1}{1+\tilde{\eta}\hat{\tau}}\hat{\tau}\left\{\nu\Phi(\chi_1) + (1-\nu)\Phi(\chi_0)\right\}$$
(43)

$$\frac{1}{2}(v^{2}+w^{2}) = \frac{1}{(1+\tilde{\eta}\hat{\tau})^{2}} \Big\{ \nu \Big( (1+1/\chi_{1}^{2})\Phi(\chi_{1}) - \phi(\chi_{1}) \Big) + (1-\nu) \Big( (1+1/\chi_{0}^{2})\Phi(\chi_{0}) - \phi(\chi_{0}) \Big) \Big\}$$
(44)

$$\hat{w} = w - \frac{\hat{\tau}}{\zeta \tau} \left( w - \mathbb{E}_{\Delta, T, Z_0, Q} \left[ Z_0 \xi \right] \right)$$
(45)

$$v(1 - \zeta \tau / \hat{\tau}) = \mathbb{E}_{\Delta, T, Z_0, Q} \Big[ Q \xi \Big]$$
(46)

$$\zeta \hat{v}^{2} = \frac{\hat{\tau}^{2}}{\tau^{2}} \mathbb{E}_{\Delta,T,Z_{0},Q} \Big[ (\xi - wZ_{0} - vQ)^{2} \Big]$$
(47)

where  $\xi$  is defined in (29),

$$\chi_0 := \frac{\alpha \hat{\tau}}{\hat{v}}, \quad \chi_1 := \frac{\alpha \hat{\tau}}{\sqrt{\hat{v}^2 + \hat{w}^2/\nu}},\tag{48}$$

and we indicated with  $\phi(x) := \exp\{-\frac{1}{2}x^2\}/\sqrt{2\pi}$  the density function and with  $\Phi(x) := \int_x^{+\infty} \phi(t) dt$  the complementary distribution function of a standard normal random variable. The numerical solution of these self-consistent equations is obtained via fixed point iteration. The integral are computed as averages of populations of size  $10^5$ .

In the next sections, the latent survival data are generated from a Log-logistic proportional hazard model

$$Y|\mathbf{X} \sim -\frac{\mathrm{d}}{\mathrm{d}t}S_0(t|\mathbf{X}), \quad S_0(t|\mathbf{X}) = \exp\{-\Lambda_0(.)\exp(\mathbf{X}'\boldsymbol{\beta}_0)\}, \ \Lambda_0(t) := \log\left(1 + \mathrm{e}_0^{\phi}t^{\rho_0}\right)$$
(49)

the censoring is uniform in the interval  $[\tau_1, \tau_2]$ , i.e.

$$C \sim \operatorname{Unif}[\tau_1, \tau_2] \tag{50}$$

and the observations  $\Delta, T$  are obtained as

$$\Delta = \mathbf{1}[Y < C], \quad Y = \min\{Y, C\}.$$
(51)

In this case the Cox model is well specified, hence it does make sense to compare what is inferred to the truth (which is available in simulations). To make the setting above concrete, we choose:

- $\theta_0 = 1, \phi_0 = -\log 2, \rho_0 = 2$  and  $\tau_1 = 2.0, \tau_2 = 3.0$ , so that the approximately 3 observations out of 10 are censored;
- $\nu = 0.3$ , so that only 3 covariates on 10 are actually associated with the survival time;
- $\rho = 0.75, 0.25$ , so that the behaviour of the elastic net regularization is more skewed toward, respectively, the lasso and ridge regularizations.

The quantities computed from simulations are always displayed as dots, indicating the averages over 100 repetitions, with errorbars, the corresponding finite sample standard deviations. The algorithm used to compute the path-wise solution for the elastic net estimator is a variant of the one used in [27] and is described in (F).

# 5 The role of regularization in inference

In this section we investigate the performance of the penalized partial likelihood estimator (PMLE) as a function of the regularization strength and of the ratio  $\zeta = p/n$ . The parameters that are estimated




Figure 1: Regression over synthetic data versus RS theory, left column  $\rho = 0.75$  (lasso -like) and right column  $\rho = 0.25$  (ridge like). Upper row:  $\hat{\kappa}_n := \hat{\beta}'_n \beta_0 / ||\beta_0||^2$ , lower row : sparsity of the estimated  $\hat{\beta}_n$  with threshold fixed at  $1.0 \cdot e^{-3}$ , against the regularization strength  $\lambda$  and for several values of  $\zeta = p/n$ .

in a Cox model are: i) the associations  $\beta$ , which are then used e.g. to compute risk scores and ii) the cumulative hazard  $\Lambda$  which is needed to estimate the survival function. In order to study the performance of  $\hat{\beta}_n$  we define the following metrics:

- $\hat{\kappa}_n := \frac{\hat{\beta}'_n \beta_0}{\|\beta_0\|^2}$ , which is the ratio between the component of  $\hat{\beta}_n$  along the direction of  $\beta_0$  normalized by the norm of  $\beta_0$ . This quantity is 1 when  $\hat{\beta}_n = \beta_0$  and 0 if  $\hat{\beta}_n \perp \beta_0$ ;
- RMSE(β̂<sub>n</sub>) := ||β̂<sub>n</sub> − β<sub>0</sub>||<sup>2</sup>/||β<sub>0</sub>||<sup>2</sup>, which is the ratio between the squared distance of β̂<sub>n</sub> from β<sub>0</sub> and the squared length of β<sub>0</sub>. This quantity is positive, it equals 1 when β̂<sub>n</sub> = 0, i.e. for large values of λ, and 0 when β̂<sub>n</sub> = β<sub>0</sub>.

The asymptotic value of these quantities can be computed starting from the solution of the replica symmetric equations and is

$$\hat{\kappa}_n \underset{n \to \infty}{\approx} w_\star / \theta_0, \quad \text{RMSE}(\hat{\beta}_n) \underset{n \to \infty}{\approx} (v_\star^2 + (w_\star - \theta_0)^2) / \theta_0^2$$
 (52)

For what concern the estimated survival function

$$\hat{S}_n(t|\mathbf{x}'\hat{\boldsymbol{\beta}}_n) := \exp\{-\hat{\Lambda}_n(t)\exp(\mathbf{x}'\hat{\boldsymbol{\beta}}_n)\},\tag{53}$$

we might consider

$$MSE(\hat{S}_n) := \frac{1}{n} \sum_{i=1}^n \left( \hat{S}_n(T_i, \mathbf{X}'_i \hat{\boldsymbol{\beta}}_n) - S_0(T_1, \mathbf{X}' \boldsymbol{\beta}_0) \right)^2,$$
(54)

which is the squared euclidian distance of the inferred survival function  $\hat{S}_n$  from the true one  $S_0$  averaged over the training data set. This quantity is 0 if the true and inferred survival curves coincide




Figure 2: Regression over synthetic data versus RS theory, left column  $\rho = 0.75$  (lasso -like) and right column  $\rho = 0.25$  (ridge like). Upper row:  $\text{RMSE}(\hat{\beta}_n) := \|\hat{\beta}_n - \beta_0\|^2 / \|\beta_0\|^2$ , lower row :  $\text{RMSE}(\hat{S}_n)$  defined in (55), against the regularization strength  $\lambda$  and for several values of  $\zeta = p/n$ .

at each training point and strictly positive otherwise. However, the MSE (54) can only be used to compared different models, hence we prefer to use

$$RMSE(\hat{S}_n) := \frac{MSE(\hat{S}_n)}{MSE(\hat{S}_{null})}$$
(55)

where  $\hat{S}_n$ ,  $\hat{S}_{null}$  are, respectively, the predicted survival functions when all the covariates and no covariates are included in the model. This metric has value 1 if the inferred model gives on average the same prediction as the null model over the training set (the simple model where no covariates are included), and value 0 is the inferred model is "perfect", i.e. has an MSE (54) equal to zero. The asymptotic value of  $MSE(\hat{S}_n)$  can be obtained via the solution of the replica symmetric equations

$$\operatorname{MSE}(\hat{S}_n) \underset{n \to \infty}{\approx} \mathbb{E}_{\Delta, T, Z_0, Q} \Big[ \Big( \exp\{-\Lambda_{\star}(T) \exp(\xi_{\star})\} - \exp\{-\Lambda_0(T) \exp(\theta_0 Z_0)\} \Big)^2 \Big]$$
(56)

and similarly we have

$$\operatorname{MSE}(\hat{S}_{null}) \approx_{n \to \infty} \mathbb{E}_{\Delta, T, Z_0, Q} \left[ \left( \exp\{-\Lambda_{null}(T)\} - \exp\{-\Lambda_0(T)\exp(\theta_0 Z_0)\} \right)^2 \right]$$
(57)

with  $\Lambda_{null}(T)$  defined as the solution of (21, 22) at  $w, v, \tau = 0$ .

The solution of the RS equations can also be used to compute the sparsity of  $\hat{\beta}_n$ , i.e. the fraction of strictly non-zero entries of  $\hat{\beta}_n$ . If we define the estimated support as

$$\widehat{\operatorname{supp}}_{n}(\epsilon) := \left\{ \mu \in \{1, \dots, p\} : |\mathbf{e}'_{\mu} \hat{\boldsymbol{\beta}}_{n}| > \epsilon \right\},$$
(58)

then the sparsity of  $\hat{\beta}_n$  is  $\lim_{\epsilon \to 0^+} \widehat{\operatorname{supp}}_n(\epsilon)$ . This quantity might be used to estimate  $\operatorname{supp}(\beta_0) = \{\mu \in \{1, \dots, p\} : |\mathbf{e}'_{\mu}\beta_0| > 0\}$ , i.e. the support of the true association vector. Chance is that

the estimated support might differ from the actual one and we can quantify the uncertainty in this estimate via:

i ) the expected fraction of false negatives, i.e. the fraction of components incorrectly excluded from the estimated support, is

$$FN(\epsilon) = \lim_{p \to \infty} \mathbb{E}_{\mathcal{D}} \left[ \frac{1}{p} \sum_{\mu=1}^{P} \boldsymbol{I} \left( \mu \notin \widehat{supp}_{n}(\epsilon) \right) \boldsymbol{I} \left( \mu \in \operatorname{supp}(\boldsymbol{\beta}_{0}) \right) \right] = \mathbb{E}_{\boldsymbol{\beta}_{0}, Z} \left[ \boldsymbol{I}(|\boldsymbol{\varphi}| < \epsilon) \boldsymbol{I}(|\boldsymbol{\beta}_{0}| > 0) \right] = \\ = \nu \left[ 1 - 2\Phi \left( \chi_{1}^{\star} \left\{ 1 + \epsilon \frac{1 + \eta \hat{\tau}}{\alpha \hat{\tau}} \right\} \right) \right];$$
(59)

ii) the expected fraction of false positives, i.e. the fraction of components incorrectly included in the estimated support, is

$$FP(\epsilon) = \lim_{p \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \frac{1}{p} \sum_{\mu=1}^{P} \boldsymbol{I} \Big( \mu \in \widehat{supp}_{n}(\epsilon) \Big) \boldsymbol{I} \Big( \mu \notin \operatorname{supp}(\boldsymbol{\beta}_{0}) \Big) \Big] = \mathbb{E}_{\boldsymbol{\beta}_{0}, Z} \Big[ \boldsymbol{I}(|\boldsymbol{\varphi}| > \epsilon) \boldsymbol{I}(|\boldsymbol{\beta}_{0}| = 0) \Big] = \\ = (1 - \nu) \Big[ 2\Phi \Big( \chi_{0}^{\star} \Big\{ 1 + \epsilon \frac{1 + \eta \hat{\tau}}{\alpha \hat{\tau}} \Big\} \Big) \Big].$$
(60)

In figure (2) we see that for small values of  $\zeta$  (e.g.  $\zeta = 0.2, 0.5, 0.8$ ), the optimal (smallest) value of the ratios  $\text{RMSE}(\hat{\beta}_n)$ ,  $\text{RMSE}(\hat{S}_n)$  is much lower than 1 (the dashed black line), where the MSE is equal to the one of the null model. However, as  $\zeta$  grows larger at fixed  $\nu = 0.3$ , the ratio of MSEs gets closer to 1. This indicates that even an optimally tuned elastic net penalization might not improve significantly the estimation accuracy when  $p \gg n$  and s is a finite fraction of p, in terms of MSE. Furthermore, as  $\zeta$  increases the curves  $\hat{k}_n$  against  $\zeta$ , upper row in (1), get more and more steep, i.e. the overlap of  $\hat{\beta}_n$  with the truth  $\beta_0$  decreases to zero as  $\lambda$  increases faster for larger  $\zeta$ . This means that at the optimal  $\lambda$  (in terms of MSE), the bias of  $\hat{\beta}_n$  due to the shrinking effect of the regularization increases with  $\zeta$ . This is due to the fact that the optimal elastic net penalization produces progressively sparser solutions as  $\zeta$  increases, as one can deduce from the lower row of fig (1). This implies that as  $\zeta$  increases at fixed  $\lambda$ , the fraction of false negative must increase, which is in agreement with fig (3), where we display the fraction of false negatives in the lower row. The upper row in fig(3) shows that the fraction of false positives decreases with  $\zeta$  at fixed  $\lambda$ , and this is again in agreement with our previous considerations: as  $\zeta$  increases the elastic net penalization shrinks all the inferred associations towards the origin, at the point that the fitted model gets very close to the one where no covariates are included (which gives a false positive fraction equal to 0).

# 6 The role of regularization in prediction

Discrimination is the ability of the model to separate (unseen) individuals with different risks scores. The rationale being that a good model should assign shorter survival times to subjects with higher risk score [23]. The disciminative ability of a model is usually evaluated via (variants of) the concordance index (c-index) [16]. One of the most used c-index estimators is Harrell's one [12], defined for a sample of  $n_{test}$  unseen subjects as follows

$$\widehat{HC}_{n_{test}} = \frac{\sum_{i=1}^{n_{test}} \Delta_i \sum_{j=1}^{n_{test}} \Theta(T_j - T_i) \mathbf{1} \Big[ \mathbf{X}'_j \hat{\boldsymbol{\beta}}_n > \mathbf{X}'_i \hat{\boldsymbol{\beta}}_n \Big]}{\sum_{i=1}^{n_{test}} \Delta_i \sum_{j=1}^{n_{test}} \Theta(T_j - T_i)}$$

This quantity is 1 if the model has perfect discrimination ability. Random guessing gives a c-index of 0.5. Notice that for a "fresh" covariates vector  $\tilde{\mathbf{X}} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p/p)$  it holds that

$$\tilde{\mathbf{X}}'\hat{\boldsymbol{\beta}}_n = \frac{\mathbf{X}'\boldsymbol{\beta}_0}{\|\boldsymbol{\beta}_0\|}\hat{w}_n + \tilde{\mathbf{X}}'\big(\boldsymbol{I} - \frac{\boldsymbol{\beta}_0\boldsymbol{\beta}'_0}{\|\boldsymbol{\beta}_0\|^2}\big)\hat{\boldsymbol{\beta}}_n \stackrel{d}{=} \hat{w}_n Z_0 + \hat{v}_n Q \underset{n \to \infty}{\approx} w_\star Z_0 + v_\star Q, \tag{61}$$

with  $Z_0, Q \sim \mathcal{N}(0, 1), Z_0 \perp Q$  and where the last approximation is because of (37). Hence we can compute the expected value of (61) as

$$\widehat{HC}_{n_{test}} \underset{n_{test}, n \to \infty}{\approx} \frac{\mathbb{E}_{\Delta, T, Z_0, Q} \left[ \Delta \mathbb{E}_{\Delta', T', Z'_0, Q'} \left[ \Theta(T' - T) \mathbf{1}(w_{\star} Z'_0 + v_{\star} Q' > w_{\star} Z_0 + v_{\star} Q) \right] \right]}{\mathbb{E}_{\Delta, T, Z_0, Q} \left[ \Delta \mathbb{E}_{\Delta', T', Z'_0, Q'} \left[ \Theta(T' - T) \right] \right]}.$$
(62)




Figure 3: Regression over synthetic data versus RS theory, **left column**  $\rho = 0.75$  (lasso -like) and **right column**  $\rho = 0.25$  (ridge like). **Upper row**: fraction of false positives (left), **lower row** : fraction of false negatives (right), against the regularization strength  $\lambda$  and for several values of  $\zeta = p/n$ .

As shown in figure (4), the optimal (highest) value of the Harrel's c-index descreases as  $\zeta$  increases at fixed  $\lambda$ . This is consistent with our previous findings, as for large  $\zeta$  the optimal  $\hat{\beta}_n$  in terms of MSE will be close to the null vector, hence the c-index will be close to that of the null model, i.e. the value 0.5. We also notice that when the elastic net penalization is more skewed toward the lasso ( $\rho = 0.25$ ), the test c-index becomes almost independent from  $\lambda$ : this can be deduced from the fact that the curves on the right column in (4) are almost flat.

# 7 Conclusion

In conclusion, we have accurately characterized the behaviour of the estimator obtained by optimization of the log-partial likelihood with an elastic net penalization when the sample size n, the number of covariates p and the number of active components s diverge proportionally. This is done via the solution of a system of six non-linear equations, which can be obtained via fixed point iteration. We deduce from our theory, which is supported by extensive numerical simulations, that the optimal value of the MSE of the estimator of the associations  $\hat{\beta}_n$  is non-vanishing, and increases as  $\zeta$  increases and the fraction of active covariates  $\nu$  is kept constant. When  $\zeta$  is "small", i.e. less than 1, the latter is significantly lower than that of the null model, but the difference gets smaller as  $\zeta$  increases. This means that, when  $p \gg n$  even an optimally tuned model might only marginally improve over the null model (where no covariates are included).

A strong limitation of the present theory is the assumption of i.i.d. Gaussian covariates with small variance  $\mathbf{X} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p/p)$ . Even though such requirement can be relaxed (as discussed in the introduction), we must have that  $\mathbf{X}'\boldsymbol{\beta}$  is well approximated by a Gaussian distribution (when  $\boldsymbol{\beta}$  is in a compact subset of  $\mathbb{R}^p$ ) for our calculation to go through.




Figure 4: Regression over synthetic data versus RS theory, left column  $\rho = 0.75$  (lasso -like) and right column  $\rho = 0.25$  (ridge like). The test Harrel's concordance index  $\hat{HC}_{n_{test}}$  against the regularization strength  $\lambda$  and for several values of  $\zeta = p/n$ .

In principle the theory developed in this manuscript can be applied to study other separable penalties. It would be interesting to study how folded concave penalties, e.g. the Smoothly Clipped Absolute Deviation (SCAD) regularization, or the  $L_q$  norm regularization affect the estimation. This will be object of future investigations. Likewise the present theoretical machinery can be directly applied to study mis-specification, and in particular what happens when the proportional hazards assumption is violated. It is also potentially interesting, at this stage, to understand what is the impact of the correlations among covariates over regularized estimation and prediction. In this regard the simplest setting to be studied is with ridge regularization, since in that case the RS equations depend on the spectrum of the (population) correlation matrix (because of the rotational invariance of the ridge penalization), and not on the distribution of the eigenvectors (as would be the case for the lasso). This is currently under investigation from the authors.

# References

- J Bradic, J Fan, and J Jiang. "Regularization for Cox's Proportional hazards model with np-dimensionality". In: *The Annals of Statistics* 39.6 (2011), pp. 3092–3120. ISSN: 00905364, 21688966. URL: http://www.jstor.org/stable/41713609 (visited on 05/02/2024).
- [2] NE Breslow. "Discussion on Professor Cox's Paper". In: Journal of the Royal Statistical Society: Series B (Methodological) 34.2 (1972), pp. 202–220. DOI: https://doi.org/ 10.1111/j.2517-6161.1972.tb00900.x.eprint: https://rss.onlinelibrary. wiley.com/doi/pdf/10.1111/j.2517-6161.1972.tb00900.x.URL: https://rss. onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1972.tb00900.x.
- [3] P Bühlmann and S van de Geer. *Statistics for High-Dimensional Data: Methods, Theory and Applications*. Springer Series in Statistics. Springer Berlin Heidelberg, 2011. ISBN: 9783642201929.
- [4] ACC Coolen et al. "Replica analysis of overfitting in generalized linear regression models". In: *Journal of Physics A: Mathematical and Theoretical* 53.36 (2020), p. 365001.
- [5] ACC Coolen et al. "Replica analysis of overfitting in regression models for time-to-event data". In: Journal of Physics A: Mathematical and Theoretical 50.37 (Aug. 2017), p. 375001. DOI: 10.1088/1751-8121/aa812f. URL: https://dx.doi.org/10.1088/1751-8121/aa812f.
- [6] DR Cox. "Regression Models and Life-Tables". In: Journal of the Royal Statistical Society. Series B (Methodological) 34.2 (1972), pp. 187–220. ISSN: 00359246. URL: http://www.jstor.org/stable/2985181 (visited on 05/07/2024).
- [7] D Donoho and A Montanari. "High dimensional robust M-estimation: asymptotic variance via approximate message passing". In: *Probability Theory and Related Fields* 166 (2013), pp. 935–969. URL: https://api.semanticscholar.org/CorpusID:10859430.

- [8] M Friedman. "Piecewise Exponential Models for Survival Data with Covariates". In: *The Annals of Statistics* 10.1 (1982), pp. 101–113. DOI: 10.1214/aos/1176345693. URL: https://doi.org/10.1214/aos/1176345693.
- [9] E Gardner and B Derrida. "Optimal storage properties of neural network models". In: Journal of Physics A: Mathematical and General 21.1 (Jan. 1988), p. 271. DOI: 10.1088/0305-4470/21/1/031. URL: https://dx.doi.org/10.1088/0305-4470/21/1/031.
- [10] SA van de Geer. "High-dimensional generalized linear models and the lasso". In: *The Annals of Statistics* 36.2 (2008), pp. 614–645. DOI: 10.1214/009053607000000929. URL: https://doi.org/10.1214/00905360700000929.
- [11] FE Harrell. Regression Modeling Strategies: With Applications to Linear Models, Logistic Regression, and Survival Analysis. Graduate Texts in Mathematics. Springer, 2001. ISBN: 9780387952321.
- [12] FE Jr Harrell et al. "Evaluating the yield of medical tests". In: (1982), pp. 1799–1823. DOI: 10.1214/23-A0S2309.
- [13] Y Kabashima, T Wadayama, and T Tanaka. "A typical reconstruction limit for compressed sensing based on Lp-norm minimization". In: *Journal of Statistical Mechanics: Theory and Experiment* 2009.09 (Sept. 2009), p. L09003. DOI: 10.1088/1742-5468/2009/09/L09003. URL: https://dx.doi.org/10.1088/1742-5468/2009/09/L09003.
- [14] JD Kalbfleisch and RL Prentice. *The Statistical Analysis of Failure Time Data*. Wiley Series in Probability and Statistics. Wiley, 2011. ISBN: 9781118031230.
- [15] S Kong and B Nan. "Non-asymptotic oracle inequalities for the high-dimensional Cox regression via Lasso". In: *Statistica Sinica* 24.1 (2014), p. 25.
- [16] E Longato, M Vettoretti, and B Di Camillo. "A practical perspective on the concordance index for the evaluation and selection of prognostic time-to-event models". In: *Journal of biomedical informatics* (Aug. 2020). DOI: 10.1088/0305-4470/21/1/031. URL: https: //dx.doi.org/10.1088/0305-4470/21/1/031.
- B Loureiro et al. "Learning curves of generic features maps for realistic datasets with a teacher-student model\*". In: *Journal of Statistical Mechanics: Theory and Experiment* 2022.11 (Nov. 2022), p. 114001. DOI: 10.1088/1742-5468/ac9825. URL: https://dx.doi.org/10. 1088/1742-5468/ac9825.
- [18] E Massa, A Mozeika, and A C C Coolen. "Replica analysis of overfitting in regression models for time to event data: the impact of censoring". In: *Journal of Physics A: Mathematical and Theoretical* 57.12 (Mar. 2024), p. 125003. DOI: 10.1088/1751-8121/ad2e40. URL: https://dx.doi.org/10.1088/1751-8121/ad2e40.
- [19] L Miolane and A Montanari. "The distribution of the Lasso: Uniform control over sparse balls and adaptive parameter tuning". In: *The Annals of Statistics* 49.4 (2021), pp. 2313–2335. DOI: 10.1214/20-A0S2038. URL: https://doi.org/10.1214/20-A0S2038.
- [20] K Okajima et al. "Average case analysis of Lasso under ultra-sparse conditions". In: International Conference on Artificial Intelligence and Statistics. 2023. URL: https://api. semanticscholar.org/CorpusID:257219909.
- [21] RT Rockafellar. *Convex Analysis*. Princeton Landmarks in Mathematics and Physics. Princeton University Press, 1997. ISBN: 9780691015866.
- [22] F Salehi, E Abbasi, and B Hassibi. "The impact of regularization on high-dimensional logistic regression". In: *Proceedings of the 33rd International Conference on Neural Information Processing Systems*. Red Hook, NY, USA: Curran Associates Inc., 2019.
- [23] EW Steyerberg et al. "Assessing the performance of prediction models: a framework for traditional and novel measures." In: *Epidemiology (Cambridge, Mass.)* (2010). DOI: 10.1097/EDE.0b013e3181c30fb2.
- [24] C Thrampoulidis, E Abbasi, and B Hassibi. "Precise Error Analysis of Regularized *M* Estimators in High Dimensions". In: *IEEE Transactions on Information Theory* 64.8 (2018), pp. 5592–5628. DOI: 10.1109/TIT.2018.2840720.
- [25] C Thrampoulidis, S Oymak, and B Hassibi. *The Gaussian min-max theorem in the Presence of Convexity*. 2015. arXiv: 1408.4837 [cs.IT].
- [26] R Tibshirani. "Regression Shrinkage and Selection via the Lasso". In: Journal of the Royal Statistical Society. Series B (Methodological) 58.1 (1996), pp. 267–288. ISSN: 00359246. URL: http://www.jstor.org/stable/2346178 (visited on 05/01/2024).

- [27] H Zou and T Hastie. "Regularization and Variable Selection via the Elastic Net". In: Journal of the Royal Statistical Society. Series B (Statistical Methodology) 67.2 (2005), pp. 301–320. ISSN: 13697412, 14679868. URL: http://www.jstor.org/stable/3647580 (visited on 05/01/2024).
- [28] H Zou and HH Zhang. "On the adaptive elastic-net with a diverging number of parameters". In: Annals of statistics 37 4 (2009), pp. 1733–1751. URL: https://api.semanticscholar.org/CorpusID:7605822.

# A Appendix / supplemental material

# **B** Replica derivation

The penalized Cox model is defined by the following energy function

$$\mathcal{H}_{n}(\boldsymbol{\beta}|\mathcal{D}) = \sum_{i=1}^{n} \Delta_{i} \left[ \log \left( \frac{1}{n} \sum_{j=1}^{n} \Theta(T_{j} - T_{i}) \mathrm{e}^{\mathbf{X}_{i}^{\prime} \boldsymbol{\beta}} \right) - \mathbf{X}_{i}^{\prime} \boldsymbol{\beta} \right] + r(\boldsymbol{\beta})$$
(63)

where we indicate with r the elastic net penalization

$$r(\mathbf{x}) = \frac{1}{2}\eta \|\mathbf{x}\|^2 + \alpha |\mathbf{x}| .$$
(64)

Equivalently we can regard the energy as a functional of the empirical distribution

$$P_n(\Delta, t, h | \boldsymbol{\beta}, \mathcal{D}) = \frac{1}{n} \sum_{i=1}^n \delta(t - T_i) \delta_{\Delta, \Delta_i} \delta(h - \mathbf{X}'_i \boldsymbol{\beta})$$
(65)

where we indicated with  $\mathcal{D}$  the data-set, i.e. the set of couples  $\{T_j, \mathbf{X}_j\}_{j=1}^n$ , which plays here the role of the disorder. Explicitly

$$\mathcal{H}\Big[P_n(.|\boldsymbol{\beta}, \mathcal{D})\Big] = n\mathcal{E}\Big[P_n(.|\boldsymbol{\beta}, \mathcal{D})\Big] + r(\boldsymbol{\beta}), \tag{66}$$

$$\mathcal{E}\Big[P_n(.|\boldsymbol{\beta}, \mathcal{D})\Big] = \sum_{\Delta=\pm 1} \int \Delta\Big(\log S_n(t|\boldsymbol{\beta}, \mathcal{D}) - h\Big) P_n(\Delta, t, h|\boldsymbol{\beta}, \mathcal{D}) \mathrm{d}h \mathrm{d}t, \qquad (67)$$

$$S_n(t|\boldsymbol{\beta}, \mathcal{D}) = \sum_{\Delta'=\pm 1} \int \Theta(t-t') \mathrm{e}^{h'} P_n(\Delta', t, h'|\boldsymbol{\beta}, \mathcal{D}) \mathrm{d}h' \mathrm{d}t' \,.$$
(68)

It is well known that the information content of inference is quantified by the free energy

$$f(\gamma) := -\lim_{n \to \infty} \frac{1}{n\gamma} \mathbb{E}_{\mathcal{D}} \Big[ \log Z_n(\gamma, \mathcal{D}) \Big], \qquad Z_n(\gamma) := \int e^{-\gamma \mathcal{H}[P_n(.|\beta, \mathcal{D})]} d\beta$$
(69)

To compute the average of the logarithm we use the replica trick

$$f(\gamma) = \lim_{n \to \infty} \lim_{r \to 0} f_n^{(r)}(\gamma), \qquad f_n^{(r)}(\gamma) := -\frac{1}{nr} \log \mathbb{E}_{\mathcal{D}} \Big[ Z_n^r(\gamma, \mathcal{D}) \Big] . \tag{70}$$

We refer to  $f_n^{(r)}(\gamma)$  as the replicated free energy. We now compute this quantity.

#### **B.1** The replicated free energy

In order to compute

$$\mathbb{E}_{\mathcal{D}}\Big[Z_n^r(\gamma, \mathcal{D})\Big] = \int e^{-\gamma \sum_{\alpha=1}^r \mathcal{H}[P_n(.|\boldsymbol{\beta}_{\alpha}, \mathcal{D})]} \prod_{\alpha=1}^r d\boldsymbol{\beta}_{\alpha} , \qquad (71)$$

We first write  $Z_n(\gamma, D)$  in a more convenient form. Let us introduce the functional delta measure

$$\delta \Big[ \mathcal{P}(.) - P_n(.|\boldsymbol{\beta}_{\alpha}, \mathcal{D}) \Big] \propto \int e^{in \left\{ \sum_{\Delta = \pm 1} \int \hat{\mathcal{P}}(\Delta, t, h) \left[ \mathcal{P}(\Delta, t, h) - P_n(\Delta, t, h|\boldsymbol{\beta}, \mathcal{D}) \right] dt dh \right\}} \mathcal{D} \hat{\mathcal{P}} = \int e^{in \left\{ \sum_{\Delta = \pm 1} \int \hat{\mathcal{P}}(\Delta, t, h) \mathcal{P}(\Delta, t, h) dt dh \right\} - i \sum_{i=1}^n \hat{\mathcal{P}}(\Delta_i, T_i, \mathbf{X}'_i \boldsymbol{\beta}_{\alpha})} \mathcal{D} \hat{\mathcal{R}} 2 \Big]$$

obtaining

$$Z_{n}(\gamma, \mathcal{D}) = \int e^{n \left\{ i \sum_{\Delta = \pm 1} \int \hat{\mathcal{P}}(\Delta, t, h) \mathcal{P}(\Delta, t, h) dt dh - \gamma \mathcal{E}\left[\mathcal{P}(.)\right] \right\}} \times \left\{ \int e^{-i \sum_{i=1}^{n} \hat{\mathcal{P}}(\Delta_{i}, T_{i}, \mathbf{X}_{i}' \boldsymbol{\beta}) - \gamma r(\boldsymbol{\beta}/\sqrt{p})} d\boldsymbol{\beta} \right\} \mathcal{D}\hat{\mathcal{P}} \mathcal{D}\mathcal{P} .$$
(73)

For integer r we then have

$$Z_{n}^{r}(\gamma, \mathcal{D}) = \int e^{n \sum_{\alpha=1}^{r} \left\{ i \sum_{\Delta=\pm 1} \int \hat{\mathcal{P}}_{\alpha}(\Delta, t, h, \gamma) \mathcal{P}_{\alpha}(\Delta, t, h, \gamma) dt dh - \gamma \mathcal{E} \left[ \mathcal{P}_{\alpha}(.) \right] \right\}} \times \left\{ \int \prod_{\alpha=1}^{r} e^{-i \sum_{i=1}^{n} \hat{\mathcal{P}}_{\alpha}(\Delta_{i}, T_{i}, \mathbf{X}_{i}' \boldsymbol{\beta}_{\alpha}) - \gamma r(\boldsymbol{\beta}_{\alpha})} d\boldsymbol{\beta}_{\alpha} \right\} \prod_{\alpha=1}^{r} \mathcal{D} \hat{\mathcal{P}}_{\alpha} \mathcal{D} \mathcal{P}_{\alpha} .$$
(74)

Taking the expectation with respect to the data-set

$$\mathbb{E}_{\mathcal{D}}\left[Z_{n}^{r}(\gamma,\mathcal{D})\right] = \int e^{n\sum_{\alpha=1}^{r}\mathcal{A}\left[\hat{\mathcal{P}}_{\alpha},\mathcal{P}_{\alpha}\right]} \mathcal{W}\left[\left\{\hat{\mathcal{P}}_{\alpha}\right\}\right] \prod_{\alpha=1}^{r} \mathcal{D}\hat{\mathcal{P}}_{\alpha} \ \mathcal{D}\mathcal{P}_{\alpha}$$
(75)

where

$$\mathcal{A}[\gamma, \hat{\mathcal{P}}_{\alpha}, \mathcal{P}_{\alpha}] = i \sum_{\Delta = \pm 1} \int \hat{\mathcal{P}}_{\alpha}(\Delta, t, h) \mathcal{P}_{\alpha}(\Delta, t, h) dt dh - \gamma \mathcal{E}[\mathcal{P}_{\alpha}(.)]$$
(76)

and

$$\mathcal{W}[\gamma, \{\hat{\mathcal{P}}_{\alpha}\}] = \int \left(\mathbb{E}_{\Delta, T, \mathbf{X}}\left[e^{-i\sum_{\alpha=1}^{r}\hat{\mathcal{P}}_{\alpha}(\Delta, T, \mathbf{X}'\boldsymbol{\beta}_{\alpha})}\right]\right)^{n} e^{-\gamma r(\boldsymbol{\beta}_{\alpha})} \prod_{\alpha=1}^{r} d\boldsymbol{\beta}_{\alpha}.$$
(77)

We notice that the expression above depends on  $\beta_{\alpha}$  only via the linear predictors  $\mathbf{X}'_i \beta_{\alpha}$ . Since  $\mathbf{X} \sim \mathcal{N}(\mathbf{0}, \frac{1}{p} \mathbf{I}_{p \times p})$ , we have that

$$\mathbf{Y} = (Y_0, Y_1, \dots, Y_r) \sim \mathcal{N}(\mathbf{0}, \mathbf{C}(\{\boldsymbol{\beta}_\alpha\})), \quad Y_\alpha := \mathbf{X}' \boldsymbol{\beta}_\alpha$$
(78)

with

$$\mathbf{C}(\{\boldsymbol{\beta}_{\alpha}\})) = \begin{bmatrix} \boldsymbol{\theta}_{0}^{2} & \mathbf{M}^{\mathrm{T}} \\ \mathbf{M} & \mathbf{R} \end{bmatrix}$$
(79)

where

$$\theta_0^2 = \|\beta_0\|^2 / p \tag{80}$$
$$\mathbf{M} = (M_1)^r + M_2 = \beta_1' \beta_1 / p \tag{81}$$

$$\mathbf{M} = (M_{\alpha})_{\alpha=1}^{r}, \quad M_{\alpha} := \beta_{0}^{r} \beta_{\alpha} / p$$

$$\mathbf{R} = (R_{\alpha,\rho})_{\alpha,\rho=1}^{r}, \quad R_{\alpha,\rho} := \beta_{\alpha}^{r} \beta_{\rho} / p = R_{\rho,\alpha} .$$
(81)
$$(82)$$

Then, introducing a matrix delta function, we can write

$$\mathcal{W}[\{\hat{\mathcal{P}}_{\alpha}\}] = \int e^{ip \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}) + p\phi(\hat{\mathbf{C}}) + n\varphi(\gamma, \mathbf{C})} \, \mathrm{d}\hat{\mathbf{C}} \mathrm{d}\mathbf{C}$$
(83)

with

$$\varphi[\gamma, \mathbf{C}, \{\hat{\mathcal{P}}_{\alpha}(.)\}_{\alpha=1}^{r}] = \log \mathbb{E}_{\Delta, T, \mathbf{Y}}^{C} \left[ e^{-i\sum_{\alpha=1}^{r} \hat{\mathcal{P}}_{\alpha}(\Delta, T, Y_{\alpha})} \right]$$
(84)

$$\phi(\gamma, \hat{\mathbf{C}}) = \frac{1}{p} \log \int e^{-ip \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}(\{\boldsymbol{\beta}_{\alpha}\})) - \gamma \frac{1}{2} \operatorname{r}(\boldsymbol{\beta}_{\alpha})} \prod_{\alpha=1}^{r} \mathrm{d}\boldsymbol{\beta}_{\alpha} .$$
(85)

Putting everything together, we have obtained a so-called "saddle point" (functional) integral

$$\mathbb{E}_{\mathcal{D}}\left[Z_{n}^{r}(\gamma,\mathcal{D})\right] = \int e^{-n\psi\left[\{\mathcal{P}_{\alpha},\hat{\mathcal{P}}_{\alpha}\}_{\alpha=1}^{r},\hat{\mathbf{C}},\mathbf{C}\right]} \prod_{\alpha=1}^{r} \mathcal{D}\hat{\mathcal{P}}_{\alpha} \,\mathcal{D}\mathcal{P}_{\alpha} \mathrm{d}\mathbf{C}\mathrm{d}\hat{\mathbf{C}}$$
(86)

where

$$-\psi [\{\mathcal{P}_{\alpha}, \hat{\mathcal{P}}_{\alpha}\}_{\alpha=1}^{r}, \hat{\mathbf{C}}, \mathbf{C}] = \sum_{\alpha=1}^{r} \mathcal{A}[\hat{\mathcal{P}}_{\alpha}, \mathcal{P}_{\alpha}] + \varphi[\gamma, \mathbf{C}, \{\hat{\mathcal{P}}_{\alpha}(.)\}_{\alpha=1}^{r}] + i\zeta \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}) + \zeta\phi(\gamma, \hat{\mathbf{C}}).$$
(87)

#### **B.2** Saddle point integration

The idea is now to evaluate the integral via the saddle point method by interchanging the limits  $n \to \infty$  and  $r \to 0$ , as customary in these calculations [5, 4, 17, 18]. We first derive the stationary conditions with respect to the functions  $\mathcal{P}_{\alpha}$  and  $\hat{\mathcal{P}}_{\alpha}$ , which are obtained via functional differentiation

$$\mathcal{P}_{\alpha}(\Delta, t, h, \gamma) = \frac{\mathbb{E}_{\mathbf{Y}}^{\mathbf{C}} \left[ f(\Delta, t | Y_0) \delta(h - Y_{\alpha}) \mathrm{e}^{-\mathrm{i} \sum_{\alpha=1}^{r} \hat{\mathcal{P}}_{\alpha}(\Delta, t, Y_{\alpha}, \gamma)} \right]}{\mathbb{E}_{\Delta, T, \mathbf{Y}}^{\mathbf{C}} \left[ \mathrm{e}^{-\mathrm{i} \sum_{\alpha=1}^{r} \hat{\mathcal{P}}_{\alpha}(\Delta, T, Y_{\alpha}, \gamma)} \right]}$$
(88)

$$i\hat{\mathcal{P}}_{\alpha}(\Delta, t, h, \gamma) = \gamma \Delta \Big[\log \mathcal{S}_{\alpha}(t, \gamma) - h\Big] + \gamma e^{h} \Lambda_{\alpha}(t, \gamma) .$$
(89)

where

$$S_{\alpha}(t,\gamma) = \sum_{\Delta'=\pm 1} \int \Theta(t-t') e^{h'} \mathcal{P}_{\alpha}(\Delta',t,h') dh' dt'$$
(90)

$$\Lambda_{\alpha}(t,\gamma) := \sum_{\Delta'=\pm 1} \int \frac{\Delta' \Theta(t-t') \mathcal{P}_{\alpha}(\Delta',t',h',\gamma)}{\mathcal{S}_{\alpha}(t',\gamma)} dt' dh' .$$
(91)

Equivalently

$$\mathcal{P}_{\alpha}(\Delta, t, h, \gamma) = \frac{\mathbb{E}_{\mathbf{Y}}^{\mathbf{C}} \Big[ f(\Delta, t | Y_0) \delta(h - Y_\alpha) \mathrm{e}^{-\sum_{\alpha=1}^r \gamma \left( \Delta \log \mathcal{S}_\alpha(t, \gamma) + \exp(Y_\alpha) \Lambda_\alpha(t, \gamma) - \Delta Y_\alpha \right)} \Big]}{\mathbb{E}_{\Delta, T, \mathbf{Y}}^{\mathbf{C}} \Big[ \mathrm{e}^{-\sum_{\alpha=1}^r \gamma \left( \Delta \log \mathcal{S}_\alpha(T, \gamma) + \exp(Y_\alpha) \Lambda_\alpha(T, \gamma) - \Delta Y_\alpha \right)} \Big]} \quad . \tag{92}$$

Furthermore at the saddle point we have

$$\varphi[\gamma, \mathbf{C}, \{\hat{\mathcal{P}}_{\alpha}(.)\}_{\alpha=1}^{r}] = \varphi(\gamma, \mathbf{C}) = \log \mathbb{E}_{\Delta, T, \mathbf{Y}}^{\mathbf{C}} \left[ e^{-\sum_{\alpha=1}^{r} \gamma \left( \Delta \log \mathcal{S}_{\alpha}(T, \gamma) + \exp(Y_{\alpha}) \Lambda_{\alpha}(T, \gamma) - \Delta Y_{\alpha} \right)} \right]$$

and hence

$$-\tilde{\psi}(\gamma, \hat{\mathbf{C}}, \mathbf{C}) := - \underset{\{\mathcal{P}_{\alpha}, \hat{\mathcal{P}}_{\alpha}\}}{\operatorname{extr}} \psi = \mathrm{i}\zeta \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}) + \zeta \phi(\gamma, \hat{\mathbf{C}}) + \varphi(\gamma, \mathbf{C}) +$$
$$+ \gamma \sum_{\Delta = \pm 1} \int \mathrm{e}^{h} \sum_{\alpha = 1}^{r} \Lambda_{\alpha}(t, \gamma) \mathcal{P}_{\alpha}(\Delta, t, h, \gamma) \mathrm{d}t \mathrm{d}h$$
(93)

With a modest amount of foresight we take the following change of variables

$$i\hat{\mathbf{C}} = \frac{1}{2}\mathbf{D} \tag{94}$$

which is expected from previous similar calculations and aids the book-keeping. In principle we could now derive the saddle point equations for the elements of the matrices  $\mathbf{C}$  nor  $\mathbf{D}$  and then take the limit  $r \to 0$ . In practice we will assume the replica symmetric ansatze

$$\mathbf{C} = \begin{pmatrix} \theta_0^2 & m & \dots & m \\ m & \rho & q & \dots & q \\ \vdots & q & \rho & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & q \\ m & q & \dots & q & \rho \end{pmatrix} \quad \mathbf{D} = \begin{pmatrix} 0 & \hat{m} & \dots & \dots & \hat{m} \\ \hat{m} & \hat{\rho} & -\hat{q} & \dots & -\hat{q} \\ \vdots & -\hat{q} & \hat{\rho} & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & -\hat{q} \\ \hat{m} & -\hat{q} & \dots & -\hat{q} & \hat{\rho} \end{pmatrix}$$
(95)

together with

$$\Lambda_{\alpha}(t,\gamma) = \Lambda(t,\gamma) \quad \forall \alpha = 1,\dots,r$$
(96)

directly from now on. Note that this directly implies that

$$\mathbf{C}^{-1} = \begin{pmatrix} \tilde{\mu}_r & \tilde{m}_r & \dots & \dots & \tilde{m}_r \\ \tilde{m}_r & \tilde{\rho}_r & \tilde{q}_r & \dots & \tilde{q}_r \\ \vdots & \tilde{q}_r & \tilde{\rho}_r & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & \tilde{q}_r \\ \tilde{m}_r & \tilde{q}_r & \dots & \tilde{q}v & \tilde{\rho}_r \end{pmatrix} .$$
(97)

After some algebra one obtains that

$$\tilde{\mu}_{r} = \frac{\rho + q(r-1)}{\theta_{0}^{2}(\rho + q(r-1)) - rm^{2}}$$
(98)

$$\tilde{m}_r = \frac{m}{rm^2 - \theta_0^2(\rho + q(r-1))}$$
(99)

$$\tilde{q}_r = \frac{1}{\rho - q} \frac{m^2 - q\theta_0^2}{\theta_0^2 (\rho + q(r-1)) - rm^2}$$
(100)

$$\tilde{\rho}_r = \frac{1}{\rho - q} \left( 1 + \frac{m^2 - q\theta_0^2}{\theta_0^2(\rho + q(r-1)) - rm^2} \right).$$
(101)

We are now in position to compute the "limit"  $r \to 0$ , i.e. to obtain the replica symmetric symmetric free energy.

#### **B.2.1** Simplification of $\phi$ within RS ansatz

Let's remind the definition of the potential  $\phi$ 

$$\phi(\gamma, \mathbf{D}) = \frac{1}{p} \log \int e^{-\frac{1}{2}p \operatorname{Tr}(\mathbf{DC}(\{\boldsymbol{\beta}_{\alpha}\})) - \gamma \mathbf{r}(\boldsymbol{\beta}_{\alpha})} \prod_{\alpha=1}^{r} \mathrm{d}\boldsymbol{\beta}_{\alpha} .$$
(102)

Using the replica symmetric ansatz we get

$$\frac{1}{2}p\operatorname{Tr}(\mathbf{DC}(\{\beta_{\alpha}\})) = \sum_{\rho=1}^{r} \left(\hat{m}\beta_{0}^{\prime}\beta_{\rho} + \frac{1}{2}\hat{\rho}\beta_{\rho}^{\prime}\beta_{\rho}\right) - \sum_{\rho,\alpha\neq\rho}^{r} \hat{q}\beta_{\rho}\beta_{\alpha} =$$
$$= \sum_{\rho=1}^{r} \left(\hat{m}\beta_{0}^{\prime}\beta_{\rho} + \frac{1}{2}(\hat{\rho}+\hat{q})\|\beta_{\rho}\|^{2}\right) - \frac{1}{2}\hat{q}\|\sum_{\rho=1}^{r}\beta_{\rho}\|^{2}$$

and via Gaussian linearization

$$\phi_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = \frac{1}{p} \log \mathbb{E}_{\mathbf{Z}} \left[ \left( \int e^{-\frac{1}{2}(\hat{\rho} + \hat{q}) \|\mathbf{x}\|^2 - (\hat{m}\boldsymbol{\beta}_0 + \sqrt{\hat{q}}\mathbf{Z})'\mathbf{x} - \gamma \mathbf{r}(\mathbf{x})} d\mathbf{x} \right)^r \right]$$

Since we are finally interested in taking the limit  $r \to 0$  it is convenient to expand the integrand for small r, thus obtaining

$$\phi_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = r \frac{1}{p} \mathbb{E}_{\mathbf{Z}} \left[ \log \int e^{-\frac{1}{2}(\hat{\rho} + \hat{q}) \|\mathbf{x}\|^2 - (\hat{m}\boldsymbol{\beta}_0 + \sqrt{\hat{q}}\mathbf{Z})'\mathbf{x} - \gamma \mathbf{r}(\mathbf{x})} d\mathbf{x} \right] + O(r^2) \,.$$

Since the integrand in the expression above factorizes over the components of x and Z, this can be further re-written in terms of the distribution of the entries of  $\beta_0$ 

$$\phi_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = r \mathbb{E}_{\beta_0, Z} \left[ \log \int e^{-\frac{1}{2}(\hat{\rho} + \hat{q})x^2 - (\hat{m}\beta_0 + \sqrt{\hat{q}}Z)x - \gamma r(x)} dx \right] + O(r^2) .$$

# **B.2.2** Simplification of $\varphi$ within RS ansatz

Inserting the replica symmetric ansatze, we obtain

$$f(y_0, y_1, \dots, y_r) \propto \exp\left\{-\left(\frac{1}{2}\tilde{\mu}y_0^2 + \sum_{\rho=1}^r \left(\tilde{m}y_0y_\rho + \frac{1}{2}(\tilde{\rho} - \tilde{q})y_\rho^2\right) + \frac{1}{2}\tilde{q}\left(\sum_{\rho=1}^r y_\rho\right)^2\right)\right\}$$
(103)

and via Gaussian linearization

$$f(y_0, y_1, \dots, y_r) \propto \mathbb{E}_Q \Big[ \exp \Big\{ -\frac{1}{2} \tilde{\mu} y_0^2 - \sum_{\rho=1}^r \Big[ \frac{1}{2} (\tilde{\rho} - \tilde{q}) y_\rho^2 + \big( \tilde{m} y_0 + i \sqrt{\tilde{q}} Q \big) y_\rho \Big] \Big\} \Big]$$
(104)

with  $Q \sim \mathcal{N}(0,1)$ . Upon setting  $\sqrt{\tilde{\mu}}y_0 = z_0$  we obtain

$$\begin{split} \varphi_{RS}^{(r)}(\gamma,\tilde{\mu}_{r},\tilde{m}_{r},\tilde{q}_{r},\tilde{\rho}_{r}) &= \\ &= \log \frac{\mathbb{E}_{\Delta,T,Z_{0},Q} \left[ \left( \int \mathrm{e}^{-\frac{1}{2}(\tilde{\rho}-\tilde{q})x^{2} - \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_{0} + \mathrm{i}\sqrt{\tilde{q}}Q\right)x - \gamma g(x,\Lambda^{(r)}(T,\gamma),\Delta) \frac{\mathrm{d}x}{\sqrt{2\pi}} \right)^{r} \right]}{\mathbb{E}_{\Delta,T,Z_{0},Q} \left[ \left( \int \mathrm{e}^{-\frac{1}{2}(\tilde{\rho}-\tilde{q})x^{2} - \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_{0} + \mathrm{i}\sqrt{\tilde{q}}Q\right)x} \right)^{r} \right]} \,. \end{split}$$

with  $Z_0 \sim \mathcal{N}(0, 1), Z_0 \perp Q$  and where we have defined

$$g(x, y, z) = \exp(x)y - zx.$$
(105)

# **B.3** Replica symmetric functional equation

Using (104) we obtain

$$\mathcal{P}^{(r)}(\Delta, t, h, \gamma) = \left( \mathbb{E}_{\Delta, T, Z_0, Q} \left[ \mathcal{Z}^r(\gamma, \Delta, T, Z_0, Q) \right] \right)^{-1} \times \\ \times \mathbb{E}_{Z_0, Q} \left[ \frac{f(\Delta, t | Z_0/\sqrt{\tilde{\mu}}) \int \delta(h-x) e^{-\left\{ \frac{1}{2} (\tilde{\rho} - \tilde{q}) x^2 + \left( \tilde{m}/\sqrt{\tilde{\mu}} Z_0 + i\sqrt{\tilde{q}} Q \right) x + \gamma g(x, \Lambda^{(r)}(T, \gamma), \Delta) \right\}}{\left( \int e^{-\left\{ \frac{1}{2} (\tilde{\rho} - \tilde{q}) x^2 + \left( \tilde{m}/\sqrt{\tilde{\mu}} Z_0 + i\sqrt{\tilde{q}} Q \right) x + \gamma g(x, \Lambda^{(r)}(T, \gamma), \Delta) \right\}} dx \right)^{1-r}} \right]$$

where we used the shorthand

$$\mathcal{Z}(\gamma, \Delta, T, Z_0, Q) = \int e^{-\gamma \left\{ \frac{1}{2} \frac{(\tilde{\rho} - \tilde{q})}{\gamma} x^2 + \frac{1}{\gamma} \left( \tilde{m} / \sqrt{\tilde{\mu}} Z_0 + i\sqrt{\tilde{q}} Q \right) x + \gamma g(x, \Lambda^{(r)}(T, \gamma), \Delta) \right\}} dx .$$
(106)

Let us stop and recall what we have achieved so far. By assuming the RS ansatz we have obtained

$$-\lim_{n \to \infty} \frac{1}{n} \log \mathbb{E}_{\mathcal{D}} \Big[ Z_n^r(\gamma, \mathcal{D}) \Big] = \operatorname{extr}_{m, q, \rho, \hat{\mu}, \hat{m}, \hat{q}, \hat{\rho}} \tilde{\psi}_{RS}^{(r)}(m, q, \rho, \hat{m}, \hat{q}, \hat{\rho})$$
(107)

with

$$-\tilde{\psi}_{RS}^{(r)}(\dots) = \zeta(\mu\hat{\mu} + rm\hat{m} + r(\rho\hat{\rho} - q\hat{q}) + r^{2}q\hat{q}) + \zeta\phi_{RS}^{(r)}(\gamma,\hat{m},\hat{q},\hat{\rho}) + \varphi_{RS}^{(r)}(\gamma,\tilde{\mu}_{r},\tilde{m}_{r},\tilde{q}_{r},\tilde{\rho}_{r}) + \gamma r\mathbb{E}_{\Delta,T,Z_{0},Q} \left[ \Lambda^{(r)}(t,\gamma) \frac{\int e^{x} e^{-\left\{\frac{1}{2}(\tilde{\rho}-\tilde{q})x^{2} + \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_{0}+i\sqrt{\tilde{q}}Q\right)x + \gamma g(x,\Lambda^{(r)}(T,\gamma),\Delta)\right\}}}{\left(\int e^{-\left\{\frac{1}{2}(\tilde{\rho}-\tilde{q})x^{2} + \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_{0}+i\sqrt{\tilde{q}}Q\right)x + \gamma g(x,\Lambda^{(r)}(T,\gamma),\Delta)\right\}} dx \right)^{1-r}} \right].$$
(108)

where now

$$\mathcal{S}^{(r)}(t',\gamma) = \mathbb{E}_{\Delta,T,Z_0,Q} \left[ \Theta(T-t') \frac{\int e^x e^{-\left\{\frac{1}{2}(\tilde{\rho}-\tilde{q})x^2 + \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_0 + i\sqrt{\tilde{q}}Q\right)x + \gamma g(x,\Lambda^{(r)}(T,\gamma),\Delta)\right\}\right\}_{\mathrm{d}x}}{\left(\int e^{-\left\{\frac{1}{2}(\tilde{\rho}-\tilde{q})x^2 + \left(\tilde{m}/\sqrt{\tilde{\mu}}Z_0 + i\sqrt{\tilde{q}}Q\right)x + \gamma g(x,\Lambda^{(r)}(T,\gamma),\Delta)\right\}\right\}_{\mathrm{d}x}\right)^{1-r}} dx}$$
$$\Lambda^{(r)}(t,\gamma) := \sum_{\Delta'=\pm 1} \int \frac{\Delta'\Theta(t-t')}{\mathcal{S}(t',\gamma)} \mathcal{P}^{(r)}(\Delta',t',h',\gamma) \mathrm{d}t' \mathrm{d}h' \,.$$

# **B.4** The limit $r \to 0$

Taking the limit  $r \to 0$ , we get the replica symmetric free energy

$$f_{RS}(\gamma) = \underset{\mu,m,q,\rho,\hat{m},\hat{q},\hat{\rho}}{\text{extr}} \tilde{f}_{RS}(\gamma,\mu,m,q,\rho,\hat{m},\hat{q},\hat{\rho})$$
(109)

with

$$\tilde{f}_{RS}(\gamma, m, q, \rho, \hat{m}, \hat{q}, \hat{\rho}) := \lim_{r \to 0} \frac{1}{\gamma r} \tilde{\psi}_{RS}^{(r)}(\gamma, m, q, \rho, \hat{m}, \hat{q}, \hat{\rho}) .$$
(110)

where

$$-\tilde{f}_{RS}(\dots) = \zeta (m\hat{m} + r(\rho\hat{\rho} - q\hat{q})) + \zeta \tilde{\phi}_{RS}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) + \tilde{\varphi}_{RS}(\gamma, m, q, \rho) + \\ + \mathbb{E}_{\Delta, T, Z_0, Q} \left[ \Lambda(t, \gamma) \frac{\int e^x e^{-\frac{1}{\rho - q} \left\{ \frac{1}{2}x^2 - \left( (m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q \right) x \right\} - \gamma g(x, \Lambda(t, \gamma), \Delta)}{\int e^{-\frac{1}{\rho - q} \left\{ \frac{1}{2}x^2 - \left( (m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q \right) x \right\} - \gamma g(x, \Lambda(t, \gamma), \Delta)} dx} \right]$$
 (111)

Above, we introduced the following definitions

$$\begin{split} \tilde{\phi}_{RS}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) &= \lim_{r \to 0} \frac{1}{\gamma r} \phi_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = \\ &= \mathbb{E}_{\beta_0, Z} \bigg[ \log \int e^{-\frac{1}{2}(\hat{\rho} + \hat{q})x^2 - (\hat{m}\beta_0 + \sqrt{\hat{q}}Z)x - \gamma r(x)} dx \bigg] \\ \tilde{\varphi}_{RS}(\gamma, m, q, \rho) &= \lim_{r \to 0} \frac{1}{\gamma r} \varphi_{RS}^{(r)}(\gamma, \tilde{\mu}_r, \tilde{m}_r, \tilde{q}_r, \tilde{\rho}_r) \\ &= \mathbb{E}_{\Delta, T, Z_0, Q} \bigg[ \log \frac{\int e^{-\frac{1}{\rho - q} \bigg\{ \frac{1}{2}x^2 - \big((m/\theta_0)Z_0 + \sqrt{q - (m/\theta_0)^2}Q\big)x \bigg\} - \gamma g(x, \Lambda(t, \gamma), \Delta)}{\int e^{-\frac{1}{\rho - q} \bigg\{ \frac{1}{2}x^2 - \big((m/\theta_0)Z_0 + \sqrt{q - (m/\theta_0)^2}Q\big)x \bigg\}} dx} \bigg] \end{split}$$

and we have used that

$$\tilde{\mu} = \lim_{r \to 0} \tilde{\mu}_r = 1/\theta_0^2 \tag{112}$$

$$\tilde{m} = \lim_{r \to 0} \tilde{m}_r = -\frac{m}{\theta_0^2(\rho - q)}$$
(113)

$$\tilde{q} = \lim_{r \to 0} \tilde{q}_r = \frac{1}{\rho - q} \frac{m^2 - q\theta_0^2}{\theta_0^2(\rho - q)}$$
(114)

$$\tilde{\rho} - \tilde{q} = \lim_{r \to 0} \tilde{\rho}_r - \tilde{q}_r = \frac{1}{\rho - q} \,. \tag{115}$$

Assuming the following scaling

$$\tau = \gamma / (\tilde{\rho} - \tilde{q}) = \gamma (\rho - q) \tag{116}$$

we obtain

$$\tilde{\varphi}_{RS}(\gamma, m, q, \rho) = \\ = \mathbb{E}_{\Delta, T, Z_0, Q} \left[ \frac{1}{\gamma} \log \frac{\int e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q) \right)^2 / \tau + g \left( x, \Lambda(T, \gamma), \Delta \right) \right\}_{\mathrm{d}x}}{\int e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q) \right)^2 / \tau \right\}_{\mathrm{d}x}} \right] (117)$$

Similarly, taking the additional rescaling

$$1/\hat{\tau} = (\hat{\rho} + \hat{q})/\gamma, \quad \hat{m} = \gamma \hat{m}, \quad \hat{q} = \gamma^2 \hat{q}$$
(118)

we have

$$\begin{split} \tilde{\phi}_{RS}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) &= \mathbb{E}_{\beta_0, Z} \left[ \frac{1}{\gamma} \log \int e^{-\gamma \left\{ \frac{1}{2\hat{\tau}} x^2 + (\hat{m}\beta_0 + \sqrt{\hat{q}}Z)x + \mathbf{r}(x) \right\}} dx \right] = \\ &= \frac{1}{2} \hat{\tau} \left( \hat{m}^2 \theta_0^2 + \hat{q} \right) + \mathbb{E}_{\beta_0, Z} \left[ \frac{1}{\gamma} \log \int e^{-\gamma \left\{ \frac{1}{2} \frac{1}{\hat{\tau}} \left\{ \tilde{x} + \hat{\tau} (\hat{m}\beta_0 + \sqrt{\hat{q}}Z) \right\}^2 + \mathbf{r}(\tilde{x})} \right\}} d\tilde{x} \right] + \text{const} \end{split}$$

Furthermore, in the limit  $r \rightarrow 0,$  the functional equation reads

$$\mathcal{P}(\Delta, t, h, \gamma) =$$

$$= \mathbb{E}_{Z_0, Q} \left[ \frac{f(\Delta, t | Z_0 / \sqrt{\tilde{\mu}}) \int \delta(h - x) e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q) \right)^2 / \tau + g \left( x, \Lambda(T, \gamma), \Delta \right) \right\}_{\mathrm{d}x}}{\int e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Q) \right)^2 / \tau + g \left( x, \Lambda(T, \gamma), \Delta \right) \right\}_{\mathrm{d}x}} \right]}$$
(119)

where

$$\mathcal{S}(t',\gamma) = \mathbb{E}_{\Delta,T,Z_0,Q} \left[ \Theta(T-t') \frac{\int e^x e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0)Z_0 + \sqrt{q - (m/\theta_0)^2}Q) \right)^2 / \tau + g\left( x, \Lambda(T,\gamma), \Delta \right) \right\} dx}{\int e^{-\gamma \left\{ \frac{1}{2} \left( x - ((m/\theta_0)Z_0 + \sqrt{q - (m/\theta_0)^2}Q) \right)^2 / \tau + g\left( x, \Lambda(T,\gamma), \Delta \right) \right\} dx} \right]}$$
  
$$\Lambda(t,\gamma) := \mathbb{E}_{\Delta',T'} \left[ \frac{\Delta' \Theta(t-T')}{\mathcal{S}(T',\gamma)} \right].$$

In the last expression we have used that

$$\int \mathcal{P}(\Delta, t, h, \gamma) \mathrm{d}h = f(\Delta, t | Z_0 / \sqrt{\tilde{\mu}}) .$$
(120)

# **B.5** The limit $\gamma \to \infty$

Via Laplace integration we see that

$$-\lim_{\gamma \to \infty} \frac{1}{\gamma} \log \int e^{-\gamma \left\{ \frac{1}{2\alpha} (z-x)^2 + b(z) \right\}} dz = \mathcal{M}_{b(.)}(x,\alpha)$$
(121)

where  $\mathcal{M}_{b(.)}$  is the Moureau envelope of a convex function  $b : \mathbb{R} \to \mathbb{R}$ , which is defined as

$$\mathcal{M}_{b(.)}(x,\alpha) = \min_{z} \left\{ \frac{1}{2\alpha} (z-x)^2 + b(z) \right\}.$$
 (122)

Using the "Laplace-Moureau" identity above (121), we obtain

$$\lim_{\gamma \to \infty} \tilde{\phi}_{RS}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = \frac{1}{2} \hat{\tau} \left( \hat{m}^2 \theta_0^2 + \hat{q} \right) - \mathbb{E}_{Z,\beta_0} \left[ \mathcal{M}_{\mathbf{r}(.)} \left( \hat{m} \beta_0 + \sqrt{\hat{q}} Z, \hat{\tau} \right) \right]$$
$$\lim_{\gamma \to \infty} \tilde{\varphi}_{RS}(\gamma, m, q, \rho) = -\mathbb{E}_{\Delta,T,Z_0,Q} \left[ \mathcal{M}_{\tau g(.,\Lambda(T),\Delta)} \left( m/\theta_0 Z_0 + \sqrt{q - (m/\theta_0)^2} Q, \tau \right) \right].$$

In the limit  $\gamma \to \infty$ , the functional equation (119) reads

$$\mathcal{P}(\Delta, t, h) = \mathbb{E}_{Z_0, Z} \left[ f(\Delta, t | Z_0) \delta \left( h - \xi(T, Z_0, Z) \right) \right]$$
(123)

$$\xi(T, Z_0, Z) := \operatorname{prox}_{g(.,\Lambda(T),\Delta)} \Big( (m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Z, \tau \Big),$$
(124)

where we used the Laplace integration method to conclude that for any "well behaved" function  $a: \mathbb{R}^d \to \mathbb{R}$ 

$$a\left(\operatorname{prox}_{b(.)}(x,\alpha)\right) = \lim_{\gamma \to \infty} \log \frac{\int e^{-\gamma \left\{\frac{1}{2\alpha}(z-x)^2 + b(z)\right\}} a(z) dz}{\int e^{-\gamma \left\{\frac{1}{2\alpha}(z-x)^2 + b(z)\right\}} dz}$$
(125)

with  $prox_{b(.)}$  the proximal mapping of the convex function b defined as

$$\operatorname{prox}_{b(.)}(x,\alpha) = \arg\min_{z} \left\{ \frac{1}{2\alpha} (z-x)^2 + b(z) \right\}.$$
 (126)

We also have

$$S(t) = \mathbb{E}_{\Delta,T,Z_0,Z} \Big[ \Theta(t-T) e^{\xi} \Big]$$
  
$$\Lambda(t) := \mathbb{E}_{\Delta',T'} \Big[ \frac{\Delta' \Theta(t-T')}{S(T')} \Big]$$

where  $\xi$  is defined in (124). Furthermore

$$\begin{split} \lim_{\gamma \to \infty} \mathbb{E}_{\Delta,T,Z_0,Z} \bigg[ \Lambda(t,\gamma) \frac{\int \mathrm{e}^x \mathrm{e}^{-\frac{1}{\rho-q} \left\{ \frac{1}{2} x^2 - \left( (m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Z \right) x \right\} - \gamma g(x,\Lambda(t,\gamma),\Delta)}}{\int \mathrm{e}^{-\frac{1}{\rho-q} \left\{ \frac{1}{2} x^2 - \left( (m/\theta_0) Z_0 + \sqrt{q - (m/\theta_0)^2} Z \right) x \right\} - \gamma g(x,\Lambda(t,\gamma),\Delta)} \mathrm{d}x} \bigg] = \\ &= \mathbb{E}_{\Delta,T} \bigg[ \frac{\Delta \mathbb{E}_{\Delta'',T'',Y_0'',Z''} \bigg[ \Theta(T'' - T) \mathrm{e}^{\xi''} \bigg]}{\mathbb{E}_{\Delta',T',Y_0',Z'} \bigg[ \Theta(T' - T) \mathrm{e}^{\xi'} \bigg]} \bigg] = \mathbb{E} \bigg[ \Delta \bigg] = \mathrm{const.} \end{split}$$

Putting all together (for the last time), we have obtained that

$$\lim_{\gamma \to \infty} -\tilde{f}_{RS}(\gamma, m, q, \rho, \hat{m}, \hat{q}, \hat{\rho}) = \mathcal{PL}(m, q, \tau, \hat{m}, \hat{q}, \hat{\tau}) = \lim_{n \to \infty} \mathcal{D}\Big[\mathcal{PL}_n(\hat{\beta}_n)\Big]$$
(127)

with

$$\mathcal{PL}(m,q,\tau,\hat{m},\hat{q},\hat{\tau}) = \zeta(m\hat{m} + \frac{1}{2}q/\hat{\tau} - \frac{1}{2}\tau\hat{q}) + \frac{1}{2}\hat{\tau}\zeta\left(\hat{m}^{2}\theta_{0}^{2} + \hat{q}\right) + -\zeta\mathbb{E}_{Z,\beta_{0}}\left[\mathcal{M}_{\mathbf{r}(.)}\left(\hat{m}\beta_{0} + \sqrt{\hat{q}}Z,\hat{\tau}\right)\right] + -\mathbb{E}_{\Delta,T,Z_{0},Q}\left[\mathcal{M}_{g(.,\Lambda(T),\Delta)}\left(m/\theta_{0}Z_{0} + \sqrt{q - (m/\theta_{0})^{2}}Q,\tau\right)\right] + \text{cons}(128)$$

It is now convenient to take additional change of variables in order to obtain "neater" formulae. Let us define

$$w = m/\theta_0, \quad v = q - m^2/\theta_0^2, \quad \hat{w} = -\hat{\tau}\hat{m}\theta_0, \quad \hat{v} = \hat{\tau}\sqrt{\hat{q}}$$
(129)

then (abusing the notation yet again)

$$\mathcal{PL}(w, v, \tau, \hat{w}, \hat{v}, \hat{\tau}) = \frac{\zeta}{2\hat{\tau}} \left( (w - \hat{w})^2 + v^2 + \hat{v}^2 (1 - \tau/\hat{\tau}) \right) + -\zeta \mathbb{E}_{Z,\beta_0} \left[ \mathcal{M}_{\mathbf{r}(.)} \left( \hat{w} \frac{\beta_0}{\theta_0} + \hat{v}Z, \hat{\tau} \right) \right] + -\mathbb{E}_{\Delta,T,Z_0,Q} \left[ \mathcal{M}_{g(.,\Lambda(T),\Delta)} \left( wZ_0 + vQ, \tau \right) \right] + \text{const}$$
(130)

# **C** Replica symmetric equations

It is convenient to define

$$\varphi(\beta_0, Z) := \operatorname{prox}_{\mathbf{r}(.)} \left( \hat{w} \frac{\beta_0}{\theta_0} + \hat{v} Z, \hat{\tau} \right)$$
(131)

$$\xi := \operatorname{prox}_{g(.,\Lambda(T),\Delta)}(wZ_0 + vQ,\tau)$$
(132)

then

$$-\frac{\partial}{\partial \hat{w}}\mathcal{PL} = \frac{\zeta}{\hat{\tau}}w - \frac{\zeta}{\hat{\tau}}\mathbb{E}_{Z,\beta_0}\left[\beta_0\varphi\right]/\theta_0 \implies w = \mathbb{E}_{Z,\beta_0}\left[\beta_0\varphi\right]/\theta_0$$
(133)

$$-\frac{\partial}{\partial \hat{v}}\mathcal{PL} = \frac{\zeta}{\hat{r}}\hat{v}\frac{\tau}{\hat{r}} - \frac{\zeta}{\hat{r}}\mathbb{E}_{Z,\beta_0}\left[Z\varphi\right] \implies \hat{v}\frac{\tau}{\hat{r}} = \mathbb{E}_{Z,\beta_0}\left[Z\varphi\right]$$
(134)

$$-\frac{\partial}{\partial \hat{\tau}} \mathcal{PL} = \frac{\zeta}{2\hat{\tau}^2} \Big( (w - \hat{w})^2 + v^2 + \hat{v^2} (1 - 2\frac{\tau}{\hat{\tau}}) \Big) - \frac{\zeta}{2\hat{\tau}^2} \mathbb{E}_{Z,\beta_0} \Big[ \big(\varphi - \hat{w}\beta_0 - \hat{v}Z\big)^2 \Big] (135)$$
$$= \frac{\zeta}{2\hat{\tau}^2} \Big( w^2 + v^2 \Big) - \frac{\zeta}{2\hat{\tau}^2} \mathbb{E}_{Z,\beta_0} \Big[ (\varphi - \hat{w}\beta_0 - \hat{v}Z\big)^2 \Big] (136)$$

$$= \frac{1}{2\hat{\tau}^2} \left( w^2 + v^2 \right) - \frac{1}{2\hat{\tau}^2} \mathbb{E}_{Z,\beta_0} \left[ \varphi^2 \right]$$

$$(136)$$

$$\mathbb{P} \left( \sum_{\alpha \in \mathcal{A}} \hat{\varphi}_{\alpha}^{(\alpha)} + \hat{\varphi}_{\alpha}^{(\alpha)} \right) = \frac{1}{2\hat{\tau}^2} \left[ \sum_{\alpha \in \mathcal{A}} \hat{\varphi}_{\alpha}^{(\alpha)} \right]$$

$$(137)$$

$$-\frac{\partial}{\partial w}\mathcal{PL} = -\zeta(w-\hat{w})/\hat{\tau} + \frac{1}{\tau}\left(w - \mathbb{E}_{\Delta,T,Z_0,Q}\left[Z_0\xi\right]\right)$$

$$(137)$$

$$\frac{\partial}{\partial v}\mathcal{PL} = -\zeta(w/\hat{\tau} + \frac{1}{\tau}\left(w - \mathbb{E}_{\Delta,T,Z_0,Q}\left[Z_0\xi\right]\right)$$

$$(138)$$

$$\frac{\partial}{\partial v} \mathcal{PL} = -\zeta v / \hat{\tau} + \frac{1}{\tau} \left( v - \mathbb{E}_{\Delta, T, Z_0, Q} \left[ Q\xi \right] \right)$$
(138)

$$-\frac{\partial}{\partial \tau} \mathcal{PL} = \zeta \frac{\hat{v}^2}{\hat{\tau}^2} - \frac{1}{\tau^2} \mathbb{E}_{\Delta, T, Z_0, Q} \left[ (\xi - wZ_0 - vQ)^2 \right].$$
(139)

Hence

$$w = \mathbb{E}_{Z,\beta_0} \Big[ \beta_0 \varphi \Big] / \theta_0 \tag{140}$$

$$\hat{v}\frac{\tau}{\hat{\tau}} = \mathbb{E}_{Z,\beta_0}\left[Z\varphi\right] \tag{141}$$

$$(w^2 + v^2) = \mathbb{E}_{Z,\beta_0} \left[ \varphi^2 \right]$$
(142)

$$\hat{w} = w - \frac{\tau}{\zeta \tau} \left( w - \mathbb{E}_{\Delta, T, Z_0, Q} \left[ Z_0 \xi \right] \right)$$
(143)

$$v(1 - \zeta \tau / \hat{\tau}) = \mathbb{E}_{\Delta, T, Z_0, Q} \left[ Q \xi \right]$$
(144)

$$\zeta \hat{v}^2 = \frac{\hat{\tau}^2}{\tau^2} \mathbb{E}_{\Delta, T, Z_0, Q} \Big[ (\xi - wZ_0 - vQ)^2 \Big]$$
(145)

# D Explicit computation for the lasso regularization

Here we assume that

$$\mathbf{e}'_{\mu}\beta_{0} \sim \mathcal{N}(0,\theta_{0}^{2}/\nu), \quad \nu := s/p \,.$$
 (146)

For the Lasso regularization

$$\operatorname{prox}_{\mathbf{r}(.)}(x,\hat{\tau}) = \operatorname{st}(x,\alpha\hat{\tau}), \qquad (147)$$

$$\operatorname{st}(x,\alpha) := \operatorname{relu}(x-\alpha) - \operatorname{relu}(-x-\alpha)$$
 (148)

with

$$\operatorname{relu}(x) = x\Theta(x) . \tag{149}$$

Before proceeding, let us remember some useful fact

$$\mathbb{E}_{Z}\left[\Theta(\sigma Z + \mu - \alpha)\right] = \Phi\left(\frac{\alpha - \mu}{\sigma}\right)$$
(150)

$$\mathbb{E}_{Z}\left[Z\Theta(\sigma Z + \mu - \alpha)\right] = \frac{e^{-\frac{1}{2}\left(\frac{\alpha - \mu}{\sigma}\right)}}{\sqrt{2\pi}}$$
(151)

$$\mathbb{E}_{Z}\left[Z^{2}\Theta(\sigma Z + \mu - \alpha)\right] = \Phi\left(\frac{\alpha - \mu}{\sigma}\right) + \frac{\alpha - \mu}{\sigma} \frac{e^{-\frac{1}{2}\left(\frac{\alpha - \mu}{\sigma}\right)^{2}}}{\sqrt{2\pi}}$$
(152)

where

$$\Phi(x) := \int_{x}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^{2}} dx$$
(153)

The identities above imply

$$\mathbb{E}_{Z}[Zrelu(\sigma Z + \mu - \alpha)] = \sigma \Phi(\frac{\alpha - \mu}{\sigma})$$
(154)

and hence

$$\mathbb{E}_{Z}[Zst(\sigma Z, \alpha)] = \sigma \Phi\left(\frac{\alpha - \mu}{\sigma}\right) + \sigma \Phi\left(\frac{\alpha + \mu}{\sigma}\right).$$
(155)

This implies that

$$\mathbb{E}_{Z,\beta_0}\left[\beta_0\varphi\right] = \hat{w}\mathbb{E}_Z\left[\Phi\left(\sqrt{\nu}\frac{\alpha\hat{\tau}-\hat{v}Z}{\hat{w}}\right) + \Phi\left(\sqrt{\nu}\frac{\alpha\hat{\tau}+\hat{v}Z}{\hat{w}}\right)\right] = 2\hat{w}\mathbb{E}_Z\left[\Phi\left(\sqrt{\nu}\frac{\alpha\hat{\tau}+\hat{v}Z}{\hat{w}}\right)\right].$$
(156)

and also

$$\mathbb{E}_{Z,\beta_0} \left[ Z\varphi \right] = \hat{v} \mathbb{E}_{\beta_0} \left[ \Phi\left( \frac{\alpha \hat{\tau} - \hat{w} \beta_0 / \theta_0}{\hat{v}} \right) + \Phi\left( \frac{\alpha \hat{\tau} + \hat{w} \beta_0 / \theta_0}{\hat{v}} \right) \right] = 2 \hat{v} \mathbb{E}_{\beta_0} \left[ \Phi\left( \frac{\alpha \hat{\tau} + \hat{w} \beta_0 / \theta_0}{\hat{v}} \right) \right].$$
(157)

Next we use that

$$\mathbb{E}_X\Big[\Phi(aX+b)\Big] = \Phi\Big(\frac{b}{\sqrt{1+a^2}}\Big)$$
(158)

and get that

$$\mathbb{E}_{Z}\left[\Phi\left(\sqrt{\nu}\frac{\alpha\hat{\tau}+\hat{v}Z}{\hat{w}}\right)\right] = \Phi\left(\frac{\alpha\hat{\tau}}{\sqrt{\hat{w}^{2}/\nu+\hat{v}^{2}}}\right)$$
(159)

and

$$\mathbb{E}_{\beta_0}\left[\Phi\left(\frac{\alpha\hat{\tau}+\hat{w}\beta_0/\theta_0}{\hat{v}}\right)\right] = \nu\Phi\left(\frac{\alpha\hat{\tau}}{\sqrt{\hat{w}^2/\nu+\hat{v}^2}}\right) + (1-\nu)\Phi\left(\frac{\alpha\hat{\tau}}{\hat{v}}\right).$$
(160)

Finally we use that

$$\mathbb{E}_{Z}[\operatorname{relu}^{2}(\sigma Z + \mu - \alpha)] = \Phi\left(\frac{\alpha - \mu}{\sigma}\right) \left(\sigma^{2} + (\mu - \alpha)^{2}\right) - \sigma(\alpha - \mu) \frac{\mathrm{e}^{-\frac{1}{2}\left(\frac{\alpha - \mu}{\sigma}\right)^{2}}}{\sqrt{2\pi}}$$
(161)

hence

$$\frac{1}{2}\mathbb{E}_{Z,\beta_{0}}\left[\varphi^{2}\right] = \nu\left\{\Phi\left(\frac{\alpha\hat{\tau}}{\sqrt{\hat{v}^{2}+\hat{w}^{2}/\nu}}\right)\left(\hat{v}^{2}+\hat{w}^{2}/\nu+\alpha^{2}\hat{\tau}^{2}\right)+ -\sqrt{\hat{v}^{2}+\hat{w}^{2}/\nu}\alpha\hat{\tau}\frac{\mathrm{e}^{-\frac{1}{2}\left(\frac{\alpha\hat{\tau}}{\sqrt{\hat{v}^{2}+\hat{w}^{2}/\nu}}\right)^{2}}}{\sqrt{2\pi}}\right\}+ \left(1-\nu\right)\left\{\Phi\left(\frac{\alpha\hat{\tau}}{\hat{v}}\right)\left(\hat{v}^{2}+\alpha^{2}\hat{\tau}^{2}\right)-\hat{v}\alpha\hat{\tau}\frac{\mathrm{e}^{-\frac{1}{2}\left(\frac{\alpha\hat{\tau}}{\hat{v}}\right)^{2}}}{\sqrt{2\pi}}\right\}.$$
(162)

If we introduce the shorthands

$$\chi_0 := \frac{\alpha \hat{\tau}}{\hat{v}}, \quad \chi_1 := \frac{\alpha \hat{\tau}}{\sqrt{\hat{v}^2 + \hat{w}^2/\nu}}$$
 (163)

then the first three RS equations take the more compact form

$$w = 2\hat{w}\Phi(\chi_1) \tag{164}$$

$$\tau = 2\hat{\tau} \left\{ \nu \Phi(\chi_1) + (1 - \nu) \Phi(\chi_0) \right\}$$
(165)

$$\frac{1}{2}(v^2 + w^2) = \nu \{ (1 + 1/\chi_1^2) \Phi(\chi_1) - G(\chi_1) \} + (1 - \nu) \{ (1 + 1/\chi_0^2) \Phi(\chi_0) - G(\chi_0) \}$$
(166)

#### D.1 Explicit computation for the elastic net regularization

For the Elastic net regularization

$$\operatorname{prox}_{\mathbf{r}(.)}(x,\hat{\tau}) = \frac{1}{1+\tilde{\eta}\hat{\tau}}\operatorname{st}(x,\alpha\hat{\tau})$$
(167)

hence the replica symmetric equations of the main section can be easily recovered.

# **E** Distributions

Here we show that the distributions

$$\mathcal{P}_{\xi}(\Delta, t, h) := \lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \frac{1}{n} \sum_{i=1}^{n} \delta_{\Delta, \Delta_{i}} \delta(t - T_{i}) \delta(h - \mathbf{X}_{i}' \hat{\boldsymbol{\beta}}) \Big],$$
(168)

$$\mathcal{P}_{\varphi}(x) := \lim_{n \to \infty} \mathbb{E}_{\mathcal{D}} \Big[ \frac{1}{p} \sum_{k=1}^{p} \delta(x - \mathbf{e}'_{k} \hat{\boldsymbol{\beta}}) \Big]$$
(169)

indeed satisfy

$$\mathcal{P}_{\xi}(\Delta, t, h) = \mathbb{E}_{\Delta', T', Z'_{0}, Q'} \Big[ \delta(t - T') \delta_{\Delta, \Delta'} \delta(h - \xi_{\star}) \Big]$$
(170)

$$\mathcal{P}_{\varphi}(x) = \mathbb{E}_{\beta_0, Z} \Big[ \delta \big( x - \varphi_{\star} \big) \Big], \tag{171}$$

as stated in the main text (33, 34). For the identity (170), observe that  $\mathcal{P}_{\xi}(\Delta, t, h)$  is the functional order parameter of the replica theory (65), hence, at the replica symmetric saddle point we have (106) and after taking the limit  $r \to 0$  and  $\gamma \to \infty$  we obtain (123), which coincides with (170) as

$$\mathcal{P}_{\xi}(\Delta, t, h) = \mathbb{E}_{Z_0, Z} \Big[ f(\Delta, t | Z_0) \delta \Big( h - \xi_{\star} \Big) \Big] = \\ = \mathbb{E}_{\Delta', T', Z'_0, Z'} \Big[ \delta_{\Delta, \Delta'} \delta(t - T') \delta \Big( h - \xi_{\star} \Big) \Big] .$$
(172)

The identity (171) requires more care. Via Laplace integration we may write

$$\frac{1}{p}\sum_{k=1}^{p}\delta(x-\mathbf{e}_{k}'\hat{\boldsymbol{\beta}}) = \lim_{\gamma \to \infty} \frac{1}{p}\sum_{i=1}^{p}\sum_{k=1}^{p}\frac{\int \mathrm{e}^{-\gamma\mathcal{H}(\boldsymbol{\beta})}\,\delta(x-\mathbf{e}_{\mu}'\boldsymbol{\beta})\,\mathrm{d}\boldsymbol{\beta}}{\int \mathrm{e}^{-\gamma\mathcal{H}(\boldsymbol{\beta})}\,\mathrm{d}\boldsymbol{\beta}}\,.$$
(173)

To compute the expectation over the data-set we use the following alternative replica identity

$$\mathcal{P}_{\varphi}(x) = \lim_{r \to 0} \lim_{\gamma \to \infty} \lim_{p \to \infty} \frac{1}{r p} \sum_{k=1}^{p} \mathbb{E}_{\mathcal{D}} \left[ \int e^{-\gamma \sum_{\alpha=1}^{r} \mathcal{H}(\boldsymbol{\beta}_{\alpha}|\mathcal{D})} \,\delta(x - \mathbf{e}_{k}^{\prime} \boldsymbol{\beta}_{1}) \prod_{\alpha=1}^{r} \mathrm{d}\boldsymbol{\beta}_{\alpha} \right].$$
(174)

Inserting the functional delta measure as in (72) and taking the expectation over  $\mathcal{D}$ , i.e. the data-set, we obtain

$$\mathcal{P}_{\varphi}(x) = \int e^{-n\psi \left[ \left\{ \mathcal{P}_{\alpha}, \hat{\mathcal{P}}_{\alpha} \right\}_{\alpha=1}^{r}, \hat{\mathbf{C}}, \mathbf{C} \right]} \mathcal{J}(\gamma, \hat{\mathbf{C}}) \prod_{\alpha=1}^{r} \mathcal{D}\hat{\mathcal{P}}_{\alpha} \mathcal{D}\mathcal{P}_{\alpha} \mathrm{d}\mathbf{C} \mathrm{d}\hat{\mathbf{C}}$$
(175)

where  $\psi$  is defined as in (87) and

$$\mathcal{J}(\gamma, \hat{\mathbf{C}}) := \frac{1}{p} \sum_{k=1}^{p} \frac{\int e^{-ip \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}(\{\boldsymbol{\beta}_{\alpha}\})) - \gamma r(\boldsymbol{\beta}_{\alpha})} \delta(x - \mathbf{e}_{k}^{\prime} \boldsymbol{\beta}_{1}) \prod_{\alpha=1}^{r} \mathrm{d}\boldsymbol{\beta}_{\alpha}}{\int e^{-ip \operatorname{Tr}(\hat{\mathbf{C}}\mathbf{C}(\{\boldsymbol{\beta}_{\alpha}\})) - \gamma r(\boldsymbol{\beta}_{\alpha})} \prod_{\alpha=1}^{r} \mathrm{d}\boldsymbol{\beta}_{\alpha}} .$$
 (176)

Now taking  $i\hat{\mathbf{C}} = \frac{1}{2}\mathbf{D}$ , and replica symmetry, we notice that the expression above can be simplified, at the saddle point, because the regularizer is separable, i.e.  $r(\mathbf{x}) = \sum_{l=1}^{p} r(x_l)$ ,

$$\begin{aligned} \mathcal{J}_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) &= \\ \frac{1}{p} \sum_{k=1}^{p} \frac{\int e^{-\sum_{\rho=1}^{r} \left( \hat{m}\beta_{0,k}\beta_{\rho,k} + \frac{1}{2}(\hat{\rho}+\hat{q})\beta_{\rho,k}^{2} \right) + \frac{1}{2}\hat{q} \left( \sum_{\rho=1}^{r} \beta_{\rho,k} \right)^{2} - \gamma r(\beta_{\alpha,k})}{\int e^{-\sum_{\rho=1}^{r} \left( \hat{m}\beta_{0,k}\beta_{\rho,k} + \frac{1}{2}(\hat{\rho}+\hat{q})\beta_{\rho,k}^{2} \right) + \frac{1}{2}\hat{q} \left( \sum_{\rho=1}^{r} \beta_{\rho,k} \right)^{2} - \gamma r(\beta_{\alpha,k})} \prod_{\alpha=1}^{r} d\beta_{\alpha,k}} . \end{aligned}$$

At this point Gaussian linearization gives

$$\mathcal{J}_{RS}^{(r)}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) = \mathbb{E}_{\beta_0} \bigg[ \mathbb{E}_Z \bigg[ \int e^{-\left(\{\hat{m}\beta_0 + \sqrt{\hat{q}}Z\}\beta + \frac{1}{2}(\hat{\rho} + \hat{q})\beta^2\right) - \gamma \mathbf{r}(\beta)} \delta(x - \beta) d\beta \times (177) \bigg]$$

$$\times \left(\int e^{-\left(\{\hat{m}\beta_{0,k}+\sqrt{\hat{q}}Z\}\beta+\frac{1}{2}(\hat{\rho}+\hat{q})\beta^{2}\right)-\gamma r(\beta)}d\beta\right)^{\gamma-1}\right] \times$$
(178)

$$\times \mathbb{E}_{Z} \left[ \left( \int \mathrm{e}^{-\left( \left\{ \hat{m}\beta_{0} + \sqrt{\hat{q}}Z \right\}\beta + \frac{1}{2}(\hat{\rho} + \hat{q})\beta^{2} \right) - \gamma \mathrm{r}(\beta)} \mathrm{d}\beta \right)^{r} \right]^{-1} \right].$$
(179)

Hence, after taking the limit  $r \to 0$ , we get

$$\mathcal{J}_{RS}(\gamma, \hat{m}, \hat{q}, \hat{\rho}) := \mathbb{E}_{Z,\beta_0} \left[ \frac{\int \mathrm{e}^{-\left(\{\hat{m}\beta_0 + \sqrt{\hat{q}}Z\}\beta + \frac{1}{2}(\hat{\rho} + \hat{q})\beta^2\right) - \gamma \mathrm{r}(\beta)} \delta(x - \beta) \mathrm{d}\beta}{\int \mathrm{e}^{-\left(\{\hat{m}\beta_0, +\sqrt{\hat{q}}Z\}\beta + \frac{1}{2}(\hat{\rho} + \hat{q})\beta^2\right) - \gamma \mathrm{r}(\beta)} \mathrm{d}\beta} \right].$$
(180)

After taking the rescaling

$$1/\hat{\tau} = (\hat{\rho} + \hat{q})/\gamma, \quad \hat{m} = \gamma \hat{m}, \quad \hat{q} = \gamma^2 \hat{q}$$
(181)

and the limit  $\gamma \to \infty$ , we get

$$\mathcal{P}_{\varphi}(x) = \mathbb{E}Z, \beta_0 \left[ \delta \left( x - \operatorname{prox}_{\mathbf{r}(.)} \left( \hat{m}_{\star} \beta_0 + \sqrt{\hat{q}_{\star}} Z, \hat{\tau}_{\star} \right) \right) \right].$$
(182)

The expression above reduces to (171) after the change of variables

$$\hat{w} = -\hat{\tau}\hat{m}\theta_0, \quad \hat{v} = \hat{\tau}\sqrt{\hat{q}},\tag{183}$$

and using the definition (30).

# F Algorithm for pathwise solution

Consider the function

$$g(\boldsymbol{\beta}, \lambda) := \frac{1}{n} \sum_{i=1}^{n} \left\{ \Lambda(T_i) e^{\mathbf{X}_i' \boldsymbol{\beta}} - \Delta_i \mathbf{X}_i' \boldsymbol{\beta} - \Delta_i \log \lambda(T_i) \right\}$$
(184)

Suppose we want to minimize the objective function

$$f(\boldsymbol{\beta}, \lambda) := g(\boldsymbol{\beta}, \lambda) + \mathbf{r}(\boldsymbol{\beta}) \tag{185}$$

with r a separable convex regularization function. In our case we will be interested in

$$\mathbf{r}(\boldsymbol{\beta}) := \alpha \left( \rho |\boldsymbol{\beta}| + \frac{1}{2} (1 - \rho) \|\boldsymbol{\beta}\|^2 \right).$$
(186)

This can be done via coordinate descent. First we minimize over  $\lambda$ , obtaining the Nelson-Aalen estimator

$$\hat{\Lambda}_n(t) = \mathrm{NA}(\{T_j, \mathbf{X}'_j \boldsymbol{\beta}\}) := \sum_{i=1}^n \frac{\Delta_i \Theta(t - T_i)}{\sum_{j=1}^n \Theta(T_j - T_i) \mathrm{e}^{\mathbf{X}'_j \boldsymbol{\beta}}} \,.$$
(187)

Then we minimize over  $\beta$  and so on an so forth, until a fixed point, i.e. the updated values for  $\beta^{t+1}$  and  $\Lambda^{t+1}(T_1), \ldots, \Lambda^{t+1}(T_n)$  are within a user defined tolerance from their previous value. To compute the minimizer in  $\beta$  at fixed  $\Lambda$ , we use the algorithm proposed in [27]. The idea is to reduce the problem to an iterative regularized least squared regression. Let us define

$$\ell(\boldsymbol{\beta}, \Lambda) := \frac{1}{n} \sum_{i=1}^{n} \left\{ \Lambda(T_i) e^{\mathbf{X}'_i \boldsymbol{\beta}} - \Delta_i \mathbf{X}'_i \boldsymbol{\beta} \right\}.$$
 (188)

A second order expansion in  $\boldsymbol{\xi}$  around  $\boldsymbol{\phi}$  gives the following approximation for  $\ell$ 

$$\tilde{\ell}(\boldsymbol{\beta}, \boldsymbol{\Lambda}, \boldsymbol{\phi}) = \ell(\boldsymbol{\phi}, \boldsymbol{\Lambda}) + \mathbf{s}(\boldsymbol{\phi}, \boldsymbol{\Lambda})'(\boldsymbol{\beta} - \boldsymbol{\phi}) + \frac{1}{2}(\boldsymbol{\beta} - \boldsymbol{\phi})'\boldsymbol{H}(\boldsymbol{\phi}, \boldsymbol{\Lambda})(\boldsymbol{\beta} - \boldsymbol{\phi})$$
(189)

where

$$\mathbf{s}(\boldsymbol{\phi}, \Lambda) = \frac{1}{n} \sum_{i=1}^{n} \left\{ \Lambda(T_i) \mathrm{e}^{\mathbf{X}'_i \boldsymbol{\phi}} - \Delta_i \right\} \mathbf{X}_i = \frac{1}{n} (\mathbf{W} - \mathrm{diag}(\boldsymbol{\Delta}) \mathbf{X}$$
(190)

$$\mathbf{M}(\boldsymbol{\phi}, \Lambda) = \frac{1}{n} \sum_{i=1}^{n} \Lambda(T_i) e^{\mathbf{X}'_i \boldsymbol{\phi}} \mathbf{X}_i \mathbf{X}'_i = \frac{1}{n} \mathbf{X}' \mathbf{W} \mathbf{X}$$
(191)

with  $\mathbf{W} := \operatorname{diag}(\Lambda(T_i) \exp{\{\mathbf{X}'_i \phi\}})$ . At this points we use a coordinate descent strategy to solve the regularized least square problem

$$\boldsymbol{\beta}^{t+1} = \operatorname*{arg\,min}_{\boldsymbol{\varphi}} \Big\{ f(\boldsymbol{\varphi}) \Big\}, \quad f(\boldsymbol{\varphi}) = \tilde{\ell}(\boldsymbol{\varphi}, \Lambda^t, \boldsymbol{\beta}^t) + r(\boldsymbol{\varphi}), \tag{192}$$

i.e. we minimize along each component keeping the remaining components fixed

$$\frac{\partial}{\partial \varphi_k} f(\boldsymbol{\varphi}) = s_k(\boldsymbol{\beta}^t, \boldsymbol{\Lambda}^t) + \varphi_k \, \mathbf{e}'_k \mathbf{M}(\boldsymbol{\beta}^t, \boldsymbol{\Lambda}^t) \mathbf{e}_k + \mathbf{e}'_k \mathbf{M}(\boldsymbol{\beta}^t, \boldsymbol{\Lambda}^t) \left\{ (\boldsymbol{I} - \mathbf{e}_k \mathbf{e}'_k) \boldsymbol{\varphi} - \boldsymbol{\beta}^t \right\} + r'(\varphi_k) = 0$$
(193)

which is solved by

$$\varphi_k^{t+1} = \operatorname{prox}_{r(.)} \left( \frac{\mathbf{e}_k' \mathbf{M}(\boldsymbol{\beta}^t, \Lambda^t) \left\{ \boldsymbol{\beta}^t - (\boldsymbol{I} - \mathbf{e}_k \mathbf{e}_k') \boldsymbol{\varphi}^t \right\} - s_k(\boldsymbol{\beta}^t, \Lambda^t)}{\mathbf{e}_k' \mathbf{M}(\boldsymbol{\beta}^t, \Lambda^t) \mathbf{e}_k}, \frac{1}{\mathbf{e}_k' \mathbf{M}(\boldsymbol{\beta}^t, \Lambda^t) \mathbf{e}_k} \right).$$
(194)

Notice that in the case of the elastic net penalization we get

$$\varphi_k^{t+1} = \frac{1}{\tau_k + \alpha(1-\rho)} \operatorname{st}(\phi_k, \alpha\rho)$$
(195)

where

$$\phi_k := \mathbf{e}'_k \mathbf{M}(\boldsymbol{\beta}^t, \Lambda^t) \left\{ \boldsymbol{\beta}^t - (\boldsymbol{I} - \mathbf{e}_k \mathbf{e}'_k) \boldsymbol{\varphi}^t \right\} - s_k(\boldsymbol{\beta}^t, \Lambda^t)$$
(196)

$$\tau_k := \mathbf{e}'_k \mathbf{M}(\boldsymbol{\beta}^t, \boldsymbol{\Lambda}^t) \mathbf{e}_k \tag{197}$$

and st is the soft thresholding operator.

A pseudocode depiction of the algorithm for the path-wise (in  $\alpha$ ) minimization of (184) with elastic net regularization (186) at fixed  $\rho$  is presented in (1).

Algorithm 1 Coordinate wise path solution for Cox model

```
 \begin{split} & \boldsymbol{\beta}^{0} \leftarrow \mathbf{0}_{p} \\ & \boldsymbol{\Lambda}^{0} \leftarrow \mathrm{NA}(\{T_{j}, \mathbf{0}_{n}\}) \\ & \boldsymbol{\alpha} \leftarrow \alpha_{\max} \\ & \mathbf{for} \ \boldsymbol{\alpha} \ \text{in path } \mathbf{do} \\ & \mathrm{err} \leftarrow 1 \\ & \mathbf{while} \ \mathrm{err} \geq \mathrm{tol} \ \mathbf{do} \\ & \mathbf{s}(\boldsymbol{\beta}^{t}, \boldsymbol{\Lambda}^{t}) \leftarrow \frac{1}{n} \sum_{i=1}^{n} \left\{ \boldsymbol{\Lambda}^{t}(T_{i}) \mathrm{e}^{\mathbf{X}'_{i}\boldsymbol{\beta}^{t}} - \boldsymbol{\Delta}_{i} \right\} \mathbf{X}_{i} = \frac{1}{n} (\mathbf{W} - \mathrm{diag}(\boldsymbol{\Delta}) \mathbf{X} \\ & \mathbf{M}(\boldsymbol{\beta}^{t}, \boldsymbol{\Lambda}^{t}) \leftarrow \frac{1}{n} \sum_{i=1}^{n} \boldsymbol{\Lambda}^{t}(T_{i}) \mathrm{e}^{\mathbf{X}'_{i}\boldsymbol{\beta}^{t}} \mathbf{X}_{i} \mathbf{X}'_{i} = \frac{1}{n} \mathbf{X}' \mathbf{W} \mathbf{X} \\ & \boldsymbol{\varphi}^{t} \leftarrow \boldsymbol{\beta}^{t} \\ & \mathbf{for} \ 1 \leq k \leq p \ \mathbf{do} \\ & \boldsymbol{\phi}_{k} \leftarrow \mathbf{e}'_{k} \mathbf{M}(\boldsymbol{\beta}^{t}, \boldsymbol{\Lambda}^{t}) \left\{ \boldsymbol{\beta}^{t} - (\mathbf{I} - \mathbf{e}_{k} \mathbf{e}'_{k}) \boldsymbol{\varphi}^{t} \right\} - s_{k}(\boldsymbol{\beta}^{t}, \boldsymbol{\Lambda}^{t}) \\ & \tau_{k} \leftarrow \mathbf{e}'_{k} \mathbf{M}(\boldsymbol{\beta}^{t}, \boldsymbol{\Lambda}^{t}) \mathrm{e}_{k} \\ & \boldsymbol{\varphi}^{t}_{k} \leftarrow \frac{1}{\tau_{k} + \alpha(1 - \lambda)} \mathrm{st}(\boldsymbol{\phi}_{k}, \boldsymbol{\alpha} \boldsymbol{\lambda}) \\ & \mathbf{end \ for} \\ & \boldsymbol{\beta}^{t+1} \leftarrow \boldsymbol{\varphi}^{t} \\ & \boldsymbol{\Lambda}^{t+1} \leftarrow \mathrm{NA}(\{T_{j}, \mathbf{X}'_{j} \boldsymbol{\beta}^{t+1}\}) \\ & \mathrm{err} \leftarrow \sqrt{\|\boldsymbol{\beta}^{t+1} - \boldsymbol{\beta}^{t}\|^{2} + \sum_{i=1}^{n} \|\boldsymbol{\Lambda}^{t+1}(T_{i}) - \boldsymbol{\Lambda}^{t}(T_{i})\|^{2}} \\ & \mathbf{end \ for} \\ & \mathbf{end \ for} \end{aligned} \right
```