

IRREDUCIBILITY IN GENERALIZED POWER SERIES

ANTONGIULIO FORNASIERO, NOA LAVI, SONIA L'INNOCENTE,
AND VINCENZO MANTOVA

ABSTRACT. A classical tool in the study of real closed fields are the fields $K((G))$ of generalized power series (i.e., formal sums with well-ordered support) with coefficients in a field K of characteristic 0 and exponents in an ordered abelian group G . In this paper we enlarge the family of ordinals α of non-additively principal Cantor degree for which $K((\mathbb{R}^{\leq 0}))$ admits irreducibles of order type α far beyond $\alpha = \omega^2$ and $\alpha = \omega^3$ known prior to this work.

1. INTRODUCTION

Rings and fields of power series are classical tools in various areas, such as valuation theory. Call **generalised power series** a formal sum $b = \sum_{\gamma} b_{\gamma} t^{\gamma}$ where the **exponents** γ vary in an ordered abelian group G , the **coefficients** b_{γ} are taken from some field K , and its **support** $\{\gamma \in G : b_{\gamma} \neq 0\}$ is well-ordered, namely every nonempty subset has a minimum. It is well known that the collection $K((G))$ of such series forms a field, when equipped with the obvious operations of sum and product [Hah07].

We are interested in the irreducible series in the subrings of the form $K((G^{\leq 0}))$ or $Z + K((G^{< 0}))$. Such rings appear in different contexts; for instance, $\mathbb{Z} + \mathbb{R}((G^{< 0}))$ is always an integer part of the field $\mathbb{R}((G))$, which in turn implies that every real closed field admits an integer part [MR93]. Regarding irreducibility, the first question was posed by Conway [Con76], who conjectured that the series $1 + \sum_n t^{-\frac{1}{n}}$ is irreducible in the ring of omnific integers, which can be written in the form $\mathbb{Z} + \mathbb{R}((G^{< 0}))$ (modulo some set-theoretic details which are irrelevant here).

Conway's conjecture was proved by Berarducci [Ber00]. A crucial part of the argument, first suggested by Gonshor [Gon86], is the reduction to the ring $K((\mathbb{R}^{\leq 0}))$. In this paper, we work exclusively in this ring. We refer the reader to [BKK06, LM24] for extensive considerations on how to use irreducibility in $K((\mathbb{R}^{\leq 0}))$ in order to find irreducibles in rings of the form $Z + K((G^{< 0}))$.

More irreducible series have been presented in the literature. Let the **order type** $\text{ot}(b)$ of a power series $b \in K((\mathbb{R}^{\leq 0}))$ be the ordinal number representing the order type of its support. Let J be the ideal of the series that are divisible by t^{γ} for some $\gamma \in \mathbb{R}^{< 0}$ (such series cannot be factored into irreducibles, since $t^{\gamma} = t^{\frac{\gamma}{2}} t^{\frac{\gamma}{2}} = \dots$). Berarducci's main result reads as follows.

Date: 27th November 2023.

2020 Mathematics Subject Classification. Primary 13F25, 13F15, secondary 13A05, 03E10.

Key words and phrases. omnific integers, surreal numbers, pre-Schreier domain, valued ring.

No data are associated with this article. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

The authors were supported by: PRIN 2017 "Mathematical logic: models, sets, computability" 2017NWTM8R (A.F, N.L, S.L.), INdAM-GNSAGA (A.F, S.L.); Engineering and Physical Sciences Research Council "Model theory of analytic functions" EP/T018461/1 (V.M.).

Theorem 1.1 ([Ber00, Thm. 10.5]). *If $b \in K((\mathbb{R}^{\leq 0})) \setminus J$ (equivalently, $b \in K((\mathbb{R}^{\leq 0}))$ not divisible by t^γ for any $\gamma < 0$) has order type ω^{ω^α} for some ordinal α , then both b and $b + 1$ are irreducible.*

Further irreducible series of order types ω^2 and ω^3 were exhibited in [PS06] and in [LM17]; in parallel, Pitteloud [Pit01] proved that for b of order type ω , b and $b + 1$ are prime, answering a question of Gonschor.

We will show that, in an appropriate sense, *most* series are irreducible for a wide class of order types, and give explicit examples of such series.

Definition 1.2. Let P_α denote the set of all series $b \in K((\mathbb{R}^{\leq 0}))$ such that $\text{ot}(b) = \omega^\alpha$ and the supremum of the support of b (also denoted by $\text{sup}(b)$) is 0. A series b in P_α is said to be **principal**.

For instance, $P_0 = K \setminus \{0\}$, and the series $\sum_n t^{-\frac{1}{n}}$ is in P_1 ; on the other hand, $t^{-1} + 1$ and $1 + \sum_n t^{-\frac{1}{n}}$ are not principal since their order types are respectively 2 and $\omega + 1$, while $\sum_n t^{-1-\frac{1}{n}}$ is also not principal, this time because the supremum of its support is -1 .

By Berarducci's analysis on principal series, all series in P_α satisfy the following properties: for all ordinals α, β, γ ,

- (1) $P_\alpha \cdot P_\beta \subseteq P_{\alpha \oplus \beta}$ ([Ber00, Cor. 9.9]);
- (2) all divisors of $b \in P_\alpha$ are principal ([LM17, Cor. 4.6]);
- (3) $R_\alpha := \{b \in P_\alpha : b \text{ is reducible}\} \stackrel{(2)}{=} \bigcup_{\beta \oplus \gamma = \alpha} P_\beta \cdot P_\gamma \subseteq P_\alpha$.

Here \oplus, \odot denote Hessenberg's natural (commutative) operations. Note that (1) is equivalent to stating that $\text{ot}(bc) = \text{ot}(b) \odot \text{ot}(c)$ for all principal series b, c . One can easily deduce that every principal b of order type ω^{ω^α} is irreducible: there are no $\beta, \gamma \neq 0$ such that $\beta \oplus \gamma = \omega^\alpha$, so $R_\alpha = \emptyset$.

Definition 1.3. We say that $b = \sum_\gamma b_\gamma t^\gamma \in K((\mathbb{R}^{\leq 0}))$ is *random* if one of the following holds:

- $\text{cl}(\text{supp}(b)) - \{0\}$ is a \mathbb{Q} -linearly independent set;
- the tuple $\langle b_\gamma : \gamma \in \text{supp}(b) \rangle$ is algebraically independent over \mathbb{Q} .

In the current work we prove the following:

Theorem 1.4. *Let α be an ordinal of the following forms:*

- (1) $\alpha = \omega^{\beta_1} + \dots + \omega^{\beta_n}$ where $\beta_1 \geq \beta_2 \geq \beta_3$ and $\beta_3 > \dots > \beta_n$.
- (2) $\alpha = \omega^{\beta_1} + \omega^{\beta_2} + k$ where $\beta_1 \geq \beta_2$ and $k \in \mathbb{N}$.

If $b \in P_\alpha$ is random, then b is irreducible, and so is $b + r$ for any principal series r of order type less than ω^α .

If α is of the form (2), then there are no principal series $p_1, \dots, p_m, q_1, \dots, q_m, r$ such that $b = \sum_{i=1}^m p_i q_i + r$, where $\text{ot}(r) < \omega^\alpha$ and for every $1 \leq i \leq m$ we have $1 < \text{ot}(p_i) \leq \text{ot}(q_i)$, $\text{ot}(p_i q_i) = \omega^\alpha$ (that is, $p_i q_i \in R_\alpha$).

Corollary 1.5. *Let b in P_n , where $n \in \mathbb{N}$. If b is random, then b is irreducible.*

Corollary 1.6. *Let α be as in Theorem 1.4(2). Then $\text{Span}_K(R_\alpha)$ is infinite co-dimensional in $\text{Span}_K(P_\alpha)$ as a K -vector space.*

Remark 1.7. As a special case, we find irreducible principal series of order types ω^2 and ω^3 , as in respectively [PS06] and [LM17].

Combining the above result with techniques from [LM24], we are also able to produce irreducible series that are not principal, at least for certain order types.

Theorem 1.8. *Let $b \in K((\mathbb{R}^{\leq 0}))$ be such that $\text{sup}(b) = 0$ and $\text{ot}(b) = m\omega^\alpha + \beta$, where α is as in Theorem 1.4, $m \in \mathbb{N}$ non-zero, and $\beta < \omega^\alpha$. If b is random, then b is irreducible and so is $b + r$ for any r such that $\text{ot}(r) < \omega^\alpha$ and $\text{sup}(b + r) = 0$.*

In Section 3 we define an independence relation which is a generalization of randomness, and can substitute it in all of the above.

We conjecture that in fact all random series are irreducible, and more precisely, we expect the following to hold.

Conjecture 1.9. *Suppose that $b \in K((\mathbb{R}^{\leq 0}))$ is random and $\text{sup}(b) = 0$. Then b is irreducible and so is $b + r$ for any series r with $\text{ot}(r) < \omega^{\text{deg}(b)}$.*

Here $\text{deg}(b)$ is the maximum ordinal α such that $\omega^\alpha \leq \text{ot}(b)$.

Conjecture 1.10. *$\text{Span}_K(R_\alpha)$ is infinite co-dimensional in $\text{Span}_K(P_\alpha)$ as a K -vector space for any $\alpha > 0$.*

2. PRELIMINARIES

We assume that the reader has familiarity with the class **On** of ordinal numbers and the classical (non-commutative) operations on them, but we will give a minimal account of Hessenberg's commutative operations.

Recall that for all $\alpha \in \mathbf{On}$, there is a maximum β such that $\omega^\beta \leq \alpha$ and a unique γ such that $\alpha = \omega^\beta + \gamma$. By repeating the argument, we find a unique finite sequence $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n \geq 0$ of ordinals such that

$$(1) \quad \alpha = \omega^{\beta_1} + \dots + \omega^{\beta_n}.$$

The expression on the right-hand side is called **Cantor normal form** of α . We let β_1 be the **Cantor degree** of α , denoted by $\text{deg}(\alpha)$.

Given $\alpha = \omega^{\gamma_1} + \omega^{\gamma_2} + \dots + \omega^{\gamma_n}$ and $\beta = \omega^{\gamma_{n+1}} + \omega^{\gamma_{n+2}} + \dots + \omega^{\gamma_{n+m}}$ in Cantor normal form, let π be a permutation of the integers $1, \dots, n+m$ such that $\gamma_{\pi(1)} \geq \dots \geq \gamma_{\pi(n+m)}$. Then $\alpha \oplus \beta$ is defined to be $\omega^{\gamma_{\pi(1)}} + \dots + \omega^{\gamma_{\pi(n+m)}}$, and $\alpha \odot \beta$ to be $\bigoplus_{1 \leq i \leq n, n+1 \leq j \leq n+m} \omega^{\gamma_i \oplus \gamma_j}$. Note that by definition $\omega^\alpha \odot \omega^\beta = \omega^{\alpha \oplus \beta}$ for all $\alpha, \beta \in \mathbf{On}$. Moreover, $\omega^{\beta_1} + \dots + \omega^{\beta_n} = \omega^{\beta_1} \oplus \dots \oplus \omega^{\beta_n}$ if (and only if) the left hand side is in Cantor normal form. These are **Hessenberg's natural sum and product**.

We remark that in Cantor normal form, Hessenberg's operations can be interpreted as sum and product for polynomials in the variables ω^{ω^α} , for $\alpha \in \mathbf{On}$. We summarise this consideration with the isomorphism

$$(\mathbf{On}, \oplus, \odot) \cong \left(\mathbb{N} \left[\omega^{\omega^0}, \omega^{\omega^1}, \dots, \omega^{\omega^\alpha}, \dots \right], +, \cdot \right).$$

An ordinal is **additively principal** if it cannot be written as a sum of two strictly smaller ordinals (equivalently, it is of the form ω^α) and **multiplicatively principal** if it cannot be written as a product of two strictly smaller ordinals (equivalently, it is of the form ω^{ω^α}).

Definition 2.1 ([LM24, Def. 3.3.6]). Given $b \in K((\mathbb{R}))$, we call the sum

$$b = b_1 t^{x_1} + \dots + b_n t^{x_n}$$

the **normal form** of b when:

- $x_1 \leq \dots \leq x_n$;
- b_i is principal for all $i = 1, \dots, n$;
- $\text{ot}(b_1) \leq \dots \leq \text{ot}(b_n)$;
- $x_i + \text{supp}(b_i) < x_{i+1} + \text{supp}(b_{i+1})$ for all $i = 1, \dots, n-1$.

By [LM24, Prop. 3.3.7], every $b \in K((\mathbb{R}^{\leq 0}))$ has a unique normal form.

Let J be the (proper) ideal of $K((\mathbb{R}^{\leq 0}))$ generated by the series of the form t^x for $x < 0$. Given $b \in K((\mathbb{R}^{\leq 0}))$, the **ordinal value** of b ([Ber00, p. 558]) is

$$v_J(b) := \begin{cases} 0 & \text{if } b \in J, \\ 1 & \text{if } b \in (J + K) \setminus J, \\ \min\{\text{ot}(c) : c \equiv b \pmod{J + K}\} & \text{otherwise.} \end{cases}$$

The keystone of [Ber00] is that v_J is a multiplicative semi-valuation.

Fact 2.2 ([Ber00, Lem. 5.5, Thm. 9.7]). For all $b, c \in K((\mathbb{R}^{\leq 0}))$ we have:

- $v_J(b + c) \leq v_J(b) \oplus v_J(c)$;
- $v_J(bc) = v_J(b) \odot v_J(c)$;
- $v_J(b) = 0$ if and only if $b \in J$.

The values of v_J are all of the form ω^α ([Ber00, Rem. 5.3]), thus we add the following notation: let $\deg_J(b)$ denote the Cantor degree of $v_J(b)$, where by convention we set $\deg_J(b) = -\infty$ when $b \in J$, and $\omega^{-\infty} = 0$. Thus we have $v_J(b) = \omega^{\deg_J(b)}$, and we may rephrase the above properties as:

- $\deg_J(b + c) \leq \max\{\deg_J(b), \deg_J(c)\}$;
- $\deg_J(bc) = \deg_J(b) \oplus \deg_J(c)$;
- $\deg_J(b) = -\infty$ if and only if $\deg_J(b) = -\infty$.

We note that to prove $\text{ot}(bc) = \text{ot}(b) \odot \text{ot}(c)$ for b, c principal, it was stated with no proof in [Ber00] that there are “few cancellation” in the product of two elements of $K((\mathbb{R}^{\leq 0}))$. We observe that not only the above is not required (and in fact not used) in order to prove the multiplicativity of v_J , it is neither true.

Example 2.3. Let $S = \{s_1, s_2, \dots\}$ be a strictly increasing sequence of \mathbb{Q} -linearly independent real numbers such that $\sup(S) = 0$. Let us define two series $b = \sum_{n \in \mathbb{N}} t^{s_n}$ and $c = \sum_{n \in \mathbb{N}} (-1)^n t^{s_n}$. If m and n have different parities, the coefficient of $t^{s_m + s_n}$ in bc is $(-1)^m + (-1)^n = 1 - 1 = 0$. Consider $T = \{s_m + s_n : m - n \text{ is odd}\}$, then for every $p \in T$ we have $p \notin \text{supp}(bc)$, thus cancellations occurs on a set of order type $\text{ot}(T) = \omega^2$, which is the same as $\text{ot}(b) \odot \text{ot}(c)$.

The following notions are fundamental in our work.

Definition 2.4. Given $b = \sum_{\beta} b_x t^x \in K((\mathbb{R}^{\leq 0}))$ and $\gamma \in \mathbb{R}^{\leq 0}$, we define:

- the **truncation** of b at γ is $b|_{\gamma} := \sum_{x \leq \gamma} b_x t^x$,
- the **translated truncation** of b at γ is $b^{|\gamma} := t^{-\gamma} b|_{\gamma}$.

The equivalence class $b^{|\gamma} + J$ is the **germ of b at γ** .

Definition 2.5. Let $b \in K((\mathbb{R}^{\leq 0})) \setminus J$. The **J -critical point** of b , denoted by $\text{crit}_J(b)$, is the minimal $\gamma \in \text{supp}(b)$ such that for every $\gamma < \delta < 0$ we have $v_J(b^{|\delta}) < v_J(b)$.

We note that if $b \in P_{\alpha}$, then $\text{crit}_J(b) = \min(\text{supp}(b))$.

Remark 2.6. One should not confuse this with the notion of *critical point* $\text{crit}(b)$, which is the minimum $\gamma \in \mathbb{R}^{\leq 0}$ such that $v_J(b^{|\gamma})$ is maximum possible [Ber00, Def. 10.2]. For example for $b = \sum_{n \in \mathbb{N}} t^{-1 - \frac{1}{n+1}} + \sum_{n \in \mathbb{N}} t^{-\frac{1}{n+1}} + \sum_{n \in \mathbb{N}} t^{-2 - \frac{1}{n+1}}$, we have that $v_J(b) = \omega = v_J(b^{|-1}) = v_J(b^{|-2})$, and one can verify that $\text{crit}(b) = -2$ and $\text{crit}_J(b) = -1$.

Given $b \in K((\mathbb{R}^{\leq 0}))$ with $v_J(b) > 1$, we know that $v_J(b)$ has the form ω^β for some ordinal $\beta > 0$. From the Cantor normal form of β it follows that $v_J(b)$ can be written uniquely as a product $\omega^{\omega^{\beta_1}} \dots \omega^{\omega^{\beta_n}}$, where $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$. We define ([Ber00, Def. 6.4]):

- (1) $v_J^p(b) := \omega^{\omega_n^\beta} =$ the **principal value** of b ,
- (2) $v_J^r(b) := \omega^{\omega^{\beta_1}} \cdots \omega^{\omega^{\beta_{n-1}}} =$ the **residual value** of b .

For instance, if $v_J(b) = \omega^3$, then $v_J^p(b) = \omega$ and $v_J^r(b) = \omega^2$. Note that $v_J(b) = v_J^r(b)v_J^p(b)$. For the sake of readability, we introduce the following notation to remove the base ω .

- (1) $\deg_J^p(b) := \omega^{\beta_n}$,
- (2) $\deg_J^r(b) := \omega^{\beta_1} + \cdots + \omega^{\beta_{n-1}}$.

We have $\deg_J(b) = \deg_J^r(b) + \deg_J^p(b)$. Note that $\deg_J^p(b)$ is **additively principal**: the ordinals strictly less than $\deg_J^p(b)$ are closed under addition.

Definition 2.7. Let $b \in K((\mathbb{R}^{\leq 0}))$. We say that $\gamma \in \text{cl}(\text{supp}(b)) - \{0\}$ is a **big point** of b if $\deg_J(b^{|\gamma|}) \geq \deg_J^r(b)$ and $\gamma > \text{crit}_J(b)$. The set of all big points of b is denoted by $\text{Big}(b)$.

More generally, let $\text{Big}^\alpha(b)$ denote all the numbers $\gamma \in \text{cl}(\text{supp}(b)) - \{0\}$ such that $\deg_J(b^{|\gamma|}) \geq \alpha$ and $\gamma > \text{crit}_J(b)$.

We say that $\gamma \in \text{cl}(\text{supp}(b)) - \{0\}$ is a **residual point** of b if $\deg_J(b^{|\gamma|}) = \deg_J^r(b)$. The set of all these point $\text{Res}(b)$ was defined in [Ber00, Def. 6.6] (where it is called $X(b)$).

Remark 2.8. By construction, the big points of a series must accumulate to 0. Moreover, they are accumulation points of $\text{cl}(\text{supp}(b))$ as soon as $\deg_J^r(b) > 0$.

It turns out that translated truncations behave like a sort of ‘generalised coefficients’, as they satisfy the following equation.

Proposition 2.9 ([Ber00, Lem. 7.5(2)]). *For all $a, b \in K((\mathbb{R}^{\leq 0}))$ and $\gamma \in \mathbb{R}^{\leq 0}$ we have:*

$$(ab)^{|\gamma|} \equiv \sum_{\delta+\varepsilon=\gamma} a^{|\delta|} b^{|\varepsilon|} \pmod{J} \quad (\text{convolution formula}).$$

And one further more may obtain a kind of a Leibniz rule.

Proposition 2.10 ([Ber00, Lem. 7.7]). *Let $b, c \in K((\mathbb{R}^{\leq 0}))$ such that $\deg_J^p(b) \leq \deg_J^p(c)$. Then for every γ sufficiently close to 0 we have $(bc)^{|\gamma|} = b^{|\gamma|}c + c^{|\gamma|}b + r$ where $\deg_J(r) < \deg_J^r(b) \oplus \deg_J(c) < \deg_J(bc)$.*

Definition 2.11. For every α we let $J_\alpha := \{b \in K((\mathbb{R}^{\leq 0})) : \deg_J(b) < \alpha\}$. J_α is a K -vector space, and for α additively principal also a ring, which would allow the use of J_α -linear combinations in our proofs.

In this work we also make use of the *degree valuation*, in order to prove irreducibility results for non-principal elements.

Definition 2.12 ([LM24, p. 5]). Given $b \in K((\mathbb{R}^{\leq 0}))$ with $b \neq 0$, let the **degree** of b , denoted by $\deg(b)$, be the Cantor degree of $\text{ot}(b)$. We also let $\deg(0) := -\infty$.

For instance, $t^{-\sqrt{2}} + t^{-1} + 1$ has degree 0, while $\sum_{n \in \mathbb{N}} t^{-1 - \frac{1}{n+1}} + \sum_{n \in \mathbb{N}} t^{-\frac{1}{n+1}}$ has degree 1 because its order type is $\omega + \omega$, and $\sum_{(m,n) \in \mathbb{N}} t^{-\frac{1}{(n+1)(m+1)}}$ has degree 2, as its order type is ω^2 . In [LM24] it is proved by the 3rd and the 4th authors that the degree is an **multiplicative valuation** in the following sense:

Fact 2.13 ([LM24, Thm. D]). For all non-zero $b, c \in K((\mathbb{R}^{\leq 0}))$,

- (1) $\deg(b + c) \leq \max\{\deg(b), \deg(c)\}$ (ultrametric inequality);
- (2) $\deg(bc) = \deg(b) \oplus \deg(c)$ (multiplicativity);
- (3) $\deg(b) = -\infty$ if and only if $b = 0$.

This makes the degree is quite similar to \deg_J , but in a sense more precise: $\deg(b) = -\infty$ only when $b = 0$, whereas $\deg_J(b) = -\infty$ for all $b \in J$.

Remark 2.14. Note that by construction $v_J(b) \leq \text{ot}(b)$, thus $\deg_J(b) \leq \deg(b)$.

The above inequality is often strict, and not just for series $b \in J$, for instance

$$b = \sum_{(n,m) \in \mathbb{N}^2} t^{-1 - \frac{1}{(n+1)(m+1)}} + \sum_{n \in \mathbb{N}} t^{-\frac{1}{n+1}}$$

has degree 2 while $\deg_J(b) = 1$.

We also remark that all well ordered subsets of \mathbb{R} are countable, thus ot , v_J , \deg_J and \deg all take values below ω_1 , the first uncountable ordinal. One can easily verify that all values below ω_1 are in the images of ot , \deg_J , \deg (but not v_J which is always of the form ω^α).

Definition 2.15. For $\nu \in \{\deg, \deg_J\}$ let $\text{rv}_\nu(b) = \text{rv}_\nu(c)$ if and only if $\nu(b - c) < \nu(b)$ or $b = c$, and let RV_ν be the quotient of $K((\mathbb{R}^{\leq 0}))$ by this equivalence relation. For the sake of notation, we write rv_J , RV_J for respectively rv_{\deg_J} , RV_{\deg_J} .

For both ν as above, RV_ν is the union over $\alpha < \omega_1$ of the quotients $\text{RV}_\nu^\alpha := \{b : \nu(b) \leq \alpha\} / \{c : \nu(c) < \alpha\}$. In particular, each RV_ν^α is naturally a K -vector space, $\text{RV}_\nu^\alpha \cap \text{RV}_\nu^\beta = \{b : \nu(b) \leq \alpha\}$ for $\alpha \neq \beta$, and notably $\text{RV}_J^\alpha = J_{\alpha+1} / J_\alpha$.

Moreover, for b, c principal, we have $\deg(b - c) < \deg(b) = \deg_J(b)$ if and only if $\deg_J(b - c) < \deg_J(b)$. This means that rv_J and rv_{\deg} agree on the principal series. On the other hand, by definition of \deg_J , for every b there is a principal c such that $\text{rv}_J(b) = \text{rv}_J(c)$: in this case we simply write $\text{rv}(b)$. Therefore, each RV_J^α embeds into RV_{\deg}^α as K -vector space and coincides with the image of the principal series under the quotient map. However, RV_{\deg}^α is much richer, for instance $\text{RV}_J^0 \cong K$ while $\text{RV}_{\deg}^0 \cong K[\mathbb{R}^{\leq}]$, the space of the series with finite support. Thanks to RV_{\deg} and its properties proved in [LM24] we are able to find irreducible elements of degree α which are not principal, as we shall see in the following sections.

3. HEREDITARY rv_J

Given b_1, \dots, b_n with $\deg_J(b_1) = \deg_J(b_i)$ for every $i = 1, \dots, n$, we give the following definition by induction on $\deg_J(b_1)$. The series b_1, \dots, b_n are said to be **hereditarily rv_J -independent**, written $\text{Q}(b_1, \dots, b_n)$, if:

- (1) $(\text{rv}_J(b_i))_{1 \leq i \leq n}$ are K -linearly independent;
- (2) when $\deg_J(b_1) \neq \deg_J^p(b_1)$, there exists some $\delta < 0$ such that for all α with $\deg_J^r(b_1) \leq \alpha < \deg_J(b_1)$, for all $\gamma_{1,1}, \dots, \gamma_{n,1}, \dots, \gamma_{n,m(n)} \geq \delta$ with $\deg_J(b_i^{|\gamma_{i,j}|}) = \alpha$ and $\gamma_{i,j} \neq \gamma_{i,j'}$ whenever $j \neq j'$, we have

$$\text{Q}(b_1^{|\gamma_{1,1}|}, \dots, b_1^{|\gamma_{1,m(1)}|}, \dots, b_n^{|\gamma_{n,1}|}, \dots, b_n^{|\gamma_{n,m(n)}|}).$$

Note that the above definition is obviously well founded.

By [LM24, Prop. 5.5.1], we know that every b has a maximal divisor in $K[\mathbb{R}^{\leq}]$ (the series with finite support), which is unique up to a multiplication by an element in K . Clearly, for every b there is a unique maximal divisor d such that the coefficient of $t^{\text{sup}(d)}$ is 1, denoted by $p(b)$.

The following allows us to obtain irreducibility for non-principal elements using irreducibility results for principal elements.

Lemma 3.1. *Let $b \in K((\mathbb{R}^{\leq 0}))$ such that $\text{sup}(b) = 0$. Suppose $b = \sum_{i=1}^m b_i t^{\gamma_i} + r$ where $b_i \in P_{\deg(b)}$ for $1 \leq i \leq m$, $\gamma_1 < \dots < \gamma_m$ and $\deg(r) < \deg(b)$. Suppose also that $\{\text{rv}(b_1), \dots, \text{rv}(b_m)\}$ are linearly independent. Then $p(b) = 1$.*

Proof. By [LM24, Prop. 5.3.1] we have that if $\{C_i\}_{i \in I}$ is a base for RV_J^α then for every $B \in \text{RV}_{\deg}^\alpha$ we have that $B = \sum q_i C_i$ where $q_i \in K[\mathbb{R}^{\leq}]$ and $q_i \neq 0$ for finitely many $i \in I$. As $\{\text{rv}(b_1), \dots, \text{rv}(b_m)\}$ is a part of a base of RV_J^α then by the algorithm

described in [LM24, Rem. 5.4.7] we have that $t^{\gamma_m} = \gcd(t^{\gamma_1}, \dots, t^{\gamma_m})$ is a maximal divisor of finite support of $\text{rv}(b)$, hence $p(\text{rv}(b)) = t^{\gamma_m}$. As $\text{rv}(p(b)) = p(b)$ it follows that $p(b)$ divides $\text{rv}(b)$, hence by maximality it divides also $p(\text{rv}(b))$. As $\text{sup}(b) = 0$, $p(b)$ must be a unit. \square

We are now ready to prove irreducibility for non-principal series:

Proposition 3.2. *Let α be such that for every $c \in P_\alpha$ with $\mathbb{Q}(c)$ we have that $c, \text{rv}(c)$ are irreducible. Let $b = \sum_{i=1}^m b_i t^{\gamma_i} + r$ where $b_1, \dots, b_m \in P_\alpha$, $\gamma_1 < \dots < \gamma_m$ and $\text{ot}(r) < \omega^\alpha$. If $\mathbb{Q}(b_1, \dots, b_m)$ holds then b is irreducible.*

Proof. By Lemma 3.1 we know that $p(b) = 1$, as $\{\text{rv}(b_1), \dots, \text{rv}(b_m)\}$ are linearly independent. By [LM24, Lem. 7.1.1] it is enough to prove that $\frac{\text{rv}(b)}{p(\text{rv}(b))}$ is irreducible. Let $B = \frac{\text{rv}(b)}{t^{\gamma_m}}$ and suppose that $AC = B$. Then there exist some a, c, b' such that $\text{rv}(a) = A, \text{rv}(b') = B, \text{rv}(c) = C$ and $ac = b'$. Then $ac = d \pmod{J}$ for some $d \in P_\alpha$ such that $\text{rv}(d) = \text{rv}(b_m)$ (equivalently, $\text{rv}_J(ac) = \text{rv}_J(b_m)$). As $\text{rv}(d)$ is irreducible, this implies that, without loss of generality, $a \in J + K$. Let $a = \sum_{i=1}^s a_i t^{\delta_i} + a_0$ be the normal form of a , then $a_0 \in K^\times$. As $p(B) = 1$ then also $p(A) = 1$. Hence, we must have also $\deg_J(a_i) = 0$ for every $1 \leq i \leq s$. Hence $a \in K[\mathbb{R}^{\leq}]$, and therefore $a \in K$. Hence, B is irreducible. \square

Remark 3.3. In the proof above we used only Axiom 1 – linear independence in RV_J . In the next section where we use induction arguments we will make heavy use of Axiom 2.

We say that $b_1, \dots, b_m \in K((\mathbb{R}^{\leq 0}))$ are *mutually random* if one of the following holds:

- $\text{cl}(\text{supp}(b_i)) \cap \text{cl}(\text{supp}(b_j)) = \{0\}$ for every $1 \leq i \neq j \leq m$ and $\bigcup_i \text{cl}(\text{supp}(b_i)) - \{0\}$ is \mathbb{Q} -linearly independent set.
- $\langle b_{i_\gamma} : 1 \leq i \leq m, \gamma \in \text{supp}(b_i) \rangle$ is algebraically independent over \mathbb{Q} .

We show now that b_1, \dots, b_m mutually random implies $\mathbb{Q}(b_1, \dots, b_m)$.

Proposition 3.4. *Let b_1, \dots, b_n be such that $\langle b_{i_\gamma} : 1 \leq i \leq n, \gamma \in \text{supp}(b_i) \rangle$ is algebraically independent over \mathbb{Q} . Suppose that $\deg_J(b_i) = \alpha \geq 1$ for every $1 \leq i \leq n$. Then $\text{rv}_J(b_1), \dots, \text{rv}_J(b_n)$ are linearly independent.*

Proof. Suppose by contradiction that we have $k_1, \dots, k_n \in K$ not all zero such that $\sum_{i=1}^n k_i b_i = r \in J_\alpha$. If we replace each coefficient b_{i_γ} with 0 for $\gamma \in \text{supp}(r)$, we find u_1, \dots, u_n all with the *same* support $\bigcup_{i=1}^n \text{supp}(b_i) \setminus \text{supp}(r)$ of order type ω^α and such that $\sum_{i=1}^n k_i u_i = 0$. Moreover, the coefficients of u_1, \dots, u_n satisfy the same algebraic independence assumption as for b_1, \dots, b_n . Let $\gamma_1, \dots, \gamma_n \in \text{supp}(u_1)$. We define $A \in M_{n \times n}(K)$ such that for every $1 \leq i, j \leq n$ we have $A[i, j] = u_{j_\gamma_i}$, and let $\bar{v} = [k_1, \dots, k_n] \neq \bar{0}$. Then we have $A\bar{v} = \bar{0}$ and hence $\det(A) = 0$. Hence, the elements of A are not algebraically independent, a contradiction. \square

Corollary 3.5. *If $\langle b_{i_\gamma} : 1 \leq i \leq m, \gamma \in \text{supp}(b_i) \rangle$ is algebraically independent over \mathbb{Q} , and $\deg_J(b_i) = \deg_J(b_1)$ for every $2 \leq i \leq n$, then $\mathbb{Q}(b_1, \dots, b_n)$ holds.*

Proposition 3.6. *If $\deg_J(b_1) = \deg_J(b_i) > 0$ for $2 \leq i \leq n$, $\text{cl}(\text{supp}(b_i)) \cap \text{cl}(\text{supp}(b_j)) = \{0\}$ for every $1 \leq i \neq j \leq m$ and $\bigcup_i \text{cl}(\text{supp}(b_i)) - \{0\}$ is \mathbb{Q} -linearly independent set, then $\mathbb{Q}(b_1, \dots, b_n)$ holds.*

Proof. For simplicity assume $b_1, \dots, b_n \in P_{\deg_J(b_1)}$. Since the closures of the supports have pairwise intersection $\{0\}$, the supports are pairwise disjoint, so $\text{rv}_J(b_1), \dots, \text{rv}_J(b_n)$ must be K -linearly independent, proving Axiom 1.

Suppose $\deg_J(b_1) \neq \deg_J^p(b_1)$. Let

$$S = \{\gamma_{1,1}, \dots, \gamma_{1,m(1)}, \dots, \gamma_{n,1}, \dots, \gamma_{n,m(n)}\}$$

as in Axiom 2, and suppose by contradiction that there are

$$k_{1,1}, \dots, k_{1,m(1)}, \dots, k_{n,1}, \dots, k_{n,m(n)} \in K$$

not all zero such that $b = \sum_{i=1}^n \sum_{j=1}^{m(i)} k_{i,j} b_i^{|\gamma_{i,j}|} \in J_{\deg_J(b_1)}$.

By comparing the ordinal values, there are i, j and $\delta < 0$ such that δ is in $\text{supp}(b_i^{|\gamma_{i,j}|})$, but not in $\text{supp}(b)$. For δ to cancel, there must be $(i', j') \neq (i, j)$ such that $\delta \in \text{supp}(b_{i'}^{|\gamma_{i',j'}|})$. But then $\beta - \gamma_{i,j} = \delta = \beta' - \gamma_{i',j'}$ for some $\beta \in \text{supp}(b_i)$ and $\beta' \in \text{supp}(b_{i'})$. If $i = i'$, then $\gamma_{i,j} \neq \gamma_{i,j'}$; if $i \neq i'$, since the closure of the supports are disjoint, then $\text{Big}(b_i) \cap \text{Big}(b_{i'}) = \emptyset$, hence again $\gamma_{i,j} \neq \gamma_{i',j'}$. Therefore, $\{\beta, \beta', \gamma_{i,j}, \gamma_{i',j'}\}$ is not linearly independent, a contradiction. \square

Note that in the above proof, we may further require δ to be isolated within the set $\bigcup_{i=1}^n \text{cl}(\text{supp}(b_i^{|\gamma_{i,j}|}))$, since the isolated points themselves have order type $v_J(b_1^{|\gamma_{1,j}|})$, thus most of them must cancel out. In particular, we can find β, β' isolated. Moreover, recall that each $\gamma_{i,j}$ is in $\text{Big}(b_i)$, thus when $\deg_J(b_1) \neq \deg_J^p(b_1)$, that is $\deg_J^r(b_1) > 0$, it is a limit point.

As the contradiction above arises from the equality $\gamma_{i,j} - \gamma_{i',j'} = \beta - \beta'$, we may weaken the condition and allow dependence between either the isolated points or the big points. In particular, to verify Axiom 2 it suffices to test the independence on the points of $\text{supp}(b_i)$ only, rather than all of $\text{cl}(\text{supp}(b_i))$. Likewise, for Axiom 1, it is sufficient to check that the supports of the b_i 's pairwise intersect only on accumulation points, or alternatively that the b_i 's do not have big points in common, again pairwise. Let $\text{Lim}(b)$ be the set of all accumulation points in $\text{cl}(\text{supp}(b))$.

Corollary 3.7. *Suppose that $\deg_J(b_1) = \deg_J(b_i) > 0$ for $2 \leq i \leq n$ and:*

- $(\text{supp}(b_i) \setminus \text{Lim}(b_i)) \cap (\text{supp}(b_j) \setminus \text{Lim}(b_j)) = \emptyset$ for $1 \leq i \neq j \leq n$;
- $\bigcup_{i=1}^n \text{supp}(b_i) \setminus \text{Lim}(b_i)$ is \mathbb{Q} -linearly independent over $\text{Span}_{\mathbb{Q}}(\bigcup_{i=1}^n \text{Big}(b_i))$.

Then $\mathbb{Q}(b_1, \dots, b_n)$ holds.

Corollary 3.8. *If $\deg_J(b_1) = \deg_J(b_i) > 0$ for $2 \leq i \leq n$ and:*

- $\text{Big}(b_i) \cap \text{Big}(b_j) = \{0\}$ for every $1 \leq i \neq j \leq n$;
- $\bigcup_{i=1}^n \text{Big}(b_i)$ is \mathbb{Q} -linearly independent over $\text{Span}_{\mathbb{Q}}(\bigcup_{i=1}^n \text{supp}(b_i) \setminus \text{Lim}(b_i))$.

Then $\mathbb{Q}(b_1, \dots, b_n)$ holds.

4. IRREDUCIBLE PRINCIPAL ELEMENTS

We obtain our results using induction on the Cantor normal form of α for elements of ordinal value ω^α .

Definition 4.1. We let A_α be the K -vector space generated by J_α and all the elements of the form bc where b, c are principal not in K and $\deg_J(bc) = \alpha$; in other words, $A_\alpha := J_\alpha + \text{Span}_K(R_\alpha)$.

Remark 4.2. As observed in the introduction, if α is additively principal, then $R_\alpha = \emptyset$, hence $A_\alpha = J_\alpha$.

We define the following property for $\alpha < \omega_1$.

- $(*)_\alpha$ For every $n \in \mathbb{N}$ and $b_1, \dots, b_n \in P_\alpha$ such that $\mathbb{Q}(b_1, \dots, b_n)$ holds we have that b_1, \dots, b_n are K -linearly independent over A_α .

Note in particular that if the above property holds, then b_1, \dots, b_n are irreducible. We shall verify that $(*)_\alpha$ implies $(*)_{\alpha+\beta}$ for certain choices of β , and thus prove that the property holds for many ordinals. First, we observe the following easy base case.

Proposition 4.3. $(*)_{\omega^\beta}$ is true for every β .

Proof. When $\alpha = \omega^\beta$, the condition $Q(b_1, \dots, b_n)$ only states that $\text{rv}_J(b_1), \dots, \text{rv}_J(b_n)$ are linearly independent, which means exactly that b_1, \dots, b_n are linearly independent over $J_\alpha = A_\alpha$. \square

4.1. The successor case.

Proposition 4.4. $(*)_\alpha \Rightarrow (*)_{\alpha+1}$.

Corollary 4.5. Let $\alpha = \omega^\beta + n$ where $n \in \mathbb{N}$.

- For every $b \in P_\alpha$ with $Q(b)$ we have that $b, \text{rv}(b)$ are irreducible and cannot be represented as a sum of reducible elements in RV_J^α .
- $\text{rv}(A_\alpha)$ has infinite co-dimension as a K -vector subspace of RV_J^α .

Remark 4.6. A special case would be $\alpha = n$.

Proof of Corollary 4.5. We show by induction that $(*)_\alpha$ holds for every $\alpha = \omega^\beta + n$. The induction base is $\alpha = \omega^\beta$, thus Proposition 4.3. The induction step is Proposition 4.4. Hence, for ever $b \in P_\alpha$ where $\alpha = \omega^\beta + n$ with $Q(b)$, we have $b \notin A_\alpha$ and for every $r \in J_\alpha$ we have $b - r \notin A_\alpha$, hence $b, \text{rv}(b)$ are irreducible. \square

Proof of Proposition 4.4. The successor stage is rather simple, because there are no ordinals between α and $\alpha + 1$.

Suppose by contradiction that there are $b_1, \dots, b_n \in P_{\alpha+1}$ and $\lambda_1, \dots, \lambda_n \in K^\times$ such that $b = \sum_{i=1}^n \lambda_i b_i \in A_{\alpha+1}$. Note that by the assumption $Q(b_1, \dots, b_n)$, we have $b \in P_{\alpha+1}$. Write $b = \sum_{i=1}^m p_i q_i + r$ where $\deg_J(q_i) \geq \deg_J(p_i) > 0$, $\deg_J(p_i q_i) = \alpha + 1$ and $r \in J_{\alpha+1}$. For every $\gamma \in \text{Res}(b)$ we have by Proposition 2.10

$$b^{|\gamma} = \sum_{i=1}^m p_i^{|\gamma} q_i + q_i^{|\gamma} p_i \pmod{J_\alpha}.$$

If $p_i^{|\gamma} \notin J + K$, then in fact $p_i^{|\gamma} q_i \in A_\alpha$. Hence, after rearranging the indices, we may write $b^{|\gamma} = \sum_{i=1}^\ell p_i^{|\gamma} q_i \pmod{A_\alpha}$ where $p_i^{|\gamma} \in J + K$ for $i \leq \ell \leq m$. Hence, $\{b^{|\gamma} : \gamma \in \text{Res}(b)\}$ is in the K -linear span of $\{q_1, \dots, q_\ell\} \cup A_\alpha$, and since $\text{Res}(b)$ is infinite, there must be a non-trivial K -linear dependence over A_α .

Pick some $\delta_1, \dots, \delta_k \in K^\times$ and $\gamma_1, \dots, \gamma_k \in \text{Res}(b)$ arbitrarily close to 0 such that $\sum_{i=1}^k \delta_i b^{|\gamma_i} \in A_\alpha$. Expanding the definition of b , we find that the tuple $\langle b_j^{|\gamma_i} \rangle_{\substack{1 \leq i \leq k \\ 1 \leq j \leq n}}$ is linearly dependent over A_α . Recall that $\deg_J(b_j^{|\gamma_i}) \leq \alpha$, and if the inequality is strict, then $b_j^{|\gamma_i} \in A_\alpha$. Thus the subtuple of the series of ordinal value ω^α is also linearly dependent over A_α . On the other hand, since $Q(b_1, \dots, b_n)$ holds, we also know that

$$Q(\langle b_j^{|\gamma_i} : \deg_J(b_j^{|\gamma_i}) = \alpha \rangle),$$

as soon as $\gamma_1, \dots, \gamma_k$ are sufficiently close to 0, which contradicts $(*)_\alpha$. \square

4.2. The case $\omega^{\alpha_1} + \omega^{\alpha_2}$.

Proposition 4.7. $(*)_{\omega^{\alpha_1} + \omega^{\alpha_2}}$ holds for all $\alpha_1 \geq \alpha_2$.

We wish to proceed as in the proof of Proposition 4.4, but we now face new obstacles. Let us retrace the proof. Suppose $b = \sum_{i=1}^n \lambda_i b_i \in A_\alpha$ for some λ_i not all zero. Write $b = \sum_{i=1}^m p_i q_i + r$ with $\deg_J(p_i) = \omega^{\alpha_1}$, $\deg_J(q_i) = \omega^{\alpha_2}$, $\deg_J(r) < \omega^{\alpha_1} + \omega^{\alpha_2}$. If we now consider $b^{|\gamma}$ for some arbitrary $\gamma \in \text{Res}(b)$, we may have $\deg_J(r^{|\gamma}) \geq \deg_J^r(b) = \omega^{\alpha_1}$.

To overcome this issue, we find a large $\Gamma \subseteq \text{Res}(b)$, namely $\text{ot}(\Gamma) = v_J^p(b) = \omega^{\omega^{\alpha_2}}$ and $\text{sup}(\Gamma) = 0$, on which we have $\deg_J(r^{|\gamma}) < \deg_J^r(b) = \omega^{\alpha_1}$.

Lemma 4.8. For every principal series c and ordinal β ,

$$\omega^\beta \cdot \text{ot}(\{\gamma : \deg_J(c^{|\gamma}) \geq \beta\}) \leq v_J(c).$$

Proof. Let c, β as in the assumptions and enumerate $\{\gamma : \deg_J(c^{|\gamma|}) \geq \beta\}$ as $\{\gamma_i : i < \alpha\}$. By construction,

$$\text{ot}(\text{supp}(c) \cap [\gamma_i, \gamma_{i+1})) \geq \deg_J(c^{|\gamma_{i+1}|}) \geq \omega^\beta.$$

It follows by [Ber00, Lem. 4.7] that

$$v_J(c) = \text{ot}(c) \geq \text{ot}(\text{supp}(c) \cap [\gamma_0, 0)) \geq \omega^\beta \cdot \alpha. \quad \square$$

It follows at once that for some $\delta < 0$ sufficiently small, $\text{Big}^{\omega^{\alpha_1}}(r) \cap [\delta, 0]$ has order type strictly smaller than $\omega^{\omega^{\alpha_2}}$, while $\text{ot}(\text{Res}(b)) = \omega^{\omega^{\alpha_2}}$ and $\text{sup}(\text{Res}(b)) = 0$ by [Ber00, Lem. 6.8]. Therefore, $\Gamma = \text{Res}(b) \setminus \text{Big}^{\omega^{\alpha_2}}(r)$ has order type $\omega^{\omega^{\alpha_2}}$ and $\text{sup}(\Gamma) = 0$. By construction, for $\gamma \in \Gamma$ we have $\deg_J(r^{|\gamma|}) < \omega^{\alpha_1}$.

We may now apply Proposition 2.10 and deduce that for every $\gamma \in \Gamma$ close enough to 0 we have

$$b^{|\gamma|} = \sum_{i=1}^m p_i^{|\gamma|} q_i + q_i^{|\gamma|} p_i \pmod{J_{\omega^{\alpha_1}}}.$$

For every $1 \leq i \leq m$ we have $p_i^{|\gamma|}, q_i^{|\gamma|} \in J_{\omega^{\alpha_1}}$. Now $J_{\omega^{\alpha_1}}$ is an integral domain, thus the series $b^{|\gamma|}$ for $\gamma \in \Gamma$ lie in a finitely generated $J_{\omega^{\alpha_1}}$ -module. It follows that for some $k \leq 2m + 1$ there exist some $\delta_1, \dots, \delta_k \in J_{\omega^{\alpha_1}}$ not all zero and $\gamma_1, \dots, \gamma_k \in \Gamma$ close enough to 0 such that $\sum_{i=1}^k \delta_i b^{|\gamma_i|} \in J_{\omega^{\alpha_1}}$. To conclude, we need to improve the coefficients δ_i so that they lie in K .

Lemma 4.9. *Let c, d, e be series such that $\deg_J(c) < \deg_J(e)$ and $\deg_J(d) < \deg_J^p(e)$. Then $\deg_J(cd) < \deg_J(e)$.*

Proof. By looking at the Cantor Normal form of $\deg_J(e)$, we observe that there is an ordinal $\epsilon > 0$ such that $\deg_J(c) \leq \deg_J^r(e) + \epsilon$ and $\epsilon < \deg_J^p(e)$. Since $\deg_J^p(e)$ is additively principal,

$$\deg_J(cd) \leq \deg_J^r(e) \oplus \epsilon \oplus \deg_J(d) < \deg_J^r(e) \oplus \deg_J^p(e) = \deg_J(e). \quad \square$$

Lemma 4.10. *Let c, d be series such that $\deg_J(d) < \deg_J^p(c)$. Then for every γ close enough to 0 we have $(cd)^{|\gamma|} = cd^{|\gamma|} \pmod{J_{\deg_J(c)}}$.*

Proof. For every $\text{crit}_J(c) < \gamma < 0$ we have $\deg_J(c^{|\gamma|}) < \deg_J(c)$, while for every $\text{crit}_J(d) < \delta \leq 0$ we also have $\deg_J(d^{|\delta|}) < \deg_J^p(c)$, thus by Lemma 4.9 we have $\deg_J(d^{|\delta|} c^{|\gamma|}) < \deg_J(c)$. Hence, for every γ sufficiently close to 0 Proposition 2.9 implies that

$$(cd)^{|\gamma|} = cd^{|\gamma|} \pmod{J_{\deg_J(c)}}. \quad \square$$

We now can show that one can find a linear combination with coefficients in K rather than just in $J_{\omega^{\alpha_1}}$.

Lemma 4.11. *Let $c_1, \dots, c_\ell \notin J$ be series such that $\deg_J(c_i) = \deg_J(c_1)$ for every $1 \leq i \leq \ell$, and let $\varepsilon_1, \dots, \varepsilon_\ell \in J_{\deg_J^p(c_1)}$. Then there exist some $\mu_1, \dots, \mu_\ell \in K$ not all zero and γ_0 arbitrarily close to 0 such that*

$$\left(\sum_{i=1}^{\ell} \varepsilon_i c_i \right)^{|\gamma_0|} = \sum_{i=1}^{\ell} \mu_i c_i \pmod{J_{\deg_J(c_1)}}.$$

Proof. Let $S = \bigcup_{i=1}^{\ell} \text{supp}(\varepsilon_i)$, and let $\gamma_0 \in S$ be an isolated point arbitrarily close to 0. Let μ_1, \dots, μ_ℓ be the coefficients $\varepsilon_{1\gamma_0}, \dots, \varepsilon_{\ell\gamma_0}$ respectively. Since γ_0 is isolated, we have $\varepsilon_i^{|\gamma_0|} = \mu_i \pmod{J}$; note in particular at least one such coefficient is not zero. Finally, by Lemma 4.10 we have $(\sum_{i=1}^{\ell} \varepsilon_i c_i)^{|\gamma_0|} = \sum_{i=1}^{\ell} \mu_i c_i \pmod{J_{\deg_J(c_1)}}$, as required. \square

Using Lemma 4.11, we obtain some γ_0 arbitrarily close to 0 and $\mu_1, \dots, \mu_k \in K$ not all zero such that

$$\left(\sum_{i=1}^k \delta_i b^{|\gamma_i} \right)^{|\gamma_0} = \sum_{i=1}^k \mu_i b^{|\gamma_i} \pmod{J_{\omega^{\alpha_1}}}.$$

Note that moreover $b^{|\gamma_i} = \sum_{j=1}^n \lambda_j b_j^{|\gamma_i}$, where $\deg_J(b^{|\gamma_i}) = \deg_J^r(b)$. It now remains to give a bound on $\deg_J(b_j^{|\gamma_i})$.

Lemma 4.12. *Let c_1, \dots, c_ℓ such that $Q(c_1, \dots, c_\ell)$ holds. Suppose $k_1, \dots, k_\ell \in K^\times$ and let $c = \sum_{i=1}^\ell k_i c_i$. Then for every $\gamma \in \text{Res}(c)$ close enough to 0 we have $\deg_J(c_i^{|\gamma}) \leq \deg_J^r(c)$ for all $i = 1, \dots, \ell$.*

Proof. Suppose by contradiction that there exists such γ for which, after reordering the b_i 's, $\deg_J(b_1^{|\gamma}) = \dots = \deg_J(b_s^{|\gamma}) > \deg_J^r(b)$ and $\deg_J(b_i^{|\gamma}) < \deg_J(b_1^{|\gamma})$ for $i > s$. Then $\deg_J(\sum_{i=1}^s k_i b_i^{|\gamma}) < \deg_J(b_1^{|\gamma})$, hence $\text{rv}(b_1^{|\gamma}), \dots, \text{rv}(b_s^{|\gamma})$ are not linearly independent, which contradicts $Q(b_1^{|\gamma}, \dots, b_s^{|\gamma})$. \square

Since we may pick the γ_i 's to be arbitrarily close to 0, we can ensure thanks to Lemma 4.12 that $\deg_J(b_j^{|\gamma_i}) \leq \deg_J^r(b)$ for every $1 \leq i \leq k$, $1 \leq j \leq n$. For each j , let $\gamma_{j,1}, \dots, \gamma_{j,m(j)}$ be an enumeration of the exponents γ_i such that $v(b_j^{|\gamma_i}) = \deg_J^r(b)$. Observe that each γ_i must appear at least once, as otherwise $\deg_J(b^{|\gamma_i}) < \deg_J^r(b)$, contradicting the fact that $\gamma_i \in \text{Res}(b)$. In particular, $\sum_j m(j) \geq n \geq 1$.

However, we have $Q(b_1^{|\gamma_{1,1}}, \dots, b_1^{|\gamma_{1,m(1)}}, \dots, b_n^{|\gamma_{n,1}}, \dots, b_n^{|\gamma_{n,m(n)}})$, thus the $b_j^{|\gamma_{i,k}}$'s cannot be K -linearly independent over $J_{\omega^{\alpha_1}}$, a contradiction. $\square_{\text{Prop. 4.7}}$

Combining with Proposition 4.4 and with Proposition 3.2 we obtain:

Corollary 4.13. *Let $\alpha = \omega^{\alpha_1} + \omega^{\alpha_2} + k$ where $\alpha_1 \geq \alpha_2$ and $k \in \mathbb{N}$.*

- For every $b \in P_\alpha$ with $Q(b)$, we have that $b, \text{rv}(b)$ are irreducible and cannot be represented as a sum of reducible elements in RV_J^α .
- $\text{rv}(A_\alpha)$ has an infinite co-dimension as a vector space in RV_J^α .
- Let $b = \sum_{i=1}^m b_i t^{\gamma_i} + r$ where $\text{sup}(b) = 0$, $b_1, \dots, b_m \in P_\alpha$, $\deg(r) < \alpha$. If $Q(b_1, \dots, b_m)$, then b is irreducible.

The above includes the conclusions of Theorems 1.4, 1.8 for the ordinals of the form $\alpha = \omega^{\alpha_1} + \omega^{\alpha_2} + k$.

4.3. The case $\omega^{\alpha_1} + \omega^{\alpha_2} + \omega^{\alpha_3}$. We show now that we may deduce irreducibility by a different induction on the Cantor normal form of α . Unlike the previous cases, in this induction we do not obtain $(*)_\alpha$, but irreducibility only.

As a result, we shall prove irreducibility for many series in P_α for $\alpha = 3\omega + 1$ but not for $\alpha = 3\omega + 2$. In fact, $3\omega + 2$ is now the smallest α for which we do not know whether there exists any irreducible element in P_α .

We start by proving a crucial lemma for the inductive step:

Lemma 4.14. *Suppose $\deg_J^p(q) > \deg_J^p(p) > 1$. Then for every γ close enough to 0 we have $(pq)^{|\gamma} = p^{|\gamma}q \pmod{J_{\deg_J^r(pq)}}$.*

Proof. Suppose $\deg_J(pq) = \sum_{i=1}^n \omega^{\alpha_i}$ where $\alpha_i \geq \alpha_{i+1}$ for $1 \leq i \leq n-1$. We have that

$$\deg_J(p) = \sum_{i \in A} \omega^{\alpha_i}, \quad \deg_J(q) = \sum_{i \in B} \omega^{\alpha_i}$$

where $A \cup B = \{1, \dots, n\}$ and $A \cap B = \emptyset$. By assumption, we must have $n \in A$ and $\deg_J^p(p) = \omega^{\alpha_n} < \deg_J^p(q) = \omega^{\alpha_\ell}$, where $\ell < n$. We arrange the enumeration so that $\alpha_{\ell+1} < \alpha_\ell$. Moreover, $\deg_J^r(pq) = \sum_{i=1}^{n-1} \omega^{\alpha_i}$.

For every γ close enough to 0 we have

$$\deg_J(q^{|\gamma|}) = \left(\sum_{i \in B \setminus \{\ell\}} \omega^{\alpha_i} \right) + \epsilon$$

where $\epsilon < \omega^{\alpha_\ell}$. It follows that

$$\deg_J(q^{|\gamma|}p) = \left(\sum_{i \neq \ell} \omega^{\alpha_i} \right) \oplus \epsilon < \sum_{i=1}^{\ell} \omega^{\alpha_i} \leq \deg_J^r(pq). \quad \square$$

We now apply the above statement in two slightly different settings. First, we show an inductive step on the Cantor Normal Form of α , under the assumption that the new term is strictly smaller than all the previous ones.

Proposition 4.15. *Let $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ be such that for every $b \in P_\alpha$ with $Q(b)$ we have that $b, \text{rv}(b)$ are irreducible. Then for every b principal with $Q(b)$ such that $\deg_J(b) = \alpha + \omega^{\alpha_{n+1}}$ where $\alpha_n > \alpha_{n+1} > 0$ we have that $b, \text{rv}(b)$ are irreducible.*

Proof. Pick b principal with $Q(b)$ and $\deg_J(b) = \alpha + \omega^{\alpha_{n+1}}$. Suppose by contradiction that $b = pq + r$ where p and q are not in $J + K$ and $\deg_J(r) < \deg_J(b)$. After possibly swapping p and q , we may assume that $\deg_J^p(q) > \deg_J^p(p)$, and so that $\deg_J^p(p) = \omega^{\alpha_{n+1}} > 1$. Here $\deg_J^r(b) = \alpha$.

Let $\Gamma = \text{Res}(b) \setminus \text{Big}^\alpha(r)$. Note that by Lemma 4.8, Γ has order type $\omega^{\alpha_{n+1}}$ and $\sup(\Gamma) = 0$. By Lemma 4.14, for every $\gamma \in \Gamma$ close enough to 0 we have $b^{|\gamma|} = p^{|\gamma|}q + r_\gamma$ where $\deg_J(r_\gamma) < \alpha$.

If $\deg_J(q) < \alpha$, for any γ as above, $p^{|\gamma|}$ is not $J+K$, thus $\text{rv}(b^{|\gamma|})$ is not irreducible. By the assumption on α , $Q(b^{|\gamma|})$ does not hold, a contradiction against $Q(b)$.

Therefore, $\deg_J(q) = \alpha$. It follows that for every $\gamma \in \Gamma$ close enough to 0 we have $\deg_J(p^{|\gamma|}) = 0$. Let $\gamma_1 \neq \gamma_2 \in \Gamma$ be two such exponents. Then $p^{|\gamma_2|}b^{|\gamma_1|} - p^{|\gamma_1|}b^{|\gamma_2|} = p^{|\gamma_2|}r_{\gamma_1} - p^{|\gamma_1|}r_{\gamma_2}$. As $\deg_J(r_{\gamma_1}), \deg_J(r_{\gamma_2}) < \deg_J(q)$, we have $\deg_J(p^{|\gamma_2|}r_{\gamma_1} - p^{|\gamma_1|}r_{\gamma_2}) < \alpha$. Hence, $\text{rv}_J(p^{|\gamma_2|}b^{|\gamma_1|}) - \text{rv}_J(p^{|\gamma_1|}b^{|\gamma_2|}) = 0$, hence $Q(b^{|\gamma_1|}, b^{|\gamma_2|})$ does not hold, another contradiction against $Q(b)$. \square

We then prove a very similar conclusion when the Cantor Normal form has length three, but without the assumption that the last term is strictly smaller than the previous ones. We do this with a preliminary lemma.

Proposition 4.16. *Suppose $b = pq + r$ where $\deg_J(q) = \deg_J^r(b) > \deg_J(p) > 0$ and $\deg_J(r) < \deg_J(b)$. Then there exist $\gamma_1 \neq \gamma_2 \in \text{Res}(b)$ arbitrarily close to 0 and $k_1, k_2 \in K$ not both 0 such that $k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|} \in A_{\deg_J(q)}$.*

Proof. Let $\Gamma = (\text{Res}(b) - \text{Big}^{\deg_J^r(b)}(r))$, which, as observed in the proof of Proposition 4.15, is infinite with supremum 0. Note moreover that $\deg_J(p) = \deg_J^p(b)$, thus in particular $\deg_J(p) \leq \deg_J^p(q)$.

If $\deg_J(p) < \deg_J^p(q)$, then as in the proof of Proposition 4.15 for every $\gamma_1 \neq \gamma_2 \in \Gamma$ close enough to 0 there exist $k_1, k_2 \in K^\times$ such that $k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|} \in J_{\deg_J(q)} \subseteq A_{\deg_J(q)}$.

Otherwise, $\deg_J(p) = \deg_J^p(q)$. For every $\gamma \in \Gamma$ close enough to 0, we have $b^{|\gamma|} = p^{|\gamma|}q + q^{|\gamma|}p + r_\gamma$ where $\deg_J(r_\gamma) < \deg_J^r(b)$ and $\deg_J(p^{|\gamma|}) < \deg_J(p) = \deg_J^p(q)$. In particular, $\deg_J(p^{|\gamma|}r_\gamma) < \deg_J(q)$ by Lemma 4.9. Therefore, for every $\gamma_1 \neq \gamma_2 \in \Gamma$ close enough to 0 we have

$$p^{|\gamma_2|}b^{|\gamma_1|} - p^{|\gamma_1|}b^{|\gamma_2|} = (q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})p \pmod{J_{\deg_J(q)}}.$$

As $\deg_J(p) = \deg_J^p(b)$, then by Lemma 4.11 there exist $k_1, k_2 \in K$ not both 0 and γ arbitrarily close to 0 such that

$$(p^{|\gamma_2|}b^{|\gamma_1|} - p^{|\gamma_1|}b^{|\gamma_2|})^{|\gamma|} = k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|} \pmod{J_{\deg_J(q)}}.$$

If $\deg_J(k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|}) < \deg_J(q)$ then we are done. Otherwise, we have

$$\deg_J(((q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})p)^{|\gamma|}) = \deg_J(q).$$

When γ is sufficiently small, for every $\varepsilon \in \text{cl}(\text{supp}(p))$ with $\gamma \leq \varepsilon < 0$ we have

$$\deg_J((q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})^{|\gamma-\varepsilon|}) < \deg_J(q), \quad \deg_J(p^{|\varepsilon|}) < \deg_J(p) = \deg_J^p(q),$$

with the former implied by Lemma 4.9. By a further application of Lemma 4.9, $\deg_J((q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})^{|\gamma-\varepsilon|}p^{|\varepsilon|}) < \deg_J(q)$. By Proposition 2.9,

$$((q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})p)^{|\gamma|} = (q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})^{|\gamma|}p \pmod{J_{\deg_J(q)}},$$

which implies that

$$\deg_J((q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})^{|\gamma|}p) = \deg_J(q).$$

As $0 < \deg_J(p) < \deg_J(q)$ we obtain that

$$(q^{|\gamma_1|}p^{|\gamma_2|} - q^{|\gamma_2|}p^{|\gamma_1|})^{|\gamma|}p \in A_{\deg_J(q)}$$

and hence $k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|} \in A_{\deg_J(q)}$ as required. \square

Corollary 4.17. *Let $\alpha = \omega^{\alpha_1} + \omega^{\alpha_2} + \omega^{\alpha_3}$ where $\alpha_1 \geq \alpha_2 \geq \alpha_3$. Let $b \in P_\alpha$ such that $Q(b)$. Then $b, \text{rv}(b)$ are irreducible.*

Proof. For $\alpha_2 > \alpha_3 > 0$ the conclusion follows from Propositions 4.7, 4.15; for $\alpha_3 = 0$, from Corollary 4.13. Therefore, we may assume that $\alpha_2 = \alpha_3 > 0$.

Suppose by contradiction that $b = pq + r$ where $\deg_J(r) < \deg_J(b)$ and $0 < \deg_J(p) \leq \deg_J(q)$. Let $\Gamma = (\text{Res}(b) - \text{Big}^{\deg_J^r(b)}(r))$. If $\deg_J^p(p) \neq \deg_J^p(q)$, then by Lemma 4.14 for every $\gamma \in \Gamma$ close enough to 0 we have $b^{|\gamma|} = p^{|\gamma|}q + r_\gamma$ or $b^{|\gamma|} = pq^{|\gamma|} + r_\gamma$, where $\deg_J(r_\gamma) < \deg_J^r(b)$, and we continue as in the proof of Proposition 4.15. Otherwise, $\alpha_2 = \alpha_3$ and $\deg_J(q) = \omega^{\alpha_1} + \omega^{\alpha_2} = \deg_J^r(b)$. By Lemma 4.16 there exist $\gamma_1 \neq \gamma_2 \in \text{Res}(b)$ close enough to 0 and $k_1, k_2 \in K$ not both 0 such that $k_1b^{|\gamma_1|} + k_2b^{|\gamma_2|} \in A_{\omega^{\alpha_1} + \omega^{\alpha_2}}$. As $Q(b^{|\gamma_1|}, b^{|\gamma_2|})$ holds this contradicts $(*)_{\omega^{\alpha_1} + \omega^{\alpha_2}}$. \square

Combining with Proposition 4.15 we obtain:

Corollary 4.18. *Let $\alpha = \sum_{i=1}^n \omega^{\alpha_i}$ where $\alpha_1 \geq \alpha_2 \geq \alpha_3$ and $\alpha_3 > \dots > \alpha_n$. Let b principal such that $\deg_J(b) = \alpha$ and $Q(b)$. Then $b, \text{rv}(b)$ are irreducible.*

By Proposition 3.2, we obtain the following:

Corollary 4.19. *Let $\alpha = \sum_{i=1}^n \omega^{\alpha_i}$ where $\alpha_1 \geq \alpha_2 \geq \alpha_3$ and $\alpha_3 > \dots > \alpha_n$. Let $b = \sum_{i=1}^m b_i t^{\gamma_i} + r$ where $\text{sup}(b) = 0$, $b_1, \dots, b_m \in P_\alpha$, $\deg(r) < \alpha$. If $Q(b_1, \dots, b_m)$ then b is irreducible.*

This concludes the proofs of Theorems 1.4, 1.8.

REFERENCES

- [Ber00] Alessandro Berarducci. Factorization in generalized power series. *Trans. Amer. Math. Soc.*, 352(2):553–577, 2000. doi:10.1090/S0002-9947-99-02172-8.
- [BKK06] Darko Biljakovic, Mikhail Kochetov, and Salma Kuhlmann. Primes and irreducibles in truncation integer parts of real closed fields. In *Logic in Tehran*, volume 26 of *Lect. Notes Log.*, pages 42–64. Assoc. Symbol. Logic, La Jolla, CA, 2006.
- [Con76] J. H. Conway. *On numbers and games*, volume No. 6 of *London Mathematical Society Monographs*. Academic Press [Harcourt Brace Jovanovich, Publishers], London-New York, 1976.

- [Gon86] Harry Gonshor. *An introduction to the theory of surreal numbers*, volume 110 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1986. doi:10.1017/CB09780511629143.
- [Hah07] H. Hahn. Über die nichtarchimedischen Größensysteme. *Wien. Ber.*, 116:601–655, 1907.
- [LM17] Sonia L’Innocente and Vincenzo Mantova. Factorisation of germ-like series. *J. Log. Anal.*, 9:Paper No. 3, 16, 2017. doi:10.4115/jla.2017.9.3.
- [LM24] Sonia L’Innocente and Vincenzo Mantova. A factorisation theory for generalised power series and omnific integers. *Advances in Mathematics*, 442:109513, 2024. doi:10.1016/j.aim.2024.109513.
- [MR93] M.-H. Mourgues and J. P. Ressayre. Every real closed field has an integer part. *J. Symbolic Logic*, 58(2):641–647, 1993. doi:10.2307/2275224.
- [Pit01] Daniel Pitteloud. Existence of prime elements in rings of generalized power series. *J. Symbolic Logic*, 66(3):1206–1216, 2001. doi:10.2307/2695102.
- [PS06] James Pommersheim and Shahriar Shahriari. Unique factorization in generalized power series rings. *Proc. Amer. Math. Soc.*, 134(5):1277–1287, 2006. doi:10.1090/S0002-9939-05-08162-1.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF FLORENCE, FLORENCE, ITALY

Email address: `antongiulio.fornasiero@unifi.it`

DEPARTMENT OF MATHEMATICS AND PHYSICS, ROMA TRE UNIVERSITY, ROME, ITALY

Email address: `noa.lavi@unicam.it`

SCHOOL OF SCIENCE AND TECHNOLOGY, MATHEMATICS DIVISION, UNIVERSITY OF CAMERINO, CAMERINO, ITALY

Email address: `sonia.linnocente@unicam.it`

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT LEEDS, UNITED KINGDOM

Email address: `v.l.mantova@leeds.ac.uk`