

Iarrobino's decomposition for self-dual modules

Maciej Wojtala

August 30, 2024

Abstract

We investigate self-dual modules and their Hilbert functions. We generalize Iarrobino's decomposition to this setup and deduce consequences for their local Hilbert functions. We classify the local Hilbert functions for small degree modules. We prove a generalization of Kunte's criterion for self-duality in terms of Macaulay's inverse systems.

1 Introduction

1.1 Main results

In our work we introduce three main results: Iarrobino's decomposition for self-dual modules, the generalization of Kunte's criterion for self-duality in terms of Macaulay's inverse systems and the classification of possible Hilbert functions of self-dual modules for degree less or equal to eight.

A. Iarrobino introduced Iarrobino's decomposition for self-dual algebras [22, 23]. It is used in the deformation theory [9], the theory of symmetric matrices [29], the classical algebraic geometry [38, 21], the theory of higher-dimensional Gorenstein algebras [13, 24]. This [22] work of A. Iarrobino is relevant for algebraic combinatorics, Lefschetz properties, theory of secant varieties and sums of powers etc. It is widely applied and cited over 110 times, including such classical books and papers as [41, 17, 18, 37, 34, 4, 20, 15, 44, 39, 31, 10]. It keeps being relevant today; it is employed for example in [25, 3, 35, 40, 16, 43, 5, 8, 26, 2, 14].

The main claim in this result is that the local Hilbert function of a self-dual algebra can be decomposed as a sum of symmetric sequences. This result introduces criteria that the local Hilbert function of a self-dual algebra must satisfy.

F. Macaulay introduced the bound on the Hilbert functions of algebras [33]. This result was generalized to modules by H. Hulett [19]. Iarrobino's decomposition in combination with Macaulay Bound is a very useful tool in showing that a sequence is not a local Hilbert function of any self-dual algebra.

We generalize the result of A.Iarrobino to the case of self-dual modules. For our best knowledge this is the first time that Iarrobino's decomposition is considered in terms of modules. Put informally, for a module M of the socle degree d , the local Hilbert function h_M is a sum

$$h_M = \sum_{i=0}^d \Delta_i,$$

where Δ_i is symmetric around $\frac{d-i}{2}$. We now introduce the necessary definitions (see also Section 3).

Let R be a local ring and let \mathfrak{m} be its maximal ideal. Let d be a maximal natural number d such that $\mathfrak{m}^d M \neq 0$. By $\text{gr } R$ we denote the associated graded ring of R , i.e.

$$\text{gr } R = \bigoplus_{k=0}^d \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}.$$

By $\text{gr } M$ we denote the associated graded module of the module M , i.e.

$$\text{gr } M = \bigoplus_{k=0}^d \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M},$$

which is a $(\text{gr } R)$ -module in a natural way. For an ideal \mathfrak{a} in R we denote by $(0 : \mathfrak{a})_M$ the annihilator of \mathfrak{a} in M , i.e. the set of all elements $x \in M$, such that $\mathfrak{a}x = 0$. The annihilator $(0 : \mathfrak{a})_M$ is a submodule of M .

Let us denote

$$Q_{k,l} = \frac{\mathfrak{m}^k M \cap (0 : \mathfrak{m}^l)_M}{\mathfrak{m}^{k+1} M \cap (0 : \mathfrak{m}^l)_M + \mathfrak{m}^k M \cap (0 : \mathfrak{m}^{l-1})_M} \text{ for all } 0 \leq k, l \leq d+1.$$

Let us also define

$$\Delta_s(t) = \begin{cases} \text{rank}_{\mathbb{K}} Q_{t, d+1-(s+t)} & \text{for } 0 \leq s \leq d, 0 \leq t \leq d-s \\ 0 & \text{otherwise.} \end{cases}$$

We have $\text{rank}_{\mathbb{K}} Q_{k,l} = \Delta_{d+1-(k+l)}(k)$.

We denote by h_M the local Hilbert function of the module M , i.e.

$$h_M(k) = \dim_{\mathbb{K}} \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M}.$$

The theorem is stated as follows:

Theorem 1.1 (Hilbert function decomposition). *If R is a local ring and M is a self-dual module, then the following statements hold true:*

1. We have the identity $Q_{k,0} = Q_{d+1,l} = 0$.
2. For $k+l > d+1$ we have the identity $A_{k,l} = A_{k,l-1}$ and so it holds that $Q_{k,l} = 0$.
3. It holds that $\Delta_s(t) = \Delta_s((d-s)-t)$. We say that Δ_s is symmetric with respect to $\frac{d-s}{2}$.
4. For $0 \leq a \leq d$ let us denote $h_a(t) = \sum_{i=0}^a \Delta_i(t)$. Then h_a is the Hilbert function of the quotient

$$\frac{\text{gr } M}{\bigoplus_{i=0}^a C_a^i},$$

where $C_a^i \subseteq \mathfrak{m}^i M / \mathfrak{m}^{i+1} M$ is the image of $A_{i,d-(i+a)}$.

5. The local Hilbert function h_M satisfies $h_M(t) = 0$ for $t > d$ and

$$h_M(t) = \sum_{i=0}^{d-t} \Delta_i(t) \text{ for } 0 \leq t \leq d.$$

As examples of applications of this theorem we show two claims that we also note in Section 5.

Example 1.2. • If h_M is a Hilbert function of a self-dual module M and k is the largest natural number such that $h_M(k) > 0$, then $h_M(0) \geq h_M(k)$.

Proof. Let us consider Iarrobino's decomposition of h_M – first row must be symmetric (from Theorem 1.1 Point 4) and must have $h_M(k)$ on the last position (since next rows are strictly shorter), so it also must have $h_M(k)$ on the first position. Since $h_M(0)$ is a sum of values on the first positions in all rows, we have $h_M(0) \geq h_M(k)$. \square

• Let h_M be a Hilbert function of a self-dual module M and let k be the largest natural number such that $h_M(k) > 0$. Then $\sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} h_M(i) \geq \sum_{i=\lceil \frac{k+1}{2} \rceil}^k h_M(i)$.

Proof. The claim states that the sum of the first half of the sequence is greater or equal to the sum of the second half. This follows from Iarrobino's decomposition (Theorem 1.1) – the sequence is a sum of symmetric parts and each of these parts contributes to the first half at least as much as it does to the second half. \square

J. Jelisiejew and K. Šivic introduced the concept of apolarity for modules, which is a very useful tool in generating examples of modules with given Hilbert function [28]. M. Kunte introduced a sufficient and necessary condition for a module to be a graded self-dual module [30]. We use the notion of apolarity to prove a generalization of Kunte's criterion to the non-graded case. However, we show the criterion with only a sufficient part, while the criterion by Kunte for the graded case was also necessary. The theorem is stated as follows: Let n be a positive natural number and let us denote $S^* = \mathbb{K}[x_1, x_2, \dots, x_n]$. Let us also fix a positive natural number r . We denote by F^* the free S^* -module $\bigoplus_{k=0}^r S^* e_k^*$. Note that each $f_k \in F^*$ is encoded by the direct sum decomposition as $\sum_{i=0}^r f_{ki} e_i^*$, where $f_{ki} \in S^*$. Thus, the tuple f_1, f_2, \dots, f_l can be viewed as a matrix $N = [f_{ki}] \in \mathbb{M}_{l \times r}(S^*)$. We call this matrix the encoding matrix of the module apolar to (f_1, f_2, \dots, f_l) .

Theorem 1.3 (Kunte's criterion). *If the square matrix $N = [f_{ki}] \in \mathbb{M}_{r \times r}(S^*)$ is symmetric, then the module apolar to $(f_k)_{k=1}^r$ is self-dual, where $f_k \in F^*$.*

A definition of the apolarity notion can be found in Section 4. The Kunte's criterion is a quite simple condition, as we only need to verify if the proper matrix is symmetric. This criterion, combined with apolarity for modules, allows us to easily generate examples of self-dual modules for a given Hilbert function. The special case of this theorem is $r = 1$. Then M becomes an algebra. Since the 1×1 matrix is trivially symmetric, it implies that every apolar algebra is self-dual.

Example 1.4. Let us take as the encoding matrix the diagonal matrix

$$\begin{bmatrix} x^2 & 0 \\ 0 & x \end{bmatrix}.$$

The obtained self-dual module has the local Hilbert function $(2, 2, 1)$. The method for computing the local Hilbert function from the encoding matrix is described in Section 4 (see Example 4.10). The reader might observe that it corresponds to the direct sum decomposition $(1, 1, 1) + (1, 1)$ (see Section 5).

We show other applications of this theorem in Example 4.10 and in the classification of the local Hilbert function $(2, 2, 3, 1)$ (see Section 5).

B. Poonen classified the local Hilbert functions and the types of isomorphisms for self-dual algebras of degree less or equal to six [36]. The corresponding problem for modules is mostly open. In our work we classify possible local Hilbert functions of self-dual modules with a small degree – we classify all possible Hilbert functions for a degree less or equal to 8.

Proposition 1.5. *The number of possible local Hilbert functions of self-dual modules for degree $m = 1, 2 \dots 8$ is exactly the following:*

Table 1: Number of possible local Hilbert functions of self-dual modules

Degree	Number of possible local Hilbert functions
$m = 1$	1
$m = 2$	2
$m = 3$	3
$m = 4$	6
$m = 5$	9
$m = 6$	16
$m = 7$	24
$m = 8$	38

The whole classification of possible local Hilbert functions of self-dual modules can be found in Section 5. This classification is interesting, as we show that some unexpected local Hilbert functions are possible to obtain from self-dual modules, for example the function $(2, 2, 3, 1)$.

1.2 Self-dual algebras and modules

Self-dual algebras (also known as Gorenstein algebras or commutative Frobenius algebras) are useful tools in algebraic complexity theory. Let us consider a tensor $t \in A \otimes B \otimes C$, where $A, B, C \cong \mathbb{C}^m$. The tensor t may be considered as a linear map $A^* \rightarrow B \otimes C$. We say that t is 1_A -generic if there exists an α such that $t(\alpha) : B^* \rightarrow C$ has full rank. We define analogically 1_B - and 1_C -genericity. We say that a tensor is 1-generic, if it is 1_A -, 1_B - and 1_C -generic. Let us assume that t is 1_A -generic. Let α be such an element that $t(\alpha)$ is invertible. Let us consider the space $t(A^*) \cdot t(\alpha)^{-1}$. We say that t satisfies the Strassen's equations, if the space $t(A^*) \cdot t(\alpha)^{-1}$ is abelian, i.e. if it consists of commuting matrices. More about Strassen's equations one can read

in [32]. A tensor is a structure tensor of an algebra A if it corresponds to the multiplication map $A \times A \rightarrow A$. The following result connects 1-generic tensors satisfying Strassen's equations and self-dual algebras:

Proposition 1.6. (*[27, Summary 2.5]*) *Let us assume that we have $A, B, C \cong \mathbb{C}^m$. A tensor $t \in A \otimes B \otimes C$ is 1-generic and satisfies Strassen's equations if and only if it is a structure tensor of a self-dual algebra.*

The self-dual algebras are also a useful tool for investigating secant varieties, see [7]. They are subjects of intensive research, see [12, 11].

In this work we investigate self-dual modules (see Definition 2.1) as they are also important in algebraic complexity theory. Deriving algebraic structure on tensors allows to analyze tensors of matrix multiplication and obtain bounds on ω [42, 1, 6, 45], which is one of the most important topics in theoretical computer science. M. Wojtala introduced the notion of structure tensors of modules [45]. The following result connects 1_A -generic tensors satisfying Strassen's equations and modules:

Proposition 1.7. (*[27, Summary 2.5], [32, Lemma 2.6], [45, Proof of Corollary 2]*) *Let us assume that we have $A, B, C \cong \mathbb{C}^m$. A tensor $t \in A \otimes B \otimes C$ is 1_A -generic and satisfies Strassen's equations if and only if it is a structure tensor of a module.*

For researchers working with tensors, it is usual to impose symmetry conditions. A tensor $t \in A \otimes A \otimes A$ is symmetric if it is invariant under the permutation of coordinates. By Proposition 1.6, one cannot hope for symmetric 1_A -generic tensors, which do not come from algebras. However, imposing the partial symmetry is actually connected to coming from self-dual modules. This connection is shown in the following proposition and can be viewed as an analogical result to Proposition 1.6 and Proposition 1.7.

Proposition 1.8. *Let us assume that we have $A, B \cong \mathbb{C}^m$. If a tensor $t \in A \otimes S^2(B)$ is 1_A -generic and satisfies Strassen's equations, then it is a structure tensor of a self-dual module.*

Proof. Proposition 1.7 implies that t is a structure tensor of some module M . Now let us observe that by symmetry of matrices from $t(A^*)$ and a reasoning from [28, Subsection 3.5] we have that M is isomorphic to M^* as modules. □

Acknowledgements

During the preparation of this publication, the author was part of the Szkoła Orłów program. The publication was created under the supervision of Joachim Jelisiejew, whose support and help were priceless.

2 Preliminaries

In this section we recall facts that will be useful for our purposes.

Let \mathbb{K} denote an algebraically closed field. Let also R denote a ring that is a finite-dimensional \mathbb{K} -vector space and let M denote an R -module that is a finite-dimensional \mathbb{K} -vector space. The *degree* of such a module M is defined to be the dimension of M as a \mathbb{K} -vector space.

On a dual vector space M^* we introduce an R -module structure by defining

$$r \cdot \phi(m) = \phi(rm).$$

Definition 2.1. A module M is self-dual if it is isomorphic with M^* as an R -module.

A case of interest is when R is a local ring and M is a self-dual module, although some results are shown without those assumptions.

Definition 2.2. For an ideal \mathfrak{a} in R we denote $\mathfrak{a}^z = (1)$ for any negative integer z .

2.1 Modules and submodules

Definition 2.3. For a submodule K of M we define K^\perp by the formula $K^\perp = \{\phi \in M^* : \phi(K) = 0\}$.

Lemma 2.4. For a module K we have that K^\perp is a submodule of M^* .

Proof. If $\phi_1, \phi_2 \in K^\perp$, then $\phi_1(K) = 0, \phi_2(K) = 0$, so $(\phi_1 + \phi_2)(K) = \phi_1(K) + \phi_2(K) = 0$, so $\phi_1 + \phi_2 \in K^\perp$. Also if $\phi \in K^\perp$, then for every $r \in R$ we have that $rK \subseteq K$ (since K is a submodule of M) and thus $(r \cdot \phi)(K) = \phi(rK) = 0$, so $r \cdot \phi \in K^\perp$. \square

Lemma 2.5. 1. Let K be a submodule of M . Then $\dim_{\mathbb{K}} K + \dim_{\mathbb{K}} K^\perp = \dim_{\mathbb{K}} M$.

2. Let K be a submodule of M . Then $(K^\perp)^\perp \cong K$.

3. Let K, L be submodules of M . Then we have $(K + L)^\perp = K^\perp \cap L^\perp$.

4. Let K, L be submodules of M . Then $(K \cap L)^\perp = K^\perp + L^\perp$.

Proof. Left to the reader. \square

Lemma 2.6. We have a canonical isomorphism $M \cong M^{**}$ given by the double dual map.

Proof. Let $\psi : M \rightarrow M^{**}$ be the double dual map, i.e. $\psi(m)(\phi) = \phi(m)$. Then ψ is an isomorphism of \mathbb{K} -linear spaces, we need to show that it is also an isomorphism of R -modules. Indeed, we have $s \cdot \psi(m)(\phi) = \psi(m)(s\phi) = (s\phi)(m) = \phi(sm) = \psi(s \cdot m)(\phi)$. \square

Lemma 2.7. Let I be a submodule of M . Then we have a canonical isomorphism $I^* \cong M^*/I^\perp$ induced by the canonical surjection.

Proof. Let $\pi : M^* \rightarrow I^*$ be the canonical surjection. Then $\ker \pi = \{\phi \in M^* : \phi(I) = 0\} = I^\perp$. So by the isomorphism theorem we have a canonical isomorphism $I^* \cong M^*/\ker \pi = M^*/I^\perp$. \square

Lemma 2.8. Let J be a submodule of I and I be a submodule of M . Then we have a canonical exact sequence $0 \rightarrow (I/J)^* \rightarrow I^* \rightarrow J^* \rightarrow 0$.

Proof. We obtain the canonical injection $(I/J)^* \rightarrow I^*$ by lifting functionals from $(I/J)^*$ to functionals from I^* , i.e. functional ϕ goes to functional $\bar{\phi}$, such that for $i \in I$, $i \equiv \bar{i} \pmod{J}$ we have $\bar{\phi}(i) = \phi(\bar{i})$. The image of this injection consists of all functionals that are zero on J . The canonical surjection $I^* \rightarrow J^*$ is obtained by restricting functionals on I to J , its kernel also consists of all functionals that are zero on J , so we constructed the desired canonical exact sequence. \square

Lemma 2.9. *Let J be a submodule of I and I be a submodule of M . Then $(I/J)^* \cong J^\perp/I^\perp$.*

Proof. Left to the reader. \square

Lemma 2.10. *Let M_1, M_2 be R -modules that are finite-dimensional \mathbb{K} -vector spaces. Let us assume that there exists a non-degenerate pairing $\langle \cdot, \cdot \rangle : M_1 \times M_2 \rightarrow \mathbb{K}$ such that for every $r \in R$, $a_1 \in M_1$, $a_2 \in M_2$, we have $\langle ra_1, a_2 \rangle = \langle a_1, ra_2 \rangle$. Then M_1 and M_2 are dual as R -modules.*

Proof. Left to the reader. \square

Definition 2.11. For an ideal \mathfrak{a} in R we denote by $(0 : \mathfrak{a})_M$ the annihilator of \mathfrak{a} in M , i.e. the set of all elements $x \in M$, such that $\mathfrak{a}x = 0$. It is easy to check that $(0 : \mathfrak{a})_M$ is a submodule of M .

Lemma 2.12. *Let \mathfrak{a} be an ideal in R . If M is self-dual, then we have the following isomorphism of R -modules: $(\mathfrak{a}M)^\perp \cong (0 : \mathfrak{a})_M$.*

Proof. We have $(\mathfrak{a}M)^\perp = \{\phi \in M^* : \phi(\mathfrak{a}M) = 0\} = \{\phi \in M^* : \forall s \in \mathfrak{a} s\phi = 0\} \cong \{l \in M : \forall s \in \mathfrak{a} sl = 0\} = (0 : \mathfrak{a})_M$, where the isomorphism follows from $M \cong M^*$. \square

2.2 Local rings

In this subsection we assume that R is a local ring and \mathfrak{m} is its only maximal ideal.

Corollary 2.13. *If M is self-dual, then we have the following isomorphisms of R -modules: $(\mathfrak{m}^k M)^\perp \cong (0 : \mathfrak{m}^k M)_M$, $((0 : \mathfrak{m}^k M)_M)^\perp \cong \mathfrak{m}^k M$.*

Proof. The first statement is just Lemma 2.12 for the ideal \mathfrak{m}^k . The second part is obtained from the first one by taking the perpendicular submodule on both sides and using Lemma 2.5 Point 2. \square

Remark 2.14. There exists the largest natural number d such that $\mathfrak{m}^d M \neq 0$.

2.3 Modularity and filtrations

Remark 2.15. Let A, B, C be submodules of M and let us assume that A is a submodule of C . Then $(A + B) \cap C = B \cap C + A$.

Corollary 2.16. *Let A_1, A_2, B_1, B_2 be submodules of M and let us also assume that A_1 is a submodule of B_1 and B_2 is a submodule of A_2 . Then $(A_1 + A_2) \cap (B_1 + B_2) = A_1 + B_2 + A_2 \cap B_1$.*

Proof. First we use Remark 2.15 for $A = A_1$, $B = A_2$, $C = (B_1 + B_2)$ and obtain $(A_1 + A_2) \cap (B_1 + B_2) = A_1 + A_2 \cap (B_1 + B_2)$. Then we use Remark 2.15 for $A = B_2$, $B = B_1$, $C = A_2$ and obtain $A_2 \cap (B_1 + B_2) = B_2 + B_1 \cap A_2$. \square

Remark 2.17. Let $F_0 \subseteq F_1 \subseteq \dots \subseteq F_c$ and $G_0 \subseteq G_1 \subseteq \dots \subseteq G_c$ be filtrations of R -modules, such that $F_k \subseteq G_k$ and $G_k \cap F_{k+1} = F_k$ for $k = 0, 1, \dots, c-1$. Then we have the filtration $\frac{G_0}{F_0} \subseteq \frac{G_1}{F_1} \subseteq \dots \subseteq \frac{G_c}{F_c}$ of R -modules.

2.4 Associated graded ring and module

In this subsection we assume that R is a local ring and \mathfrak{m} is its only maximal ideal. We denote by d the largest natural number such that $\mathfrak{m}^d \neq 0$, as in Remark 2.14.

Definition 2.18. By $\text{gr } R$ we denote the associated graded ring of R , i.e.

$$\text{gr } R = \bigoplus_{k=0}^d \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}.$$

The multiplication in this ring is defined as follows: for $a_k \in \mathfrak{m}^k$ let $\overline{a_k}$ denote the image of a_k in $\frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}$; then $\overline{a_i} \cdot \overline{a_j}$ is defined as $\overline{a_i \cdot a_j}$, i.e. it is the image of $a_i \cdot a_j$ in $\frac{\mathfrak{m}^{i+j}}{\mathfrak{m}^{i+j+1}}$.

By $\text{gr } M$ we denote the associated graded module of the module M , i.e.

$$\text{gr } M = \bigoplus_{k=0}^d \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M},$$

which is a $(\text{gr } R)$ -module in a natural way.

Definition 2.19. We denote by h_M the local Hilbert function of the module M , i.e.

$$h_M(k) = \dim_{\mathbb{K}} \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M}.$$

Let us denote $\mathfrak{n} = \bigoplus_{k>0} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}} \subseteq \text{gr } R$. Clearly \mathfrak{n} is an ideal in $\text{gr } R$. The following lemma shows that is the unique maximal ideal in $\text{gr } R$.

Lemma 2.20. *The associated graded ring $\text{gr } R$ is a local ring and \mathfrak{n} is its unique maximal ideal.*

Proof. We have

$$\frac{\text{gr } R}{\mathfrak{n}} = \frac{\text{gr } R}{\bigoplus_{k>0} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}} \cong \frac{R}{\mathfrak{m}}.$$

Since $\frac{R}{\mathfrak{m}}$ is a field, we obtain that \mathfrak{n} is a maximal ideal. Moreover, $\mathfrak{n}^{d+1} = \left(\bigoplus_{k>0} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}\right)^{d+1} = 0$, so since maximal ideals are prime, every maximal ideal must contain \mathfrak{n} . It follows that \mathfrak{n} is the only maximal ideal in $\text{gr } R$. \square

Lemma 2.21. *It holds that $\mathfrak{n}^l = \bigoplus_{k \geq l} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}$.*

Proof. Clearly $\mathfrak{n}^l = \left(\bigoplus_{k \geq 1} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}\right)^l \subseteq \bigoplus_{k \geq l} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}$. To show the opposite inclusion, let us fix an element $a \in \frac{\mathfrak{m}^{l+c}}{\mathfrak{m}^{l+c+1}}$, where $c \geq 0$. We can write a as a finite sum $\sum_j a_{1,j} \cdot a_{2,j} \cdot \dots \cdot a_{l+c,j} + \mathfrak{m}^{l+c+1}$,

where $a_{i,j} \in \mathfrak{m}$. Then we have $a = \sum_j a_{1,j} \cdot a_{2,j} \cdot \dots \cdot a_{l+c,j} + \mathfrak{m}^{l+c+1} = \sum_j (a_{1,j} + \mathfrak{m}^2) \cdot (a_{2,j} + \mathfrak{m}^2) \cdot \dots \cdot (a_{l+c,j} + \mathfrak{m}^2) \in (\frac{\mathfrak{m}}{\mathfrak{m}^2})^{l+c} \subseteq (\bigoplus_{k \geq 1} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}})^{l+c} \subseteq (\bigoplus_{k \geq 1} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}})^l$. \square

Corollary 2.22. *Let $C = \bigoplus_{k=0}^d \frac{C_k + \mathfrak{m}^{k+1}M}{\mathfrak{m}^{k+1}M}$ be a graded submodule of $\text{gr } M$, where $C_k \subseteq \mathfrak{m}^k M$ is an R -submodule. Then the local Hilbert function h_q of the quotient $\frac{\text{gr } M}{C}$ satisfies $h_q(k) = \dim_{\mathbb{K}} \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1}M + C_k}$ and the local Hilbert function h_C of the module C satisfies $h_C(k) = \dim_{\mathbb{K}} \frac{C_k + \mathfrak{m}^{k+1}M}{\mathfrak{m}^{k+1}M}$.*

Proof. by Lemma 2.21 it follows that

$$\frac{\mathfrak{m}^l}{\mathfrak{m}^{l+1}} = \frac{\bigoplus_{k \geq l} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}}{\bigoplus_{k \geq l+1} \frac{\mathfrak{m}^k}{\mathfrak{m}^{k+1}}} = \frac{\mathfrak{m}^l}{\mathfrak{m}^{l+1}}.$$

So the direct sum decompositions of $\text{gr } M$ coming from $(\frac{\mathfrak{m}^l}{\mathfrak{m}^{l+1}})_{l=0}^d$ and $(\frac{\mathfrak{m}^l}{\mathfrak{m}^{l+1}})_{l=0}^d$ are identical, so values of the Hilbert function can be obtained from the latter one, which we wanted to show. \square

The following remark states that the associated graded ring $\text{gr } R$ satisfies the same assumptions as R and $\text{gr } M$ satisfies the same assumptions as M (we do not assume that M is self-dual here). It implies that we can apply Macaulay's Bound to $\text{gr } M$. Note that $\text{gr } M$ in general might not be self-dual, so we cannot apply Iarrobino's decomposition to $\text{gr } M$.

Remark 2.23. If R is a local ring, then the associated graded ring $\text{gr } R$ is a local \mathbb{K} -algebra and is a finite-dimensional \mathbb{K} -vector space. The associated graded module $\text{gr } M$ is a finite-dimensional \mathbb{K} -vector space.

2.5 Macaulay's Bound

In this subsection we assume that R is a local ring. We recall now a criterion (thanks to Macaulay) that the local Hilbert functions of modules have to satisfy. Unlike Iarrobino's decomposition, this criterion is also valid for modules that are not self-dual. Before we recall the aforesaid Macaulay's Bound, we need to introduce one representation system.

Definition 2.24. Let a, s be positive integers. Then there exist uniquely determined positive integers $a_s > a_{s-1} > a_{s-2} > \dots > a_k \geq k > 0$, such that we have the equality

$$a = \binom{a_s}{s} + \binom{a_{s-1}}{s-1} + \binom{a_{s-2}}{s-2} + \dots + \binom{a_k}{k}.$$

We denote

$$a^{(s)} = \binom{a_s + 1}{s+1} + \binom{a_{s-1} + 1}{s} + \binom{a_{s-2} + 1}{s-1} + \dots + \binom{a_k + 1}{k+1}.$$

Theorem 2.25 (Macaulay's Bound, [19]). *We have the following bound on the local Hilbert function:*

$$h_M(r+1) \leq h_M(r)^{(r)}$$

for $r \geq 1$. The bound holds true for M that are not self-dual modules.

Corollary 2.26. *Let us assume that it holds $h_M(r) \leq r$. Then we have $h_M(r+1) \leq h_M(r)$.*

Proof. If $r = 0$, then the assumption states that $h_M(0) = 0$, so $M = 0$ and the claim holds true. Now let us assume that $r \geq 1$. Since we have $h_M(r) \leq r$, the representation of $h_M(r)$ is $\binom{r}{r} + \binom{r-1}{r-1} + \binom{r-2}{r-2} + \dots + \binom{r-h_M(r)+1}{r-h_M(r)+1}$. Thus $h_M(r)^{\langle r \rangle} = h_M(r)$ and by Macaulay's Bound 2.25 we obtain $h_M(r+1) \leq h_M(r)^{\langle r \rangle} = h_M(r)$. \square

Corollary 2.27. *Let us assume that it holds $h_M(r) \leq r+1$ and $r \geq 1$. Then we have $h_M(r+1) \leq r+2$.*

Proof. If $h_M(r) \leq r$, then from Corollary 2.26 we obtain the claim. If $h_M(r) = r+1$, then the representation of $h_M(r)$ is $\binom{r+1}{r}$ and thus $h_M(r)^{\langle r \rangle} = r+2$. So by Macaulay's Bound 2.25 we obtain $h_M(r+1) \leq h_M(r)^{\langle r \rangle} = r+2$. \square

3 Iarrobino's decomposition

In this section we introduce a new result – Iarrobino's decomposition for modules. Iarrobino's decomposition was known for algebras, but as far as we know, this is the first time when Iarrobino's decomposition has been considered for modules. With properly defined objects, the generalization from algebras to modules is direct, but there are some subtle differences, see Example 3.3. We assume in this section that R is a local ring and \mathfrak{m} is its only maximal ideal. Additionally, we assume that M is self-dual, i.e. M and M^* are isomorphic as R -modules.

Proposition 3.1. *If R is a local ring and M is a self-dual module, then we have the following isomorphism of \mathbb{K} -vector spaces*

$$\left(\frac{\mathfrak{m}^i M \cap (0 : \mathfrak{m}^j)_M}{\mathfrak{m}^{i+1} M \cap (0 : \mathfrak{m}^j)_M + \mathfrak{m}^i M \cap (0 : \mathfrak{m}^{j-1})_M} \right)^* \cong \frac{(0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M}{(0 : \mathfrak{m}^i)_M \cap \mathfrak{m}^{j-1} M + (0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^j M}.$$

Proof. Using Lemma 2.9, Lemma 2.5 Point 3 and Lemma 2.5 Point 4 respectively, we obtain

$$\begin{aligned} \left(\frac{\mathfrak{m}^i M \cap (0 : \mathfrak{m}^j)_M}{\mathfrak{m}^{i+1} M \cap (0 : \mathfrak{m}^j)_M + \mathfrak{m}^i M \cap (0 : \mathfrak{m}^{j-1})_M} \right)^* &\cong \frac{(\mathfrak{m}^{i+1} M \cap (0 : \mathfrak{m}^j)_M + \mathfrak{m}^i M \cap (0 : \mathfrak{m}^{j-1})_M)^\perp}{(\mathfrak{m}^i M \cap (0 : \mathfrak{m}^j)_M)^\perp} \cong \\ &\cong \frac{(\mathfrak{m}^{i+1} M \cap (0 : \mathfrak{m}^j)_M)^\perp \cap (\mathfrak{m}^i M \cap (0 : \mathfrak{m}^{j-1})_M)^\perp}{(\mathfrak{m}^i M \cap (0 : \mathfrak{m}^j)_M)^\perp} \cong \\ &\cong \frac{\left((\mathfrak{m}^{i+1} M)^\perp + ((0 : \mathfrak{m}^j)_M)^\perp \right) \cap \left((\mathfrak{m}^i M)^\perp + ((0 : \mathfrak{m}^{j-1})_M)^\perp \right)}{(\mathfrak{m}^i M)^\perp + ((0 : \mathfrak{m}^j)_M)^\perp}. \end{aligned} \tag{1}$$

Now from Corollary 2.13 we have

$$\begin{aligned} \frac{\left((\mathfrak{m}^{i+1} M)^\perp + ((0 : \mathfrak{m}^j)_M)^\perp \right) \cap \left((\mathfrak{m}^i M)^\perp + ((0 : \mathfrak{m}^{j-1})_M)^\perp \right)}{(\mathfrak{m}^i M)^\perp + ((0 : \mathfrak{m}^j)_M)^\perp} &\cong \\ &\cong \frac{((0 : \mathfrak{m}^{i+1})_M + \mathfrak{m}^j M) \cap ((0 : \mathfrak{m}^i)_M + \mathfrak{m}^{j-1} M)}{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M}. \end{aligned} \tag{2}$$

From Corollary 2.16 we obtain

$$\begin{aligned} & \frac{((0 : \mathfrak{m}^{i+1})_M + \mathfrak{m}^j M) \cap ((0 : \mathfrak{m}^i)_M + \mathfrak{m}^{j-1} M)}{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M} \cong \\ & \cong \frac{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M + (0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M}{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M}. \end{aligned} \quad (3)$$

Now we can rewrite

$$\begin{aligned} & \frac{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M + (0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M}{(0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M} \cong \\ & \cong \frac{(0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M}{((0 : \mathfrak{m}^i)_M + \mathfrak{m}^j M) \cap ((0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M)} \cong \\ & \cong \frac{(0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^{j-1} M}{(0 : \mathfrak{m}^i)_M \cap \mathfrak{m}^{j-1} M + (0 : \mathfrak{m}^{i+1})_M \cap \mathfrak{m}^j M}. \end{aligned} \quad (4)$$

□

Let d be a maximal natural number d such that $\mathfrak{m}^d M \neq 0$ (d exists as we observe in Remark 2.14). Let us denote

$A_{k,l} = \mathfrak{m}^k M \cap (0 : \mathfrak{m}^l)_M$ for all integers k, l (we follow the convention from Definition 2.2),

$$Q_{k,l} = \frac{A_{k,l}}{A_{k+1,l} + A_{k,l-1}} \text{ for all } 0 \leq k, l \leq d+1.$$

Let us also define

$$\Delta_s(t) = \begin{cases} \text{rank}_{\mathbb{K}} Q_{t, d+1-(s+t)} & \text{for } 0 \leq s \leq d, 0 \leq t \leq d-s \\ 0 & \text{otherwise.} \end{cases}$$

Remark 3.2. We have $\text{rank}_{\mathbb{K}} Q_{k,l} = \Delta_{d+1-(k+l)}(k)$.

Theorem (Theorem 1.1 – Hilbert function decomposition). *If R is a local ring and M is a self-dual module, then the following statements hold true:*

1. We have the identity $Q_{k,0} = Q_{d+1,l} = 0$.
2. For $k+l > d+1$ we have the identity $A_{k,l} = A_{k,l-1}$ and so it holds that $Q_{k,l} = 0$.
3. It holds that $\Delta_s(t) = \Delta_s((d-s)-t)$. We say that Δ_s is symmetric with respect to $\frac{d-s}{2}$.
4. For $0 \leq a \leq d$ let us denote $h_a(t) = \sum_{i=0}^a \Delta_i(t)$. Then h_a is the Hilbert function of the quotient

$$\frac{\text{gr } M}{\bigoplus_{i=0}^a C_a^i},$$

where $C_a^i \subseteq \mathfrak{m}^i M / \mathfrak{m}^{i+1} M$ is the image of $A_{i,d-(i+a)}$.

5. The local Hilbert function h_M satisfies $h_M(t) = 0$ for $t > d$ and

$$h_M(t) = \sum_{i=0}^{d-t} \Delta_i(t) \text{ for } 0 \leq t \leq d.$$

Proof of Theorem 1.1. 1. We have $(0 : \mathfrak{m}^0)_M = (0 : \mathfrak{m}^{-1})_M$, so $A_{k,0} = A_{k,-1}$ and thus $Q_{k,0} = 0$. Also, since $\mathfrak{m}^{d+1} = 0$, we have that $A_{d+1,l} = 0$, so $Q_{d+1,l} = 0$.

2. Clearly, we have $A_{k,l} \subseteq A_{k,l-1}$, so let us prove the other inclusion. Let us assume that $x \in A_{k,l-1}$. Then $x \in \mathfrak{m}^k M$, so $\mathfrak{m}^{l-1} x \in \mathfrak{m}^{k+l-1} M$. Since by the assumption $k+l-1 \geq d+1$, it holds that $\mathfrak{m}^{k+l-1} = 0$, so $\mathfrak{m}^{l-1} x = 0$. That proves that $x \in (0 : \mathfrak{m}^{l-1})_M$, so indeed $A_{k,l-1} \subseteq A_{k,l}$.

3. Proposition 3.1 states that $Q_{k,l}^* \cong Q_{l-1,k+1}$, so we have

$$\Delta_s(t) = \text{rank}_{\mathbb{K}} Q_{t,d+1-(s+t)} = \text{rank}_{\mathbb{K}} Q_{t,d+1-(s+t)}^* = \text{rank}_{\mathbb{K}} Q_{d-s-t,t+1} = \Delta_s((d-s)-t).$$

4. Let us denote $C_a = \bigoplus_{i=0}^a C_a^i$. First we need to argue that C_a is actually a submodule of $\text{gr } M$. Fix i . We have that C_a^i is the image of $A_{i,d-(i+a)}$. For any $k, i \geq 0$ we have

$$\begin{aligned} \mathfrak{m}^k \cdot A_{i,d-(i+a)} &= \mathfrak{m}^k \cdot (\mathfrak{m}^i M \cap (0 : \mathfrak{m}^{d-(i+a)})_M) = \mathfrak{m}^{k+i} M \cap (\mathfrak{m}^k \cdot (0 : \mathfrak{m}^{d-(i+a)})_M) \subseteq \\ &\subseteq \mathfrak{m}^{k+i} M \cap (0 : \mathfrak{m}^{d-(i+k+a)})_M = A_{i+k,d-((i+k)+a)}, \end{aligned}$$

so indeed we obtain a submodule of the associated graded module.

Denote $e = d - (k + a)$. Observe that we have

$$\begin{aligned} \sum_{i=0}^a \Delta_i(k) &= \sum_{i=0}^{d+1-k-(e+1)} \Delta_i(k) = \sum_{i=0}^{d+1-k-(e+1)} \text{rank}_{\mathbb{K}} Q_{k,d+1-(i+k)} = \\ &= \sum_{l=e+1}^{d+1-k} \text{rank}_{\mathbb{K}} Q_{k,l}. \end{aligned}$$

Using Point 2, we have

$$\sum_{l=e+1}^{d+1-k} \text{rank}_{\mathbb{K}} Q_{k,l} = \sum_{l=e+1}^{d+1} \text{rank}_{\mathbb{K}} Q_{k,l},$$

so we need to show that it holds

$$h_a(k) = \sum_{l=e+1}^{d+1} \text{rank}_{\mathbb{K}} Q_{k,l}.$$

We have the filtrations

$$A_{k,e} \subseteq A_{k,e+1} \subseteq A_{k,e+2} \subseteq \dots \subseteq A_{k,d} \subseteq A_{k,d+1} = \mathfrak{m}^k M,$$

$$A_{k+1,e} + A_{k,e} \subseteq A_{k+1,e+1} + A_{k,e} \subseteq \dots \subseteq A_{k+1,d+1} + A_{k,e} = \mathfrak{m}^{k+1}M + A_{k,e}.$$

We have that $A_{k+1,e+c} + A_{k,e} \subseteq A_{k,e+c}$ for $c = 0, 1, \dots, d+1-e$. Also Remark 2.15 gives us $A_{k,e+c} \cap (A_{k+1,e+c+1} + A_{k,e}) = A_{k,e+c} \cap A_{k+1,e+c+1} + A_{k,e}$. We can then rewrite $A_{k,e+c} \cap A_{k+1,e+c+1} + A_{k,e} = \mathfrak{m}^k M \cap (0 : \mathfrak{m}^{e+c})_M \cap \mathfrak{m}^{k+1} M \cap (0 : \mathfrak{m}^{e+c+1})_M + A_{k,e} = \mathfrak{m}^{k+1} M \cap (0 : \mathfrak{m}^{e+c})_M + A_{k,e} = A_{k+1,e+c} + A_{k,e}$ for $c = 0, 1, \dots, d-e$. Thus by Remark 2.17 we have the filtration

$$0 = \frac{A_{k,e}}{A_{k+1,e} + A_{k,e}} \subseteq \frac{A_{k,e+1}}{A_{k+1,e+1} + A_{k,e}} \subseteq \dots \subseteq \frac{A_{k,d+1}}{A_{k+1,d+1} + A_{k,e}} = \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M + A_{k,e}}.$$

So after using Corollary 2.22 we have

$$\begin{aligned} h_a(k) &= \dim_{\mathbb{K}} \frac{\mathfrak{m}^k M}{\mathfrak{m}^{k+1} M + A_{k,e}} = \sum_{c=0}^{d-e} \dim_{\mathbb{K}} \left(\frac{A_{k,e+c+1}}{A_{k+1,e+c+1} + A_{k,e}} \right) / \left(\frac{A_{k,e+c}}{A_{k+1,e+c} + A_{k,e}} \right) = \\ &= \sum_{c=0}^{d-e} \dim_{\mathbb{K}} \frac{A_{k,e+c+1}}{A_{k+1,e+c+1} + A_{k,e+c}} = \sum_{c=0}^{d-e} \dim_{\mathbb{K}} Q_{k,e+c+1} = \sum_{l=e+1}^{d+1} \dim_{\mathbb{K}} Q_{k,l}, \end{aligned}$$

which we wanted to obtain.

5. We use Point 4 with $a = d$. We only need to show that $\bigoplus_{i=0}^a C_a^i = 0$. We know that each C_a^i is the image of $A_{i,d-(i+d)} = A_{i,-i} = \mathfrak{m}^i M \cap (0 : \mathfrak{m}^{-i})_M$. Since $-i$ is a non-positive integer, $\mathfrak{m}^{-i} = (1)$, so $(0 : \mathfrak{m}^{-i})_M = 0$. Thus $A_{i,-i} = 0$, which follows that each C_a^i is zero. \square

We show a handful of examples of applications of Iarrobino's decomposition in Section 5, but now we want to show a slight difference to the case of self-dual algebras.

Example 3.3. Iarrobino's decomposition for algebras asserts also that $Q_{0,l} = 0$ for $l < d+1$. This is not the case for modules. Let us consider the sequence $(2, 1)$. We show in Section 5 that there exists a self-dual module with a local Hilbert function equal to this sequence. The decomposition of this function must be $(1, 1) + (1)$, which gives us $1 = \Delta_1(0) = \text{rank}_{\mathbb{K}} Q_{0,1}$, so $Q_{0,1} \neq 0$ and $1 < d+1 = 2$.

Proposition 3.4. *Let us define the vector space*

$$D_s = \bigoplus_{0 \leq t \leq d-s} Q_{t,d+1-(s+t)} = \bigoplus_{k+l=d+1-s} Q_{k,l}.$$

Then D_s is a self-dual module over the associated graded ring $\text{gr } R$.

Proof. First, we prove that D_s is indeed a module. For a given $r \in \mathfrak{m}^i$ multiplication by r induces a linear function from $A_{k,l}$ to $A_{k+i,l-i}$. After dividing by $A_{k+1,l} + A_{k,l-1}$ it induces a linear function from $Q_{k,l}$ to $Q_{k+i,l-i}$. Moreover, multiplication by \mathfrak{m}^{i+1} acts zero on $Q_{k,l}$, so we obtain a linear function from $\frac{\mathfrak{m}^i}{\mathfrak{m}^{i+1}} \times Q_{k,l}$ to $Q_{k+i,l-i}$. Suming over i we obtain a linear function from $\text{gr } R \times Q_{k,l}$ to $\bigoplus_{i \geq 0} Q_{k+i,l-i}$. Let us fix s . Suming over $k+l = d+1-s$ we obtain a

linear function from $\text{gr } R \times D_s$ to D_s . It is straightforward that this function satisfies the module axioms as it comes from multiplication by r . Proposition 3.1 states that $Q_{k,l}^* \cong Q_{l-1,k+1}$ as \mathbb{K} -vector spaces. Suming over $k+l = d+1-s$ we obtain isomorphism $D_s \cong D_s^*$ of \mathbb{K} -vector spaces. Moreover, the isomorphism from Proposition 3.1 is consistent with multiplication (in the proof we use two isomorphism: the canonical isomorphism from Lemma 2.9 and the R -modules isomorphism from Corollary 2.13, both of them are consistent with the multiplication), so we obtain the $\text{gr } R$ -modules isomorphism. \square

Remark 3.5. Taking in Theorem 1.1 Point 4 $a = 0$, we obtain a surjection from $\text{gr } M$ to D_0 . Moreover, if Iarrobino's decomposition is trivial (i.e. we have only one row of the decomposition), we have $\dim_{\mathbb{K}} \frac{m^k M}{m^{k+1} M} = h_0(k) = \dim_{\mathbb{K}} Q_{k,d-k}$ and the surjection becomes an isomorphism.

4 Apolarity

In this section we recall the definition of an apolar module. Then we prove the new result – Kunte's criterion for self-duality. The criterion was previously known in the graded case, we show a generalization to the non-graded case. However, our result is only the sufficient statement, while in the graded case the criterion was sufficient and necessary.

Let n be a positive natural number and let us denote $S = \mathbb{K}[y_1, y_2, \dots, y_n]$ and $S^* = \mathbb{K}[x_1, x_2, \dots, x_n]$. Let us also fix a positive natural number r . We denote by F^* the free S^* -module $\bigoplus_{k=0}^r S^* e_k^*$ and by F the free S -module $\bigoplus_{k=0}^r S e_k$. We introduce the S -module structure on F^* in the following manner: for $y_1^{a_1} y_2^{a_2} \dots y_n^{a_n} \in S$ and $x_1^{b_1} x_2^{b_2} \dots x_n^{b_n} \in S^*$ we define

$$y_1^{a_1} y_2^{a_2} \dots y_n^{a_n} \cdot x_1^{b_1} x_2^{b_2} \dots x_n^{b_n} = \begin{cases} 0, & \text{if there exists } k \text{ such that } b_k < a_k \\ x_1^{b_1-a_1} x_2^{b_2-a_2} \dots x_n^{b_n-a_n} & \text{in other cases.} \end{cases}$$

Note that in this definition multiplication behaves as a derivation, where $y_k^{a_k}$ encodes differentiating a_k times over x_k and we normalize the multiplication constant.

Now for $f_k \in S^*$ we define

$$y_1^{a_1} y_2^{a_2} \dots y_n^{a_n} \cdot \left(\sum_{k=1}^r f_k e_k^* \right) = \sum_{k=1}^r (y_1^{a_1} y_2^{a_2} \dots y_n^{a_n} \cdot f_k) e_k^*,$$

and we extend this definition to multiplication by the whole S using linearity.

We also introduce a bilinear form $\langle \cdot, \cdot \rangle : F \times F^* \rightarrow \mathbb{K}$ by defining

$$\langle y_1^{a_1} y_2^{a_2} \dots y_n^{a_n} e_s, x_1^{b_1} x_2^{b_2} \dots x_n^{b_n} e_l^* \rangle = \begin{cases} 1, & \text{if } s = l \text{ and for all } k \text{ we have } a_k = b_k \\ 0 & \text{in other cases} \end{cases}$$

and extending it to the bilinear form by linearity. We define \perp with respect to this form, i.e. for an S -submodule $I \subseteq F$ we have that I^\perp is an S^* -submodule of F^* consisting of all elements e , such that $\langle I, e \rangle = 0$. Analogically, for an S^* -submodule $J \subseteq F^*$ we have that J^\perp is an S -submodule of F consisting of all elements e , such that $\langle e, J \rangle = 0$.

Definition 4.1. For an element $f \in F^*$ we define the module apolar to f by $\frac{F}{(Sf)^\perp}$, where \perp refers to the introduced bilinear form (i.e. $(Sf)^\perp$ is an S^* -submodule consisting of all elements e such that $\langle e, Sf \rangle = 0$).

Similarly, for a tuple of elements $f_1, f_2, \dots, f_l \in F^*$ we define the module apolar to (f_1, f_2, \dots, f_l) by $\frac{F}{(Sf_1+Sf_2+\dots+Sf_l)^\perp}$.

Theorem 4.2. *Every finitely-generated module M can be obtained as an apolar module.*

Proof. Since the module is finitely-generated, we can write $M = \frac{F^r}{K}$ for some module K . Let us take K^\perp . It is also finitely-generated, so we can take its generators f_1, f_2, \dots, f_s . Now by Lemma 2.5 Point 2 we have $(K^\perp)^\perp \cong K$, so we have $M = \frac{F^r}{K} \cong \frac{F^r}{(Sf_1+Sf_2+\dots+Sf_s)^\perp}$. \square

Note that each $f_k \in F^*$ is encoded by the direct sum decomposition as $\sum_{i=0}^r f_{ki}e_i^*$, where $f_{ki} \in S^*$. Thus the tuple f_1, f_2, \dots, f_l can be viewed as a matrix $N = [f_{ki}] \in \mathbb{M}_{l \times r}(S^*)$. We call this matrix the encoding matrix of the module apolar to (f_1, f_2, \dots, f_l) . Note also that we are interested in the case of self-dual modules, so we would like to have a tool that would allow us to determine if the obtained apolar module is self-dual. It turns out that there is a very simple criterion for that in the case $l = r$, so if N is square. This criterion was introduced by Kunte in the graded case [30, Theorem 1.1], we generalize it to the non-graded case. However, our condition is only sufficient, while the one from the graded case was also necessary.

Theorem (Kunte's criterion). *If $r = l$ and the square matrix $N = [f_{ki}] \in \mathbb{M}_{r \times r}(S^*)$ is symmetric, then the module apolar to $(f_k)_{k=1}^r$ is self-dual.*

Before proving the theorem, we show a more general result. Let us recall that we have fixed a natural number r and defined free modules $F = \bigoplus_{k=0}^r Se_k$ and $F^* = \bigoplus_{k=0}^r S^*e_k^*$. Let us also fix a natural number l and define $G = \bigoplus_{k=0}^l Se_k$, $G^* = \bigoplus_{k=0}^l S^*e_k^*$. Let us also recall that we are considering a tuple $f_1, f_2, \dots, f_l \in F^*$ and we are using a decomposition $f_k = \sum_{i=1}^r f_{ki}e_i^*$ and encoding this tuple as a matrix $N = [f_{ki}]$. Let us now define a tuple $f_1^T, f_2^T, \dots, f_r^T \in G^*$ such that $f_i^T = \sum_{k=1}^l f_{ki}e_k^*$. Then the matrix encoding the tuple $f_1^T, f_2^T, \dots, f_r^T$ is N^T . We show that the modules apolar to (f_1, f_2, \dots, f_l) and $(f_1^T, f_2^T, \dots, f_r^T)$ are dual. It shows that the dual module of an apolar module is obtained by transposing the encoding matrix. In particular, if the matrix N is symmetric, then this matrix is invariant under transposition, which implies that the module is self-dual.

Theorem 4.3. *The modules apolar to (f_1, f_2, \dots, f_l) and $(f_1^T, f_2^T, \dots, f_r^T)$ are dual.*

The result follows from two simpler lemmas.

Lemma 4.4. *We have an S -modules isomorphism $\frac{F}{(Sf_1+Sf_2+\dots+Sf_l)^\perp} \cong (Sf_1+Sf_2+\dots+Sf_l)^*$.*

Proof. Let $c > \max_i \deg f_i$. Then $Sf_1+Sf_2+\dots+Sf_l \subseteq F_{\leq c-1}$ and thus $(Sf_1+Sf_2+\dots+Sf_l)^\perp \supseteq F_{\geq c}$, so we have

$$\frac{F}{(Sf_1+Sf_2+\dots+Sf_l)^\perp} \cong \frac{\frac{F}{F_{\geq c}}}{(Sf_1+Sf_2+\dots+Sf_l)^\perp}.$$

Now let us observe that the pairing $F \times F^* \rightarrow \mathbb{K}$ induces the non-degenerate pairing $\frac{F}{F_{\geq c}} \times F_{\leq c-1}^* \rightarrow \mathbb{K}$, so we have an S -module isomorphism $\frac{F}{F_{\geq c}} \cong (F_{\leq c-1}^*)^*$. Since also $Sf_1 + Sf_2 + \dots + Sf_l$ is a submodule of $F_{\leq c-1}^*$, we can use Lemma 2.7 for the module $F_{\leq c-1}^*$ and the submodule $Sf_1 + Sf_2 + \dots + Sf_l$, obtaining

$$\frac{(F_{\leq c-1}^*)^*}{(Sf_1 + Sf_2 + \dots + Sf_l)^\perp} \cong (Sf_1 + Sf_2 + \dots + Sf_l)^*.$$

Since we have $(F_{\leq c-1}^*)^* \cong \frac{F}{F_{\geq c}}$ we obtain the isomorphism

$$\frac{\frac{F}{F_{\geq c}}}{(Sf_1 + Sf_2 + \dots + Sf_l)^\perp} \cong (Sf_1 + Sf_2 + \dots + Sf_l)^*.$$

As we have already shown, it also holds $\frac{F}{(Sf_1 + Sf_2 + \dots + Sf_l)^\perp} \cong \frac{\frac{F}{F_{\geq c}}}{(Sf_1 + Sf_2 + \dots + Sf_l)^\perp}$, which implies the claim. \square

Lemma 4.5. *We have an S -modules isomorphism $(Sf_1 + Sf_2 + \dots + Sf_l) \cong (Sf_1^T + Sf_2^T + \dots + Sf_r^T)^*$.*

Proof. Let f be a column vector of f_i and let f_T be a column vector of (f_i^T) . By Lemma 2.10 it is sufficient to show that there exists a non-degenerate pairing

$$\langle \cdot, \cdot \rangle : (Sf_1 + Sf_2 + \dots + Sf_l) \times (Sf_1^T + Sf_2^T + \dots + Sf_r^T) \rightarrow \mathbb{K},$$

such that for every $s \in S$, $\sigma_1 \in S^l$, $\sigma_2 \in S^l$ we have $\langle s \cdot \sigma_1^T f, \sigma_2^T f_T \rangle = \langle \sigma_1^T f, s \cdot \sigma_2^T f_T \rangle$.

Let N be a matrix encoding f_1, f_2, \dots, f_l , i.e. $N = [f_{ki}]$. Let $\pi : S^* \rightarrow \mathbb{K}$ be a functional that sends 1 in S^* to 1 in \mathbb{K} . For elements $\sigma_1 = (\sigma_{11}, \sigma_{12}, \dots, \sigma_{1l}) \in S^l$ and $\sigma_2 = (\sigma_{21}, \sigma_{22}, \dots, \sigma_{2r}) \in S^r$, we consider the pairing

$$\langle \sigma_{11}f_1 + \sigma_{12}f_2 + \dots + \sigma_{1l}f_l, \sigma_{21}f_1^T + \sigma_{22}f_2^T + \dots + \sigma_{2r}f_r^T \rangle = \pi \left(\sum_{i,k} \sigma_{1i}\sigma_{2k} f_{ik} \right),$$

which can be written in a more compact form as

$$\langle \sigma_1^T f, \sigma_2^T f_T \rangle = \pi(\sigma_1^T N \sigma_2).$$

Then for $s \in S$ we have $\langle s \cdot \sigma_1^T f, \sigma_2^T f_T \rangle = \langle (s \cdot \sigma_1^T) f, \sigma_2^T f_T \rangle = \pi(s \cdot \sigma_1^T N \sigma_2)$. Similarly, $\langle \sigma_1^T f, s \cdot \sigma_2^T f_T \rangle = \langle \sigma_1^T f, (s \cdot \sigma_2^T) f_T \rangle = \pi(\sigma_1^T N \cdot s \cdot \sigma_2) = \pi(s \cdot \sigma_1^T N \sigma_2)$, where in the last equality we used commutativity of S . We show now that this pairing is also non-degenerate.

Let us observe that the rows of the matrix N are vectors f_k and its columns are vectors f_i^T . Thus we have $\sigma_1^T N = \sigma_1^T f$ and $N \sigma_2 = f_T^T \sigma_2 = (\sigma_2^T f_T)^T$. So if it holds that $\sigma_1^T f \neq 0$, then it also holds that $\sigma_1^T N \neq 0$. Thus there exists such an element σ_2 that $(\sigma_1^T N) \sigma_2 = 1$: we can take such an index j that $(\sigma_1^T N)_j \neq 0$, then there exists an element $a \in S$ such that $a \cdot (\sigma_1^T N)_j = 1$ and we can take $\sigma_{2j} = a$ and $\sigma_{2s} = 0$ for $s \neq j$. Then $\pi(\sigma_1^T N \sigma_2) = \pi(1) = 1$. Similarly if it holds that $\sigma_2^T f \neq 0$, then it also holds that $N \sigma_2 \neq 0$. Thus there exists such an element σ_1 that

$\sigma_1^T(N\sigma_2) = 1$ (analogical argument as above). Then $\pi(\sigma_1^T N \sigma_2) = \pi(1) = 1$. So the pairing is indeed non-degenerate. \square

Proof of Theorem 4.3. We want to show duality between $\frac{F}{(Sf_1+Sf_2+\dots+Sf_l)^\perp}$ and $\frac{G}{(Sf_1^T+Sf_2^T+\dots+Sf_r^T)^\perp}$. Using Lemma 4.4 for the tuple (f_1, f_2, \dots, f_l) and the module F and for the tuple $(f_1^T, f_2^T, \dots, f_r^T)$ and the module G we obtain $\frac{F}{(Sf_1+Sf_2+\dots+Sf_l)^\perp} \cong (Sf_1+Sf_2+\dots+Sf_l)^*$ and $\frac{G}{(Sf_1^T+Sf_2^T+\dots+Sf_r^T)^\perp} \cong (Sf_1^T+Sf_2^T+\dots+Sf_r^T)^*$. So we want to show $(Sf_1+Sf_2+\dots+Sf_l)^** \cong (Sf_1^T+Sf_2^T+\dots+Sf_r^T)^*$. By Lemma 2.6 we have $(Sf_1+Sf_2+\dots+Sf_l)** \cong Sf_1+Sf_2+\dots+Sf_l$, so we need to show $(Sf_1+Sf_2+\dots+Sf_l) \cong (Sf_1^T+Sf_2^T+\dots+Sf_r^T)^*$, which is true thanks to Lemma 4.5. \square

Proof of Theorem 1.3. If N is symmetric, then N is invariant under transposition, which by Theorem 4.3 implies that the module apolar to $(f_k)_{k=1}^r$ is self-dual. \square

Let us now introduce a tool that simplifies the computation of the local Hilbert function of an apolar module. Formally, for an apolar module M we consider it as a module over a ring $S_M = \frac{S}{\text{Ann}(M)}$, which is a local ring with the unique maximal ideal $S_{\geq 1} \cdot S_M$. Clearly, if M is self-dual as an S -module, then it is also self-dual as an S_M -module. By definition we have $h_M(k) = \dim_{\mathbb{K}} \frac{S_{\geq k} M}{S_{\geq k+1} M}$. However, computing the local Hilbert function from this formula is usually inconvenient and thus we propose another approach. First we need to introduce one definition.

Definition 4.6. For an element $f = \sum_{i=1}^r f_i e_i^* \in F$ we define a degree of f by taking $\deg f = \max_{i=1, 2, \dots, r} \deg f_i$.

Remark 4.7. For every non-negative integer c and every element $f \in F^*$ we have the following equivalence: $\deg f < c \iff S_{\geq c} f = 0$.

Proposition 4.8. *The local Hilbert function h_M of an apolar finite-dimensional self-dual module M satisfies the following formula:*

$$h_M(k) = \dim_{\mathbb{K}} \{f \in M : \deg f \leq k\} - \dim_{\mathbb{K}} \{f \in M : \deg f \leq k-1\}.$$

Proof. Remark 4.7 implies that we have the equality $\{f \in M : \deg f \leq k\} = (0 : S_{\geq k+1})_M$. From Corollary 2.13 we obtain that $(0 : S_{\geq k+1})_M \cong (S_{\geq k+1})_M^\perp$, where \perp refers to Definition 2.3. Now we can rewrite using Lemma 2.5 Point 1: $\dim_{\mathbb{K}} \{f \in M : \deg f \leq k\} - \dim_{\mathbb{K}} \{f \in M : \deg f \leq k-1\} = \dim_{\mathbb{K}} (S_{\geq k+1})_M^\perp - \dim_{\mathbb{K}} (S_{\geq k})_M^\perp = (\dim_{\mathbb{K}} M - \dim_{\mathbb{K}} (S_{\geq k+1})_M) - (\dim_{\mathbb{K}} M - \dim_{\mathbb{K}} (S_{\geq k})_M) = \dim_{\mathbb{K}} (S_{\geq k})_M - \dim_{\mathbb{K}} (S_{\geq k+1})_M = \dim_{\mathbb{K}} \frac{S_{\geq k} M}{S_{\geq k+1} M} = h_M(k)$. \square

Now we show two examples of applications of Proposition 4.8 and Theorem 1.3 to computations of the local Hilbert functions of self-dual modules. The first example is the case when M is an algebra. It is a special case of Theorem 1.3, when the encoding matrix is 1×1 , so it is trivially symmetric. The second example is for 2×2 encoding matrix.

Example 4.9. In this example we show that the module (algebra) M apolar to $x^3 + y^3 + z^2$ is self-dual and compute its local Hilbert function. The fact that the module M is self-dual follows

directly from Theorem 1.3, as explained above. Now let us compute the local Hilbert function using the formula from Proposition 4.8.

Let us denote $f = x^3 + y^3 + z^2$, $Z_k = \{f \in M : \deg f \leq k\}$. Let W_k be some \mathbb{K} -linear space satisfying $Z_k = W_k \oplus Z_{k-1}$, where we take $Z_{-1} = 0$ (we are interested in the dimension of W_k over \mathbb{K} , so we just want to find some space W_k satisfying this condition). Then we can take as W_3, W_2, W_1, W_0 the \mathbb{K} -linear subspaces spanned by respectively $(x^3 + y^3 + z^2)$, (x^2, y^2) , (x, y, z) , (1). Now we have $h_M(k) = \dim_{\mathbb{K}} Z_k - \dim_{\mathbb{K}} Z_{k-1} = \dim_{\mathbb{K}} W_k$, so the local Hilbert function is (1, 3, 2, 1).

Example 4.10. In this example we show that the module apolar to the tuple $((x^2 + y^2)e_1^* + (x^3 + y^2)e_2^*, (x^3 + y^2)e_1^* + (x^5 + y^4)e_2^*)$ is self-dual and compute its local Hilbert function. The fact that the module is self-dual follows from Theorem 1.3, as its encoding matrix

$$N = \begin{bmatrix} x^2 + y^2 & x^3 + y^2 \\ x^3 + y^2 & x^5 + y^4 \end{bmatrix}$$

is symmetric. Now we compute the local Hilbert function using the formula from Proposition 4.8. We follow a very similar approach to the one from Example 4.9.

Let us denote $f_1 = (x^2 + y^2)e_1^* + (x^3 + y^2)e_2^*$, $f_2 = (x^3 + y^2)e_1^* + (x^5 + y^4)e_2^*$, $Z_k = \{f \in M : \deg f \leq k\}$. Let D_k be a \mathbb{K} -linear subspace such that $Z_k = Z_{k-1} + D_k$, where we take $Z_{-1} = 0$. Observe that we now have $\dim_{\mathbb{K}} Z_k - \dim_{\mathbb{K}} Z_{k-1} = \dim_{\mathbb{K}} D_k - \dim_{\mathbb{K}} D_k \cap Z_{k-1}$. Let $E_k = D_k \cap Z_{k-1}$. We can choose the pairs of \mathbb{K} -linear subspaces (D_k, E_k) for $k = 5, 4, 3, 2, 1, 0$ as follows:

$$k = 5: D_k = \text{span}\{(x^3 + y^2)e_1^* + (x^5 + y^4)e_2^*\}, E_k = 0;$$

$$k = 4: D_k = \text{span}\{x^2e_1^* + x^4e_2^*\}, E_k = 0;$$

$$k = 3: D_k = \text{span}\{xe_1^* + x^3e_2^*, ye_1^* + y^3e_2^*, (x^2 + y^2)e_1^* + (x^3 + y^2)e_2^*\}, E_k = \text{span}\{(x^2 + y^2 - x)e_1^* + y^2e_2^*\};$$

$$k = 2: D_k = \text{span}\{e_1^* + x^2e_2^*, e_1^* + y^2e_2^*, xe_1^* + x^2e_2^*, (x^2 + y^2 - x)e_1^* + y^2e_2^*\}, E_k = \text{span}\{(x - 1)e_1^*\};$$

$$k = 1: D_k = \text{span}\{xe_2^*, ye_2^*, ye_1^* + ye_2^*, e_1^* + xe_2^*, (x - 1)e_1^*\}, E_k = \text{span}\{e_1^*\};$$

$$k = 0: D_k = \text{span}\{e_1^*, e_2^*\}, E_k = 0;$$

We have $h_M(k) = \dim_{\mathbb{K}} Z_k - \dim_{\mathbb{K}} Z_{k-1} = \dim_{\mathbb{K}} D_k - \dim_{\mathbb{K}} E_k$, so the local Hilbert function is (2, 4, 3, 2, 1, 1).

5 Classification for small degrees

In this section we classify all possible local Hilbert functions for self-dual modules of small degree (so we assume here that R is local and M is self-dual) using tools we have introduced before – Iarrobino’s decomposition, Macaulay’s Bound and the apolarity. As far as we know, this is the first such classification. B. Poonen classified the local Hilbert functions and types of isomorphisms,

but only for algebras of degree less or equal to six [36]. First we introduce a few useful lemmas that simplify the task of classification (some of them do not assume that M is self-dual – we note it then in the statements).

Proposition 5.1. *Let M_1, M_2 be self-dual modules and h_{M_1}, h_{M_2} be their Hilbert functions. Then the module $M_1 \oplus M_2$ is a self-dual module with Hilbert function $h_{M_1} + h_{M_2}$ (addition is element-wise).*

Proof. We have $(M_1 \oplus M_2)^* \cong M_1^* \oplus M_2^* \cong M_1 \oplus M_2$, so the module $M_1 \oplus M_2$ is indeed self-dual. Then the unique maximal ideals in $M_1, M_2, M_1 \oplus M_2$ are respectively $\mathfrak{m}M_1, \mathfrak{m}M_2, \mathfrak{m}(M_1 \oplus M_2)$. Now we have $h_{M_1 \oplus M_2}(k) = \dim_{\mathbb{K}} \frac{\mathfrak{m}^k(M_1 \oplus M_2)}{\mathfrak{m}^{k+1}(M_1 \oplus M_2)} = \dim_{\mathbb{K}} \left(\frac{\mathfrak{m}^k M_1}{\mathfrak{m}^{k+1} M_1} \oplus \frac{\mathfrak{m}^k M_2}{\mathfrak{m}^{k+1} M_2} \right) = \dim_{\mathbb{K}} \left(\frac{\mathfrak{m}^k M_1}{\mathfrak{m}^{k+1} M_1} \right) + \dim_{\mathbb{K}} \left(\frac{\mathfrak{m}^k M_2}{\mathfrak{m}^{k+1} M_2} \right) = h_{M_1}(k) + h_{M_2}(k)$. \square

Corollary 5.2. *Let M_1, M_2, \dots, M_s be self-dual modules and $h_{M_1}, h_{M_2}, \dots, h_{M_s}$ be their Hilbert functions. Then the module $M_1 \oplus M_2 \oplus \dots \oplus M_s$ is a self-dual module with Hilbert function $h_{M_1} + h_{M_2} + \dots + h_{M_s}$ (addition is element-wise).*

Proof. It directly follows from Proposition 5.1 by induction. \square

Lemma 5.3. *The sequence $(1, 1, \dots, 1)$ is a Hilbert function of some self-dual module for every positive length of the sequence.*

Proof. Let us denote the length of the sequence of ones by m . Then $\frac{\mathbb{K}[x]}{x^m}$ is a self-dual module with Hilbert function $(1, 1, \dots, 1)$. \square

Lemma 5.4. *If h_M is a Hilbert function of a module M (not necessarily self-dual) and $h_M(k) = 0$ for some k , then $h_M(l) = 0$ for $l > k$.*

Proof. We prove by induction that for $c \geq 0$ we have $h_M(k + c) = 0$. For $c = 0$ it is true thanks to the assumption. Now if $h_M(k + c) = 0$, then we have that $h_M(k + c) \leq k + c$ and thus, thanks to Corollary 2.26, we have $h_M(k + c + 1) \leq h_M(k + c) = 0$. \square

Lemma 5.5. *If h_M is a Hilbert function of a module M (not necessarily self-dual) and $h_M(k) = 1$ for some $k \geq 1$, then $h_M(l) \leq 1$ for $l > k$.*

Proof. We prove by induction that for $c \geq 0$ we have $h_M(k + c) \leq 1$. For $c = 0$ it is true thanks to the assumption. Now if $h_M(k + c) \leq 1$, then since $k \geq 1$ we have that $h_M(k + c) \leq k + c$ and thus, thanks to Corollary 2.26, we have $h_M(k + c + 1) \leq h_M(k + c) \leq 1$. \square

Lemma 5.6. *If a sequence l is non-increasing, then it is a Hilbert function of some self-dual module.*

Proof. We prove it by induction on the sum of elements in the sequence. If the sequence is $(1, 1, \dots, 1)$, then it is a Hilbert function of some self-dual module, thanks to Lemma 5.3. In particular it holds for (1) , which is a base of the induction.

Now observe that if sequence l is non-increasing, then it can be decomposed into a sum $(1, 1, \dots, 1) + l'$, where l' is non-increasing. If $l \neq (1, 1, \dots, 1)$, then l' is non-empty and thus,

by the induction hypothesis, it is a Hilbert function of some self-dual module. Now observe that since $(1, 1, \dots, 1)$ and l' are both Hilbert functions of self-dual modules, then by Proposition 5.1 their sum is also a Hilbert function of some self-dual module. \square

Lemma 5.7. *If h_M is a Hilbert function of a self-dual module M and k is the largest natural number such that $h_M(k) > 0$, then $h_M(0) \geq h_M(k)$.*

Proof. Let us consider Iarrobino's decomposition of h_M – first row must be symmetric (from Theorem 1.1 Point 4) and must have $h_M(k)$ on the last position (since next rows are strictly shorter), so it also must have $h_M(k)$ on the first position. Since $h_M(0)$ is a sum of values on the first positions in all rows, we have $h_M(0) \geq h_M(k)$. \square

Lemma 5.8. *Let h_M be a Hilbert function of a self-dual module M and let k be the largest natural number such that $h_M(k) > 0$. Then $\sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} h_M(i) \geq \sum_{i=\lceil \frac{k+1}{2} \rceil}^k h_M(i)$.*

Proof. The claim states that the sum of the first half of the sequence is greater or equal to the sum of the second half. This follows from Iarrobino's decomposition (Theorem 1.1) – the sequence is a sum of symmetric parts and each of these parts contributes to the first half at least as much as it does to the second half. \square

Lemma 5.9. *Let h_M be a Hilbert function of a self-dual module M and let us assume that h_M is symmetric (i.e. $h_M(i) = h_M(k - i)$, where k is the largest natural number such that $h_M(k) > 0$). Then Iarrobino's decomposition of h_M has only one row.*

Proof. We prove the claim for decompositions of symmetric sequences (so in the proof we assume only that h_M is a symmetric sequence). We prove by induction of the length s of h_M . For $s = 1$ and $s = 2$ the claim is clear. Let the first row of Iarrobino's decomposition be $l_{0:k}$. Then we have $l_0 = l_k = h_M(k) = h_M(0)$. Let us consider $h_{M_{1:k-1}}$. This sequence is symmetric, so its decomposition has only one row. On the other hand, the decomposition of h_M induces the decomposition of $h_{M_{1:k-1}}$. So positions from 1 to $k - 1$ are fulfilled by only one row. Since $l_0 = l_k = h_M(k) = h_M(0)$, positions 0 and k are fulfilled by the first row. Thus, the whole decomposition has only one row. \square

Lemma 5.10. *Let l be a sequence such that $l_0 = 1$ and l_1 is non-increasing. Then the sequence l is the Hilbert function of some self-dual module (algebra).*

Proof. We construct a polynomial w such that l is the Hilbert function of the algebra apolar to w . Let us denote the polynomial w as a sum of monomials $w = \sum_{i, j \in Z} a_{ij} x_i^j$, where Z is a set we will determine later. We take $a_{ij} = 1$ for all $i, j \in Z$. Now we determine the set Z consisting of non-zero coefficients of w . Let t be the index of the last non-zero element of l (i.e. $l_t \neq 0$ and $l_{t+1} = 0$). Let us denote $c_n = l_n - l_{n+1}$. Since l_1 is non-increasing, we have that $l_1 = c_1 + c_2 + \dots + c_t$. Now we take l_1 variables x_1, x_2, \dots, x_{l_1} . For each variable we take exactly one power for which the coefficient is non-zero (i.e. for each i there exists a unique j such that $i, j \in Z$), so let us denote $j(i)$. We set $j(1) = t$. Let s be a variable that is the greatest i with set $j(i)$ (so at the beginning we have $s = 1$). Let us observe that by Lemma 5.7 we have $c_t = 1$ and from the assumption that l_1 is non-increasing we have that c_1, c_2, \dots, c_t are

non-negative. Now we iterate over the sequence $c_{t-1}, c_{t-2}, \dots, c_1$ and if c_n is non-zero, then we set $j(s+1), j(s+2), \dots, j(s+c_n) = n+1$ and we increase the variable s by assigning $s := s + c_n$. Thanks to Proposition 4.8 we obtain that the above algorithm creates an apolar algebra A with the Hilbert function h_A satisfying $h_A(n) = l_n$ for $n \geq 1$. Clearly $h_A(0) = 1 = l_0$, so we have constructed the desired self-dual algebra.

For given i we choose as j the greatest number such that $c_1 + c_2 + \dots + c_{j-1} < i$ □

Lemma 5.11. *Let us assume that there exists a self-dual module with the Hilbert function that has a trivial Iarrobino's decomposition (i.e. the decomposition has only one row). Then there exists a graded apolar self-dual module with the same Hilbert function.*

Proof. From Remark 3.5 we have the gr R -modules isomorphism $\text{gr } M \cong D_0$. From Proposition 3.4 D_0 is a graded self-dual module. From Theorem 4.2 we can also assume that it is apolar. □

Now we classify possible local Hilbert functions of self-dual modules given the (small) degree. We omit the sequences that trivially are not local Hilbert functions of self-dual modules due to the previous lemmas. We denote by m the degree of considered modules.

m = 1: By Lemma 5.6 possible is:

$$(1). \tag{5}$$

m = 2: By Lemma 5.6 possible are:

$$(1, 1), (2). \tag{6}$$

m = 3: By Lemma 5.6 possible are:

$$(1, 1, 1), (2, 1), (3). \tag{7}$$

m = 4: By Lemma 5.6 possible are:

$$(1, 1, 1, 1), (2, 1, 1), (2, 2), (3, 1), (4). \tag{8}$$

(1, 2, 1) : possible – module (algebra) apolar to $x^2 + y^2$.

m = 5: By Lemma 5.6 possible are:

$$(1, 1, 1, 1, 1), (2, 1, 1, 1), (2, 2, 1), (3, 1, 1), (3, 2), (4, 1), (5). \tag{9}$$

(1, 2, 1, 1) : possible – module (algebra) apolar to $x^3 + y^2$,

(1, 3, 1) : possible – module (algebra) apolar to $x^2 + y^2 + z^2$.

m = 6: By Lemma 5.6 possible are:

$$(1, 1, 1, 1, 1, 1), (2, 1, 1, 1, 1), (2, 2, 1, 1), (2, 2, 2), \quad (10)$$

$$(3, 1, 1, 1), (3, 2, 1), (3, 3), (4, 1, 1), (4, 2), (5, 1), (6). \quad (11)$$

(1, 2, 1, 1, 1) : possible – module (algebra) apolar to $x^4 + y^2$,

(1, 2, 2, 1) : possible – module (algebra) apolar to $x^3 + y^3$,

(1, 3, 1, 1) : possible – module (algebra) apolar to $x^3 + y^2 + z^2$,

(1, 4, 1) : possible – module (algebra) apolar to $x^2 + y^2 + z^2 + w^2$,

(2, 3, 1) : possible – direct sum (1, 1) + (1, 2, 1).

m = 7: By Lemma 5.6 possible are:

$$(1, 1, 1, 1, 1, 1, 1), (2, 1, 1, 1, 1, 1), (2, 2, 1, 1, 1), (2, 2, 2, 1), \quad (12)$$

$$(3, 1, 1, 1, 1), (3, 2, 1, 1), (3, 2, 2), (3, 3, 1), \quad (13)$$

$$(4, 1, 1, 1), (4, 2, 1), (4, 3), (5, 1, 1), (5, 2), (6, 1), (7). \quad (14)$$

(1, 2, 1, 1, 1, 1) : possible – module (algebra) apolar to $x^5 + y^2$,

(1, 2, 2, 1, 1) : possible – module (algebra) apolar to $x^4 + y^3$,

(1, 3, 1, 1, 1) : possible – module (algebra) apolar to $x^4 + y^2 + z^2$,

(1, 3, 2, 1) : possible – module (algebra) apolar to $x^3 + y^3 + z^2$,

(1, 4, 1, 1) : possible – module (algebra) apolar to $x^3 + y^2 + z^2 + w^2$,

(1, 5, 1) : possible – module (algebra) apolar to $x^2 + y^2 + z^2 + w^2 + u^2$,

(2, 3, 1, 1) : possible – direct sum (1, 1) + (1, 2, 1, 1),

(2, 3, 2) : possible – direct sum (1, 1, 1) + (1, 2, 1),

(2, 4, 1) : possible – direct sum (1, 1) + (1, 3, 1).

m = 8: By Lemma 5.6 possible are:

$$(1, 1, 1, 1, 1, 1, 1, 1), (2, 1, 1, 1, 1, 1, 1), (2, 2, 1, 1, 1, 1), (2, 2, 2, 1, 1), (2, 2, 2, 2), \quad (15)$$

$$(3, 1, 1, 1, 1, 1), (3, 2, 1, 1, 1), (3, 2, 2, 1), (3, 3, 1, 1), (3, 3, 2) \quad (16)$$

$$(4, 1, 1, 1, 1), (4, 2, 1, 1), (4, 2, 2), (4, 3, 1), (4, 4) \quad (17)$$

$$(5, 1, 1, 1), (5, 2, 1), (5, 3), (6, 1, 1), (6, 2), (7, 1), (8). \quad (18)$$

(1, 2, 1, 1, 1, 1, 1) : possible – module (algebra) apolar to $x^6 + y^2$,

- (1, 2, 2, 1, 1, 1) : possible – module (algebra) apolar to $x^5 + y^3$,
- (1, 2, 2, 2, 1) : possible – module (algebra) apolar to $x^4 + y^4$,
- (1, 2, 3, 1, 1) : impossible due to Iarrobino’s decomposition – let us assume that there exists a decomposition; then the first row must be (1, a, b, a, 1) (due to the last position) and so the second is (0, c, c, 0) and the third is (0, d, 0) (due to the first position); now analyzing the fourth position gives us $a = 1$; since from Theorem 1.1 Point 4 the first row is a Hilbert function of some module and thus thanks to Lemma 5.5 we obtain $b = 1$; so analyzing the second position gives us $2 = 1 + c + d$ and analyzing the third position gives us $3 = 1 + c$, which implies $d < 0$ – contradiction,
- (1, 3, 1, 1, 1, 1) : possible – module (algebra) apolar to $x^5 + y^2 + z^2$,
- (1, 3, 2, 1, 1) : possible – module (algebra) apolar to $x^4 + y^3 + z^2$,
- (1, 3, 3, 1) : possible – module (algebra) apolar to $x^3 + y^3 + z^3$,
- (1, 4, 1, 1, 1) : possible – module (algebra) apolar to $x^4 + y^2 + z^2 + w^2$,
- (1, 4, 2, 1) : possible – module (algebra) apolar to $x^3 + y^3 + z^2 + w^2$,
- (1, 5, 1, 1) : possible – module (algebra) apolar to $x^3 + y^2 + z^2 + w^2 + u^2$,
- (1, 6, 1) : possible – module (algebra) apolar to $x^2 + y^2 + z^2 + w^2 + u^2 + v^2$,
- (2, 2, 3, 1) : possible – module apolar to $(x^3 + y^3)e_1^* + xye_2^*, xye_1^* + ye_2^*$, this module is self-dual from Theorem 1.3 with encoding matrix

$$\begin{bmatrix} x^3 + y^3 & xy \\ xy & y \end{bmatrix},$$

- (2, 3, 1, 1, 1) : possible – direct sum (1, 1) + (1, 2, 1, 1, 1),
- (2, 3, 2, 1) : possible – direct sum (1, 1, 1, 1) + (1, 2, 1),
- (2, 4, 1, 1) : possible – direct sum (1, 1) + (1, 3, 1, 1),
- (2, 4, 2) : possible – direct sum (1, 1, 1) + (1, 3, 1),
- (2, 5, 1) : possible – direct sum (1, 1) + (1, 4, 1),
- (3, 2, 3) : impossible: Let us assume that there exists a self-dual module with the Hilbert function (3, 2, 3). Then by Lemma 5.9 Iarrobino’s decomposition must be trivial (i.e. has only one row). Thus by Lemma 5.11 there exists a self-dual graded module with the Hilbert function (3, 2, 3). We have $h(1) = 2$, so the space of the first order derivatives has a degree 2. We have $h(2) = 3$, so we have three polynomials of degree 3. If all these polynomials have one-dimensional subspaces of derivatives, then they are linearly dependent modulo constants – contradiction. So there exists a polynomial of degree 2 that has two-dimensional subspace of

derivatives. From $h(1) = 2$ the subspace of the polynomials of degree 1 is two-dimensional, so it is generated by at most two variables. Since the module is graded, it follows that we only have two variables in all degrees. Let us take the polynomial that has two-dimensional subspace of derivatives. It has a form $v_1x_1^2 + v_2x_2^2 + v_3x_1x_2$, where $v_i \in \mathbb{R}^3$. Now let us take the subspace of the polynomials of degree 2 – they have a form $w_1x_1^2 + w_2x_2^2 + w_3x_1x_2$. Their derivatives are in the space $(v_1x_1 + v_3x_2; v_2x_2 + v_3x_1)$. From derivation with respect to x_1 we obtain $w_1 = k_1v_1 + l_1v_3, w_3 = k_1v_3 + l_1v_2$ and from derivation with respect to x_2 we obtain $w_2 = k_2v_3 + l_2v_2, w_3 = k_2v_1 + l_2v_3$, where k_1, k_2, l_1, l_2 are constants. So we obtain the equation $k_2v_1 + l_2v_3 = k_1v_3 + l_1v_2$. Without losing generality let us assume $v_1 \neq 0$. Let us observe that the subspace of degree 2 has dimension 3, so it is spanned by k_1, l_1, l_2 . We have $k_2v_{1i} = k_1v_{3i} + l_1v_{2i} - l_2v_{3i}$. Taking $l_1 = 0, l_2 = 0, k_1 = 1$ and $k_1 = 0, l_2 = 0, l_1 = 1$ we obtain $v_{3i} = bv_{1i}, v_{2i} = av_{1i}$, so $v_2 = av_1, v_3 = bv_1$. Thus we have that our polynomial that has two-dimensional subspace of derivatives is of the form $v_1x_1^2 + av_1x_2^2 + bv_1x_1x_2$. The subspace of the polynomials of degree 1 is then $(v_1x_1 + bv_1x_2; av_1x_2 + bv_1x_1)$ and the subspace of degree 0 is (v_1) . From self-duality there are exactly three generators and we already have three of them (the polynomials of degree 2), so there is only one polynomial of degree 0, which contradicts $h(0) = 3$.

(3, 4, 1) : possible – direct sum (1, 1) + (2, 3, 1).

References

- [1] Josh Alman and Virginia Vassilevska Williams. “A Refined Laser Method and Faster Matrix Multiplication”. In: *SODA*. 2021.
- [2] Jarod Alper and Alexander Isaev. “Associated forms and hypersurface singularities: the binary case”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 2018.745 (2018), pp. 83–104.
- [3] Ivan Arzhantsev and Yulia Zaitseva. “Equivariant completions of affine spaces”. In: *arXiv preprint arXiv:2008.09828* (2020).
- [4] Alessandra Bernardi and Kristian Ranestad. “On the cactus rank of cubic forms”. In: *Journal of Symbolic Computation* 50 (2013), pp. 291–297.
- [5] Alessandra Bernardi et al. “On polynomials with given Hilbert function and applications”. In: *Collectanea mathematica* 69 (2018), pp. 39–64.
- [6] Markus Bläser and Vladimir Lysikov. “Slice Rank of Block Tensors and Irreversibility of Structure Tensors of Algebras”. In: *45th International Symposium on Mathematical Foundations of Computer Science (MFCS 2020)*. Ed. by Javier Esparza and Daniel Král'. Vol. 170. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2020, 17:1–17:15. ISBN: 978-3-95977-159-7. DOI: [10.4230/LIPIcs.MFCS.2020.17](https://drops.dagstuhl.de/opus/volltexte/2020/12686). URL: <https://drops.dagstuhl.de/opus/volltexte/2020/12686>.

- [7] Weronika Buczyńska and Jarosław Buczyński. “Secant varieties to high degree Veronese reembeddings, catalecticant matrices and smoothable Gorenstein schemes”. In: *J. Algebraic Geom.* 23.1 (2014), pp. 63–90. ISSN: 1056-3911. DOI: [10.1090/S1056-3911-2013-00595-0](https://doi.org/10.1090/S1056-3911-2013-00595-0). URL: <https://doi.org/10.1090/S1056-3911-2013-00595-0>.
- [8] Jarosław Buczyński et al. “Constructions of k -regular maps using finite local schemes”. In: *Journal of the European Mathematical Society* 21.6 (2019), pp. 1775–1808.
- [9] Gianfranco Casnati, Joachim Jelisiejew, and Roberto Notari. “Irreducibility of the Gorenstein loci of Hilbert schemes via ray families”. In: *Algebra & Number Theory* 9 (2014), pp. 1525–1570. URL: <https://api.semanticscholar.org/CorpusID:56308660>.
- [10] Gianfranco Casnati, Joachim Jelisiejew, and Roberto Notari. “Irreducibility of the Gorenstein loci of Hilbert schemes via ray families”. In: *Algebra & Number Theory* 9.7 (2015), pp. 1525–1570.
- [11] Gianfranco Casnati, Joachim Jelisiejew, and Roberto Notari. “Irreducibility of the Gorenstein loci of Hilbert schemes via ray families”. In: *Algebra Number Theory* 9.7 (2015), pp. 1525–1570. ISSN: 1937-0652. DOI: [10.2140/ant.2015.9.1525](https://doi.org/10.2140/ant.2015.9.1525). URL: <https://doi.org/10.2140/ant.2015.9.1525>.
- [12] Gianfranco Casnati and Roberto Notari. “On the Gorenstein locus of some punctual Hilbert schemes”. In: *J. Pure Appl. Algebra* 213.11 (2009), pp. 2055–2074. ISSN: 0022-4049. DOI: [10.1016/j.jpaa.2009.03.002](https://doi.org/10.1016/j.jpaa.2009.03.002). URL: <https://doi.org/10.1016/j.jpaa.2009.03.002>.
- [13] J. Elias and M.E. Rossi. “The structure of the inverse system of Gorenstein k -algebras”. In: *Advances in Mathematics* 314 (2017), pp. 306–327. ISSN: 0001-8708. DOI: <https://doi.org/10.1016/j.aim.2017.03.002>. URL: <https://www.sciencedirect.com/science/article/pii/S0001870815305247>.
- [14] Juan Elias. “Inverse systems of local rings”. In: *Commutative Algebra and its Interactions to Algebraic Geometry: VIASM 2013–2014* (2018), pp. 119–163.
- [15] Juan Elias and M Rossi. “Isomorphism classes of short Gorenstein local rings via Macaulay’s inverse system”. In: *Transactions of the American Mathematical Society* 364.9 (2012), pp. 4589–4604.
- [16] Gioia Failla, Zachary Flores, and Chris Peterson. “On the weak Lefschetz property for vector bundles on P^2 ”. In: *Journal of Algebra* 568 (2021), pp. 22–34.
- [17] Tadahito Harima et al. *Lefschetz properties*. Springer, 2013.
- [18] Tadahito Harima et al. “The weak and strong Lefschetz properties for Artinian K -algebras”. In: *Journal of Algebra* 262.1 (2003), pp. 99–126.
- [19] Heather A. Hulett. “A generalization of Macaulay’s theorem”. In: *Communications in Algebra* 23.4 (1995), pp. 1249–1263. DOI: [10.1080/00927879508825278](https://doi.org/10.1080/00927879508825278). eprint: <https://doi.org/10.1080/00927879508825278>. URL: <https://doi.org/10.1080/00927879508825278>.
- [20] Craig Huneke. “Hyman Bass and ubiquity: Gorenstein rings”. In: *Algebra, K-theory, groups, and education*. 1999, pp. 55–78.
- [21] Yu. I. Zaitseva I. V. Arzhantsev. “Equivariant completions of affine spaces”. In: *Uspekhi Mat. Nauk* 77 (2022), pp. 3–90. DOI: <https://doi.org/10.4213/rm10046>.

- [22] Anthony Iarrobino. “Associated Graded Algebra of a Gorenstein Artin Algebra”. In: 1994. URL: <https://api.semanticscholar.org/CorpusID:121059708>.
- [23] Anthony Iarrobino and Pedro Macias Marques. “Symmetric decomposition of the associated graded algebra of an Artinian Gorenstein algebra”. In: *Journal of Pure and Applied Algebra* 225.3 (2021), p. 106496. ISSN: 0022-4049. DOI: <https://doi.org/10.1016/j.jpaa.2020.106496>. URL: <https://www.sciencedirect.com/science/article/pii/S0022404920301973>.
- [24] Joachim Jelisiejew. “Classifying local Artinian Gorenstein algebras”. In: *Collectanea Mathematica* 68 (2015), pp. 101–127. URL: <https://api.semanticscholar.org/CorpusID:119171782>.
- [25] Joachim Jelisiejew. “Open problems in deformations of Artinian algebras, Hilbert schemes and around”. In: *arXiv preprint arXiv:2307.08777* (2023).
- [26] Joachim Jelisiejew. “VSPs of cubic fourfolds and the Gorenstein locus of the Hilbert scheme of 14 points on A^6 ”. In: *Linear Algebra and its Applications* 557 (2018), pp. 265–286.
- [27] Joachim Jelisiejew, Joseph Landsberg, and Arpan Pal. “Concise tensors of minimal border rank”. In: *Mathematische Annalen* (2024). DOI: <https://doi.org/10.1007/s00208-023-02569-y>.
- [28] Joachim Jelisiejew and Klemen Šivic. “Components and singularities of Quot schemes and varieties of commuting matrices”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 2022 (2021), pp. 129–187. URL: <https://api.semanticscholar.org/CorpusID:23562381>.
- [29] Tomaz Kosir and Polona Oblak. “On pairs of commuting nilpotent matrices”. In: *Transformation Groups* 14 (Dec. 2007), pp. 175–182. DOI: [10.1007/s00031-008-9045-6](https://doi.org/10.1007/s00031-008-9045-6).
- [30] Michael Kunte. “Gorenstein modules of finite length”. In: *Mathematische Nachrichten* 284 (2008).
- [31] Andrew Kustin, Claudia Polini, and Bernd Ulrich. “The equations defining blowup algebras of height three Gorenstein ideals”. In: *Algebra & Number Theory* 11.7 (2017), pp. 1489–1525.
- [32] Joseph M Landsberg and Mateusz Michałek. “Abelian tensors”. In: *Journal de Mathématiques Pures et Appliquées* 108.3 (2017), pp. 333–371.
- [33] F. S. Macaulay. “Some Properties of Enumeration in the Theory of Modular Systems”. In: *Proceedings of the London Mathematical Society* s2-26.1 (1927), pp. 531–555. DOI: <https://doi.org/10.1112/plms/s2-26.1.531>. eprint: <https://londmathsoc.onlinelibrary.wiley.com/doi/abs/10.1112/plms/s2-26.1.531>. URL: <https://londmathsoc.onlinelibrary.wiley.com/doi/abs/10.1112/plms/s2-26.1.531>.
- [34] Toshiaki Maeno and Junzo Watanabe. “Lefschetz elements of Artinian Gorenstein algebras and Hessians of homogeneous polynomials”. In: *Illinois Journal of Mathematics* 53.2 (2009), pp. 591–603.
- [35] Chris McDaniel et al. “Free extensions and Lefschetz properties, with an application to rings of relative coinvariants”. In: *Linear and Multilinear Algebra* 69.2 (2021), pp. 305–330.
- [36] Bjorn Poonen. “Isomorphism types of commutative algebras of finite rank over an algebraically closed field”. In: *Contemp. Math.* 463 (Jan. 2008). DOI: [10.1090/conm/463/09050](https://doi.org/10.1090/conm/463/09050).

- [37] Kristian Ranestad and Frank-Olaf Schreyer. “Varieties of sums of powers”. In: *Journal für die Reine und Angewandte Mathematik* (2000), pp. 147–182.
- [38] Frank-Olaf Schreyer. “Geometry and Algebra of Prime Fano 3-folds of Genus 12”. In: *Compositio Mathematica* 127 (1999), pp. 297–319. URL: <https://api.semanticscholar.org/CorpusID:1631728>
- [39] Frank-Olaf Schreyer. “Geometry and algebra of prime Fano 3-folds of genus 12”. In: *Compositio Mathematica* 127.3 (2001), pp. 297–319.
- [40] David Ssevviiri. “Applications of reduced and coreduced modules I”. In: *International Electronic Journal of Algebra* (2024), pp. 1–21.
- [41] Richard P Stanley. *Combinatorics and commutative algebra*. Vol. 41. Springer Science & Business Media, 2007.
- [42] Volker Strassen. “Gaussian elimination is not optimal”. In: *Numerische mathematik* 13.4 (1969), pp. 354–356.
- [43] Robert Szafarczyk. “New elementary components of the Gorenstein locus of the Hilbert scheme of points”. In: *Communications in Algebra* 51.6 (2023), pp. 2688–2704.
- [44] Attila Wiebe. “The Lefschetz property for componentwise linear ideals and Gotzmann ideals”. In: *Communications in Algebra* 32.12 (2004), pp. 4601–4611.
- [45] Maciej Wojtala. “Irreversibility of structure tensors of modules”. In: *Collectanea Mathematica* (Apr. 2022). ISSN: 2038-4815. DOI: [10.1007/s13348-022-00361-w](https://doi.org/10.1007/s13348-022-00361-w). URL: <https://doi.org/10.1007/s13348-022-00361-w>