DIMENSIONS OF RIESZ PRODUCTS AND PLURIHARMONIC MEASURES

EVGUENI DOUBTSOV

ABSTRACT. We estimate the energy and Hausdorff dimensions of the Riesz products on the unit sphere of \mathbb{C}^n , $n \geq 2$. Also, we obtain similar results for the pluriharmonic measures on the torus.

1. INTRODUCTION

The present paper is motivated by the following general question: how does the spectrum of a measure μ affect the size of support of μ ? We primarily consider Riesz product measures and related objects on the unit sphere $S = S_n = \{\zeta \in \mathbb{C}^n : |\zeta| = 1\}, n \geq 2$, therefore, we start by introducing the spaces $H(p,q), (p,q) \in \mathbb{Z}_+^2$.

1.1. Complex spherical harmonics. Let $\mathcal{U}(n)$ denote the group of unitary operators on the Hilbert space \mathbb{C}^n , $n \geq 2$. Observe that $S_n = \mathcal{U}(n)/\mathcal{U}(n-1)$, hence, S_n is a homogeneous space. General constructions of abstract harmonic analysis are explicitly implemented on S_n in terms of the spaces H(p,q), $(p,q) \in \mathbb{Z}^2_+$.

Definition 1.1. Fix a dimension $n, n \ge 2$. Let H(p,q) = H(p,q;n) denote the space of all homogeneous harmonic polynomials of bidegree $(p,q) \in \mathbb{Z}_+^2$. By definition, this means that the polynomials under consideration have degree p in z_1, z_2, \ldots, z_n , degree q in $\overline{z_1}, \overline{z_2}, \ldots, \overline{z_n}$, and have total degree p + q.

For the restriction of H(p,q) on S, one uses the same symbol. The elements of H(p,q) are often called *complex* spherical harmonics.

1.2. Riesz products on the sphere. Zygmund's dichotomy [15] suggests that the homogeneous holomorphic polynomials introduced by Ryll and Wojtaszczyk [14] could provide examples of singular Riesz-type products on the sphere. We say that $\{R_i\}_{i=1}^{\infty}$ is a Ryll–Wojtaszczyk sequence with a constant $\delta \in (0, 1)$ if

- $R_j \in H(j,0)$, that is, R_j is a homogeneous holomorphic polynomial of degree j,
- $||R_j||_{L^{\infty}(S)} = 1$,
- $||R_j||_{L^2(S)} \ge \delta$ for all j = 1, 2, ...

Definition 1.2. Let $R = \{R_j\}_{j=1}^{\infty}$ be a Ryll–Wojtaszczyk sequence, $J = \{j_k\}_{k=1}^{\infty} \subset \mathbb{N}, j_{k+1}/j_k \geq 3$, and $a = \{a_k\}_{k=1}^{\infty} \subset \mathbb{D}$. Then (R, J, a) is called a Riesz triple.

²⁰²⁰ Mathematics Subject Classification. Primary 42A55; Secondary 28A78, 31C10, 43A85. Key words and phrases. Riesz product, energy dimension, Hausdorff dimension, pluriharmonic measure.

Each Riesz triple (R, J, a) generates a (standard) Riesz product. Namely, the Riesz product $\Pi(R, J, a)$ is defined by the formal equality

$$\Pi(R, J, a) = \prod_{k=1}^{\infty} \left(\frac{\overline{a}_k \overline{R}_{j_k}}{2} + 1 + \frac{a_k R_{j_k}}{2} \right),$$

where the product converges in the weak*-sense (see [7], [6] for details).

There exists a fairly rich set of singular Riesz products on the sphere. Indeed, if (R, J, a) is a Riesz pair with $a \notin \ell^2$, then Corollary 1 from [6] provides a sequence $U = \{U_j\}_{j=1}^{\infty}, U_j \in \mathcal{U}(n)$, such that $\Pi(R \circ U, J, a)$ is singular with respect to Lebesgue measure on S_n .

For a singular Riesz product, it is natural to ask about its dimension. In the present paper, we estimate the energy and Hausdorff dimensions of the Riesz products on the sphere. Let $\dim_{\mathcal{H}} \Pi$ denote the Hausdorff dimension of a measure Π . We obtain, in particular, the following result.

Theorem 1.3. Let (R, J, a) be a Riesz triple. Then

$$\dim_{\mathcal{H}} \Pi(R, J, a) \ge 2n - 1 - \limsup_{k \to \infty} \left(\frac{1}{2 \log j_k} \sum_{\ell=1}^{k-1} |a_\ell|^2 \right).$$

1.3. Organization of the paper. Auxiliary results, including definitions and basic facts of harmonic analysis on S_n , are collected in Section 2. Estimates of the energy and Hausdorff dimensions of the Riesz products on the unit sphere are obtained in Section 3. A related problem about Hausdorff dimensions of the pluriharmonic measures on the torus is discussed in Section 4.

1.4. Notation. For F, G > 0, we write $F \leq G$ provided that $F \leq CG$ for a constant C > 0. If $F \leq G$ and $G \leq F$, then we write $F \approx G$.

2. Auxiliary results

2.1. Basics of harmonic analysis on S_n . Let $\sigma = \sigma_n$ denote the normalized Lebesgue measure on the unit sphere S_n . Observe that

(2.1)
$$L^{2}(\sigma) = \bigoplus_{(p,q) \in \mathbb{Z}^{2}_{+}} H(p,q).$$

Specific aspects of the harmonic analysis on S are illustrated by the following multiplication rule for the spaces H(p,q): if $f \in H(p,q)$ and $g \in H(r,s)$, then

$$fg \in \sum_{\ell=0}^{L} H(p+r-\ell, q+s-\ell),$$

where $L = \min(p, s) + \min(q, r)$. See [13, Chapter 12] for the proofs of the above facts and for further results.

Let $M(S_n)$ denote the space of complex Borel measures on the unit sphere S_n . Let $K_{p,q}(z,\zeta)$ denote the reproducing kernel for the Hilbert space $H(p,q) \subset L^2(S)$. The polynomial

$$u_{p,q}(z) = \int_{S} K_{p,q}(z,\zeta) \, d\mu(\zeta), \quad z \in S,$$

is called the H(p,q)-projection of $\mu \in M(S)$. For $\mu \in M(S)$, the spectrum spec (μ) in terms of H(p,q) is defined by

$$\operatorname{spec}(\mu) = \left\{ (p,q) \in \mathbb{Z}_+^2 : \mu_{p,q} \neq \mathbf{0} \right\}.$$

2.2. Disintegration into slice-products. Let $\mathbb{T} = S_1$ denote the unit circle. Given a $\xi \in S_n$, the slice-product

$$\Pi_{\xi}(R, J, a)(\lambda) := \Pi(R(\lambda\xi), J, a), \quad \lambda \in \mathbb{T},$$

is the classical Riesz product

(2.2)
$$\prod_{k=1}^{\infty} \left(\frac{\overline{a_k \overline{R}_{j_k}(\xi)} \overline{\lambda}^{j_k}}{2} + 1 + \frac{a_k R_{j_k}(\xi) \lambda^{j_k}}{2} \right), \quad \lambda \in \mathbb{T}.$$

In particular, $\Pi_{\xi}(R, J, a)$ is a correctly defined probability measure.

Let \mathbb{CP}^{n-1} denote the complex projective space of dimension n-1, that is, the collection of all one-dimensional linear subspaces of \mathbb{C}^n . Let $\pi = \pi_n$ denote the canonical projection from S_n onto \mathbb{CP}^{n-1} . Observe that $\Pi_{\lambda\xi}(w) = \Pi_{\xi}(\lambda w)$ for $\xi \in S_n$, $\lambda, w \in \mathbb{T}$. Therefore, the probability slice-measure Π_{ζ} is correctly defined for $\zeta \in \mathbb{CP}^{n-1}$ as an element of $M_+(S_n)$ supported by the unit circle $\pi^{-1}(\zeta) \subset S_n$.

Let $\hat{\sigma}_n$ denote the unique probability measure on \mathbb{CP}^{n-1} invariant with respect to all unitary transformations of \mathbb{C}^n . The following lemma shows that $\Pi(R, J, a)$ is the integral of its slices.

Lemma 2.1 ([6, Lemma 1]). Let (R, J, a) be a Riesz triple on the sphere S_n , $n \ge 2$, and let $\Pi(R, J, a)$ denote the corresponding Riesz product. Then

$$\Pi(R,J,a) = \int_{\mathbb{CP}^{n-1}} \Pi_{\zeta}(R,J,a) \, d\widehat{\sigma}_n(\zeta)$$

in the following weak sense:

$$\int_{S_n} f \, d\Pi(R, J, a) = \int_{\mathbb{CP}^{n-1}} \int_{S_n} f \, d\Pi_{\zeta}(R, J, a) \, d\widehat{\sigma}_n(\zeta)$$

for all $f \in C(S_n)$.

3. Dimensions of Riesz products

3.1. Energy dimension. Let $M_+(S_n)$ denote the set of all finite positive Borel measures on S_n . Given a measure $\mu \in M_+(S)$, its *t*-energy $I_t(\mu)$, t > 0, is defined as

$$I_t(\mu) := \int_S \int_S \frac{d\mu(x)d\mu(y)}{|x-y|^t}$$

We say that d is the energy dimension of μ and we write $\dim_{\mathcal{E}} \mu = d$ if

$$d = \sup\left\{t : I_t(\mu) < \infty\right\}.$$

To estimate the energy dimension of $\Pi(R, J, a)$, we need the following theorem.

Theorem 3.1 ([12, Theorem 3.1]). For each t, 0 < t < 2n - 1, there are constants $C_1, C_2 > 0$ such that

$$C_1 I_t(\mu) \le \|\mu_{0,0}\|_2^2 + \sum_{j=1}^\infty j^{t-2n+1} \sum_{p+q=j} \|\mu_{p,q}\|_2^2 \le C_2 I_t(\mu)$$

for all $\mu \in M_+(S_n)$ with $n \ge 2$.

In [11], an analog of Theorem 3.2 for the unit circle is used to compute the energy dimension of a classical Riesz product on \mathbb{T} . In the following result, we apply Theorem 3.1 to estimate from below the energy dimension of a Riesz product on the sphere. In fact, modulo technical details, the principal argument below is similar to that from the proof of Theorem 3.1 from [11].

Theorem 3.2. Let (R, J, a) be a Riesz triple. Then

$$\dim_{\mathcal{E}} \Pi(R, J, a) \ge 2n - 1 - \alpha_0,$$

where

(3.1)
$$\alpha_0 = \max\left[0, \ \limsup_{k \to \infty} \left(\frac{\log \frac{|a_k|^2}{2} + \sum_{\ell=1}^{k-1} \log\left(1 + \frac{|a_\ell|^2}{2}\right)}{\log j_k}\right)\right].$$

In particular,

$$\dim_{\mathcal{E}} \Pi(R, J, a) \ge 2n - 1 - \limsup_{k \to \infty} \left(\frac{1}{2 \log j_k} \sum_{\ell=1}^{k-1} |a_\ell|^2 \right).$$

Proof. Let $\Pi := \Pi(R, J, a)$. For k = 1, 2, ..., let

$$\Gamma_k := \left\{ \pm j_k + \sum_{\ell=1}^{k-1} \varepsilon_\ell j_\ell : \varepsilon_\ell = 0, \pm 1 \right\}.$$

Suppose that $\gamma \in \Gamma_k$. Without loss of generality, assume that $\gamma > 0$. Then there exists exactly one representation

$$\gamma = j_k + \sum_{\ell=1}^{k-1} \varepsilon_\ell j_\ell.$$

By the multiplication rule for the spherical harmonics (see Section 2.1),

$$\sum_{p-q=\gamma} \prod_{p,q} = R_{j_k} \prod_{\ell=1}^{k-1} R_{j_\ell}(\varepsilon_\ell),$$

where

$$R_{j_{\ell}}(\varepsilon_{\ell}) = R_{j_{\ell}}, \quad \varepsilon_{\ell} = 1,$$

$$R_{j_{\ell}}(\varepsilon_{\ell}) = 1, \quad \varepsilon_{\ell} = 0,$$

$$R_{j_{\ell}}(\varepsilon_{\ell}) = \overline{R}_{j_{\ell}}, \quad \varepsilon_{\ell} = -1.$$

Applying (2.1), we obtain

$$\sum_{p-q=\gamma} \|\Pi_{p,q}\|_{2}^{2} = \left\| R_{j_{k}} \prod_{\ell=1}^{k-1} R_{j_{\ell}}(\varepsilon_{\ell}) \right\|_{2}^{2} \le \left(\frac{|a_{k}|}{2} \prod_{\ell: \varepsilon_{\ell} \neq 0} \frac{|a_{\ell}|}{2} \right)^{2},$$

where the empty product is equal to one.

Now, fix an $\alpha > \alpha_0$. Since $p + q \ge p - q$, we have

(3.2)
$$\sum_{\gamma \in \Gamma_k} |p+q|^{-\alpha} \sum_{p-q=\gamma} \|\Pi_{p,q}\|_2^2 \le \sum_{\gamma \in \Gamma_k} |\gamma|^{-\alpha} \sum_{p-q=\gamma} \|\Pi_{p,q}\|_2^2.$$

4

If $\gamma \in \Gamma_k$, then $|\gamma| \approx j_k$. Thus,

(3.3)
$$\sum_{\gamma \in \Gamma_k} |\gamma|^{-\alpha} \sum_{p-q=\gamma} \|\Pi_{p,q}\|_2^2 \lesssim j_k^{-\alpha} \frac{|a_k|^2}{2} \prod_{\ell=1}^{k-1} \left(1 + \frac{|a_\ell|^2}{2}\right).$$

Combining (3.2) and (3.3), we obtain

(3.4)
$$I_{2n-1-\alpha}(\Pi) \lesssim 1 + j_k^{-\alpha} \frac{|a_k|^2}{2} \prod_{\ell=1}^{k-1} \left(1 + \frac{|a_\ell|^2}{2}\right)$$

by Theorem 3.1.

We have $\log j_k \ge (k-1)\log 3$, thus, selecting an A, 0 < A < 1, sufficiently close to 1, we deduce from the inequality $\alpha > \alpha_0$ that

$$\alpha \ge \frac{\log \frac{|a_k|^2}{2} + \sum_{\ell=1}^{k-1} \log \left(1 + \frac{|a_\ell|^2}{2}\right) + k |\log A|}{\log j_k}$$

or, equivalently,

$$j_k^{-\alpha} \frac{|a_k|^2}{2} \prod_{\ell=1}^{k-1} \left(1 + \frac{|a_\ell|^2}{2}\right) \le A^k.$$

Combining the above property and (3.4), we conclude that $I_{2n-1-\alpha}(\Pi) < \infty$. Since $\alpha > \alpha_0$ is arbitrary, we conclude that $\dim_{\mathcal{E}} \Pi \ge 2n - 1 - \alpha_0$, as required. \Box

3.2. Hausdorff dimension. For a Borel set E, let $\dim_{\mathcal{H}} E$ denote its Hausdorff dimension. The Hausdorff dimension of a measure $\mu \in M_+(S_n)$ is defined as

$$\dim_{\mathcal{H}} \mu = \inf \{ \dim_{\mathcal{H}} E : E \text{ is a Borel set with } \mu(E) > 0 \}.$$

For properties of the Hausdorff dimension of a measure see [9, Chapter 10].

Given a Riesz product $\Pi = \Pi(R, J, a)$, direct inspection shows that there exists a sufficiently small $\varepsilon > 0$ such that $\Pi_{p,q} = \mathbf{0}$ provided that

$$\left|\frac{p}{q}-1\right| < \varepsilon, \quad (p,q) \neq (0,0).$$

Therefore, Theorem 1.1 from [3] applies to Π and guarantees that $\dim_{\mathcal{H}} \Pi \geq 2n-2$. In fact, applying Theorem 3.2, we obtain a more precise estimate.

3.2.1. Application of Theorem 3.2. If $I_t(\mu) < \infty$, then $\dim_{\mathcal{H}} \mu > t$ (cf. [8, Section 4.3]); thus, the Hausdorff dimension of a measure is always at least the energy dimension.

Corollary 3.3. Let (R, J, a) be a Riesz triple. Then

$$\dim_{\mathcal{H}} \Pi(R, J, a) \ge 2n - 1 - \alpha_0,$$

where α_0 is defined by (3.1).

Proof. Since $\dim_{\mathcal{H}} \Pi \geq \dim_{\mathcal{E}} \Pi$, it suffices to apply Theorem 3.2.

Remark 3.4. Clearly, Theorem 1.3 is a particular case of the above corollary.

3.2.2. Reduction to slice-products. An alternative approach to Corollary 3.3 is to consider the slice-products of $\Pi(R, J, a)$. We need the following theorem which is a consequence of [10, Sections 2.10.2 and 2.10.17].

Theorem 3.5. Let $K \subset \mathbb{CP}^{n-1} \times \mathbb{T}$ be a compact set and $K_{\zeta} = \{w \in \mathbb{T} : (\zeta, w) \in K\}$. Assume that $\dim_{\mathcal{H}} K_{\zeta} > \beta$ for $\zeta \in X \subset \mathbb{CP}^{n-1}$. If $\widehat{\sigma}_n(X) > 0$, then

$$\dim_{\mathcal{H}} K \ge 2n - 2 + \beta.$$

Fix an $\alpha > \alpha_0$. It suffices to prove that

(3.5)
$$\dim_{\mathcal{H}} \Pi(R, J, a) \ge 2n - 1 - \alpha$$

Assume that the above estimate does not hold. Then there exists a compact $E \subset S_n$ such that $\Pi(E) > 0$ and

$$\dim_{\mathcal{H}} E < 2n - 1 - \alpha$$

We identify S_n and $\mathbb{CP}^{n-1} \times \mathbb{T}$, and we consider the sets $E_{\zeta} = \{ w \in \mathbb{T} : (\zeta, w) \in E \}$. We claim that

(3.6)
$$\dim_{\mathcal{H}} E_{\zeta} \leq 1 - \alpha \quad \text{for } \widehat{\sigma}_n \text{-almost all } \zeta \in \mathbb{CP}^{n-1}.$$

Indeed, if this is not the case, then $\dim_{\mathcal{H}} E_{\zeta} > 1 - \alpha$ for $\zeta \in X$, where $\widehat{\sigma}_n(X) > 0$. Applying Theorem 3.5, we obtain $\dim_{\mathcal{H}} E \geq 2n - 1 - \alpha$, a contradiction. Thus, (3.6) holds.

Since $\Pi(E) > 0$, Lemma 2.1 guarantees that $\Pi_{\zeta}(E_{\zeta}) > 0$ for $\zeta \in Y_0$ with $\widehat{\sigma}_n(Y_0) > 0$. Combining this fact and (3.6), we obtain

(3.7)
$$\dim_{\mathcal{H}} \Pi_{\zeta} \leq 1 - \alpha \quad \text{for } \zeta \in Y \text{ with } \widehat{\sigma}_n(Y) > 0$$

Now, recall that $\Pi_{\zeta}(R, J, a)$ is the classical Riesz product defined by (2.2). Therefore, $\dim_{\mathcal{H}} \Pi_{\zeta}(R, J, a) \geq 1 - \alpha_0 > 1 - \alpha$ by [11, Theorem 3.1]. This contradicts (3.7). Thus, (3.5) holds true for any $\alpha > \alpha_0$, as required.

In other words, consideration of slice-products allows to recover Corollary 3.3 from [11, Theorem 3.1], the corresponding result for the classical Riesz products.

4. HAUSDORFF DIMENSIONS OF PLURIHARMONIC MEASURES

This section is motivated by analogs of Lemma 2.1 and their applications in the setting of pluriharmonic measures.

4.1. Pluriharmonic measures on the unit sphere. A measure $\mu \in M(S_n)$ is called pluriharmonic if $\operatorname{spec}(\mu) \subset \{(p,q) \in \mathbb{Z}^2_+ : pq = 0\}$. An analog of Lemma 2.1 is known for the pluriharmonic measures on S_n . This fact implies that $\dim_{\mathcal{H}} \mu \geq 2n-2$ for any pluriharmonic measure $\mu \in M(S_n)$, see [1, 3] for further details. However, to the best of the author's knowledge, the sharpness of this estimate is an open problem.

4.2. Pluriharmonic measures on the torus. A measure $\mu \in M(\mathbb{T}^n)$ is called pluriharmonic if $\hat{\mu}(k_1, \ldots, k_n) = 0$ for $(k_1, \ldots, k_n) \in \mathbb{Z}^n \setminus (\mathbb{Z}^n_- \cup \mathbb{Z}^n_+)$. It is well known that $\mu \in M(\mathbb{T}^n)$ is pluriharmonic if and only if the Poisson integral of μ is a pluriharmonic function in the polydisk \mathbb{D}^n . Therefore, an analog of Lemma 2.1 holds true for every pluriharmonic measure μ on \mathbb{T}^n . See [2, Proposition 2.1] for the case of μ on the unit sphere S_n , see also [5], where pluriharmonic measures on

$$\mathbf{6}$$

the Shilov boundary of a bounded symmetric domain are considered. Similar to the case of S_n , we conclude that

(4.1)
$$\dim_{\mathcal{H}} \mu \ge n-1;$$

cf. [5, Corollary 3.3]. For n = 2, estimate (4.1) and its sharpness were earlier proved in [4]. Simple examples show that estimate (4.1) is sharp for all $n \ge 2$. Indeed, put

$$\mu = \delta_1(\xi_1) \otimes m(\xi_2) \otimes \cdots \otimes m(\xi_n),$$

where δ_1 denotes the Dirac measure at $1 \in \mathbb{T}$ and $m = \sigma_1$ denotes the normalized Lebesgue measure on \mathbb{T} . Clearly, μ is a pluriharmonic measure and $\dim_{\mathcal{H}} \mu = n - 1$.

References

- A. B. Aleksandrov, Function theory in the ball, Current problems in mathematics. Fundamental directions, Vol. 8, Itogi Nauki i Tekhniki, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1985, pp. 115–190, 274 (Russian); English transl., Encyclopaedia Math. Sci., vol. 8, Springer–Verlag, Berlin, 1994, pp. 107–178.
- [2] A. B. Aleksandrov and E. Doubtsov, Clark measures on the complex sphere, J. Funct. Anal. 278 (2020), no. 2, 108314, 30.
- [3] R. Ayoush and M. Wojciechowski, Microlocal approach to the Hausdorff dimension of measures, Adv. Math. 395 (2022), Paper No. 108088, 11pp.
- [4] L. Bergqvist, Necessary conditions on the support of RP-measures, arXiv:2304.03072.
- [5] M. Calzi, Clark measures on bounded symmetric domains, arXiv:2403.05429.
- [6] E. Doubtsov, Mutual singularity of Riesz products on the unit shpere, arXiv:2404.02652.
- [7] E. Doubtsov, Henkin measures, Riesz products and singular sets, Ann. Inst. Fourier (Grenoble) 48 (1998), no. 3, 699–728.
- [8] K. Falconer, Fractal geometry, John Wiley & Sons, Ltd., Chichester, 1990. MR 1102677
- K. Falconer, Techniques in fractal geometry, John Wiley & Sons, Ltd., Chichester, 1997. MR 1449135
- [10] H. Federer, Geometric measure theory, Die Grundlehren der mathematischen Wissenschaften, vol. Band 153, Springer-Verlag New York, Inc., New York, 1969. MR 257325
- [11] K. E. Hare and M. Roginskaya, A Fourier series formula for energy of measures with applications to Riesz products, Proc. Amer. Math. Soc. 131 (2003), no. 1, 165–174. MR 1929036
- [12] K. E. Hare and M. Roginskaya, Multipliers of spherical harmonics and energy of measures on the sphere, Ark. Mat. 41 (2003), no. 2, 281–294. MR 2011922
- [13] W. Rudin, Function theory in the unit ball of \mathbb{C}^n , Grundlehren der Mathematischen Wissenschaften, vol. 241, Springer-Verlag, New York-Berlin, 1980.
- [14] J. Ryll and P. Wojtaszczyk, On homogeneous polynomials on a complex ball, Trans. Amer. Math. Soc. 276 (1983), no. 1, 107–116. MR 684495
- [15] A. Zygmund, Trigonometric series. Vol. I, II, third ed., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2002. MR 1963498

St. Petersburg Department of Steklov Mathematical Institute, Fontanka 27, St. Petersburg 191023, Russia

Email address: dubtsov@pdmi.ras.ru