EINSTEIN MANIFOLDS WITH OPTICAL GEOMETRIES OF KERR TYPE

MASOUD GANJI, CRISTINA GIANNOTTI, GERD SCHMALZ, AND ANDREA SPIRO

ABSTRACT. We classify the Ricci flat Lorentzian *n*-manifolds satisfying three particular conditions, encoding and combining some crucial features of the Kerr metrics and the Robinson-Trautman optical structures. We prove that: (a) If n > 4, there is no Lorentzian manifold satisfying the considered Kerr type conditions, in unexpected contrast with what occurs for the metrics satisfying (very similar) Taub-NUT type conditions; (b) If n = 4 there are two large classes of such Kerr type manifolds. Each class consists of manifolds fibering over open Riemann surfaces, equipped with a metric of constant Gaussian curvature $\kappa = 1$ or $\kappa = -1$. The first class includes a three parameter family of metrics admitting real analytic extensions to $(\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R} = (S^2 \times \mathbb{R}_+) \times \mathbb{R}$ and a large class of other metrics not admitting this kind of extensions. The metrics of this first class admitting such extensions are all isometric to the well known Kerr metrics, with the three parameters corresponding to the three space-like components of the angular momentum of the gravitational field. The second class contains a subclass of metrics defined on $(\mathbb{D} \times \mathbb{R}_+) \times \mathbb{R}$, where \mathbb{D} is the Lobachevsky Poincaré disc. This subclass is in bijection with the holomorphic functions on \mathbb{D} satisfying an appropriate open condition. These and other results are consequences of a very simple way to construct totally explicit examples of Ricci flat Lorentzian manifolds.

1. INTRODUCTION

The main result of this paper is the classification of the even dimensional Lorentzian manifolds, admitting families of Ricci flat metrics that are compatible with a twisting optical structure of Robinson-Trautman type and satisfying certain constraints, which encode some of the most important features of the 4-dimensional Kerr metrics. The results and their proofs have two noticeable consequences: (1) a no-go theorem for higher dimensional Einstein manifolds with twisting optical structures and (2) a simple method of construction of a very large class of explicit Ricci flat 4-dimensional Lorentzian manifolds, parameterised by the solutions of a linear elliptic p.d.e. in two variables.

Before going into the details of our results, we need to recall a few known facts on Kerr metrics. These metrics are Lorentzian metrics, depending on real parameters m and a, which are defined on the 4-dimensional manifold $M^{\text{Kerr}} = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R} = (S^2 \times \mathbb{R}_+) \times \mathbb{R}$

²⁰²⁰ Mathematics Subject Classification. 53B30, 83C05.

Key words and phrases. Einstein equations; Ricci flat Lorentzian manifolds; Optical geometry; Kerr metrics.

and having the following coordinate expressions:

$$g^{\text{Kerr}} = -\frac{1}{2}(\rho^2 + a^2\cos^2\xi)(d\xi^2 + \sin^2\xi d\psi^2) - (dv + a\sin^2\xi d\psi) \vee (d\rho + a\sin^2\xi d\psi) + \frac{1}{2}\left(1 - \frac{2m\rho}{\rho^2 + a^2\cos^2\xi}\right)(dv + a\sin^2\xi d\psi)^2.$$
(1.1)

Here ψ , ξ are the usual spherical coordinates of S^2 , ρ , v are the standard Cartesian coordinates of $(0, +\infty) \times \mathbb{R}$ and m, a are the parameters which are physically interpreted as the Schwarzschild mass of the gravitational field, and the ratio $a = \mathcal{J}/m$ between the angular momentum $\mathcal{J} \neq 0$ of the gravitational field and the mass m (see e.g. [17] (¹)).

There is however an alternative and very useful presentation of g^{Kerr} or, more precisely, of its restriction to the open subset

$$M_{(+)}^{\text{Kerr}} = (S_{(+)}^2 \times \mathbb{R}_+) \times \mathbb{R} \subset M^{\text{Kerr}}, S_{(+)}^2 := \text{upper hemisphere} = \left\{ (\xi, \psi) : \xi \in \left[0, \frac{\pi}{2}\right) \right\} ,$$

that makes manifest that g^{Kerr} is compatible with a particular twisting optical structure.

The notion of optical structure was introduced by Robinson and Trautman to study Lorentzian manifolds admitting electromagnetic plane waves (or appropriate higher dimensional generalisations) propagating along a prescribed foliation by null geodesics(see [29, 30, 33] – see also [10, 2, 3, 4] and references therein for coordinate-free presentations and discussions of the equivalences among the diverse presentations of the same objects). In detail, an *optical structure* on a manifold M is a pair $\Omega = (\mathcal{K}, \{g\})$ given by

- a 1-dimensional distribution $\mathcal{K} \subset TM$,
- the collection $\{g\}$ of all Lorentzian metrics g, such that: (1) the curves that are tangent to \mathcal{K} determine a geodesic shearfree null congruence for M; (2) the g-orthogonal distribution \mathcal{K}^{\perp_g} is a fixed distribution $\mathcal{W} = \mathcal{K}^{\perp_g} \subset TM$, independent of g (²).

The metrics in the class $\{g\}$ are said to be *compatible* with the optical structure Q.

An important class of optical structures is given by those on a special class of bundles, which we now very briefly review (they are extensively discussed in [3], where they are called *Robinson-Trautman shearfree structures*). Consider a 2k-dimensional quantisable Kähler manifold (N, J, g_o) . This means that there exists: (a) a principal bundle $\pi : S \to N$, with 1-dimensional structure group $A = \mathbb{R}_+ \simeq \mathbb{R}$ or S^1 and (b) a connection 1-form $\theta : TS \to \text{Lie}(A) = \mathbb{R}$ such that $d\theta = \pi^* \omega_o$, where $\omega_o := g_o(J, \cdot)$ is the Kähler form of (N, J, g_o) . Then take the trivial principal \mathbb{R} -bundle $\pi^S : M = S \times \mathbb{R} \to S$ over S and denote by ϑ and p_o the pulled-back 1-form $\vartheta = \pi^{S*}\theta$ on M and the vertical vector field $p_o = \frac{\partial}{\partial u}$ generating the right \mathbb{R} -action on $M = S \times \mathbb{R}$, respectively. It is known (see [3, 4]

¹Note that (1.1) is actually a metric which is a re-scaling of the one in [16] by the factor $-\frac{1}{2}$. It is a normalisation that better fits with our purposes. Notice also that we are considering the signature (-, -, -, +) instead of the signature (+, +, +, -) used in [16].

²A geodesic shearfree congruence is a foliation by null geodesics that are tangent to a vector field p_o , whose flow leaves invariant the g-orthogonal distribution $\mathcal{W} = \mathcal{K}^{\perp_g}$ and the conformal class of $g|_{\mathcal{W}\times\mathcal{W}}$.

and §2.2 of this paper) that the pair $\Omega = (\mathcal{K}, \{g\})$ given by the 1-dimensional distribution $\mathcal{K} = \langle \mathbf{p}_o \rangle \subset TM$ and the class $\{g\}$ of all Lorentzian metrics on M having the form

$$g = \sigma(\pi \circ \pi^{\delta})^* g_o + \vartheta \vee \left(\mathbf{p}_o^* + \mathbf{\mathfrak{d}} + \frac{\tilde{\beta}}{2} \vartheta \right) , \quad \text{where}$$
(1.2)

- (i) p_o^* is \mathbb{R} -invariant, satisfies $p_o^*(p_o) = 1$ and is such that $\mathcal{H} := \ker p_o^* \cap \ker \vartheta$ is the horizontal distribution in $TS \subset TM \simeq TS +_M \mathbb{R}$,
- (ii) $\sigma, \tilde{\beta}$ are freely specifiable smooth functions with σ nowhere vanishing, and \mathfrak{d} is a freely specifiable 1-form with ker \mathfrak{d} containing the vertical distribution of $\pi \circ \pi^{\mathfrak{S}} : M \to N$,

is an optical structure on $M = \mathcal{S} \times \mathbb{R}$. For such an optical structure, the common *g*orthogonal distribution $\mathcal{W} = \mathcal{K}^{\perp_g}$ is the kernel distribution $\mathcal{W} = \ker \vartheta$ and it is such that dim ker $d\vartheta|_{\mathcal{W}\times\mathcal{W}} = 1$. Note that 1 is the minimal possible value that might occur for the dimensions of the kernels ker $d\alpha_x|_{\mathcal{W}\times\mathcal{W}}$, $x \in M$, for an arbitrary 1-form α that vanishes identically on \mathcal{W} . The optical structures for which this property occurs are called *twisting* ([29, 30, 3, 10]).

Coming back to the Kerr metrics, we can immediately observe that:

- (a) $M_{(+)}^{\text{Kerr}}$ is a trivial \mathbb{R} -bundle over the \mathbb{R}_+ -bundle $\pi : \mathcal{S} = S_{(+)}^2 \times \mathbb{R}_+ \to S^{(+)};$
- (b) the restricted Kerr metric $g^{\text{Kerr}(+)} := g^{\text{Kerr}}|_{M_{(+)}}$ has the form (1.2) with $\mathfrak{d} \equiv 0$, the function σ , $\tilde{\beta}$ given by

$$\sigma = -\frac{\rho^2}{4a\cos\xi} - \frac{a\cos\xi}{4} , \qquad \tilde{\beta} = -1 - \frac{2m\rho}{\rho^2 + a^2\cos^2\xi} , \qquad (1.3)$$

and the Kähler metric g_o and the 1-forms ϑ , p_o^* given by

$$g_o := 2a\cos\xi (d\xi^2 + \sin^2\xi d\psi^2) , \qquad \vartheta := dv + a\sin^2\xi d\psi , \qquad \mathbf{p}_o^* := dv - d\rho .$$
(1.4)

It follows that for any choice of a, m, the corresponding (restricted) Kerr metric $g^{\text{Kerr}(+)}$ on $M_{(+)}^{Kerr} = (S_{(+)}^2 \times \mathbb{R}_+) \times \mathbb{R}$ is compatible with the twisting optical structure Ω constructed as above on an \mathbb{R} -bundle over the quantisable Kähler manifold

$$\left(S_{(+)}^2, g_o := 2a\cos\xi(d\xi^2 + \sin^2\xi d\psi^2)\right)$$
.

The Kerr metrics (1.1) are the unique real analytic extensions of such Ω -compatible metrics on $M_{(+)}^{\text{Kerr}}$ to the whole manifold $M^{\text{Kerr}} = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$ and are Ricci flat, as is well known.

At this point it is important to recall that another well-known class of Q-compatible metrics on $M_{(+)}^{\text{Kerr}}$ is given by the classical 4-dimensional Taub-NUT metrics. Actually, as it has been shown in [13], the classical Kerr and Taub-NUT metrics can be locally combined into a single smoothly parameterised Q-compatible family of Ricci flat metrics on $M_{(+)}^{\text{Kerr}}$.

Looking for new Einstein manifolds with optical geometries and inspired by the method of construction in [3] of the higher dimensional versions of the Taub-NUT metrics [5, 26, 8, 1], we decided to focus on a class of manifolds with optical structures which naturally includes $M_{(+)}^{\text{Kerr}}$, namely the connected Lorentzian manifolds $M = S \times I$, $I \subset \mathbb{R}$, where S is the total space of an \mathbb{R} -bundle $\pi : S = N \times \mathbb{R} \to N$ (³) over a quantisable Kähler manifold (N, J, g_o) , equipped with the optical structure Q determined by the above recipe. We call them *Kerr manifolds* (see below, Definition 2.3).

In this paper we tackle the classification problem of the Kerr manifolds admitting 1parameter families $g^{(m)}$ of compatible Einstein metrics, parameterised by a real number mand for which the data $\mathfrak{d}, \sigma, \tilde{\beta}$ appearing in (1.2) have the following form

- (1) $\mathfrak{d} \equiv 0$,
- (2) $\tilde{\beta} = B + m\tilde{\beta}_o$ for some constant $B \neq 0$ and a smooth *m*-independent function $\tilde{\beta}_o$,
- (3) σ is independent of m

and are constrained by the condition $\mathbb{T}(\sigma) = \mathbb{T}(\widetilde{\beta}_o) = 0$, with $\mathbb{T} = \mathbf{q}_o - B\mathbf{p}_o$ where \mathbf{q}_o , \mathbf{p}_o are null and vertical with respect to the projection $\pi \circ \pi^{\mathbb{S}} : M \to N$ and such that $g(\mathbf{q}_o, \mathbf{p}_o) = 1 = \vartheta(\mathbf{q}_o)$. This constraint is equivalent to impose that each metric $g^{(m)}$ is invariant with respect to \mathbb{T} , a condition which generalises the time-invariance condition satisfied by the classical Kerr metrics. We call the families of metrics $g^{(m)}$ of this kind *Kerr families of gravitational fields*. For each of them, the Einstein metric $g^{(m=0)}$ is called *background (of Kerr type)*. Such a name is motivated by the fact that, for any fixed $a \neq 0$, the Kerr metric corresponding to m = 0 is

$$g^{\text{Kerr}} = -\frac{1}{2} (\rho^2 + a^2 \cos^2 \xi) (d\xi^2 + \sin^2 \xi d\psi^2) - (dv + a \sin^2 \xi d\psi) \vee (d\rho + a \sin^2 \xi d\psi) + \frac{1}{2} (dv + a \sin^2 \xi d\psi)^2 , \qquad (1.5)$$

a metric that can be transformed into the standard expression of the flat Minkowski metric under an appropriate change of coordinates. This is not a coincidence: in Proposition 2.9 we prove that *any background metric on any 4-dimensional Kerr manifold is flat*, so that it can be reasonably considered as a metric for a background geometry in a physical sense.

We stress the fact that the conditions that characterise the Kerr families are very close to those that define the Taub-NUT metrics and their higher dimensional versions: To pass from one set of conditions to the other it suffices to set B = 0 and remove the requirement of linear dependence of $\tilde{\beta}$ from a "mass" parameter m. In fact, by the classification in [3, Thm. 5.2], the class of the Einstein metrics satisfying these modified conditions are precisely the above mentioned generalisations of the 4-dimensional Taub-NUT metrics for arbitrary dimensions and with arbitrary Einstein constants.

In this paper we succeed in classifying the Kerr manifolds admitting Kerr families of *Ricci flat* gravitational fields and explicitly determine coordinate expressions for such Ricci flat Kerr families. It is reasonable to expect that a similar classification can be reached by the same tools also for the case of non-Ricci flat Einstein metrics. The following theorem summarises our main results.

³ We recall that, by a classical result on principal bundles (see [31, Cor. 12.3]), each \mathbb{R} -bundle is trivial.

Theorem 1.1. Let $M = \mathbb{S} \times I$, $I \subset \mathbb{R}$, be a Kerr manifold of dimension $n = 2k + 2 \ge 4$. We denote by $\pi^{\mathbb{S}} : M = \mathbb{S} \times I \to \mathbb{S}$ the standard projection onto the total space of the \mathbb{R} bundle $\pi : \mathbb{S} = N \times \mathbb{R} \to N$ over a Kähler manifold (N, J, g_o) , equipped with a connection 1-form θ such that $\pi^* \omega_o = d\theta$. We also denote by ϑ the pull-back $\vartheta := (\pi^{\mathbb{S}})^* \theta$ on M and by \mathfrak{Q}_o the canonical optical structure on $\mathbb{S} \times \mathbb{R}$ described above. Then:

- (i) If n > 4, there is no background and no Kerr family of gravitational fields on M;
- (ii) If n = 4, there exists a Kerr family $g^{(m)}$ of Ricci flat gravitational fields on M if and only if
 - (a) (N, J) admits a Kähler metric \tilde{g}_o of constant Gaussian curvature $\kappa = \pm 1$;
 - (b) there is a globally defined Kähler potential $\varphi : N \to \mathbb{R}$ for g_o (i.e. a real function such that $g_o = dd^c \varphi(\cdot, J \cdot)$) satisfying the constraint $\operatorname{sign}(\varphi) \equiv -\kappa$ and the linear elliptic equation

$$dd^c \varphi(\cdot, J \cdot) + 2 \kappa \varphi \,\widetilde{g}_o = 0 ; \qquad (1.6)$$

(c) modulo a bundle equivalence of $\pi^{\mathbb{S}} : M = \mathbb{S} \times I \to \mathbb{S}$, there is a choice for the coordinate r for the factor $I \subset \mathbb{R}$ of $M = \mathbb{S} \times I$ and a (locally defined) coordinate set (x, y, v, r) with (x, y) coordinates for N and v for the fiber of $\pi : \mathbb{S} = N \times \mathbb{R} \to N$, such that the 1-form ϑ and the Lorentzian metrics $g^{(m)}$ take the form

$$\vartheta = dv + (\pi \circ \pi^{\delta})^* (d^c \varphi)$$

$$g^{(m)} = -\frac{\kappa}{2} \left(r^2 + \varphi^2 \right) (\pi \circ \pi^{\delta})^* \widetilde{g}_o + \vartheta \vee \left(d(v+r) + \frac{1}{2} \left(-1 + \frac{\mathbf{k}m r}{r^2 + \varphi^2} \right) \vartheta \right) , \qquad (1.7)$$

where \mathbf{k} is just a rescaling constant for the parameter m.

Claim (i) is a consequence of a no-go theorem (see Corollary 3.2) according to which no Kerr manifold of dimension greater than 4 admits an Einstein metric that is compatible with its twisting optical structure Q. We have to admit that such a result was a surprising outcome to us, being somehow in contrast with the above mentioned abundance of higher dimensional analogs of the Taub-NUT metrics (⁴). For what concerns claim (ii), it is a combination of various results that we prove for the 4-dimensional Kerr manifolds – see Corollary 4.3, Theorem 5.1, Lemma 2.7 and formula (6.7).

The geometrical constraints provided by the claims (i), (ii-a) and (ii-b) of Theorem 1.1 are crucially rooted on the "twisting" property of the considered optical structures and on the invariance under the vector field $\mathbb{T} = \mathbf{q}_o - B\mathbf{p}_o$ with $B \neq 0$. The existence of such constraints is in apparent contrast with the following well-known facts: (1) The classical Kerr metrics are well defined over the whole bundle $M^{\text{Kerr}} = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R} \to S^2$ – such a bundle fibers over the *compact* Kähler surface S^2 , which therefore cannot admit any globally defined Kähler potential; (2) There do exist higher dimensional analogs of Kerr black holes ([22, 15]). However, there is no actual contradiction – everything is consistent. In fact the metrics in (1) and (2) are simply not *globally* compatible with any twisting optical structures. More precisely: (a) The classical Kerr metrics are compatible with a

⁴However, one can notice that our no-go theorem is consistent with Taghavi-Chabert's classification of higher dimensional Ricci flat metrics with mWAND structures [32] – see also [24, 25].

prescribed optical structure on $M^{\text{Kerr}} = S^2 \times \mathbb{R}_+ \times \mathbb{R}$, but such an optical structure is twisting only if it is restricted to $M^{\text{Kerr}}_{(+)} = S^2_{(+)} \times \mathbb{R}_+ \times \mathbb{R}$ (or to $M^{\text{Kerr}}_{(-)} = S^2_{(-)} \times \mathbb{R}_+ \times \mathbb{R}$); (b) In [27] (see also [22, 11]) it is proven that the geodesic null congruences of Mayer and Perry's higher dimensional black holes are not shearfree and are therefore not compatible with any (neither twisting nor with no twist) optical structure as in the above definition.

We thus think that the two conditions, given by the compatibility with a prescribed (twisting) optical structure and the invariance under a vector field $\mathbb{T} = \mathbf{q}_o - B\mathbf{p}_o$ with $B \neq 0$, are somehow conflicting in dimensions greater than 4. In fact, removing the condition $B \neq 0$ allows the construction of the higher dimensional Taub-NUT metrics, while removing the compatibility requirement with a prescribed optical structure opens the way to the construction of Mayer and Perry's higher dimensional black holes.

Despite of the just discussed limitations, claim (ii) of Theorem 1.1 shows the existence of a very large class of families of 4-dimensional Ricci flat gravitational fields of Kerr type, being such families easily constructible for any choice of a non-compact Riemann surface (N, J). In fact, they are in bijections with the solutions φ to (1.6) with the prescribed sign property. Using isothermal coordinates for \tilde{g}_o , in which the metric is given by the classical formula $\tilde{g}_o = \frac{4(dx^2+dy^2)}{(1+\kappa(x^2+y^2))^2}$, such conditions on φ take the form

$$\Delta \varphi + \kappa \frac{8\varphi}{\left(1 + \kappa (x^2 + y^2)\right)^2} = 0 \qquad \& \qquad \kappa \varphi < 0 . \tag{1.8}$$

Classical results on second order elliptic equations yield that the solutions to (1.8) in a disk $\mathbb{D}(0, \mathfrak{r}_o) \subset \mathbb{R}^2$ of radius $\mathfrak{r}_o < 1$, are real analytic and in bijection with the Fourier series of the non-negative real analytic functions on the boundary of such a disk. Using this, in §6 we succeed in writing explicitly all solutions to (1.8) over a closed disk as sums of series. Moreover, considering the complex coordinate z = x + iy, we get that the solutions to (1.8) over a closed disk $\overline{\mathbb{D}(0, \mathfrak{r}_o)}$ coincide with the functions of the form

$$\varphi(z) = \operatorname{Re}\left(F(z) - \frac{2|z|^2}{1+|z|^2} \frac{1}{z} \int_0^z F(\zeta) d\zeta\right) \text{ in case } \kappa = 1 \text{ or}$$

$$\varphi(z) = \operatorname{Re}\left(\frac{1+|z|^2}{1-|z|^2} F(z) + z \frac{dF(z)}{dz}\right) \text{ in case } \kappa = -1 ,$$
(1.9)

for some holomorphic function F constrained just by the sign condition $\kappa\varphi < 0$. Notice that F may be defined as a holomorphic function on any subset of N. In this regard we would like to mention that the contents of Theorem 1.1 (ii) are very likely related with Debney, Kerr and Schild's classification of Ricci flat algebraic degenerate metrics with non-zero complex expansion rate [9]. Indeed, on the one hand, as a consequence of our Proposition 2.9, any of our metrics (1.7) admits the presentation (2.16) and is therefore a *Kerr-Schild metric* (see e.g. [34, Ch. 32] for the definition and references to the vast literature on this class of Lorentzian metrics). On the other hand, the Ricci flat metrics considered in [9] are proved to be not only Kerr-Schild and compatible with an appropriate geodesic shearfree null congruence, but also including a distinguished subclass of metrics with a one-dimensional isometry group and parameterised by the real parts of the solutions of a certain ordinary complex differential equation, a result which is strongly reminiscent of ours. Investigations on the relation between these two noticeable classes of metrics are left to future activities.

As it is natural to expect, among the solutions to (1.8) for $\kappa = +1$, there are potentials φ which, under an appropriate change of coordinates, take the form $\varphi(\xi, \psi) = a \cos \xi$, $a \in \mathbb{R}$, and are thus corresponding to the classical Kerr metrics (1.1). Moreover, the metrics which satisfy the conditions:

- (i) the corresponding potential φ is as in (1.9) with F polynomial,
- (ii) they admit real analytic extensions to the whole $M^{\text{Kerr}} = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$,

are precisely the same metrics that are obtained from (1.1) through a space-like rotation of the flat background. We expect that (i) is not a really necessary condition and that only the Kerr metrics (and their rotated versions) are metrics of the form (1.7) that admit real analytic extensions to the whole M^{Kerr} , all the others being forced to develop curvature singularities of various kinds. It would be consistent with the known uniqueness theorems for the Kerr metrics (see e.g. [28]). Investigations in this regard are left to future works.

For what concerns the case $\kappa = -1$, a very large number of local functions satisfying (1.8) exist and, in contrast with the case $\kappa = +1$, a huge part of them (namely, those corresponding to complex functions F that are holomorphic over the whole unit disk $\mathbb{D} := \mathbb{D}(0,1)$ and for which the function φ defined in (1.9) is positive on at least a subset of \mathbb{D}) admit unique real analytic extensions over the whole 2-dimensional Lobachevsky space $(\mathbb{D}, \tilde{g}_o)$. The associated Ricci flat metrics are well defined on the Kerr manifold $M^{\text{Lob}} := (\mathbb{D}^2 \times \mathbb{R}) \times \mathbb{R}$ and, at least on the regions in which $\sup |\varphi|$ is sufficiently small, they can be considered as deformations of a Lorentzian metric that has $\mathbb{R} \times SO_{1,2}^o$ as connected component of the isometry group (this mirrors the fact that the Kerr metrics can be considered as deformations of the Schwarzschild metric, i.e. of a metric for which the connected component of the isometric group is $\mathbb{R} \times SO_3$). It follows that most of the metrics of this second class are not even locally isometric to the classical Kerr metrics.

We expect that, by appropriately choosing the holomorphic function F that determines the potential φ by (1.9), the corresponding metric has an isometry group of dimension not greater than 1 (i.e., with connected component given just by the flow of the vector field \mathbb{T}) and that it is not isometric to any of the previously known exact solutions to the Einstein equations. Investigations on this issue and on diverse aspects of the geometries of the Ricci flat Lorentzian manifolds given in this paper are left to future work. However, the coordinate-free presentations of these manifolds and of their compatible metrics seem to us to be quite novel and suggest diverse intriguing questions. For instance, it should be noticed that the Ricci flat metrics on $M^{\text{Lob}} = (\mathbb{D} \times \mathbb{R}) \times \mathbb{R}$ described above are not static in time and not asymptotically spherically symmetric. Therefore they are not suited to model stationary gravitational fields generated by localised masses. Nonetheless, they might be considered as non-static cosmological models or, more precisely, as limit forms (for stress-energy tensors tending to 0) of non-static cosmological models, quite different from the most commonly considered cosmological models. As a concluding remark, we would like to emphasise that some points of our proofs are crucially based on long sequences of straightforward computations. We are aware this makes the verification difficult. In order to avoid any possible errors, we scrupulously rechecked each sequence of computations with the help of the symbolic manipulation program *Maple*. Note that, with the help of this or similar software, everybody can quickly verify *a posteriori* the Ricci flatness of any of the 4-dimensional metrics given in our classification. Shorter proofs than ours are surely desirable and we hope that they will come up in the near future.

The structure of the paper is as follows. The section §2 is devoted to a brief review of some facts on optical structures, to giving a convenient parameterisation of the compatible metrics of an optical structure and to introducing the definitions and first properties of Kerr manifolds, backgrounds of Kerr type and Kerr families of gravitational fields. In §3, we prove our no-go theorem for Einstein Kerr manifolds in higher dimensions and a corollary that implies (i) of Theorem 1.1. In §4 and §5, we classify the 4-dimensional Kerr manifolds admitting backgrounds of Kerr type and Kerr families of Ricci flat gravitational fields, proving in this way (ii) of Theorem 1.1. In section §6, explicit expressions for the potentials of the metrics of our classification are determined and the above mentioned discussion of the sets of solutions corresponding to $\kappa = \pm 1$ is given in greater detail. In two appendices, we provide the details of some computations for the proofs in §3 and §4.

Acknowledgments. We are very grateful to Marcello Ortaggio for his invaluable assistance in providing us with several pieces of information and constructive feedback.

Notation. For any pair of 1-forms α, β on a manifold M, the symbol $\alpha \vee \beta$ stands for the symmetric tensor product $\alpha \vee \beta := \frac{1}{2} (\alpha \otimes \beta + \beta \otimes \alpha)$. Given a vector field X and a pseudo-Riemannian metric g on M, we denote $X^{\flat} := g(X, \cdot)$.

2. KERR FAMILIES OF GRAVITATIONAL FIELDS ON KERR TYPE BACKGROUNDS

2.1. **Optical geometries.** Let (M, g) be a Lorentzian *n*-manifold. A null congruence on M is a foliation of the manifold by curves, which are tangent to a nowhere vanishing null vector field at all points. The following objects are naturally associated with a null congruence:

- The class {p} of all null vector fields, which are tangent to the curves of the foliation;
- The codimension one distribution $\mathcal{W} := \mathbf{p}_o^{\perp}$ which is *g*-orthogonal to some (hence, to all) nowhere vanishing vector field $\mathbf{p}_o \in \{\mathbf{p}\}$;
- The degenerate metric $h = g|_{\mathcal{W} \times \mathcal{W}}$ induced on \mathcal{W} .

The null congruence is called *shearfree* if there is a vector field p in the class $\{p\}$, whose (local) flow preserves \mathcal{W} and the conformal class [h] of h, i.e.

 $\mathcal{L}_{p}X \in \mathcal{W}$ for any $X \in \mathcal{W}$ and $\mathcal{L}_{p}h = fh$ for some function $f \in \mathcal{F}(M)$. (2.1)

This is in turn equivalent to the condition (see e.g. [3, Lemma 2.1])

$$\mathcal{L}_{\mathbf{p}}g = fg + \mathbf{p}^{\flat} \lor \eta$$
 for some function f and a 1-form η (2.2)

and, if this is the case, then p is geodesic and all curves of the congruence are geodesics ([3, Prop. 2.4]). Due to this, often the shearfree null congruences are alternatively called *geodesic shearfree null congruences*.

The property of a foliation by curves to be a shearfree congruence with respect to some metric g is encoded in the following definition, which have appeared (in possibly different but all equivalent ways) in [29, 3, 4]. Actually, the following is a very mild generalisation of the previous, because it deals with Lorentzian metrics that might have any of the two possible types of signatures, the mostly plus and the mostly minus.

Definition 2.1. An optical geometry on M is a quadruple $\mathfrak{Q} = (\mathcal{W}, [h]_{\pm}, \mathcal{K}, \{g\})$, given by

- (i) a codimension one distribution $\mathcal{W} \subset TM$,
- (ii) the union $[h]_{\pm} = [h] \cup [-h]$ of two conformal classes [h], [-h], one consisting of semipositive degenerate metrics h on \mathcal{W} with one-dimensional kernels, the other made of their opposite metrics -h,
- (iii) the 1-dimensional distribution \mathcal{K} , given by the kernels $\mathcal{K}|_x = \ker h_x \subset \mathcal{W}_x, x \in M$, of the degenerate metrics h,
- (iv) the class $\{g\}$ of all Lorentzian metrics (with either mostly plus or mostly minus signs) inducing on \mathcal{W} one of the degenerates metrics in $[h]_{\pm}$,

satisfying the following conditions: the class $\{g\}$ is not empty and there exists a vector field p in \mathcal{K} such that (2.1) holds. The metrics in $\{g\}$ are said to be compatible with \mathcal{Q} .

Remark 2.2. If $Q = (W, [h], \mathcal{K}, \{g\})$ is an optical geometry, both pairs (W, [h]) and $(\mathcal{K}_h = \ker h, \{g\})$ provide sufficient information to recover the other terms of the quadruple. Thus, an optical geometry can be equivalently defined as a pair of the form (W, [h]) or of the form $(\mathcal{K}, \{g\})$, provided that its two terms satisfy appropriate conditions. The pairs (W, [h]) of such a kind have been named *shearfree structures* in [3], while the corresponding pairs $(\mathcal{K}, \{g\})$ are *Robinson and Trautman's optical geometries* considered in [29].

2.2. Kerr manifolds.

Definition 2.3. A Kerr structure of dimension n = 2k is a quadruple

$$\mathcal{M} = (M = \mathcal{S} \times \mathbb{R}, \pi : \mathcal{S} \to N, (J, g_o), \mathcal{H}) ,$$

given by

- (a) An *n*-dimensional manifold M, identifiable with a Cartesian product $M = \mathbb{S} \times \mathbb{R}$ between \mathbb{R} and a total space \mathbb{S} of an \mathbb{R} -bundle $\pi : \mathbb{S} \to N$ as in (b);
- (b) A principal \mathbb{R} -bundle $\pi: \mathbb{S} \to N$ over a Kähler manifold N;
- (c) A Kähler structure (J, g_o) on N;
- (d) A connection $\mathcal{H} \subset TS$ on $\pi : S \to N$, which is the kernel distribution of an \mathbb{R} -invariant connection 1-form θ^{S} satisfying the condition $d\theta^{S} = \pi^{*}\omega_{o}$, in which $\omega_{o} := g_{o}(J \cdot, \cdot)$ is the Kähler form of (N, J, g_{o}) .

A manifold M as in (a), equipped with a Kerr structure $\mathcal{M} = (M = \mathbb{S} \times \mathbb{R}, \pi : \mathbb{S} \to N, (J, g_o), \mathcal{H})$, is called *Kerr manifold*. By a small abuse of language, in what follows we are

going to call "Kerr manifolds" also any connected open subset $M' = \mathbb{S} \times I$, $I \subset \mathbb{R}$, of a Kerr manifold $M = \mathbb{S} \times \mathbb{R}$ as defined above.

Remark 2.4. The identification $M = S \times \mathbb{R}$, given by a Kerr structure on a manifold M, defines a natural \mathbb{R} -action on such a manifold and, consequently, a local \mathbb{R} -action on any connected open subset $M' = S \times I$, $I \subsetneq \mathbb{R}$. On the other hand, any such $M' \subset M$ is clearly diffeomorphic to $S \times \mathbb{R}$ and hence it can be equipped with an \mathbb{R} -action of its own. However such \mathbb{R} -action is essentially different from the local \mathbb{R} -action defined above. So, in what follows, we write $M' = S \times I$, $I \subset \mathbb{R}$, whenever we need to stress that the (local) \mathbb{R} -action to be considered on M' is the one inherited from the \mathbb{R} -action of the larger manifold $M = S \times \mathbb{R}$.

A Kerr manifold $M = \mathbb{S} \times \mathbb{R}$ is naturally equipped with the following optical geometry. Let $\frac{\partial}{\partial u}$ be the standard coordinate vector field of \mathbb{R} and denote by $p_o(\equiv \frac{\partial}{\partial u})$ the naturally corresponding vector field on $M = \mathbb{S} \times \mathbb{R}$. Let also $q_o^{\mathbb{S}}$ be the fundamental vertical vector field of the principal bundle $\pi^{\mathbb{S}} : \mathbb{S} \to N$ corresponding to the standard basis "1" of $Lie(\mathbb{R}) = \mathbb{R}$ (i.e. such that the flow $\Phi_s^{q_o}$, $s \in \mathbb{R}$, is the family of diffeomorphisms, determined by the right actions on the \mathbb{R} -bundle \mathbb{S} of the real numbers $s \in \mathbb{R}$) and denote by $q_o(\equiv q_o^{\mathbb{S}})$ the corresponding vector field on the Cartesian product $M = \mathbb{S} \times \mathbb{R}$. Finally, let $\mathcal{W}_o, \mathcal{K}_o, h_o$ and \check{g}_o be the two distributions, the semi-positive scalar metric on \mathcal{W}_o and the Lorentzian metric that are defined at the points $x = (y, u) \in M = \mathbb{S} \times \mathbb{R}$ by

$$\begin{aligned} \mathcal{W}_{o}|_{(y,u)} &:= \mathcal{H}_{y} + \langle \mathbf{p}_{o}|_{u} \rangle , \qquad \mathcal{K}_{o}|_{(y,u)} := \langle \mathbf{p}_{o}|_{u} \rangle , \\ h_{o}(X + \lambda \mathbf{p}_{o}, Y + \mu \mathbf{p}_{o}) &:= g_{o}(\pi_{*}^{S}(X), \pi_{*}^{S}(Y)) , \qquad X, Y \in \mathcal{H} , \ \lambda, \mu \in \mathbb{R} , \\ \check{g}_{o}(Z + \lambda \mathbf{q}_{o} + \lambda' \mathbf{p}_{o}, Z' + \mu \mathbf{q}_{o} + \mu' \mathbf{p}_{o}) &:= h_{o}(X, Y) + \frac{1}{2}(\lambda \mu' + \lambda' \mu) , \qquad Z, Z' \in \mathcal{H} , \\ \lambda, \lambda', \mu, \mu' \in \mathbb{R} . \end{aligned}$$

$$(2.3)$$

Since \check{g}_o induces h_o on \mathcal{W}_o , the class $\{g\}$ of the metrics inducing an element of $[h_o]_{\pm} = \{\sigma h_o, \sigma > 0 \text{ or } \sigma < 0\}$ on \mathcal{W}_o , is not empty. Moreover, by construction, $\mathcal{L}_{\mathbf{p}_o} X \in \mathcal{W}_o$ for any $X \in \mathcal{W}, \mathcal{L}_{\mathbf{p}_o} h_o = 0$ and $\mathcal{K}_o|_x = \ker h_o|_x, x \in M$. We therefore have that $\mathcal{Q}_o = (\mathcal{W}_o, [h_o]_{\pm}, \mathcal{K}_o, \{g\})$ is an optical geometry.

Definition 2.5. Let M be a Kerr manifold, equipped with a Kerr structure $\mathcal{M} = (M = S \times \mathbb{R}, \pi : S \to N, (J, g_o), \mathcal{H})$. The above defined quadruple \mathcal{Q}_o is called \mathcal{M} -canonical optical geometry of M. The optical geometries of this kind are called of Kerr type.

Remark. The shearfree structures corresponding to optical geometries of Kerr type are typical examples of *shearfree structures of Kähler-Sasaki type* in the sense of [3].

2.3. Parameterisations of compatible metrics and adapted frame fields. Consider a Kerr manifold M, equipped with the Kerr structure $\mathcal{M} = (M = \mathcal{S} \times \mathbb{R}, \pi : \mathcal{S} \rightarrow N, (J, g_o), \mathcal{H})$ and the corresponding \mathcal{M} -canonical optical geometry $\mathcal{Q}_o = (\mathcal{W}_o, [h_o], \mathcal{K}_o, \{g\})$.

Let also \mathbf{p}_o and \mathbf{q}_o be the vector fields defined above and $\check{\pi} = \pi \circ \pi^{\mathbb{S}} : M \to N$ the projection onto N, determined by composing the natural projection $\pi^{\mathbb{S}} : M = \mathbb{S} \times \mathbb{R} \to \mathbb{S}$ with the \mathbb{R} -bundle projection $\pi : \mathbb{S} \to N$. Each tangent space $T_x M$, x = (y, u), admits the following natural direct sum decomposition

$$T_{(y,u)}M = \mathcal{H}_y + \mathbb{R}\mathbf{p}_o|_u + \mathbb{R}\mathbf{q}_o|_y = \mathcal{W}_o + \mathbb{R}\mathbf{q}_o|_y .$$
(2.4)

We remark that a Lorentzian metric g on M is compatible with Ω_o (for simplicity, later, we call it just *compatible*) if and only if $g|_{W \times W} = \sigma \check{\pi}^* g_o$ for a smooth real function $\sigma > 0$ or $\sigma < 0$ at all points. This means that a compatible metric g is uniquely determined by the following four objects: the function $\sigma \neq 0$, the two functions

$$x \longmapsto g_x(\mathbf{p}_o, \mathbf{q}_o) , \qquad x \longmapsto g_x(\mathbf{q}_o, \mathbf{q}_o)$$

and the tensor field in \mathcal{H}^* that is defined at each point x by $v \in \mathcal{H}_x \longmapsto g_x(\mathbf{q}_o, v)$. Note that the function $x \longmapsto g_x(\mathbf{p}_o, \mathbf{q}_o)$ is a nowhere vanishing function (otherwise the metric would be no longer non-degenerate). These observations motivate the following

Definition 2.6. The *canonical datum* for a compatible metric g is a quadruple $(\sigma, \alpha, \beta, \mathfrak{d})$ given by three smooth real functions σ , α , β and a tensor field $\mathfrak{d} \in \mathcal{H}^*$, with σ , α nowhere vanishing. Given a canonical datum, there is a uniquely associated compatible metric g, namely the Lorentzian metric defined by

$$g|_{W \times W} = \sigma h_o$$
, $g(\mathbf{q}_o, \mathbf{p}_o) = \frac{\sigma \alpha}{2}$, $g(\mathbf{q}_o, \mathbf{q}_o) = \frac{\sigma \beta}{2}$, $g(\mathbf{q}_o, \cdot)|_{\mathcal{H}} = \frac{\sigma \mathfrak{d}}{2}$. (2.5)

Identifying $\mathfrak{d} \in \mathfrak{H}^*$ with the 1-form of M that coincides with \mathfrak{d} on \mathfrak{H}^* and vanishes identically on the complementary spaces $\langle \mathbf{p}_o|_x, \mathbf{q}_o|_x \rangle$, and denoting by ϑ and \mathbf{p}_o^* the 1-forms defined by

$$\vartheta_x(\mathbf{q}_o|_x) = 1$$
, $\vartheta_x|_{\mathcal{W}_x} = 0$, $\mathbf{p}_{ox}^*(\mathbf{p}_o|_x) = 1$, $\mathbf{p}_{ox}^*|_{\mathcal{H}_x + \mathbb{R}\mathbf{q}_o|_x} = 0$ for any $x \in M$, (2.6)

the compatible Lorentzian metric corresponding to $(\sigma, \alpha, \beta, \mathfrak{o})$ is

$$g = \sigma \breve{\pi}^* g_o + \vartheta \vee \left(\sigma \alpha \mathbf{p}_o^* + \sigma \mathfrak{d} + \frac{\sigma \beta}{2} \vartheta \right) = \sigma \left(\breve{\pi}^* g_o + \vartheta \vee \left(\alpha \mathbf{p}_o^* + \mathfrak{d} + \frac{\beta}{2} \vartheta \right) \right) .$$
(2.7)

Note that any compatible metric has this form.

Now, for any (local) vector field X on the Kähler manifold (N, J) there is a unique vector field in the distribution \mathcal{H} on $M = \mathcal{S} \times \mathbb{R}$, determined by the spaces $\mathcal{H}_y = \mathcal{H}_{(y,u)}$ of the decompositions (2.4), projecting onto X. We denote such a vector field by \hat{X} and call it the lifted vector field corresponding to X.

The following lemma shows how to locally express each lift X in terms of the corresponding vector field X on N and a potential for the Kähler metric g_o (we recall that, if $\omega_o = g_o(J, \cdot)$ is the Kähler form of g and d^c is the differential operator on functions defined by $d^c f = -df \circ J$, a potential is a (local) function φ such that $\omega_o = dd^c \varphi$).

Lemma 2.7. There exists

(i) an open cover $\{\mathcal{V}_{\mathcal{A}}\}$ of the Kähler manifold (N, J) of the Kerr structure \mathcal{M} ,

(ii) an associated family of trivialisations $\check{\pi}^{-1}(\mathcal{V}_{\mathcal{A}}) \simeq \mathcal{V}_{\mathcal{A}} \times \mathbb{R} \times \mathbb{R}$ of $\check{\pi}: M \to N$,

(iii) a corresponding family of Kähler potentials $\{\varphi_A : \mathcal{V}_A \to \mathbb{R}\}$ on (N, J, g_o) ,

such that for any vector field X on a set \mathcal{V}_A , the corresponding lifted vector field \hat{X} on $\check{\pi}^{-1}(\mathcal{V}) \simeq \mathcal{V} \times \mathbb{R} \times \mathbb{R}$ is given by

$$\hat{X} = X - d^c \varphi(X) q_o = X + d\varphi(JX) q_o .$$
(2.8)

Proof. We recall that the local sections $f: \mathcal{V} \to \check{\pi}^{-1}(\mathcal{V})$ of the \mathbb{R}^2 -bundle $\check{\pi}: M \to N$ are in one-to-one correspondence with the trivialisations $\xi: \check{\pi}^{-1}(\mathcal{V}) \to \mathcal{V} \times \mathbb{R} \times \mathbb{R}$ of M. It is the correspondence that associates f with the unique $(\mathbb{R} \times \mathbb{R})$ -equivariant map $\xi^{(f)}: \check{\pi}^{-1}(\mathcal{V}) \to \mathcal{V} \times \mathbb{R} \times \mathbb{R}$ satisfying $(\xi^{(f)} \circ f)(y) = (y, 0, 0), y \in \mathcal{V}$. Since M is the Cartesian product $M = \mathcal{S} \times \mathbb{R}$, any local section f has the form

$$f(y) = (f^{\mathbb{S}}(y), f^{\mathbb{R}}(y))$$
 for a local section $f^{\mathbb{S}} : \mathcal{V} \to \pi^{-1}(\mathcal{V}) \subset \mathbb{S}$ and a map $f^{\mathbb{R}} : \mathcal{V} \to \mathbb{R}$.

We consider only the local sections with $f^{\mathbb{R}} = 0$. For any such section, the corresponding trivialisation identifies any point $x \in M$ with a triple $(y, \rho, u = 0)$ in $(\mathcal{V} \times \mathbb{R}) \times \mathbb{R}$ and any tangent space $T_x M$ with the direct sum $T_x M \simeq T_y \mathcal{V} + T_\rho \mathbb{R} + T_0 \mathbb{R} = T_y \mathcal{V} + \mathbb{R} + \mathbb{R}$. Under this identification, for any $x \in \check{\pi}^{-1}(\mathcal{V})$, the vectors $\mathbf{q}_o|_x$ and $\mathbf{p}_o|_x$ are represented by the unit elements $\mathbf{1}^{(I)}$ and $\mathbf{1}^{(II)}$ of the Lie algebras $\mathbb{R}^{(I)}$, $\mathbb{R}^{(II)}$ (both isomorphic to \mathbb{R}) of the two copies of the Lie group \mathbb{R} . By the definition of the distribution \mathcal{H} of $M = \mathcal{S} \times \mathbb{R}$, the lift $\hat{X} \in \mathcal{H}$ of a vector field X of \mathcal{V} is identifiable with a vector field of $\mathcal{V} \times \mathbb{R} \times \mathbb{R}$ of the form $\hat{X} = X + \kappa^f(X)\mathbf{1}^{(I)}$ where κ^f is the 1-form on \mathcal{V} (determined by the section $f = (f^{\mathcal{S}}, 0)$), which is the potential 1-form $\kappa^f := -f^{\mathcal{S}*}\theta$ on \mathcal{V} for the connection 1-form θ . In particular,

$$d\kappa^f = -\omega_o \ . \tag{2.9}$$

For any y_o in N, we may consider a simply connected neighbourhood \mathcal{V} , in which there are a trivialisation for $\check{\pi}^{-1}(\mathcal{V}) \subset M$ and a Kähler potential φ . In this neighbourhood, considering the local section $f(y) = (f^{\mathcal{S}}(y), 0)$ that determines the trivialisation and due to (2.9), we have that the corresponding κ^f satisfies $d(\kappa^f + d^c\varphi) = -\omega_o + \omega_o = 0$. Hence there exists a function $\psi : \mathcal{V} \to \mathbb{R}$ such that $\kappa^f + d^c\varphi = -d\psi$ and the modified section $f' = (f^{\mathcal{S}} + \psi, 0)$ is such that, in the corresponding trivialisation, each lifted vector field has the form $\hat{X} = X + (\kappa^f(X) + d\psi(X))q_o = X - d^c\varphi(X)q_o$, as desired.

For any trivialisable open subset $\check{\pi}^{-1}(\mathcal{V}) \simeq \mathcal{V} \times \mathbb{R} \times \mathbb{R}$, associated with an open subset \mathcal{V} of the cover $\{\mathcal{V}_A\}_{A \in \mathcal{J}}$ of the Lemma 2.7, we may consider a local frame field (E_1, \ldots, E_{n-2}) on $\mathcal{V} \subset N$ and the uniquely associated frame field $(X_A)_{A=1,\ldots,n}$ on $\check{\pi}^{-1}(\mathcal{V})$ given by

$$\begin{split} X_i &= \hat{E}_i = E_i - d^c \varphi(E_i) \mathbf{q}_o = E_i + J_i^j E_j(\varphi) \mathbf{q}_o \ , \ 1 \leqslant i \leqslant n-2 \ , \\ X_{n-1} &= \mathbf{p}_o \ , \end{split} \quad X_n = \mathbf{q}_o \ , \end{split}$$

where J_i^j denote the components of the complex structure J in the frame field (E_i) . The frame fields of this kind are called *adapted to the* \mathcal{M} -canonical optical structure (or just *adapted*, for short). The dual coframe fields of the frame field (E_i) on N and of the

frame field $(X_A) = (X_i = \hat{E}_i, X_{n-1} = \mathbf{p}_o, X_n = \mathbf{q}_o)$ on M, are denoted by (E^i) and $(X^i = \hat{E}^i, X^{n-1} = \mathbf{p}_o^*, X_n = \mathbf{q}_o^*)$, respectively.

2.4. Kerr families of gravitational fields on Kerr type backgrounds.

Definition 2.8. A background of Kerr type is a Kerr manifold M, equipped with a Kerr structure $\mathcal{M} = (M = \mathcal{S} \times \mathbb{R}, \pi : \mathcal{S} \to N, (J, g_o), \mathcal{H})$ and an Einstein metrics η , which is compatible with the \mathcal{M} -canonical optical geometry and associated with a canonical datum of the form $(\sigma, \alpha = \frac{1}{\sigma}, \beta = \frac{B}{\sigma}, \mathfrak{d} = 0)$ for a constant $B \neq 0$ and a function $\sigma \neq 0$ satisfying the constraint $\mathbb{T}(\sigma) = 0$ for $\mathbb{T} := q_o - Bp_o$. In other words, η is an Einstein metric of the form

$$\eta = \sigma \,\check{\pi}^* g_o + \vartheta \vee \left(\mathbf{p}_o^* + \frac{B}{2} \vartheta \right) \,, \qquad \text{with} \qquad \check{\pi} = \pi \circ \pi^{\mathbb{S}} : M \to N \,, \tag{2.10}$$

with $B \neq 0$ constant and $\sigma \neq 0$ nowhere vanishing function satisfying the above condition.

The reason for considering Lorentzian manifolds of this kind comes from the following proposition, according to which the 4-dimensional backgrounds of Kerr type are just flat Lorentzian manifolds admitting an optical structure of Kerr type. We recall that, according to the narration of R. P. Kerr of the origin of his celebrated metrics [18] the classical Kerr metrics were discovered studying deformations of flat metrics compatible with optical structures of such a kind.

Proposition 2.9. Let M be a 4-dimensional Kerr manifold, with Kerr manifold structure \mathcal{M} . Any background metric $\eta = \sigma \breve{\pi}^* g_o + \vartheta \lor (\mathbf{p}_o^* + \frac{B}{2}\vartheta)$ of Kerr type on M has identically vanishing curvature.

Proof. Let (N, J, g_o) be the 2-dimensional Kähler manifold, which occurs in the definition of the Kerr manifold structure of M and $\mathcal{V} \subset N$ one of the open sets of the cover $\{\mathcal{V}_A\}_{A \in \mathcal{J}}$, with associated trivialisations $\check{\pi}^{-1}(\mathcal{V}_A) \simeq \mathcal{V}_A \times \mathbb{R} \times \mathbb{R}$ and Kähler potentials $\varphi_A : \mathcal{V} \to \mathbb{R}$, as in Lemma 2.7. If (v, u) are coordinates on the standard fiber $\mathbb{R} \times \mathbb{R}$ of $\check{\pi}^{-1}(\mathcal{V})$, the 1-forms p_o^* and ϑ have the form

$$\mathbf{p}_o^* = du , \qquad \vartheta = dv + d^c \varphi ,$$

Consider the modified fiber coordinates

$$R = \frac{1}{\sqrt{2|B|}} u + \text{sign}(B) \sqrt{\frac{|B|}{2}} v , \quad T = -\frac{1}{\sqrt{2|B|}} u .$$
 (2.11)

In these coordinates, we have that $\mathbb{T} = \operatorname{sign}(B)\sqrt{\frac{|B|}{2}} \frac{\partial}{\partial T}$ (so that, since $\mathbb{T}(\sigma) = 0$, the function σ is independent of T) and

$$p_o^* = du = -\sqrt{2|B|}dT , \qquad \vartheta = dv + d^c \varphi = \operatorname{sign}(B)\sqrt{\frac{2}{|B|}}dR + \operatorname{sign}(B)\sqrt{\frac{2}{|B|}}dT + d^c \varphi ,$$

$$\vartheta \vee (p_o^* + \frac{B}{2} \vartheta) =$$

$$= \frac{B}{2} \left(\sqrt{\frac{2}{|B|}}dR + \operatorname{sign}(B)d^c \varphi\right) \vee \left(\sqrt{\frac{2}{|B|}}dR + \operatorname{sign}(B)d^c \varphi\right) - \operatorname{sign}(B)dT \vee dT .$$

(2.12)

We therefore see that

$$\eta = \sigma \left((\pi^{\delta} \circ \pi)^* g_o \right) + \vartheta \vee \left(p_o^* + \frac{B}{2} \vartheta \right) =$$

$$= \operatorname{sign}(B) \left\{ \left(\operatorname{sign}(B) \sigma \left((\pi^{\delta} \circ \pi)^* g_o \right) + \frac{|B|}{2} (\sqrt{\frac{2}{|B|}} dR + \operatorname{sign}(B) d^c \varphi) \vee (\sqrt{\frac{2}{|B|}} dR + \operatorname{sign}(B) d^c \varphi) \right) - dT \vee dT \right\}.$$

$$(2.13)$$

This expression for η and the fact that σ is *T*-independent show that the 4-dimensional Lorentzian manifold $(\check{\pi}^{-1}(\mathcal{V}), \operatorname{sign}(B)\eta)$ is locally isometric to the Cartesian product of the 1-dimensional Minkowski space $\mathbb{R}^{0,1} = (\mathbb{R}, -dT \vee dT)$ and a totally geodesic 3-dimensional submanifold (which is either Riemannian or Lorentzian, depending on whether $B\sigma > 0$ or $B\sigma < 0$). Let us denote such a totally geodesic submanifold by $(\check{\mathfrak{S}}, g^{\check{\mathfrak{S}}})$. Being

$$(\check{\pi}^{-1}(\mathcal{V}), \operatorname{sign}(B)\eta) = (\check{\mathfrak{S}} \times \mathbb{R}, g^{\check{\mathfrak{S}}} + (-dT \lor dT))$$
(2.14)

a Cartesian product of two geodesic submanifolds, one of which is 1-dimensional and thus flat, the Ricci tensor of this 4-dimensional metric satisfies an Einstein equation $\operatorname{Ric}^{(\operatorname{sign}(B)\eta)} = \Lambda \operatorname{sign}(B)\eta$ for some constant Λ if and only if

- (a) the Einstein constant Λ is 0 (because $\operatorname{Ric}^{(\operatorname{sign}(B)\eta)}\left(\frac{\partial}{\partial T}, \frac{\partial}{\partial T}\right) = 0$ since $(\mathbb{R}, -dT \vee dT)$ is totally geodesic and flat) and
- (b) the Ricci tensor $\operatorname{Ric}^{(g^{\check{S}})}$ of $(\check{S}, g^{\check{S}})$ is identically vanishing (this is because \check{S} is a totally geodesic submanifold and the Einstein constant is $\Lambda = 0$, by (a)).

By a classical fact on 3-dimensional pseudo-Riemannian manifolds (see e.g. [6, Prop. 1.120]), the Riemann curvature tensor $R^{(g^{\check{S}})}$ of $(\check{S}, g^{\check{S}})$ is uniquely determined by its Ricci tensor $\operatorname{Ric}^{(g^{\check{S}})}$ and the latter vanishes if and only if $R^{(g^{\check{S}})} = 0$. This means that $(\check{S}, g^{\check{S}})$ is flat and the curvature of (2.14) vanishes identically. This proves the claim.

We are now ready to introduce the class of Lorentzian metrics, on which we focus and that can be considered as natural generalisations of the classical Kerr metrics.

Definition 2.10. Let (M, \mathcal{M}, η) be a background of Kerr type, hence with η of the form (2.10). A Kerr family of gravitational fields on (M, \mathcal{M}, η) (or in any of its connected open subsets $M' = \mathcal{S} \times I$, $I \subset \mathbb{R}$) is a one-parameter family $g^{(m)}$ of compatible Einstein Lorentzian metrics on M, which is determined by canonical data $(\sigma^{(m)} := \sigma, \alpha^{(m)} := \frac{1}{\sigma}, \beta^{(m)} := \frac{B + m \tilde{\beta}_o}{\sigma}, \mathfrak{d}^{(m)} := 0)$, i.e. of metrics of the form

$$g^{(m)} = \sigma \breve{\pi}^* g_o + \vartheta \lor \left(\mathbf{p}_o^* + \frac{B + m\widetilde{\beta}_o}{2} \vartheta \right) = \eta + \frac{m\widetilde{\beta}_o}{2} \vartheta \lor \vartheta , \qquad (2.15)$$

where $\tilde{\beta}_o$ is an *m*-independent function satisfying the condition $\mathbb{T}(\tilde{\beta}_o) = 0$.

From (2.15), we see that a Kerr family $g^{(m)}$ is a one-parameter deformation, linearly depending on m, of the background metric η :

$$g^{(m)} = \eta + \delta g^{(m)}$$
, with $\delta g^{(m)} := m \frac{\beta_o}{2} \vartheta \vee \vartheta$. (2.16)

Motivated by a well-known terminology for the classical Kerr metrics, we call the decomposition $g^{(m)} = \eta + \frac{m\tilde{\beta}_o}{2}\vartheta \lor \vartheta$ the Kerr-Schild presentation of the family $g^{(m)}$.

Remark 2.11. According to (2.13), a 4-dimensional background metric η locally decomposes into $\eta = -(\operatorname{sign}(B)dT \vee dT) + \operatorname{sign}(B)g^{\check{\delta}}$, where $\operatorname{sign}(B)g^{\check{\delta}}$ is the induced metric of any totally geodesic hypersurface $\check{\delta} = \{T = \operatorname{const.}\}$. This decomposition shows that the vector field $\mathbb{T} = \operatorname{sign}(B)\sqrt{\frac{|B|}{2}}\frac{\partial}{\partial T}$ has a constant squared η -norm $-\frac{B}{2}$ and that its nature changes from time-like to space-like according to whether $\operatorname{sign}(B)g^{\check{\delta}}$ is (positive/negative) defined or is Lorentzian. In the cases in which $B \cdot \sigma > 0$, the metric $\operatorname{sign}(B)g^{\check{\delta}}$ is either positive or negative defined and in both cases \mathbb{T} is time-like. In the other cases, $g^{\check{\delta}}$ has a Lorentzian type signature and \mathbb{T} is space-like. Thus, in coordinates $(x^i)_{i=0,\dots,3}$ in which $\mathbb{T} = \frac{\partial}{\partial x^0}$, the coordinate x^0 can be considered as "time" whenever $B \cdot \sigma > 0$ and as a "space-like coordinate" otherwise. Accordingly, the constraints $\mathbb{T}(\sigma) = 0$, $\mathbb{T}(\beta_o) = 0$ for a Kerr family of metrics correspond to imposing that the metrics $g^{(m)}$ are time invariant or invariant under (certain) space-like translations. For what concerns the classical Kerr metrics, (1.3) shows that they correspond to a background $\eta = g^{(m=0)}$ in which $B \cdot \sigma = (-1)\sigma > 0$ and $\sigma < 0$, meaning that the classical Kerr metrics are "time invariant", as is well known.

3. A NO-GO THEOREM FOR THE EINSTEIN METRICS THAT ARE COMPATIBLE WITH OPTICAL GEOMETRIES OF KERR TYPE

3.1. A dimensional constraint for Kerr manifolds with Kerr type backgrounds. The purpose of this section is to prove (i) of Theorem 1.1. As we mentioned in the Introduction, that claim is actually a corollary of the following more general result. **Theorem 3.1** (Dimensional constraint). Let M be a Kerr manifold, equipped with the Kerr structure $\mathcal{M} = (M = \mathbb{S} \times \mathbb{R}, \pi : \mathbb{S} \to N, (J, g_o), \mathcal{H})$ and the corresponding \mathcal{M} -canonical optical structure Ω_o . If $n = \dim M$ is greater than 4, then there is no Ω_o -compatible Lorentzian metric g satisfying the following two conditions:

(a) It has the form

$$g = \sigma \,\check{\pi}^* g_o + \vartheta \vee \left(\mathbf{p}_o^* + \frac{\hat{\beta}}{2} \vartheta \right) \,, \qquad \text{with} \qquad \check{\pi} = \pi \circ \pi^{\delta} : M \to N \tag{3.1}$$

for some $\sigma, \widetilde{\beta}: M \to \mathbb{R}$, where σ is nowhere vanishing and such that $\mathbb{T}(\sigma) = \mathbb{T}(\widetilde{\beta}) = 0$ for the vector field $\mathbb{T} := \mathbf{q}_o - B\mathbf{p}_o$ with a constant $B \neq 0$;

(b) Its Ricci curvature satisfies

$$\operatorname{Ric}(\mathbf{p}_o, X) = 0 \qquad \text{for any vector field } X \in \mathcal{W} \;. \tag{3.2}$$

In particular, if dim M > 4, then there is no Einstein metric with the properties in (a).

The proof is given in the next subsection. As a corollary, here is the mentioned result:

Corollary 3.2 (No-go Theorem). No Kerr manifold of dimension greater than 4 admits a background metric of Kerr type.

Proof. A background metric η of Kerr type is Einstein and it satisfies the property (a) of Theorem (3.1) with $\tilde{\beta} = B$. Thus no such metric exists if dim M > 4.

3.2. Proof of the dimensional constraint. Consider the restriction of a metric (3.1)to an open subset $\mathcal{U} \subset M$ of the form $\mathcal{U} = \check{\pi}^{-1}(\mathcal{V}) \simeq \mathcal{V} \times \mathbb{R} \times \mathbb{R}$, where \mathcal{V} is an open subset of N of the open cover $\{\mathcal{V}_A\}_{A\in\mathcal{J}}$ described in Lemma 2.7. On such open set, let us consider an adapted frame field $(X_A)_{A=1,\ldots,n} = (\hat{E}_1,\ldots,\hat{E}_{n-2},\mathbf{p}_o,\mathbf{q}_o)$, with \hat{E}_i lifts of vector fields E_i on $\mathcal{V} \subset N$. The corresponding dual coframe field is denoted by $(X^A) = (X^i =$ $\hat{E}^i, X^{n-1} = p_o^*, X_n = q_o^*$. We organise the proof Theorem 3.1 for $g|_{\mathcal{U}}$ into two steps. We first translate the conditions $\operatorname{Ric}(\mathbf{p}_o,\mathbf{p}_o) = 0$ and $\operatorname{Ric}(\mathbf{p}_o,\widehat{E}_i) = 0$ (which are equivalent to (3.2)) into equations on the potential φ for g_o , on the constant B and on the function σ in (3.1). Then we prove that such equations have no solution if $\dim M > 4$.

3.2.1. Notational issues. Before going into the details of the proof, we need to fix some convenient notation.

- We denote by c^k_{ij}: V → ℝ the functions given by the relations [E_i, E_j] = c^k_{ij}E_k.
 The components of g_o, ω_o = g_o(J·, ·) and J in the frame field (E_i) on V ⊂ N, are denoted by

$$g_{ij} := g(E_i, E_j) , \quad \omega_{ij} := \omega_o(E_i, E_j) = g_o(JE_i, E_j) , \quad J_i^j = E^j(J(E_i)) = g^{jk}\omega_{ik} .$$

• Whenever a tensor component refers to the vector fields $X_{n-1} = p_o$ or $X_n = q_o$ or to the corresponding 1-forms $X^{n-1} = p_o^*$ and $X^n = q_o^*$, we mark it with indices " p_o " or " q_o " instead of n-1 or n. This is to make formulas more readable (for instance, the component $\operatorname{Ric}(X_{n-1}, X_n)$ of the Ricci tensor is denoted by $\operatorname{Ric}_{p_0 q_0}$ instead of $\operatorname{Ric}_{n-1 n}$).

- We denote by Γ_{ij}^m , $1 \leq i, j, m \leq n-2$, the Christoffel symbols of the Levi-Civita connection ∇^{g_o} of (\mathcal{V}, g_o) with respect to the frame field (E_i) and we denote by Γ_{AB}^{C} , $1 \leq A, B, C \leq n$, the Christoffel symbols of the Levi-Civita connection ∇^g of $(\mathcal{U} = \check{\pi}^{-1}(\mathcal{V}), g)$ with respect to the adapted frame field (X_A) .
- \mathbf{R}_{ABC}^{D} are the curvature components of $(\mathcal{U} = \check{\pi}^{-1}(\mathcal{V}), g)$ in the frame field (X_A) (⁵).
- v, u are coordinates for the fiber $\mathbb{R} \times \mathbb{R}$ of $\mathcal{U} = \check{\pi}^{-1}(\mathcal{V})$, in which \mathbf{p}_o and \mathbf{q}_o have the form

$$\mathbf{p}_o = \frac{\partial}{\partial u} , \qquad \mathbf{q}_o = \frac{\partial}{\partial v} .$$
 (3.3)

• Given a real function $f: \mathcal{V} \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$, we denote $f_{\ell} := E_{\ell}(f), 1 \leq \ell \leq n-2$. In this way, by (2.8), the vector fields \hat{E}_i take the form

$$\widehat{E}_i = E_i - d^c \varphi(E_i) \mathbf{q}_o = E_i + J_i^j E_j(\varphi) \mathbf{q}_o = E_i + J_i^j \varphi_j \mathbf{q}_o .$$
(3.4)

$$\mathbf{R}_{A_oB_oC}^{\ \ D} = X_{A_o}(\mathbf{\Gamma}_{B_oC}^{\ \ D}) - X_{B_o}(\mathbf{\Gamma}_{A_oC}^{\ \ D}) - \mathbf{\Gamma}_{A_oC}^{\ \ F}\mathbf{\Gamma}_{B_oF}^{\ \ D} + \mathbf{\Gamma}_{B_oC}^{\ \ F}\mathbf{\Gamma}_{A_oF}^{\ \ D}.$$
(3.5)

Since

$$[\widehat{E}_i, \widehat{E}_j] = [\widehat{E_i, E_j}] - \omega_o(E_i, E_j)\mathbf{q}_o = c_{ij}^k \widehat{E}_k - \omega_{ij}\mathbf{q}_o , \qquad [\widehat{E}_i, \mathbf{p}_o] = [\widehat{E}_i, \mathbf{q}_o] = [\mathbf{p}_o, \mathbf{q}_o] = 0 ,$$

the formula (3.5) holds unless $1 \leq A_o = i, B_o = j \leq n-2$. For such pairs, instead of (3.5), one has to use

$$\mathbf{R}_{ijC}^{\ \ D} = \hat{E}_i(\boldsymbol{\Gamma}_{jC}^{\ \ D}) - \hat{E}_j(\boldsymbol{\Gamma}_{iC}^{\ \ D}) - \boldsymbol{\Gamma}_{iC}^{\ F}\boldsymbol{\Gamma}_{jF}^{\ \ D} + \boldsymbol{\Gamma}_{jC}^{\ F}\boldsymbol{\Gamma}_{iF}^{\ \ D} - c_{ij}^k \boldsymbol{\Gamma}_{kC}^{\ \ D} + \omega_{ij} \boldsymbol{\Gamma}_{\mathbf{q}_oC}^{\ \ D} .$$
(3.6)

3.2.2. The Christoffel symbols of the metric (3.1). In [4] (see also [3, Appendix]), the complete list of the Christoffel symbols of the Levi-Civita connection of a compatible metric of a manifold of Kähler-Sasaki type is given. Since any metric (3.1) is of such kind, we may use the results of that paper to determine the Christoffel symbols $\Gamma_{AB}^{\ \ C}$. Actually, since (3.1) is a compatible metric of a particularly special form (in fact, its associated canonical datum has the form $(\sigma, \alpha = \frac{1}{\sigma}, \beta = \frac{\tilde{\beta}}{\sigma}, \mathfrak{d} = 0)$ with σ and $\tilde{\beta}$ such that $\mathbb{T}(\sigma) = \mathbb{T}(\tilde{\beta}) = 0$)

⁵Following [19], we assume that R is defined by $R_{XY}Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z$.

the list in [4, Prop. 3.1] radically simplifies and we get:

$$\Gamma_{ij}^{m} = g^{mk}g_{o}(\nabla_{E_{i}}^{o}E_{j},E_{k}) + \frac{1}{2\sigma}\hat{E}_{i}(\sigma)\delta_{j}^{m} + \frac{1}{2\sigma}\hat{E}_{j}(\sigma)\delta_{i}^{m} - \frac{1}{2\sigma}g_{ij}g^{mk}\hat{E}_{k}(\sigma) =$$

$$= \Gamma_{ij}^{m} + \frac{1}{2\sigma}\sigma_{i}\delta_{j}^{m} + \frac{1}{2\sigma}\sigma_{j}\delta_{i}^{m} - \frac{1}{2\sigma}g_{ij}g^{mk}\sigma_{k} +$$

$$+ B\frac{p_{o}(\sigma)\varphi_{\ell}}{2\sigma}\left(J_{i}^{\ell}\delta_{j}^{m} + J_{j}^{\ell}\delta_{i}^{m} - g_{ij}g^{mk}J_{k}^{\ell}\right) , \qquad (3.7)$$

$$\Gamma_{ij}^{\mathbf{p}_o} = -g_{ij} \left(B \mathbf{p}_o(\sigma) - \widetilde{\beta} \mathbf{p}_o(\sigma) \right) , \quad \Gamma_{ij}^{\mathbf{q}_o} = -\frac{\omega_{ij}}{2} - g_{ij} \mathbf{p}_o(\sigma) , \qquad (3.8)$$

$$\Gamma_{ip_o}^{\ m} = \Gamma_{p_o i}^{\ m} = \frac{g^{mk}\omega_{ik}}{4\sigma} + \frac{\delta_i^m}{2\sigma}p_o(\sigma) , \qquad \Gamma_{ip_o}^{\ p_o} = \Gamma_{p_o i}^{\ p_o} = \Gamma_{p_o i}^{\ q_o} = \Gamma_{p_o i}^{\ q_o} = 0 , \qquad (3.9)$$

$$\Gamma_{i\mathbf{q}_{o}}^{m} = \Gamma_{\mathbf{q}_{o}i}^{m} = \frac{g^{mk}\omega_{ik}(\widetilde{\beta}-B)}{4\sigma} + \left(B\frac{g^{mk}\omega_{ik}}{4\sigma} + B\frac{\delta_{i}^{m}}{2\sigma}\mathbf{p}_{o}(\sigma)\right) = \frac{J_{i}^{m}}{4\sigma}(\widetilde{\beta}-B) + B\Gamma_{i\mathbf{p}_{o}}^{m},$$

$$= \frac{J_{i}^{m}}{4\sigma}(\widetilde{\beta}-B) = \frac{J_{i}^{m}}{4\sigma}$$

$$\Gamma_{i\mathbf{q}_{o}}^{\mathbf{p}_{o}} = \Gamma_{\mathbf{q}_{o}i}^{\mathbf{p}_{o}} = \frac{1}{2}\widetilde{\beta}_{i} - \frac{1}{2}J_{i}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\widetilde{\beta}), \quad \Gamma_{i\mathbf{q}_{o}}^{\mathbf{q}_{o}} = \Gamma_{\mathbf{q}_{o}i}^{\mathbf{q}_{o}} = 0, \quad (3.10)$$

$$\Gamma_{\mathbf{p}_o\mathbf{p}_o}^{\ m} = \Gamma_{\mathbf{p}_o\mathbf{p}_o}^{\ p_o} = \Gamma_{\mathbf{p}_o\mathbf{p}_o}^{\ q_o} = 0 , \qquad (3.11)$$

$$\Gamma_{\mathbf{p}_{o}\mathbf{q}_{o}}^{m} = \Gamma_{\mathbf{q}_{o}\mathbf{p}_{o}}^{m} = 0 , \qquad \Gamma_{\mathbf{p}_{o}\mathbf{q}_{o}}^{p} = \Gamma_{\mathbf{q}_{o}\mathbf{p}_{o}}^{p} = \frac{1}{2}\mathbf{p}_{o}(\widetilde{\beta}) , \qquad \Gamma_{\mathbf{p}_{o}\mathbf{q}_{o}}^{q} = \Gamma_{\mathbf{q}_{o}\mathbf{p}_{o}}^{q} = 0 , \qquad (3.12)$$

$$\Gamma_{\mathbf{q}_{o}\mathbf{q}_{o}}^{m} = -\frac{g^{mk}}{4\sigma}\widehat{E}_{k}(\widetilde{\beta}) = -\frac{g^{mk}}{4\sigma}\widetilde{\beta}_{k} - B\frac{g^{mk}}{4\sigma}J_{k}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\widetilde{\beta}) ,$$

$$\Gamma_{\mathbf{q}_{o}\mathbf{q}_{o}}^{p} = \frac{\mathbf{q}_{o}(\widetilde{\beta})}{2} + \frac{\mathbf{p}_{o}(\widetilde{\beta})\widetilde{\beta}}{2} , \ \Gamma_{\mathbf{q}_{o}\mathbf{q}_{o}}^{q} = -\frac{1}{2}\mathbf{p}_{o}(\widetilde{\beta}) .$$
(3.13)

Combing these formulas with (3.5) and (3.6), we are ready to translate the conditions $\operatorname{Ric}(\mathbf{p}_o,\mathbf{p}_o) = \operatorname{Ric}(\mathbf{p}_o,\hat{E}_i) = 0$ into equations for σ , B and φ .

3.2.3. The condition $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$ as an equation on σ . We recall that $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = \operatorname{Ric}_{\mathbf{p}_o\mathbf{p}_o} = \operatorname{R}_{m\mathbf{p}_o\mathbf{p}_o}^m + \operatorname{R}_{\mathbf{q}_o\mathbf{p}_o\mathbf{p}_o}$. Thus, writing the curvature components in terms of the Christoffel symbols (3.7) – (3.13) and neglecting all trivially vanishing terms, we get

$$\operatorname{Ric}(\mathbf{p}_{o},\mathbf{p}_{o}) = -\operatorname{p}_{o}(\mathbf{\Gamma}_{m\mathbf{p}_{o}}^{m}) - \mathbf{\Gamma}_{m\mathbf{p}_{o}}^{\ell}\mathbf{\Gamma}_{\mathbf{p}_{o}\ell}^{m} = = -\frac{n-2}{2}\operatorname{p}_{o}\left(\frac{\operatorname{p}_{o}(\sigma)}{\sigma}\right) - \left(\frac{J_{m}^{\ell}}{4\sigma} + \frac{\operatorname{p}_{o}(\sigma)\delta_{m}^{\ell}}{2\sigma}\right)\left(\frac{J_{\ell}^{m}}{4\sigma} + \frac{\operatorname{p}_{o}(\sigma)\delta_{\ell}^{m}}{2\sigma}\right) = = \frac{n-2}{4\sigma^{2}}\left(-2\sigma\operatorname{p}_{o}(\operatorname{p}_{o}(\sigma)) + \operatorname{p}_{o}(\sigma)^{2} + \frac{1}{4}\right) .$$

$$(3.14)$$

Since $n \ge 4$, the condition $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$ is equivalent to the equation

$$-2\sigma p_o(p_o(\sigma)) + (p_o(\sigma))^2 + \frac{1}{4} = 0.$$
(3.15)

Differentiating along p_o yields that any solution of (3.15) satisfies also the equation

$$\mathbf{p}_{o}\left(\mathbf{p}_{o}\left(\mathbf{p}_{o}(\sigma)\right)\right) = \frac{\partial^{3}\sigma}{\partial u^{3}} = 0$$

Together with $0 = \mathbb{T}(\sigma) = -B\frac{\partial\sigma}{\partial u} + \frac{\partial\sigma}{\partial v}$, we get that σ has necessarily the form

$$\sigma = C_0 + C_1(u + Bv) + C_2(u + Bv)^2$$
(3.16)

for some smooth functions $C_{\alpha} : \mathcal{V} \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}, \alpha = 0, 1, 2$, which are independent of the fiber coordinates u and v (hence, identifiable with functions $C_{\alpha} : \mathcal{V} \subset N \to \mathbb{R}$). Plugging (3.16) into (3.15), we see that the C_{α} satisfy the algebraic equation

$$(C_1 + 2C_2(u + Bv))^2 - 4C_2 \left(C_0 + C_1(u + Bv) + C_2(u + Bv)^2\right) + \frac{1}{4} = C_1^2 - 4C_0C_2 + \frac{1}{4} = 0. \quad (3.17)$$

In particular, $C_2, C_0 : \mathcal{V} \to \mathbb{R}$ are nowhere vanishing. If for each given solution σ to (3.15), determined by the functions $C_{\alpha} : \mathcal{V} \to \mathbb{R}$, we introduce the associated function $r : \mathcal{V} \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ defined by r(y, v, u) := u + Bv + D(y) with $D(y) = \frac{C_1(y)}{2C_2(y)}$, we conclude that σ takes the form

$$\sigma = C_0 + C_1 r - \frac{C_1^2}{2C_2} + C_2 r^2 - \frac{C_1 C_2}{C_2} r + \frac{C_1^2 C_2}{4C_2^2} \stackrel{(3.17)}{=} C_2 r^2 + \frac{1}{16C_2} .$$
(3.18)

3.2.4. The conditions $\operatorname{Ric}(\widehat{E}_i, \mathbf{p}_o) = 0$ as equations on φ , B and σ . Utilising the Ricci tensor definition and expressing curvature components with Christoffel symbols (and removing any term that is manifestly zero according to (3.7)–(3.13)) we get

$$\operatorname{Ric}(\widehat{E}_{i}, \mathbf{p}_{o}) = \operatorname{Ric}_{i\mathbf{p}_{o}} = \operatorname{R}_{mi\mathbf{p}_{o}}^{m} + \operatorname{R}_{\mathbf{p}_{o}i\mathbf{p}_{o}}^{\mathbf{p}_{o}} + \operatorname{R}_{\mathbf{q}_{o}i\mathbf{p}_{o}}^{\mathbf{q}_{o}} = \\ = \widehat{E}_{m}(\mathbf{\Gamma}_{i\mathbf{p}_{o}}^{m}) - \widehat{E}_{i}(\mathbf{\Gamma}_{m\mathbf{p}_{o}}^{m}) - \mathbf{\Gamma}_{m\mathbf{p}_{o}}^{\ell}\mathbf{\Gamma}_{i\ell}^{m} + \mathbf{\Gamma}_{i\mathbf{p}_{o}}^{\ell}\mathbf{\Gamma}_{m\ell}^{m} - c_{mi}^{r}\mathbf{\Gamma}_{r\mathbf{p}_{o}}^{m} .$$
(3.19)

We may now plug in (3.7) - (3.13) into (3.19) and simplify the expression using the fact that $\nabla^{g_o} J = 0$ (because (N, J, g_o) is Kähler) and that ∇ is torsion free. After a few straightforward (but quite tedious) computations, we get

$$\operatorname{Ric}(\widehat{E}_{i}, \mathbf{p}_{o}) = \frac{n-3}{8\sigma^{2}} \left(\frac{n-6}{n-3} J_{i}^{\ell} \sigma_{\ell} - 4\sigma E_{i} \left(\mathbf{p}_{o}(\sigma) \right) + 4\mathbf{p}_{o}(\sigma)\sigma_{i} - \frac{n-6}{n-3} B\varphi_{i} \mathbf{p}_{o} \left(\sigma \right) - 4\sigma B J_{i}^{\ell} \varphi_{\ell} \mathbf{p}_{o} \left(\mathbf{p}_{o}(\sigma) \right) + 4B J_{i}^{\ell} \varphi_{\ell} \left(\mathbf{p}_{o}(\sigma) \right)^{2} \right).$$
(3.20)

The reader is referred to Appendix A for the details of those computations. Having $n \ge 4$,

it follows that each equation $\operatorname{Ric}(\hat{E}_i, \mathbf{p}_o) = 0$ is equivalent to

$$\frac{n-6}{n-3}J_i^{\ell}\sigma_{\ell} - 4\sigma E_i\left(\mathbf{p}_o(\sigma)\right) + 4\mathbf{p}_o(\sigma)\sigma_i - \frac{n-6}{n-3}B\varphi_i\mathbf{p}_o\left(\sigma\right) - 4\sigma B J_i^{\ell}\varphi_{\ell}\mathbf{p}_o\left(\mathbf{p}_o(\sigma)\right) + 4B J_i^{\ell}\varphi_{\ell}\left(\mathbf{p}_o\left(\sigma\right)\right)^2 = 0. \quad (3.21)$$

3.2.5. The proof that $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$ and $\operatorname{Ric}(\mathbf{p}_o, \widehat{E}_i) = 0$ are not compatible if n > 4. Assume that g is such that $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$, i.e. that σ has the form (3.18). Let us also adopt the short-hand notation $A := C_2$, $D = \frac{C_1}{2C_2}$, $A_i := E_i(A)$, $D_i := E_i(D)$ and $\sigma_i = E_i(\sigma)$. Since r = u + Bv + D, the following relations hold:

$$\sigma_{\ell} = E_{\ell} \left(Ar^2 + \frac{1}{16A} \right) = A_{\ell}r^2 + 2AD_{\ell}r - \frac{1}{16A^2}A_{\ell} ,$$

$$p_o(\sigma) = \frac{\partial\sigma}{\partial u} = 2Ar\frac{\partial r}{\partial u} = 2Ar , \quad p_o(p_o(\sigma)) = 2A , \quad E_i(p_o(\sigma)) = 2A_ir + 2AD_i .$$
(3.22)

If we now assume that g is also a solution to $\operatorname{Ric}(\hat{E}_i, \mathbf{p}_o) = 0, 1 \leq i \leq n-2$, plugging (3.18) and (3.22) into (3.21) we get

$$0 = \frac{n-6}{n-3} J_i^{\ell} \left(A_{\ell} r^2 + 2Ar D_{\ell} - \frac{1}{16A^2} A_{\ell} \right) -$$

$$-4\left(Ar^{2} + \frac{1}{16A}\right)\left(2A_{i}r + 2AD_{i}\right) + 8Ar\left(A_{i}r^{2} + 2ArD_{i} - \frac{1}{16A^{2}}A_{i}\right) - \frac{n-6}{n-3}B\varphi_{i}2Ar + 8\left(Ar^{2} - \frac{1}{16A}\right)BJ_{i}^{\ell}\varphi_{\ell}A + 16BJ_{i}^{\ell}\varphi_{\ell}A^{2}r^{2} = 0$$

$$= \frac{n-6}{n-3}J_{i}^{\ell}A_{\ell}r^{2} + 2\frac{n-6}{n-3}J_{i}^{\ell}AD_{\ell}r - \frac{n-6}{n-3}J_{i}^{\ell}\frac{1}{16A^{2}}A_{\ell} - \frac{1}{2}A_{i}r^{3} - 8A^{2}D_{i}r^{2} - \frac{A_{i}}{2A}r - \frac{1}{2}D_{i} + 8AA_{i}r^{3} + 16A^{2}r^{2}D_{i} - \frac{A_{i}}{2A}r - \frac{1}{2}BJ_{i}^{\ell}\varphi_{\ell}r^{2} - \frac{1}{2}BJ_{i}^{\ell}\varphi_{\ell}A^{2}r^{2} . \quad (3.23)$$

The right hand side of the last equality is a second order polynomial in the function r = r(y, v, u) (here, $y = (y^i)$ stands for coordinates on $\mathcal{V} \subset N$). By definition of such a function, for any fixed value of y, the values of r run unconstrained in an unbounded interval of \mathbb{R} . This implies that (3.23) is satisfied if and only if the coefficients of the monomials r^2 , r and 1 vanish identically, i.e. if and only if the following equations are

satisfied for any $1 \leq i \leq n-2$:

$$\frac{n-6}{n-3}J_i^{\ell}A_{\ell} + 8A^2 \left(D_i + BJ_i^{\ell}\varphi_{\ell}\right) = 0,$$

$$2A\left(\frac{n-6}{n-3}J_i^{\ell}D_{\ell} - \frac{n-6}{n-3}B\varphi_i - \frac{A_i}{2A^2}\right) = 0,$$

$$-\frac{1}{16A^2}\frac{n-6}{n-3}J_i^{\ell}A_{\ell} - \frac{1}{2}\left(D_i + BJ_i^{\ell}\varphi_{\ell}\right) = 0.$$
(3.24)

Here, the third line is equivalent to the first and we may neglect it. Moreover, by multiplying the first line by J_m^i and contracting for *i*, the system (3.24) becomes equivalent to

$$\begin{pmatrix} -1 & 8\frac{n-3}{n-6} \\ -1 & 2\frac{n-6}{n-3} \end{pmatrix} \begin{pmatrix} \frac{A_i}{A^2} \\ J_i^{\ell} D_{\ell} \end{pmatrix} = \begin{pmatrix} 8\frac{n-3}{n-6} \\ 2\frac{n-6}{n-3} \end{pmatrix} B\varphi_i .$$

$$(3.25)$$

Note that the matrix $\begin{pmatrix} -1 & \frac{n-3}{n-6} \\ -1 & \frac{2n-6}{n-3} \end{pmatrix}$ is invertible if and only if

$$8\frac{n-3}{n-6} \neq 2\frac{n-6}{n-3} \qquad \Longleftrightarrow \qquad 4(n-3)^2 \neq (n-6)^2 \qquad \Longleftrightarrow \qquad n \neq 4 \ .$$

Hence, if n > 4, for a given $1 \le i \le n-2$ the equations (3.23) are equivalent to

$$\begin{pmatrix} \frac{A_i}{A^2} \\ J_i^{\ell} D_{\ell} \end{pmatrix} = \begin{pmatrix} -1 & 8\frac{n-3}{n-6} \\ -1 & 2\frac{n-6}{n-3} \end{pmatrix}^{-1} \begin{pmatrix} 8\frac{n-3}{n-6} \\ 2\frac{n-6}{n-3} \end{pmatrix} B\varphi_i = \begin{pmatrix} 0 \\ 1 \end{pmatrix} B\varphi_i .$$
(3.26)

This means that $B\varphi_i J_\ell^i = -D_\ell$ or, equivalently, that $Bd^c\varphi = -dD$. From this it follows that $0 = -d^2D = Bdd^c\varphi = B\omega_o$. This is impossible, because ω_o is non-degenerate and $B \neq 0$. From this contradiction, Theorem 3.1 follows.

4. Classification of Kerr 4-manifolds admitting Kerr type backgrounds

4.1. The fundamental constraints on the underlying Kähler surface (N, J, g_o) . From now on we focus on the 4-dimensional case. Let M be a Kerr 4-manifold, equipped with a Kerr structure $\mathcal{M} = (M = S \times \mathbb{R}, \pi : S \to N, (J, g_o), \mathcal{H})$ and the associated \mathcal{M} -canonical optical structure $\mathcal{Q}_o = (\mathcal{W}_o, [h_o]_{\pm}, \mathcal{K}_o, \{g\})$. The following theorem provides (topological and metrical) conditions on the 2-dimensional Kähler manifold (N, J, g_o) that are necessary for M to admit a Kerr type background metric. In the subsequent section we prove that such necessary conditions are also sufficient and a detailed description of all background metrics of a Kerr manifold is given. According to Proposition 2.9, all such background metrics are locally isometric to each other (they are all flat). However, as we will shortly see, they all have different presentations as compatible metrics and correspond to different canonical data. **Theorem 4.1.** Let M be a Kerr 4-manifold with Kerr structure \mathcal{M} and \mathcal{M} -canonical optical structure \mathcal{Q}_o . The Ricci tensor of a metric η as in (a) of Theorem 3.1 is such that

$$\operatorname{Ric}(X,Y) = 0$$
 for any pair of vector fields $X, Y \in W_o$ (4.1)

if and only if the following three conditions hold:

- (1) The Riemann surface (N, J) admits a Riemannian metric \tilde{g}_o of constant Gaussian curvature $\kappa = \pm 1$, which is Kähler with respect to the complex structure J (i.e., in real terms, there exists an atlas of real coordinate charts, each of them simultaneously isothermal for g_o and \tilde{g}_o);
- (2) There exists a Kähler potential $\varphi : N \to \mathbb{R}$ for g_o (i.e. such that $\omega_o = dd^c \varphi$) satisfying

$$\operatorname{sign} \varphi = -\kappa , \qquad g_o + 2\kappa\varphi \,\widetilde{g}_o = 0 ; \qquad (4.2)$$

(3) The Riemann surface N is not compact and, up to a bundle automorphism of the trivial bundle $\pi^{\mathbb{S}}: M = \mathbb{S} \times \mathbb{R} \to \mathbb{S}$, there is a globally defined coordinate r for the fiber of $\pi^{\mathbb{S}}: M \to \mathbb{S}$ such that

$$\sigma = -\frac{r^2}{4B\varphi} - \frac{B\varphi}{4} \ . \tag{4.3}$$

The proof is divided into two steps and it is given in the following subsections. We first show that a sub-system of the equations corresponding to the (4.1) implies that σ has locally the form (4.3) for some potential φ for the metric g_o . Second, we prove that the remaining equations corresponding to (4.1) are satisfied if and only if there is an open cover $\{\mathcal{V}_A\}$ for N and a corresponding family of local potentials on the sets \mathcal{V}_A , each of them satisfying (4.2) for a metric $\tilde{g}_o|_{\mathcal{V}_A}$ with constant scalar curvature $\kappa = \pm 1$. From these facts it directly follows that: (i) The metrics $\tilde{g}_o|_{\mathcal{V}_A}$ combine into a globally defined metric of constant Gaussian curvature over N with the same isothermal coordinates of g_o ; (ii) There exists a function $\varphi: N \to \mathbb{R}$, which is well defined at each $p \in N$ by

$$\varphi(p) := -\kappa \frac{g_o(X_p, X_p)}{2\tilde{g}_o(X_p, X_p)}$$

for a freely specifiable non-trivial local vector field X on a neighbourhood of p; (iii) for any open set \mathcal{V}_A of the above open cover of N, the restriction $\varphi|_{\mathcal{V}_A}$ is a potential for $g_o|_{\mathcal{V}_A}$ and, consequently, the function φ is a potential for g_o , which is globally defined over N. From (i) –(iii), the claims (1) – (2) follow. Moreover:

- The Kähler manifold (N, J, g_o) is not compact (because it admits a global potential);
- Since φ and σ are globally defined on M, the relation (4.3) (which was originally shown to be valid just in appropriate coordinates sets) defines a global real function $r: M \to \mathbb{R}$, that provides a global coordinate for the fiber of $\pi^{\mathbb{S}}: M = \mathbb{S} \times \mathbb{R} \to \mathbb{S}$.

From these two observations, (3) follows.

We also stress the following two facts:

(a) If $\mathcal{V} \subset N$ is an open set with coordinates (x, y, v, r) on $\check{\pi}^{-1}(\mathcal{V}) = \mathcal{V} \times \mathbb{R} \times \mathbb{R}$ with (x, y) selected in such a way that the coordinate components of \tilde{g}_o have the classical

expressions $\tilde{g}_{o\,ij} = \frac{4\delta_{ij}}{(1+\kappa(x^2+y^2))^2}$, then (2) holds if and only if $\varphi|_{\mathcal{V}}$ satisfies one of the following pairs of conditions:

$$\Delta \varphi + \frac{8\varphi}{\left(1 + x^2 + y^2\right)^2} = 0 \qquad \& \qquad \varphi < 0 \qquad \text{in case } \kappa = 1 \quad \text{or}$$

$$\Delta \varphi - \frac{8\varphi}{\left(1 - x^2 - y^2\right)^2} = 0 \qquad \& \qquad \varphi > 0 \qquad \text{in case } \kappa = -1 .$$

$$(4.4)$$

(b) For both possibilities for κ , we have that $\operatorname{sign}(\sigma) = \operatorname{sign}(B\kappa)$ and the signature of g is mostly plus or mostly minus, according to whether $\operatorname{sign}(B\kappa)$ is equal to -1 or +1.

These facts will greatly assist in the discussions of the following sections.

4.1.1. Step 1 of the Proof of Theorem 4.1. Consider an open cover $\{\mathcal{V}_A\}_{A\in\mathcal{J}}$ of N as in Lemma 2.7, and pick an open set \mathcal{V} in this cover and a (local) potential $\varphi: \mathcal{V} \to \mathbb{R}$ for g_o . We adopt the notational conventions of §3.2.1. With no loss of generality, we also assume that $\mathcal{V} \subset N$ admits isothermal coordinates (x, y) for g_o and that $(E_1 := \frac{\partial}{\partial x}, E_2 := \frac{\partial}{\partial y})$. In this way the components J_i^j , ω_{ij} and g_{ij} of the complex structure, the Kähler form and the Kähler metric have the form

$$(J_i^j) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

$$(\omega_{ij}) = (dd^c \varphi(E_i, E_j)) = (-E_i (J_j^\ell \varphi_\ell) + E_j (J_i^\ell \varphi_\ell)) = \begin{pmatrix} 0 & \lambda \\ -\lambda & 0 \end{pmatrix},$$

$$(g_{ij}) = (g(E_i, E_j)) = (\omega_{im} J_j^m) = \lambda \, \delta_{ij}, \quad \text{where we set } \lambda := \Delta \varphi.$$
(4.5)

Finally, as usual, we denote by $(X_A) = (\hat{E}_1, \hat{E}_2, \mathbf{p}_o, \mathbf{q}_o)$ the adapted frame field on $\check{\pi}^{-1}(\mathcal{V}) \simeq \mathcal{V} \times \mathbb{R}^2$, corresponding to the frame field (E_1, E_2) .

By the results of §3.2.3, the condition $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$ (which is a consequence of (4.1)) is satisfied on the open set $\check{\pi}^{-1}(\mathcal{V})$ if and only if $\sigma = \sigma(x, y, v, u)$ has the form $\sigma = Cr^2 + \frac{1}{16C}$ for a nowhere vanishing function $C : \mathcal{V} \to \mathbb{R}$ and r := u + Bv + D(x, y), where B is the constant appearing in the Definition 2.8 and D = D(x, y) is an arbitrary function. Replacing the fibre coordinate u by r = u + Bv + D(x, y) yields new coordinates (x, y, v, r)on $\check{\pi}^{-1}(\mathcal{V})$, in which the vector fields \mathbf{p}_o , \mathbf{q}_o take the form $\mathbf{p}_o = \frac{\partial}{\partial r}$, $\mathbf{q}_o = \frac{\partial}{\partial v} + B\frac{\partial}{\partial r}$ and $\sigma = Cr^2 + \frac{1}{16C}$ becomes a function of the new (independent) variables (x, y, v, r).

The computations in §3.2.5 (which hold in *any* dimension) show that $\operatorname{Ric}(\mathbf{p}_o, \hat{E}_i) = 0$, i = 1, 2, hold true if and only if the system (3.24) is satisfied. Since we are now in the case n = 4 and we are in a coordinates system, in which σ has no summand that is linear with respect to the coordinate r (meaning that we may assume $D = \frac{C_1}{2C_2}$ to be zero), the first two equations of (3.24) are equivalent each other and both equivalent to

$$J_i^{\ell}\left(\frac{C_{\ell}}{C^2} - 4B\varphi_{\ell}\right) = 0 \quad \left(\iff d\left(\frac{1}{C} + 4B\varphi\right) = 0 \iff \frac{1}{C} = -4B\varphi + \text{const.} \right)$$
(4.6)

Since the potentials are determined up to constants, we may assume $\frac{1}{C} = -4B\varphi$ and conclude that $\operatorname{Ric}(\mathbf{p}_o, X) = 0$ for any $X \in \mathcal{W}_o$ if and only if σ has the form (4.3) for an appropriate Kähler potential φ on \mathcal{V} .

4.1.2. Step 2 of the Proof of Theorem 4.1. Let us now assume that σ has the form (4.3) and hence that $\operatorname{Ric}(\mathbf{p}_o, X) = 0$ for any $X \in \mathcal{W}_o$. Then (4.1) holds if and only if $\operatorname{Ric}_{ij} = \operatorname{Ric}(\hat{E}_i, \hat{E}_j) = 0$ for any $1 \leq i, j \leq 2$. Writing the curvature in terms of the Christoffel symbols (3.7) – (3.13)) (see (3.5), (3.6)) and recalling that the $E_i = \frac{\partial}{\partial x^i}$ commute, we get the following (here, as in the previous section, $\lambda := \Delta \varphi$):

Since $\mathbf{p}_o = \frac{\partial}{\partial r}$, $\mathbf{q}_o = \frac{\partial}{\partial v} + B \frac{\partial}{\partial r}$, and no Christoffel symbol depends on v, the differential operator \mathbf{q}_o acts on the $\mathbf{\Gamma}_{AB}^{\ \ C}$ just as $B \frac{\partial}{\partial r} = B \mathbf{p}_o$. Thus $\operatorname{Ric}_{ij} = \operatorname{Ric}(\hat{E}_i, \hat{E}_j) = 0$ reduce to

$$\begin{aligned} \operatorname{Ric}_{11} &= \left(E_m + J_m^{\ell} \varphi_{\ell} B_{\overline{\partial} r}^{\frac{\partial}{\partial r}} \right) \left(\Gamma_{11}^{\ m} \right) - \left(\frac{\partial}{\partial x} + B \varphi_y \frac{\partial}{\partial r} \right) \left(\Gamma_{m1}^{\ m} \right) - \Gamma_{m1}^{\ \ell} \Gamma_{1\ell}^{\ m} - \Gamma_{m1}^{\ q_o} \Gamma_{1q_o}^{\ m} + \\ &+ \Gamma_{11}^{\ \ell} \Gamma_{m\ell}^{\ m} + \Gamma_{11}^{\ q_o} \Gamma_{mq_o}^{\ m} - \lambda \Gamma_{q_o}^{\ 2} + B \frac{\partial}{\partial r} (\Gamma_{11}^{\ q_o}) - \Gamma_{q_o}^{\ \ell} \Gamma_{1\ell}^{\ q_o} = 0 \;, \end{aligned} \tag{4.10} \\ \operatorname{Ric}_{12} &= \left(E_m + J_m^{\ell} E_{\ell}(\varphi) B_{\overline{\partial} r}^{\frac{\partial}{\partial r}} \right) \left(\Gamma_{12}^{\ m} \right) - \left(\frac{\partial}{\partial x} + B \varphi_y \frac{\partial}{\partial r} \right) \left(\Gamma_{m2}^{\ m} \right) - \Gamma_{m2}^{\ \ell} \Gamma_{1\ell}^{\ m} - \Gamma_{m2}^{\ q_o} \Gamma_{1q_o}^{\ m} + \\ &+ \Gamma_{1\ell}^{\ \ell} \Gamma_{m\ell}^{\ m} + \Gamma_{12}^{\ q_o} \Gamma_{mq_o}^{\ m} - \lambda \Gamma_{q_o}^{\ 2} + B \frac{\partial}{\partial r} (\Gamma_{12}^{\ q_o}) - \Gamma_{q_o}^{\ \ell} \Gamma_{1\ell}^{\ q_o} = 0 \;, \end{aligned} \tag{4.11} \\ \operatorname{Ric}_{22} &= \left(E_m + J_m^{\ell} E_{\ell}(\varphi) B_{\overline{\partial} r}^{\frac{\partial}{\partial r}} \right) \left(\Gamma_{22}^{\ m} \right) - \left(\frac{\partial}{\partial y} - B \varphi_x \frac{\partial}{\partial r} \right) \left(\Gamma_{m2}^{\ m} \right) - \Gamma_{m2}^{\ \ell} \Gamma_{2\ell}^{\ m} - \Gamma_{m2}^{\ q_o} \Gamma_{2q_o}^{\ m} + \\ &+ \Gamma_{2\ell}^{\ \ell} \Gamma_{m\ell}^{\ m} + \Gamma_{22}^{\ q_o} \Gamma_{mq_o}^{\ m} + \lambda \Gamma_{q_o}^{\ 1} + B \frac{\partial}{\partial r} (\Gamma_{22}^{\ q_o}) - \Gamma_{q_o}^{\ \ell} \Gamma_{2\ell}^{\ q_o} = 0 \;. \end{aligned} \tag{4.12}$$

Now, plugging the condition $\tilde{\beta} = B$ into the expressions (3.7) – (3.13), we get that the only non-zero Christoffel symbols $\Gamma_{AB}^{\ \ C}$ of η are

$$\Gamma_{11}^{11} = \Gamma_{11}^{11} + \frac{1}{2\sigma}\sigma_x + B\frac{\sigma_t\varphi_y}{2\sigma} \qquad \Gamma_{11}^{2} = \Gamma_{11}^{2} - \frac{1}{2\sigma}\sigma_y + B\frac{\sigma_t\varphi_x}{2\sigma} \qquad (4.13)$$

$$\Gamma_{12}^{11} = \Gamma_{21}^{11} = \Gamma_{12}^{11} + \frac{1}{2\sigma}\sigma_y - B\frac{\sigma_t\varphi_x}{2\sigma} \qquad \Gamma_{12}^{2} = \Gamma_{21}^{2} = \Gamma_{12}^{2} + \frac{1}{2\sigma}\sigma_x + B\frac{\sigma_t\varphi_y}{2\sigma} \qquad (4.14)$$

$$\Gamma_{22}^{\ 1} = \Gamma_{22}^{\ 1} - \frac{1}{2\sigma}\sigma_x - B\frac{\sigma_t\varphi_y}{2\sigma} \qquad \qquad \Gamma_{22}^{\ 2} = \Gamma_{22}^{\ 2} + \frac{1}{2\sigma}\sigma_y - B\frac{\sigma_t\varphi_x}{2\sigma} \qquad (4.15)$$

$$\Gamma_{11}^{q_o} = -\lambda \sigma_t , \qquad \Gamma_{12}^{q_o} = -\frac{\lambda}{2} , \qquad \Gamma_{21}^{q_o} = \frac{\lambda}{2} , \qquad \Gamma_{22}^{q_o} = -\lambda \sigma_t , \qquad (4.16)$$

$$\Gamma_{12}^{1} = \Gamma_{r_o}^{-1} = \frac{\sigma_t}{2} , \qquad \Gamma_{12}^{2} = \Gamma_{r_o}^{-2} = \frac{1}{2} , \qquad (4.17)$$

$$\Gamma_{1p_{o}} - \Gamma_{p_{o}1} - \frac{1}{2\sigma}, \qquad \Gamma_{1p_{o}} - \Gamma_{p_{o}1} - \frac{1}{4\sigma}, \qquad (4.17)$$

$$\Gamma_{2p_{o}}^{1} = \Gamma_{p_{o}2}^{1} = -\frac{1}{4\sigma}, \qquad \Gamma_{2p_{o}}^{2} = \Gamma_{p_{o}2}^{2} = \frac{\sigma_{t}}{2\sigma}, \qquad (4.18)$$

$$\Gamma_{1q_o}^{1} = \Gamma_{q_o1}^{1} = B \frac{\sigma_t}{2\sigma}, \qquad \qquad \Gamma_{1q_o}^{2} = \Gamma_{q_o1}^{2} = B \frac{1}{4\sigma}, \qquad (4.19)$$

$$\Gamma_{2q_o}^{\ 1} = \Gamma_{q_o^2}^{\ 1} = -B \frac{1}{4\sigma} , \qquad \qquad \Gamma_{2q_o}^{\ 2} = \Gamma_{q_o^2}^{\ 2} = B \frac{\sigma_t}{2\sigma} . \qquad (4.20)$$

Using this, a sequence of tedious but straightforward computations and simplifications lead to the following

Lemma 4.2. The equation (4.11) is identically satisfied, while (4.10) and (4.12) are both equivalent to the equation

$$\sigma^{2}\operatorname{Ric}_{11}^{N} - \frac{1}{2}\sigma_{xx}\sigma + \frac{1}{2}\sigma_{x}^{2} - \frac{1}{2}\sigma_{yy}\sigma + \frac{1}{2}\sigma_{y}^{2} - \frac{B}{4}(\varphi_{xx} + \varphi_{yy})\sigma - B(\varphi_{xx} + \varphi_{yy})\sigma^{2}\sigma_{rr} - \frac{B^{2}}{2}\sigma\sigma_{rr}\left((\varphi_{y})^{2} + (\varphi_{x})^{2}\right) + \frac{B}{2}\sigma_{y}\varphi_{y} + \frac{B}{2}\sigma_{x}\varphi_{x} + \frac{B^{2}}{8}\left((\varphi_{x})^{2} + (\varphi_{y})^{2}\right) = 0 \quad (4.21)$$

where Ric^N denotes the Ricci tensor of the Kähler metric g_o of N and $\operatorname{Ric}_{ij}^N := \operatorname{Ric}(E_i, E_j)$.

We give the details of the proof of this lemma in Appendix B. Plugging the expressions

$$\sigma = -\frac{1}{4B\varphi}r^2 - \frac{B\varphi}{4}, \qquad \sigma_r = -\frac{1}{2B\varphi}r, \qquad \sigma_{rr} = -\frac{1}{2B\varphi},$$

$$\sigma_x = \frac{\varphi_x}{4B\varphi^2}r^2 - \frac{B\varphi_x}{4}, \qquad \sigma_y = \frac{\varphi_y}{4B\varphi^2}r^2 - \frac{B\varphi_y}{4}, \qquad (4.22)$$

$$\sigma_{xx} = \frac{\varphi_{xx}}{4B\varphi^2}r^2 - \frac{\varphi_x^2}{2B\varphi^3}r^2 - \frac{B\varphi_{xx}}{4}, \qquad \sigma_{yy} = \frac{\varphi_{yy}}{4B\varphi^2}r^2 - \frac{\varphi_y^2}{2B\varphi^3}r^2 - \frac{B\varphi_{yy}}{4}.$$

into (4.21), we see that the left hand side is an even polynomial of degree 4 in the variable r. The equation is therefore satisfied if and only if the coefficients of the monomials r^4 , r^2 and r^0 are identically vanishing. These three conditions are

$$\frac{1}{16B^2\varphi^2} \left(\operatorname{Ric}_{11}^N - \frac{\varphi_x^2 + \varphi_y^2}{2\varphi^2} + \frac{\varphi_{xx} + \varphi_{yy}}{\varphi} \right) = 0 , \quad \frac{1}{8} \left(\operatorname{Ric}_{11}^N - \frac{\varphi_x^2 + \varphi_y^2}{2\varphi^2} + \frac{\varphi_{xx} + \varphi_{yy}}{\varphi} \right) = 0 ,$$

$$\frac{\varphi^2}{16^2} \left(\operatorname{Ric}_{11}^N - \frac{\varphi_x^2 + \varphi_y^2}{2\varphi^2} + \frac{\varphi_{xx} + \varphi_{yy}}{\varphi} \right) = 0 ,$$

$$(4.23)$$

which are manifestly equivalent each other and all to the single equation

$$\operatorname{Ric}_{11}^{N} - \frac{\varphi_x^2 + \varphi_y^2}{2\varphi^2} + \frac{\Delta\varphi}{\varphi} = 0 \left(\iff -\frac{1}{2}\Delta(\log\Delta\varphi) + \frac{\Delta\varphi}{2\operatorname{sign}(\varphi)|\varphi|} + \frac{1}{2}\Delta\log|\varphi| = 0 \right)$$
(4.24)

(here we used the classical formula for curvatures in isothermal coordinates). Introducing the auxiliary function $\psi := \frac{\Delta \varphi}{|\varphi|} > 0$, the equation (4.24) becomes equivalent to the system

$$\Delta \varphi - \operatorname{sign}(\varphi)\varphi\psi = 0, \qquad -\frac{1}{2}\Delta(\log\psi) + \operatorname{sign}(\varphi)\frac{1}{2}\psi = 0.$$
(4.25)

If we consider the metric $\tilde{g}_o|_{\mathcal{V}}$ on \mathcal{V} with components $(\tilde{g}_{o\,ij}) = \begin{pmatrix} \frac{\psi}{2} & 0\\ 0 & \frac{\psi}{2} \end{pmatrix}$, the second equation in (4.25) simply means that, in the regions where $\operatorname{sign}(\varphi)$ is constant, the Gaussian curvature κ of $\tilde{g}_o|_{\mathcal{V}}$ is constant and equal to $\kappa = -\operatorname{sign}(\varphi) = \mp 1$. At the same time, since (x, y) are isothermal coordinates for both metrics, the first equation in (4.25) can be rewritten as $g_o|_{\mathcal{V}} + 2\kappa\varphi\tilde{g}_o|_{\mathcal{V}}$. This yields the above described results for the second step.

4.2. Classification of backgrounds of Kerr type. As we previously planned, we can now prove the main classification result of this section.

Corollary 4.3. Let M be a 4-dimensional Kerr manifold, equipped with a Kerr structure \mathcal{M} and the corresponding \mathcal{M} -canonical optical structure \mathcal{Q}_o , as in Theorem 4.1. Such a Kerr manifold admits a background metric η of Kerr type if and only if the following hold:

(1) The Kähler surface (N, J, g_o) satisfies: (a) it is non-compact, (b) it admits a Riemannian metric \tilde{g}_o of constant Gaussian curvature $\kappa = \pm 1$, which is Kähler for the complex manifold (N, J) and (c) there exists a global Kähler potential $\varphi : N \to \mathbb{R}$ for g_o satisfying

$$\operatorname{sign}\varphi = -\kappa , \qquad g_o + 2\kappa\varphi \,\widetilde{g}_o = 0. \tag{4.26}$$

(2) The \mathbb{R} -bundle $\pi: \mathbb{S} \to N$ is trivial and, up to a bundle automorphism of M, there is a standard coordinate r for the fiber of $\pi^{\mathbb{S}}: M = \mathbb{S} \times \mathbb{R} \to \mathbb{S}$ such that $\sigma = -\frac{r^2}{4B\varphi} - \frac{B\varphi}{4}$.

If M satisfies the above conditions, for any φ satisfying $\omega_o = dd^c \varphi$ and (4.2), there is a family of background metrics η , parameterised by B, with σ as in (2).

Proof. By Proposition 2.9, any background metric η on M is flat and satisfies (4.1). Thus, the necessity of the above conditions is a consequence of Theorem 4.1. Conversely, assume that there are \tilde{g}_o and φ as in the hypotheses and consider an open cover $\{\mathcal{V}_A\}_{A\in\mathcal{J}}$ as in Lemma 2.7 and on each corresponding trivialisable set $\mathcal{U} = \check{\pi}^{-1}(\mathcal{V}) = \mathcal{V} \times \mathbb{R} \times \mathbb{R}$, let $\sigma: \mathcal{V}_A \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ be as (4.3). By condition (2) these locally defined functions combine into a global function $\sigma: M \to \mathbb{R}$. Let \mathcal{V} be one of the sets of the open cover $\{\mathcal{V}_A\}$ and denote by $(\hat{E}_1, \hat{E}_2, \mathbf{p}_o, \mathbf{q}_o)$ the adapted frame field on $\check{\pi}^{-1}(\mathcal{V})$ in which \hat{E}_1, \hat{E}_2 are lifts of the two coordinate vector fields, given by g_o -isothermal coordinates (x, y) on \mathcal{V} . By Theorem 4.1, the Ricci tensor of a metric η as in (a) of Theorem 3.1 with $\tilde{\beta} = B \neq 0$ being constant and σ determined as above, the equation (4.1) vanishes on the section \mathcal{W}_o . Therefore, in order to prove that such η is Einstein on each open set $\check{\pi}^{-1}(\mathcal{V})$, it suffices to show that on any such open set it satisfies the Einstein equations that are not included in (4.1), that is the equations

$$\operatorname{Ric}(\mathbf{q}_{o},\mathbf{q}_{o}) = \operatorname{Ric}(\mathbf{q}_{o},\mathbf{p}_{o}) = \operatorname{Ric}(\mathbf{q}_{o},\hat{E}_{1}) = \operatorname{Ric}(\mathbf{q}_{o},\hat{E}_{2}) = 0 .$$
(4.27)

Let us start with the first two equations, $\operatorname{Ric}(\mathbf{q}_o, \mathbf{q}_o) = 0$ and $\operatorname{Ric}(\mathbf{q}_o, \mathbf{p}_o) = 0$. By definition, $\operatorname{Ric}(\mathbf{q}_o, \mathbf{q}_o) = \operatorname{R}_{m\mathbf{q}_o\mathbf{q}_o}{}^m + \operatorname{R}_{\mathbf{p}_o\mathbf{q}_o\mathbf{q}_o}{}^{\mathbf{p}_o}$ and $\operatorname{Ric}(\mathbf{p}_o, \mathbf{q}_o) = \operatorname{R}_{m\mathbf{p}_o\mathbf{q}_o}{}^m + \operatorname{R}_{\mathbf{q}_o\mathbf{p}_o\mathbf{q}_o}{}^{\mathbf{q}_o}$. Expressing the components of the Riemann curvature in terms of the Christoffel symbols (3.7) – (3.13) (with $\tilde{\beta} := B$!) and omitting all terms that are manifestly zero, the two equations can be written as

$$\operatorname{Ric}_{q_{o}q_{o}} = -q_{o}(\boldsymbol{\Gamma}_{mq_{o}}^{m}) - \boldsymbol{\Gamma}_{mq_{o}}^{\ell} \boldsymbol{\Gamma}_{q_{o}\ell}^{m} = -B^{2} p_{o}(\boldsymbol{\Gamma}_{mp_{o}}^{m}) - B^{2} \boldsymbol{\Gamma}_{mp_{o}}^{\ell} \boldsymbol{\Gamma}_{p_{o}\ell}^{m} = \\ = B^{2} \left(-p_{o} \left(\frac{J_{m}^{m}}{4\sigma} + \frac{p_{o}(\sigma)\delta_{m}^{m}}{2\sigma} \right) - \left(\frac{J_{m}^{\ell}}{4\sigma} + \frac{p_{o}(\sigma)\delta_{m}^{\ell}}{2\sigma} \right) \left(\frac{J_{\ell}^{m}}{4\sigma} + \frac{p_{o}(\sigma)\delta_{\ell}^{m}}{2\sigma} \right) \right) = \\ = B^{2} \left(-2 \frac{p_{o}(p_{o}(\sigma))}{2\sigma} + 2 \frac{p_{o}(\sigma)^{2}}{4\sigma^{2}} + \frac{2}{16\sigma^{2}} \right) = \\ = \frac{B^{2}}{\sigma^{2}} \left(-2\sigma p_{o}(p_{o}(\sigma)) + p_{o}(\sigma)^{2} + \frac{1}{4} \right) = 0 , \qquad (4.28)$$

Ric_{p,q,e} = - $p_{o}(\boldsymbol{\Gamma}_{mq}^{m}) - \boldsymbol{\Gamma}_{m\ell}^{m\ell} \boldsymbol{\Gamma}_{n}^{m}$

Since they are both equivalent to (3.15), they are identically satisfied because η satisfies $\operatorname{Ric}(\mathbf{p}_o, \mathbf{p}_o) = 0$ by (4.1).

For what concerns the equations $\operatorname{Ric}(\widehat{E}_i, \mathbf{q}_o) = 0$, i = 1, 2, using once again the expressions (3.5) and (3.6) and neglecting the terms that are trivial, according to (3.7) - (3.13), the two equations reduce to

$$\operatorname{Ric}(\widehat{E}_{i},\mathbf{q}_{o}) = \operatorname{R}_{miq_{o}}^{m} + \operatorname{R}_{\mathbf{p}_{o}iq_{o}}^{\mathbf{p}_{o}} + \operatorname{R}_{\mathbf{q}_{o}iq_{o}}^{\mathbf{q}_{o}} = \\ = \widehat{E}_{m}(\boldsymbol{\Gamma}_{iq_{o}}^{m}) - \widehat{E}_{i}(\boldsymbol{\Gamma}_{mq_{o}}^{m}) - \boldsymbol{\Gamma}_{mq_{o}}^{\ell}\boldsymbol{\Gamma}_{i\ell}^{m} + \boldsymbol{\Gamma}_{iq_{o}}^{\ell}\boldsymbol{\Gamma}_{m\ell}^{m} - \boldsymbol{\Gamma}_{\mathbf{p}_{o}q_{o}}^{\mathbf{q}_{o}}\boldsymbol{\Gamma}_{iq_{o}}^{\mathbf{p}_{o}} = 0 . \quad (4.30)$$

Using once again the explicit expressions for the Christoffel symbols (3.7) - (3.13) with $\tilde{\beta} := B$, a straightforward check shows that these two equations are equivalent to the equations $\operatorname{Ric}(\hat{E}_i, \mathbf{p}_o) = 0$, which we discussed in §4.1.1. Since (4.1) holds, we conclude that also the equations (4.30) are identically satisfied.

5. KERR FAMILIES OF RICCI FLAT GRAVITATIONAL FIELDS

As in the previous section, M is a 4-dimensional Kerr manifold, with a Kerr structure $\mathcal{M} = (M = \mathcal{S} \times \mathbb{R}, \pi : \mathcal{S} \to N, (J, g_o), \mathcal{H})$ and associated \mathcal{M} -canonical optical structure $\mathcal{Q}_o = (\mathcal{W}_o, [h_o]_{\pm}, \mathcal{K}_o, \{g\})$. The next theorem together with Corollaries 3.2 and 4.3 represent our main result. In §6 we will show that it can be re-stated as in Theorem 1.1.

Theorem 5.1. Given M, M and Q, assume that M admits a background metric η of Kerr type, determined by a constant $B \neq 0$ and functions $\varphi : N \to \mathbb{R}$, $\sigma : M \to \mathbb{R}$ as in Corollary 4.3. Then there exists a constant $m_o \neq 0$ and a Ricci flat compatible metric $g^{(m=m_o)} \neq \eta$ of the form (2.15) in a connected open subset $M' = S \times I$, $I \subset \mathbb{R}$, of M if and only if the following two conditions hold:

- (1) B = -1;
- (2) Using the coordinate r for the fiber of $\pi^{\mathbb{S}} : M = \mathbb{S} \times \mathbb{R} \to \mathbb{S}$, so that $p_o = \frac{\partial}{\partial r}$ and σ takes the form (4.3), the function $\tilde{\beta}_o$ takes the form

$$\widetilde{\beta}_o = \frac{\mathbf{k}r}{r^2 + \varphi^2} , \qquad (5.1)$$

for some constant $\mathbf{k} \neq 0$.

In this case, all metrics $g^{(m)}$ of the form (2.15), with B = -1 and $\tilde{\beta}_o$ as in (5.1), are Ricci flat and constitute a Kerr family of gravitational fields.

The proof is essentially just a straightforward consequence of imposing the condition $\operatorname{Ric}^{(m)} = 0$ on the Ricci tensor of a metric $g^{(m)}$ having the form (2.15) with the same constant B and the same function σ of the background η . Considering coordinates (x, y, v, r) and an adapted frame field $\mathcal{B} = \left(\hat{E}_1, \hat{E}_2, \mathbf{p}_o = \frac{\partial}{\partial r}, \mathbf{q}_o = \frac{\partial}{\partial v} + B\frac{\partial}{\partial r}\right)$ in an open set $\check{\pi}^{-1}(\mathcal{V}_A) \subset M$ of Lemma 2.7, with $E_1 = \frac{\partial}{\partial x}, E_2 = \frac{\partial}{\partial y}$, the Ricci flatness condition is equivalent to the three following systems of equations:

$$\operatorname{Ric}^{(m)}(\mathbf{p}_{o}, \mathbf{p}_{o}) = 0 , \qquad \operatorname{Ric}^{(m)}(\mathbf{p}_{o}, \widehat{E}_{i}) = 0 , \qquad (5.2)$$

$$\operatorname{Ric}^{(m)}(\mathbf{p}_{o},\mathbf{q}_{o}) = 0$$
, $\operatorname{Ric}^{(m)}(\widehat{E}_{i},\widehat{E}_{j}) = 0$, (5.3)

$$\operatorname{Ric}^{(m)}(\widehat{E}_i, \mathbf{q}_o) = 0$$
, $\operatorname{Ric}^{(m)}(\mathbf{q}_o, \mathbf{q}_o) = 0$. (5.4)

The proof is divided into three steps, which we are going to perform in §5.1, 5.2 and 5.3, respectively. First, we show that, since $\eta = g^{(m=0)}$ is a background metric, the equations (5.2) are identically satisfied by any metric of the form (2.15) (regardless of what are the constant B and the function $\tilde{\beta}_o$). Second, we prove that there exists an $m_o \neq 0$ such that $g^{(m_o)}$ satisfies (5.3) if and only if $\tilde{\beta}_o$ has the form (5.1) for a freely specifiable function $\mathbf{k} = \mathbf{k}(x, y)$ on the open set $\mathcal{V}_A \subset N$. In the last step, we show that, assuming that (5.3) is satisfied (and hence that $\tilde{\beta}_o$ has the form (5.1) for some function \mathbf{k}), the remaining equations (5.4) are satisfied if and only if $\mathbf{k} = \mathbf{k}(x, y)$ is constant and B = -1. This will prove the first claim and imply the last claim as well, because it will show that all metrics $g^{(m)}$ of the form (2.15) satisfy (1) and (2) if and only if those conditions are satisfied by just one value $m = m_o \neq 0$.

5.1. First step. From now on, we use the notation Γ_{AB}^{C} and $\widetilde{\Gamma_{AB}^{C}}$ for the Christoffel symbols with respect to the adapted frame field \mathcal{B} of the background metric $\eta = g^{(m=0)}$

and of the metric $g^{(m_o)}$, respectively. We also denote $\mathbf{H}_{AB}^{\ C} := \widetilde{\mathbf{\Gamma}_{AB}^{\ C}} - \mathbf{\Gamma}_{AB}^{\ C}$, so that we may write that $\widetilde{\mathbf{\Gamma}_{AB}^{\ C}} = \mathbf{\Gamma}_{AB}^{\ C} + \mathbf{H}_{AB}^{\ C}$. From (3.7) – (3.13) we have (recall that $\mathbf{q}_o(\widetilde{\beta}_o) = B\mathbf{p}_o(\widetilde{\beta}_o)$!)

$$\mathbf{H}_{ij} = \mathbf{0} \tag{(3.3)}$$

$$\mathbf{H}_{ij}^{\mathbf{p}_o} = m_o g_{ij} \beta_o \mathbf{p}_o(\sigma) , \quad \mathbf{H}_{ij}^{\mathbf{q}_o} = 0 , \qquad (5.6)$$

$$\mathbf{H}_{ip_{o}}^{m} = \mathbf{H}_{p_{o}i}^{m} = 0 , \qquad \mathbf{H}_{ip_{o}}^{p_{o}} = \mathbf{H}_{p_{o}i}^{p_{o}} = \mathbf{H}_{ip_{o}}^{q_{o}} = \mathbf{H}_{p_{o}i}^{q_{o}} = 0 , \qquad (5.7)$$

$$\mathbf{H}_{i\mathbf{q}_{o}}^{r} = \mathbf{H}_{\mathbf{q}_{o}i}^{r} = m_{o}\frac{J_{i}}{4\sigma}\widetilde{\beta}_{o} ,$$

$$\mathbf{H}^{\mathbf{p}_{o}} = \mathbf{H}^{\mathbf{p}_{o}} = \frac{m_{o}}{6}\widetilde{\beta} , \qquad \frac{m_{o}}{2}J_{i}^{\ell}(\mathbf{q},\mathbf{p},i) \quad \mathbf{H}^{\mathbf{q}_{o}} = \mathbf{H}^{\mathbf{q}_{o}} = 0$$
(5.8)

$$\mathbf{H}_{i\mathbf{q}_{o}}^{\mathbf{p}_{o}} = \mathbf{H}_{\mathbf{q}_{o}i}^{\mathbf{p}_{o}} = \frac{m_{o}}{2}\widetilde{\beta}_{oi} - \frac{m_{o}}{2}J_{i}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\widetilde{\beta}_{o}) , \quad \mathbf{H}_{i\mathbf{q}_{o}}^{\mathbf{q}_{o}} = \mathbf{H}_{\mathbf{q}_{o}i}^{\mathbf{q}_{o}} = 0 , \quad (5.8)$$

$$\mathbf{H}_{\mathbf{p}_{o}\mathbf{p}_{o}}^{m} = \mathbf{H}_{\mathbf{p}_{o}\mathbf{p}_{o}}^{\mathbf{p}_{o}} = \mathbf{H}_{\mathbf{p}_{o}\mathbf{p}_{o}}^{\mathbf{q}_{o}} = 0 , \qquad (5.9)$$

$$\mathbf{H}_{\mathbf{p}_{o}\mathbf{q}_{o}}^{m} = \mathbf{H}_{\mathbf{q}_{o}\mathbf{p}_{o}}^{m} = 0 , \qquad \mathbf{H}_{\mathbf{p}_{o}\mathbf{q}_{o}}^{p} = \mathbf{H}_{\mathbf{q}_{o}\mathbf{p}_{o}}^{p} = \frac{m_{o}}{2}\mathbf{p}_{o}(\widetilde{\beta}_{o}) , \qquad \mathbf{H}_{\mathbf{p}_{o}\mathbf{q}_{o}}^{q} = \mathbf{H}_{\mathbf{q}_{o}\mathbf{p}_{o}}^{q} = 0 , \quad (5.10)$$

$$\mathbf{H}_{\mathbf{q}_{o}\mathbf{q}_{o}}^{r} = -m_{o}\frac{g^{r\kappa}}{4\sigma}\widetilde{\beta}_{ok} - m_{o}B\frac{g^{r\kappa}}{4\sigma}J_{k}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\widetilde{\beta}_{o}) , \\
\mathbf{H}_{\mathbf{q}_{o}\mathbf{q}_{o}}^{\mathbf{p}_{o}} = \frac{m_{o}B}{2}\mathbf{p}_{o}(\widetilde{\beta}_{o}) + m_{o}\frac{\mathbf{p}_{o}(\widetilde{\beta}_{o})(B+m_{o}\widetilde{\beta}_{o})}{2} , \ \mathbf{H}_{\mathbf{q}_{o}\mathbf{q}_{o}}^{\mathbf{q}_{o}} = -\frac{m_{o}}{2}\mathbf{p}_{o}(\widetilde{\beta}_{o}) .$$
(5.11)

As announced above, in this step we just observe that the equations $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{p}_o) = \operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \hat{E}_i) = 0$ are identically satisfied for any choice of β_o and B. Indeed, after expanding $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{p}_o)$ in terms of the Christoffel symbols $\Gamma_{AB}^{\ C}$, one can immediately see that $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{p}_o)$ is independent of the parameter m_o and is therefore identically equal to 0, because $\eta = g^{(m=0)}$ is Ricci flat. For instance, using the fact that $c_{mj}^r \equiv 0$, we immediately get

$$\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \hat{E}_i) = \hat{E}_m(\mathbf{\Gamma}_{i\mathbf{p}_o}^m) + \hat{E}_m(\mathbf{H}_{i\mathbf{p}_o}^m) - \hat{E}_i(\mathbf{\Gamma}_{m\mathbf{p}_o}^m) - \hat{E}_i(\mathbf{H}_{m\mathbf{p}_o}^m) - \left(\mathbf{\Gamma}_{m\mathbf{p}_o}^{\ell} + \mathbf{H}_{m\mathbf{p}_o}^{\ell}\right) - \left(\mathbf{\Gamma}_{i\mathbf{p}_o}^{\ell} + \mathbf{H}_{i\mathbf{p}_o}^{\ell}\right) + \left(\mathbf{\Gamma}_{i\mathbf{p}_o}^{\ell} + \mathbf{H}_{i\mathbf{p}_o}^{\ell}\right) \left(\mathbf{\Gamma}_{m\ell}^{m} + \mathbf{H}_{m\ell}^{m}\right) = \operatorname{Ric}^{(m=0)}(\mathbf{p}_o, \hat{E}_i) = 0$$

Similar computations hold for $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{p}_o) = 0.$

5.2. Second step. Let us expand the left hand side of the first equation in (5.3) (i.e. the Ricci component $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{q}_o)$) in terms of the Christoffel symbols $\widetilde{\Gamma_{AB}} = \Gamma_{AB}^{\ \ C} + \mathbf{H}_{AB}^{\ \ C}$. Using the Ricci flatness of η , the equation reduces to

$$\begin{split} \operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{q}_o) &= -\operatorname{p}_o(\mathbf{\Gamma}_{r\mathbf{q}_o}^r + \mathbf{H}_{r\mathbf{q}_o}^r) - \left(\mathbf{\Gamma}_{r\mathbf{q}_o}^\ell + \mathbf{H}_{r\mathbf{q}_o}^\ell\right)\left(\mathbf{\Gamma}_{\mathbf{p}_o\ell}^r + \mathbf{H}_{\mathbf{p}_o\ell}^r\right) + \\ &+ \left(\mathbf{\Gamma}_{\mathbf{p}_o\mathbf{q}_o}^{\mathbf{p}_o} + \mathbf{H}_{\mathbf{p}_o\mathbf{q}_o}^{\mathbf{p}_o}\right)\left(\mathbf{\Gamma}_{r\mathbf{p}_o}^r + \mathbf{H}_{r\mathbf{p}_o}^r\right) - \operatorname{p}_o(\mathbf{\Gamma}_{\mathbf{q}_o\mathbf{q}_o}^{\mathbf{q}_o} + \mathbf{H}_{\mathbf{q}_o\mathbf{q}_o}^{\mathbf{q}_o}) = \\ &= \operatorname{Ric}_{\mathbf{p}_o\mathbf{q}_o} - \mathbf{H}_{r\mathbf{q}_o}^\ell \mathbf{\Gamma}_{\mathbf{p}_o\ell}^r + \mathbf{H}_{\mathbf{p}_o\mathbf{q}_o}^{\mathbf{p}_o} \mathbf{\Gamma}_{r\mathbf{p}_o}^r - \operatorname{p}_o(\mathbf{H}_{\mathbf{q}_o\mathbf{q}_o}^{\mathbf{q}_o}) = \end{split}$$

EINSTEIN MANIFOLDS WITH OPTICAL GEOMETRIES OF KERR TYPE

$$= -m_o \frac{J_r^{\ell}}{4\sigma} \widetilde{\beta}_o \left(\frac{J_\ell^r}{4\sigma} + \frac{\delta_\ell^r}{2\sigma} \mathbf{p}_o(\sigma) \right) + m_o \mathbf{p}_o(\widetilde{\beta}_o) \frac{\mathbf{p}_o(\sigma)}{2\sigma} + \frac{m_o}{2} \mathbf{p}_o(\mathbf{p}_o(\widetilde{\beta}_o)) =$$
$$= \frac{m_o}{2} \left(\mathbf{p}_o(\mathbf{p}_o(\widetilde{\beta}_o)) + \mathbf{p}_o(\widetilde{\beta}_o) \frac{\mathbf{p}_o(\sigma)}{\sigma} + \frac{\widetilde{\beta}_o}{4\sigma^2} \right) = 0 .$$
(5.12)

Recalling that $p_o = \frac{\partial}{\partial r}$ and that σ has the form $\sigma(x, y, r) = -\frac{r^2}{4B\varphi(x,y)} - \frac{B\varphi(x,y)}{4}$, the equation (5.12) can be written as

$$\sigma \tilde{\beta}_{o\,rr} + \tilde{\beta}_{o\,r} \sigma_r + \frac{\tilde{\beta}_o}{4\sigma} = 0.$$
(5.13)

The general solution of (5.13) is

$$\widetilde{\beta}_{o}(x,y,r) = \mathbf{k}_{1}(x,y) \frac{\left(B^{2}\varphi^{2}(x,y) - r^{2}\right)}{r^{2} + B^{2}\varphi^{2}(x,y)} + \mathbf{k}_{2}(x,y) \frac{r}{r^{2} + B^{2}\varphi^{2}(x,y)} , \qquad (5.14)$$

where $\mathbf{k}_1 = \mathbf{k}_1(x, y), \mathbf{k}_2 = \mathbf{k}_2(x, y)$ are two freely specifiable functions, independent of r. Therefore, if $g^{(m_o)}$ is Ricci flat, on each set $\check{\pi}^{-1}(\mathcal{V}_A)$ such a metric is completely determined by the constant B, the potential φ and a free choice of the functions $\mathbf{k}_1, \mathbf{k}_2 : \mathcal{V}_A \to \mathbb{R}$, provided that σ is as in (4.3) and $\check{\beta}_o$ is as in (5.14). From now on, we assume all this, so that we automatically have $\operatorname{Ric}^{(m_o)}(\mathbf{p}_o, \mathbf{q}_o) = 0$. Let us now consider the remaining equations in (5.3). For any $1 \leq i, j \leq 2$, we have

$$\begin{aligned} \operatorname{Ric}^{(m_o)}(\widehat{E}_i, \widehat{E}_j) &= \overbrace{\operatorname{Ric}^{(m=0)}(\widehat{E}_i, \widehat{E}_j)}^{(m_o} - \operatorname{H}_{mj}^{p_o} \Gamma_{ip_o}^{m} - \Gamma_{mj}^{q_o} \operatorname{H}_{iq_o}^{m} + \\ &+ \operatorname{H}_{ij}^{p_o} \Gamma_{mp_o}^{m} + \overbrace{\Gamma_{ij}}^{q_o} \operatorname{H}_{mq_o}^{m} + \omega_{mi} \operatorname{H}_{q_oj}^{m} + \\ &+ \operatorname{p}_o(\operatorname{H}_{ij}^{p_o}) - \Gamma_{p_oj}^{\ell} \operatorname{H}_{i\ell}^{p_o} + \Gamma_{ij}^{q_o} \operatorname{H}_{p_oq_o}^{p_o} - \operatorname{H}_{q_oj}^{\ell} \Gamma_{i\ell}^{q_o} + \Gamma_{ij}^{q_o} \operatorname{H}_{q_oq_o}^{q_o} = \\ &= -\frac{m_o \widetilde{\beta}_o \omega_{ij}}{4\sigma} \operatorname{p}_o(\sigma) - \frac{m_o \widetilde{\beta}_o g_{ij}}{2\sigma} (\operatorname{p}_o(\sigma))^2 - \frac{m_o \widetilde{\beta}_o g_{ij}}{8\sigma} + \frac{m_o \widetilde{\beta}_o \omega_{ij}}{4\sigma} \operatorname{p}_o(\sigma) + \\ &+ \frac{m_o \widetilde{\beta}_o g_{ij}}{\sigma} (\operatorname{p}_o(\sigma))^2 - \frac{m_o \widetilde{\beta}_o g_{ji}}{4\sigma} + m_o g_{ij} \operatorname{p}_o(\widetilde{\beta}_o) \operatorname{p}_o(\sigma) + m_o \widetilde{\beta}_o g_{ij} \operatorname{p}_o(\sigma)) + \\ &+ \frac{m_o \widetilde{\beta}_o \omega_{ij}}{4\sigma} \operatorname{p}_o(\sigma) - \frac{m_o \widetilde{\beta}_o g_{ij}}{2\sigma} (\operatorname{p}_o(\sigma))^2 - \frac{m_o \omega_{ij}}{4} \operatorname{p}_o(\widetilde{\beta}_o) - \frac{m_o g_{ij}}{2} \operatorname{p}_o(\widetilde{\beta}_o) \operatorname{p}_o(\sigma) + \\ &+ \frac{m_o \widetilde{\beta}_o g_{ji}}{4\sigma} + \frac{m_o \widetilde{\beta}_o \omega_{ji}}{4\sigma} \operatorname{p}_o(\sigma) + \frac{m_o \omega_{ij}}{4} \operatorname{p}_o(\widetilde{\beta}_o) + \frac{m_o g_{ij}}{2} \operatorname{p}_o(\widetilde{\beta}_o) \operatorname{p}_o(\sigma) = \\ &= m_o g_{ij} \left(- \frac{\widetilde{\beta}_o}{4\sigma} + \operatorname{p}_o(\widetilde{\beta}_o) \operatorname{p}_o(\sigma) + \widetilde{\beta}_o \operatorname{p}_o(\sigma)) \right) \end{aligned}$$

If we now plug the expressions for $\tilde{\beta}_o$ and σ into this formula, we get that

$$\operatorname{Ric}^{(m_o)}(\hat{E}_i, \hat{E}_j) = \frac{m_o \mathbf{k}_1 g_{ij}}{2B\varphi} .$$
(5.15)

Since $m_o \neq 0$, this means that the equations $\operatorname{Ric}^{(m_o)}(\widehat{E}_i, \widehat{E}_j) = 0$, i, j = 1, 2, are satisfied if and only if the function $\mathbf{k}_1 = \mathbf{k}_1(x, y)$ is identically zero $\widetilde{\beta}_o = \widetilde{\beta}_o(x, y, r)$ has the form

$$\widetilde{\beta}_o(x,y,r) = \mathbf{k}(x,y)\frac{r}{r^2 + B^2\varphi^2(x,y)} , \qquad (5.16)$$

for a freely specifiable function $\mathbf{k} = \mathbf{k}(x, y)$.

5.3. Third step. From now on, we assume not only that σ is as in (4.3) but also that $\tilde{\beta}_o$ is as in (5.16) so that all equations in (5.2) and (5.3) are satisfied. Let us expand the component $\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \hat{E}_i), i = 1, 2$, in terms of the Christoffel symbols $\widetilde{\Gamma_{AB}} = \Gamma_{AB}^{\ \ C} + \mathbf{H}_{AB}^{\ \ C}$. We recall that, according to the hypotheses on the coordinates (x, y) and on the frame field (E_1, E_2) , all functions c_{ij}^k are zero and the components J_i^j are equal to either ± 1 or 0 at all points. Therefore, $\operatorname{Ric}^{(m_o)}(\hat{E}_i, \mathbf{q}_o)$ is equal to

$$\begin{split} \operatorname{Ric}^{(m_{o})}(\hat{E}_{i},\mathbf{q}_{o}) &= \overline{\operatorname{Ric}}^{(m_{o})}(\hat{E}_{i},\overline{\mathbf{q}_{o}}) + \hat{E}_{r}(\mathbf{H}_{iq_{o}}^{r}) - \mathbf{H}_{q_{o}}^{\ell}\mathbf{r}_{o}^{r} - \mathbf{H}_{q_{o}}^{p}\mathbf{r}_{o}^{r}\mathbf{r}_{o}^{s} + \mathbf{H}_{iq_{o}}^{\ell}\mathbf{\Gamma}_{r\ell}^{r} + \\ &+ \mathbf{H}_{iq_{o}}^{p}\mathbf{\Gamma}_{rp_{o}}^{r} + \omega_{ri}\mathbf{H}_{q_{o}q_{o}}^{r} - \mathbf{p}_{o}(\mathbf{H}_{iq_{o}}^{p}) - \hat{E}_{i}(\mathbf{H}_{p,q_{o}}^{p}) - \hat{E}_{i}(\mathbf{H}_{q,q_{o}}^{q}) - \mathbf{H}_{q,q_{o}}^{\ell}\mathbf{L}_{o}^{r}\mathbf{\Gamma}_{i\ell}^{r} + \\ &+ \mathbf{H}_{iq_{o}}^{p}\mathbf{\Gamma}_{rp_{o}}^{r} + \omega_{ri}\mathbf{H}_{q,q_{o}}^{r} + \mathbf{p}_{o}(\mathbf{H}_{iq_{o}}^{p}) - \hat{E}_{i}(\mathbf{H}_{q,q_{o}}^{q}) - \mathbf{H}_{q,q_{o}}^{\ell}\mathbf{L}_{o}^{r}\mathbf{\Gamma}_{i\ell}^{r} + \\ &+ \mathbf{H}_{iq_{o}}^{p}\mathbf{\Gamma}_{r\ell}^{r} + \frac{\sigma_{\ell}}{4\sigma^{2}}\hat{\beta}_{o} + m_{o}g_{i}B\frac{\mathbf{p}_{o}(\sigma)}{4\sigma^{2}}\hat{\beta}_{o} - m_{o}\varphi_{i}\frac{\mathbf{B}_{p}(\tilde{\beta})}{4\sigma} - \\ &- m_{o}\frac{1}{4\sigma}\tilde{\beta}_{o}\left(J_{r}^{\ell}\mathbf{\Gamma}_{i\ell}^{r} + \frac{\sigma_{\ell}}{\sigma}J_{i}^{\ell} - B\frac{\mathbf{p}_{o}(\sigma)\varphi_{i}}{\sigma}\right) - \left(\frac{m_{o}}{2}\tilde{\beta}_{or} - \frac{m_{o}}{2}J_{r}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\tilde{\beta})\right)\left(\frac{J_{i}}{4\sigma} + \frac{\delta_{i}^{r}}{2\sigma}\mathbf{p}_{o}(\sigma)\right) + \\ &+ m_{o}\frac{\delta_{o}}{4\sigma^{2}}\left(\mathbf{\Gamma}_{r\ell}^{r} + \frac{\sigma_{\ell}}{\sigma} + B\frac{\varphi_{i}J_{i}^{l}p_{o}(\sigma)}{\sigma}\right) + \left(\frac{m_{o}}{2}\tilde{\beta}_{oi} - \frac{m_{o}}{2}J_{i}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\tilde{\beta})\right)\frac{\mathbf{p}_{o}(\sigma)}{\sigma} + \\ &+ m_{o}\frac{J_{i}^{k}}{4\sigma}\tilde{\beta}_{ok} - m_{o}B\frac{1}{4\sigma}\varphi_{i}\mathbf{p}_{o}(\tilde{\beta}_{o}) + \frac{m_{o}}{2}\mathbf{p}_{o}\left(\tilde{\beta}_{oi} - J_{i}^{\ell}\mathbf{p}_{o}(\tilde{\beta}_{o})\varphi_{i}\right) - \frac{m_{a}}{2}\tilde{E}_{i}(\mathbf{p}_{o}(\tilde{\beta}))\right) + \\ &+ \frac{m_{a}}{2}\tilde{E}_{i}(\mathbf{p}_{o}(\tilde{\beta})) - \left(m_{o}\frac{g^{\ell k}}{4\sigma^{2}}\tilde{\beta}_{ok} + m_{o}B\frac{g^{\ell k}}{4\sigma^{2}}J_{s}^{k}\varphi_{s}\mathbf{p}_{o}(\tilde{\beta}_{o})\right)\left(\frac{\omega_{i\ell}}{2} + g_{i\ell}\mathbf{p}_{o}(\sigma)\right) = \\ &= -m_{o}J_{i}\left(\frac{\sigma_{\ell}}{4\sigma^{2}}\tilde{\beta}_{o} - m_{o}J_{i}\left(\frac{1}{4\sigma}\tilde{\beta}_{o\ell}}\right) + \frac{m_{o}\varphi_{i}}{4\sigma^{2}}\tilde{\beta}_{o}} - m_{o}g_{i}(\tilde{\beta}_{o})\right) - \left(m_{o}\frac{g^{\ell k}}{4\sigma^{2}}\tilde{\beta}_{o}} - m_{o}g_{i}\frac{g}{4\sigma^{2}}\tilde{\beta}_{o}} - m_{o}B\varphi_{i}\frac{g}{4\sigma^{2}}}\right) - \\ - \frac{m_{o}J_{i}}\left(\frac{\sigma_{\ell}}{4\sigma^{2}}\tilde{\beta}_{o}} - m_{o}J_{i}\left(\frac{1}{4\sigma}\tilde{\beta}_{o}}\right) + m_{o}J_{i}^{\ell}\varphi_{\ell}\mathbf{p}_{o}(\tilde{\beta})\right) + \frac{m_{o}}g_{i}(\tilde{\beta}_{o})} + \\ &= \left(m_{o}\frac{1}{4\sigma}\tilde{\beta}_{o}\left(-J_{r}^{r}\Gamma_{i}r_{i}^{r} + J_{i}^{\ell}\Gamma_{r}r_{i}^{r}\right) - m_{o}\tilde{\beta}_{o}} + \frac{\sigma_{\ell}}}{2}\tilde{\beta}_{o} + m_{o}J_{i}\left(\frac{1}{4\sigma^{2$$

(5.17)

We now plug the explicit formulas for σ and β_o into the above expressions. Such a replacement is straightforward, but very arduous and it is convenient to use a symbolic manipulation program, such as *Maple*, to perform it. In this way the equations $\operatorname{Ric}^{(m_o)}(\hat{E}_i, \mathbf{q}_o) = 0$, i = 1, 2, become equivalent to the following equations on $\mathbf{k} = \mathbf{k}(x, y)$:

$$\left(B\mathbf{k}\varphi(B+1)(B\varphi-r)(B\varphi+r)\frac{\partial\varphi}{\partial x}+5r(B+1)\mathbf{k}\left(B^{2}\varphi^{2}-\frac{r^{2}}{5}\right)\frac{\partial\varphi}{\partial y}+\left((B^{2}\varphi^{2}-r^{2})\frac{\partial\mathbf{k}}{\partial x}-2Br\varphi\frac{\partial\mathbf{k}}{\partial y}\right)(B^{2}\varphi^{2}+r^{2})\right)\frac{1}{2(B^{2}\varphi^{2}+r^{2})^{3}}=0,\quad(5.18)$$

$$\left(B\mathbf{k}\varphi(B+1)(B\varphi-r)(B\varphi+r)\frac{\partial\varphi}{\partial y} - 5(B+1)\mathbf{k}r\left(B^{2}\varphi^{2} - \frac{r^{2}}{5}\right)\frac{\partial\varphi}{\partial x} + \left(\frac{\partial\mathbf{k}}{\partial y}(B^{2}\varphi^{2} - r^{2}) + 2Br\varphi\frac{\partial\mathbf{k}}{\partial x}\right)(B^{2}\varphi^{2} + r^{2})\right)\frac{1}{2(B^{2}\varphi^{2} + r^{2})^{3}} = 0. \quad (5.19)$$

The numerators of the left hand sides of these equations are polynomials of order 4 in the variable r. The monomials in r of highest degree in these equations have coefficients $-\frac{\partial \mathbf{k}}{\partial x}$ and $\frac{\partial \mathbf{k}}{\partial y}$, respectively. Since the equations are satisfied for any value of r only if such coefficients vanish identically, it follows that they are satisfied only if $\mathbf{k}(x, y)$ is constant. If we now assume that \mathbf{k} is a non-zero constant ($\mathbf{k} \neq 0$ because otherwise $\tilde{\beta}_o$ would be 0 and $g^{(m_o)} = \eta$), the equations (5.18) and (5.19) reduce to

$$(B+1)\mathbf{k}\left(B\varphi(B^{2}\varphi^{2}-r^{2})\frac{\partial\varphi}{\partial x}+r\left(5B^{2}\varphi^{2}-r^{2}\right)\frac{\partial\varphi}{\partial y}\right)=0,$$

$$(B+1)\mathbf{k}\left(B\varphi(B^{2}\varphi^{2}-r^{2})\frac{\partial\varphi}{\partial y}-r\left(5B^{2}\varphi^{2}-r^{2}\right)\frac{\partial\varphi}{\partial x}\right)=0.$$
(5.20)

The left hand sides of these equations are now polynomials in r of order 3 and they vanish identically if and only if either $\varphi = \varphi(x, y)$ is constant (but this cannot be because φ is a potential of a Kähler metric) or B = -1. We conclude that all equations in (5.2), (5.3) and (5.4) with the only exception of $\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o) = 0$, are satisfied by a $g^{(m_o)} \neq \eta$ if and only if $\tilde{\beta}_o(x, y)$ is as in (5.1) and B = -1.

It remains to check that if $\tilde{\beta}_o$ and B are as above, the equation $\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o) = 0$ is identically satisfied. As usual, let us expand $\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o)$ in terms of the $\widetilde{\Gamma}_{AB}^{\ \ C} = \Gamma_{AB}^{\ \ C} + \mathbf{H}_{AB}^{\ \ C}$. We get

Since (5.12) implies $p_o(p_o(\tilde{\beta}_o)) + p_o(\tilde{\beta}_o) \frac{p_o(\sigma)}{\sigma} + \frac{\tilde{\beta}_o}{4\sigma^2} = 0$, the sum of the three terms in (5.21) that are proportional to m_o^2 vanishes identically, while the sum of all other terms in the last two lines of that expression reduces to $m_o B \frac{\tilde{\beta}_o}{8\sigma^2}$. We therefore get

$$\begin{split} \operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o) &= \\ &= -m_o \hat{E}_r \left(\frac{g^{rk}}{4\sigma} \tilde{\beta}_{ok} + B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_{\ell} \mathbf{p}_o(\tilde{\beta}_o) \right) - m_o \left(\frac{g^{\ell k}}{4\sigma} \tilde{\beta}_{ok} + B \frac{g^{\ell k}}{4\sigma} J_k^t \varphi_{\ell} \mathbf{p}_o(\tilde{\beta}_o) \right) \mathbf{\Gamma}_{r\ell}^r + m_o B \frac{\tilde{\beta}_o}{8\sigma^2} = \\ &= -m_o \frac{\tilde{\beta}_{ok}}{4\sigma} E_r(g^{rk}) + \overline{m_o g^{rk}} \frac{\sigma_r}{4\sigma^2} \tilde{\beta}_{ok} + m_o B g^{rk} J_r^* \varphi_s \frac{\mathbf{p}_o(\sigma)}{4\sigma^2} \tilde{\beta}_{ok} - m_o \frac{g^{rk}}{4\sigma^2} \tilde{\beta}_{okr} - m_o B \frac{g^{rk}}{4\sigma} J_r^s \varphi_s \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B E_r(g^{rk}) \frac{J_k^\ell}{4\sigma} \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) + m_o B \frac{g^{rk}}{4\sigma^2} J_r^* \varphi_s \mathcal{F}_{d}(\tilde{\beta}_o) + m_o B^2 \frac{g^{rk}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\sigma) J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_{\ell} \mathbf{p}_o(\tilde{\beta}_o) - m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell E_r \left(\mathbf{p}_o(\tilde{\beta}_o) \right) - m_o B^2 \frac{g^{rk}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\sigma) J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) - m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell E_r \left(\mathbf{p}_o(\tilde{\beta}_o) \right) - m_o B^2 \frac{g^{rk}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\sigma) - \\ &- m_o B \frac{g^{\ell k}}{4\sigma} \tilde{\beta}_{ok} \Gamma_r^r - \overline{m_o B} \frac{g^{\ell k}}{4\sigma^2} \tilde{\beta}_{ok} - m_o B \frac{g^{\ell k}}{4\sigma^2} J_\ell^* \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{\ell k}}{4\sigma} J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) \Gamma_r^r - m_o B \frac{g^{\ell k}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\tilde{\beta}_o) \sigma_\ell - \overline{m_o B^2 \varphi_\ell J_\ell^*} \frac{g^{\ell k}}{4\sigma^2} J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) \Gamma_r^r - m_o B \frac{g^{\ell k}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\tilde{\beta}_o) \sigma_\ell - m_o B^2 \frac{g^{rk}}{4\sigma^2} J_k^\ell \varphi_\ell \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell \mathbf{p}_o(\tilde{\beta}_o) \Gamma_r^r - m_o B \frac{g^{rk}}{4\sigma^2} J_r^* \varphi_s \mathbf{p}_o(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_k^\ell \varphi_\ell \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{rk}}{4\sigma} J_r^\ell \varphi_\sigma \mathbf{p}_\sigma(\tilde{\beta}_o) - \\ &- m_o B \frac{g^{r$$

We now recall that, for each $k, r, \ell = 1, 2$, the expression $E_r(g^{rk}) + g^{\ell k} \Gamma_{r\ell}^{r}$ is identically 0. One way (out of many others) to check this is by using the isothermal coordinates (x, y) on \mathcal{V} , with $E_1 = \frac{\partial}{\partial x}$, $E_2 = \frac{\partial}{\partial y}$, and $g^{rs} = \frac{1}{\lambda} \delta^{rs}$, $\lambda = \Delta \varphi$. The Christoffel symbols $\Gamma_{r\ell}^{s}$ of the metric g_o in these coordinates are given by (B.2). Using this information, a straightforward check shows that $E_r(g^{rk}) + g^{\ell k} \Gamma_{r\ell}^{\ r} = 0$, which we plug into into (5.22) to get

$$\operatorname{Ric}^{(m_{o})}(\mathbf{q}_{o},\mathbf{q}_{o}) = \\ = -m_{o}\frac{g^{rk}}{4\sigma}\widetilde{\beta}_{okr} - m_{o}B\frac{g^{rk}}{2\sigma}J_{r}^{s}\varphi_{s}\mathbf{p}_{o}(\widetilde{\beta}_{ok}) - m_{o}B\frac{g^{rk}}{4\sigma}J_{k}^{\ell}\varphi_{\ell r}\mathbf{p}_{o}(\widetilde{\beta}_{o}) - m_{o}B^{2}\frac{g^{rk}}{4\sigma}J_{k}^{\ell}J_{r}^{s}\varphi_{\ell}\varphi_{s}\mathbf{p}_{o}\left(\mathbf{p}_{o}(\widetilde{\beta}_{o})\right) + \frac{m_{o}B}{8\sigma^{2}}\widetilde{\beta}_{o} = \\ = -\frac{m_{o}}{4\lambda\sigma}\widetilde{\beta}_{oxx} - \frac{m_{o}}{4\lambda\sigma}\widetilde{\beta}_{oyy}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_{2\lambda\sigma}\underbrace{-\frac{m_{o}B}{2\lambda\sigma}}_$$

We may now use the explicit expressions (4.3) and (5.16) for σ and $\tilde{\beta}_o$ in terms of $\varphi(x, y)$ and **k**. With the help of a symbolic manipulation program such as *Maple*, one can immediately check that these substitutions yield the identity

$$\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o) = m_o \frac{2r\varphi^2(B+1)}{(r^2 + \varphi^2)^3} .$$
(5.24)

Since B = -1, it follows that $\operatorname{Ric}^{(m_o)}(\mathbf{q}_o, \mathbf{q}_o) \equiv 0$, as we needed to prove. The computations of the three steps show also that there is no constraint on m_o and that if a metric $g^{(m_o)}$ is Ricci flat, then automatically any other metric $g^{(m)}$, $m \neq m_o$, is Ricci flat.

6. EXPLICIT EXPRESSIONS OF RICCI FLAT KERR FAMILIES AND REAL ANALYTIC EXTENSIONS OF SOME OF SUCH METRICS

6.1. Potentials for Kerr type backgrounds and corresponding Kerr families. By Corollary 4.3, if M is a Kerr 4-manifold admitting a Kerr type background, then the underlying Riemann surface (N, J) is non-compact, admitting a Kähler metric \tilde{g}_o of constant Gaussian curvature $\kappa = \pm 1$ and equipped with a Kähler metric g_o with a global potential $\varphi : N \to \mathbb{R}$ satisfying (4.2). In suitable isothermal coordinates (x, y) for \tilde{g}_o and g_o in an open subset $\mathcal{V} \subset N$, the conditions (4.2) are equivalent to either

$$\Delta \varphi + \frac{8\varphi}{(1 + (x^2 + y^2))^2} = 0 \quad \& \quad \varphi < 0 \qquad \text{or to}$$
 (6.1)

$$\Delta \varphi - \frac{8\varphi}{\left(1 - (x^2 + y^2)\right)^2} = 0 \quad \& \quad \varphi > 0 \;. \tag{6.2}$$

The solutions to these equations on bounded open sets have the following properties.

Lemma 6.1. Let \mathcal{V} be a relatively compact region in $\mathbb{D} = \{|z|^2 < 1\} \subset \mathbb{C} = \mathbb{R}^2$ with boundary of class $\mathscr{C}^{k,\alpha}$, $k \ge 2$, $\alpha \in (0,1)$. Any solution to (6.1) (resp. (6.2)) on $\overline{\mathcal{V}}$ is real analytic on \mathcal{V} and the correspondence

$$\varphi \longmapsto f = \varphi|_{\partial \mathcal{V}}$$

is a bijection between the solutions φ that are in $\mathscr{C}^{\omega}(\mathcal{V}) \cap \mathscr{C}^{k,\alpha}(\overline{\mathcal{V}})$ and the non identically vanishing functions $f \leq 0$ (resp. $f \geq 0$) that are in $\mathscr{C}^{k,\alpha}(\partial \mathcal{V})$.

Proof. By [12, Thms. 3.5, 6.13 & 6.19] and by [14] (see also [21]), there exists a unique solution to (6.2) on $\overline{\mathcal{V}}$ of class $\mathscr{C}^{\omega}(\mathcal{V}) \cap \mathscr{C}^{k,\alpha}(\overline{\mathcal{V}})$ for any prescribed boundary value $f = \varphi|_{\mathcal{V}}$ in $\mathscr{C}^{k,\alpha}(\partial \mathcal{V})$ and such a solution satisfies the Maximum Principle. This implies the claim for the conditions (6.2). This argument does not work for (6.1) because, in this case, the zero order term is positive and [12, Thms. 3.5 & 6.13] do not apply. Nonetheless, we may consider the special negative solution to (6.1)

$$\varphi_o : \mathbb{D} \to (0, +\infty) , \qquad \varphi_o(x, y) := -\frac{1 - x^2 - y^2}{1 + x^2 + y^2}$$
 (6.3)

and observe that the correspondences $\varphi \mapsto \widetilde{\varphi} := \frac{\varphi}{\varphi_o}$ and $\widetilde{\varphi} \mapsto \varphi = \widetilde{\varphi}\varphi_o$ establish a bijection between the solutions to (6.1) and the solutions to

$$\Delta \widetilde{\varphi} + \frac{2}{\varphi_o} \left(\frac{\partial \varphi_o}{\partial x} \ \frac{\partial \widetilde{\varphi}}{\partial x} + \frac{\partial \varphi_o}{\partial y} \ \frac{\partial \widetilde{\varphi}}{\partial y} \right) = 0 \tag{6.4}$$

on $\overline{\mathcal{V}} \subset \mathbb{D}$. Since the previous argument now works for (6.4), the lemma is proved for (6.1) as well.

Lemma 6.1, Corollary 4.3 and Theorem 5.1 immediately yield the following

Corollary 6.2. Any background metric η of Kerr type and any metric $g^{(m)}$, $m \neq 0$, of a Kerr family of gravitational fields on a Kerr 4-manifold M are real analytic in the coordinates (x, y, v, r) considered in Corollary 4.3 and Theorem 5.1. In particular, each such metric is uniquely determined by its restriction to any open subset $\mathcal{V} \subset N$ of the open cover $\{\mathcal{V}_A\}$ of N, described in Corollary 4.3.

We note that, as a direct consequence of [23], the potentials that satisfy (6.1) (resp. (6.2)) on a disk $\overline{\mathbb{D}}(0, \mathfrak{r}_o) := \{|z| \leq \mathfrak{r}_o\}, 0 < \mathfrak{r}_o < 1$, are real analytic up to the boundary, thus in oneto-one correspondence with the Fourier series of real analytic boundary data on $\partial \mathbb{D}(0, \mathfrak{r}_o)$. In fact, using polar coordinates (\mathfrak{r}, w) with $(x = \mathfrak{r} \cos w, y = \mathfrak{r} \sin w)$ and considering the Fourier expansion $f(w) = a_0 + \sum_{n=1}^{\infty} (a_n \cos(nw) + b_n \sin(nw))$ of a non-negative real analytic function f(w) on $\partial \mathbb{D}(0, \mathfrak{r}_o), \mathfrak{r}_o < 1$, one can check that the corresponding solution φ to (6.1) (resp. (6.2)) is the sum of the series

$$\varphi(\mathbf{r}, w) = -\left(a_0 \frac{\varphi_0(\mathbf{r})}{\varphi_0(\mathbf{r}_o)} + \sum_{n=1}^{\infty} \frac{\varphi_n(\mathbf{r})}{\varphi_n(\mathbf{r}_o)} (a_n \cos(nw) + b_n \sin(nw))\right) ,$$

where $\varphi_n(\mathbf{r}) := \left(1 - \frac{2\mathbf{r}^2}{(1+\mathbf{r}^2)} \frac{1}{n+1}\right) \mathbf{r}^n$ (6.5)
(resp. $\varphi_n(\mathbf{r}) := \left(\frac{1+\mathbf{r}^2}{1-\mathbf{r}^2} + n\right) \mathbf{r}^n \right) .$

Each of such solutions can be conveniently expressed in terms of a single holomorphic function as follows. Using the complex coordinate $z = \mathbf{r}e^{iw}$ and setting $c_0 := \frac{-a_0}{\varphi_0(\mathbf{r}_0)}$ and

$$c_n := \frac{-(a_n - ib_n)}{\varphi_0(\mathfrak{r}_o)}, \ n \ge 1, \text{ we have that } (6.5) \text{ can be re-written as}$$
$$\varphi(z) = \operatorname{Re}\left(\sum_{n=0}^{\infty} \left(1 - \frac{2|z|^2}{(1+|z|^2)} \frac{1}{n+1}\right) c_n z^n\right) \qquad \text{if } \kappa = 1 ,$$

$$\varphi(z) = \operatorname{Re}\left(\sum_{n=0}^{\infty} \left(\frac{1+|z|^2}{1-|z|^2}+n\right)c_n z^n\right) \quad \text{if } \kappa = -1$$

Considering the holomorphic function $F(z) := \sum_{n=0}^{\infty} c_n z^n$, we get that

$$\varphi(z) = \operatorname{Re}\left(F(z) - \frac{2|z|^2}{1+|z|^2} \frac{1}{z} \int_0^z F(\zeta) d\zeta\right) \quad \text{if } \kappa = 1 ,$$

$$\varphi(z) = \operatorname{Re}\left(\frac{1+|z|^2}{1-|z|^2} F(z) + z \frac{dF(z)}{dz}\right) \quad \text{if } \kappa = -1 .$$
(6.6)

Conversely, for any holomorphic function F, for which the corresponding function in (6.6) is non-positive or non-negative (according to $\kappa = 1$ or -1), determines a solution $\varphi(z)$ to (6.1) or (6.2), respectively.

6.2. Explicit examples of Ricci flat Kerr families of gravitational fields. Let $M = (N \times \mathbb{R}) \times \mathbb{R}$ (or $M = (N \times \mathbb{R}) \times I$, $I \subset \mathbb{R}$) be a Kerr 4-manifold admitting a Kerr family of Ricci flat gravitational fields $g^{(m)}$. As usual, we denote by (N, J, g_o) the (non-compact) Riemann surface with Kähler metric g_o which underlies the Kerr manifold.

Note that, denoting by \tilde{g}_o the Kähler metric on (N, J) with constant Gaussian curvature $\kappa = \pm 1$ that corresponds to the family $g^{(m)}$ by Theorem 5.1, using the relation $g_o = -2\kappa\varphi\tilde{g}_o$, the metrics of the given family $g^{(m)}$ have locally the form

$$g^{(m)} = -\frac{\kappa}{2} \left(r^2 + \varphi^2\right) \widetilde{g}_o + \left(dv + d^c \varphi\right) \vee \left(d(v+r) + \frac{1}{2} \left(-1 + \frac{\mathbf{k}mr}{r^2 + \varphi^2}\right) \left(dv + d^c \varphi\right)\right), \quad (6.7)$$

where:

- **k** is a scaling constant (for the Kerr metrics (1.1), such a constant is $\mathbf{k} = 2$);
- φ is the global Kähler potential for g_o , defined in claim (2) of Theorem 4.1;
- r is a coordinate for the standard fiber of the trivial bundle $\pi^{\mathbb{S}}: M = \mathbb{S} \times \mathbb{R} \to N \times \mathbb{R};$
- $v : \mathcal{U} \subset M \to \mathbb{R}$ is a locally defined function, which completes a triple (x, y, r), given by a pair of local coordinates (x, y) for $\mathcal{V} \subset N$ and the globally defined function r, to a coordinate chart $(x, y, v, r) : \mathcal{U} \subset M \to \mathbb{R}^4$, in which $q_o = \frac{\partial}{\partial v} + B \frac{\partial}{\partial r}$.

Note also that the real function $v : \mathcal{U} \to \mathbb{R}$ is a local potential for the globally defined closed 1-form $\vartheta - d^c \varphi$ (i.e., $dv = \vartheta - d^c \varphi$). Thus, in case (N, J) (and consequently also $M = N \times \mathbb{R} \times \mathbb{R}$) is simply connected, such local potential v can be extended to a globally defined potential $v : N \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$. On the other hand, if (N, J) is *not* simply connected, several non equivalent connection 1-forms θ satisfying $d\theta = \pi^S \omega_o$ might occur, each of them corresponding to non equivalent local potentials v. For an exhaustive discussion of such non-equivalent possibilities, see e.g. [20]. Plugging explicit coordinate expressions for solutions φ to (6.1) and (6.2) into (6.7), one gets a huge number of explicit coordinate expressions for Lorentzian Ricci flat metrics $g^{(m)}$. In the next two subsections, we briefly discuss a few features of these metrics.

6.2.1. A quick look at the metrics of the Kerr families with $\kappa = 1$. Consider the expressions (6.5) for possible potentials φ in case $\kappa = 1$. There, they are given in terms of the polar coordinates (\mathfrak{r}, w) associated with a pair of isothermal coordinates $(x = \mathfrak{r} \cos w, y = \mathfrak{r} \sin w)$ in which \tilde{g}_o takes the form $\tilde{g}_o = \frac{4}{(1+(x^2+y^2))^2}(dx^2+dy^2)$. Replacing (\mathfrak{r}, w) by coordinates (ξ, ψ) defined by the relations $\mathfrak{r} = \tan\left(\frac{\xi}{2}\right), w = \psi$, the metric \tilde{g}_o and the potentials φ take the form

$$\widetilde{g}_{o} = d\xi^{2} + \sin^{2}\xi d\psi^{2} ,$$

$$\varphi(\xi, \psi) = \widetilde{a}_{0}\cos\xi + \widetilde{a}_{1}\sin\xi\cos\psi + \widetilde{b}_{1}\sin\xi\sin\psi +$$

$$+ \sum_{n=2}^{\infty} \frac{1 + \frac{n-1}{n+1}\tan^{2}\left(\frac{\xi}{2}\right)}{1 + \tan^{2}\left(\frac{\xi}{2}\right)} \tan^{n}\left(\frac{\xi}{2}\right) \left(\widetilde{a}_{n}\cos(n\psi) + \widetilde{b}_{n}\sin(n\psi)\right)$$
(6.8)

for appropriate constants $\tilde{a}_n, \tilde{b}_n \ge 0$. In this way, \tilde{g}_o takes the usual form of the standard metric of S^2 in spherical coordinates and φ can be (locally) identified with a function on an open subset of the unit sphere.

Let us now introduce the following notation. Given a pair of real numbers (α_1, α_2) , let $R^{(\alpha_1, \alpha_2)}$ be the isometry of (S^2, \tilde{g}_o) given by the double rotation

$$R^{(\alpha_1,\alpha_2)} : S^2 \longrightarrow S^2 ,$$

$$R^{(\alpha_1,\alpha_2)} \begin{pmatrix} y^1 \\ y^2 \\ y^3 \end{pmatrix} := \begin{pmatrix} \cos \alpha_2 & -\sin \alpha_2 & 0 \\ \sin \alpha_2 & \cos \alpha_2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \cos \alpha_1 & 0 & -\sin \alpha_1 \\ 0 & 1 & 0 \\ \sin \alpha_1 & 0 & \cos \alpha_1 \end{pmatrix} \cdot \begin{pmatrix} y^1 \\ y^2 \\ y^3 \end{pmatrix} .$$

We may thus consider the corresponding "rotated" spherical coordinates $(\xi' := \xi(\alpha_1, \alpha_2), \psi' := \psi(\alpha_1, \alpha_2))$ of S^2 , i.e. the coordinates that are related with the usual spherical coordinates (ξ, ψ) by

$$\begin{pmatrix} \sin\xi'\cos\psi'\\ \sin\xi'\sin\psi'\\ \cos\xi' \end{pmatrix} = R^{(\alpha_1,\alpha_2)} \begin{pmatrix} \sin\xi\cos\psi\\ \sin\xi\sin\psi\\ \cos\xi \end{pmatrix}.$$
(6.9)

In different words, (ξ', ψ') are just new spherical coordinates, which one gets after an appropriate replacement of the north-south axis and of the reference meridian $\psi = 0$.

A straightforward check shows that for any $\varphi(\xi, \psi) = \tilde{a}_0 \cos \xi + \tilde{a}_1 \sin \xi \cos \psi + \tilde{b}_1 \sin \xi \sin \psi$ (i.e. a potential (6.8) with $\tilde{a}_n, \tilde{b}_n = 0, n \ge 2$), there exists a choice of "rotated" spherical coordinates $(\xi' := \xi(\alpha_1^o, \alpha_2^o), \psi' := \psi(\alpha_1^o, \alpha_2^o))$ in which φ takes the form

$$\varphi(\xi',\psi') = a\cos\xi' , \quad \text{with} \quad a = \sqrt{\widetilde{a}_0^2 + \widetilde{a}_1^2 + \widetilde{b}_1^2} . \quad (6.10)$$

Plugging (6.10) into (6.7), we see that the 3-parameter family of (local) potentials φ with $\tilde{a}_n = \tilde{b}_n = 0$, $n \ge 2$, correspond to Lorentzian metrics that are locally isometric to the classical Kerr metrics (1.1). The parameters of such a family of potentials (i.e. the constants $\tilde{a}_0, \tilde{a}_1, \tilde{b}_1$) correspond to the three space-like components (with respect to a fixed frame) of the total angular momentum $\vec{\mathcal{J}} = m\vec{a}$ of the gravitational field.

From this we get that, whenever \tilde{a}_n , \tilde{b}_n , $n \ge 2$, are equal to 0, the corresponding metrics – which a priori are defined just on an open subset of $S^2_{(+)} \times \mathbb{R}_+ \times \mathbb{R}$ – extend in a real analytic way to the whole $(S^2 \times \mathbb{R}_+) \times \mathbb{R} = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$. All such extensions are isometric to the classical Kerr metrics for appropriate values for m and $\mathcal{J} = m |\vec{a}|$. On the contrary, for nonzero choices of a finite collection of constants $\tilde{a}_{n_\ell}, \tilde{b}_{n_\ell}, 2 \le n_1 < \ldots < n_s$, by just looking at the terms corresponding to the highest integer n_s , one realises that the corresponding metrics $g^{(m)}$ do not extend to the whole $(\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$. Actually, for a few examples of this kind, the explicit computations of the Carminati-McLenaghan invariants of the Riemann curvature [7] indicate that such metrics develop intrinsic curvature singularities in approaching the south pole of S^2 and it is reasonable to expect that this occurs for all of them.

These remarks give evidence that the potentials (6.8) provide a very large number of Ricci flat metrics that are not locally isometric to any of the classical Kerr metrics. It would be interesting to study their geometric properties and explore the possibilities to generalise and/or glue them with appropriate solutions of the Einstein equations in order to obtain new solutions to the Einstein equations, representing gravitational fields with non-zero total angular momentum and determined by stress-energy tensors that vanish on large (but not spherically symmetric) regions of the space-time. We hope to address these issues in the near future.

As we mentioned in the Introduction, it is reasonable to expect that also when there is an *infinite* collection of non-vanishing coefficients $\tilde{a}_{n_{\ell}}$, $\tilde{b}_{n_{\ell}} \neq 0$, $n_{\ell} \geq 2$, the corresponding metric admits no real analytic extension to $(\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$. A proof of this quite sensible conjecture would give a new characterisation of the classical Kerr metrics (they would be the only metrics (6.7) that are extendible to $(\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}$), with interesting relations with the so far known uniqueness theorems for the Kerr metrics (see e.g. [28]).

6.2.2. A quick look at the metrics of the Kerr families with $\kappa = -1$. Let us now focus on the potentials in (6.5) for $\kappa = -1$. Replacing the polar coordinates (\mathfrak{r}, w) by the coordinates (ξ, ψ) given by the relations $\mathfrak{r} = \tanh\left(\frac{\xi}{2}\right)$, $w = \psi$, the metric \tilde{g}_o and the potentials φ take the form

$$\widetilde{g}_{o} = d\xi^{2} + \sinh^{2} \xi d\psi^{2} ,$$

$$\varphi(\xi, \psi) = \widetilde{a}_{0} \cosh \xi + \widetilde{a}_{1} \sinh \xi \cos \psi + \widetilde{b}_{1} \sinh \xi \sin \psi +$$

$$+ \sum_{n=2}^{\infty} \tanh^{n} \left(\frac{\xi}{2}\right) \left(\cosh \xi + n - 1\right) \left(\widetilde{a}_{n} \cos(n\psi) + \widetilde{b}_{n} \sin(n\psi)\right) .$$
(6.11)

Note also that, if the coefficients \tilde{a}_{ℓ} are taken sufficiently small so that φ is sufficiently close to 0 on some relatively compact subset of \mathbb{D} , the corresponding Lorentzian metric is close to the metric (6.7) with $\varphi = 0$, namely to

$$g^{(m)} = \frac{r^2}{2} (d\xi^2 + \sinh^2 \xi d\psi^2) + dv \vee \left(dr + \frac{r+m}{2r} dv \right) , \qquad (6.12)$$

This metric can be taken as an analog of the classical Schwarzschild metric, but with the crucial difference of having the group $\mathbb{R} \times SO_{1,2}^o$ instead of $\mathbb{R} \times SO_3$ as connected component of the group of isometries. This fact gives evidence that for potentials sufficiently close to $\varphi = 0$, if $\kappa = -1$, the corresponding Lorentzian metrics cannot be locally isometric to the classical Kerr metrics. A rigorous proof would however demand a detailed study of the isometric invariants of those metrics, a task which goes beyond the scope of this paper.

For what concerns the geometric properties, the Ricci flat metrics (6.7) with $\kappa = -1$ mirror certain crucial features of those with $\kappa = 1$. Indeed, in both settings, each metric is invariant under the action of the vector field $\mathbb{T} := q_o - Bp_o$. By the discussion in Remark 2.11, with respect to the background metric η , the vector field \mathbb{T} is time-like if $\kappa = 1$, while it is space-like if $\kappa = -1$. Moreover, passing from the case $\kappa = 1$ to the case $\kappa = -1$, according to (2.13) the signature of η changes from (-, -, -, +) to (+, +, -, +). This means that the roles of the last two coordinates appearing in (2.13) switch from space-like to time-like and vice versa.

This phenomenon greatly helps in guessing features of the metrics with $\kappa = -1$. For instance, let us first recall that the classical Kerr metrics (i.e. the cases with $\kappa = 1$ and $\tilde{a}_n = \tilde{b}_n = 0, n \ge 2$) determine distinguished hypersurfaces (characterised by an equation in the space-like coordinate r) which are time-invariant and have the fundamental role of being *event horizons*. According to the above described switch between time-like and spacelike roles, it is reasonable to expect that the analogs in the class with $\kappa = -1$ determine distinguished hypersurfaces (characterised by an equation in the *time-like* coordinate r), which are invariant under a space-like vector field and with features similar to those of the event horizons, provided that the meanings of the words "time" and "radius" are switched at all places.

Guided by this kind of intuition, while the Kerr metrics tend to stationary spherically symmetric weak gravitational fields at large distances, the newly determined Ricci flat metrics on $M^{\text{Lob}} := (\mathbb{D} \times \mathbb{R}) \times \mathbb{R}$ are expected to tend to radially-independent small deformations of the Minkowski metrics for large values of the "time". Being not asymptotically spherically symmetric, these Ricci flat metrics are not appropriate to model the gravity generated by a localised mass. Nonetheless, they might be considered as non-static cosmological models or, more precisely, as geometric limits (for the stress-energy tensors tending to 0) of a large class of non-static cosmological models.

Appendix A. Details of the computations that yield (3.20)

Let us recall the following properties and relations between the Christoffel symbols and components J_i^i of the complex structure:

- (a) $J_m^m = 0$ (it is the trace of a complex structure);
- (b) $\Gamma_{ij}^r \Gamma_{ji}^r c_{ij}^r = 0$ (Γ_{ij}^r are Christoffel symbols of the torsion free connection ∇^{g_o}); (c) $E_m(J_i^\ell) = -\Gamma_{mr}^{\ \ell} J_i^r + \Gamma_{mi}^{\ r} J_r^\ell$, (in fact, since (N, J, g) is Kähler and hence $\nabla^{g_o} J = 0$, we have that

$$E_m(J_i^\ell) = \left(\nabla_{E_m}^{g_o} E^\ell\right)(J(E_i)) + E^\ell\left(\left(\nabla_{E_m}^{g_o} J\right)(E_i)\right) + E^\ell\left(J(\nabla_{E_m}^{g_o} E_i)\right) = -\Gamma_{mr}^{\ell} J_i^r + \Gamma_{mi}^r J_r^\ell\right).$$

Using (3.4), (3.7) - (3.13) and the above relations (a) – (c), the Ricci curvature components (3.19) can be written as

$$\begin{split} \operatorname{Ric}(\hat{E}_{i},\mathbf{p}_{o}) &= \hat{E}_{i}(\mathbf{\Gamma}_{mp_{o}}^{m}) - \mathbf{\Gamma}_{mp_{o}}^{k} \mathbf{\Gamma}_{i}^{m} + \mathbf{\Gamma}_{tp_{o}}^{k} \mathbf{\Gamma}_{m}^{m} - \mathbf{c}_{mi}^{m} \mathbf{\Gamma}_{p_{o}}^{m} = \\ &= E_{m}\left(\frac{J_{i}^{m}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{i}^{m}}{2\sigma}\right) + BJ_{m}^{\ell}\varphi_{\ell}\mathbf{p}_{o}\left(\frac{J_{i}^{m}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{m}^{m}}{2\sigma}\right) - \\ &\quad - E_{i}\left(\frac{J_{m}^{m}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{m}^{m}}{2\sigma}\right) \mathbf{\Gamma}_{i}^{m} + \left(\frac{J_{i}^{\ell}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{i}^{m}}{2\sigma}\right) \mathbf{\Gamma}_{m\ell}^{m} - \mathbf{c}_{mi}^{r}\mathbf{\Gamma}_{p_{o}}^{m} = \\ &\quad - \left(\frac{J_{m}^{m}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{m}^{m}}{2\sigma}\right) \mathbf{\Gamma}_{i}^{m} + \left(\frac{J_{i}^{\ell}}{4\sigma} + \mathbf{p}_{o}(\sigma)\frac{\delta_{i}^{j}}{2\sigma}\right) \mathbf{\Gamma}_{m\ell}^{m} - \mathbf{c}_{mi}^{r}\mathbf{\Gamma}_{p_{o}}^{m} = \\ &= \frac{E_{m}(J_{i}^{m})}{4\sigma} - \frac{J_{i}^{m}\sigma_{m}}{4\sigma^{2}} + \frac{1}{2\sigma}E_{i}\left(\mathbf{p}_{o}(\sigma)\right) - \frac{1}{2\sigma^{2}}\mathbf{p}_{o}(\sigma)- \\ &\quad - BJ_{i}^{m}J_{m}^{\ell}\varphi_{\ell}\frac{D\rho_{o}^{2}}{4\sigma^{2}} + BJ_{i}^{\ell}\varphi_{\ell}\frac{1}{2\sigma}\mathbf{p}_{o}\left(\mathbf{p}_{o}(\sigma)\right) - BJ_{i}^{\ell}\varphi_{\ell}\frac{1}{2\sigma^{2}}\left(\mathbf{p}_{o}(\sigma)\right)^{2} - \\ &\quad - E_{i}\left(\mathbf{p}_{o}(\sigma)\right)\frac{(n-2)}{2\sigma} + \sigma_{i}\mathbf{p}_{o}(\sigma)\frac{(n-2)}{2\sigma^{2}} - \\ &\quad - BJ_{i}^{t}\varphi_{\ell}\left(\frac{n-2}{2\sigma}\right)\mathbf{p}_{o}\left(\mathbf{p}_{o}(\sigma)\right) + BJ_{i}^{\ell}\varphi_{\ell}\left(\frac{n-2}{2\sigma^{2}}\right)\left(\mathbf{p}_{o}(\sigma)\right)^{2} + \\ &\quad - \frac{J_{m}^{\ell}}{4\sigma}\mathbf{\Gamma}_{i}^{m} - \mathbf{p}_{o}(\sigma)\frac{1}{2\sigma}\mathbf{\Gamma}_{i}^{m} + \frac{J_{i}^{\ell}}{4\sigma}\mathbf{\Gamma}_{m}^{m} + \mathbf{p}_{o}(\sigma)\frac{1}{2\sigma}\mathbf{\Gamma}_{mi}^{m} - \mathbf{c}_{mi}^{r}\mathbf{\Gamma}_{p_{o}}^{m} = \\ &= \underbrace{-\frac{V_{mp}T_{i}}{4\sigma}\mathbf{\Gamma}_{i}^{T}} - \frac{J_{i}^{t}J_{i}^{T}}{4\sigma} - \underbrace{-\frac{J_{i}^{\ell}}{4\sigma^{2}} - (n-3)\frac{E_{i}\left(\mathbf{p}_{o}(\sigma)\right)}{2\sigma}\mathbf{\Gamma}_{i} + (n-3)\frac{P_{o}(\sigma)\sigma_{i}}{2\sigma^{2}} + \\ &\quad - \frac{J_{m}^{\ell}}{4\sigma^{2}}\mathbf{T}_{i}^{\ell} - \frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell} - \frac{J_{m}^{\ell}}{8\sigma^{2}} + \underbrace{-\frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell}} - \underbrace{-\frac{J_{m}^{\ell}}{3\sigma^{2}}\mathbf{T}_{i}^{\ell} - \frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell}} - \underbrace{-\frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell}} - \underbrace{-\frac{J_{m}^{\ell}}{4\sigma}\mathbf{T}_{i}^{\ell} - \frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell}} + \frac{Bp_{o}(\sigma)\varphi_{i}}{8\sigma^{2}} + \underbrace{-\frac{J_{m}^{\ell}}{8\sigma^{2}}\mathbf{T}_{i}^{\ell}} - \underbrace{-\frac{J_{m}^{\ell}}{4\sigma^{2}}\mathbf{T}_{i}^{\ell}} + \underbrace{-\frac{J_{m}^{\ell}}{4\sigma^{2}}\mathbf{T}_{i}^{\ell}} + \underbrace{-\frac{J_{m}^{\ell}}{4\sigma^{2}}\mathbf{T}_{i}^{\ell}} + \underbrace{-\frac{J_{m}^{\ell}}{4\sigma^{2}}\mathbf{T}_{i}^{\ell}} + \underbrace{-\frac{J_{m}^{\ell}}{4\sigma}\mathbf{T}_{i}^{\ell}} + \underbrace{-\frac{J_{m}^{\ell$$

$$+ \frac{J_{i}^{\ell} \Gamma_{mi}}{4\sigma} + \frac{J_{i}^{\ell} \sigma_{\ell}}{8\sigma^{2}} + \frac{(n-2)J_{i}^{\ell} \sigma_{\ell}}{8\sigma^{2}} - \frac{J_{i}^{\ell} \sigma_{\ell}}{8\sigma^{2}} - \frac{(n-2)Bp_{o}(\sigma)\varphi_{i}}{8\sigma^{2}} + \\ + \frac{p_{o}(\sigma)\Gamma_{mi}}{2\sigma} + \frac{p_{o}(\sigma)\sigma_{i}}{4\sigma^{2}} + \frac{p_{o}(\sigma)\sigma_{i}(n-2)}{4\sigma^{2}} - \frac{p_{o}(\sigma)\sigma_{i}}{4\sigma^{2}} + \\ + \frac{(n-2)\cdot(p_{o}(\sigma))^{2}BJ_{i}^{\ell}\varphi_{\ell}}{4\sigma^{2}} - c_{mi}^{r}\frac{J_{r}^{m}}{4\sigma} - \frac{c_{mi}^{m}p_{o}(\sigma)}{2\sigma} = \\ = \frac{J_{\ell}}{4\sigma} \underbrace{\Gamma_{mi}}_{\ell} + \frac{e_{\ell}}{c_{mi}} + \frac{e_{\ell}}{c_{mi}} + (n-6)\frac{J_{i}^{\ell}\sigma_{\ell}}{8\sigma^{2}} - \frac{(n-3)E_{i}\left(p_{o}(\sigma)\right)}{2\sigma} + \frac{(n-3)p_{o}(\sigma)\sigma_{i}}{2\sigma^{2}} + \\ - \frac{(n-6)B\varphi_{i}p_{o}(\sigma)}{8\sigma^{2}} - \frac{(n-3)BJ_{i}^{\ell}\varphi_{\ell}}{2\sigma} p_{o}\left(p_{o}(\sigma)\right) + \frac{(n-3)BJ_{i}^{\ell}\varphi_{\ell}}{2\sigma^{2}}\left(p_{o}(\sigma)\right)^{2} + \\ + \frac{J_{\ell}}{4\sigma} \underbrace{\Gamma_{mi}}_{\ell} + \frac{e_{\ell}}{c_{mi}} - \frac{e_{mi}}{c_{mi}} - \frac{p_{o}(\sigma)}{2\sigma} \underbrace{\Gamma_{mi}}_{mi} + \frac{e_{mi}}{c_{mi}} - \frac{e_{mi}}{c_{mi}}$$

From the last equality and factoring out $\frac{n-3}{8\sigma^2}$, (3.20) follows.

Appendix B. The proof of Lemma 4.2

In this appendix we provide the details of the computations that prove Lemma 4.2. In what follows we denote $\lambda = \Delta \varphi$ and we use the notation $\delta \Gamma_{ij}^{\ k}$ for the differences $\delta \Gamma_{ij}^{\ k} = \Gamma_{ij}^{\ k} - \Gamma_{ij}^{\ k}$. Using the symmetry $\Gamma_{ij}^{\ k} = \Gamma_{ji}^{\ k}$ (because the E_i commute) and the identity

$$0 = E_i \left(g(E_j, E_k) \right) - g(\nabla_{E_i} E_j, E_k) - g(E_j, \nabla_{E_i} E_k) =$$

= $E_i(\lambda)\delta_{jk} - \lambda \Gamma_{ij}^{\ell} \delta_{\ell k} - \lambda \Gamma_{ik}^{\ell} \delta_{j\ell} = E_i(\lambda)\delta_{jk} - \lambda \Gamma_{ij}^{k} - \lambda \Gamma_{ik}^{j}$, (B.1)

we obtain the following expressions for the $\Gamma_{ij}^{\ k}$ and for the differences $\delta \Gamma_{ij}^{\ k}$

$$\Gamma_{11}^{11} = \frac{1}{2} \frac{\lambda_x}{\lambda} , \qquad \Gamma_{22}^{2} = \frac{1}{2} \frac{\lambda_y}{\lambda} , \qquad \Gamma_{21}^{11} = \Gamma_{22}^{22} , \qquad \Gamma_{12}^{2} = \Gamma_{11}^{11} ,$$

$$\Gamma_{12}^{11} = \Gamma_{22}^{22} , \qquad \Gamma_{11}^{2} = -\Gamma_{22}^{22} , \qquad \Gamma_{21}^{2} = \Gamma_{11}^{11} , \qquad \Gamma_{22}^{11} = -\Gamma_{11}^{11} .$$

$$\delta\Gamma_{11}^{11} = \frac{\sigma_x}{2\sigma} + B \frac{\sigma_t \varphi_y}{2\sigma} , \qquad \delta\Gamma_{22}^{22} = \frac{1}{2\sigma} \sigma_y - B \frac{\sigma_t \varphi_x}{2\sigma} , \qquad \delta\Gamma_{21}^{11} = \delta\Gamma_{22}^{22} , \qquad \delta\Gamma_{12}^{12} = \delta\Gamma_{11}^{11} ,$$

$$\delta\Gamma_{12}^{11} = \delta\Gamma_{22}^{22} , \qquad \delta\Gamma_{12}^{22} = -\delta\Gamma_{11}^{11} .$$

$$(B.2)$$

$$\delta\Gamma_{12}^{11} = \delta\Gamma_{22}^{22} , \qquad \delta\Gamma_{12}^{22} = -\Gamma_{11}^{11} .$$

$$\delta\Gamma_{12}^{11} = \delta\Gamma_{22}^{22} , \qquad \delta\Gamma_{12}^{22} = \delta\Gamma_{11}^{11} , \qquad \delta\Gamma_{22}^{12} = -\delta\Gamma_{11}^{11} .$$

$$(B.3)$$

Note that each of the Christoffel symbols Γ_{ij}^{k} (resp. Γ_{ij}^{k}) is (up to a sign) equal either to $\Gamma_{11}^{11} = \frac{\lambda_x}{\lambda}$ (resp. Γ_{11}^{11}) or to $\Gamma_{22}^{2} = \frac{\lambda_y}{\lambda}$ (resp. Γ_{22}^{2}). Let us now focus on the equation (4.11). Expanding the summations, the left hand side becomes (in the formula, pairs of terms that cancel out are marked with equal underlying boxed symbols)

$$\begin{split} \operatorname{Ric}_{12} = \underbrace{\left(\underbrace{F_1} + J_1^{*} \underbrace{F_2^{*} \varphi_1^{*} \varphi_2^{*}}{\partial r} \right) (\Gamma_{12}^{+})}_{\left[\underbrace{A} \right]} - \underbrace{\left(\underbrace{\widehat{A}} + B \varphi_2^{*} \underbrace{\widehat{C}} \right) (\Gamma_{12}^{+})}_{\left[\underbrace{A} \right]} - \underbrace{\Gamma_{12}^{*} \Gamma_{12}^{+} - \Gamma_{12}^{*} \Gamma_{12}^{+} - \Gamma_{12}^{*} \Gamma_{12}^{*} + \Gamma_{12}^{*} \Gamma_{12}^{+} - \Gamma_{12}^{*} \Gamma_{12}^{*} + \Gamma_{12}^{*} \Gamma_{12}^{*} - \Gamma_{12}^{*} \Gamma_{12}^{*} + \Gamma_{12}^{*} + \Gamma_{12}^{*} + \Gamma$$

=

This means that (4.11) is equivalent to the equation $\operatorname{Ric}^{N}(E_{1}, E_{2}) = 0$. The latter is identically satisfied because, by assumption, g_{o} is Kähler and $E_{2} = JE_{1}$. We now need to show that the equations (4.10) and (4.12) are both equivalent to (4.21). Expanding the summations in the indices m = 1, 2, the left hand sides of those equations become

$$\begin{split} \operatorname{Ric}_{11} &= \underbrace{\frac{\partial}{\partial x}} \left(\mathbf{F}_{41}^{(1)} + J_{1}^{\ell} E_{\ell}(\varphi) B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{(1)}) - \underbrace{\frac{\partial}{\partial x}} \left(\mathbf{F}_{41}^{(1)} - B \varphi_{y} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{(1)}) - \mathbf{\Gamma}_{11}^{\ell} \mathbf{F}_{1t}^{+} - \mathbf{\Gamma}_{1t}^{-\alpha_{0}} \mathbf{F}_{1ty}^{+} + \\ &+ \mathbf{\Gamma}_{11}^{\ell} \mathbf{F}_{1t}^{+} + \mathbf{\Gamma}_{1ty}^{+\alpha_{0}} \mathbf{F}_{1ty}^{+} - \lambda \mathbf{\Gamma}_{q_{0}}^{-2} + B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{1}^{0}) - \mathbf{\Gamma}_{q_{0}}^{\ell} \mathbf{\Gamma}_{1\ell}^{-q_{0}} + \\ &+ \frac{\partial}{\partial y} (\mathbf{\Gamma}_{11}^{2}) + J_{2}^{\ell} E_{\ell}(\varphi) B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{2}) - \frac{\partial}{\partial x} (\mathbf{\Gamma}_{21}^{2}) - B \varphi_{y} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{21}^{2}) - \mathbf{\Gamma}_{21}^{\ell} \mathbf{\Gamma}_{1\ell}^{2} - \mathbf{\Gamma}_{21}^{0} \mathbf{\Gamma}_{1q_{0}}^{-2} + \mathbf{\Gamma}_{11}^{\ell} \mathbf{\Gamma}_{2\ell}^{2} + \mathbf{\Gamma}_{11}^{q_{0}} \mathbf{\Gamma}_{2q_{0}}^{2} \\ &= \underbrace{\varphi_{y}} B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{41}^{-1}) - B \varphi_{y} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{41}^{-1}) - \lambda \mathbf{\Gamma}_{q_{0}}^{-2} + B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{-1}) - \mathbf{\Gamma}_{q_{0}}^{-1} \mathbf{F}_{1ty}^{-\alpha_{0}} - \mathbf{\Gamma}_{21}^{2} \mathbf{F}_{1ty}^{-2} + \frac{\partial}{\partial y} (\mathbf{\Gamma}_{11}^{-2}) - \varphi_{x} B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{12}^{-1}) \\ &= \underbrace{B} \\ \end{array}$$

$$= \underbrace{\varphi_{y}} B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{41}^{-1}) - B \varphi_{y} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{41}^{-1}) - \lambda \mathbf{\Gamma}_{q_{0}}^{-2} + B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{-1}) - \mathbf{\Gamma}_{q_{0}}^{-1} \mathbf{F}_{1ty}^{-\alpha_{0}} - \mathbf{\Gamma}_{q_{0}}^{-2} \mathbf{F}_{1ty}^{-\alpha_{0}} + \frac{\partial}{\partial y} (\mathbf{\Gamma}_{11}^{-2}) - \varphi_{x} B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{11}^{-2}) \\ &= \underbrace{B} \\ - \frac{\partial}{\partial x} (\mathbf{\Gamma}_{21}^{-2}) - B \varphi_{y} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{21}^{-2}) - \mathbf{\Gamma}_{21}^{-1} \mathbf{\Gamma}_{12}^{-2} - \mathbf{\Gamma}_{21}^{-2} \mathbf{F}_{1ty}^{-2} + \mathbf{\Gamma}_{11}^{-1} \mathbf{\Gamma}_{21}^{-2} + \mathbf{\Gamma}_{11}^{-\alpha_{0}} \mathbf{F}_{2ty}^{-\alpha_{0}} \\ &= \underbrace{B} \\ \\ \operatorname{Ric}_{22} = \frac{\partial}{\partial x} (\mathbf{\Gamma}_{21}^{-2}) + J_{1}^{\ell} E_{\ell} (\varphi) B \frac{\partial}{\partial r} (\mathbf{\Gamma}_{21}^{-2}) - \frac{\partial}{\partial y} (\mathbf{\Gamma}_{12}^{-1}) + B \varphi_{x} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{22}^{-1}) - \mathbf{\Gamma}_{12}^{\ell} \mathbf{\Gamma}_{21}^{-2} - \mathbf{\Gamma}_{12}^{-2} \mathbf{\Gamma}_{2ty}^{-\alpha_{0}} + \\ &+ \mathbf{\Gamma}_{22}^{\ell} \mathbf{\Gamma}_{1\ell}^{-1} + \mathbf{\Gamma}_{22}^{-\alpha_{0}} \mathbf{\Gamma}_{1q_{0}}^{-2} + \lambda \mathbf{\Gamma}_{q_{0}}^{-2} + \mathbf{F}_{22}^{-\alpha_{0}} \mathbf{\Gamma}_{22}^{-\alpha_{0}} + \frac{\partial}{\partial r} (\mathbf{\Gamma}_{22}^{-2}) - \\ &- \frac{\partial}{\partial y} (\mathbf{F}_{22}^{-2}) + B \varphi_{x} \frac{\partial}{\partial r} (\mathbf{\Gamma}_{22}^{-2}) - \mathbf{\Gamma}_{22}^{\ell} \mathbf{F}_{2t}^{-\alpha_{0}} - \mathbf{\Gamma}_{2t}^{-2} \mathbf{F}_{2t}^{-\alpha_{0}} + \\ &+ \mathbf{\Gamma}_{2}^{\ell} \mathbf{\Gamma}_{2}^{-\alpha_{0}}$$

After all cancellations, the expressions reduce to

$$\operatorname{Ric}_{11} = -\lambda \Gamma_{q_o}^{2} + B \frac{\partial}{\partial r} (\Gamma_{11}^{q_o}) + \frac{\partial}{\partial y} (\Gamma_{11}^{2}) - B \varphi_x \frac{\partial}{\partial r} (\Gamma_{11}^{2}) - \frac{\partial}{\partial x} (\Gamma_{21}^{2}) - B \varphi_y \frac{\partial}{\partial r} (\Gamma_{21}^{2}) - \Gamma_{21}^{1} \Gamma_{11}^{2} - \Gamma_{21}^{2} \Gamma_{12}^{2} + \Gamma_{11}^{1} \Gamma_{21}^{2} + \Gamma_{12}^{2} \Gamma_{22}^{2}$$

$$\operatorname{(B.4)}$$

$$\operatorname{Ric}_{22} = \lambda \Gamma_{q_o}^{1} + B \frac{\partial}{\partial r} (\Gamma_{22}^{q_o}) + \frac{\partial}{\partial x} (\Gamma_{21}^{1}) + B \varphi_y \frac{\partial}{\partial r} (\Gamma_{22}^{1}) - \frac{\partial}{\partial y} (\Gamma_{12}^{1}) + B \varphi_x \frac{\partial}{\partial r} (\Gamma_{12}^{1}) - \Gamma_{12}^{1} \Gamma_{21}^{1} - \Gamma_{12}^{2} \Gamma_{22}^{1} + \Gamma_{21}^{1} \Gamma_{11}^{1} + \Gamma_{22}^{2} \Gamma_{11}^{1} .$$

$$\operatorname{(B.5)}$$

Let us now expand these expressions using the decompositions $\Gamma_{ij}^{\ k} = \Gamma_{ij}^{\ k} + \delta \Gamma_{ij}^{\ k}$. We get:

$$\begin{split} \operatorname{Ric}_{11} &= \frac{\partial}{\partial y} (\Gamma_{11}^{\ 2}) - \frac{\partial}{\partial x} (\Gamma_{21}^{\ 2}) - \Gamma_{21}^{\ 1} \Gamma_{11}^{\ 2} - \Gamma_{21}^{\ 2} \Gamma_{12}^{\ 2} + \Gamma_{11}^{\ 1} \Gamma_{21}^{\ 2} + \Gamma_{11}^{\ 2} \Gamma_{22}^{\ 2} + \frac{\partial}{\partial y} \delta \Gamma_{11}^{\ 2} - \frac{\partial}{\partial x} \delta \Gamma_{21}^{\ 2} - \\ &- \Gamma_{21}^{\ 1} \delta \Gamma_{11}^{\ 2} - \Gamma_{21}^{\ 2} \delta \Gamma_{12}^{\ 2} + \Gamma_{11}^{\ 1} \delta \Gamma_{21}^{\ 2} + \Gamma_{11}^{\ 2} \delta \Gamma_{22}^{\ 2} - \delta \Gamma_{21}^{\ 1} \Gamma_{11}^{\ 2} - \delta \Gamma_{21}^{\ 2} \Gamma_{12}^{\ 2} + \delta \Gamma_{11}^{\ 1} \Gamma_{21}^{\ 2} + \delta \Gamma_{11}^{\ 2} \Gamma_{22}^{\ 2} + \\ &- \delta \Gamma_{21}^{\ 1} \delta \Gamma_{11}^{\ 2} - \delta \Gamma_{21}^{\ 2} \delta \Gamma_{12}^{\ 2} + \delta \Gamma_{11}^{\ 1} \delta \Gamma_{21}^{\ 2} + \delta \Gamma_{11}^{\ 2} \delta \Gamma_{22}^{\ 2} - \lambda \Gamma_{q_0}^{\ 2} + B \frac{\partial}{\partial r} \left(\Gamma_{11}^{\ q_0} - \varphi_y \Gamma_{21}^{\ 2} - \varphi_x \Gamma_{11}^{\ 2} \right) \ , \\ \operatorname{Ric}_{22} &= \frac{\partial}{\partial x} (\Gamma_{21}^{\ 2}) - \frac{\partial}{\partial y} (\Gamma_{12}^{\ 1}) - \Gamma_{12}^{\ 2} \Gamma_{21}^{\ 1} - \Gamma_{12}^{\ 2} \Gamma_{21}^{\ 2} + \Gamma_{22}^{\ 1} \Gamma_{11}^{\ 1} + \Gamma_{22}^{\ 2} \Gamma_{12}^{\ 1} + \frac{\partial}{\partial x} (\delta \Gamma_{21}^{\ 2}) - \frac{\partial}{\partial y} (\delta \Gamma_{12}^{\ 1}) - \\ &- \delta \Gamma_{12}^{\ 1} \Gamma_{21}^{\ 1} - \delta \Gamma_{12}^{\ 2} \Gamma_{21}^{\ 1} + \delta \Gamma_{22}^{\ 2} \Gamma_{11}^{\ 1} + \delta \Gamma_{22}^{\ 2} \Gamma_{11}^{\ 1} + \Gamma_{22}^{\ 2} \delta \Gamma_{12}^{\ 1} + \Gamma_{22}^{\ 2} \delta \Gamma_{11}^{\ 1} + \Gamma_{22}^{\ 2} \delta \Gamma_{12}^{\ 1} - \\ &- \delta \Gamma_{12}^{\ 1} \delta \Gamma_{21}^{\ 1} - \delta \Gamma_{12}^{\ 2} \delta \Gamma_{21}^{\ 1} + \delta \Gamma_{22}^{\ 2} \delta \Gamma_{12}^{\ 1} + \delta \Gamma_{22}^{\ 2} \delta \Gamma_{2$$

In both expansions, the first six summands give the components $\operatorname{Ric}_{11}^N$ and $\operatorname{Ric}_{22}^N$ of the Ricci tensor of the Kähler metric g_o of (N, J). Hence, using the expressions for the Γ_{ij}^k

and $\delta \Gamma_{ij}^{\ k}$ given in (B.2) and the fact that the functions φ and $\Gamma_{ij}^{\ k}$ are independent of the coordinate r, the equations $\operatorname{Ric}_{ii} = 0$, i = 1, 2, become

$$\begin{aligned} \operatorname{Ric}_{11} &= \operatorname{Ric}_{11}^{N} - \frac{\partial}{\partial y} (\delta\Gamma_{22}^{\ 2}) - \frac{\partial}{\partial x} (\delta\Gamma_{11}^{\ 1}) + \Gamma_{22}^{\ 2} \delta\Gamma_{22}^{\ 2} - \Gamma_{11}^{\ 1} \delta\Gamma_{14}^{\ +} + \Gamma_{11}^{\ 1} \delta\Gamma_{14}^{\ +} - \Gamma_{22}^{\ 2} \delta\Gamma_{22}^{\ 2} + \\ &+ \delta\Gamma_{22}^{\ 2} \Gamma_{22}^{\ 2} - \delta\Gamma_{11}^{\ 1} \Gamma_{14}^{\ +} + \delta\Gamma_{11}^{\ 1} \Gamma_{14}^{\ +} - \delta\Gamma_{22}^{\ 2} \Gamma_{22}^{\ 2} + \delta\Gamma_{22}^{\ 2} \delta\Gamma_{22}^{\ 2} - \delta\Gamma_{11}^{\ 1} \delta\Gamma_{44}^{\ +} + \delta\Gamma_{11}^{\ 1} \delta\Gamma_{44}^{\ +} - \delta\Gamma_{22}^{\ 2} \delta\Gamma_{22}^{\ 2} + \\ &- \lambda\Gamma_{q_{o}1}^{\ 2} + B \frac{\partial}{\partial r} \left(\Gamma_{11}^{\ q_{o}} - \varphi_{y}\Gamma_{11}^{\ 1} + \varphi_{x}\Gamma_{22}^{\ 2}\right) \end{aligned} \tag{B.6}$$

$$\operatorname{Ric}_{22} &= \operatorname{Ric}_{22}^{N} - \frac{\partial}{\partial x} (\delta\Gamma_{11}^{\ 1}) - \frac{\partial}{\partial x} (\delta\Gamma_{22}^{\ 2}) - \delta\Gamma_{22}^{\ 2} \Gamma_{22}^{\ 2} + \delta\Gamma_{11}^{\ 1} \Gamma_{44}^{\ +} - \delta\Gamma_{11}^{\ 1} \Gamma_{44}^{\ +} + \delta\Gamma_{22}^{\ 2} \Gamma_{22}^{\ 2} - \\ \end{aligned}$$

$$= \frac{1}{22} - \frac{1}{22} \frac{\partial^2 g}{\partial x^2} + \frac{1}{210} \frac{\partial^2 g}{\partial x^2} + \frac{1}{210} \frac{\partial^2 g}{\partial x^2} + \frac{1}{220} \frac{\partial^2 g}{\partial x^2} + \frac{1}{220} \frac{\partial^2 g}{\partial x^2} + \frac{1}{220} \frac{\partial^2 g}{\partial x^2} + \frac{1}{200} \frac{\partial^2$$

We now recall that $\Gamma_{q_o2}^{\ \ 1} = -\Gamma_{q_o1}^{\ \ 2}$, $\Gamma_{22}^{\ q_o} = \Gamma_{11}^{\ q_o}$ and that $\operatorname{Ric}_{22}^N = \operatorname{Ric}_{11}^N$ (because g_o is Kähler). Inserting these identities into (B.7) we get (B.6), showing that the equations (4.10) and (4.12) are equivalent.

Let us now focus on the equation (4.10) using (B.6). Using the property that φ and λ are independent of the coordinate r, the equation can be written as

$$\begin{split} \operatorname{Ric}_{11}^{N} &- \frac{\partial}{\partial x} \left(\frac{\sigma_{x}}{2\sigma} + B \frac{\sigma_{t}\varphi_{y}}{2\sigma} \right) - \frac{\partial}{\partial y} \left(\frac{1}{2\sigma} \sigma_{y} - B \frac{\sigma_{t}\varphi_{x}}{2\sigma} \right) - \lambda B \frac{1}{4\sigma} + \\ &+ B \frac{\partial}{\partial r} \left(-\lambda \sigma_{t} - \varphi_{y} \left(\Gamma_{11}^{-1} + \frac{1}{2\sigma} \sigma_{x} + B \frac{\sigma_{t}\varphi_{y}}{2\sigma} \right) + \varphi_{x} \left(\Gamma_{22}^{-2} + \frac{1}{2\sigma} \sigma_{y} - B \frac{\sigma_{t}\varphi_{x}}{2\sigma} \right) \right) = \\ &= \operatorname{Ric}_{11}^{N} - \frac{1}{2} (\log \sigma)_{xx} - \frac{B}{2} (\log \sigma)_{tx} \varphi_{y} - \frac{B}{2} (\log \sigma)_{t\varphi_{xy}} - \frac{1}{2} (\log \sigma)_{yy} + \frac{B}{2} (\log \sigma)_{ty} \varphi_{x} + \frac{B}{2} (\log \sigma)_{t\varphi_{xy}} - \\ &- \frac{\lambda B}{4\sigma} - B \lambda \sigma_{tt} - \varphi_{y} \frac{B}{2} (\log \sigma)_{xt} - B^{2} \frac{(\log \sigma)_{tt} (\varphi_{y})^{2}}{4} + \frac{B}{2} (\log \sigma)_{ty} \varphi_{x} - B^{2} \frac{(\log \sigma)_{tt} (\varphi_{x})^{2}}{2} = \\ &= \operatorname{Ric}_{11}^{N} - \frac{1}{2} (\log \sigma)_{xx} - \frac{1}{2} (\log \sigma)_{yy} - B (\log \sigma)_{tx} \varphi_{y} + B (\log \sigma)_{ty} \varphi_{x} - \frac{\lambda B}{4\sigma} - B \lambda \sigma_{tt} - \\ &- B^{2} \frac{(\log \sigma)_{tt} ((\varphi_{x})^{2} + (\varphi_{y})^{2})}{4} = 0 \;. \end{split}$$

Since

$$(\log \sigma)_x = \frac{\sigma_x}{\sigma} , \quad (\log \sigma)_{xx} = \frac{\sigma_{xx}}{\sigma} - \frac{\sigma_x^2}{\sigma^2} , \quad (\log \sigma)_y = \frac{\sigma_y}{\sigma} , \quad (\log \sigma)_{yy} = \frac{\sigma_{yy}}{\sigma} - \frac{\sigma_y^2}{\sigma^2} , \\ (\log \sigma)_{tx} = \frac{\sigma_{tx}}{\sigma} - \frac{\sigma_t \sigma_x}{\sigma^2} , \quad (\log \sigma)_{ty} = \frac{\sigma_{ty}}{\sigma} - \frac{\sigma_t \sigma_y}{\sigma^2} , \quad (\log \sigma)_{tt} = \frac{\sigma_{tt}}{\sigma} - \frac{\sigma_t^2}{\sigma^2} ,$$

the same equation can be also written as

$$\operatorname{Ric}_{11}^{N} - \frac{\sigma_{xx}}{2\sigma} + \frac{\sigma_{x}^{2}}{2\sigma^{2}} - \frac{\sigma_{yy}}{2\sigma} + \frac{\sigma_{y}^{2}}{2\sigma^{2}} - B\frac{\sigma_{tx}}{\sigma}\varphi_{y} + B\frac{\sigma_{t}\sigma_{x}}{\sigma^{2}}\varphi_{y} + B\frac{\sigma_{ty}}{\sigma}\varphi_{x} - B\frac{\sigma_{t}\sigma_{y}}{\sigma}\varphi_{x} - B\frac{\sigma_{t}\sigma_{y}}{\sigma}\varphi_{x} - B\frac{\lambda}{4\sigma} - B\lambda\sigma_{tt} - B^{2}\left(\frac{\sigma_{tt}}{2\sigma} - \frac{\sigma_{t}^{2}}{2\sigma^{2}}\right)\left((\varphi_{x})^{2} + (\varphi_{y})^{2}\right) = 0.$$

Note that $\frac{\partial}{\partial r} = p_o$. Hence, re-arranging the summands in an appropriate way, we may

write this equation as

$$\operatorname{Ric}_{11}^{N} - \frac{\sigma_{xx}}{2\sigma} + \frac{\sigma_{x}^{2}}{2\sigma^{2}} - \frac{\sigma_{yy}}{2\sigma} + \frac{\sigma_{y}^{2}}{2\sigma^{2}} - B\frac{\lambda}{4\sigma} - B\lambda p_{o}(p_{o}(\sigma)) - \\ - B\left(\frac{E_{1}(p_{o}(\sigma))}{\sigma} - \frac{p_{o}(\sigma)E_{1}(\sigma)}{\sigma^{2}} + B\frac{p_{o}(p_{o}(\sigma))}{2\sigma}\varphi_{y} - B\frac{(p_{o}(\sigma))^{2}}{2\sigma^{2}}\varphi_{y}\right)\varphi_{y} + \\ + B\left(\frac{E_{2}(p_{o}(\sigma))}{\sigma} - \frac{p_{o}(\sigma)E_{2}(\sigma)}{\sigma^{2}} - B\frac{p_{o}(p_{o}(\sigma))}{2\sigma}\varphi_{x} + B\frac{(p_{o}(\sigma))^{2}}{2\sigma^{2}}\varphi_{x}\right)\varphi_{x} = 0.$$
(B.8)

We now recall n = 4 and φ and σ satisfy (3.20), which is the same as the system

$$-\frac{1}{2\sigma^2}E_2(\sigma) - \frac{1}{\sigma}E_1(\mathbf{p}_o(\sigma)) + \frac{1}{\sigma^2}\mathbf{p}_o(\sigma)E_1(\sigma) + \frac{B}{2\sigma^2}\varphi_x\mathbf{p}_o(\sigma) - \frac{B}{\sigma}\varphi_y\mathbf{p}_o(\mathbf{p}_o(\sigma)) + \frac{B}{\sigma^2}\varphi_y(\mathbf{p}_o(\sigma))^2 = 0 ,$$

$$\frac{1}{2\sigma^2}E_1(\sigma) - \frac{1}{\sigma}E_2(\mathbf{p}_o(\sigma)) + \frac{1}{\sigma^2}\mathbf{p}_o(\sigma)E_2(\sigma) + \frac{B}{2\sigma^2}\varphi_y\mathbf{p}_o(\sigma) + \frac{B}{\sigma}\varphi_x\mathbf{p}_o(\mathbf{p}_o(\sigma)) - \frac{B}{\sigma^2}\varphi_x(\mathbf{p}_o(\sigma))^2 = 0 ,$$

or, equivalently, to

$$\frac{E_{1}\left(\mathbf{p}_{o}(\sigma)\right)}{\sigma} - \frac{\mathbf{p}_{o}(\sigma)E_{1}\left(\sigma\right)}{\sigma^{2}} = -\frac{1}{2\sigma^{2}}E_{2}\left(\sigma\right) + \frac{B}{2\sigma^{2}}\varphi_{x}\mathbf{p}_{o}\left(\sigma\right) - \frac{B}{\sigma}\varphi_{y}\mathbf{p}_{o}\left(\mathbf{p}_{o}(\sigma)\right) + \frac{B}{\sigma^{2}}\varphi_{y}\left(\mathbf{p}_{o}\left(\sigma\right)\right)^{2} ,$$

$$\frac{E_{2}\left(\mathbf{p}_{o}(\sigma)\right)}{\sigma} - \frac{\mathbf{p}_{o}(\sigma)E_{2}\left(\sigma\right)}{\sigma^{2}} = \frac{1}{2\sigma^{2}}E_{1}\left(\sigma\right) + \frac{B}{2\sigma^{2}}\varphi_{y}\mathbf{p}_{o}\left(\sigma\right) + \frac{B}{\sigma}\varphi_{x}\mathbf{p}_{o}\left(\mathbf{p}_{o}(\sigma)\right) - \frac{B}{\sigma^{2}}\varphi_{x}\left(\mathbf{p}_{o}\left(\sigma\right)\right)^{2} .$$

Replacing this in (B.8), we get

$$\operatorname{Ric}_{11}^{N} - \frac{\sigma_{xx}}{2\sigma} + \frac{\sigma_{x}^{2}}{2\sigma^{2}} - \frac{\sigma_{yy}}{2\sigma} + \frac{\sigma_{y}^{2}}{2\sigma^{2}} - B\frac{\lambda}{4\sigma} - B\lambda p_{o}(\mathbf{p}_{o}(\sigma)) - B\left(-\frac{1}{2\sigma^{2}}\sigma_{y} + \frac{B}{2\sigma^{2}}\varphi_{x}p_{o}(\sigma) - \frac{B}{2\sigma}\varphi_{y}p_{o}(\mathbf{p}_{o}(\sigma)) + \frac{B}{2\sigma^{2}}\varphi_{y}\left(\mathbf{p}_{o}(\sigma)\right)^{2}\right)\varphi_{y} + B\left(\frac{1}{2\sigma^{2}}\sigma_{x} + \frac{B}{2\sigma^{2}}\varphi_{y}p_{o}(\sigma) + \frac{B}{2\sigma}\varphi_{x}p_{o}\left(\mathbf{p}_{o}(\sigma)\right) - \frac{B}{2\sigma^{2}}\varphi_{x}\left(\mathbf{p}_{o}(\sigma)\right)^{2}\right)\varphi_{x} = 0.$$
(B.9)

Since $-2\sigma p_o(p_o(\sigma)) + p_o(\sigma)^2 + \frac{1}{4} = 0$ (which allows us to replace $-\frac{B}{2\sigma^2} p_o(\sigma)^2 = -\frac{B}{\sigma} p_o(p_o(\sigma)) + \frac{B}{8\sigma^2}$ in all terms), multiplying left and right hand side by σ^2 , the equation transforms into (4.21), as we needed to prove.

DECLARATIONS

The authors declare that they have no conflict of interest.

References

- A. Awad and A. Chamblin, A Bestiary of Higher Dimensional Taub-NUT-AdS Spacetimes, Class. Quant. Grav. 19 (2002), 2051–2062.
- [2] D. V. Alekseevsky, M. Ganji and G. Schmalz, *CR-geometry and shearfree Lorentzian geometry*, pp. 11–22, in "Geometric Complex Analysis", Springer, Singapore.
- [3] D. V. Alekseevsky, M. Ganji, G. Schmalz and A. Spiro, Lorentzian manifolds with shearfree congruences and Kähler-Sasaki geometry, Differential Geom. Appl., 75 (2021), Paper No. 101724, 32
- [4] D. V. Alekseevsky, M. Ganji, G. Schmalz and A. Spiro, The Levi-Civita connections of Lorentzian manifolds with prescribed optical geometries, preprint on arXiv (2023) – to appear on CUBO.

- [5] F. A. Bais and P. Batenburg, A new class of higher-dimensional Kaluza-Klein monopole and instanton solutions, Nuclear Phys. B, 253 (1985), 162–172.
- [6] A. L. Besse, Einstein manifolds, Springer-Verlag, Berlin, 2008.
- [7] J. Carminati and R. G. McLenaghan, Algebraic invariants of the Riemann tensor in a four-dimensional Lorentzian space, J. Math. Phys. 32 (1991), 3135–3140.
- [8] A. Chamblin and G.W. Gibbons, *Topology and Time Reversal*, in "Erice 1995, String gravity and physics at the Planck energy scale" (N. Sanchez and A. Zichichi ed.), pp. 233–253, *Kluwer Academic Publishers, Dordrecht*, 1996.
- G. C. Debney, R. P. Kerr and A. Schild, Solutions of the Einstein and Einstein-Maxwell equations, J. Mathematical Phys. 10 (1969), 1842–1854.
- [10] A. Fino, T. Leistner and A. Taghavi-Chabert, Optical Geometries, to appear on Ann. Sc. Norm., 2022.
- [11] V. Frolov and D. Stojković, Particle and light motion in a space-time of a five-dimensional rotating black hole, Phys. Rev. D (3), 68 (2003), 064011-1–8.
- [12] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer-Verlag, Berlin, 2001.
- [13] D. C. Hill, J. Lewandowski and P. Nurowski, Einstein's equations and the embedding of 3-dimensional CR manifolds, Indiana Univ. Math. J., 57 (2008), 3131–3176.
- [14] E. Hopf, Über den funktionalen, insbesondere den analytischen Charakter der Lösungen elliptischer Differentialgleichungen zweiter Ordnung, Math. Z. 34 (1932), 194–233.
- [15] G. T. Horowitz, Higher dimensional generalizations of the Kerr black hole, in "The Kerr spacetime", pp. 332–343, Cambridge Univ. Press, Cambridge, 2009.
- [16] R. P. Kerr, Scalar invariants and groups of motions in a V_n with positive definite metric tensor, Tensor (N.S.) **12** (1962), 74–83.
- [17] R. P. Kerr, Gravitational field of a spinning mass as an example of algebraically special metrics, Phys. Rev. Lett. 11 (1963), 237–238.
- [18] R. P. Kerr, Discovering the Kerr and Kerr-Schild metrics, in "The Kerr spacetime", pp. 38–72, Cambridge Univ. Press, Cambridge, 2009.
- [19] S. Kobayashi and N. Nomizu, Foundations of differential geometry. Vol I, Interscience Publishers, York-London, 1963.
- [20] B. Kostant, Quantization and unitary representations. I. Prequantization, in Lectures in Modern Analysis and Applications III, Lecture Notes in Math., Vol. 170, pp. 87–208, Springer, Berlin-New York, 1970.
- [21] F. John, On linear partial differential equations with analytic coefficients, Comm. Pure Appl. Math. 2 (1949), 209–253
- [22] R. C. Myers and M. J. Perry, Black holes in higher-dimensional space-times, Ann. Physics 172, (1986), 304–347.
- [23] C. B. Morrey Jr. and L. Nirenberg, On the Analyticity of the Solutions of Linear Elliptic Systems of Partial Differential Equations, Comm. Pure Appl. Math. 10 (1957), 271–290.
- [24] M. Ortaggio, On the uniqueness of the Myers-Perry spacetime as a type II(D) solution in six dimensions, J. High Energy Phys. (2017) n. 6, pp. 042, front matter+39.
- [25] M. Ortaggio and A. Srinivasan. Charging Kerr-Schild spacetimes in higher dimensions, preprint posted on arXiv:2309.02900 (2023).
- [26] D. N. Page and C. N. Pope, Inhomogeneous Einstein metrics on complex line bundles, Classical Quantum Gravity, 4, (1987), 213–225,
- [27] V. Pravda, A. Pravdová, and M. Ortaggio, Type D Einstein spacetimes in higher dimensions, Classical Quantum Gravity, 24 (2007), 4407–4428.
- [28] D. C. Robinson, Four decades of black hole uniqueness theorems, in "The Kerr spacetime", pp. 115–143, Cambridge Univ. Press, Cambridge, 2009.
- [29] I. Robinson and A. Trautman, Conformal geometry of flows in n dimensions, J. Math. Phys. 24 (1983), 1425–1429.

- [30] I. Robinson and A. Trautman, Optical geometry in "Warsaw symposium on elementary particle physics: new theories in physics, Kazimierz (Poland), 1988", pp. 454–497, World Scientific Pub. Co., Teaneck, NJ (USA), 1989.
- [31] N. Steenrod, Topology of Fiber Bundles, Princeton University Press, Princeton, 1950.
- [32] A. Taghavi-Chabert, Twisting non-shearing congruences of null geodesics, almost CR structures and Einstein metrics in even dimensions, Ann. Mat. Pura Appl. (4), **201** (2022), 655–693.
- [33] A. Trautman, Cauchy-Riemann structures in optical geometry, in "Proceedings of the fourth Marcel Grossmann meeting on general relativity, Part A, B (Rome, 1985)", pp. 317–324, North-Holland, Amsterdam, 1986.
- [34] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Herlt, Exact solutions of Einstein's field equations, *Cambridge University Press, Cambridge-New York*, 2003.

MASOUD GANJI AND GERD SCHMALZ SCHOOL OF SCIENCE AND TECHNOLOGY UNIVERSITY OF NEW ENGLAND, ARMIDALE NSW 2351 AUSTRALIA

E-mail: mganjia2@une.edu.au E-mail: schmalz@une.edu.au Cristina Giannotti and Andrea Spiro Scuola di Scienze e TecnolSogie Università di Camerino I-62032 Camerino (Macerata) Italy

E-mail: cristina.giannotti@unicam.it *E-mail*: andrea.spiro@unicam.it