

SPECTRAL RADII FOR SUBSETS OF HILBERT C^* -MODULES AND SPECTRAL PROPERTIES OF POSITIVE MAPS

B V RAJARAMA BHAT, BISWARUP SAHA AND PRAJAKTA SAHASRABUDDHE

ABSTRACT. The notions of joint and outer spectral radii are extended to the setting of Hilbert C^* -bimodules. A Rota-Strang type characterisation is proved for the joint spectral radius. In this general setting, an approximation result for the joint spectral radius in terms of the outer spectral radius has been established.

This work leads to a new proof of the Wielandt-Friedland's formula for the spectral radius of positive maps. Following an idea of J. E. Pascoe, a positive map called the *maximal part* has been associated to any positive map with non-zero spectral radius, on finite dimensional C^* -algebras. This provides a constructive treatment of the Perron-Frobenius theorem. It is seen that the maximal part of a completely positive map has a very simple structure and it is irreducible if and only if the original map is irreducible.

It is observed that algebras generated by tuples of matrices can be determined and their dimensions can be computed by realizing them as linear span of Choi-Kraus coefficients of some easily computable completely positive maps.

1. INTRODUCTION

Throughout this paper M_m denotes the C^* -algebra of $m \times m$ complex matrices where m is a natural number and \mathcal{A} denotes a general unital C^* -algebra. It is well-known that the spectral radius of an element $a \in \mathcal{A}$, denoted $r(a)$, can be obtained by the Gelfand formula:

$$r(a) = \lim_{n \rightarrow \infty} \|a^n\|^{\frac{1}{n}}.$$

Taking a cue from this, the notion of *joint spectral radius* for collections of unital normed algebra elements was first introduced by Rota and Strang [35]. For any bounded subset S of \mathcal{A} , they define the joint spectral radius of S by

$$\rho(S) := \lim_{n \rightarrow \infty} \sup_{a_1, \dots, a_n \in S} \|a_1 \cdots a_n\|^{1/n}. \quad (1.1)$$

They characterize this quantity as follows. Let $\mathfrak{N}_{\mathcal{A}}$ be the family of all norms on \mathcal{A} , which are equivalent to the original norm and under which \mathcal{A} is a normed algebra. With this notation ([35, Proposition 1]):

$$\rho(S) = \inf_{N \in \mathfrak{N}_{\mathcal{A}}} \sup_{a \in S} N(a). \quad (1.2)$$

The problem of understanding the convergence behaviour of products of matrices appears in various mathematical projects. It is only natural that the notion of joint spectral radius plays a crucial role in such situations. Research advancements in the last few decades have led to further exploration of the concept of joint spectral radius, and the literature is vast. We cite investigations by Daubechies and Lagarias [9], Berger and Wang [3] and Elsner [10]. Computational challenges in computing the joint spectral radius spurred the development of efficient approximation methods. See, for instance, Parrilo

2020 *Mathematics Subject Classification.* 46L08, 47A10, 46L57.

Key words and phrases. spectral radius, joint spectral radius, Hilbert modules, completely positive maps.

and Jadbabaie [25], and Blondel and Nesterov [6]. Recently, Pascoe ([27], [28]) examined a special approximation procedure for the joint spectral radius by introducing a notion of *outer spectral radius* for p -tuple (A_1, A_2, \dots, A_p) of $m \times m$ matrices via the formula

$$\widehat{\rho}(A_1, \dots, A_p) = \sqrt{r \left(\sum_{i=1}^p \overline{A_i} \otimes A_i \right)}, \quad (1.3)$$

where $A \otimes B$ denotes the usual Kronecker product of two matrices and \overline{A} denotes the complex conjugate of A .

Pascoe's notion of the outer spectral radius unveils connections to completely positive (CP) maps, offering insights into the quantum dynamics on M_m . The author did notice this connection. However, all his computations are mostly based on the formula 1.3. This amounts to taking a matrix representation of the completely positive map and using vectorization and other similar methods that use coordinates. This restricts the applicability of the theory to completely positive maps on finite dimensions. In this article, we consider positive maps and completely positive maps on general C^* -algebras. This necessitates a more abstract approach. In particular, we are led to defining a joint spectral radius for subsets of Hilbert C^* -modules and an outer spectral radius for sequences of elements from Hilbert C^* -modules.

Here is a brief outline of the contents and the organisation of this article. In Section 2, we collect some definitions and important results from the literature essential for our discussion. In Section 3.1, we begin by proving some basic results for general positive maps on unital C^* -algebras. These are motivated by similar results obtained by Popescu in [34]. His focus was w^* -continuous positive maps on $\mathcal{B}(\mathcal{H})$, where \mathcal{H} is a possibly infinite dimensional Hilbert space and his methods make use of this setup. We modify the proofs suitably to suit our situation. Well-known Perron-Frobenius theory tells us that spectral radii of non-negative matrices are eigenvalues. In Theorem 3.6 we show that the spectral radius is a spectral point (but not necessarily an eigenvalue) for positive maps on general C^* -algebras. This is a known classical result (Karlin (see [21, Theorem 4]), Bonsall (see [7, Lemma 5]) and Schaefer (see [36, Proposition 1])), however, we have it here by fairly elementary methods.

In Section 3.2, we generalize the notion of the joint spectral radius and the outer spectral radius to the setup of Hilbert C^* -bimodules. Let E denote any Hilbert \mathcal{A} - \mathcal{A} -module, and let G be any bounded subset of E . Taking inspiration from the definition of the joint spectral radius of a bounded subset of a Banach algebra, we define the joint spectral radius of G , using the interior tensor product \odot on E by

$$\rho_E(G) := \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \dots \odot \xi_n\|^{\frac{1}{n}}. \quad (1.4)$$

We show that this is well-defined and has analogous properties. Further, to define the outer spectral radius, we first observe that with any p -tuple $A = (A_1, \dots, A_p)$ of $m \times m$ matrices, one can associate a completely positive (CP) map $\tau_A : M_m \rightarrow M_m$ defined by $\tau_A(X) = \sum_{i=1}^p A_i^* X A_i$. Clearly, $\sum_{i=1}^p \overline{A_i} \otimes A_i$ is the matrix of τ_A^* where $\tau_A^*(X) = \sum_{i=1}^p A_i X A_i^*$ and $\widehat{\rho}(A_1, \dots, A_p) = \sqrt{r(\tau_A^*)} = \sqrt{r(\tau_A)}$, that is, the outer spectral radius defined in (1.3) is nothing but the square root of the spectral radius of the associated CP map τ_A . Taking motivation from this, consider a finite or countably infinite sequence $G = (\xi_1, \xi_2, \dots)$ of elements of E for which $\tau_G(a) := \sum_i \langle \xi_i, a \xi_i \rangle$ defines a CP map on \mathcal{A} and define the outer spectral radius of G by

$$\widehat{\rho}_E(G) := \sqrt{r(\tau_G)}. \quad (1.5)$$

Note that the joint spectral radius is defined for bounded subsets of Hilbert C^* -modules, whereas, the outer spectral radius is naturally defined for (finite or infinite) sequences of elements of Hilbert C^* -modules. Here, repeating some elements does make a difference. However, for any such sequence, the associated joint spectral radius, by definition, is the joint spectral radius of the set $\{\xi : i \geq 1\}$, and here repetitions of terms will not make any difference.

With definitions of joint spectral radius and outer spectral radius in place, we can study and compare them. In Theorem 3.9, we give a Rota-Strang type characterisation for the joint spectral radius. The main difference from classical Rota-Strang theory is that we are forced to consider equivalent norms on the full Fock module generated by the Hilbert C^* -module E , with some sub-multiplicative properties. Previously, it was enough to consider norms on the original algebra. This is natural in view of the definition of the joint spectral radius in (1.4).

While developing this theory of spectral radii, we observe connections with some well-known results. A famous theorem of Wielandt [42] provides a simple way of computing the spectral radius of an irreducible entry-wise positive matrix $A = [a_{ij}]_{1 \leq i, j \leq m}$ by showing

$$r(A) = \min_{x > 0} \max_{1 \leq i \leq m} \frac{(Ax)_i}{x_i}.$$

This result was generalized considerably by Friedland [14]. He showed that the spectral radius of a positive map φ on a unital C^* -algebra \mathcal{A} is given by

$$r(\varphi) = \inf\{r(v^{-1}\varphi(v)) : v \in \mathcal{A} \text{ is strictly positive}\}. \quad (1.6)$$

We obtain these results as simple consequences of our analysis of spectral radii for subsets of Hilbert C^* -modules in Section 3. We obtain inequalities between the joint and outer spectral radii, culminating in an approximation theorem for the joint spectral radius of finite sets in terms of outer spectral radius (See Theorem 3.15):

$$\rho_E(\xi_1, \dots, \xi_d) = \lim_{k \rightarrow \infty} \widehat{\rho}_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k})^{1/k}.$$

This is a generalization of a similar result for matrices proved by Pascoe [28].

In Section 4, we focus on the study of positive maps on finite dimensional C^* -algebras and their spectral properties. We return to Perron-Frobenius theory. Perron ([30], [31]) and Frobenius ([15], [16]) discovered that the spectral radius of matrices with positive entries is an eigenvalue with a corresponding eigenvector of positive entries. In fact, they established that if the matrix is irreducible, then its spectral radius is a simple eigenvalue with unique positive eigenvectors (up to scalar multiplication). Evans and Høegh-Krohn [11] extended this classical result to finite-dimensional C^* -algebras by broadening the notion of irreducibility to positive maps on such algebras: A positive map φ on \mathcal{A} is said to be irreducible if there is no non-trivial projection $p \in \mathcal{A}$ for which the hereditary C^* subalgebra $p\mathcal{A}p$ is invariant under φ . Further observations by Albeverio-Høegh-Krohn [1], Groh ([18], [19]), Schrader [37] and Lagro-Yang-Xiong [23] highlighted the intriguing spectral properties possessed by strictly positive or irreducible positive maps.

For a positive map on a C^* -algebra, we will call any positive eigenvector, with spectral radius as the eigenvalue, a Perron-Frobenius eigenvector. Pascoe [28] introduced the concept of maximal spectrum for matrices, as comprising eigenvalues with maximum modulus and maximal degeneracy index. Using Jordan decomposition, he demonstrated that the spectral radius of a matrix of the form $\sum_i \overline{A_i} \otimes A_i$ lies within the maximal spectrum. Inspired by Pascoe's approach, we formally introduce a notion of maximal part of a positive map φ with strictly positive spectral radius, on a finite dimensional

C^* -algebra \mathcal{A} . It is defined as the positive map

$$\widehat{\varphi} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\varphi^n}{\|\varphi^n\|_{\beta}}, \quad (1.7)$$

where β is a Jordan basis of φ . Analyzing this map we see that (Theorem 4.9) the spectral radius is in the maximal spectrum of φ , that is, it has Jordan blocks of maximum size among eigenvalues of the same modulus (see Definition 4.5). Next in Theorem 4.10, we observe that $\widehat{\varphi}$ has remarkable commutation properties with φ and $\widehat{\varphi}(1)$ serves as a Perron-Frobenius eigenvector of φ . This is significantly different from the work Evans and Høegh-Krohn [11], as we now have got a constructive method to obtain the Perron-Frobenius eigenvector.

In the last Section, we restrict our attention to completely positive (CP) maps on finite dimensional C^* algebras. We tackle two issues described below.

Given a p -tuple of matrices, say A_1, A_2, \dots, A_p in M_m , the first question is how to get a basis for the algebra (not $*$ -algebra) generated by them. In [27], Pascoe developed an efficient method for the same. We show that his method has a very natural interpretation on viewing the tuple as Choi-Kraus coefficients of the CP maps $\tau(X) = \sum_{i=1}^p A_i^* X A_i$ on M_m . The algebra we are interested can be constructed as the space of Choi-Kraus coefficients of some other CP maps constructed out of τ . Theorem 5.1 provides an ample supply of such CP maps. Pascoe [27] obtained one such map, namely, $\gamma_1 = (1 - b\tau)^{-1} - 1$ where b is a scalar such that $br(\tau) < 1$. In fact, using the finite dimensionality in Theorem 5.5, we show that there is a polynomial p of degree not exceeding m^2 , with strictly positive scalar coefficients such that the Choi-Kraus coefficient space of $p(\tau)$ is equal to the (unital) subalgebra of M_m generated by A_i 's.

The second issue is about obtaining a characterization of irreducibility of CP maps in terms of their Choi-Kraus coefficients. It is well-known that a CP map on M_m is irreducible if and only if its Choi-Kraus coefficients do not admit any non-trivial common invariant subspace ([12], [13]). This condition is no longer necessary for irreducibility of CP maps on general finite dimensional C^* -algebras (See Example 5.6). To resolve this for a CP map τ on a unital C^* -subalgebra \mathcal{A} of M_m , we introduce a notion of the canonical extension $\widetilde{\tau}$ of τ to M_m and in Proposition 5.9, we establish equivalence between irreducibility of τ and $\widetilde{\tau}$. Hence, the desired necessary and sufficient criterion for the irreducibility of τ can be described in terms of the Choi-Kraus decompositions of $\widetilde{\tau}$.

Finally, in Theorem 5.17 we study consequences of irreducibility condition on the maximal part. This is possible as irreducibility implies that the map has strictly positive spectral radius (see Remark 4.11). The main observation is that the maximal part of an irreducible CP map on a finite dimensional C^* -algebra has the form:

$$X \mapsto \text{trace}(RX) L$$

with a strictly positive density matrix R and a strictly positive element L . Two examples are given to illustrate this.

2. PRELIMINARIES AND BACKGROUND MATERIAL

For the basic theory of C^* -algebras, positive and completely positive (CP) maps, we refer to ([24], [29], [4]). Typically if a map we are studying is assumed to be just positive we denote it by φ and if it has to be CP then by τ . We recall the well-known Choi-Kraus decomposition ([8], [5]) of CP maps on matrix algebras. A linear map $\tau : M_m \rightarrow M_m$ is

CP if and only if there exist $A_1, \dots, A_p \in M_m$, for some p , such that

$$\tau(X) = \sum_{i=1}^p A_i^* X A_i, \quad \forall X \in M_m. \quad (2.1)$$

The matrices A_i 's are known as Choi-Kraus coefficients of the CP map τ . This is decomposition for τ is not unique. However, if

$$\tau(X) = \sum_{j=1}^q B_j^* X B_j, \quad \forall X \in M_m,$$

is another such decomposition, then every A_i is a linear combination of B_j 's and conversely every B_j is a linear combination of A_i 's. In particular,

$$\mathcal{M}_\tau := \text{span}\{A_i : 1 \leq i \leq p\}$$

is a subspace of M_m , independent of the decomposition. The space \mathcal{M}_τ is said to be the *Choi-Kraus coefficient space* (or, *coefficient space* in short) of τ . Dimension of \mathcal{M}_τ is called the Choi rank of the CP map τ . W. Arveson [2] considers the adjoint of this space (called the metric operator space of τ) and defines a metric on it, which is not relevant for this article. A CP map η is said to be *dominated* by τ if $\tau - \eta$ is also CP. By considering Choi-Kraus representations of η and $\tau - \eta$ it is clear that if η is dominated by τ then $\mathcal{M}_\eta \subseteq \mathcal{M}_\tau$.

We are interested in determining the algebra (not the $*$ -algebra) generated by elements of \mathcal{M}_τ . In this context, the following observation is useful:

Proposition 2.1. *Let $\tau : M_n \rightarrow M_n$ be a CP map. Then $A \in M_n$ is in the metric space \mathcal{M}_τ if and only if there exists $q > 0$ such that α_A is dominated by $q\tau$, where $\alpha_A(X) = A^* X A$.*

Proof. Suppose that τ has the Choi-Kraus decomposition as in (2.1). Let us assume that $q\tau - \alpha_A$ is CP for some $q > 0$. Then there are matrices $C_1, \dots, C_r \in M_n$ such that $q\tau(X) - \alpha_A(X) = \sum_{j=1}^r C_j^* X C_j$ for all $X \in M_n$. Let σ be the CP map on M_n defined by $\sigma(X) = A^* X A + \sum_{j=1}^r C_j^* X C_j$. Observe that σ has two Choi-Kraus expressions,

$$\sigma(X) = \sum_{i=1}^p (\sqrt{q}A_i)^* X (\sqrt{q}A_i) = A^* X A + \sum_{j=1}^r C_j^* X C_j, \quad \forall X \in M_n.$$

This ensures that A is a linear combination of A_i 's, that is, $A \in \mathcal{M}_\tau$.

Conversely, assume that $A \in \mathcal{M}_\tau$. Let $A = \sum_{i=1}^p \lambda_i A_i$ for some scalars λ_i . Consider the positive matrix $V = (\overline{\lambda_i} \lambda_j) \in M_n$ and choose $q > \|V\|$. Then,

$$q\tau(X) - \alpha_A(X) = q \sum_{i=1}^p A_i^* X A_i - A^* X A = \sum_{i,j} (q\delta_{ij} - \overline{\lambda_i} \lambda_j) A_i^* X A_j, \quad \forall X \in M_n,$$

Since $qI - V = (q\delta_{ij} - \overline{\lambda_i} \lambda_j)$ is a positive matrix, $q\tau - \alpha_A$ is CP. \square

For a linear map τ on M_m , we define the Choi matrix of τ as

$$C_\tau := \sum_{i,j} E_{ij} \otimes \tau(E_{ij}) \in M_m \otimes M_m.$$

Observe that the association $\tau \mapsto C_\tau$ is continuous. It is well known fact that a linear map τ on M_m is CP if and only if its Choi matrix $C_\tau \in M_{m^2}$ is positive. Define a linear map $\text{vec} : M_m \rightarrow \mathbb{C}^{m^2}$ by $\text{vec}(E_{ij}) = e_{i+(j-1)m}$. For $A \in M_m$, $\text{vec} A$ is called the vectorization of A . If τ is an elementary CP map α_A , then $i + (j - 1)m^{\text{th}}$ column of the Choi matrix C_τ is $a_{ij}\text{vec}(A)$ where $A = [a_{ij}]_{1 \leq i, j \leq m}$ for all i, j . We recall the following interpretation of coefficient spaces using Choi matrices.

Proposition 2.2. *Let τ be a CP map on M_m . Then, the vectorization map vec establishes a one-to-one correspondence between the coefficient space of τ and the range of its conjugated Choi matrix. In particular, the Choi rank of τ is same as the rank of the Choi matrix C_τ .*

We now recall the notion of irreducible positive maps and a couple of basic results from Evans and Høegh-Krohn ([11]). A positive map φ on a finite dimensional C^* -algebra \mathcal{A} is said to be irreducible if there is no non-trivial projection $p \in \mathcal{A}$, such that the hereditary C^* -subalgebra $p\mathcal{A}p$ is invariant under φ . Evans and Høegh-Krohn ([11, Lemma 2.1]) show that if \mathcal{A} is a subalgebra of M_n , then φ is irreducible if and only if $(id + \varphi)^{n-1}$ is strictly positive, that is, it sends non-zero positive elements to strictly positive elements.

Irreducible CP maps on the full matrix algebra M_m have several interesting properties.

Proposition 2.3 (Theorem 2, [13]). *Let $\tau : M_m \rightarrow M_m$ be a completely positive map with a Choi-Kraus decomposition as in 2.1. Then the following are equivalent:*

- (1) *The positive map τ is irreducible;*
- (2) *The matrices $\{A_1, \dots, A_p\}$ have no non-trivial common invariant subspace;*
- (3) *The algebra generated by $\{A_1, \dots, A_p\}$ is whole of M_m .*

Evans and Høegh-Krohn have also proved the following *Perron-Frobenius type theorem* for irreducible positive maps on finite dimensional C^* -algebras.

Theorem 2.4 (Theorem 2.3, [11]). *Let φ be a (irreducible) positive linear map on a finite dimensional C^* -algebra. Then the spectral radius $r(\varphi)$ is an (simple) eigenvalue of φ possessing a (invertible) positive eigenvector.*

Now, we recall some of the required definitions and results about Hilbert C^* -modules. For further details, we refer to [22, 26, 38, 39].

Definition 2.5. Let \mathcal{A} be a unital C^* -algebra. A Hilbert \mathcal{A} -module is a right \mathcal{A} -module (compatible with scalar multiplication) together with a \mathcal{A} -valued inner product, which is complete with respect to the associated norm $\|\xi\| = \sqrt{\|\langle \xi, \xi \rangle\|}$. For $\xi \in E$, $|\xi|$ denotes the positive element $\langle \xi, \xi \rangle^{1/2}$.

Certainly, we can talk about bounded operators on Hilbert C^* -modules. However, unlike bounded maps on Hilbert spaces, they may not possess an adjoint.

Definition 2.6. Let E, F be Hilbert \mathcal{A} -modules. The space of all *adjointable* maps from E to F , denoted by $\mathcal{B}^a(E, F)$, is the collection of all right linear maps $t : E \rightarrow F$ such that there exists a map $t^* : F \rightarrow E$ satisfying $\langle \xi, t\eta \rangle = \langle t^*\xi, \eta \rangle$ for all $\xi, \eta \in E$.

It is evident that every element of $\mathcal{B}^a(E, F)$ is bounded (right) \mathcal{A} -linear map. When $E = F$, we denote $\mathcal{B}^a(E, F)$ as $\mathcal{B}^a(E)$, constituting a C^* -algebra. Now, let us revisit the definition of two sided Hilbert C^* -modules.

Definition 2.7. A Hilbert \mathcal{A} - \mathcal{B} -module E is a Hilbert \mathcal{B} -module which possess a non-degenerate left action of \mathcal{A} on E , that is, there exists a $*$ -homomorphism $\pi : \mathcal{A} \rightarrow$

$\mathcal{B}^a(E)$ such that $E = \overline{\text{span}}\{\pi(a)\xi : a \in \mathcal{A}, \xi \in E\}$. For simplicity, denote $\pi(a)\xi$ by $a\xi$, establishing the left \mathcal{A} -action.

For Hilbert \mathcal{A} - \mathcal{B} -module E , the set of all adjointable \mathcal{A} - \mathcal{B} -linear operators on E forms a C^* -algebra and it is denoted by $\mathcal{B}^{a,bil}(E)$. Analogous to Hilbert spaces, we can define the direct sum of Hilbert bimodules.

Definition 2.8. Let $(E_i)_{i \in \mathbb{I}}$ be a family of Hilbert \mathcal{A} - \mathcal{B} -modules. The direct sum $E = \bigoplus_{i \in \mathbb{I}} E_i = \{(\xi_i) : \sum_{i \in \mathbb{I}} \langle \xi_i, \xi_i \rangle \text{ converges in } \mathcal{A}\}$ which forms a \mathcal{A} - \mathcal{B} -module in a natural way. Define the inner product $\langle (\xi_i), (\eta_i) \rangle = \sum_{i \in \mathbb{I}} \langle \xi_i, \eta_i \rangle$ which turns E into a Hilbert \mathcal{A} - \mathcal{B} -module.

This Hilbert module E is said to be direct sum of Hilbert modules E_i 's.

Now, we recall the construction of the exterior and interior tensor product of Hilbert bimodules. For C^* -algebras, we denote by $\mathcal{A} \otimes \mathcal{B}$ the completion of the algebraic tensor product $\mathcal{A} \otimes_{\text{alg}} \mathcal{B}$ with respect to the spatial or minimal C^* -norm. Since we are working with Hilbert \mathcal{A} - \mathcal{B} -modules, where $\mathcal{A} = \mathcal{B}$, we discuss these concepts in this specific case only.

Definition 2.9. Let E and F be Hilbert \mathcal{A} - \mathcal{A} -modules. Consider the algebraic tensor product $E \otimes_{\text{alg}} F$ with the natural $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$ - $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$ -action $(\xi \otimes \eta)(a \otimes b) = \xi a \otimes \eta b$ and $(a \otimes b)(\xi \otimes \eta) = a\xi \otimes b\eta$. Define

$$\langle \xi_1 \otimes \eta_1, \xi_2 \otimes \eta_2 \rangle = \langle \eta_1, \eta_2 \rangle \otimes \langle \xi_1, \xi_2 \rangle \text{ for all } \xi_1, \xi_2 \in E \text{ and } \eta_1, \eta_2 \in F.$$

This sesquilinear form $\langle \cdot, \cdot \rangle$ makes $E \otimes_{\text{alg}} F$ into an inner product $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$ - $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$ -module. Performing double completion, we obtain $E \otimes F$ as a Hilbert $\mathcal{A} \otimes \mathcal{A}$ - $\mathcal{A} \otimes \mathcal{A}$ -module. We call $E \otimes F$ the *exterior tensor product* of E and F .

Definition 2.10. Let E and F be Hilbert \mathcal{A} - \mathcal{A} -modules. Consider the algebraic tensor product $E \otimes_{\text{alg}} F$ with the natural right \mathcal{A} -action $(\xi \otimes \eta)a = \xi \otimes \eta a$ and define

$$\langle \xi_1 \otimes \eta_1, \xi_2 \otimes \eta_2 \rangle = \langle \eta_1, \langle \xi_1, \xi_2 \rangle \eta_2 \rangle \text{ for all } \xi_1, \xi_2 \in E \text{ and } \eta_1, \eta_2 \in F.$$

This sesquilinear form $\langle \cdot, \cdot \rangle$ makes $E \otimes_{\text{alg}} F$ into a semi-inner product \mathcal{A} -module. Let $N = \{\zeta \in E \otimes_{\text{alg}} F : \langle \zeta, \zeta \rangle = 0\}$. The *interior tensor product* of E and F is defined to be the completion of $E \otimes_{\text{alg}} F/N$ and it is denoted by $E \odot F$.

We denote the equivalence class of $\xi \otimes \eta$ in $E \odot F$ by $\xi \odot \eta$. Observe that $E \odot F$ is a Hilbert \mathcal{A} - \mathcal{A} -module with the natural left \mathcal{A} -action $a(\xi \odot \eta) = a\xi \odot \eta$. It is interesting to note that the set N in the previous definition is nothing but the subspace generated by the elements of the form $\xi \otimes a\eta - \xi a \otimes \eta$.

In [26, Theorem 5.2], Paschke has described the structure of CP maps on unital C^* -algebras generalizing the famous Stinespring's theorem ([41]). Here we provide a brief overview of the construction. Let τ be a CP map on a unital C^* -algebra \mathcal{A} . Consider the algebraic tensor product $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$. For $a, b, c, d \in \mathcal{A}$, define $\langle a \otimes b, c \otimes d \rangle = b^* \tau(a^* c) d$. Then $\langle \cdot, \cdot \rangle$ is a semi-inner product on $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}$. Let $N = \{z \in \mathcal{A} \otimes_{\text{alg}} \mathcal{A} : \langle z, z \rangle = 0\}$ and E be the completion of $\mathcal{A} \otimes_{\text{alg}} \mathcal{A}/N$. Then E is the Hilbert \mathcal{A} - \mathcal{A} -module in a natural way. Let $\xi = \mathbf{1} \otimes \mathbf{1} + N$. It follows that $\tau(a) = \langle \xi, a\xi \rangle$ for all $a \in \mathcal{A}$. Moreover, ξ is cyclic, meaning, $E = \overline{\text{span}} \mathcal{A}\xi\mathcal{A}$. The pair (E, ξ) is referred to as the GNS-construction of τ , with E being the GNS-module of τ . The CP map τ corresponding to the GNS-construction (E, ξ) is denoted by τ_ξ and plays a central role in our discussion of joint and outer spectral radii. We record the following two properties of τ_ξ which will be used frequently: For $\xi, \eta \in E$, $\|\tau_\xi\| = \|\xi\|^2$ and $\tau_\xi \circ \tau_\eta = \tau_{\eta \odot \xi}$.

Definition 2.11. Let τ be a CP on unital C^* -algebra \mathcal{A} and E be a Hilbert \mathcal{A} - \mathcal{A} -module with an element $\xi \in E$. We call the pair (E, ξ) to be a *GNS-representation* of τ if $\tau(a) = \langle \xi, a\xi \rangle$ for all $a \in \mathcal{A}$. It is said to be *minimal* if $E = \overline{\text{span}} \mathcal{A}\xi\mathcal{A}$.

Note that the GNS-module in the GNS-construction is minimal. For two minimal GNS-representations (E, ξ) and (F, η) of a CP map τ , the mapping $\xi \mapsto \eta$ extends as an \mathcal{A} - \mathcal{A} -linear unitary transformation from E to F . Consequently, the GNS-representation is unique up to (unitary) isomorphism.

We conclude our discussion on Hilbert modules with the definition of full fock module over a Hilbert bimodule, originally introduced by Pimsner [32] and Speicher [40]. It assumes a pivotal role in our discussion on joint and outer spectral radii and their characterisation. For detailed theory, refer to [39, Chapter 6].

Definition 2.12. Let \mathcal{A} be a unital C^* -algebra and E be a Hilbert \mathcal{A} - \mathcal{A} -module. The *full fock module* over E is defined to be the Hilbert \mathcal{A} - \mathcal{A} -module $\mathcal{F}(E) = \bigoplus_{n=0}^{\infty} E^{\odot n}$ where $E^{\odot 0} = \mathcal{A}$ and $\omega = 1 \in E^{\odot 0}$.

3. JOINT AND OUTER SPECTRAL RADII ON HILBERT C^* MODULES

Before coming to defining spectral radii for subsets of Hilbert C^* -modules we prove some results on positive maps. Elements of Hilbert C^* -modules generate completely positive maps. However, in the following subsection, we can deal with general positive maps. We begin with generalizing some of Popescu's results (see [33, Theorem 5.9]) to positive maps on general C^* -algebras. We also show that spectral radius is a spectral point for positive maps.

3.1. Positive maps. Popescu considered only w^* -continuous positive maps on the algebra $\mathcal{B}(\mathcal{H})$ of all bounded operators on a Hilbert space and some of his proofs does use w^* -continuity. We need these results for general positive maps on arbitrary C^* -algebras. So we provide the proofs.

To begin with note that, φ is a positive map on a unital C^* -algebra \mathcal{A} , then φ^n is positive for all $n \in \mathbb{N}$ and we have $\|\varphi^n\| = \|\varphi^n(1)\|$. Hence, the spectral radius of φ is

$$r(\varphi) = \lim_{n \rightarrow \infty} \|\varphi^n(1)\|^{\frac{1}{n}}. \quad (3.1)$$

Recall that, for $v \in \mathcal{A}$, we denote the map $a \mapsto v^*av$ by α_v . With this notation we have the following definition.

Definition 3.1. A positive map φ on \mathcal{A} is said to be *elementarily similar* to a positive map ψ on \mathcal{A} if there exists an invertible $a \in \mathcal{A}$ such that

$$\varphi = \alpha_a^{-1} \circ \psi \circ \alpha_a,$$

where α_a denotes the elementary CP map: $\alpha_a(x) = a^*xa$ for all $x \in \mathcal{A}$.

Now, we have some of the results of Popescu [Theorem 5.9, [33]] in our setting.

Theorem 3.2. *Let φ be a positive map on a unital C^* -algebra \mathcal{A} and s be any positive real number. Then the following statements are equivalent:*

- (1) $r(\varphi) < s$;
- (2) *There exists $w \geq 0$ such that $\varphi(w) = s(w - 1)$;*
- (3) *There exists strictly positive $v \in \mathcal{A}$ such that $\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| < s$;*
- (4) *There exists a positive map ψ on \mathcal{A} which is elementarily similar to φ such that $\|\psi\| < s$.*

Proof. (1) \Rightarrow (2): From (1), we have $r(\varphi/s) < 1$, which implies $\lim_{n \rightarrow \infty} \|(\frac{\varphi}{s})^n\|^{1/n} < 1$. Therefore, the power series $\sum_{n \geq 0} (\frac{\varphi}{s})^n z^n$ has radius of convergence greater than 1. Hence the series $\sum_{n \geq 0} (\frac{\varphi}{s})^n$ converges. Let $w = \sum_{n \geq 0} (\frac{\varphi}{s})^n (1)$. Since $\varphi^n(1) \geq 0$ for all $n \geq 0$, we have $w \geq 0$. Further,

$$w - 1 = \frac{\varphi}{s} \left(\sum_{n \geq 0} \left(\frac{\varphi}{s} \right)^n (1) \right) = \frac{\varphi}{s}(w) \implies \varphi(w) = s(w - 1).$$

This proves (2).

(2) \Rightarrow (3): Since $w = 1 + \frac{\varphi}{s}(w) \geq 1$, w is invertible. Consider $v = w^{1/2}$. Then

$$\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| = \|\alpha_v^{-1} \circ \varphi \circ \alpha_v(1)\| = \|s(1 - w^{-1})\| < s,$$

where the last inequality follows from $1 - w^{-1} < 1$. This proves (3).

(3) \Rightarrow (4): Let us take $\psi = \alpha_v^{-1} \circ \varphi \circ \alpha_v$. Then the implication follows trivially.

(4) \Rightarrow (1): Since ψ is elementarily similar to φ , we have $\psi = \alpha^{-1} \circ \varphi \circ \alpha$ for some elementary, invertible, CP map α . Then

$$\|\psi^n\| = \|\alpha^{-1} \varphi^n \alpha\| \leq \|\alpha^{-1}\| \|\alpha\| \|\varphi^n\| \implies \|\psi^n\|^{1/n} \leq \|\alpha^{-1}\|^{1/n} \|\alpha\|^{1/n} \|\varphi^n\|^{1/n}.$$

Taking limit as $n \rightarrow \infty$, we get $r(\psi) \leq r(\varphi)$. Similarly, we can obtain $r(\varphi) \leq r(\psi)$. Therefore, $r(\varphi) = r(\psi) \leq \|\psi\| < s$, which proves (1). \square

We have the following immediate corollary. This will be useful in characterizing outer spectral radius.

Corollary 3.3. *Let φ be a positive map on a unital C^* -algebra \mathcal{A} . Then*

$$\begin{aligned} r(\varphi) &= \inf\{\|\psi\| : \psi \text{ is elementarily similar to } \varphi\} \\ &= \inf\{\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| : v \in \mathcal{A}_+ \text{ is invertible}\}. \end{aligned} \quad (3.2)$$

Proof. For any ψ which is elementary similar to φ , there is an elementary, invertible, CP map α on \mathcal{A} such that $\psi = \alpha^{-1} \varphi \alpha$. Therefore, $\|\psi\| \geq r(\psi) = r(\alpha^{-1} \varphi \alpha) = r(\varphi)$. Now for any $s > r(\varphi)$, Theorem 3.2 guarantees existence of a CP map ψ which is elementarily similar to φ such that $\|\psi\| < s$. Hence,

$$r(\varphi) = \inf\{\|\psi\| : \psi \text{ is elementarily similar to } \varphi\}.$$

Similarly, $r(\varphi) = \inf\{\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| : v \in \mathcal{A}_+ \text{ is invertible}\}$ can be proved using ((1) \Rightarrow (3)) part of Theorem 3.2. \square

With the help of the Corollary 3.3, we can directly establish the following result, which was originally proved for entry-wise positive matrices by Wielandt [42] and for positive maps on C^* -algebras by Friedland ([42]) by entirely different methods.

Theorem 3.4. *Let φ be a positive map on a unital C^* -algebra \mathcal{A} . Then*

$$r(\varphi) = \inf\{r(w^{-1} \varphi(w)) : w \in \mathcal{A} \text{ is strictly positive}\}.$$

Proof. Since norm of a positive element is equal to its spectral radius and $r(ab) = r(ba)$ for all $a, b \in \mathcal{A}$, for any strictly positive $v \in \mathcal{A}$, we infer that

$$\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| = \|\alpha_v^{-1} \circ \varphi \circ \alpha_v(1)\| = \|v^{-1} \varphi(v^2) v^{-1}\| = r(v^{-1} \varphi(v^2) v^{-1}) = r(w^{-1} \varphi(w)),$$

where $w = v^2$. Now the result follows by (3.2). \square

It is a natural question as to when is the infimum in (3.2) is attained, that is, when can we find a positive map ψ which is elementarily similar to φ with norm equal to $r(\varphi)$. We give a pair of equivalent conditions that yield an affirmative answer. We will revisit this inquiry in Section 4.

Theorem 3.5. *Let φ be a positive map on a unital C^* -algebra \mathcal{A} with $r = r(\varphi)$. Then the following statements are equivalent:*

- (1) *There is a positive map ψ on \mathcal{A} which is elementarily similar to φ with $\|\psi\| = r$;*
- (2) *There exists a strictly positive $w \in \mathcal{A}$ such that $\varphi(w) \leq r w$;*
- (3) *There is a strictly positive $v \in \mathcal{A}$ such that $\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| = r$.*

Proof. (1) \Rightarrow (2): Since ψ is elementarily similar to φ with $\|\psi\| = r$, there is $a \in \mathcal{A}$ such that $\psi = \alpha_a^{-1} \circ \varphi \circ \alpha_a$. Let $w = a^*a$. Clearly, $w \in \mathcal{A}$ is positive and invertible. Further,

$$r w - \varphi(w) = r a^*a - \varphi(a^*a) = a^*(r 1 - \alpha_a^{-1} \varphi(a^*a) \alpha_a^{-1})a = a^*(r 1 - \psi(1))a.$$

Now as $\|\psi(1)\| = \|\psi\| = r$ and $\psi(1)$ is positive, we have

$$r w - \varphi(w) = a^*(r 1 - \psi(1))a \geq 0.$$

(2) \Rightarrow (3): Suppose there is a positive, invertible $w \in \mathcal{A}$ such that $\varphi(w) \leq r w$. Let $v = w^{1/2} \in \mathcal{A}$. Clearly, v is positive, invertible and we have

$$\|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| = \|(\alpha_v^{-1} \circ \varphi \circ \alpha_v)(1)\| = \|v^{-1} \varphi(v^2) v^{-1}\| = \|w^{-1/2} \varphi(w) w^{-1/2}\|.$$

Since $\varphi(w) \leq r w$, $w^{-1/2} \varphi(w) w^{-1/2} \leq r 1$ and hence

$$r = r(\varphi) = r(\alpha_v^{-1} \circ \varphi \circ \alpha_v) \leq \|\alpha_v^{-1} \circ \varphi \circ \alpha_v\| = \|w^{-1/2} \varphi(w) w^{-1/2}\| \leq r.$$

This proves (3).

(3) \Rightarrow (1): Let $\psi = \alpha_v^{-1} \circ \varphi \circ \alpha_v$, then the implication follows trivially. \square

We conclude this subsection with a compelling application of Theorem 3.2. Recall that due to the Perron-Frobenius theorem, as generalized by Evans and Høegh-Krohn, the spectral radius is an eigenvalue for positive maps on finite dimensional C^* -algebras. A positive map on an infinite dimensional C^* -algebra may not possess any eigenvalue. For instance, consider the unital C^* -algebra $\mathcal{B}(\ell^2)$ and the positive map defined by the right shift operator V on ℓ^2 , denoted as $\varphi(X) = VXV^*$ for all $X \in \mathcal{B}(\ell^2)$. Using Gelfand's formula for the spectral radius it is not hard to see that the spectral radius of φ is 1. Here, φ is injective but $\varphi^n(X)$ converges to zero in the weak operator topology as n tends to infinity. Therefore, φ has no non-trivial eigenvalue. In particular, it has no non-trivial fixed point. So, the spectral radius is not in the point spectrum.

We show that the spectral radius of a positive map on a unital C^* -algebra is an element in the spectrum, though it need not be an eigenvalue. It can be considered as an infinite dimensional extension of Perron-Frobenius theorem for positive maps on finite dimensional C^* -algebras (Theorem 2.4).

Theorem 3.6. *Let φ be a positive map on a unital C^* -algebra \mathcal{A} . Then $r(\varphi) \in \sigma(\varphi)$.*

Proof. If $r(\varphi) = 0$, there is nothing to prove. Therefore, without loss of generality, we assume $r(\varphi) = 1$. Suppose that 1 does not belong to $\sigma(\varphi)$. Then $1 - t\varphi$ is invertible for all $t \in [0, 1]$. Let us define $w_t = (1 - t\varphi)^{-1}(1) \in \mathcal{A}$. Since $x \mapsto x^{-1}$ is continuous on $\mathcal{G}(\mathcal{A})$, the assignment $t \mapsto w_t$ defines a continuous map from $[0, 1] \rightarrow \mathcal{A}$. For $0 \leq t < 1$, $r(t\varphi) = tr(\varphi) = t < 1$, hence we have the power series expression

$$w_t = 1 + (t\varphi)(1) + (t\varphi)^2(1) + \dots$$

This shows that $w_t \geq 0$ and $w_t - 1 = t\varphi(w_t)$ for all $t \in [0, 1)$. Therefore, by continuity, we have $\varphi(w_1) = w_1 - 1$. Thus, there exists $w_1 \geq 0$ in \mathcal{A} such that $\varphi(w_1) = w_1 - 1$, which is equivalent to $r(\varphi) < 1$ by Theorem 3.2. This contradicts $r(\varphi) = 1$. \square

3.2. Subsets of Hilbert C^* -modules. Here we introduce and study notions of joint spectral radius and outer spectral radius for subsets of Hilbert C^* -modules. Let E be a Hilbert \mathcal{A} - \mathcal{A} -module where \mathcal{A} is a unital C^* -algebra. Let G be any bounded subset of E . Notice that \mathcal{A} itself is a Hilbert \mathcal{A} - \mathcal{A} -module such that $\|a_1 \odot \cdots \odot a_n\| = \|a_1 \dots a_n\|$ for any $a_1, \dots, a_n \in \mathcal{A}$. This is because

$$\|a_1 \odot \cdots \odot a_n\|^2 = \|\tau_{a_1 \odot \cdots \odot a_n}\|^2 = \|\tau_{a_n} \circ \cdots \circ \tau_{a_1}\|^2 = \|\tau_{a_1 \dots a_n}\|^2 = \|a_1 \dots a_n\|^2.$$

Hence, taking inspiration from the notion of joint spectral radius as in (1.1), it is natural to define the joint spectral radius of any subset G of E using the interior tensor product as follows.

Definition 3.7. For any subset G of a Hilbert \mathcal{A} - \mathcal{A} -module E , the *joint spectral radius* of \mathbf{G} , denoted by $\rho_E(\mathbf{G})$, is defined by

$$\rho_E(G) = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \cdots \odot \xi_n\|^{1/n}.$$

Now the question is, “Why does the limit in the definition exists?”. To answer this we observe that for any such bounded CP subset G of E , we have the corresponding bounded set $S := \mathfrak{T}_G := \{\tau_\xi : \xi \in G\}$ of CP maps in the unital Banach algebra $\mathcal{B}(\mathcal{A})$, where τ_ξ is the map $a \mapsto \langle \xi, a\xi \rangle$ on \mathcal{A} . Hence, using the Rota-Strang theory of the joint spectral radius in particular for this $S \subseteq \mathcal{B}(\mathcal{A})$, we have

$$\rho(S) = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in S} \|\tau_{\xi_n} \circ \cdots \circ \tau_{\xi_1}\|^{1/n}.$$

Since for a Hilbert bimodule E with $\xi, \eta \in E$, $\tau_\xi \circ \tau_\eta = \tau_{\eta \odot \xi}$ and $\|\tau_\xi\| = \|\xi\|_E^2$, it follows that

$$\sup_{\xi_1, \dots, \xi_n \in S} \|\tau_{\xi_n} \circ \cdots \circ \tau_{\xi_1}\|^{1/n} = \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \cdots \odot \xi_n\|^{2/n}.$$

Therefore, $\lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \cdots \odot \xi_n\|^{2/n}$ exists and

$$\rho_E(G)^2 = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \cdots \odot \xi_n\|^{2/n} = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in S} \|\tau_{\xi_n} \circ \cdots \circ \tau_{\xi_1}\|^{1/n} = \rho(S).$$

Hence, $\rho_E(G)$ is well defined and it is the square root of the joint spectral radius of the subset \mathfrak{T}_G of $\mathcal{B}(\mathcal{A})$, that is, we have

$$\rho_E(G) = \sqrt{\rho(\mathfrak{T}_G)}. \quad (3.3)$$

If $G = \{\xi_1, \dots, \xi_d\}$ consists of finitely many elements, then we denote $\rho_E(G)$ by $\rho_E(\xi_1, \dots, \xi_d)$.

Remark 3.8. Given distinct elements $\xi_1, \dots, \xi_d \in E$, we can talk about the joint spectral radius of the set $G = \{\xi_1, \dots, \xi_d\} \subseteq E$ and the joint spectral radius of the singleton set $\{(\xi_1, \dots, \xi_d)\}$ of E^d . For a Hilbert bimodule E^d with $(\xi_1, \dots, \xi_d) \in E$, we will use the notation $\rho_{E^d}(\xi_1, \dots, \xi_d)$ instead of $\rho_{E^d}((\xi_1, \dots, \xi_d))$ to denote the joint spectral radius of the singleton subset $\{(\xi_1, \dots, \xi_d)\}$ of E^d . Notice that, it follows from the Definition 3.7 that,

$$\rho_E(\xi_1, \dots, \xi_d) = \lim_{n \rightarrow \infty} \sup_{1 \leq i_1, \dots, i_n \leq d} \|\xi_{i_1} \odot \cdots \odot \xi_{i_n}\|^{1/n}$$

and

$$\rho_{E^d}(\xi_1, \dots, \xi_d) = \lim_{n \rightarrow \infty} \left\| \sum_{i_1, \dots, i_n} |\xi_{i_1} \odot \cdots \odot \xi_{i_n}|^2 \right\|^{1/2n}.$$

Clearly, the two quantities in general need not be equal. Hence, we should be conscious that $\rho_E(\xi_1, \dots, \xi_d)$ and $\rho_{E^d}(\xi_1, \dots, \xi_d)$ are different. This also justifies keeping the Hilbert bimodule under consideration as a suffix for ρ .

Recall the Rota-Strang characterization of joint spectral radius (1.2). We now establish a similar characterisation for joint spectral radius of a subset of Hilbert C^* -module.

Theorem 3.9. *Let \mathcal{A} be a unital C^* -algebra and $G \subseteq E$ be any non-empty bounded subset of a Hilbert \mathcal{A} - \mathcal{A} -module E . Let \mathfrak{N} denote the family of all norms \mathcal{N} on the full Fock module $\mathcal{F}(E)$ which are equivalent to the usual norm and satisfy the inequality $\mathcal{N}(X \odot Y) \leq \mathcal{N}(X)\mathcal{N}(Y)$ for all $X \in E^{\odot n}$ and $Y \in E^{\odot m}$, with $m, n \geq 0$. Then*

$$\rho_E(G) = \inf_{\mathcal{N} \in \mathfrak{N}} \sup_{\xi \in G} \mathcal{N}(\xi). \quad (3.4)$$

Proof. Let $\mathcal{N} \in \mathfrak{N}$ be arbitrary. Then there is $k > 0$ such that $k^{-1}\|\cdot\|_{\mathcal{F}(E)} \leq \mathcal{N}(\cdot) \leq k\|\cdot\|_{\mathcal{F}(E)}$. For any $n \in \mathbb{N}$ and $\xi_1, \dots, \xi_n \in G$,

$$\|\xi_1 \odot \dots \odot \xi_n\|_{\mathcal{N}}^{\frac{1}{n}} \leq k^{\frac{1}{n}} \mathcal{N}(\xi_1 \odot \dots \odot \xi_n)^{\frac{1}{n}} \leq k^{\frac{1}{n}} (\mathcal{N}(\xi_1) \dots \mathcal{N}(\xi_n))^{\frac{1}{n}} \leq k^{\frac{1}{n}} \sup_{\xi \in G} \mathcal{N}(\xi).$$

Therefore,

$$\sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \dots \odot \xi_n\|_{\mathcal{N}}^{\frac{1}{n}} \leq k^{\frac{1}{n}} \sup_{\xi \in G} \mathcal{N}(\xi).$$

Taking limit as $n \rightarrow \infty$, it follows from the Definition 3.7, that $\rho_E(G) \leq \sup_{\xi \in G} \mathcal{N}(\xi)$ for all $\mathcal{N} \in \mathfrak{N}$. Hence, we have

$$\rho_E(G) \leq \inf_{\mathcal{N} \in \mathfrak{N}} \sup_{\xi \in G} \mathcal{N}(\xi).$$

Now, from (1.2) and (3.3) it follows that

$$\rho_E(G) = \sqrt{\rho(\mathfrak{T}_G)} = \sqrt{\inf_{N \in \mathfrak{N}_{\mathcal{B}(\mathcal{A})}} \sup_{\tau \in \mathfrak{T}_G} N(\tau)} = \inf_{N \in \mathfrak{N}_{\mathcal{B}(\mathcal{A})}} \sup_{\tau \in \mathfrak{T}_G} \sqrt{N(\tau)},$$

where $\mathfrak{T}_G = \{\tau_\xi : \xi \in G\}$. Let $N \in \mathfrak{N}_{\mathcal{B}(\mathcal{A})}$ be arbitrary. Then it can easily be verified that,

$$\mathcal{N}_F((X_0, X_1, \dots)) = \sqrt{\sum_{n=0}^{\infty} N(\tau_{X_n})} \text{ for all } X_n \in E^{\odot n}$$

defines a norm on $\mathcal{F}(E)$, which is Clearly, induced by the given N . Since N is equivalent to the operator norm on $\mathcal{B}(\mathcal{A})$, there is $j > 0$ such that $j^{-1}\|\cdot\| \leq N(\cdot) \leq j\|\cdot\|$. Thus, for any $(X_n) \in \mathcal{F}(E)$, we have

$$j^{-1}\|(X_n)\|^2 = j^{-1} \sum_{n=0}^{\infty} \|X_n\|^2 = \sum_{n=0}^{\infty} j^{-1} \|\tau_{X_n}\| \leq \sum_{n=0}^{\infty} N(\tau_{X_n}) = \mathcal{N}_F((X_n))^2.$$

Similarly,

$$\mathcal{N}_F((X_n))^2 \leq j\|(X_n)\|^2.$$

Hence, \mathcal{N}_F is equivalent to the usual norm on $\mathcal{F}(E)$. Now let $X \in E^{\odot n}$ and $Y \in E^{\odot m}$ be arbitrary. Then $X \odot Y \in E^{\odot n+m}$ and

$$\mathcal{N}_F(X \odot Y) = \sqrt{N(\tau_{X \odot Y})} = \sqrt{N(\tau_Y \circ \tau_X)} \leq \sqrt{N(\tau_Y)} \sqrt{N(\tau_X)} = \mathcal{N}_F(X) \mathcal{N}_F(Y).$$

Therefore, \mathcal{N}_F belongs to the family \mathfrak{N} . This shows that

$$\inf_{\mathcal{N} \in \mathfrak{N}} \sup_{\xi \in G} \mathcal{N}(\xi) \leq \sup_{\xi \in G} \mathcal{N}_F(\xi) = \sup_{\xi \in G} \sqrt{N(\tau_\xi)},$$

for all $N \in \mathfrak{N}_{\mathcal{B}(\mathcal{A})}$. Thus,

$$\inf_{N \in \mathfrak{N}} \sup_{\xi \in G} \mathcal{N}(\xi) \leq \inf_{N \in \mathfrak{N}_{\mathcal{B}(\mathcal{A})}} \sup_{\xi \in G} \sqrt{N(\tau_\xi)} = \rho_E(G).$$

Hence the proof. \square

Taking inspiration from the notion of outer spectral radius for tuples of matrices, we define the same for sequences of elements of Hilbert C^* -bimodules.

Definition 3.10. Let $G = (\xi_1, \xi_2, \dots)$ be a finite or countably infinite sequence of elements from a Hilbert \mathcal{A} - \mathcal{A} -module E such that

$$\tau_G(a) := \sum_{i=1}^{\infty} \langle \xi_i, a \xi_i \rangle, \quad \forall a \in \mathcal{A} \quad (3.5)$$

defines a CP map on \mathcal{A} . Then, the *outer spectral radius* of G , denoted by $\widehat{\rho}_E(\mathbf{G})$, is defined as $\widehat{\rho}_E(G) = \sqrt{r(\tau_G)}$.

It is to be noted that the outer spectral radius is defined for sequences, instead of subsets, as repetition of same elements would make a difference.

It is easy to see that for $G = (\xi_1, \xi_2, \dots)$, τ_G as in 3.5 defines a CP map if and only if $\sum_{i=1}^{\infty} |\xi_i|^2$ is convergent in \mathcal{A} . It is clear that countability of G is a necessity here. When $G = (\xi_1, \dots, \xi_d)$, we write $\widehat{\rho}_E(G)$ as $\widehat{\rho}_E(\xi_1, \dots, \xi_d)$.

Proposition 3.11. Let $G = (\xi_1, \xi_2, \dots)$ be a sequence of elements from a Hilbert \mathcal{A} - \mathcal{A} -module E such that $\sum_{i=1}^{\infty} |\xi_i|^2$ converges. Then

$$\widehat{\rho}_E(G) = \lim_{n \rightarrow \infty} \left\| \sum_{i_1, \dots, i_n \in \mathbb{N}} |\xi_{i_1} \odot \dots \odot \xi_{i_n}|^2 \right\|^{1/2n} \quad (3.6)$$

$$= \lim_{n \rightarrow \infty} \sup \left\{ \left\| \sum_{\xi \in \mathcal{F}} |\xi|_{E^{\odot n}}^2 \right\|^{1/2} : \mathcal{F} \subseteq \mathcal{P}_n(G) \text{ is finite} \right\}^{1/n} \quad (3.7)$$

Proof. We have $G = (\xi_1, \xi_2, \dots)$ such that $\sum_{i=1}^{\infty} |\xi_i|^2$ converges. Clearly, G is associated to a CP map τ_G on \mathcal{A} defined by $\tau_G(a) = \sum_{i=1}^{\infty} \langle \xi_i, a \xi_i \rangle$ for all $a \in \mathcal{A}$. Consider the tuple $\xi = (\xi_1, \xi_2, \dots)$ from the Hilbert \mathcal{A} - \mathcal{A} -module $\bigoplus_{\mathbb{N}} E$. Observe that, the given CP map τ_G can be expressed as $\tau_G = \tau_\xi = \sum_{i=1}^{\infty} \tau_{\xi_i}$. Then we have

$$\widehat{\rho}(G)^2 = r(\tau_G) = \lim_{n \rightarrow \infty} \|\tau_G^n(1)\|^{1/n} = \left\| \sum_{i_1, \dots, i_n \in \mathbb{N}} |\xi_{i_1} \odot \dots \odot \xi_{i_n}|^2 \right\|^{1/n}.$$

Hence the proof. \square

The expression in (3.6) provides us with a Gelfand type formula for the outer spectral radius of G . Given a tuple $G = (\xi_1, \xi_2, \dots) \in \bigoplus_{\mathbb{N}} E$, by applying the identity (3.2) and utilizing Friedland's result (Theorem 3.4) with the CP map τ_G , we can drive the following two characterisations for the outer spectral radius of a set of elements in a Hilbert bimodule.

Proposition 3.12. *Let $G = (\xi_1, \xi_2, \dots)$ be a sequence of elements from a Hilbert \mathcal{A} - \mathcal{A} -module E such that $\sum_{i=1}^{\infty} |\xi_i|^2$ converges. Then,*

$$\begin{aligned} \widehat{\rho}_E(G) &= \inf \left\{ \left\| \sum_{i=1}^{\infty} |v \xi_i v^{-1}|^2 \right\|^{1/2} : v \in \mathcal{A} \text{ is strictly positive} \right\} \\ &= \inf \left\{ r \left(\sum_{i=1}^{\infty} \langle w \xi_i w^{-1}, \xi_i \rangle \right)^{1/2} : w \in \mathcal{A} \text{ is strictly positive} \right\}. \end{aligned} \quad (3.8)$$

The following Proposition compares the joint spectral radius with the outer spectral radius.

Proposition 3.13. *Let E be a Hilbert \mathcal{A} - \mathcal{A} module. Then, for any $G = (\xi_1, \xi_2, \dots)$ satisfying $\sum_{i=1}^{\infty} |\xi_i|^2 < \infty$, we have $\rho_E(G) \leq \widehat{\rho}_E(G)$ and in particular if $G = (\xi_1, \dots, \xi_d)$,*

$$d^{-1/2} \widehat{\rho}_E(\xi_1, \dots, \xi_d) \leq \rho_E(\xi_1, \dots, \xi_d) \leq \widehat{\rho}_E(\xi_1, \dots, \xi_d).$$

Proof. From (3.6) and the definition of joint spectral radius, we observe that $\rho_E(G) \leq \widehat{\rho}_E(G)$. Further, it follows from the Proposition 3.11 that,

$$\begin{aligned} \widehat{\rho}_E(\xi_1, \dots, \xi_d) &= \lim_{n \rightarrow \infty} \left\| \sum_{1 \leq i_1, \dots, i_n \leq d} |\xi_{i_1} \odot \dots \odot \xi_{i_n}|^2 \right\|^{1/2n} \\ &\leq \sqrt{d} \lim_{n \rightarrow \infty} \sup_{1 \leq i_1, \dots, i_n \leq d} |\xi_{i_1} \odot \dots \odot \xi_{i_n}|^{1/n} \\ &= \sqrt{d} \rho_E(\xi_1, \dots, \xi_d). \end{aligned}$$

This completes the proof. \square

The following lemma is useful in proving an approximation theorem for the joint spectral radius.

Lemma 3.14. *Let E be a Hilbert \mathcal{A} - \mathcal{A} module with a bounded set $G \subset E$. Then for each $k \in \mathbb{N}$, $\rho_E(G^{(k)}) = \rho_E(G)^k$ where $G^{(k)} = \{\xi^{\otimes k} : \xi \in G\}$ is a bounded subset of Hilbert $\mathcal{A}^{\otimes k}$ - $\mathcal{A}^{\otimes k}$ -module $E^{\otimes k}$.*

Proof. For any $\xi_1, \dots, \xi_n \in E$, we have

$$\|\xi_1^{\otimes k} \odot \dots \odot \xi_n^{\otimes k}\| = \|\tau_{\xi_1^{\otimes k} \odot \dots \odot \xi_n^{\otimes k}}(1 \otimes \dots \otimes 1)\|^{1/2} = \|(\tau_{\xi_1 \odot \dots \odot \xi_n}(1))^{\otimes k}\|^{1/2} = \|\xi_1 \odot \dots \odot \xi_n\|^k.$$

Therefore, we infer that

$$\rho_E(G^{(k)}) = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1^{\otimes k} \odot \dots \odot \xi_n^{\otimes k}\|^{1/n} = \lim_{n \rightarrow \infty} \sup_{\xi_1, \dots, \xi_n \in G} \|\xi_1 \odot \dots \odot \xi_n\|^{k/n} = \rho_E(G)^k. \quad \square$$

In particular when G is finite, we can approximate the joint spectral radius by the outer spectral radius as follows.

Theorem 3.15. *Let E be a Hilbert \mathcal{A} - \mathcal{A} module with $\xi_1, \dots, \xi_d \in E$. Then,*

$$d^{-1/2k} \widehat{\rho}_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k})^{1/k} \leq \rho_E(\xi_1, \dots, \xi_d) \leq \widehat{\rho}_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k})^{1/k},$$

for every $k \geq 1$ and

$$\rho_E(\xi_1, \dots, \xi_d) = \lim_{k \rightarrow \infty} \widehat{\rho}_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k})^{1/k}. \quad (3.9)$$

Proof. By Lemma 3.14, $\rho_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k}) = \rho_E(\xi_1, \dots, \xi_d)^k$. Now, using Proposition 3.13, we have $d^{-1/2} \widehat{\rho}_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k}) \leq \rho_E(\xi_1, \dots, \xi_d)^k \leq \rho_E(\xi_1^{\otimes k}, \dots, \xi_d^{\otimes k})$. This proves the inequality and taking limit as $k \rightarrow \infty$, we get our desired approximation of the joint spectral radius. \square

We explain the notions described above in the particular case, where the Hilbert C^* -module E is equal to \mathcal{A} , with standard inner product: $\langle a, b \rangle = a^*b$ for a, b in \mathcal{A} . Let \mathcal{A} be a unital C^* -algebra with $a_1, \dots, a_d \in \mathcal{A}$. Consider the Hilbert \mathcal{A} - \mathcal{A} -module $E = \mathcal{A}$ and let $G = \{a_1, \dots, a_d\}$ be any subset of \mathcal{A} . Clearly,

$$\|a_1 \odot \dots \odot a_n\|^2 = \|\tau_{a_1 \odot \dots \odot a_n}\| = \|\tau_{a_n} \circ \dots \circ \tau_{a_1}\| = \|\tau_{a_1 \dots a_n}\| = \|a_1 \dots a_n\|^2.$$

Hence, the Definition 3.7 gives us that,

$$\rho_E(G) = \rho_{\mathcal{A}}(a_1, \dots, a_d) = \lim_{n \rightarrow \infty} \sup_{1 \leq i_1, \dots, i_n \leq d} \|a_{i_1} \dots a_{i_n}\|^{1/n}, \quad (3.10)$$

which is the familiar joint spectral radius as in (1.1). For $G = \{a_1, \dots, a_d\} \subseteq \mathcal{A}$, $\tau_G : \mathcal{A} \rightarrow \mathcal{A}$ will be of the form

$$\tau_G(x) = \sum_{i=1}^d a_i^* x a_i. \quad (3.11)$$

Observe that, by induction we obtain

$$\tau_G^n(x) = \sum_{1 \leq i_1, \dots, i_n \leq d} (a_{i_1} \dots a_{i_n})^* x (a_{i_1} \dots a_{i_n}) = \sum_I a_I^* x a_I.$$

Here, the multi-index I is running over the set $\{(i_1, \dots, i_n) : 1 \leq i_1, \dots, i_n \leq d\}$ and for $I = (i_1, \dots, i_n)$, we denote $a_I = a_{i_1} a_{i_2} \dots a_{i_n}$. Therefore, $\|\tau_G^n\| = \|\tau_G^n(1)\| = \left\| \sum_I a_I^* a_I \right\|$. Hence, the outer spectral radius of G has the expression

$$\widehat{\rho}_{\mathcal{A}}(G) = \sqrt{r(\tau_G)} = \lim_{n \rightarrow \infty} \|\tau_G^n\|^{1/2n} = \lim_{n \rightarrow \infty} \left\| \sum_I a_I^* a_I \right\|^{1/2n}. \quad (3.12)$$

Theorem 3.16. *Let a_1, a_2, \dots, a_d be elements in a unital C^* -algebra \mathcal{A} . Then,*

$$\widehat{\rho}_{\mathcal{A}}(a_1, \dots, a_d) = \inf \{ \|(va_1 v^{-1}, \dots, va_d v^{-1})\| : v \in \mathcal{A} \text{ is strictly positive} \} \quad (3.13)$$

$$= \inf \left\{ \left\| \sum_{i=1}^d |va_i v^{-1}|^2 \right\|^{1/2} : v \in \mathcal{A} \text{ is strictly positive} \right\}, \quad (3.14)$$

$$= \inf \left\{ r \left(\sum_{i=1}^{\infty} (a_i v^{-1})^* (va_i) \right)^{1/2} : v \in \mathcal{A} \text{ is strictly positive} \right\} \quad (3.15)$$

Furthermore,

$$\rho_{\mathcal{A}}(a_1, \dots, a_d) = \lim_{k \rightarrow \infty} \widehat{\rho}_{\mathcal{A}}(a_1^{\otimes k}, \dots, a_d^{\otimes k})^{1/k}. \quad (3.16)$$

Proof. Take $E = \mathcal{A}$ in Proposition 3.12 and Theorem 3.15. \square

Remark 3.17. For a tuple of $m \times m$ matrices (A_1, \dots, A_d) ,

$$\widehat{\rho}_{M_m}(A_1, \dots, A_d) = \sqrt{r \left(\sum_{i=1}^d \overline{A_i} \otimes A_i \right)} = \sqrt{r \left(\sum_{i=1}^d \overline{A_i^*} \otimes A_i^* \right)} = \widehat{\rho}_{M_m}(A_1^*, \dots, A_d^*).$$

Furthermore, if \mathcal{A} is a finite dimensional C^* -algebra, then for a tuple (a_1, a_2, \dots, a_d) of elements of \mathcal{A} , $\widehat{\rho}_{\mathcal{A}}(a_1, \dots, a_d) = \widehat{\rho}_{\mathcal{A}}(a_1^*, \dots, a_d^*)$. However, this equality does not generally hold if \mathcal{A} is not finite dimensional. For example, consider the unital C^* -algebra $\mathcal{A} = \mathcal{B}(\ell^2)$ and the isometries V, W on ℓ^2 defined as $V(e_n) = e_{2n+1}$ and $W(e_n) = e_{2n}$ on the standard basis. Let $\tau(X) = V^*XV + W^*XW$ and $\sigma(X) = VXV^* + WXW^*$. Then, $\widehat{\rho}_{\mathcal{A}}(V, W) = \sqrt{r(\tau)} = 2$, whereas $\widehat{\rho}_{\mathcal{A}}(V^*, W^*) = \sqrt{r(\sigma)} = 1$.

4. POSITIVE MAPS ON FINITE DIMENSIONAL ALGEBRAS

For the rest of our discussion, we focus on finite dimensional C^* -algebras. To begin with we look at general positive maps on finite dimensional algebras. We study the maximal part of a general positive map, and its analysis allows us to construct a positive eigenvector with Perron-Frobenius eigenvalue.

Let \mathcal{A} be a unital C^* -algebra with $a \in \mathcal{A}$. Consider the elementary CP map $\alpha_a(x) = a^*xa$, $x \in \mathcal{A}$. Applying Corollary 3.3, we obtain the following well-known formula for the spectral radius $r(a)$ of a :

$$r(a) = \inf\{\|b\| : b \in \mathcal{A} \text{ is similar to } a\} \quad (4.1)$$

$$= \inf\{\|vav^{-1}\| : v \in \mathcal{A}_+ \text{ is invertible}\}. \quad (4.2)$$

It is a natural question as to when the infimum in (4.1) can be attained. With help from the following Lemma, we show that in finite dimensions the boundedness of the set $\{(a/r(a))^n : n \geq 0\}$ ensures this.

Lemma 4.1. *Let $A \in M_m$ be a matrix with $r(A) = 1$. If there is $M > 0$ such that $\|A^n\| \leq M$ for all $n \in \mathbb{N}$, then there exists $P \in GL_m$ such that $\|PAP^{-1}\| = 1$.*

Proof. Let J be the Jordan form of the matrix A . Then $A = SJS^{-1}$ for some $S \in GL_m$. Let $\lambda \in \mathbb{C}$ be an eigenvalue of A and J_λ be the Jordan block corresponding to λ of size k . First, suppose $|\lambda| = 1$. Note that, $J_\lambda = \lambda I_k + N$ where N is the $k \times k$ nilpotent matrix with nilpotency k . Observe that,

$$J_\lambda^n = \lambda^n I_k + \binom{n}{1} \lambda^{n-1} N + \binom{n}{2} \lambda^{n-2} N^2 + \dots + \binom{n}{k-1} \lambda^{n-k+1} N^{k-1}.$$

When $k \geq 2$, since $\|A^n\| \leq M$, we have

$$n = |n\lambda^{n-1}| = |\langle e_1, J_\lambda^n e_2 \rangle| \leq \|J_\lambda^n\| \leq \|J_\lambda^n\| \leq \|S\| \|S^{-1}\| \|A^n\| \leq \|S\| \|S^{-1}\| M \text{ for all } n,$$

which is paradoxical. So, we must have $k = 1$ and hence for $|\lambda| = 1$, J_λ is of size 1. Thus,

the Jordan form of A looks like $J = \begin{bmatrix} D & & & \\ & J_1 & & \\ & & J_2 & \\ & & & \ddots \\ & & & & J_l \end{bmatrix}$ where D is a diagonal matrix with

diagonal entries of modulus 1 and each J_i is a Jordan block corresponding to eigenvalue of modulus less than 1.

Now, consider $|\lambda| < 1$. Choose $\epsilon > 0$ such that $|\lambda| + (k-1)\epsilon < 1$ and let $C = \text{diag}(c_1, \dots, c_k)$ where $c_j = \epsilon^{k-j+1} \forall j \in \{1, \dots, k\}$. We infer that

$$\|C J_\lambda C^{-1}\| = \left\| \lambda I_k + \epsilon \sum_{j=2}^k E_{(j-1)j} \right\| \leq |\lambda| + (k-1)\epsilon < 1.$$

where $\{E_{ij} : 1 \leq i, j \leq k\}$ is the standard basis of M_k consisting of elementary matrices. Define

$$T = \begin{bmatrix} I & & & \\ & C_1 & & \\ & & C_2 & \\ & & & \ddots \\ & & & & C_l \end{bmatrix} \text{ and } P = TS^{-1}.$$

Then, $\|PAP^{-1}\| = \|TJT^{-1}\| = \max\{\|D\|, \|C_j J_j C_j^{-1}\| : j \in \{1, \dots, l\}\} = 1$. \square

We now answer the above query in form of the following theorem.

Theorem 4.2. *Let \mathcal{A} be a finite dimensional C^* -algebra and $a \in \mathcal{A}$ with $r = r(a) > 0$. Then the following statements are equivalent:*

- (1) *There is $M > 0$ such that $\|\frac{a^n}{r^n}\| \leq M$ for all $n \in \mathbb{N}$.*
- (2) *There is $b \in \mathcal{A}$ which is similar to a and $\|b\| = r$.*
- (3) *There is strictly positive $v \in \mathcal{A}$ such that $\|v^{-1}av\| = r$.*

Proof. As \mathcal{A} is a finite dimensional C^* -algebra, by Structure Theorem, we have $\mathcal{A} = M_{n_1} \oplus \dots \oplus M_{n_d}$. Therefore, for given $a \in \mathcal{A}$, a is of the form $a = X_1 \oplus \dots \oplus X_d$ where each $X_i \in M_{n_i}$. Without loss of generality, we assume that $r = 1$.

(1) \Rightarrow (2): Suppose, for given $a \in \mathcal{A}$ with $r(a) = 1$, there is $M > 0$ such that $\|a^n\| \leq M$ for all $n \in \mathbb{N}$. Let $I = \{i : r(X_i) = 1, 1 \leq i \leq d\}$ and $K = \{i : r(X_i) < 1, 1 \leq i \leq d\}$. Since $r = \max\{r(X_i) : i \in \{1, \dots, d\}\} = 1$, we have $I \neq \emptyset$ and $I \cup K = \{1, \dots, d\}$. By Corollary 3.3,

$$\forall i \in K, \exists P_i \in GL_{n_i} \text{ such that } \|P_i X_i P_i^{-1}\| < 1$$

Now, for any $i \in I$, spectral radius of X_i is 1 and $\|X_i^n\| \leq \|a^n\| \leq M$ for all $n \in \mathbb{N}$. Then by Lemma 4.1

$$\exists P_i \in GL_{n_i} \text{ such that } \|P_i X_i P_i^{-1}\| = 1.$$

Define

$$p = \begin{bmatrix} P_1 & & & \\ & P_2 & & \\ & & \ddots & \\ & & & P_d \end{bmatrix} \text{ and } b = pap^{-1} \in \mathcal{A}.$$

Then, $\|b\| = \max\{\|P_i X_i P_i^{-1}\| : i \in \{1, \dots, d\}\} = 1$.

(2) \Rightarrow (3): Assume that there is $b \in \mathcal{A}$ which is similar to a with $\|b\| = 1$. Thus, there is an invertible $p \in \mathcal{A}$ such that $\|b\| = \|p^{-1}ap\| = 1$. Let τ be the elementary CP map α_a and $w = p^*p$. Then

$$w - \tau(w) = p^*p - a^*p^*pa = p^*(1 - p^{*-1}a^*p^*pap^{-1})p = p^*(1 - (p^{-1}ap)^*(p^{-1}ap))p \geq 0$$

Therefore, by Theorem 3.5, there exists positive, invertible $v \in \mathcal{A}$ such that $\|\alpha_v^{-1} \circ \tau \circ \alpha_v\| = 1$. Clearly,

$$\|\alpha_v^{-1} \circ \tau \circ \alpha_v\| = \|\alpha_{v^{-1}av}\| = \|v^{-1}av\|^2.$$

Thus there is a positive, invertible $v \in \mathcal{A}$ such that $\|v^{-1}av\| = 1$.

(3) \Rightarrow (1): Let $b = v^{-1}av$ and we have $\|b\| = 1$. Then for all $n \in \mathbb{N}$,

$$\|a^n\| = \|vb^n v^{-1}\| \leq \|v\| \|v^{-1}\| \|b\|^n = \|v\| \|v^{-1}\|.$$

This proves (1). \square

For a CP map τ on finite dimensional C^* -algebra, it is evident that boundedness of the sequence $\left\| \frac{\tau^n}{r(\tau)^n} \right\|$ is a necessary condition for existence of an elementarily similar CP map σ with $\|\sigma\| = r(\tau)$. But it is not sufficient, as evidenced by the following example.

Example 4.3. Let $\tau : M_2 \rightarrow M_2$ be a CP map defined by

$$\tau \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = \begin{bmatrix} a+d & 0 \\ 0 & 0 \end{bmatrix}.$$

It's easy to see that $\tau^n(I) = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}$. Therefore, $\|\tau^n\| = \|\tau^n(I)\| = 2$ for all $n \in \mathbb{N}$.

This shows that $r(\tau) = 1$ and $\left\| \frac{\tau^n}{r(\tau)^n} \right\|$ is bounded. Now, if there is a CP map $\tilde{\tau}$ which is elementarily similar to τ such that $\|\tilde{\tau}\| = r(\tau)$, then due to Theorem 3.5, there is a strictly positive $V = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2$ such that $\tau(V) \leq V$. But

$$\begin{bmatrix} a+d & 0 \\ 0 & 0 \end{bmatrix} \leq \begin{bmatrix} a & b \\ c & d \end{bmatrix} \implies \begin{bmatrix} -d & b \\ c & d \end{bmatrix} \geq 0 \implies d \leq 0,$$

which contradicts the strict positivity of d .

In their seminal paper [11], Evans and Høegh-Krohn explored Perron-Frobenius theory for positive maps on C^* -algebras. They showed that for any positive map φ on a finite dimensional C^* -algebra \mathcal{A} , the spectral radius $r(\varphi)$ is an eigenvalue of φ with a positive eigenvector. However, their proof is completely existential. We adopt a different approach and explicitly construct the positive eigenvector.

Note that when $r(\varphi) = 0$, there is n such that $\varphi^n = 0$ and $\varphi^{n-1} \neq 0$, hence $\varphi^{n-1}(1)$ is a positive eigenvector with eigenvalue equal to the spectral radius. Therefore, in the following we assume $r(\varphi) > 0$. The key to our construction is the ‘maximal part of a positive map’ defined as follows. This map was introduced by Pascoe [28] using matrices. The justification for the nomenclature will be clear from the discussion below.

Definition 4.4. Let φ be a positive map on a finite dimensional C^* -algebra \mathcal{A} with $r(\varphi) > 0$. The *maximal part* of φ is defined as

$$\widehat{\varphi} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\varphi^n}{\|\varphi^n\|_\beta}, \quad (4.3)$$

where β is a Jordan basis of φ and $\|\varphi^n\|_\beta := \|[\varphi^n]_\beta\|_\infty$.

The series in equation 4.3 is not well-defined if $r(\varphi)$ equals 0, as then φ becomes nilpotent. It is for this reason that we have assumed $r(\varphi) > 0$. Now we need to show that the series in 4.3 is convergent. Before doing that let us explain the norm appearing in the denominator. Suppose $m := \dim(\mathcal{A})$, and $[\varphi]_\gamma \in M_m$ is the matrix of φ with respect to a basis γ of \mathcal{A} . Now $\|\cdot\|_\infty$ denotes the supremum norm on M_m , that is,

$$\|(a_{ij})\|_\infty = \sup\{|a_{ij}| : i, j \in \{1, \dots, m\}\}.$$

As \mathcal{A} is a finite dimensional C^* -algebra, $\|\cdot\|_\gamma$ is equivalent to the usual operator norm $\|\cdot\|$ of the Banach algebra $\mathcal{B}(\mathcal{A})$. Currently, we are choosing Jordan basis of φ , meaning that on this basis the matrix of φ is a direct sum of Jordan blocks. By uniqueness of the Jordan decomposition the norm $\|\varphi^n\|_\beta := \|[\varphi^n]_\beta\|_\infty$ is independent of the choice of the Jordan basis. Consequently, the maximal part of φ is also independent of the choice of Jordan basis.

Let $x \in \mathcal{A}$ be a vector in a Jordan basis β of φ . Suppose that it is a generalized eigenvector of φ of rank k corresponding to eigenvalue λ . Then, taking powers of the

Jordan block, we see that for any $n \in \mathbb{N}$,

$$\varphi^n(x) = \sum_{j=0}^{k-1} \binom{n}{j} \lambda^{n-j} (\varphi - \lambda)^j x, \quad (4.4)$$

and $(\varphi - \lambda)^j x$ are Jordan basis vectors. Using this expression, it is possible to compute $\|\varphi^n\|_\beta$. Here it is convenient to have the following definition.

Definition 4.5. Let φ be a positive map on a finite dimensional C^* -algebra. The degeneracy index $d_{\varphi, \lambda}$ of an eigenvalue λ is defined to be the size of the largest Jordan block corresponding to λ , or equivalently, the power of the $(z - \lambda)$ in the factorization of the minimal polynomial $m(z)$ of φ . The *maximal degeneracy index* of φ , denoted by d_φ , is defined as

$$d_\varphi = \max\{d_{\varphi, \lambda} : |\lambda| = r(\varphi)\}.$$

Accordingly, the *maximal spectrum* of φ is set as

$$\{\lambda \in \sigma(\varphi) : |\lambda| = r(\varphi) \text{ and } d_{\varphi, \lambda} = d_\varphi\}.$$

Now from equation (4.4), since the binomial coefficients $\binom{n}{j}$ are increasing for $0 \leq j \leq [n/2]$, for large n , $\|[\varphi^n]_\beta\|_\infty$ occurs at the top right hand corner of the powers of Jordan blocks with maximal degeneracy index of maximal modulus spectral values. Then it is evident that for any Jordan basis β of φ , we have

$$\|\varphi^n\|_\beta = \|[\varphi^n]_\beta\|_\infty = \binom{n}{d_\varphi - 1} r(\varphi)^{n-d_\varphi+1}, \quad (4.5)$$

for all but finitely many n . To demonstrate that for a positive map φ on \mathcal{A} with $r(\varphi) > 0$, $\widehat{\varphi}$ is a well defined non-zero positive map, we need the following lemma.

Lemma 4.6. *Let $\{V_n\}$ be a sequence of positive $m \times m$ matrices such that*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N V_n = 0.$$

Then there exists a sub-sequence $\{V_{n_k}\}$ converging to 0.

Proof. Let $W_N = \frac{1}{N} \sum_{n=1}^N V_n$. Clearly, $\{W_N\}$ is a sequence of positive matrices converging to 0, in particular $\text{Tr}(W_N) \rightarrow 0$. Let $\liminf \text{Tr}(V_n) = t$. Given any $\epsilon > 0$, there is $k \in \mathbb{N}$ such that $\text{Tr}(V_n) > t - \epsilon$ for all $n > k$. Then for $N > k$,

$$\text{Tr}(W_N) = \frac{1}{N} \sum_{n=1}^k \text{Tr}(V_n) + \frac{1}{N} \sum_{n=k+1}^N \text{Tr}(V_n) > \frac{1}{N} \sum_{n=1}^k \text{Tr}(V_n) + (t - \epsilon)(1 - k/N)$$

Taking limit as $N \rightarrow \infty$, we obtain $(t - \epsilon)1 \leq 0$ for all $\epsilon > 0$. Therefore, $t = 0$ and hence there is a sub-sequence $\{\text{Tr}(V_{n_k})\}$ converging to 0. Since V_{n_k} 's are positive matrices, we conclude that $\{V_{n_k}\}$ converges to 0. \square

We now prove that the maximal part of φ defines a non-zero positive map.

Proposition 4.7. *Let φ be a positive map on a finite dimensional C^* -algebra \mathcal{A} with $r(\varphi) > 0$. Then $\widehat{\varphi}$ is a non-zero positive map on \mathcal{A} . If φ is CP, then so is $\widehat{\varphi}$.*

Proof. Recall from (4.3) that $\widehat{\varphi} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\varphi^n}{\|\varphi^n\|_\beta}$ where β is a Jordan basis of φ . To prove the well-definedness of $\widehat{\varphi}$, that is, the convergence of the above limit, we first consider the case $r(\varphi) = 1$.

Now, since \mathcal{A} is finite dimensional, it is enough to ensure pointwise convergence on Jordan basis vectors. Let $x \in \beta$ be any Jordan basis vector. Then x is a generalized eigenvector. Suppose it is of rank k corresponding to eigenvalue λ , that is, $(\varphi - \lambda)^k x = 0$ and $(\varphi - \lambda)^{k-1} x \neq 0$. Then, there is a Jordan chain $\{x_{k-1}, x_{k-2}, \dots, x_0\} \subseteq \beta$ where $x_j = (\varphi - \lambda)^j x$ for all $0 \leq j \leq (k-1)$. Observe that, for any $n \in \mathbb{N}$,

$$\varphi^n(x) = \varphi^n(x_0) = \sum_{j=0}^{k-1} \binom{n}{j} \lambda^{n-j} x_j \quad (4.6)$$

and there is $N_0 \in \mathbb{N}$ such that $\|\varphi^n\|_\beta = \binom{n}{d_\varphi-1}$ for all $n \geq N_0$. Note that for $|\lambda| = 1$, $d_{\varphi, \lambda} \leq d_\varphi$, by the definition of d_φ . Thus cardinality of $\{x_0, \dots, x_{k-1}\}$ corresponding to $|\lambda| = 1$ is at most d_φ . Hence, it suffices to look at the following three cases.

Case I. First assume that $|\lambda| < 1$. Then, for any $p \in \mathbb{N}$, we have

$$\lim_{n \rightarrow \infty} \frac{\binom{n}{p}}{\|\varphi^n\|_\beta} \lambda^n = \lim_{n \rightarrow \infty} \frac{\binom{n}{p}}{\binom{n}{d_\varphi-1}} \lambda^n = 0.$$

Therefore, using (4.6), we infer that

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\varphi^n(x_0)}{\|\varphi^n\|_\beta} = \sum_{j=0}^{k-1} \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{j}}{\|\varphi^n\|_\beta} \lambda^{n-j} x_j = 0.$$

Hence, when $|\lambda| < 1$, $\widehat{\varphi}(x) = 0$.

Case II. Let $|\lambda| = 1$ and $k < d_\varphi$. Therefore, by (4.6),

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\varphi^n(x_0)}{\|\varphi^n\|_\beta} = \sum_{j=0}^{k-1} \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{j}}{\|\varphi^n\|_\beta} \lambda^{n-j} x_j.$$

For $0 \leq j \leq k-1 < d_\varphi-1$, we observe that

$$\lim_{n \rightarrow \infty} \left| \frac{\binom{n}{j}}{\|\varphi^n\|_\beta} \lambda^{n-j} \right| = \lim_{n \rightarrow \infty} \frac{\binom{n}{j}}{\binom{n}{d_\varphi-1}} = 0.$$

Hence, we have

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{j}}{\binom{n}{d_\varphi-1}} \lambda^{n-j} x_j = 0,$$

that is, $\widehat{\varphi}(x) = 0$.

Case III. Now let $|\lambda| = 1$ and $k = d_\varphi$. Then for $0 \leq j < d_\varphi-1$, we similarly have

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{j}}{\|\varphi^n\|_\beta} \lambda^{n-j} x_j = 0.$$

Therefore,

$$\widehat{\varphi}(x) = \sum_{j=0}^{d_\varphi-1} \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{j}}{\|\varphi^n\|_\beta} \lambda^{n-j} x_j = \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{d_\varphi-1}}{\|\varphi^n\|_\beta} \lambda^{n-d_\varphi+1} x_{d_\varphi-1}. \quad (4.7)$$

Now, for $\lambda \neq 1$,

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m \frac{\binom{n}{d_\varphi-1}}{\|\varphi^n\|_\beta} \lambda^{n-d_\varphi+1} = \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=N_0}^m \frac{\binom{n}{d_\varphi-1}}{\|\varphi^n\|_\beta} \lambda^{n-d_\varphi+1} = \lambda^{1-d_\varphi} \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=N_0}^m \lambda^n = 0.$$

Thus, for $|\lambda| = 1$, $k = d_\varphi$ and $\lambda \neq 1$ we have, $\widehat{\varphi}(x) = 0$.

For $\lambda = 1$, by (4.7),

$$\widehat{\varphi}(x) = \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=N_0}^m 1^{n-d_\varphi+1} x_{d_\varphi-1} = x_{d_\varphi-1}.$$

Therefore, we have shown the convergence of the limit at all points of β . This proves the existence of $\widehat{\varphi}$ when $r(\varphi) = 1$.

For general positive map φ with $r = r(\varphi) > 0$, let $\psi = \varphi/r$ and β' be a Jordan basis of ψ . Then ψ is a non-zero positive map on \mathcal{A} of spectral radius 1. From the above discussion, $\widehat{\psi}$ exists. We claim that φ and ψ have same maximal degeneracy index. From the minimal polynomials of φ and ψ , we observe that $d_{\varphi,\lambda} = d_{\psi,\lambda/r}$ for all $\lambda \in \sigma(\varphi)$. Hence, we have

$$d_\varphi = \max_{\lambda \in \sigma(\varphi)} \{d_{\varphi,\lambda} : |\lambda| = r\} = \max_{\lambda \in \sigma(\varphi)} \{d_{\psi,\lambda/r} : |\lambda| = r\} = \max_{\mu \in \sigma(\psi)} \{d_{\psi,\mu} : |\mu| = 1\} = d_\psi.$$

For simplicity, we denote $d = d_\varphi = d_\psi$. Recall from (4.5) that there is $N_1 \in \mathbb{N}$ such that $\|\psi^n\|_{\beta'} = \binom{n}{d-1}$ and $\|\varphi^n\|_\beta = \binom{n}{d-1} r^{n-d+1}$ for all $n \geq N_1$. Now, it is evident that

$$\left(\frac{\widehat{\varphi}}{r}\right) = \widehat{\psi} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\psi^n}{\|\psi^n\|_{\beta'}} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=N_1}^N \frac{\varphi^n}{\binom{n}{d-1} r^n} = \frac{\widehat{\varphi}}{r^{d-1}}. \quad (4.8)$$

This completes the proof of existence of $\widehat{\varphi}$ for all positive map φ with $r(\varphi) > 0$. Being a limit of positive maps, $\widehat{\varphi}$ is positive. If $\widehat{\varphi} = 0$, then

$$\widehat{\varphi}(1) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\varphi^n(1)}{\|\varphi^n\|_\beta} = 0.$$

By Lemma 4.6, there exists a subsequence $\frac{\varphi^{n_k}(1)}{\|\varphi^{n_k}\|_\beta}$ converging to 0 in norm of \mathcal{A} . But since φ^{n_k} is a positive map and $\|\cdot\|_\beta$ is equivalent to $\|\cdot\|$, there is some $c > 0$ such that

$$\|\varphi^{n_k}(1)\| = \|\varphi^{n_k}\| \geq c \|\varphi^{n_k}\|_\beta.$$

This contradicts the fact that $\frac{\varphi^{n_k}(1)}{\|\varphi^{n_k}\|_\beta} \rightarrow 0$. Therefore, $\widehat{\varphi}$ is a well defined, non-zero, positive map as needed. From the definition of $\widehat{\varphi}$, it is clear that if φ is CP, so is $\widehat{\varphi}$. \square

Remark 4.8. From the proof above, it is clear that, if x is a generalized eigenvector of φ corresponding to the eigenvalue $r(\varphi)$, then

$$\widehat{\varphi}(x) = (\varphi - r(\varphi))^{d_\varphi-1} x. \quad (4.9)$$

On the other hand, if x is any other generalized eigenvector, then $\widehat{\varphi}(x) = 0$.

Theorem 4.9. *Let φ be a non-zero positive map on a finite dimensional C^* -algebra. Then the spectral radius $r(\varphi)$ belongs to the maximal spectrum of φ .*

Proof. If $r(\varphi) = 0$, there is nothing to prove. Observe from Remark 4.8 that $\widehat{\varphi}$ is non-zero only when $r(\varphi)$ is an element of the maximal spectrum of φ . Since $\widehat{\varphi}$ is non-zero by Proposition 4.7, $r(\varphi)$ belongs to the maximal spectrum of φ . \square

Now, we explore some useful properties of the maximal part of φ in the following theorem. It tells us as to how to explicitly obtain an eigenvector with spectral radius as the eigenvalue.

Theorem 4.10. *Let φ be a positive map on a finite dimensional C^* -algebra \mathcal{A} with $r(\varphi) > 0$. Then the maximal part $\widehat{\varphi}$ is a non-zero positive map satisfying the following properties:*

- (1) $\varphi\widehat{\varphi} = \widehat{\varphi}\varphi = r(\varphi)\widehat{\varphi}$.
- (2) $\widehat{\varphi}(1)$ is a positive eigenvector of φ corresponding to the eigenvalue $r(\varphi)$.
- (3) There are two possibilities with $\widehat{\varphi}$ which are:

$$\widehat{\varphi}^2 = \widehat{\varphi} \iff d_\varphi = 1$$

and

$$\widehat{\varphi}^2 = 0 \iff d_\varphi > 1.$$

- (4) If $\widehat{\varphi}^2 = \widehat{\varphi}$, then for any element x , $\varphi(x) = r(\varphi)x$ if and only if $\widehat{\varphi}(x) = x$.

Proof. For notational simplicity, we take $d = d_\varphi$.

- (1) Recall from the equation (4.8) that $\widehat{(\varphi/r)} = \widehat{\varphi}/r^{d-1}$. Hence, without loss of generality, we assume $r(\varphi) = 1$ and show that $\varphi\widehat{\varphi} = \widehat{\varphi}\varphi = \widehat{\varphi}$. From Theorem 4.9, we know that 1 belongs to the maximal spectrum of φ . Since β is a Jordan basis of φ , $[\varphi]_\beta$ is of the form

$$[\varphi]_\beta = [\oplus_k(I + N_d)] \oplus A$$

where N_d is the $d \times d$ Jordan nilpotent matrix of nilpotency d so that $(I + N_d)$ is the largest Jordan block of φ corresponding to eigenvalue 1, and k indicates its multiplicity in the Jordan form of φ , and A consists of the remaining Jordan blocks. Then, from Remark 4.8, we get that

$$[\widehat{\varphi}]_\beta = [\oplus_k N_d^{d-1}] \oplus 0.$$

Since $(I + N_d)N_d^{d-1} = N_d^{d-1}(I + N_d) = N_d^{d-1}$, we have $\varphi\widehat{\varphi} = \widehat{\varphi}\varphi = \widehat{\varphi}$.

- (2) Since $\widehat{\varphi}$ is a non-zero positive map, $\widehat{\varphi}(1)$ is non-zero, positive element of \mathcal{A} . From (1), we obtain $\varphi(\widehat{\varphi}(1)) = r(\varphi)\widehat{\varphi}(1)$.
- (3) We observe that

$$\widehat{\varphi}^2 = \widehat{\varphi} \iff N_d^{d-1}N_d^{d-1} = N_d^{d-1} \iff d = 1 \text{ and}$$

$$\widehat{\varphi}^2 = 0 \iff N_d^{d-1}N_d^{d-1} = 0 \iff d > 1.$$

- (4) If $\widehat{\varphi}(x) = x$, then

$$\varphi(x) = \varphi\widehat{\varphi}(x) = r\widehat{\varphi}(x) = rx$$

If $\varphi(x) = rx$, then $\varphi^n(x) = r^n x$ for all n . Since $\widehat{\varphi}^2 = \widehat{\varphi}$, it follows from (3) that $d = 1$. Thus, $\|\varphi^n\|_\beta = \binom{n}{d_\varphi-1} r^{n-d+1} = r^n$ for all but finitely many n 's. Therefore,

$$\widehat{\varphi}(x) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\varphi^n(x)}{\|\varphi^n\|_\beta} = x.$$

□

Theorem 4.9 and part (2) of Theorem 4.10 provide an alternate treatment to the Perron-Frobenius theorem.

Remark 4.11. If φ is an irreducible positive map on a finite dimensional C^* -algebra, then Evan and Høegh-Krohn ([11]) has proved the equality $r(\varphi) = \sup_{x \geq 0} \inf \{s > 0 : \varphi(x) \leq sx\}$, which in particular implies that $r(\varphi) > 0$. Therefore, we can define its maximal part. Theorem 2.4 asserts that the eigenspace corresponding to eigenvalue $r(\varphi)$ is of dimension 1 and is spanned by a strictly positive element $L \in \mathcal{A}$. Since by Theorem 4.10, $\widehat{\varphi}(1)$ is a positive eigenvector of φ corresponding to r , without loss of generality, we can assume that $L = \widehat{\varphi}(1)$. Also, by part (2) of Proposition 4.7, it is evident that $\widehat{\varphi}(X)$ is a Perron-Frobenius eigenvector of φ for all $X \in \mathcal{A}$. Therefore, there exists a state ψ on \mathcal{A} such that $\widehat{\varphi}(X) = \psi(X)L$ for all $X \in \mathcal{A}$, with $\varphi(L) = rL$.

5. ALGEBRA GENERATED BY CHOI-KRAUS COEFFICIENTS

Given a p -tuple (A_1, A_2, \dots, A_p) of $m \times m$ matrices how to compute a basis for the algebra generated by them in M_m is an important problem in Linear Algebra. Apparently, this has practical applications. In Pascoe [27], the author shows that this can be done by looking at the vectorization of a certain matrix. The p -tuple can be considered as the Choi-Kraus coefficients of the CP map τ on M_m defined by

$$\tau(X) = \sum_{i=1}^p A_i^* X A_i, \quad X \in M_m.$$

Recasting the work of Pascoe and going further, we observe that the elements of the algebra generated by these coefficients can be obtained as Choi-Kraus coefficients of certain other CP maps constructed out of τ . There are many ways we can do this. We also look at irreducibility of completely positive maps, in light of the study of the maximal part done in the previous section.

As observed in Proposition 2.3, the irreducibility property of a CP map on M_m is characterised in terms of the algebra generated by its Choi coefficients. For this purpose, we want to study the algebra generated by some family of matrices in terms of the coefficient space and Choi rank of some associated CP maps.

Theorem 5.1. *Let $\mathcal{S} = \{A_1, \dots, A_p\}$ be a p -tuple of $m \times m$ matrices for some $p \geq 1$ and let $\langle \mathcal{S} \rangle_0$ be the unital sub-algebra of M_m generated by A_i 's and let $\langle \mathcal{S} \rangle_1$ be the (not-necessarily unital) algebra generated by A_i 's. Let $\tau(X) = \sum_{i=1}^p A_i^* X A_i$ be the CP map associated with \mathcal{S} . Let $\{a_n : n \geq 0\}$ be a sequence of strictly positive scalars such that $\sum_{n=0}^{\infty} a_n \tau^n$ converges in norm. Take*

$$\sigma_0 = \sum_{n=0}^{\infty} a_n \tau^n, \quad \sigma_1 = \sum_{n=1}^{\infty} a_n \tau^n.$$

Then for $j = 0, 1$, the coefficient space of σ_j is equal to $\langle \mathcal{S} \rangle_j$. Consequently, the dimension of $\langle \mathcal{S} \rangle_j$ is equal to the Choi rank of σ_j and $A \in \langle \mathcal{S} \rangle_j$ iff there exists a scalar $q > 0$ such that the map $\sigma_j - q\alpha_A$ is CP.

Proof. Note that $\tau^0(X) = X$ and for $n \geq 1$,

$$\tau^n(X) = \sum_{i_1, \dots, i_n} (A_{i_1} A_{i_2} \cdots A_{i_n})^* X (A_{i_1} A_{i_2} \cdots A_{i_n}).$$

Therefore, we have

$$\sigma_0(X) = \sum_{n=0}^{\infty} a_n \sum_{i_1, \dots, i_n} (A_{i_1} A_{i_2} \cdots A_{i_n})^* X (A_{i_1} A_{i_2} \cdots A_{i_n}). \quad (5.1)$$

Let \mathcal{M}_0 be the coefficient space of σ_0 . Then from expression (5.1), it follows that $\mathcal{M}_0 = \text{span}\{I, A_{i_1}A_{i_2} \cdots A_{i_n} : i_1, \dots, i_n \in \{1, \dots, p\}, n \geq 1\}$ which is nothing but the algebra $\langle \mathcal{S} \rangle_0$. Therefore, we can conclude that the unital sub-algebra of M_m generated by A_i 's is same as the coefficient space of σ_0 . Similar argument works for σ_1 . The remaining statements are clear from 2.1. \square

This theorem provides an abundant supply of CP maps which can be used to determine the algebras generated by Choi-Kraus coefficients of CP maps. We single out two particularly useful examples. In a matrix form, the CP map γ_0 was considered by Pascoe [27].

Corollary 5.2. *Consider the set up of Theorem 5.1. Suppose $b > 0$ is a scalar such that $br(\tau) < 1$ and let $d > 0$. Then*

$$\gamma_0 := (1 - b\tau)^{-1}, \quad \eta_0 := \exp(d\tau)$$

are CP maps with their coefficient spaces equal to the unital algebra $\langle \mathcal{S} \rangle_0$ and

$$\gamma_1 = \gamma_0 - 1, \quad \eta_1 = \eta_0 - 1$$

are CP maps with their coefficient spaces equal to the algebra $\langle \mathcal{S} \rangle_1$.

Proof. This is clear from Theorem 5.1, by considering power series expansions:

$$\gamma_0 = \sum_{n=0}^{\infty} b^n \tau^n, \quad \eta_0 = \sum_{n=0}^{\infty} \frac{d^n}{n!} \tau^n.$$

Note that the power series for γ_0 converges as $r(b\tau) < 1$. \square

Remark 5.3. Theorem 5.1 and Corollary 5.2 are stated for CP maps in finite dimensions. However, it is clear that they can be extended to normal completely positive maps on $\mathcal{B}(\mathcal{H})$, with separable infinite dimensional Hilbert space \mathcal{H} , where the power series under consideration are assumed to be convergent in strong operator topology.

Remark 5.4. If one is interested in analyzing the unital or non-unital $*$ -algebra generated by a tuple A_1, \dots, A_p of matrices, one can consider the CP map $X \mapsto \sum_{i=1}^p (A_i^* X A_i + A_i X A_i^*)$ and the methods of Theorem 5.1 are applicable.

Here in Theorem 5.1, we use power series with infinitely many terms, and examples in Corollary 5.2 could be useful due to computational advantages. However, due to dimension considerations, it is always enough to consider finitely many terms. This can be seen as follows.

Once again, consider the set up of Theorem 5.1. For $k \geq 0$, let \mathcal{M}_k be the coefficient space of $\sum_{n=0}^k a_n \tau^n$. Then, it is clear that $\mathcal{M}_0 \subseteq \mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \cdots$ and all of them are subspaces of M_m . We claim that this increasing sequence stabilizes after a few steps. Suppose $\mathcal{M}_k = \mathcal{M}_{k+1}$ for some k . Then since any $(k+1)$ -length product $A_{j_1} A_{j_2} \cdots A_{j_{k+1}}$ is in the coefficient space \mathcal{M}_{k+1} , we get

$$A_{j_1} A_{j_2} \cdots A_{j_{k+1}} = \text{span}\{A_{i_1} A_{i_2} \cdots A_{i_r} : 0 \leq r \leq k, 1 \leq i_1, \dots, i_r \leq p\}.$$

It follows that then any $(k+2)$ -length product $A_{j_1} A_{j_2} \cdots A_{j_{k+2}}$ is a linear combination of $(k+1)$ -length products of A_j 's. In other words, $\mathcal{M}_{k+2} = \mathcal{M}_{k+1}$. By induction, $\mathcal{M}_k = \mathcal{M}_l$ for all $k \leq l$. If $\mathcal{M}_j \subsetneq \mathcal{M}_{j+1}$, then their dimensions have to differ by at least 1 and all of them are subspaces of M_m , which has dimension m^2 , implies that the stabilization has to take place by index m^2 . Consequently, we have proved the following theorem.

Theorem 5.5. *Consider the set up of Theorem 5.1. Then for $j = 0, 1$ there exists natural numbers $N_j \leq m^2$ such that the coefficient space of $\sum_{n=j}^{N_j} a_n \tau^n$ is equal to $\langle \mathcal{S} \rangle_j$ and the Choi rank of $\sum_{n=j}^{N_j} a_n \tau^n$ is equal to the dimension of $\langle \mathcal{S} \rangle_j$.*

To check whether a CP map τ on M_m is irreducible, one can construct a convenient CP map σ_1 as described in Theorem 5.1. The combination of Theorem 5.1 and Proposition 2.3 ensures that τ is irreducible if and only if σ_1 has a Choi rank of m^2 .

Now, we proceed with our desired characterisation for irreducible CP maps on general finite dimensional C^* -algebras and a description of their maximal parts. Compared to maps on full matrix algebra (see Proposition 2.3), the situation is a bit more complicated now. Consider a CP map τ with a specified Choi-Kraus type decomposition. Whenever these coefficients do not admit a proper common invariant subspace, τ is irreducible. However, as the following example illustrates, this condition is not always necessary.

Example 5.6. Let $\mathcal{A} = M_1 \oplus M_1$ and let τ be the CP map defined by $\tau(X) = 2 \text{trace}(X) I$ on $\mathcal{A} = M_1 \oplus M_1$. We can express

$$\tau(X) = A^* X A + B^* X B \text{ for all } X \in \mathcal{A},$$

where $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$. Observe that τ is strictly positive, and therefore, it is irreducible. However, it exhibits a Choi-Kraus type expression involving A and B which share span $\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ as a common invariant subspace.

Consider a finite dimensional C^* -algebra \mathcal{A} . With out loss of generality, $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ for some natural numbers n_1, n_2, \dots, n_d . Clearly, \mathcal{A} has a naturally embedding in M_m where $m = n_1 + \cdots + n_d$. Throughout we will be considering this embedding of \mathcal{A} . Now, we can canonically extend any CP map on \mathcal{A} to a CP map on M_m as follows.

Definition 5.7. For a CP map τ on a finite dimensional C^* -algebra, $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$, where $n_1 + \cdots + n_d = m$, if $\iota : \mathcal{A} \hookrightarrow M_m$ is the canonical embedding of \mathcal{A} into M_m , for each $k = 1, \dots, d$, P_k is the orthogonal projection of M_m to M_k and $\phi : M_m \rightarrow \mathcal{A}$ is defined by $\phi(X) = \sum_{k=1}^d P_k^* X P_k$ then the *canonical extension* of τ to M_m denoted by $\tilde{\tau}$ is defined by $\tilde{\tau} = \iota \circ \tau \circ \phi$.

It is natural to expect a connection between the irreducibility of τ and $\tilde{\tau}$. To establish this, we need the following lemma. Recall that a positive map is called strictly positive if it sends all non-zero positive elements to strictly positive elements. The lemma tells us that for a positive map having a block matrix decomposition, the diagonals can be perturbed suitably to get a strictly positive map.

Lemma 5.8. *Let $\{\psi_{ij} : M_{n_j} \rightarrow M_{n_i} \mid i, j \in \{1, \dots, d\}\}$ be a collection of positive maps with $1 + \psi_{ii} > 0$ for all i and $\psi_{ij} > 0$ for all $i \neq j$ where $n_1, \dots, n_d \in \mathbb{N}$. Let $X = (X_{ij}) \in M_m$ be a non-zero positive (block) matrix where $X_{ij} \in M_{n_i \times n_j}$ and $m = n_1 + \dots + n_d$. Then the matrix $Y = (Y_{ij}) \in M_m$ defined as*

$$Y_{ij} = \begin{cases} X_{ii} + \sum_{k=1}^d \psi_{ik}(X_{kk}), & \text{if } i = j \\ X_{ij}, & \text{if } i \neq j \end{cases}$$

is strictly positive.

Proof. Observe that, we can write $Y = X + Z$ where Z is a block diagonal positive matrix with diagonal entries $Z_{ii} = \sum_{k=1}^d \psi_{ik}(X_{kk})$. If all Z_{ii} 's are strictly positive, then Z , hence

Y , becomes strictly positive matrix. If some Z_{ll} is not strictly positive, then $X_{kk} = 0$ for all $k \neq l$. Then X turns into a block diagonal matrix with only non-zero entry X_{ll} . Then we have

$$Y_{ij} = \begin{cases} \psi_{il}(X_{ll}), & \text{if } i = j \neq l \\ (1 + \psi_{ll})(X_{ll}), & \text{if } i = j = l. \\ 0, & \text{if } i \neq j \end{cases}$$

Thus, Y becomes a block diagonal matrix with strictly positive diagonal entries. Therefore, Y is a strictly positive matrix in either cases. \square

Proposition 5.9. *Let τ be a CP map on finite dimensional C^* -algebra $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ and let $\tilde{\tau}$ be its canonical extension to M_m where $m = n_1 + \cdots + n_d$. Then τ is irreducible if and only if $\tilde{\tau}$ is irreducible. The spectral radii of τ and $\tilde{\tau}$ are equal.*

Proof. First assume that, $\tilde{\tau}$ is irreducible. Then by Lemma 2.1 of [11], $(1 + \tilde{\tau})^{m-1}$ is strictly positive. For any non-zero positive $X \in \mathcal{A}$, we infer that $(1 + \tau)^{m-1}(X) = (1 + \tau)^{m-1}(\phi(X)) = (1 + \tilde{\tau})^{m-1}(\phi(X)) > 0$, proving irreducibility of τ .

Conversely, assume that τ is irreducible. Since τ is a positive map, for any $A = A_1 \oplus \cdots \oplus A_d \in \mathcal{A}$, we have $(1 + \tau)^{m-1}(A) - A \geq 0$ and its i^{th} diagonal block entry then must look like sum of the positive maps corresponding to each $j = 1, \dots, d$ that take $A_j \in M_{n_j}$ to some matrix in M_{n_i} . That is, there is a collection of positive maps $\{\psi_{ij} : M_{n_j} \rightarrow M_{n_i} \mid i, j \in \{1, \dots, d\}\}$ such that

$$((1 + \tau)^{m-1}(A))_{ii} = A_i + \sum_{j=1}^d \psi_{ij}(A_j) \quad \text{for all } X \in \mathcal{A}.$$

We claim that this collection of ψ_{ij} 's satisfy the hypothesis of Lemma 5.8. Fix some $i \in \{1, \dots, d\}$ arbitrarily and let $B \in M_{n_i}$ be any non-zero positive matrix. Define $A \in \mathcal{A}$ to be the block diagonal matrix whose i^{th} diagonal block is B and all other are zero. It follows from the irreducibility of τ that $(1 + \tau)^{m-1}$ is strictly positive. Thus, we have $((1 + \tau)^{m-1}(A))_{kk} > 0$ for each k . Hence, in particular

$$((1 + \tau)^{m-1}(A))_{ii} = (1 + \psi_{ii})(B) > 0$$

and for $j \neq i$,

$$((1 + \tau)^{m-1}(A))_{jj} = \psi_{ji}(B) > 0.$$

As B is arbitrary non-zero positive matrix in M_{n_i} , we get $1 + \psi_{ii} > 0$ and $\psi_{ij} > 0$ for all $i \neq j$. Since i, j are arbitrary, this proves the required claim. Now, let $X \in M_m$ be non-zero positive. A simple computation shows that,

$$(1 + \tilde{\tau})^{m-1}(X) = (1 + \iota \circ \tau \circ \phi)^{m-1}(X) = (X - \phi(X)) + (1 + \tau)^{m-1}(\phi(X)).$$

Then,

$$((1 + \tilde{\tau})^{m-1}(X))_{ij} = \begin{cases} X_{ii} + \sum_{k=1}^d \psi_{ik}(X_{kk}), & \text{if } i = j \\ X_{ij}, & \text{if } i \neq j \end{cases}.$$

Hence, by Lemma 5.8, we infer that $(1 + \tilde{\tau})^{m-1}(X) > 0$ for all non-zero positive $X \in M_m$. Therefore, $\tilde{\tau}$ is irreducible. To see the second part, observe that $\tilde{\tau}^n = \iota \circ \tau^n \circ \phi$ for all $n \in \mathbb{N}$ and

$$r(\tilde{\tau}) = \lim_{n \rightarrow \infty} \|\tilde{\tau}^n(1)\|^{1/n} = \lim_{n \rightarrow \infty} \|\tau^n(1)\|^{1/n} = r(\tau).$$

\square

The following theorem says that if we are looking at a Choi-Kraus decomposition $\tau(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in \mathcal{A}$, which satisfies also $\tilde{\tau}(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in M_m$, then the irreducibility of τ can be read of from A_i 's. Note that this was not the case in Example 5.6.

Theorem 5.10. *Let τ be a CP map on finite dimensional C^* -algebra $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ and $\tilde{\tau}$ be its canonical extension to M_m where $m = n_1 + \cdots + n_d$. Then the following statements are equivalent:*

- (1) τ is irreducible;
- (2) $\tilde{\tau}$ is irreducible;
- (3) *There exists matrices $A_1, \dots, A_p \in M_m$ which generates M_m as algebra such that $\tilde{\tau}(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in M_m$. In particular, $\tau(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in \mathcal{A}$;*
- (4) *There exist matrices $A_1, \dots, A_p \in M_m$ which generates M_m as algebra such that $\tau(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in \mathcal{A}$.*

Proof. We have already established the equivalence between (1) and (2) in Proposition 5.9, and the implication of (3) leading to (4) is self-evident. Now, we prove the remaining implications.

(2) \Rightarrow (3): Suppose $\tilde{\tau}$ has Choi-Kraus expression $\tilde{\tau}(X) = \sum_{i=1}^p A_i^* X A_i$ for all $X \in M_m$ where $\{A_1, \dots, A_p\} \subseteq M_m$. Since τ is irreducible, by Proposition 2.3, the algebra generated by A_i 's is whole of M_m . In particular, for $X = \bigoplus_{j=1}^d X_j \in \mathcal{A}$, we have

$$\tau(X) = \iota \circ \tau \circ \phi(X) = \tilde{\tau}(X) = \sum_{i=1}^p A_i^* X A_i.$$

(4) \Rightarrow (1): Let us assume on contrary that τ is reducible. Then there is a non-trivial projection $P \in \mathcal{A}$ such that $\tau(P) \leq \lambda P$ for some $\lambda > 0$. We show that $\text{Ker}(P)$ is invariant under all A_i 's. For $h \in \text{Ker}(P)$, we have

$$\|PA_i h\|^2 = \langle A_i^* P A_i h, h \rangle \leq \lambda \langle P h, h \rangle = 0, \text{ for all } i.$$

Therefore, $A_i(\text{Ker}(P)) \subseteq \text{Ker}(P)$ for all i , hence, A_i 's possess a non-trivial common invariant subspace $\text{Ker}(P)$. This contradicts to the fact that A_1, \dots, A_p generate the algebra M_m . \square

Let us now give two examples of CP maps on finite dimensional C^* -algebras whose irreducibility can be verified using the analysis done so far. For instance, one may use Corollary 5.2 for $\tilde{\tau}$.

Example 5.11. Let $\mathcal{A} = M_2 \oplus M_1$ and $\tau : \mathcal{A} \rightarrow \mathcal{A}$ be defined as

$$\tau \left(\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & e \end{bmatrix} \right) = \begin{bmatrix} a+e & 0 & 0 \\ 0 & a+e & 0 \\ 0 & 0 & d \end{bmatrix}.$$

Then τ is irreducible.

Example 5.12. Let $\mathcal{A} = M_1 \oplus M_1 \oplus M_1$ and $\tau : \mathcal{A} \rightarrow \mathcal{A}$ be defined as

$$\tau \left(\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \right) = \begin{bmatrix} b & 0 & 0 \\ 0 & a+c & 0 \\ 0 & 0 & b \end{bmatrix}.$$

Then τ is irreducible.

In Section 4, we have defined and studied the maximal part $\widehat{\varphi}$ for any positive map φ with $r(\varphi) > 0$, on finite dimensional C^* -algebras. For CP maps, there is more to explore. We observe that one can obtain a Choi-Kraus decomposition for a CP map τ in such a way that the Choi-Kraus coefficients of $\widehat{\tau}$ span a non-zero ideal in the algebra generated by the Choi-Kraus coefficients of τ . To proceed, we require the following technical lemma.

Lemma 5.13. *Let $\{\tau_n\}$ be a sequence of CP maps on M_m converging to a CP map τ . Suppose that there is a subspace \mathcal{V} of M_m such that $\mathcal{M}_{\tau_n} \subseteq \mathcal{M}_{\tau_{n+1}} \subseteq \mathcal{V}$ for all n . Then, $\mathcal{M}_{\tau} \subseteq \mathcal{V}$.*

Proof. Let C_n and C be the conjugated Choi matrices of the CP maps τ_n and τ respectively. Since $\lim \tau_n = \tau$, we have $\lim C_n = C$. Now, let P_n and P be the $m^2 \times m^2$ projection matrices with $\text{range}(P_n) = \text{range}(C_n)$ and $\text{range}(P) = \text{range}(C)$. As $\mathcal{M}_{\tau_n} \subseteq \mathcal{M}_{\tau_{n+1}} \subseteq \mathcal{V}$, by Proposition 2.2, we have $\text{range}(P_n) \subseteq \text{range}(P_{n+1}) \subseteq \text{vec}(\mathcal{V})$. Let Q be the projection onto the subspace $\text{vec}(\mathcal{V}) \subseteq \mathbb{C}^{m^2}$. Then, $P_n \leq P_{n+1} \leq Q$ for all n . For any $h \in \mathbb{C}^m$, there is $h' \in \mathbb{C}^m$ such that $Ph = Ch'$. We infer that

$$\|P_n Ph - Ph\| = \|P_n Ch' - Ch'\| \leq \|P_n(Ch' - C_n h')\| + \|C_n h' - Ch'\| \leq 2\|C_n h' - Ch'\|.$$

Taking limit as n tends to infinity, this implies that $\lim_n P_n P = P$. Also, since $\{P_n\}$ is an increasing sequence of projections with $P_n \leq Q$, there exists a projection $R \in M_{m^2}$ such that P_n converges to R and $R \leq Q$. Then, $RP = \lim P_n P = P$. Consequently, $\text{range}(P) \subseteq \text{range}(R)$, and thus, $\text{range}(P) \subseteq \text{range}(R) \subseteq \text{range}(Q) = \text{vec}(\mathcal{V})$. Therefore, we have $\mathcal{M}_{\tau} = \text{vec}^{-1}(\text{range}(P)) \subseteq \mathcal{V}$. \square

Remark 5.14. In the above lemma, the family of coefficient spaces \mathcal{M}_{τ_n} may not increase to \mathcal{M}_{τ} , as demonstrated by the following simple example. Let τ_n be the CP map on M_2 defined as

$$\tau_n(X) = \text{trace}(X) \begin{bmatrix} 1 & 1 - \frac{1}{n} \\ 1 - \frac{1}{n} & 1 \end{bmatrix}, \quad \forall X \in M_2.$$

It converges to the CP map $\tau(X) = \text{trace}(X) \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, $X \in M_2$. Observing that the rank of each Choi matrix of τ_n is 4, we have $\mathcal{M}_{\tau_n} = M_2$ for all n . However, the rank of the Choi matrix of τ is 2, indicating that $\mathcal{M}_{\tau} \subsetneq M_2$.

Theorem 5.15. *Let τ be a CP map on a finite dimensional C^* -algebra $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ with $r(\tau) > 0$. Consider canonical extensions of $\tau, \widehat{\tau}$. Then*

$$\widetilde{\widehat{\tau}} = \widehat{\widetilde{\tau}}. \quad (5.2)$$

For any two Choi-Kraus decompositions:

$$\widetilde{\tau}(X) = \sum_{i=1}^p A_i^* X A_i, \quad \widetilde{\widehat{\tau}}(X) = \sum_{j=1}^q B_j^* X B_j, \quad X \in M_m$$

of $\widetilde{\tau}, \widetilde{\widehat{\tau}}$ respectively, the vector space $\text{span}\{B_1, \dots, B_q\}$ is a subalgebra of M_m and it is a non-zero ideal in the algebra generated by A_1, \dots, A_p .

Proof. Observe that $\tau : \mathcal{A} \rightarrow \mathcal{A}$, $\widetilde{\tau} : M_m \rightarrow M_m$ and \mathcal{A} is a subspace of M_m . With respect to the Hilbert-Schmidt inner product, we decompose M_m as $M_m = \mathcal{A} \oplus \mathcal{A}^\perp$ and with respect to this decomposition, $\widetilde{\tau} = \tau \oplus 0$. Now, it is clear that given a Jordan basis β for τ (basis elements span \mathcal{A}), we can extend it to a Jordan basis β' for $\widetilde{\tau}$ (basis elements span M_m) such that elements of $\beta' \setminus \beta$ form a basis of \mathcal{A}^\perp . In particular, $\widehat{\widetilde{\tau}}(X) = 0$ for all

$X \in \beta' \setminus \beta$. Then it is easy to see that $\|\tilde{\tau}^n\|_{\beta'} = \|\tau^n\|_{\beta}$ for all $n \in \mathbb{N}$. From Proposition 5.9, $r(\tilde{\tau}) = r(\tau) > 0$ and hence

$$\widehat{\tilde{\tau}} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\tilde{\tau}^n}{\|\tilde{\tau}^n\|_{\beta'}} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\iota \circ \tau^n \circ \phi}{\|\tau^n\|_{\beta}} = \iota \circ \widehat{\tau} \circ \phi = \widehat{\tilde{\tau}}.$$

This proves the first part.

For the second part, let $\tau_n = \frac{1}{N} \sum_{k=1}^N \frac{\tilde{\tau}^k}{\|\tilde{\tau}^k\|_{\beta'}}$ and \mathcal{V} be the algebra generated by A_i 's. Then, $\lim \tau_n = \widehat{\tilde{\tau}}$. Observe that $\mathcal{M}_{\widehat{\tilde{\tau}}} = \text{span}\{B_j : 1 \leq j \leq q\}$ and $\mathcal{M}_{\tau_n} \subseteq \mathcal{M}_{\tau_{n+1}} \subseteq \mathcal{V}$ for all n . Therefore, by Lemma 5.13, it is evident that $\text{span}\{B_j : 1 \leq j \leq q\}$ is a subspace of the algebra generated by A_i 's. From Theorem 4.10, we know that either $\widehat{\tilde{\tau}}^2 = \widehat{\tilde{\tau}}$ or $\widehat{\tilde{\tau}}^2 = 0$. In the first case, the expression

$$\sum_{j,k} B_k^* B_j^* X B_j B_k = \sum_j B_j^* X B_j, \quad X \in M_m$$

represents two different Choi-Kraus decompositions of $\widehat{\tilde{\tau}}$, implying $B_j B_k \in \text{span}\{B_j : 1 \leq j \leq q\}$. In the second case,

$$\sum_{j,k} B_k^* B_j^* X B_j B_k = 0, \quad X \in M_m$$

and hence $B_j B_k = 0$ for all j, k . Therefore, in either case $\text{span}\{B_j : 1 \leq j \leq q\}$ is an algebra. Similarly, from Theorem 4.10, we have $\tilde{\tau} \widehat{\tilde{\tau}} = \widehat{\tilde{\tau}} \tilde{\tau} = r(\tau) \widehat{\tilde{\tau}}$. Thus,

$$\sum_{i,j} (A_i B_j)^* X (A_i B_j) = \sum_{j,i} (B_j A_i)^* X (B_j A_i) = r(\tau) \sum_{j=1}^q B_j^* X B_j \quad \text{for all } X \in M_m.$$

Hence, by Proposition 2.1 each of $A_i B_j$ and $B_j A_i$ belong to the span of B_j 's, which implies $\text{span}\{B_j : 1 \leq j \leq q\}$ forms an ideal of the algebra generated by A_i 's. It is non-zero because $\widehat{\tilde{\tau}}$ is a non-zero map. \square

Now, we look at irreducible CP maps on finite dimensional C^* -algebras. We need a technical lemma.

Lemma 5.16. *Let W_1, W_2 be two linearly independent, positive operators on a Hilbert space \mathcal{H} . Then there exists a non-zero $W_3 \in \text{span}\{W_1, W_2\}$ which is positive and singular.*

Proof. If any one of W_1 and W_2 is not invertible, we can take W_3 to be that. So, we can assume both W_1 and W_2 are invertible. Let d be the spectral radius of the positive operator $V = W_1^{-1/2} W_2 W_1^{-1/2}$ and $W_3 = W_1 - d^{-1} W_2$ which is non-zero due to the linear independence of W_1, W_2 . Since $\sigma(V) = \sigma(W_1^{-1/2} W_2 W_1^{-1/2}) = \sigma(W_1^{-1} W_2)$, we have $d \in \sigma(W_1^{-1} W_2)$. This implies $W_1^{-1} W_2 - dI$ is singular which is equivalent to the singularity of $W_3 = d^{-1} W_1 (dI - W_1^{-1} W_2)$. We need to verify the positivity of W_3 . For $x \in \mathcal{H}$, take $y = W_1^{1/2} x$. Then

$$\langle x, (W_1 - d^{-1} W_2)x \rangle = \langle y, y \rangle - d^{-1} \langle y, W_1^{-1/2} W_2 W_1^{-1/2} y \rangle = d^{-1} (d \langle y, y \rangle - \langle y, V y \rangle)$$

Since $V \leq dI$, we infer that $\langle x, W_3 x \rangle \geq 0$ for all $x \in \mathcal{H}$, proving positivity of W_3 . \square

We have observed in Remark 4.11 that if a positive map on a finite dimensional C^* -algebra is irreducible, then its spectral radius is strictly positive. So, we can talk about its maximal part. Here, we connect the irreducibility of a CP map and with that of its maximal part.

Theorem 5.17. *Let τ be a CP map on a finite dimensional C^* -algebra $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ and $m = n_1 + \cdots + n_d$. Then τ is irreducible if and only if $\hat{\tau}$ is irreducible. In such a case, the following properties hold:*

- (1) *The maximal degeneracy index of τ is 1.*
- (2) *The map $\hat{\tau}$ is a strictly positive map of rank 1.*
- (3) *The CP map $\hat{\tau}$ has Choi rank m^2 . In particular, there exists $m \times m$ matrices $\{B_1, \dots, B_q\}$ which span M_m and*

$$\hat{\tau}(X) = \sum_{j=1}^q B_j^* X B_j, \quad \forall X \in \mathcal{A}.$$

- (4) *There is a faithful state ψ on \mathcal{A} and a strictly positive $L \in \mathcal{A}$ such that*

$$\hat{\tau}(X) = \psi(X) L \quad \text{for all } X \in \mathcal{A}$$

and (a) $\tau(L) = r L$, (b) $\psi(L) = 1$, and (c) $\psi \circ \tau = r(\tau) \psi$.

- (5) *There are no positive generalized eigenvectors of τ corresponding to eigenvalues of modulus less than $r(\tau)$. In other words, the set $\{X \geq 0 : \lim \tau^n(X) = 0\}$ only contains 0.*

Proof. The irreducibility of $\hat{\tau}$ ensures the irreducibility of τ since any hereditary subspace invariant under τ remains invariant under τ^n , and therefore under $\hat{\tau}$ as well. Assume that τ is irreducible. Suppose that the CP maps $\tilde{\tau}$ and $\hat{\tau}$ have the Choi-Kraus decompositions

$$\tilde{\tau}(X) = \sum_{i=1}^N A_i^* X A_i, \quad \hat{\tau}(X) = \sum_{j=1}^q B_j^* X B_j \quad \text{for all } X \in M_m.$$

According to Theorem 5.15, B_j 's span a non-zero ideal in the algebra generated by the A_i 's. Since τ is irreducible, it follows from Theorem 5.10 that the Choi-Kraus coefficients A_i 's of the canonical extension $\tilde{\tau}$ generate the full matrix algebra M_m . Since M_m is simple, we have $\text{span}\{B_j : 1 \leq j \leq q\} = M_m$. Therefore, $\mathcal{M}_{\tilde{\tau}} = M_m$, in other words, $\hat{\tau}$ has Choi rank m^2 . This proves part (3). By Proposition 2.1, we infer that α_I is dominated by $s \hat{\tau}$ for some $s > 0$ where $\alpha_I(X) = X$ for all $X \in M_m$. This implies in particular that for all positive $X \in \mathcal{A}$, $\hat{\tau}(X) \geq s^{-1} X$, thus confirming the faithfulness of $\hat{\tau}$.

Let $X \in \mathcal{A}$ be non-zero positive. Then, by Theorem 4.10, $\tau(\hat{\tau}(X)) = r(\tau)\hat{\tau}(X)$. For any $h \in \text{Ker}(\hat{\tau}(X))$, we infer that

$$\|\hat{\tau}(X)^{1/2} A_i h\|^2 \leq \sum_k \langle \hat{\tau}(X) A_k h, A_k h \rangle = \langle \tau(\hat{\tau}(X)) h, h \rangle = r(\tau) \langle \hat{\tau}(X) h, h \rangle = 0 \quad \text{for all } i.$$

Therefore, $\hat{\tau}(X) A_i h = 0$ for all $h \in \text{Ker}(\hat{\tau}(X))$, consequently implying $A_i(\text{Ker}(\hat{\tau}(X))) \subseteq \text{Ker}(\hat{\tau}(X))$ for all p . Since A_i 's generate the full matrix algebra M_m and $\hat{\tau}(X)$ is non-zero, $\text{Ker}(\hat{\tau}(X)) = 0$, proving strict positivity of $\hat{\tau}(X)$. Hence, $\hat{\tau}$ is strictly positive and, in particular, irreducible.

Now, as per Theorem 4.10, either $\hat{\tau}^2 = 0$ or $\hat{\tau}^2 = \hat{\tau}$. Since $\hat{\tau}$ is strictly positive, we must have $\hat{\tau}^2 = \hat{\tau}$, hence from Theorem 4.10, $d_\tau = 1$. This proves part (1). Let us assume on contrary that $\hat{\tau}$ has rank greater than 1. Then, since \mathcal{A} is span of its positive elements, there are positive elements V_1, V_2 in \mathcal{A} such that $\hat{\tau}(V_1), \hat{\tau}(V_2)$ are linearly independent. Let $W_i = \hat{\tau}(V_i)$ for $i = 1, 2$. Then W_1 and W_2 are positive, linearly independent with $\hat{\tau}(W_i) = W_i$ for $i = 1, 2$. By Lemma 5.16, there exists a non-zero positive $W_3 \in \text{span}\{W_1, W_2\}$ which is singular. But $W_3 = \hat{\tau}(W_3)$ has to be strictly positive, leading to a contradiction. Therefore, $\hat{\tau}$ has rank 1, validating part (2).

Furthermore, the range of $\widehat{\tau}$ is spanned by the strictly positive element $L = \widehat{\tau}(1) = \sum B_j B_j^*$. Therefore,

$$\forall X \in \mathcal{A}, \exists \psi(X) \in \mathbb{C} \text{ such that } \widehat{\tau}(X) = \psi(X) L. \quad (5.3)$$

Since $\widehat{\tau}$ is linear and positive, ψ is a positive linear functional. As $\widehat{\tau}$ is faithful, so is ψ . Furthermore, as $L = \widehat{\tau}(1) = \psi(1)L$, we have $\psi(1) = 1$. Therefore, ψ is unital and hence a state. Again, recall from Theorem 4.10 that $\widehat{\tau}\tau = r(\tau)\widehat{\tau}$. We infer that $\tau(L) = \tau\widehat{\tau}(1) = r\widehat{\tau}(1) = rL$. Since $\widehat{\tau}^2 = \widehat{\tau}$, using (5.3), we have

$$\widehat{\tau}^2(X) = \widehat{\tau}(\widehat{\tau}(X)) = \widehat{\tau}(\psi(X)L) = \psi(X)\widehat{\tau}(L) = \psi(X)\psi(L)L = \widehat{\tau}(X)\psi(L).$$

Thus, we must have $\psi(L) = 1$. Finally, it is evident that for all $X \in \mathcal{A}$,

$$\widehat{\tau}\tau(X) = r(\tau)\widehat{\tau}(X) \implies \psi(\tau(X))L = r(\tau)\psi(X)L \implies \psi(\tau(X)) = r(\tau)\psi(X).$$

Altogether, it establishes part (4).

Recall from remark 4.8 that if $X \in \mathcal{A}$ is a generalized eigenvector of τ corresponding to an eigenvalue of modulus less $r(\tau)$, then $\widehat{\tau}(X) = 0$. Since $\widehat{\tau}$ is faithful here, there is no such element which is positive. This proves part (5). \square

We also note that a faithful state as in this theorem means that there exists a strictly positive density matrix R such that $\psi(X) = \text{trace}(RX)$ for all $X \in \mathcal{A}$.

Corollary 5.18. *Let τ be a CP map on a finite dimensional C^* -algebra \mathcal{A} with $r(\tau) > 0$. If τ is irreducible, then there is a CP map σ on \mathcal{A} which is elementarily similar to τ such that $\|\sigma\| = r(\tau)$.*

Proof. From Theorem 5.17, we have strictly positive $L \in \mathcal{A}$ such that $\tau(L) = rL$. Due to Theorem 3.5, there exists a CP map σ on \mathcal{A} which is elementarily similar to τ such that $\|\sigma\| = r(\tau)$. \square

The converse of Corollary 5.18 does not hold true in general, as shown by the following counter example.

Example 5.19. Let $\mathcal{A} = M_{n_1} \oplus \cdots \oplus M_{n_d}$ be a finite dimensional C^* -algebra and choose $A \in \mathcal{A}$ such that $\left\| \frac{A^n}{r^n} \right\|$ forms a bounded sequence where $r = r(A)$. Let us consider the CP map $\tau = \alpha_A$ on \mathcal{A} . If \mathcal{A} is a proper sub-algebra of M_m ($m = n_1 + \cdots + n_d$), A will not generate the full matrix algebra M_m . First, we will prove that there is σ which is elementarily similar to τ with $\|\sigma\| = r(\tau)$. Then, we will show that it does not satisfy the hypothesis of Corollary 5.18.

Due to Theorem 4.2, there exists positive, invertible $V \in \mathcal{A}$ such that $\|V^{-1}AV\| = r$. This implies

$$r^2 I - (V^{-1}AV)^*(V^{-1}AV) \geq 0 \implies V^{-1}A^*V^2AV^{-1} \leq r^2 I \implies A^*V^2A \leq r^2 V^2 \quad (5.4)$$

Now let $L = V^2$ which is a positive and invertible element of \mathcal{A} . Recall that for elementary CP map $\tau = \alpha_A$, we have $r(\tau) = r(A)^2 = r^2$. Therefore, (5.4) shows that $\tau(L) \leq r(\tau)L$. Using Theorem 3.5, we get that there exists a CP map σ which is elementarily similar to τ such that $\|\sigma\| = r(\tau)$.

Now, since $A \in M_{n_1} \oplus \cdots \oplus M_{n_d}$, the canonical extension has the same Choi-Kraus decomposition $\widetilde{\tau}(X) = A^*XA$, $X \in M_m$. As A does not generate the algebra M_m , it follows from Proposition 2.3 that $\widetilde{\tau}$ is reducible; hence, so is τ by Proposition 5.9.

For the irreducible CP maps in Examples 5.11 and 5.12, we now proceed to compute $\widehat{\tau}$, ψ , and the strictly positive eigenvector as outlined in Theorem 5.17. Additionally, we determine the elementarily similar CP map σ as described in Corollary 5.18.

Example 5.20. Let us recall the irreducible CP map τ on $\mathcal{A} = M_2 \oplus M_1$ from Example 5.11, which is defined as follows:

$$\tau \left(\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & e \end{bmatrix} \right) = \begin{bmatrix} a+e & 0 & 0 \\ 0 & a+e & 0 \\ 0 & 0 & d \end{bmatrix}.$$

Since τ is irreducible, it is noted in Theorem 5.17 that $d_\tau = 1$, implying $\|\tau^n\|_\beta = r^n$ for all but finitely many n where r is the spectral radius of τ . Thus, $\hat{\tau} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\tau^n}{r(\tau)^n}$. By induction, we can prove that for all $n \geq 4$,

$$\tau^n \left(\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & e \end{bmatrix} \right) = \begin{bmatrix} x_{n-2}(a+e) + x_{n-3}d & 0 & 0 \\ 0 & x_{n-2}(a+e) + x_{n-3}d & 0 \\ 0 & 0 & x_{n-3}(a+e) + x_{n-4}d \end{bmatrix}.$$

where x_n denotes the n^{th} term of the Fibonacci sequence with $x_0 = 1, x_1 = 2$. Recall that, $x_n = \frac{p^{n+2} - q^{n+2}}{\sqrt{5}}$ where $p = \frac{1+\sqrt{5}}{2}$ and $q = \frac{1-\sqrt{5}}{2}$. Then, $r = \lim_{n \rightarrow \infty} \|\tau^n(I)\|^{1/n} = \lim_{n \rightarrow \infty} x_n^{1/n} = p$. Since $\lim_{n \rightarrow \infty} \frac{x_n}{r^n} = \frac{r^2}{\sqrt{5}}$ and $r^2 - r - 1 = 0$ (which implies in particular $r^{-1} = r - 1, r^{-2} = 2 - r$), we infer that $\frac{\tau^n}{r^n}$ converges as n tends to infinity and hence $\hat{\tau} = \lim_{n \rightarrow \infty} \frac{\tau^n}{r^n}$. Then direct computation yields,

$$\hat{\tau} \left(\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & e \end{bmatrix} \right) = \frac{1}{\sqrt{5}}(a + (r-1)d + e) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & (r-1) \end{bmatrix}.$$

In particular, $\hat{\tau}$ has the form $X \mapsto \psi(X)L$ for a faithful state ψ and

$$L = \hat{\tau}(I) = \frac{1+r}{\sqrt{5}} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & r-1 \end{bmatrix} = \frac{1}{\sqrt{5}} \begin{bmatrix} r^2 & 0 & 0 \\ 0 & r^2 & 0 \\ 0 & 0 & r \end{bmatrix},$$

which is positive, invertible and belongs to the eigenspace of τ corresponding to the eigenvalue r . Let $\sigma = \alpha_V^{-1} \circ \tau \circ \alpha_V$ where $V = L^{1/2}$. Then,

$$\sigma \left(\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & e \end{bmatrix} \right) = \begin{bmatrix} a + (r-1)e & 0 & 0 \\ 0 & a + (r-1)e & 0 \\ 0 & 0 & rd \end{bmatrix}.$$

We can easily verify that $\|\sigma\| = \|\sigma(I)\| = r = r(\tau)$.

Example 5.21. Let us revisit the irreducible CP map τ on $\mathcal{A} = M_1 \oplus M_1 \oplus M_1$, considered in Example 5.12. It is defined as follows:

$$\tau \left(\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \right) = \begin{bmatrix} b & 0 & 0 \\ 0 & a+c & 0 \\ 0 & 0 & b \end{bmatrix}.$$

It is easy to see that τ is diagonalizable with eigenvalues $\{0, \pm\sqrt{2}\}$. Therefore τ has spectral radius $\sqrt{2}$ and $\|\tau^n\|_\beta = 2^{\frac{n}{2}}$ for all but finitely many n . Now, $\hat{\tau}$ can be computed using its definition. It maps into the eigenspace of τ with eigenvalue $\sqrt{2}$ and is given by

$$\hat{\tau} \left(\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \right) = \frac{1}{2}(a + b\sqrt{2} + c) \begin{bmatrix} 1 & 0 & 0 \\ 0 & \sqrt{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Once again $\hat{\tau}$ is of the form described in Theorem 5.17.

ACKNOWLEDGMENTS

Bhat gratefully acknowledges funding from SERB(India) through J C Bose Fellowship No. JBR/2021/000024. Sahasrabudhe thanks the Indian Statistical Institute for hosting her through Visiting Scientist Fellowship and through J C Bose Fellowship of Bhat mentioned above.

REFERENCES

- [1] S. Albeverio and R. Høegh-Krohn, Frobenius theory for positive maps of von Neumann algebras, *Commun.Math. Phys.* **64** (1978), 83 – 94.
- [2] W. Arveson, The index of a quantum dynamical semigroup, *J. Funct. Anal.* **146** (1997), 557 – 588.
- [3] M. A. Berger and Y. Wang, Bounded semigroups of matrices, *Linear Algebra Appl.* **166** (1992), 21 – 27.
- [4] B.V.R. Bhat and T. Bhattacharyya, Dilations, Completely Positive Maps and Geometry, Text and Readings in Mathematics, *Hindustan Book Agency*, 2023.
- [5] R. Bhatia, Positive definite matrices, Princeton Series in Applied Mathematics, *Princeton University Press, Princeton, NJ*, 2007.
- [6] V. D. Blondel and Y. Nesterov, Computationally efficient approximations of the joint spectral radius, *SIAM J. Matrix Anal. Appl.* **27** (2005), 256 – 272.
- [7] F. F. Bonsall, Endomorphisms of partially ordered vector spaces, *J. London Math. Soc.* **30** (1955), 133 – 144.
- [8] M. D. Choi, Completely Positive Linear Maps on Complex Matrices, *Linear Algebra Appl.* **10** (1975), 285 – 290.
- [9] I. Daubechies and J. C. Lagarias, Sets of matrices all infinite products of which converge, *Linear Algebra Appl.* **161** (1992), 227 – 263.
- [10] L. Elsner, The generalized spectral-radius theorem: an analytic-geometric proof, *Linear Algebra Appl.* **220** (1995), 151 – 159.
- [11] D. E. Evans and R. Høegh-Krohn, Spectral properties of positive maps on C^* -algebras, *J. London Math. Soc.* **17** (1978), 345 – 355.
- [12] F. Fagnola, R. Pellicer, Irreducible and periodic positive maps, *Commun. Stoch. Anal.* **3** (2009), 407 – 418.
- [13] D. R. Farenick, Irreducible positive linear maps on operator algebras, *Proc. Amer. Math. Soc.* **124** (1996), 3381 – 3390.
- [14] S. Friedland, Characterizations of Spectral Radius of Positive Operators on C^* -Algebras, *J. Funct. Anal.* **97** (1991), 64 – 70.
- [15] V. G. Frobenius, Uber Matrizen aus positiven Elementen I, II, *S. -B. Preuss. Akad. Wiss. Berlin* (1908) 471 – 476; (1909) 514 – 518.
- [16] V. G. Frobenius, Uber Matrixen aus nicht negativen Elementen, *S. -B. Preuss. Akad. Wiss. Berlin* (1912), 456 – 477.

- [17] M. Goldberg and G. Zwas, On matrices having equal spectral radius and spectral norm, *Linear Algebra Appl.* **8** (1974), 427 – 434.
- [18] U. Groh, The peripheral point spectrum of Schwarz operators on C^* -algebra, *Math. Z.* **176** (1981), 311 – 318.
- [19] U. Groh, Some observations on the spectra of positive operators on finite-dimensional C^* -algebras, *Linear Algebra Appl.* **42** (1982), 213 – 222.
- [20] I. Halperin and P. Rosenthal, Burnside’s theorem on algebras of matrices, *Amer. Math. Monthly* **87** (1980), 810.
- [21] S. Karlin, Positive operators, *J. Math. Mech.* **8** (1959), 907 – 937.
- [22] E. C. Lance, Hilbert C^* -Modules: A Toolkit for Operator Algebraists, London Mathematical Society Lecture Note Series, *Cambridge University Press*, 1995.
- [23] M. Largo, W. S. Yang and S. Xiong, A Perron-Frobenius type of theorem for quantum operations, *J. Stat. Phys.* **169** (2017), 38 – 62.
- [24] G. J. Murphy, C^* -Algebras and Operator Theory, *Academic Press, Inc., Boston, MA*, 1990.
- [25] P. A. Parrilo and A. Jadbabaie, Approximation of the joint spectral radius using sum of squares, *Linear Algebra Appl.* **10** (2008), 2385 – 2402.
- [26] W. L. Paschke, Inner Product modules over B^* -algebras, *Trans. Amer. Math. Soc.* **182** (1973), 443 – 468.
- [27] J. E. Pascoe, An elementary method to compute the algebra generated by some given matrices and its dimension, *Linear Algebra Appl.* **571** (2019), 132 – 142.
- [28] J. E. Pascoe, The Outer Spectral Radius and dynamics of CP maps, *Israel J. Math.* **244** (2021), 945 – 969.
- [29] V. Paulsen, Completely Bounded Maps and Operator Algebras, Cambridge studies in advanced Mathematics, *Cambridge University Press*, 2003.
- [30] O. Perron, Grundtagen fiir eine Theorie der Jacobischen Kettenbruchalgorithmen, *Math. Ann.* **64** (1907), 1 – 76.
- [31] O. Perron, Zur Theorie der Matrices, *Math. Ann.* **64** (1907), 248 – 263.
- [32] M. .V. Pimsner, A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbf{Z} , *Free probability theory (Waterloo, ON, 1995)*, *Amer. Math. Soc., Providence, RI*, **12** (1997), 189 – 212.
- [33] G. Popescu, Similarity and ergodic theory of positive linear maps, *J. Reine Angew. Math.* **561** (2003), 87 – 129.
- [34] G. Popescu, Similarity problems in noncommutative polydomains, *J. Funct. Anal.* **267** (2014), 4446 – 4498.
- [35] G. C. Rota, G. Strang, A note on the joint spectral radius, *Indag. Math.* **22** (1960), 379 – 381.
- [36] H. Schaefer, Some spectral properties of positive linear operators, *Pacific J. Math.* **10** (1960), 1009 – 1019.

- [37] R. Schrader, Perron-Frobenius theory for positive maps on trace ideals, *Mathematical physics in mathematics and physics (Siena, 2000)*, *Fields Inst. Commun.* 30, Amer. Math. Soc., Providence, RI (2001) , 361 – 378.
- [38] M. Skeide, Quantum Stochastic Calculus on Full Fock Modules, *J. Funct. Anal.* **173** (2000), 401 – 452.
- [39] M. Skeide, Hilbert Modules and Applications in Quantum Probability, Habilitationsschrift, Cottbus, 2001. Available at https://www.math.tu-cottbus.de/INSTITUT/lswas/_MS/downloads/habil.pdf.
- [40] R. Speicher, Combinatorial theory of the free product with amalgamation and operator-valued free probability theory, *Mem. Amer. Math. Soc.* **132** (1998).
- [41] W. F. Stinespring, Positive function on C^* -algebras, *Proc. Amer. Math. Soc.* **6** (1955), 211 – 216.
- [42] H. Wielandt, Unzerlegbare, nicht negative Matrizen, *Mathematische Zeitschrift* **52** (1950), 642 – 648.

(Statistics and Mathematics Unit, Indian Statistical Institute, R V College Post, Bangalore - 560059, India)

Email address: bvajaramabhat@gmail.com

Email address: brsaha27700@gmail.com

Email address: praju1093@gmail.com