

# A FINITENESS RESULT FOR COMMON ZEROS OF ITERATES OF RATIONAL MAPS

CHATCHAI NOYTAPTIM AND XIAO ZHONG

**ABSTRACT.** Answering a question asked by Hsia and Tucker in their paper on the finiteness of greatest common divisors of iterates of polynomials, we prove that if  $f, g \in \mathbb{C}(X)$  are compositionally independent rational functions and  $c \in \mathbb{C}(X)$ , then there are at most finitely many  $\lambda \in \mathbb{C}$  with the property that there is an  $n$  such that  $f^n(\lambda) = g^n(\lambda) = c(\lambda)$ , except for a few families of  $f, g \in \text{Aut}(\mathbb{P}_{\mathbb{C}}^1)$  which gives counterexamples.

## 1. INTRODUCTION

**1.1. Historical background.** Bugeaud, Corvaja, and Zannier [BCZ03] provided an upper bound on the greatest common divisor (GCD) of two integer sequences using tools from Diophantine approximation. More precisely, they showed that for any multiplicatively independent integers  $a, b \geq 2$  and for any  $\epsilon > 0$ , then

$$\gcd(a^n - 1, b^n - 1) < \exp(\epsilon n)$$

for sufficiently large  $n \geq 1$ .

In 2004, Ailon and Rudnick [AR04] obtained the first instance of GCD-type result over function field of characteristic 0. In fact, they proved that for any nonconstant multiplicatively independent polynomials  $a, b \in \mathbb{C}[x]$  there exists a polynomial  $h \in \mathbb{C}[x]$  such that  $\gcd(a^n - 1, b^n - 1) | h$  for all  $n \geq 1$ . Ostafe [Os16] pointed out that the result of Ailon-Rudnick holds generally for any pair  $(m, n) \in \mathbb{N} \times \mathbb{N}$ . Ostafe also provided an upper bound on the degree of  $\gcd(a^m - 1, b^n - 1)$  in terms of degrees of  $a$  and  $b$ .

There are many generalizations and extensions of GCD-type results in various contexts, see for example [CZ13], [GHT17], [GHT18], [Gr20], [GW20], [Hu20], [Le19], [Lu05], [Si04a], [Si04b], [Si05] and [Za12].

Motivated by taking multiplicative powers of polynomials, one can study GCD-type results of iterates of rational maps. Let  $f(x) \in \mathbb{C}(x)$  be a rational map with complex coefficients. Here and what follows,  $f^n$  denotes the  $n$ -fold composition of  $f$  with itself.

**Definition 1.1.** Two rational maps  $f$  and  $g$  are said to be compositionally independent if there exist positive integers  $i_1, \dots, i_s, j_1, \dots, j_s, l_1, \dots, l_t, m_1, \dots, m_t$  such that

$$f^{i_1} \circ g^{j_1} \circ \dots \circ f^{i_s} \circ g^{j_s} = f^{l_1} \circ g^{m_1} \circ \dots \circ f^{l_t} \circ g^{m_t}$$

---

*Date:* May 27, 2024.

*2020 Mathematics Subject Classification.* 37P05, 37P30.

*Key words and phrases.* arithmetic dynamics, iterated rational maps, arithmetic equidistribution, common zeros.

The second author was supported in part by NSERC grant RGPIN-2022-02951.

then  $s = t$  and  $i_k = l_k, j_k = m_k$  for  $1 \leq k \leq s$ . In other words, the semigroup generated by  $f$  and  $g$  is isomorphic to the free semigroup of two generators.

Hsia and Tucker [HT17] first addressed the question posed by Ostafe [Os16, §4.2] in dynamical setting. Under suitable compositional assumptions, they showed that for any polynomial maps  $f, g$  of degree  $\geq 2$  with complex coefficients and  $c(x) \in \mathbb{C}[x]$  there are only finitely many irreducible factors of

$$\gcd(f^m(x) - c(x), g^n(x) - c(x))$$

for some positive integers  $m$  and  $n$  (cf. [HT17, Theorem 1]). Interestingly, Theorem 2 in [HT17] can be viewed as a compositional analog result of Ailon-Rudnick for linear polynomials. However, for nonlinear polynomials, their result holds conditionally on the assumption that  $c(x)$  is not a complex constant which is in ramified cycle of both  $f$  and  $g$  (cf. [HT17, Theorem 4]).

**1.2. Statement of main results.** In this article, we answer the following question posed by Hsia and Tucker.

**Question 1.2.** [HT17, Question 18] Let  $f, g$  be two nonconstant, compositionally independent rational maps with complex coefficients. Let  $c$  be any rational map with complex coefficients. Is it true that there must be at most finitely many  $\lambda \in \mathbb{C}$  such that  $f^n(\lambda) = g^n(\lambda) = c(\lambda)$  for some positive integer  $n$ ?

It turns out that Question 1.2 is not true in general. This can be seen from the counter-examples provided in Remark (3.6), Remark (3.7) and Remark (3.8).

However, after excluding these families of rational maps  $f$  and  $g$ , we obtain a finiteness result of common zeros of iterated rational maps in a similar spirit to Question 1.2. More precisely, we state our main results as follows.

**Theorem 1.3.** *Let  $f$  and  $g$  be nonconstant automorphisms on  $\mathbb{P}^1(\mathbb{C})$  such that  $f$  and  $g$  are compositionally independent. Let  $c$  be any rational function with complex coefficients. Suppose that  $f(x)$  and  $g(x)$  are not conjugate simultaneously by an automorphisms on  $\mathbb{P}^1(\mathbb{C})$  to one of the following:*

- (1)  $\alpha X + \beta, X/(\gamma X + \delta)$ , with some  $\alpha, \beta, \gamma, \delta \in \mathbb{C}^*$  such that  $\alpha/\delta$  is a root of unity;
- (2)  $\alpha X + \beta, \delta X + \gamma$ , with some  $\alpha, \delta \in \mathbb{C}^*$  such that  $\alpha$  and  $\delta$  are not roots of unity and  $\alpha/\delta$  is a root of unity other than 1;

Then there are only finitely many  $\lambda \in \mathbb{C}$  such that

$$f^n(\lambda) = g^n(\lambda) = c(\lambda)$$

for some  $n \geq 1$ .

When at most one of rational maps  $f$  and  $g$  is of degree one, we can relax some conditions on  $(m, n) \in \mathbb{N} \times \mathbb{N}$ . Also, the finiteness result on the number zeros of both equations  $f^m(x) - c(x) = 0$  and  $g^n(x) - c(x) = 0$  holds under natural assumptions. This is reminiscent of polynomial maps counterpart (cf. [HT17, Theorem 1]).

**Theorem 1.4.** *Let  $f$  and  $g$  be two compositionally independent rational functions with complex coefficients, at least one of which has degree greater than one. Suppose that  $c(x) \in \mathbb{C}(x)$  is not a compositional power of  $f$  or  $g$ . Then there are only finitely many  $\lambda \in \mathbb{C}$  such that*

$$f^m(\lambda) = g^n(\lambda) = c(\lambda)$$

for some positive integers  $m$  and  $n$ .

Notice that the same statement allowing distinct  $n$  and  $m$  for the case that  $f$  and  $g$  are both automorphisms on  $\mathbb{P}^1(\mathbb{C})$  doesn't hold. There is a counter-example in the beginning of Section 3 of [HT17].

**1.3. Proof strategy.** Hsia and Tucker [HT17] combine various tools and techniques from Diophantine geometry, complex dynamics, and number theory. Our approach is akin to their method. There are technical difficulties to overcome for rational maps. We briefly explain our ideas of proving Theorem 1.3 and Theorem 1.4.

For automorphisms  $f$  and  $g$  defined over  $\overline{\mathbb{Q}}$ , we heavily hinge on tools from Diophantine geometry (e.g., Roth's theorem, Siegel's theorem, and  $S$ -unit theorem). After suitable excluding  $f, g$  (see §1.2), we show that  $f$  and  $g$  must share a common fixed point (see Lemma 3.2). Then we dynamically conjugate by moving such fixed point to  $\infty$ , we may assume that  $f(x) = \alpha x + \beta$  and  $g(x) = \delta x + \gamma$  with  $\alpha, \beta, \gamma, \delta \in \overline{\mathbb{Q}}^*$ . We break our computation into several cases depending on whether  $\alpha$  and  $\delta$  is a root of unity to show that if there are infinitely many solutions to

$$f^n(x) = g^n(x) = c(x)$$

where  $c(x)$  is any rational map with complex coefficients, then  $f$  and  $g$  are not compositionally independent. In other words, the semigroup generated by  $f$  and  $g$  under composition is not free (see Theorem 3.5). Using specialization argument, we obtain the desired result over  $\mathbb{C}$ .

When exactly one of rational maps  $f, g \in \mathbb{C}(x)$  has degree greater than one, the proof follows verbatim in the polynomial maps setting as a consequence of Northcott's property explained in [HT17, Proposition 9].

In the case both rational maps  $f, g \in \mathbb{C}(x)$  are of degree greater than 1 and  $c(x) \in \mathbb{C}(x)$ . Under suitable compositional assumptions, we show that the infinite sequence of solutions to

$$f^{m_i}(x_i) = g^{n_i}(x_i) = c(x_i)$$

as  $i \rightarrow +\infty$  (so  $m, n \rightarrow +\infty$ ) form a sequence of small dynamical height (Lemma 4.1). We, then, apply the arithmetic equidistribution theorem of small points (in reminiscence of [BD11], [BD13], [GHT13], [GHT15], and [PST12]) to deduce that dynamical height attached to  $f$  and  $g$  must be identical (Proposition 4.2) and hence  $f$  and  $g$  must share the same set of preperiodic points (i.e., points of finite forward orbit under iteration). Unlike polynomial maps setting, one cannot appeal to the classification of polynomials with the same Julia set of [BE87] and [SS95]. Alternatively, we apply the Tits alternative for rational maps of [BHPT] to deduce that  $f$  and  $g$  are not compositionally independent.

## 2. PRELIMINARIES

**2.1. Absolute values over global fields.** Here and throughout this section,  $K$  denote a global field (unless otherwise stated) equipped with a set of inequivalent absolute values (places) satisfying the product formula. For example,  $K$  could be a number field or a function field of positive transcendence degree over another field  $k$ . It was pointed out in [Art06] that a global field which has an archimedean place must be a number field.

If  $K$  is a number field, we let  $M_K$  to be the set of all absolute values of  $K$  which extends the absolute values of  $\mathbb{Q}$ . If  $K$  is a function field of finite transcendence degree  $m \geq 1$  defined over a field  $k$ , then  $M_K$  is the set of all absolute values on  $K$  associated to the irreducible hypersurfaces of the projective space  $\mathbb{P}_k^m$  (whose function field is  $K' := k(t_1, \dots, t_m)$  where  $t_1, \dots, t_m$  are algebraically independent over  $k$ -functions in  $K$ ). Thus there exist  $N_v \geq 1$  such that the product formula  $\prod_{v \in M_K} |x|_v^{N_v} = 1$  holds for any nonzero  $x \in K$ . (see [BG06] and [La83]).

Let  $K_v$  be the completion of  $K$  with respect to absolute value  $|\cdot|_v$ . Then  $\mathbb{C}_v$  is the completion of the algebraic closure of  $K_v$  and it is algebraically closed and complete. Note that, by abuse notation, we still denote the absolute value on  $\mathbb{C}_v$  which extends uniquely from  $|\cdot|_v$  on  $K$  by  $|\cdot|_v$ .

**2.2. Dynamical height and preperiodic points.** Let  $S$  be the finite set of Galois conjugates  $\text{Gal}(\overline{K}/K) \cdot x$  of  $x$  in  $\mathbb{P}^1(\overline{K})$ . Then we define the Weil height  $h : \mathbb{P}^1(\overline{K}) \rightarrow [0, +\infty)$  by

$$h(x) := \frac{1}{|S|} \sum_{x \in S} \sum_{v \in M_K} N_v \log \max\{1, |x|_v\}.$$

In light of Baker-Rumely's construction [BR10, §7.9], our Weil height is defined as the average over Galois conjugate of points in  $\overline{K}$ . It was pointed out that this definition does not depend on the choice of embedding  $\overline{K} \hookrightarrow \mathbb{C}_v$  as  $S$  is Galois stable subset of  $\mathbb{P}^1(\overline{K})$ .

Let  $f$  be a rational map defined over a global field  $K$  of degree  $\geq 2$ . Then the Call-Silverman canonical height  $\hat{h}_f : \mathbb{P}^1(\overline{K}) \rightarrow [0, +\infty)$  is defined by

$$\hat{h}_f(x) = \lim_{n \rightarrow \infty} \frac{1}{(\deg f)^n} h(f^n(x)).$$

We record important properties of Weil height and dynamical canonical height as follows.

**Proposition 2.1.** ([Si07, §3], [CS93]) *Let  $f$ ,  $h$  and  $\hat{h}_f$  be as above. Then*

- (i)  $h(f(x)) = (\deg f)h(x) + O(1)$ .
- (ii)  $\hat{h}_f(x) = h(x) + O(1)$ .
- (iii)  $\hat{h}_f(f(x)) = (\deg f)\hat{h}_f(x)$ .

Over a number field, dynamical height is used to detect preperiodic points in the following sense.

**Proposition 2.2.** ([Si07, Theorem 3.22]) *Let  $f(x) \in K(x)$  be a rational map of degree  $\geq 2$  defined over a number field  $K$ . Then  $x \in \mathbb{P}^1(\overline{K})$  is  $f$ -preperiodic if and only if  $\hat{h}_f(x) = 0$ .*

The situation is different over function field. The dynamical height of a function defined on a function field (over an infinite field) fails to properly detect preperiodic points. As an interesting example shown by Benedetto [Be05], we consider  $f(x) = x^2 \in K[x]$  defined over a function field  $K = \overline{\mathbb{Q}}(T)$  of transcendence degree 1 with constant field  $\overline{\mathbb{Q}}$ . Notice that all points in  $\mathbb{P}^1(\overline{\mathbb{Q}})$  have dynamical height zero but  $0, \infty$  and roots of unity in  $\overline{\mathbb{Q}}$  are the only  $f$ -preperiodic points. In fact,  $f$  is defined over the constant field of  $K$ .

Recall that a rational map  $f(x) \in K(x)$  defined over a function field  $K$  is called **isotrivial** if there exists a finite extension  $K'/K$  and a Möbius transformation  $\phi \in \mathrm{PGL}(2, K')$  such that  $\phi^{-1} \circ f \circ \phi$  defined over the constant field of  $K'$ . More related results, see [Ba09], [Be05], and [PST09].

After excluding maps with isotriviality, the dynamical height works as expected. The following result is due to Benedetto (for polynomial maps) and Baker (for rational maps) over  $\mathbb{P}^1$ .

**Theorem 2.3.** ([Ba09], [Be05]) *Let  $f(x) \in K(x)$  be a rational map of degree  $\geq 2$  defined over a function field  $K$ . Suppose that  $f$  is not isotrivial. Then  $x \in \mathbb{P}^1(\overline{K})$  is  $f$ -preperiodic if and only if  $\hat{h}_f(x) = 0$ .*

**2.3. Arakelov-Green's function.** The dynamical height attached to rational maps can be decomposed in terms of Arakelov-Green's function. We follow the construction in Baker-Rumely [BR10, §10.3]. Suppose that  $f \in K(x)$  is a rational function of degree  $\geq 2$  defined over a global field  $K$ . The  $v$ -adic Arakelov-Green's function of  $f$  is a function

$$G_{f,v} : \mathbb{P}_{\mathrm{Berk},v}^1 \times \mathbb{P}_{\mathrm{Berk},v}^1 \rightarrow \mathbb{R} \cup \{\infty\}$$

which is symmetric, finite and continuous off the diagonal in  $\mathbb{P}^1(\mathbb{C}_v) \times \mathbb{P}^1(\mathbb{C}_v)$ . Here  $\mathbb{P}_{\mathrm{Berk},v}^1$  is the Berkovich projective line, we refer the reader to [Be19] and [BR10] for detailed exposition. For each  $y \in \mathbb{P}_{\mathrm{Berk},v}^1$ , the function  $G_{f,v}(x, y)$  is bounded differential variation on  $\mathbb{P}_{\mathrm{Berk},v}^1$  and it satisfies the differential equation

$$(2.1) \quad \Delta_x G_{f,v}(x, y) = \mu_{f,v}(x) - \delta_y(x)$$

where  $\mu_{f,v}$  is the equilibrium measure characterized by  $f^*(\mu_{f,v}) = (\deg f)\mu_{f,v}$  and  $G_{f,v}$  is normalized so that

$$\iint_{\mathbb{P}_{\mathrm{Berk},v}^1 \times \mathbb{P}_{\mathrm{Berk},v}^1} G_{f,v}(x, y) d\mu_{f,v}(x) d\mu_{f,v}(y) = 0.$$

Our Laplacian is the negative of the one introduced by Baker-Rumely [BR06, §3.5]. For any  $x, y \in \mathbb{P}^1(\overline{K})$  such that  $x \neq y$ , we have

$$\hat{h}_f(x) + \hat{h}_f(y) = \sum_{v \in M_K} N_v G_{f,v}(x, y)$$

where the positive number  $N_v$  is the same normalization as appeared in the product formula.

**2.4. Results from Diophantine geometry.** Here we collect some Diophantine geometry results that will be used in the proof.

**Theorem 2.4.** [EG15, Theorem 6.1.3] *Let  $K$  be a field of characteristic 0, let  $n \geq 2$ , let  $a_1, \dots, a_n \in K^*$  and let  $\Gamma$  be a subgroup of  $(K^*)^n$  of finite rank  $r$ . Then the number of non-degenerate solutions of*

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 1$$

*in  $(x_1, x_2, \dots, x_n) \in \Gamma$  is finite.*

**Remark 2.5.** We call a solution to the equation  $a_1x_1 + a_2x_2 + \dots + a_nx_n = 1$  like above non-degenerate if

$$\sum_{i \in I} a_i x_i \neq 0$$

for any proper subset  $I \subset \{1, 2, \dots, n\}$ .

**Proposition 2.6.** [HT17, Proposition 12] *Let  $W$  be a one-dimensional subtorus in  $\mathbb{G}_m^2$  defined over a number field  $K$  and let  $S$  be a finite set of places of  $K$  containing all the archimedean places. Let*

$$\Phi(X, Y) = P(X, Y)/Q(X, Y)$$

*where  $P, Q \in K[X, Y]$  are two relatively prime polynomials neither of which is divisible by  $X$  or  $Y$ . Assume that  $\Phi$  restricts to a nonconstant rational function  $\phi$  on  $W$  with at least a pole in  $W(\bar{K})$ . Let  $\Gamma$  be a finitely generated subgroup of  $W(K)$ . Then, there are at most finitely many points  $q \in \Gamma$  such that  $\Phi(q)$  is an  $S$ -integer.*

**Lemma 2.7.** [HT17, Lemma 14] *Let  $w_1$  and  $w_2$  be two multiplicatively independent elements of a number field  $K$ , neither of which is a root of unity, and let  $y$  be a nonzero element of  $K$ . Let  $S$  be a finite set of places of  $K$  including all the archimedean places. Then for all sufficiently large  $n$ , there is a  $v \notin S$  such that  $|w_1^n - y|_v < |w_2^n - y|_v \leq 1$ .*

## 2.5. The Ping-pong Lemma.

**Lemma 2.8.** *Let  $G$  be a group of maps under compositions. Let  $S = \langle f_1, \dots, f_r \rangle$ ,  $r \geq 2$  and  $f_i \in G$ , for  $i \in \{1, 2, \dots, r\}$ , be a semigroup of  $G$ . If there exists a collection of non-empty mutually non-intersecting sets  $I_1, \dots, I_r$  such that  $f_i(\bigcup_{j=1}^r I_j) = I_i$  for all  $1 \leq i \leq r$ , then  $S$  is a free semigroup of rank  $r$  with free basis  $f_1, \dots, f_r$ .*

*Proof.* This is basically the Ping-pong Lemma stated in [KT21, The Ping-Pong Lemma]. The only difference is that it is stated for  $G$  being the group of affine linear maps on  $\mathbb{R}$ . However the proof doesn't rely on this assumption.  $\square$

**2.6. Asymptotic notation.** We clarify the asymptotic notation here:

**Definition 2.9.** Consider two functions  $f(n)$  and  $g(n)$  with  $n \in \mathbb{N}$  and take values in  $\mathbb{C}$ . We defined the asymptotic notation when  $n$  goes large as follows:

- (1) We write  $f(n) \sim g(n)$  if there exists a non-zero constant  $k \in \mathbb{C}$  such that

$$\lim_{n \rightarrow \infty} f(n)/g(n) = k;$$

(2) We write  $f(n) = o(g(n))$  if

$$\lim_{n \rightarrow \infty} f(n)/g(n) = 0;$$

(3) We write  $f(n) = O(g(n))$  if

$$\limsup_{n \rightarrow \infty} f(n)/g(n) < \infty.$$

Furthermore, for two functions  $f(Z)$  and  $g(Z)$  take values in  $\mathbb{C}$  with  $Z \in \mathbb{C}$ . We defined the asymptotic notation when  $Z$  is close to 0 similarly as above but taking limits with respect to  $Z \rightarrow 0$ .

### 3. PROOF OF THEOREM 1.3

**Lemma 3.1.** *Let*

$$X_n = \left( -t_n - \sqrt{t_n^2 - 4\alpha^n \delta^n \frac{(1-\alpha^n)(1-\delta^n)}{(1-\alpha)(1-\delta)} \beta \gamma} \right) / \left( 2\alpha^n \frac{1-\delta^n}{1-\delta} \gamma \right),$$

$$t_n = \frac{(1-\alpha^n)(1-\delta^n)}{(1-\alpha)(1-\delta)} \beta \gamma + \alpha^n \delta^n - 1,$$

where  $\beta, \gamma \in \overline{\mathbb{Q}}^*$ ,  $\alpha, \delta$  and  $\alpha/\delta$  are not roots of unity in  $\overline{\mathbb{Q}}^*$ . Suppose for infinitely many  $n \in \mathbb{Z}^+$ , we have

$$(3.1) \quad \alpha^n X_n + \frac{1-\alpha^n}{1-\alpha} \beta = A(X_n)/B(X_n),$$

where  $A$  and  $B$  are two co-prime polynomials. Then  $\deg(A) = \deg(B) + 1$ .

*Proof.* Let  $K$  be a number field such that  $\alpha, \delta, \beta$  and  $\gamma$  are all defined in  $K$  and  $A(X)$  and  $B(X)$  are defined over  $K$ . Let  $S$  be a finite set of places of  $K$ , such that for any  $v \notin S$  we have

$$|\alpha|_v = |\beta|_v = |\delta|_v = |\gamma|_v = 1$$

and all the coefficients of  $A$  and  $B$  are of norm 1 with respect to  $|\cdot|_v$ .

Suppose there are infinitely many  $X_n$  satisfies the Equation (3.1), by Lemma 2.7, except for finitely many  $n \in \mathbb{Z}^+$ , there exists a  $v \notin S$  such that

$$(3.2) \quad \left| \frac{1-\alpha^n}{1-\delta^n} \right|_v > 1.$$

Fix a  $v \notin S$  such that (3.2) and Equation (3.1) hold for some  $n \in \mathbb{Z}^+$ . Fix such a  $n$  and then

$$\begin{aligned} & \sqrt{t_n^2 - 4\alpha^n \delta^n \frac{(1-\alpha^n)(1-\delta^n)}{(1-\alpha)(1-\delta)} \beta \gamma} / \left( 2\alpha^n \frac{1-\delta^n}{1-\delta} \gamma \right) \\ &= \frac{(1-\delta)\delta^n}{2\alpha^n \gamma} \frac{1-\alpha^n}{1-\delta^n} + O(1), \end{aligned}$$

here  $O(1)$  means the terms with absolute value not greater than 1 and the absolute value is associated to the valuation  $v$ . Thus,

$$X_n = -2 \frac{(1-\delta)\delta^n}{2\alpha^n \gamma} \frac{1-\alpha^n}{1-\delta^n} + O(1),$$

and from this we have

$$|X_n|_v > 1.$$

Therefore, Equation (3.1) implies that

$$|X_n|_v^{\deg(A)} = |X_n|_v^{\deg(B)+1}$$

and thus

$$\deg(A) = \deg(B) + 1.$$

□

**Lemma 3.2.** *Let  $f(X), g(X)$  be automorphisms on  $\mathbb{P}^1$  defined over  $\mathbb{C}$  and  $c(X)$  be a rational function defined over  $\mathbb{C}$ . If there are infinitely many  $n \in \mathbb{Z}^+$  such that*

$$(3.3) \quad c(X) = f^n(X) = g^n(X)$$

*has solutions, and  $f(X), g(X)$  are not simultaneously conjugated to  $\alpha X + \beta, X/(\gamma X + \delta)$  respectively, with some  $\alpha, \delta, \beta, \gamma \in \mathbb{C}^*$  satisfy that either at least one of  $\alpha, \delta$  is root of unity other than 1 or  $\alpha/\delta$  is a root of unity, then  $f$  and  $g$  share a common fixed point.*

*Proof.* Notice that if  $\gamma$  or  $\beta$  is zero, then certainly  $f(X)$  and  $g(X)$  share a common fixed point. Let's first assume that  $f(X), g(X)$  and  $c(X)$  are all defined over  $\overline{\mathbb{Q}}$  and  $\beta, \gamma \in \overline{\mathbb{Q}}^*$ . Suppose  $f$  and  $g$  don't share a fixed point. After conjugation, we assume that

$$\begin{aligned} f(X) &= \alpha X + \beta, \\ g(X) &= \frac{X}{\gamma X + \delta}. \end{aligned}$$

We always assume that  $\alpha$  and  $\delta$  are non-zero as otherwise it will result in a constant map. We break the discussion into two cases depending on  $\alpha$  and  $\delta$ .

**Case I.** Suppose  $\alpha$  is not a root of unity and  $\delta = 1$ .

**Lemma 3.3.** *There are only finitely many  $n \in \mathbb{Z}^+$  such that*

$$c(X) = f^n(X) = g^n(X)$$

*has solutions.*

The same result for  $\alpha = 1$  and  $\delta$  is not a root of unity holds. We just need to further conjugate  $c(X), f(X)$  and  $g(X)$  by  $1/X$  to swap the role of  $\alpha$  and  $\delta$ . Then the same argument will give the result. This concludes these cases.

**Case II.** Now, suppose both  $\alpha$  and  $\beta$  are not roots of unity and  $\alpha/\delta$  is not a root of unity. Then let

$$X_n^\pm = \left( -t_n \pm \sqrt{t_n^2 - 4\alpha^n \delta^n \frac{(1-\alpha^n)(1-\delta^n)}{(1-\alpha)(1-\delta)} \beta \gamma} \right) / \left( 2\alpha^n \frac{1-\delta^n}{1-\delta} \gamma \right),$$

be the solution to the  $f^n(X) = g^n(X)$ , where

$$t_n = \frac{(1-\alpha^n)(1-\delta^n)}{(1-\alpha)(1-\delta)} \beta \gamma + \alpha^n \delta^n - 1.$$

We break the proof into steps:



- **Step I:** we will show that  $c(X) = f^n(X) = g^n(X)$  has solution for infinitely many  $n$  will pose a restriction on  $\alpha$  and  $\delta$ : there exists a non-trivial polynomial in two variables  $F(X, Y)$  such that  $F(\alpha^n, \delta^n) = 0$  holds for infinitely many  $n \in \mathbb{Z}^+$ .
- **Step II:** we use S-unit Theorem to argue that this implies  $\alpha$  and  $\delta$  are multiplicatively dependent.
- **Step III:** we deduce that then  $c(X) = f^n(X) = g^n(X)$  will only have solution for infinitely many  $n$  if  $f(X)$  and  $g(X)$  share a common fixed point.

**Step I.** Suppose there are infinitely many  $n \in \mathbb{Z}^+$  such that either

$$(3.4) \quad c(X_n^-) = f^n(X_n^-)$$

or

$$(3.5) \quad c(X_n^+) = f^n(X_n^+).$$

We claim that the following equation, which is obtained from above by replacing  $\alpha^n$  with a variable  $Z$  and  $\delta^n$  with a variable  $W$ ,

$$(3.6) \quad c(X^\pm(Z, W)) = ZX^\pm(Z, W) + \frac{1-Z}{1-\alpha}\beta$$

doesn't hold constantly for every  $Z, W \in \overline{\mathbb{Q}}$ , where

$$X^\pm(Z, W) = \left( -t(Z, W) \pm \sqrt{t(Z, W)^2 - 4ZW \frac{(1-Z)(1-W)}{(1-\alpha)(1-\delta)}\beta\gamma} \right) / \left( 2Z \frac{1-W}{1-\delta}\gamma \right),$$

$$t(Z, W) = \frac{(1-Z)(1-W)}{(1-\alpha)(1-\delta)}\beta\gamma + ZW - 1.$$

To see this, if Equation (3.5) holds for infinitely many  $n$ , then, taking a specialization at  $W = 0$ , we have

$$X^+(Z, W)|_{W=0} = 0,$$

and Equation (3.6) implies

$$c(0) = \frac{1-Z}{1-\alpha}\beta$$

which only holds for finitely many values of  $Z$ . So,

$$c(X^+(Z, W)) = ZX^+(Z, W) + \frac{1-Z}{1-\alpha}\beta$$

doesn't hold constantly for very value of  $Z$  and  $W$ .

On the other hand, if Equation (3.4) holds for infinitely many  $n$ , then we have

$$X_0^-(Z) = X^-(Z, W)|_{W=0} = \frac{\beta}{1-\alpha} + \left( \frac{1-\delta}{\gamma} - \frac{\beta}{1-\alpha} \right) \frac{1}{Z}.$$

If

$$\frac{1-\delta}{\gamma} = \frac{\beta}{1-\alpha}$$

then  $f(X)$  and  $g(X)$  share a common fixed point and we are done. So, we assume

$$\frac{1-\delta}{\gamma} \neq \frac{\beta}{1-\alpha}.$$

Then Equation (3.6) implies

$$(3.7) \quad c(X_0^-(Z)) = A(X_0^-(Z))/B(X_0^-(Z)) = ZX_0^-(Z) + \frac{1-Z}{1-\alpha}\beta = \frac{1-\delta}{\gamma}.$$

However, by Lemma 3.1, we have  $\deg(A) = \deg(B) + 1$ . Thus, we have

$$\text{val}_Z(A(X_0^-(Z))/B(X_0^-(Z))) < 0,$$

but

$$\text{val}_Z(ZX_0^-(Z) + \frac{1-Z}{1-\alpha}\beta) \geq 0,$$

which implies that Equation (3.7) only holds for finitely many values of  $Z$  as

$$A(X_0^-(Z))/B(X_0^-(Z)) - ZX_0^-(Z) - \frac{1-Z}{1-\alpha}\beta$$

is a non-constant rational function with variable  $Z$ . Thus

$$c(X^-(Z, W)) = f^n(X^-(Z, W))$$

doesn't hold constantly.

**Step II.** Now, either

$$c(X^+(Z, W)) = (X^+(Z, W)) + \frac{1-Z}{1-\alpha}\beta$$

or

$$c(X^-(Z, W)) = (X^-(Z, W)) + \frac{1-Z}{1-\alpha}\beta$$

is a non-trivial equation on variables  $Z$  and  $W$  with  $(Z, W) = (\alpha^n, \delta^n)$  as solutions for infinitely many  $n \in \mathbb{Z}^+$ . Thus, this gives a polynomial, by clearing the denominators and possible square roots on both sides of the equation,

$$(3.8) \quad \sum_{1 \leq i \leq k_1, 1 \leq j \leq k_2} c_{i,j} \alpha^{in} \delta^{jn} = 0$$

for infinitely many  $n \in \mathbb{Z}^+$ . Then, we want to apply the S-unit theorem Theorem 2.4 to show that  $\alpha$  and  $\delta$  are multiplicatively dependent. Let  $G \subset \overline{\mathbb{Q}}^*$  be the finitely generated subgroup generated by  $\alpha$  and  $\delta$ . Then Equation (3.8) can be rewritten as

$$(3.9) \quad \sum_{1 \leq i \leq m} c_i W_i = 0,$$

for some positive integer  $m$ ,  $c_i \in \overline{\mathbb{Q}}^*$  and  $W_i \in G$  for each  $i = 1, 2, \dots, m$ , has solutions

$$(W_1, W_2, \dots, W_m) = (\alpha^{s_1 n} \delta^{t_1 n}, \alpha^{s_2 n} \delta^{t_2 n}, \dots, \alpha^{s_m n} \delta^{t_m n}) \in G^m,$$

for infinitely many  $n \in \mathbb{Z}^+$ , where  $(s_i, t_i)$ 's are distinct pairs of integers. The S-unit theorem 2.4 says that there are only finitely many non-degenerate solutions to Equation (3.9). After an easy induction argument on numbers of terms involved in Equation (3.8) we can get that there exists infinitely many  $n \in \mathbb{Z}^+$  such that

$$c_{i_1} \alpha^{s_{i_1} n} \delta^{t_{i_1} n} + c_{i_2} \alpha^{s_{i_2} n} \delta^{t_{i_2} n} = 0,$$

where  $i_1$  and  $i_2$  are distinct and in  $\{1, 2, \dots, m\}$ . Since  $\alpha$  and  $\delta$  are not zero and not roots of unity, this is only possible if  $s_{i_1} \neq s_{i_2}$  and  $t_{i_1} \neq t_{i_2}$  and  $c_{i_1}/c_{i_2}$  is a root of unity. This implies that  $\alpha$  and  $\delta$  are multiplicative dependent. In another word, there exist non-zero integers  $k_1$  and  $k_2$  such that  $\alpha^{k_1} = \delta^{k_2}$ .

**Step III.** Let  $k = k_1/k_2 \in \mathbb{Q}^*$  such that  $\delta = \xi\alpha^k$ , where  $\xi$  is a root of unity of order  $k_2 \in \mathbb{Z}^+$  and  $\alpha^k$  is understood as a particular choice of  $k_2$ -th root of  $\alpha^{k_1}$ . Then our assumption implies that there exists an  $i \in \{1, 2, \dots, k_2\}$  such that either the system of equations

$$(3.10) \quad A(X^+(Z, \xi^i Z^k)) = B(X^+(Z, \xi^i Z^k))(ZX^+(Z, \xi^i Z^k) + \frac{1-Z}{1-\alpha}\beta),$$

(3.11)

$$A(X^+(Z, \xi^i Z^k))\left(\frac{1-\xi^i Z^k}{1-\delta}\gamma X^+(Z, \xi^i Z^k) + \xi^i Z^k\right) = B(X^+(Z, \xi^i Z^k))X^+(Z, \xi^i Z^k)$$

or the system of equations

$$(3.12) \quad A(X^-(Z, \xi^i Z^k)) = B(X^-(Z, \xi^i Z^k))(ZX^-(Z, \xi^i Z^k) + \frac{1-Z}{1-\alpha}\beta),$$

(3.13)

$$A(X^-(Z, \xi^i Z^k))\left(\frac{1-\xi^i Z^k}{1-\delta}\gamma X^-(Z, \xi^i Z^k) + \xi^i Z^k\right) = B(X^-(Z, \xi^i Z^k))X^-(Z, \xi^i Z^k)$$

has infinitely many values of  $Z$ , given by  $Z = \alpha^{nk_2+i}$  for infinitely many  $n \in \mathbb{Z}^+$ , as solutions. This cannot happen unless the system of equations holds constantly for every possible values of  $Z$  as it only has one variable. From now on, we denote

$$X_i^\pm(Z) = X^\pm(Z, \xi^i Z^k).$$

Notice that

$$(3.14) \quad X_i^+(Z)X_i^-(Z) = \frac{\xi^i Z^k(1-Z)(1-\delta)\beta}{Z(1-\xi^i Z^k)(1-\alpha)\gamma},$$

$$(3.15) \quad X_i^+(Z) + X_i^-(Z) = -2t(Z)/(2Z\frac{1-\xi^i Z^k}{1-\delta}\gamma),$$

for every  $i \in \{1, 2, \dots, k_2\}$ .

Let's first suppose the system of Equation (3.12) and (3.13) have infinitely many zeros. That is saying they hold constantly for every  $Z$ .

We do the power series expansion of  $X_i^-(Z)$  around  $Z = 0$  and get that if  $k > 0$

$$X_i^-(Z) = \frac{1-\delta}{2c}Z^{-1}\left(-2\frac{\beta\gamma}{(1-\alpha)(1-\delta)} + 2 + O(Z)\right);$$

if  $k < 0$  then

$$X_i^-(Z) = -\frac{1-\delta}{2c\xi^i}Z^{-1-k}\left(2\frac{\beta\gamma}{(1-\alpha)(1-\delta)}\xi^i Z^k + O(1)\right).$$

In both cases, we have that

$$\text{val}_Z(X_i^-(Z)) = -1.$$

Also, by Lemma 3.1, we have

$$\deg(A) = \deg(B) + 1.$$

However, Equation (3.12) implies that

$$\text{val}_Z(A(X_i^-(Z))) = -\deg(A) = \text{val}_Z(B(X_i^-(Z))(ZX_i^-(Z) + \frac{1-Z}{1-\alpha}\beta)) \geq -\deg(B).$$

This is a contradiction.

Next, suppose that  $k > 0$  and the system of Equation (3.10) and (3.11) have infinitely many zeros. Since  $\text{val}_Z(X_i^-(Z)) = -1$ , by Equation (3.14), we have

$$\text{val}_Z(X_i^+(Z)) = k > 0.$$

Then Equation (3.10) implies that

$$(3.16) \quad A(X_i^+(Z))/B(X_i^+(Z)) = \frac{\beta}{1-\alpha} - \frac{\beta}{1-\alpha}Z + o(Z).$$

If  $k > 1$ , we will have

$$A(X_i^+(Z))/B(X_i^+(Z)) = A(0)/B(0) + O(Z^k),$$

which contradicts Equation (3.16). Similarly, if  $0 < k < 1$ , we conjugate  $f(X)$ ,  $g(X)$  and  $c(X)$  by  $1/X$  to swap the role of  $\alpha$  and  $\delta$  and we get  $k > 1$  again. The same argument shows that this case is also not possible. Thus, we left with the case  $k = 1$ . However, our assumption implies that  $k \neq 1$  since otherwise  $\alpha/\delta$  is a root of unity. So we are done in this case.

On the other hand, if  $k < 0$  and system of Equation (3.12) and (3.13) have infinitely many zeros, we have again by Equation (3.14) and the fact that  $\text{val}_Z(X_i^-(Z)) = -1$  that  $\text{val}_Z(X_i^+(Z)) = 0$ . This implies that

$$X_i^+(Z) = c_0 + o(1)$$

for some non-zero constant  $c_0$  when  $Z$  is closed to 0. Since  $A(X)$  and  $B(X)$  don't share a common zero, we have, by Equation (3.10) and (3.11) that

$$\frac{1-\delta}{\gamma} = A(c_0)/B(c_0) = \frac{\beta}{1-\alpha}.$$

This implies that  $f$  and  $g$  share a common fixed point.

Now we have proved the lemma with the assumption that everything is algebraic. The following lemma concludes the proof.

**Lemma 3.4.** *It is enough to prove the Lemma 3.2 with the assumption that everything is defined over  $\overline{\mathbb{Q}}$ .*

□

*Proof of Lemma 3.3.* Suppose there are infinitely many  $n \in \mathbb{Z}^+$  such that

$$c(X) = f^n(X) = g^n(X)$$

has solutions. Let's denote

$$X_n^\pm = \frac{-t_n \pm \sqrt{t_n^2 - 4\alpha^n n \gamma (1 - \alpha^n) \beta / (1 - \alpha)}}{2n\alpha^n \gamma},$$

$$t_n = \alpha^n + n\gamma \frac{1 - \alpha^n}{1 - \alpha} \beta - 1,$$

where  $X_n^\pm$  satisfies  $f^n(X_n^\pm) = g^n(X_n^\pm)$ .

Let's first suppose  $|\alpha| > 1$  and there are infinitely many  $n \in \mathbb{Z}^+$  such that

$$(3.17) \quad c(X_n^+) = f^n(X_n^+)$$

holds. Let's denote  $c(X) = A(X)/B(X)$  where  $A(X)$  and  $B(X)$  are two polynomials. Since

$$(3.18) \quad \begin{aligned} t_n^2 - 4\alpha^n n \gamma \frac{1 - \alpha^n}{1 - \alpha} \beta &= \left( \frac{n\gamma\alpha^n\beta}{1 - \alpha} \right)^2 + \frac{2n\gamma\beta\alpha^{2n}}{1 - \alpha} + \alpha^{2n} - \frac{2n^2\gamma^2\alpha^n\beta^2}{(1 - \alpha)^2} \\ &- \frac{2\gamma\beta}{1 - \alpha} n\alpha^n - 2\alpha^n + \frac{\gamma^2\beta^2}{(1 - \alpha)^2} n^2 - \frac{2n\gamma\beta}{1 - \alpha} + o(n), \end{aligned}$$

we have

$$(3.19) \quad X_n^\pm = \frac{\left( \frac{\gamma\beta}{1 - \alpha} n\alpha^n - \alpha^n - \frac{\gamma\beta}{1 - \alpha} n + 1 \pm \left( -\frac{\gamma\beta}{1 - \alpha} n\alpha^n - \alpha^n + \frac{n\gamma\beta}{1 - \alpha} + 1 - 2\alpha^{-n} + o(\alpha^{-n}) \right) \right)}{2n\alpha^n\gamma}.$$

Therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} |X_n^+| &= 0, \\ \lim_{n \rightarrow \infty} |\alpha^n X_n^+| &= \infty. \end{aligned}$$

Since we have

$$X_n^+ \sim 1/n$$

when  $n$  goes large, we have similarly

$$\begin{aligned} A(X_n^+)/B(X_n^+) &\sim n^k, \\ \alpha^n X_n^+ + \frac{1 - \alpha^n}{1 - \alpha} \beta &\sim \alpha^n \end{aligned}$$

when  $n$  is large with some integer  $k$ , which contradicts the Equation (3.17).

Suppose  $|\alpha| > 1$  and there are infinitely many  $n \in \mathbb{Z}^+$  such that

$$(3.20) \quad c(X_n^-) = f^n(X_n^-)$$

holds. Notice that

$$X_n^- = \frac{\beta}{1 - \alpha} - \frac{\beta}{1 - \alpha} \alpha^{-n} + \gamma^{-1} n^{-1} \alpha^{-2n} + o(n^{-1} \alpha^{-2n}),$$

when  $n$  becomes large. Therefore,

$$\begin{aligned} f^n(X_n^-) &\sim n^{-1} \alpha^{-2n} \\ A(X_n^-)/B(X_n^-) &\sim \alpha^{-nm} \end{aligned}$$

for some integer  $m$  when  $n$  becomes large. These contradicts the assumption that Equation (3.20) has solutions for infinitely many  $n \in \mathbb{Z}^+$ .

Now suppose  $|\alpha| < 1$  and there are infinitely many  $n \in \mathbb{Z}^+$  such that Equation (3.20) holds. Now, similarly, since

$$(3.21) \quad t_n^2 - 4\alpha^n n \gamma \beta \frac{1 - \alpha^n}{1 - \alpha} = \frac{\gamma^2\beta^2}{(1 - \alpha)^2} n^2 - 2\frac{\beta\gamma}{1 - \alpha} n + 1 - 2\frac{\gamma^2\beta^2}{(1 - \alpha)^2} n^2 \alpha^n - 2\alpha^n + o(\alpha^n)$$

when  $n$  is large, we have

$$X_n^\pm = \frac{\left( \frac{\gamma\beta}{1 - \alpha} n\alpha^n - \alpha^n - \frac{\beta\gamma}{1 - \alpha} n + 1 \pm \left( \frac{\beta\gamma}{1 - \alpha} n - 1 - \frac{\gamma\beta}{1 - \alpha} n\alpha^n - \alpha^n + o(\alpha^n) \right) \right)}{2n\alpha^n\gamma}.$$

So,

$$X_n^- = -\frac{\beta}{1-\alpha}\alpha^{-n} + \frac{1}{\gamma n \alpha^n} + o(n^{-1}\alpha^{-n}),$$

and we have

$$A(X_n^-)/B(X_n^-) \sim \alpha^{kn}$$

for some integer  $k$  when  $n$  is large. However,

$$\alpha^n X_n^- + \frac{1-\alpha^n}{1-\alpha}\beta \sim \frac{1}{n}$$

contradicting that the Equation (3.20) has solution for infinitely many  $n \in \mathbb{Z}^+$ .

Suppose  $|\alpha| < 1$  and there are infinitely many  $n \in \mathbb{Z}^+$  such that Equation (3.17) holds. Then  $X_n^+ \sim \frac{1}{n}$  when  $n$  is large. Thus,

$$A(X_n^+)/B(X_n^+) \sim n^k$$

for some integer  $k$  and

$$\alpha^n X_n^+ + \frac{1-\alpha^n}{1-\alpha}\beta = \frac{\beta}{1-\alpha} - \frac{\beta}{1-\alpha}\alpha^n + o(\alpha^n)$$

when  $n$  is large. If  $k \neq 0$ , then these contradict the assumption that the Equation (3.17) has solutions for infinitely many  $n \in \mathbb{Z}^+$ . If  $k = 0$ , then

$$A(X_n^+)/B(X_n^+) = A(0)/B(0) + O(1/n^m)$$

for some positive integer  $m$ . Still, these contradicts the assumption that the Equation (3.17) has solution for infinitely many  $n \in \mathbb{Z}^+$  by comparing the rates they approach constants when  $n$  is large.

Now, suppose  $|\alpha| = 1$  but not a root of unity and there are infinitely many  $n \in \mathbb{Z}^+$  such that Equation (3.20) holds. After eliminating denominators, combining terms and taking squares, we found that this is equivalent to that there exists a polynomial  $F(Z, W)$  in two variables such that  $F(n, \alpha^n) = 0$  for infinitely many  $n \in \mathbb{Z}^+$ . By [BGT, Proposition 2.5.1.4], we have  $\{a_n = F(n, \alpha^n)\}_{n \in \mathbb{N}}$  satisfies a linear recurrence. Then, by [BGT, Theorem 2.5.4.1], the set  $\{n \in \mathbb{Z}^+ : a_n = 0\}$  is a finite union of arithmetic progressions and, since we assumed this set is infinite, we have that there exists  $s_1$  and  $s_2 \in \mathbb{Z}^+$  such that  $a_n = 0$  for every  $n \in \{s_1 m + s_2 : m \in \mathbb{Z}^+\}$ . This is equivalent to say that Equation (3.20) holds for any  $n \in \{s_1 m + s_2 : m \in \mathbb{Z}^+\}$ . Then, we consider  $n$  within an infinite subsequence  $I$  of  $\{s_1 m + s_2 : m \in \mathbb{Z}^+\}$  such that  $|1 - \alpha^n|$  is bounded from below independently from  $n$  and not converging. Thus, we have

$$(3.22) \quad X_n^+ = -\frac{1}{n\gamma} + o\left(\frac{1}{n}\right),$$

when  $n \in I$  becomes large. Then Equation (3.20) implies

$$(3.23) \quad c\left(-\frac{1}{n\gamma} + o\left(\frac{1}{n}\right)\right) = \alpha^n\left(-\frac{1}{n\gamma} + o\left(\frac{1}{n}\right)\right) + \frac{1-\alpha^n}{1-\alpha}\beta.$$

However,

$$(3.24) \quad c\left(-\frac{1}{n\gamma} + o\left(\frac{1}{n}\right)\right) \sim (1/n)^k$$

for some integer  $k$  and

$$(3.25) \quad \alpha^n \left( -\frac{1}{n\gamma} + o\left(\frac{1}{n}\right) \right) + \frac{1 - \alpha^n}{1 - \alpha} \beta \sim (1 - \alpha^n)$$

when  $n$  is large. This gives a contradiction. Notice that if  $k = 0$ , then  $c(X_n^+)$  converges to a constant when  $n$  is large. But

$$\alpha^n \left( -\frac{1}{n\gamma} + o\left(\frac{1}{n}\right) \right) + \frac{1 - \alpha^n}{1 - \alpha} \beta$$

is not converging as  $1 - \alpha^n$  doesn't converge.

Now, suppose  $|\alpha| = 1$  but not a root of unity and there are infinitely many  $n \in \mathbb{Z}^+$  such that Equation (3.17) holds. Similarly as above, there exists  $s_3, s_4 \in \mathbb{Z}^+$  such that Equation (3.17) holds for any  $n \in \{s_3m + s_4 | m \in \mathbb{Z}^+\}$ . We consider  $n$  within an infinite subsequence of  $\{s_3m + s_4 | m \in \mathbb{Z}^+\}$  such that  $|1 - \alpha^n|$  is bounded from below independent from  $n$  and not converging. Thus, we have

$$(3.26) \quad X_n^- = -\frac{\beta}{1 - \alpha} \frac{1 - \alpha^n}{\alpha^n} + \frac{1}{n\alpha^n\gamma} + o\left(\frac{1}{n}\right).$$

Therefore, Equation (3.17) implies that

$$(3.27) \quad c(X_n^-) = \frac{1}{n\gamma} + o\left(\frac{1}{n}\right).$$

However, for a non-zero rational function  $c(X)$ ,

$$(3.28) \quad c\left(-\frac{\beta}{1 - \alpha} \frac{1 - \alpha^n}{\alpha^n} + \frac{1}{n\alpha^n\gamma} + o\left(\frac{1}{n}\right)\right)$$

cannot converge to 0 when  $n$  becomes large since  $1 - \alpha^n$  is not converging. This gives a contradiction.  $\square$

*proof of Lemma 3.4.* Suppose  $f(X)$ ,  $g(X)$ ,  $c(X)$  are defined over  $\mathbb{C}$ . If the condition in the statement of Lemma 3.2 holds, then we can use the specialization argument as in the proof of [HT17, Theorem 2] to show that it is enough to only consider the algebraic case. Suppose Lemma 3.2 holds when everything is defined over  $\overline{\mathbb{Q}}$ . If  $\alpha$ ,  $\delta$ ,  $\gamma$  and  $\beta$  are all in  $\overline{\mathbb{Q}}$ , then it is done by assumption. Also, suppose  $\alpha/\delta$  is not a root of unity, and none of  $\alpha$  and  $\delta$  is a root of unity other than 1. If one of  $\alpha$  or  $\delta$  is 1, then the argument in Lemma 3.3 can be applied directly to prove that Equation 3.29 admit solutions for only finitely many  $n \in \mathbb{Z}^+$ , which is a contradiction. So we assume both  $\alpha$  and  $\beta$  are not roots of unity. We further assume that  $\gamma \neq 0$ ,  $\beta \neq 0$ ,  $\gamma\beta/((1 - \alpha)(1 - \gamma)) \neq 1$ , since otherwise it can be seen directly that  $f(X)$  and  $g(X)$  share a common fixed point. Let  $R$  be the ring generated over  $\mathbb{Z}$  by  $\alpha$ ,  $\delta$ ,  $\beta$ ,  $\gamma$  and coefficients of  $c(X)$ . Take  $K = \text{Frac}(R)$ .

We do induction on the transcendence degree,  $n$ , of  $K$ . The base case is that  $n = 0$ , which means everything is algebraic and then we showed from the discussion above that Equation (3.29) can only have solution for finitely many  $n \in \mathbb{Z}^+$ . Now, assume that if the transcendence degree is  $n - 1$ , then the above assumptions on  $\gamma$ ,  $\beta$ ,  $\alpha$ ,  $\delta$ ,  $\alpha/\delta$  and  $\beta\gamma/((1 - \alpha)(1 - \delta))$  implies that the equation only admits solutions for finitely many  $n \in \mathbb{N}^+$ . Take  $L$  as a subfield of  $K$  of transcendence degree  $n - 1$ . Applying [CS93, Theorem

4.1], we have that for all specialization  $s : R \rightarrow \bar{L}$  of sufficiently large height, we have  $\hat{h}_{\bar{L},x^2}(\alpha_s)$ ,  $\hat{h}_{\bar{L},x^2}(\delta_s)$ ,  $\hat{h}_{\bar{L},x^2}(\alpha_s/\delta_s)$  and  $\hat{h}_{\bar{L},x^2}(\beta_s\gamma_s/((1-\alpha_s)(1-\delta_s)))$  are all greater than 0 and  $\hat{h}_{\bar{L},x^2}(\gamma_s)$  and  $\hat{h}_{\bar{L},x^2}(\beta_s)$  are greater than 0 if  $\beta$  and  $\gamma$  are not roots of unity. This implies, in particular, that none of  $\alpha_s$ ,  $\delta_s$ ,  $\alpha_s/\delta_s$  and  $\beta_s\gamma_s/((1-\alpha_s)(1-\delta_s))$  is a root of unity or zero and  $\beta_s$  and  $\gamma_s$  are nonzero.

Let  $f_s(X) = \alpha_s X + \beta_s$ ,  $g_s(X) = X/(\gamma_s X + \delta_s)$  and  $c_s(X)$  denote the rational functions obtained by specializing all the coefficients by  $s$ . Then, the inductive hypothesis implies that

$$f_s^n(X) = g_s^n(X) = c_s(X)$$

has solution for only finitely many  $n \in \mathbb{Z}^+$ . Thus,

$$f^n(X) = g^n(X) = c(X)$$

has solution for only finitely many  $n \in \mathbb{Z}^+$ . Therefore, we complete the inductive step and this concludes the proof.  $\square$

Now we can prove Theorem 1.3 which is equivalent to the following as the contrapositive statement:

**Theorem 3.5.** *Let  $f(X)$ ,  $g(X)$  be automorphisms on  $\mathbb{P}^1$  defined over  $\mathbb{C}$  and  $c(X)$  be a rational function defined over  $\mathbb{C}$ . If there are infinitely many  $n \in \mathbb{Z}^+$  such that*

$$(3.29) \quad c(X) = f^n(X) = g^n(X)$$

*has solutions, and  $f(X)$ ,  $g(X)$  are not simultaneously conjugated by an automorphism on  $\mathbb{P}^1(\mathbb{C})$  to one of the following:*

- (1)  $\alpha X + \beta$ ,  $X/(\gamma X + \delta)$ , with some  $\alpha, \delta, \gamma, \beta \in \mathbb{C}^*$  such that  $\alpha$  and  $\delta$  are not roots of unity other than 1 and  $\alpha/\delta$  is a root of unity;
- (2)  $\alpha X + \beta$ ,  $\delta X + \gamma$ , with some  $\alpha, \delta \in \mathbb{C}^*$  such that  $\alpha$  and  $\delta$  are not roots of unity,  $\gamma$  and  $\beta$  are not both 0 and  $\alpha/\delta$  is a root of unity other than 1;

*then the semigroup generated by  $f$  and  $g$  under composition is not free.*

Before we start the proof, we gather here three counter-examples in Remark (3.6), (3.7) and (3.8) corresponding to the two exceptional cases mentioned in the statement of Theorem 3.5. Therefore, Theorem 3.5 is the best we can expect.

**Remark 3.6.** Suppose  $f(X) = X + \beta$ ,  $g(X) = X/(\gamma X + 1)$  and  $c(X) = -\beta/(\gamma X)$ , with some  $\beta, \gamma \in \mathbb{C}^*$ , we have that there are infinitely many  $n \in \mathbb{Z}^+$  such that Equation (3.29) has solutions. One of the solutions is given by

$$x_n = \left( -n\beta\gamma + \sqrt{n^2\beta^2\gamma^2 - 4\beta\gamma} \right) / (2\gamma).$$

However,  $f(X)$  and  $g(X)$  may still generate a free semigroup. For example, when  $f(X) = X + 2$  and  $g(X) = X/(2X + 1)$ , we denote

$$V_1 = \{x \in \mathbb{R}^+ : x > 1\},$$

$$V_2 = \{x \in \mathbb{R}^+ : x < 1\}.$$



Then we have

$$\begin{aligned} f(V_1) &\subseteq V_1, \\ f(V_2) &\subseteq V_1, \\ g(V_2) &\subseteq V_2, \\ g(V_1) &\subseteq V_2. \end{aligned}$$

Thus, the Ping-pong Lemma 2.8 implies  $\langle f, g \rangle$  is a free semigroup under composition.

**Remark 3.7.** Suppose  $f(X) = \alpha X + \beta$  and  $g(X) = X/(\gamma X + \alpha)$ , where  $\gamma$  and  $\beta$  are non-zero and  $\beta/(1 - \alpha) \neq (1 - \alpha)/\gamma$ . Denote

$$\begin{aligned} K_1 &= \frac{\beta}{1 - \alpha} + \frac{1 - \alpha}{\gamma}, \\ K_2 &= \frac{\beta}{1 - \alpha} - \frac{1 - \alpha}{\gamma}, \\ K_3 &= \frac{\beta}{\gamma}. \end{aligned}$$

Let

$$c(X) = \frac{K_2 X^2}{-2K_3 - 2X^2 + K_1 X} + \frac{\beta}{1 - \alpha} - \frac{\beta}{1 - \alpha} \frac{K_2 X}{-2K_3 - 2X^2 + K_1 X}.$$

One can directly verify that

$$c(X) = f^n(X) = g^n(X)$$

has

$$X_n = \frac{-K_1 + K_2 \alpha^{-n} - \sqrt{(-K_1 + K_2 \alpha^{-n})^2 - 4}}{2}$$

as a solution for each  $n \in \mathbb{Z}^+$ . Further more, suppose  $\gamma = 1$ ,  $\beta = 1$  and  $\alpha = 2$  for example. Then, similarly as in Remark (3.6), one can use the Ping-pong Lemma 2.8 to show that  $\langle f(X), g(X) \rangle$  is a free semigroup.

**Remark 3.8.** Suppose  $f(X) = \alpha X + \beta$  and  $g(X) = \delta X + \gamma$ , where  $\alpha$  and  $d$  are not roots of unity but  $\xi = \delta/\alpha$  is a root of unity of order  $l > 1$ . For each  $i \in \{1, 2, \dots, l - 1\}$ , denote

$$\begin{aligned} K_{1,i} &= \frac{1}{1 - \xi^i} \left( \frac{\gamma}{1 - \delta} - \frac{\beta}{1 - \alpha} \right), \\ K_{2,i} &= \frac{1}{1 - \xi^i} \left( \frac{\xi^i \gamma}{1 - \delta} - \frac{\beta}{1 - \alpha} \right). \end{aligned}$$

Let

$$c_i(X) = K_{1,i} + \frac{\beta}{1 - \alpha} - \frac{K_{1,i}}{X + K_{2,i}} \left( K_{2,i} + \frac{\beta}{1 - \alpha} \right).$$

We have that

$$X_{n,i} = K_{1,i} \alpha^{-nl-i} - K_{2,i}$$

is a solution to

$$f^{nl+i}(X) = g^{nl+i}(X) = c_i(X)$$

for every  $n \in \mathbb{Z}^+$  and  $i \in \{1, 2, \dots, l - 1\}$ . For an explicit example, if we take  $f(X) = 2x + 1$  and  $g(X) = -2x$  then  $\langle f, g \rangle$  is a free semigroup by [KT21, Theorem 2] as  $f \circ g \neq g \circ f$ . So this provides a counter-example such that

the finiteness of solutions to  $f^n(X) = g^n(X) = c(X)$  doesn't hold even when  $f$  and  $g$  generate a free semigroup.

*Proof of Theorem 3.5.* Let's first suppose that  $f$  and  $g$  don't share a fixed point. After conjugation, we assume that

$$\begin{aligned} f(X) &= \alpha X + \beta, \\ g(X) &= \frac{X}{\gamma X + \delta}. \end{aligned}$$

We always assume that  $\alpha$  and  $\delta$  are non-zero as otherwise it will result in a constant map.

Suppose that least one of  $\alpha$  and  $\delta$  is a root of unity other than 1. Assume without loss of generality it is  $\delta$ . Then there exists a positive integer  $m$  such that  $\delta^m = 1$  and

$$g^m(X) = \frac{X}{\delta^m} = X.$$

Therefore,  $\langle g^m(X), f(X) \rangle$  is not a free semigroup under composition and there is nothing to show.

Now, we assume that none of  $\alpha$  and  $\delta$  are roots of unity other than 1. By Lemma 3.2, we can assume that  $f$  and  $g$  share at least one common fixed point. Now we first assume that  $c(X)$ ,  $f(X)$  and  $g(X)$  are all defined over  $\overline{\mathbb{Q}}^*$ .

If  $f$  and  $g$  share two fixed points, then after conjugation by an automorphism, they are both scaling maps and certainly not generating a free semigroup under composition.

If  $f$  and  $g$  share only one common fixed point. Then after conjugating  $c(X)$ ,  $f(X)$  and  $g(X)$  by a common automorphism, we assume that  $f(X) = \alpha X + \beta$  and  $g(X) = \delta X + \gamma$ , where  $\alpha, \delta \in \overline{\mathbb{Q}}^*$ ,  $\beta, \gamma \in \overline{\mathbb{Q}}$  and  $\beta, \gamma$  are not both 0. Now our discussion is divided into several cases:

**Case I.** Suppose at least one of  $\delta$  and  $\alpha$  is a root of unity other than 1. Without loss of generality we assume it is  $\delta$ . Then there exists  $m$  such that  $g^m(x) = x$  and we have  $g^m \circ f = f \circ g^m$ . Thus,  $\langle f, g \rangle$  is not a free semigroup. Same argument works if  $\alpha$  is a root of unity other than 1 and  $\delta$  is not a root of unity.

**Case II.** Suppose  $\delta = 1$  and  $\alpha$  is not a root of unity.

**Lemma 3.9.** *There are only finitely many  $n \in \mathbb{Z}^+$  such that  $c(X) = f^n(X) = g^n(X)$  has solution.*

This concludes this case.

**Case III.** Suppose both  $\alpha$  and  $\delta$  are not roots of unity and  $\alpha/\delta$  is not a root of unity. We have

$$X_n = \left( \frac{1 - \delta^n}{1 - \delta} \gamma - \frac{1 - \alpha^n}{1 - \alpha} \beta \right) / (\alpha^n - \delta^n)$$

is the solution to

$$f^n(X) = g^n(X).$$

If there are infinitely many  $n \in \mathbb{Z}^+$  such that the Equation (3.29) has solutions, then

$$c(X_n) = f^n(X_n)$$

for infinitely many  $n \in \mathbb{Z}^+$ . It means that

$$A(X_n) = \alpha^n B(X_n) X_n + \frac{1 - \alpha^n}{1 - \alpha} \beta B(X_n)$$

holds for infinitely many  $n \in \mathbb{Z}^+$ . This is equivalent to say  
(3.30)

$$F(Z, W) = A(X(Z, W)) - ZB(X(Z, W))X(Z, W) - \frac{1 - Z}{1 - \alpha} \beta B(X(Z, W)) = 0$$

holds for infinitely many  $\{(\alpha^n, \delta^n) : n \in \mathbb{Z}^+\}$ , where

$$X(Z, W) = \left( \frac{1 - W}{1 - \delta} \gamma - \frac{1 - Z}{1 - \alpha} \beta \right) / (Z - W).$$

We first claim that the rational function  $F(Z, W)$  is not constantly zero. To prove this, we suppose on the contrary that Equation (3.30) holds constantly for any choice of values of  $Z$  and  $W$ . Let's denote  $m_1 = \deg(A)$  and  $m_2 = \deg(B)$ . Then choose a specialization  $Z = W + 1$ , we have

$$X(Z, Z - 1) = \left( \frac{\beta}{1 - \alpha} - \frac{\gamma}{1 - \delta} \right) Z + \frac{2\gamma}{1 - \delta} - \frac{\beta}{1 - \alpha}$$

in this case. Then Equation (3.30) holds constantly implies that  $m_1 = m_2 + 2$  unless

$$(3.31) \quad \frac{\beta}{1 - \alpha} = \frac{\gamma}{1 - \delta}.$$

While Equation (3.31) implies  $f(X)$  and  $g(X)$  share two fixed points and this case has already been handled. So, we assume Equation (3.31) doesn't hold and thus Equation (3.30) doesn't hold constantly for every values of  $Z$  and  $W$ .

Let's then write

$$F(Z, W) = \sum_{0 \leq i \leq k_1, 0 \leq j \leq k_2} c_{i,j} Z^i W^j,$$

for some positive integers  $k_1$  and  $k_2$  after clearing the denominators. Then we use the same argument as in Lemma 3.2 by applying S-unit Theorem to conclude that  $\alpha$  and  $\delta$  are multiplicatively dependent. Thus, there exists  $k \in \mathbb{Q}^*$  such that  $\xi \alpha^k = \delta$ , for some root of unity  $\xi$  of order  $k_3 \in \mathbb{Z}^+$ .

Therefore, we have

$$X_n = \left( \frac{1 - \xi^n \alpha^{nk}}{1 - \xi \alpha^k} \gamma - \frac{1 - \alpha^n}{1 - \alpha} \beta \right) / (\alpha^n - \xi^n \alpha^{nk}).$$

Then following the same idea as how we argue in the proof of Lemma 3.2 (Case **II**, Step **III**), it is enough to show the claim that

$$(3.32) \quad A(X_i(Z)) = (Z X_i(Z) + \frac{1 - Z}{1 - \alpha} \beta) B(X_i(Z))$$

doesn't hold constantly for any  $i \in \{1, 2, \dots, k_3\}$  unless  $f$  and  $g$  share two common fixed points, where

$$X_i(Z) = \left( \frac{1 - \xi^i Z^k}{1 - \xi \alpha^k} \gamma - \frac{1 - Z}{1 - \alpha} \beta \right) / (Z - \xi^i Z^k).$$

This is because, similarly as in Lemma 3.2, Equation (3.32) is a equation on one variable and having infinitely many roots implies it is constantly zero.

To prove the claim, we can assume that  $k \neq 0$  since then  $\delta$  is a root of unity and we back to the case that we have discussed. We first suppose  $k > 0$  and suppose the contrary that there exists a

$$i \in \{1, 2, \dots, k_3\}$$

such that Equation (3.32) holds constantly. Notice that in the beginning of Case **III** we assume that  $\alpha/\delta$  is not a root of unity, which is equivalent to  $k \neq 1$ . Then if  $f$  and  $g$  only share one common fixed point, then

$$\frac{\gamma}{1 - \delta} \neq \frac{\beta}{1 - \alpha}$$

and

$$\begin{aligned} \text{val}_Z(X_i(Z)) &= -\min\{1, k\}, \\ \text{val}_Z(A(X_i(Z))) &= -m_1 \min\{1, k\}, \\ \text{val}_Z\left(\left(ZX_i(Z) + \frac{1 - Z}{1 - \alpha} \beta\right)B(X_i(Z))\right) &\geq 1 - (m_2 + 1) \min\{1, k\}. \end{aligned}$$

Suppose, Equation (3.32) holds constantly for every values of  $Z$ , then

$$m_2 + 1 > m_1.$$

However, by Proposition 2.6, there exists a finite set of places  $S$ , and a place  $v \notin S$ , such that

$$|X_i(\alpha^n)|_v > 1$$

for some  $n \in \mathbb{Z}^+$  such that  $X_i(\alpha^n)$  is a solution to the equation  $f^n(X) = g^n(X) = C(X)$  and in particular a solution to Equation (3.32). Then, if  $m_2 + 1 > m_1$ , we have

$$|A(X_i(\alpha^n))|_v = |X_i(\alpha^n)|_v^{m_1} < |(\alpha^n X_i(\alpha^n) + \frac{1 - \alpha^n}{1 - \alpha} \beta)B(X_i(\alpha^n))|_v = |X_i(\alpha^n)|_v^{m_2 + 1},$$

which is a contradiction to Equation (3.32). Thus, the claim is proved for  $k > 0$ .

Now suppose  $k < 0$ . Then we have

$$(3.33) \quad X_i(Z) = \frac{\gamma}{1 - \delta} + o(1)$$

when  $Z$  is closed to 0 and

$$(3.34) \quad A(X_i(Z)) = \left( ZX_i(Z) + \frac{1 - Z}{1 - \alpha} \beta \right) B(X_i(Z)),$$

$$(3.35) \quad A(X_i(Z)) = \left( ZX_i(Z) + \frac{1 - Z}{1 - \delta} \gamma \right) B(X_i(Z)),$$

hold constantly imply that

$$A\left(\frac{\gamma}{1 - \delta}\right) = \frac{\beta}{1 - \alpha} B\left(\frac{\gamma}{1 - \delta}\right),$$

$$A\left(\frac{\gamma}{1-\delta}\right) = \frac{\gamma}{1-\delta} B\left(\frac{\gamma}{1-\delta}\right).$$

By the fact that  $A(X)$  and  $B(X)$  don't share common root, we have

$$\frac{\gamma}{1-\delta} = \frac{\beta}{1-\alpha},$$

which implies that  $f$  and  $g$  share two common fixed points.

**Case IV.** Lastly, suppose  $\alpha = \delta$ , then  $f^n(X) = g^n(X)$  only has a solution if  $f = g$  and certainly  $\langle f, g \rangle$  is not free in this case.

Now, consider the general case that  $f(X)$ ,  $g(X)$  and  $c(X)$  are all defined over  $\mathbb{C}$ . We use the same specialization argument as in Lemma 3.2. We assume  $\alpha$ ,  $\delta$  and  $\alpha/\delta$  are not roots of unity and  $\gamma/(1-\delta) - \beta/(1-\alpha) \neq 0$ , since otherwise the argument above directly shows that  $\langle f, g \rangle$  is not free. We construct a ring extension  $R$  over  $\mathbb{Z}$  generated by coefficients of  $f(X)$ ,  $g(X)$  and  $c(X)$  and apply [CS93, Theorem 4.1] and do the induction on the transcendental degree of  $\text{Frac}(R)$  over  $\mathbb{Q}$  to get a specialization  $s$  with height large enough so that  $\alpha_s$ ,  $\delta_s$  and  $\alpha_s/\delta_s$  are not roots of unity and  $\gamma_s/(1-\delta_s) - \beta_s/(1-\alpha_s) \neq 0$ . Then the above argument shows that

$$f_s^n(X) = g_s^n(X) = c_s(X)$$

has solution for only finitely many  $n \in \mathbb{Z}^+$  and thus

$$f^n(X) = g^n(X) = c(X)$$

has solution for only finitely many  $n \in \mathbb{Z}^+$ . □

*Proof of Lemma 3.9.* We have that

$$f^n(X) = \alpha^n X + \frac{1-\alpha^n}{1-\alpha} \beta$$

$$g^n(X) = X + n\gamma.$$

If  $\gamma = 0$ , then  $\langle f, g \rangle$  is obviously not free, so we assume that  $\gamma \neq 0$ . Then

$$X_n = \frac{\beta}{1-\alpha} - \frac{n\gamma}{1-\alpha^n}$$

is the solution to

$$f^n(X_n) = g^n(X_n).$$

Denote  $c(X) = A(X)/B(X)$ , for some polynomials  $A(X)$  and  $B(X)$ . Now suppose the equation

$$(3.36) \quad A(X_n) = B(X_n) \left( \alpha^n X_n + \frac{1-\alpha^n}{1-\alpha} \beta \right)$$

is satisfied by infinitely many  $n \in \mathbb{Z}^+$ . If  $|\alpha| > 1$ , then

$$A(X_n)/B(X_n) \sim (n/\alpha^n)^k$$

for some integer  $k$  when  $n$  is large. However,

$$\left( \alpha^n X_n + \frac{1-\alpha^n}{1-\alpha} \beta \right) \sim \gamma n$$

when  $n$  is large, which contradicts the assumption that the Equation (3.36) holds for infinitely many  $n \in \mathbb{Z}^+$ .

On the other hand, suppose  $|\alpha| < 1$ . Now, take a set of places  $S$  such that for any  $v \notin S$ , we have  $A(X)$  and  $B(X)$  has coefficients with norm not greater than 1 with respect to  $v$  and also

$$|1 - \alpha|_v = |\alpha|_v = |\beta|_v = |\gamma|_v = 1,$$

if  $\beta \neq 0$ . Then, by the proof of [HT17, Proposition 15, **Case II**], when  $n$  is large enough, there is a  $v$  outside of  $S$  such that

$$|n/(1 - \alpha^n)|_v > 1.$$

Thus, the Equation (3.36) implies that

$$(3.37) \quad \deg(A) = \deg(B) + 1$$

since  $|X_n|_v > 1$  with the choice of  $n$  and  $v$  such that  $|n/(1 - \alpha^n)|_v > 1$ . Now, since

$$\lim_{n \rightarrow \infty} |\alpha^n X_n| = 0,$$

we have

$$\lim_{n \rightarrow \infty} |A(X_n)| = \frac{\beta}{1 - \alpha} \lim_{n \rightarrow \infty} |B(X_n)|$$

by Equation (3.36). This together with

$$\lim_{n \rightarrow \infty} |X_n| = \infty,$$

imply that

$$\deg(A) = \deg(B)$$

which contradicts Equation (3.37).

Now, suppose  $|\alpha| = 1$ . By the same argument from the last case, we have  $\deg(A) = \deg(B) + 1$ . Let

$$F(n, \alpha^n) = (A(X_n) - B(X_n)(\alpha^n X_n + \frac{1 - \alpha^n}{1 - \alpha} \beta))(1 - \alpha^n)^{\deg(A)}$$

be a polynomial in  $n$  and  $\alpha^n$ . Then by [BGT, Proposition 2.5.1.4] we have  $\{F(n, \alpha^n)\}_{n \in \mathbb{Z}^+}$  satisfies a linear recurrence. Thus, by [BGT, Theorem 2.5.4.1], the set  $\{n \in \mathbb{Z}^+ : F(n, \alpha^n) = 0\}$  is a finite union of arithmetic progressions. Since we assumed that  $F(n, \alpha^n) = 0$  holds for infinitely many  $n$ , there exists  $s_1, s_2 \in \mathbb{Z}^+$  such that  $F(n, \alpha^n) = 0$  for any  $n \in \{s_1 m + s_2 : m \in \mathbb{Z}^+\}$ . Now, we consider  $n$  within an infinite subsequence  $I$  of  $\{s_1 m + s_2 : m \in \mathbb{Z}^+\}$  such that  $|1 - \alpha^n|$  is bounded from below independent from  $n$  and not converging. Thus

$$\lim_{n \rightarrow \infty, n \in I} F(n, \alpha^n) = 0,$$

where  $n$  ranges in the chosen subsequence  $I$ , implies that

$$\lim_{n \rightarrow \infty, n \in I} \alpha^n = a/b$$

where  $a$  and  $b$  are the leading coefficients of  $A(X)$  and  $B(X)$  respectively. This gives a contradiction as  $1 - \alpha^n$  doesn't converge.  $\square$

## 4. PROOF OF THEOREM 1.4

The following lemma asserts that an infinite sequence  $\{x_n\}$  of solutions to the equation  $f^n(x_n) = c(x_n)$  for all  $n \in \mathbb{N}$  is a sequence of small dynamical height.

**Lemma 4.1.** *Let  $K$  be a global field. Let  $\{x_n\}$  be a sequence in  $\mathbb{P}^1(\overline{K})$  so that  $f^n(x_n) = c(x_n)$  for all  $n \in \mathbb{N}$  where  $f, c \in K(x)$  and  $\deg(f) > 1$ . Then*

$$\lim_{n \rightarrow \infty} \hat{h}_f(x_n) = 0.$$

*Proof.* For each  $x \in \mathbb{P}^1(\overline{K})$ , we have by Theorem 2.1 that

$$\hat{h}_f(c(x)) = h(c(x)) + O(1) = (\deg c)h(x) + O(1) = (\deg c)\hat{h}_f(x) + O(1)$$

where the implied constants do not depend on  $x$  (but coefficients of  $f$ ). For each  $n \in \mathbb{N}$ , the sequence  $\{x_n\}_{n \in \mathbb{N}}$  satisfies  $f^n(x_n) = c(x_n)$  and the functorial property of canonical height yields

$$\hat{h}_f(c(x_n)) = \hat{h}_f(f^n(x_n)) = (\deg f)^n \hat{h}_f(x_n).$$

This implies

$$(\deg f)^n \hat{h}_f(x_n) = (\deg c)\hat{h}_f(x_n) + O(1)$$

where the implied constant does not depend on  $x_n$  and so does not depend on  $n$ . Thus the quantity  $((\deg f)^n - \deg c)\hat{h}_f(x_n)$  is bounded by a constant independent of  $n$ . As the term  $((\deg f)^n - \deg c)$  becomes arbitrary large as  $n$  gets large because  $\deg f > 1$ , it follows that  $\hat{h}_f(x_n)$  must become arbitrary small and approaches to 0 as  $n \rightarrow +\infty$ .  $\square$

The following result is well-known in arithmetic dynamics and it asserts that if two rational maps share the same (non-repeating) sequence of points of small height, their dynamical height must be identical. The strategy of the proof is similar to that of Baker-DeMarco [BD11, Theorem 1.2] but we include the proof here for completeness and for the convenience of the reader. For different proof (in the language of metrics on line bundle of  $\mathbb{P}^1$ ) and ingredient we refer the reader to [PST12, Corollary 14].

**Proposition 4.2.** *Let  $K$  be a global field and let  $f, g \in K(x)$  be rational maps of degree greater than 1. Suppose that there exists an infinite distinct sequence  $\{x_n\}_{n \in \mathbb{N}} \subset \mathbb{P}^1(\overline{K})$  such that*

$$\lim_{n \rightarrow \infty} \hat{h}_f(x_n) = \lim_{n \rightarrow \infty} \hat{h}_g(x_n) = 0.$$

*Then  $\hat{h}_f = \hat{h}_g$ .*

*Proof.* Suppose that  $\{x_n\}_{n \geq 1}$  is a non-repeating sequence in  $\mathbb{P}^1(\overline{K})$  such that  $\hat{h}_f(x_n) \rightarrow 0$  and  $\hat{h}_g(x_n) \rightarrow 0$  as  $n \rightarrow +\infty$ .

**Case I.** Suppose that  $f, g \in \overline{\mathbb{Q}}(x)$ . Suppose that  $L$  is a number field for which both  $f$  and  $g$  are defined. Thus  $\{x_n\}$  must be a sequence of algebraic points in  $\mathbb{P}^1(\overline{L})$  for all  $n \in \mathbb{N}$ . Since  $\{x_n\}$  is a sequence of point of small dynamical height with respect to both  $f$  and  $g$ , so the sequence of probability measure supported equally on the sets  $\text{Gal}(\overline{L}/L) \cdot x_n$  of conjugates of  $x_n$  converges weakly to the canonical measures  $\mu_{f,v}$  and  $\mu_{g,v}$  at all places

$v$  of  $L$  in the Berkovich projective line  $\mathbb{P}_{\text{Berk},v}^1$  by the arithmetic equidistribution theorem of Baker-Rumely [BR06], Favre-Rivera-Letelier [FRL06], and Chambert-Loir [CL06]. Thus  $\mu_{f,v} = \mu_{g,v}$ . From the definition of the Laplacian (2.1), for all  $v \in M_L$  and all  $x, y \in \mathbb{P}_{\text{Berk},v}^1$ , we have that the  $v$ -adic Arakelov Green's functions of  $f$  and  $g$  must differ by a constant  $C_v(y) \in \mathbb{R}$  ([BR10, Proposition 5.28])

$$(4.1) \quad G_{\mu_{f,v}}(x, y) = G_{\mu_{g,v}}(x, y) + C_v(y).$$

Using the fact that the Arakelov-Green's function is symmetric, fixing  $x$  and viewing as a function of  $y$ , we can deduce that  $C_v(y) = C_v \in \mathbb{R}$  is independent of  $y$ . Thus the equation (4.1) becomes

$$(4.2) \quad G_{\mu_{f,v}}(x, y) = G_{\mu_{g,v}}(x, y) + C_v.$$

Summing equation (4.2) over all places  $v \in M_L$  with appropriate local degree, it follows that

$$\hat{h}_f(x) + \hat{h}_f(y) = \hat{h}_g(x) + \hat{h}_g(y) + C$$

where  $C = \sum_{v \in M_L} N_v C_v$  and is independent of  $y$ . Setting  $y = x_n$  and letting  $n \rightarrow \infty$ , we have

$$\hat{h}_f(x) = \hat{h}_g(x) + C.$$

Taking  $x$  to be  $f$ -preperiodic, we obtain  $C = -\hat{h}_g(x) \leq 0$ . Similarly, taking  $x$  to be  $g$ -preperiodic, it yields  $C = \hat{h}_f(x) \geq 0$ . Hence  $C = 0$  and the desired result follows immediately.

**Case II.** Let  $L$  be a finitely generated field extension of  $\overline{\mathbb{Q}}$  which is generated by all coefficients of  $f$  and  $g$ . Suppose that neither  $f$  nor  $g$  is isotrivial (i.e., they are not conjugate to a rational map defined over  $\overline{\mathbb{Q}}$ ). Thus the sequence  $\{x_n\}_{n \in \mathbb{N}}$  is defined over  $\mathbb{P}^1(\overline{L})$ . Then the proof follows along similar lines to that of Case I. The last step to conclude that  $C = 0$ , we instead use the Theorem 2.3 for function field.

**Case III.** Suppose that at least one of  $f$  and  $g$  is isotrivial. Without loss of generality, assume that  $f$  is isotrivial (i.e., it is conjugate to a rational map defined over  $\overline{\mathbb{Q}}$ ). Then we follow the argument used in Baker-DeMarco [BD11, Theorem 2] to deduce that  $g$  must also be isotrivial. Hence both  $f$  and  $g$  are conjugate to a rational map defined over  $\overline{\mathbb{Q}}$  and we are back in Case I. Therefore, the desired result is established.  $\square$

Recall that

$$\text{Prep}(f) := \{x \in \mathbb{C} \mid f^n(x) = f^m(x), \text{ for } m, n \in \mathbb{N}\}$$

the collection of all preperiodic points of  $f$ . It is infinite and countable [Bea91, §6.2].

**Proposition 4.3.** *Let  $f, g \in \mathbb{C}(x)$  be two rational maps of degree greater than 1. Suppose that the semigroup generated by  $f$  and  $g$  under composition is free. Then there are finitely many  $\lambda \in \mathbb{C}$  such that there are  $m, n \in \mathbb{N}$  with the following properties:*

- (i)  $f^m(x) \neq c(x)$
- (ii)  $g^n(x) \neq c(x)$
- (iii)  $f^m(\lambda) = g^n(\lambda) = c(\lambda)$ .



*Proof.* Let  $K$  be a field generated by all coefficients of  $f, g$ , and  $c$  over  $\mathbb{Q}$ . Thus  $K$  is either a number field or a function field of finite transcendence degree with constant field  $K \cap \overline{\mathbb{Q}}$ .

Suppose, on the contrary, that there is an infinite distinct sequence  $\{\lambda_i\}_{i \in \mathbb{N}}$  such that for every  $m_i, n_i \in \mathbb{N}$  we have  $f^{m_i}(x) \neq c(x), g^{n_i}(x) \neq c(x)$ , and  $\lambda_i$  is solution to both  $f^{m_i}(x) = c(x)$  and  $g^{n_i}(x) = c(x)$ . Note that  $m_i, n_i \rightarrow +\infty$  as  $i \rightarrow +\infty$ . It follows from Lemma 4.1 that

$$\lim_{i \rightarrow \infty} \hat{h}_f(\lambda_i) = \lim_{i \rightarrow \infty} \hat{h}_g(\lambda_i) = 0.$$

It is, then, immediately from Proposition 4.2 that  $\hat{h}_f = \hat{h}_g$ .

**Case I.** Suppose that  $K$  is a number field. Proposition 2.2 asserts that the canonical heights  $\hat{h}_f$  and  $\hat{h}_g$  vanish precisely on the set of  $f$ -preperiodic points and  $g$ -preperiodic points, respectively. Then it follows that  $\text{Prep}(f) = \text{Prep}(g)$  (i.e.,  $f$  and  $g$  share the same set of preperiodic points). Applying the result of Bell-Haung-Peng-Tucker [BHPT, Corollary 4.12] to deduce that the semigroup generated by  $f$  and  $g$  under composition is not free. Hence it contradicts the assumption.

**Case II.** Suppose that  $K$  is a function field of finite transcendence degree and neither  $f$  nor  $g$  is isotrivial. Thus  $f$  and  $g$  must share the same set of preperiodic points by Theorem 2.3. Again, applying the result of [BHPT], we have that  $\langle f, g \rangle$  is not free.

**Case III.** Assume that at least one of  $f$  and  $g$  is isotrivial. Without loss of generality, assume that  $f$  is isotrivial. We claim that  $g$  must also be isotrivial. This follows immediately from a weak Northcott result of Baker [Ba09, Theorem 1.6]. Thus both  $f$  and  $g$  are conjugate to a rational map defined over  $\overline{\mathbb{Q}}$ . We are back to Case I.  $\square$

Next, we treat the case when one of rational maps  $f$  and  $g$  is of degree 1. Here we briefly discuss proof idea due to Hsia-Tucker [HT17] and it goes through for rational maps  $f$  and  $g$  without any changes.

**Proposition 4.4.** *Let  $f, g \in \mathbb{C}(x)$  be two rational maps such that  $\deg(f) > 1$  and  $\deg(g) = 1$ . Suppose that the semigroup generated by  $f$  and  $g$  under composition is free. Then there are finitely many  $\lambda \in \mathbb{C}$  such that there are  $m, n \in \mathbb{N}$  with the following properties:*

- (i)  $f^m(x) \neq c(x)$
- (ii)  $g^n(x) \neq c(x)$
- (iii)  $f^m(\lambda) = g^n(\lambda) = c(\lambda)$ .

*Proof.* Again,  $K$  is a field generated by all coefficients of  $f, g$  and  $c$  over  $\mathbb{Q}$ . Thus  $K$  is either a number field or a function field of finite transcendence degree.

**Case I.** Suppose that  $K$  is a number field. We follow closely the argument of Hsia-Tucker [HT17, Proposition 9]. For any infinite distinct sequence of  $\{\lambda_i\}$  and  $\{m_i\}$  such that  $f^{m_i}(\lambda_i) = c(\lambda_i)$  for every  $i \in \mathbb{N}$ , we have that  $\{\lambda_i\}$  is a sequence of small height with respect to  $f$  by Lemma 4.1. Also, the condition (iii) yields that any solution  $\lambda \in \mathbb{C}$  to the equation  $g^n(\lambda) = c(\lambda)$  is of bounded degree (i.e., degree at most  $\deg(c)$  since  $\deg(g) = 1$ ) over  $K$ . Hence the Northcott's property implies that there are only finitely many  $\lambda \in \mathbb{C}$  satisfying (i), (ii), and (iii).

**Case II.** Suppose that  $K$  is a function field of finite transcendence degree. We refer the reader to Hsia-Tucker [HT17, Proposition 9] as the proof is verbatim to the polynomial maps setting.  $\square$

We are now ready to prove the main result of this section.

**Theorem 4.5.** *Let  $f(x)$  and  $g(x)$  be rational maps in  $\mathbb{C}(x)$ , at least one of which has degree greater than one. Suppose that semigroup generated by  $f$  and  $g$  under composition is free and  $c(x)$  is not a compositional power of  $f$  or  $g$ . Then there are only finitely many  $\lambda \in \mathbb{C}$  such that*

$$f^m(\lambda) = g^n(\lambda) = c(\lambda).$$

for some  $m, n \in \mathbb{N}$ .

*Proof.* The proof divides into two cases:

**Case I.** If  $\deg(f), \deg(g) > 1$ , then the desired result follows immediately from Proposition 4.3.

**Case II.** If one of rational maps  $f$  and  $g$  is of degree one (without loss of generality, assume that  $\deg(f) > 1$  and  $\deg(g) = 1$ ), then the desired conclusion follows from Proposition 4.4.  $\square$

#### ACKNOWLEDGEMENTS

We thank Jason Bell for helpful suggestions and conversations and Nathan Grieve for useful comments concerning the exposition. The second author was supported in part by NSERC grant RGPIN-2022-02951.

#### REFERENCES

- [AR04] N. Ailon and Z. Rudnick. *Torsion points on curves and common divisors of  $a^k - 1$  and  $b^k - 1$* . Acta Arith. **113** (2004), no. 1, 31–38.
- [Art06] E. Artin. *Algebraic numbers and algebraic functions*. (Reprint of the 1967 original). AMS Chelsea Publishing, Providence, RI, 2006.
- [BCZ03] Y. Bugeaud, P. Corvaja, and U. Zannier. *An upper bound for the G.C.D. of  $a^n - 1$  and  $b^n - 1$* . Math. Z. **243** (2003), no. 1, 79–84.
- [Ba09] M. Baker. *A finiteness theorem for canonical heights attached to rational maps over function fields*. J. Reine Angew. Math. **626** (2009), 205–233.
- [BD11] M. Baker and L. DeMarco. *Preperiodic points and unlikely intersections*. Duke Math. J. **159** (2011), no. 1, 1–29.
- [BD13] M. Baker and L. DeMarco. *Special curves and postcritically finite polynomials*. Forum Math. Pi. **1** (2013), e3, 35.
- [Bea91] A. F. Beardon. *Iteration of rational functions*. Graduate Texts in Mathematics, Vol. 132. Springer-Verlag, New York, 1991.
- [Be05] R. L. Benedetto. *Heights and preperiodic points of polynomials over function fields*. Int. Math. Res. Not. (2005), no. 62, 3855–3866.
- [Be19] R. L. Benedetto. *Dynamics in one non-archimedean variable*. Graduate Studies in Mathematics, Vol. 198. American Mathematical Society, Providence, RI, 2019.
- [BE87] I. N. Baker and A. Erëmenko. *A problem on Julia sets*. Ann. Acad. Sci. Fenn. Ser. A I Math. **12** (1987), no. 2, 229–236.
- [BG06] E. Bombieri and Walter Gubler. *Heights in Diophantine geometry*. New Mathematical Monographs, Vol. 4. Cambridge University Press, Cambridge, 2006.
- [BGT] J. Bell, D. Ghioca, and T. Tucker. *The dynamical Mordell-Lang conjecture*. Mathematical Surveys and Monographs, Vol. 210. American Mathematical Society, Providence, RI, 2016.

- [BHPT] J. Bell, K. Haug, W. Peng, and T. Tucker. *A Tits alternative for endomorphisms of the projective line*. J. Eur. Math. Soc. (2023), to appear.
- [BR06] M. H. Baker and R. Rumely. *Equidistribution of small points, rational dynamics, and potential theory*. Ann. Inst. Fourier (Grenoble). **56** (2006), no. 3, 625–688.
- [BR10] M. Baker and R. Rumely. *Potential theory and dynamics on the Berkovich projective line*. Mathematical Surveys and Monographs, Vol. 159. American Mathematical Society, Providence, RI, 2010.
- [CL06] A. Chambert-Loir. *Mesures et équidistribution sur les espaces de Berkovich*. J. Reine Angew. Math. **59** (2006), no. 3, 215–235.
- [CS93] G. S. Call and J. H. Silverman. *Canonical heights on varieties with morphisms*. Compositio Math. **89** (1993), no. 2, 163–205.
- [CZ13] P. Corvaja and U. Zannier. *Greatest common divisors of  $u - 1$ ,  $v - 1$  in positive characteristic and rational points on curves over finite fields*. J. Eur. Math. Soc. (JEMS). **15** (2013), no. 5, 1927–1942.
- [EG15] J.-H. Evertse and K. Györy, *Unit equations in Diophantine number theory*. Cambridge Studies in Advanced Mathematics, Vol. 146. Cambridge University Press, Cambridge, 2015.
- [FRL06] C. Favre and J. Rivera-Letelier. *Équidistribution quantitative des points de petite hauteur sur la droite projective*. Math. Ann. **335** (2006), no. 2, 311–361.
- [GHT13] D. Ghioca, L.-C. Hsia and T. J. Tucker. *Preperiodic points for families of polynomials*. Algebra Number Theory. **7** (2013), no. 3, 701–732.
- [GHT15] D. Ghioca, L.-C. Hsia and T. J. Tucker. *Preperiodic points for families of rational maps*. Proc. Lond. Math. Soc. (3). **110** (2015), no. 2, 395–427.
- [GHT17] D. Ghioca, L.-C. Hsia and T. J. Tucker. *On a variant of the Ailon-Rudnick theorem in finite characteristic*. New York J. Math. **23** (2017), 213–225.
- [GHT18] D. Ghioca, L.-C. Hsia and T. J. Tucker. *A variant of a theorem by Ailon-Rudnick for elliptic curves*. Pacific J. Math. **295** (2018), no. 1, 1–15.
- [Gr20] N. Grieve. *Generalized GCD for toric Fano varieties*. Acta Arith. **195** (2020), no. 4, 415–428.
- [GW20] N. Grieve and J. T.-Y. Wang. *Greatest common divisors with moving targets and consequences for linear recurrence sequences*. Trans. Amer. Math. Soc. **373** (2020), no. 2, 8095–8126.
- [HT17] L.-C. Hsia and T. J. Tucker. *Greatest common divisors of iterates of polynomials*. Algebra Number Theory. **11** (2017), no. 6, 1437–1459.
- [Hu20] K. Huang. *Generalized greatest common divisors for orbits under rational functions*. Monatsh. Math. **191** (2020), no. 1, 103–123.
- [KT21] A. Kolpakov and A. Talambutsa. *On free semigroups of affine maps on the real line*. Proc. Amer. Math. Soc. **150** (2021), no. 6, 2301–2307.
- [La83] S. Lang. *Fundamentals of Diophantine geometry*. Springer-Verlag, New York, 1983.
- [Le19] A. Levin. *Greatest common divisors and Vojta’s conjecture for blowups of algebraic tori*. Invent. Math. **215** (2019), no. 2, 493–533.
- [Lu05] F. Luca. *On the greatest common divisor of  $u - 1$  and  $v - 1$  with  $u$  and  $v$  near  $S$ -units*. Monatsh. Math. **146** (2005), no. 3, 239–256.
- [Os16] A. Ostafe. *On some extensions of the Ailon-Rudnick theorem*. Monatsh. Math. **181** (2016), no. 2, 451–471.
- [PST09] C. Petsche, L. Szpiro and M. Tepper. *Isotriviality is equivalent to potential good reduction for endomorphisms of  $\mathbb{P}^N$  over function fields*. J. Algebra. **322** (2009), no. 9, 3345–3365.
- [PST12] C. Petsche, L. Szpiro and T. J. Tucker. *A dynamical pairing between two rational maps*. Trans. Amer. Math. Soc. **364** (2012), no. 4, 1687–1710.
- [Si04a] J. H. Silverman. *Common divisors of  $a^n - 1$  and  $b^n - 1$  over function fields*. New York J. Math. **10** (2004), 37–43.
- [Si04b] J. H. Silverman. *Common divisors of elliptic divisibility sequences over function fields*. Manuscripta Math. **114** (2004), no. 4, 431–446.

- [Si05] J. H. Silverman. *Generalized greatest common divisors, divisibility sequences, and Vojta's conjecture for blowups*. *Monatsh. Math.* **145** (2005), no. 4, 333–350.
- [Si07] J. H. Silverman. *The arithmetic of dynamical systems*. Graduate Texts in Mathematics, Vol. 241. Springer, New York, 2007.
- [SS95] W. Schmidt and N. Steinmetz. *The polynomials associated with a Julia set*. *Bull. London Math. Soc.* **27** (1995), no. 3, 239–241.
- [Za12] U. Zannier. *Some problems of unlikely intersections in arithmetic and geometry*. (With appendixes by David Masser). *Annals of Mathematics Studies*, Vol. 181. Princeton University Press, Princeton, NJ, 2012.

UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO, CANADA N2L 3G1

*Email address:* `cnoytaptim@uwaterloo.ca`, `chatchai.noytaptim@gmail.com`

UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO, CANADA N2L 3G1

*Email address:* `x48zhong@uwaterloo.ca`