Well Posed Origin Anywhere Consistent Systems in Celestial Mechanics

Harry Gingold and Jocelyn Quaintance

West Virginia University, Morgantown, WV, USA; gingold@math.wvu.edu University of Pennsylvania, Philadelphia, PA, USA; jocelynq@seas.upenn.edu

May 27, 2024

Abstract

Certain measurements in celestial mechanics necessitate having the origin O of a Cartesian coordinate system (CCS) coincide with a point mass. For the two and three body problems we show mathematical inadequacies in Newton's celestial mechanics equations (NCME) when the origin of a coordinate system coincides with a point mass. A certain system of equations of relative differences implied by NCME is free of these inadequacies and is invariant with respect to any CCS translation. A new constant of motion is derived for the relative system. It shows that the universe of relative differences of the N-body problem is "restless".

Keywords: Cartesian coordinate system (CCS); Newton's celestial mechanics equations (NCME); *N*-body problem; body centered origin system (BCOS); well posed anywhere consistent; relative difference; origin invariant; constant of motion.

AMS Classification: Primary 70F15; Secondary 85A04, 70K99.

1 Introduction

Observations and measurements from earth or from other moving bodies viewed theoretically as point masses require the utilization of a Cartesian coordinate system (CCS) in which the origin *O* coincides with the center of mass of one of these point masses. Their acceleration is modeled by Newton's celestial mechanics equations (NCME), namely

$$r_i''(t) = \sum_{j \neq i} \frac{Gm_j(r_j(t) - r_i(t))}{\|r_j(t) - r_i(t)\|^3}, \ 1 \le i \le N,$$
(1.1)

where $r_i(t)$ is the position vector (in \mathbb{R}^3) of the *i*th body (which has mass m_i), $\| \|$ represents the Euclidean norm, and G is the gravitational constant. Ideally we want initial value problems of these systems to possess unique solutions no matter where the origin of the coordinate system is in space. The following theorem, whose proof is found on Pages 1-7 of [2], applies.

Theorem 1.1. Given any CCS, for $1 \le j \le N$, let $r_j(t) \in \mathbb{R}^3$ and $r'_j(t) = dr_j/dt \in \mathbb{R}^3$ be the position and velocity of the *j*th point mass respectively. Assume that the accelerations $r''_i(t)$ satisfy (1.1). Let $1 \le i, j \le i$

N, with $i \neq j$. Then the initial value problem

$$r_j''(t) = \sum_{\substack{i=1\\i\neq j}}^{N} \frac{Gm_i(r_i(t) - r_j(t))}{\|r_i(t) - r_j(t)\|^3}, \qquad r_i(t_0), r_i'(t_0) \in \mathbb{R}^3, \quad r_i(t_0) - r_j(t_0) \neq \overrightarrow{0}, \tag{1.2}$$

possesses unique solutions $r_j(t) \in C^2([a,b])$, a < b, on some interval $[a,b] \subseteq \mathbb{R}$, where $t_0 \in [a,b]$.

Inconsistencies in NCME occur when the origin *O* of a CCS coincides with a point mass. This is discussed in Section 2. We propose in Section 3 to replace NCME by a related system of equations that is free of inconsistencies. In Section 4 we derive a new constant of motion for the relative difference equations. In Section 5 we show by means of Section 4 that our universe is "restless". Conclusions in Section 6 provide an overview of our study.

2 Inconsistencies in Body-Centered Origin Systems (BCOS) for the 2 and 3-Body Problems

Given a two body problem with two point masses m_1 and m_2 , suppose that an origin O of a CCS coincides with the point mass m_1 . Let the 3×1 vectors $r_1(t), r_2(t)$, or r_1, r_2 , denote the position vectors of m_1, m_2 respectively. The NCME are

$$r_1''(t) = \frac{Gm_2(r_2 - r_1)}{\|r_2 - r_1\|^3}, \qquad r_2''(t) = \frac{Gm_1(r_1 - r_2)}{\|r_1 - r_2\|^3},$$
(2.1)

where $||r(t)|| = \sqrt{r^T r}$. Measurements made from the point O require that we set in (2.1)

$$r_1(t) \equiv \overrightarrow{0} \Longrightarrow r'_1(t) \equiv r''_1(t) \equiv \overrightarrow{0}, \qquad (2.2)$$

where $\overrightarrow{0} = [0, 0, 0]^T$. The second equation of (2.1) becomes

$$r_2''(t) = -\frac{Gm_1r_2}{\|r_2\|^3}.$$
(2.3)

But what happens to the first equation of (2.1)? It implies

$$\overrightarrow{0} = \frac{Gm_2r_2}{\|r_2\|^3} \Longrightarrow 0 = \left\|\overrightarrow{0}\right\| = \left\|\frac{Gm_2r_2}{\|r_2\|^3}\right\| = \frac{Gm_2}{\|r_2\|^2} \Longleftrightarrow \|r_2\| = \infty.$$
(2.4)

Evidently, this is a contradiction to a well known solution $r_2(\theta) = \frac{ed}{1+e\cos\theta}$ to (2.3). One may explain the inconstancy by claiming that the origin O is accelerating and try to introduce fictitious forces to avoid this. In contrast one may blame the inconsistency on overdetermination inherent in BCOS. Therefore we subtract the two equations in (2.1) and propose one origin independent relative system

$$(r_1 - r_2)'' = -\frac{G(m_1 + m_2)(r_1 - r_2)}{\|r_1 - r_2\|^3}.$$
(2.5)

Thus, when $r_1(t) \equiv \vec{0}$, i.e. when m_1 corresponds to origin O, Equation (2.5) becomes a modified Kepler problem

$$-r_2''(t) = \frac{G(m_1 + m_2)r_2}{\|r_2\|^3}.$$
(2.6)

Now we analyze a BCOS model for the 3-body problem. From a fixed origin O we record the position of three point masses, m_1, m_2 , and m_3 as the 3×1 vectors $r_1(t), r_2(t)$, and $r_3(t)$ respectively. Then NCME give rise to the following system of three nonlinear second order differential equations:

$$r_1''(t) = \frac{Gm_2(r_2 - r_1)}{\|r_2 - r_1\|^3} + \frac{Gm_3(r_3 - r_1)}{\|r_3 - r_1\|^3}$$
$$r_2''(t) = \frac{Gm_1(r_1 - r_2)}{\|r_1 - r_2\|^3} + \frac{Gm_3(r_3 - r_2)}{\|r_3 - r_2\|^3}$$
$$r_3''(t) = \frac{Gm_1(r_1 - r_3)}{\|r_1 - r_3\|^3} + \frac{Gm_2(r_2 - r_3)}{\|r_2 - r_3\|^3}.$$

Assume m_1 corresponds to O, i.e. $r_1 \equiv \overrightarrow{0}$. Since $r_1 \equiv \overrightarrow{0}$ implies that $r'_1 \equiv r''_1 \equiv \overrightarrow{0}$, the preceding system becomes

$$\overrightarrow{0} = \frac{Gm_2r_2}{\|r_2\|^3} + \frac{Gm_3r_3}{\|r_3\|^3}$$

$$r_2''(t) = -\frac{Gm_1r_2}{\|r_2\|^3} + \frac{Gm_3(r_3 - r_2)}{\|r_3 - r_2\|^3}$$

$$r_3''(t) = -\frac{Gm_1r_3}{\|r_3\|^3} + \frac{Gm_2(r_2 - r_3)}{\|r_2 - r_3\|^3}.$$
(S1)

The first equation of (S1) implies that

$$\frac{m_2 r_2}{\|r_2\|^3} = -\frac{m_3 r_3}{\|r_3\|^3} \iff r_2 = -\frac{m_3 \|r_2\|^3}{m_2 \|r_3\|^3} r_3.$$
(2.7)

If we take the norm of (2.7) we find that

$$||r_2||^{-2} = \frac{m_3}{m_2} ||r_3||^{-2} \iff ||r_2|| = \sqrt{\frac{m_2}{m_3}} ||r_3||.$$
(2.8)

By substituting (2.8) into (2.7) we obtain the relation

$$r_2 = -\sqrt{\frac{m_2}{m_3}} r_3. \tag{2.9}$$

We now place (2.7) and (2.9) into the second equation of (S1) to obtain

$$r_3'' = -G\left(\frac{m_3}{m_2}\right)^{\frac{3}{2}} \left[\frac{m_1(\sqrt{m_3} + \sqrt{m_2})^2 + m_2m_3}{(\sqrt{m_3} + \sqrt{m_2})^2}\right] \frac{r_3}{\|r_3\|^3}.$$
 (2.10)

However, if we substitute (2.9) into the third equation of (S1) we obtain

$$r_3'' = -G\left[\frac{m_1(\sqrt{m_3} + \sqrt{m_2})^2 + m_2m_3}{(\sqrt{m_3} + \sqrt{m_2})^2}\right]\frac{r_3}{\|r_3\|^3}.$$
(2.11)

Since (2.10) must equal (2.11) we have the relation

$$\left(\frac{m_3}{m_2}\right)^{\frac{3}{2}} \left[\frac{m_1(\sqrt{m_3} + \sqrt{m_2})^2 + m_2 m_3}{(\sqrt{m_3} + \sqrt{m_2})^2}\right] \frac{r_3}{\|r_3\|^3} = \left[\frac{m_1(\sqrt{m_3} + \sqrt{m_2})^2 + m_2 m_3}{(\sqrt{m_3} + \sqrt{m_2})^2}\right] \frac{r_3}{\|r_3\|^3}.$$
 (2.12)

In order for (2.12) to be valid, either one of two possibilities occurs. First $\frac{r_3}{\|r_3\|^3} \equiv \vec{0}$, which implies that $\|r_3\| = \infty$, a contradiction to the assumption of finite initial conditions, or

$$\left(\frac{m_3}{m_2}\right)^{\frac{3}{2}} = 1 \iff m_3 = m_2. \tag{2.13}$$

Thus if (S1) does not satisfy (2.13), NCEM are inconsistent with respect to a BCOS model. However (S1) does satisfy (2.13), we can substitute (2.13) into (2.9) and (2.8) to obtain

$$r_2(t) = -r_3(t)$$
 and $||r_2(t)|| = ||r_3(t)||.$ (2.14)

Then (S1) reduces to

$$r_1 \equiv \overrightarrow{0}$$
 and $r_2'' = -r_3'' = G\left[m_1 + \frac{m_3}{4}\right] \frac{r_3}{\|r_3\|^3},$ (2.15)

and the second equation of (2.15) is a modified Kepler equation which has a conic section curve as a solution. Furthermore, if $r_1 \equiv \vec{0}$ and r_2 and r_3 satisfy (2.14), we see that the center of mass of the three bodies coincides with m_1 .

$$\frac{m_1r_1 + m_2r_2 + m_3r_3}{m_1 + m_2 + m_3} = \overrightarrow{0} = r_1.$$

The above discussion shows that we can not freely choose the origin O to be centered on a point mass m_1 , since unless the other point masses obey "antipodal" symmetry conditions, the choice of $r_1 \equiv \overrightarrow{0}$ in (S1) leads to the conclusion of m_2 and m_3 escaping to "infinity". To avoid this "seeming" inconsistency, we propose to use an origin invariant model of relative differences, namely

$$(r_1 - r_2)'' = -\frac{G(m_1 + m_2)(r_1 - r_2)}{\|r_1 - r_2\|^3} - \frac{Gm_3(r_1 - r_3)}{\|r_1 - r_3\|^3} + \frac{Gm_3(r_2 - r_3)}{\|r_2 - r_3\|^3}$$
$$(r_1 - r_3)'' = -\frac{Gm_2(r_1 - r_2)}{\|r_1 - r_2\|^3} - \frac{G(m_1 + m_3)(r_1 - r_3)}{\|r_1 - r_3\|^3} - \frac{Gm_2(r_2 - r_3)}{\|r_2 - r_3\|^3}.$$

By setting in $r_1(t) \equiv r_1'(t) \equiv r_1''(t) \equiv \overrightarrow{0}$, we obtain

$$r_{2}'' = -\frac{G(m_{1} + m_{2})r_{2}}{\|r_{2}\|^{3}} - \frac{Gm_{3}r_{3}}{\|r_{3}\|^{3}} - \frac{Gm_{3}(r_{2} - r_{3})}{\|r_{2} - r_{3}\|^{3}}$$
$$-r_{3}'' = \frac{Gm_{2}r_{2}}{\|r_{2}\|^{3}} + \frac{G(m_{1} + m_{3})r_{3}}{\|r_{1} - r_{3}\|^{3}} - \frac{Gm_{2}(r_{2} - r_{3})}{\|r_{2} - r_{3}\|^{3}}.$$

If m_2 and m_3 are small compared to m_1 , the preceding equations are perturbations of the second and third equations of (S1).

3 Well Posed Origin Anywhere Consistent Systems (WPOACS)

As the examples of the previous section demonstrate, it is desirable to have a system of differential equations for the N-body problem that has the following properties:

- i) The system of differential equations is consistent with a CCS with any origin O.
- ii) A unique solution is guaranteed.

We call and denote such systems as WPOACS. We now generalize the origin invariant model of the relative differences to N point masses m_i , where $1 \le i \le N$.

Theorem 3.1. Let *O* be any origin and $r_i(t) \in \mathbb{R}^3$ be the position of m_i . Denote N - 1 dependent variables by

$$r_{1k}(t) := r_1(t) - r_k(t), \qquad 2 \le k \le N.$$
 (3.1)

Then the following system

$$(r_1 - r_k)'' = G \sum_{i=2}^{N} \frac{m_i(r_i - r_1)}{\|r_i - r_1\|^3} - G \sum_{\substack{i=1\\i \neq k}}^{N} \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3}$$
$$= G \sum_{i=2}^{N} \frac{m_i(r_i - r_1)}{\|r_i - r_1\|^3} - G \sum_{\substack{i=1\\i \neq k}}^{N} \frac{m_i[r_1 - r_k - (r_1 - r_i)]}{\|r_1 - r_k - (r_1 - r_i)\|^3}, \qquad 2 \le k \le N,$$
(RS1)

is invariant with respect to an arbitrary CCS translation and is WPOACS.

Proof: For any $c(t) \in C^2(\mathbb{R})$, observe that

$$\begin{aligned} r_1'' - r_k'' &= (r_1 + c(t))'' - (r_k + c(t))'' \\ &= \sum_{i=2}^N \frac{Gm_i(r_i + c(t) - (r_1 + c(t)))}{\|r_i + c(t) - (r_1 + c(t))\|^3} - \sum_{\substack{i=1\\i \neq k}}^N \frac{Gm_i(r_i + c(t) - (r_k + c(t)))}{\|r_i(t) + c(t) - (r_k + c(t))\|^3} \\ &= G\sum_{i=2}^N \frac{m_i(r_i - r_1)}{\|r_i - r_1\|^3} - G\sum_{\substack{i=1\\i \neq k}}^N \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3}.\end{aligned}$$

Thus our system is invariant under an arbitrary translation. Next we impose on (RS1) the following initial conditions $r_{1k}(t_0)$, $r'_{1k}(t_0) \in \mathbb{R}^3$ subject to

$$r_{1k}(t_0) \neq \overrightarrow{0}, \qquad r_{1k}(t_0) - r_{1j}(t_0) \neq \overrightarrow{0}, \qquad 1 \le k, j \le N, \ k \ne j.$$
 (3.2)

By Pages 1-7 of [2] there exists an interval $[a, b] \subseteq \mathbb{R}$, where $t_0 \in [a, b]$, on which a solution $r_{1k}(t) \in C^2([a, b])$ to (RS1) exists. \Box

Observe the perfect match of certain constraints on the initial conditions in (1.1) to certain constraints on the initial conditions in (3.2).

$$r_k(t_0) - r_j(t_0) \neq \overrightarrow{0} \iff r_{1k}(t_0) \neq \overrightarrow{0}, \text{ and } r_{1k}(t_0) - r_{1j}(t_0) \neq \overrightarrow{0}, \text{ when } k \neq j,$$
 (3.3)

where we implicitly assumed that $2 \le k, j \le N$. Moreover, the substitution $r_1(t) \equiv r'_1(t) \equiv \vec{r'_1}(t) \equiv \vec{0}$ in (*RS1*) results in a system of equations free of inconsistencies

$$-r_k'' = G\sum_{i=2}^N \frac{m_i r_i}{\|r_i\|^3} + \frac{Gm_1 r_k}{\|r_k\|^3} - G\sum_{\substack{i=2\\i\neq k}}^N \frac{m_i (r_i - r_k)}{\|r_i - r_k\|^3}, \qquad 2 \le k \le N.$$
(3.4)

Another system closely related to (RS1) and invariant under arbitrary translations c(t) is

$$(r_j - r_k)'' = G \sum_{\substack{i=1\\i \neq j}}^N \frac{m_i(r_i - r_j)}{\|r_i - r_j\|^3} - G \sum_{\substack{i=1\\i \neq k}}^N \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3}, \qquad 1 \le j < k \le N,$$
(RS2)

where the dependent variables are

$$r_{jk} := r_j(t) - r_k(t), \qquad 1 \le j < k \le N.$$
 (3.5)

System (RS2) consists of $\binom{N}{2}$ equations and is generated by the N-1 equations $(r_1 - r_p)'', 2 \le p \le N$ since

$$(r_j - r_k)'' = (r_1 - r_k)'' - (r_1 - r_j)''$$
 whenever $j \neq 1$. (3.6)

Because the right side of the equations of (1.1) involves differences of the form $r_i - r_j$, (RS2) could be technically preferable to (RS1).

4 A New Origin Invariant Constant of Motion

In this section we derive a new constant of motion for (RS2) which has several ramifications that we later discuss.

Theorem 4.1. Assume that $r_j(t) - r_k(t) \in \mathbb{R}^3$, where $1 \le j < k \le N$, are solutions on (a, b) to the system of equations

$$(r_j - r_k)'' = G \sum_{\substack{i=1\\i \neq j}}^{N} \frac{m_i(r_i - r_j)}{\|r_i - r_j\|^3} - G \sum_{\substack{i=1\\i \neq k}}^{N} \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3}$$
$$= \frac{-G(m_j + m_k)(r_j - r_k)}{\|r_j - r_k\|^3} + G \sum_{\substack{i=1\\i \neq j,k}}^{N} m_i \left[\frac{r_i - r_j}{\|r_i - r_j\|^3} - \frac{r_i - r_k}{\|r_i - r_k\|^3} \right].$$
(4.1)

Let $M := \sum_{i=1}^{N} m_i$. Then the following identity holds

$$\sum_{1 \le j < k \le N} m_j m_k (r_j - r_k) \cdot (r_j - r_k)'' = -GM \sum_{1 \le j < k \le N} \frac{m_j m_k}{\|r_j - r_k\|} < 0.$$
(4.2)

Proof: Take each equation in (4.1) and form the dot product with the vector $m_i m_k (r_i - r_k)$ to obtain

$$m_j m_k (r_j - r_k) \cdot (r_j - r_k)'' = -\frac{G m_j m_k (m_j + m_k)}{\|r_j - r_k\|} + T(j,k),$$
(4.3)

where

$$T(j,k) := Gm_j m_k \sum_{\substack{i=1\\i \neq j,k}}^N m_i \left[\frac{(r_j - r_k) \cdot (r_i - r_j)}{\|r_i - r_j\|^3} - \frac{(r_j - r_k) \cdot (r_i - r_k)}{\|r_i - r_k\|^3} \right].$$
(4.4)

Next sum both sides of (4.3) with respect to all pairs of indices (j, k) to obtain

$$\sum_{1 \le j < k \le N} m_j m_k (r_j - r_k) \cdot (r_j - r_k)'' = -\sum_{1 \le j < k \le N} \frac{Gm_j m_k (m_j + m_k)}{\|r_j - r_k\|} + \sum_{1 \le j < k \le N} T(j, k),$$
(4.5)

where

$$\sum_{1 \le j < k \le N} T(j,k) = G \sum_{1 \le j < k \le N} m_j m_k (r_j - r_k) \cdot \sum_{\substack{i=1\\i \ne j,k}}^N m_i \left[\frac{r_i - r_j}{\|r_i - r_j\|^3} - \frac{r_i - r_k}{\|r_i - r_k\|^3} \right].$$
(4.6)

The goal is to show that

$$\sum_{1 \le j < k \le N} T(j,k) = -G \sum_{1 \le j < k \le N} \frac{(M - m_j - m_k)m_jm_k}{\|r_j - r_k\|}.$$
(4.7)

In order to prove (4.7), temporarily fix an index pair (j, k) and recall that we are summing the $\binom{N}{2}$ equations $m_{\ell}m_p(r_{\ell}-r_p) \cdot (r_{\ell}-r_p)''$, where $1 \leq \ell . Look at the sum of the <math>\binom{N}{2} - 1$ equations arising from $(r_{\ell}-r_p)''$, where $1 \leq \ell and <math>(j,k) \neq (\ell,p)$, and add together the terms which have a factor of $\frac{r_j-r_k}{\|r_j-r_k\|^3}$. This process is equivalent to interchanging the order of summation on the right side of (4.6). Observe that in order to obtain $\frac{r_j-r_k}{\|r_j-r_k\|^3}$ either $\ell = j, k$ or p = j, k. The number of such ordered pairs $(\ell, p) \neq (j, k)$ for which either $\ell = j, k$ or p = j, k is $\binom{N}{2} - \binom{N-2}{2} - 1 = 2N - 4$, where $\binom{N-2}{2}$ counts those ordered pairs which are independent of j and k. These 2N - 4 ordered pairs will be paired up with opposite signs. To explain how the pairing occurs we analyze four mutually exclusive cases.

Case 1: $\ell = j$ and $p \neq k, j$. This implies that p > j. Then $(r_{\ell} - r_p)'' = (r_j - r_p)''$ where

$$(r_j - r_p)'' = \frac{-G(m_j + m_p)(r_j - r_p)}{\|r_j - r_p\|^3} + G\sum_{\substack{i=1\\i\neq j,p}}^N \frac{m_i(r_i - r_j)}{\|r_i - r_j\|^3} - G\sum_{\substack{i=1\\i\neq j,p}}^N \frac{m_i(r_i - r_p)}{\|r_i - r_p\|^3}.$$
 (4.8)

Since $p \neq j, k$, only i = k in second summand on the right side of (4.8) gives rise to $r_j - r_k$, in which case

we obtain the term

$$\frac{Gm_k(r_k - r_j)}{\|r_k - r_j\|^3} = -\frac{Gm_k(r_j - r_k)}{\|r_j - r_k\|^3}$$

We then form the dot product with the vector $m_j m_p (r_j - r_p)$ to obtain a summand of the form

$$-\frac{Gm_j m_k m_p (r_j - r_k) \cdot (r_j - r_p)}{\|r_j - r_k\|^3}.$$
(4.9)

Case 2: $\ell = k$ and $p \neq j, k$. This implies that p > k. Then $(r_{\ell} - r_p)'' = (r_k - r_p)''$ where

$$(r_k - r_p)'' = \frac{-G(m_k + m_p)(r_k - r_p)}{\|r_k - r_p\|^3} + G\sum_{\substack{i=1\\i \neq k, p}}^N \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3} - G\sum_{\substack{i=1\\i \neq k, p}}^N \frac{m_i(r_i - r_p)}{\|r_i - r_p\|^3}.$$
 (4.10)

Since $p \neq j, k$, only i = j in second summand on the right side of (4.10) gives rise to $r_j - r_k$, which after taking the dot product with $m_k m_p (r_k - r_p)$, leads to the summand

$$\frac{Gm_j m_k m_p (r_k - r_p) \cdot (r_j - r_k)}{\|r_j - r_k\|^3}.$$
(4.11)

Case 3: $\ell \neq k, j$ and p = j. This implies $\ell < j$. Then $(r_{\ell} - r_p)'' = (r_{\ell} - r_j)''$ where

$$(r_{\ell} - r_j)'' = \frac{-G(m_{\ell} + m_j)(r_{\ell} - r_j)}{\|r_{\ell} - r_j\|^3} + G\sum_{\substack{i=1\\i \neq \ell, j}}^N \frac{m_i(r_i - r_\ell)}{\|r_i - r_\ell\|^3} - G\sum_{\substack{i=1\\i \neq \ell, j}}^N \frac{m_i(r_i - r_j)}{\|r_i - r_j\|^3}.$$
(4.12)

Since $\ell \neq j, k$, only i = k in the third summand on the right side of (4.12) gives rise to $r_j - r_k$, which after taking the dot product with $m_\ell m_j (r_\ell - r_j)$, leads to the summand

$$\frac{Gm_j m_\ell m_k (r_\ell - r_j) \cdot (r_j - r_k)}{\|r_j - r_k\|^3}.$$
(4.13)

Case 4: $\ell \neq k, j$ and p = k. This implies $\ell < k$. Then $(r_{\ell} - r_p)'' = (r_{\ell} - r_k)''$ where

$$(r_{\ell} - r_k)'' = \frac{-G(m_{\ell} + m_k)(r_{\ell} - r_k)}{\|r_{\ell} - r_j\|^3} + G\sum_{\substack{i=1\\i \neq \ell,k}}^N \frac{m_i(r_i - r_\ell)}{\|r_i - r_\ell\|^3} - G\sum_{\substack{i=1\\i \neq \ell,k}}^N \frac{m_i(r_i - r_k)}{\|r_i - r_k\|^3}.$$
 (4.14)

Since $\ell \neq j, k$, only i = j in the third summand on the right side of (4.12) gives rise to $r_j - r_k$, which after taking the dot product with the vector $m_\ell m_k (r_\ell - r_k)$, lead to the summand

$$\frac{Gm_j m_\ell m_k (r_\ell - r_k) \cdot (r_j - r_k)}{\|r_j - r_k\|^3}.$$
(4.15)

In all four cases, as evidenced by (4.9), (4.11), (4.13), and (4.15), there is a factor of the form $m_j m_k m_{\alpha}$, where α is either ℓ or p. We want to pairwise combine via the value of α . The above four cases imply that $\alpha \neq j, k$. However, α is free to be any other value from the set $\{1, ..., N\}$. This leads to following considerations: $\alpha < j, j < \alpha < k$, and $\alpha > k$. Suppose $\alpha > k$. This occurs for all of the p in Case 2 and those p in Case 1 for which p > k. We pairwise add (4.9) and (4.11) to obtain

$$-\frac{Gm_jm_km_p(r_j - r_p) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} + \frac{Gm_jm_km_p(r_k - r_p) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} = -\frac{Gm_jm_km_p}{\|r_j - r_k\|}.$$
 (4.16)

Next consider $\alpha < j$. This occurs for all of the ℓ in Case 3 and those ℓ in Case 4 for which $\ell < j$. We pairwise add (4.13) and (4.15) to obtain

$$\frac{Gm_j m_\ell m_k (r_\ell - r_j) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} - \frac{Gm_j m_\ell m_k (r_\ell - r_k) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} = -\frac{Gm_j m_k m_p}{\|r_j - r_k\|},$$
(4.17)

where in the last term we relabeled ℓ to p. Finally we have to consider when $j < \alpha < k$. This occurs in the remaining p and ℓ of Cases 1 and 4 not covered by (4.16) and (4.17) respectively. We can pairwise add (4.9) to (4.15), where in (4.15) we have renamed ℓ to p, to obtain

$$-\frac{Gm_jm_km_p(r_j - r_p) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} - \frac{Gm_jm_pm_k(r_p - r_k) \cdot (r_j - r_k)}{\|r_j - r_k\|^3} = -\frac{Gm_jm_km_p}{\|r_j - r_k\|}.$$
(4.18)

If we add (4.16) through (4.18) together, we get $-G(M - m_j - m_k)m_jm_k/||r_j - r_k||$. The above term is true for an arbitrary yet fixed (j, k), where $1 \le j < k \le N$. By summing over $1 \le j < k \le N$, we obtain (4.7).

By combining (4.5) with (4.7) we obtain

$$\sum_{1 \le j < k \le N} m_j m_k (r_j - r_k) \cdot (r_j - r_k)'' = -GM \sum_{1 \le j < k \le N} \frac{m_j m_k}{\|r_j - r_k\|} < 0. \qquad \Box \qquad (4.19)$$

5 The Restless Universe

Below we list some consequences of Inequality (4.2). These consequences reflect the fact that $\sum_{1 \le j < k \le N} m_j m_k (r_j - r_k) \cdot (r_j - r_k)''$ is always negative and that for $1 \le j < k \le N$ we made the implicit assumption of $r_j - r_k \ne 0$ on [a, b]. This last condition is tantamount to the collision free model.

- **Corollary 5.1.** 1. Under the conditions of Theorem 4.1, the relative system (4.1) has no critical points (constant solutions).
 - 2. For each $t \in [a, b]$ there exists an index set (j, k) with $1 \le j < k \le N$ such that $(r_j(t) r_k(t))'' \ne \overrightarrow{0}$.
 - 3. For each $t \in [a, b]$ there exists an index set (j, k) with $1 \le j < k \le N$ such that $r_j(t) r_k(t)$ is not orthogonal to $(r_j(t) r_k(t))''$.
 - 4. There exists an index set (j,k) such that $(r_j r_k)'' \not\equiv \overrightarrow{0}$ on [a,b]. Hence, there exists an index set (j,k) such that $r_j r_k$ is not identically constant on [a,b]; this implies that one of the two position vectors associated with the index set (j,k) is not identically constant on [a,b].

Corollary 5.1, Part 2, implies that $\left[r_1''(t) - r_2''(t), \dots, r_i''(t) - r_j''(t), \dots, r_{N-1}''(t) - r_N''(t)\right]^T \neq \vec{0}$ for all $t \in [a, b]$. However, as evidenced by the following corollary, we can say more.

Corollary 5.2. For $N \ge 3$, under the conditions of Theorem 4.1, there are two distinct pairs of indices (j_1, j_2) and (j_3, j_4) from $\{(\ell, p)\}_{1 \le \ell such that <math>(r_{j_1} - r_{j_2})''$ and $(r_{j_3} - r_{j_4})''$ are both not identically zero.

Proof: Without loss of generality, after relabeling if necessary, we can assume via (4.19) that $j_1 = 1$ and $j_2 = 2$ and that $(r_1 - r_2)'' \neq \vec{0}$. Since

$$\sum_{i=1}^{N-1} (r_i'' - r_{i-1}'') + (r_N'' - r_1'') = (r_1'' - r_2'') + (r_2'' - r_3'') + \dots + (r_{N-1}'' - r_N) + (r_N'' - r_1'') \equiv \overrightarrow{0},$$

and since $(r_1 - r_2)'' \not\equiv \overrightarrow{0}$, then

$$(r_2'' - r_3'') + \dots + (r_{N-1}'' - r_N) + (r_N'' - r_1'') \equiv -(r_1 - r_2)'' \not\equiv \vec{0},$$

and we can choose (j_3, j_4) to be any (i, i+1) where $2 \le i \le N-1$. \Box

By combining Corollaries 5.1 and 5.2, we deduce the following.

Corollary 5.3. Let $N \ge 3$. Under the conditions of Theorem 4.1, for each $t \in [a, b]$, at least two bodies accelerate.

Remark 5.1. Assume that the $(r_j(t))_{j=1}^N$ associated with Theorem 1.1 are real analytic functions on [c,d] where $a \le c < d \le b$. Furthermore assume that $r''_j - r''_k \not\equiv \overrightarrow{0}$ on [c,d]. Then $r''_j(t) - r''_k(t) = \overrightarrow{0}$ for only a finite number of $t \in [c,d]$.

Proof By Contradiction: Define $h(t) := r''_j(t) - r''_k(t)$ on [c, d] and assume that $h(t_j) = \overrightarrow{0}$ for a countable number of $t_j \in [c, d]$. There exists a subsequence (t_{j_ℓ}) of (t_j) such that $t_{j_\ell} \to t_0 \in [c, d]$ and $h(t_{j_\ell}) = \overrightarrow{0}$. The analyticity of h at t_0 implies that $h(t) = h(t_0) + \sum_{n=1}^{\infty} a_n(t-t_0)^n$ for some neighborhood of $|t - t_0| < \rho$ in the complex plane. By construction $\lim_{\ell \to \infty} h(t_{j_\ell}) = \overrightarrow{0} = h(t_0)$. Hence we deduce that

$$h(t) = \sum_{n=1}^{\infty} a_n (t - t_0)^n = (t - t_0) \left[a_1 + \sum_{n=2}^{\infty} a_n (t - t_0)^{n-1} \right], \qquad |t - t_0| < \rho.$$
(5.1)

If we substitute $t_{j_{\ell}} \neq t_0$ into (5.1), since $t_{j_{\ell}} \neq t_0$ and $h(t_{j_{\ell}}) = \overrightarrow{0}$, we deduce that

$$\overrightarrow{0} = a_1 + \sum_{n=2}^{\infty} a_n (t_{j_\ell} - t_0)^{n-1}$$
(5.2)

Suppose $a_1 \neq \overrightarrow{0}$. By analyticity of h at t_0 there exists $\hat{\rho}$ such that

$$0 < \left| \sum_{n=2}^{\infty} a_n (t - t_0)^{n-1} \right| \le \frac{|a_1|}{2}, \qquad 0 < |t - t_0| < \hat{\rho} < \rho.$$
(5.3)

The triangle inequality, when combined with (5.3) implies that

$$0 < \frac{|a_1|}{2} = |a_1| - \frac{|a_1|}{2} \le |a_1| - \left|\sum_{n=2}^{\infty} a_n (t - t_0)^{n-1}\right| \le \left|a_1 + \sum_{n=2}^{\infty} a_n (t - t_0)^{n-1}\right|,$$
(5.4)

whenever $0 < |t - t_0| < \hat{\rho} < \rho$. Since $t_{j_\ell} \to t_0$, Inequality (5.4) holds for those $t_{j_\ell} \in |t - t_0| < \hat{\rho}$ and implies that $a_1 + \sum_{n=2}^{\infty} a_n (t_{j_\ell} - t_0)^{n-1} \not\equiv \vec{0}$, a contradiction of (5.2). Hence $a_1 = \vec{0}$. By repeating this

argument, we show there is a neighborhood of t_0 in the complex plane such that all $a_n \equiv \vec{0}$, i.e $h(t) \equiv \vec{0}$ for $|t - t_0| < \rho_1$. Then by analytic continuation we conclude that $h(t) \equiv \vec{0}$ for $t \in [c, d]$, a contradiction of the initial assumption. \Box

6 Conclusions

This article is motivated by the necessity to have a framework of equations for celestial mechanics which support measurements from any point mass in the universe whether accelerating or not. In Section 2 we showed that NCME for the two and three body problems do not satisfy this requirements. In order to construct a framework of origin invariant equations, we derive from NCME a relative system of equations, either System (RS1) or System (RS2), that is consistent in any CCS. System (RS1) (respectively System (RS2)) is shown to be WPOACS, to accommodate BCOS, and be invariant with respect to an arbitrary time dependent translation of coordinates of the form $r_i(t) = \tilde{r_i}(t) - c(t)$, $c''(t) \neq \vec{0}$, $t \in [a, b]$; see Theorem 3.1. We recall that NCME are invariant only with respect to inertial translations where $c''(t) \equiv \vec{0}$. NCME and (RS1) are related but are not equivalent. However, for the earth-sun Kepler problem, the NCME and (RS1) are approximately the same. The adoption of the relative system comes with an uncertainty advisory. Given an N-body problem we can be certain about the relative position of N - 1-point masses but we must stay uncertain about the position of one point mass. Finally we observe that we need a substitute for the resultant of forces acting on the binary mass (m_j, m_k) associated with the relative accelerations $(r_j - r_k)''$ of (RS1) (respectively (RS2)). The integral of motion calculations in [4] suggests the use of a factor $\mu m_j m_k$, where μ is a certain cosmological constant to be determined by experiment.

We end this article with a foundational proposition which underlies the calculations of Section 2.

Proposition 6.1. Given a unique CCS whose origin O corresponds to a point mass m, if $r(t) \in \mathbb{R}^3$ is defined to be the position of m as measured from O, and if $A(t) \in C^2(\mathbb{R})$, then the initial value problem

$$r''(t) = A(t), \qquad r(t_0), r'(t_0) \in \mathbb{R}^3$$

(6.1)

does not have a solution unless $A(t) \equiv \overrightarrow{0}$.

The proof follows from the fact the assumptions imply that $r(t) \equiv r'(t) \equiv r''(t) \equiv \vec{0}$. The inconsistencies shown in Section 2 are a result of the assumptions of Proposition 6.1. A main result of this article is employing relative systems as a means of avoiding the contradiction inherent in Proposition 6.1. These relative difference systems utilize only one CCS with an origin O that coincides with a point mass m_j and are shown to be WPOACS. If $A(t) \neq \vec{0}$, then a solution to (6.1) exists if and only if the assumptions of Proposition 6.1 are invalid. The negation of these assumptions logically corresponds to the following statement: either there is a unique CCS whose origin O does not correspond to a point mass or there is more than one CCS. This leads to the following corollary.

Corollary 6.1. Given a unique CCS whose origin O does not correspond to any point mass m, if $r(t) \in \mathbb{R}^3$ is defined to be the position of m as measured from O, and if $A(t) \in C^2(\mathbb{R})$, then the initial value problem

$$r''(t) = A(t) \not\equiv \overrightarrow{0}, \qquad r(t_0), r'(t_0) \in \mathbb{R}^3$$
(6.2)

has a unique solution given by $r(t) = r(t_0) + r'(t_0)(t-t_0) + \int_{t_0}^t \int_{t_0}^s A(u) du ds$.

Corollary 6.2. Given two distinct CCSs, one with origin O and the other with origin $\tilde{O} \neq O$, set r(t) := OOO(t). Then the initial value problem (6.2) has a unique solution.

The existence of more than one CCS provides another means, independent of the relative difference WPOACS, in which to have the origin of a coordinate system coincide with a point mass in a manner which avoids the contradictions of Proposition 6.1. Assume we want our equations formulated in a CCS whose origin \tilde{O} coincides with the point mass m_1 . We start with NCME (1.1) that are formulated in a CCS with origin O that does not coincide with any of the N point masses. Theorem 1.1 guarantees a solution on some interval [a, b]. If we let $\tilde{r}_i(t)$ denote the position of m_i with respect to \tilde{O} , since $r_i(t) = r_1(t) + \tilde{r}_i(t)$, we transform (1.2) into

$$r_1''(t) + \tilde{r}_1''(t) = \sum_{j=2}^N \frac{Gm_j(r_j(t) - r_1(t))}{\|r_j(t) - r_1(t)\|^3} \equiv \sum_{j=2}^N \frac{Gm_j\tilde{r}_j(t)}{\|\tilde{r}_j(t)\|^3},$$
(6.3)

and

$$r_1''(t) + \tilde{r}_i''(t) = \sum_{j \neq i} \frac{Gm_j(r_j(t) - r_i(t))}{\|r_j(t) - r_i(t)\|^3} \equiv \sum_{j \neq i} \frac{Gm_j(\tilde{r}_j(t) - \tilde{r}_i(t))}{\|\tilde{r}_j(t) - \tilde{r}_i(t)\|^3}, \ 2 \le i \le N,$$
(6.4)

which is a different set of equations than the relative equations in of Section 3. Since $r_1''(t) = \sum_{j=2}^{N} \frac{Gm_j(r_j(t) - r_1(t))}{\|r_j(t) - r_1(t)\|^3}$, Equation (6.3) is equivalent to $\tilde{r}_1(t) = 0$. The system of equations provided by (6.3) is equivalent to postulating that the N-body problem is governed by a system of equations

$$r_i''(t) + c''(t) = \sum_{\substack{j=1\\j\neq i}}^N \frac{Gm_j(r_j(t) - r_i(t))}{\|r_j(t) - r_i(t)\|^3}, \ 1 \le i \le N,$$

where $c(t) \in C^2([a, b])$ is a fictitious acceleration that is yet to be determined, and where [a, b] is an interval of existence of the desired solutions. This will allow for non inertial translations, but in a manner different than that provided by Theorem 3.1.

References

- [1] E. Bannikova and M. Capaccioli, *Foundations of Celestial Mechanics*, Graduate Texts in Physics, 1st Edition, Springer, 2022.
- [2] P. F. Hsieh and Y. Sibuya, *Basic Theory of Ordinary Differential Equations*, Universitext, Springer-Verlag, New York, 1999.
- [3] H. Pollard, *Celestial Mechanics*, Carus Mathematical Monographs, No. 18, Mathematical Association of America, 1976.
- [4] J. Quaintance and H. Gingold, *Relative Difference System Involving Newton's Equations of Celestial Mechanics*, Preprints 2022, 2022100277.
- [5] A. Wintner, The Analytical Foundation of Celestial Mechanics, Princeton University Press, 1941.