Theoretical guarantees for lifted samplers

Philippe Gagnon¹, Florian Maire¹

May 28, 2024

¹Department of Mathematics and Statistics, Université de Montréal.

Abstract

Lifted samplers form a class of Markov chain Monte Carlo methods which has drawn a lot attention in recent years due to superior performance in challenging Bayesian applications. A canonical example of such sampler is the one that is derived from a random walk Metropolis algorithm for a totally-ordered state space such as the integers or the real numbers. The lifted sampler is derived by splitting into two the proposal distribution: one part in the increasing direction, and the other part in the decreasing direction. It keeps following a direction, until a rejection, upon which it flips the direction. In terms of asymptotic variances, it outperforms the random walk Metropolis algorithm, regardless of the target distribution, at no additional computational cost. Other studies show, however, that beyond this simple case, lifted samplers do not always outperform their Metropolis counterparts. In this paper, we leverage the celebrated work of Tierney (1998) to provide an analysis in a general framework encompassing a broad class of lifted samplers. Our finding is that, essentially, the asymptotic variances cannot increase by a factor of more than 2, regardless of the target distribution, the way the directions are induced, and the type of algorithm from which the lifted sampler is derived (be it a Metropolis–Hastings algorithm, a reversible jump algorithm, etc.). This result indicates that, while there is potentially a lot to gain from lifting a sampler, there is not much to lose.

Keywords: asymptotic variances; Bayesian statistics; Markov chain Monte Carlo; Metropolis–Hastings; Non-reversible Markov chains.

1 Introduction

1.1 Context

Let us consider the situation where one is interested in sampling from π , a probability distribution defined on a measurable space (X, X), where $X \subset \mathbb{R}^d$ is assumed to correspond to the support of π , with d a positive integer, and X is a sigma-algebra on X. In a sampling context, π is often referred to as the *target distribution*. Markov chain Monte Carlo (MCMC) methods can be used to sample from π . We consider in the first part of this paper that one wants to compare the performance of a Metropolis–Hastings (MH, (Metropolis *et al.*, 1953; Hastings, 1970)) algorithm with that of a *lifted* version (Horowitz, 1991; Gustafson, 1998; Chen *et al.*, 1999; Diaconis *et al.*, 2000). Recently, Andrieu and Livingstone (2021) proved that the latter outperforms the former in terms of asymptotic variances in specific situations. Empirically, it has been shown not to be the case in other situations (Gagnon and Maire, 2024). In this paper, we fill some gaps in the theory by providing general guarantees.

In the rest of this section, we set the scene for a comparison between MH and its lifted version by recalling how a MH algorithm works and by introducing its lifted version through an intermediate sampler. We next present in Section 2 a theoretical result about performance guarantees and in Section 3 empirical results for an illustration. The MH framework of Sections 1-3 is explicit and, thus, easy to understand. The presentation of this framework will facilitate the understanding of the content of Section 4, which is a theoretical result analogous to that in Section 2 under a more general framework. To introduce this framework, we define a generalized MH algorithm following Tierney (1998) and, similarly as below, its lifted version through an intermediate sampler. This generalized MH algorithm includes as special cases algorithms such as reversible jump (Green, 1995), Multiple-try Metropolis (Liu *et al.*, 2000) and pseudo-marginal (Beaumont, 2003). The result of Section 4 can be seen as a complement to those about lifted versions of sophisticated sampling schemes such as simulated tempering (Sakai and Hukushima, 2016b), parallel tempering (Syed *et al.*, 2022) and reversible jump for Bayesian nested model selection (Gagnon and Doucet, 2021).

1.2 MH algorithm and an auxiliary sampler

A MH algorithm proceeds at each iteration by first proposing a state $\mathbf{y} \in X$ using a *proposal distribution* $Q(\mathbf{x}, \cdot)$, where $\mathbf{x} \in X$ is the current state of the Markov chain, and next by deciding whether or not the proposal is accepted. If the proposal is accepted, the next state of the Markov chain is \mathbf{y} , otherwise, the chain remains at \mathbf{x} . The decision to accept or reject the proposal is random and is based on an acceptance probability, denoted by $\alpha(\mathbf{x}, \mathbf{y})$, whose form is made explicit below.

We assume that π and $Q(\mathbf{x}, \cdot)$ admit positive densities with respect to a common dominating measure, denoted for simplicity dy. With an abuse of notation, we also denote these densities by π and $Q(\mathbf{x}, \cdot)$. Let us denote by $\mathbf{N}(\mathbf{x})$ the support of $Q(\mathbf{x}, \cdot)$, where $\{\mathbf{N}(\mathbf{x})\}$ yields a neighbourhood structure on \mathcal{X} . We assume that $\mathbf{x} \notin \mathbf{N}(\mathbf{x})$. This implies that, strictly speaking, $\mathbf{N}(\mathbf{x})$ is not a neighbourhood in a topological sense; we nevertheless carry on with this abuse of terminology. The restriction $\mathbf{x} \notin \mathbf{N}(\mathbf{x})$ is not necessary but it simplifies the presentation of the auxiliary and lifted samplers below. We also assume that $\mathbf{N}(\mathbf{x})$ is not empty and of positive $Q(\mathbf{x}, \cdot)$ -measure, for all \mathbf{x} . Finally, we assume that: for any $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$,

$$\mathbf{y} \in \mathbf{N}(\mathbf{x})$$
 if and only if $\mathbf{x} \in \mathbf{N}(\mathbf{y})$.

For instance, when X is the set of integers, $\{N(x)\}$ can be such that $N(x) = \{x - 1, x + 1\}$, and when $X = \mathbb{R}$, $\{N(x)\}$ can be such that $N(x) = \mathbb{R} \setminus \{x\}$. In the case where $\{x\}$ is of null measure, for all x, like with continuous random variables, one would typically include $\{x\}$ in N(x), but in practice, including it or not makes no difference in the algorithm.

Under the framework just described, the acceptance probability has the following form: for all $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$, $\alpha(\mathbf{x}, \mathbf{y}) = 1 \wedge r(\mathbf{x}, \mathbf{y})$, where $x \wedge y = \min\{x, y\}$, $r(\mathbf{x}, \mathbf{y}) = 0$ whenever $\mathbf{y} \notin \mathbf{N}(\mathbf{x})$, and otherwise,

$$r(\mathbf{x}, \mathbf{y}) = \frac{\pi(\mathbf{y}) Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) Q(\mathbf{x}, \mathbf{y})}.$$

The Markov kernel associated to the MH algorithm, denoted by P_{MH} , is such that, for any $\mathbf{x} \in \mathcal{X}$,

$$P_{\rm MH}(\mathbf{x}, d\mathbf{y}) = Q(\mathbf{x}, d\mathbf{y}) \,\alpha(\mathbf{x}, \mathbf{y}) + \delta_{\mathbf{x}}(d\mathbf{y}) \int (1 - \alpha(\mathbf{x}, \mathbf{u})) \,Q(\mathbf{x}, d\mathbf{u}),$$

where δ is the Dirac measure. We consider that

$$Q(\mathbf{x}, \mathbf{y}) = \frac{q(\mathbf{x}, \mathbf{y})}{c(\mathbf{x})}, \quad \mathbf{y} \in \mathbf{N}(\mathbf{x}),$$

where $q(\mathbf{x}, \cdot)$ represents the unnormalized density, and $c(\mathbf{x}) > 0$ is the normalizing constant, that is

$$c(\mathbf{x}) = \int_{\mathbf{N}(\mathbf{x})} q(\mathbf{x}, \mathbf{y}) \,\mathrm{d}\mathbf{y}.$$

The MH Markov kernel is π -reversible. We now present a variant of the MH algorithm which uses a notion of *direction* and will also be seen to be π -reversible. Let us define a *direction* variable $\nu \in$ $\{-1, +1\}$ and *directional* neighbourhoods $\mathbf{N}_{-1}(\mathbf{x})$ and $\mathbf{N}_{+1}(\mathbf{x})$ that are such that $\mathbf{N}_{-1}(\mathbf{x}) \cup \mathbf{N}_{+1}(\mathbf{x}) = \mathbf{N}(\mathbf{x})$ and $\mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x}) = \emptyset$ for all \mathbf{x} . Analogously to the MH case, we assume that: for any $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$ and $\nu \in \{-1, +1\}$,

$$\mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x}) \text{ if and only if } \mathbf{x} \in \mathbf{N}_{-\nu}(\mathbf{y}).$$
 (1)

If **x** would belong to $N(\mathbf{x})$, we would need to include **x** in both $N_{-1}(\mathbf{x})$ and $N_{+1}(\mathbf{x})$ to satisfy the condition in (1), and we would have $N_{-1}(\mathbf{x}) \cap N_{+1}(\mathbf{x}) = \{\mathbf{x}\}$ instead of $N_{-1}(\mathbf{x}) \cap N_{+1}(\mathbf{x}) = \emptyset$; the result presented in Section 2 would hold even in this case. We define a notion of *boundary* of X as $\partial X := \{\mathbf{x} \in X :$ $N_{-1}(\mathbf{x}) = \emptyset\} \cup \{\mathbf{x} \in X : N_{+1}(\mathbf{x}) = \emptyset\}$. To simplify the presentation of the variant of MH, we assume for now that $\partial X = \emptyset$. This is the case for example when $X = \mathbb{Z}$ and $N(\mathbf{x}) = \{\mathbf{x} - 1, \mathbf{x} + 1\}$ with $N_{-1}(\mathbf{x}) = \{\mathbf{x} - 1\}$ and $N_{+1}(\mathbf{x}) = \{\mathbf{x} + 1\}$.

The variant of MH proceeds similarly to MH, the main difference being that it starts by picking uniformly at random a direction $v \in \{-1, +1\}$. Next, it samples a proposal y using the conditional distribution given that the proposal belongs to $\mathbf{N}_{v}(\mathbf{x})$ that we denote by $Q_{v}(\mathbf{x}, \cdot)$. This conditional distribution admits a density with respect to dy, which is denoted by $Q_{v}(\mathbf{x}, \cdot)$ as well to simplify and given by

$$\mathbf{y} \in \mathcal{X}, \quad Q_{\nu}(\mathbf{x}, \mathbf{y}) = \frac{q(\mathbf{x}, \mathbf{y})}{c_{\nu}(\mathbf{x})} \, \mathbb{1}(\mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})),$$

where

$$c_{\nu}(\mathbf{x}) = \int_{\mathbf{N}_{\nu}(\mathbf{x})} q(\mathbf{x}, \mathbf{y}) \,\mathrm{d}\mathbf{y},$$

and 1 is the indicator function. We refer to the conditional distribution $Q_{\nu}(\mathbf{x}, \cdot)$ as the *directional* proposal distribution. The last step of the variant of MH is a random decision of whether or not the proposal \mathbf{y} is accepted, based on an acceptance probability α_{ν} defined as follows: for all $(\mathbf{x}, \mathbf{y}) \in X^2$, $\alpha_{\nu}(\mathbf{x}, \mathbf{y}) = 1 \wedge r_{\nu}(\mathbf{x}, \mathbf{y})$, where $r_{\nu}(\mathbf{x}, \mathbf{y}) = 0$ whenever $\mathbf{y} \notin \mathbf{N}_{\nu}(\mathbf{x})$, and otherwise,

$$r_{\nu}(\mathbf{x}, \mathbf{y}) = \frac{\pi(\mathbf{y}) \, Q_{-\nu}(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{y})}$$

The resulting Markov kernel, denoted by $P_{rev.}$, is such that, for any $\mathbf{x} \in \mathcal{X}$,

$$P_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) = \sum_{\nu \in \{-1, +1\}} \frac{1}{2} Q_{\nu}(\mathbf{x}, d\mathbf{y}) \alpha_{\nu}(\mathbf{x}, \mathbf{y}) + \delta_{\mathbf{x}}(d\mathbf{y}) \sum_{\nu \in \{-1, +1\}} \frac{1}{2} \int (1 - \alpha_{\nu}(\mathbf{x}, \mathbf{u})) Q_{\nu}(\mathbf{x}, d\mathbf{u}).$$
(2)

It can be readily verified that this Markov kernel is π -reversible by viewing $P_{\text{rev.}}$ as a MH Markov kernel with a proposal kernel given by $(1/2)Q_{-1} + (1/2)Q_{+1}$.

4 P. Gagnon and F. Maire

When there exists a boundary, meaning that $\partial X \neq \emptyset$, and it is decided to try to go in the direction ν while at some state $\mathbf{x} \in \partial X$ with $\mathbf{N}_{\nu}(\mathbf{x}) = \emptyset$, then the next state of the Markov chain is directly set to the current state \mathbf{x} . Note that in this case $\mathbf{N}_{-\nu}(\mathbf{x}) = \mathbf{N}(\mathbf{x})$. It is possible to have empty directional neighbourhoods when, for instance, X is a bounded subset of the integers (in particular, when the current state \mathbf{x} corresponds to the upper bound, $\mathbf{N}(\mathbf{x}) = \{\mathbf{x} - 1\}$ and $\mathbf{N}_{+1}(\mathbf{x}) = \emptyset$). We will use $P_{\text{rev.}}$ to denote the Markov kernel of the variant of MH in the general case, that is including the special case where $\partial X \neq \emptyset$; its general expression, including an acceptance probability similar as above, is provided in Appendix A. We can thus think of the expression in (2) as that of the Markov kernel in the special case.

Proposition 1. Under the framework presented in this subsection, $P_{rev.}$ is π -reversible.

Proof. By Lemma 2 in Appendix A, $P_{rev.}$ is the standard MH Markov kernel (in the sense of Tierney (1998)) with π as invariant distribution and proposal distribution

$$Q_{\text{rev.}}(\mathbf{x},\cdot) := \begin{cases} (1/2)\delta_{\mathbf{x}} + (1/2)Q(\mathbf{x},\cdot) & \text{if } \mathbf{x} \in \partial \mathcal{X}, \\ (1/2)Q_{-1}(\mathbf{x},\cdot) + (1/2)Q_{+1}(\mathbf{x},\cdot) & \text{otherwise.} \end{cases}$$

Thus, $P_{rev.}$ is π -reversible and has the optimal acceptance probability (in the sense of minimizing the asymptotic variances).

1.3 A lifted version

While using a notion of direction, $P_{rev.}$ does not really exploit it as it resamples the direction variable at the beginning of each iteration. We now present a *lifted* sampler which does exploit that notion. To simplify the presentation as in Section 1.2, we assume for now that $\partial X = \emptyset$. The main difference with $P_{rev.}$ is that the lifted sampler does *not* resample the direction variable at the beginning of each iteration; it keeps going in the same direction, until a rejection, upon which it starts going in the reverse direction. The lifted sampler thus operates on a *lifted* (i.e., extended) state-space $X \times \{-1, +1\}$. The direction variable *guides* the Markov chain so as to avoid backtracking, a behaviour often exhibited by MH or its variants that tends to increases the autocorrelation of the process. The lifting technique can be seen as a way to equip the resulting stochastic process with some memory of its past (the direction it comes from), while retaining the Markov property. Let us define the associated Markov kernel: for any $(\mathbf{x}, \mathbf{v}) \in \mathbf{X} \times \{-1, +1\}$,

$$P_{\text{Lifted}}((\mathbf{x},\nu), \mathbf{d}(\mathbf{y},\nu')) := Q_{\nu}(\mathbf{x}, \mathbf{d}\mathbf{y}) \,\alpha_{\nu}(\mathbf{x}, \mathbf{y}) \,\delta_{\nu}(\mathbf{d}\nu') + \delta_{(\mathbf{x},-\nu)}(\mathbf{d}(\mathbf{y},\nu')) \int (1 - \alpha_{\nu}(\mathbf{x}, \mathbf{u})) \,Q_{\nu}(\mathbf{x}, \mathbf{d}\mathbf{u}). \tag{3}$$

It can be shown that it satisfies a skewed detailed balance (Andrieu and Livingstone, 2021), implying that the product measure $\pi \otimes \mathcal{U}\{-1, +1\}$ is an invariant distribution, where $\mathcal{U}\{-1, +1\}$ denotes the uniform distribution over the set $\{-1, +1\}$. Also, it is *non-reversible*. For completeness, the definition of skewed detailed balance is presented in Appendix A.

In the case where there exists a boundary and the current state is (\mathbf{x}, ν) with $\mathbf{x} \in \partial X$ and $\mathbf{N}_{\nu}(\mathbf{x}) = \emptyset$, then, analogously to the variant of MH in Section 1.2, the next state of the Markov chain is directly set to $(\mathbf{x}, -\nu)$. We will use P_{Lifted} to denote the Markov kernel in the general case, that is including the special case where $\partial X \neq \emptyset$. The general definition is presented in Appendix A; it generalizes that in (3). We now present a proposition about P_{Lifted} in the general case whose proof can be found in Appendix A.

Proposition 2. Under the framework presented in Section 1.2, P_{Lifted} satisfies a skewed detailed balance.

Note that the definition of lifted sampler presented above is in fact specific. Andrieu and Livingstone (2021) present a more general definition in which Q_{-1} and Q_{+1} are not necessarily defined through splits of neighbourhoods as in Section 1.2; they can be any proposal kernels. In such generality, it can be shown that it is not possible to establish a comparison with the MH algorithm (see Appendix B for a demonstration). At least, it is not possible using the route proposed in Section 2 that allows for a comparison when considering the definition of Q_{-1} and Q_{+1} in Section 1.2. Note that most applications of the lifting technique in practice use this definition.

2 A comparison between the MH algorithm and its lifted version

Lifting is usually thought of as a technique that reduces the auto-correlation of the Markov chain and increase the efficiency of the MCMC estimators. Some important works indeed point in this direction, whether theoretically (Diaconis *et al.*, 2000; Andrieu and Livingstone, 2021) or empirically (Sakai and Hukushima, 2016a; Vucelja, 2016). In particular, Andrieu and Livingstone (2021) prove that lifting a one-dimensional random walk MH sampler with unbounded stationary distribution, referred to as *a guided walk* (Gustafson, 1998), can only decrease the asymptotic variances of the MCMC estimators. However, there exists some counterexamples (Gagnon and Maire, 2024, Example 1) showing that the lifting technique may increase the asymptotic variances. In this section, we show that, under the framework presented in Section 1, a general result exists. It essentially guarantees that the asymptotic variances cannot increase by a factor of more than 2, which is shown to be the strongest possible result in such generality.

Let us recall that, for an homogeneous Markov chain $\{X_k\}$ of ergodic transition kernel *P* leaving π invariant started in stationarity, the asymptotic variance of a MCMC estimator of πf , the expectation of $f(\mathbf{X})$ under $\mathbf{X} \sim \pi$, is defined as

$$\operatorname{var}(f, P) := \lim_{T \to \infty} T \operatorname{\mathbb{V}ar}\left[\frac{1}{T} \sum_{k=1}^{T} f(\mathbf{X}_0)\right] = \operatorname{\mathbb{V}ar}[f(\mathbf{X}_0)] + 2 \sum_{k=1}^{\infty} \operatorname{\mathbb{C}ov}[f(\mathbf{X}_0), f(\mathbf{X}_k)],$$
(4)

where the second equality holds provided that P is π -reversible and $\pi(f^2) < \infty$ (Häggström and Rosenthal, 2007, Theorem 4). A standard route to compare the asymptotic variances of MCMC estimators produced by two different samplers (e.g., P_{Lifted} and P_{MH}) is by establishing a Peskun-type ordering (Peskun, 1973; Tierney, 1998) between the Markov kernels. In our context, a challenge is that P_{Lifted} and P_{MH} are not defined on the same state-space and thus cannot be ordered. The strategy proposed by Andrieu and Livingstone (2021) to compare the asymptotic variances is to see P_{rev} as a Markov kernel that can also be defined on the extended state-space $X \times \{-1, +1\}$ and to consider only estimators of expectations of functions of $X \sim \pi$. When focusing on functions f of solely the first argument, extending the state-space to include the direction variable in P_{rev} is superfluous. Let us be more specific and consider such a function $f : X \times \{-1, +1\} \rightarrow \mathbb{R}$ with $f(\mathbf{x}, v) = g(\mathbf{x})$. For such a function f, the asymptotic variance of an estimator produced by P_{rev} , but when the latter is defined on the extended state-space, is equivalent to the asymptotic variance for g of an estimator produced by P_{rev} defined on X (as in Section 1.2). The kernel P_{rev} is thus used as an intermediate kernel through which comparison of the asymptotic variances of P_{MH} and P_{Lifted} is possible. With an abuse of notation, we will use f to denote both a function acting on $X \times \{-1, +1\}$ of solely the first argument and a function acting on X.

6 P. Gagnon and F. Maire

The proof of our theoretical result is thus based on establishing a Peskun-type ordering between P_{MH} and $P_{\text{rev.}}$. The fact that the MCMC estimators produced by $P_{\text{rev.}}$ cannot have smaller asymptotic variances than those produced by P_{Lifted} (Andrieu and Livingstone, 2021, Theorem 7) concludes the argument. Intuitively, this latter fact can be seen as following from the observation that $P_{\text{rev.}}$ proceeds as P_{Lifted} with only *one* difference: $P_{\text{rev.}}$ resamples the direction variable at the beginning of each iteration instead of keeping it fixed until a rejection. The Markov kernel $P_{\text{rev.}}$ is thus seen as a reversible counterpart to P_{Lifted} .

In our theoretical result, we use the notion of λ -asymptotic variance (as in Andrieu and Livingstone (2021)) defined, for an homogeneous Markov chain {**X**_{*k*}} of ergodic transition kernel *P* leaving π invariant started in stationarity, as

$$\operatorname{var}_{\lambda}(f, P) := \operatorname{\mathbb{V}ar}[f(\mathbf{X}_0)] + 2\sum_{k=1}^{\infty} \lambda^k \operatorname{\mathbb{C}ov}[f(\mathbf{X}_0), f(\mathbf{X}_k)], \quad \lambda \in [0, 1),$$

whenever $\pi(f^2) < \infty$. Note that $\lim_{\lambda \to 1} \operatorname{var}_{\lambda}(f, P) = \operatorname{var}(f, P)$ under some conditions (not requiring reversibility). A general condition is that $\sum_{k=1}^{\infty} |\mathbb{C}\operatorname{ov}[f(\mathbf{X}_0), f(\mathbf{X}_k)]| < \infty$, which can be established under fairly mild assumptions (Andrieu and Livingstone, 2021, Corollary 3). A more concrete (but stronger) condition is uniform ergodicity (Gagnon and Maire, 2024). Note that, when *P* is reversible, the limit always holds, whether $\operatorname{var}(f, P)$ is finite or not (Tierney, 1998).

We are now ready to present our result.

Theorem 1. For any $\mathbf{x} \in \mathcal{X}$ and $A \in \mathbf{X}$,

$$P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \geq \frac{1}{2} P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\}).$$

Therefore, for any f such that $\pi(f^2) < \infty$ and whenever $\lim_{\lambda \to 1} \operatorname{var}_{\lambda}(f, P_{\operatorname{Lifted}}) = \operatorname{var}(f, P_{\operatorname{Lifted}})$,

$$\operatorname{var}(f, P_{\operatorname{Lifted}}) \le \operatorname{var}(f, P_{\operatorname{rev.}}) \le 2\operatorname{var}(f, P_{\operatorname{MH}}) + \operatorname{Var}[f(\mathbf{X})], \quad \mathbf{X} \sim \pi$$

Proof. We first establish the order between P_{MH} and $P_{\text{rev.}}$. Let $A \in \mathbf{X}$ be an event. Define $B := A \setminus \{\mathbf{x}\}$, **x** being the current state. Let us consider for now the case where $\partial \mathbf{X} = \emptyset$. For any **x**, we have that

$$P_{\text{rev.}}(\mathbf{x}, B) = P_{\text{rev.}}(\mathbf{x}, B \cap \mathbf{N}(\mathbf{x}))$$

$$= \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} \frac{1}{2} Q_{\nu}(\mathbf{x}, d\mathbf{y}) \left(1 \wedge \frac{\pi(\mathbf{y}) Q_{-\nu}(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) Q_{\nu}(\mathbf{x}, \mathbf{y})} \right)$$

$$= \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} \frac{1}{2} \frac{q(\mathbf{x}, d\mathbf{y})}{c_{\nu}(\mathbf{x})} \left(1 \wedge \frac{\pi(\mathbf{y}) q(\mathbf{y}, \mathbf{x})/c_{-\nu}(\mathbf{y})}{\pi(\mathbf{x}) q(\mathbf{x}, \mathbf{y})/c_{\nu}(\mathbf{x})} \right)$$

$$= \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} \frac{q(\mathbf{x}, d\mathbf{y}) c(\mathbf{x})/2}{c(\mathbf{x}) c(\mathbf{x}) c(\mathbf{x}) c(\mathbf{x})} \left(1 \wedge \frac{\pi(\mathbf{y}) q(\mathbf{y}, \mathbf{x})/c(\mathbf{y})}{\pi(\mathbf{x}) q(\mathbf{x}, \mathbf{y})/c(\mathbf{x})} \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y}) c(\mathbf{x})/2} \right) (5)$$

$$\geq \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} \frac{q(\mathbf{x}, d\mathbf{y}) c(\mathbf{x})/2}{c(\mathbf{x}) c(\mathbf{x}) c(\mathbf{x}) c(\mathbf{x})} \left(1 \wedge \frac{\pi(\mathbf{y}) q(\mathbf{y}, \mathbf{x})/c(\mathbf{y})}{\pi(\mathbf{x}) q(\mathbf{x}, \mathbf{y})/c(\mathbf{x})} \right) \left(1 \wedge \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y}) c(\mathbf{x})/2} \right)$$

$$= \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} Q(\mathbf{x}, d\mathbf{y}) \alpha(\mathbf{x}, \mathbf{y}) \left(\frac{c(\mathbf{x})/2}{c_{\nu}(\mathbf{x})} \wedge \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} \right)$$

$$(6)$$

$$\geq \frac{1}{2} \sum_{\nu \in \{-1,+1\}} \int_{B \cap \mathbf{N}_{\nu}(\mathbf{x})} Q(\mathbf{x}, d\mathbf{y}) \alpha(\mathbf{x}, \mathbf{y}) = \frac{1}{2} P_{\mathrm{MH}}(\mathbf{x}, B \cap \mathbf{N}(\mathbf{x})) = \frac{1}{2} P_{\mathrm{MH}}(\mathbf{x}, B),$$

using in the first inequality that

$$\frac{\pi(\mathbf{y}) q(\mathbf{y}, \mathbf{x})/c(\mathbf{y})}{\pi(\mathbf{x}) q(\mathbf{x}, \mathbf{y})/c(\mathbf{x})} \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} \frac{c_{\nu}(\mathbf{x})}{c(\mathbf{x})/2} \ge \frac{\pi(\mathbf{y}) q(\mathbf{y}, \mathbf{x})/c(\mathbf{y})}{\pi(\mathbf{x}) q(\mathbf{x}, \mathbf{y})/c(\mathbf{x})} \left(1 \wedge \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} \frac{c_{\nu}(\mathbf{x})}{c(\mathbf{x})/2} \right),$$

and that, for any a > 0 and $b \in (0, 1]$, we have $1 \land ab \ge b(1 \land a)$; in the second inequality, we used that

$$\frac{c(\mathbf{x})/2}{c_{\nu}(\mathbf{x})} \wedge \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} = \frac{c_{\nu}(\mathbf{x})/2 + c_{-\nu}(\mathbf{x})/2}{c_{\nu}(\mathbf{x})} \wedge \frac{c_{\nu}(\mathbf{y})/2 + c_{-\nu}(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} = \frac{1}{2} + \frac{1}{2} \left(\frac{c_{-\nu}(\mathbf{x})}{c_{\nu}(\mathbf{x})} \wedge \frac{c_{\nu}(\mathbf{y})}{c_{-\nu}(\mathbf{y})} \right) \ge \frac{1}{2}.$$

In the case where $\partial X \neq \emptyset$, we can obtain the same bound using the same strategy as above, but we have to use the definition of the proposal distribution given at the end of Section 1.2 and that of the acceptance probability given in Appendix A. For instance, if $\mathbf{x} \notin \partial X$, we can proceed exactly as above, but it is possible that, while integrating over $B \cap \mathbf{N}_{\nu}(\mathbf{x})$, we obtain $\mathbf{y} \in \partial X$ in a certain area. For such a value of $\mathbf{y}, Q_{-\nu}(\mathbf{y}, \cdot) = Q(\mathbf{y}, \cdot)$ and

$$\frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})} = \frac{1}{2}$$

If $\mathbf{x} \in \partial X$, there is one term in the sum in the mathematical development above (that with v such that $\mathbf{N}_{v}(\mathbf{x}) = \mathbf{N}(\mathbf{x})$), and, while integrating over $B \cap \mathbf{N}_{v}(\mathbf{x})$, we have to separate the cases where $\mathbf{y} \notin \partial X$ to those where $\mathbf{y} \in \partial X$.

From the order between $P_{\rm MH}$ and $P_{\rm rev.}$, we can deduce that

$$\operatorname{var}(f, P_{\operatorname{rev.}}) \leq 2\operatorname{var}(f, P_{\operatorname{MH}}) + \operatorname{Var}[f(\mathbf{X})], \quad \mathbf{X} \sim \pi,$$

using Lemma 33 in Andrieu et al. (2018). Theorem 7 in Andrieu and Livingstone (2021) states that

$$\operatorname{var}_{\lambda}(f, P_{\operatorname{Lifted}}) \leq \operatorname{var}_{\lambda}(f, P_{\operatorname{rev.}}).$$

The proof is concluded by taking the limit $\lambda \to 1$ on both side of this inequality.

We now make a few remarks about Theorem 1. Firstly, the result reflects a notion of *universality*: in addition to being stated in a context of general state-spaces, it is valid for *any* distribution π , *any* proposal mechanism (represented by the unnormalized density $q(\mathbf{x}, \cdot)$), and for *any* way of inducing directions (through the definition of $\{\mathbf{N}_{-1}(\mathbf{x})\}$ and $\{\mathbf{N}_{+1}(\mathbf{x})\}$).

Secondly, we observe in the proof that the factor

$$\left(\frac{c(\mathbf{x})/2}{c_{\nu}(\mathbf{x})} \wedge \frac{c(\mathbf{y})/2}{c_{-\nu}(\mathbf{y})}\right)$$

in (6) is an explanation of the difference between $P_{\text{rev.}}$ and P_{MH} . It is a measure of whether the directional neighbourhoods $\mathbf{N}_{-1}(\mathbf{x})$ and $\mathbf{N}_{+1}(\mathbf{x})$ (and $\mathbf{N}_{-1}(\mathbf{y})$ and $\mathbf{N}_{+1}(\mathbf{y})$) are well balanced or not. When $c_{\nu}(\mathbf{x}) = c_{-\nu}(\mathbf{x})$ and $c_{\nu}(\mathbf{y}) = c_{-\nu}(\mathbf{y})$ for all $\mathbf{y} \in B \cap \mathbf{N}_{\nu}(\mathbf{x})$ and $\nu \in \{-1, +1\}$, that factor is always equal to 1 in the integral in (6). Therefore, under this condition, we can conclude that $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$ from (6). When this condition holds for all $\mathbf{x} \in X$ and $A \in \mathbf{X}$, we say that we are in a situation of *perfect balance* and we have a guarantee that the lifted sampler outperforms the MH one, that is $\operatorname{var}(f, P_{\text{Lifted}}) \le \operatorname{var}(f, P_{\text{MH}})$ whenever $\lim_{\lambda \to 1} \operatorname{var}_{\lambda}(f, P_{\text{Lifted}}) = \operatorname{var}(f, P_{\text{Lifted}})$. It is for instance the case with the guided walk, by symmetry of the proposal distribution in the random walk MH (implying that $c_{\nu}(\mathbf{x}) = c_{-\nu}(\mathbf{x}) = 1/2$ for all $\mathbf{x} \in X$). Note that before introducing the first inequality in the mathematical development in the proof, we had that $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) = P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$ under the condition

 $c_{\nu}(\mathbf{x}) = c_{-\nu}(\mathbf{x})$ and $c_{\nu}(\mathbf{y}) = c_{-\nu}(\mathbf{y})$ for all $\mathbf{y} \in B \cap \mathbf{N}_{\nu}(\mathbf{x})$ and $\nu \in \{-1, +1\}$ (see (5)). This indicates that the first inequality in the mathematical development in the proof is tight because, under this condition, the inequality yields $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$ but we know that $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) = P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$.

Thirdly, in a context that fits within the framework of Theorem 1, Gagnon and Maire (2024) prove that $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\})/P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \rightarrow 1/2$ for some \mathbf{x} and A, as a dimension parameter grows without bounds (Proposition 2 in that paper). We can therefore conclude that our result is (essentially) optimal, in the sense that it is (essentially) not possible to obtain a better bound without additional assumptions.

Finally, in a context of comparison between two samplers, the magnitude of the asymptotic variances, which is proportional to $Var[f(\mathbf{X})]$, is irrelevant. Thus, it is appropriate to consider standardized functions. When $Var[f(\mathbf{X})] = 1$, the expression in (4) corresponds to the integrated autocorrelation time. In this case, the factor 2 and the additive term in the bound on the asymptotic variances (and thus on the integrated autocorrelation times) are independent of all problem parameters, whether it is the dimension of the state-space, the sample size in Bayesian statistics contexts, etc. This implies that the lifted sampler is *at worst* comparable to the MH algorithm, in the sense that the asymptotic variances can be smaller, but at worst, the factor 2 and the additive term explaining the difference between the asymptotic variances do not deteriorate when a problem parameter changes. All that suggests the following practical suggestion: *when one as a way of inducing directions which does not significantly increase the computational cost of the algorithm, better do it as there is not much to lose, but there is potentially a lot to gain (as observed many times in the past).*

3 Examples

In this section, we illustrate Theorem 1 by presenting empirical results and instances of lifted samplers in two concrete situations: one where the state-space is discrete and partially ordered (Section 3.1), and one where the state-space is the real numbers, and thus, totally ordered (Section 3.2). We use this illustration to highlight that, even in situations where it is suspected that the lifted sampler will not offer a great performance, the asymptotic variances are at worst comparable to those of MH, as guaranteed by Theorem 1. In particular, we show that, in very unfavourable situations, the upper bound in Theorem 1 is attained. The code to reproduce our numerical results is available online (see ancillary files on the arXiv page of the paper).

3.1 Example 1: Partially-ordered discrete state-space

In this example, we consider that X is a finite state-space which admits a partial order. This partial order will be exploited to induce directions to follow by the lifted sampler. We consider more precisely that $\mathbf{x} = (x_1, \ldots, x_n)$, where each component x_i can be of two types. Many statistical contexts fit within this framework, such as the modelling of binary data using networks or graphs and in variable selection. Indeed, for the former, X can be parameterized such that $X = \{-1, +1\}^n$, where for example for an Ising model, $x_i \in \{-1, +1\}$ represents the state of a spin. For variable selection, $X = \{0, 1\}^n$ and $x_i \in \{0, 1\}$ indicates whether or not the *i*-th covariate is included in the model employed.

The use of lifted algorithms to sample from distributions defined on partially-ordered discrete statespaces has been thoroughly studied in Gagnon and Maire (2024). In particular, the Ising model is studied. We here provide numerical results that are complementary to those presented there. The specific model that we study is the two-dimensional Ising model. For this model, the state-space (V_{η}, E_{η}) is a $\eta \times \eta$ square lattice regarded here as a square matrix in which each element takes either the value -1 or +1. We write each state as a vector as above: $\mathbf{x} = (x_1, \dots, x_n)$, where $n = \eta^2$. The states can be encoded as follows: the values of the components on the first line are x_1, \dots, x_η , those on the second line $x_{\eta+1}, \dots, x_{2\eta}$, and so on. The probability mass function is given by

$$\pi(\mathbf{x}) = \frac{1}{Z} \exp\left(\sum_{i} \alpha_{i} x_{i} + \lambda \sum_{\langle ij \rangle} x_{i} x_{j}\right),$$

where $\alpha_1, \ldots, \alpha_n \in \mathbb{R}$ and $\lambda > 0$ are fixed parameters, *Z* is the normalizing constant and the notation $\langle ij \rangle$ indicates that sites *i* and *j* are nearest neighbours. The notion of neighbourhood on (V_η, E_η) should not be confused with that on *X* on which the samplers rely. The neighbourhood of a site $i \in V_\eta$ comprises, when they exist, its North-South-East-West neighbours on the lattice. The parameter $\alpha := (\alpha_1, \ldots, \alpha_n)$ is often referred to as the *external field* and λ plays the role of a *spatial correlation parameter*. The external field tends to polarize each spin, while the spatial correlation tends to create patches of identical spin states.

To simulate an Ising model, the MH algorithm typically proceeds by proposing to flip a single bit at each iteration. More formally, the neighbourhood structure $\{N(x)\}$ is given by $N(x) = \{y \in X : \sum_i |x_i - y_i| = 2\}$. A uniform proposal distribution is often employed. This strategy has been shown to be often inefficient in Zanella (2020), who proposed a generic approach to sample from distributions defined on discrete state-spaces. The approach leverages local target-distribution information and is thus less naive than a uniform approach. It possesses appealing properties and is referred to as *locally balanced*. The term *locally balanced* comes form the fact that, in the limit, when the state-space becomes larger and larger (but the neighbourhoods have a fixed size and proposed moves are thus local), there is no need for an accept-reject step anymore; the proposal distribution leaves the distribution π invariant. The locally-balanced proposal distribution that we use in this example is the *Barker proposal distribution*, with

$$q(\mathbf{x}, \mathbf{y}) = \frac{\pi(\mathbf{y})/\pi(\mathbf{x})}{1 + \pi(\mathbf{y})/\pi(\mathbf{x})},$$

for $\mathbf{x} \in X$ and \mathbf{y} in N(x), where the name is in reference to Barker (1965)'s acceptance probability choice.

To explicitly define the directional neighbourhoods that are used in the lifted sampler, we leverage that X admits a partial order. Indeed, an inclusion-based partial order on X can be defined through a set:

$$\mathcal{R} := \{ (\mathbf{x}, \mathbf{y}) \in \mathcal{X} \times \mathcal{X} : \{ i : x_i = +1 \} \subset \{ i : y_i = +1 \} \}.$$

Pairs $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$ with $\mathbf{x} \neq \mathbf{y}$ are said to be *comparable* when either $(\mathbf{x}, \mathbf{y}) \in \mathcal{R}$ or $(\mathbf{y}, \mathbf{x}) \in \mathcal{R}$ and are said *incomparable* otherwise. The existence of incomparable pairs (\mathbf{x}, \mathbf{y}) with $\mathbf{x} \neq \mathbf{y}$ represents the difference with a totally-ordered set such as \mathbb{N} or \mathbb{R} in which every pair of different elements is comparable. We denote $\mathbf{x} \prec \mathbf{y}$ whenever $(\mathbf{x}, \mathbf{y}) \in \mathcal{R}$ and $\mathbf{x} \neq \mathbf{y}$. Hence, setting $\mathbf{N}(\mathbf{x}) = \{\mathbf{y} \in \mathcal{X}_n : \sum_i |x_i - y_i| = 2\}$ yields neighbourhoods with states \mathbf{y} that are all comparable with \mathbf{x} . For the lifter sampler, we can set $\mathbf{N}_{-1}(\mathbf{x}) := \{\mathbf{y} \in \mathbf{N}(\mathbf{x}) : \mathbf{y} \prec \mathbf{x}\}$ and $\mathbf{N}_{+1}(\mathbf{x}) := \{\mathbf{y} \in \mathbf{N}(\mathbf{x}) : \mathbf{x} \prec \mathbf{y}\}$. States \mathbf{y} in $\mathbf{N}_{+1}(\mathbf{x})$, for instance, are equal to \mathbf{x} except for one component which is flipped from -1 to +1.

We are now ready to present simulation results. They are about a common problem in statistical physics which is to estimate the average *magnetisation* of an Ising model, the magnetisation being defined as the mapping $\mathbf{x} \mapsto \sum_{i=1}^{n} x_i$. The numerical results that we present are more specifically about the asymptotic variances of estimators of the expectation of the standardized version of the magnetisation. We consider a base target distribution for which $n = 10^2$ and the spatial correlation is moderate and

more precisely $\lambda = 0.5$. We set the α_i as follows: $\alpha_i = -\mu + \epsilon_i$ if the column index is smaller than or equal to $\lfloor \eta/2 \rfloor$ and $\alpha_i = \mu + \epsilon_i$ otherwise, where $\mu = 1$, the ϵ_i are independent uniform random variables on the interval (-0.1, +0.1) and $\lfloor \cdot \rfloor$ is the floor function. We can thus think of the external field as a matrix with negative values on the left, and positive ones on the right. The base target is moderately rough, in the sense that it concentrates on a subset of the state-space, but with directional neighbourhoods that have a smoothly varying mass on this subset (implying similar $c_{\nu}(\mathbf{x})$ and $c_{-\nu}(\mathbf{x})$).

The base target distribution described above is favourable for lifted samplers with locally-balanced proposal distributions which leverage variations in the target distribution and create persistent movement (as seen in Figure 1 by looking at the starting points on the left of the plots). In Figure 1 (a), we observe the impact of dealing with larger systems for a moderately rough target, by increasing η from 10 to 50. Increasing η in this case leads to longer paths along which the state-space can be explored, which is again favourable for lifted samplers. In Figure 1 (a), we observe that the lifted sampler scales better with the system size than the MH algorithm. When comparing the latter with the reversible counterpart to the lifted sampler, we observe a stable absolute difference of asymptotic variances, which translates into a diminishing relative difference. With Figure 1 (b), we consider an opposite situation which is increasingly unfavourable for lifted samplers, with a fixed system size of $\eta = 10$ but an increasing μ yielding an increasingly rougher target distribution with an increasing level of concentration. As the level of concentration increases, the mass of the directional neighbourhoods becomes increasingly variable (implying increasingly different $c_{\nu}(\mathbf{x})$ and $c_{-\nu}(\mathbf{x})$). Also, when the mass is concentrated on few configurations, it leaves not much room for persistent movement for the lifted sampler, and it thus loses its advantage. Such a situation corresponds to one where it is suspected that the lifted sampler will not offer a great performance. When the lifted sampler is at the mode and leaves it, it "wastes" an iteration because it tries continuing in the same direction whereas the MH sampler has the possibility to return to the mode the following iteration. In Figure 1 (b), we also observe an increasing difference in asymptotic variance between $P_{rev.}$ and P_{MH} , with that of $P_{rev.}$ reaching the upper bound provided by Theorem 1 of $2var(f, P_{MH}) + 1$. Note that the numerical results are based on 100 independent runs of 1,000,000 iterations for each algorithm and each value of η and μ , with burn-ins of 100,000.



Figure 1. Asymptotic variance of the MCMC estimator of the standardized version of the mapping $\mathbf{x} \mapsto \sum_{i=1}^{n} x_i$ for the MH algorithm, the lifted sampler and the reversible counterpart of the latter, all using the Barker proposal distribution when: (a) η increases from 10 to 50 and the other parameters are kept fixed ($\mu = 1$ and $\lambda = 0.5$); (b) μ increases from 1 to 3.5 and the other parameters are kept fixed ($\eta = 10$ and $\lambda = 0.5$); in (b), the upper bound provided by Theorem 1 is also presented.

3.2 Example 2: Totally-ordered continuous state-space

In this section, we study the simple case where $X = \mathbb{R}$. When the MH sampler is the random walk Metropolis algorithm, the lifted version described in Section 1.3 corresponds to the *guided walk* (Gustafson, 1998). The latter has essentially the same computation cost as the random walk Metropolis algorithm, and has been proved to outperform the latter for any target distribution π (Andrieu and Livingstone, 2021, Theorem 7). This is a consequence of the symmetry of the normal distribution used in the random walk Metropolis. This symmetry leads to a situation of perfect balance, as described in Section 2, where, $c_v(\mathbf{x}) = c_{-v}(\mathbf{x}) = 1/2$, for all $\mathbf{x} \in \mathbb{R}$.

But what if the proposal distribution is not symmetrical? This is the case that we now study. In this case, one may suspect that the lifted sampler will not offer a great performance because of possible significant differences between $c_{\nu}(\mathbf{x})$ and $c_{-\nu}(\mathbf{x})$. In our example, we consider that the proposal distribution is the continuous version of the Barker proposal distribution described in Section 3.1. It has recently been proposed in Livingstone and Zanella (2022) and has the following density:

$$Q(\mathbf{x}, \mathbf{y}) = 2 \frac{\varphi_{\sigma}(\mathbf{y} - \mathbf{x})}{1 + \exp\left(-(\mathbf{y} - \mathbf{x})\nabla\log\pi(\mathbf{x})\right)}, \quad \mathbf{x} \in \mathbb{R}, \mathbf{y} \in \mathbf{N}(\mathbf{x}) = \mathbb{R} \setminus \{\mathbf{x}\},$$

where φ_{σ} is the density of a normal distribution with a mean of 0 and a standard deviation of $\sigma > 0$, and $\nabla \log \pi$ is the gradient of the log-target density. The gradient injects skewness: the more substantial is $\nabla \log \pi(\mathbf{x})$, the more asymmetrical is the density $Q(\mathbf{x}, \cdot)$. The scale parameter σ is a tuning parameter.

We study the natural case where the target distribution is a standard normal. We present empirical results in Table 1 and Figure 2. The directional neighbourhoods are given by $N_{-1}(x) = \{y : y < x\}$ and $N_{+1}(x) = \{y : y > x\}$. In Table 1, we present results for the MH sampler, its lifted version and the reversible counterpart of the latter. We more precisely present acceptance rates and asymptotic variances of the MCMC estimators of πf when f is the identity mapping $\mathbf{x} \mapsto \mathbf{x}$, for different values of σ . Note that the identity mapping is already standardized given that the target distribution is a standard normal. The values of σ in Table 1 represent optimal values for the different samplers (at least according to our grid search). For the MH algorithm, the optimal value is 2.5, associated with an acceptance rate of 62%. When using this algorithm, the asymptotic variance of the MCMC estimator of the mean is 1.94. The lifted sampler suffers from instability among directional-neighbourhood mass $c_{\nu}(\mathbf{x})$ and $c_{-\nu}(\mathbf{x})$. To compensate for smaller acceptance rates, σ has to be reduced. The optimal value for this sampler is 2.0 and is associated with an acceptance rate of 46%. When using this algorithm, the asymptotic variance of the MCMC estimator of the mean is 2.31. We observe in Figure 2 that the reason why the lifted sampler manages to offer a performance that is not so bad (despite significantly smaller σ and acceptance rate) is persistent movement. Nevertheless, the impact of persistent movement is not significant enough. In addition, the computational cost associated to sampling from conditional distributions $Q_{\nu}(\mathbf{x}, \cdot)$ and computing the normalizing constant of the latter is quite high in this case: runtime is about 130 times larger.¹ Regarding the reversible counterpart to the lifted sampler, the optimal value of σ is 2.2, which is in between those of the two other samplers. When using this value, the asymptotic variance is 4.08, which is not so far below the bound provided by Theorem 1 of 4.99. The smallest difference between the bound and the asymptotic variance among those calculated from Table 1 is 0.76 when $\sigma = 2.5$.

¹We do not claim optimality of our implementation, but we employed commonly used techniques for sampling from the conditional distributions $Q_{\nu}(\mathbf{x}, \cdot)$ and computing the normalizing constant of the latter.

	MH sampler		Lifted sampler		Reversible counterpart		Upper bound
	Acc. rate	Asymp. var.	Acc. rate	Asymp. var.	Acc. rate	Asymp. var.	Asymp. var.
$\sigma = 2.0$	71%	2.10	46%	2.31	46%	4.17	5.20
$\sigma = 2.2$	67%	2.00	43%	2.35	43%	4.08	4.99
$\sigma = 2.5$	62%	1.94	38%	2.47	38%	4.13	4.89

Table 1. Acceptance rates and asymptotic variances for a MH sampler with a Barker proposal distribution, its lifted version and the reversible counterpart of the latter, along with the upper bound on the asymptotic variances provided by Theorem 1, for different values of σ .



Figure 2. Trace plots for a MH algorithm on the left panel and a lifted sampler on the right panel, both with a Barker proposal distribution.

4 Generalized MH algorithm and its lifted version: A comparison

In Section 4.1, we present a generalized version of the MH algorithm. It essentially includes any π -reversible MCMC algorithm which proceeds with a proposal distribution $Q(\mathbf{x}, \cdot)$ and an accept/reject step; π and $Q(\mathbf{x}, \cdot)$ thus do not necessarily admit densities with respect to a common dominating measure. This represents a significant difference, both conceptually and technically, to the special case where π and $Q(\mathbf{x}, \cdot)$ do admit densities with respect to a common dominating measure treated in detail in Section 1.2. As mentioned, additionally to the usual MH algorithm presented in Section 1.2, this framework includes MCMC algorithms such as reversible jump, Multiple-try Metropolis and pseudomarginal. For the presentation of the generalized MH algorithm, we follow Tierney (1998). In Section 4.2, we present the generalized auxiliary sampler through which the lifted one is defined. In Section 4.3, we present a generalized version of Theorem 1. The proofs of all theoretical results presented in this section are deferred to Appendix A.

4.1 Generalized MH algorithm

As before, we consider that π , the target, is a probability distribution on (X, X), and that $\{Q(\mathbf{x}, \cdot) : \mathbf{x} \in X\}$ is a collection of (proposal) probability distributions on (X, X), however, here the support of π and $Q(\mathbf{x}, \cdot)$ is left unspecified. Let μ and μ^T be two probability measures on the product space $(X^2, X \otimes X)$

13

defined as:

$$\mu(A \times B) = \int_{A} \pi(\mathbf{d}\mathbf{x}) Q(\mathbf{x}, B), \qquad \mu^{T}(A \times B) = \int_{B} \pi(\mathbf{d}\mathbf{x}) Q(\mathbf{x}, A).$$

Also, let $v := \mu + \mu^T$. From these definitions, we conclude that both μ and μ^T are absolutely continuous with respect to v and that, from the Radon–Nikodym theorem, there exist two (essentially unique) measurable functions $h : \mathcal{X}^2 \to [0, \infty)$ and $h^T : \mathcal{X}^2 \to [0, \infty)$ such that

$$A \in \mathbf{X}, \quad \mu(A) = \int_A h(\mathbf{x}, \mathbf{y}) \, \nu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}), \quad \mu^T(A) = \int_A h^T(\mathbf{x}, \mathbf{y}) \, \nu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}),$$

with the usual notation $d\mu/d\nu := h$ and $d\mu^T/d\nu := h^T$. As observed by Tierney (1998), $h^T(\mathbf{x}, \mathbf{y}) = h(\mathbf{y}, \mathbf{x})$ for all (\mathbf{x}, \mathbf{y}) given that $\nu(d\mathbf{x}, d\mathbf{y}) = \nu(d\mathbf{y}, d\mathbf{x})$. We also define the set

$$R := \{ (\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : h(\mathbf{x}, \mathbf{y}) > 0 \text{ and } h(\mathbf{y}, \mathbf{x}) > 0 \},\$$

which is symmetric given that $(\mathbf{x}, \mathbf{y}) \in R$ if and only if $(\mathbf{y}, \mathbf{x}) \in R$. Finally, we define a function $r : R \to (0, \infty)$ that verifies

$$r(\mathbf{x}, \mathbf{y}) = \frac{\mathrm{d}\mu^T}{\mathrm{d}\mu}(\mathbf{x}, \mathbf{y}) = \frac{\mathrm{d}\mu^T/\mathrm{d}\nu(\mathbf{x}, \mathbf{y})}{\mathrm{d}\mu/\mathrm{d}\nu(\mathbf{x}, \mathbf{y})} = \frac{h(\mathbf{y}, \mathbf{x})}{h(\mathbf{x}, \mathbf{y})}, \quad \text{for all } (\mathbf{x}, \mathbf{y}) \in R,$$
(7)

meaning that for any bounded measurable function $\phi : \mathcal{X}^2 \to \mathbb{R}$,

$$\int_R \phi \, r \, \mathrm{d}\mu = \int_R \phi \, \mathrm{d}\mu^T.$$

The (generalized) MH kernel, $P_{\rm MH}$, is defined for all $\mathbf{x} \in \boldsymbol{X}$ as

$$P_{\rm MH}(\mathbf{x}, \mathbf{dy}) = Q(\mathbf{x}, \mathbf{dy}) \,\alpha(\mathbf{x}, \mathbf{y}) + \delta_{\mathbf{x}}(\mathbf{dy}) \int (1 - \alpha(\mathbf{x}, \mathbf{u})) \,Q(\mathbf{x}, \mathbf{du}),$$

where $\alpha : \mathcal{X}^2 \to [0, 1]$; note that we reused the same notation as in Section 1.2 to simplify. Theorem 2 in Tierney (1998) states that P_{MH} is π -reversible if and only if the function α verifies: $\alpha(\mathbf{x}, \mathbf{y}) = 0$ if $(\mathbf{x}, \mathbf{y}) \notin R$, and

$$\alpha(\mathbf{x}, \mathbf{y}) = r(\mathbf{x}, \mathbf{y}) \,\alpha(\mathbf{y}, \mathbf{x}) \quad \text{if} \quad (\mathbf{x}, \mathbf{y}) \in R.$$
(8)

We restrict our attention here to the case where α is defined as follows:

$$\alpha(\mathbf{x}, \mathbf{y}) = \begin{cases} \varphi(r(\mathbf{x}, \mathbf{y})) & \text{if } (\mathbf{x}, \mathbf{y}) \in R, \\ 0 & \text{otherwise,} \end{cases}$$
(9)

with $\varphi : (0, \infty) \to [0, 1]$ verifying $\varphi(r) = r\varphi(1/r)$ for all $r \in (0, \infty)$. Such functions φ appeared in several sampling contexts, for instance as efficient *balancing functions* in Zanella (2020). In our context, this class of functions allows for the usual acceptance probability $\varphi(r) = 1 \wedge r$ and Barker's acceptance probability $\varphi(r) = r/(1 + r)$ (Barker, 1965). If we assume that, on *R*, the function α in (8) must involve $r(\mathbf{x}, \mathbf{y})$ in some way, then the special case with $\alpha(\mathbf{x}, \mathbf{y}) = \varphi(r(\mathbf{x}, \mathbf{y}))$ represents a natural one. Indeed, if we consider that $r : \mathbf{X}^2 \to (0, \infty)$ (instead of $r : R \to (0, \infty)$), defined as in (7) on *R* and equal to 1 otherwise, then we can consider that α is a function with a third argument, that is $\alpha : \mathbf{X}^2 \times (0, \infty) \to [0, 1]$, and such that $\alpha(\mathbf{x}, \mathbf{y}, r(\mathbf{x}, \mathbf{y})) = 0$ if $(\mathbf{x}, \mathbf{y}) \notin R$, and

$$\alpha(\mathbf{x}, \mathbf{y}, r(\mathbf{x}, \mathbf{y})) = r(\mathbf{x}, \mathbf{y}) \, \alpha(\mathbf{y}, \mathbf{x}, r(\mathbf{y}, \mathbf{x})) = r(\mathbf{x}, \mathbf{y}) \, \alpha(\mathbf{y}, \mathbf{x}, 1/r(\mathbf{x}, \mathbf{y})) \quad \text{if} \quad (\mathbf{x}, \mathbf{y}) \in R.$$

From this perspective, the special case considered in (9) with $\varphi(r) = r\varphi(1/r)$ corresponds to that where $\alpha(\mathbf{x}, \mathbf{y}, r(\mathbf{x}, \mathbf{y})) = \varphi(r(\mathbf{x}, \mathbf{y}))$ if $(\mathbf{x}, \mathbf{y}) \in R$, and thus to a function that does not use its first two arguments **x** and **y**, but only its third one $r(\mathbf{x}, \mathbf{y})$ which can be seen as a *summary statistic*.

Such a function φ has interesting properties, as stated in Lemma 1, which will be useful to prove a generalized version of Theorem 1 in Section 4.3.

Lemma 1. For any function $\varphi : (0, \infty) \to [0, 1]$ verifying $\varphi(r) = r\varphi(1/r)$ for all r, we have that:

- 1. φ is non-decreasing,
- 2. φ is continuous on $(0, \infty)$,
- *3.* $\varphi(r) \downarrow 0$ when $r \downarrow 0$,
- 4. φ is either a null function, that is $\varphi(r) = 0$ for all r, or it is a positive function,
- 5. for any $a > 0, b > 0, \varphi(ab) \ge \varphi(a)\varphi(b)$,
- 6. for any a > 0, $b \in (0, 1]$, $\varphi(ab) \ge b\varphi(a)$.

What can be difficult in practice to implement a generalized MH algorithm is the identification of explicit forms for the function r and the set R (when available). The results in Andrieu *et al.* (2020) can help in this task.

4.2 Generalized auxiliary sampler and the lifted one

We now present the generalized auxiliary sampler, constructed from the generalized MH algorithm presented in Section 4.1. The generalized auxiliary sampler will in turn be used to construct the generalized lifted sampler. The presentation of the generalized auxiliary and lifted samplers consists essentially of definitions; it will allow to understand how a lifted sampler can be constructed under the general framework of Section 4.1.

For each $x \in X$, we define two measurable sets $N_{-1}(x)$ and $N_{+1}(x)$. The collection of sets is assumed to satisfy the following properties:

- for any $\mathbf{x} \in \mathcal{X}$, $Q(\mathbf{x}, A) = Q(\mathbf{x}, A \cap \mathbf{N}_{-1}(\mathbf{x})) + Q(\mathbf{x}, A \cap \mathbf{N}_{+1}(\mathbf{x}))$ for all $A \in \mathbf{X}$,
- for any $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$ and $\nu \in \{-1, +1\}$, $\mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})$ *if and only if* $\mathbf{x} \in \mathbf{N}_{-\nu}(\mathbf{y})$, and for such a related couple (\mathbf{x}, \mathbf{y}) , $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) > 0$ and $Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) > 0$.

The sets $\mathbf{N}_{-1}(\mathbf{x})$ and $\mathbf{N}_{+1}(\mathbf{x})$ can be constructed as in Section 1.2 by splitting into two the support of $Q(\mathbf{x}, \cdot)$ that can be seen as defining a neighbourhood $\mathbf{N}(\mathbf{x})$. From the properties of the directional neighbourhoods $\{\mathbf{N}_{-1}(\mathbf{x})\}$ and $\{\mathbf{N}_{+1}(\mathbf{x})\}$, we have that if $\mathbf{x} \in \mathbf{N}_{\nu}(\mathbf{x})$ then $\mathbf{x} \in \mathbf{N}_{-\nu}(\mathbf{x})$, and from this, we can deduce that $Q(\mathbf{x}, \{\mathbf{x}\}) = 2Q(\mathbf{x}, \{\mathbf{x}\}) = 0$ for all \mathbf{x} . We now define a notion of *interior* of \mathcal{X} (that depends on the choice of directional neighbourhoods $\{\mathbf{N}_{-1}(\mathbf{x})\}$ and $\{\mathbf{N}_{+1}(\mathbf{x})\}$) as $\mathring{\mathcal{X}} := \{\mathbf{x} \in \mathcal{X} : Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}))Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) > 0\}$. For instance, $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}))Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 0$ when one of the two directional neighbourhoods is empty. Let $\partial \mathcal{X} := \mathcal{X} \setminus \mathring{\mathcal{X}}$. For all $\mathbf{x} \in \mathring{\mathcal{X}}$ and $\nu \in \{-1, +1\}$, we denote the conditional proposal distribution given that the proposal belongs to $\mathbf{N}_{\nu}(\mathbf{x})$ as $Q_{\nu}(\mathbf{x}, \cdot)$ which is such that

$$Q_{\nu}(\mathbf{x},A) = \frac{Q(\mathbf{x},A \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x},\mathbf{N}_{\nu}(\mathbf{x}))}, \quad A \in \mathbf{X}.$$

Note that, when $\mathbf{x} \in \partial X$, there exists $\nu \in \{-1, +1\}$ such that $Q(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x})) = 0$, which implies that $Q(\mathbf{x}, X) = Q(\mathbf{x}, X \cap \mathbf{N}_{\nu}(\mathbf{x})) = Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1$ and that $Q(\mathbf{x}, \cdot) = Q_{\nu}(\mathbf{x}, \cdot)$.

We define

$$Q_{\text{rev.}}(\mathbf{x}, \cdot) := \begin{cases} (1/2)Q_{-1}(\mathbf{x}, \cdot) + (1/2)Q_{+1}(\mathbf{x}, \cdot) & \text{if } \mathbf{x} \in \mathring{X}, \\ (1/2)\delta_{\mathbf{x}} + (1/2)Q(\mathbf{x}, \cdot) & \text{otherwise.} \end{cases}$$

Similarly as in Section 4.1, we define

$$\mu_{\text{rev.}}(A \times B) := \int_A \pi(\mathrm{d}\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, B) \,, \qquad \mu_{\text{rev.}}^T(A \times B) := \int_B \pi(\mathrm{d}\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, A).$$

We also define the following sets: $R_{\nu} := \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})\}$ for $\nu \in \{-1, +1\}$, and $\Delta := \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{x} = \mathbf{y}\}$. Finally, we introduce a function $\beta : \mathcal{X}^2 \to [0, \infty)$:

$$\beta(\mathbf{x}, \mathbf{y}) = \begin{cases} 0 & \text{if } (\mathbf{x}, \mathbf{y}) \in R^{c}, \\ Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))/Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) & \text{if } (\mathbf{x}, \mathbf{y}) \in R_{\nu} \cap R^{c}_{-\nu} \cap \Delta^{c} \cap R \text{ for some } \nu \in \{-1, +1\}, \\ 1 & \text{otherwise.} \end{cases}$$

We now present a result which will allow to define a valid acceptance probability for a sampler using Q_{rev} as proposal kernel.

Proposition 3. *The restriction of the measures* $\mu_{\text{rev.}}$ *and* $\mu_{\text{rev.}}^T$ *to R are mutually absolutely continuous and, for all* $(\mathbf{x}, \mathbf{y}) \in R$ *,*

$$r_{\text{rev.}}(\mathbf{x},\mathbf{y}) := \frac{\mathrm{d}\mu_{\text{rev.}}^T}{\mathrm{d}\mu_{\text{rev.}}}(\mathbf{x},\mathbf{y}) = r(\mathbf{x},\mathbf{y})\beta(\mathbf{x},\mathbf{y}).$$

Following Tierney (1998), this result allows to state that $P_{rev.}$ defined as

$$P_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) = Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \,\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) + \delta_{\mathbf{x}}(d\mathbf{y}) \int (1 - \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{u})) \,Q_{\text{rev.}}(\mathbf{x}, d\mathbf{u}), \quad \text{for all } \mathbf{x} \in \mathcal{X},$$

with the function $\alpha_{rev.}$ defined as

$$(\mathbf{x}, \mathbf{y}) \in \boldsymbol{X}^2, \quad \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = \begin{cases} \varphi(r_{\text{rev.}}(\mathbf{x}, \mathbf{y})) & \text{if } (\mathbf{x}, \mathbf{y}) \in R, \\ 0 & \text{otherwise,} \end{cases}$$

is in fact a generalized MH kernel, but one that uses $Q_{rev.}$ as proposal kernel. It is thus π -reversible.

From the properties of the directional neighbourhoods $\{N_{-1}(x)\}$ and $\{N_{+1}(x)\}$, we can deduce that, for all x,

$$Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})) = 2Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})) = 0,$$

and also,

$$Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x})^{c} \cap \mathbf{N}_{+1}(\mathbf{x})^{c}) = \sum_{\nu \in \{-1, +1\}} Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{c} \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \cap \mathbf{N}_{\nu}(\mathbf{x})) = 0,$$

the latter implying that $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cup \mathbf{N}_{+1}(\mathbf{x})) = 1$. Therefore, for $\mu_{\text{rev.}}$ -almost all (\mathbf{x}, \mathbf{y}) with $(\mathbf{x}, \mathbf{y}) \in R$, the acceptance probability $\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})$ is equal to

$$\varphi(r_{\text{rev.}}(\mathbf{x}, \mathbf{y})) = \varphi\left(r(\mathbf{x}, \mathbf{y}) \frac{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))}\right)$$

for some $v \in \{-1, +1\}$, except if $\mathbf{y} = \mathbf{x}$, in which case $\varphi(r_{rev.}(\mathbf{x}, \mathbf{y})) = 1$.

From the generalized auxiliary sampler, we can introduce the generalized lifted sampler. For all $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$, let $T_{\nu}(\mathbf{x}, \cdot)$ be a measure on $(\mathcal{X}, \mathbf{X})$ defined as:

$$A \in \mathbf{X}, \quad T_{\nu}(\mathbf{x}, A) = \begin{cases} \int_{A} \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) Q_{\nu}(\mathbf{x}, d\mathbf{y}) & \text{if } \mathbf{x} \in \mathring{\mathcal{X}} \text{ or } \mathbf{x} \in \partial \mathcal{X} \text{ with } Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let P_{Lifted} be defined as: for all $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$,

$$P_{\text{Lifted}}((\mathbf{x},\nu), \mathbf{d}(\mathbf{y},\nu')) = \delta_{\nu}(\nu') T_{\nu}(\mathbf{x}, \mathbf{dy}) + \delta_{-\nu}(\nu') \delta_{\mathbf{x}}(\mathbf{dy}) (1 - T_{\nu}(\mathbf{x}, \boldsymbol{\mathcal{X}})).$$

Note that, if (\mathbf{x}, ν) is such that $\mathbf{x} \in \overset{\circ}{\mathcal{X}}$ or $\mathbf{x} \in \partial \mathcal{X}$ with $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1$,

$$Q_{\nu}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \cap \{\mathbf{x}\}^{c}) = 1,$$

given that

$$Q_{\nu}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{c} \cup \mathbf{N}_{-\nu}(\mathbf{x}) \cup \{\mathbf{x}\}) \leq \frac{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{c} \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} + \frac{Q(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x}) \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} + \frac{Q(\mathbf{x}, \{\mathbf{x}\} \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} = 0,$$

by the union bound. Therefore, if (\mathbf{x}, ν) is such that $\mathbf{x} \in \overset{\circ}{\mathcal{X}}$ or $\mathbf{x} \in \partial \mathcal{X}$ with $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1$, the lifted sampler proceeds with a proposal $\mathbf{y} \sim Q_{\nu}(\mathbf{x}, \cdot)$ such that, with probability 1, the acceptance probability $\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})$ is equal to

$$\varphi(r_{\text{rev.}}(\mathbf{x}, \mathbf{y})) = \varphi\left(r(\mathbf{x}, \mathbf{y}) \frac{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))}\right)$$

if $(\mathbf{x}, \mathbf{y}) \in R$ or 0 if $(\mathbf{x}, \mathbf{y}) \notin R$.

We now present a result ensuring that P_{Lifted} leaves the product measure $\pi \otimes \mathcal{U}\{-1, +1\}$ invariant.

Proposition 4. Under the framework of this section, P_{Lifted} satisfies a skewed detailed balance.

4.3 The comparison

With the definitions and results of Sections 4.1 and 4.2, we are ready to present the generalized version of Theorem 1.

Theorem 2. For any $\mathbf{x} \in \mathcal{X}$ and $A \in \mathbf{X}$,

$$P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \geq \frac{1}{2} P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\}).$$

Therefore, for any f such that $\pi(f^2) < \infty$ and whenever $\lim_{\lambda \to 1} \operatorname{var}_{\lambda}(f, P_{\text{Lifted}}) = \operatorname{var}(f, P_{\text{Lifted}})$,

$$\operatorname{var}(f, P_{\operatorname{Lifted}}) \le \operatorname{var}(f, P_{\operatorname{rev.}}) \le 2\operatorname{var}(f, P_{\operatorname{MH}}) + \operatorname{Var}[f(\mathbf{X})], \quad \mathbf{X} \sim \pi$$

Under the general framework of this section, there exists a situation of perfect balance which is analogous to that described in Section 2 (it is described in the proof of Theorem 2). Under this situation of perfect balance, $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$ for all $\mathbf{x} \in \mathcal{X}$ and $A \in \mathcal{X}$. Therefore, for any f such that $\pi(f^2) < \infty$ and whenever $\lim_{\lambda \to 1} \operatorname{var}_{\lambda}(f, P_{\text{Lifted}}) = \operatorname{var}(f, P_{\text{Lifted}})$,

$$\operatorname{var}(f, P_{\operatorname{Lifted}}) \leq \operatorname{var}(f, P_{\operatorname{rev.}}) \leq \operatorname{var}(f, P_{\operatorname{MH}}).$$

This implies that Theorem 2 generalizes Theorem 7 in Andrieu and Livingstone (2021) as the latter is stated under the general framework of this section, but with the restriction to the situation of perfect balance and to the function $\varphi(r) = 1 \wedge r$.

References

- Andrieu, C., Lee, A. and Livingstone, S. (2020) A general perspective on the Metropolis–Hastings kernel. arXiv:2012.14881.
- Andrieu, C., Lee, A. and Vihola, M. (2018) Uniform ergodicity of the iterated conditional SMC and geometric ergodicity of particle Gibbs samplers (supplemental content). *Bernoulli*, **24**, 842–872.
- Andrieu, C. and Livingstone, S. (2021) Peskun–Tierney ordering for Markovian Monte Carlo: Beyond the reversible scenario. *Ann. Statist.*, **49**, 1958 1981.
- Barker, A. A. (1965) Monte Carlo calculations of the radial distribution functions for a proton-electron plasma. *Austral. J. Phys.*, **18**, 119–134.
- Beaumont, M. A. (2003) Estimation of population growth or decline in genetically monitored populations. *Genetics*, **164**, 1139–1160.
- Chen, F., Lovász, L. and Pak, I. (1999) Lifting Markov chains to speed up mixing. In *Proceedings of* the thirty-first annual ACM symposium on Theory of computing, 275–281.
- Diaconis, P., Holmes, S. and Neal, R. M. (2000) Analysis of a nonreversible Markov chain sampler. *Ann. Appl. Probab.*, 726–752.
- Gagnon, P. and Doucet, A. (2021) Nonreversible jump algorithms for Bayesian nested model selection. *J. Comput. Graph. Statist.*, **30**, 312–323. ArXiv:1911.01340.
- Gagnon, P. and Maire, F. (2024) An asymptotic Peskun ordering and its application to lifted samplers. *Bernoulli*, **30**, 2301 2325.
- Green, P. J. (1995) Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, **82**, 711–732.
- Gustafson, P. (1998) A guided walk Metropolis algorithm. Stat. Comput., 8, 357-364.
- Hastings, W. K. (1970) Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, **57**, 97–109.
- Häggström, O. and Rosenthal, J. S. (2007) On Variance Conditions for Markov Chain CLTs. *Electron. Commun. Probab.*, **12**, 454 464.
- Horowitz, A. M. (1991) A generalized guided Monte Carlo algorithm. Phys. Lett. B, 268, 247-252.
- Liu, J. S., Liang, F. and Wong, W. H. (2000) The multiple-try method and local optimization in Metropolis sampling. *J. Amer. Statist. Assoc.*, **95**, 121–134.
- Livingstone, S. and Zanella, G. (2022) The Barker proposal: combining robustness and efficiency in gradient-based MCMC. J. R. Stat. Soc. Ser. B. Stat. Methodol., 84, 496–523.
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H. and Teller, E. (1953) Equation of state calculations by fast computing machines. *J. Chem. Phys.*, **21**, 1087.
- Peskun, P. (1973) Optimum Monte-Carlo sampling using Markov chains. *Biometrika*, **60**, 607–612.

- Roberts, G. O. and Rosenthal, J. S. (2004) General state space Markov chains and MCMC algorithms. *Probab. Surv.*, **1**, 20–71.
- Sakai, Y. and Hukushima, K. (2016a) Eigenvalue analysis of an irreversible random walk with skew detailed balance conditions. *Phys. Rev. E*, **93**, 043318.

- (2016b) Irreversible simulated tempering. J. Phys. Soc. Jpn., 85, 104002.

- Syed, S., Bouchard-Côté, A., Deligiannidis, G. and Doucet, A. (2022) Non-reversible parallel tempering: A scalable highly parallel MCMC scheme. J. R. Stat. Soc. Ser. B. Stat. Methodol., 84, 321–350.
- Tierney, L. (1998) A note on Metropolis-Hastings kernels for general state spaces. *Ann. Appl. Probab.*, **8**, 1–9.
- Vucelja, M. (2016) Lifting-a nonreversible Markov chain Monte Carlo algorithm. *Amer. J. Phys.*, **84**, 958–968.
- Zanella, G. (2020) Informed proposals for local MCMC in discrete spaces. J. Amer. Statist. Assoc., 115, 852–865.

A Lemmas and proofs

In this section, we present proofs of results in the same order as these results appeared in the paper. We start in Section A.1 with proofs of results in Section 1. In Section A.1, we also present Lemma 2 which is used to prove Proposition 1. In Section A.2, we present proofs of results in Section 4.

A.1 Proofs of results in Section 1 and Lemma 2

Before presenting Lemma 2, which is about the reversibility of $P_{\text{rev.}}$, we introduce definitions. To introduce those definitions, it is useful to recall the definitions presented in Section 1.2. We first introduce $\mathring{X} := X \setminus \partial X$ and define $\tilde{Q}(\mathbf{x}, \cdot) := (1/2)Q_{-1}(\mathbf{x}, \cdot) + (1/2)Q_{+1}(\mathbf{x}, \cdot)$ for all $\mathbf{x} \in X$. We use $\alpha_{\text{rev.}}$ to denote the acceptance probability in the variant of MH in the general case which is defined as: for all $(\mathbf{x}, \mathbf{y}) \in X^2$, $\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = 1 \wedge r_{\text{rev.}}(\mathbf{x}, \mathbf{y})$, where $r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = 0$ whenever $\mathbf{y} \notin \mathbf{N}(\mathbf{x})$ and

$$r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = \begin{cases} \pi(\mathbf{y})\tilde{Q}(\mathbf{y}, \mathbf{x})/[\pi(\mathbf{x})\tilde{Q}(\mathbf{x}, \mathbf{y})] & \text{if } (\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \mathring{X}, \\ \pi(\mathbf{y})Q(\mathbf{y}, \mathbf{x})/[2\pi(\mathbf{x})\tilde{Q}(\mathbf{x}, \mathbf{y})] & \text{if } (\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \partial X, \\ 2\pi(\mathbf{y})\tilde{Q}(\mathbf{y}, \mathbf{x})/[\pi(\mathbf{x})Q(\mathbf{x}, \mathbf{y})] & \text{if } (\mathbf{x}, \mathbf{y}) \in \partial X \times \mathring{X}, \\ \pi(\mathbf{y})Q(\mathbf{y}, \mathbf{x})/[\pi(\mathbf{x})Q(\mathbf{x}, \mathbf{y})] & \text{if } (\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X, \end{cases}$$
(10)

whenever $\mathbf{y} \in \mathbf{N}(\mathbf{x})$. In the general case, $P_{\text{rev.}}$ is defined as: for any $\mathbf{x} \in \mathcal{X}$,

$$P_{\text{rev.}}(\mathbf{x}, \mathbf{dy}) = Q_{\text{rev.}}(\mathbf{x}, \mathbf{dy}) \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) + \delta_{\mathbf{x}}(\mathbf{dy}) \int (1 - \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{u})) Q_{\text{rev.}}(\mathbf{x}, \mathbf{du})$$

Lemma 2. Let $\mu_{\text{rev.}}$ and $\mu_{\text{rev.}}^T$ be two probability measures on the product space $(X^2, X \otimes X)$ defined as:

$$\mu_{\text{rev.}}(A \times B) = \int_A \pi(\mathbf{d}\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, B) \,, \qquad \mu_{\text{rev.}}^T(A \times B) = \int_B \pi(\mathbf{d}\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, A).$$

The set $R := \{(\mathbf{x}, \mathbf{y}) : \mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathbf{N}(\mathbf{x})\}$ is symmetric. The measures $\mu_{\text{rev.}}$ and $\mu_{\text{rev.}}^T$ are mutually absolutely continuous (they are positive on R and null on the complement of R, R^c). For all $(\mathbf{x}, \mathbf{y}) \in R$,

$$r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = \frac{\mathrm{d}\mu_{\text{rev.}}^T}{\mathrm{d}\mu_{\text{rev.}}}(\mathbf{x}, \mathbf{y}),$$

and we have that $0 < r_{rev.}(\mathbf{x}, \mathbf{y}) < \infty$ and $r_{rev.}(\mathbf{x}, \mathbf{y}) = 1/r_{rev.}(\mathbf{y}, \mathbf{x})$ for all $(\mathbf{x}, \mathbf{y}) \in R$.

An implication of Lemma 2 is that, by definition, $P_{\text{rev.}}$ is the standard MH Markov kernel (in the sense of Tierney (1998)) with π as invariant distribution and proposal distribution $Q_{\text{rev.}}(\mathbf{x}, \cdot)$. Thus, $P_{\text{rev.}}$ is π -reversible.

Proof of Lemma 2. We have that $(\mathbf{x}, \mathbf{y}) \in R$ if and only if $(\mathbf{y}, \mathbf{x}) \in R$ (using that $\mathbf{y} \in \mathbf{N}(\mathbf{x})$ if and only if $\mathbf{x} \in \mathbf{N}(\mathbf{y})$). Therefore, *R* is symmetric.

We have that $\mu_{\text{rev.}}(A \times B) = 0$ if, for all $(\mathbf{x}, \mathbf{y}) \in A \times B$, $\mathbf{y} \notin \mathbf{N}(\mathbf{x})$, meaning that $A \times B \subset R^c$. Therefore, for all such $(\mathbf{x}, \mathbf{y}), \mathbf{x} \notin \mathbf{N}(\mathbf{y})$, and $\mu_{\text{rev.}}^T(A \times B) = 0$. Note that, for the same reasons, we have that $\mu_{\text{rev.}}^T(A \times B) = 0$ implies that $\mu_{\text{rev.}}(A \times B) = 0$. Consequently, $\mu_{\text{rev.}}$ and $\mu_{\text{rev.}}^T$ are mutually absolutely continuous (they are positive on *R* and null on R^c).

Now, we prove that, for all $(\mathbf{x}, \mathbf{y}) \in R$,

$$r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = \frac{\mathrm{d}\mu_{\text{rev.}}^T}{\mathrm{d}\mu_{\text{rev.}}}(\mathbf{x}, \mathbf{y})$$

To prove this, we show that for any bounded measurable function $\phi: \mathcal{X}^2 \to \mathbb{R}$,

$$\int_R \phi r_{\text{rev.}} \, \mathrm{d}\mu = \int_R \phi \, \mathrm{d}\mu^T.$$

The equivalence between these two previous statements can be derived from Proposition 3 in Roberts and Rosenthal (2004). For all $(\mathbf{x}, \mathbf{y}) \in R$, we have that

$$\begin{aligned} r_{\text{rev.}}(\mathbf{x},\mathbf{y}) &= \frac{\pi(\mathbf{y})\,\tilde{Q}(\mathbf{y},\mathbf{x})}{\pi(\mathbf{x})\,\tilde{Q}(\mathbf{x},\mathbf{y})}\,\mathbb{1}((\mathbf{x},\mathbf{y})\in\mathring{X}\times\mathring{X}) + \frac{\pi(\mathbf{y})\,Q(\mathbf{y},\mathbf{x})}{2\,\pi(\mathbf{x})\,\tilde{Q}(\mathbf{x},\mathbf{y})}\,\mathbb{1}((\mathbf{x},\mathbf{y})\in\mathring{X}\times\partial\mathcal{X}) \\ &+ \frac{2\,\pi(\mathbf{y})\,\tilde{Q}(\mathbf{y},\mathbf{x})}{\pi(\mathbf{x})\,Q(\mathbf{x},\mathbf{y})}\,\mathbb{1}((\mathbf{x},\mathbf{y})\in\partial\mathcal{X}\times\mathring{X}) + \frac{\pi(\mathbf{y})\,Q(\mathbf{y},\mathbf{x})}{\pi(\mathbf{x})\,Q(\mathbf{x},\mathbf{y})}\,\mathbb{1}((\mathbf{x},\mathbf{y})\in\partial\mathcal{X}\times\partial\mathcal{X}). \end{aligned}$$

We thus write $\int_{R} \phi r_{rev} d\mu$ as a sum of 4 integrals and analyze each of them. First,

$$\int_{R} \phi(\mathbf{x}, \mathbf{y}) \frac{\pi(\mathbf{y}) \, \tilde{Q}(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, \tilde{Q}(\mathbf{x}, \mathbf{y})} \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \mathring{X}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\mathrm{rev.}}(\mathbf{x}, \mathrm{d}\mathbf{y})$$
$$= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, \tilde{Q}(\mathbf{y}, \mathrm{d}\mathbf{x}) \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \mathring{X})$$
$$= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, Q_{\mathrm{rev.}}(\mathbf{y}, \mathrm{d}\mathbf{x}) \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \mathring{X}).$$

Second,

$$\int_{R} \phi(\mathbf{x}, \mathbf{y}) \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{2 \, \pi(\mathbf{x}) \, \tilde{Q}(\mathbf{x}, \mathbf{y})} \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \partial X) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\mathrm{rev.}}(\mathbf{x}, \mathrm{d}\mathbf{y})$$

20 P. Gagnon and F. Maire

$$= \int_{R} \phi(\mathbf{x}, \mathbf{y}) (1/2) \pi(\mathrm{d}\mathbf{y}) Q(\mathbf{y}, \mathrm{d}\mathbf{x}) \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \partial X)$$

$$= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \pi(\mathrm{d}\mathbf{y}) \left[(1/2) Q(\mathbf{y}, \mathrm{d}\mathbf{x}) + (1/2) \delta_{\mathbf{y}}(\mathrm{d}\mathbf{x}) \right] \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \partial X)$$

$$= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \pi(\mathrm{d}\mathbf{y}) Q_{\mathrm{rev.}}(\mathbf{y}, \mathrm{d}\mathbf{x}) \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \mathring{X} \times \partial X).$$

Third,

$$\begin{split} &\int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{2 \, \pi(\mathbf{y}) \, \tilde{Q}(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \partial \mathcal{X} \times \mathring{\mathcal{X}}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\mathrm{rev.}}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, \tilde{Q}(\mathbf{y}, \mathrm{d}\mathbf{x}) \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \partial \mathcal{X} \times \mathring{\mathcal{X}}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, Q_{\mathrm{rev.}}(\mathbf{y}, \mathrm{d}\mathbf{x}) \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \partial \mathcal{X} \times \mathring{\mathcal{X}}). \end{split}$$

Fourth,

$$\begin{split} &\int_{R} \phi(\mathbf{x}, \mathbf{y}) \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \pi(d\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbb{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, (\mathbbm{1}(\mathbf{x} = \mathbf{y}) + \mathbbm{1}(\mathbf{x} \neq \mathbf{y})) \, \pi(d\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbbm{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} = \mathbf{y}) \, \pi(d\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \\ &+ \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbbm{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(d\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbbm{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} = \mathbf{y}) \, \pi(d\mathbf{x}) \, (1/2) \delta_{\mathbf{x}}(d\mathbf{y}) \\ &+ \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbbm{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(d\mathbf{x}) \, (1/2) Q(\mathbf{x}, d\mathbf{y}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{x}, \mathbf{y})} \, \mathbbm{1}((\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(d\mathbf{x}) \, (1/2) Q(\mathbf{y}, d\mathbf{x}) \\ &+ \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \frac{\pi(\mathbf{y}) \, Q(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q(\mathbf{y}, \mathbf{x})} \, \mathbbm{1}(\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(d\mathbf{x}) \, (1/2) Q(\mathbf{y}, d\mathbf{x}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \mathbbm{1}(\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \mathbbm{1}(\mathbf{x}) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, (1/2) Q(\mathbf{y}, d\mathbf{x}) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(\mathbf{y}) \, (1/2) Q(\mathbf{y}, d\mathbf{x}) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \mathbbm{1}(\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \pi(d\mathbf{y}) \, \mathbbm{1}(\mathbf{x} = \mathbf{y}) \, (1/2) \delta_{\mathbf{y}}(d\mathbf{x}) + \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, (1/2) Q(\mathbf{y}, d\mathbf{x}) \right) \\ &= \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \mathbbm{1}(\mathbf{x}, \mathbf{y}) \in \partial X \times \partial X) \, \pi(d\mathbf{y}) \, \mathbbm{1}(\mathbf{x} = \mathbf{y}) \, \pi(\mathbf{x}) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(\mathbf{x}) \, \mathbbm{1}(\mathbf{x} \neq \mathbf{y}) \, \pi(\mathbf{x}) \, \mathbbm{1}(\mathbf{x}) \,$$

Putting all those results together yields

$$\int_{R} \phi r_{\text{rev.}} \, \mathrm{d}\mu = \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\text{rev.}}(\mathbf{x}, \mathrm{d}\mathbf{y}) = \int_{R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, Q_{\text{rev.}}(\mathbf{y}, \mathrm{d}\mathbf{x}) = \int_{R} \phi \, \mathrm{d}\mu^{T},$$

which concludes the proof.

Before presenting the proof of Proposition 2, we introduce definitions. We use $\alpha_{\text{Lifted},\nu}$ to denote the acceptance probability in the lifted sampler in the general case which is defined as: for all $(\mathbf{x}, \mathbf{y}) \in \mathbf{X}^2$

21

and $v \in \{-1, +1\}$, let $\alpha_{\text{Lifted},v}(\mathbf{x}, \mathbf{y}) := 1 \land r_{\text{Lifted},v}(\mathbf{x}, \mathbf{y})$, where $r_{\text{Lifted},v}(\mathbf{x}, \mathbf{y}) = 0$ whenever $\mathbf{N}_{v}(\mathbf{x}) = \emptyset$ or $\mathbf{y} \notin \mathbf{N}_{v}(\mathbf{x})$ and

$$r_{\text{Lifted},\nu}(\mathbf{x},\mathbf{y}) = \frac{\pi(\mathbf{y}) \, Q_{-\nu}(\mathbf{y},\mathbf{x})}{\pi(\mathbf{x}) \, Q_{\nu}(\mathbf{x},\mathbf{y})}$$

whenever $\mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})$. For all $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$, let $T_{\nu}(\mathbf{x}, \cdot)$ be a measure on $(\mathcal{X}, \mathbf{X})$ with a density with respect to dy, which is denoted by $T_{\nu}(\mathbf{x}, \cdot)$ as well to simplify and defined as:

$$T_{\nu}(\mathbf{x}, \mathbf{y}) = \begin{cases} 0 & \text{if } \mathbf{N}_{\nu}(\mathbf{x}) = \emptyset \text{ or } \mathbf{y} \notin \mathbf{N}_{\nu}(\mathbf{x}), \\ \alpha_{\text{Lifted}, \nu}(\mathbf{x}, \mathbf{y}) Q_{\nu}(\mathbf{x}, \mathbf{y}) & \text{if } \mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x}). \end{cases}$$

In the general case, P_{Lifted} is defined as: for any $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$,

$$P_{\text{Lifted}}((\mathbf{x}, \nu), \mathbf{d}(\mathbf{y}, \nu')) = \delta_{\nu}(\nu') T_{\nu}(\mathbf{x}, \mathbf{dy}) + \delta_{-\nu}(\nu') \delta_{\mathbf{x}}(\mathbf{dy}) (1 - T_{\nu}(\mathbf{x}, \mathcal{X})).$$

Note that this definition is equivalent to: for any $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$ such that $\mathbf{N}_{\nu}(\mathbf{x}) \neq \emptyset$,

$$P_{\text{Lifted}}((\mathbf{x}, \nu), \mathbf{d}(\mathbf{y}, \nu')) = Q_{\nu}(\mathbf{x}, \mathbf{dy}) \alpha_{\nu}(\mathbf{x}, \mathbf{y}) \delta_{\nu}(\mathbf{d\nu'}) + \delta_{(\mathbf{x}, -\nu)}(\mathbf{d}(\mathbf{y}, \nu')) \int (1 - \alpha_{\nu}(\mathbf{x}, \mathbf{u})) Q_{\nu}(\mathbf{x}, \mathbf{du}),$$

and for any $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$ such that $\mathbf{N}_{\nu}(\mathbf{x}) = \emptyset$,

$$P_{\text{Lifted}}((\mathbf{x}, \nu), \mathbf{d}(\mathbf{y}, \nu')) = \delta_{(\mathbf{x}, -\nu)}(\mathbf{d}(\mathbf{y}, \nu')).$$

Following Andrieu and Livingstone (2021), P_{Lifted} is said to satisfy a skewed detailed balance if the part of the kernel associated to a move of the **X** component, that is $T_{\nu}(\mathbf{x}, d\mathbf{y})$, satisfies

$$\pi(\mathbf{d}\mathbf{x}) T_{+1}(\mathbf{x}, \mathbf{d}\mathbf{y}) = \pi(\mathbf{d}\mathbf{y}) T_{-1}(\mathbf{y}, \mathbf{d}\mathbf{x}).$$

More formally, it corresponds to proving that the two following measures on the product space ($X^2, X \otimes X$) are equal:

$$\mu_{+1}(A \times B) = \int_A \pi(\mathbf{d}\mathbf{x}) T_{+1}(\mathbf{x}, B) \text{ and } \mu_{-1}(A \times B) = \int_B \pi(\mathbf{d}\mathbf{x}) T_{-1}(\mathbf{x}, A).$$

From the skewed detailed balance, we can conclude that P_{Lifted} leaves the distribution $\pi \otimes \mathcal{U}\{-1, +1\}$ invariant. Indeed, considering an event $A \in \mathbf{X}$ and $\nu' \in \{-1, +1\}$,

$$\sum_{\nu \in \{-1,+1\}} \int \pi(\mathbf{d}\mathbf{x}) \frac{1}{2} P_{\text{Lifted}}((\mathbf{x},\nu), A \times \nu')$$

$$= \frac{1}{2} \sum_{\nu \in \{-1,+1\}} \left(\mathbb{1}(\nu' = \nu) \int \pi(\mathbf{d}\mathbf{x}) T_{\nu}(\mathbf{x}, A) + \mathbb{1}(\nu' = -\nu) \int \pi(\mathbf{d}\mathbf{x}) \int_{A} \delta_{\mathbf{x}}(\mathbf{d}\mathbf{y}) (1 - T_{\nu}(\mathbf{x}, X)) \right)$$

$$= \frac{1}{2} \sum_{\nu \in \{-1,+1\}} \left(\mathbb{1}(\nu' = \nu) \int_{A} \pi(\mathbf{d}\mathbf{x}) T_{-\nu}(\mathbf{x}, X) + \mathbb{1}(\nu' = -\nu) \int_{A} \pi(\mathbf{d}\mathbf{x}) (1 - T_{\nu}(\mathbf{x}, X)) \right)$$

$$= \frac{1}{2} \int_{A} \pi(\mathbf{d}\mathbf{x}) (T_{-\nu'}(\mathbf{x}, X) + 1 - T_{-\nu'}(\mathbf{x}, X)) = \frac{1}{2} \int_{A} \pi(\mathbf{d}\mathbf{x}).$$

Therefore, the probability to reach $A \times v'$ in one step, if the chain is in stationarity (under $\pi \otimes \mathcal{U}\{-1, +1\}$), is equal to the probability of $A \times v'$ under the stationary distribution $\pi \otimes \mathcal{U}\{-1, +1\}$.

Proof of Proposition 2. For $v \in \{-1, +1\}$, let $R_v := \{(\mathbf{x}, \mathbf{y}) : \mathbf{x} \in X, \mathbf{y} \in \mathbf{N}_v(\mathbf{x})\}$. Note that $(\mathbf{x}, \mathbf{y}) \in R_v$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-v}$, using that $\mathbf{y} \in \mathbf{N}_v(\mathbf{x})$ if and only if $\mathbf{x} \in \mathbf{N}_{-v}(\mathbf{y})$.

We now prove that μ_{+1} and μ_{-1} are equal. To achieve this, we prove that for any bounded measurable function $\phi : X^2 \to \mathbb{R}$,

$$\int \phi \, \mathrm{d}\mu_{+1} = \int \phi \, \mathrm{d}\mu_{-1}$$

We have that

$$\int \phi \, \mathrm{d}\mu_{+1} = \int_{R_{+1}} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, \alpha_{\mathrm{Lifted}, +1}(\mathbf{x}, \mathbf{y}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}) \, \mathbb{1}(\mathbf{y} \in \mathbf{N}_{+1}(\mathbf{x}))$$
$$= \int_{R_{+1}} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, \alpha_{\mathrm{Lifted}, -1}(\mathbf{y}, \mathbf{x}) \, Q_{-1}(\mathbf{y}, \mathrm{d}\mathbf{x}) \, \mathbb{1}(\mathbf{x} \in \mathbf{N}_{-1}(\mathbf{y}))$$
$$= \int \phi \, \mathrm{d}\mu_{-1}.$$

A.2 **Proofs of results in Section 4**

Proof of Lemma 1. We prove the results in the same order as they appear in Lemma 1.

Proof of Result 1. Let r > 0 and $\epsilon > 0$, then $\varphi(r + \epsilon) = r\varphi(1/(r + \epsilon)) + \epsilon\varphi(1/(r + \epsilon))$. Assume that φ is decreasing, that is r < r' implies that $\varphi(r) > \varphi(r')$. Then, $\varphi(1/(r + \epsilon)) > \varphi(1/r)$ and

$$\varphi(r+\epsilon) = r\varphi(1/(r+\epsilon)) + \epsilon\varphi(1/(r+\epsilon)) > r\varphi(1/r) = \varphi(r),$$

which leads to a contradiction.

Proof of Result 2. Let r > 0. Take $\epsilon > 0$ and note that $\varphi(r + \epsilon) - \varphi(r) \ge 0$ (by Result 1). Moreover,

$$\varphi(r+\epsilon) = (r+\epsilon)\varphi(1/(r+\epsilon)) \le (r+\epsilon)\varphi(1/r) \le \varphi(r) + \epsilon,$$

using again Result 1 and the fact that $\varphi \leq 1$. Thus, $\varphi(r) \leq \varphi(r + \epsilon) \leq \varphi(r) + \epsilon$ and φ is continuous from the right. A symmetric argument shows that φ is also continuous from the left, thus φ is continuous.

Proof of Result 3. We have that $0 \le \varphi(r) = r\varphi(1/r) \le r$, which allows to conclude by taking $r \downarrow 0$.

Proof of Result 4. The null function verifies $\varphi(r) = r\varphi(1/r) = 0$ for all $r \in (0, \infty)$. Now consider that φ is not the null function. Then, $\varphi(r) > 0$ for some r and $\varphi(r') > 0$ for all $r' \ge r$ by Result 1. For any r' as small as we want, we also have $\varphi(r') > 0$ given that $\varphi(r') = r'\varphi(1/r') > 0$. Result 1, thus allows to confirm that $\varphi(r) > 0$ for all $r \in (0, \infty)$.

Proof of Result 5. If φ is the null function, then $0 = \varphi(ab) \ge \varphi(a)\varphi(b) = 0$ for any a > 0, b > 0. Now consider that φ is not the null function; thus, it is a positive function. We consider two cases. First, we consider that $a \le 1, b > 0$. Thus, $1/ab \ge 1/b$ and $\varphi(1/ab) \ge \varphi(1/b)$ from Result 1. Then, using that $\varphi(1/a) \le 1$, we have that

$$\varphi(ab) = ab\varphi(1/ab) \ge ab\varphi(1/b) \ge ab\varphi(1/b)\varphi(1/a) = \varphi(a)\varphi(b).$$

Now, consider that a > 1, b > 0. Given that $b = \varphi(b)/\varphi(1/b)$ and $a = \varphi(a)/\varphi(1/a)$, we have that

$$\varphi(ab) = \frac{\varphi(a)}{\varphi(1/a)} \frac{\varphi(b)}{\varphi(1/b)} \varphi(1/ab) = \varphi(a)\varphi(b) \frac{\varphi(1/ab)}{\varphi(1/a)\varphi(1/b)} \ge \varphi(a)\varphi(b) \,,$$

where the last inequality follows from the fact that $\varphi(1/ab) \ge \varphi(1/a)\varphi(1/b)$ using the result of the first case: $\varphi(a'b') \ge \varphi(a')\varphi(b')$ with $a' = 1/a \le 1$ and b' = 1/b > 0.

Proof of Result 6. We have that $1/ab \ge 1/a$ so $\varphi(1/ab) \ge \varphi(1/a)$ (Result 1), but $\varphi(ab) = ab\varphi(1/ab) \ge ba\varphi(1/a) = b\varphi(a)$.

Before presenting the proof of Proposition 3, we present and prove one lemma that will be useful. Lemma 3. Let $g : X^2 \to \mathbb{R}$ be such that

$$g(\mathbf{x}, \mathbf{y}) = \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{-1} \cap R_{+1}^{c}) + \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{+1} \cap R_{-1}^{c})$$

for all $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$. Then, $g = 1 \mu$ -almost everywhere and μ^T -almost everywhere. In particular, for any measurable function $\vartheta : \mathcal{X}^2 \to \mathbb{R}$,

$$\int \vartheta \, \mathrm{d}\mu = \sum_{\nu \in \{-1,+1\}} \int_{R_{\nu} \cap R_{-\nu}^{\mathsf{c}}} \vartheta \, \mathrm{d}\mu$$

and

$$\int \vartheta \, \mathrm{d} \mu^T = \sum_{\nu \in \{-1,+1\}} \int_{R_{\nu} \cap R_{-\nu}^{\mathsf{c}}} \vartheta \, \mathrm{d} \mu^T.$$

Proof. For $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2$,

$$g(\mathbf{x}, \mathbf{y}) = \begin{cases} 0 & \text{if} \qquad (\mathbf{x}, \mathbf{y}) \in (R_{-1}^{c} \cap R_{+1}^{c}) \cup (R_{+1} \cap R_{+1}^{c}) \cup (R_{-1} \cap R_{-1}^{c}) \cup (R_{+1} \cap R_{-1}), \\ 2 & \text{if} \qquad (\mathbf{x}, \mathbf{y}) \in R_{-1} \cap R_{-1}^{c} \cap R_{+1} \cap R_{+1}^{c}, \\ 1 & \text{otherwise.} \end{cases}$$

Clearly,

$$\mu(R_{-1} \cap R_{-1}^{c} \cap R_{+1} \cap R_{+1}^{c}) = 0.$$

Also,

$$\mu((R_{-1}^{c} \cap R_{+1}^{c}) \cup (R_{+1} \cap R_{+1}^{c}) \cup (R_{-1} \cap R_{-1}^{c}) \cup (R_{+1} \cap R_{-1}))$$

$$\leq \mu(R_{-1}^{c} \cap R_{+1}^{c}) + \mu(R_{+1} \cap R_{+1}^{c}) + \mu(R_{-1} \cap R_{-1}^{c}) + \mu(R_{+1} \cap R_{-1}).$$

Clearly,

$$\mu(R_{+1} \cap R_{+1}^{c}) = \mu(R_{-1} \cap R_{-1}^{c}) = 0.$$

By definition,

$$\mu(R_{-1} \cap R_{+1}) = \int \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{-1} \cap R_{+1}) \, \mu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) = \int \pi(\mathrm{d}\mathbf{x}) \, Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})),$$

and

$$\mu(R_{-1}^{c} \cap R_{+1}^{c}) = \int \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{-1}^{c} \cap R_{+1}^{c}) \,\mu(d\mathbf{x}, d\mathbf{y}) = \int \pi(d\mathbf{x}) \, Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x})^{c} \cap \mathbf{N}_{+1}(\mathbf{x})^{c}),$$

given that $R_{\nu}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathbf{X}^{2} : \mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})\}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathbf{X}^{2} : \mathbf{y} \notin \mathbf{N}_{\nu}(\mathbf{x})\}$. But, we have that $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})) = 0$ and $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x})^{c} \cap \mathbf{N}_{+1}(\mathbf{x})^{c}) = 0$ for all $\mathbf{x} \in \mathbf{X}$, as proved in Section 4.2. This implies that $g = 1 \mu$ -almost everywhere. We also have that $g = 1 \mu^{T}$ -almost everywhere given that g is symmetric (using that $\mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})$ if and only if $\mathbf{x} \in \mathbf{N}_{-\nu}(\mathbf{y})$).

Proof of Proposition 3. To prove that for all $(\mathbf{x}, \mathbf{y}) \in R$,

$$r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) := \frac{d\mu_{\text{rev.}}^T}{d\mu_{\text{rev.}}}(\mathbf{x}, \mathbf{y}) = r(\mathbf{x}, \mathbf{y})\beta(\mathbf{x}, \mathbf{y}),$$

we will prove that

$$\int_{R} \phi r_{\text{rev.}} \, \mathrm{d}\mu_{\text{rev.}} = \int_{R} \phi \, \mathrm{d}\mu_{\text{rev.}}^{T},$$

for any bounded measurable function $\phi : \mathcal{X}^2 \to \mathbb{R}$. This will allow to conclude that the restriction of the measures $\mu_{\text{rev.}}$ and $\mu_{\text{rev.}}^T$ to *R* are mutually absolutely continuous given that $r_{\text{rev.}}$ is positive on *R*. For any bounded measurable function $\phi : \mathcal{X}^2 \to \mathbb{R}$,

$$\begin{split} \int_{\{\hat{X}\times\hat{X}\}\cap R}\phi\,r_{\mathrm{rev.}}\,\mathrm{d}\mu_{\mathrm{rev.}} &= \sum_{\nu\in\{-1,+1\}}\int_{\{\hat{X}\times\hat{X}\}\cap R}\phi(\mathbf{x},\mathbf{y})\,r_{\mathrm{rev.}}(\mathbf{x},\mathbf{y})\,\pi(\mathrm{d}\mathbf{x})\,(1/2)\,Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \sum_{\nu\in\{-1,+1\}}\int_{\{\hat{X}\times\hat{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{\mathsf{c}}\cap\Delta^{\mathsf{c}}}\phi(\mathbf{x},\mathbf{y})\,r_{\mathrm{rev.}}(\mathbf{x},\mathbf{y})\,\pi(\mathrm{d}\mathbf{x})\,(1/2)\,Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}), \end{split}$$

using that $\pi(\mathbf{d}\mathbf{x}) Q_{\nu}(\mathbf{x}, \mathbf{d}\mathbf{y})$ assigns null measure to $\{\mathbf{\mathcal{X}} \times \mathbf{\mathcal{X}}\} \cap (R_{\nu} \cap R_{-\nu}^{\mathsf{c}})^{\mathsf{c}}$ and $\{\mathbf{\mathcal{X}} \times \mathbf{\mathcal{X}}\} \cap \Delta$, for $\nu \in \{-1, +1\}$. Indeed,

$$\begin{split} &\int_{\{\mathring{X}\times\mathring{X}\}\cap(R_{\nu}^{\mathsf{c}}\cup R_{-\nu})} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{\nu}^{\mathsf{c}} \cup R_{-\nu}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &\leq \int_{\{\mathring{X}\times\mathring{X}\}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{\nu}^{\mathsf{c}}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathrm{d}\mathbf{y}) + \int_{\{\mathring{X}\times\mathring{X}\}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{-\nu}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{\mathsf{c}} \cap \mathring{X}) + \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x}) \cap \mathring{X}) \\ &\leq \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{\mathsf{c}}) + \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x})) \\ &= \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, \frac{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{\mathsf{c}} \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})} + \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, \frac{Q(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x}) \cap \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})} = 0, \end{split}$$

and

$$\begin{split} \int_{\{\mathring{X}\times\mathring{X}\}\cap\Delta} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) &= \int_{\{\mathring{X}\times\mathring{X}\}} \mathbbm{1}(\mathbf{x}=\mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) = \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, Q_{\nu}(\mathbf{x},\{\mathbf{x}\}) \\ &= \int_{\mathring{X}} \pi(\mathrm{d}\mathbf{x}) \, \frac{Q(\mathbf{x},\{\mathbf{x}\})}{Q(\mathbf{x},\mathbf{N}_{\nu}(\mathbf{x}))} = 0, \end{split}$$

using the definition of R_{ν} , that $R_{\nu}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^{2} : \mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})\}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^{2} : \mathbf{y} \notin \mathbf{N}_{\nu}(\mathbf{x})\}$, and that $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})) = 0$ and $Q(\mathbf{x}, \{\mathbf{x}\}) = 0$ for all \mathbf{x} . Therefore, for $\nu \in \{-1, +1\}$ and for any measurable function $\vartheta : \chi^2 \to \mathbb{R}$,

$$\int_{\{\mathring{X}\times\mathring{X}\}\cap(R_{\nu}\cap R_{-\nu}^{c})^{c}}\vartheta(\mathbf{x},\mathbf{y})\,\pi(\mathrm{d}\mathbf{x})\,Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) = \int_{\{\mathring{X}\times\mathring{X}\}\cap\Delta}\vartheta(\mathbf{x},\mathbf{y})\,\pi(\mathrm{d}\mathbf{x})\,Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) = 0.$$
(11)

Also,

$$\begin{split} &\int_{\{\mathring{X}\times\mathring{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) r_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \pi(\mathrm{d}\mathbf{x}) (1/2) Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{c}\cap\Delta^{c}} \frac{\phi(\mathbf{x},\mathbf{y})}{2Q(\mathbf{y},\mathbf{N}_{-\nu}(\mathbf{y}))} r(\mathbf{x},\mathbf{y}) \mu(\mathrm{d}\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{c}\cap\Delta^{c}} \frac{\phi(\mathbf{x},\mathbf{y})}{2Q(\mathbf{y},\mathbf{N}_{-\nu}(\mathbf{y}))} \mu^{T}(\mathrm{d}\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \pi(\mathrm{d}\mathbf{y}) (1/2) Q_{-\nu}(\mathbf{y},\mathrm{d}\mathbf{x}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}\cap R} \phi(\mathbf{x},\mathbf{y}) \pi(\mathrm{d}\mathbf{y}) (1/2) Q_{-\nu}(\mathbf{y},\mathrm{d}\mathbf{x}). \end{split}$$

In the first equality, we used the definition of $r_{rev.}$. In the second equality, we used that $\int_R \phi r d\mu = \int_R \phi d\mu^T$ for all ϕ bounded measurable. Note that $\phi(\mathbf{x}, \mathbf{y})/(2Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})))$ is not necessarily bounded, but we can limit ourselves to the case where $Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))$ is bounded from below by an arbitrarily small constant and use a limiting argument with the monotone convergence theorem (where we separate the cases i) $\phi \ge 0$ and ii) $\phi < 0$). In the third equality, we used that $(\mathbf{x}, \mathbf{y}) \in R_{\nu}$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-\nu}$. In the fourth equality, we used the same arguments as before to obtain an equality by omitting the intersection with Δ^c and $R_{\nu} \cap R_{-\nu}^c$, recalling that $(\mathbf{x}, \mathbf{y}) \in R_{\nu}$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-\nu}$.

$$\begin{split} \int_{\{\mathring{X}\times\mathring{X}\}\cap R} \phi \, r_{\text{rev.}} \, \mathrm{d}\mu_{\text{rev.}} &= \sum_{\nu\in\{-1,+1\}} \int_{\{\mathring{X}\times\mathring{X}\}\cap R} \phi(\mathbf{x},\mathbf{y}) \, r_{\text{rev.}}(\mathbf{x},\mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) \, Q_{\nu}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \sum_{\nu\in\{-1,+1\}} \int_{\{\mathring{X}\times\mathring{X}\}\cap R} \phi(\mathbf{x},\mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, (1/2) \, Q_{-\nu}(\mathbf{y},\mathrm{d}\mathbf{x}) \\ &= \int_{\{\mathring{X}\times\mathring{X}\}\cap R} \phi \, \mathrm{d}\mu_{\text{rev.}}^T. \end{split}$$

Using similar arguments as above, we obtain that

$$\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R}\phi\,r_{\mathrm{rev.}}\,\mathrm{d}\mu_{\mathrm{rev.}}=\sum_{\nu\in\{-1,+1\}}\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{c}\cap\Delta^{c}}\frac{\phi(\mathbf{x},\mathbf{y})}{2Q(\mathbf{y},\mathbf{N}_{-\nu}(\mathbf{y}))}\,r(\mathbf{x},\mathbf{y})\,\mu(\mathrm{d}\mathbf{x},\mathrm{d}\mathbf{y}).$$

But, by the properties of the directional neighbourhoods, for any $(\mathbf{x}, \mathbf{y}) \in \{ \mathring{X} \times \partial X \} \cap R_{\nu}$, we have that $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) > 0$, $Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) > 0$ and $Q(\mathbf{y}, \mathbf{N}_{-1}(\mathbf{y}))Q(\mathbf{y}, \mathbf{N}_{+1}(\mathbf{y})) = 0$, which, together with $1 = Q(\mathbf{y}, \mathbf{X}) = Q(\mathbf{y}, \mathbf{N}_{-1}(\mathbf{y})) + Q(\mathbf{y}, \mathbf{N}_{+1}(\mathbf{y}))$, implies that $Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) = 1$. Therefore,

$$\begin{split} &\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R}\phi\,r_{\mathrm{rev.}}\,\mathrm{d}\mu_{\mathrm{rev.}} \\ &=\sum_{\nu\in\{-1,+1\}}\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{\mathsf{c}}\cap\Delta^{\mathsf{c}}}\frac{\phi(\mathbf{x},\mathbf{y})}{2}\,\mu^{T}(\mathrm{d}\mathbf{x},\mathrm{d}\mathbf{y}) \\ &=\sum_{\nu\in\{-1,+1\}}\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R\cap R_{\nu}\cap R_{-\nu}^{\mathsf{c}}\cap\Delta^{\mathsf{c}}}\phi(\mathbf{x},\mathbf{y})\,\pi(\mathrm{d}\mathbf{y})\,(1/2)\,Q(\mathbf{y},\mathrm{d}\mathbf{x}) \\ &=\int_{\{\mathring{X}\times\partial\mathfrak{X}\}\cap R}\phi(\mathbf{x},\mathbf{y})\,(\mathbbm{1}((\mathbf{x},\mathbf{y})\in R_{-1}\cap R_{+1}^{\mathsf{c}})+\mathbbm{1}((\mathbf{x},\mathbf{y})\in R_{+1}\cap R_{-1}^{\mathsf{c}}))\,\pi(\mathrm{d}\mathbf{y})\,(1/2)\,Q(\mathbf{y},\mathrm{d}\mathbf{x}) \end{split}$$

$$= \int_{\{\mathring{\boldsymbol{X}} \times \partial \boldsymbol{X}\} \cap R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, (1/2) \, Q(\mathbf{y}, \mathrm{d}\mathbf{x})$$

$$= \int_{\{\mathring{\boldsymbol{X}} \times \partial \boldsymbol{X}\} \cap R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, ((1/2) \, Q(\mathbf{y}, \mathrm{d}\mathbf{x}) + (1/2)\delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}))$$

$$= \int_{\{\mathring{\boldsymbol{X}} \times \partial \boldsymbol{X}\} \cap R} \phi \, \mathrm{d}\mu_{\mathrm{rev.}}^{T}.$$

In the first equality, we used that $\int_R \phi r \, d\mu = \int_R \phi \, d\mu^T$ for all ϕ bounded measurable. In the third equality, we used that $\pi(d\mathbf{y}) Q(\mathbf{y}, d\mathbf{x})$ assigns null measure to Δ and an argument similar as in (11) (recall that $Q(\mathbf{y}, \{\mathbf{y}\}) = 0$ for all \mathbf{y}). In the fourth equality, we applied Lemma 3.

Symmetrically,

$$\begin{split} \int_{\partial X \times \hat{X} \cap R} \phi \, r_{\text{rev.}} \, d\mu_{\text{rev.}} &= \int_{\partial X \times \hat{X} \cap R} \frac{\phi(\mathbf{x}, \mathbf{y})}{2} r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \mu(d\mathbf{x}, d\mathbf{y}) \\ &= \int_{\partial X \times \hat{X} \cap R \cap \Delta^{\circ}} \frac{\phi(\mathbf{x}, \mathbf{y})}{2} r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \mu(d\mathbf{x}, d\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\partial X \times \hat{X} \cap R \cap R_{\nu} \cap R_{\nu}^{\circ} \cap \Delta^{\circ}} \frac{\phi(\mathbf{x}, \mathbf{y})}{2} \frac{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))}{Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))} \, r(\mathbf{x}, \mathbf{y}) \, \mu(d\mathbf{x}, d\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\partial X \times \hat{X} \cap R \cap R_{\nu} \cap R_{\nu}^{\circ} \cap \Delta^{\circ}} \frac{\phi(\mathbf{x}, \mathbf{y})}{2} \frac{1}{Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))} \, \mu^{T}(d\mathbf{x}, d\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\partial X \times \hat{X} \cap R \cap R_{\nu} \cap R_{\nu}^{\circ} \cap \Delta^{\circ}} \phi(\mathbf{x}, \mathbf{y}) \, \pi(d\mathbf{y}) \, (1/2) Q_{-\nu}(\mathbf{y}, d\mathbf{x}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\partial X \times \hat{X} \cap R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(d\mathbf{y}) \, (1/2) Q_{-\nu}(\mathbf{y}, d\mathbf{x}) \\ &= \int_{\partial X \times \hat{X} \cap R} \phi(\mathbf{x}, \mathbf{y}) \, \pi(d\mathbf{y}) \, ((1/2) Q_{-1}(\mathbf{y}, d\mathbf{x}) + (1/2) Q_{+1}(\mathbf{y}, d\mathbf{x})) \\ &= \int_{\partial X \times \hat{X} \cap R} \phi \, d\mu_{\text{rev.}}^{T}. \end{split}$$

In the second equality, we used that $\mu(d\mathbf{x}, d\mathbf{y}) = \pi(d\mathbf{x}) Q(\mathbf{x}, d\mathbf{y})$ assigns null measure to Δ and an argument similar as in (11). In the third equality, we applied Lemma 3 and used the definition of the function $r_{\text{rev.}}$. In the fourth equality, we used that, similarly as in the previous case, $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1$ for any $(\mathbf{x}, \mathbf{y}) \in \{\partial \mathbf{X} \times \mathbf{X}\} \cap R_{\nu}$. We also used that $\int_{R} \phi r d\mu = \int_{R} \phi d\mu^{T}$ for all ϕ bounded measurable (again we can use a limiting argument with the monotone convergence theorem). In the fifth equality, we used that $(\mathbf{x}, \mathbf{y}) \in R_{\nu}$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-\nu}$. In the sixth equality, we used that $\pi(d\mathbf{y}) Q_{-\nu}(\mathbf{y}, d\mathbf{x})$ assigns null measure to $\{\partial \mathbf{X} \times \mathbf{X}\} \cap (R_{\nu} \cap R_{-\nu}^{c})^{c}$ and $\{\partial \mathbf{X} \times \mathbf{X}\} \cap \Delta$ and a similar argument as in (11) (recall that $(\mathbf{x}, \mathbf{y}) \in R_{\nu}$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-\nu}$).

Finally, using the definition of $r_{rev.}$,

$$\begin{split} \int_{\{\partial \mathcal{X} \times \partial \mathcal{X}\} \cap R} \phi \, r_{\text{rev.}} \, \mathrm{d}\mu_{\text{rev.}} &= \int_{\{\partial \mathcal{X} \times \partial \mathcal{X}\} \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) \, r(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) \, \delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}) \\ &+ \int_{\{\partial \mathcal{X} \times \partial \mathcal{X}\} \cap R \cap \Delta^{\circ}} \phi(\mathbf{x}, \mathbf{y}) \, r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) \, Q(\mathbf{x}, \mathrm{d}\mathbf{y}), \end{split}$$

27

because

$$\int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \pi(\mathrm{d}\mathbf{x}) (1/2) \,\delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}) = 0,$$

and

$$\int_{\partial X \times \partial X \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \pi(\mathrm{d}\mathbf{x}) (1/2) Q(\mathbf{x}, \mathrm{d}\mathbf{y}) = 0,$$

given that $\mu(d\mathbf{x}, d\mathbf{y}) = \pi(d\mathbf{x}) Q(\mathbf{x}, d\mathbf{y})$ assigns null measure to Δ and using an argument similar as in (11). Using that $r(\mathbf{x}, \mathbf{y}) = 1$ when $\mathbf{y} = \mathbf{x}$,

$$\int_{\partial \mathcal{X} \times \partial \mathcal{X} \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) \, r(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) \, \delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}) = \int_{\partial \mathcal{X} \times \partial \mathcal{X} \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) \, \delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}) \\ = \int_{\partial \mathcal{X} \times \partial \mathcal{X} \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, (1/2) \, \delta_{\mathbf{y}}(\mathrm{d}\mathbf{x}).$$

Also,

$$\begin{split} &\int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, (1/2) Q(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, (1/2) \mu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\{\partial X \times \partial X\} \cap R \cap R_{\nu} \cap R_{-\nu}^{c} \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, r_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, (1/2) \mu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\{\partial X \times \partial X\} \cap R \cap R_{\nu} \cap R_{-\nu}^{c} \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, r(\mathbf{x}, \mathbf{y}) \, (1/2) \mu(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \sum_{\nu \in \{-1, +1\}} \int_{\{\partial X \times \partial X\} \cap R \cap R_{\nu} \cap R_{-\nu}^{c} \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, (1/2) \, \mu^{T}(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, (1/2) \, \mu^{T}(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{y}) \, (1/2) \, Q(\mathbf{y}, \mathrm{d}\mathbf{x}). \end{split}$$

In the second and fifth equalities, we applied Lemma 3. In the third equality, we used that all $(\mathbf{x}, \mathbf{y}) \in$ $\{\partial X \times \partial X\} \cap R \cap R_{\nu} \cap R_{-\nu}^{c} \cap \Delta^{c}, Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1 \text{ and } Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1 \text{ and } Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y})) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (similarly as in the previous } \mathcal{O}(\mathbf{x}, \mathbf{y}) = 1 \text{ (sin the previous } \mathcal{O}(\mathbf{x}, \mathbf{$ cases). Therefore, for all $(\mathbf{x}, \mathbf{y}) \in \{\partial X \times \partial X\} \cap R \cap R_{\nu} \cap R_{-\nu}^{c} \cap \Delta^{c}, \beta(\mathbf{x}, \mathbf{y}) = 1$. In the fourth equality, we used that $\int_{R} \phi r \, d\mu = \int_{R} \phi \, d\mu^{T}$ for all ϕ bounded measurable. Putting those results together yields

$$\int_{\{\partial X \times \partial X\} \cap R} \phi \, r_{\text{rev.}} \, d\mu_{\text{rev.}} = \int_{\{\partial X \times \partial X\} \cap R \cap \Delta} \phi(\mathbf{x}, \mathbf{y}) \, \pi(d\mathbf{y}) \, (1/2) \, \delta_{\mathbf{y}}(d\mathbf{x}) \\ + \int_{\{\partial X \times \partial X\} \cap R \cap \Delta^{c}} \phi(\mathbf{x}, \mathbf{y}) \, \pi(d\mathbf{y}) \, (1/2) \, Q(\mathbf{y}, d\mathbf{x}) = \int_{\{\partial X \times \partial X\} \cap R} \phi \, d\mu_{\text{rev.}}^{T}$$

using similar arguments as above.

This concludes the proof by additivity after having noted that

$$\mathcal{X} \times \mathcal{X} = \{ \mathring{\mathcal{X}} \times \mathring{\mathcal{X}} \} \cup \{ \mathring{\mathcal{X}} \times \partial \mathcal{X} \} \cup \{ \partial \mathcal{X} \times \mathring{\mathcal{X}} \} \cup \{ \partial \mathcal{X} \times \partial \mathcal{X} \}.$$

Proof of Proposition 4. As mentioned in Section A.1, to prove that P_{Lifted} satisfies a skewed detailed balance, we need to show that the two following measures on the product space ($X^2, X \otimes X$) are equal:

$$\mu_{+1}(A \times B) = \int_A \pi(\mathbf{d}\mathbf{x}) T_{+1}(\mathbf{x}, B) \text{ and } \mu_{-1}(A \times B) = \int_B \pi(\mathbf{d}\mathbf{x}) T_{-1}(\mathbf{x}, A),$$

which ensures that P_{Lifted} leaves the distribution $\pi \otimes \mathcal{U}\{-1, +1\}$ invariant. To achieve this, we prove that for any bounded measurable function $\phi : \mathcal{X}^2 \to \mathbb{R}$,

$$\int \phi \, \mathrm{d}\mu_{+1} = \int \phi \, \mathrm{d}\mu_{-1}$$

Let us define

$$G_{+1} := \{ (\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{x} \in \mathring{\mathcal{X}} \text{ or } \mathbf{x} \in \partial \mathcal{X} \text{ with } Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 1 \},\$$

and

$$G_{-1} := \{ (\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{y} \in \mathring{\mathcal{X}} \text{ or } \mathbf{y} \in \partial \mathcal{X} \text{ with } Q(\mathbf{y}, \mathbf{N}_{-1}(\mathbf{y})) = 1 \}.$$

We have that

$$\int \phi \, \mathrm{d}\mu_{+1} = \int_{G_{+1}} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y})$$
$$= \int_{G_{+1} \cap R \cap R_{+1} \cap R_{-1}^{\mathsf{c}} \cap \Delta^{\mathsf{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}).$$

In the first equality, we used the definitions of μ_{+1} and T_{+1} . In the second equality, we used that $\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = 0$ when $(\mathbf{x}, \mathbf{y}) \notin R$, that $\pi(d\mathbf{x}) Q_{+1}(\mathbf{x}, d\mathbf{y})$ assigns null measure to $G_{+1} \cap (R_{+1} \cap R_{-1}^{c})^{c}$ and $G_{+1} \cap \Delta$, as we now prove, and a similar argument as in (11). Let us recall that it has been shown in Section 4.2 that, if (\mathbf{x}, ν) is such that $\mathbf{x} \in \mathring{X}$ or $\mathbf{x} \in \partial X$ with $Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})) = 1$, then $Q_{\nu}(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x})^{c}) = 0$, $Q_{\nu}(\mathbf{x}, \mathbf{N}_{-\nu}(\mathbf{x})) = 0$ and $Q_{\nu}(\mathbf{x}, \{\mathbf{x}\}) = 0$. Let

$$G_{+1}^{\mathbf{x}} := \{ \mathbf{x} \in \mathcal{X} : \mathbf{x} \in \mathcal{X} \text{ or } \mathbf{x} \in \partial \mathcal{X} \text{ with } Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 1 \}.$$

Recall the definitions of R_{ν} and Δ . We have that

$$\begin{split} &\int_{G_{+1}\cap(R_{+1}\cap R_{-1}^{c})^{c}} \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{+1}^{c} \cup R_{-1}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &\leq \int_{G_{+1}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{+1}^{c}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}) + \int_{G_{+1}} \mathbb{1}((\mathbf{x}, \mathbf{y}) \in R_{-1}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1}^{\mathbf{x}}} \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})^{c}) + \int_{G_{+1}^{\mathbf{x}}} \pi(\mathrm{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x})) = 0, \end{split}$$

given that $R_{\nu}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^{2} : \mathbf{y} \in \mathbf{N}_{\nu}(\mathbf{x})\}^{c} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^{2} : \mathbf{y} \notin \mathbf{N}_{\nu}(\mathbf{x})\}.$ Also,

$$\int_{G_{+1}\cap\Delta} \pi(\mathbf{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathbf{d}\mathbf{y}) = \int_{G_{+1}} \mathbb{1}(\mathbf{x} = \mathbf{y}) \, \pi(\mathbf{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \mathbf{d}\mathbf{y}) = \int_{G_{+1}^{\mathbf{x}}} \pi(\mathbf{d}\mathbf{x}) \, Q_{+1}(\mathbf{x}, \{\mathbf{x}\}) = 0.$$

Let $G_{+1}^1 := \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{x} \in \mathring{\mathcal{X}}\}$ and $G_{+1}^2 := \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}^2 : \mathbf{x} \in \partial \mathcal{X} \text{ with } Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 1\}$ with $G_{+1}^1 \cup G_{+1}^2 = G_{+1}$ and $G_{+1}^1 \cap G_{+1}^2 = \emptyset$. We have that

$$\begin{split} &\int_{G_{+1}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,Q_{+1}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1}^{1}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,Q_{+1}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &+ \int_{G_{+1}^{2}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,Q_{+1}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1}^{1}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,(Q_{+1}(\mathbf{x},\mathrm{d}\mathbf{y}) + Q_{-1}(\mathbf{x},\mathrm{d}\mathbf{y})) \\ &+ \int_{G_{+1}^{2}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,(Q_{+1}(\mathbf{x},\mathrm{d}\mathbf{y}) + \delta_{\mathbf{x}}(\mathrm{d}\mathbf{y})) \\ &= \int_{G_{+1}^{1}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,2Q_{\mathrm{rev.}}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &+ \int_{G_{+1}^{2}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,\pi(\mathrm{d}\mathbf{x}) \,2Q_{\mathrm{rev.}}(\mathbf{x},\mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1}\cap R\cap R_{+1}\cap R_{-1}^{c}\cap\Delta^{c}} \phi(\mathbf{x},\mathbf{y}) \,\alpha_{\mathrm{rev.}}(\mathbf{x},\mathbf{y}) \,2\mu_{\mathrm{rev.}}(\mathrm{d}\mathbf{x},\mathrm{d}\mathbf{y}), \end{split}$$

using the definitions of $Q_{\rm rev.}$ and $\mu_{\rm rev.}$. In the second equality, we used that

$$\int_{G_{+1}^1 \cap R \cap R_{+1} \cap R_{-1}^c \cap \Delta^c} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, Q_{-1}(\mathbf{x}, \mathrm{d}\mathbf{y}) = 0,$$

and

$$\int_{G_{+1}^2 \cap R \cap R_{+1} \cap R_{-1}^c \cap \Delta^c} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) \, \pi(\mathrm{d}\mathbf{x}) \, \delta_{\mathbf{x}}(\mathrm{d}\mathbf{y}) = 0,$$

where the latter follows directly from the definition of Δ^c . The former follows from the fact that, for any $\mathbf{x} \in \mathring{X}$, $Q_{-1}(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 0$ and a similar argument as above.

Therefore,

$$\begin{split} \int \phi \, \mathrm{d}\mu_{+1} &= \int_{G_{+1} \cap R \cap R_{+1} \cap R_{-1}^{\mathrm{c}} \cap \Delta^{\mathrm{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y}) \, 2\mu_{\mathrm{rev.}}(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{G_{+1} \cap R \cap R_{+1} \cap R_{-1}^{\mathrm{c}} \cap \Delta^{\mathrm{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \varphi(1/r_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y})) \, 2 \, r_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y}) \, \mu_{\mathrm{rev.}}(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{G_{-1} \cap R \cap R_{+1} \cap R_{-1}^{\mathrm{c}} \cap \Delta^{\mathrm{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \varphi(1/r_{\mathrm{rev.}}(\mathbf{x}, \mathbf{y})) \, 2 \, \mu_{\mathrm{rev.}}^{T}(\mathrm{d}\mathbf{x}, \mathrm{d}\mathbf{y}) \\ &= \int_{G_{-1} \cap R \cap R_{+1} \cap R_{-1}^{\mathrm{c}} \cap \Delta^{\mathrm{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \varphi(r_{\mathrm{rev.}}(\mathbf{y}, \mathbf{x})) \, 2 \, \pi(\mathrm{d}\mathbf{y}) \, Q_{\mathrm{rev.}}(\mathbf{y}, \mathrm{d}\mathbf{x}) \\ &= \int_{G_{-1} \cap R \cap R_{+1} \cap R_{-1}^{\mathrm{c}} \cap \Delta^{\mathrm{c}}} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\mathrm{rev.}}(\mathbf{y}, \mathbf{x}) \, \pi(\mathrm{d}\mathbf{y}) \, Q_{-1}(\mathbf{y}, \mathrm{d}\mathbf{x}) \\ &= \int_{G_{-1}} \phi(\mathbf{x}, \mathbf{y}) \, \alpha_{\mathrm{rev.}}(\mathbf{y}, \mathbf{x}) \, \pi(\mathrm{d}\mathbf{y}) \, Q_{-1}(\mathbf{y}, \mathrm{d}\mathbf{x}) = \int \phi \, \mathrm{d}\mu_{-1}. \end{split}$$

29

In the second equality, we used the definitions of $\alpha_{rev.}$ and φ . In the third equality, we used Proposition 3. In the fourth equality, we used that, for all $(\mathbf{x}, \mathbf{y}) \in R \cap R_{+1} \cap R_{-1}^{c} \cap \Delta^{c}$, $(\mathbf{y}, \mathbf{x}) \in R \cap R_{-1} \cap R_{+1}^{c} \cap \Delta^{c}$ and

$$\frac{1}{r_{\text{rev.}}(\mathbf{x},\mathbf{y})} = \frac{1}{r(\mathbf{x},\mathbf{y})} \frac{Q(\mathbf{y},\mathbf{N}_{-1}(\mathbf{y}))}{Q(\mathbf{x},\mathbf{N}_{+1}(\mathbf{x}))} = r(\mathbf{y},\mathbf{x})\beta(\mathbf{y},\mathbf{x}) = r_{\text{rev.}}(\mathbf{y},\mathbf{x}).$$

We also used that, for all $(\mathbf{x}, \mathbf{y}) \in G_{+1} \cap R \cap R_{+1} \cap R_{-1}^c \cap \Delta^c$, $Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) > 0$ and $Q(\mathbf{y}, \mathbf{N}_{-1}(\mathbf{y})) > 0$ according to the properties of the directional neighbourhoods $\{\mathbf{N}_{+1}(\mathbf{x})\}$ and $\{\mathbf{N}_{-1}(\mathbf{x})\}$, implying that $G_{+1} \cap R \cap R_{+1} \cap R_{-1}^c \cap \Delta^c = G_{-1} \cap R \cap R_{+1} \cap R_{-1}^c \cap \Delta^c$. Finally, we used the definition of μ_{rev}^T . In the fifth and sixth equalities, we used symmetrical arguments as above by recalling that $(\mathbf{x}, \mathbf{y}) \in R_{\nu}$ if and only if $(\mathbf{y}, \mathbf{x}) \in R_{-\nu}$. In the final equality, we used the definitions of μ_{-1} and T_{-1} . This concludes the proof.

Proof of Theorem 2. With some abuse of notation, let, for $\mathbf{x} \in \mathcal{X}$, $R_{\mathbf{x}} := {\mathbf{y} \in \mathcal{X} : (\mathbf{x}, \mathbf{y}) \in R}$. Let us first consider the case where $\mathbf{x} \in \mathcal{X}$. For $A \in \mathbf{X}$, we have that

$$2P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) = 2 \int_{A \setminus \{\mathbf{x}\}} Q_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) \,\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = \sum_{\nu \in \{-1, +1\}} \int_{A \setminus \{\mathbf{x}\}} Q_{\nu}(\mathbf{x}, d\mathbf{y}) \,\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})$$
$$= \sum_{\nu \in \{-1, +1\}} \int_{A \cap R_{\mathbf{x}} \cap \mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \setminus \{\mathbf{x}\}} Q_{\nu}(\mathbf{x}, d\mathbf{y}) \,\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})$$
$$= \sum_{\nu \in \{-1, +1\}} \int_{A \cap R_{\mathbf{x}} \cap \mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \setminus \{\mathbf{x}\}} Q(\mathbf{x}, d\mathbf{y}) \,\frac{\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))}$$

using the definitions of $P_{\text{rev.}}$ and $Q_{\text{rev.}}(\mathbf{x}, \cdot)$, that, for any $\mathbf{x} \in \mathring{X}$, $Q_{\nu}(\mathbf{x}, \cdot)$ assigns null measure outside $\mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \setminus \{\mathbf{x}\}$ (see Section 4.2) with an argument similar as in (11), and that $\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y}) = 0$ for all $(\mathbf{x}, \mathbf{y}) \notin R$.

For all (\mathbf{x}, \mathbf{y}) with $\mathbf{x} \in \overset{\circ}{\mathcal{X}}$ and $\mathbf{y} \in R_{\mathbf{x}} \cap \mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{\mathsf{c}} \setminus \{\mathbf{x}\},$

$$\begin{aligned} \frac{\alpha_{\text{rev.}}(\mathbf{x}, \mathbf{y})}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} &= \frac{\varphi(r(\mathbf{x}, \mathbf{y})\beta(\mathbf{x}, \mathbf{y}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} \geq \frac{\varphi(r(\mathbf{x}, \mathbf{y})(1 \land \beta(\mathbf{x}, \mathbf{y})))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} \\ &\geq \frac{\varphi(r(\mathbf{x}, \mathbf{y}))(1 \land \beta(\mathbf{x}, \mathbf{y}))}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} \\ &= \varphi(r(\mathbf{x}, \mathbf{y})) \left(\frac{1}{Q(\mathbf{x}, \mathbf{N}_{\nu}(\mathbf{x}))} \land \frac{1}{Q(\mathbf{y}, \mathbf{N}_{-\nu}(\mathbf{y}))}\right) \\ &\geq \varphi(r(\mathbf{x}, \mathbf{y})) = \alpha(\mathbf{x}, \mathbf{y}), \end{aligned}$$

using the definitions of $\alpha_{rev.}$ and β , and Results 1 and 6 of Lemma 1.

Therefore,

$$2P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge \sum_{\nu \in \{-1, +1\}} \int_{A \cap R_{\mathbf{x}} \cap \mathbf{N}_{\nu}(\mathbf{x}) \cap \mathbf{N}_{-\nu}(\mathbf{x})^{c} \setminus \{\mathbf{x}\}} Q(\mathbf{x}, d\mathbf{y}) \alpha(\mathbf{x}, \mathbf{y})$$

=
$$\int_{A \cap R_{\mathbf{x}} \setminus \{\mathbf{x}\}} \left(\mathbbm{1}(\mathbf{y} \in \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})^{c}) + \mathbbm{1}(\mathbf{y} \in \mathbf{N}_{+1}(\mathbf{x}) \cap \mathbf{N}_{-1}(\mathbf{x})^{c}) \right) Q(\mathbf{x}, d\mathbf{y}) \alpha(\mathbf{x}, \mathbf{y})$$

=
$$\int_{A \setminus \{\mathbf{x}\}} Q(\mathbf{x}, d\mathbf{y}) \alpha(\mathbf{x}, \mathbf{y}) = P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\}),$$

using in the first equality that, for any $\mathbf{x} \in \mathcal{X}$,

$$\mathbb{1}(\mathbf{y} \in \mathbf{N}_{-1}(\mathbf{x}) \cap \mathbf{N}_{+1}(\mathbf{x})^{c}) + \mathbb{1}(\mathbf{y} \in \mathbf{N}_{+1}(\mathbf{x}) \cap \mathbf{N}_{-1}(\mathbf{x})^{c}) = 1 \quad \text{for} \quad Q(\mathbf{x}, \cdot) \text{-almost all } \mathbf{y}$$

which can be proved in a similar fashion as Lemma 3, and in the second equality that $\alpha(\mathbf{x}, \mathbf{y}) = 0$ for all $(\mathbf{x}, \mathbf{y}) \notin R$.

The case $\mathbf{x} \notin \mathbf{X}$ is done similarly after noticing that, in this case, there exists $v \in \{-1, +1\}$ such that $Q(\mathbf{x}, \mathbf{N}_{-v}(\mathbf{x})) = 0$ and $Q(\mathbf{x}, \mathbf{N}_{v}(\mathbf{x})) = 1$, implying that $Q(\mathbf{x}, A) = Q_{v}(\mathbf{x}, A)$.

Under the general framework of Section 4, the situation of perfect balance corresponds to that where $Q(\mathbf{x}, \mathbf{N}_{-1}(\mathbf{x})) = Q(\mathbf{x}, \mathbf{N}_{+1}(\mathbf{x})) = 1/2$ for all $\mathbf{x} \in \mathcal{X}$. In this case,

$$\frac{\alpha_{\text{rev.}}(\mathbf{x},\mathbf{y})}{Q(\mathbf{x},\mathbf{N}_{\nu}(\mathbf{x}))} \geq 2\alpha(\mathbf{x},\mathbf{y}),$$

from which we can deduce that $P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\})$.

The result

$$P_{\text{rev.}}(\mathbf{x}, A \setminus \{\mathbf{x}\}) \ge \frac{1}{2} P_{\text{MH}}(\mathbf{x}, A \setminus \{\mathbf{x}\}), \text{ for any } \mathbf{x} \in \mathcal{X} \text{ and } A \in \mathbf{X},$$

implies that, for any *f* such that $\pi(f^2) < \infty$,

$$\operatorname{var}(f, P_{\operatorname{rev.}}) \leq 2\operatorname{var}(f, P_{\operatorname{MH}}) + \operatorname{Var}[f(\mathbf{X})], \quad \mathbf{X} \sim \pi,$$

by Lemma 33 in Andrieu et al. (2018).

The proof of

$$\operatorname{var}_{\lambda}(f, P_{\operatorname{Lifted}}) \leq \operatorname{var}_{\lambda}(f, P_{\operatorname{rev.}}),$$

in the case where $\varphi(r) = 1 \wedge r$ follows from Theorem 7 in Andrieu and Livingstone (2021). It can be readily verified that the result holds for any function φ considered here using the same proof technique as in Andrieu and Livingstone (2021). The proof is concluded by taking the limit $\lambda \to 1$ on both side of the inequality.

B No Peskun ordering in all generality

In this section, we show using an example that it is not possible to obtain a Peskun ordering in all generality when considering the general definition of lifted samplers in Andrieu and Livingstone (2021). The Peskun ordering that we consider is the generalized version given in Lemma 33 in Andrieu *et al.* (2018). Let us consider two ergodic reversible transition kernels P_1 and P_2 operating on the same space (X, X) and leaving π invariant. We say that there exists a (generalized) Peskun ordering between P_1 and P_2 if there exists some M > 0 such that for each $B \in X$,

$$P_2(\mathbf{x}, B \setminus \{\mathbf{x}\}) \ge MP_1(\mathbf{x}, B \setminus \{\mathbf{x}\}),$$

for all $\mathbf{x} \in \mathcal{X}$. Conversely, we say that P_1 and P_2 do not admit a Peskun ordering if for all M > 0 (as small as we want), there exists $B_M \in \mathbf{X}$ such that $P_2(\mathbf{x}, B_M \setminus \{\mathbf{x}\}) < MP_1(\mathbf{x}, B_M \setminus \{\mathbf{x}\})$ for some $\mathbf{x} \in \mathcal{X}$.

We now present the example. Let $X = \mathbb{R}$ and $\pi = \mathcal{N}(0, 1)$. Let $Q_{+1}(\mathbf{x}, \cdot) = \mathcal{N}(\mathbf{x}, \sigma^2)$ and $Q_{-1}(\mathbf{x}, \cdot) = \mathcal{N}(\mathbf{x}, 1/\sigma^2)$ with $\sigma \in (0, 1)$. Let $Q = (1/2)Q_{-1} + (1/2)Q_{+1}$. For any $B \in \mathbf{X}$ such that $0 \notin B$, we have that

$$P_{\rm MH}(0,B) = \int_{B} \left(\frac{1}{2} Q_{-1}(0,\mathbf{y}) + \frac{1}{2} Q_{+1}(0,\mathbf{y})\right) \left(1 \wedge \exp\left(-\frac{\mathbf{y}^{2}}{2}\right)\right) d\mathbf{y}$$

32 P. Gagnon and F. Maire

$$= \int_{B} \left(\frac{1}{2} Q_{-1}(0, \mathbf{y}) + \frac{1}{2} Q_{+1}(0, \mathbf{y}) \right) \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) d\mathbf{y},$$

using the symmetry of Q.

We can define a lifted version similarly as in Section A.1. For all $\mathbf{x} \in \mathcal{X}$ and $\nu \in \{-1, +1\}$, let

$$T_{\nu}(\mathbf{x}, \mathrm{d}\mathbf{y}) := Q_{\nu}(\mathbf{x}, \mathbf{y}) \left(1 \wedge \frac{\pi(\mathbf{y}) \, Q_{-\nu}(\mathbf{y}, \mathbf{x})}{\pi(\mathbf{x}) \, Q_{\nu}(\mathbf{x}, \mathbf{y})} \right) \mathrm{d}\mathbf{y}.$$

Note that the definition does not involve splits of neighbourhoods. As in Section A.1, we can prove that $\mu_{+1}(A \times B) = \int_A \pi(d\mathbf{x}) T_{+1}(\mathbf{x}, B)$ and $\mu_{-1}(A \times B) = \int_B \pi(d\mathbf{x}) T_{-1}(\mathbf{x}, A)$ are equal, which implies that

$$P_{\text{Lifted}}((\mathbf{x},\nu), \mathbf{d}(\mathbf{y},\nu')) := \delta_{\nu}(\nu') T_{\nu}(\mathbf{x}, \mathbf{d}\mathbf{y}) + \delta_{-\nu}(\nu') \delta_{\mathbf{x}}(\mathbf{d}\mathbf{y}) (1 - T_{\nu}(\mathbf{x}, \mathbf{X}))$$

leaves the distribution $\pi \otimes \mathcal{U}\{-1, +1\}$ invariant.

The reversible counterpart to the lifted sampler is defined as

$$P_{\text{rev.}}(\mathbf{x}, d\mathbf{y}) := \frac{1}{2} T_{+1}(\mathbf{x}, d\mathbf{y}) + \frac{1}{2} T_{-1}(\mathbf{x}, d\mathbf{y}) + \delta_{\mathbf{x}}(d\mathbf{y}) \left(1 - \frac{1}{2} T_{+1}(\mathbf{x}, \mathbf{X}) - \frac{1}{2} T_{-1}(\mathbf{x}, \mathbf{X}) \right).$$

This definition can be seen as being similar to that in Section A.1 and, like in that section, it can be proved that $P_{\text{rev.}}$ is π -reversible.

For the same $B \in \mathbf{X}$ as above with $0 \notin B$, we have that

$$P_{\text{rev.}}(0, B) = \int_{B} \frac{1}{2} Q_{+1}(0, \mathbf{y}) \left(1 \wedge \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \frac{Q_{-1}(\mathbf{y}, 0)}{Q_{+1}(0, \mathbf{y})} \right) d\mathbf{y} + \int_{B} \frac{1}{2} Q_{-1}(0, \mathbf{y}) \left(1 \wedge \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \frac{Q_{+1}(\mathbf{y}, 0)}{Q_{-1}(0, \mathbf{y})} \right) d\mathbf{y}$$

We have that

$$\exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{-1}(\mathbf{y},0)}{Q_{+1}(0,\mathbf{y})} = \sigma^2 \exp\left(\frac{\mathbf{y}^2}{2}\left(\frac{1}{\sigma^2} - \sigma^2 - 1\right)\right),$$

and

$$\exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{+1}(\mathbf{y},0)}{Q_{-1}(0,\mathbf{y})} = \frac{1}{\sigma^2}\exp\left(-\frac{\mathbf{y}^2}{2}\left(\frac{1}{\sigma^2}+1-\sigma^2\right)\right).$$

Note that

$$\frac{1}{\sigma^2} + 1 - \sigma^2 > 0$$

as $\sigma \in (0, 1)$. We can choose σ small enough so that

$$\frac{1}{\sigma^2} - \sigma^2 - 1 > 0$$

We choose σ accordingly. Therefore, we can choose $B = \{\mathbf{y} \in \mathbb{R} : |\mathbf{y}| > k\}$ with k a large enough constant, so that, for all $\mathbf{y} \in B$, we have that

$$\exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{-1}(\mathbf{y},0)}{Q_{+1}(0,\mathbf{y})} \ge 1$$
 and $\exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{+1}(\mathbf{y},0)}{Q_{-1}(0,\mathbf{y})} \le 1.$

Note that $0 \notin B$. With such a set *B*, we have that

$$P_{\text{rev.}}(0,B) = \int_{B} \frac{1}{2} Q_{+1}(0,\mathbf{y}) \, \mathrm{d}\mathbf{y} + \int_{B} \frac{1}{2} Q_{+1}(0,\mathbf{y}) \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \mathrm{d}\mathbf{y}.$$

Let M > 0. We have that

$$\begin{aligned} P_{\text{rev.}}(0,B) &- MP_{\text{MH}}(0,B) \\ &= \int_{B} \frac{1}{2} \, Q_{+1}(0,\mathbf{y}) \, \mathrm{d}\mathbf{y} + \int_{B} \frac{1}{2} \, Q_{+1}(0,\mathbf{y}) \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \mathrm{d}\mathbf{y} \\ &- M \int_{B} \left(\frac{1}{2} \, Q_{-1}(0,\mathbf{y}) + \frac{1}{2} \, Q_{+1}(0,\mathbf{y})\right) \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \mathrm{d}\mathbf{y} \\ &= \frac{1}{2} \int_{B} Q_{+1}(0,\mathbf{y}) \left(1 + \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) - M \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \frac{Q_{-1}(0,\mathbf{y})}{Q_{+1}(0,\mathbf{y})} - M \exp\left(-\frac{\mathbf{y}^{2}}{2}\right)\right) \mathrm{d}\mathbf{y} \\ &\leq \frac{1}{2} \int_{B} Q_{+1}(0,\mathbf{y}) \left(1 + \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) - M \exp\left(-\frac{\mathbf{y}^{2}}{2}\right) \frac{Q_{-1}(0,\mathbf{y})}{Q_{+1}(0,\mathbf{y})}\right) \mathrm{d}\mathbf{y} \end{aligned}$$

Note that

$$\exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{-1}(0,\mathbf{y})}{Q_{+1}(0,\mathbf{y})} = \exp\left(-\frac{\mathbf{y}^2}{2}\right)\frac{Q_{-1}(\mathbf{y},0)}{Q_{+1}(0,\mathbf{y})} = \sigma^2 \exp\left(\frac{\mathbf{y}^2}{2}\left(\frac{1}{\sigma^2} - \sigma^2 - 1\right)\right).$$

Therefore, regardless the value of M, we could have chosen k such that

$$\left(1 + \exp\left(-\frac{\mathbf{y}^2}{2}\right) - M \exp\left(-\frac{\mathbf{y}^2}{2}\right) \frac{Q_{-1}(0, \mathbf{y})}{Q_{+1}(0, \mathbf{y})}\right) < 0.$$

Consequently, $P_{rev.}(0, B) - MP_{MH}(0, B) < 0$, which concludes the example.