

Orbital dynamics in galactic potentials under mass transfer

Eduárd Illés^{1,2}, Dániel János^{2,3}, and Tamás Kovács^{4,5}

- ¹ Lund University, Division of Astrophysics, Department of Physics, Box 43, Lund SE-221 00, Sweden
² Department of Theoretical Physics, Eötvös Loránd University, Pázmány Péter sétány 1A, H-1117 Budapest, Hungary
e-mail: ed8687il-s@student.lu.se
³ HUN-REN Institute of Earth Physics and Space Science, Csatkai Endre utca 6-8, H-9400 Sopron, Hungary
e-mail: daniel.janos@ttk.elte.hu
⁴ Department of Atomic Physics, Eötvös Loránd University, Pázmány Péter sétány 1A, H-1117 Budapest, Hungary
⁵ HUN - REN – ELTE Extragalactic Astrophysics Research Group, Pázmány P.s. 1A, H-1117, Budapest, Hungary
e-mail: tamas.kovacs@ttk.elte.hu

Received -; accepted -

ABSTRACT

Context. Time-dependent potentials are common in galactic systems that undergo significant evolution, interactions, or encounters with other galaxies, or when there are dynamic processes like star formation and merging events. Recent studies show that an ensemble approach along with the so-called snapshot framework in dynamical system theory provide a powerful tool to analyze time dependent dynamics.

Aims. In this work, we aim to explore and quantify the phase space structure and dynamical complexity in time-dependent galactic potentials consisting of multiple components.

Methods. We apply the classical method of Poincaré-surface of section to analyze the phase space structure in a chaotic Hamiltonian system subjected to parameter drift. This, however, makes sense only when the evolution of a large ensemble of initial conditions is followed. Numerical simulations explore the phase space structure of such ensembles while the system undergoes a continuous parameter change. The pair-wise average distance of ensemble members allows us to define a generalized Lyapunov-exponent, that might also be time dependent, to describe the system stability.

Results. We revise the system parameters for the Milky Way galaxy and provide a comprehensive dynamical analysis of the system under circumstances where linear mass transfer undergoes between the disk and bulge components of the model.

Key words. galaxies: kinematics and dynamics – methods : numerical – chaos

1. Introduction

The theory of stellar orbits in a galactic potential is a crucial aspect of galactic dynamics and astrophysics. It involves understanding how stars and other celestial objects move under the influence of the combined gravitational forces within a galaxy (Binney & Tremaine 2008; Contopoulos 2002). The motion of stars is influenced not only by the central mass concentration (such as a supermassive black hole in the case of many galaxies) but also by the combined gravitational influence of all the stars and dark matter. This leads to complex and chaotic orbital motions. Numerically solving the equations of motion for stars in this potential allows to model and predict the overall behavior of the stellar population and interstellar medium from a dynamical standpoint. Galaxies can be classified based on the symmetry of their mass distribution, usually as axisymmetric (Toomre 1963; Miyamoto & Nagai 1975) and non-axisymmetric potentials (Binney & Spergel 1982; Contopoulos & Seimenis 1990; Cincotta et al. 1996). The shape of the potential significantly affects the stellar orbits. The study of stellar orbits is closely related to the measurement of a galaxy's rotation curve, which plots the orbital velocity of stars as a function of distance from the galactic center. The rotation curve can provide valuable information

about the mass distribution within the galaxy (Verheijen & Sancisi 2001; González et al. 2010).

In a galactic potential, the orbits of stars and other celestial objects can be broadly classified into two categories: regular orbits and chaotic orbits (Lichtenberg & Leiberman 1992; Contopoulos 2002). These classifications depend on the initial conditions of the orbits and the underlying dynamics governed by the gravitational forces within the galaxy. The study of regular and chaotic orbits is a fundamental aspect of galactic dynamics and provides valuable insights into the overall structure, formation, and evolution of galaxies. Small perturbations from nearby stars or external influences can cause deviations in stellar orbits over time. Understanding the stability of orbits is essential in explaining the observed structures and dynamics of galaxies. (Hénon & Heiles 1964; Benettin et al. 1980; Skokos 2010; Froeschlé et al. 1997; Cincotta & Simó 2000; Laskar 1990; Gottwald & Melbourne 2004).

The understanding of the global structure of galaxies has rapidly gained ground with the development of many studies of rotation curves and mass density distributions calculated from observational data, which are relatively easy to interpret and decompose into different galaxy components. The very first models were built from observational data from our own galaxy, the Milky Way (Schmidt

1956; Toomre 1963; Kuzmin 1956), followed by various milestones in radio astronomy and more by the Hipparcos project (Bernacca 1985), several updated mass density models (Bahcall & Soneira 1980; Caldwell & Ostriker 1981; Miyamoto & Nagai 1975), based on data from stars in and near our galaxy and other galaxies. These models are usually completed in the form of a potential, describing the gravitational interactions in the interior of galaxies in a compressed form, thus easily modelling the motions of different stars, clusters and other stellar objects.

In the recent years the re-evaluation and refinement of these models was possible due to more sophisticated and accurate observational tools, of which one of the most important and ambitious projects was the Gaia Survey (Gaia Collaboration et al. 2016), of which one of the goals was to map out the stellar population and obtain valuable parameters, including kinematic data, in great accuracy in order to receive an even more realistic picture of our own galaxy (Katz et al. 2018).

Chaotic orbits arise when the gravitational potential is more complex, such as in regions with irregular mass distributions or significant gravitational perturbations from nearby massive objects. Furthermore, chaotic orbits can arise due to orbital resonances induced by different gravitational influences and other kinematic characteristics of the orbits in a given system, which result in orbital periods related by simple integer ratios between the objects, and in the grand schemes, may lead to the formation of structures in the phase space which give valuable information about the motion of an ensemble in certain regions of the system (Malhotra 1998).

Motion in a time-dependent galactic potential refers to the study of how stars and other celestial objects move in a galaxy when the gravitational potential is not static but changes with time. Time-dependent potentials describe galactic systems that undergo significant evolution, interactions, or encounters with other galaxies, or when dynamic processes like star formation and merging events take place inside certain regions. Analyzing motion in a time-dependent galactic potential can be more challenging than in a static potential because the gravitational forces experienced by objects change over time (Caranicolas (1996); Caranicolas & Papadopoulos (2003); Manos et al. (2013); Manos & Machado (2014); Zotos (2011, 2012, 2014); Zotos et al. (2020)). As a result, the orbits of stars can become more complex and less predictable in larger timescales.

There are many studies in recent past that deal with mass exchange between different parts of a galaxy. For example Caranicolas & Papadopoulos (2003); Zotos (2012) investigated stellar dynamics in cases where low angular momentum stars are scattered through the nucleus into the halo. This process transfers mass from the disk into the nucleus, while the total mass of the galaxy remains unchanged. Their model involves exponential mass transfer as time dependent parameter change in potential. Manos et al. (2013) modeled the results of self-consistent N-body simulations by introducing an analytical time-dependent potential with linear mass transfer between the disk and the bar. They showed that an increase in the mass of the bar leads to a more chaotic dynamics.

When a Hamiltonian system is subjected to a parameter shift, it means that some parameters in the Hamiltonian function are changed with time. These parameter shifts can occur due to various reasons, such as external influences

or changing physical conditions in the system. The specific form of the parameter shift will determine how the Hamiltonian depends on time and this, in turn, affects the system's dynamics (Lichtenberg & Lieberman 1992). Different parameter shifts can lead to various interesting phenomena in Hamiltonian systems such as adiabatic and non-adiabatic parameter change, resonances, and chaos.

A substantial work was done in the past on galactic systems with potentials containing components undergoing perturbations. One of such works was the analysis done by Pichardo et al. (2003), who looked at changes in the systems where an axisymmetric galactic potential was perturbed by three-dimensional galactic spiral arms, which were added as a superposition of inhomogeneous oblate spheroids, introducing a periodic change into the system. This resulted in the emergence of nonlinear effects in the orbits of test particles which showed significant changes in the overall behavior of the system, which can mainly be seen in the Poincaré Surface of Sections shown in the work mentioned above.

An active research area in chaos theory is the study of systems subject to monotonic non-adiabatic parameter changes. In this context, a comprehensive picture of this type of dissipative (i.e. frictional) systems has emerged over the last three decades, motivated by climate change. These systems can be considered as simple models of the Earth's climate when a parameter (e.g. carbon dioxide concentration) varies monotonically. Here, it was shown that a long-time study of a single trajectory is not representative, instead the relevant phase space structure, the so-called snapshot chaotic attractor, is revealed by tracing an extended trajectory ensemble (Yu et al. 1990; Sommerer & Ott 1993; Kovács 2020). In contrast, the theory of chaotic Hamiltonian (or conservative) systems subject to monotonic parameter changes has been a completely unknown field, but in the last years some discoveries have been made in this area, developing several methods to describe such systems (Jánosi & Tél 2019, 2021, 2022). One of the most important of these is the study of so-called snapshot tori, which are the relevant ensembles in the phase space. These are generalizations of the so-called Kolmogorov-Arnold-Moser (KAM) tori of stationary Hamiltonian systems, and can be defined as closed curves in the phase space of systems with varying parameters, but their shape is not constant and over time their dynamics can become chaotic through a break-up process.

In this work, we aim to explore and quantify the phase space structure and dynamical instability in a time-dependent galactic disk potential. We also aim to demonstrate the efficiency of the recently proposed ensemble description and snapshot method in time-varying Hamiltonian dynamics.

The paper is organized as follows. Section 2 presents the model we use and the parameters obtained from radial velocity fits. Section 3 is devoted to the phase space analysis of the steady-state and time-dependent stellar dynamics. In Section 4 we summarize our results and draw conclusions on the implications.

2. The model

2.1. Potential profiles for the components

In this work the analysis was done on Milky-way like galactic potential models. These models can be described as a combination of three separate potentials, each representing a prominent component of a galactic system: a galactic disk, a central bulge and a dark matter halo component.

2.1.1. Galactic disk

For several decades the most well-known and commonly used galactic potential model was the one described by Miyamoto & Nagai (1975), which is a mathematically simple, yet good approximation when it comes to the numerical analysis of such systems. For the disk components for our constructed models an extended Miyamoto-Nagai potential was chosen, which is derived from the Poisson equation (Vogt & Letelier 2005), and was used in the analysis of 3-dimensional axisymmetric potential models for the Milky Way by Barros et al. (2016).

The equation for the extended Miyamoto-Nagai disk potential is as follows:

$$\Phi_d(R, z) = -\frac{A}{\sqrt{R^2 + (a + \bar{z})^2}} \left(1 + \frac{a(a + \bar{z})}{R^2 + (a + \bar{z})^2} - \frac{1}{3} \frac{a^2 [R^2 - 2(a + \bar{z})^2]}{[R^2 + (a + \bar{z})^2]^2} \right), \quad (1)$$

where R and z are the cylindrical coordinates, A is the mass of the disk component multiplied by the gravitational constant G , while $\bar{z} = \sqrt{z^2 + b^2}$. The a parameter is the radial scaling length, while the b parameter is the scaling height, as it is commonly denoted in the Miyamoto-Nagai model.

2.1.2. Bulge

For the central bulge, which is commonly approximated as a spheroidal component, we adopted the Hernquist potential as a simple alternative (Hernquist 1990).

The potential takes the following form:

$$\Phi_b(R, z) = -\frac{C}{\sqrt{R^2 + z^2 + c}}, \quad (2)$$

where C again is a mass multiplied by the gravitational constant G , while c is the scaling radius of the spheroidal bulge.

2.1.3. Dark matter halo

For the dark matter halo component a Navarro-Frenk-White profile (Navarro et al. 1996) was chosen, written in terms of the scaling radius r_s and the enclosed mass M_s (Sanderson et al. 2017), instead of the commonly used form determined by the scale density ρ_s or virial radius r_{vir} and mass M_{vir} , in order to keep the potential components mathematically consistent. The equation for the scaling mass is as follows:

$$M_s = 4\pi\rho_s r_s^3 (\ln 2 - 1/2), \quad (3)$$

and the NFW profile in terms of M_s can be written in the following form using the cylindrical coordinate system:

$$\Phi_{DM}(R, z) = -\frac{D}{\ln 2 - 1/2} \frac{\ln(1 + \sqrt{R^2 + z^2}/d)}{\sqrt{R^2 + z^2}}, \quad (4)$$

where $D = GM_s$ is the enclosed mass multiplied by the gravitational constant G and $d = r_s$.

The overall model of a Milky Way-like disk galaxy is simply the sum of the disk, bulge and dark matter halo potentials written in Eqs. (1, 2, 4):

$$\Phi(R, z) = \Phi_d + \Phi_b + \Phi_{DM}. \quad (5)$$

2.2. Equations of motion

The motion of a test particle (star or interstellar object) can be described by the following Lagrangian, with $L_z = p_\phi$ being a constant representing the angular momentum component around the z -axis since the models under consideration exhibit axial symmetry (more precisely, cylindrical symmetry):

$$\mathcal{L} = \frac{1}{2} (\dot{R}^2 + \dot{z}^2) - \Phi(R, z), \quad (6)$$

where (R, z) represent the cylindrical coordinates, and we incorporate the summed potential given by Eq. (5), into the expression for $\Phi(R, z)$. The corresponding generalized momenta:

$$p_R = \dot{R}, \quad p_z = \dot{z}, \quad (7)$$

and thus, the Hamiltonian function can be written as:

$$\mathcal{H} = \frac{1}{2} (p_R^2 + p_z^2) + \Phi_{\text{eff}}(R, z), \quad (8)$$

where the effective potential Φ_{eff} is given by:

$$\Phi_{\text{eff}}(R, z) = \frac{L_z^2}{2R^2} + \Phi(R, z), \quad (9)$$

Thus, out of the original 6 generalized coordinates, only 4 remain, allowing us to define 4 differential equations describing the motion:

$$\begin{aligned} \dot{R} &= p_R, \quad \dot{z} = p_z, \\ \dot{p}_R &= \frac{L_z^2}{R^3} - \frac{\partial \Phi(R, z)}{\partial R}, \quad \dot{p}_z = -\frac{\partial \Phi(R, z)}{\partial z}. \end{aligned} \quad (10)$$

2.3. Fit of radial velocity profile

The rotation curve equation can be obtained by the following general relation:

$$v_c(R) = \left(R \frac{\partial \Phi(R, z)}{\partial R} \Big|_{z=0} \right)^{\frac{1}{2}}. \quad (11)$$

This equation is needed for determining the parameters a, b, c, d, A, C and D . Two main sources of data were used as a combined data set: one is the grand rotation curve of the Milky Way compiled by Sofue (2012, 2013), while the other one is the collection of rotational velocity data

obtained using Classical Cepheids by Mróz et al. (2019), ensuring that the inner parts of the rotation curve around the range of $R = 0 - 20$ kpc is well-captured. After that the parameter space can be explored to find the best-fitting parameters for constructing a model representing a Milky Way-like disk galaxy.

The procedure to find the best parameters was done in two steps:

- First, the rotation curve Eq. (11) was fitted using Truncated Newtonian Method, or TNC, (Dembo & Steihaug 1983) for all the parameters to receive approximate values in preparation for the next step. The initial parameter guesses of scaling distances and masses for the TNC fit were obtained from the DR3+ data analysis results of Labini et al. (2023), while the lower and upper limits for the parameter space were implemented from McMillan (2011). The fits were done using the *lmfit* Python library (Newville et al. 2015).
- After that, the fitted parameters from the TNC were used as initial conditions for an Affine Invariant Markov Chain Monte Carlo, or MCMC model (Goodman & Weare 2010). The scaling height b was excluded from this step due to the rotation curve having no information of the exact distribution of the measurements in that space, keeping the obtained parameter from the TNC constant. The software of choice used for performing the MCMC samplings was *emcee* (Foreman-Mackey et al. 2013). The number of walkers were chosen to be 1000, with a sampling time of $t_{smp} = 10^6$ and a burning time of 5 times the time of highest auto-correlation time $t_{ac} \approx 200$. The same lower and upper constraints were implemented for the parameter space in this analysis as in the previous step. The results of this analysis can be seen in Fig. 1, while the obtained parameters are summarised in Table 1.

The rotation curve resulting from the fitting with the parameters of Table 1 is shown in Fig. 2. The fitting was done on the joint datasets of Sofue (2012, 2013) (blue) and Mróz et al. (2019) (orange), and shows quite good agreement with the observations.

3. Results

3.1. Time-independent case

The energy of a trajectory initiated from a particular set of initial conditions can be determined by evaluating the Hamiltonian-function (8) using the given initial conditions. Assuming a drift-free case, the energy of the trajectory is conserved at all times. Therefore, we can express it as follows:

$$\mathcal{H}(R, z, p_R, p_z) = \mathcal{H}(R_0, z_0, p_{R0}, p_{z0}) = H. \quad (12)$$

Merely tracing trajectories is far from sufficient for an extensive examination of a system, where a deeper understanding of a specific galaxy model's morphology and the structures within it can be obtained. Investigating trajectories of a system with multiple initial conditions using physical and phase space plots, depending on the number of initial conditions, may not be efficient or even feasible. Therefore, it is required to continue the investigations using




Fig. 1: The MCMC results on the gathered rotational velocity data illustrated in a corner plot. As expected, the highest correlations can be seen between the scaling distances and mass components corresponding to the same potential components. The red dots in the two-dimensional projection, along with the red lines in the one-dimensional projection, show the Maximum Likelihood estimates for the parameters, along with the black lines showing the median and 1σ confidence interval results.




Fig. 2: Rotation curve fitted to the observational data indicated in the legend, using parameters obtained from the MCMC analysis of Fig. 1 and are shown in Table 1.

the *Poincaré surface of section* method (Ott 1993; Teschl 2012). Each intersection is registered under the conditions $z = 0$ and $p_z > 0$.

In order to obtain the initial conditions for the simulations we firstly need to find the circular orbit at $R_{circ} =$

Table 1: Table of the parameters obtained from the TNC and MCMC methods. The final parameters used in the fitting of the rotation curve are indicated in green.

	Initial guess	TNC	MCMC	
			Median	MLE
a [kpc]	4.5	8.640	$7.300^{+0.0435}_{-0.0433}$	7.304
b [kpc]	0.2	1.865	-	-
c [kpc]	0.25	0.284	$0.266^{+0.00301}_{-0.00298}$	0.267
d [kpc]	183	182	$109^{+10.7}_{-9.69}$	106
A [kpc km s ⁻¹]	$3.0 \cdot 10^{10} G$	$3.77 \cdot 10^5$	$(2.73^{+0.0246}_{-0.0247}) \cdot 10^5$	$2.72 \cdot 10^5$
C [kpc km s ⁻¹]	$2.0 \cdot 10^{10} G$	$7.37 \cdot 10^4$	$(6.63^{+0.0540}_{-0.0537}) \cdot 10^4$	$6.63 \cdot 10^4$
D [kpc km s ⁻¹]	$6.5 \cdot 10^{11} G$	$7.79 \cdot 10^6$	$(9.02^{+1.67}_{-1.39}) \cdot 10^6$	$8.66 \cdot 10^6$

8.249 kpc and $z_{circ} = 0.0208$ kpc, which is the estimated location of our Sun in galactocentric coordinate system from [GRAVITY Collaboration et al. \(2021\)](#), and calculate the radial velocity $V_R \approx 226.77$ km s⁻¹ from the derived rotation curve, which then can be used to acquire the z -component of the angular momentum $L_z = V_R R$. After that the reference Hamiltonian energy for a circular orbit $H_0 = -4.222 \cdot 10^5$ can be calculated using Eq. (8), with $p_{R0}, p_{z0} = 0$.

By lowering the L_z parameter it is possible to create ensemble of orbits which have their Hamiltonian energies distributed to the p_R, p_z velocity components, resulting in orbits which are misaligned with the galactic plane. Stellar streams can have such orbits, and in the last couple of years they have been proven to be useful tools for measuring the Milky Way's gravitational potential ([Sanderson et al. 2017](#); [Walder et al. 2024](#)). Figure A.1 of the Appendix illustrates the effect of a smaller L_z .

It is possible to find initial parameter sets by setting $p_{R0} = 0$ and calculating the maximum permitted value for the velocity in the z -direction $p_{z_{max}}$ using Eq. 8, with known H and chosen L_z , and scanning the phase space with an arbitrary resolution in the range of $p_{z0} = [0; p_{z_{max}}]$. By simultaneously going through the range of radial distances $R_0 = [0; R_{circ}]$ it is possible to find all the possible orbits in the phase space with a given H_0 and L_z parameter values.

A total of 400 initial conditions were launched in each chosen L_z, H values. The numerical integrations were performed using the NumbaLSODA integrator ([Wogan 2021](#)), which uses the LSODA algorithm with automatically switching between stiff and nonstiff class of methods during the integration depending on which of the two can more likely solve the equation in most efficient manner, which can especially be beneficial in cases where the nature of the problem is unknown or may change drastically during the chosen time intervals ([Petzold 1983](#)). The simulations in the time-independent case were run for $t = 5000$ with a resolution of $dt = 10^{-4}$, which is sufficient to determine the intersections within a certain time interval, to obtain detailed Poincaré SoSs.

Figure 3 show the phase space plots in the (R, p_R) plane, with a colormap representing the value of p_z at each point, obtained from Eq. (12) with $z = 0$. With a chosen reference of $L_z = 100$ we illustrate the phase portrait for four different values of constant H . The red curves are highlighted KAM tori of initial conditions $R_0 = 0.7211$, $p_{R0} = -59.3423$ (Fig. 3a) and $R_0 = 1.9632$, $p_{R0} = -80.5212$ (Fig. 3c). The analysis of the time-dependent dynamics to follow in the next section will be done on these curves.




Fig. 3: Phase portraits of Eq. (10) for different values of H with constant $L_z = 100$. The trajectories are plotted on the (R, p_R) plane, following the Poincaré cut at $z = 0$, and the value of p_z obtained from (12), indicated at each point by the colormap. The red KAM tori belong to initial conditions $R_0 = 0.7211$, $p_{R0} = -59.3423$ (panel a) and $R_0 = 1.9632$, $p_{R0} = -80.5212$ (panel c).

One can observe that the system possesses a typical divided phase space, with the coexistence of four types of dynamics. Quasi-periodic motion occurs on the closed curves, i.e. the KAM tori, where trajectories densely fill up these shapes. These tori form concentric islands, where the radius of the shapes depends on the potential energy of the trajectories, which is conserved within a curve. In the center of such islands one finds so-called elliptic fixed points, representing periodic trajectories.

The third type of dynamics is chaos, which manifests in this illustration as dense bands of irregularly mixed trajectories, called the chaotic sea, observable along the edge of the phase space, between the large central and several smaller islands. Motion within this region is ergodic, meaning that trajectories uniformly cover the chaotic sea, and have equal probability to reach any parts of it. Here the dynamics is unpredictable, and trajectories, even if they were very close initially, tend to diverge at an exponential rate.

Although the shown phase portraits do not capture this explicitly, it is known from chaos theory (see e.g. [Ott](#)

(1993)) that a peculiar phenomenon occurs on the border between the island of KAM tori and the chaotic sea. In a narrow band outside the outermost KAM torus the trajectories do not form any comprehensive pattern, they are scattered just like in the chaotic sea, however their dynamics is not yet chaotic, they do not diverge exponentially. Orbits are able to detach from this region into the chaotic sea (and vice versa), however the transport of trajectories in this direction was found to follow a power-law (Channon & Lebowitz 1980; Karney 1983; Meiss et al. 1983; Cristadoro & Ketzmerick 2008; Altmann et al. 2013), which is much slower than the ergodic wandering in the chaotic sea. Because of this, chaotic trajectories can be "trapped" inside these regions and "stick" to the outermost torus for long times. This is the fourth type of motion present in the phase space, and is often referred to as sticky dynamics.

3.2. Parameter drift and EAPD-curves

The simulations conducted in Section 3.1 were calculated by keeping the parameters received from the model fittings fixed, assuming that the physical properties associated with these parameters, as well as those derived from them, do not change over time. The changes occurring in these systems are generally considered relatively slow, making these assumptions reasonable approximations for extrapolating behaviors of certain galaxies, which can be a considered isolated and stable for a longer period of time when it comes to the stage of stellar evolution.

In this section, we assign a time dependence to the A and C parameters by introducing linear drifts as

$$A(t) = A(0) - \mu t, \quad C(t) = C(0) + \mu t, \quad (13)$$

that is, the decrease in $A(t)$ and the increase in $C(t)$ has the *same* μ rate, i.e. we simulate a *mass transfer* from the disk to the bulge, so that the overall mass of the galaxy is conserved. The $A(0), C(0)$ initial values are read from Table 1, and the rate is chosen to be $\mu = \frac{3}{5}A(0) \cdot \frac{1}{T_r}$, with $T_r = 200$ Gyr.

Following the methods used in János & Tél (2019, 2021); János et al. (2021); János & Tél (2022), we have to accept that simulating individual trajectories is no longer reliable. Instead, one has to follow an *ensemble* of trajectories in order to obtain sufficient statistics. Furthermore, in Hamiltonian systems the shape of this ensemble is not arbitrary, they have to be initiated on e.g. KAM tori of the stationary system. In our case, we will focus on the two red tori highlighted in Fig. 3. The analysis includes studying the temporal evolution of the tori, whose shapes become time-dependent, and are called snapshot tori. It was found that at some point, the dynamics of these snapshot tori becomes chaotic, indicated by a strong stretching in their shape. This is called the break-up of snapshot tori, and it ends with them becoming entrained into what is called the snapshot chaotic sea, the time-dependent image of the stationary chaotic sea, whose shape is also changing in time.

It should also be noted that the snapshot picture taken on consecutive Poincaré-sections implies intersections at various time instants, which are in different stages of the parameter drift. Consequently, the energy does not remain constant. Therefore, what we depict is a projection of the (R, p_R, p_z) phase space onto the (R, p_R) plane, which has evolved from a stationary case Poincaré-section.

The most straightforward approach to elucidating our simulation results is through visualization. In Figs. 4-7, the time-dependent simulations showcase the time evolution of snapshot tori in the galactic model previously derived and discussed. From such images we can easily acquire geometrical information about the general behavior the evolution of the selected ensemble throughout the integration time.

In Fig. 4 we follow the snapshot torus whose initial image is the red KAM torus shown in Fig. 3c. Its evolution is similar to those shown in János & Tél (2019, 2021, 2022): for some time it does not appear to show any peculiar change in its dynamics, which can persist for longer times, like we see here at $n = 687$, although one can observe slight deformations in its shape, particularly on the left. However, this suddenly changes when the torus strongly stretches in multiple directions ($n = 760$), followed by disintegration and in the end fully chaotic dynamics ($n = 1634$).

For a quantitative analysis, we turn to the *ensemble-averaged pairwise distance* (EAPD) method (János & Tél 2019, 2021, 2022), which allows measuring the dynamical instability of time-dependent chaos. This method can be seen as a generalized form of the concept of Lyapunov-exponents, where the fundamental behavior of the system (or significant parts of it) can be deduced from the ensemble average of distances between pairs of points in the phase space. In most analysis methods, these distances are evaluated on the projections of phase space. The definition of EAPD is as follows:

$$\rho_n = \langle \ln r_n \rangle, \quad (14)$$

where the bracket denotes the average over the ensemble, $\ln r_n$ represents the natural logarithm at the n -th Poincaré section of the phase space distance r_n between one point in the snapshot torus, and its assigned pair of initial distance $r_0 = \sqrt{2} \cdot 10^{-12}$, which is not on the snapshot torus itself but very close to it, which turns out to be a good approximation for our purposes. If the temporal change of the distances follows an exponential trend, we can fit a linear function defined on a natural semi-logarithmic scale, where the slope of the function is the so-called instantaneous Lyapunov exponent denoted by λ . In the above case λ is constant, while in general, the instantaneous Lyapunov exponent is given as the derivative of the EAPD function,

$$\lambda_n = \dot{\rho}_n, \quad (15)$$

where the dot denotes the time-derivative evaluated in discrete time. This approach allows making assumptions about the system in a manner similar to the traditional counterpart. If the r_n distances diverge linearly during the time evolution, it indicates regular or periodic behavior, whereas in chaotic cases, it results in exponential distance growth.

During the calculations of distances in the phase space at the moments of intersections it is important to take the differences in units into account, since it is evident that the velocity components p_R, p_z would overbear the distances in R . In order to remedy that problem all the phase space dimensions are standardized using z-score transform, so that the standard deviations and means of phase space coordinates are comparable to each other. This also means, that the distance growths in the following EAPD curves are not direct indications of the distances on the Poincaré sections, rather it is a normalized approach which, however, does capture the overall form of the evolution of distances in the actual phase space.

$$L_z = 100, H(t=0) = 1.05 \times H_0$$




Fig. 4: Time evolution of the red torus on Fig. 3c. For around 700 iterations it only deforms slightly, but at $n = 760$ it starts to break up, resulting in the dynamics of more and more trajectories of the snapshot becoming chaotic.

The EAPD function evaluated for the snapshot torus of Fig. 4 is shown in Fig. 5. This means that the average denoted by the bracket in (14) was evaluated on the ensemble representing this torus. One can clearly see the sudden increase in distance starting from $n = 760$. The slope of the linear fit is $\lambda = 0.20198 \pm 0.00203$. After this, the curve goes into saturation, since the available phase space is of fi-

nite volume, thus once all trajectories of the snapshot torus become chaotic, their average distance cannot grow any further. This can be thought of as the typical behavior of snapshot tori in Hamiltonian systems subjected to parameter drift, at least in cases where the dynamics tends towards stronger chaos as the consequence of the drift, which seems to be the case here. For example, notice that in Fig. 3c the red torus is in the middle of the island, surrounded by other KAM tori. However, by the end of the parameter drift scenario at $n = 1634$ it has dissolved into a chaotic sea that spans the whole outer region of the phase space, meaning that all of the tori that were outside the red one had to have broken up at some point.

Note that during the period when the dynamics is still regular (i.e. before $n = 760$) the distance between point pairs does grow, although slowly. An explanation for this could be that since the assigned pairs for each point are not exactly on the snapshot torus, during the slowly deforming phase they drift slightly further apart from the torus. Also, there is a short jump at the very beginning of the curve, which likely happens because the initial assigned pairs technically correspond to different tori (since they continuously cover the island), possessing different winding numbers from the one investigated.




Fig. 5: EAPD curve of Eq. (14), evaluated for the snapshot torus of Fig. 4. The Lyapunov exponent obtained from the fitting is $\lambda_{n_r} = 0.20198 \pm 0.00203$.

The chosen form of the parameter drift in (13) representing a mass transfer makes for a non-trivial dynamics even compared to previous studies (János & Tél (2019, 2021, 2022)), where a simple linear drift in only one parameter was applied, typically obtaining results similar to Figs. 4 and 5. In our case however, we found that the inclusion of just one more time-dependent parameter, even with the same drift rate, leads to more diverse dynamics on a broad scale of parameters H and L_z . We provide an example of such peculiar dynamics in Fig. 6 by following the snapshot torus initiated from the red KAM torus in Fig. 3a. Another example is shown in Figs. A.2 and A.3 of the Appendix.

The first stage of its evolution, until about $n = 111$, is characterized not by deformations in its shape, but by small crinkles developing on its left side, the shape of which

quickly dissolve, creating a narrow band (panel c). Next, the corners of the torus extend, until about $p_R = \pm 400$, while some trajectories remain on the previous border of the torus, essentially closing in parts of the phase space. A similar thing happens around $p_R = 0$ on the left. The created three "closed curves" then seem to be evolving on their own (panels d,e), inside the outer points of the snapshot torus, until about $n = 166$, when they experience stretching, as if they were separate snapshot tori themselves, and start to dissolve. By around $n = 559$ the whole snapshot torus breaks up.

The behavior of the snapshot torus between $n = 132$ and $n = 166$ can be explained if one takes a look at the initial phase space of Fig. 3a. Outside of the red torus, about $p_R = \pm 400$ there are two smaller islands. Within these islands there are KAM tori just like in the largest one, and if we were to choose these tori as initial ensembles, they would become snapshot tori as well. In this respect, in the time-dependent scenario both islands move and deform in the phase space, initially on the outside of the snapshot torus in Fig. 6. However, what ends up happening is that points of the large snapshot torus *wrap around* the two smaller islands (as well as a third one in the middle). This state holds until $n = 166$ when the snapshot tori within the small islands start to break up, taking with them the points of the large snapshot torus around them.

The EAPD curve for this snapshot torus is shown in Fig 7, and it is clear that in this case *there is no clear section of exponential divergence* of the trajectories, like the one we saw in Fig. 5. The jump in the beginning can again be explained by the initial difference in winding numbers, but for the rest of the curve the distance only grows slowly, never reaching above $\rho = -10$. This is despite the fact that by the end of the simulation ($n = 559$) the trajectories are spread out in an extended region of the phase space. This means that the dynamics of this snapshot torus is never (fully) chaotic, there has to be some other mechanisms at play.

Let us remember that in time-independent Hamiltonian phase spaces on the border of islands there exist the so-called sticky regions, which exhibit non-chaotic dynamics, outside of which one finds the chaotic sea. For systems subjected to parameter drift, it was found¹ that the picture is even more intricate. It turns out, that even when one chooses a stationary chaotic sea as initial ensemble, when it is evolved in time as a snapshot chaotic sea, some regions in it could exhibit non-chaotic dynamics. These regions are termed time-dependent non-chaotic regions, and trajectories in them can either become chaotic quickly, or stay non-chaotic for longer times. The subsets of time-dependent non-chaotic regions where the slow dynamics is persistent are called time-dependent non-hyperbolic regions, and such latter sets are thought to be the time-dependent analogs of sticky regions. Since on the EAPD curve of Fig. 7 no chaotic section was found, and because in Fig. 6 points of the snapshot torus was observed to wrap around islands of other snapshot tori, we can speculate that the time-dependent non-hyperbolic regions described above, existing around the




Fig. 6: Time evolution of the red torus on Fig. 3a. At $n = 132$ its points start to wrap around the snapshot tori initially belonging to the islands at $p_R = \pm 400$, until these small snapshot tori break up starting at $n = 166$. Points of the large snapshot torus are completely dispersed by instant $n = 559$.

¹ Publication of these results will be included in the review paper "Overview of the advances in understanding chaos in low-dimensional dynamical systems subjected to parameter drift" by D. János and T. Tél, invited for publication in Physics Reports.

smaller islands, might be responsible for slowing down the dynamics of the large snapshot torus.




Fig. 7: EAPD curve of Eq. (14) evaluated for the snapshot torus of Fig. 6. The inset shows an oscillating behavior of small amplitudes after $n = 166$.

3.3. Time-dependent energy

Changing the parameter in a Hamiltonian system results in a violation of energy conservation. The dynamics of the system can be described by the well-known adiabatic invariant for sufficiently slow parameter shifts, which is represented by the volume enclosed by a periodic trajectory in phase space. In this framework, instead of total energy conservation, an adiabatic invariant can be considered a constant of motion, provided that the time scale of the periodic orbit is much smaller than that of the parameter change. The conservation of the action variable depends on the slowness of the parameter variation. [Lichtenberg & Leiberman \(1992\)](#)

In this study, the speed of parameter variation is chosen such that the system evolves beyond the adiabatic limit. That is, we do not expect even the action variable to remain constant during the motion. This fact is also evident from the performed numerical simulations, where not only the geometric structure but also the size of KAM islands changes over time in phase portraits. During the mass exchange between the two parts of the galaxy, Eq. (13), the total energy of the system is monitored. It turns out that - see Figures 8 and 9 - the system becomes increasingly bound as the median energy decreases. The energy curves are plotted alongside the respective EAPD curves of Figs. 5 and 7, in order to compare and highlight relevant parts of the processes. The relative change in total energy averaged over the ensemble is less than 3% by the end of each integration. In Fig. 8, along the energy curve of the snapshot torus shown in Fig. 4, a slight break is visible at the time of torus break-up. This characteristic becomes even more pronounced when considering the energy dispersion (σ) evaluated over the ensemble. The sudden change in σ around the disappearance of the torus can be regarded as an alternative indicator of the dramatic structural evolution of KAM islands in phase space during parameter changes. For the case of the snapshot torus belonging to Fig. 6, slight oscillations around a non-trivial curve can be observed in the

median energy, while the dispersion grows monotonously, with some larger fluctuations around $n = 150$, see Fig. 9.




Fig. 8: Median energy of the ensemble of the snapshot torus shown in Fig. 4 compared to the EAPD curve of Fig. 5. The 1σ dispersion is shown as a light red band around the energy curve, while the vertical dashed line marks $n = 760$, the start of the torus break-up.

The Hamiltonian function (12) determines conservative dynamics while the parameters remain unchanged over time. However, if we allow the mass between the disk and the bulge to evolve, the internal dynamics experience time-dependent perturbation, which affects the initial energy of the system, as shown here. In other words, manually setting the parameters appears as an external disturbance in the dynamics. Consequently, we model stellar dynamics in a time-dependent potential without employing any concrete physical mechanisms. However, similar results are expected if we had taken into account an additional term explicitly in the effective potential Eq. (5) controlling mass transfer within the galaxy.

4. Conclusions

In this paper we have numerically studied ensembles of stellar orbits for the Milky Way with different potential profiles for the galactic components: the extended Myamoto-Nagai potential for the disk, the Hernquist potential for the bulge and the Navarro-Frenk-White potential for the




Fig. 9: Median energy of the ensemble of the snapshot torus shown in Fig. 6 compared to the EAPD curve of Fig. 7. The 1σ dispersion is shown as a light red band around the energy curve, while the vertical dashed lines mark $n = 132, 149, 166$, during the process of the snapshot torus wrapping around smaller islands.

dark matter halo. We simulated a mass transfer from the disk to the bulge by heuristically introducing a decreasing and increasing parameter drift of the same rate in the mass parameters of the disk and the bulge, respectively. For simplicity, for the form of the drift we chose a linear function, but in general cases any monotonous form could be used, obtained from the observations. The chosen rate results in 60% of the disk mass transferred to the bulge over time scales which can be deemed reasonably long compared to the time scales of our Universe (Spergel et al. 1997).

The stellar orbits were illustrated on Poincaré surface of sections in the case of constant mass, and were shown to exhibit a mixed phase space with coexisting periodic, quasi-periodic, chaotic and sticky dynamics. When the parameter drift is introduced, as an appropriate ensemble of initial conditions we chose quasi-periodic KAM tori of the stationary case. In the time-dependent dynamics they become snapshot tori, which change their shape in the phase space and in one of the investigated cases their dynamics becomes chaotic through a break-up process. The torus break-up results in an exponential separation of nearby points on the torus. This is characterized by the so-called EAPD

curve, that describes the evolution of the average distance of point pairs. The slope of this curve can be thought of as a new type of finite-time Lyapunov exponent that characterizes the fate of the torus. In the case mentioned above, this Lyapunov exponent is approximately constant and strictly positive through the break-up process. However, we found that in other cases the EAPD curve does not contain any exponential parts, the dynamics of the torus never turns chaotic. This slow dynamics results in non-trivial Lyapunov-exponents throughout the whole process. We note that the two presented cases are far from fully covering the rich dynamics of the problem, further studies are needed for its comprehensive discovery.

As it was noted in Sec. 3.3, the snapshot pictures comprised of different intersections with various time instants of the parameter drift can result in changing trajectory energies. Future studies will be necessary to further explore the emerging changes induced by different parameter changes in order to better understand the dynamics of time-dependent systems. The change in trajectory energies may play a significant role in inducing local, or even global, changes in the morphology of these galaxy models, like the emergence of more dense star forming regions with distinct structures by purely dynamical causes (Daffern-Powell & Parker (2020); Rieder et al. (2021); Arnold et al. (2022)).

The general aim of this paper is to give, through the presented example, a universal tool for the description of problems arising galactic and celestial mechanics where a parameter drift is present. The method originates in chaos theory, and is a currently actively developed area of research. Some other concepts that could be useful from this method include time-dependent stable and unstable manifolds and foliations, doubly transient chaos (Motter et al. 2013), as well as snapshot attractors in dissipative cases János & Tél (2019); János et al. (2021); János & Tél (2022). Some further methods that could be used for augmenting the presented methods include e.g. developing new ways of phase space projections dependent on trajectory energies, or by expanding the EAPD-analysis with mapping methods. Furthermore, it can be also beneficial to expand the analysis into more composite and realistic N-body simulation models, where the axisymmetric bulge and dark halo components and the non-axisymmetric spiral arms are also included in the potential (Barros et al. (2016); Pouliasis et al. (2016); Trick et al. (2017); Michtchenko et al. (2018)).

Acknowledgements. We are grateful to Tamás Tél and Santi Roca-Fàbrega for their valuable comments, which significantly improved the interpretation of the results. This work was supported by the Hungarian National Research, Development and Innovation Office, under Grants No. KDP-2023 C2262591, TKP-2021 BME-NVA-02 (D.J.) and TKP2021-NKTA-64 (T.K.), financed by the Ministry of Culture and Innovation of Hungary.

References

- Altmann, E. G., Portela, J. S. E., & Tél, T. 2013, *Reviews of Modern Physics*, 85, 869
- Arnold, B., Wright, N., & Parker, R. 2022, *Monthly Notices of the Royal Astronomical Society*, 515
- Bahcall, J. N. & Soneira, R. M. 1980, *The Astrophysical Journal Letters*, 238, L17
- Barros, D. A., Lépine, J. R. D., & Dias, W. S. 2016, *A&A*, 593, A108
- Benettin, G., Galgani, L., Giorgilli, A., & Strelcyn, J. M. 1980, *Mechanica*, 15, 9
- Bernacca, P. L. 1985, *Ap&SS*, 110, 21
- Binney, J. & Spergel, D. 1982, *ApJ*, 252, 308

- Binney, J. & Tremaine, S. 2008, *Galactic Dynamics: Second Edition* (Princeton, USA: Princeton Univ. Press)
- Caldwell, J. A. R. & Ostriker, J. P. 1981, *The Astrophysical Journal Letters*, 251, 61
- Caranicas, N. D. 1996, *Ap&SS*, 246, 15
- Caranicas, N. D. & Papadopoulos, N. J. 2003, *A&A*, 399, 957
- Channon, S. R. & Lebowitz, J. L. 1980, *Annals of the New York Academy of Sciences*, 357, 108
- Cincotta, P. M., Nunez, J. A., & Muzzio, J. C. 1996, *ApJ*, 456, 274
- Cincotta, P. M. & Simó, C. 2000, *A&AS*, 147, 205
- Contopoulos, G. 2002, *Order and chaos in dynamical astronomy* (Berlin, Germany: Springer)
- Contopoulos, G. & Seimenis, J. 1990, *A&A*, 227, 49
- Cristadoro, G. & Ketzmerick, R. 2008, *Phys. Rev. Lett.*, 100, 184101
- Daffern-Powell, E. & Parker, R. 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 4925
- Dembo, R. S. & Steihaug, T. 1983, *Mathematical Programming*, 26, 190
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *Publications of the Astronomical Society of the Pacific*, 125, 306–312
- Froeschlé, C., Lega, E., & Gonzi, R. 1997, *Celestial Mechanics and Dynamical Astronomy*, 67, 41
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1
- González, G. A., Plata-Plata, S. M., & Ramos-Caro, J. 2010, *MNRAS*, 404, 468
- Goodman, J. & Weare, J. 2010, *Communications in Applied Mathematics and Computational Science*, 5, 65
- Gottwald, G. A. & Melbourne, I. 2004, *Proceedings of the Royal Society of London Series A*, 460, 603
- GRAVITY Collaboration, Abuter, R., Amorim, A., et al. 2021, *A&A*, 647, A59
- Henon, M. & Heiles, C. 1964, *AJ*, 69, 73
- Hernquist, L. 1990, *ApJ*, 356, 359
- Jánosi, D., Károlyi, G., & Tél, T. 2021, *Nonlinear Dynamics*, 106, 2781–2805
- Jánosi, D. & Tél, T. 2019, *Chaos*, 29, 121105
- Jánosi, D. & Tél, T. 2021, *Chaos*, 31, 033142
- Jánosi, D. & Tél, T. 2022, *Physical Review E*, 105, L062202
- Karney, C. F. 1983, *Physica D: Nonlinear Phenomena*, 8, 360
- Katz, D., Antoja, T., Romero-Gómez, M., et al. 2018, *Astronomy I& Astrophysics*, 616, A11
- Kovács, T. 2020, *J. R. Soc. Interface.*, 17, 20200648
- Kuzmin, G. 1956, *Astron. Zh.*, 33, 27
- Labini, F. S., Žofia Chrobáková, Capuzzo-Dolcetta, R., & López-Corredoira, M. 2023, *The Astrophysical Journal*, 945, 3
- Laskar, J. 1990, *Icarus*, 88, 266
- Lichtenberg, A. & Leiberman, M. 1992, *Regular and Chaotic Dynamics*, 1st edn. (New York, NY, USA: Springer)
- Malhotra, R. 1998, in *Astronomical Society of the Pacific Conference Series*, Vol. 149, *Solar System Formation and Evolution*, ed. D. Lazzaro, R. Vieira Martins, S. Ferraz-Mello, & J. Fernandez, 37
- Manos, T., Bountis, T., & Skokos, C. 2013, *Journal of Physics A Mathematical General*, 46, 254017
- Manos, T. & Machado, R. E. G. 2014, *MNRAS*, 438, 2201
- McMillan, P. J. 2011, *Monthly Notices of the Royal Astronomical Society*, 414, 2446
- Meiss, J. D., Cary, J. R., Grebogi, C., et al. 1983, *Physica D: Nonlinear Phenomena*, 6, 375
- Michtchenko, T., Lepine, J., Barros, D., & S. S. Vieira, R. 2018, *Astronomy I& Astrophysics*, 615
- Miyamoto, M. & Nagai, R. 1975, *Publications of the Astronomical Society of Japan*, 27, 533
- Motter, A. E., Gruiz, M., Károlyi, G., & Tél, T. 2013, *Phys. Rev. Lett.*, 111, 194101
- Mróz, P., Udalski, A., Skowron, D. M., et al. 2019, *ApJ*, 870, L10
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- Newville, M., Stensitzki, T., Allen, D. B., & Ingargiola, A. 2015
- Ott, E. 1993, *Chaos in dynamical systems*, 1st edn. (Cambridge, UK: Cambridge University Press)
- Petzold, L. 1983, *SIAM Journal on Scientific and Statistical Computing*, 4, 136
- Pichard, B., Martos, M., Moreno, E., & Espresate, J. 2003, *ApJ*, 582, 230
- Pouliasis, E., Matteo, P., & Haywood, M. 2016, *Astronomy I& Astrophysics*, 598
- Rieder, S., Dobbs, C., Bending, T., Liow, K. Y., & Wurster, J. 2021, *Monthly Notices of the Royal Astronomical Society*, 509
- Sanderson, R. E., Hartke, J., & Helmi, A. 2017, *The Astrophysical Journal*, 836, 234
- Schmidt, M. 1956, *Bulletin of the Astronomical Institutes of the Netherlands*, 13, 15
- Skokos, C. 2010, in *Lecture Notes in Physics*, Berlin Springer Verlag, ed. J. Souchay & R. Dvorak, Vol. 790, 63–135
- Sofue, Y. 2012, *Publications of the Astronomical Society of Japan*, 64, 75
- Sofue, Y. 2013, 985
- Sommerer, J. C. & Ott, E. 1993, *Science*, 259, 335
- Spergel, D. N., Bolte, M., & Freedman, W. 1997, *Proceedings of the National Academy of Sciences*, 94, 6579
- Teschl, G. 2012, *Ordinary Differential Equations and Dynamical Systems*, Graduate studies in mathematics (American Mathematical Soc.)
- Toomre, A. 1963, *The Astrophysical Journal*, 138, 385
- Trick, W., Bovy, J., D’Onghia, E., & Rix, H.-W. 2017, *The Astrophysical Journal*, 839
- Verheijen, M. A. W. & Sancisi, R. 2001, *Astronomy & Astrophysics*, 370, 765
- Vogt, D. & Letelier, P. S. 2005, *Publications of the Astronomical Society of Japan*, 57, 871
- Walder, M., Erkal, D., Collins, M., & Martinez-Delgado, D. 2024, *Probing the dark matter haloes of external galaxies with stellar streams*
- Wogan, N. 2021, *NumbaLSODA*, <https://github.com/Nicholaswogan/numbalsoda>
- Yu, L., Ott, E., & Chen, Q. 1990, *Phys. Rev. Lett.*, 65, 2935
- Zotos, E. E. 2011, *Chaos Solitons and Fractals*, 44, 501
- Zotos, E. E. 2012, *New A*, 17, 576
- Zotos, E. E. 2014, *Baltic Astronomy*, 23, 151
- Zotos, E. E., Dubeibe, F. L., Steklain, A. F., & Saeed, T. 2020, *A&A*, 643, A33

Appendix A: Illustrating another example of peculiar dynamics

Figure A.1 show the phase space plots in the (R, p_R) plane, where we illustrate the phase portrait for four different values of L_z with the calculated H_0 value. A second




Fig. A.1: Phase portraits of Eq. (10) for different values of L_z with constant $H = H_0$. The trajectories are plotted on the (R, p_R) plane, following the Poincaré cut at $z = 0$, and the value of p_z obtained from (12), indicated at each point by the colormap. The red KAM torus belongs to the initial condition $R_0 = 0.7211$, $p_{R0} = -170.7053$ (panel a).

case of even more peculiar dynamics is shown in Fig. A.2 by following the snapshot torus shown in red in Fig. A.1a, similarly to the previous two cases. In this case an ensemble of orbits is chosen which have significantly lower angular momentum L_z , hence also having significantly low radial velocities compared to the cases presented before. The first stage of its evolution is similar to the torus presented in Fig. 4, where the torus develops stretches, the two differences being the number and the orientation of them, which results in the deformation of the torus. However, the torus does not break up like the one in Fig. 4, and a structure of continuous lines remains up until $n = 694$, from which point the structure goes through a secondary evolution phase. This time the deformation begins from three points, in both the outer and inner direction, which results in the formation of two structures at $p_R \approx \pm 300$ and three continuously stretching elongated structures in the inner parts of the phase space. The rest of the trajectories scatter into the snapshot chaotic sea. The two structures on the left are similar to time-dependent non-hyperbolic regions, trapping trajectories into a smaller parts of the phase space. The behavior of the peculiar structures penetrating inside the large island, present in the snapshots of $n = 761, 874$, are not yet understood and further, more extended research is needed. The EAPD curve for this torus is shown in Fig. A.3.

In Fig. A.4, a monotonous growth in energy dispersion can be seen, similarly to the case in Fig. 6, and after $n = 694$ the mean curve in energy shows a slight break.

$$L_z = 1, H(t=0) = H_0$$




Fig. A.2: Time evolution of the red torus on Fig. A.1a. At $n = 100$ we can see the initial torus starts to deform significantly, while a significantly low fraction of trajectories scatter out into the chaotic sea after $n = 177$. After $n = 694$ the inner structure goes through a second evolutionary phase, with the larger portion of orbits scattering out into the chaotic sea by $n = 761$.




Fig. A.3: EAPD curve of Eq. (14) evaluated for the snapshot torus of Fig. A.2. The inset shows an oscillating behavior of small amplitudes after $n = 100$, similarly to the case in Fig. 7. A breakup phase is also present on the EAPD curve after $n = 761$, for which the Lyapunov exponent obtained from the fitting is $\lambda_{n_r} = 0.04514 \pm 0.00026$.




Fig. A.4: Median energy of the ensemble of snapshot tori shown in Fig. A.2 compared to the EAPD curve of Fig. A.3. The 1σ dispersion is shown as a light red band around the energy curve, while the vertical dashed lines mark $n = 100, 694, 761$, where the first shows the warping of the initial torus, while the latter two show the start of the secondary evolution and breakup phase.