

A Mathematical Theory of Integer Quantum Hall Effect in Photonics*

Jiayu Qiu[†], Hai Zhang[‡]

May 29, 2024

Abstract

This paper investigates interface modes in a square lattice of photonic crystal composed of gyromagnetic particles with C_{4v} point group symmetry. The study shows that Dirac or linear degenerate points cannot occur at the three high symmetry points in the Brillouin zone where two Bloch bands touch. Instead, a touch point at the M-point has a quadratic degeneracy in the generic case. It is further proved that when a magnetic field is applied to the two sides of an interface in opposite directions, two interface modes that are supported along that interface can be bifurcated from the quadratic degenerate point. The results provide a mathematical foundation for the first experiment realization of the integer quantum Hall effect in the context of photonics.

Key words: Integer quantum Hall effect, square lattice of photonic crystals, interface modes, quadratic degenerate point, topological photonics

1 Introduction

1.1 Background

Topological photonics and phononics are rapidly growing fields that apply the principles of topological phases of matter, originally discovered in solid-state physics for electronic systems, to the context of optics and phononics. The study of topological phases of matter in condensed-matter systems began with the discovery of the integer quantum Hall effect (IQHE) in 1980 [35, 54]. Thouless et al. discovered in 1982 that the integer in the quantized Hall conductance is related to a topological invariant of the system, the Chern number, which is determined by the Bloch eigenfunctions over the Brillouin zone [52]. The analog of IQHE in photonics was first proposed by Haldane and Raghu in their seminal work in 2008 [24], while the first experimental realization was made one year later by Wang et al. [55]. Since then, there have been great activities in the study of a variety of photonic and phononic systems realizing

*This work was partially supported by Hong Kong RGC grant GRF 16304621 and NSFC grant 12371425.

[†]Department of Mathematics, HKUST, Clear Water Bay, Kowloon, Hong Kong S.A.R., China. jqiuj@connect.ust.hk.

[‡]Department of Mathematics, HKUST, Clear Water Bay, Kowloon, Hong Kong S.A.R., China. haizhang@ust.hk.

band structures with nontrivial topological invariants, leading to the emerging research field of topological photonics and phononics [42, 34].

An important feature of topological insulators or topological phases is the existence of gapless edge/interface modes that are spatially localized at the boundary of the bulk insulator or the interface separating it with another bulk insulator that has different topological phases. This is known as the bulk-edge/interface correspondence [28, 27, 26, 45]. A rigorous justification of the bulk-edge/interface correspondence is one of the most interesting and challenging mathematical problems in the study of topological materials. In 1993, Hatsugai established the first proof for the Harper model via transfer matrix and Riemann sheets [27, 26]. Since then, much progress has been made for discrete models in electronic systems by the K-theory approach [30, 15, 7, 9, 36] and functional analysis approach [16, 23, 22, 5], etc. Unlike discrete models, there has been relatively little progress made in continuous models where partial differential equations are involved. Most research in continuous models focuses on electronic systems [32, 50, 13, 12, 31, 8, 14], and few are concerned with the photonic/phononic systems [51]. With rapid experiment developments, an applicable mathematical theory for photonic systems that illustrates the relation between the existence of interface modes and topological phases of bulk materials is highly desirable.

As a special case of the bulk-interface correspondence, the existence of interface modes can be proved using the idea of bifurcation of Dirac points. A Dirac point is a special degenerate point in the spectral bands of a periodic operator where the dispersion curves or surfaces of two bands intersect in a linear or conic manner [2, 20, 41, 6, 1, 4]. Dirac points occur in photonic graphene, electronic systems, and photonic/phononic systems with a honeycomb structure. Generally, a topological phase transition occurs near a Dirac point, and an in-gap eigenvalue can be generated by applying proper perturbations to the periodic operator. The bifurcation of eigenvalues from Dirac points was rigorously analyzed for one-dimensional Schrodinger operators [18], two-dimensional Schrodinger operators [19], and two-dimensional elliptic operators with smooth coefficients [39], all using domain wall models. It is worth mentioning that the original proposal of Raghu and Haldane on IQHE in photonics can be interpreted as a result of the bifurcation of Dirac points. The need for a Dirac point led them to focus on TE modes in a triangular lattice of gyromagnetic particles. However, such a proposal was not adopted in the experiment [55], since the gyromagnetic effect that can break the time-reversal symmetry in realistic materials is too weak, resulting in a too-small band gap that is not robust against disorder. Instead, a 2D square lattice of gyromagnetic particles whose band diagram contains a quadratic degenerate M-point was used. We note that a quadratic degenerate point cannot be treated as in the linear case of Dirac points; see [10] for discussion on those two types of degenerate points. This suggests the need for a new framework to study the interface modes that bifurcate from the quadratic degenerate points, as is established in this paper.

This paper aims to develop a mathematical theory to explain the first experiment realization of IQHE in the context of photonics [55]. To be precise, we prove rigorously the existence of interface modes when an appropriate perturbation is applied to a 2D square lattice of gyromagnetic particles. The major contribution of this paper is the development of a mathematical framework to reveal the mechanism of creating interface modes from a quadratic degenerate point. Our analysis shows that a topological phase transition occurs at the degenerate point. This phase transition is characterized by the change of parity of the Bloch eigenfunctions if one switches the direction of the magnetic field applied to the system. Consequently, we prove that

such a topological phase transition leads to the emergence of interface modes that bifurcate from the degenerate point. We note that in the original paper [55], the argument to support the existence of the interface mode is based on Hatsugai's relation between edge states and Chern numbers and an interesting relationship that allows to map the photon states in gyromagnetic crystals to electron wave functions in a family of quantum Hall systems. However, it is worth noting that Hatsugai's relation was derived using a lattice tight-binding model. It remains unclear whether the continuous photonics system can be reduced to such a model due to the long-range interactions in electromagnetic waves. We also point out that the Chern number argument provides little information for structures with the time-reversal symmetry it is trivial in that case. In contrast, the method developed in this paper can be applied to systems with time-reversal symmetry and be used to study the existence of interface modes, as is done in [46, 40].

1.2 Model description and main results

We start with the following periodic elliptic operator in $L^2(\mathbf{R}^2)$

$$\mathcal{L}^A : D(\mathcal{L}^A) \subset L^2(\mathbf{R}^2) \rightarrow L^2(\mathbf{R}^2), \quad u \mapsto -\operatorname{div}(A\nabla)u, \quad (1.1)$$

where

$$D(\mathcal{L}^A) = \{u \in H^1(\mathbf{R}^2) : -\operatorname{div}(A\nabla)u \in L^2(\mathbf{R}^2)\},$$

and the coefficient matrix $A = A(\mathbf{x})$ satisfies the following assumption.

Assumption 1.1. $A(\mathbf{x}) = (1 + a(\mathbf{x})) \cdot I_{2 \times 2}$, where $a(\mathbf{x}) = c \cdot \sum_{n_1, n_2 \in \mathbf{Z}} \chi_{D_{n_1, n_2}}(\mathbf{x})$. Here $c > 0$ is a positive constant, χ is the indicator function and $D_{n_1, n_2} := D + n_1 \mathbf{e}_1 + n_2 \mathbf{e}_2$ are inclusions. We assume that $D \subset (0, 1) \times (0, 1)$ is simply connected and the structure is symmetric under the C_{4v} point group of the square lattice in the sense that

$$\mathbf{x} \in \cup_{n_1, n_2 \in \mathbf{Z}} D_{n_1, n_2} \implies R\mathbf{x}, M_2\mathbf{x} \in \cup_{n_1, n_2 \in \mathbf{Z}} D_{n_1, n_2},$$

where $R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ($\pi/2$ -rotation about the origin) and $M_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ (reflection about the x_1 -axis).

The operator \mathcal{L}^A models the propagation of time-harmonic TE polarized electromagnetic waves in a 2D square lattice of gyromagnetic particles without applied magnetic field (see Figure 1 where the inclusions are disks, as in the original setting of [55]). It can be equivalently defined through the following sesquilinear form on $L^2(\mathbf{R}^2)$

$$\mathfrak{a}^A(u, v) = \int_{\mathbf{R}^2} (A(\mathbf{x})\nabla u(\mathbf{x})) \cdot \overline{\nabla v(\mathbf{x})} d\mathbf{x}, \quad u, v \in H^1(\mathbf{R}^2) \subset L^2(\mathbf{R}^2). \quad (1.2)$$

Assumption 1.1 implies that \mathcal{L}^A is invariant under translation, rotation and reflection in the sense that

$$\mathcal{L}^A \mathcal{T}_i u = \mathcal{T}_i \mathcal{L}^A u, \quad \mathcal{L}^A \mathcal{R} u = \mathcal{R} \mathcal{L}^A u, \quad \mathcal{L}^A \mathcal{M}_2 u = \mathcal{M}_2 \mathcal{L}^A u, \quad u \in D(\mathcal{L}^A), \quad (1.3)$$

where

$$(\mathcal{T}_i u)(\mathbf{x}) := u(\mathbf{x} + \mathbf{e}_i) \quad (i = 1, 2), \quad (\mathcal{R} u)(\mathbf{x}) := u(R\mathbf{x}), \quad (\mathcal{M}_2 u)(\mathbf{x}) := u(M_2\mathbf{x}).$$




Figure 1: Left: structure of the problem in the case the inclusions are taken as disks. Right: illustration of the unit cell.

The spectral theory of periodic operators tells that

$$\sigma(\mathcal{L}^A) = \bigcup_{\boldsymbol{\kappa} \in Y^*} \sigma(\mathcal{L}^A(\boldsymbol{\kappa})),$$

where $Y_* := [0, 2\pi]^2$ denotes the Brillouin zone of the square lattice, $\mathcal{L}^A(\boldsymbol{\kappa})$ is the restriction of \mathcal{L}^A on the space

$$L_{\boldsymbol{\kappa}}^2(\mathbf{R}^2) := \{u \in L_{loc}^2(\mathbf{R}^2) : u(\mathbf{x} + n_1 \mathbf{e}_1 + n_2 \mathbf{e}_2) = e^{i(n_1 \boldsymbol{\kappa}_1 + n_2 \boldsymbol{\kappa}_2)} u(\mathbf{x})\}.$$

The following identities hold for all $\boldsymbol{\kappa}$, which describe the symmetry in momentum space

$$\mathcal{L}^A(\boldsymbol{\kappa} + n_1 \cdot 2\pi \mathbf{e}_1 + n_2 \cdot 2\pi \mathbf{e}_2) = \mathcal{L}^A(\boldsymbol{\kappa}), \quad n_1, n_2 \in \mathbf{Z}, \quad (1.4)$$

$$\mathcal{R} \mathcal{L}^A(\boldsymbol{\kappa}) \mathcal{R}^{-1} = \mathcal{L}^A(R^{-1} \boldsymbol{\kappa}), \quad (1.5)$$

$$\mathcal{M}_2 \mathcal{L}^A(\boldsymbol{\kappa}) \mathcal{M}_2^{-1} = \mathcal{L}^A(M_2 \boldsymbol{\kappa}). \quad (1.6)$$

Let $\{\lambda_n(\boldsymbol{\kappa}), u_n(\mathbf{x}; \boldsymbol{\kappa})\}_{n \geq 1}$ be the Floquet-Bloch eigenpair of $\mathcal{L}^A(\boldsymbol{\kappa})$. It's known that $\lambda_n(\boldsymbol{\kappa})$ and $u_n(\mathbf{x}; \boldsymbol{\kappa})\}_{n \geq 1}$ are piecewisely smooth on $\boldsymbol{\kappa} \in Y^*$ [37]. The possible singular points appear as the degenerate points in the spectrum $\sigma(\mathcal{L}^A(\boldsymbol{\kappa}))$. The most common degenerate points are the linear ones which are referred to as Dirac points defined below.

Definition 1.2 (Linear degenerate point or Dirac point in square lattice). *A point $(\boldsymbol{\kappa}_*, \lambda_*) \in Y^* \times \mathbf{R}$ is defined as a Dirac point if there exist $n_* \in \mathbf{Z}$ such that*

- (1) $\lambda_{n_*}(\boldsymbol{\kappa}_*) = \lambda_{n_*+1}(\boldsymbol{\kappa}_*) = \lambda_*$;
- (2) $\lambda_{n_*}(\boldsymbol{\kappa}) \neq \lambda_*$, $\lambda_{n_*+1}(\boldsymbol{\kappa}) \neq \lambda_*$ for any $\boldsymbol{\kappa} \in Y^* \setminus \{\boldsymbol{\kappa}_*\}$, and $\lambda_n(\boldsymbol{\kappa}) \neq \lambda_*$ for any $n \notin \{n_*, n_* + 1\}$ and $\boldsymbol{\kappa} \in Y^*$ (spectral no-fold condition);
- (3) For $|\boldsymbol{\kappa} - \boldsymbol{\kappa}_*| \ll 1$, there exists $\alpha_* > 0$ such that

$$\begin{aligned} \lambda_{n_*}(\boldsymbol{\kappa}) - \lambda_{n_*}(\boldsymbol{\kappa}_*) &= -\alpha_* |\boldsymbol{\kappa} - \boldsymbol{\kappa}_*| + \mathcal{O}(|\boldsymbol{\kappa} - \boldsymbol{\kappa}_*|^2), \\ \lambda_{n_*+1}(\boldsymbol{\kappa}) - \lambda_{n_*}(\boldsymbol{\kappa}_*) &= \alpha_* |\boldsymbol{\kappa} - \boldsymbol{\kappa}_*| + \mathcal{O}(|\boldsymbol{\kappa} - \boldsymbol{\kappa}_*|^2). \end{aligned} \quad (1.7)$$

The band structure with a Dirac point is shown in Figure 2(a) (plotted along the slice $\boldsymbol{\kappa}(\kappa_1) = \boldsymbol{\kappa}_* - \mathbf{e}_1 + \kappa_1 \mathbf{e}_1$). Dirac points usually appear in *triangular lattices* structures [25, 6, 19]. In contrast, we prove the following theorem on the nonexistence of linear degenerate points in the *square lattices* on the three high symmetry points in Y_* , i.e. $\boldsymbol{\kappa}^{(1)} := (0, 0)$, $\boldsymbol{\kappa}^{(2)} := (\pi, \pi)$ and $\boldsymbol{\kappa}^{(3)} := (0, \pi)$, see Section 2. We note that $\boldsymbol{\kappa}^{(2)}$ is also called the M-point. Those high symmetry points arise naturally as they are the only fixed points of the symmetry operation in momentum space (see (1.4)-(1.6)).




Figure 2: (a) Band structure supposed in Definition 1.2. A Dirac point exists between the first and second bands; (b) Band structure supposed in Assumption 1.4. The first and second bands touch at $\boldsymbol{\kappa}^{(2)}$ quadratically; (c) Band structure of the parity-broken crystal (described by (1.10)). The perturbation lifts the quadratic degeneracy and opens a band gap.

Theorem 1.3. *The linear degenerate points can't appear at $\boldsymbol{\kappa}^{(i)}$ ($i = 1, 2, 3$).*

As shown in the proof of Theorem 1.3, when two bands touch at $\boldsymbol{\kappa}^{(i)}$ ($i = 1, 2, 3$), the symmetry of the degenerate Bloch modes can be classified by the representation of C_{4v} point group. In physics literature, the degenerate point is called *accidental* when the two degenerate Bloch modes belong to two copies of one-dimensional representations of C_{4v} (a perturbation can lift the degeneracy without breaking the C_{4v} -symmetry). On the opposite, the degenerate point is called *generic* when the two degenerate Bloch modes belong to the two-dimensional representation of C_{4v} (in that case, a symmetry-preserving perturbation cannot lift the degeneracy). Following Theorem 1.3, we focus on the case where the first and second bands touch at $\boldsymbol{\kappa} = \boldsymbol{\kappa}^{(2)}$ quadratically and in a generic way.

Assumption 1.4 (Generic quadratic degenerate point at $\boldsymbol{\kappa}^{(2)}$). *We assume that*

- (1) $\lambda_1(\boldsymbol{\kappa}^{(2)}) = \lambda_2(\boldsymbol{\kappa}^{(2)}) = \lambda_*$;
- (2) $\lambda_1(\boldsymbol{\kappa}) \neq \lambda_*$, $\lambda_2(\boldsymbol{\kappa}) \neq \lambda_*$ for any $\boldsymbol{\kappa} \in Y^* \setminus \{\boldsymbol{\kappa}^{(2)}\}$, and $\lambda_n(\boldsymbol{\kappa}) \neq \lambda_*$ for any $n \geq 3$ and $\boldsymbol{\kappa} \in Y^*$;
- (3) For $|\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}| \ll 1$, there exist a positive definite matrix H_* such that

$$\begin{aligned} \lambda_1(\boldsymbol{\kappa}) - \lambda_1(\boldsymbol{\kappa}^{(2)}) &= -\frac{1}{2}(\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)})^T H_* (\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}) + \mathcal{O}(|\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}|^3), \\ \lambda_2(\boldsymbol{\kappa}) - \lambda_2(\boldsymbol{\kappa}^{(2)}) &= \frac{1}{2}(\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)})^T H_* (\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}) + \mathcal{O}(|\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}|^3). \end{aligned} \quad (1.8)$$

- (4) The Bloch eigenfunctions at $\boldsymbol{\kappa} = \boldsymbol{\kappa}^{(2)}$ are partners of the 2D irreducible representation of C_{4v} in the sense that (see also (2.5))

$$\begin{aligned} u_1(R\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}) &= i \cdot u_2(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}), & u_1(M_2\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}) &= u_1(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}), \\ u_2(R\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}) &= i \cdot u_1(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}), & u_2(M_2\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}) &= -u_2(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)}). \end{aligned} \quad (1.9)$$

A picture of the quadratic degenerate point is shown in Figure 2(b). Now we consider the following perturbed operator

$$\mathcal{L}^{A+\delta \cdot B} : D(\mathcal{L}^{A+\delta \cdot B}) \subset L^2(\mathbf{R}^2) \rightarrow L^2(\mathbf{R}^2), \quad u \mapsto -\operatorname{div}((A + \delta \cdot B)\nabla)u, \quad (1.10)$$

where the coefficient matrix B describes the parity-breaking permeability induced by an external magnetic field. We assume

Assumption 1.5. $B(\mathbf{x}) = b(\mathbf{x}) \cdot \sigma_2$, where $b(\mathbf{x}) = \sum_{n_1, n_2 \in \mathbf{Z}} \chi_{D_{n_1, n_2}}(\mathbf{x})$ and $\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

The operator $\mathcal{L}^{A+\delta \cdot B}$ is defined in the weak sense by the sesquilinear form (1.2) with the replacement $A \rightarrow A + \delta \cdot B$. The coefficient matrix B breaks the reflection symmetry while preserving the rotation symmetry, i.e. the following identities (compared with (1.3)) hold

$$\mathcal{L}^B \mathcal{R}u = \mathcal{R} \mathcal{L}^B u, \quad \mathcal{L}^B \mathcal{M}_2 u = -\mathcal{M}_2 \mathcal{L}^B u, \quad u \in D(\mathcal{L}^B).$$

We note that B also breaks the time-reversal symmetry. Such a symmetry-breaking perturbation can lift the quadratic degeneracy at $\kappa^{(2)}$. To be precise, we have

Theorem 1.6 (Band gap opening at $\kappa_2 = \pi$). *Let $0 < c_0 < 1$ be a constant and let $L_\pi^2(\Omega) = \{u \in L_{loc}^2(\mathbf{R}^2) \cap L^2(\Omega) : u(\mathbf{x} + n_2 \mathbf{e}_2) = e^{in_2 \pi} u(\mathbf{x})\}$ with $\Omega := \mathbf{R} \times (-\frac{1}{2}, \frac{1}{2})$. Assume that t_* is defined in (4.2) is nonzero. Then, for $|\delta|$ being sufficiently small and nonzero, the operator $\mathcal{L}^{A+\delta \cdot B} \Big|_{L_\pi^2(\Omega)}$ has a spectral gap $\mathcal{I}_\delta = (\lambda_* - c_0 |t_* \delta|, \lambda_* + c_0 |t_* \delta|)$ near $\lambda = \lambda_*$.*

Remark 1.7. *The assumption $t_* \neq 0$ can be verified numerically.*

Theorem 1.6 follows directly from Theorem 4.1. Note that in Theorem 4.1, it is also proved that the Bloch eigenspace of $\mathcal{L}^{A+\delta \cdot B} \Big|_{L_\pi^2(\Omega)}$ near $\lambda = \lambda_*$ exchanges its parity for $\delta > 0$ and $\delta < 0$ (see Remark 4.2). This is the so-called *band-inversion phenomenon* in the physics literature, which can be seen as a topological phase transition [53]. More importantly, when such a phase transition occurs, one expects the existence of localized modes at the interface between the two lattices associated with the two operators $\mathcal{L}^{A+\delta \cdot B}$ and $\mathcal{L}^{A-\delta \cdot B}$. This is indeed the case as is proved in this paper. To be precise, we consider the following operator

$$\mathcal{L}^{inter} : D(\mathcal{L}^{inter}) \subset L^2(\mathbf{R}^2) \rightarrow L^2(\mathbf{R}^2), \quad u(\mathbf{x}) \mapsto \begin{cases} (\mathcal{L}^{A-\delta \cdot B} u)(\mathbf{x}), & x_1 < 0, \\ (\mathcal{L}^{A+\delta \cdot B} u)(\mathbf{x}), & x_1 > 0. \end{cases} \quad (1.11)$$

The main result of this paper establishes that

Theorem 1.8 (Interface modes at $\kappa_2 = \pi$). *Assume that t_* is defined in (4.2) is nonzero. For $|\delta|$ being sufficiently small and nonzero, there exist exactly two eigenvalues $\lambda_n^*(\pi)$ ($n = 1, 2$) of $\mathcal{L}^{inter} \Big|_{L_\pi^2(\Omega)}$ inside the band gap \mathcal{I}_δ , with the corresponding eigenfunction $u_n^*(\mathbf{x}; \pi) \in L_\pi^2(\Omega)$.*

Remark 1.9. *With Theorem 1.8, one can immediately prove there exist exactly two eigenvalues $\lambda_n^*(\kappa_2)$ ($n = 1, 2$) of $\mathcal{L}^{inter} \Big|_{L_{\kappa_2}^2(\Omega)}$ inside the band gap for $|\kappa_2 - \pi| \ll 1$ by a standard perturbation argument. One can also apply the tools developed in this paper to calculate the slope $(\lambda_n^*)'(\pi)$ of the two dispersion curves for the interface modes, as in [40].*

Remark 1.10. *The IQHE in photonics is featured by robust unidirectional interface or edge modes that can propagate along the interface or boundary without backscattering. In this paper, we only established the existence of the interface modes. The unidirectionality of the modes can be established by studying the slope of the two dispersion curves of the two family*

of interface modes, see Remark 1.9. On the other hand, the robustness of the interface modes to impurities of the structure is a very subtle issue that needs further investigation, although certain stability follows from the stability of eigenvalues for Hermitian systems. One way to establish the stability is to prove the bulk-interface correspondence which links the Chern numbers of the two bulk structures on the two sides of the interface with the number of interface modes. This is a fascinating yet open problem that requires future work.

Remark 1.11. *We analyzed the approach of using gyromagnetic materials to generate the IQHE in photonics. Some proposals do not rely on gyromagnetic materials. This includes: 1). Suitably designed temporal modulation of the coupling in optical resonator lattices that creates effective magnetic fields for light [17]; 2). Waveguiding geometries with longitudinal refractive index modulations [47]. The analysis of these proposals is beyond the scope of this paper.*

1.3 Outline

This paper is structured as follows:

In Section 2, we prove the nonexistence of the Dirac points on the three high symmetry points in Y_* , i.e. $\boldsymbol{\kappa}^{(1)} := (0, 0)$, $\boldsymbol{\kappa}^{(2)} := (\pi, \pi)$ and $\boldsymbol{\kappa}^{(3)} := (0, \pi)$. This result naturally leads to the consideration of quadratic degenerate points, as supposed in Assumption 1.4;

In Section 3, we first briefly review the Floquet theory of periodic operators, which is basic for our analysis. Second, we study the limiting absorption principle at the energy level of the quadratic degenerate point. In particular, we obtain the asymptotic expansion of the Green function in a strip in the presence of a quadratic degenerate point (see Theorem 3.3), which plays a critical role in our proof of the main result. This result also generalizes the previous studies of the limiting absorption principle in the whole space [44, 38, 33] or in a strip without the quadratic degenerate point [21]. Finally, we calculate the energy flux in a strip between different modes of the same eigenvalue λ_* , which leads to an orthogonal decomposition of the interface function spaces (see (5.9)).

In Section 4, we obtain the asymptotic expansion of the Bloch eigenvalues and eigenfunctions near the degenerate point when we apply the perturbation; see Theorem 4.1. Those results indicate two key points: (1) a common band gap is opened in the band structure of the two systems at two sides of the interface (see Theorem 1.6); (2) there is a parity-changing phenomenon in the perturbation (see Remark 4.2). These two points imply the existence of interface modes;

In Section 5, we prove the existence of the interface modes, which is the main result of this paper (Theorem 1.8). Our method is based on the layer-potential and integral equation theory. Roughly speaking, we transform the eigenvalue problem of \mathcal{L}^{inter} to a characteristic value equation imposed on the interface by using the layer-potential expression; see (5.10) and (5.11). Then, by choosing a suitable scaling, we study the limit of the characteristic value equation; see Proposition 5.1. Those results pave the way for applying the Gohberg-Sigal theory to solve the characteristic value equation, as we did in Section 5.3. In particular, the results in Section 5.3 clearly show that the interface modes are bifurcated from the degenerate point in the band structure.

1.4 Notations

1.4.1 Operators and relations

Throughout, $R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $M_1 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$, $M_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ denote the $\pi/2$ -rotation, x_2 -axis reflection and x_1 -axis reflection matrix in the Euclidean space \mathbf{R}^2 , respectively. Their counterparts in the function space are given by

$$\mathcal{R} : u(\mathbf{x}) \mapsto u(R\mathbf{x}), \quad \mathcal{M}_1 : u(\mathbf{x}) \mapsto u(M_1\mathbf{x}), \quad \mathcal{M}_2 : u(\mathbf{x}) \mapsto u(M_2\mathbf{x}).$$

We denote $u \sim v$ if the two functions u and v are unitarily equivalent, i.e. there exists a unit $c \in \mathbf{C}$ such that $u = c \cdot v$.

1.4.2 Geometry

$$\begin{aligned} Y &= [-\frac{1}{2}, \frac{1}{2}]^2, \quad Y^* = [0, 2\pi]^2; \\ \Omega &= \mathbf{R} \times (-\frac{1}{2}, \frac{1}{2}); \\ \Omega^{\text{right}} &:= \Omega \cap (\mathbf{R}^+ \times \mathbf{R}), \quad \Omega^{\text{left}} := \Omega \cap (\mathbf{R}^- \times \mathbf{R}); \\ \Gamma &= \{0\} \times (-\frac{1}{2}, \frac{1}{2}) \subset \Omega; \\ \Gamma^{\text{right}} &:= \partial(\Omega^{\text{right}}), \quad \Gamma^{\text{left}} := \partial(\Omega^{\text{left}}); \\ \Gamma^+ &= \mathbf{R} \times \{\frac{1}{2}\}, \quad \Gamma^- = \mathbf{R} \times \{-\frac{1}{2}\}. \end{aligned}$$

1.4.3 Function spaces and brackets

$L^2_{\kappa}(\mathbf{R}^2) := \{u \in L^2_{loc}(\mathbf{R}^2) : u(\mathbf{x} + n_1\mathbf{e}_1 + n_2\mathbf{e}_2) = e^{i(n_1\kappa_1 + n_2\kappa_2)}u(\mathbf{x})\}$, equipped with $L^2(Y)$ -inner product.

$H^1_{\kappa}(\mathbf{R}^2) := \{u \in H^1_{loc}(\mathbf{R}^2) : \partial_{\alpha}u \in L^2_{\kappa}(\mathbf{R}^2) (|\alpha| \leq 1)\}$, equipped with $H^1(Y)$ -inner product;

$L^2_{\kappa_2}(\Omega) := \{u \in L^2_{loc}(\mathbf{R}^2) : u|_{\Omega} \in L^2(\Omega), u(\mathbf{x} + n_2\mathbf{e}_2) = e^{in_2\kappa_2}u(\mathbf{x})\}$, equipped with $L^2(\Omega)$ -inner product.

$H^1_{\kappa_2}(\Omega) := \{u \in H^1_{loc}(\mathbf{R}^2) : u|_{\Omega} \in H^1(\Omega), u(\mathbf{x} + n_2\mathbf{e}_2) = e^{in_2\kappa_2}u(\mathbf{x}), \partial_{x_2}u(\mathbf{x} + n_2\mathbf{e}_2) = e^{in_2\kappa_2}\partial_{x_2}u(\mathbf{x})\}$.

$(H^1(U))^*$ ($U \subset \mathbf{R}^2$ is an open set): dual of $H^1(U)$ under the dual pairing induced by the $L^2(U)$ -inner product.

$H^{\frac{1}{2}}(\Gamma) := \{u = U|_{\Gamma} : U \in H^{\frac{1}{2}}(\Gamma^{\text{right}})\}$, where $H^{\frac{1}{2}}(\Gamma^{\text{right}})$ is defined in the standard way.

$\tilde{H}^{-\frac{1}{2}}(\Gamma) := \{u = U|_{\Gamma} : U \in H^{-\frac{1}{2}}(\Gamma^{\text{right}}) \text{ and } \text{supp}(U) \subset \bar{\Gamma}\}$, where $H^{-\frac{1}{2}}(\Gamma^{\text{right}})$ is the dual of $H^{\frac{1}{2}}(\Gamma^{\text{right}})$ under the dual pair $\langle \varphi, \phi \rangle := \int_{\Gamma^{\text{right}}} \varphi(\cdot)\phi(\cdot)$.

(\cdot, \cdot) : $L^2(Y)$ -inner product and dual pairing induced by the $L^2(Y)$ -inner product.

$(\cdot, \cdot)_{\Omega}$: $L^2(\Omega)$ -inner product and dual pairing induced by the $L^2(\Omega)$ -inner product.

$\langle \cdot, \cdot \rangle$: the $\tilde{H}^{-\frac{1}{2}}(\Gamma) - H^{\frac{1}{2}}(\Gamma)$ dual pairing.

2 Absence of linear degeneracies

In this section, we prove Theorem 1.3 for $\kappa = \kappa^{(2)}$ by following the argument in [6]. We point out that the proof for $\kappa = \kappa^{(1)}$ and $\kappa^{(3)}$ is similar.

2.1 Representation theory of C_{4v} point group and symmetry in momentum space

We briefly summarize the representation theory of C_{4v} point group that is used in this paper. Let \mathcal{L} be a self-adjoint operator acting on a separable Hilbert space \mathcal{X} . Let \mathcal{S} be a finite group of unitary operators that acts on \mathcal{X} and which commutes with \mathcal{L} . It's well-known that there is an isotypic decomposition of \mathcal{X} into a finite orthogonal sum of subspaces, each carrying copies of an irreducible representation ρ of \mathcal{S} [6]. To be specific,

$$\mathcal{X} = \bigoplus_{\rho} \mathcal{X}_{\rho},$$

where for any two vectors $u_1, u_2 \in \mathcal{X}_{\rho}$, there is an isomorphism between the spaces

$$\mathcal{S}u_1 = \text{span}\{Su_1 : S \in \mathcal{S}\} \text{ and } \mathcal{S}u_2 = \text{span}\{Su_2 : S \in \mathcal{S}\}$$

and the isomorphism commutes with all $S \in \mathcal{S}$. The dimension of $\mathcal{S}u$ is equal to the dimension of the representation ρ .

Since \mathcal{L} commutes with \mathcal{S} , each isotypic component \mathcal{X}_{ρ} is invariant with respect to \mathcal{L} . Moreover, if \mathcal{L} has a discrete spectrum, then the restriction of \mathcal{L} to \mathcal{X}_{ρ} has eigenvalues with multiplicities divisible by the dimension of ρ . Indeed, if u is an eigenvector of \mathcal{L} , then the subspace $\mathcal{S}u$ belongs to the eigenspace of \mathcal{L} with the same eigenvalue.

We now restrict to the case $\mathcal{S} = C_{4v}$. It's known that \mathcal{S} is generated by \mathcal{R} ($\pi/2$ -rotation) and \mathcal{M}_2 (x_1 -reflection), with the following rules

$$\mathcal{R}^4 = \mathcal{M}_2^2 = I, \quad \mathcal{M}_2\mathcal{R}^{-1} = \mathcal{R}\mathcal{M}_2.$$

Note that \mathcal{S} is invariant on the space $L_{\kappa^{(i)}}^2$ ($i = 1, 2$) (since the quasi-periodicity of u is preserved under \mathcal{S} if $u \in L_{\kappa^{(i)}}^2$). The group representation theory dictates that \mathcal{S} has four one-dimensional representations and a unique two-dimensional representation on $L_{\kappa^{(i)}}^2$, as listed below.

$$\rho_1 : \quad \mathcal{R} \mapsto (1), \quad \mathcal{M}_2 \mapsto (1); \tag{2.1}$$

$$\rho_2 : \quad \mathcal{R} \mapsto (1), \quad \mathcal{M}_2 \mapsto (-1); \tag{2.2}$$

$$\rho_3 : \quad \mathcal{R} \mapsto (-1), \quad \mathcal{M}_2 \mapsto (1); \tag{2.3}$$

$$\rho_4 : \quad \mathcal{R} \mapsto (-1), \quad \mathcal{M}_2 \mapsto (-1); \tag{2.4}$$

$$\rho_5 : \quad \mathcal{R} \mapsto i\sigma_3 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \mathcal{M}_2 \mapsto \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{2.5}$$

On the other hand, note that $\mathcal{L}^A(\kappa^{(2)})$ commutes with \mathcal{S} . Therefore, the following holds

Proposition 2.1. *Suppose $\{\lambda, u\} \in \mathbf{R} \times L_{\kappa^{(2)}}^2$ is an eigenpair of $\mathcal{L}^A(\kappa^{(2)})$. Then there exists a unique ρ_k ($1 \leq k \leq 5$) such that $u \in L_{\kappa^{(2)}, \rho_k}^2$. Here $L_{\kappa^{(2)}, \rho_k}^2$ denotes the isotypic component of $L_{\kappa^{(2)}}^2$ associated with the representation ρ_k .*

2.2 Proof of Theorem 1.3

We prove Theorem 1.3 in this subsection by contradiction. Without loss of generality, we assume that the first and second spectral bands of the operator \mathcal{L}^A touch at $(\boldsymbol{\kappa}^{(2)}, \lambda_*)$ and that $(\boldsymbol{\kappa}^{(2)}, \lambda_*)$ is a linear degenerate point, i.e. the conditions in Definition 1.2 hold for $n_* = 1$ and $(\boldsymbol{\kappa}_*, \lambda_*) = (\boldsymbol{\kappa}^{(2)}, \lambda_*)$. We denote $u_1 = u_1(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)})$ and $u_2 = u_2(\boldsymbol{x}; \boldsymbol{\kappa}^{(2)})$ for ease of notations. By Proposition 2.1, there are two cases:

- Case 1. $u_1, u_2 \in L^2_{\boldsymbol{\kappa}^{(2)}, \rho_5}$;
- Case 2. $u_1, u_2 \in \bigoplus_{k=1}^4 L^2_{\boldsymbol{\kappa}^{(2)}, \rho_k}$.

We shall derive contradictions in both cases to prove Theorem 1.3.

Proof for Case 1: Without loss of generality, we set $\mathcal{R}u_1 = iu_1$, $\mathcal{R}u_2 = -iu_2$. In the first-order perturbation argument (i.e. ignoring the $\mathcal{O}(|\boldsymbol{\kappa} - \boldsymbol{\kappa}^{(2)}|^2)$ term in (1.7)), the dispersion surface near $(\boldsymbol{\kappa}^{(2)}, \lambda_*)$ can be obtained by solving the following equation (See [46, 18, 19])

$$\det(\delta\kappa_1 \cdot h_1 + \delta\kappa_2 \cdot h_2 - \delta\lambda) = 0,$$

where $\delta\kappa_i := (\boldsymbol{\kappa}_i - \boldsymbol{\kappa}_i^{(2)}) \cdot \boldsymbol{e}_i$, $\delta\lambda := \lambda - \lambda_*$, and

$$h_i = \begin{pmatrix} (u_1, \frac{\partial \mathcal{L}^A}{\partial \kappa_i}(\boldsymbol{\kappa}^{(2)})u_1) & (u_1, \frac{\partial \mathcal{L}^A}{\partial \kappa_i}(\boldsymbol{\kappa}^{(2)})u_2) \\ (u_2, \frac{\partial \mathcal{L}^A}{\partial \kappa_i}(\boldsymbol{\kappa}^{(2)})u_1) & (u_2, \frac{\partial \mathcal{L}^A}{\partial \kappa_i}(\boldsymbol{\kappa}^{(2)})u_2) \end{pmatrix}, \quad i = 1, 2.$$

Here (\cdot, \cdot) denotes the $L^2(Y)$ -inner product. We claim that $h_1 = h_2 = 0$. Then the first-order terms in the asymptotic expansion of the dispersion function $\lambda = \lambda(\boldsymbol{\kappa})$ near $\boldsymbol{\kappa} = \boldsymbol{\kappa}^{(2)}$ vanishes, and this contradicts to the definition of Dirac points. We now prove the claim by exploiting the symmetry of u_1 and u_2 . By taking $\boldsymbol{\kappa} = \boldsymbol{\kappa}^{(2)} + \delta\kappa_1 \cdot \boldsymbol{e}_1 + \delta\kappa_2 \cdot \boldsymbol{e}_2$ in (1.5), we have

$$\begin{aligned} \mathcal{R}\mathcal{L}^A(\boldsymbol{\kappa}^{(2)} + \delta\kappa_1 \cdot \boldsymbol{e}_1 + \delta\kappa_2 \cdot \boldsymbol{e}_2)\mathcal{R}^{-1} &= \mathcal{L}^A\left(R^{-1}(\boldsymbol{\kappa}^{(2)} + \delta\kappa_1 \cdot \boldsymbol{e}_1 + \delta\kappa_2 \cdot \boldsymbol{e}_2)\right) \\ &= \mathcal{L}^A(\boldsymbol{\kappa}^{(2)} - 2\pi\boldsymbol{e}_2 + \delta\kappa_2 \cdot \boldsymbol{e}_1 - \delta\kappa_1 \cdot \boldsymbol{e}_2). \end{aligned}$$

The equality (1.4) gives

$$\mathcal{R}\mathcal{L}^A(\boldsymbol{\kappa}^{(2)} + \delta\kappa_1 \cdot \boldsymbol{e}_1 + \delta\kappa_2 \cdot \boldsymbol{e}_2)\mathcal{R}^{-1} = \mathcal{L}^A(\boldsymbol{\kappa}^{(2)} + \delta\kappa_2 \cdot \boldsymbol{e}_1 - \delta\kappa_1 \cdot \boldsymbol{e}_2).$$

By expanding the equation to the first order, we arrive at

$$\mathcal{R}\left(\delta\kappa_1 \cdot \frac{\partial \mathcal{L}^A}{\partial \kappa_1}(\boldsymbol{\kappa}^{(2)}) + \delta\kappa_2 \cdot \frac{\partial \mathcal{L}^A}{\partial \kappa_2}(\boldsymbol{\kappa}^{(2)})\right)\mathcal{R}^{-1} = -\delta\kappa_1 \cdot \frac{\partial \mathcal{L}^A}{\partial \kappa_2}(\boldsymbol{\kappa}^{(2)}) + \delta\kappa_2 \cdot \frac{\partial \mathcal{L}^A}{\partial \kappa_1}(\boldsymbol{\kappa}^{(2)}). \quad (2.6)$$

Note that the matrix of \mathcal{R} under the basis $\{u_1, u_2\}$ is $i\sigma_3$ by (2.5). Thus, by conjugating both sides of (2.6) with u_i, u_j ($i, j \in \{1, 2\}$), we obtain

$$(-i\sigma_3)(\delta\kappa_1 \cdot h_1 + \delta\kappa_2 \cdot h_2)(i\sigma_3) = -\delta\kappa_1 \cdot h_2 + \delta\kappa_2 \cdot h_1.$$

Separating the variables $\delta\kappa_1$ and $\delta\kappa_2$ yields

$$(-i\sigma_3)h_1(i\sigma_3) = -h_2, \quad (-i\sigma_3)h_2(i\sigma_3) = h_1. \quad (2.7)$$

Hence

$$h_1 = (-i\sigma_3)h_2(i\sigma_3) = -(-i\sigma_3)^2 h_1 (i\sigma_3)^2 = -h_1.$$

Thus $h_1 = 0$, and consequently, $h_2 = -(-i\sigma_3)h_1(i\sigma_3) = 0$.

Proof for Case 2: The proof is similar to Case 1. The main idea is to prove that in all subcases, equation (2.7) holds, which then implies that $h_1 = h_2 = 0$, and hence leads to a contradiction. For instance, when $u_1 \in L^2_{\kappa^{(2),\rho_1}}$ and $u_2 \in L^2_{\kappa^{(2),\rho_3}}$, the matrix of \mathcal{R} under the basis $\{u_1, u_2\}$ is σ_3 since $\mathcal{R}u_1 = u_1$, $\mathcal{R}u_2 = -u_2$. Then $\sigma_3 h_1 \sigma_3 = -h_2$, $\sigma_3 h_2 \sigma_3 = h_1$, i.e. equation (2.7) holds. The other cases can be proved similarly.

3 Preliminaries

We present some preliminaries for the proof of Theorem 4.1 and 1.8.

3.1 Floquet theory

Consider the strip domain $\Omega := \mathbf{R} \times (-\frac{1}{2}, \frac{1}{2})$. We define

$$\mathcal{L}_{\Omega,\pi}^A : D(\mathcal{L}_{\Omega,\pi}^A) \subset L^2_\pi(\Omega) \rightarrow L^2_\pi(\Omega), \quad u \mapsto -\operatorname{div}(A\nabla)u,$$

where

$$L^2_\pi(\Omega) := \{u \in L^2_{loc}(\mathbf{R}^2) \cap L^2(\Omega) : u(\mathbf{x} + \mathbf{e}_2) = e^{i\pi}u(\mathbf{x})\}.$$

$\mathcal{L}_{\Omega,\pi}^A$ is the section of $\mathcal{L}^A(\boldsymbol{\kappa})$ along the line $\kappa_2 = \pi$. The spectrum of $\mathcal{L}_{\Omega,\pi}^A$ can be decomposed by its Floquet components, i.e.

$$\sigma(\mathcal{L}_{\Omega,\pi}^A) = \cup_{0 \leq \kappa_1 < 2\pi} \sigma(\mathcal{L}_{\Omega,\pi}^A(\kappa_1)),$$

where $\mathcal{L}_{\Omega,\pi}^A(\kappa_1) = \mathcal{L}^A((\kappa_1, \pi))$. Consequently,

$$\sigma(\mathcal{L}_{\Omega,\pi}^A) = \cup_{n \geq 1} \{\lambda_n(\kappa_1; \pi), 0 \leq \kappa_1 < 2\pi\} = \cup_{n \geq 1} \{\lambda_n((\kappa_1, \pi)), 0 \leq \kappa_1 < 2\pi\},$$

where $\{\lambda_n(\kappa_1; \pi) : n \geq 1\}$ are the Floquet-Bloch eigenvalues of $\mathcal{L}_{\Omega,\pi}^A$. On the other hand, since $\mathcal{L}_{\Omega,\pi}^A(\kappa_1)$ depends analytically on κ_1 , the Kato-Rellich theorem [29] indicates that there exist a family of analytic functions $\{\mu_n(\kappa_1; \pi)\}_{n \geq 1}$ which forms a rearrangement of $\{\lambda_n(\kappa_1; \pi)\}_{n \geq 1}$, i.e.

$$\cup_{n \geq 1} \{\mu_n(\kappa_1; \pi)\} = \cup_{n \geq 1} \{\lambda_n(\kappa_1; \pi)\}, \quad 0 \leq \kappa_1 < 2\pi. \quad (3.1)$$

We call $\{\mu_n(\kappa_1; \pi)\}_{n \geq 1}$ the analytic labeling of Floquet-Bloch eigenvalues of $\mathcal{L}_{\Omega,\pi}^A$. In the sequel, we shall abbreviate $\mu_n(\kappa_1) = \mu_n(\kappa_1; \pi)$ if no confusion arises. The normalized Bloch eigenfunctions associated with $\mu_n(\kappa_1)$ can be chosen analytical in κ_1 , which is denoted by $v_n(\mathbf{x}; \kappa_1)$.

Proposition 3.1 ([21]). *For each $n \geq 1$, there exists a complex neighborhood $\mathcal{D}_n \supset \mathbf{R}$, and two analytic maps*

$$\kappa_1 \in \mathcal{D}_n \mapsto \mu_n(\kappa_1) \in \mathbf{C}, \quad \kappa_1 \in \mathcal{D}_n \mapsto v_n(\mathbf{x}; \kappa_1) \in H^1_{(\kappa_1, \pi)}(\mathbf{R}^2),$$

such that $(\mu_n(\kappa_1), v_n(\mathbf{x}; \kappa_1))$ ($n \geq 1$) represent all Floquet-Bloch eigenpairs of $\mathcal{L}_{\Omega,\pi}^A(\kappa_1)$ for $0 \leq \kappa_1 < 2\pi$.

Proposition 3.2. *Under Assumption 1.4, the first two branches of analytic Floquet-Bloch eigenvalues, i.e. $\mu_n(\kappa_1)$ for $n = 1$ and 2 , can be chosen so that they meet at (π, λ_*) near $\kappa_1 = \pi$, i.e.*

$$\begin{aligned}\mu_1(\kappa_1) - \lambda_* &= -\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^3), \\ \mu_2(\kappa_1) - \lambda_* &= \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^3),\end{aligned}\tag{3.2}$$

where $\gamma_* > 0$ denotes the first diagonal element of the matrix H_* in (1.8). Moreover,

$$\mu_n(\kappa_1) = \mu_n(2\pi - \kappa_1), \quad \text{for } n = 1, 2.\tag{3.3}$$

Proof. The existence of two branches of analytic Floquet-Bloch eigenvalues near $(\kappa_1, \lambda) = (\pi, \lambda_*)$ follows from Proposition 3.1. We denote them by $\mu_n(\kappa_1)$ for $n = 1, 2$. Consequently,

$$\{\mu_1(\kappa_1), \mu_2(\kappa_1)\} = \{\lambda_1(\kappa_1; \pi), \lambda_2(\kappa_1; \pi)\}, \quad \text{for } |\kappa_1 - \pi| \ll 1.$$

To prove (3.2), it's sufficient to exclude the ‘‘cross-branch’’ case where

$$\mu_1(\kappa_1) = \begin{cases} \lambda_1(\kappa_1; \pi), & \kappa_1 < \pi, |\kappa_1 - \pi| \ll 1, \\ \lambda_2(\kappa_1; \pi), & \kappa_1 > \pi, |\kappa_1 - \pi| \ll 1. \end{cases}\tag{3.4}$$

However, (3.4) implies that the second-order derivative of $\mu_1(\kappa_1)$ is discontinuous at $\kappa_1 = \pi$, which contradicts to the analyticity of μ_1 . Hence (3.2) holds.

Finally, we point out that (3.3) follows from the reflection symmetry $[\mathcal{L}^A, \mathcal{M}_2] = 0$. Indeed, for each κ_1 with $|\kappa_1 - \pi| \ll 1$, the reflection symmetry implies that $\mu_1(\kappa_1)$ is also a Floquet-Bloch eigenvalue of $\mathcal{L}_{\Omega, \pi}^A(2\pi - \kappa_1)$, i.e. $\mu_1(\kappa_1) \in \{\mu_n(2\pi - \kappa_1)\}_{n \geq 1}$. Since there are exactly two branches of analytic Floquet-Bloch eigenvalues near (π, λ_*) , i.e. μ_1 and μ_2 , and $\mu_2(\kappa_1) > \lambda_* > \mu_1(\kappa_1)$ for $|\kappa_1 - \pi| \ll 1$ by (3.2), it holds that $\mu_1(\kappa_1) = \mu_1(2\pi - \kappa_1)$. Hence (3.3) is proved for $n = 1$. The proof of $n = 2$ is similar. \square

3.2 Asymptotics of Green function $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$

In this section, we consider the Green function $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$ which satisfies the following equations

$$\begin{cases} (\operatorname{div}(A\nabla) + \lambda_* + i\epsilon)G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon) = \delta(\mathbf{x} - \mathbf{y}), & \mathbf{x}, \mathbf{y} \in \Omega, \\ G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)|_{\Gamma^+} = e^{i\pi}G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)|_{\Gamma^-}, \\ \frac{\partial G}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)|_{\Gamma^+} = e^{i\pi}\frac{\partial G}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)|_{\Gamma^-}. \end{cases}$$

Using the Floquet transform, for each $\epsilon > 0$, we can write

$$G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon) = \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 1} \frac{v_n(\mathbf{x}; \kappa_1) \overline{v_n(\mathbf{y}; \kappa_1)}}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1.$$

Define the integral operator

$$\mathcal{G}(\lambda_* + i\epsilon)f := \int_{\Omega} G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)f(\mathbf{y})d\mathbf{y}.$$

The main result of this section is the following asymptotics of $\mathcal{G}(\lambda_* + i\epsilon)$ as $\epsilon \rightarrow 0^+$.

Theorem 3.3. For each $f \in (H^1(\Omega))^*$ with compact support, we have in $H_{loc}^1(\Omega)$

$$\begin{aligned}
\mathcal{G}(\lambda_* + i\epsilon)f &= (\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}})) \cdot \left(\frac{1-i}{2\gamma_*^{\frac{1}{2}}} v_1(\mathbf{x}; \pi) (f(\cdot), v_1(\cdot; \pi))_{\Omega} - \frac{1+i}{2\gamma_*^{\frac{1}{2}}} v_2(\mathbf{x}; \pi) (f(\cdot), v_2(\cdot; \pi))_{\Omega} \right) \\
&\quad + \mathcal{G}_0(\lambda_*)f \\
&\quad - \frac{2}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(2(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) (f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_{\Omega} - 2(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) (f(\cdot), (\partial_{\kappa_1} v_2)(\cdot; \pi))_{\Omega} \right. \\
&\quad \quad + (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi) (f(\cdot), v_1(\mathbf{y}; \pi))_{\Omega} + v_1(\mathbf{x}; \pi) (f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_{\Omega} \\
&\quad \quad \left. - (\partial_{\kappa_1}^2 v_2)(\mathbf{x}; \pi) (f(\cdot), v_2(\cdot; \pi))_{\Omega} - v_2(\mathbf{x}; \pi) (f(\cdot), (\partial_{\kappa_1}^2 v_2)(\cdot; \pi))_{\Omega} \right) \\
&\quad + \mathcal{O}(\epsilon^{\frac{1}{3}}),
\end{aligned} \tag{3.5}$$

where $\mathcal{G}_0(\lambda_*)$ is the integral operator associated with the kernel function

$$\begin{aligned}
G_0(\mathbf{x}, \mathbf{y}; \lambda_*) &:= \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1) \overline{v_n(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 \\
&\quad + \lim_{\epsilon \rightarrow 0^+} \left(\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup [\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1) \overline{v_1(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 - \epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} v_1(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} \right. \\
&\quad \left. + \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup [\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_2(\mathbf{x}; \kappa_1) \overline{v_2(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 + \epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} v_2(\mathbf{x}; \pi) \overline{v_2(\mathbf{y}; \pi)} \right).
\end{aligned} \tag{3.6}$$

Proof. See Appendix A. \square

Proposition 3.4. The kernel function $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ defined in (3.6) is the fundamental solution to the following equations

$$\begin{cases}
(\operatorname{div}(A\nabla)u + \lambda_*)G_0(\mathbf{x}, \mathbf{y}; \lambda_*) = \delta(\mathbf{x} - \mathbf{y}), & x, y \in \Omega, \\
G_0(\mathbf{x}, \mathbf{y}; \lambda_*)|_{\Gamma^+} = e^{i\pi} G_0(\mathbf{x}, \mathbf{y}; \lambda_*)|_{\Gamma^-}, \\
\frac{\partial G_0}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda_*)|_{\Gamma^+} = e^{i\pi} \frac{\partial G_0}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda_*)|_{\Gamma^-}.
\end{cases} \tag{3.7}$$

Moreover, for $\mathbf{x}, \mathbf{y} \in \Omega$,

$$G_0(\mathbf{x}, \mathbf{y}; \lambda_*) = G_0(\mathcal{M}_1 \mathbf{x}, \mathcal{M}_1 \mathbf{y}; \lambda_*), \tag{3.8}$$

$$G_0(\mathbf{x}, \mathbf{y}; \lambda_*) = \overline{G_0(\mathbf{y}, \mathbf{x}; \lambda_*)}. \tag{3.9}$$

In addition, for fixed $\mathbf{y} \in \Omega$, $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ has the following decomposition

$$\begin{aligned}
G_0(\mathbf{x}, \mathbf{y}; \lambda_*) &= G_0^+(\mathbf{x}, \mathbf{y}; \lambda_*) + \frac{i}{\gamma_*} v_1(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_1)(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} \\
&\quad - \frac{i}{\gamma_*} v_2(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_2)(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \overline{v_2(\mathbf{y}; \pi)},
\end{aligned} \tag{3.10}$$

and

$$\begin{aligned}
G_0(\mathbf{x}, \mathbf{y}; \lambda_*) &= G_0^-(\mathbf{x}, \mathbf{y}; \lambda_*) - \frac{i}{\gamma_*} v_1(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_1)(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} \\
&\quad + \frac{i}{\gamma_*} v_2(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_2)(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \overline{v_2(\mathbf{y}; \pi)},
\end{aligned} \tag{3.11}$$

where $G_0^+(\mathbf{x}, \mathbf{y}; \lambda_*)$ ($G_0^-(\mathbf{x}, \mathbf{y}; \lambda_*)$, resp.) decays exponentially as $x_1 \rightarrow \infty$ ($x_1 \rightarrow -\infty$, resp.).

Proof. See Appendix A. \square

We introduce the following layer potential operators associated with $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$

$$\begin{aligned}
\mathcal{S}(\lambda_*; G_0) &: \tilde{H}^{-\frac{1}{2}}(\Gamma) \rightarrow H_{loc}^1(\Omega), \quad \varphi \mapsto \int_{\Gamma} G_0(\mathbf{x}, \mathbf{y}; \lambda_*) \varphi(\mathbf{y}) ds(\mathbf{y}), \\
\mathcal{D}(\lambda_*; G_0) &: H^{\frac{1}{2}}(\Gamma) \rightarrow H_{loc}^1(\Omega), \quad \phi \mapsto \int_{\Gamma} \frac{\partial G_0}{\partial y_1}(\mathbf{x}, \mathbf{y}; \lambda_*) \phi(\mathbf{y}) ds(\mathbf{y}), \\
\mathcal{N}(\lambda_*; G_0) &: H^{\frac{1}{2}}(\Gamma) \rightarrow \tilde{H}^{-\frac{1}{2}}(\Gamma), \quad \phi \mapsto \int_{\Gamma} \frac{\partial^2 G_0}{\partial x_1 \partial y_1}(\mathbf{x}, \mathbf{y}; \lambda_*) \phi(\mathbf{y}) ds(\mathbf{y}).
\end{aligned} \tag{3.12}$$

In the above notations, we use $\mathcal{S}(\lambda; G)$ to denote the single-layer potential operator associated with the Green function G at energy λ and adopt similar notations for \mathcal{D} and \mathcal{N} . The following jump formulas hold.

Proposition 3.5. *For $\mathbf{x} \in \Gamma$, we have*

$$\lim_{t \rightarrow 0} \left(\mathcal{S}(\lambda_*; G_0)[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x}) \tag{3.13}$$

$$\lim_{t \rightarrow 0} \frac{\partial}{\partial x_1} \left(\mathcal{D}(\lambda_*; G_0)[\phi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mathcal{N}(\lambda_*; G_0)[\phi](\mathbf{x}) \tag{3.14}$$

$$\lim_{t \rightarrow 0^\pm} \frac{\partial}{\partial x_1} \left(\mathcal{S}(\lambda_*; G_0)[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \pm \frac{1}{2} \varphi(\mathbf{x}), \tag{3.15}$$

$$\lim_{t \rightarrow 0^\pm} \left(\mathcal{D}(\lambda_*; G_0)[\phi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mp \frac{1}{2} \phi(\mathbf{x}), \tag{3.16}$$

Proof. See Appendix A. \square

3.3 Bloch modes and energy flux

We introduce the following sesquilinear form on $H_{loc}^1(\mathbf{R}^2)$:

$$\mathbf{q}(u, v; \Gamma_s) = \int_{\Gamma_s} \left(\frac{\partial u}{\partial x_1} \bar{v} - u \frac{\partial \bar{v}}{\partial x_1} \right) dx_2, \quad \Gamma_s := \{s\} \times \left(-\frac{1}{2}, \frac{1}{2}\right) \subset \Omega.$$

When $s = 0$, we write $\mathbf{q}(u, v; \Gamma) = \mathbf{q}(u, v; \Gamma_s)$. Theorem 3 in [21] gives that

Proposition 3.6. *For all $0 \leq \kappa_1 < 2\pi$, and $n \geq 1$,*

$$\mathbf{q}(v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma) = i\mu'(\kappa_1). \tag{3.17}$$

When $\lambda_n(\kappa_1) = \lambda_{n'}(\kappa_1')$, we have

$$\mathbf{q}(v_n(\cdot; \kappa_1), v_{n'}(\cdot; \kappa_1'); \Gamma) = 0, \quad \text{if } n \neq n' \text{ or } \kappa_1 \neq \kappa_1'. \tag{3.18}$$

Proof. We sketch the proof of (3.17) in [21] here since the arguments are used several times in this section.

By differentiating $(-div(A\nabla) - \mu_n(\kappa_1))v_n(\cdot; \kappa_1) = 0$ with respect to κ_1 , we have

$$(-div(A\nabla) - \mu_n(\kappa_1))\partial_{\kappa_1}v_n(\cdot; \kappa_1) = \mu'_n(\kappa_1)v_n(\cdot; \kappa_1). \quad (3.19)$$

Similarly, differentiating $v_n(\mathbf{x} + \mathbf{e}_1; \kappa_1) = e^{i\kappa_1}v_n(\mathbf{x}; \kappa_1)$ and $\partial_{x_1}v_n(\mathbf{x} + \mathbf{e}_1; \kappa_1) = e^{i\kappa_1}\partial_{x_1}v_n(\mathbf{x}; \kappa_1)$ yields

$$\partial_{\kappa_1}v_n(\mathbf{x} + \mathbf{e}_1; \kappa_1) = e^{i\kappa_1}\partial_{\kappa_1}v_n(\mathbf{x}; \kappa_1) + ie^{i\kappa_1}v_n(\mathbf{x}; \kappa_1), \quad (3.20)$$

and

$$\partial_{x_1}(\partial_{\kappa_1}v_n)(\mathbf{x} + \mathbf{e}_1; \kappa_1) = e^{i\kappa_1}\partial_{x_1}(\partial_{\kappa_1}v_n)(\mathbf{x}; \kappa_1) + ie^{i\kappa_1}\partial_{x_1}v_n(\mathbf{x}; \kappa_1). \quad (3.21)$$

By the Green's formula, for any $\lambda \in \mathbf{C}$,

$$\begin{aligned} & \int_Y \left((-div(A\nabla) - \lambda)u \cdot \bar{v} - (-div(A\nabla) - \lambda)\bar{v} \cdot u \right) \\ &= \int_{\Gamma} (1 + a(\mathbf{x})) \left(\frac{\partial u}{\partial x_1} \bar{v} - u \frac{\partial \bar{v}}{\partial x_1} \right) dx_2 - \int_{\Gamma_1} (1 + a(\mathbf{x})) \left(\frac{\partial u}{\partial x_1} \bar{v} - u \frac{\partial \bar{v}}{\partial x_1} \right) dx_2. \end{aligned}$$

Following Assumption 1.1, $a(\mathbf{x})$ vanishes on $\Gamma \cup \Gamma_1$. Hence,

$$\mathbf{q}(u, v; \Gamma) - \mathbf{q}(u, v; \Gamma_1) = \int_Y \left((-div(A\nabla) - \lambda)u \cdot \bar{v} - (-div(A\nabla) - \lambda)\bar{v} \cdot u \right). \quad (3.22)$$

By letting $u = \partial_{\kappa_1}v_n(\cdot; \kappa_1)$, $v = v_n(\cdot; \kappa_1)$, $\lambda = \mu_n(\kappa_1)$ in (3.22) and utilizing (3.19), we have

$$\mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma) - \mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma_1) = \mu'_n(\kappa_1)(v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1)). \quad (3.23)$$

On the other hand, by using (3.20)-(3.21), a direct calculation gives that

$$\mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma) - \mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma_1) = -i\mathbf{q}(v_n(\cdot; \kappa_1), v_n(\cdot; \kappa_1); \Gamma). \quad (3.24)$$

One then concludes the proof of (3.17) by combining (3.23) and (3.24). \square

Proposition 3.7. *If $\mu_n(\kappa_1) \neq \mu_m(\kappa_1)$,*

$$\mathbf{q}(v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma) = 0. \quad (3.25)$$

Proof. By taking $u = \partial_{\kappa_1}v_n(\cdot; \kappa_1)$, $v = v_m(\cdot; \kappa_1)$, $\lambda = \mu_n(\kappa_1)$ in (3.22) and utilizing (3.19), we have

$$\begin{aligned} & \mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma) - \mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma_1) \\ &= \mu'_n(\kappa_1)(v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1)) + (\mu_m(\kappa_1) - \mu_n(\kappa_1))(v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1)). \end{aligned}$$

When $\mu_n(\kappa_1) \neq \mu_m(\kappa_1)$, the Bloch eigenfunctions are orthogonal:

$$(v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1)) = 0.$$

Thus,

$$\mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma) - \mathbf{q}(\partial_{\kappa_1}v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma_1) = 0. \quad (3.26)$$

By combining (3.26) and (3.24), we obtain $\mathbf{q}(v_n(\cdot; \kappa_1), v_m(\cdot; \kappa_1); \Gamma) = 0$. \square

Next, we investigate the parity of $v_n(\mathbf{x}; \kappa_1)$ ($n = 1, 2$). Note that the analytic Floquet-Bloch eigenvalues $\mu_n(\kappa_1)$ are identified with $\lambda_n(\kappa_1; \pi)$ for $n = 1, 2$ and $|\kappa_1 - \pi| \ll 1$, as in Proposition 3.2. Hence Assumption 1.4(4) gives parity of $v_n(\mathbf{x}; \pi)$

$$(\mathcal{M}_1 v_1)(\cdot; \pi) = -v_1(\cdot; \pi), \quad (\mathcal{M}_1 v_2)(\cdot; \pi) = v_2(\cdot; \pi), \quad (3.27)$$

Note that (3.27) is gauge-independent in the sense that it holds whenever $v_n(\mathbf{x}; \kappa_1)$ is multiplied by a smooth phase function $e^{i\eta(\kappa_1)}$. Remarkably, we prove that one can choose a special class of gauge so that the *derivative* of the Floquet-Bloch eigenfunction, i.e. $\partial_{\kappa_1} v_n(\mathbf{x}; \pi)$, has the opposite parity to $v_n(\mathbf{x}; \pi)$.

Proposition 3.8. *When the following holds*

$$\text{Im}(\partial_{\kappa_1} v_n(\cdot; \pi), v_n(\cdot; \pi)) = 0, \quad n = 1, 2, \quad (3.28)$$

we have

$$(\mathcal{M}_1 \partial_{\kappa_1} v_1)(\cdot; \pi) = (\partial_{\kappa_1} v_1)(\cdot; \pi), \quad (\mathcal{M}_1 \partial_{\kappa_1} v_2)(\cdot; \pi) = -(\partial_{\kappa_1} v_2)(\cdot; \pi). \quad (3.29)$$

Proof. We prove (3.29) for $n = 1$. The proof of $n = 2$ is similar. By Proposition 3.2, $\mu_1(\kappa_1) = \mu_1(2\pi - \kappa_1)$. By the reflection symmetry $[\mathcal{L}^A, \mathcal{M}_1] = 0$, for each κ_1 with $|\kappa_1 - \pi| \ll 1$, $(\mathcal{M}_1 v_1)(\cdot; \kappa_1)$ is a Bloch eigenfunction of $\mathcal{L}_{\Omega, \pi}^A(2\pi - \kappa_1)$ associated with the eigenvalue $\mu_1(2\pi - \kappa_1)$. Consequently, there exists $c(\kappa_1) \in \mathbf{C}$ with $|c(\kappa_1)| = 1$ such that

$$(\mathcal{M}_1 v_1)(\cdot; \kappa_1) = c(\kappa_1) \cdot v_1(\cdot; 2\pi - \kappa_1). \quad (3.30)$$

By letting $\kappa_1 = \pi$ in (3.30), we see that $v_1(\cdot; \pi)$ is an eigenfunction of the reflection operator \mathcal{M}_1 with eigenvalue $c(\pi)$. Hence, by (3.27), we know $c(\pi) = -1$.

By conjugating the equation (3.30) with $v_1(\cdot; 2\pi - \kappa_1)$, we have

$$c(\kappa_1) = ((\mathcal{M}_1 v_1)(\cdot; \kappa_1), v_1(\cdot; 2\pi - \kappa_1)). \quad (3.31)$$

The analyticity of $v_1(\cdot; \kappa_1)$ in κ_1 implies that $c(\kappa_1)$ is smooth near $\kappa_1 = \pi$. By differentiating (3.31), we obtain

$$\begin{aligned} c'(\pi) &= (\mathcal{M}_1(\partial_{\kappa_1} v_1)(\cdot; \pi), v_1(\mathbf{x}; \pi)) - (\mathcal{M}_1 v_1(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi)) \\ &= (\partial_{\kappa_1} v_1(\cdot; \pi), \mathcal{M}_1 v_1(\mathbf{x}; \pi)) - (\mathcal{M}_1 v_1(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi)). \end{aligned}$$

Using (3.27), we further obtain

$$c'(\pi) = -(\partial_{\kappa_1} v_1(\cdot; \pi), v_1(\mathbf{x}; \pi)) + (v_1(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi)) = -2i \text{Im}(\partial_{\kappa_1} v_1(\cdot; \pi), v_1(\cdot; \pi)) = 0. \quad (3.32)$$

Hence (3.28) implies $c'(\pi) = 0$. By differentiating (3.30) at $\kappa_1 = \pi$, we have

$$(\mathcal{M}_1 \partial_{\kappa_1} v_n)(\cdot; \pi) = -c(\pi) \cdot (\partial_{\kappa_1} v_n)(\cdot; \pi) + c'(\pi) \cdot v_n(\cdot; \pi) = (\partial_{\kappa_1} v_n)(\cdot; \pi),$$

which concludes the proof of (3.29) for $n = 1$. □

Remark 3.9. *An explicit construction of $v_n(\mathbf{x}; \kappa_1)$ for $|\kappa_1 - \pi| \ll 1$ that satisfies (3.28) is given in the proof of Theorem 4.1 by using the perturbation argument; see (4.27).*

The parity property (3.29) greatly simplifies the calculation of energy fluxes involving $v_n(\cdot; \pi)$ and $(\partial_{\kappa_1} v_n)(\cdot; \pi)$ ($n = 1, 2$), as is illustrated below. We first note that by (3.27) and (3.29),

$$v_1(\mathbf{x}; \pi) = \partial_{\kappa_1} v_2(\mathbf{x}; \pi) = 0, \quad \frac{\partial v_2}{\partial x_1}(\mathbf{x}; \pi) = \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) = 0, \quad \mathbf{x} \in \Gamma. \quad (3.33)$$

Proposition 3.10. *Assume (3.29) holds. Then we have*

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \cdot \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)} = -\frac{i}{2}\gamma_*, \quad (3.34)$$

$$\int_{\Gamma} \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\cdot; \pi) \cdot \overline{v_2(\cdot; \pi)} = \frac{i}{2}\gamma_*. \quad (3.35)$$

Moreover,

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \cdot \overline{v_2(\cdot; \pi)} = 0, \quad (3.36)$$

$$\int_{\Gamma} \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\cdot; \pi) \cdot \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)} = 0. \quad (3.37)$$

Proof. We first prove (3.34). We point out that the proof of (3.35) is similar. By (3.17),

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \kappa_1) \overline{v_1(\cdot; \kappa_1)} - v_1(\cdot; \kappa_1) \overline{\frac{\partial v_1}{\partial x_1}(\cdot; \kappa_1)} = i\mu'_1(\kappa_1). \quad (3.38)$$

Using the time-reversal symmetry, there exists $c(\kappa_1) \in \mathbf{C}$ such that $|c(\kappa_1)| = 1$ and

$$v_1(\mathbf{x}; \kappa_1) = c(\kappa_1) \cdot \overline{v_1(\mathbf{x}; 2\pi - \kappa_1)}.$$

Then (3.38) implies that

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \kappa_1) \overline{v_1(\cdot; \kappa_1)} - \overline{v_1(\cdot; 2\pi - \kappa_1)} \frac{\partial v_1}{\partial x_1}(\cdot; 2\pi - \kappa_1) = i\mu'_1(\kappa_1).$$

By taking derivative at $\kappa_1 = \pi$ and applying (3.33), we obtain

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)} + \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) = i\mu''_1(\pi) = -i\gamma_*,$$

which gives (3.34).

We next prove (3.36). By Proposition 3.7, we have

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \kappa_1) \overline{v_2(\cdot; \kappa_1)} - v_1(\cdot; \kappa_1) \overline{\frac{\partial v_2}{\partial x_1}(\cdot; \kappa_1)} = 0, \quad \kappa_1 \neq \pi.$$

Since the functions inside the integral are smooth in κ_1 , we can let $\kappa_1 \rightarrow \pi$ to obtain

$$\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \overline{v_2(\cdot; \pi)} - v_1(\cdot; \pi) \overline{\frac{\partial v_2}{\partial x_1}(\cdot; \pi)} = 0.$$

By (3.33), we have $\int_{\Gamma} v_1(\cdot; \pi) \overline{\frac{\partial v_2}{\partial x_1}(\cdot; \pi)} = 0$. Thus $\int_{\Gamma} \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \overline{v_2(\cdot; \pi)} = 0$, which is (3.36).

Finally, we prove (3.37). Notice that (3.19) gives that $(\operatorname{div}(A\nabla) + \lambda_*)\partial_{\kappa_1}v_1(\cdot; \pi) = 0$ since $\mu'_1(\pi) = 0$. Thus, we can apply the Green's formula to $\partial_{\kappa_1}v_1$ and the fundamental solution G_0 in (3.7) inside the domain $(0, N) \times (-\frac{1}{2}, \frac{1}{2}) \subset \Omega$ to get

$$\begin{aligned} \overline{\partial_{\kappa_1}v_1(\mathbf{y}; \pi)} &= \int_{(0, N) \times (-\frac{1}{2}, \frac{1}{2})} \left((\operatorname{div}(A\nabla) + \lambda_*)G_0(\mathbf{x}, \mathbf{y}; \lambda_*) \cdot \overline{\partial_{\kappa_1}v_1(\mathbf{x}; \pi)} \right. \\ &\quad \left. - (\operatorname{div}(A\nabla) + \lambda_*)\overline{\partial_{\kappa_1}v_1(\mathbf{x}; \pi)} \cdot G_0(\mathbf{x}, \mathbf{y}; \lambda_*) \right) \\ &= \int_{\Gamma_N} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*)\overline{\partial_{\kappa_1}v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\mathbf{x}; \pi)} \\ &\quad - \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*)\overline{\partial_{\kappa_1}v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\mathbf{x}; \pi)}, \end{aligned} \quad (3.39)$$

where in the last line above we used the fact that $A(\mathbf{x}) = I$ for $\mathbf{x} \in \Gamma$ and $\mathbf{x} \in \Gamma_N$. By (3.10),

$$\begin{aligned} &\lim_{N \rightarrow \infty} \int_{\Gamma_N} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*)\overline{\partial_{\kappa_1}v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\mathbf{x}; \pi)} \\ &= \frac{i}{\gamma_*} \left(\lim_{N \rightarrow \infty} \mathbf{q}(v_1(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma_N) \right) \cdot \overline{\partial_{\kappa_1}v_1(\mathbf{y}; \pi)} \\ &\quad + \frac{i}{\gamma_*} \left(\lim_{N \rightarrow \infty} \mathbf{q}(\partial_{\kappa_1}v_1(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma_N) \right) \cdot \overline{v_1(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \left(\lim_{N \rightarrow \infty} \mathbf{q}(v_2(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma_N) \right) \cdot \overline{\partial_{\kappa_1}v_2(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \left(\lim_{N \rightarrow \infty} \mathbf{q}(\partial_{\kappa_1}v_2(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma_N) \right) \cdot \overline{v_2(\mathbf{y}; \pi)}. \end{aligned} \quad (3.40)$$

Since all the energy fluxes in (3.40) are independent of N (a consequence of the fact that $(\operatorname{div}(A\nabla) + \lambda_*)\partial_{\kappa_1}v_n(\cdot; \pi) = (\operatorname{div}(A\nabla) + \lambda_*)v_n(\cdot; \pi) = 0$ for $n = 1, 2$),

$$\begin{aligned} &\lim_{N \rightarrow \infty} \int_{\Gamma_N} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*)\overline{\partial_{\kappa_1}v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\mathbf{x}; \pi)} \\ &= \frac{i}{\gamma_*} \mathbf{q}(v_1(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1}v_1(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1}v_1(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma) \cdot \overline{v_1(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \mathbf{q}(v_2(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1}v_2(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1}v_2(\cdot; \pi), \partial_{\kappa_1}v_1(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)}. \end{aligned} \quad (3.41)$$

On the other hand, (3.13)-(3.16) and (3.9) imply that

$$\begin{aligned} &\lim_{\mathbf{y} \rightarrow \Gamma} \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*)\overline{\partial_{\kappa_1}v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\cdot; \pi)} \\ &= -\frac{1}{2} \overline{\partial_{\kappa_1}v_1(\mathbf{y}; \pi)} - \int_{\Gamma} G_0(\mathbf{y}, \cdot; \lambda_*)\overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1}v_1)(\cdot; \pi)} \\ &= -\frac{1}{2} \overline{\partial_{\kappa_1}v_1(\mathbf{y}; \pi)}, \end{aligned}$$

where we used (3.33) in the last equality. By letting $N \rightarrow \infty$ and $\mathbf{y} \rightarrow \Gamma$ in (3.39), we obtain

$$\begin{aligned} \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} &= \frac{i}{\gamma_*} \mathbf{q}(v_1(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_1(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{v_1(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \mathbf{q}(v_2(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_2(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)} \\ &\quad + \frac{1}{2} \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)}. \end{aligned} \tag{3.42}$$

Using (3.33) and (3.34), we further obtain

$$\overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} = \frac{1}{2} \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} + \frac{1}{2} \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)}.$$

It follows that

$$\mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)} = 0, \quad \mathbf{y} \in \Gamma.$$

Since $v_2(\cdot; \pi)|_{\Gamma} \neq 0$ (otherwise we have a contradiction to (3.35)), we conclude that

$$\mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), \partial_{\kappa_1} v_1(\cdot; \pi); \Gamma) = 0.$$

Then the proof of (3.37) is complete by using (3.33) again. \square

Proposition 3.10 and the identities (3.33) immediately lead to

Corollary 3.11. *Assume (3.29) holds. Then*

$$\mathbf{q}(v_1(\cdot; \pi), (\partial_{\kappa_1} v_1)(\cdot; \pi); \Gamma) = -\frac{i}{2} \gamma_*, \quad \mathbf{q}(v_2(\cdot; \pi), (\partial_{\kappa_1} v_2)(\cdot; \pi); \Gamma) = \frac{i}{2} \gamma_*.$$

Moreover,

$$\begin{aligned} \mathbf{q}(v_1(\cdot; \pi), v_2(\cdot; \pi); \Gamma) &= 0, \\ \mathbf{q}(v_1(\cdot; \pi), (\partial_{\kappa_1} v_2)(\cdot; \pi); \Gamma) &= 0, \quad \mathbf{q}(v_2(\cdot; \pi), (\partial_{\kappa_1} v_1)(\cdot; \pi); \Gamma) = 0, \\ \mathbf{q}((\partial_{\kappa_1} v_n)(\cdot; \pi), (\partial_{\kappa_1} v_m)(\cdot; \pi); \Gamma) &= 0, \quad n, m \in \{1, 2\}. \end{aligned}$$

4 Band-gap opening at the quadratic degenerate point

In this section, we derive asymptotic expansion of the Bloch eigen-pairs $(\mu_{n,\delta}(\kappa_1), v_{n,\delta}(\mathbf{x}; \kappa_1))$ of the operator $\mathcal{L}_{\Omega,\pi}^{A+\delta B}(\kappa_1)$ for $n = 1, 2$ and $0 < |\delta| \ll 1$. Consequently, it's clear that the perturbation lifts quadratic degeneracy at $\kappa_1 = \pi$ is lifted by the perturbation, and a band gap is opened as described in Theorem 1.6. The main result of this section reads as follows.

Theorem 4.1. *Suppose $|\delta| \ll 1$ and $|\kappa_1| \ll 1$. Then*

$$\begin{aligned} \mu_{1,\delta}(\kappa_1) &= \lambda_* - \sqrt{\frac{1}{4} \gamma_*^2 (\kappa_1 - \pi)^4 + t_*^2 \delta^2} \cdot (1 + \mathcal{O}(|\kappa_1 - \pi| + |\delta|)), \\ \mu_{2,\delta}(\kappa_1) &= \lambda_* + \sqrt{\frac{1}{4} \gamma_*^2 (\kappa_1 - \pi)^4 + t_*^2 \delta^2} \cdot (1 + \mathcal{O}(|\kappa_1 - \pi| + |\delta|)), \end{aligned} \tag{4.1}$$

where

$$t_* := -(\mathcal{L}_{\Omega,\pi}^B(\pi)v_1(\cdot; \pi), v_2(\cdot; \pi)). \quad (4.2)$$

In addition, there exist normalized Bloch eigenfunctions $v_i(\mathbf{x}; \kappa_1)$ ($i = 1, 2$) of the operator $\mathcal{L}_{\Omega,\pi}^A(\kappa_1)$ which are analytic in κ_1 and satisfy (3.28), such that

$$\begin{aligned} v_{1,\delta}(\mathbf{x}; \kappa_1) &= (1 + \mathcal{O}(|\kappa_1 - \pi|))v_1(\mathbf{x}; \pi) \\ &+ \left(\frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \mathcal{O}(|\kappa_1 - \pi|) \right) \cdot v_2(\mathbf{x}; \pi) \\ &+ (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) + \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_1(\mathbf{x}) \\ &+ R_1(\mathbf{x}; \kappa_1, \delta), \end{aligned} \quad (4.3)$$

$$\begin{aligned} v_{2,\delta}(\mathbf{x}; \kappa_1) &= \left(\frac{-t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \mathcal{O}(|\kappa_1 - \pi|) \right) \cdot v_1(\mathbf{x}; \pi) \\ &+ (1 + \mathcal{O}(|\kappa_1 - \pi|))v_2(\mathbf{x}; \pi) \\ &+ (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) + \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_2(\mathbf{x}) \\ &+ R_2(\mathbf{x}; \kappa_1, \delta). \end{aligned} \quad (4.4)$$

Here $r_i(\mathbf{x}), R_i(\mathbf{x}; \kappa_1, \delta) \in H^1(Y)$, $i = 1, 2$, have the following estimate

$$\|r_i(\mathbf{x})\|_{H^1(Y)} = \mathcal{O}(1), \quad \|R_i(\mathbf{x}; \kappa_1, \delta)\|_{H^1(Y)} = \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2).$$

Remark 4.2. Suppose $\delta > 0$ and $t_* \neq 0$. From (4.3) and (4.4), we see that

$$\begin{aligned} v_{1,\delta}(\mathbf{x}; \pi) &= v_1(\mathbf{x}; \pi) + \operatorname{sgn}(t_*)v_1(\mathbf{x}; \pi) + \mathcal{O}(\delta), & v_{2,\delta}(\mathbf{x}; \pi) &= -\operatorname{sgn}(t_*)v_1(\mathbf{x}; \pi) + v_1(\mathbf{x}; \pi) + \mathcal{O}(\delta), \\ v_{1,-\delta}(\mathbf{x}; \pi) &= v_1(\mathbf{x}; \pi) - \operatorname{sgn}(t_*)v_1(\mathbf{x}; \pi) + \mathcal{O}(\delta), & v_{2,-\delta}(\mathbf{x}; \pi) &= \operatorname{sgn}(t_*)v_1(\mathbf{x}; \pi) + v_1(\mathbf{x}; \pi) + \mathcal{O}(\delta). \end{aligned}$$

This implies the following relation when we drop the $\mathcal{O}(\delta)$ terms

$$v_{1,\delta}(\mathbf{x}; \pi) \sim v_{2,-\delta}(\mathbf{x}; \pi), \quad v_{2,\delta}(\mathbf{x}; \pi) \sim v_{1,-\delta}(\mathbf{x}; \pi).$$

In other words, $\mathcal{L}_{\Omega,\pi}^{A+\delta \cdot B}(\pi)$ and $\mathcal{L}_{\Omega,\pi}^{A-\delta \cdot B}(\pi)$ exchange their eigenspaces near the energy λ_* . One can view this band-inversion phenomenon as a kind of topological phase transition. This observation plays an important role in our proof of Theorem 1.8 (especially, in the calculation of $\mathbb{E}(h)$ operator in Proposition 5.1).

4.1 Preliminaries

We make some preparations for the proof of Theorem 4.1 in this subsection. We introduce the following operator

$$\begin{aligned} \tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1) : H &\rightarrow H^*, \\ u &\mapsto e^{-i\kappa_1 x_1} \circ \mathcal{L}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1) \circ e^{i\kappa_1 x_1} u \\ &= -(\nabla + i\kappa_1 \mathbf{e}_1) \cdot (A + \delta \cdot B)(\nabla + i\kappa_1 \mathbf{e}_1)u, \end{aligned} \quad (4.5)$$

where $H = H^1_{(0,\pi)}(\mathbf{R}^2)$ and H^* denotes the dual of H . The $H - H^*$ pairing is the natural extension of the $L^2(Y)$ -inner product, which is still denoted as (\cdot, \cdot) . For each $u \in H$, the following holds for all $v \in H$

$$(\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)u, v) = \mathfrak{a}_{\kappa_1}^{A+\delta B}(u, v) := \int_Y (A(\mathbf{x})(\nabla + i\kappa_1 \mathbf{e}_1)u(\mathbf{x})) \cdot \overline{(\nabla + i\kappa_1 \mathbf{e}_1)v(\mathbf{x})} d\mathbf{x}. \quad (4.6)$$

It is clear that the eigen-pairs of the operator $\mathcal{L}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)$ for $|(\kappa_1, \lambda) - (\pi, \lambda_*)| \ll 1$ can be obtained by solving those of $\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)$. Moreover, the latter family of operators is more suitable for perturbation arguments since their domain H is independent of both κ_1 and δ . We expand $\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)$ near $\kappa_1 = \pi$ and $\delta = 0$ to obtain

$$\begin{aligned} \tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1) &= -(\nabla + i\pi \mathbf{e}_1) \cdot A(\nabla + i\pi \mathbf{e}_1) \\ &\quad + (\kappa_1 - \pi)(-i(\nabla + i\pi \mathbf{e}_1) \cdot A\mathbf{e}_1 - i\mathbf{e}_1 \cdot A(\nabla + i\pi \mathbf{e}_1)) \\ &\quad + \delta(-(\nabla + i\pi \mathbf{e}_1) \cdot B(\nabla + i\pi \mathbf{e}_1)) \\ &\quad + (\kappa_1 - \pi)^2(\mathbf{e}_1 \cdot A\mathbf{e}_1) \\ &\quad + \delta(\kappa_1 - \pi)(-i(\nabla + i\pi \mathbf{e}_1) \cdot B\mathbf{e}_1 - i\mathbf{e}_1 \cdot B(\nabla + i\pi \mathbf{e}_1)) \\ &\quad + \delta(\kappa_1 - \pi)^2(\mathbf{e}_1 \cdot B\mathbf{e}_1) \\ &=: \tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) + (\kappa_1 - \pi)\tilde{\mathcal{L}}_1^A + \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + (\kappa_1 - \pi)^2\tilde{\mathcal{L}}_{11}^A \\ &\quad + \delta(\kappa_1 - \pi)\tilde{\mathcal{L}}_1^B + \delta(\kappa_1 - \pi)^2\tilde{\mathcal{L}}_{11}^B. \end{aligned} \quad (4.7)$$

Let

$$v_n(\mathbf{x}) := u_n(\mathbf{x}; \boldsymbol{\kappa}^{(2)}), \quad \tilde{v}_n(\mathbf{x}) := e^{-i\pi x_1}v_n(\mathbf{x}), \quad n = 1, 2$$

where $u_n(\mathbf{x}; \boldsymbol{\kappa}^{(2)})$ are the Bloch eigenfunctions at $\boldsymbol{\kappa} = \boldsymbol{\kappa}^{(2)}$ introduced in (1.9). Before we solve for the eigen-pairs of $\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)$, we first present some useful identities. See Appendix B for their proofs.

Lemma 4.3.

$$\left((\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi)\tilde{v}_n, \tilde{v}_m) \right)_{1 \leq n, m \leq 2} = \begin{pmatrix} 0 & -t_* \\ -t_* & 0 \end{pmatrix}, \quad t_* \in \mathbf{R},$$

where t_* is defined in (4.2).

Lemma 4.4.

$$\left((\tilde{\mathcal{L}}_1^A\tilde{v}_m, \tilde{v}_n) \right)_{1 \leq n, m \leq 2} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Lemma 4.5. *Let the projection Q_\perp be defined as in (4.12). Then*

$$\left(\left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1}Q_\perp\tilde{\mathcal{L}}_1^A)\tilde{v}_n, \tilde{v}_m \right) \right)_{1 \leq n, m \leq 2} = \begin{pmatrix} -\frac{1}{2}\gamma_* & 0 \\ 0 & \frac{1}{2}\gamma_* \end{pmatrix},$$

where γ_* is introduced in Proposition 3.2.

Lemma 4.6.

$$(ix_1v_n(\mathbf{x}), v_n(\mathbf{x})) = 0, \quad n = 1, 2.$$

The following property follows from the Fredholm alternative of second-order elliptic operators.

Lemma 4.7. $\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) : H \rightarrow H^*$ is a Fredholm operator with zero index. Moreover, $\ker(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*) = \text{span}\{\tilde{v}_1(\mathbf{x}), \tilde{v}_2(\mathbf{x})\}$.

4.2 Proof of Theorem 4.1

The goal is to solve the following eigenvalue problem by perturbation arguments

$$(\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1) - \mu)\tilde{v} = 0. \quad (4.8)$$

Step 0. The solvability of (4.8) for $|\kappa_1 - \pi| \ll 1$, $|\mu - \lambda_*| \ll 1$ and $|\delta| \ll 1$ follows from the fact that (4.8) is solvable when $\kappa_1 = \pi, \mu = \lambda_*$ and $\delta = 0$. Indeed, the expansion (4.7) shows that

$$\|\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1) - \tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi)\|_{\mathcal{B}(H,H^*)} \ll 1, \quad \text{for } |\kappa_1 - \pi|, |\delta| \ll 1. \quad (4.9)$$

Since λ_* is an isolated eigenvalue of $\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi)$ with multiplicity equal to two, the stability theorem of eigenvalues (see Chapter VI of [29]) guarantees that $\tilde{\mathcal{L}}_{\Omega,\pi}^{A+\delta \cdot B}(\kappa_1)$ has two isolated eigenvalues near λ_* for $|\kappa_1 - \pi| \ll 1$ and $|\delta| \ll 1$.

Step 1. To solve (4.8) for $|\kappa_1 - \pi| \ll 1$ and $|\delta| \ll 1$, we write

$$\mu = \lambda_* + \mu^{(1)}, \quad \kappa_1 = \pi + p, \quad \tilde{v} = \tilde{v}^{(0)} + \tilde{v}^{(1)} \quad (4.10)$$

with

$$|\mu^{(1)}|, |p| \ll 1, \quad \tilde{v}^{(0)} = a \cdot \tilde{v}_1(\mathbf{x}) + b \cdot \tilde{v}_2(\mathbf{x}) \in H_1, \quad \tilde{v}^{(1)} \in H_2,$$

where $H_1 := \ker(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)$ and H_2 is the orthogonal complement of H_1 in the Hilbert space H . Plugging (4.10) and (4.7) into (4.8), we get

$$\begin{aligned} (\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)\tilde{v}^{(1)} &= \left(\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - p^2\tilde{\mathcal{L}}_{11}^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + \mathcal{O}(|p\delta| + |p^2\delta|) \right) \tilde{v}^{(0)} \\ &+ \left(\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - p^2\tilde{\mathcal{L}}_{11}^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + \mathcal{O}(|p\delta| + |p^2\delta|) \right) \tilde{v}^{(1)}. \end{aligned} \quad (4.11)$$

Step 2. We solve (4.11) by following a Lyapunov-Schmidt reduction argument. To do so, we introduce the following projector

$$\begin{aligned} Q_{\perp} : H^* &\rightarrow \text{Ran}(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*), \\ f &\mapsto f - (f, \tilde{v}_1(\mathbf{x}))\tilde{v}_1(\mathbf{x}) - (f, \tilde{v}_2(\mathbf{x}))\tilde{v}_2(\mathbf{x}). \end{aligned} \quad (4.12)$$

By applying Q_{\perp} to (4.11), we obtain

$$\begin{aligned} (\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)\tilde{v}^{(1)} &= Q_{\perp} \left(\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - p^2\tilde{\mathcal{L}}_{11}^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + \mathcal{O}(|p\delta| + |p^2\delta|) \right) \tilde{v}^{(0)} \\ &+ Q_{\perp} \left(\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - p^2\tilde{\mathcal{L}}_{11}^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + \mathcal{O}(|p\delta| + |p^2\delta|) \right) \tilde{v}^{(1)}. \end{aligned}$$

By Lemma 4.7, $(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1} \in \mathcal{B}(\text{Ran}(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*), H_2)$. Hence the above equation can be rewritten as

$$(I - T)\tilde{v}^{(1)} = T\tilde{v}^{(0)}, \quad (4.13)$$

where

$$T = T(\delta, p, \mu^{(1)}) := (\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} (\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - p^2\tilde{\mathcal{L}}_{11}^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) + \mathcal{O}(|p\delta| + |p^2\delta|)).$$

For $\delta, p, \mu^{(1)}$ sufficiently small, $(I - T)^{-1} \in \mathcal{B}(H_2)$. Hence, (4.13) indicates that

$$\begin{aligned} \tilde{v}^{(1)} &= T\tilde{v}^{(0)} + ((I - T)^{-1} - I)T\tilde{v}^{(0)} \\ &= -p \cdot (\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} \tilde{\mathcal{L}}_1^A \tilde{v}^{(0)} + \mathcal{O}(|\delta| + |p|^2 + |\mu^{(1)}|^2) \tilde{v}^{(0)}. \end{aligned} \quad (4.14)$$

By plugging (4.14) into (4.11) and taking dual pairs with $\tilde{v}_n(\mathbf{x})$ ($n = 1, 2$), we obtain

$$\begin{aligned} & \left((\mu^{(1)} - p\tilde{\mathcal{L}}_1^A - \delta\tilde{\mathcal{L}}_{\Omega,\pi}^B(\pi) - p^2(\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1}Q_\perp\tilde{\mathcal{L}}_1^A))\tilde{v}^{(0)}(\mathbf{x}), \tilde{v}_n(\mathbf{x}) \right) \\ & + \mathcal{O}(|p\delta| + \delta^2 + |\mu^{(1)}|^3 + |p|^3), \quad n = 1, 2. \end{aligned} \quad (4.15)$$

By applying Lemma 4.3-4.5, (4.15) reduces to

$$\left(\mathcal{M}^{(0)}(\mu^{(1)}, p, \delta) + \mathcal{M}^{(1)}(\mu^{(1)}, p, \delta) \right) \begin{pmatrix} a \\ b \end{pmatrix} = 0, \quad (4.16)$$

where

$$\mathcal{M}^{(0)}(\mu^{(1)}, p, \delta) = \begin{pmatrix} \mu^{(1)} + \frac{1}{2}\gamma_*p^2 & t_*\delta \\ t_*\delta & \mu^{(1)} - \frac{1}{2}\gamma_*p^2 \end{pmatrix},$$

and

$$\mathcal{M}^{(1)}(\mu^{(1)}, p, \delta) = \begin{pmatrix} m_{11}^{(1)}(\mu^{(1)}, p, \delta) & m_{12}^{(1)}(\mu^{(1)}, p, \delta) \\ m_{21}^{(1)}(\mu^{(1)}, p, \delta) & m_{22}^{(1)}(\mu^{(1)}, p, \delta) \end{pmatrix}, \quad m_{ij}^{(1)} = \mathcal{O}(|p\delta| + \delta^2 + |p|^3 + |\mu^{(1)}|^3). \quad (4.17)$$

Step 3. We prove (4.1). Note that for each $\kappa_2 = \pi + p$, $\mu = \lambda_* + \mu^{(1)}$ solves the eigenvalue problem (4.8) if and only if $\mu^{(1)}$ solves

$$F(\mu^{(1)}, p, \delta) := \det \left(\mathcal{M}^{(0)} + \mathcal{M}^{(1)} \right) = (\mu^{(1)})^2 - \frac{1}{4}\gamma_*^2p^4 - t_*^2\delta^2 + \rho(\mu^{(1)}, p, \delta) = 0, \quad (4.18)$$

where $\rho(\mu^{(1)}, p, \delta)$ is smooth in its variables and satisfies that

$$\begin{aligned} \rho(\mu^{(1)}, p, \delta) & \lesssim \|\mathcal{M}^{(0)}\|_{max} \cdot \|\mathcal{M}^{(1)}\|_{max} + \|\mathcal{M}^{(1)}\|_{max}^2 \\ & = \mathcal{O}(|\mu^{(1)}p\delta| + |\mu^{(1)}\delta^2| + |p\delta^2| + |\delta^3|) \\ & \quad + \mathcal{O}(|\mu^{(1)}|^4 + |\delta^4| + |\mu^{(1)}p^3| + |p^3\delta| + |\mu^{(1)}|^3|\delta| + |p^5|). \end{aligned} \quad (4.19)$$

The partial derivative $\frac{\partial \rho}{\partial \mu^{(1)}}$ can be estimated similarly. We have

$$\frac{\partial \rho}{\partial \mu^{(1)}}(\mu^{(1)}, p, \delta) = \mathcal{O}(|p\delta| + |\mu^{(1)}|^2 + |\delta^2| + |p^3|). \quad (4.20)$$

We then solve $\mu^{(1)} = \mu^{(1)}(p, \delta)$ from (4.18) for each p and δ . First, note that $\pm\sqrt{\frac{1}{4}\gamma_*^2p^4 + t_*^2\delta^2}$ give two branches of solutions if we drop the remainder ρ from (4.18). Thus, we seek a solution to (4.18) in the following form

$$\mu^{(1)}(p, \delta) = x \cdot \sqrt{\frac{1}{4}\gamma_*^2p^4 + t_*^2\delta^2} \quad (4.21)$$

with $|x|$ close to 1. By substituting (4.21) into (4.18), we obtain the following equation of x (with p and δ being regarded as parameters)

$$H(x; p, \delta) := \frac{1}{\frac{1}{4}\gamma_*^2p^4 + t_*^2\delta^2} F(x \cdot \sqrt{\frac{1}{4}\gamma_*^2p^4 + t_*^2\delta^2}, p, \delta) = x^2 - 1 + \rho_1(x; p, \delta) = 0, \quad (4.22)$$

where

$$\rho_1(x; p, \delta) = \frac{\rho(x \cdot \sqrt{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2}, p, \delta)}{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2}.$$

Now we consider the solution to (4.22) with $|x - 1| \ll 1$. By (4.19), one can check that the following estimate holds uniformly in x when $|x - 1| \ll 1$

$$\rho_1(x; p, \delta) = \frac{\rho(x \cdot \sqrt{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2}, p, \delta)}{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2} = \mathcal{O}(|p| + |\delta|).$$

Moreover, (4.20) gives that

$$\frac{\partial H}{\partial x}(1; p, \delta) = 2 + \frac{\partial \rho_1}{\partial x}(1; p, \delta) = 2 + \mathcal{O}(|p| + |\delta|) \neq 0.$$

Thus, the implicit function theorem indicates that there exists a unique solution $x_s(p, \delta)$ to (4.22) with $\|x_s(p, \delta) - 1\| \ll 1$. The expansion of $x_s(p, \delta)$ to linear order in p is

$$x_s(p, \delta) = 1 + r_+ \cdot p + \mathcal{O}(|p|^2 + |\delta|), \quad r_+ \in \mathbf{R}.$$

It follows from (4.21) that

$$\mu_+^{(1)}(p, \delta) = x_s(p, \delta) \cdot \sqrt{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2} = \sqrt{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2} \cdot (1 + r_+ \cdot p + \mathcal{O}(|p|^2 + |\delta|)).$$

Similarly, the other solution to (4.18) has the expansion

$$\mu_-^{(1)}(p, \delta) = -\sqrt{\frac{1}{4}\gamma_*^2 p^4 + t_*^2 \delta^2} \cdot (1 + r_- \cdot p + \mathcal{O}(|p|^2 + |\delta|)).$$

Note that $\mu_{1,\delta}(\kappa_1) = \lambda_* + \mu_-^{(1)}(\kappa_1 - \pi, \delta)$ and $\mu_{2,\delta}(\kappa_1) = \lambda_* + \mu_+^{(1)}(\kappa_1 - \pi, \delta)$. This proves (4.1).

Step 4. We solve (4.16) and prove (4.3)-(4.4). By substituting $\mu^{(1)} = \mu_-^{(1)}(\kappa_1 - \pi, \delta)$ into (4.16), we obtain the following eigenvector

$$\begin{pmatrix} a(\kappa_1, \delta) \\ b(\kappa_1, \delta) \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{t_* \delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} \end{pmatrix} + \begin{pmatrix} \frac{-m_{22}^{(1)}(\mu_-^{(1)}(\kappa_1 - \pi, \delta), \kappa_1 - \pi, \delta) + r_- \cdot (\kappa_1 - \pi) \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} \\ \frac{m_{21}^{(1)}(\mu_-^{(1)}(\kappa_1 - \pi, \delta), \kappa_1 - \pi, \delta)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} \end{pmatrix}. \quad (4.23)$$

Using the estimate (4.17), we have

$$\begin{aligned} \frac{m_{22}^{(1)}(\mu_-^{(1)}(\kappa_1 - \pi, \delta), \kappa_1 - \pi, \delta)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} &= \frac{m_{22,1}^{(1)} \cdot (\kappa_1 - \pi)\delta + m_{22,2}^{(1)} \cdot (\kappa_1 - \pi)^3}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2), \\ \frac{m_{21}^{(1)}(\mu_-^{(1)}(\kappa_1 - \pi, \delta), \kappa_1 - \pi, \delta)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} &= \frac{m_{21,1}^{(1)} \cdot (\kappa_1 - \pi)\delta + m_{21,2}^{(1)} \cdot (\kappa_1 - \pi)^3}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2 \delta^2}} + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2), \end{aligned}$$

where $m_{22,1}^{(1)}, m_{22,2}^{(1)}, m_{21,1}^{(1)}, m_{21,2}^{(1)}$ are constants. By substituting (4.23) into $\tilde{v}^{(0)} = a(\kappa_1; \delta) \cdot \tilde{v}_1(\mathbf{x}) + b(\kappa_1; \delta) \cdot \tilde{v}_2(\mathbf{x})$ and utilizing (4.14), we obtain

$$\begin{aligned}
\tilde{v} &= \tilde{v}^{(0)} + \tilde{v}^{(1)} \\
&= \left(1 + \frac{-m_{22,1}^{(1)} \cdot (\kappa_1 - \pi)\delta - m_{22,2}^{(1)} \cdot (\kappa_1 - \pi)^3 + r_- \cdot (\kappa_1 - \pi) \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \right) \tilde{v}_1(\mathbf{x}) \\
&\quad + \left(\frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \frac{m_{21,1}^{(1)} \cdot (\kappa_1 - \pi)\delta + m_{21,2}^{(1)} \cdot (\kappa_1 - \pi)^3}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \right) \tilde{v}_2(\mathbf{x}) \\
&\quad - (\kappa_1 - \pi) \cdot (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} \tilde{\mathcal{L}}_1^A \tilde{v}_1(\mathbf{x}) \\
&\quad - \frac{t_*\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} \tilde{\mathcal{L}}_1^A \tilde{v}_2(\mathbf{x}) \\
&\quad + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2).
\end{aligned}$$

Let $v_{1,\delta}(\mathbf{x}; \kappa_1) = e^{i\kappa_1 x_1} \tilde{v}$. Then

$$\begin{aligned}
v_{1,\delta}(\mathbf{x}; \kappa_1) &= (e^{i\pi x_1} + i(\kappa_1 - \pi)x_1 e^{i\pi x_1} + \mathcal{O}(|\kappa_1 - \pi|^2)) \cdot \tilde{v} \\
&= \left(1 + \frac{-m_{22,1}^{(1)} \cdot (\kappa_1 - \pi)\delta - m_{22,2}^{(1)} \cdot (\kappa_1 - \pi)^3 + r_- \cdot (\kappa_1 - \pi) \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \right) v_1(\mathbf{x}) \\
&\quad + (\kappa_1 - \pi) \cdot i x_1 v_1(\mathbf{x}) \\
&\quad + \left(\frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \frac{m_{21,1}^{(1)} \cdot (\kappa_1 - \pi)\delta + m_{21,2}^{(1)} \cdot (\kappa_1 - \pi)^3}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \right) v_2(\mathbf{x}) \\
&\quad + \frac{t_*\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} i x_1 v_2(\mathbf{x}) \\
&\quad - (\kappa_1 - \pi) \cdot e^{i\pi x_1} (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} \tilde{\mathcal{L}}_1^A \tilde{v}_1(\mathbf{x}) \\
&\quad - \frac{t_*\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot e^{i\pi x_1} (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_{\perp} \tilde{\mathcal{L}}_1^A \tilde{v}_2(\mathbf{x}) \\
&\quad + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2).
\end{aligned}$$

It follows that

$$\begin{aligned}
v_{1,\delta}(\mathbf{x}; \kappa_1) &= v_1(\mathbf{x}) + \frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot v_2(\mathbf{x}) \\
&\quad + (\kappa_1 - \pi) \cdot r_{11,\delta}(\mathbf{x}; \kappa_1) \\
&\quad + \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_1(\mathbf{x}) + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2),
\end{aligned} \tag{4.24}$$

where

$$\begin{aligned}
r_{11,\delta}(\mathbf{x}; \kappa_1) &= \left(-\frac{m_{22,2}^{(1)} \cdot (\kappa_1 - \pi)^2 - r_- \cdot \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}v_1(\mathbf{x}) \right) \\
&\quad + \left(ix_1v_1(\mathbf{x}) - e^{i\pi x_1}(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1}Q_{\perp}\tilde{\mathcal{L}}_1^A\tilde{v}_1(\mathbf{x}) \right. \\
&\quad \left. + \frac{m_{21,2}^{(1)} \cdot (\kappa_1 - \pi)^2}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}v_2(\mathbf{x}) \right) \\
&=: r_{11,\delta}^{\parallel}(\mathbf{x}; \kappa_1) + r_{11,\delta}^{\perp}(\mathbf{x}; \kappa_1), \\
r_1(\mathbf{x}) &= -m_{22,1}^{(1)}v_1(\mathbf{x}) + m_{21,1}^{(1)}v_2(\mathbf{x}) \\
&\quad + it_*x_1v_2(\mathbf{x}) - t_*e^{i\pi x_1}(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1}Q_{\perp}\tilde{\mathcal{L}}_1^A\tilde{v}_2(\mathbf{x})
\end{aligned}$$

Step 5. We reduce (4.24) to (4.3) by simplifying the term $r_{11,\delta}(\mathbf{x}; \kappa_1)$. Note that when $\delta = 0$, (4.24) becomes

$$\begin{aligned}
v_{1,0}(\mathbf{x}; \kappa_1) &= v_1(\mathbf{x}) + (\kappa_1 - \pi)r_{11,0}(\mathbf{x}; \pi) + \mathcal{O}((\kappa_1 - \pi)^2), \\
&= \left(1 + \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*}(\kappa_1 - \pi) \right) v_1(\mathbf{x}) + (\kappa_1 - \pi)r_{11,0}^{\perp}(\mathbf{x}; \pi) + \mathcal{O}((\kappa_1 - \pi)^2)
\end{aligned} \tag{4.25}$$

where

$$r_{11,0}^{\perp}(\mathbf{x}; \pi) = ix_1v_1(\mathbf{x}) - e^{i\pi x_1}(\tilde{\mathcal{L}}_{\Omega,\pi}^A(\pi) - \lambda_*)^{-1}Q_{\perp}\tilde{\mathcal{L}}_1^A\tilde{v}_1(\mathbf{x}) + \frac{m_{21,2}^{(1)}}{\gamma_*}v_2(\mathbf{x})$$

It is clear that $v_{1,0}(\mathbf{x}; \kappa_1)$ defined above is a Bloch eigenfunction of the operator $\mathcal{L}_{\Omega,\pi}^A(\kappa_1)$ that is smooth in κ_1 .

We claim that

$$(r_{11,0}^{\perp}(\cdot; \pi), v_1(\cdot)) = 0. \tag{4.26}$$

Indeed, note that $\tilde{\mathcal{L}}_1^A\tilde{v}_1 \in L^2(Y)$. The spectral expansion of $(\tilde{\mathcal{L}}^A(\pi) - \lambda_*)^{-1}Q_{\perp}$ indicates that \tilde{v}_1 is orthogonal to $\text{Ran}((\tilde{\mathcal{L}}^A(\pi) - \lambda_*)^{-1}Q_{\perp})$ in L^2 -sense. Therefore

$$(e^{i\pi x_1}(\tilde{\mathcal{L}}^A(\pi) - \lambda_*)^{-1}Q_{\perp}\tilde{\mathcal{L}}_1^A\tilde{v}_1, v_1) = ((\tilde{\mathcal{L}}^A(\pi) - \lambda_*)^{-1}Q_{\perp}\tilde{\mathcal{L}}_1^A\tilde{v}_1, \tilde{v}_1) = 0,$$

On the other hand, Lemma 4.6 gives that

$$(ix_1v_1(\mathbf{x}), v_1(\mathbf{x})) = 0.$$

These combined with the identity $(v_2, v_1) = 0$ yield (4.26).

By (4.25) and (4.26),

$$\|v_{1,0}(\mathbf{x}; \kappa_1)\|_{L^2(Y)} = \sqrt{\left(1 + \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*}(\kappa_1 - \pi) \right)^2 + \mathcal{O}((\kappa_1 - \pi)^2)}.$$

Thus we obtain the following normalized Bloch eigenfunction of $\mathcal{L}_{\Omega,\pi}^A(\kappa_1)$:

$$v_1(\mathbf{x}; \kappa_1) = \frac{\left(1 + \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*}(\kappa_1 - \pi) \right) v_1(\mathbf{x}) + (\kappa_1 - \pi)r_{11,0}^{\perp}(\mathbf{x}; \pi)}{\sqrt{\left(1 + \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*}(\kappa_1 - \pi) \right)^2 + \mathcal{O}((\kappa_1 - \pi)^2)}} + \mathcal{O}((\kappa_1 - \pi)^2). \tag{4.27}$$

Note that $v_1(\mathbf{x}; \kappa_1)$ is analytic in κ_1 for $|\kappa_1 - \pi| \ll 1$. Its analytic continuation to a neighborhood of \mathbf{R} , which is guaranteed by Proposition 3.1, gives the first analytic branch of Bloch eigenfunctions of $\mathcal{L}_{\Omega, \pi}^A(\kappa_1)$. A direct calculation shows that

$$(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) = \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*} v_1(\mathbf{x}) + r_{11,0}^\perp(\mathbf{x}; \pi) - \frac{-2m_{22,2}^{(1)} + r_- \gamma_*}{2\gamma_*} v_1(\mathbf{x}) = r_{11,0}^\perp(\mathbf{x}; \pi),$$

which implies that $((\partial_{\kappa_1} v_1)(\cdot; \pi), v_1(\cdot; \pi)) = 0$ by (4.26). Hence (3.28) holds. With the normalized Bloch eigenfunction $v_1(\mathbf{x}; \kappa_1)$ defined in (4.27), we can rewrite $r_{11,\delta}(\mathbf{x}; \kappa_1)$ as

$$\begin{aligned} r_{11,\delta}(\mathbf{x}; \kappa_1) &= r_{11,\delta}^\parallel(\mathbf{x}; \kappa_1) + (r_{11,\delta}^\perp(\mathbf{x}; \kappa_1) - r_{11,0}^\perp(\mathbf{x}; \kappa_1)) + r_{11,0}^\perp(\mathbf{x}; \kappa_1) \\ &= \frac{-m_{22,2}^{(1)} \cdot (\kappa_1 - \pi)^2 + r_- \cdot \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} v_1(\mathbf{x}; \pi) \\ &\quad + \left(\frac{m_{21,2}^{(1)} \cdot (\kappa_1 - \pi)^2}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} - \frac{m_{21,2}^{(1)}}{\gamma_*} \right) v_2(\mathbf{x}; \pi) \\ &\quad + (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) \\ &= \mathcal{O}(1) \cdot v_1(\mathbf{x}; \pi) + \mathcal{O}(1) \cdot v_2(\mathbf{x}; \pi) + (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi). \end{aligned}$$

Hence (4.24) becomes

$$\begin{aligned} v_{1,\delta}(\mathbf{x}; \kappa_1) &= (1 + \mathcal{O}(|\kappa_1 - \pi|)) \cdot v_1(\mathbf{x}; \pi) \\ &\quad + \left(\frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \mathcal{O}(|\kappa_1 - \pi|) \right) \cdot v_2(\mathbf{x}; \pi) \\ &\quad + (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) + \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_1(\mathbf{x}) \\ &\quad + \mathcal{O}(|\delta| + |\kappa_1 - \pi|^2). \end{aligned}$$

This concludes the proof of (4.3). The proof of (4.4) is similar.

Step 6. We finally show the analyticity of $v_{1,\delta}(\mathbf{x}; \kappa_1)$ in κ_1 . Indeed, for $|\kappa_1 - \pi| \ll 1$, this follows from the identity

$$v_{1,\delta}(\mathbf{x}; \kappa_1) = e^{i\kappa_1 x_1} (\tilde{v}^{(0)} + \tilde{v}^{(1)}) = e^{i\kappa_1 x_1} (1 + (I - T)^{-1}T)(a(\kappa_1; \delta) \cdot \tilde{v}_1(\mathbf{x}) + b(\kappa_1; \delta) \cdot \tilde{v}_2(\mathbf{x})),$$

and the fact that $a(\kappa_1; \delta), b(\kappa_1; \delta)$ and the operator $T = T(\delta, p, \mu^{(1)})$ are analytic in κ_1 . The analyticity of $v_{1,\delta}(\mathbf{x}; \kappa_1)$ for κ_1 away from π follows from the fact that $v_{1,\delta}(\mathbf{x}; \kappa_1)$ is an isolated eigenfunction for the operator $\tilde{\mathcal{L}}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1)$ which is analytical in κ_1 .

5 Proof of Theorem 1.8

In this section, we prove Theorem 1.8 on the existence of the interface modes at $\kappa_2 = \pi$.

5.1 Boundary integral equation formulation of the interface eigenvalue problem

We start with formulating the interface eigenvalue problem in Theorem 1.8 using boundary integral equations. In the sequel, we reparametrize the band gap near λ_* by

$$\lambda = \lambda_* + \delta \cdot h, \quad h \in \mathcal{J} := \{z \in \mathbf{C} : |z| < c_0 |t_*|\}.$$

We first introduce the following Green functions G^δ for the perturbed operator $\mathcal{L}^{A+\delta \cdot B}$ on Ω

$$\begin{cases} \left(\operatorname{div}((A + \delta \cdot B)\nabla) + \lambda \right) G^\delta(\mathbf{x}, \mathbf{y}; \lambda) = \delta(\mathbf{x} - \mathbf{y}), & \mathbf{x}, \mathbf{y} \in \Omega, \\ G^\delta(\mathbf{x}, \mathbf{y}; \lambda)|_{\Gamma^+} = e^{i\pi} G^\delta(\mathbf{x}, \mathbf{y}; \lambda)|_{\Gamma^-}, \\ \frac{\partial G^\delta}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda)|_{\Gamma^+} = e^{i\pi} \frac{\partial G^\delta}{\partial x_2}(\mathbf{x}, \mathbf{y}; \lambda)|_{\Gamma^-}. \end{cases}$$

By Floquet transform,

$$G^\delta(\mathbf{x}, \mathbf{y}; \lambda) = \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 1} \frac{v_{n,\delta}(\mathbf{x}; \kappa_1) \overline{v_{n,\delta}(\mathbf{y}; \kappa_1)}}{\lambda - \mu_{n,\delta}(\kappa_1)} d\kappa_1.$$

We then introduce the following layer potential operators associated with G^δ .

$$\begin{aligned} \mathcal{S}(\lambda; G^\delta) : \tilde{H}^{-\frac{1}{2}}(\Gamma) &\rightarrow H_{loc}^1(\Omega), & \varphi &\mapsto \int_{\Gamma} G^\delta(\mathbf{x}, \mathbf{y}; \lambda) \varphi(\mathbf{y}) ds(\mathbf{y}), \\ \mathcal{D}(\lambda; G^\delta) : H^{\frac{1}{2}}(\Gamma) &\rightarrow H_{loc}^1(\Omega), & \phi &\mapsto \int_{\Gamma} \frac{\partial G^\delta}{\partial y_1}(\mathbf{x}, \mathbf{y}; \lambda) \phi(\mathbf{y}) ds(\mathbf{y}), \\ \mathcal{K}^*(\lambda; G^\delta) : \tilde{H}^{-\frac{1}{2}}(\Gamma) &\rightarrow \tilde{H}^{-\frac{1}{2}}(\Gamma), & \varphi &\mapsto p.v. \int_{\Gamma} \frac{\partial G^\delta}{\partial x_1}(\mathbf{x}, \mathbf{y}; \lambda) \varphi(\mathbf{y}) ds(\mathbf{y}), \\ \mathcal{K}(\lambda; G^\delta) : H^{\frac{1}{2}}(\Gamma) &\rightarrow H^{\frac{1}{2}}(\Gamma), & \phi &\mapsto p.v. \int_{\Gamma} \frac{\partial G^\delta}{\partial y_1}(\mathbf{x}, \mathbf{y}; \lambda) \phi(\mathbf{y}) ds(\mathbf{y}), \\ \mathcal{N}(\lambda; G^\delta) : H^{\frac{1}{2}}(\Gamma) &\rightarrow \tilde{H}^{-\frac{1}{2}}(\Gamma), & \phi &\mapsto \int_{\Gamma} \frac{\partial^2 G^\delta}{\partial x_1 \partial y_1}(\mathbf{x}, \mathbf{y}; \lambda) \phi(\mathbf{y}) ds(\mathbf{y}). \end{aligned} \tag{5.1}$$

We have the following jump formulas, which are analogous to those in Proposition 3.5.

$$\lim_{t \rightarrow 0} \left(\mathcal{S}(\lambda; G^\delta)[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mathcal{S}(\lambda; G^\delta)[\varphi](\mathbf{x}), \tag{5.2}$$

$$\lim_{t \rightarrow 0} \frac{\partial}{\partial x_1} \left(\mathcal{D}(\lambda; G^\delta)[\phi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mathcal{N}(\lambda; G^\delta)[\phi](\mathbf{x}), \tag{5.3}$$

$$\lim_{t \rightarrow 0^\pm} \frac{\partial}{\partial x_1} \left(\mathcal{S}(\lambda; G^\delta)[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \left(\pm \frac{1}{2} + \mathcal{K}^*(\lambda; G^\delta) \right) \varphi(\mathbf{x}), \tag{5.4}$$

$$\lim_{t \rightarrow 0^\pm} \left(\mathcal{D}(\lambda; G^\delta)[\phi] \right) (\mathbf{x} + t\mathbf{e}_1) = \left(\mp \frac{1}{2} + \mathcal{K}(\lambda; G^\delta) \right) \phi(\mathbf{x}), \tag{5.5}$$

where $\mathbf{x} \in \Gamma$. Now we fix $\delta > 0$. We construct an interface mode of (1.11) in the following form

$$u(\mathbf{x}; \lambda) = \begin{cases} -\mathcal{D}(\lambda; G^\delta)[\phi](\mathbf{x}) + \mathcal{S}(\lambda; G^\delta)[\varphi](\mathbf{x}), & x_1 > 0, \\ \mathcal{D}(\lambda; G^{-\delta})[\phi](\mathbf{x}) - \mathcal{S}(\lambda; G^{-\delta})[\varphi](\mathbf{x}), & x_1 < 0. \end{cases} \tag{5.6}$$

By the jump formulas, for $\mathbf{x} \in \Gamma$, we have

$$\lim_{t \rightarrow 0^+} \begin{pmatrix} u(\mathbf{x} + t\mathbf{e}_1) \\ \frac{\partial u}{\partial x_1}(\mathbf{x} + t\mathbf{e}_1) \end{pmatrix} = \begin{pmatrix} \frac{1}{2} - \mathcal{K}(\lambda; G^\delta) & \mathcal{S}(\lambda; G^\delta) \\ -\mathcal{N}(\lambda; G^\delta) & \frac{1}{2} + \mathcal{K}^*(\lambda; G^\delta) \end{pmatrix} \begin{pmatrix} \phi \\ \varphi \end{pmatrix},$$

and

$$\lim_{t \rightarrow 0^-} \begin{pmatrix} u(\mathbf{x} + t\mathbf{e}_1) \\ \frac{\partial u}{\partial x_1}(\mathbf{x} + t\mathbf{e}_1) \end{pmatrix} = \begin{pmatrix} \frac{1}{2} + \mathcal{K}(\lambda; G^{-\delta}) & -\mathcal{S}(\lambda; G^{-\delta}) \\ \mathcal{N}(\lambda; G^{-\delta}) & \frac{1}{2} - \mathcal{K}^*(\lambda; G^{-\delta}) \end{pmatrix} \begin{pmatrix} \phi \\ \varphi \end{pmatrix}.$$

Therefore, (5.6) is an interface mode if and only if

$$\lim_{t \rightarrow 0^+} \begin{pmatrix} u(\mathbf{x} + t\mathbf{e}_1) \\ \frac{\partial u}{\partial x_1}(\mathbf{x} + t\mathbf{e}_1) \end{pmatrix} = \lim_{t \rightarrow 0^-} \begin{pmatrix} u(\mathbf{x} + t\mathbf{e}_1) \\ \frac{\partial u}{\partial x_1}(\mathbf{x} + t\mathbf{e}_1) \end{pmatrix} \neq 0.$$

This leads to the following system of equations

$$\left(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda) \right) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} = 0, \quad (5.7)$$

$$\left(\frac{1}{2} + \mathbb{T}^\delta(\lambda) \right) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} \neq 0, \quad (5.8)$$

where

$$\mathbb{T}^\delta \in \mathcal{B}(H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma)), \quad \mathbb{T}^\delta \begin{pmatrix} \phi \\ \varphi \end{pmatrix} := \begin{pmatrix} -\mathcal{K}(\lambda; G^\delta) & \mathcal{S}(\lambda; G^\delta) \\ -\mathcal{N}(\lambda; G^\delta) & \mathcal{K}^*(\lambda; G^\delta) \end{pmatrix} \begin{pmatrix} \phi \\ \varphi \end{pmatrix}.$$

We point out that any interface mode can be expressed in the form (5.6). Hence (5.7) and (5.8) give an equivalent formulation of the interface mode problem.

We next introduce the following projections on $H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma)$

$$\Pi_1 : \begin{pmatrix} \phi \\ \varphi \end{pmatrix} \mapsto \frac{\int_\Gamma \phi(\cdot) \overline{\frac{\partial v_1}{\partial x_1}(\cdot; \pi)}}{i\gamma_*/2} \begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix} + \frac{\int_\Gamma \varphi(\cdot) \overline{v_2(\cdot; \pi)}}{i\gamma_*/2} \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \end{pmatrix},$$

$$\Pi_2 : \begin{pmatrix} \phi \\ \varphi \end{pmatrix} \mapsto \frac{\int_\Gamma \varphi(\cdot) \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)}}{-i\gamma_*/2} \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi) \end{pmatrix} + \frac{\int_\Gamma \phi(\cdot) \overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\cdot; \pi)}}{-i\gamma_*/2} \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix},$$

and

$$\mathbb{Q} = \mathbb{I}_{2 \times 2} - \Pi_1 - \Pi_2.$$

Here the Bloch eigenfunctions $v_n(\mathbf{x}; \kappa_1)$ are the ones constructed in Theorem 4.1. By Proposition 3.10, Π_1 and Π_2 are orthogonal in the sense that

$$\Pi_1 \cdot \Pi_2 = \Pi_2 \cdot \Pi_1 = 0.$$

We have the following decomposition

$$H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma) = X \oplus Y \oplus Y^\perp, \quad X = \text{Ran}(\mathbb{Q}), Y = \text{Ran}(\Pi_1), Y^\perp = \text{Ran}(\Pi_2). \quad (5.9)$$

Then (ϕ, φ) solves (5.7) if and only if

$$\begin{pmatrix} \mathbb{Q}(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\mathbb{Q} & \mathbb{Q}(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_1 & \mathbb{Q}(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_2 \\ \Pi_2(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\mathbb{Q} & \Pi_2(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_1 & \Pi_2(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_2 \\ \Pi_1(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\mathbb{Q} & \Pi_1(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_1 & \Pi_1(\mathbb{T}^\delta(\lambda) + \mathbb{T}^{-\delta}(\lambda))\Pi_2 \end{pmatrix} \begin{pmatrix} \Psi \\ \Phi^{(1)} \\ \Phi^{(1)} \end{pmatrix} = 0, \quad (5.10)$$

where $\Psi = \mathbb{Q} \begin{pmatrix} \phi \\ \varphi \end{pmatrix}$, $\Phi^{(1)} = \Pi_1 \begin{pmatrix} \phi \\ \varphi \end{pmatrix}$, $\Phi^{(1)} = \Pi_2 \begin{pmatrix} \phi \\ \varphi \end{pmatrix}$. By multiplying $\begin{pmatrix} 1 & 0 & 0 \\ 0 & \delta^{\frac{1}{4}} & 0 \\ 0 & 0 & \delta^{-\frac{1}{12}} \end{pmatrix}$ to both sides of the matrix in (5.10), we further obtain

$$\left(\mathbb{M}^\delta(\lambda) + \mathbb{M}^{-\delta}(\lambda) \right) \begin{pmatrix} \Psi \\ \Phi^{(1)} \\ \Phi^{(1)} \end{pmatrix} = 0, \quad (5.11)$$

where

$$\mathbb{M}^\delta(\lambda) := \begin{pmatrix} \mathbb{Q}\mathbb{T}^\delta(\lambda)\mathbb{Q} & \delta^{\frac{1}{4}}\mathbb{Q}\mathbb{T}^\delta(\lambda)\Pi_1 & \delta^{-\frac{1}{12}}\mathbb{Q}\mathbb{T}^\delta(\lambda)\Pi_2 \\ \delta^{\frac{1}{4}}\Pi_2\mathbb{T}^\delta(\lambda)\mathbb{Q} & \delta^{\frac{1}{2}}\Pi_2\mathbb{T}^\delta(\lambda)\Pi_1 & \delta^{\frac{1}{6}}\Pi_2\mathbb{T}^\delta(\lambda)\Pi_2 \\ \delta^{-\frac{1}{12}}\Pi_1\mathbb{T}^\delta(\lambda)\mathbb{Q} & \delta^{\frac{1}{6}}\Pi_1\mathbb{T}^\delta(\lambda)\Pi_1 & \delta^{-\frac{1}{6}}\Pi_1\mathbb{T}^\delta(\lambda)\Pi_2 \end{pmatrix}. \quad (5.12)$$

On the other hand, (5.8) is equivalent to

$$\left(\frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \delta^{\frac{1}{6}} \\ 0 & \delta^{\frac{1}{4}} & 0 \end{pmatrix} + \mathbb{M}^\delta(\lambda) \right) \begin{pmatrix} \Psi \\ \Phi^{(1)} \\ \Phi^{(1)} \end{pmatrix} \neq 0. \quad (5.13)$$

We remark that the scaled matrix $\mathbb{M}^\delta(\lambda)$ is a normalized version of the matrix on the left side of (5.10). This will be illustrated in the next section.

5.2 Properties of $\mathbb{M}^\delta(\lambda)$

We present asymptotic properties of the operator $\mathbb{M}^\delta(\lambda)$ in this subsection.

Proposition 5.1. *The following convergence holds uniformly for $h \in \overline{\mathcal{J}}$ as $\delta \rightarrow 0^+$:*

$$\mathbb{M}^\delta(\lambda_* + \delta \cdot h) \xrightarrow{\|\cdot\|_{\mathcal{B}(X \oplus Y \oplus Y^\perp, X \oplus Y^\perp \oplus Y)}} \begin{pmatrix} \mathbb{Q}\mathbb{T}^0(\lambda_*)\mathbb{Q} & 0 & 0 \\ 0 & \mathbb{E}(h) & 0 \\ 0 & 0 & \mathbb{F} \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathbb{E}^\times(h) & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (5.14)$$

where

$$\mathbb{T}^0(\lambda_*) \in \mathcal{B}(H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma)), \quad \mathbb{T}^0(\lambda_*) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} := \begin{pmatrix} 0 & \mathcal{S}(\lambda_*; G_0) \\ -\mathcal{N}(\lambda_*; G_0) & 0 \end{pmatrix} \begin{pmatrix} \phi \\ \varphi \end{pmatrix},$$

$$\begin{aligned} \mathbb{E}(h) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} &:= -\frac{1}{2\gamma_*} \cdot \left(\frac{\frac{h}{\gamma_*}}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{3}{4}}} + \frac{1}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{1}{4}}} \right) \cdot \left(\int_\Gamma \varphi(\cdot) \overline{v_2(\cdot; \pi)} \right) \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix} \\ &+ \frac{1}{2\gamma_*} \cdot \left(\frac{\frac{h}{\gamma_*}}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{3}{4}}} - \frac{1}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{1}{4}}} \right) \cdot \left(\int_\Gamma \phi(\cdot) \overline{\frac{\partial v_1}{\partial x_1}(\cdot; \pi)} \right) \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \end{pmatrix}, \end{aligned}$$

$$\mathbb{E}^\times(h) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} := -\frac{1}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{3}{4}}} \left[\left(\int_\Gamma \phi(\cdot) \overline{\frac{\partial v_1}{\partial x_1}(\cdot; \pi)} \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix} - \left(\int_\Gamma \varphi(\cdot) \overline{v_2(\cdot; \pi)} \right) \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\cdot; \pi) \end{pmatrix} \right), \right.$$

$$\mathbb{F} \begin{pmatrix} \phi \\ \varphi \end{pmatrix} := \frac{4}{\pi\gamma_*} \left(\int_\Gamma \varphi(\cdot) \overline{(\partial_{\kappa_1} v_1)(\cdot; \pi)} \right) \begin{pmatrix} (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) \\ 0 \end{pmatrix} - \frac{4}{\pi\gamma_*} \left(\int_\Gamma \phi(\cdot) \overline{\frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\cdot; \pi)} \right) \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_1)(\cdot; \pi) \end{pmatrix}.$$

Moreover,

$$\mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h) \xrightarrow{\|\cdot\|_{\mathcal{B}(X \oplus Y \oplus Y^*, X \oplus Y^* \oplus Y)}} \begin{pmatrix} \mathbb{Q}\mathbb{T}^0(\lambda_*)\mathbb{Q} & 0 & 0 \\ 0 & \mathbb{E}(h) & 0 \\ 0 & 0 & \mathbb{F} \end{pmatrix} - \begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathbb{E}^\times(h) & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (5.15)$$

Proof. See Appendix C. □

Proposition 5.2. $\mathbb{T}^0(\lambda_*) \in \mathcal{B}(H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma))$ is a Fredholm operator with zero index. Moreover,

$$\ker \mathbb{T}^0(\lambda_*) = \text{span} \left\{ \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi) \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \end{pmatrix} \right\}. \quad (5.16)$$

Proof. See Appendix D. □

5.3 Proof of Theorem 1.8

We now prove Theorem 1.8 in this subsection. First, using (5.14)-(5.15), the characteristic values of the operator $\lim_{\delta \rightarrow 0} \left(\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h) \right)$ in \mathcal{J} can be obtained by solving $\mathbb{E}(h)\Phi = 0$. This yields the following two characteristic values

$$h_1 = \frac{1}{\sqrt{2}}|t_*|, \quad h_2 = -\frac{1}{\sqrt{2}}|t_*|.$$

Their associated eigenfunctions are $(\Psi_1, \Phi_1^{(1)}, \Phi_1^{(2)})^T$ and $(\Psi_2, \Phi_2^{(1)}, \Phi_2^{(2)})^T$, respectively, where

$$\Psi_1 = 0, \quad \Phi_1^{(1)} = \begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \quad \Phi_1^{(2)} = 0, \quad (5.17)$$

$$\Psi_2 = 0, \quad \Phi_2^{(1)} = \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \end{pmatrix}, \quad \Phi_2^{(2)} = 0. \quad (5.18)$$

Next, note the following facts:

(i) $\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h)$ is Fredholm with zero index for any $h \in \overline{\mathcal{J}}$. Indeed, since $\mathbb{T}^0(\lambda_*)$ is Fredholm, we know the limiting operator of $\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h)$ is Fredholm with zero index because it differs from $2\mathbb{T}^0(\lambda_*)$ by finite-rank operators. Consequently, $\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h)$ is Fredholm by the convergence (5.14) and (5.15);

(ii) $\mathbb{M}^{\pm\delta}(\lambda_* + \delta \cdot h)$ are analytic for $h \in \mathcal{J}$ because the Green function $G^\delta(\mathbf{x}, \mathbf{y}; \lambda)$ is analytic when λ lies in the band gap;

(iii) $\lim_{\delta \rightarrow 0} \left(\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h) \right)$ is invertible for $h \in \overline{\mathcal{J}} \setminus \{h_1, h_2\}$. This follows from the facts that $\lim_{\delta \rightarrow 0} \left(\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h) \right)$ is Fredholm and it's injective for $h \neq h_1$ and $h \neq h_2$ as we proved earlier.

Hence, the generalized Rouché theorem (cf. Chapter 1 of [3]) concludes that $\mathbb{M}^\delta(\lambda_* + \delta \cdot h) + \mathbb{M}^{-\delta}(\lambda_* + \delta \cdot h)$ has the following two characteristic values

$$h_1(\delta) = \frac{1}{\sqrt{2}}|t_*| + o(1), \quad h_1(\delta) = -\frac{1}{\sqrt{2}}|t_*| + o(1), \quad (5.19)$$

with respective eigenvectors

$$\begin{pmatrix} \Psi_1(\delta) \\ \Phi_1^{(1)}(\delta) \\ \Phi_1^{(2)}(\delta) \end{pmatrix} = \begin{pmatrix} \Psi_1 \\ \Phi_1^{(1)} \\ \Phi_1^{(2)} \end{pmatrix} + o(1), \quad \begin{pmatrix} \Psi_2(\delta) \\ \Phi_2^{(1)}(\delta) \\ \Phi_2^{(2)}(\delta) \end{pmatrix} = \begin{pmatrix} \Psi_2 \\ \Phi_2^{(1)} \\ \Phi_2^{(2)} \end{pmatrix} + o(1). \quad (5.20)$$

Finally, we check (5.13). By substituting $h_1(\delta)$ and $(\Psi_1(\delta), \Phi_1^{(1)}(\delta), \Phi_1^{(2)}(\delta))^T$ into (5.13), we obtain the following Π_2 -component

$$\frac{1}{2}\delta^{\frac{1}{6}}\Phi_1^{(2)}(\delta) + \Pi_2\mathbb{M}^\delta(\lambda_* + \delta \cdot h_1(\delta)) \begin{pmatrix} \Psi_1 \\ \Phi_1^{(1)} \\ \Phi_1^{(2)} \end{pmatrix}.$$

By using (5.14), (5.19) and (5.20), the Π_2 -component is further equal to

$$\mathbb{E}^\times(h_1)\Phi_1^{(1)} + o(1) = -i \cdot \operatorname{sgn}(t_*) \left(\frac{1}{2}\right)^{5/4} \left(\frac{|t_*|}{\gamma_*}\right)^{-1/2} \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix} + o(1) \neq 0.$$

Hence $h_1(\delta)$ and its eigenfunction satisfy (5.13). The proof of $h_2(\delta)$ is similar. This completes the proof of Theorem 1.8.

Appendix

Appendix A: Proof of Theorem 3.3, Proposition 3.4 and 3.5

Proof of Theorem 3.3

By Proposition 3.1, the Floquet-Bloch eigenpairs $(\mu_n(\kappa_1), v_n(\mathbf{x}; \kappa_1))$ ($n = 1, 2$) are analytic in $\mathcal{D} = \mathcal{D}_1 \cap \mathcal{D}_2$ which contains a closed rectangle $\mathcal{R}_\nu = \{z \in \mathbf{C} : 0 \leq \operatorname{Re} z \leq 2\pi, -\nu \leq \operatorname{Im} z \leq \nu\}$ for some $\nu > 0$. By Assumption 1.4, $\kappa_1 = \pi$ is the unique root of $\mu_n(\kappa_1) = \lambda_*$ ($n = 1, 2$) for $\kappa_1 \in [0, 2\pi]$. Thus, by taking ν to be sufficiently small, we can assume that $\kappa_1 = \pi$ is the unique root of $\mu_n(\kappa_1) = \lambda_*$ ($n = 1, 2$) for $\kappa_1 \in \mathcal{D}$.

We decompose the operator $\mathcal{G}(\lambda_* + i\epsilon)$ as

$$\mathcal{G}(\lambda_* + i\epsilon) = \mathcal{G}^{\epsilon, prop}(\lambda_* + i\epsilon) + \mathcal{G}^{\epsilon, evan}(\lambda_* + i\epsilon),$$

where

$$\mathcal{G}^{\epsilon, prop}(\lambda_* + i\epsilon)f := \frac{1}{2\pi} \sum_{n=1,2} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1,$$

$$\begin{aligned}
\mathcal{G}^{\epsilon, \text{evan}}(\lambda_* + i\epsilon)f &:= \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1 \\
&+ \frac{1}{2\pi} \sum_{n=1,2} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1.
\end{aligned} \tag{A.1}$$

Then Theorem 3.3 follows from the two lemmas below.

Lemma A.1 (Asymptotics of $\mathcal{G}^{\epsilon, \text{prop}}(\lambda_* + i\epsilon)$).

$$\begin{aligned}
&\frac{1}{2\pi} \int_{\pi - \epsilon^{\frac{1}{6}}}^{\pi + \epsilon^{\frac{1}{6}}} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} d\kappa_1 \\
&= (\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}})) \cdot \frac{1-i}{2\gamma_*^{\frac{1}{2}}} v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega \\
&+ \frac{1}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(2(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right. \\
&+ \left. (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi)(f(\cdot), v_1(\mathbf{y}; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_\Omega \right) + \mathcal{O}(\epsilon^{\frac{1}{3}}).
\end{aligned} \tag{A.2}$$

$$\begin{aligned}
&\frac{1}{2\pi} \int_{\pi - \epsilon^{\frac{1}{6}}}^{\pi + \epsilon^{\frac{1}{6}}} \frac{v_2(\mathbf{x}; \kappa_1)(f(\cdot), v_2(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_2(\kappa_1)} d\kappa_1 \\
&= -(\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}})) \cdot \frac{1+i}{2\gamma_*^{\frac{1}{2}}} v_2(\mathbf{x}; \pi)(f(\cdot), v_2(\cdot; \pi))_\Omega \\
&+ \frac{1}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(-2(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_2)(\cdot; \pi))_\Omega \right. \\
&- \left. (\partial_{\kappa_1}^2 v_2)(\mathbf{x}; \pi)(f(\cdot), v_2(\mathbf{y}; \pi))_\Omega - v_2(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_2)(\cdot; \pi))_\Omega \right) + \mathcal{O}(\epsilon^{\frac{1}{3}}).
\end{aligned} \tag{A.3}$$

Lemma A.2 (Asymptotics of $\mathcal{G}^{\epsilon, \text{evan}}(\lambda_* + i\epsilon)$).

$$\frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1 = \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 + \mathcal{O}(\epsilon). \tag{A.4}$$

$$\begin{aligned}
& \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} d\kappa_1 \\
&= (\epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} + \mathcal{O}(\epsilon^{\frac{1}{6}})) \cdot v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega \\
&\quad + \mathcal{G}_0^1(\lambda_*)f \\
&\quad - \frac{1}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(2(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right. \\
&\quad \left. + (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_\Omega \right) + \mathcal{O}(\epsilon^{\frac{1}{3}}),
\end{aligned} \tag{A.5}$$

where $\mathcal{G}_0^1(\lambda_*)$ is associated with the kernel function

$$G_0^1(\mathbf{x}, \mathbf{y}; \lambda_*) = \lim_{\epsilon \rightarrow 0^+} \left(\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1) \overline{v_1(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 - \epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} v_1(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} \right).$$

Similarly,

$$\begin{aligned}
& \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_2(\mathbf{x}; \kappa_1)(f(\cdot), v_2(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_2(\kappa_1)} d\kappa_1 \\
&= (-\epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} + \mathcal{O}(\epsilon^{\frac{1}{6}})) \cdot v_2(\mathbf{x}; \pi)(f(\cdot), v_2(\cdot; \pi))_\Omega \\
&\quad + \mathcal{G}_0^2(\lambda_*)f \\
&\quad + \frac{1}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(2(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_2)(\cdot; \pi))_\Omega \right. \\
&\quad \left. + (\partial_{\kappa_1}^2 v_2)(\mathbf{x}; \pi)(f(\cdot), v_2(\cdot; \pi))_\Omega + v_2(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_2)(\cdot; \pi))_\Omega \right) + \mathcal{O}(\epsilon^{\frac{1}{3}}),
\end{aligned} \tag{A.6}$$

where $\mathcal{G}_0^2(\lambda_*)$ is associated with the kernel function

$$G_0^2(\mathbf{x}, \mathbf{y}; \lambda_*) = \lim_{\epsilon \rightarrow 0^+} \left(\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_2(\mathbf{x}; \kappa_1) \overline{v_2(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 + \epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} v_2(\mathbf{x}; \pi) \overline{v_2(\mathbf{y}; \pi)} \right).$$

Proof of Lemma A.1. Here we only present the proof of (A.2). The one of (A.3) is similar. The analyticity of $v_1(\mathbf{x}; \kappa_1)$ and $\mu_1(\kappa_1)$ implies that

$$\begin{aligned}
v_1(\mathbf{x}; \kappa_1) &= v_1(\mathbf{x}; \pi) + (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi) + \frac{1}{2}(\kappa_1 - \pi)^2 \cdot (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi) + \mathcal{O}(|\kappa_1 - \pi|^3), \\
\mu_1(\kappa_1) &= \lambda_* - \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^4).
\end{aligned} \tag{A.7}$$

It follows that

$$\frac{1}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} = \frac{1 + \mathcal{O}(|\kappa_1 - \pi|^2)}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2}. \tag{A.8}$$

Using (A.7)-(A.8), we have

$$\begin{aligned}
& \frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} d\kappa_1 \\
&= \left(\frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{1 + \mathcal{O}(|\kappa_1 - \pi|^2)}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 \right) \cdot v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega \\
&+ \left(\frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{\kappa_1 - \pi + \mathcal{O}(|\kappa_1 - \pi|^3)}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 \right) \cdot \left(\partial_{\kappa_1} v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right) \\
&+ \left(\frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^4)}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 \right) \cdot \left((\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right. \\
&\quad \left. + \frac{1}{2}(\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi)(f(\cdot), v_1(\mathbf{y}; \pi))_\Omega + \frac{1}{2}v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_\Omega \right) \\
&+ \mathcal{O}\left(\int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{|\kappa_1 - \pi|^3}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 \right)
\end{aligned} \tag{A.9}$$

Note that $\int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 = 0$ and

$$\begin{aligned}
& \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \left| \frac{(\kappa_1 - \pi)^2}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} \right| d\kappa_1 \lesssim \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} 1 \cdot d\kappa_1 = \mathcal{O}(\epsilon^{\frac{1}{6}}), \\
& \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \left| \frac{(\kappa_1 - \pi)^3}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} \right| d\kappa_1 \lesssim \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} |\kappa_1 - \pi| d\kappa_1 = \mathcal{O}(\epsilon^{\frac{1}{3}}).
\end{aligned}$$

One further obtain

$$\begin{aligned}
& \frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} d\kappa_1 \\
&= \left(\frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{1}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 + \mathcal{O}(\epsilon^{\frac{1}{6}}) \right) \cdot v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega \\
&+ \left(\frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 \right) \cdot \left(2(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right. \\
&\quad \left. + (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi)(f(\cdot), v_1(\mathbf{y}; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_\Omega \right) \\
&+ \mathcal{O}(\epsilon^{\frac{1}{3}})
\end{aligned} \tag{A.10}$$

By direct calculation

$$\begin{aligned} \frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{1}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 &= \frac{1-i}{2\gamma_*^{\frac{1}{2}}} \epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}}), \\ \frac{1}{2\pi} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2} d\kappa_1 &= \frac{1}{\pi\gamma_*} \int_{\pi-\epsilon^{\frac{1}{6}}}^{\pi+\epsilon^{\frac{1}{6}}} \left(1 - \frac{i\epsilon}{i\epsilon + \frac{1}{2}\gamma_*(\kappa_1 - \pi)^2}\right) d\kappa_1 = \frac{2}{\pi\gamma_*} \epsilon^{\frac{1}{6}} + \mathcal{O}(\epsilon^{\frac{1}{2}}). \end{aligned} \quad (\text{A.11})$$

Thus, (A.2) follows by substituting (A.11) into (A.10). \square

Proof of Lemma A.2. Step 1. We first prove (A.4). Define

$$\mathbb{P}_n(\kappa_1)g = (g(\cdot), v_n(\cdot; \overline{\kappa_1}))v_n(\cdot; \kappa_1) \quad (n = 1, 2), \quad \mathbb{Q}(\kappa_1) = I - \mathbb{P}_1(\kappa_1) - \mathbb{P}_2(\kappa_1). \quad (\text{A.12})$$

Then, by Floquet transform,

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1 &= \frac{1}{2\pi} \int_0^{2\pi} (\lambda_* + i\epsilon - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))^{-1} \mathbb{Q}(\kappa_1) \hat{f}(\kappa_1) d\kappa_1, \\ \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 &= \frac{1}{2\pi} \int_0^{2\pi} (\lambda_* - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))^{-1} \mathbb{Q}(\kappa_1) \hat{f}(\kappa_1) d\kappa_1, \end{aligned} \quad (\text{A.13})$$

where $\hat{f}(\kappa_1)$ denotes the Floquet transform of f . Notice that

$$\sigma(\mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1)) = \{0\} \cup \{\mu_n(\kappa_1)\}_{n \geq 3}.$$

Consequently, $\lambda_* \notin \sigma(\mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))$ for any $\kappa_1 \in [0, 2\pi)$ by Assumption 1.4(2). Thus, for ϵ being sufficiently small, the resolvent $(\lambda_* + i\epsilon - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))^{-1} : (H^1(Y))^* \rightarrow H^1(Y)$ is uniformly bounded. Hence (A.13) and the resolvent identity shows that

$$\begin{aligned} &\frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_n(\kappa_1)} d\kappa_1 - \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\cdot), v_n(\cdot; \kappa_1))_\Omega}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 \\ &= i\epsilon \cdot \frac{1}{2\pi} \int_0^{2\pi} \left[(\lambda_* + i\epsilon - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))^{-1} \cdot (\lambda_* - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1))^{-1} \right] \mathbb{Q}(\kappa_1) \hat{f}(\kappa_1) d\kappa_1 \\ &= \mathcal{O}(\epsilon), \end{aligned}$$

which concludes the proof of (A.4).

Step 2. We prove (A.5). We point out that the proof of (A.6) is similar. First, by Assumption 1.4,

$$|\lambda_* - \mu_1(\kappa_1)| \gtrsim \epsilon^{\frac{1}{3}} \quad \text{for } |\kappa_1 - \pi| > \epsilon^{\frac{1}{6}}.$$

Thus

$$\begin{aligned} &\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* + i\epsilon - \mu_1(\kappa_1)} d\kappa_1 \\ &= \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{-i\epsilon}{(\lambda_* + i\epsilon - \mu_1(\kappa_1))(\lambda_* - \mu_1(\kappa_1))} v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega d\kappa_1 \\ &= I + \mathcal{O}(\epsilon^{\frac{1}{3}}). \end{aligned} \quad (\text{A.14})$$




Figure 3: Integral contour used in the proof.

where

$$I = \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1)(f(\cdot), v_1(\cdot; \kappa_1))_\Omega}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 = \frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{\mathbb{P}_1(\kappa_1) \hat{f}(\kappa_1)}{\lambda - \mu_1(\kappa_1)} d\kappa_1.$$

Note that the projection $\mathbb{P}_1(\kappa_1)$ and $\mu_1(\kappa_1)$ are analytic in $\kappa_1 \in \mathcal{D} \supset \mathcal{R}_\nu = \{z \in \mathbf{C} : 0 \leq \operatorname{Re} z \leq 2\pi, -\nu \leq \operatorname{Im} z \leq \nu\}$. The Cauchy theorem gives

$$\begin{aligned} I &= \frac{1}{2\pi} \int_{C_\epsilon} \frac{\mathbb{P}_1(\kappa_1) \hat{f}(\kappa_1)}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 \\ &+ \frac{1}{2\pi} \int_0^\nu \frac{\mathbb{P}_1(it) \hat{f}(it)}{\lambda_* - \mu_1(it)} idt - \frac{1}{2\pi} \int_0^\nu \frac{\mathbb{P}_1(2\pi + it) \hat{f}(2\pi + it)}{\lambda_* - \mu_1(2\pi + it)} idt \\ &+ \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathbb{P}_1(\kappa_1 + i\nu) \hat{f}(\kappa_1 + i\nu)}{\lambda_* - \mu_1(\kappa_1 + i\nu)} d\kappa_1, \end{aligned} \quad (\text{A.15})$$

where $C_\epsilon := \{\pi + \epsilon^{\frac{1}{6}} e^{i\theta} : 0 \leq \theta \leq \pi\}$. The shifted contour is illustrated in Figure 3(a). Note that ν is independent of ϵ . By the Taylor expansion (A.7), we obtain (similar to (A.9))

$$\begin{aligned} &\frac{1}{2\pi} \int_{C_\epsilon} \frac{\mathbb{P}_1(\kappa_1) \hat{f}(\kappa_1)}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 \\ &= (\epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} + \mathcal{O}(\epsilon^{\frac{1}{6}})) \cdot v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega \\ &+ \frac{i}{\gamma_*} \left(\partial_{\kappa_1} v_1(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right) \\ &- \frac{1}{\pi\gamma_*} \epsilon^{\frac{1}{6}} \left(2(\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1} v_1)(\cdot; \pi))_\Omega \right. \\ &\left. + (\partial_{\kappa_1}^2 v_1)(\mathbf{x}; \pi)(f(\cdot), v_1(\cdot; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\cdot), (\partial_{\kappa_1}^2 v_1)(\cdot; \pi))_\Omega \right) + \mathcal{O}(\epsilon^{\frac{1}{3}}) \end{aligned} \quad (\text{A.16})$$

On the other hand, by calculating the asymptotics of $\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1) \overline{v_1(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1$ using a similar contour integral approach, we obtain

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0^+} \left(\frac{1}{2\pi} \int_{[0, \pi - \epsilon^{\frac{1}{6}}] \cup (\pi + \epsilon^{\frac{1}{6}}, 2\pi]} \frac{v_1(\mathbf{x}; \kappa_1) \overline{v_1(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_1(\kappa_1)} d\kappa_1 - \epsilon^{-\frac{1}{6}} \cdot \frac{2}{\pi \gamma_*} v_1(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} \right) \\ &= \frac{1}{2\pi} \int_0^\nu \frac{\mathbb{P}_1(it) \hat{f}(it)}{\lambda_* - \mu_1(it)} idt - \frac{1}{2\pi} \int_0^\nu \frac{\mathbb{P}_1(2\pi + it) \hat{f}(2\pi + it)}{\lambda_* - \mu_1(2\pi + it)} idt \\ & \quad + \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathbb{P}_1(\kappa_1 + i\nu) \hat{f}(\kappa_1 + i\nu)}{\lambda_* - \mu_1(\kappa_1 + i\nu)} d\kappa_1. \end{aligned} \quad (\text{A.17})$$

Combining (A.14)-(A.17), the proof of (A.5) is complete. \square

Proof of Proposition 3.4

Step 1: We first prove $(\text{div}(A\nabla) + \lambda_*)G_0(\mathbf{x}, \mathbf{y}; \lambda_*) = \delta(\mathbf{x} - \mathbf{y})$. It's equivalent to proving the following weak formulation

$$\mathbf{a}_\Omega^A(\mathcal{G}_0(\lambda_*)f, g) - \lambda_* \cdot (\mathcal{G}_0(\lambda_*)f, g)_\Omega = (f, g)_\Omega, \quad \text{for any } f, g \in C_c^\infty(\Omega) \quad (\text{A.18})$$

where the form $\mathbf{a}_\Omega^A(\cdot, \cdot)$ on $H^1(\Omega)$ is defined similarly as (1.2). From (3.5), we see

$$\begin{aligned} & \mathbf{a}_\Omega^A(\mathcal{G}_0(\lambda_*)f, g) - \lambda_* \cdot (\mathcal{G}_0(\lambda_*)f, g)_\Omega \\ &= \frac{(1-i) \cdot (\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}}))}{2\gamma_*^{\frac{1}{2}}} (f(\cdot), v_1(\cdot; \pi)) \cdot \left(\mathbf{a}^A(v_1(\cdot; \pi), g(\cdot)) - \lambda_* \cdot (v_1(\cdot; \pi), g(\cdot))_\Omega \right) \\ & \quad - \frac{(1+i) \cdot (\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}}))}{2\gamma_*^{\frac{1}{2}}} (f(\cdot), v_2(\cdot; \pi)) \cdot \left(\mathbf{a}^A(v_2(\cdot; \pi), g(\cdot)) - \lambda_* \cdot (v_2(\cdot; \pi), g(\cdot))_\Omega \right) \\ & \quad + \mathbf{a}_\Omega^A(\mathcal{G}(\lambda_* + i\epsilon)f, g) - \lambda_* \cdot (\mathcal{G}(\lambda_* + i\epsilon)f, g)_\Omega + \mathcal{O}(\epsilon^{\frac{1}{6}}). \end{aligned} \quad (\text{A.19})$$

In addition, the weak formulation of $(\text{div}(A\nabla) + \lambda_*)v_n(\mathbf{x}; \pi) = 0$ ($n = 1, 2$) and $(\text{div}(A\nabla) + \lambda_* + i\epsilon)G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon) = \delta(\mathbf{x} - \mathbf{y})$ yield

$$\mathbf{a}_\Omega^A(v_n(\cdot; \pi), g(\cdot)) - \lambda_* \cdot (v_n(\cdot; \pi), g(\cdot))_\Omega = 0, \quad n = 1, 2 \quad (\text{A.20})$$

and

$$\mathbf{a}_\Omega^A(\mathcal{G}(\lambda_* + i\epsilon)f, g) - (\lambda_* + i\epsilon) \cdot (\mathcal{G}(\lambda_* + i\epsilon)f, g)_\Omega = (f, g)_\Omega, \quad \text{for any } f, g \in C_c^\infty(\Omega). \quad (\text{A.21})$$

Substituting (A.20) and (A.21) into (A.19), we obtain that

$$\mathbf{a}_\Omega^A(\mathcal{G}_0(\lambda_*)f, g) - \lambda_* \cdot (\mathcal{G}_0(\lambda_*)f, g)_\Omega = (f, g)_\Omega + i\epsilon \cdot (\mathcal{G}(\lambda_* + i\epsilon)f, g)_\Omega + \mathcal{O}(\epsilon^{\frac{1}{6}}). \quad (\text{A.22})$$

Note that by (3.5),

$$\epsilon \cdot (\mathcal{G}(\lambda_* + i\epsilon)f, g)_\Omega = \mathcal{O}(\epsilon^{\frac{1}{2}}).$$

Hence, by letting $\epsilon \rightarrow 0^+$ in (A.22), we obtain (A.18). The boundary condition of $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ as stated in (3.7) can be checked directly.

Step 2: We prove (3.8). By the reflection symmetry of the operator \mathcal{L}^A , i.e. $[\mathcal{L}^A, \mathcal{M}_1] = 0$, the Bloch eigenvalues and eigenfunctions of \mathcal{L}^A have the following properties:

$$\mu_n(\kappa_1) = \mu_n(2\pi - \kappa_1), \quad v_n(\mathbf{x}; \kappa_1) \sim v_n(\mathcal{M}_1\mathbf{x}; 2\pi - \kappa_1), \quad n = 1, 2,$$

and for each $n \geq 3$, there exists $n' \geq 3$ such that

$$\mu_n(\kappa_1) = \mu_{n'}(2\pi - \kappa_1), \quad v_n(\mathbf{x}; \kappa_1) \sim v_{n'}(\mathcal{M}_1\mathbf{x}; 2\pi - \kappa_1).$$

Therefore, the formula (3.6) is invariant under the substitution $\mathbf{x} \rightarrow \mathcal{M}_1\mathbf{x}$, $\mathbf{y} \rightarrow \mathcal{M}_1\mathbf{y}$, which implies $G_0(\mathbf{x}, \mathbf{y}; \lambda_*) = G_0(\mathcal{M}_1\mathbf{x}, \mathcal{M}_1\mathbf{y}; \lambda_*)$.

We point out that (3.9) follows from a similar argument by using the time-reversal symmetry of the operator \mathcal{L}^A .

Step 3: We prove (3.10). We note that $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ has the following contour integral expression, which is obtained from the proof of Lemma A.1 and A.2,

$$\begin{aligned} G_0(\mathbf{x}, \mathbf{y}; \lambda_*) &= \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1) \overline{v_n(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 + G_0^1(\mathbf{x}, \mathbf{y}; \lambda_*) + G_0^2(\mathbf{x}, \mathbf{y}; \lambda_*) \\ &= G_0^+(\mathbf{x}, \mathbf{y}; \lambda_*) + \frac{i}{\gamma_*} \left(\partial_{\kappa_1} v_1(\mathbf{x}; \pi) \overline{v_1(\mathbf{y}; \pi)} + v_1(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_1)(\mathbf{y}; \pi)} \right) \\ &\quad - \frac{i}{\gamma_*} \left(\partial_{\kappa_1} v_2(\mathbf{x}; \pi) \overline{v_2(\mathbf{y}; \pi)} + v_2(\mathbf{x}; \pi) \overline{(\partial_{\kappa_1} v_2)(\mathbf{y}; \pi)} \right), \end{aligned} \quad (\text{A.23})$$

where

$$\begin{aligned} G_0^+(\mathbf{x}, \mathbf{y}; \lambda_*) &:= \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1) \overline{v_n(\mathbf{y}; \kappa_1)}}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_0^\nu \sum_{n=1,2} \frac{v_n(\mathbf{x}; it) \overline{v_n(\mathbf{y}; it)}}{\lambda_* - \mu_n(it)} idt - \frac{1}{2\pi} \int_0^\nu \sum_{n=1,2} \frac{v_n(\mathbf{x}; 2\pi + it) \overline{v_n(\mathbf{y}; 2\pi + it)}}{\lambda_* - \mu_n(2\pi + it)} idt \\ &\quad + \frac{1}{2\pi} \int_0^{2\pi} \sum_{n=1,2} \frac{v_n(\mathbf{x}; \kappa_1 + i\nu) \overline{v_n(\mathbf{y}; \kappa_1 + i\nu)}}{\lambda_* - \mu_n(\kappa_1 + i\nu)} d\kappa_1. \end{aligned}$$

Since $G_0^+(\mathbf{x}, \mathbf{y}; \lambda_*)$ decays exponentially as $x_1 \rightarrow \infty$ (see Theorem 7 in [21]), (3.10) follows from (A.23). (3.11) can be proved similarly by replacing the integral contour in (A.23) with the one in Figure 3(b). The details are omitted here.

Proof of Proposition 3.5

Here we only prove (3.13) and (3.15). The proof of (3.14) and (3.16) is similar. The strategy is to show first that the jump formula (3.13) and (3.15) holds for the Green function $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$, and then obtain the jump formula for $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ by letting $\epsilon \rightarrow 0^+$.

To show the jump formula (3.13) and (3.15) holds $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$, we first fix a neighborhood of Γ , say, $U := (-\frac{1-2r_0}{4}, \frac{1-2r_0}{4}) \times (-\frac{1}{2}, \frac{1}{2})$, where $r_0 := \text{diam}(V)$. Let

$$G^{\text{empty}}(\mathbf{x}, \mathbf{y}; \lambda) = \frac{1}{2\pi} \int_0^{2\pi} \sum_{\mathbf{n} \in \mathbf{Z}^2} \frac{e^{i(2\pi\mathbf{n} + (\kappa_1, \pi)) \cdot (\mathbf{x} - \mathbf{y})}}{\lambda - |2\pi\mathbf{n} + (\kappa_1, \pi)|} d\kappa_1$$

be the quasi-periodic Green function for the Laplacian operator in Ω . Since $U \cap (\cup_{n_1, n_2 \in \mathbf{Z}} V_{n_1, n_2} = \emptyset$, $\mathcal{L}^A = -\Delta$ in U . Hence

$$\mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x}) - \mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi](\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in U \quad (\text{A.24})$$

for some function $f \in H^1(U)$. From Section 2 of [11], the following jump relations hold for $\mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi](\mathbf{x})$

$$\lim_{t \rightarrow 0} \left(\mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi](\mathbf{x}) \quad (\text{A.25})$$

$$\lim_{t \rightarrow 0^\pm} \frac{\partial}{\partial x_1} \left(\mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \pm \frac{1}{2} \varphi(\mathbf{x}) + \frac{1}{2} \mathcal{K}^{empty,*}[\varphi](\mathbf{x}), \quad (\text{A.26})$$

where

$$\mathcal{K}^{empty,*}(\lambda) : \tilde{H}^{-\frac{1}{2}}(\Gamma) \rightarrow \tilde{H}^{-\frac{1}{2}}(\Gamma), \quad \varphi \mapsto p.v. \int_{\Gamma} \frac{\partial G^{empty}}{\partial n_{\mathbf{x}}}(\mathbf{x}, \mathbf{y}; \lambda) \varphi(\mathbf{y}) ds(\mathbf{y}).$$

Thus, by (A.24) and (A.25), we have

$$\begin{aligned} \lim_{t \rightarrow 0^\pm} \mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x} + t\mathbf{e}_1) &= \lim_{t \rightarrow 0^\pm} \mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi](\mathbf{x} + t\mathbf{e}_1) + f(\mathbf{x}) \\ &= \mathcal{S}(\lambda_* + i\epsilon; G^{empty})[\varphi](\mathbf{x}) + f(\mathbf{x}) \\ &= \mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x}). \end{aligned} \quad (\text{A.27})$$

Thus (3.13) holds for $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$. On the other hand, the reflection symmetry implies the following identities (analogous to (3.8))

$$G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon) = G(\mathcal{M}_1 \mathbf{x}, \mathcal{M}_1 \mathbf{y}; \lambda_* + i\epsilon), \quad G^{empty}(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon) = G^{empty}(\mathcal{M}_1 \mathbf{x}, \mathcal{M}_1 \mathbf{y}; \lambda_* + i\epsilon). \quad (\text{A.28})$$

Hence we have $f(\mathbf{x}) = f(\mathcal{M}_1 \mathbf{x})$ by (A.24) and $\mathcal{K}^{empty,*}(\lambda_* + i\epsilon) = 0$ by the definition of $\mathcal{K}^{empty,*}$. Thus (A.24) and (A.26) yield

$$\lim_{t \rightarrow 0^\pm} \frac{\partial}{\partial x_1} \left(\mathcal{S}(\lambda_* + i\epsilon; G)[\varphi] \right) (\mathbf{x} + t\mathbf{e}_1) = \pm \frac{1}{2} \varphi(\mathbf{x}) + \frac{\partial f}{\partial x_1}(\mathbf{x}) = \pm \frac{1}{2} \varphi(\mathbf{x}), \quad (\text{A.29})$$

where we used the fact that $\frac{\partial f}{\partial x_1}|_{\Gamma} = 0$ (since $f(\mathbf{x}) = f(\mathcal{M}_1 \mathbf{x})$). Hence (3.15) holds for $G(\mathbf{x}, \mathbf{y}; \lambda_* + i\epsilon)$.

We now prove (3.13) and (3.15) by using (A.27) and (A.29). Note that (3.5) implies

$$\mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x}) - \mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x}) = f_\epsilon(\mathbf{x}) \in H^1(U), \quad (\text{A.30})$$

with

$$\begin{aligned} f_\epsilon(\mathbf{x}) &= -(\epsilon^{-\frac{1}{2}} + o(\epsilon^{-\frac{1}{2}})) \cdot \left(\frac{(1-i) \cdot \int_{\Gamma} \varphi(\cdot) \overline{v_1(\cdot; \pi)}}{2\gamma_*^{\frac{1}{2}}} v_1(\mathbf{x}; \pi) - \frac{(1+i) \cdot \int_{\Gamma} \varphi(\cdot) \overline{v_2(\cdot; \pi)}}{2\gamma_*^{\frac{1}{2}}} v_2(\mathbf{x}; \pi) \right) \\ &\quad + \mathcal{O}(\epsilon^{\frac{1}{6}}). \end{aligned} \quad (\text{A.31})$$

Then, by (A.27),

$$\begin{aligned}\lim_{t \rightarrow 0^\pm} \mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x} + t\mathbf{e}_1) &= \lim_{t \rightarrow 0^\pm} \left(\mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x} + t\mathbf{e}_1) + f_\epsilon(\mathbf{x} + t\mathbf{e}_1) \right) \\ &= \mathcal{S}(\lambda_* + i\epsilon; G)[\varphi](\mathbf{x}) + f_\epsilon(\mathbf{x}) \\ &= \mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x}),\end{aligned}$$

which gives (3.13). For (3.15), note that the parity of $v_n(\mathbf{x}; \pi)$ ($n = 1, 2$) in (3.27) implies that

$$v_1(\mathbf{x}; \pi) = 0, \quad \frac{\partial v_2}{\partial x_2}(\mathbf{x}; \pi) = 0, \quad \mathbf{x} \in \Gamma.$$

Thus (A.31) indicates that

$$\frac{\partial f_\epsilon}{\partial x_1}(\mathbf{x}) = \mathcal{O}(\epsilon^{\frac{1}{6}}), \quad \mathbf{x} \in \Gamma.$$

By (A.29) and (A.30), we have

$$\lim_{t \rightarrow 0^\pm} \frac{\partial}{\partial x_1} \left(\mathcal{S}(\lambda_*; G_0)[\varphi] \right)(\mathbf{x} + t\mathbf{e}_1) = \pm \frac{1}{2} \varphi(\mathbf{x}) + \frac{\partial f_\epsilon}{\partial x_1}(\mathbf{x}) = \pm \frac{1}{2} \varphi(\mathbf{x}) + \mathcal{O}(\epsilon^{\frac{1}{6}}).$$

Hence, by letting $\epsilon \rightarrow 0^+$, we conclude the proof of (3.15).

Appendix B: Proof of Lemma 4.3, 4.4, 4.5 and 4.6

Proof of Lemma 4.3

By the relation $\tilde{v}_n(\mathbf{x}) = e^{i\pi x_1} v_n(\mathbf{x})$ and (4.6), it suffices to prove the following identities

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_2(\mathbf{x})} d\mathbf{x} = \int_Y (B\nabla v_2(\mathbf{x})) \cdot \overline{\nabla v_1(\mathbf{x})} d\mathbf{x}, \quad (\text{B.1})$$

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_2(\mathbf{x})} d\mathbf{x} \in \mathbf{R}, \quad (\text{B.2})$$

$$\int_Y (B\nabla v_n(\mathbf{x})) \cdot \overline{\nabla v_n(\mathbf{x})} d\mathbf{x} = 0, \quad n = 1, 2. \quad (\text{B.3})$$

Step 1: We first prove (B.1). Note that (1.9) in Assumption 1.4 and the relationship $v_n(\mathbf{x}) := u_n(\mathbf{x}; \boldsymbol{\kappa}^{(2)})$ give that

$$\mathcal{R}v_1 = iv_2, \quad \mathcal{R}v_2 = iv_1. \quad (\text{B.4})$$

Thus,

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_2(\mathbf{x})} d\mathbf{x} = \int_Y (B\nabla(v_2(R\mathbf{x}))) \cdot \overline{\nabla(v_1(R\mathbf{x}))} d\mathbf{x}. \quad (\text{B.5})$$

For any functions $f(\mathbf{x})$, it holds that

$$\nabla(f(R\mathbf{x})) = R^T(\nabla f)(R\mathbf{x}). \quad (\text{B.6})$$

Hence, (B.5) gives that

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_2(\mathbf{x})} d\mathbf{x} = \int_Y (BR^T(\nabla v_2)(R\mathbf{x})) \cdot \overline{R^T(\nabla v_1)(R\mathbf{x})} d\mathbf{x}$$

By Assumption 1.5, B commutes with R^T . Thus

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_2(\mathbf{x})} d\mathbf{x} = \int_Y B(\mathbf{x})(\nabla v_2)(R\mathbf{x}) \cdot \overline{(\nabla v_1)(R\mathbf{x})} d\mathbf{x}$$

Then (B.1) follows by a change of variable $R\mathbf{x} \rightarrow \mathbf{x}$ and the identity $B(\mathbf{x}) = B(R\mathbf{x})$.

Step 2: (B.2) follows from (B.1) and the fact that B is Hermitian.

Step 3: We prove (B.3) for $n = 1$. We point out that $n = 2$ can be treated similarly. The proof exploits the reflection asymmetry of the system, namely $BM_2 = -M_2B$.

Note that (1.9) in Assumption 1.4 and the relationship $v_n(\mathbf{x}) := u_n(\mathbf{x}; \boldsymbol{\kappa}^{(2)})$ gives that $\mathcal{M}_2 v_1 = v_1$. Thus,

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_1(\mathbf{x})} d\mathbf{x} = \int_Y (B\nabla(v_1(M_2\mathbf{x}))) \cdot \overline{\nabla(v_1(M_2\mathbf{x}))} d\mathbf{x}. \quad (\text{B.7})$$

For any functions $f(\mathbf{x})$, it holds that

$$\nabla(f(M_2\mathbf{x})) = M_2(\nabla f)(M_2\mathbf{x}). \quad (\text{B.8})$$

Hence, (B.7) gives that

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_1(\mathbf{x})} d\mathbf{x} = \int_Y (BM_2(\nabla v_1)(M_2\mathbf{x})) \cdot \overline{M_2(\nabla v_1)(M_2\mathbf{x})} d\mathbf{x}.$$

Recall that by Assumption 1.5, B anti-commutes with M_2 , i.e. $BM_2 = -M_2B$. Thus,

$$\int_Y (B\nabla v_1(\mathbf{x})) \cdot \overline{\nabla v_1(\mathbf{x})} d\mathbf{x} = - \int_Y B(\mathbf{x})(\nabla v_1)(M_2\mathbf{x}) \cdot \overline{\nabla v_1(M_2\mathbf{x})} d\mathbf{x}.$$

Then (B.3) for $n = 1$ follows by a change of variable $M_2\mathbf{x} \rightarrow \mathbf{x}$ and the identity $B(\mathbf{x}) = B(M_2\mathbf{x})$.

Proof of Lemma 4.4

It's a direct consequence of Theorem 1.3. As shown in Section 2.2, the first-order terms vanish when we apply the perturbation argument near $(\kappa_1, \kappa_2) = (\pi, \pi)$. Here the first-order terms are given by $(\tilde{\mathcal{L}}_1^A \tilde{v}_m, \tilde{v}_n)$ ($n, m = 1, 2$).

Proof of Lemma 4.5

Step 1: We show that

$$[\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A, \mathcal{M}_2] = 0. \quad (\text{B.9})$$

Note that $[\mathcal{L}_{\Omega, \pi}^A(\kappa_1), \mathcal{M}_2] = 0$ by (1.6) and (1.4). Thus $[\tilde{\mathcal{L}}_{\Omega, \pi}^A(\kappa_1), \mathcal{M}_2] = 0$ since $[e^{\pm i\kappa_1 x_1}, \mathcal{M}_2] = 0$. Consequently, the expansion (4.7) yields

$$[\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi), \mathcal{M}_2] = 0, \quad [\tilde{\mathcal{L}}_1^A, \mathcal{M}_2] = 0, \quad [\tilde{\mathcal{L}}_{11}^A, \mathcal{M}_2] = 0. \quad (\text{B.10})$$

On the other hand, the equality $\tilde{v}_n(\mathbf{x}) = e^{-i\pi x_1} u_n(\mathbf{x}; \boldsymbol{\kappa}^{(2)})$ and the parity (1.9) in Assumption 1.4 give that

$$\mathcal{M}_2 \tilde{v}_1 = \tilde{v}_1, \quad \mathcal{M}_2 \tilde{v}_2 = -\tilde{v}_2. \quad (\text{B.11})$$

Recall $Q_\perp f = f - \sum_{n=1,2} (f, \tilde{v}_n) \tilde{v}_n$. Hence $[Q_\perp, \mathcal{M}_2] = 0$. Consequently, by (B.10),

$$[Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp, \mathcal{M}_2] = 0. \quad (\text{B.12})$$

Then (B.9) follows from (B.10) and (B.12).

Step 2: We prove that

$$\left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \tilde{v}_1 \right) = 0. \quad (\text{B.13})$$

Indeed,

$$\begin{aligned} & \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \tilde{v}_1 \right) \\ &= - \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \mathcal{M}_2 \tilde{v}_2, \mathcal{M}_2 \tilde{v}_1 \right) \\ &= - \left(\mathcal{M}_2 (\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \mathcal{M}_2 \tilde{v}_1 \right) \\ &= - \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \tilde{v}_1 \right), \end{aligned}$$

where we applied (B.9) in the second equality above. Then (B.13) follows.

The proof of $\left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_1, \tilde{v}_2 \right) = 0$ is similar.

Step 3: We prove

$$\begin{aligned} \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_1, \tilde{v}_1 \right) &= -\frac{1}{2} \gamma_*, \\ \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \tilde{v}_2 \right) &= \frac{1}{2} \gamma_*. \end{aligned} \quad (\text{B.14})$$

Let $\alpha_1 := \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_1, \tilde{v}_1 \right)$ and $\alpha_2 := \left((\tilde{\mathcal{L}}_{11}^A - \tilde{\mathcal{L}}_1^A Q_\perp (\tilde{\mathcal{L}}_{\Omega, \pi}^A(\pi) - \lambda_*)^{-1} Q_\perp \tilde{\mathcal{L}}_1^A) \tilde{v}_2, \tilde{v}_2 \right)$. Then, by applying the perturbation argument in Section 4 with $\delta = 0$, we see (4.16) becomes

$$\left(\begin{pmatrix} \mu^{(1)} - \alpha_1 p^2 & 0 \\ 0 & \mu^{(1)} - \alpha_2 p^2 \end{pmatrix} + \mathcal{O}(|p^3|) \right) \begin{pmatrix} a \\ b \end{pmatrix} = 0, \quad (\text{B.15})$$

where we've used Lemma 4.4 and (B.13) to simplify the form of $\mathcal{M}^{(0)}(\mu^{(1)}, p, 0)$. Based on (B.15), we can follow the calculation in Section 4 to obtain two branches of Bloch eigenpairs

$$\begin{aligned} \mu_1(\kappa_1) &= \lambda_* + \alpha_1 (\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^3), & v_1(\mathbf{x}; \kappa_1) &= v_1(\mathbf{x}) + \mathcal{O}(|\kappa_1 - \pi|), \\ \mu_2(\kappa_1) &= \lambda_* + \alpha_2 (\kappa_1 - \pi)^2 + \mathcal{O}(|\kappa_1 - \pi|^3), & v_2(\mathbf{x}; \kappa_1) &= v_2(\mathbf{x}) + \mathcal{O}(|\kappa_1 - \pi|). \end{aligned}$$

In accordance with (3.2), it holds that $\alpha_1 = -\frac{1}{2} \gamma_*$ and $\alpha_2 = \frac{1}{2} \gamma_*$, which concludes the proof of (B.14).

Proof of Lemma 4.6

By (3.27), $ix_1 v_n(\mathbf{x}) \overline{v_n(\mathbf{x})}$ is an odd function in x_1 for $n = 1, 2$. Hence, $(ix_1 v_n(\mathbf{x}), v_n(\mathbf{x})) = 0$.

Appendix C: Proof of Proposition 5.1

We only prove (5.14). The proof of (5.15) is similar. To check the convergence of the interface integral operators $\mathcal{S}(\lambda; G^\delta), \mathcal{K}(\lambda; G^\delta), \mathcal{N}(\lambda; G^\delta), \mathcal{K}^*(\lambda; G^\delta)$, we consider the associated volume integral operators and apply the following result.

Lemma C.1. *Let $K_n(\mathbf{x}, \mathbf{y})$ ($n \geq 1$) and $K(x, y)$ be integral kernels such that $K_n(\mathbf{x}, \mathbf{y}) = \overline{K_n(\mathbf{y}, \mathbf{x})}$ and the layer potential operators $\mathcal{S}(K_n), \mathcal{D}(K_n)$ and $\mathcal{S}(K), \mathcal{D}(K)$ (defined similarly as in (5.1)) satisfy the jump conditions (5.2)-(5.5). Let $U \subset \Omega$ be a neighborhood of Γ . Suppose that the following holds uniformly for all $f \in (H^1(U))^*$ with $\|f\|_{(H^1(U))^*} = 1$ and $\text{supp} f \subset\subset U$*

$$\int_U K_n(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y} \xrightarrow{H^1(U)} \int_U K(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y}. \quad (\text{C.1})$$

Then the integral operators $\mathcal{S}(K_n) \in \mathcal{B}(\tilde{H}^{-\frac{1}{2}}(\Gamma), H^{\frac{1}{2}}(\Gamma))$, $\mathcal{N}(K_n) \in \mathcal{B}(H^{\frac{1}{2}}(\Gamma), \tilde{H}^{-\frac{1}{2}}(\Gamma))$, $\mathcal{K}(K_n) \in \mathcal{B}(H^{\frac{1}{2}}(\Gamma), H^{\frac{1}{2}}(\Gamma))$ and $\mathcal{K}^*(K_n) \in \mathcal{B}(\tilde{H}^{-\frac{1}{2}}(\Gamma), \tilde{H}^{-\frac{1}{2}}(\Gamma))$ converge to $\mathcal{S}(K), \mathcal{N}(K), \mathcal{K}(K)$ and $\mathcal{K}^*(K)$ in operator norm, respectively.

Proof. Here we prove $\mathcal{S}(K_n) \rightarrow \mathcal{S}(K)$, $\mathcal{K}^*(K_n) \rightarrow \mathcal{K}^*(K)$ and $\mathcal{K}(K_n) \rightarrow \mathcal{K}(K)$. One can prove $\mathcal{N}(K_n) \rightarrow \mathcal{N}(K)$ by a similar argument.

We first prove $\mathcal{S}(K_n) \rightarrow \mathcal{S}(K)$. For each $\varphi \in \tilde{H}^{-\frac{1}{2}}(\Gamma)$, (C.1) implies that

$$\text{Tr} \int_U K_n(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y} \xrightarrow{H^{\frac{1}{2}}(\Gamma)} \text{Tr} \int_U K(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y}, \quad (\text{C.2})$$

where $\text{Tr} : H^1(U) \rightarrow H^{\frac{1}{2}}(\Gamma)$ is the trace operator, and Tr^* denotes its adjoint. Note that $\mathcal{S}(K_n)[\varphi](\mathbf{x}) = \text{Tr} \int_U K_n(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y}$ and $\mathcal{S}(K)[\varphi](\mathbf{x}) = \text{Tr} \int_U K(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y}$. Thus it's clear that $\mathcal{S}(K_n) \rightarrow \mathcal{S}(K)$ by (C.2).

We next prove $\mathcal{K}^*(K_n) \rightarrow \mathcal{K}^*(K)$. For each $\varphi \in \tilde{H}^{-\frac{1}{2}}(\Gamma)$, (C.1) implies that

$$\frac{\partial}{\partial x_1} \int_U K_n(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y} \xrightarrow{\tilde{H}^{-\frac{1}{2}}(\Gamma)} \frac{\partial}{\partial x_1} \int_U K(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y}, \quad \mathbf{x} \in \Gamma.$$

By the jump condition (5.4), it is equivalent to

$$p.v. \int_\Gamma \frac{\partial}{\partial x_1} K_n(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y} \xrightarrow{\tilde{H}^{-\frac{1}{2}}(\Gamma)} p.v. \int_\Gamma \frac{\partial}{\partial x_1} K(\mathbf{x}, \mathbf{y}) (\text{Tr}^* \varphi)(\mathbf{y}) d\mathbf{y}, \quad (\text{C.3})$$

which gives $\mathcal{K}^*(K_n) \rightarrow \mathcal{K}^*(K)$.

We finally prove $\mathcal{K}(K_n) \rightarrow \mathcal{K}(K)$. The condition $K_n(\mathbf{x}, \mathbf{y}) = \overline{K_n(\mathbf{y}, \mathbf{x})}$ and (C.3) implies that the following convergence holds uniformly for all $\phi \in H^{\frac{1}{2}}(\Gamma)$ and $\varphi \in \tilde{H}^{-\frac{1}{2}}(\Gamma)$ with unit norm

$$\begin{aligned} \langle \varphi(\mathbf{y}), p.v. \int_\Gamma \frac{\partial}{\partial x_1} K_n(\mathbf{y}, \mathbf{x}) \phi(\mathbf{x}) d\mathbf{x} \rangle &= \overline{\langle \phi(\mathbf{x}), p.v. \int_\Gamma \frac{\partial}{\partial x_1} K_n(\mathbf{x}, \mathbf{y}) \varphi(\mathbf{y}) d\mathbf{y} \rangle} \\ &\rightarrow \overline{\langle \phi(\mathbf{x}), p.v. \int_\Gamma \frac{\partial}{\partial x_1} K(\mathbf{x}, \mathbf{y}) \varphi(\mathbf{y}) d\mathbf{y} \rangle} \\ &= \langle \varphi(\mathbf{y}), p.v. \int_\Gamma \frac{\partial}{\partial x_1} K(\mathbf{y}, \mathbf{x}) \phi(\mathbf{x}) d\mathbf{x} \rangle, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the $\tilde{H}^{-\frac{1}{2}} - H^{\frac{1}{2}}$ pairing. Thus $\mathcal{K}(K_n) \rightarrow \mathcal{K}(K)$ is proved. \square

We decompose $G^\delta(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) = G^{\delta, \text{evan}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) + G^{\delta, \text{prop}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h)$, where

$$\begin{aligned} G^{\delta, \text{evan}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) &= \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_{n, \delta}(\mathbf{x}; \kappa_1) \overline{v_{n, \delta}(\mathbf{y}; \kappa_1)}}{\lambda_* + \delta \cdot h - \mu_{n, \delta}(\kappa_1)} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_{[0, \pi - \delta^{\frac{1}{6}}] \cup [\pi + \delta^{\frac{1}{6}}, 2\pi]} \frac{v_{1, \delta}(\mathbf{x}; \kappa_1) \overline{v_{1, \delta}(\mathbf{y}; \kappa_1)}}{\lambda_* + \delta \cdot h - \mu_{1, \delta}(\kappa_1)} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_{[0, \pi - \delta^{\frac{1}{6}}] \cup [\pi + \delta^{\frac{1}{6}}, 2\pi]} \frac{v_{2, \delta}(\mathbf{x}; \kappa_1) \overline{v_{2, \delta}(\mathbf{y}; \kappa_1)}}{\lambda_* + \delta \cdot h - \mu_{2, \delta}(\kappa_1)} d\kappa_1, \end{aligned}$$

and

$$G^{\delta, \text{prop}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) = \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{1, \delta}(\mathbf{x}; \kappa_1) \overline{v_{1, \delta}(\mathbf{y}; \kappa_1)}}{\lambda_* + \delta \cdot h - \mu_{1, \delta}(\kappa_1)} d\kappa_1 + \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{2, \delta}(\mathbf{x}; \kappa_1) \overline{v_{2, \delta}(\mathbf{y}; \kappa_1)}}{\lambda_* + \delta \cdot h - \mu_{2, \delta}(\kappa_1)} d\kappa_1.$$

As in the proof of Proposition 3.5, we take a neighborhood of Γ , i.e. $U := (-\frac{1-2r_0}{4}, \frac{1-2r_0}{4}) \times (-\frac{1}{2}, \frac{1}{2})$, where $r_0 := \text{diam}(V)$. Then we claim the following results:

Lemma C.2. *There exist $f_{n, \delta}^{\text{evan}}, g_{n, \delta}^{\text{evan}}, f_{n, \delta}^{\text{prop}}, g_{n, \delta}^{\text{prop}} \in H^1(U)$ ($n = 1, 2$) such that the following asymptotics holds uniformly for all $f \in (H^1(U))^*$ with unit norm*

$$\begin{aligned} &\int_U G^{\delta, \text{evan}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) f(\mathbf{y}) d\mathbf{y} \\ &= (\delta^{-\frac{1}{6}} \cdot \frac{2}{\pi\gamma_*} + o(\delta^{-\frac{1}{6}})) \cdot (v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega - v_2(\mathbf{x}; \pi)(f(\mathbf{y}), v_2(\mathbf{y}; \pi))_\Omega) \\ &\quad + \mathcal{G}_0(\lambda_*) f \\ &\quad + \frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_1)(\mathbf{y}; \pi))_\Omega - \frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} \cdot (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_2)(\mathbf{y}; \pi))_\Omega \\ &\quad + \sum_{n=1,2} (v_n(\mathbf{x}; \pi)(f(\mathbf{y}), f_{n, \delta}^{\text{evan}}(\mathbf{y}))_\Omega + g_{n, \delta}^{\text{evan}}(\mathbf{x})(f(\mathbf{y}), v_n(\mathbf{y}; \pi))_\Omega) + \mathcal{O}(\delta^{\frac{1}{3}}), \end{aligned} \tag{C.4}$$

$$\begin{aligned} &\int_U G^{\delta, \text{prop}}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) f(\mathbf{y}) d\mathbf{y} \\ &= - \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \left(\frac{\frac{h}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} - \frac{1}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{1}{4}}} \right) + o(\delta^{-\frac{1}{2}}) \right) \cdot v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega \\ &\quad - \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \left(\frac{\frac{h}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} + \frac{1}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{1}{4}}} \right) + o(\delta^{-\frac{1}{2}}) \right) \cdot v_2(\mathbf{x}; \pi)(f(\mathbf{y}), v_2(\mathbf{y}; \pi))_\Omega \\ &\quad + \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} + o(\delta^{-\frac{1}{2}}) \right) \cdot (v_2(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega + v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_2(\mathbf{y}; \pi))_\Omega) \\ &\quad + \left(\frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} + o(\delta^{\frac{1}{6}}) \right) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_1)(\mathbf{y}; \pi))_\Omega \\ &\quad + \left(-\frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} + o(\delta^{\frac{1}{6}}) \right) \cdot (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_2)(\mathbf{y}; \pi))_\Omega \\ &\quad + \sum_{n=1,2} (v_n(\mathbf{x}; \pi)(f(\mathbf{y}), f_{n, \delta}^{\text{prop}}(\mathbf{y}))_\Omega + g_{n, \delta}^{\text{prop}}(\mathbf{x})(f(\mathbf{y}), v_n(\mathbf{y}; \pi))_\Omega) + \mathcal{O}(\delta^{\frac{1}{3}}), \end{aligned} \tag{C.5}$$

and $\|f_{n,\delta}^{evan}\|_{H^1(U)}, \|g_{n,\delta}^{evan}\|_{H^1(U)}, \|f_{n,\delta}^{prop}\|_{H^1(U)}, \|g_{n,\delta}^{prop}\|_{H^1(U)} = \mathcal{O}(\delta^{\frac{1}{6}})$.

By Lemma C.1 and Lemma C.2, we obtain the convergence of interface integral operator $\mathbb{M}(\lambda_* + \delta \cdot h)$ in (5.12). We point out that the scaling in $\mathbb{M}(\lambda_* + \delta \cdot h)$ is critical for the convergence. In fact, as seen from (C.4) and (C.5), the scaling we choose for each subspace of $H^{\frac{1}{2}}(\Gamma) \times \tilde{H}^{-\frac{1}{2}}(\Gamma)$ balances the δ -order of the corresponding operator exactly. This guarantees the uniform convergence of each component of $\mathbb{M}(\lambda_* + \delta \cdot h)$. We also point out that the major terms (of which we give detailed expressions) in (C.4) and (C.5) correspond to the nonzero terms in $\lim_{\delta \rightarrow 0^+} \mathbb{M}(\lambda_* + \delta \cdot h)$, while the minor terms (of which we only estimate the order) correspond to the zeros in $\lim_{\delta \rightarrow 0^+} \mathbb{M}(\lambda_* + \delta \cdot h)$. This completes the proof of (5.14).

We now prove (C.4) and (C.5) separately below.

Proof of (C.4)

We claim that

$$\frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_{n,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{n,\delta}(\mathbf{y}; \kappa_1))}{\lambda_* + \delta \cdot h - \mu_{n,\delta}(\kappa_1)} d\kappa_1 = \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_n(\mathbf{y}; \kappa_1))}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 + \mathcal{O}(\delta), \quad (\text{C.6})$$

$$\begin{aligned} & \frac{1}{2\pi} \int_{[0, \pi - \delta^{\frac{1}{6}}] \cup [\pi + \delta^{\frac{1}{6}}, 2\pi]} \frac{v_{n,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{n,\delta}(\mathbf{y}; \kappa_1))_{\Omega}}{\lambda_* + \delta \cdot h - \mu_{n,\delta}(\kappa_1)} d\kappa_1 \\ &= \frac{1}{2\pi} \int_{[0, \pi - \delta^{\frac{1}{6}}] \cup [\pi + \delta^{\frac{1}{6}}, 2\pi]} \frac{v_n(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_n(\mathbf{y}; \kappa_1))_{\Omega}}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 + \mathcal{O}(\delta^{\frac{1}{3}}), \quad n = 1, 2. \end{aligned} \quad (\text{C.7})$$

By (C.6) and (C.7), we have

$$\int_U G^{\delta, evan}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) f(\mathbf{y}) d\mathbf{y} = \mathcal{G}^{\delta, evan}(\lambda_*) f + \mathcal{O}(\delta^{\frac{1}{3}}), \quad (\text{C.8})$$

where $\mathcal{G}^{\delta, evan}(\lambda_*)$ is defined in (A.1) (the parameter ϵ in the original definition is replaced by δ). Then (C.4) follows from (C.8) and Lemma A.2.

In the sequel, we prove (C.6) and (C.7).

Step 1. The proof of (C.6) follows the same strategy as the one of (A.4). We define the following projections

$$\mathbb{P}_{n,\delta}(\kappa_1)g = (g(\cdot), v_{n,\delta}(\cdot; \overline{\kappa_1}))v_{n,\delta}(\cdot; \kappa_1) \quad (n = 1, 2) \quad \mathbb{Q}_{\delta}(\kappa_1) = I - \mathbb{P}_{1,\delta}(\kappa_1) - \mathbb{P}_{2,\delta}(\kappa_1).$$

Then, by Floquet transform, we can rewrite the integral operator in (C.6) as

$$\begin{aligned} & \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_{n,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{n,\delta}(\mathbf{y}; \kappa_1))_{\Omega}}{\lambda_* + \delta \cdot h - \mu_{n,\delta}(\kappa_1)} d\kappa_1 - \frac{1}{2\pi} \int_0^{2\pi} \sum_{n \geq 3} \frac{v_n(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_n(\mathbf{y}; \kappa_1))_{\Omega}}{\lambda_* - \mu_n(\kappa_1)} d\kappa_1 \\ &= \frac{1}{2\pi} \int_0^{2\pi} (\lambda_* + \delta \cdot h - \mathcal{L}^{A+\delta \cdot B}(\kappa_1, \pi) \mathbb{Q}_{\delta}(\kappa_1))^{-1} \mathbb{Q}_{\delta}(\kappa_1) \hat{f}(\kappa_1) d\kappa_1 \\ & \quad - \frac{1}{2\pi} \int_0^{2\pi} (\lambda_* - \mathcal{L}^A(\kappa_1, \pi) \mathbb{Q}(\kappa_1))^{-1} \mathbb{Q}(\kappa_1) \hat{f}(\kappa_1) d\kappa_1. \end{aligned} \quad (\text{C.9})$$

Note that

$$\sigma(\mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1) \mathbb{Q}_{\delta}(\kappa_1)) = \{0\} \cup \{\mu_{n,\delta}(\kappa_1)\}_{n \geq 3}.$$

Thus, for $h \in \mathcal{J}$, the resolvent $\lambda_* + \delta \cdot h - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1)$ is uniformly bounded. Consequently, as in the proof of Lemma A.2, the resolvent identity shows that

$$\left\| \left(\lambda_* + \delta \cdot h - \mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1)\mathbb{Q}_\delta(\kappa_1) \right)^{-1} - \left(\lambda_* - \mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1)\mathbb{Q}_\delta(\kappa_1) \right)^{-1} \right\|_{\mathcal{B}((H^1(Y))^*, H^1(Y))} = \mathcal{O}(\delta). \quad (\text{C.10})$$

On the other hand, the estimates $\|\mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1) - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\|_{\mathcal{B}(\mathcal{B}(H^1(Y), (H^1(Y))^*))} = \mathcal{O}(\delta)$ and $\|\mathbb{Q}_\delta(\kappa_1) - \mathbb{Q}(\kappa_1)\|_{\mathcal{B}(H^1(Y), (H^1(Y))^*)} = \mathcal{O}(\delta)$ indicate

$$\|\mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1)\mathbb{Q}_\delta(\kappa_1) - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1)\|_{\mathcal{B}(H^1(Y), (H^1(Y))^*)} = \mathcal{O}(\delta).$$

This implies the estimate of resolvents

$$\left\| \left(\lambda_* - \mathcal{L}_{\Omega, \pi}^{A+\delta \cdot B}(\kappa_1)\mathbb{Q}_\delta(\kappa_1) \right)^{-1} - \left(\lambda_* - \mathcal{L}_{\Omega, \pi}^A(\kappa_1)\mathbb{Q}(\kappa_1) \right)^{-1} \right\|_{\mathcal{B}((H^1(Y))^*, H^1(Y))} = \mathcal{O}(\delta). \quad (\text{C.11})$$

By taking (C.10) and (C.11) inside (C.9), we obtain (C.6).

Step 2. We prove (C.7) for $n = 1$. The proof of the $n = 2$ case is similar. Note that Assumption 1.4 indicates that for $\delta^{\frac{1}{6}} \leq |\kappa_1 - \pi| \leq \pi$, we have

$$|\mu_1(\kappa_1) - \lambda_*| \geq |\mu_1(\pi + \delta^{\frac{1}{6}}) - \lambda_*| \gtrsim \delta^{\frac{1}{3}}. \quad (\text{C.12})$$

The following estimates for $\delta^{\frac{1}{6}} \leq |\kappa_1 - \pi| \leq \pi$ follow from a similar perturbation argument as we did in Theorem 4.1,

$$|\mu_{1, \delta}(\kappa_1) - \mu_1(\kappa_1)| = \mathcal{O}(\delta), \quad (\text{C.13})$$

$$\|v_{1, \delta}(\mathbf{x}; \kappa_1) - v_1(\mathbf{x}; \kappa_1)\|_{H^1(Y)} = \mathcal{O}(\delta^{\frac{2}{3}}). \quad (\text{C.14})$$

(C.12) and (C.13) imply that

$$\frac{1}{\lambda_* + \delta \cdot h - \mu_{1, \delta}(\kappa_1)} = \mathcal{O}(\delta^{-\frac{1}{3}}), \quad \forall h \in \mathcal{J}. \quad (\text{C.15})$$

By (C.15) and (C.14), one can check that for $\delta^{\frac{1}{6}} \leq |\kappa_1 - \pi| \leq \pi$

$$\frac{v_{1, \delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{1, \delta}(\mathbf{y}; \kappa_1))_\Omega}{\lambda_* + \delta \cdot h - \mu_{1, \delta}(\kappa_1)} - \frac{v_1(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_1(\mathbf{y}; \kappa_1))_\Omega}{\lambda_* - \mu_1(\kappa_1)} = \mathcal{O}(\delta^{\frac{1}{3}}).$$

Hence (C.7) follows.

Proof of (C.5)

Note that

$$\begin{aligned} & \int_U G^{\delta, prop}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) f(\mathbf{y}) d\mathbf{y} \\ &= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{1, \delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{1, \delta}(\mathbf{y}; \kappa_1))_\Omega}{\lambda_* + \delta \cdot h - \mu_{1, \delta}(\kappa_1)} \frac{1}{N_{1, \delta}^2(\kappa_1)} d\kappa_1 \\ & \quad + \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{2, \delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{2, \delta}(\mathbf{y}; \kappa_1))_\Omega}{\lambda_* + \delta \cdot h - \mu_{2, \delta}(\kappa_1)} \frac{1}{N_{2, \delta}^2(\kappa_1)} d\kappa_1, \end{aligned}$$

where $v_{n,\delta}(\mathbf{x}; \kappa_1)$ admits the asymptotic expansion as given in Theorem 4.1, and $N_{n,\delta}(\kappa_1)$ is its normalization factor. Moreover, by Theorem 4.1,

$$\begin{aligned}\lambda_* + \delta \cdot h - \mu_{1,\delta}(\kappa_1) &= \left(\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2} \right) \cdot (1 + \mathcal{O}(|\kappa_1 - \pi| + \delta)), \\ \lambda_* + \delta \cdot h - \mu_{2,\delta}(\kappa_1) &= \left(\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2} \right) \cdot (1 + \mathcal{O}(|\kappa_1 - \pi| + \delta)), \\ N_{n,\delta}(\kappa_1) &= \left(1 + f_\delta(\kappa_1)^2 \right)^{\frac{1}{2}} \cdot (1 + \mathcal{O}(|\kappa_1 - \pi| + \delta)),\end{aligned}$$

where

$$f_\delta(\kappa_1) := \frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}.$$

Thus,

$$\int_U G^{\delta,prop}(\mathbf{x}, \mathbf{y}; \lambda_* + \delta \cdot h) f(\mathbf{y}) d\mathbf{y} = I + II,$$

where

$$\begin{aligned}I &= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{1,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{1,\delta}(\mathbf{y}; \kappa_1))_\Omega}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{2,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{2,\delta}(\mathbf{y}; \kappa_1))_\Omega}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1,\end{aligned}\tag{C.16}$$

and

$$\begin{aligned}II &= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{1,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{1,\delta}(\mathbf{y}; \kappa_1))_\Omega}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{\mathcal{O}(|\kappa_1 - \pi| + \delta)}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ &\quad + \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{2,\delta}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{2,\delta}(\mathbf{y}; \kappa_1))_\Omega}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{\mathcal{O}(|\kappa_1 - \pi| + \delta)}{1 + f_\delta(\kappa_1)^2} d\kappa_1.\end{aligned}\tag{C.17}$$

We will compute (C.16) in detail and show that it contributes to all the major terms on the right-hand side of (C.5). Following the same lines, (C.17) contributes to only the minor terms on the right-hand side of (C.5). This proof is omitted here.

By the asymptotic expansion (4.3), we write

$$v_{1,\delta}(\mathbf{x}; \kappa_1) = \sum_{i=1}^5 v_{1,\delta}^{(i)}(\mathbf{x}; \kappa_1), \quad v_{2,\delta}(\mathbf{x}; \kappa_1) = \sum_{i=1}^5 v_{2,\delta}^{(i)}(\mathbf{x}; \kappa_1),$$

where

$$\begin{aligned}
v_{1,\delta}^{(1)}(\mathbf{x}; \kappa_1) &:= (1 + \mathcal{O}(|\kappa_1 - \pi|))v_1(\mathbf{x}; \pi), \\
v_{1,\delta}^{(2)}(\mathbf{x}; \kappa_1) &:= \left(\frac{t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \mathcal{O}(|\kappa_1 - \pi|) \right) \cdot v_2(\mathbf{x}; \pi), \\
v_{1,\delta}^{(3)}(\mathbf{x}; \kappa_1) &:= (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi), \\
v_{1,\delta}^{(4)}(\mathbf{x}; \kappa_1) &:= \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_1(\mathbf{x}), \\
v_{1,\delta}^{(5)}(\mathbf{x}; \kappa_1) &:= R_1(\mathbf{x}; \kappa_1, \delta),
\end{aligned} \tag{C.18}$$

and

$$\begin{aligned}
v_{2,\delta}^{(1)}(\mathbf{x}; \kappa_1) &:= \left(\frac{-t_*\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} + \mathcal{O}(|\kappa_1 - \pi|) \right) \cdot v_1(\mathbf{x}; \pi), \\
v_{2,\delta}^{(2)}(\mathbf{x}; \kappa_1) &:= (1 + \mathcal{O}(|\kappa_1 - \pi|))v_2(\mathbf{x}; \pi), \\
v_{2,\delta}^{(3)}(\mathbf{x}; \kappa_1) &:= (\kappa_1 - \pi) \cdot (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi), \\
v_{2,\delta}^{(4)}(\mathbf{x}; \kappa_1) &:= \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \cdot r_2(\mathbf{x}), \\
v_{2,\delta}^{(5)}(\mathbf{x}; \kappa_1) &:= R_2(\mathbf{x}; \kappa_1, \delta).
\end{aligned} \tag{C.19}$$

Then (C.16) is rewritten as

$$I = \sum_{1 \leq i, j \leq 5, 1 \leq k \leq 2} I_k^{i,j}, \tag{C.20}$$

where

$$\begin{aligned}
I_1^{i,j} &:= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{1,\delta}^{(i)}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{1,\delta}^{(j)}(\mathbf{y}; \kappa_1))\Omega}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1, \\
I_2^{i,j} &:= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{v_{2,\delta}^{(i)}(\mathbf{x}; \kappa_1)(f(\mathbf{y}), v_{2,\delta}^{(j)}(\mathbf{y}; \kappa_1))\Omega}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1.
\end{aligned}$$

Among the 50 terms in (C.20), we will show that only 10 terms contribute to the major terms on the right-hand side of (C.5). The detail is presented in Steps 1-4 in the sequel. The remaining 40 terms are estimated by elementary integral estimation (presented in Lemma C.3) as shown in Steps 5-7. The following lemma are frequently recalled in the proof.

Lemma C.3. *For $|h| < c_0|t_*|$, we have*

$$\int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = 0, \tag{C.21}$$

$$\int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{|\kappa_1 - \pi|}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = \mathcal{O}(\ln \delta), \tag{C.22}$$

$$\int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = \frac{4}{\gamma_*} \delta^{\frac{1}{6}} + o(\delta^{\frac{1}{6}}), \quad (\text{C.23})$$

$$\int_{\pi}^{\pi+\delta^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = \mathcal{O}(\delta^{\frac{1}{3}}). \quad (\text{C.24})$$

Moreover,

$$\begin{aligned} & \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{1}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ & + \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{f_\delta(\kappa_1)^2}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ & = -\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \left(\frac{\frac{h}{\gamma_*}}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{3}{4}}} - \frac{1}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{1}{4}}} \right) + o(\delta^{-\frac{1}{2}}), \end{aligned} \quad (\text{C.25})$$

$$\begin{aligned} & \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{f_\delta(\kappa_1)}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ & + \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{-f_\delta(\kappa_1)}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ & = \frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{\left(\left(\frac{t_*}{\gamma_*}\right)^2 - \left(\frac{h}{\gamma_*}\right)^2\right)^{\frac{3}{4}}} + o(\delta^{-\frac{1}{2}}), \end{aligned} \quad (\text{C.26})$$

Proof. (C.21) is obvious because the integral is an odd function in $\kappa_1 - \pi$.

For (C.22), we note that $\frac{1}{1+f_\delta(\kappa_1)^2} = \mathcal{O}(1)$ and

$$\frac{\sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} = \mathcal{O}(1), \quad \text{for } |h| < c_0|t_*|,$$

where we used the fact that $c_0 < 1$. Hence

$$\begin{aligned} & \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{|\kappa_1 - \pi|}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ & \lesssim \int_{\pi}^{\pi+\delta^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{\sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} d\kappa_1 \lesssim \int_0^{\delta^{\frac{1}{3}}} \frac{dp}{\sqrt{p^2 + \delta^2}} = \mathcal{O}(\ln \delta). \end{aligned}$$

For (C.23), a change of variable $p = \sqrt{\frac{\gamma_*}{2}} \frac{\kappa_1 - \pi}{\sqrt{\delta}}$ leads to

$$\begin{aligned} & \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ &= 2\sqrt{\delta} \cdot \left(\frac{2}{\gamma_*}\right)^{\frac{3}{2}} \int_0^{\sqrt{\frac{\gamma_*}{2}}\delta^{-\frac{1}{3}}} \frac{p^2}{h + \sqrt{p^4 + t_*^2}} \frac{(p^2 + \sqrt{p^4 + t_*^2})^2}{(p^2 + \sqrt{p^4 + t_*^2})^2 + t_*^2} dp. \end{aligned} \quad (\text{C.27})$$

The integrand is uniformly bounded and has the following asymptotics as $p \rightarrow \infty$

$$\frac{p^2}{h + \sqrt{p^4 + t_*^2}} \frac{(p^2 + \sqrt{p^4 + t_*^2})^2}{(p^2 + \sqrt{p^4 + t_*^2})^2 + t_*^2} = 1 + \mathcal{O}\left(\frac{1}{p^2}\right).$$

Hence

$$\int_0^{\sqrt{\frac{\gamma_*}{2}}\delta^{-\frac{1}{3}}} \frac{p^2}{h + \sqrt{p^4 + t_*^2}} \frac{(p^2 + \sqrt{p^4 + t_*^2})^2}{(p^2 + \sqrt{p^4 + t_*^2})^2 + t_*^2} dp = \sqrt{\frac{\gamma_*}{2}}\delta^{-\frac{1}{3}} + o(\delta^{-\frac{1}{3}}). \quad (\text{C.28})$$

Then (C.23) follows from (C.27) and (C.28).

For (C.24), we first note the following estimate for $\pi \leq \kappa_1 \leq \pi + \delta^{\frac{1}{6}}$

$$\frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} = \mathcal{O}(\sqrt{\delta}). \quad (\text{C.29})$$

When $\kappa_1 = \pi$, (C.29) holds trivially. When $\kappa_1 > \pi$, we have

$$\begin{aligned} \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} &= \frac{\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi) + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^2 + \frac{t_*^2\delta^2}{(\kappa_1 - \pi)^2}}} \\ &\leq \frac{\delta}{\frac{1}{2}\gamma_*(\kappa_1 - \pi) + \sqrt{\gamma_* t_* \delta}} \lesssim \sqrt{\delta}, \end{aligned}$$

which implies (C.29). By using (C.29) and (C.22), we obtain that

$$\begin{aligned} & \int_{\pi}^{\pi + \delta^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\ &\lesssim \sqrt{\delta} \cdot \int_{\pi}^{\pi + \delta^{\frac{1}{6}}} \frac{|\kappa_1 - \pi|}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = \mathcal{O}(\sqrt{\delta} \ln \delta), \end{aligned}$$

which concludes the proof of (C.24).

For (C.25), we can directly calculate that

$$\begin{aligned} LHS &= \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{\delta \cdot h}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1 \\ &\quad - \frac{1}{2\pi} \int_{\pi - \delta^{\frac{1}{6}}}^{\pi + \delta^{\frac{1}{6}}} \frac{\sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} \frac{1 - f_\delta(\kappa_1)^2}{1 + f_\delta(\kappa_1)^2} d\kappa_1. \end{aligned} \quad (\text{C.30})$$

Note that

$$\frac{1 - f_\delta(\kappa_1)^2}{1 + f_\delta(\kappa_1)^2} = \frac{\gamma_*(\kappa_1 - \pi)^2}{2\sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}.$$

Thus

$$\begin{aligned} LHS &= \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\delta \cdot h}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1 \\ &\quad - \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1. \end{aligned} \quad (\text{C.31})$$

By direct calculation, we have

$$\begin{aligned} \frac{\delta^{\frac{1}{2}}}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\delta \cdot h}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1 &\xrightarrow{\delta \rightarrow 0} -\frac{1}{2\gamma_*} \frac{\frac{h}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}}, \\ \frac{\delta^{\frac{1}{2}}}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1 &\xrightarrow{\delta \rightarrow 0} -\frac{1}{2\gamma_*} \frac{1}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{1}{4}}}, \end{aligned}$$

whence (C.25) follows.

Similarly, for (C.26), we can directly calculate that

$$LHS = \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{-2\sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} \frac{f_\delta(\kappa_1)}{1 + f_\delta(\kappa_1)^2} d\kappa_1. \quad (\text{C.32})$$

Note that

$$\frac{f_\delta(\kappa_1)}{1 + f_\delta(\kappa_1)^2} = \frac{t_*\delta}{2\sqrt{\frac{\delta \cdot h}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)}}}.$$

Hence

$$LHS = \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{-t_*\delta}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1. \quad (\text{C.33})$$

By direct calculation, we have

$$\frac{\delta^{\frac{1}{2}}}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{-t_*\delta}{\delta^2 \cdot h^2 - (\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2)} d\kappa_1 \xrightarrow{\delta \rightarrow 0} \frac{1}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}},$$

whence (C.26) follows. □

Step 1 (2 terms are considered). In this step, we show that

$$I_1^{1,1} + I_2^{1,1} = -\left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \left(\frac{\frac{h}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} - \frac{1}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{1}{4}}} \right) + o(\delta^{-\frac{1}{2}}) \right) \cdot v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega. \quad (\text{C.34})$$

Indeed, by (C.22),

$$\begin{aligned}
& \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{|1 + \mathcal{O}(|\kappa_1 - \pi|)|^2 - 1}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\
&= \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\mathcal{O}(|\kappa_1 - \pi|)}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \\
&= \mathcal{O}(\ln \delta) = o(\delta^{-\frac{1}{2}}).
\end{aligned}$$

Similarly,

$$\frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{|f_\delta(\kappa_1) + \mathcal{O}(|\kappa_1 - \pi|)|^2 - f_\delta^2(\kappa_1)}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 = o(\delta^{-\frac{1}{2}}).$$

Hence the definition of $I_1^{1,1}, I_2^{1,1}$ indicates

$$\begin{aligned}
& I_1^{1,1} + I_2^{1,1} \\
&= \left(\frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{1}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \right. \\
&\quad \left. + \frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{f_\delta(\kappa_1)^2}{\delta \cdot h - \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 + o(\delta^{-\frac{1}{2}}) \right) v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega.
\end{aligned}$$

Then (C.34) follows by applying (C.25).

Step 2 (2 terms are considered). Similar as Step 1, we have

$$I_1^{2,2} + I_2^{2,2} = - \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \left(\frac{\frac{h}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} + \frac{1}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{1}{4}}} \right) + o(\delta^{-\frac{1}{2}}) \right) \cdot v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega.$$

Step 3 (4 terms are considered). By (C.18) and (C.22) and (C.26), we can show that

$$I_1^{1,2} + I_2^{1,2} = \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} + o(\delta^{-\frac{1}{2}}) \right) \cdot v_1(\mathbf{x}; \pi)(f(\mathbf{y}), v_2(\mathbf{y}; \pi))_\Omega.$$

Similarly,

$$I_1^{2,1} + I_2^{2,1} = \left(\frac{\delta^{-\frac{1}{2}}}{2\gamma_*} \frac{\frac{t_*}{\gamma_*}}{((\frac{t_*}{\gamma_*})^2 - (\frac{h}{\gamma_*})^2)^{\frac{3}{4}}} + o(\delta^{-\frac{1}{2}}) \right) \cdot v_2(\mathbf{x}; \pi)(f(\mathbf{y}), v_1(\mathbf{y}; \pi))_\Omega.$$

Step 4 (2 terms are considered). By (C.18),

$$I_1^{3,3} = \left(\frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{(\kappa_1 - \pi)^2}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_\delta(\kappa_1)^2} d\kappa_1 \right) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_1)(\mathbf{y}; \pi))_\Omega.$$

Hence, using (C.23), we obtain

$$I_1^{3,3} = \left(\frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} + o(\delta^{\frac{1}{6}}) \right) \cdot (\partial_{\kappa_1} v_1)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_1)(\mathbf{y}; \pi))_{\Omega}. \quad (\text{C.35})$$

Similarly,

$$I_2^{3,3} = \left(-\frac{2}{\pi\gamma_*} \delta^{\frac{1}{6}} + o(\delta^{\frac{1}{6}}) \right) \cdot (\partial_{\kappa_1} v_2)(\mathbf{x}; \pi)(f(\mathbf{y}), (\partial_{\kappa_1} v_2)(\mathbf{y}; \pi))_{\Omega}.$$

Step 5 (6 terms are considered). We estimate the integral $I_1^{1,3}$. By (C.18),

$$I_1^{1,3} = \left(\frac{1}{2\pi} \int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{(\kappa_1 - \pi) + \mathcal{O}(|\kappa_1 - \pi|^2)}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_{\delta}(\kappa_1)^2} d\kappa_1 \right) \cdot v_1(\mathbf{x}; \pi)(f(\mathbf{y}), \partial_{\kappa_1} v_1(\mathbf{x}; \pi))_{\Omega}. \quad (\text{C.36})$$

By (C.23), we have

$$\int_{\pi-\delta^{\frac{1}{6}}}^{\pi+\delta^{\frac{1}{6}}} \frac{\mathcal{O}(|\kappa_1 - \pi|^2)}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_{\delta}(\kappa_1)^2} d\kappa_1 = \mathcal{O}(\delta^{\frac{1}{6}}). \quad (\text{C.37})$$

Hence, by (C.36), (C.21) and (C.37), we conclude that $I_1^{1,3}$ can be expressed in the form $v_1(\mathbf{x}; \pi)(f(\mathbf{y}), f_{1,\delta}^{prop}(\mathbf{y}))_{\Omega}$ where $\|f_{1,\delta}^{prop}\|_{H^1(U)} = \mathcal{O}(\delta^{\frac{1}{6}})$.

Similar arguments work for the integrals

$$I_1^{1,4}, I_1^{1,5}, I_2^{1,3}, I_2^{1,4}, I_2^{1,5}.$$

Step 6 (18 terms are considered). Similar to Step 5, we can write

$$\begin{aligned} I_1^{3,1} + I_1^{4,1} + I_1^{5,1} + I_2^{3,1} + I_2^{4,1} + I_2^{5,1} &= g_{1,\delta}^{prop}(\mathbf{x})(f(\mathbf{y}), v_1(\mathbf{y}; \pi)), \\ I_1^{2,3} + I_1^{2,4} + I_1^{2,5} + I_2^{2,3} + I_2^{2,4} + I_2^{2,5} &= v_2(\mathbf{x}; \pi)(f(\mathbf{y}), f_{2,\delta}^{prop}(\mathbf{x})), \\ I_1^{3,2} + I_1^{4,2} + I_1^{5,2} + I_2^{3,2} + I_2^{4,2} + I_2^{5,2} &= g_{2,\delta}^{prop}(\mathbf{x})(f(\mathbf{y}), v_2(\mathbf{y}; \pi)), \end{aligned}$$

with $\|g_{1,\delta}^{prop}\|_{H^1(U)}, \|f_{2,\delta}^{prop}\|_{H^1(U)}, \|g_{2,\delta}^{prop}\|_{H^1(U)} = \mathcal{O}(\delta^{\frac{1}{6}})$.

Step 7 (16 terms are considered). We show that $I_1^{3,4} = \mathcal{O}(\delta^{\frac{1}{3}})$. Indeed, by (C.18), $I_1^{3,4}$ is proportional to

$$\int_{\pi}^{\pi+\delta^{\frac{1}{6}}} \frac{\kappa_1 - \pi}{\delta \cdot h + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{\delta(\kappa_1 - \pi)}{\frac{1}{2}\gamma_*(\kappa_1 - \pi)^2 + \sqrt{\frac{1}{4}\gamma_*^2(\kappa_1 - \pi)^4 + t_*^2\delta^2}} \frac{1}{1 + f_{\delta}(\kappa_1)^2} d\kappa_1.$$

Hence, by (C.24), $I_1^{3,4} = \mathcal{O}(\delta^{\frac{1}{3}})$. Similarly, one can check that the following integrals are of the order $\mathcal{O}(\delta^{\frac{1}{3}})$

$$I_1^{3,5}, I_1^{4,3}, I_1^{4,4}, I_1^{4,5}, I_1^{5,3}, I_1^{5,4}, I_1^{5,5}, I_2^{3,4}, I_2^{3,5}, I_2^{4,3}, I_2^{4,4}, I_2^{4,5}, I_2^{5,3}, I_2^{5,4}, I_2^{5,5}.$$

Appendix D: Proof of Proposition 5.2

Note that Γ is a bounded closed smooth curve when Ω is identified with a cylinder by the periodicity in \mathbf{e}_2 . Consequently, both $\mathcal{S}(\lambda_*; G_0)$ and $\mathcal{N}(\lambda_*; G_0)$ are Fredholm with zero index because the Green function $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ is the fundamental solution in the sense of (3.7) [43, 48]. Thus $\mathbb{T}^0(\lambda_*)$ is Fredholm.

In the following paragraph, we prove (5.16). We first show that

$$\begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix} \in \ker \mathbb{T}^0(\lambda_*). \quad (\text{D.1})$$

By applying the Green's formula between $v_2(\mathbf{x})$ and $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$ inside $(0, N) \times (-\frac{1}{2}, \frac{1}{2})$, as we did in the proof of Proposition 3.10, we have

$$\begin{aligned} \overline{v_2(\mathbf{y})} &= \frac{i}{\gamma_*} \mathbf{q}(v_1(\cdot; \pi), v_2(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_1(\cdot; \pi), v_2(\cdot; \pi); \Gamma) \cdot \overline{v_1(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \mathbf{q}(v_2(\cdot; \pi), v_2(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_2(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), v_2(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)} \quad (\text{D.2}) \\ &\quad - \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*) \overline{v_2(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*) \frac{\partial v_2}{\partial x_1}(\mathbf{x}; \pi). \end{aligned}$$

By Corollary 3.11, (D.2) becomes

$$\overline{v_2(\mathbf{y})} = \frac{1}{2} \overline{v_2(\mathbf{y}; \pi)} - \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*) \overline{v_2(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*) \frac{\partial v_2}{\partial x_1}(\mathbf{x}; \pi).$$

Taking the normal derivative for $\mathbf{y} \in \Gamma$, the jump formula (3.14), (3.15) and the symmetry (3.9) show that

$$\frac{\partial v_2}{\partial y_1}(\mathbf{y}) = \frac{1}{2} \frac{\partial v_2}{\partial y_1}(\mathbf{y}) - \overline{\mathcal{N}(\lambda_*; G_0)[v_2(\cdot; \pi)](\mathbf{y})} + \frac{1}{2} \frac{\partial v_2}{\partial y_1}(\mathbf{y}),$$

which implies $\mathcal{N}(\lambda_*; G_0)[v_2(\cdot; \pi)] = 0$, and (D.1). Similarly, we can prove

$$\begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix} \in \ker \mathbb{T}^0(\lambda_*). \quad (\text{D.3})$$

Now we show that

$$\begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\mathbf{x}) \end{pmatrix} \in \ker \mathbb{T}^0(\lambda_*). \quad (\text{D.4})$$

By applying the Green's formula between $v_1(\mathbf{x})$ and $G_0(\mathbf{x}, \mathbf{y}; \lambda_*)$, (D.2) becomes

$$\begin{aligned} \overline{v_1(\mathbf{y})} &= \frac{i}{\gamma_*} \mathbf{q}(v_1(\cdot; \pi), v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_1(\mathbf{y}; \pi)} + \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_1(\cdot; \pi), v_1(\cdot; \pi); \Gamma) \cdot \overline{v_1(\mathbf{y}; \pi)} \\ &\quad - \frac{i}{\gamma_*} \mathbf{q}(v_2(\cdot; \pi), v_1(\cdot; \pi); \Gamma) \cdot \overline{\partial_{\kappa_1} v_2(\mathbf{y}; \pi)} - \frac{i}{\gamma_*} \mathbf{q}(\partial_{\kappa_1} v_2(\cdot; \pi), v_1(\cdot; \pi); \Gamma) \cdot \overline{v_2(\mathbf{y}; \pi)} \\ &\quad - \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*) \overline{v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*) \frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi). \end{aligned}$$

With 3.11, we obtain

$$\overline{v_1(\mathbf{y})} = \frac{1}{2} \overline{v_1(\mathbf{y}; \pi)} - \int_{\Gamma} \frac{\partial G_0}{\partial x_1}(\cdot, \mathbf{y}; \lambda_*) \overline{v_1(\cdot; \pi)} - G_0(\cdot, \mathbf{y}; \lambda_*) \overline{\frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi)}.$$

By pushing $\mathbf{y} \rightarrow \Gamma$, the jump formula (3.13), (3.16) and the symmetry (3.9) show that

$$\overline{v_1(\mathbf{y})} = \frac{1}{2} \overline{v_1(\mathbf{y}; \pi)} - \left(-\frac{1}{2} \overline{v_1(\mathbf{y}; \pi)} \right) + \overline{\mathcal{S}(\lambda_*; G_0) \left[\frac{\partial v_1}{\partial x_1}(\cdot; \pi) \right] (\mathbf{y})}.$$

which implies $\mathcal{S}(\lambda_*; G_0) \left[\frac{\partial v_1}{\partial x_1}(\cdot; \pi) \right] = 0$, and (D.4). Similarly, we can prove

$$\left(\begin{array}{c} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}) \end{array} \right) \in \ker \mathbb{T}^0(\lambda_*). \quad (\text{D.5})$$

Thus, from (D.1) and (D.3)-(D.4), we conclude that

$$\ker \mathbb{T}^0(\lambda_*) \supset \text{span} \left\{ \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi) \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \end{pmatrix} \right\}. \quad (\text{D.6})$$

We next prove

$$\ker \mathbb{T}^0(\lambda_*) \subset \text{span} \left\{ \begin{pmatrix} v_2(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} \partial_{\kappa_1} v_1(\mathbf{x}; \pi) \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial v_1}{\partial x_1}(\mathbf{x}; \pi) \end{pmatrix}, \begin{pmatrix} 0 \\ \frac{\partial}{\partial x_1}(\partial_{\kappa_1} v_2)(\mathbf{x}; \pi) \end{pmatrix} \right\}. \quad (\text{D.7})$$

By the decomposition (5.9), in order to prove (D.7), it's sufficient to show that $\begin{pmatrix} \phi \\ \varphi \end{pmatrix} = 0$ for any $\begin{pmatrix} \phi \\ \varphi \end{pmatrix} \in X$ which satisfies $\mathbb{T}^0(\lambda_*) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} = 0$. Suppose the contrary that $\begin{pmatrix} \phi \\ \varphi \end{pmatrix} \neq 0$. We define the following function in Ω

$$u(\mathbf{x}; \lambda_*) := \begin{cases} -\mathcal{D}(\lambda_*; G_0)[\phi](\mathbf{x}) + \mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x}), & x_1 > 0, \\ \mathcal{D}(\lambda_*; G_0)[\phi](\mathbf{x}) - \mathcal{S}(\lambda_*; G_0)[\varphi](\mathbf{x}), & x_1 < 0. \end{cases}$$

Since $\begin{pmatrix} \phi \\ \varphi \end{pmatrix} \in X$, the decomposition (3.10) and (3.11) indicate that $u(\mathbf{x}; \lambda_*)$ exponentially decays as $|x_1| \rightarrow \infty$. On the other hand, the jump formula (3.13)-(3.16) and the assumption $\mathbb{T}^0(\lambda_*) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} = 0$ show that for $\mathbf{x} \in \Gamma$,

$$\left(\begin{array}{c} \lim_{t \rightarrow 0^+} u(\mathbf{x}; \lambda_*) \\ \lim_{t \rightarrow 0^+} \frac{\partial u}{\partial x_1}(\mathbf{x}; \lambda_*) \end{array} \right) = \left(\begin{array}{c} \frac{1}{2} \phi \\ \frac{1}{2} \varphi \end{array} \right) = \left(\begin{array}{c} \lim_{t \rightarrow 0^-} u(\mathbf{x}; \lambda_*) \\ \lim_{t \rightarrow 0^-} \frac{\partial u}{\partial x_1}(\mathbf{x}; \lambda_*) \end{array} \right) \neq \mathbf{0}.$$

Finally, (3.7) indicates that

$$(\mathcal{L}^A - \lambda_*)u = 0 \quad \text{in } \Omega.$$

Hence, $u(\mathbf{x}; \lambda_*)$ is a localized mode for the periodic operator \mathcal{L}^A in Ω , which contradicts to the absolute continuity of the spectrum \mathcal{L}^A [49]. This concludes the proof of (D.7).

By (D.6) and (D.7), the proof of (5.16) is complete.

References

- [1] Habib Ammari, Bryn Davies, Erik Orvehed Hiltunen, and Sanghyeon Yu. Topologically protected edge modes in one-dimensional chains of subwavelength resonators. *Journal de Mathématiques Pures et Appliquées*, 144:17–49, 2020.
- [2] Habib Ammari, Brian Fitzpatrick, Erik Orvehed Hiltunen, Hyundae Lee, and Sanghyeon Yu. Honeycomb-lattice minnaert bubbles. *SIAM Journal on Mathematical Analysis*, 52(6):5441–5466, 2020.
- [3] Habib Ammari, Brian Fitzpatrick, Hyeonbae Kang, Matias Ruiz, Sanghyeon Yu, and Hai Zhang. *Mathematical and computational methods in photonics and phononics*, volume 235. American Mathematical Society, 2018.
- [4] Habib Ammari, Erik Orvehed Hiltunen, and Sanghyeon Yu. A high-frequency homogenization approach near the dirac points in bubbly honeycomb crystals. *Archive for Rational Mechanics and Analysis*, 238(3):1559–1583, Dec 2020.
- [5] Julio Cesar Avila, Hermann Schulz-Baldes, and Carlos Villegas-Blas. Topological invariants of edge states for periodic two-dimensional models. *Mathematical Physics, Analysis and Geometry*, 16(2):137–170, Jun 2013.
- [6] Gregory Berkolaiko and Andrew Comech. Symmetry and dirac points in graphene spectrum. *Journal of Spectral Theory*, 8(3):1099–1147, 2018.
- [7] Chris Bourne, Johannes Kellendonk, and Adam Rennie. The k-theoretic bulk–edge correspondence for topological insulators. In *Annales Henri Poincaré*, volume 18, pages 1833–1866. Springer, 2017.
- [8] Chris Bourne and Adam Graham Rennie. Chern numbers, localisation and the bulk-edge correspondence for continuous models of topological phases. *Mathematical Physics, Analysis and Geometry*, 21:1–62, 2016.
- [9] Maxim Braverman. Spectral flows of toeplitz operators and bulk-edge correspondence. *Letters in Mathematical Physics*, 109:2271–2289, 2019.
- [10] Jonah Chaban and Michael I. Weinstein. Instability of quadratic band degeneracies and the emergence of dirac points, 2024.
- [11] David Colton and Rainer Kress. *Integral equation methods in scattering theory*. SIAM, 2013.
- [12] Jean-Michel Combes and François Germinet. Edge and impurity effects on quantization of hall currents. *Communications in Mathematical Physics*, 256(1):159–180, May 2005.
- [13] Alexis Drouot. The bulk-edge correspondence for continuous honeycomb lattices. *Communications in Partial Differential Equations*, 44:1406 – 1430, 2019.
- [14] Alexis Drouot. Microlocal analysis of the bulk-edge correspondence. *Communications in Mathematical Physics*, 383(3):2069–2112, May 2021.
- [15] Peter Elbau and Gian Michele Graf. Equality of bulk and edge hall conductance revisited. *Communications in Mathematical Physics*, 229:415–432, 2002.
- [16] A. Elgart, G. M. Graf, and J. H. Schenker. Equality of the bulk and edge hall conductances in a mobility gap. *Communications in Mathematical Physics*, 259(1):185–221, Oct 2005.
- [17] Kejie Fang, Zongfu Yu, and Shanhui Fan. Realizing effective magnetic field for photons by controlling the phase of dynamic modulation. *Nature Photonics*, 6:782 – 787, 2012.
- [18] Charles Fefferman, J Lee-Thorp, and Michael I Weinstein. *Topologically protected states in one-dimensional systems*, volume 247. American Mathematical Society, 2017.
- [19] Charles L Fefferman, James P Lee-Thorp, and Michael I Weinstein. Edge states in honeycomb structures. *Annals of PDE*, 2(2):1–80, 2016.
- [20] Charles. L. Fefferman and Michael. I. Weinstein. Honeycomb lattice potentials and dirac points. *Journal of the American Mathematical Society*, 25(4):1169–1220, 2012.

- [21] Sonia Fliss and Patrick Joly. Solutions of the time-harmonic wave equation in periodic waveguides: asymptotic behaviour and radiation condition. *Archive for Rational Mechanics and Analysis*, 219(1):349–386, 2016.
- [22] Gian Michele Graf and Marcello Porta. Bulk-edge correspondence for two-dimensional topological insulators. *Communications in Mathematical Physics*, 324:851–895, 2012.
- [23] Gian Michele Graf and Jacob Shapiro. The bulk-edge correspondence for disordered chiral chains. *Communications in Mathematical Physics*, 363(3):829–846, Nov 2018.
- [24] F. D. M. Haldane and S. Raghu. Possible realization of directional optical waveguides in photonic crystals with broken time-reversal symmetry. *Phys. Rev. Lett.*, 100:013904, Jan 2008.
- [25] M. Z. Hasan and C. L. Kane. Colloquium: Topological insulators. *Rev. Mod. Phys.*, 82:3045–3067, Nov 2010.
- [26] Yasuhiro Hatsugai. Chern number and edge states in the integer quantum hall effect. *Phys. Rev. Lett.*, 71:3697–3700, Nov 1993.
- [27] Yasuhiro Hatsugai. Edge states in the integer quantum hall effect and the riemann surface of the bloch function. *Phys. Rev. B*, 48:11851–11862, Oct 1993.
- [28] R. Jackiw and C. Rebbi. Solitons with fermion number $\frac{1}{2}$. *Phys. Rev. D*, 13:3398–3409, Jun 1976.
- [29] Tosio Kato. *Perturbation theory for linear operators*, volume 132. Springer Science & Business Media, 2013.
- [30] J. KELLENDONK, T. RICHTER, and H. SCHULZ-BALDES. Edge current channels and chern numbers in the integer quantum hall effect. *Reviews in Mathematical Physics*, 14(01):87–119, 2002.
- [31] J. Kellendonk and H. Schulz-Baldes. Boundary maps for c^* -crossed products with with an application to the quantum hall effect. *Communications in Mathematical Physics*, 249(3):611–637, Aug 2004.
- [32] Johannes Kellendonk and Hermann Schulz-Baldes. Quantization of edge currents for continuous magnetic operators. *Journal of Functional Analysis*, 209(2):388–413, 2004.
- [33] Minh Kha, Peter Kuchment, and Andrew Raich. Green’s function asymptotics near the internal edges of spectra of periodic elliptic operators. Spectral gap interior. *J. Spectr. Theory*, 7(4):1171–1233, 2017.
- [34] Alexander B. Khanikaev and Gennady Shvets. Two-dimensional topological photonics. *Nature Photonics*, 11(12):763–773, Dec 2017.
- [35] K. v. Klitzing, G. Dorda, and M. Pepper. New method for high-accuracy determination of the fine-structure constant based on quantized hall resistance. *Phys. Rev. Lett.*, 45:494–497, Aug 1980.
- [36] Yosuke Kubota. Controlled topological phases and bulk-edge correspondence. *Communications in Mathematical Physics*, 349(2):493–525, 2017.
- [37] Peter Kuchment. An overview of periodic elliptic operators. *Bulletin of the American Mathematical Society*, 53(3):343–414, 2016.
- [38] Peter Kuchment and Andrew Raich. Green’s function asymptotics near the internal edges of spectra of periodic elliptic operators. Spectral edge case. *Math. Nachr.*, 285(14-15):1880–1894, 2012.
- [39] James P Lee-Thorp, Michael I Weinstein, and Yi Zhu. Elliptic operators with honeycomb symmetry: Dirac points, edge states and applications to photonic graphene. *Archive for Rational Mechanics and Analysis*, 232(1):1–63, 2019.
- [40] Wei Li, Junshan Lin, Jiayu Qiu, and Hai Zhang. Interface modes in honeycomb topological photonic structures with broken reflection symmetry, 2024.
- [41] Wei Li, Junshan Lin, and Hai Zhang. Dirac points for the honeycomb lattice with impenetrable obstacles. *SIAM Journal on Applied Mathematics*, 83(4):1546–1571, 2023.
- [42] Ling Lu, John D Joannopoulos, and Marin Soljačić. Topological photonics. *Nature photonics*, 8(11):821–829, 2014.

- [43] William Charles Hector McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge university press, 2000.
- [44] Minoru Murata and Tetsuo Tsuchida. Asymptotics of green functions and martin boundaries for elliptic operators with periodic coefficients. *Journal of Differential Equations*, 195(1):82–118, 2003.
- [45] Xiao-Liang Qi, Yong-Shi Wu, and Shou-Cheng Zhang. Topological quantization of the spin hall effect in two-dimensional paramagnetic semiconductors. *Phys. Rev. B*, 74:085308, Aug 2006.
- [46] Jiayu Qiu, Junshan Lin, Peng Xie, and Hai Zhang. Mathematical theory for the interface mode in a waveguide bifurcated from a dirac point, 2023.
- [47] Mikael C. Rechtsman, Julia M. Zeuner, Yonatan Plotnik, Yaakov Lumer, Daniel Podolsky, Felix Dreisow, Stefan Nolte, Mordechai Segev, and Alexander Szameit. Photonic floquet topological insulators. *Nature*, 496(7444):196–200, Apr 2013.
- [48] Jukka Saranen and Gennadi Vainikko. *Periodic integral and pseudodifferential equations with numerical approximation*. Springer Science & Business Media, 2001.
- [49] Alexander V. Sobolev and Jonathan Walthoe. Absolute continuity in periodic waveguides. *Proceedings of the London Mathematical Society*, 85(3):717–741, 2002.
- [50] Amal Taarabt. Equality of bulk and edge hall conductances for continuous magnetic random schrödinger operators, 2014.
- [51] Guo Chuan Thiang and Hai Zhang. Bulk-interface correspondences for one-dimensional topological materials with inversion symmetry. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 479(2270):20220675, 2023.
- [52] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs. Quantized hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.*, 49:405–408, Aug 1982.
- [53] David Vanderbilt. *Berry phases in electronic structure theory: electric polarization, orbital magnetization and topological insulators*. Cambridge University Press, 2018.
- [54] Klaus von Klitzing. The quantized hall effect. *Rev. Mod. Phys.*, 58:519–531, Jul 1986.
- [55] Zheng Wang, Yidong Chong, John Joannopoulos, and Marin Soljacić. Observation of unidirectional backscattering-immune topological electromagnetic states. *Nature*, 461:772–5, 10 2009.

This figure "dirac_cone.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/2405.17200v2>

This figure "quadratic_degenerate_point.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/2405.17200v2>