

THE LIMIT SPACE OF SELF-SIMILAR GROUPS AND SCHREIER GRAPHS

BOZORGMEHR VAZIRI AND FARHAD RAHMATI

ABSTRACT. The present paper investigates the limit G -space \mathcal{J}_G generated by the self-similar action of automatic groups on a regular rooted tree. The limit space \mathcal{J}_G is the Gromov-Hausdorff limit of the family of Schreier graphs Γ_n ; therefore, \mathcal{J}_G can be approximated by Schreier graphs on level n -th when n tends to infinity. We propose a computer program whose code is written in Wolfram language computes the adjacency matrix of Schreier graph Γ_n at each specified level of the regular rooted tree. In this paper, the Schreier graphs corresponding to each automatic group is computed by applying the program to some collection of automata groups, including classic automatic groups.

1. INTRODUCTION

The idea of the boundary at infinity is due to Gromov, who attached to a proper hyperbolic metric space (X, d) a compact metrizable space ∂X whose points are classes of asymptotic geodesic rays in X , reflecting all coarse geometry (i.e., quasi-isometric invariants) of the metric space X . This attachment has played a central role in recent advances in geometry, geometric group theory, ergodic theory, and dynamical systems called *limit space* (*boundary* or *boundary at infinity*). When a discrete group G acts geometrically (co-compactly properly discontinuously by isometry) on the metric space X , besides the geometric structure of the space X , the boundary ∂X also reflects the asymptotic structure of the group G (like the growth rate, the complexity of the word problem and the Hausdorff dimension of the limit space), since the G -action $G \curvearrowright X$ extends to a homeomorphic G -action on the boundary $G \curvearrowright \partial X$. When X is a complete n -dimensional Riemannian manifold of negative sectional curvature, then the boundary ∂X is homeomorphic to the $(n-1)$ -sphere, such that for given a base point $x_0 \in X$, there is a homeomorphism by the map $T_{x_0}X \rightarrow \partial X$ which associates to each tangent unit vector u at the base point the class of the geodesic ray $c : [0, \infty) \rightarrow X$ issues from the base point with velocity vector u . Furthermore, there is a G -invariant measure μ on the boundary called harmonic measure, making ∂X a dynamical system $(\partial X, G, \mu)$.

The limit space associated with a negatively curved metric space possesses impressive rich geometry, topology, and dynamics, which rised the attractions from different realms of mathematics and modern physics in the last two decades. For instance, the author [35] uses ergodic theory at infinity of an arbitrary discrete isometry group Γ acting on any Hadamard manifold \mathbb{H} of pinched variable negative curvature to measure the thickness of the galaxy. In addition, the dynamic of the geodesic flow on a negatively curved manifold consider an ideal chaotic behavior in quantum dynamical systems. The limit space also is used to model the long-time

Key words and phrases. automaton ,Schreier graph, self-similar group, limit space, boundary at infinity, dynamical systems.

behavior of the chaotic physical systems to estimate the thermodynamic equilibrium (see [13],[4],[26]).

The notation of limit space (or asymptotic boundary) in geometry group theory faced a significant advance after impressive works of Gromov [22][21][24], also since 1990 the theory of automatic groups acting on a homogeneous rooted tree and their limit space extensively developed by Grigorchuk, Bartholdi, Nekrashevich, Sidiki, Bondarenko ([5][19][16][18][31] [6][11]) To each self-similar contracting group, [31] associates a limit-space is defined as a quotient of the space $A^{-\omega}$ of the left-infinite words over a finite alphabet A , modulo a G -invariant asymptotic equivalence relation, that can describe using the Moore diagram of the automaton generating the group. This limit space is approximate by a sequence of finite graphs, namely the finite Schreier graphs associated with the automorphism action of the automatic group G on the regular rooted tree A^* . It is shown when G acts by contracting on A^* , then G is a word hyperbolic group by the word metric, and there is an algorithm to decide whether two automatic fractal groups, G_1 and G_2 , are isomorphic subgroups in the automorphism group $Aut(A^*)$. Accordingly, the contracting condition prepares an algorithmic way to make the algebraic structure of automatic groups computable. These additional structures (self-similarity and contracting) provide a significant advantage; it paves the way for bringing the automatic groups and associated limit space \mathcal{J}_G into computer science and computational geometry. Moreover [31] proves the limit space \mathcal{J}_G has finite topological dimension (i.e., complexity) if and only if G has a finite non-empty nucleus (contracting).

Authors [6] explore the connections between automata, groups, limit spaces of self-similar actions, and tilings. In particular, they show how a group acting “nicely” on a tree gives rise to a self-covering of a topological groupoid and how the group can be reconstructed from the groupoid and its covering and The connection is via finite-state automata. V. Nekrashevych and A. Teplyaev [32] investigate the relation between the Analysis of fractals and self-similar Groups. The random walks on a crystal or quasi-crystal are formed by the direct limit of the random walks on a sequence of metric spaces studied by [28][29].

The current research considers the limit space \mathcal{J}_G , where G is a contracting fractal group, and tries to supply a computational mechanism in the form of a computer program to approximate \mathcal{J}_G by its corresponding Schreier graphs. In this direction, the package AutomGrp in the GAP program environment is created by Muntyan and Savchuk [30]. It provides methods or computations with groups and semigroups generated by finite automata or given by wreath recursion, as well as with their finitely generated subgroups and elements. The other is a computer program presented in the Ph.D. thesis [34] written by Perez in 2019 that produces Schreier graphs of automatic groups.

The limit dynamical system (\mathcal{J}_G, s, μ) is naturally used to model chaotic systems with self-similar structures; that is why the limit-space is highly capable of applying to different areas of science. Before anything, we require to approximate the incidence structure of \mathcal{J}_G . Fortunately, the Schreier graphs of a self-similar action (G, A) make this possible. In other words, Schreier graphs provide an algorithmic way to approximate the limit space by an arrangement of finite graphs (see figure 1).

We consider the demand for computational tools and propose a procedure for calculating the adjacency structure of Schreier graphs for the arbitrary n -th level




FIGURE 2. The Schreier graph of the first Grigorchuk group at level 3-th

The asymptotic equivalence relation on the boundary $\mathcal{X}_G := A^{-\omega}$ is determined by the infinite directed paths in the Moor diagram of automata. It is not hard to see that the contracting assumption yields a G -invariant closed asymptotic equivalence relation \sim on the boundary \mathcal{X}_G , making the limit-space \mathcal{J}_G a compact Hausdorff quotient space $\mathcal{X}_G = A^{-\omega} / \sim$ with finite topological dimension $\leq |\mathcal{N}| - 1$ and the embedding

$$\mathcal{J}_G \hookrightarrow \mathbb{R}^{2|\mathcal{N}|-1}$$

Nekrashevich [31] proves that when G is a contractive and self-replacing automatic group, G is generated by the finite nucleus \mathcal{N} , and the limit space \mathcal{J}_G is a path-connected and locally path-connected compact metric space accompanied by the G -bundle \mathcal{X}_G and projecting map $\pi : \mathcal{X}_G \rightarrow \mathcal{J}_G$, that provides a maximal orbifold structure on the limit space. Indeed, the self-similar structure on limit spaces is produced by extended self-similar action of the countable automatic group G on the spherically homogenous rooted tree A^* .

The geometric action of G on the *self-similarity graph* $\Sigma(G, S, X)$ (see definition), which is a proper hyperbolic space demonstrated by [31], implies that the group G to be a word hyperbolic group in the sense of Gromov [23]. Consequently, the limit space \mathcal{J}_G can interpret as the boundary of a proper hyperbolic space $\Sigma(G, S, X)$ with homeomorphism $\partial G \rightarrow \partial \Sigma(G, S, X)$ induced by the quasi-isometry between G and $\Sigma(G, S, X)$. For more detail about boundary of hyperbolic groups see [27]

Example 1.1. [15] The Basilica group is a torsion free weakly branch group G without free subgroups defined by was introduced by R. Grigorchuk and A. Zuk in [20], where they show that it does not belong to the closure of the set of groups of subexponential growth under the operations of group extension and direct limit. L. Bartholdi [7] showed it to be amenable, making Basilica the first example of an amenable but not subexponentially amenable group, and there exists therefore a natural way to associate to it a compact limit space homeomorphic to the well-known Basilica fractal (see 4). Let G be group generated by finite automata with contracting and fractal action on binary tree

$$a = \sigma(b, id), b = (a, id), id = (id, id)$$

It is shown that the *iterated monodromy group* $IMG(z^2 - 1)$ which it's limit G -space is homeomorphic to the Julia set of polynomial $z^2 - 1$ [31].

The self-similar contracting groups appear naturally in studying expanding (partial) self-coverings of topological spaces and orbispaces as their iterated monodromy groups. When the group is the iterated monodromy group of an expanding partial self covering map f , [31] shows the limit space is homeomorphic to the Julia set $J(f)$. Finite Schreier graphs corresponding to the group G , form a sequence of

```
{ (00000000000 - 101010101010, a), (00000000001 - 101010101011, a), (00000000010 - 101010101000, a), (00000000011 - 101010101001, a),
(00000000100 - 101010101100, a), (00000000101 - 101010101101, a), (00000000110 - 101010101110, a), (00000000111 - 101010101111, a),
(00000001000 - 101010100000, a), (00000001001 - 101010100001, a), (00000001010 - 101010100010, a), (00000001011 - 101010100011, a),
(00000001100 - 101010100100, a), (00000001101 - 101010100101, a), (00000001110 - 101010100110, a), (00000001111 - 101010100111, a),
(111111110001 - 111111110001, b), (111111110010 - 111111110010, b), (111111110011 - 111111110011, b), (111111110100 - 111111110100, b), (111111110101 - 111111110101, b),
(111111110110 - 111111110110, b), (111111110111 - 111111110111, b), (111111111000 - 111111111000, b), (111111111001 - 111111111001, b), (111111111010 - 111111111010, b),
(111111111011 - 111111111011, b), (111111111100 - 111111111100, b), (111111111101 - 111111111101, b), (111111111110 - 111111111110, b), (111111111111 - 111111111111, b) }
```

FIGURE 3. The calculation of the adjacency structure of the Basilica group $IMG(z^2 - 1)$ at level 12




FIGURE 4. The Schreier graph of Basilica group $IMG(z^2 - 1)$ at level 12




FIGURE 5. The Schreier graph of the Lamplighter group $\mathbb{Z}_2 \wr \mathbb{Z}$ generated with the wreath recursion $a = \sigma(b, a), b = (b, a)$.

combinatorial approximations to the limit space. If the open conditions satisfy, the limit-space is homeomorphic to a post-critically finite self-similar fractal, being generated by bounded automaton guarantees that the limit space is post-critically finite. Authors [6] explore the connections between automata, groups, limit spaces of self-similar actions, and tilings. In particular, they show how a group acting nicely on a tree gives rise to a self-covering of a topological groupoid and how the

group can be reconstructed from the groupoid and its covering via finite-state automata. V. Nekrashevych and A. Teplyaev [32] investigate the relationship between the Analysis of fractals and self-similar Groups. The random walks on a crystal or quasi-crystal are formed by the direct limit of the random walks on a sequence of metric spaces studied by [28][29]

The structure of the present paper is as follows: The introduction1 briefly describes the theory of contracting self-similar groups acting on a spherically homogeneous rooted tree and their limit G-space. In the second section2, the definitions and known results about the Schreier graphs and limit-space are given. In section 3, the program is applied to the collection of automata and explores the incidence structure of the family of Schreier graphs. The collection contains the first Grigorchuk group, Basilica group, the free group \mathbb{F}_3 of rank 3, Sierpinski group, the lamplighter group mother groups, longe range group, Hanoi tower group, and some automatic groups that are expressed in [16][12]. In the final section, the code written in Wolfram Mathematica language is presented, which is used to construct the Schreier graphs and calculate their adjacency structure.

2. PRELIMINARY

This section reviews the definitions and theorems related to the self-similar groups generated by closed finite-state automata, their Schreier graphs, and the limit space. We have left theorems and propositions without proof, guiding the reader to the references mentioned. The main resources that are used in this section are [5][6][31][32].

2.0.1. *Group generated by automaton.* Let A is a alphabet set, the free monoid A^* with empty set consider as root tree and A^ω denote the *end (boundary)* of rooted tree consist of geodesic paths issuing from a point $x \in A^*$ in the tree.

Definition 2.1. The *automata* Π is a tuple (Q, A, τ, ν) where Q is the set of states and A alphabet set and ;

The *transition* map $\tau : Q \times A \rightarrow Q$ and *output* map $\nu : Q \times A \rightarrow A$ obey the inductive rules:

$$\begin{aligned} q|_\emptyset &= q, q(\emptyset) = \emptyset, \\ q|_{xv} &= q|_x|_v, q(xv) = q(x)q|_x(v), \end{aligned}$$

for $x \in A, v \in A^*$ and $q \in Q$ and $q|_x = \tau(q, x), q(x) = \nu(q, x)$.

By the composition low of automatas, define the automata Π^2 over alphabet set A and state set $Q^2 = Q \times Q$ such that

$$\begin{aligned} (q_1q_2, v) &\mapsto q_2|_{q_1(v)}q_1|_v, \\ (q_1q_2, v) &\mapsto q_1(q_2(v)) \end{aligned}$$

for $v \in A^*$ and $q_1, q_2 \in Q$. However by induction one can obtains automaton Π^* , over alphabet set A with state set $Q^* = \bigsqcup_{\mathbb{N}} Q$. The states Q^* of the automata Π^* with empty set produce semi-group $G = \langle \Pi^* \rangle$.

Note that the transition τ and output ν map can be extended to continuous functions $\hat{\tau} : Q \times A^\omega \rightarrow Q$ and $\hat{\nu} : Q \times A^\omega \rightarrow A^\omega$ respect to the pro-discrete topology.

The automaton Π is *invertible* if, for all $q \in \Pi$, the transformation $\tau(q, \cdot) : A \rightarrow A$ is invertible equivalently q acts by permutation on A . Consequently the states of an invertible automata $\Pi = \Pi^{-1}$ generate a group is called *automata group* denoted

by $G = \langle \Pi \rangle$ which is a subgroup $G \leq \text{Aut}(A^*)$ of the automorphism group of the rooted tree. An automaton Π can be represented by its *Moore diagram*, a directed labeled graph whose vertices are identified with the states of Π . For every state $q \in \Pi$ and every letter $x \in A$, the diagram has an arrow from q to $q|_x$ labeled by $x|q(x)$. This graph contains complete information about the automaton, and we will identify the automaton with its Moore diagram.

The faithful action of a group G on A^* (or on A^ω) is said to be *self-similar* if for every $g \in G$ and every $x \in A$ there exist $h \in G$ and $y \in A$ such that

$$g(x.w) = y.h(w)$$

for every $w \in A^*$ (resp. $w \in A^\omega$). The pair (h, y) is uniquely determined by the pair (g, x) , since the action is faithful. Hence we get an automaton with the set of states G and with the output and transition functions

$$g.x = y.h$$

i.e., $y = g(x)$ and $h = g|_x$. This automaton is called the *complete automaton* of the self-similar action

In other words, the action of a group G on A^* is self-similar if there exists an automaton (G, A) such that the action of $g \in G$ on A^* coincides with the action of the state g of the automaton (G, A) .

The graph of automaton $\mathcal{D}(\Pi)$ is a directed labeled graph with vertices set $V(\mathcal{D}) := Q$ and directed edge $q \rightarrow q|_x$ labeled $x|q(x)$ equipped with *source* map $s : E(\mathcal{D}) \rightarrow V(\mathcal{D})$ and *target* map $t : E(\mathcal{D}) \rightarrow V(\mathcal{D})$. Let Π and Π' be two automaton on a same alphabet A , an *automaton homomorphism* is a map $f : \Pi \rightarrow \Pi'$ such that $f : Q \rightarrow Q$, $\nu'(x, f(q)) = \nu(x, q)$ and $\tau'(x, f(q)) = f(\tau(x, q))$ for all $x \in A, q \in Q$. Furthermore, there is label preserving homomorphism of graphs $\phi : \mathcal{D}(\Pi) \rightarrow \mathcal{D}(\Pi')$ if and only if $\phi : \Pi \rightarrow \Pi'$ is automaton homomorphism.

$$\psi : G \rightarrow \text{Aut}(\mathfrak{M}_G) \cong \text{Sym}(A) \wr G, \psi(g)(m) \in \text{Aut}(\mathfrak{M}), m \in \mathfrak{M}.$$

Definition 2.2. A self-similar action (G, A) is called *contracting* (or *hyperbolic*) if there exists a finite set $\mathcal{N} \subset G$ such that for every $g \in G$ there exists $k \in \mathbb{N}$ such that $g|_v \in \mathcal{N}$ for all words $v \in A^*$ of length $\geq k$. The minimal set with this property is called the *nucleus* of the self-similar action. The nucleus of contracting action is unique and is defined by:

$$\mathcal{N} := \bigcup_{g \in G} \bigcap_{n \geq 0} \{g|_v; v \in A^*, |v| \geq n\}$$

if $g|_v = g$ for some $v \in A^* \setminus \emptyset$, then g belongs to the nucleus by definition.

Lemma 2.3. [31] *A self-similar action of a group with a generating set $S = S^{-1}$, $1 \in S$ is contracting if and only if there exists a finite set \mathcal{N} and a number $k \in \mathbb{N}$ such that*

$$(S \cup \mathcal{N})^2|_{A^k} \subseteq \mathcal{N}$$

The self-similar action of the group G on a rooted tree A^* is spherically transitive and faithful if the action of G on each level A^n is transitive and faithful. Moreover, the action extends to a continuous transitive faithful action on the boundary $\partial \mathcal{T}$ (see [9]).

Definition 2.4. [5][31] A self-similar action is said to be *self-replacing* (recurrent or *fractal*) if it is transitive on the first level A of the tree A^* and the map $g \mapsto g|_x$ from the stabilizer $\text{Stab}_G(x)$ to the group G is surjective for some (every) letter $x \in A$. It can be shown that a self-replicating group acts transitively on X_n for

every $n \geq 1$. It is also easy to see ([31], Proposition 2.11.3) that if a finitely generated contracting group is self-replicating then its nucleus \mathcal{N} is a generating set.

It is easy to see that when the action is recurrent the transivity on the first level A results the transivity on all levels A^n of rooted tree A^* .

Remark 2.5. [31] Even though it is usually challenging to determine whether the automata generate a contracting group. It takes polynomial time to determine the nucleus of self-similar group action.

2.0.2. Schreier graphs.

Definition 2.6. Let G be a group generated by a finite set S and let H be a subgroup of G . The *Schreier graph* $\Gamma(G, S, H)$ of the group G is the graph whose vertices are the right cosets $G/H = \{Hg : g \in G\}$, and two vertices Hg_1 and Hg_2 are adjacent if there exists $s \in S$ such that $g_2 = g_1.s$ or $g_1 = g_2.s$. suppose the finitely generated group G acts on a set M . Then the corresponding *Schreier graph* $\Gamma(G, S, M)$ is the graph with the set of vertices M and set of arrows $S \times M$, where the arrow $(s, v), v \rightarrow s(v)$ starts in v and ends in $s(v)$. The *simplicial Schreier graph* $\bar{\Gamma}(G, S, M)$ remembers only the vertex adjacency:(see figur) its set of vertices is M and two vertices are adjacent if and only if one is an image of another under the action of a generator $s \in S$. If (G, A) is a self-similar action and G is generated by a finite set S , then we get a sequence $\Gamma_n = \Gamma(S, A^n)$ of Schreier graphs of the action of G on the levels A^n of the rooted tree A^* (and the sequence $\bar{\Gamma}_n = \bar{\Gamma}(S, A^n)$ of the respective simplicial Schreier graphs). If (G, A) is generated by a finite automaton $\Pi(Q, A)$, then the graphs $\Gamma(Q, A^n)$ coincide with the dual Moore diagrams of the automata $\Pi'(Q, A)$.

If the action of G is transitive, then the Schreier graph $\Gamma(G, S, M)$ is isomorphic to the Schreier graph $\Gamma(G, S, St_G(x))$ of the group where $St_G(x)$ is the stablizer of some point $x \in A$.

Let G be a self-similar group generated by a finite set S . The sets A^n are invariant under the action of G , and we denote the associated Schreier graphs by $\Gamma_n = \Gamma_n(G, S, A^n)$. For a point $\xi \in A^\omega$ we consider the action of the group G on the G -orbit of ξ , and the associated Schreier graph is called *orbital Schreier graph* denoted $\Gamma_\xi = \Gamma_\xi(G, S)$. For every $\xi \in A^\omega$ we have $St_G(\xi) = \bigcap_{n \geq 1} St_G(\xi_n)$, where ξ_n denotes the prefix of length n of the infinite word ξ . The connected component of the rooted graph (Γ_n, ξ_n) around the root ξ_n is exactly the Schreier graph of G with respect to the stabilizer of ξ_n . It follows immediately that the graphs (Γ_n, ξ_n) converge to the graph $(\partial\Gamma, \xi)$ in the pointed Gromov-Hausdorff topology.

2.0.3. *Limit space and the ends of trees.* The limit space space can be defined in different ways: as a quotient of the Cantor set by an asymptotic equivalence relation (Definition 9.1), as a inverse limit of finite Schreier graphs (Theorem 9.6) or as the boundary of a δ -hyperbolic graph.

Assume G to be an infinite finitely generated group with finite generating set S . There is a natural geometric left action of G on the Cayley graph $\Gamma(G, S)$ equipped with word metric. One can obtain a natural compactification $\bar{\Gamma}(G, S)$ of X by corresponding to X the inverse limit of a system of closed balls in the Cayley graph concentrated on an arbitrary point $x_0 \in \Gamma(G, S)$ with radius r and projection map $p_r : \bar{B}_\Gamma(x_0, r') \rightarrow \bar{B}_\Gamma(x_0, r)$ for $r' \geq r$. There is natural continuous




FIGURE 6. The Schreier graph of $IMG\left(\frac{z-1^2}{z+1}\right) \cong$ group generated with the wreath recursion $a = \sigma(c, c), b = \sigma(b, a), c = (c, c)$.

(respect to the inverse topology) bijection $\phi(x_0) : \bar{\Gamma}(G, S) \rightarrow \varprojlim \bar{B}_\Gamma(x_0, r)$ which induces a topology on $\bar{\Gamma}(G, S)$ sometimes called *cone topology* [14]. The induced topology coincides with the topology of uniform convergence on compact subsets. Thus $\varprojlim \bar{B}_\Gamma(x_0, r)$ is a compactification of the Cayley graph Γ ; it may be viewed as a subspace of the set of continuous maps, which associated to each point $\xi \in \partial\Gamma = \bar{\Gamma} \setminus \Gamma$, a geodesic ray $c : [0, \infty) \rightarrow \Gamma(G, S)$ that issues from x_0 that converge to is referred to *ideal point* or *point at infinity*. That $\partial\Gamma$ is a closed subspace of $\bar{\Gamma}$ and $\bar{\Gamma}$ is compact. Notice that the asymptotic geometry of Cayley graph for group G is independent of the choice of the generator set, so the notation $\partial\Gamma(G, S)$ can relax to ∂G .

Let $G = \langle \Pi \rangle$ be an automatic group generated by the states of an automata over alphabet set A , which acts self-similarly on spherically homogeneous rooted tree A^* . The *ends* (boundary at infinity) of the tree denoted by $A^{-\omega}$ consists of the left infinite sequence of symbols. The group action admits the asymptotic equivalence relation \sim on $A^{-\omega}$. Two left-infinite sequences $\cdots x_2 x_1$ and $\cdots y_2 y_1$ are equivalence if there is left infinite path in the Moor diagram labeled $\cdots (x_2, y_2)(x_1, y_1)$. The limit space of the self-similar action denoted by \mathcal{J}_G is the quotient of the topological space $A^{-\omega}$ by the asymptotic equivalence relation \sim .

H.Bass [9] shows that if an element $g \in \text{Aut}(A^*)$ acts spherically transitively and faithful on each level of the tree, this action extends to a continuous faithful transitive on the boundary of the tree.

Theorem 2.7. [31] *Let G be a finitely generated group with a recurrent contracting action; then the limit G -space \mathcal{J}_G is connected and locally connected.*

One can immediately reach If the action of a finitely generated group is contracting and recurrent; then the limit space is path-connected and locally path-connected.

Despite the homeomorphic action of the group G on the boundary $\partial\Sigma(G, S, X)$, the limit space \mathcal{J}_G is a hyperbolic boundary. For a definition of hyperbolic boundary and more details, the reader is referred to [25][27][14].

Definition 2.8. Let G is a groups with generating set S with self-similar action on A . The *self-similarity graph* $\Sigma(G, S, A)$ is a graph with A^* as vertices set and to vertices $v_1, v_2 \in A^*$ are adjacent if they are connectet either by a vetrical edge if $v_1 = xv_2$ for some $x \in A$, or horizontal edge if $s(v_i) = v_j$, $i, j \in 1, 2$ for some $s \in S$.

The following theorem determine some important aspect of the limit space when the group G acts by contraction on the rooted tree A^* .

Theorem 2.9. [31] *The limit space \mathcal{J}_G of a contracting action of a finitely generated group G is homeomorphic to the hyperbolic boundary $\partial\Sigma(G, S, X)$ of the self-similarity graph $\Sigma(G, S, X)$. Moreover, there exists a homeomorphism $F : \mathcal{J}_G \rightarrow \partial\Sigma(G, S, X)$, such that $D = F \circ \pi$, were $\pi : X^{-\omega} \rightarrow \mathcal{J}_G$ is the canonical projection and $D : X^{-\omega} \rightarrow \partial\Sigma(G, S, X)$ carries every sequence $\cdots x_2x_1 \in X^{-\omega}$ to its limit*

$$\lim_{\infty} x_n \cdots x_2x_1 \in \partial\Sigma(G, S, X)$$

The asymptotic equivalence relation is closed and has finite equivalence classes. The limit space \mathcal{J}_G is a compact metrizable with finite topological dimension $\leq |\mathcal{N}|$.

Example 2.10 (Sierpinski gasket). Let $A = \{0, 1, 2\}$ be the alphabet set and consider the group G generated by wreath recursion:

$$\begin{aligned} a &= \sigma_1(e, a, e), b = \sigma_2(e, e, b), c = \sigma_2(c, e, e), e = (e, e, e) \\ \sigma_1 &= (0, 2), \sigma_2 = (0, 1), \sigma_3 = (1, 2) \end{aligned}$$

where $\sigma_i, i = 1, 2, 3$ are permutation of A .

3. APPROXIMATING WITH THE SCHREIER GRAPHS

In this section, we use the proposed program to explore the incidence structure of Schreier graphs related to groups generated with finite automaton and assemble the approximation of the limit space $\mathcal{J}_{\langle \Pi \rangle}$ by their Schreier graphs. The program is applied to a set of automata which contains the first Grigorchuk group, Basilica group, the free group \mathbb{F}_3 of rank 3, Sierpinski group, the lamplighter group, the mother groups, the longe range group, Hanoi tower group, and some automatic groups that are represented in [16][12].

- 1- The automata by the wreath recursion $a = \sigma(b, c)b = \sigma(c, b)c = (a, a)$ get the free group of rank tree first time introduced by Aleshin The associated limit space has not have a finite Hausdorff dimension (Figure 8) .
- 2- Let $G = \langle \Pi \rangle$ be an automata group generated by Wreath recursion:




FIGURE 7. Sierpinski group action on 3-regular rooted tree at level 8-th

$$a = \sigma(b, c), b = \sigma(c, b), c = (a, a).$$

Aleshin [1] introduced this automaton. [33] has proved that group G is a free group of rank three. The automaton is the smallest automaton among all automata over a 2-letter alphabet, generating a free non-abelian group.




FIGURE 8. The Schreier graph of the \mathbb{F}_3 free group of rank three at level $n = 7, 10$

- 3- The automata 882, produced by the wreath recursion [12]:

$$a = \sigma(c, c), b = (b, c), c = (b, a).$$

The element $(ca^{-1}cb^{-1})^2$ stabilizes the vertex 00 and its section at this vertex is equal to $ca^{-1}cb^{-1}$. Hence, $ca^{-1}cb^{-1}$ has infinite order.

- 4- the case of self-similar actions of the free abelian group \mathbb{Z}^n the limit space $\mathcal{X}_{\mathbb{Z}^n}$ is homeomorphic to \mathbb{R}^n and the tile \mathcal{T} is an integral self-affine tile (Figure 10),

The last recursion defines $IMG(1 - \frac{1}{z^2})$ for more details the reader are referred to[8]. Thus, $\langle \Pi_{878} \rangle \cong C_2 \times IMG(1 - \frac{1}{z^2})$.




FIGURE 11. The simplicial Schreier graph of $IMG(1 - \frac{1}{z^2})$ generated by wreath recursion $a = \sigma(b, b), b = (b, c), c = (b, a)$.

6- Let G be a group generated by a finite state automata with wreath recursion:

$$a = \sigma(b, b), b = (c, b), c = (c, a)$$

The action of group G on the binary tree is conjugated with the half-Basilica group $C_2 \times IMG(z^2 - 1)$ (see figure 12).




FIGURE 12. The Schreier graph generated by automata $a = \sigma(b, b), b = (c, b), c = (c, a)$ isomorphic to the half-Basilica group $C_2 \times IMG(z^2 - 1)$

- 7- [2] The long-range group, an interesting group generate by the wreath recursion $a = (a, b), b = \sigma(b, 1)$. where $\sigma = (01)$ is permutation of $\{0, 1\}$ The




FIGURE 13. The Schreier graph of the long-range group at level 12

Schreier graphs of long-range group (Figure 13) were studied by Benjamini and Hoffman (2005)[10] in the context of long-range percolation theory.

- 8- The *Hanoi towers* group, the group of possible moves in the Hanoi Towers game on three pegs, a game introduced by ‘ Edouard Lucas in 1883. Its level Schreier graphs are discrete Sierpinski gaskets (see[17],[18]).




FIGURE 14. The Schreier graph Hanoi tower group at level 6

- 9- The linear-activity automaton groups is named *Mother group*(Figure 15 is used by [7][2] to prove the amenability of large class of bounded automata groups. The mother group, denoted $\mathfrak{M}_{d,m}$, is defined as the automaton group generated by wreath product [3]:

$$\begin{aligned} a_{k,\sigma} &= \langle a_{k,\sigma}, a_{k-1,\sigma}, 1, \dots, 1 \rangle \text{ for } 0 \leq k \leq d, \\ a_{-1,\sigma} &= \sigma \\ b_{k,\rho} &= \langle b_{k,\rho}, b_{k-1,\rho}, 1, \dots, 1 \rangle \text{ for } 1 \leq k \leq d, \\ b_{0,\rho} &= \langle b_{0,\rho}, 1, \dots, 1 \rangle \end{aligned}$$




FIGURE 15. The Schreier graph of the mother group $\mathfrak{M}_{d,m}$ for $d = 2, m = 2$ at level 12

4. PROGRAM AND CODES

In this section, the program code written in Wolfram language by the first author is presented. to compute the adjacency matrixes of Schreier graphs $\Gamma_n(G, \Pi, A^n)$, where G is the group generated by finite closed state automata Π and $n \in \mathbb{N}$. We approximate limit G -space \mathcal{J}_G . By The comparison of two cases, G be self-similar contracting fractal and self-similar non-contracting fractal groups, in some examples, one can see the vital role of contracting condition influences on limit G -space \mathcal{J}_G to have a charming asymptotic behavior in geometry, topology, and ergodicity.

The computer program introduced in the Ph.D. thesis [34] written by Perez in 2019 generates Schreier graphs of automatic groups with almost six states and 6-letter of the alphabet. Perez's program gives coincident output with the program proposed in present paper while the above conditions are satisfied. In contrast, the algorithm proposed in this paper has no constraint on the number of states or alphabets. However, the code does its task up to the processing capacity. Indeed this code is a part of a program that builds the Schreier graphs. Furthermore, it computes the spectrum of random walks on Schreier graphs on different levels. Note that this paper is devoted only to the first part of a program that produces Schreier graphs.

4.0.1. *automata codes.* The first part of the program is *Input* section conducted by pice of the code that setups the states of automata and the adjacency structure of the Moor diagram. More percisly The *Input* sector of the program gets:

- 1 - A finite set of alphabet A .
- 2 - The adjacency structure of Moor diagram associated to the automaton.
- 4 - The permutation action of each state $q \in Q$ of automata on alphabet set A
- 3 - Integer $n > 0$.

The *Output* give the adjacency matrix of the Schreier graph at n -th level. That is a $|A|^n \times |A|^n$ matrix, such that (i, j) -th entry is an elements of the symbol set $Q \cup \{0\}$, where Q is state set of automata $\Pi(Q, A)$.

for example in following the automata generated by wreath recursion:

$$a = \sigma(b, e), b = (a, e), e = (e, e)$$

```
ClearAll
state[A_, B_] :=
Function[{u, x},
  If[x == 0, Lookup[<|a -> b, b -> a , e -> e|>, u],
  Lookup[{a -> e, b -> e, e -> e}, u]][A, B]
G = Graph[{a -> b, b -> a , e -> e, a -> e, b -> e, e -> e},
  VertexLabels -> "Name"]
S = VertexList[G]
a[x_] := PermutationReplace[x + 1, Cycles[{{1, 2}}]] - 1
b[x_] := PermutationReplace[x, Cycles[{}]]
e[x_] := PermutationReplace[x, Cycles[{}]]
```

If the automata have an identity element by use of the following code should remove otherwise, all vertices are adjacent to a trivial edge, $x \mapsto e.x$.

```
For[i = 1, i <= Dimensions[S][[1]], i++, If[S[[i]] == e, identity = i]]
identity
S = Delete[S, identity]
```

Next, generate the vertices of the rooted tree on a given level; in this example, the vertices of the binary rooted tree at level 12 are generated.

```
generator = Tuples[{0, 1}, 12];
Clear[u, y, st, H]
H = Table[{}, {i, 1, Dimensions[S][[1]]};
For[st = 1, st <= Dimensions[S][[1]], st++,
  For[j = 1; y := {}, j <= Dimensions[generator][[1]], j++,
    For[i = 1; t = S[[st]]; u = {S[[st]]},
      i <= Dimensions[generator][[2]], i++,
      t = state[t, generator[[j, i]]]; u = Append[u, t]; {i, j};
    y = Append[y, u]; H[[st]] = y]
```

The last part is the main part that produces the Schreier graph at a given level.

```
Table[H[[st]][[i, j]][generator[[i, j]]], {st, 1,
  Dimensions[S][[1]]}, {i, 1, Dimensions[generator][[1]]}, {j, 1,
  Dimensions[generator][[2]]};
Schreiergraph =
Flatten[Table[
  Table[{StringJoin[
    Table[ToString[generator[[i, j]]], {j, 1,
      Dimensions[generator][[2]]}] ->
    StringJoin[
      Table[ToString[H[[st]][[i, j]][generator[[i, j]]], {j, 1,
        Dimensions[generator][[2]]}], S[[st]], {i, 1,
        Dimensions[generator][[1]]}], {st, 1, Dimensions[S][[1]]}, 1]
GraphPlot[Schreiergraph, DirectedEdges -> False,
```



```

EdgeRenderingFunction -> Automatic,
VertexRenderingFunction -> ({Black, Text[#2, #1]} &)]
Clear[edgespace]
edgespace = {};
For[st = 1, st <= Dimensions[S][[1]], st++,
  For[j = 1, j <= Dimensions[generator][[1]], j++,
    For[k = 1, k <= Dimensions[generator][[2]], k++,
      If[H[[st]][[j, k]][generator[[j, k]]] - generator[[j, k]] != 0,
        edgespace =
          Append[edgespace, {Table[
            H[[st]][[j, s]][generator[[j, s]]], {s, 1,
              Dimensions[generator][[2]]}],
            Table[generator[[j, s]], {s, 1,
              Dimensions[generator][[2]]}}}}];
edgespace;
ShreierEdge =
  Union[Table[
    edgespace[[x]][[2]] \[DirectedEdge] edgespace[[x]][[1]], {x, 1,
      Dimensions[edgespace][[1]]}]];
simplicialSchreieredge =
  Union[Table[
    StringJoin[
      Table[ToString[edgespace[[x]][[2]][[i]]], {i, 1,
        Dimensions[generator][[2]]}] <->
      StringJoin[
        Table[ToString[edgespace[[x]][[1]][[i]]], {i, 1,
          Dimensions[generator][[2]]}], {x, 1,
        Dimensions[edgespace][[1]]}]];
gr = Graph[
  Table[Labeled[Schreiergraph[[i, 1]], Schreiergraph[[i, 2]]], {i, 1,
    Dimensions[Schreiergraph][[1]]}]]
simplicialSchreieredge = SimpleGraph[simplicialSchreieredge]

```

REFERENCES

1. Stanislav V Aleshin, *Finite automata and burnside's problem for periodic groups*, Mathematical Notes of the Academy of Sciences of the USSR **11** (1972), no. 3, 199–203.
2. Gideon Amir, Omer Angel, and Bálint Virág, *Amenability of linear-activity automaton groups*, Journal of the European Mathematical Society **15** (2013), no. 3, 705–730.
3. Gideon Amir and Bálint Virág, *Positive speed for high-degree automaton groups*, Groups, Geometry, and Dynamics **8** (2014), no. 1, 23–38.
4. Nalini Anantharaman, *Entropy and localization of eigenfunctions*, Séminaire Équations aux dérivées partielles (Polytechnique) dit aussi " Séminaire Goulaouic-Schwartz" (2007), 1–17.
5. Laurent Bartholdi, Rostislav Grigorchuk, and Volodymyr Nekrashevych, *From fractal groups to fractal sets*, Fractals in Graz 2001, Springer, 2003, pp. 25–118.
6. Laurent Bartholdi, André G Henriques, and Volodymyr V Nekrashevych, *Automata, groups, limit spaces, and tilings*, Journal of Algebra **305** (2006), no. 2, 629–663.
7. Laurent Bartholdi, Vadim A Kaimanovich, and Volodymyr V Nekrashevych, *On amenability of automata groups*, Duke Mathematical Journal **154** (2010), no. 3, 575–598.
8. Laurent Bartholdi and Volodymyr Nekrashevych, *Thurston equivalence of topological polynomials*, Acta mathematica **197** (2006), no. 1, 1–51.

9. Hyman Bass, Maria V Otero-Espinar, Daniel Rockmore, and Charles Tresser, *Cyclic renormalization and automorphism groups of rooted trees*, Springer, 2006.
10. Itai Benjamini and Chris Hoffman, *omega-periodic graphs*, arXiv preprint math/0308092 (2003).
11. Ievgen Bondarenko, Daniele D'Angeli, and Tatiana Nagnibeda, *Ends of schreier graphs of self-similar groups*, preprint (2011).
12. Ievgen Bondarenko, Rostislav Grigorchuk, Rostyslav Kravchenko, Yevgen Muntyan, Volodymyr Nekrashevych, Dmytro Savchuk, and Zoran Sunic, *Classification of groups generated by 3-state automata over a 2-letter alphabet*, arXiv preprint arXiv:0803.3555 (2008).
13. Jean Bourgain and Elon Lindenstrauss, *Entropy of quantum limits*, Communications in mathematical physics **233** (2003), no. 1, 153–171.
14. Martin R Bridson and André Haefliger, *Metric spaces of non-positive curvature*, vol. 319, Springer Science & Business Media, 2013.
15. Daniele D'angeli, Alfredo Donno, Michel Matter, and Tatiana Nagnibeda, *Schreier graphs of the basilica group*, arXiv preprint arXiv:0911.2915 (2009).
16. Rostislav Grigorchuk and Dmytro Savchuk, *Self-similar groups acting essentially freely on the boundary of the binary rooted tree*, Group theory, combinatorics, and computing **611** (2014), 9–48.
17. Rostislav Grigorchuk and Zoran Sunic, *Schreier spectrum of the hanoi towers group on three pegs*, arXiv preprint arXiv:0711.0068 (2007).
18. Rostislav Grigorchuk and Zoran Šunik, *Asymptotic aspects of schreier graphs and hanoi towers groups*, Comptes Rendus Mathématique **342** (2006), no. 8, 545–550.
19. Rostislav I Grigorchuk, *Some topics in the dynamics of group actions on rooted trees*, Proceedings of the Steklov Institute of Mathematics **273** (2011), no. 1, 64–175.
20. Rostislav I Grigorchuk and Andrzej Żuk, *On a torsion-free weakly branch group defined by a three state automaton*, International Journal of Algebra and Computation **12** (2002), no. 01n02, 223–246.
21. Michael Gromov, *Groups of polynomial growth and expanding maps (with an appendix by jacques tits)*, Publications Mathématiques de l'IHÉS **53** (1981), 53–78.
22. Mikhael Gromov, *Hyperbolic manifolds, groups and actions*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, NY, 1978), vol. 97, 1980, pp. 183–213.
23. ———, *Hyperbolic groups*, Essays in group theory, Springer, 1987, pp. 75–263.
24. ———, Tech. report, P00001028, 1992.
25. Misha Gromov, *Topological invariants of dynamical systems and spaces of holomorphic maps: I*, Mathematical Physics, Analysis and Geometry **2** (1999), no. 4, 323–415.
26. Sébastien Jezouin, FD Parmentier, A Anthore, U Gennser, A Cavanna, Yong Jin, and F Pierre, *Quantum limit of heat flow across a single electronic channel*, Science **342** (2013), no. 6158, 601–604.
27. Ilya Kapovich and Nadia Benakli, *Boundaries of hyperbolic groups*, arXiv preprint math/0202286 (2002).
28. Jun Kigami, *Analysis on fractals*, no. 143, Cambridge University Press, 2001.
29. Tom Lindstrom and Tom Lindstrøm, *Brownian motion on nested fractals*, no. 420, American Mathematical Soc., 1990.
30. Y. Muntyan and D. Savchuk, *AutomGrp, automata groups, Version 1.3.2*, <https://gap-packages.github.io/automgrp>, Sep 2019, Refereed GAP package.
31. Volodymyr Nekrashevych, *Self-similar groups*, no. 117, American Mathematical Soc., 2005.
32. Volodymyr Nekrashevych and Alexander Teplyaev, *Groups and analysis on fractals*, Analysis on graphs and its applications **77** (2008), 143–180.
33. Volodymyr V Nekrashevych, *Free subgroups in groups acting on rooted trees*, Groups, Geometry, and Dynamics **4** (2010), no. 4, 847–862.
34. Aitor Perez Perez, *Structural and spectral properties of schreier graphs of spinal groups*, Ph.D. thesis, University of Geneva, 2020.
35. Chengbo Yue, *The ergodic theory of discrete isometry groups on manifolds of variable negative curvature*, Transactions of the American Mathematical Society **348** (1996), no. 12, 4965–5005.