

# Synchronization on circles and spheres with nonlinear interactions

Christopher Criscitiello    Quentin Rebjock    Andrew D. McRae    Nicolas Boumal

Institute of Mathematics

EPFL

Lausanne, Switzerland

{christopher.criscitiello, quentin.rebjock,  
andrew.mcrae, nicolas.boumal}@epfl.ch

May 29, 2024

## Abstract

We consider the dynamics of  $n$  points on a sphere in  $\mathbb{R}^d$  ( $d \geq 2$ ) which attract each other according to a function  $\varphi$  of their inner products. When  $\varphi$  is linear ( $\varphi(t) = t$ ), the points converge to a common value (i.e., synchronize) in various connectivity scenarios: this is part of classical work on Kuramoto oscillator networks. When  $\varphi$  is exponential ( $\varphi(t) = e^{\beta t}$ ), these dynamics correspond to a limit of how idealized transformers process data, as described by [Geshkovski et al. \(2024\)](#). Accordingly, they ask whether synchronization occurs for exponential  $\varphi$ .

In the context of consensus for multi-agent control, [Markdahl et al. \(2018\)](#) show that for  $d \geq 3$  (spheres), if the interaction graph is connected and  $\varphi$  is increasing and convex, then the system synchronizes. What is the situation on circles ( $d = 2$ )? First, we show that  $\varphi$  being increasing and convex is no longer sufficient. Then we identify a new condition (that the Taylor coefficients of  $\varphi'$  are decreasing) under which we do have synchronization on the circle. In so doing, we provide some answers to the open problems posed by [Geshkovski et al. \(2024\)](#).

## 1 Introduction

Motivated by open questions [Geshkovski et al. \(2024\)](#) raised in their paper “*A mathematical perspective on transformers*,” (specifically, Problems 4 and 5) we consider gradient flow to maximize<sup>1</sup> the following function defined for  $n$  points on the unit sphere in  $\mathbb{R}^d$ :

$$f(x_1, \dots, x_n) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n w_{ij} \varphi(x_i^\top x_j), \quad \|x_1\| = \dots = \|x_n\| = 1. \quad (\text{P})$$

The real numbers  $w_{ij} \geq 0$  are the weights of a graph ( $w_{ij} = w_{ji}$ ) and the function  $\varphi: [-1, 1] \rightarrow \mathbb{R}$  is twice continuously differentiable. We assume  $\varphi$  is strictly increasing, so that the global maxima correspond to synchronized states:  $x_1 = \dots = x_n$ . The question is: under what conditions does gradient flow reliably converge to such a state?

This is well studied in the *linear* case ( $\varphi(t) = t$ ) as it is equivalent to synchronization of Kuramoto networks of phases ([Kuramoto, 1975](#)) and (by extension) on spheres. Synchronization

---

<sup>1</sup>In contrast, running *negative* gradient flow may or may not take us to a minimum of  $f$ . That problem is also interesting (and quite different) as it leads to configurations of points that are well spread out on the sphere. This is related to Thomson’s problem and Smale’s 7th problem with  $\varphi(t) = -\log(1 - t)$ .

questions also go by the name “consensus” in the context of multi-agent control (Sarlette and Sepulchre, 2009a). From that literature (see Section 2), we expect markedly different behavior between synchronization on circles ( $d = 2$ ) and spheres ( $d \geq 3$ ). The graph structure matters greatly on circles whereas, for higher-dimensional spheres, gradient flow converges to a synchronized state from almost every initial configuration, as long as the graph is connected. The key difference between the two cases is that spheres are simply connected but circles are not (Markdahl, 2021). The results in this paper also reflect this dichotomy.

In the paper of Geshkovski et al. (2024, Problems 4 and 5), problem (P) arises for complete graph with unit weights ( $w_{ij} = 1$  for all  $i, j$ ) and *nonlinear*  $\varphi$  set to be

$$\varphi_\beta(t) = \frac{1}{\beta} e^{\beta t}, \tag{1}$$

where  $\beta > 0$  is an “inverse temperature” in the language of statistical physics. The limit  $\beta \rightarrow 0$  corresponds to the linear case  $\varphi(t) = t$ .

That setting materializes through their study of an idealized model of how input data (tokens) are processed in infinitely deep neural networks with an architecture inspired by transformers. In that context, the interacting particles on the sphere correspond to tokens, and time for the gradient flow corresponds to depth in the network.<sup>2</sup> This step of modelling discrete layers as continuous time variables is in line with previous literature on modeling residual neural networks as neural ODEs (Chen et al., 2018; E, 2017; Haber and Ruthotto, 2017). Neural ODEs to study transformers were first proposed in (Lu et al., 2020; Dutta et al., 2021; Sander et al., 2022).

For random initial configurations, Geshkovski et al. (2024) observed in their setting that the points always converge to a global maximum of  $f$ , namely,  $x_1 = \dots = x_n$ . This is sensible since  $\varphi(x_i^\top x_j)$  is maximal when  $x_i = x_j$ , but it is not a foregone conclusion given that spheres are nonconvex.

Appendix G includes Matlab code for readers who wish to explore the landscape of (P) and the associated dynamical system with various choices of parameters. It requires the Manopt toolbox (Boumal et al., 2014), which is under GNU GPLv3 license.

## Via landscapes

To analyze the typical asymptotic behavior of this dynamical system, it is sufficient to understand the structure of some critical points of  $f$ , rather than tracking the entire dynamics. Indeed, if  $\varphi$  (hence  $f$ ) is real-analytic, then gradient flow converges to a critical point (Łojasiewicz, 1965). Assuming a uniformly random initialization,<sup>3</sup> the Hessian at that critical point is almost surely negative semidefinite owing to the center-stable manifold theorem (Shub, 1987, Thm. III.7, Ex. III.3). (Though classical, this argument is rarely spelled out: see (Geshkovski et al., 2024, App. B) for welcome details.)

Thus, to confirm that gradient flow almost surely converges to a synchronized state, it is sufficient to show that points where the gradient of  $f$  is zero and the Hessian is negative semidefinite are in fact synchronized (in particular, that they are global maxima). The gradient and Hessian are defined with respect to the usual Riemannian metric on the sphere: see Appendix A for explicit formulas.

---

<sup>2</sup>Geshkovski et al. (2024) obtain gradient flow on  $f$  with  $\varphi = \varphi_\beta$  by modeling deep networks composed of self-attention and layer-normalization layers. In the self-attention layers, replacing the exponential of the softmax with  $\varphi'$  gives, in exactly the same way, the gradient flow on  $f$  with *general*  $\varphi$  (the object of our study).

<sup>3</sup>Or any other absolutely continuous probability measure.

## Positive answers, obstructions and remaining open questions

To set the scene, we first state the following theorem to handle  $d \geq 3$ . In particular, it positively answers the questions of [Geshkovski et al. \(2024\)](#) for  $d \geq 3$  (see Corollary 2). This follows directly from ([Markdahl et al., 2018](#), Thm. 13). We give a short proof in Appendix C based on a randomized selection of tangent vectors plugged into the Hessian quadratic form, paralleling the proof in ([McRae and Boumal, 2024](#), §4) for synchronization of rotations.

**Theorem 1** (Spheres, [Markdahl et al. \(2018\)](#), Thm. 13)). *Fix  $n \geq 1$ . Assume*

1. *the ambient dimension  $d$  satisfies  $d \geq 3$ ,*
2. *the undirected graph defined by the weights  $w_{ij} \geq 0$  is connected, and*
3.  *$\varphi'(t) > 0$  and  $\varphi''(t) \geq 0$  for all  $t \in [-1, 1]$ .*

*Then, critical points of  $f$  where the Hessian is negative semidefinite are global maxima of  $f$ . In particular: local maxima are global maxima, they are the synchronized states ( $x_1 = \dots = x_n$ ), and (if  $\varphi$  is real-analytic) gradient flow converges to a synchronized state from almost every initialization.*

**Corollary 2.** *Theorem 1 applies for  $\varphi(t) = t$  and for  $\varphi(t) = \varphi_\beta(t)$  with all  $\beta > 0$ .*

Let us go through the assumptions in Theorem 1. If  $\varphi'$  is not positive ( $\varphi$  not monotonously increasing) there may be global maxima that are not synchronized: see  $\varphi(t) = t^2$  analyzed in Appendix F. If  $\varphi''$  is not nonnegative ( $\varphi$  not convex) there may be non-global local maxima: consider a tetrahedron ( $d = 3, n = 4$ ) with  $\varphi(t) = t^3$ . If the graph is not connected, then the claims of Theorem 1 still apply to each connected component separately.

The assumption  $d \geq 3$  is more interesting—and indeed necessary. As mentioned above, in the linear case ( $\varphi(t) = t$ ) there are counterexamples when  $d = 2$ . However, these are for *incomplete* graphs (the simplest example is a cycle graph; see, e.g., [Townsend et al. \(2020\)](#) for many more). In contrast, the setting in ([Geshkovski et al., 2024](#)) centers on complete graphs.

Remarkably, with nonlinear  $\varphi$  and  $d = 2$ , even a *complete* graph with *unit* weights can harbor spurious local maxima. In Section 5, we construct a *single* function  $\varphi$  which satisfies the assumptions of Theorem 1 yet for which, with  $d = 2$ , the conclusion of that theorem fails for *all*  $n \geq 5$ .

**Theorem 3** (Circles). *Let  $d = 2$ , and consider the complete graph with  $w_{ij} = 1$  for all  $i, j$ . For  $\varepsilon > 0$  and  $\tau \in (-1, 1)$ , define*

$$\varphi_{\varepsilon, \tau}(t) = \varepsilon \log\left(1 + e^{(t-\tau)/\varepsilon}\right),$$

*which is real-analytic and satisfies  $\varphi'_{\varepsilon, \tau}(t), \varphi''_{\varepsilon, \tau}(t) > 0$  for all  $t$ . There exist  $\varepsilon$  and  $\tau$  such that, for all  $n \geq 5$ , gradient flow on (P) with  $\varphi = \varphi_{\varepsilon, \tau}$  and uniformly random initialization converges to a spurious local maximum (not synchronized) with positive probability.*

Intuitively, Theorem 3 relies on the fact that there are functions  $\varphi$  and configurations  $x_1, \dots, x_n$  such that (P) *locally* behaves much like an incomplete graph with linear  $\varphi$  (in particular, a circulant graph in a “twisted state” configuration, which is a well-known source of spurious local maxima in the linear case; see, e.g., [Wiley et al. \(2006\)](#); [Townsend et al. \(2020\)](#)).

Still, Theorem 3 does not exclude the possibility that the landscape is benign for  $d = 2$  when  $\varphi = \varphi_\beta$ .<sup>4</sup> We show that this is indeed true for  $0 < \beta \leq 1$  in Section 4. This follows as a corollary of our next theorem, which requires the Taylor expansion coefficients of  $\varphi'$  to be nonincreasing. That corollary improves on ([Geshkovski et al., 2024](#), Thm. 5.3) which requires  $\beta \lesssim 1/n$ .

<sup>4</sup>As an example, for fixed  $n$ , [Sarlette and Sepulchre \(2009b\)](#), §6) construct a  $\varphi$  such that the  $n$  particles synchronize as long as the graph is connected, but it is different from  $\varphi_\beta$  and it does not work for all  $n$ .

**Theorem 4.** Fix  $d \geq 2$ . Assume  $\varphi'(t) > 0$  and  $\varphi''(t) \geq 0$  for all  $t \in [-1, 1]$ , and also that

$$\varphi'(t) = \sum_{\ell=0}^{\infty} a_{\ell} t^{\ell} \quad \text{for all } t \in [-1, 1] \quad \text{with } a_0 > a_1 \geq a_2 \geq a_3 \geq \dots$$

Let the weights of the graph satisfy  $w_{ij} = q_i q_j$  for some  $q \in \mathbb{R}^n$  with positive entries (in particular, the graph is complete). Then all the same conclusions as in Theorem 1 apply. The assumption  $a_0 > a_1$  can be relaxed to  $a_0 \geq a_1$  if  $\varphi''(t) > 0$  for all  $t \in [-1, 1]$ .

**Corollary 5** (Small  $\beta$ ). Theorem 4 applies for  $\varphi(t) = t$  and for  $\varphi(t) = \varphi_{\beta}(t)$  with  $0 < \beta \leq 1$ . It also applies for  $\varphi(t) = -\log(1 - t + \epsilon)$  for all  $\epsilon > 0$  (Thomson’s problem corresponds to  $\epsilon = 0$ ).

Geshkovski et al. (2024) also argue that synchronization occurs almost surely for  $\beta \gtrsim n^2$ . We repeat their argument in Appendix E with minor changes to handle an arbitrary connected graph  $W$  and more general  $\varphi$ , and to make all quantities explicit.

**Theorem 6.** Fix  $n \geq 1$  and  $d = 2$ . Assume for all  $t \in [-1, 1]$  that  $\varphi'(t) > 0$  and also that

$$t\varphi'(t) - (1 - t^2)\varphi''(t) \geq 0 \quad \implies \quad t \geq \cos\left(\frac{\pi}{n}\right). \quad (2)$$

Assume the graph given by weights  $w_{ij} \geq 0$  is connected. Then all the same conclusions as in Theorem 1 apply. Condition (2) holds in particular when  $\varphi''(t) \geq \frac{n^2}{\pi^2}\varphi'(t)$  for all  $t \in [-1, 1]$ .

**Corollary 7** (large  $\beta$ , Geshkovski et al. (2024, Thm. 5.3)). Fix  $n \geq 1$  and  $d \geq 2$ . Let  $\varphi = \varphi_{\beta}$  with  $\beta \geq \frac{n^2}{\pi^2}$ , and consider a connected graph with weights  $w_{ij} \geq 0$ . Then all the same conclusions as in Theorem 1 apply.

It remains open whether the same holds for values of  $\beta$  between 1 and  $\frac{n^2}{\pi^2}$  when  $d = 2$ .

Finally, we use a different proof in Appendix F to show that the landscape is benign in the quadratic case ( $\varphi(t) = t^2$ ), although the global maxima are synchronized only up to sign.

**Remark 8.** To model transformers, Geshkovski et al. (2024) consider two dynamical systems: SA (for self-attention) and USA. Problem (P) corresponds to USA. In their §3.4 and Remark C.1, they note that the two models correspond to gradient flow on the same energy (namely,  $f$  with  $\varphi = \varphi_{\beta}$ ) only with two different Riemannian metrics on the product of spheres. As the Riemannian metric has no effect on the limit points of gradient flow, our landscape analyses apply directly to both models.

## 2 Related work

Problem (P) is closely connected to the Kuramoto model for a network of coupled oscillators (Kuramoto, 1975; Acebrón et al., 2005), which has deep roots in the dynamical systems literature. The “homogeneous” variant considers the following dynamics for  $n$  time-varying angles  $\theta_1, \dots, \theta_n$ :

$$\dot{\theta}_i = - \sum_{j=1}^n w_{ij} \sin(\theta_i - \theta_j). \quad (3)$$

This is precisely the gradient flow of (P) in the linear case ( $\varphi(t) = t$ ) with  $d = 2$  under the change of variable  $x_i = (\cos \theta_i, \sin \theta_i)$ . A basic question is *which graphs* have the property that the system converges to the synchronized state  $\theta_1 = \dots = \theta_n$  from almost every initial configuration. The

literature is vast: see the references below and the survey by [Dörfler and Bullo \(2014\)](#) (particularly §5). For our purposes, the key results are the following.

For complete graphs and, more generally, sufficiently dense or expander-like graphs, the dynamical system (3) synchronizes from almost every initialization ([Sepulchre et al., 2007](#); [Taylor, 2012](#); [Kassabov et al., 2021](#); [Abdalla et al., 2022](#)). In contrast, for sparse or structured connected graphs, the dynamics (3) have stable equilibria other than the synchronized state (equivalently, (P) with  $d = 2$  and linear  $\varphi$  has spurious local maxima). A rich source of such spurious configurations consists of “twisted states” on a circulant or otherwise ring-like graph ([Wiley et al., 2006](#); [Canale and Monzón, 2015](#); [Townsend et al., 2020](#); [Yoneda et al., 2021](#)). One can also construct more exotic counterexamples, including graphs with manifolds of stable equilibria of arbitrary dimension ([Sclosa, 2023](#)).

In higher dimensions ( $d \geq 3$ ), less attention has been given to this problem. Relevant works for synchronization on spheres include ([Olfati-Saber, 2006](#); [Li and Spong, 2014](#); [Caponigro et al., 2015](#); [Lageman and Sun, 2016](#)), though none of these guarantee synchronization almost surely. The key work for us is by [Markdahl et al. \(2018\)](#), who show that a broad class of consensus algorithms on the sphere succeed as long as the interaction graph is connected. Hence, synchronization/consensus on spheres is fundamentally simpler than on circles. One may also entertain synchronization on more general manifolds ([Sarlette and Sepulchre, 2009a](#); [Markdahl, 2021](#)). The rotation groups are of particular interest in applications (and they constitute another way to generalize circles).

If we minimize rather than maximize  $f$ , we obtain a packing problem. These have been extensively studied in the literature (e.g., [Cohn and Kumar \(2007\)](#)). Packing on the circle or sphere is closely related to Smale’s 7th problem and Thomson’s problem, which ask for the minimal energy configurations of charges constrained to lie on a circle or sphere. Among this line of work, most relevant to us is the work of [Cohn \(1960\)](#), who considers Thomson’s problem on the circle. Using Morse theory, [Cohn \(1960\)](#) completely characterizes the minima and critical points of  $f$ , and their signatures, when  $d = 2$  and  $\varphi(t) = -\log(1 - t)$  (and for other similar  $\varphi$ ). It is unclear to us how to apply the techniques of [Cohn \(1960\)](#) to the present setting: crucially with  $\varphi(t) = -\log(1 - t)$ , the Hessian of  $f$  on the circle corresponds to a Laplacian with all *nonpositive weights*. This is a substantial departure from our setting: see the open questions in Section 6.

### 3 Riemannian geometry tools and optimality conditions

Endow  $\mathbb{R}^d$  with the inner product  $\langle u, v \rangle = u^\top v$ . The unit sphere  $\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : \|x\| = 1\}$  is a Riemannian submanifold of  $\mathbb{R}^d$ : the tangent space  $\mathbb{T}_x \mathbb{S}^{d-1} = \{\dot{x} \in \mathbb{R}^d : x^\top \dot{x} = 0\}$  inherits the Euclidean inner product as a subspace of  $\mathbb{R}^d$ . Formally,  $f$  in (P) is defined on the product manifold

$$\mathcal{M} = (\mathbb{S}^{d-1})^n = \{X \in \mathbb{R}^{d \times n} : \text{diag}(X^\top X) = \mathbf{1}\}$$

with the product Riemannian structure. In this matrix notation, the points  $x_1, \dots, x_n$  are arranged as the columns of  $X$ ,  $\text{diag}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^n$  extracts the diagonal of a matrix, and  $\mathbf{1} \in \mathbb{R}^n$  is the all-ones vector (sometimes denoted  $\mathbf{1}_n$  if we want to emphasize the dimension). The cost function is

$$f(X) = \frac{1}{2} \langle W, \varphi(X^\top X) \rangle,$$

where  $\langle A, B \rangle = \text{Tr}(A^\top B)$  is the Frobenius inner product, and  $\varphi$  applies entrywise ( $\varphi(A)_{ij} = \varphi(a_{ij})$ ). The symmetric matrix  $W \in \mathbb{R}^{n \times n}$  holds the graph weights  $w_{ij} \geq 0$ .

Based on these choices, we can derive expressions for the Riemannian gradient and Hessian of  $f: \mathcal{M} \rightarrow \mathbb{R}$ , and deduce necessary optimality conditions for (P). These are standard computations: see Appendix A.

Since our results in Theorems 3 and 4 only require proofs for  $d = 2$ , we only spell out the conditions for that case here. This is simpler in part because the tangent space of a circle is one-dimensional, so that the Riemannian Hessian can be expressed as an ordinary  $n \times n$  matrix.

The Riemannian Hessian for  $d = 2$  exhibits a Laplacian structure, defined as follows. Given a symmetric matrix  $M \in \mathbb{R}^{n \times n}$ , the Laplacian of the associated graph (where we think of  $m_{ij}$  as the weight between nodes  $i$  and  $j$ ) is

$$L(M) = \text{diag}(M\mathbf{1}) - M, \quad (4)$$

where  $\text{diag}: \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$  forms a diagonal matrix. As a quadratic form, it is well known that  $\alpha^\top L(M)\alpha = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n m_{ij}(\alpha_i - \alpha_j)^2$ . In particular, if the weights  $m_{ij}$  are nonnegative, then  $L(M) \succeq 0$ . If, furthermore, the graph is connected, then  $\ker L(M) = \text{span}(\mathbf{1})$ .

We now characterize second-order critical points. In the following,  $\text{ddiag}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$  sets all off-diagonal entries of a matrix to zero,  $\odot$  denotes the entrywise (Hadamard) matrix product, and  $M \mapsto M^{\odot 2}$  denotes entrywise squaring.

**Lemma 9.** *Let  $d = 2$ . Assume  $\varphi$  is twice continuously differentiable. The eigenvalues of the Hessian of  $f$  at  $X \in \mathcal{M}$  are equal to the eigenvalues of  $-L(M)$ , where  $L(M)$  is the Laplacian (4) for the graph with weights*

$$m_{ij} = w_{ij}h(x_i^\top x_j) \quad \text{where} \quad h(t) = t\varphi'(t) - (1 - t^2)\varphi''(t). \quad (5)$$

*The Riemannian gradient at  $X$  is zero (i.e.,  $X$  is critical) and the Riemannian Hessian at  $X$  is negative semidefinite if and only if*

$$X\text{ddiag}(X^\top X A) = X A \quad \text{and} \quad \text{ddiag}(X^\top X A) - A \odot X^\top X \succeq L(K), \quad (6)$$

$$\text{where} \quad A = W \odot \varphi'(X^\top X) \quad \text{and} \quad K = W \odot \varphi''(X^\top X) \odot (\mathbf{1}\mathbf{1}^\top - (X^\top X)^{\odot 2}). \quad (7)$$

## 4 A sufficient condition for synchronization on the circle: Theorem 4

Given  $q \in \mathbb{R}^n$  with positive entries, define the complete graph  $W = qq^\top$ . (The notation  $q$  echoes classical work where each  $x_i$  is a particle with charge  $q_i$  and  $f$  is their associated potential (Cohn, 1960).) From Theorem 1, we know that in dimension  $d \geq 3$  it is sufficient for  $\varphi$  to be a strictly increasing convex function to rule out spurious local maxima for  $f$ . From Theorem 3, we also know that this is not sufficient when  $d = 2$ . Yet, experimentally, even with  $d = 2$  we do not know of spurious maximizers when  $\varphi = \varphi_\beta$  (1). Thus, it appears that  $\varphi_\beta$  has additional favorable properties.

In this section, we show as much for  $\beta$  up to 1. Specifically, we prove Theorem 4. Our argument relies crucially on the *Schur product theorem* (Horn and Johnson, 1991, Thm. 5.2.1).

To prove Theorem 1 for spheres (see Appendix C), we chose a *random* direction  $\gamma$ , and then moved all points  $x_i$  in that same direction  $\gamma$  using  $\hat{x}_i = (I - x_i x_i^\top)\gamma$ . This approach works for spheres, but falls short when applied to circles. It is natural to try to choose the common direction  $\gamma$  more purposefully. Albeit indirectly, the proof below moves all the points in the direction of their *weighted mean*  $\gamma = Xq = \sum_{i=1}^n q_i x_i$ . See Remark 11 at the end of this section.

We start with a lemma showing that we are done if we can show  $x_1, \dots, x_n$  lie in a common (closed) hemisphere (we also use this in the proof of Theorem 6). This strengthens existing results for an *open* hemisphere by Markdahl et al. (2018, Prop. 12) and, in their setting, Geshkovski et al. (2024, Lem. 4.2). The rank-deficient case (which is treated separately in the proof) resembles familiar results for Burer–Monteiro relaxations (see, for example, Journée et al. (2010, Thm. 7)), but the standard arguments used in that literature do not directly apply because we are *maximizing* a convex function (and not minimizing).

**Lemma 10.** *Assume  $\varphi'(t) > 0$  for all  $t \in [-1, 1]$  and that the graph with weights  $w_{ij} \geq 0$  is connected. Let  $X$  be a critical point of  $f$  where the Hessian is negative semidefinite. If there exists a nonzero  $v \in \mathbb{R}^d$  such that  $X^\top v \geq 0$  (entrywise; that is, the points  $x_1, \dots, x_n$  lie in a common closed hemisphere), then  $X$  is a global maximum. Such a vector  $v$  exists if  $\text{rank}(X) < d$ .*

See Appendix B for a proof. With this, we can prove the main result of this section.

*Proof of Theorem 4.* Fix  $d = 2$  (the case  $d \geq 3$  holds by Theorem 1). Lemma 9 provides

$$X \text{ddiag}(X^\top X A) = X A \quad \text{and} \quad \text{ddiag}(X^\top X A) - A \odot X^\top X \succeq L(K),$$

where  $L(K)$  is the Laplacian (4) of the graph with weights  $K_{ij} = w_{ij} \varphi''(x_i^\top x_j) (1 - (x_i^\top x_j)^2)$  and  $A = W \odot \varphi'(X^\top X)$  (7). Since  $\varphi'' \geq 0$ , it follows that  $K_{ij} \geq 0$  and so  $L(K) \succeq 0$ .

Multiply the matrix inequality left and right by  $X$  and  $X^\top$  to compress it to a  $2 \times 2$  matrix inequality. Crucially, this allows us to use the first-order condition:

$$\begin{aligned} X L(K) X^\top &\preceq X \left( \text{ddiag}(X^\top X A) - A \odot X^\top X \right) X^\top \\ &= X (A - A \odot X^\top X) X^\top \\ &= X (A \odot (\mathbf{1}\mathbf{1}^\top - X^\top X)) X^\top \\ &= X \left( W \odot \varphi'(X^\top X) \odot (\mathbf{1}\mathbf{1}^\top - X^\top X) \right) X^\top. \end{aligned} \tag{8}$$

By the assumption on  $\varphi'$ , it holds that

$$\varphi'(t)(1-t) = a_0 - (a_0 - a_1)t - (a_1 - a_2)t^2 - \dots$$

for all  $t \in [-1, 1]$ , with  $a_\ell - a_{\ell+1} \geq 0$  for all  $\ell \geq 0$ . In particular,

$$\varphi'(X^\top X) \odot (\mathbf{1}\mathbf{1}^\top - X^\top X) = a_0 \mathbf{1}\mathbf{1}^\top - (a_0 - a_1) X^\top X - (a_1 - a_2) (X^\top X)^{\odot 2} - \dots$$

Plugging this back into (8) yields

$$a_0 X W X^\top \succeq X L(K) X^\top + X \left( (a_0 - a_1) W \odot X^\top X + (a_1 - a_2) W \odot (X^\top X)^{\odot 2} + \dots \right) X^\top.$$

The Schur product theorem states that  $M, N \succeq 0 \implies M \odot N \succeq 0$ . Using  $X^\top X \succeq 0$  and the assumption that  $W = qq^\top \succeq 0$  plus the assumption that  $a_1 - a_2 \geq 0, a_2 - a_3 \geq 0$  etc., we deduce

$$a_0 X W X^\top \succeq (a_0 - a_1) X (W \odot X^\top X) X^\top + X L(K) X^\top.$$

Since  $W = qq^\top$ , we also have  $W \odot X^\top X = \text{diag}(q) X^\top X \text{diag}(q)$ . With  $Z = X \text{diag}(q)^{1/2}$ , the above can be restated as:

$$a_0 (Xq)(Xq)^\top \succeq (a_0 - a_1) (ZZ^\top)^2 + X L(K) X^\top \succeq 0. \tag{9}$$

The matrix on the left-hand side has rank 1. The matrix in the middle is positive semidefinite since  $a_0 - a_1 \geq 0$  and  $L(K) \succeq 0$ .

If  $a_0 > a_1$ , then the matrix in the middle of inequality (9) must have rank equal to  $\text{rank}(Z) = \text{rank}(X)$ , using  $q_i > 0$  for all  $i$ . The result then follows, because inequality (9) therefore implies  $\text{rank}(X) < 2$ , and Lemma 10 handles  $\text{rank}(X) < 2$ .

For  $a_0 = a_1$ , we assume  $\varphi'' > 0$  and take a closer look at  $L(K)$ . Since  $w_{ij} = q_i q_j > 0$  and  $\varphi''(x_i^\top x_j) > 0$ , it follows that  $K_{ij} = w_{ij} \varphi''(x_i^\top x_j) (1 - (x_i^\top x_j)^2) = 0$  if and only if  $x_i^\top x_j = \pm 1$ . We consider two cases.

First, suppose  $\dim \ker(L(K)) > 1$ . This means the graph of  $K$  is disconnected. For  $i, j$  in different connected components,  $K_{ij} = 0$ , so  $x_i = \pm x_j$ . Since  $K$  has at least two nonempty connected components, this implies  $x_i = \pm x_j$  for *all*  $i, j$  (simply observe that each  $x_k$  is in a component that is different from that of  $x_i$  or  $x_j$ ). Thus,  $\text{rank}(X) = 1$  and the result again follows from Lemma 10.

Second, suppose  $\ker(L(K)) = \text{span}(\mathbf{1})$ . Choose a unit  $v \in \mathbb{R}^2$  orthogonal to  $Xq$ . Then (9) implies

$$0 \geq (X^\top v)^\top L(K) (X^\top v) \geq 0.$$

Then we must have  $X^\top v \in \text{span}(\mathbf{1})$ . In particular, negating  $v$  if necessary, we have  $X^\top v \geq 0$  entrywise, so the result once again follows from Lemma 10.  $\square$

**Remark 11.** *The proof above moves all points in the direction of their weighted mean  $Xq$ . To see this, note that all cases at the end of the proof are handled (explicitly or implicitly) by hitting (9) with an appropriately chosen vector  $v$  proportional to  $JXq$ , where  $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ . One can verify (see the proof of Lemma 9 in Appendix A) that this translates to tangent vectors  $\dot{x}_i = (I - x_i x_i^\top) Xq$ .*

## 5 Obstacles to synchronization on the circle: Theorem 3

The circle ( $d = 2$ ) stands out among “spheres”  $\mathbb{S}^{d-1}$  in that it is not simply connected: the circle itself (a closed loop) cannot be continuously collapsed into a single point while staying on the circle, whereas, say, the equator can be collapsed to a single point on a sphere. Accordingly, to construct non-synchronizing scenarios on the circle, it makes sense to entertain configurations of points that “go around” the circle. The simplest such configuration is when all points  $x_1, \dots, x_n$  lie on a regular  $n$ -gon as follows (see Figure 1):

$$x_i = (\cos \theta_i, \sin \theta_i) \quad \text{with} \quad \theta_i = 2\pi \frac{i-1}{n} \quad \text{for} \quad i = 1, \dots, n. \quad (10)$$

In this section we prove Theorem 3: when  $d = 2$  there exists a real-analytic  $\varphi$  with  $\varphi' > 0$ ,  $\varphi'' > 0$  such that  $f(X) = \frac{1}{2} \langle \mathbf{1}_n \mathbf{1}_n^\top, \varphi(X^\top X) \rangle$  has a spurious local maximum for all  $n \geq 5$ . Let us sketch the proof. As a first step, we show that for *each*  $n$  there is a  $\varphi_n$  such that the regular  $n$ -gon is a spurious local maximum of  $f$  with  $\varphi = \varphi_n$  (Lemma 13 below); this follows from a well-known result for linear synchronization on the circle (Lemma 12). In order to build a single  $\varphi$  which works for all  $n$  as in Theorem 3, we then fix some integer  $m$  (e.g.,  $m = 5$ ) and distribute  $n$  points on the regular  $m$ -gon with roughly  $n/m$  points at each vertex. This configuration may not be critical for  $f$  with  $\varphi = \varphi_m$  (e.g., if  $n$  is not a multiple of  $m$ ), but we argue that it is close to a spurious critical point (and hence that one exists) provided  $n$  is large enough (combine Lemmas 14 and 15 below). Finally, we exhibit a  $\varphi$  that covers all  $n \geq 5$ . Now let us proceed to the full proof.

For the linear synchronization problem on the circle ( $\varphi(t) = t, d = 2$ ), it is well known that the regular  $n$ -gon configuration (10) (with large enough  $n$ ) is a spurious local maximum when the weight matrix  $W$  corresponds to a (circulant) nearest-neighbor graph in which every node is connected to at most 68% of its nearest neighbors. This observation is due to Wiley et al. (2006).

In our setting, we can see this as taking the complete graph  $W = \mathbf{1}\mathbf{1}^\top$  and the ReLU-type function

$$\varphi(t) = \max\{0, t - \tau\} \quad (11)$$






Figure 1: A regular  $n$ -gon on the circle ( $n = 13, d = 2$ ). In Section 5, we construct the function  $\varphi$  by smoothing a ReLU such that  $\varphi'(\cos(\theta_i - \theta_j))$  is about 1 when  $\cos(\theta_i - \theta_j) > \tau$  and about 0 otherwise. Points are color-coded by  $\varphi'(\cos(\theta_i - 0))$ .

with  $\tau$  at least  $\cos(0.68\pi)$ : see Figure 1. This is valid for large  $n$ . To handle all  $n \geq 5$ , we reduce connectivity and require  $\tau \geq \cos(.50\pi) = 0$ . We also require  $\tau < \cos(\frac{2\pi}{5})$  to ensure that each point interacts with at least its two nearest neighbors. The following lemma can be checked with formulas by Ling et al. (2019, §5.3), who obtain the eigenvalues of the Hessian at the regular  $n$ -gon in closed form using a Fourier transform.

**Lemma 12.** [Wiley et al. (2006), Ling et al. (2019, §5.3)] Given  $n \geq 5$ , let  $x_1, \dots, x_n$  lie on a regular  $n$ -gon (10). Fix  $\tau \in [0, \cos(\frac{2\pi}{5})]$  such that  $\tau \notin \{\cos(2\pi\frac{i-1}{n}) : i = 1, \dots, n\}$ .<sup>5</sup> With  $\varphi$  as in (11) and  $W = \mathbf{1}_n \mathbf{1}_n^\top$ , the point  $X$  is critical for  $f$ , and  $\text{Hess}f(X)$  is negative<sup>6</sup> definite.

The next step is simple: we smooth the ReLU (11) and, by continuity, the regular  $n$ -gon remains a spurious local maximum. In order to ensure  $\varphi$  is real-analytic, we apply the log-sum-exp smoothing

$$\varphi_{\varepsilon, \tau}(t) = \varepsilon \log(1 + e^{(t-\tau)/\varepsilon}) \quad (12)$$

with parameter  $\varepsilon > 0$ .

**Lemma 13.** Let  $n \geq 5$  and fix  $\tau$  as in Lemma 12. There exists  $\varepsilon_{n, \tau} > 0$  such that, for all  $\varepsilon \in (0, \varepsilon_{n, \tau}]$ , the regular  $n$ -gon configuration  $X$  is critical for  $f$  with  $W = \mathbf{1}_n \mathbf{1}_n^\top$  and  $\varphi = \varphi_{\varepsilon, \tau}$  (12). Moreover,  $\text{Hess}f(X)$  is negative<sup>6</sup> definite.

*Proof of Lemma 13.* From Lemma 9, one can easily verify by symmetry that the regular  $n$ -gon  $X$  is critical for  $f$  for any  $\varphi$  such that  $f$  is differentiable in a neighborhood of  $X$ . We now study  $\text{Hess}f_{\varepsilon, \tau}(X)$ , which is well defined (even at  $\varepsilon = 0$ ) since we assume  $\tau \neq \cos(2\pi\frac{i-1}{n})$  for  $i = 1, \dots, n$ . Lemma 12 gives that  $\text{Hess}f_{0, \tau}(X)$  is negative<sup>6</sup> definite. Hence, it is enough to show that the curve  $\varepsilon \mapsto \text{Hess}f_{\varepsilon, \tau}(X)$  is continuous from the right at  $\varepsilon = 0$ . We can then conclude using the continuity of eigenvalues. (More formally, one should remove the trivial zero eigenvalue of the Hessian by fixing one of the points, or passing to the quotient, and then applying this argument.) The function

<sup>5</sup>This condition ensures that none of the inner products  $x_i^\top x_j$  for the  $n$ -gon lie on the kink of the ReLU.

<sup>6</sup>Throughout Section 5, “negative definite Hessian” means all eigenvalues of the Hessian are negative except for the single zero eigenvalue which appears due to the global rotation symmetry of the problem.

$\varphi$  only appears in  $\text{Hess}f(X)$  through  $\varphi'$  and  $\varphi''$  (see Lemma 16 in Appendix A). Therefore, it is enough to show that for any fixed  $t \neq \tau$ , both

$$\varepsilon \mapsto \varphi'_{\varepsilon,\tau}(t) = 1 - (1 + e^{(t-\tau)/\varepsilon})^{-1} \quad \text{and} \quad \varepsilon \mapsto \varphi''_{\varepsilon,\tau}(t) = (2\varepsilon + 2\varepsilon \cosh((t-\tau)/\varepsilon))^{-1}$$

are continuous from the right at  $\varepsilon = 0$ . This is readily apparent from their expressions.  $\square$

Lemma 13 does not give a  $\varphi$  which works for all  $n$  as in Theorem 3 because, as  $n$  grows, it seems that  $\varepsilon_{n,\tau}$  must decrease to zero. However, we can circumvent this issue by the following observation. Fix  $m \geq 5$  and let  $n = Nm$  be an integer multiple of  $m$ . Consider the configuration of points on the circle where  $N = n/m$  points are placed at the vertices of a regular  $m$ -gon—we call this configuration the “repeated  $m$ -gon.” As the  $m$ -gon is a spurious local maximum for  $f$  on  $m$  points with  $\varphi = \varphi_{\varepsilon_m,\tau}$ , it stands to reason that the repeated  $m$ -gon should be a spurious local maximum on  $n$  points with  $\varphi = \varphi_{\varepsilon_m,\tau}$  (when  $n$  is a multiple of  $m$ ). This is indeed true. We give a proof of the following (more general) statement in Appendix D. (Allowing arbitrary  $q$ , as opposed to just a multiple of  $\mathbf{1}$ , shall be useful for handling the cases where  $n$  is not a multiple of  $m$ .)

**Lemma 14.** *Assume  $\varphi'(1) > 0$ . Let  $q \in \mathbb{R}^m$  have positive integer entries and  $n = q_1 + \dots + q_m$ . Given  $X = (x_1, \dots, x_m) \in \mathbb{R}^{2 \times m}$ , define*

$$\tilde{X} = \underbrace{(x_1, \dots, x_1)}_{q_1 \text{ times}}, \dots, \underbrace{(x_m, \dots, x_m)}_{q_m \text{ times}} \in \mathbb{R}^{2 \times n}. \quad (13)$$

*If  $X$  is critical with negative<sup>6</sup> definite Hessian for  $f_m(X) = \frac{1}{2}\langle qq^\top, \varphi(X^\top X) \rangle$ , then  $\tilde{X}$  is critical with negative<sup>6</sup> definite Hessian for  $f_n(\tilde{X}) = \frac{1}{2}\langle \mathbf{1}_n \mathbf{1}_n^\top, \varphi(\tilde{X}^\top \tilde{X}) \rangle$ .*

For fixed  $m \geq 5$ , this shows that  $f$  has a spurious local maximum with  $\varphi = \varphi_{\varepsilon_m,\tau}$  for  $n$  points, where  $n$  is a multiple of  $m$  (and  $\tau$  is chosen as in Lemma 12 with  $n$  replaced by  $m$ ). To handle a number of points  $n = Nm + r$  that is not a multiple of  $m$  (i.e.,  $r \in [1, m-1]$ ), first place  $N$  points on each vertex of the regular  $m$ -gon, then distribute the extraneous  $r$  points arbitrarily on the  $m$  vertices. This configuration is unlikely to be a critical point; however, it is close to one: if the total number of points is sufficiently large, the few extra points should only cause a minor perturbation from what was a (strict) spurious local maximum.

In Lemma 15, this intuition is made rigorous via *the implicit function theorem*. In the language of Lemma 14, we let  $X$  be the regular  $m$ -gon, and we take each  $q_i$  to be an integer close to  $N$ ; consequently  $\frac{1}{N}q$  is close to  $\mathbf{1}_m$ . We rescale by  $\frac{1}{N}$  so that  $q$  itself becomes close  $\mathbf{1}_m$  (this scaling does not affect the landscape of  $f$ ). The proof is in Appendix D.

**Lemma 15.** *Fix  $m \geq 5$  odd. There exists  $\delta > 0$  such that if  $\varepsilon, \tau$  and  $q \in \mathbb{R}^m$  satisfy*

$$\|q - \mathbf{1}_m\| \leq \delta, \quad 0 \leq \varepsilon \leq \delta \quad \text{and} \quad -\delta \leq \tau \leq \delta,$$

*then  $f(X) = \frac{1}{2}\langle qq^\top, \varphi_{\varepsilon,\tau}(X^\top X) \rangle$  has a spurious critical point with negative<sup>6</sup> definite Hessian.*

We can now combine Lemmas 13, 14 and 15 to prove Theorem 3.

*Proof of Theorem 3.* Fix  $m = 5$ . Write  $n = Nm + r$  with remainder  $r \in [0, m-1]$ . Let  $q = (1 + r/N, 1, 1, \dots, 1)^\top \in \mathbb{R}^m$ . Note that if  $n$  is sufficiently large then  $q$  is arbitrarily close to  $\mathbf{1}_m$ . Invoke Lemma 15 to select  $\delta > 0$ . Set  $n_0 \geq 5$  such that  $\|q - \mathbf{1}_m\| \leq \delta$  for all  $n > n_0$ . If

$$n > n_0, \quad 0 \leq \varepsilon \leq \delta, \quad \text{and} \quad -\delta \leq \tau \leq \delta, \quad (14)$$

then  $f(X) = \frac{1}{2}\langle qq^\top, \varphi_{\varepsilon, \tau}(X^\top X) \rangle$  has a spurious critical point with negative<sup>6</sup> definite Hessian: call it  $X_{q, \varepsilon, \tau}$ .

Let  $\tilde{X}_{q, \varepsilon, \tau}$  be the configuration of  $n$  points where  $Nq_i$  points are placed at point  $i$  of  $X_{q, \varepsilon, \tau}$ , as in equation (13). By Lemma 14 and our choice of  $q$ , this configuration on  $n$  points is a spurious local maximum for  $f(X) = \frac{1}{2}\langle \mathbf{1}_n \mathbf{1}_n^\top, \varphi_{\varepsilon, \tau}(X^\top X) \rangle$ . In other words, we have shown that for all  $n, \varepsilon, \tau$  satisfying (14),  $f$  has a spurious local maximum with  $\varphi = \varphi_{\varepsilon, \tau}$  and the graph  $W = \mathbf{1}_n \mathbf{1}_n^\top$ .

It remains to choose  $\varepsilon$  and  $\tau$  to accommodate  $m \leq n \leq n_0$ . Select any  $\tau \in [0, \min\{\delta, \cos(\frac{2\pi}{5})\})$  such that  $\tau \neq \cos(2\pi \frac{i-1}{n})$  for all integers  $n \in [m, n_0]$  and  $i \in [1, n]$ . Further define

$$\varepsilon = \min\{\delta, \varepsilon_{m, \tau}, \varepsilon_{m+1, \tau}, \dots, \varepsilon_{n_0, \tau}\}$$

where  $\varepsilon_{n, \tau}$  are provided by Lemma 13. Then, by Lemma 13, we conclude that  $f$  with  $\varphi = \varphi_{\varepsilon, \tau}$  has a spurious local maximum for all  $n \geq m$ .  $\square$

## 6 Conclusions and perspectives

We conclude with a list of open questions for  $d = 2$ ,  $\varphi = \varphi_\beta$  and  $W = \mathbf{1}\mathbf{1}^\top$  (complete graph).

- **(All  $\beta$ )** Do we have synchronization for  $\beta \in (1, \frac{n^2}{\pi^2})$ ?
- **(Minimal configurations)** Is the regular  $n$ -gon the minimal configuration of  $f$ ? Cohn (1960) shows that this is true for  $\varphi(t) = -\log(1-t)$ . On the other hand, Theorem 3 shows that there exist nice  $\varphi$  for which the  $n$ -gon is not minimal.
- **(Critical configurations)** We say a configuration consists of *clusters* if there exist  $i \neq j$  such that  $x_i = x_j$ , i.e., at least two points overlap. The number of clusters is the number of distinct values of the points  $x_i$ . Do all critical configurations of  $f$ , apart from the regular  $n$ -gon, consist of clusters? Numerically this appears to be the case. Cohn (1960) shows that this is true for  $\varphi(t) = -\log(1-t)$ . Note that for  $d \geq 3$  there do exist fully non-clustered critical configurations which are not minimal (e.g., an  $n$ -gon on the equator of a sphere).
- **(Signatures of critical configurations)** The signature of a critical configuration is the number of positive eigenvalues of the Hessian at that configuration. Is the signature of every critical configuration at least the number of clusters minus one? Numerically, this appears to be true. Cohn (1960) shows that for  $\varphi(t) = -\log(1-t)$  the signature always *equals* the number of clusters minus one—this is not true in general for  $\varphi = \varphi_\beta$  (e.g., for  $n = 9, \beta = 1$ , and a critical configuration with four clusters of sizes 1, 2, 2, 4).

By Lemma 18 (which is used to prove Lemma 14 above), to show that the signature is at least the number of clusters minus one, it is enough to show the following: if  $q$  has positive entries and  $X$  is a critical configuration of  $f(X) = \frac{1}{2}\langle qq^\top, \varphi_\beta(X^\top X) \rangle$  with distinct points, then all eigenvalues of  $\text{Hess}f(X)$  are positive (except for the single zero eigenvalue).

- **(Dynamics of gradient flow)** Geshkovski et al. (2024) are not only interested in the asymptotic behavior of gradient flow on  $f$  but also its dynamics. Numerically, gradient flow on  $f$  often slows down near saddles consisting of only a few clusters (so-called meta-stable states (Cohn, 1960; Erber and Hockney, 1991)). Also, gradient flow seems to jump between such saddles (akin to saddle-to-saddle dynamics, see (Jacot et al., 2022; Berthier, 2023; Pesme and Flammarion, 2023) and references therein). Can one characterize these dynamics? A first step would be to compute the maximal eigenvalue of the Hessian at clustered critical configurations, as this eigenvalue controls the escape time of gradient flow. How does that eigenvalue depend on the number of clusters?

## Acknowledgments

We thank Pedro Abdalla, Afonso Bandeira and Cyril Letrouit for insightful discussions.

This work was supported by the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number MB22.00027.

## References

- P. Abdalla, A. S. Bandeira, M. Kassabov, V. Souza, S. H. Strogatz, and A. Townsend. Expander graphs are globally synchronising. Oct. 2022.
- P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, Princeton, NJ, 2008. ISBN 978-0-691-13298-3.
- J. A. Acebrón, L. L. Bonilla, C. J. Pérez Vicente, F. Ritort, and R. Spigler. The Kuramoto model: A simple paradigm for synchronization phenomena. *Reviews of Modern Physics*, 77(137):137–185, 2005. doi:[10.1103/revmodphys.77.137](https://doi.org/10.1103/revmodphys.77.137).
- R. Berthier. Incremental learning in diagonal linear networks. *Journal of Machine Learning Research*, 24(171):1–26, 2023. URL <http://jmlr.org/papers/v24/22-1395.html>.
- N. Boumal. *An introduction to optimization on smooth manifolds*. Cambridge University Press, 2023. doi:[10.1017/9781009166164](https://doi.org/10.1017/9781009166164). URL <https://www.nicolasboumal.net/book>.
- N. Boumal, B. Mishra, P.-A. Absil, and R. Sepulchre. Manopt, a Matlab toolbox for optimization on manifolds. *Journal of Machine Learning Research*, 15(42):1455–1459, 2014. URL <https://www.manopt.org>.
- E. A. Canale and P. Monzón. Exotic equilibria of Harary graphs and a new minimum degree lower bound for synchronization. *Chaos*, 25(2), 2015. doi:[10.1063/1.4907952](https://doi.org/10.1063/1.4907952).
- M. Caponigro, A. Chiara Lai, and B. Piccoli. A nonlinear model of opinion formation on the sphere. *Discrete & Continuous Dynamical Systems*, 35(9):4241–4268, 2015. doi:[10.3934/dcds.2015.35.4241](https://doi.org/10.3934/dcds.2015.35.4241).
- R. T. Q. Chen, Y. Rubanova, J. Bettencourt, and D. K. Duvenaud. Neural ordinary differential equations. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL [https://proceedings.neurips.cc/paper\\_files/paper/2018/file/69386f6bb1dfed68692a24c8686939b9-Paper.pdf](https://proceedings.neurips.cc/paper_files/paper/2018/file/69386f6bb1dfed68692a24c8686939b9-Paper.pdf).
- H. Cohn. Global equilibrium theory of charges on a circle. *The American Mathematical Monthly*, 67(4):338–343, 1960. doi:[10.1080/00029890.1960.11989502](https://doi.org/10.1080/00029890.1960.11989502).
- H. Cohn and A. Kumar. Universally optimal distribution of points on spheres. *Journal of the American Mathematical Society*, 20(1):99–148, 2007.
- F. Dörfler and F. Bullo. Synchronization in complex networks of phase oscillators: A survey. *Automatica*, 50(6):1539–1564, 2014. ISSN 0005-1098. doi:[10.1016/j.automatica.2014.04.012](https://doi.org/10.1016/j.automatica.2014.04.012).

- S. Dutta, T. Gautam, S. Chakrabarti, and T. Chakraborty. Redesigning the transformer architecture with insights from multi-particle dynamical systems. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 5531–5544. Curran Associates, Inc., 2021. URL [https://proceedings.neurips.cc/paper\\_files/paper/2021/file/2bd388f731f26312bfc0fe30da009595-Paper.pdf](https://proceedings.neurips.cc/paper_files/paper/2021/file/2bd388f731f26312bfc0fe30da009595-Paper.pdf).
- W. E. A proposal on machine learning via dynamical systems. *Communications in Mathematics and Statistics*, 5(1):1–11, March 2017. doi:[10.1007/s40304-017-0103-z](https://doi.org/10.1007/s40304-017-0103-z).
- T. Erber and G. Hockney. Equilibrium configurations of N equal charges on a sphere. *Journal of Physics A: Mathematical and General*, 24(23):L1369–L1377, 1991. doi:[10.1088/0305-4470/24/23/008](https://doi.org/10.1088/0305-4470/24/23/008).
- B. Geshkovski, C. Letrouit, Y. Polyanskiy, and P. Rigollet. A mathematical perspective on transformers. *arXiv preprint arXiv:2312.10794*, 2024.
- E. Haber and L. Ruthotto. Stable architectures for deep neural networks. *Inverse Problems*, 34(1):014004, dec 2017. doi:[10.1088/1361-6420/aa9a90](https://doi.org/10.1088/1361-6420/aa9a90).
- R. Horn and C. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1991.
- A. Jacot, F. Ged, B. Şimşek, C. Hongler, and F. Gabriel. Saddle-to-saddle dynamics in deep linear networks: Small initialization training, symmetry, and sparsity. *arXiv preprint arXiv:2106.15933*, 2022. URL <https://arxiv.org/abs/2106.15933>.
- M. Journée, F. Bach, P.-A. Absil, and R. Sepulchre. Low-rank optimization on the cone of positive semidefinite matrices. *SIAM Journal on Optimization*, 20(5):2327–2351, 2010. doi:[10.1137/080731359](https://doi.org/10.1137/080731359).
- M. Kassabov, S. H. Strogatz, and A. Townsend. Sufficiently dense Kuramoto networks are globally synchronizing. *Chaos*, 31(7), 2021. doi:[10.1063/5.0057659](https://doi.org/10.1063/5.0057659).
- S. Krantz and H. Parks. *The Implicit Function Theorem*. Springer New York, 2013. doi:[10.1007/978-1-4614-5981-1](https://doi.org/10.1007/978-1-4614-5981-1).
- Y. Kuramoto. Self-entrainment of a population of coupled non-linear oscillators. In *Proceedings of the International Symposium on Mathematical Problems in Theoretical Physics*, pages 420–422, Kyoto, Japan, Jan. 1975. doi:[10.1007/bfb0013365](https://doi.org/10.1007/bfb0013365).
- C. Lageman and Z. Sun. Consensus on spheres: Convergence analysis and perturbation theory. In *Proceedings of the IEEE 55th Conference on Decision and Control (CDC)*, Las Vegas, NV, USA, Dec. 2016. doi:[10.1109/cdc.2016.7798240](https://doi.org/10.1109/cdc.2016.7798240).
- W. Li and M. W. Spong. Unified cooperative control of multiple agents on a sphere for different spherical patterns. *IEEE Transactions on Automatic Control*, 59(5):1283–1289, 2014. doi:[10.1109/tac.2013.2286897](https://doi.org/10.1109/tac.2013.2286897).
- S. Ling, R. Xu, and A. S. Bandeira. On the landscape of synchronization networks: A perspective from nonconvex optimization. *SIAM Journal on Optimization*, 29(3):1879–1907, 2019. doi:[10.1137/18m1217644](https://doi.org/10.1137/18m1217644).
- S. Łojasiewicz. Ensembles semi-analytiques. *Lecture Notes IHES (Bures-sur-Yvette)*, 1965.

- Y. Lu, Z. Li, D. He, Z. Sun, B. Dong, T. Qin, L. Wang, and T.-Y. Liu. Understanding and improving transformer from a multi-particle dynamic system point of view. *International Conference on Learning Representations 2020 Workshop on Integration of Deep Neural Models and Differential Equations*, *arXiv 1906.02762*, 2020.
- J. Markdahl. Synchronization on Riemannian manifolds: Multiply connected implies multistable. *IEEE Transactions on Automatic Control*, 66(9):4311–4318, 2021. doi:[10.1109/tac.2020.3030849](https://doi.org/10.1109/tac.2020.3030849).
- J. Markdahl, J. Thunberg, and J. Goncalves. Almost global consensus on the  $n$ -sphere. *IEEE Transactions on Automatic Control*, 63(6):1664–1675, 2018. doi:[10.1109/tac.2017.2752799](https://doi.org/10.1109/tac.2017.2752799).
- A. McRae and N. Boumal. Benign landscapes of low-dimensional relaxations for orthogonal synchronization on general graphs. *SIAM Journal on Optimization*, 34(2):1427–1454, 2024. doi:[10.1137/23M1584642](https://doi.org/10.1137/23M1584642).
- R. Olfati-Saber. Swarms on sphere: A programmable swarm with synchronous behaviors like oscillator networks. In *Proceedings of the 45th IEEE Conference on Decision and Control*, San Diego, CA, USA, Dec. 2006. doi:[10.1109/cdc.2006.376811](https://doi.org/10.1109/cdc.2006.376811).
- S. Pesme and N. Flammarion. Saddle-to-saddle dynamics in diagonal linear networks. In A. Oh, T. Naumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 7475–7505. Curran Associates, Inc., 2023. URL [https://proceedings.neurips.cc/paper\\_files/paper/2023/file/17a9ab4190289f0e1504bbb98d1d111a-Paper-Conference.pdf](https://proceedings.neurips.cc/paper_files/paper/2023/file/17a9ab4190289f0e1504bbb98d1d111a-Paper-Conference.pdf).
- M. E. Sander, P. Ablin, M. Blondel, and G. Peyré. Sinkformers: Transformers with doubly stochastic attention. In G. Camps-Valls, F. J. R. Ruiz, and I. Valera, editors, *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pages 3515–3530. PMLR, 28–30 Mar 2022. URL <https://proceedings.mlr.press/v151/sander22a.html>.
- A. Sarlette and R. Sepulchre. Consensus optimization on manifolds. *SIAM Journal on Control and Optimization*, 48(1):56–76, 2009a.
- A. Sarlette and R. Sepulchre. Synchronization on the circle. *arXiv preprint arXiv:0901.2408*, 2009b.
- D. Sclosa. Kuramoto networks with infinitely many stable equilibria. *SIAM Journal on Applied Dynamical Systems*, 22(4):3267–3283, 2023. doi:[10.1137/23m155400x](https://doi.org/10.1137/23m155400x).
- R. Sepulchre, D. A. Paley, and N. E. Leonard. Stabilization of planar collective motion: All-to-all communication. *IEEE Transactions on Automatic Control*, 52(5):811–824, 2007. doi:[10.1109/tac.2007.898077](https://doi.org/10.1109/tac.2007.898077).
- M. Shub. *Global Stability of Dynamical Systems*. Springer New York, 1987. doi:[10.1007/978-1-4757-1947-5](https://doi.org/10.1007/978-1-4757-1947-5).
- R. Taylor. There is no non-zero stable fixed point for dense networks in the homogeneous Kuramoto model. *Journal of Physics A: Mathematical and Theoretical*, 45(5):055102, 2012. doi:[10.1088/1751-8113/45/5/055102](https://doi.org/10.1088/1751-8113/45/5/055102).
- A. Townsend, M. Stillman, and S. H. Strogatz. Dense networks that do not synchronize and sparse ones that do. *Chaos*, 30(8), 2020. doi:[10.1063/5.0018322](https://doi.org/10.1063/5.0018322).

D. A. Wiley, S. H. Strogatz, and M. Girvan. The size of the sync basin. *Chaos*, 16, 2006. doi:10.1063/1.2165594.

R. Yoneda, T. Tatsukawa, and J. Teramae. The lower bound of the network connectivity guaranteeing in-phase synchronization. *Chaos*, 31(6), 2021. doi:10.1063/5.0054271.

## A Riemannian Gradient and Hessian for general $d$

In this section, we give additional Riemannian geometry background and derive the gradient and Hessian of problem (P) for general  $d \geq 2$ . From this, we can derive the simplified criticality conditions for  $d = 2$  as they appear in Lemma 9 in Section 3. See, for example, (Absil et al., 2008; Boumal, 2023) for details of such computations in the context of Riemannian optimization.

For convenience, we repeat some definitions from Section 3. We endow  $\mathbb{R}^d$  with the inner product  $\langle u, v \rangle = u^\top v$ . The unit sphere  $\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : \|x\| = 1\}$  is a Riemannian submanifold of  $\mathbb{R}^d$ : the tangent space  $\mathbb{T}_x \mathbb{S}^{d-1} = \{\dot{x} \in \mathbb{R}^d : x^\top \dot{x} = 0\}$  inherits the Euclidean inner product as a subspace of  $\mathbb{R}^d$ . Formally,  $f$  is defined on the product manifold

$$\mathcal{M} = (\mathbb{S}^{d-1})^n = \{X \in \mathbb{R}^{d \times n} : \text{diag}(X^\top X) = \mathbf{1}\}$$

with the product Riemannian structure. In this matrix notation, the points  $x_1, \dots, x_n$  are arranged as the columns of  $X$ ,  $\text{diag}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^n$  extracts the diagonal of a matrix,  $\mathbf{1} \in \mathbb{R}^n$  is the all-ones vector, and

$$f(X) = \frac{1}{2} \langle W, \varphi(X^\top X) \rangle,$$

where  $\langle A, B \rangle = \text{Tr}(A^\top B)$  is the Frobenius inner product and  $\varphi$  applies entrywise:  $\varphi(A)_{ij} = \varphi(a_{ij})$ . The symmetric matrix  $W \in \mathbb{R}^{n \times n}$  holds the graph weights  $w_{ij} \geq 0$ .

An important object for studying  $\mathcal{M}$  is the orthogonal projection from  $\mathbb{R}^{d \times n}$  to the tangent space

$$\mathbb{T}_X \mathcal{M} = \{\dot{X} \in \mathbb{R}^{d \times n} : \text{diag}(X^\top \dot{X}) = 0\}, \quad (15)$$

which has the following formula:

$$\text{Proj}_X(Z) = Z - X \text{ddiag}(X^\top Z), \quad (16)$$

where  $\text{ddiag}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$  sets all off-diagonal entries of a matrix to zero. Indeed, the  $i$ th column of  $\text{Proj}_X(Z)$  is  $z_i - (x_i^\top z_i)x_i$ , which is the projection of  $z_i$  to the tangent space  $\mathbb{T}_{x_i} \mathbb{S}^{d-1}$ .

With this, we can give the general formulas for the gradient and Hessian of  $f$  on  $\mathcal{M}$ :

**Lemma 16.** *Assume  $\varphi$  is twice continuously differentiable. Given  $X \in \mathcal{M}$ , let*

$$S = \text{ddiag}(X^\top X A) - A \quad \text{with} \quad A = W \odot \varphi'(X^\top X), \quad (17)$$

where  $\odot$  is the entrywise (Hadamard) product and  $\varphi'$  applies entrywise. The Riemannian gradient of  $f: \mathcal{M} \rightarrow \mathbb{R}$  at  $X \in \mathcal{M}$  is given by

$$\text{grad} f(X) = -XS. \quad (18)$$

The Riemannian Hessian of  $f$  at  $X$  is a self-adjoint linear map defined by the quadratic form

$$\langle \dot{X}, \text{Hess} f(X)[\dot{X}] \rangle = -\langle \dot{X}^\top \dot{X}, S \rangle + \frac{1}{2} \langle (X^\top \dot{X} + \dot{X}^\top X)^{\odot 2}, W \odot \varphi''(X^\top X) \rangle \quad (19)$$

for all  $\dot{X}$  in the tangent space  $\mathbb{T}_X \mathcal{M}$  (15), where  $M \mapsto M^{\odot 2}$  denotes entrywise squaring.

*Proof.* Given  $\dot{X} \in T_X \mathcal{M}$ , one calculates that the directional derivative of  $f$  at  $X$  along  $\dot{X}$  is

$$\begin{aligned} Df(X)[\dot{X}] &= \frac{1}{2} \left\langle W, \varphi'(X^\top X) \odot (\dot{X}^\top X + X^\top \dot{X}) \right\rangle \\ &= \left\langle X \left( W \odot \varphi'(X^\top X) \right), \dot{X} \right\rangle = \langle XA, \dot{X} \rangle. \end{aligned}$$

Thus, the Riemannian gradient is the orthogonal projection (16) of  $XA$  to  $T_X \mathcal{M}$  (Boumal, 2023, Prop. 3.61), that is,

$$\text{grad}f(X) = \text{Proj}_X(XA) = XA - X \text{ddiag}(X^\top XA) = -XS.$$

The directional derivative of  $\text{grad}f(X)$  along  $\dot{X} \in T_X \mathcal{M}$  is given by

$$D\text{grad}f(X)[\dot{X}] = -\dot{X}S - X\dot{S},$$

where  $\dot{S}$  is the derivative of  $S$  (17) at  $X$  along  $\dot{X}$ . More explicitly,  $\dot{S} = \dot{D} - \dot{A}$ , where  $\dot{D}$  is diagonal and  $\dot{A}$  is the derivative of  $A$  (17) at  $X$  along  $\dot{X}$ . The Riemannian Hessian at  $X$  as a quadratic form is given by (Boumal, 2023, Cor. 5.16)

$$\begin{aligned} \langle \dot{X}, \text{Hess}f(X)[\dot{X}] \rangle &= \langle \dot{X}, D\text{grad}f(X)[\dot{X}] \rangle \\ &= -\langle \dot{X}^\top \dot{X}, S \rangle - \langle X^\top \dot{X}, \dot{S} \rangle \\ &= -\langle \dot{X}^\top \dot{X}, S \rangle + \langle X^\top \dot{X}, \dot{A} \rangle, \end{aligned}$$

where the last equality uses  $\langle X^\top \dot{X}, \dot{D} \rangle = 0$ , which follows from the fact that  $\text{diag}(X^\top \dot{X}) = 0$  for any tangent vector  $\dot{X}$ .

Finally, note that  $\dot{A} = W \odot \varphi''(X^\top X) \odot (X^\top \dot{X} + \dot{X}^\top X)$ . This is symmetric, so  $\langle X^\top \dot{X}, \dot{A} \rangle = \frac{1}{2} \langle X^\top \dot{X} + \dot{X}^\top X, \dot{A} \rangle$ . Thus we obtain (19).  $\square$

In Sections 4 and 5, we are primarily interested in the case  $d = 2$ . It is then convenient to particularize Lemma 16 to  $d = 2$ , where the Hessian has a simpler matrix form. Using Lemma 16, let us prove Lemma 9.

*Proof of Lemma 9.* Start from Lemma 16 and recall the definitions  $A = W \odot \varphi'(X^\top X)$  and  $S = \text{ddiag}(X^\top XA) - A$  (17). The first-order condition is  $0 = \text{grad}f(X) = -XS = XA - X \text{ddiag}(X^\top XA)$ , as stated in (6).

For the second-order condition, we consider the eigenvalues of the Hessian quadratic form (repeated here from (19)):

$$\langle \dot{X}, \text{Hess}f(X)[\dot{X}] \rangle = -\langle \dot{X}^\top \dot{X}, S \rangle + \frac{1}{2} \left\langle (X^\top \dot{X} + \dot{X}^\top X)^{\odot 2}, W \odot \varphi''(X^\top X) \right\rangle. \quad (20)$$

With  $d = 2$ , all tangent vectors  $\dot{X} \in T_X \mathcal{M}$  (15) are of the form

$$\dot{X} = JX \text{diag}(\alpha) \quad \text{with} \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \alpha \in \mathbb{R}^n.$$

This is because  $Jx_i$  rotates  $x_i$  by  $\pi/2$ , making it a basis for the tangent space to the circle at  $x_i$ . This corresponds to expanding tangent vectors in an orthonormal basis with coordinates  $\alpha$ , which has no effect on eigenvalues of quadratic forms, so we now express (20) in terms of  $\alpha$ .



Using  $\dot{X}^\top \dot{X} = \text{diag}(\alpha)X^\top X \text{diag}(\alpha) = (X^\top X) \odot \alpha \alpha^\top$  and also  $\text{diag}(X^\top X) = \mathbf{1}$ , we compute

$$\langle S, \dot{X}^\top \dot{X} \rangle = \langle S, (X^\top X) \odot \alpha \alpha^\top \rangle = \alpha^\top (S \odot X^\top X) \alpha = \alpha^\top \left( \text{ddiag}(X^\top X A) - A \odot X^\top X \right) \alpha.$$

Since  $A$  and  $X^\top X$  are symmetric, it holds that  $\text{diag}(X^\top X A) = (A \odot X^\top X) \mathbf{1}$ . Thus, we have found  $\langle S, \dot{X}^\top \dot{X} \rangle = \alpha^\top L(A \odot X^\top X) \alpha$  where  $L$  is the graph Laplacian (4).

Furthermore,  $(X^\top \dot{X})_{ij} = \alpha_j x_i^\top J x_j$  and  $(\dot{X}^\top X)_{ij} = \alpha_i (J x_i)^\top x_j = -\alpha_i x_i^\top J x_j$  so that

$$(X^\top \dot{X} + \dot{X}^\top X)_{ij} = (\alpha_j - \alpha_i) x_i^\top J x_j.$$

Since  $J$  rotates vectors in  $\mathbb{R}^2$  by  $\pi/2$ , it is easy to check that  $x_i^\top J x_j$  is the sine of the angle between  $x_i$  and  $x_j$ , whereas  $x_i^\top x_j$  is the cosine of that angle. It then follows from  $\sin(\theta)^2 = 1 - \cos(\theta)^2$  that  $(x_i^\top J x_j)^2 = 1 - (x_i^\top x_j)^2$ . Thus,

$$\frac{1}{2} \left\langle (X^\top \dot{X} + \dot{X}^\top X)^{\odot 2}, W \odot \varphi''(X^\top X) \right\rangle = \frac{1}{2} \sum_{i,j} w_{ij} \varphi''(x_i^\top x_j) (1 - (x_i^\top x_j)^2) (\alpha_i - \alpha_j)^2.$$

From the comments after eq. (4), we recognize a Laplacian structure. We have

$$\frac{1}{2} \left\langle (X^\top \dot{X} + \dot{X}^\top X)^{\odot 2}, W \odot \varphi''(X^\top X) \right\rangle = \alpha^\top L(K) \alpha,$$

with weights  $K_{ij} = w_{ij} \varphi''(x_i^\top x_j) (1 - (x_i^\top x_j)^2)$  as in (7). Overall, we found

$$-\langle \dot{X}, \text{Hess}f(X)[\dot{X}] \rangle = \alpha^\top \left( \text{ddiag}(X^\top X A) - A \odot X^\top X \right) \alpha - \alpha^\top L(K) \alpha \quad (21)$$

$$= \alpha^\top L(A \odot X^\top X - K) \alpha = \alpha^\top L(M) \alpha. \quad (22)$$

The Hessian quadratic form is negative semidefinite if and only if the right-hand side quadratic form (21) is nonnegative for all  $\alpha \in \mathbb{R}^n$ , which is equivalent to the matrix inequality in (6). From (22), we also see that the eigenvalues of  $\text{Hess}f(X)$  equal those of  $-L(M)$ .  $\square$

## B Benign landscape in a hemisphere

In this section, we provide a proof of Lemma 10, which we use in the proofs of Theorems 4 and 6. Markdahl et al. (2018, Prop. 12) (and Geshkovski et al. (2024, Lem. 4.2) in their setting) showed that if all the points  $x_1, \dots, x_n$  lie in a common *open* hemisphere, then first-order criticality implies global optimality (this is classical for linear  $\varphi$ ). We improve this slightly to allow a general connected graph and to allow the points to lie in a *closed* hemisphere by using second-order conditions.

*Proof of Lemma 10.* Because  $\varphi'$  is positive,  $X$  is a global maximum if and only if  $x_1 = \dots = x_n$ . Since  $X$  is critical, Lemma 16 provides  $XA - X \text{ddiag}(X^\top X A) = 0$ , where  $A = W \odot \varphi'(X^\top X)$ . (That is, the gradient at  $X$  is zero.) Transposing and multiplying by  $v$  yields, entrywise,

$$0 = (AX^\top v)_i - (X^\top X A)_{ii} (X^\top v)_i = \sum_{j=1}^n w_{ij} \varphi'(x_i^\top x_j) \left( (x_j^\top v) - (x_i^\top x_j) (x_i^\top v) \right) \quad \text{for all } i.$$

Select  $i$  such that  $x_i^\top v \geq 0$  is smallest among  $\{x_1^\top v, \dots, x_n^\top v\}$ . Then each term in the sum is nonnegative and therefore must be zero. Choose  $j$  distinct from  $i$  such that  $w_{ij} > 0$  (which exists since the graph is connected). By the assumption  $\varphi'(x_i^\top x_j) > 0$ , we obtain

$$x_i^\top v \leq x_j^\top v = (x_i^\top x_j) (x_i^\top v) \leq x_i^\top v.$$

Thus, the inequalities are equalities.

If  $x_i^\top v > 0$ , we deduce  $x_i^\top x_j = 1$  (using  $(x_i^\top x_j)(x_i^\top v) = x_i^\top v$ ), and so  $x_i = x_j$ . Using that  $x_j^\top v = x_i^\top v$  is minimal among  $\{x_1^\top v, \dots, x_n^\top v\}$ , we can repeat this argument across a spanning tree of positive weights to conclude that  $x_1 = \dots = x_n$ .

Otherwise, if  $x_i^\top v = 0$ , then  $x_j^\top v = 0$  as well (since  $x_i^\top v = x_j^\top v$ ). Repeat this argument across a spanning tree of positive weights to deduce that  $X^\top v = 0$ . Thus  $\text{rank}(X) < d$ .

To handle this rank-deficient case, we use second-order conditions, namely, that the Riemannian Hessian is negative semidefinite. Assume, without loss of generality, that  $v$  is unit-norm (we still have  $X^\top v = 0$ ). Set  $\dot{X} = v\mathbf{1}^\top \in \mathbb{R}^{d \times n}$ , and note that  $X^\top \dot{X} = 0$  (which guarantees that  $\dot{X}$  is a valid tangent vector) and that  $\dot{X}^\top \dot{X} = \mathbf{1}\mathbf{1}^\top$ .

With the Hessian as given in Lemma 16, we have, again using the fact that  $X^\top \dot{X} = 0$ ,

$$\begin{aligned} 0 &\geq \langle \dot{X}, \text{Hess}f(X)[\dot{X}] \rangle = -\langle \dot{X}^\top \dot{X}, S \rangle \\ &= \langle \mathbf{1}\mathbf{1}^\top, A - \text{ddiag}(X^\top X A) \rangle = \sum_{i,j=1}^n w_{ij} \varphi'(x_i^\top x_j) (1 - x_i^\top x_j), \end{aligned}$$

where  $S = \text{ddiag}(X^\top X A) - A$  and, again,  $A = W \odot \varphi'(X^\top X)$  as given in (17). The sum is over nonnegative terms, hence they must each be equal to zero. For  $i, j$  such that  $w_{ij} > 0$ , we deduce from  $\varphi'(x_i^\top x_j) > 0$  that  $x_i = x_j$ . Applying this to a spanning tree of positive weights confirms that  $x_1 = \dots = x_n$ .  $\square$

## C Synchronization on spheres ( $d \geq 3$ ): a direct proof of Theorem 1

In this section, we provide a proof for Theorem 1, adapting the proof technique in (McRae and Boumal, 2024, Thm. 1.1). That paper considers synchronization on more general Stiefel manifolds (beyond circles and spheres) with general connected graphs. It is limited to what here would be a linear  $\varphi$ , but the extension to increasing, convex  $\varphi$  is easy on spheres.

To exploit second-order criticality conditions, we should perturb the points  $x_1, \dots, x_n$ . Rather than selecting these perturbations deterministically, it is convenient to choose them at random. Moreover, as the goal is to achieve synchrony, we perturb the points toward the *same* direction. Explicitly, with  $\gamma \in \mathbb{R}^d$  random, we move  $x_i$  in the direction  $\dot{x}_i = \gamma - (x_i^\top \gamma)x_i$ —the projection of  $\gamma$  to the tangent space of the sphere at  $x_i$ . The same  $\gamma$  is used for all points. We make this precise below.

Assume  $X$  is a critical point for  $f$  where the Hessian is negative semidefinite. Lemma 16 provides

$$XS = 0 \quad \text{and} \quad \langle S, \dot{X}^\top \dot{X} \rangle \geq 0 \quad (23)$$

for all  $\dot{X} \in T_X \mathcal{M}$ , with  $S = \text{ddiag}(X^\top X A) - A$  and  $A = W \odot \varphi'(X^\top X)$ ; we have simplified the Hessian expression (19), because the assumption  $\varphi''(t) \geq 0$  for all  $t \in [-1, 1]$  implies that the second term in (19) is nonnegative.

Since the inequality holds for all tangent  $\dot{X}$ , we can also allow  $\dot{X}$  to be random in  $T_X \mathcal{M}$  and claim the inequality holds in expectation, that is,  $\langle S, \mathbb{E}\{\dot{X}^\top \dot{X}\} \rangle \geq 0$ .

Let  $\gamma \sim \mathcal{N}(0, I_d)$  be a random vector in  $\mathbb{R}^d$  with i.i.d. entries following a standard normal distribution. We use it to build a random tangent vector at  $X$  as follows, with  $\mathbf{1} \in \mathbb{R}^n$  and  $\text{Proj}_X$  (16) the orthogonal projector to the tangent space at  $X$ :

$$\dot{X} = \text{Proj}_X(\gamma\mathbf{1}^\top) = \gamma\mathbf{1}^\top - X \text{diag}(X^\top \gamma). \quad (24)$$

Then

$$\begin{aligned}\dot{X}^\top \dot{X} &= \|\gamma\|^2 \mathbf{1}\mathbf{1}^\top - \mathbf{1}\gamma^\top X \text{diag}(X^\top \gamma) - \text{diag}(X^\top \gamma) X^\top \gamma \mathbf{1}^\top + \text{diag}(X^\top \gamma) X^\top X \text{diag}(X^\top \gamma) \\ &= \|\gamma\|^2 \mathbf{1}\mathbf{1}^\top - \mathbf{1}(\gamma^\top X)^{\odot 2} - (X^\top \gamma)^{\odot 2} \mathbf{1}^\top + (X^\top X) \odot (X^\top \gamma \gamma^\top X).\end{aligned}$$

Taking expectations, we have  $\mathbb{E}\{\|\gamma\|^2\} = d$ ,  $\mathbb{E}\{\gamma\gamma^\top\} = I_d$  and  $\mathbb{E}\{(X^\top \gamma)_i^2\} = \mathbb{E}\{x_i^\top \gamma \gamma^\top x_i\} = x_i^\top x_i = 1$ , so that

$$\mathbb{E}\{\dot{X}^\top \dot{X}\} = (d-2)\mathbf{1}\mathbf{1}^\top + (X^\top X)^{\odot 2}.$$

We decompose the second term as follows, in order to isolate the discrepancy between  $x_i^\top x_j$  and 1 (its target value):

$$(x_i^\top x_j)^2 = (1 - x_i^\top x_j)^2 - 1 + 2x_i^\top x_j.$$

In matrix notation,  $(X^\top X)^{\odot 2} = (\mathbf{1}\mathbf{1}^\top - X^\top X)^{\odot 2} - \mathbf{1}\mathbf{1}^\top + 2X^\top X$ . Therefore,

$$\mathbb{E}\{\dot{X}^\top \dot{X}\} = (d-3)\mathbf{1}\mathbf{1}^\top + (\mathbf{1}\mathbf{1}^\top - X^\top X)^{\odot 2} + 2X^\top X.$$

Exploiting the first-order condition  $XS = 0$ , we further obtain

$$0 \leq \langle S, \mathbb{E}\{\dot{X}^\top \dot{X}\} \rangle = (d-3)\langle S, \mathbf{1}\mathbf{1}^\top \rangle + \langle S, (\mathbf{1}\mathbf{1}^\top - X^\top X)^{\odot 2} \rangle.$$

Recall that  $S = \text{ddiag}(X^\top X A) - A$  with  $A = W \odot \varphi'(X^\top X)$ . Thus

$$\langle S, \mathbf{1}\mathbf{1}^\top \rangle = \text{Tr}(X^\top X A) - \langle A, \mathbf{1}\mathbf{1}^\top \rangle = \langle A, X^\top X - \mathbf{1}\mathbf{1}^\top \rangle \leq 0$$

owing to the fact that each entry of  $X^\top X$  is in the interval  $[-1, 1]$  and the assumption  $\varphi'(t) \geq 0$  for  $t \in [-1, 1]$  ensures the entries of  $A$  are nonnegative. Moreover,  $d-3 \geq 0$ , and the diagonal of  $\mathbf{1}\mathbf{1}^\top - X^\top X$  is zero, so

$$\begin{aligned}0 &\leq (d-3)\langle S, \mathbf{1}\mathbf{1}^\top \rangle + \langle S, (\mathbf{1}\mathbf{1}^\top - X^\top X)^{\odot 2} \rangle \\ &\leq -\langle A, (\mathbf{1}\mathbf{1}^\top - X^\top X)^{\odot 2} \rangle \\ &= -\sum_{i,j} w_{ij} \varphi'(x_i^\top x_j) (1 - x_i^\top x_j)^2 \leq 0,\end{aligned}\tag{25}$$

where the last inequality follows once again from  $\varphi'(t) \geq 0$  for all  $t \in [-1, 1]$ . As a result, the final sum in (25) is equal to zero, so each individual term is equal to zero. If nodes  $i$  and  $j$  are connected by an edge ( $w_{ij} > 0$ ), then the (stricter) assumption  $\varphi'(t) > 0$  for  $t \in [-1, 1]$  forces  $1 - x_i^\top x_j = 0$ , that is,  $x_i = x_j$ . As we have assumed the graph is connected, apply the same argument along the edges of a spanning tree to deduce that  $x_1 = \dots = x_n$ . (If the graph is not connected, apply the same reasoning to a spanning forest to deduce synchrony in each connected component.) This concludes the proof of Theorem 1.

**Remark 17.** To establish (23), we simplified the conclusions of Lemma 16 by using the assumption  $\varphi''(t) \geq 0$ . That is the only place where that assumption is used. Alternatively, we could keep the full expression for the Hessian and compute the expectation of  $(X^\top \dot{X} + \dot{X}^\top X)_{ij}^2$ . This yields a more refined final inequality which could be used to relax the assumptions on  $\varphi$  or the assumption  $d \geq 3$ . However, it is unclear to us how to improve the results for  $\varphi_\beta$  with  $d = 2$  in this way.

## D Proofs from Section 5: Lemmas 14 and 15

### D.1 Lemma 14: trading weights for repetitions

To prove Lemma 14, we first prove the following finer statement:

**Lemma 18.** *Let  $d = 2$ , and let  $m$  be any positive integer. Let  $q \in \mathbb{R}^m$  have positive integer entries, let  $n = q_1 + \dots + q_m$ , and define the  $n \times n$  diagonal matrix*

$$\tilde{Q} = \begin{pmatrix} q_1 I_{q_1} & & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & q_m I_{q_m} \end{pmatrix}.$$

Given  $X = (x_1, \dots, x_m) \in \mathbb{R}^{2 \times m}$ , define

$$\tilde{X} = \underbrace{(x_1, \dots, x_1)}_{q_1 \text{ times}}, \dots, \underbrace{(x_m, \dots, x_m)}_{q_m \text{ times}} \in \mathbb{R}^{2 \times n}.$$

If  $X$  is critical for  $f_m(X) = \frac{1}{2} \langle qq^\top, \varphi(X^\top X) \rangle$ , then  $\tilde{X}$  is critical for  $f_n(\tilde{X}) = \frac{1}{2} \langle \mathbf{1}_n \mathbf{1}_n^\top, \varphi(\tilde{X}^\top \tilde{X}) \rangle$ .

Further, let  $M$ ,  $m_{ij} = q_i q_j h(x_i^\top x_j)$ , be the adjacency matrix corresponding to  $\text{Hess} f_m(X)$ , as described in equation (5) of Lemma 9, and define  $D = \text{diag}(M \mathbf{1}_m)$ . Likewise define  $\tilde{M}, \tilde{D}$  for  $\text{Hess} f_n(\tilde{X})$ .

Let  $\lambda_1, \dots, \lambda_{m-1}, 0$  be the eigenvalues of the Laplacian  $L(M)$ . Then  $\tilde{Q}^{1/2} L(\tilde{M}) \tilde{Q}^{1/2}$  has eigenvalues  $\lambda_1, \dots, \lambda_{m-1}, 0$  too, and also has eigenvalues  $d_{ii}$ ,  $i = 1, \dots, m$ , each with multiplicity  $q_i - 1$  (where  $d_{ii}$  is the  $i$ th diagonal entry of  $D$ ). This covers all  $n$  eigenvalues of  $\tilde{Q}^{1/2} L(\tilde{M}) \tilde{Q}^{1/2}$ .

*Proof of Lemma 18.* Define  $Q = \text{diag}(q) \in \mathbb{R}^{m \times m}$ , and define the  $n \times m$  matrix

$$E = \begin{pmatrix} \mathbf{1}_{q_1} & 0 & 0 & \dots & 0 \\ 0 & \mathbf{1}_{q_2} & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \mathbf{1}_{q_m} \end{pmatrix}.$$

Note the identities

$$\tilde{X} = X E^\top, \quad E^\top E = Q, \quad \tilde{Q} E = E Q. \quad (26)$$

Following equation (7) of Lemma 9 (for both  $f_m$  and  $f_n$ ), define  $A = qq^\top \odot \varphi'(X^\top X) \in \mathbb{R}^{m \times m}$  and  $\tilde{A} = \varphi'(\tilde{X}^\top \tilde{X}) \in \mathbb{R}^{n \times n}$ . Defining  $A' = \varphi'(X^\top X)$ , we obtain the identities

$$A = Q A' Q, \quad \tilde{A} = E A' E^\top. \quad (27)$$

Since  $X$  is critical for  $f_m$ , Lemma 9 gives  $X A = X \text{diag}(X^\top X A)$ . Hence, using equations (26) and (27), we have

$$\begin{aligned} \tilde{X} \text{diag}(\tilde{X}^\top \tilde{X} \tilde{A}) &= X E^\top \text{diag}(E X^\top X Q A' E^\top) \\ &= X E^\top \text{diag}(E X^\top X A Q^{-1} E^\top) \\ &\stackrel{(1)}{=} X \text{diag}(X^\top X A Q^{-1}) E^\top \\ &= X \text{diag}(X^\top X A) Q^{-1} E^\top \\ &= X A Q^{-1} E^\top \\ &= X Q A' Q Q^{-1} E^\top \\ &= X E^\top E A' E^\top \\ &= \tilde{X} \tilde{A}, \end{aligned}$$

where equality  $\stackrel{(1)}{=}$  follows from the identity  $E^\top \text{diag}(EBE^\top) = \text{diag}(B)E^\top$  for  $B \in \mathbb{R}^{m \times m}$ . We conclude that  $\tilde{X}$  is critical for  $f_n$ , again by Lemma 9.

Let us move on to the second-order condition. Our goal is to express the eigenvalues of  $L(\tilde{M}) = \tilde{D} - \tilde{M}$  in terms of those of  $L(M) = D - M$ . Towards this end, define the  $m \times m$  matrices  $M' = Q^{-1}MQ^{-1}$  and  $D' = Q^{-1}DQ^{-1}$ , which have entries

$$m'_{ij} = (q_i q_j)^{-1} m_{ij} = h(x_i^\top x_j) \quad \text{and} \quad d'_{ii} = q_i^{-2} d_{ii}.$$

By the definition of  $\tilde{M}$ , we have

$$\tilde{M} = \begin{pmatrix} m'_{11} \mathbf{1}_{q_1} \mathbf{1}_{q_1}^\top & m'_{12} \mathbf{1}_{q_1} \mathbf{1}_{q_2}^\top & \cdots & m'_{1m} \mathbf{1}_{q_1} \mathbf{1}_{q_m}^\top \\ \vdots & \vdots & \ddots & \vdots \\ m'_{m1} \mathbf{1}_{q_m} \mathbf{1}_{q_1}^\top & m'_{m2} \mathbf{1}_{q_m} \mathbf{1}_{q_2}^\top & \cdots & m'_{mm} \mathbf{1}_{q_m} \mathbf{1}_{q_m}^\top \end{pmatrix} = EM'E^\top.$$

Moreover,  $\tilde{D} = \text{diag}(\tilde{M} \mathbf{1}_n)$  is diagonal with diagonal blocks  $\tilde{D}_i$  given by:

$$\tilde{D} = \begin{pmatrix} \tilde{D}_1 & & \\ & \ddots & \\ & & \tilde{D}_m \end{pmatrix}, \quad \tilde{D}_i = q_i^{-1} d_{ii} I_{q_i} = q_i d'_{ii} I_{q_i}. \quad (28)$$

We can decompose  $L(\tilde{M})$  as

$$\begin{aligned} L(\tilde{M}) &= \tilde{D} - \tilde{M} \\ &= [\tilde{D} - ED'E^\top] + E(D' - M')E^\top \\ &= [\tilde{D} - ED'E^\top] + EQ^{-1}L(M)Q^{-1}E^\top. \end{aligned}$$

Using  $\tilde{Q}E = EQ$ , we get

$$\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2} = \underbrace{\tilde{Q}^{1/2}[\tilde{D} - ED'E^\top]\tilde{Q}^{1/2}}_{\text{left term}} + \underbrace{EQ^{-1/2}L(M)Q^{-1/2}E^\top}_{\text{right term}}. \quad (29)$$

Crucially, the decomposition (29) reveals the eigendecomposition of  $\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2}$ . Indeed,  $EQ^{-1/2}$  is orthonormal since  $E^\top E = Q$ , and the column spaces of the left and right terms of (29) are orthogonal. This latter observation is apparent from the fact that  $\tilde{Q}^{1/2}[\tilde{D} - ED'E^\top]\tilde{Q}^{1/2}$  is *block diagonal* with  $i$ -th diagonal block given by

$$q_i (\tilde{D}_i - d'_{ii} \mathbf{1}_{q_i} \mathbf{1}_{q_i}^\top) = d_{ii} (I_{q_i} - q_i^{-1} \mathbf{1}_{q_i} \mathbf{1}_{q_i}^\top), \quad (30)$$

where we have used (28).

We conclude that  $\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2}$  has  $m$  eigenvalues equal to the eigenvalues of  $L(M)$ , due to the right term of (29). Due to the left term of (29) and the form of the blocks (30),  $\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2}$  also has eigenvalues  $d_{ii}, i = 1, \dots, m$ , each with multiplicity  $q_i - 1$ .

To summarize, we have identified all  $m + \sum_{i=1}^m (q_i - 1) = n$  eigenvalues of  $\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2}$ .  $\square$

Lemma 14 now follows as a special case of Lemma 18:

*Proof of Lemma 14.* We use the notation from Lemma 18. Since we assume  $\text{Hess} f_m(X)$  is negative definite (up to the trivial eigenvalue due to symmetry, as usual), Lemma 9 implies  $\lambda_i > 0$  for  $i = 1, \dots, m - 1$ , and  $D - M \succeq 0$ . Thus,

$$d_{ii} \geq m_{ii} = q_i^2 h(1) = q_i^2 \varphi'(1) > 0.$$

So invoking Lemma 18, we have shown that  $\tilde{Q}^{1/2}L(\tilde{M})\tilde{Q}^{1/2}$  has

$$(m-1) + \sum_{i=1}^m (q_i - 1) = n - 1$$

positive eigenvalues (and of course has one remaining zero eigenvalue). So the same is true for  $L(\tilde{M})$ . Lemma 9 then tells us that  $\text{Hess}f_n(\tilde{X})$  is also negative definite.  $\square$

## D.2 Lemma 15: strict spurious points do not vanish under small perturbations

*Proof of Lemma 15.* In order to ensure that  $\varepsilon$  is nonnegative, we reparameterize  $\varepsilon = \alpha^2$  for  $\alpha \in \mathbb{R}$ . Consider the function

$$f_{q,\alpha,\tau}(X) := \frac{1}{2} \langle qq^\top, \varphi_{\alpha^2,\tau}(X^\top X) \rangle.$$

We think of  $f_{q,\alpha,\tau}$  as a perturbation of the function  $f_{\mathbf{1}_m,0,0}$ , which is simply  $f$  with  $W$  the all-ones matrix and the ReLU  $\varphi(t) = \max\{0, t\}$ . Let  $X$  be the regular  $m$ -gon (10).

We want to apply the implicit function theorem (Krantz and Parks, 2013, Thm. 3.3.1) to the map

$$F: ((q, \alpha, \tau), Y) \mapsto \text{grad}f_{q,\alpha,\tau}(Y)$$

at  $q = \mathbf{1}_m, \alpha = 0, \tau = 0$  and  $Y = X$ . In order to do this, we need:

- (a)  $F$  to be continuously differentiable in a neighborhood of  $q = \mathbf{1}_m, \alpha = 0, \tau = 0, Y = X$ .
- (b)  $0 = F((\mathbf{1}_m, 0, 0), X) = \text{grad}f_{\mathbf{1}_m,0,0}(X)$ . This is true by Lemma 12 (and the assumption  $m$  is odd).
- (c) The differential of  $Y \mapsto F((\mathbf{1}_m, 0, 0), Y)$  at  $Y = X$  to be invertible. That differential is exactly  $\text{Hess}f_{\mathbf{1}_m,0,0}(X)$ , which is negative definite (and so invertible) by Lemma 12.<sup>7</sup>

For item (a): by Lemma 16,  $\varphi_{\alpha^2,\tau}$  appears in  $F$  only through  $\varphi'_{\alpha^2,\tau}$ . So it is enough to verify that

$$(\alpha, \tau, t) \mapsto \varphi'_{\alpha^2,\tau}(t) = 1 - \frac{1}{1 + e^{(t-\tau)/\alpha^2}}$$

is continuously differentiable in a neighborhood of  $(0, 0, t_0)$  when  $t_0 \neq \tau$ .<sup>8</sup> It is straightforward to do this by explicitly computing the differential of  $(\alpha, \tau, t) \mapsto \varphi'_{\alpha^2,\tau}(t)$ ; we omit the details.

The hypotheses of the implicit function theorem are satisfied, and that theorem yields that there is a  $\delta > 0$  such that if  $\|q - \mathbf{1}_m\| \leq \delta$ ,  $\alpha \in [-\sqrt{\delta}, \sqrt{\delta}]$  and  $\tau \in [-\delta, \delta]$ , then there is an  $X_{q,\alpha,\tau} \in \mathcal{M}$  near the  $m$ -gon  $X$  such that

$$0 = F((q, \alpha, \tau), X_{q,\alpha,\tau}) = \text{grad}f_{q,\alpha,\tau}(X_{q,\alpha,\tau}).$$

Since  $X_{q,\alpha,\tau}$  is near  $X$ , by continuity  $X_{q,\alpha,\tau}$  is also spurious and has negative definite Hessian (except for the single zero eigenvalue), possibly after making  $\delta$  smaller.  $\square$

<sup>7</sup>Strictly speaking, the Hessian is not negative definite since it has a single zero eigenvalue due to the global rotation symmetry. So to apply the implicit function theorem we must first mod out that zero eigenvalue, e.g., by fixing one of the points or passing to the quotient. Then the Hessian becomes truly negative definite.

<sup>8</sup>Since  $m$  is odd, none of the inner products  $x_i^\top x_j$  of the regular  $m$ -gon lie on the kink of the ReLU; this is why we only require continuous differentiability when  $t_0 \neq \tau$ .

## E Synchronization with $\beta \geq \frac{n^2}{\pi^2}$ : Theorem 6

We now give a proof for Theorem 6. That result and the proof below are due to [Geshkovski et al. \(2024, Thm. 5.3\)](#), with only slight improvements as outlined in the introduction.

*Proof of Theorem 6.* Assume  $X$  is critical for  $f$  with negative semidefinite Hessian. From Lemma 9, we know the Laplacian (4)  $L(M) = \text{diag}(M\mathbf{1}) - M$  is positive semidefinite, with

$$m_{ij} = w_{ij}h(x_i^\top x_j) \quad \text{and} \quad h(t) = t\varphi'(t) - (1-t^2)\varphi''(t).$$

By condition (2), we know that  $h(t) \geq 0$  implies  $t \geq \cos(\frac{\pi}{n})$ .

Pick a nonempty proper subset  $S \subset \{1, \dots, n\}$ . Let  $\alpha_i = 1$  if  $i \in S$  and  $\alpha_i = 0$  if  $i \notin S$ . Since the Laplacian  $L$  is positive semidefinite we find

$$\alpha^\top L \alpha = \frac{1}{2} \sum_{i,j} m_{ij} (\alpha_i - \alpha_j)^2 = \sum_{i \in S, j \notin S} w_{ij} h(x_i^\top x_j) \geq 0.$$

There exist indices  $i \in S, j \notin S$  such that  $w_{ij} > 0$  (as otherwise all the weights between  $S$  and its complement would be zero, and the graph would be disconnected). For at least one of those pairs,  $h(x_i^\top x_j) \geq 0$  (as otherwise the sum would be negative), i.e.,  $x_i^\top x_j \geq \cos(\frac{\pi}{n})$  by condition (2). With  $\text{dist}$  denoting the distance on the circle (in radians), this gives  $\text{dist}(x_i, x_j) \leq \cos^{-1}(\cos(\frac{\pi}{n})) = \frac{\pi}{n}$ .

Starting with  $S = \{1\}$ , apply the argument above to identify a node in the complement. Add it to  $S$  and repeat. This gradually grows a spanning tree over the  $n$  nodes, and it satisfies  $\text{dist}(x_i, x_j) \leq \frac{\pi}{n}$  for each edge of the tree. The radius of a tree is at most  $n/2$ , hence we can select a central node  $x_i$  such that  $\text{dist}(x_i, x_j) \leq \frac{n}{2} \cdot \frac{\pi}{n} = \frac{\pi}{2}$  for all  $j$ . In other words,  $x_1, \dots, x_n$  lie in the same half circle: we can then apply Lemma 10 to conclude, proving the main part of Theorem 6.

It remains to verify that if  $\varphi''(t) \geq \frac{n^2}{\pi^2} \varphi'(t) > 0$  for all  $t$ , then condition (2) holds. We can of course assume  $n \geq 2$  (because if  $n = 1$ , the state is already trivially synchronized). Assume  $h(t) \geq 0$ ; we want to show  $t \geq \cos(\frac{\pi}{n})$ . First, note that the assumption on  $\varphi$  gives

$$0 \leq h(t) = t\varphi'(t) - (1-t^2)\varphi''(t) \leq \left(t - \frac{n^2}{\pi^2}(1-t^2)\right)\varphi'(t).$$

Since  $\varphi' > 0$ , we conclude  $t - \frac{n^2}{\pi^2}(1-t^2) \geq 0$ , which implies<sup>9</sup>  $\frac{t}{1-t^2} \geq \frac{n^2}{\pi^2}$ . Using the inequality  $\frac{\cos(s)}{1-\cos(s)^2} \leq \frac{1}{s^2}$  for all  $s \in (0, \frac{\pi}{2}]$ , we have  $\frac{t}{1-t^2} \geq \frac{n^2}{\pi^2} \geq \frac{\cos(\pi/n)}{1-\cos(\pi/n)^2}$ . This implies  $t \geq \cos(\frac{\pi}{n})$ , as  $t \mapsto \frac{t}{1-t^2}$  is increasing on  $[-1, 1]$ .  $\square$

## F The quadratic case $\varphi(t) = \frac{1}{2}t^2$

Consider the quadratic case  $\varphi(t) = \frac{1}{2}t^2$  with a complete graph  $w_{ij} = 1$  for all  $i, j$ . Notice that  $\varphi'(t) = t$  can be negative on  $[-1, 1]$ , hence this falls outside the scope of the main theorems in this paper. Since  $\varphi$  has two maxima on the interval  $[-1, 1]$ , namely,  $\pm 1$ , the global maxima of  $f$  correspond to points  $x_1, \dots, x_n$  all equal *up to sign*. We now show that for  $f(X) = \frac{1}{4}\|X^\top X\|_F^2$  (Frobenius norm), the problem (P) has no spurious local maxima. We use tools that are different from the other cases.

<sup>9</sup>We already have  $t \geq 0$  (as otherwise  $h(t) < 0$ ), and we can assume  $t < 1$  (as otherwise we are done).

**Theorem 19.** Fix  $d \geq 2$  and  $n \geq 1$ . Let  $\varphi(t) = \frac{1}{2}t^2$  and consider a complete graph with unit weights  $w_{ij} = 1$ . Then, critical points of  $f$  where the Hessian is negative semidefinite are global maxima of  $f$ . In particular: local maxima are global maxima, they are synchronized up to sign ( $x_i = \pm x_j$  for all  $i, j$ ), and gradient flow converges to such a state from almost every initialization.

We prove Theorem 19 in this section via three lemmas.

**Lemma 20.** Within the context of Theorem 19, if  $X \in \mathcal{M}$  is a critical point of  $f$  then

$$(XX^\top)X = XD \quad \text{where} \quad D = \text{ddiag}(X^\top XX^\top X). \quad (31)$$

If the Hessian at  $X$  is negative semidefinite then, for all  $\dot{X} \in \mathbb{T}_X \mathcal{M}$ ,

$$\langle \dot{X}^\top \dot{X}, \text{ddiag}(X^\top XX^\top X) \rangle \geq \langle XX^\top, \dot{X} \dot{X}^\top \rangle + \langle X \dot{X}^\top, \dot{X} X^\top \rangle + \|\dot{X} X^\top\|_{\mathbb{F}}^2. \quad (32)$$

*Proof.* Consider Lemma 16 with  $\varphi'(t) = t$  and  $\varphi''(t) = 1$ . We see that  $A = XX^\top$  and  $S = \text{ddiag}(X^\top XX^\top X) - X^\top X$ . The first-order condition  $XS = 0$  provides (31). The second-order conditions provide

$$0 \geq -\langle \dot{X}^\top \dot{X}, S \rangle + \frac{1}{2} \|X^\top \dot{X} + \dot{X}^\top X\|_{\mathbb{F}}^2$$

for all  $\dot{X}$  in the tangent space at  $X$ . The right-most term expands into

$$\frac{1}{2} \|X^\top \dot{X} + \dot{X}^\top X\|_{\mathbb{F}}^2 = \frac{1}{2} \langle X^\top \dot{X} + \dot{X}^\top X, X^\top \dot{X} + \dot{X}^\top X \rangle = \langle XX^\top, \dot{X} \dot{X}^\top \rangle + \langle X \dot{X}^\top, \dot{X} X^\top \rangle,$$

whereas  $\langle \dot{X}^\top \dot{X}, S \rangle = \langle \dot{X}^\top \dot{X}, \text{ddiag}(X^\top XX^\top X) \rangle - \langle \dot{X}^\top \dot{X}, X^\top X \rangle$ . Combine to confirm (32).  $\square$

The first-order condition (31) is formatted to highlight the following key fact: each column  $x_i$  of  $X$  is an eigenvector of  $XX^\top \in \mathbb{R}^{d \times d}$ , with eigenvalue  $D_{ii}$ . Since  $XX^\top$  is symmetric, the spectral theorem tells us that if  $x_i, x_j$  are associated to distinct eigenvalues ( $D_{ii} \neq D_{jj}$ ) then  $x_i$  and  $x_j$  are orthogonal. The next step is to show that this does not happen.

**Lemma 21.** With  $X$  as in Lemma 20, no two columns of  $X$  are orthogonal.

*Proof.* For contradiction, assume  $x_i^\top x_j = 0$  for some  $i, j$ . Then,  $x_j$  is in the tangent space to the sphere at  $x_i$ , and vice versa. Consequently,

$$\dot{X} = x_i e_j^\top + x_j e_i^\top$$

is in the tangent space to  $\mathcal{M}$  at  $X$ , where  $e_k \in \mathbb{R}^n$  is the  $k$ th column of the identity matrix of size  $n$ . By computation,  $\dot{X}^\top \dot{X} = e_i e_i^\top + e_j e_j^\top$ . Also,  $\dot{X} \dot{X}^\top = x_i x_i^\top + x_j x_j^\top$  and  $X \dot{X}^\top = x_i x_j^\top + x_j x_i^\top = \dot{X} X^\top$ . Plugging these into the second-order conditions (32) reveals that

$$\langle e_i e_i^\top + e_j e_j^\top, \text{ddiag}(X^\top XX^\top X) \rangle \geq \langle XX^\top, x_i x_i^\top + x_j x_j^\top \rangle + 2 \|x_i x_j^\top + x_j x_i^\top\|_{\mathbb{F}}^2.$$

The left-hand side and the first term on the right-hand side are equal. Also,  $\|x_i x_j^\top + x_j x_i^\top\|_{\mathbb{F}}^2 = 2$  since  $x_i$  and  $x_j$  are orthogonal. Overall, we have found  $0 \geq 4$ : a contradiction indeed.  $\square$

Thus, we know that  $x_1, \dots, x_n$  are all eigenvectors for  $XX^\top$  with the *same* eigenvalue. If those vectors span all of  $\mathbb{R}^d$ , this imposes severe constraints on  $XX^\top$ —too strong, in fact.

**Lemma 22.** With  $X$  as in Lemma 20, it holds that  $\text{rank}(X) < d$ .



*Proof.* For contradiction, assume  $\text{rank}(X) = d$ . Then, we may select  $d$  linearly independent columns among those of  $X$ . These form a basis for  $\mathbb{R}^d$  and (by the reasoning above) are eigenvectors of  $XX^\top$  for the same eigenvalue  $\lambda$ . It follows that  $XX^\top$  has a single eigenvalue, and so  $XX^\top = \lambda I_d$ . Plug this and  $\text{diag}(X^\top X) = \mathbf{1}$  into the second-order conditions (32) to reveal that

$$\lambda \|\dot{X}\|_F^2 \geq \lambda \|\dot{X}\|_F^2 + \langle X\dot{X}^\top, \dot{X}X^\top \rangle + \|\dot{X}X^\top\|_F^2 \quad (33)$$

for all tangent  $\dot{X}$ . To contradict this, let  $\dot{X} = ue_1^\top$  where  $e_1$  is the first column of the identity matrix  $I_n$ , and  $u$  is an arbitrary unit-norm vector in the tangent space to the sphere at  $x_1$ . (Such a vector exists as long as  $d \geq 2$ .) Then,  $\dot{X}X^\top = ux_1^\top$ . Since  $x_1^\top u = 0$ , the above inequality becomes  $0 \geq 1$ : a contradiction indeed.  $\square$

We now know that  $X$  is rank deficient, and we wish to deduce that  $X$  is optimal. Notice that we cannot use Lemma 10 here, since  $\varphi'(t) = t$  is not positive on  $[-1, 1]$ . Thus, we resort to a different argument.

Rotate the points such that the last row of  $X$  is zero (formally: let  $X = U\Sigma V^\top$  be an SVD of  $X$ , and apply  $U^\top$  to  $X$ ). The new  $X$  still satisfies first- and second-order conditions since  $f$  is invariant to such rotations. For now, discard the last row of  $X$ , producing a new matrix  $\tilde{X} \in \mathbb{R}^{(d-1) \times n}$ . In particular,  $X^\top X = \tilde{X}^\top \tilde{X}$ . This new matrix also satisfies first- and second-order conditions for  $f$  in the new dimension  $d - 1$  (because all we did was remove potential directions for improvement). Moreover, the rank did not change:  $\text{rank}(\tilde{X}) = \text{rank}(X)$ . If  $d - 1 \geq 2$ , Lemma 22 applies and so  $\tilde{X}$  too is rank deficient. We may repeat this operation until we reach the conclusion that the original  $X$  must have had rank 1.

That implies that all columns of  $X$  are colinear, hence that they are equal up to sign. This is indeed maximal for  $f$  since  $\varphi(t)$  is maximal if and only if  $t = \pm 1$ : the proof of Theorem 19 is complete.

## G Matlab script for numerical exploration

```
clear; clf; clc;

d = 2;
n = 16;

phichoice = 'expbeta';
switch phichoice
    case 'expbeta'
        beta = 6;
        phi = @(t) exp(beta*(t-1));
        dphi = @(t) beta*exp(beta*(t-1));
        d2phi = @(t) beta^2*exp(beta*(t-1));
    case 'linear'
        phi = @(t) t;
        dphi = @(t) 1;
        d2phi = @(t) 0;
    case 'quadratic'
        phi = @(t) (1/2)*t.^2;
        dphi = @(t) t;
        d2phi = @(t) 1;
    case 'cubic'
        phi = @(t) (1/3)*t.^3;
        dphi = @(t) t.^2;
        d2phi = @(t) 2*t;
    case 'logsumexp'
        m = n; %
        tau = cos((3/2)*(2*pi/m));
        eps = 1/m^3;
        phi = @(t) eps*log(1 + exp((t-tau)/eps));
        dphi = @(t) 1./(1 + exp(-(t-tau)/eps));
        d2phi = @(t) (sech((t-tau)/(2*eps)).^2)./(4*eps);
end

graph = 'complete';
switch graph
    case 'complete'
        W = ones(n, n);
    case 'cycle'
        I = 1:n;
        J = circshift(I, 1);
        W = zeros(n, n);
        W(sub2ind([n, n], I, J)) = 1;
        W = W+W';
end
```

```

problem.M = obliquefactory(d, n, false, false);
sgn = -1; %
inner = @(A, B) A(:).'*B(:);
problem.cost = @(X) sgn*(1/n^2)*inner(W, phi(X'*X));
if exist('dphi', 'var') && exist('d2phi', 'var')
    problem.egrad = @(X) sgn*(2/n^2)*X*(W.*dphi(X'*X));
    problem.ehess = @(X, Xdot) ...
        sgn*(2/n^2)*(Xdot*(W.*dphi(X'*X)) + ...
            X*(W.*d2phi(X'*X).*(Xdot'*X + X'*Xdot)));
else
    problem = manoptAD(problem);
end

init = 'random';
switch init
    case 'random'
        X0 = problem.M.rand();
    case 'ngon'
        X0 = zeros(d, n);
        t = linspace(0, 2*pi, n+1);
        t = t(1:end-1);
        X0(1, :) = cos(t);
        X0(2, :) = sin(t);
    case 'qngon'
        X0 = zeros(d, n);
        q = 5;
        t = linspace(0, 2*pi, q+1);
        t = repmat(t(1:q), 1, ceil(n/q));
        X0(1, :) = cos(t(1:n));
        X0(2, :) = sin(t(1:n));
    case 'tetrahedron'
        assert(d == 3 && n == 4);
        X0 = [1      -1/3      -1/3      -1/3
              0  sqrt(8/9) -sqrt(2/9) -sqrt(2/9)
              0          0  sqrt(6/9) -sqrt(6/9)];
end

hdots = plot_dots(X0); title('Initial X0');

options.maxiter = 300;
options.tolgradnorm = 1e-12;
options.statsfun = @(problem, X, stats) ...
    update_dots(hdots, X, .1, stats);

optimizationplots = 'gradientdynamics';
switch optimizationplots
    case 'fast'
        X = trustregions(problem, X0, options);

```

```

        case 'gradientdynamics'
            X = steepestdescent(problem, X0, options);
        case 'none'
            X = X0;
    end
    update_dots(hdots, X); title('Final X');

    gnorm = problem.M.norm(X, getGradient(problem, X));
    hspec = hessianspectrum(problem, X);
    fprintf('Gradient norm at X: %
    fprintf('Eigenvalues of Hessian at X:\n');
    disp(hspec');

function hdots = plot_dots(X)
    hold all;
    dotscolors = X(1, :) + X(2, :);
    if size(X, 1) == 3
        [Sx, Sy, Sz] = sphere(50);
        surf(Sx, Sy, Sz, 'FaceAlpha', .25, ...
            'FaceColor', 2* [.1, .2, .3]);
        hdots = scatter3(X(1, :)', X(2, :)', X(3, :)', ...
            200, dotscolors, 'filled');
    elseif size(X, 1) == 2
        t = linspace(0, 2*pi, 501);
        plot(cos(t), sin(t), 'k-', 'LineWidth', 2);
        hdots = scatter(X(1, :)', X(2, :)', 200, ...
            dotscolors, 'filled');
    end
    axis equal; axis off;
    set(gcf, 'Color', 'w');
end

function stats = update_dots(hdots, X, time, stats)
    set(hdots, 'XData', X(1, :));
    set(hdots, 'YData', X(2, :));
    if size(X, 1) == 3
        set(hdots, 'ZData', X(3, :));
    end
    drawnow;
    if exist('time', 'var') && ~isempty(time)
        pause(time);
    end
    if ~exist('stats', 'var') %
        stats = [];
    end
end
end

```