Motor Imagery Task Alters Dynamics of Human Body Posture

Fatemeh Delavari, Seyyed Mohammad Reza Hashemi Golpayegani, Mohammad Ali Ahmadi-Pajouh

Biomedical Engineering Department, Amirkabir University of Technology, Tehran, Iran

Fatemeh.delavari@uconn.edu,{m.r.hashemig, pajouh}@aut.ac.ir

Abstract

Motor Imagery (MI) is gaining traction in both rehabilitation and sports settings, but its immediate influence on human postural control is not yet clearly understood. The focus of this study is to examine the effects of MI on the dynamics of the Center of Pressure (COP), a crucial metric for evaluating postural stability.

In the experiment, thirty healthy young adults participated in four different scenarios: normal standing with both open and closed eyes, and kinesthetic motor imagery focused on mediolateral (ML) and anteroposterior (AP) sway movements. A mathematical model was developed to characterize the nonlinear dynamics of the COP and to assess the impact of MI on these dynamics.

Our results show a statistically significant increase (p-value<0.05) in variables such as COP path length (PL) and Long-Range Correlation (LRC) during MI compared to the closed-eye and normal standing conditions. These observations align well with psycho-neuromuscular theory, which suggests that imagining a specific movement activates neural pathways, consequently affecting postural control.

This study presents compelling evidence that motor imagery not only has a quantifiable impact on COP dynamics but also that changes in the Center of Pressure (COP) are directionally consistent with the imagined movements. This finding holds significant implications for the field of rehabilitation science, suggesting that motor imagery could be strategically utilized to induce targeted postural adjustments. Nonetheless, additional research is required to fully understand the complex mechanisms that underlie this relationship and to corroborate these results across a more diverse set of populations.

Key Words— Postural Control, Motor Imagery, Center of Pressure, Nonlinear dynamics, Balance

Introduction

Postural control, integral to most daily activities, is a multifaceted task dependent on the synergy of various systems, notably the visual and proprioceptive systems (1). Motor imagery, defined as the mental reconstruction of movement without its actual execution (2), can be categorized into two strategies: kinesthetic and visual. In kinesthetic motor imagery (MI), individuals visualize themselves executing the movement (first person perspective). In contrast, visual MI involves picturing another person or oneself performing the action from a thirdperson viewpoint (3). Research has identified overlapping mechanisms between motor imagery and actual movement execution, particularly in the realm of kinesthetic motor imagery (4). Moreover, MI has shown promise in rehabilitation and sports domains due to its benefits (5-12). Motor imagery is grounded in two primary theories: the peripheral (often called psycho-neuromuscular) and the central theories. The peripheral theory posits that imagining a specific movement results in sub-threshold muscle activations related to that movement, effectively simulating the neural pathways activated during actual movement (13-15). Conversely, other research underscores the enhancement in motor performance post MI sessions without noticeable muscle activity, accentuating the shared brain regions' activation during both imagined and actual movements (central theory). This central viewpoint indicates overlapping neural pathways for both MI and execution, especially within areas like the premotor and primary motor cortex (16).

The relationship between motor imagery and postural sway has been the focus of several studies (17-22). Some have posited that MI can induce observable changes in postural control and sway. Yet, the underlying mechanisms remain elusive. Is it cognitive distraction, neural resource allocation, inhibitory circuits, the dual mechanism, or another factor responsible for the changes in postural sway during MI? This study endeavors to probe the relation between the direction of actual postural sway and the imagined one. There are four possible hypotheses: 1) No discernible change in postural sway occurs during MI and improvements in postural control after MI sessions are attributed either to the activation of brain regions common between MI and movement execution (central theory) or to mentally rehearsing movement steps (symbolic learning theory). 2) Imagining a movement stimulates identical motor circuits as executing the movement, albeit to a lesser degree. Hence, visualizing a specific sway direction might elicit a minimal yet perceptible postural muscle response in that direction, aligning with the psycho-neuromuscular theory. 3) If maintaining stable posture is prioritized over the imagined sway, and/or if inhibitory circuits are hyperactive during MI, the actual sway could be counteractive. 4) Changes in postural sway might be observed, but these alterations might not correspond to the imagined sway direction. Such a phenomenon could stem from cognitive distractions or a lack of feedback mechanisms governing postural sway during MI. Deciphering the relationship between the direction of imagined and actual postural sway provides crucial insights for rehabilitation. These insights are particularly valuable in

crafting therapy sessions that leverage MI and in monitoring the progress of these sessions, especially when addressing issues of compromised balance and postural control.

Balance assessment techniques span from rudimentary clinical tests to advanced devices like force plates, electromyograms, and cameras. Force plates, instrumental for biomechanical evaluations of gait and posture, record the center of pressure (COP) (23). Within the force plate's context, the COP is the singular point where the pressure field's cumulative sum acts. This study employs COP's movement and positioning as a postural control indicator.

This research delves into the effects of motor imagery on COP dynamics by studying thirty healthy individuals during upright standing. Further, we suggest a COP dynamics model to illustrate these effects. Although numerous models exist for human posture (24-31), many view the human body as a linear system like an inverted pendulum—with attributes such as time delay, feedback, and a proportional derivative controller. However, Khanian et. al 's model (32) accounts for the nonlinear dynamics of COP and can better explain and quantify the COP changes. Consequently, we adapt this model in this study to illustrate postural control during regular standing and MI and discuss its potential implications for future occupational therapy and physiotherapy sessions.

Methods

1. Participants

Thirty healthy young adults, comprising twenty females and ten males, participated in the study. These individuals had no history of neurological, musculoskeletal, or vestibular disorders. They refrained from taking any medications 24 hours prior to the experiment, though the use of vitamins and dietary supplements was permissible. Participation was voluntary, and all participants provided informed consent. They were instructed to self-report their levels of mental and physical fatigue, as well as their attention, before the experiment and after each testing block. After each block, they also reported any discomfort experienced. Due to excessive fatigue or diminished attention, six participants were excluded from the analysis. The characteristics of the remaining 24 participants, including age, height, and weight, can be found in Table 1.

 Table 1: Range, mean and standard deviation of participants' age, height, and weight.

 Age (year)
 Height (cm)
 Weight (kg)

	${\bf Age} ({\bf year})$	${ m Height} \ ({ m cm})$	Weight (kg)	
$\begin{array}{l} \text{Mean} \pm \text{SD} \\ \text{Range} \end{array}$	24.6 ± 3.5 22-34	166.1 ± 6.9 152-179	60.6 ± 10.9 42-92	

Block 1		Block 2		Block 3		Block 4
Normal standing (open and closed eyes)	Rest	Motor imagery of AP sway	Rest	Motor imagery of ML sway	Rest	Normal standing (open and closed eyes)
2 min	1 min	4 min	1 min	4 min	1 min	2 min

Figure 1: An overview of the protocol. The experiment consisted of 4 blocks with a total duration of 15 minutes. In the first and final blocks, postural control of normal standing was recorded in two conditions of open and closed eyes, each for one minute. In the second and third blocks, the participants were asked to have motor imagery of sway around the ankle joint in ML and AP directions. In these two blocks, three minutes of training were considered, in which participants became familiar with the task by performing the sway with open eyes and then practicing the imagination of the sway with closed eyes. The order of performing the open and closed eyes tasks, as well as the blocks 2 and 3, were random for preventing the effects of precedence on the results. One minute of rest was considered between the blocks.

2. Experimental apparatus and procedure

We utilized the Zebris FDM-S force plate to capture the COP data at a sampling frequency of 120 Hz. Each recording session lasted 60 seconds. During the experiments, participants stood barefoot, positioning their feet together and keeping their arms by their sides. They were instructed to look straight ahead, maintain an erect posture for the neck and shoulders, and refrain from speaking.

The experiment consisted of four blocks, with a total duration of 15 minutes, as shown in Figure 1. In both the initial and final blocks, participants' postural control was assessed during standard standing under two conditions: with eyes open and eyes closed, each lasting one minute. For blocks 2 and 3, participants engaged in motor imagery of mediolateral (ML) and anteroposterior (AP) sways around the ankle joint. Within these blocks, a three-minute familiarization session was integrated. During this period, participants first executed the sway with their eyes open and then practiced the imagery of the sway with eyes closed. If the participant needed more practice, the training duration was extended by a minute. Participants were instructed to envision swaying around their ankle joint at the maximum angle where they felt comfortable. The ankle joint was selected due to its critical role in both postural control and gait. To mitigate any order effects on the outcomes, the sequence in which participants performed the eyes-open and eyes-closed tasks, as well as the tasks in blocks 2 and 3, was randomized. Participants were allotted a one-minute rest interval between blocks. Motor imagery was conducted with eyes closed to eliminate potential visual feedback that might influence visualization and to emphasize the role of proprioception. To ascertain that the effects noted in the MI blocks weren't simply a result of having the eyes closed, we compared postural control during motor imagery to postural control in both eyes-closed and eyes-open conditions.

3. Data analysis

The COP time series were analyzed in both the AP and ML directions, using the features outlined below.

PL is one of the most common features for analyzing the changes in the COP (33) and is calculated as:

Path length =
$$\sum_{i=1}^{N} \sqrt{(x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2}$$
 (1)

In equation 1, i represents the sample number, and x and y are the COP coordinates in the ML and AP directions, respectively.

2) Correlation dimension (CD)

Earlier research indicates that the center of pressure (COP) in a standing individual displays chaotic (34, 35). The correlation dimension is a feature commonly used in chaotic signal analysis. It quantifies the fractality of a signal by determining the average number of points within a specific radial distance (R) from each point in the phase space (36, 37). Through this method, we aim to delve into the nonlinear dynamics and chaotic attributes of the COP.

$$P_i(R) = \frac{1}{N-1} \sum_{j=1, j \neq i}^{N} \Theta(R - \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2})$$
(2)

$$C(R) =$$
(3)
$$\frac{1}{N(N-1)} \sum_{i=1}^{N} \sum_{j=1}^{N} \sum_{i=1}^{N} \sum_{j=1}^{N} \frac{\Theta\left(R - \sqrt{(x_j - x_j)^2 + (y_j - y_j)^2}\right)}{\Theta\left(R - \sqrt{(x_j - x_j)^2 + (y_j - y_j)^2}\right)}$$

$$D_c = \lim_{R \to 0} \frac{\log C(R)}{\log R}$$

$$(4)$$

In equations 2 and 3, x and y show the COP coordinates in the phase space. R is a small radius, Θ is a Heaviside function, and N denotes the total number of points. $P_i(R)$ is the possibility of points to lie in the given distance (R) from the i-th point. In equations 3 and 4, C(R) is the mean of $P_i(R)$ and, finally, D_c is the correlation dimension.

3) Mean velocity (MV)

MV of COP shifts in each direction was determined by averaging the displacement for each time interval, as follows:

$$Mean \ velocity = F \times mean \ |x_{i+1} - x_i| \tag{5}$$

In equation 5, x indicates the COP in ML or AP direction and i represents the sample number. F is the sampling frequency, which was 120 Hz in this study.

4) Root mean square (RMS)

We calculated the RMS for the mediolateral and anteroposterior COP signal.

$$RMS = \sqrt{\frac{x_1^2 + x_2^2 + \ldots + x_n^2}{n}} \tag{6}$$

In equation 6, x and n indicate the COP in each of ML or AP directions at each timepoint and the number of samples, respectively.

5) Range

Range was defined as the difference between the maximum and minimum amount of the COP during the test period. This was computed independently for both the mediolateral and anteroposterior COP signals.

$$Range = x_{\max} - x_{\min} \tag{7}$$

In equation 7, x indicates the COP in either direction.

6) Long-range correlation (LRC)

Detrended Fluctuation Analysis (DFA) gives a measure of long-range correlation in chaotic signals (38). In DFA, at first, the infinite function X is obtained from the x time series. Then, the resulting time series is divided into N nonoverlapping windows. The Y line is fitted on each window by the least squares method. The detrended time series for each window (i.e., F(n) in equations 9 and 10) is calculated as the difference between the time series values and the fitted line on each piece. This amount is calculated for different window lengths (i.e., t in equations 8 and 9). Finally, the slope of the logarithmic diagram of the detrended time series versus the window length is considered as α , which indicates the degree of self-similarity.

$$X_t = \sum_{i=1}^t \left(x_i - mean(x) \right) \tag{8}$$

$$F(n) = \sqrt{\frac{1}{N} \sum_{t=1}^{N} (X_t - Y_t)^2}$$

$$F(n) \propto n^{\alpha}$$
(9)
(10)

4. Statistical analysis

The statistical test was performed using repeated measures analysis of variance (ANOVA). The Mauchly test was used to examine the sphericity of the data, and in case of non-spherical data, the Greenhouse-Geisser adjustment corrected the degrees of freedom. After examining the results of the ANOVA test with the significance level of 0.05, which was obtained in all cases, Tukey's honest significant difference performed post hoc comparisons.

5. Model

Model description

We propose a COP dynamics model to illustrate postural control system and the effects of motor imagery on this system. We adapt Khanian et. al 's model for the human posture stability system (32) which captures the nonlinear dynamics of COP and can properly explain and quantify the COP changes. The proposed model is a two-dimensional discrete-time chaotic map. The process equation which is a model for creative processes (39) is used as the basis of the COP dynamics model (equation 11).

$$x_{k+1} = x_k + g \times \sin x_k \tag{11}$$

This map has a biotic behavior for a wide range of the parameter g with a lot of periodic and instability windows. Bios is an expansive pattern (40) and can model many structures, such as population dynamics, air temperature and visual perception (41, 42). Since the biotic pattern exists in the COP signal, this equation is appropriate for modeling the COP signal in each of AP and ML directions. Considering the interdependence of COP changes in both ML and AP directions, the equations for these directions were coupled. As a result, equation 12 emerges as the model for COP dynamics.

$$x_{k+1} = a \times e^{\frac{k}{q}} \times (x_k + b \times \sin y_k)$$
$$y_{k+1} = c \times e^{\frac{k}{q}} \times (y_k + d \times \sin x_k)$$
(12)

In this discrete-time map, x and y are the changes in the COP signal in the ML and AP directions, respectively. Other parameters such as a, b, c, d, and q are constants which should be identified using optimization methods. The COP's position at each time step is influenced by its preceding location. Moreover, postural control may decline due to physiological factors like fatigue and attention. This results in an increased range of COP changes, which is accounted for in this map through the exponential terms.

The model parameters are set as follows:

- 1) The bifurcation diagram was plotted for all parameters and the acceptable span for each of the parameters was determined. The range of COP changes in the recorded data was considered as a criterion for determining the acceptable span of parameters.
- 2) The genetic algorithm was used in order to set the parameters. In this method, the Euclidean distance was considered as the cost function. This method sets the parameters in a way that enhances the similarity between data and model phase spaces.

Effects of MI on COP dynamics

During the imagery of front-back sway, parameter d exhibits a greater increase than parameter b. Conversely, during lateral sway imagery, parameter b rises more than parameter d. Indeed, the dependence of the COP changes on the direction of the motor imagery is captured in the proposed model.

Model validation

To validate the model, we compared the range of COP changes between the model and the data, as illustrated in Table 2. The range of COP changes expands under closed eyes and motor imagery conditions. Further, when imagining the ML and AP sways, the range of changes in the ML and AP directions increases respectively. The model aptly captures this trend.

Table 2: A comparison between the range of changes in the model and the data. For both the model and the data, range of changes increased in the closed eyes and the motor imagery conditions. Plus, the range of changes in ML and AP directions was higher while imagining the ML and AP sways, respectively. NS: normal standing with open eyes. CE: normal standing with closed eyes. ML MI: mediolateral motor imagery. AP MI: anteroposterior motor imagery.

		\mathbf{NS}	CE	AP MI	ML MI
Range (Model)	Х	27.7	31.6	34.6	52.9
(Millimeter)	Y	21.3	29.8	46.6	30.7
Range (Data)	\mathbf{X}	26.8	33.45	35.2	54.8
(Millimeter)	Y	24.7	30.5	42.6	37.6

For the normal standing (NS) condition, we took the average of the range of changes from NS1 (normal standing with open eyes during the first block) and NS2 (normal standing with open eyes during the final block). Similarly, for the closed eyes (CE) condition, we considered the average range of changes between CE1 (normal standing with closed eyes in the initial block) and CE2 (normal standing with closed eyes in the concluding block).



Figure 2: A comparison between the phase space of COP for the data and the model. A) Phase space of COP for the data of a representative subject in the normal standing condition. B) Phase space of COP for the model with parameters a=0.8, b=4.4, c=0.7, d=5.8 and q=4999.7.

Figure 2 presents a comparison of the COP phase space between the data and the model. The data presented is the COP signal obtained from a representative subject.

Results

Relative to standing with closed eyes, the motor imagery condition exhibited an increase in MV, range, RMS, LRC, and PL, all pointing to an augmented COP displacement. A similar trend was observed when comparing standing with closed eyes to standing with open eyes. In the CE condition, the absence of visual feedback leads to greater COP displacement. Additionally, a decrease in the correlation dimension was noted when comparing the MI condition to both the CE and NS conditions, as well as when comparing CE to NS, indicating a reduction in the system's chaotic behavior. Typical postural control exhibits a certain level of variability, reflecting the body's adaptability and responsiveness to slight disturbances or environmental shifts. An overly regular postural control (i.e., reduced chaos) may hint at a diminished capacity for adaptation. Conversely, excessive chaotic behavior might imply instability. The observed decrease in chaotic behavior during CE and MI conditions in the results suggests a deviation from the normal postural control in these scenarios. Table 3 provides detailed information on the mean and standard deviation for each measure across all conditions.

Table 3: Mean of the measures in different conditions. MV is reported in millimeters per sec. Range and Path length are reported in Millimeters. NS1: normal standing with open eyes in the first block. NS2: normal standing with open eyes in the last block. CE1: normal standing with closed eyes in the first block. CE2: normal standing with closed eyes in the last block. ML MI: mediolateral motor imagery. AP MI: anteroposterior motor imagery.

		$\mathbf{NS1}$	CE1	APMI	MLMI	CE2	NS2
N / N /	Χ	58.4	84.8	78.8	139.3	81.7	57.7
IVI V	Y	48.1	70.3	0.3 104.9 83.4 7	70.5	48.8	
Range	\mathbf{X}	27.2	32.1	35.2	54.9	35.7	26.4
	Y	23.5	29.2	42.6	37.6	32.0	26.0
DMC	\mathbf{X}	4.8	5.8	6.9	10.8	6.8	5.2
RIVIS	Y	4.4	5.2	8.8	7.5	5.9	5.4
\mathbf{PL}		492.1	716.1	857.7	1043.5	705.04	493.6
CD		0.53	0.34	0.21	0.19	0.36	0.53
LRC	\mathbf{X}	0.07	0.14	0.13	0.29	0.14	0.09
	Y	0.10	0.19	0.38	0.27	0.19	0.12

MV in both the AP and ML directions, PL, correlation dimension, and LRC values in both AP and ML directions consistently show significant differences in the ML and CE scenarios compared to normal standing (p-value < 0.05). In the MI condition, PL is also consistently and significantly different from that in the CE condition. Furthermore, LRC in the ML direction is significantly different when performing ML MI, and not when performing AP MI, compared to standing with closed eyes. Similarly, LRC in the AP direction during AP MI shows a significant difference compared to standing with closed eyes.

Discussion

The primary objective of our study was to scrutinize the immediate effects of motor imagery (MI) on Center of Pressure (COP) dynamics within a population of healthy adults. Employing the model initially proposed by Khanian et al. (32), we formulated a specialized COP dynamics model that showcased the nonlinear intricacies of postural control during MI activities. Our results strongly validate the psycho-neuromuscular theory, revealing notable changes in variables like COP PL and Long-Range Correlation (LRC) during MI sessions as compared to closed eyes (CE) and normal standing (NS) conditions. The observed enhancements in COP displacement underline MI's substantial impact on postural control mechanisms.

Our findings have critical implications for rehabilitation, particularly for therapies centered on enhancing balance and postural stability. The model's adaptability for capturing individual postural dynamics holds promise for the development of patient-specific treatment protocols. This could be particularly beneficial for patients with compromised balance, such as stroke survivors, in optimizing MIbased therapeutic strategies. Beyond rehabilitation, our model parameters could function as viable biomarkers for not only gauging the efficacy of MI-centered interventions but also for evaluating postural control among varying populations, including athletes and the elderly.

Our findings indicate that alterations in the body's center of pressure (COP) align directionally with the imagined movements. Thus, the observed increase in COP changes during motor imagery is not attributable to mental load or the lack of a feedback mechanism. This insight has important implications for rehabilitation science, as it suggests that motor imagery can be strategically employed to elicit specific postural adjustments.

The results of this study align well with previous research, particularly the study conducted by Rodrigues et al. (17), who focused on motor imagery of bilateral plantar flexions. Rodrigues and colleagues observed that kinesthetic imagery significantly increased COP oscillations. These findings are consistent with other studies that have demonstrated marked changes in postural sway during MI tasks. Specifically, there is a noticeable increase in the center of pressure (COP) displacement when standing subjects engage in kinesthetic imagery, imagining themselves rising on their tiptoes (43). The congruence between our results and prior research further validates the influence of motor imagery on postural control and opens avenues for its application in rehabilitative settings.

While our findings offer substantial evidence supporting MI's influence on COP dynamics, they are not without limitations. One major limitation is the demographic homogeneity of our study group, comprising solely of healthy young adults. Future studies could enrich these findings by involving a more diverse participant pool, like the elderly or individuals with neurodegenerative disorders. Additionally, our study did not thoroughly account for external factors like fatigue and attention, which were only self-reported by participants. Subsequent research should aim to objectively measure these variables.

Looking ahead, it would be useful to reconcile the psycho-neuromuscular and central theories by examining the neural foundations of observed COP modifications. Longitudinal studies could further elucidate the long-term benefits of MI training on postural control, thereby cementing its therapeutic potential.

To sum up, our study provides robust evidence underscoring the role of motor imagery in affecting postural control, as demonstrated through measurable changes in COP dynamics. Leveraging a validated model, we have quantitatively corroborated these shifts, thus reinforcing MI's potential impact on enhancing postural stability. These findings lay a crucial foundation for future research and offer a substantive rationale for the integration of MI into rehabilitation programs, particularly those aiming at improving balance and postural control.

The ramifications of our work are especially noteworthy for the field of rehabilitative medicine. Given MI's non-invasive character and its proven efficacy in modifying COP dynamics, it emerges as a promising supplement to existing therapeutic strategies for improving postural stability, especially among those grappling with balance-related disorders or conditions. However, while the study sheds new light on MI's influence over postural control, it also indicates the need for more in-depth research to unpack the complex mechanisms governing this relationship. Enhanced understanding of these processes could streamline the optimization of MI-based treatments. Furthermore, broadening the scope to include various demographics is imperative for verifying the universal applicability and adaptability of MI-centered therapies.

In summary, our research marks a significant advancement in the application of motor imagery for rehabilitative therapies focused on balance and postural stability. It offers a robust framework, backed by empirical data, which future studies can refine and extend to understand the underlying mechanisms better and to assess MI's utility across diverse populations.

Implications on Physiotherapy Practice

The study demonstrates that motor imagery (MI) significantly impacts the dynamics of the Center of Pressure (COP), a key metric in evaluating postural stability. This finding can be directly applied in physiotherapy practices, especially in designing rehabilitation protocols for patients with balance and postural control issues. Physiotherapists can incorporate MI tasks into their treatment plans to induce specific postural adjustments, thus enhancing overall stability and balance in patients.

The research shows that MI can alter COP dynamics in a directionally consistent manner with the imagined movements. This insight is particularly beneficial for rehabilitation programs aimed at patients suffering from balance disorders, such as those recovering from stroke or neurological injuries. Tailoring MI exercises to the specific needs of these patients can aid in quicker and more effective recovery of postural control.

Given the non-invasive nature of MI, it emerges as a promising adjunctive tool in physiotherapy. It can be used alongside conventional physical therapies, especially in cases where physical movement might be limited or painful. This makes it an excellent option for early-stage rehabilitation or for patients with severe mobility restrictions.

The adaptability of the MI approach, as demonstrated by the study's model, suggests that physiotherapists can develop individualized treatment protocols. These personalized plans can be more effective as they can be designed to target the specific postural and balance issues of each patient, considering their unique COP dynamics.

The study's findings provide a quantitative basis for understanding the changes in postural control due to MI. This can help physiotherapists not only in planning treatment but also in monitoring progress. The measurable changes in COP dynamics offer a way to objectively assess the effectiveness of the MI-based interventions over time.

The study highlights the need for further research, particularly involving diverse demographics and long-term effects. This opens up avenues for physiotherapists to engage in or follow up on ongoing research, keeping abreast of the latest developments in the field which can then be translated into practice.

In conclusion, this study's findings on the effects of motor imagery on postural control offer valuable insights and practical applications for physiotherapy practice, particularly in enhancing balance and postural stability in various patient populations.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT in order to improve the readability of the manuscript. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

Acknowledgments

The authors thank Dr. Nasser Fatouraee and Arman Gholizadeh from the Biomechanics department of Amirkabir University of Technology, Tehran, Iran

Conflict of interest

None declared.

References

1. Massion J. Postural control system. Current opinion in neurobiology. 1994;4(6):877-87.

2. Dickstein R, Deutsch JE. Motor imagery in physical therapist practice. Physical therapy. 2007;87(7):942-53.

3. Grangeon M, Guillot A, Collet C. Postural control during visual and kinesthetic motor imagery. Applied psychophysiology and biofeedback. 2011;36:47-56.

4. Mulder T. Motor imagery and action observation: cognitive tools for rehabilitation. Journal of neural transmission. 2007;114:1265-78.

5. de Souza NS, Martins ACG, da Silva Canuto K, Machado D, Teixeira S, Orsini M, et al. Postural control modulation during motor imagery tasks: A systematic review. International Archives of Medicine. 2015;8.

6. Hecker JE, Kaczor LM. Application of imagery theory to sport psychology: Some preliminary findings. Journal of Sport and Exercise Psychology. 1988;10(4):363-73.

7. Morris T, Spittle M, Watt AP. Imagery in sport: Human Kinetics; 2005.

8. Oh D-S, Choi J-D. The effect of motor imagery training for trunk movements on trunk muscle control and proprioception in stroke patients. Journal of physical therapy science. 2017;29(7):1224-8.

9. Cho H-y, Kim J-s, Lee G-C. Effects of motor imagery training on balance and gait abilities in post-stroke patients: a randomized controlled trial. Clinical rehabilitation. 2013;27(8):675-80.

10. Fujikawa S, Ohsumi C, Ushio R, Tamura K, Sawai S, Yamamoto R, Nakano H. Potential Applications of Motor Imagery for Improving Standing Posture Balance in Rehabilitation. Neurorehabilitation and Physical Therapy: IntechOpen; 2022.

11. Sarasso E, Agosta F, Piramide N, Gardoni A, Canu E, Leocadi M, et al. Action observation and motor imagery improve dual task in Parkinson's disease: a clinical/fMRI study. Movement Disorders. 2021;36(11):2569-82.

12. Chiacchiero M, Cagliostro P, DeGenaro J, Giannina C, Rabinovich Y. Motor imagery improves balance in older adults. Topics in Geriatric Rehabilitation. 2015;31(2):159-63.

13. Borah B, Yadav A. Effect of Psychoneuromuscular Theory and Visualization Technique in Reducing Anxiety Levels of Female Soccer Players in Competition Situations. International Journal of Behavioral Social and Movement Sciences. 2017;6(2):1-6.

14. Livesay JR, Samaras MR. Covert neuromuscular activity of the dominant forearm during visualization of a motor task. Perceptual and motor skills. 1998;86(2):371-4.

15. Bakker FC, Boschker MSJ, Chung T. Changes in muscular activity while imagining weight lifting using stimulus or response propositions. Journal of Sport and Exercise Psychology. 1996;18(3):313-24.

16. Krüger B, Hettwer M, Zabicki A, de Haas B, Munzert J, Zentgraf K. Practice modality of motor sequences impacts the neural signature of motor imagery. Scientific reports. 2020;10(1):19176-.

17. Rodrigues EC, Lemos T, Gouvea B, Volchan E, Imbiriba LA, Vargas CD. Kinesthetic motor imagery modulates body sway. Neuroscience. 2010;169(2):743-50.

18. Stins JF, Schneider IK, Koole SL, Beek PJ. The influence of motor imagery on postural sway: differential effects of type of body movement and person perspective. Advances in Cognitive Psychology. 2015;11(3):77-.

19. Stins JF, Michielsen ME, Roerdink M, Beek PJ. Sway regularity reflects attentional involvement in postural control: Effects of expertise, vision and cognition. Gait & posture. 2009;30(1):106-9.

20. Mouthon A, Ruffieux J, Mouthon M, Hoogewoud HM, Annoni JM, Taube W. Age-related differences in cortical and subcortical activities during observation

and motor imagery of dynamic postural tasks: an fMRI study. Neural plasticity. 2018;2018.

21. Jahn K, Deutschländer A, Stephan T, Strupp M, Wiesmann M, Brandt T. Brain activation patterns during imagined stance and locomotion in functional magnetic resonance imaging. Neuroimage. 2004;22(4):1722-31.

22. Taube W, Mouthon M, Leukel C, Hoogewoud H-M, Annoni J-M, Keller M. Brain activity during observation and motor imagery of different balance tasks: an fMRI study. cortex. 2015;64:102-14.

23. Alvarenga R, Porto F, Braga R, Cantreva R, Espinosa G, Itaborahy A, et al., editors. Construction and calibration of a low-cost force plate for human balance evaluation. ISBS-Conference Proceedings Archive; 2011.

24. Pasma JH, Boonstra TA, Van Kordelaar J, Spyropoulou VV, Schouten AC. A sensitivity analysis of an inverted pendulum balance control model. Frontiers in computational neuroscience. 2017;11:99-.

25. Maurer C, Peterka RJ. A new interpretation of spontaneous sway measures based on a simple model of human postural control. Journal of neurophysiology. 2005;93(1):189-200.

26. Kot A, Nawrocka A, editors. Modeling of human balance as an inverted pendulum. Proceedings of the 2014 15th International Carpathian Control Conference (ICCC); 2014: IEEE.

27. Mergner T, Maurer C, Peterka RJ. A multisensory posture control model of human upright stance. Progress in brain research. 2003;142:189-201.

28. Milton J, Cabrera JL, Ohira T, Tajima S, Tonosaki Y, Eurich CW, Campbell SA. The time-delayed inverted pendulum: implications for human balance control. Chaos: An Interdisciplinary Journal of Nonlinear Science. 2009;19(2).

29. Asai Y, Tasaka Y, Nomura K, Nomura T, Casadio M, Morasso P. A model of postural control in quiet standing: robust compensation of delay-induced instability using intermittent activation of feedback control. PLoS One. 2009;4(7):e6169-e.

30. Hilts WW, Szczecinski NS, Quinn RD, Hunt AJ, editors. Simulation of human balance control using an inverted pendulum model. Biomimetic and Biohybrid Systems: 6th International Conference, Living Machines 2017, Stanford, CA, USA, July 26–28, 2017, Proceedings 6; 2017: Springer.

31. Suzuki Y, Nomura T, Casadio M, Morasso P. Intermittent control with ankle, hip, and mixed strategies during quiet standing: a theoretical proposal based on a double inverted pendulum model. Journal of theoretical biology. 2012;310:55-79.

32. Khanian MYA, Golpayegni SMRH, Rostami M. A new multi-attractor model for the human posture stability system aimed to follow self-organized dynamics. Biocybernetics and Biomedical Engineering. 2020;40(1):162-72.

33. Donker SF, Roerdink M, Greven AJ, Beek PJ. Regularity of center-ofpressure trajectories depends on the amount of attention invested in postural control. Experimental brain research. 2007;181:1-11.

34. Yamada N. Chaotic swaying of the upright posture. Human movement science. 1995;14(6):711-26.

35. Pascolo PB, Marini A, Carniel R, Barazza F. Posture as a chaotic system and an application to the Parkinson's disease. Chaos, Solitons & Fractals. 2005;24(5):1343-6.

36. Cimolin V, Galli M, Rigoldi C, Grugni G, Vismara L, Mainardi L, Capodaglio P. Fractal dimension approach in postural control of subjects with Prader-Willi Syndrome. Journal of neuroengineering and rehabilitation. 2011;8:1-6.

37. Nygård JF, Glattre E. Fractal analysis of time series in epidemiology: Is there information hidden in the noise? Norsk Epidemiologi. 2003;13(2).

38. González-Salas JS, Shbat MS, Ordaz-Salazar FC, Simón J. Analyzing chaos systems and fine spectrum sensing using detrended fluctuation analysis algorithm. Mathematical Problems in Engineering. 2016;2016.

39. Sabelli H. Bios: a study of creation (With CD-ROM): World Scientific; 2005.

40. Sabelli H, Kovacevic L. Biotic complexity of population dynamics. Complexity. 2008;13(4):47-55.

41. Patel M, Sabelli H. Autocorrelation and frequency analysis differentiate cardiac and economic bios from 1/f noise. Kybernetes. 2003;32(5/6):692-702.

42. Beigzadeh M, Golpayegani SMRH, editors. A macroscopic chaotic model of visual perception. 2016 23rd Iranian Conference on Biomedical Engineering and 2016 1st International Iranian Conference on Biomedical Engineering (ICBME); 2016: IEEE.

43. Lemos T, Rodrigues EC, Vargas CD. Motor imagery modulation of postural sway is accompanied by changes in the EMG–COP association. Neuroscience Letters. 2014;577:101-5.