

The Isospectral Problem for p -widths: An Application of Zoll Metrics

Jared Marx-Kuo

May 2024

Abstract

We pose the isospectral problem for the p -widths: Is a riemannian manifold (M^n, g) uniquely determined by its p -widths, $\omega_p(M, g)$? We construct many counterexamples on S^2 using Zoll metrics and the fact that geodesic p -widths are given by unions of immersed geodesics.

1 Introduction

In [Gro06], Gromov introduced the p -widths, $\{\omega_p\}$, of a riemannian manifold, (M^{n+1}, g) , as an analogue of the spectrum of the laplacian. Intuitively, one replaces the rayleigh quotient in the definition of eigenvalues with the area functional in the definition of ω_p - see [Gro10] [Gro02] [Gut09] [FNT⁺20] for more background. The p -widths have proven to be extremely useful due to the Almgren–Pitts/Marques–Neves Morse theory program for the area functional (see [MN14] [MMN23] [MN15] [MN21] [AJ62] [Zho20]). In [LMN18], Liokumovich–Marqueves–Neves demonstrated that $\{\omega_p\}$ satisfy a Weyl–law, which later lead to the resolution of Yau’s conjecture: on any closed manifold (M^{n+1}, g) , there exist infinitely many minimal hypersurfaces (see [CM20] [Son18] [MNS19] [MN17] [IMN18] [Li23]). We refer the reader to the introduction of [CM23] for a more thorough overview of the history and applications of the p -widths.

Recall the isospectral problem for the laplacian

Problem 1. *Is a closed manifold, (M^n, g) , uniquely determined by its discrete spectrum, $\{\lambda_k\}$?*

Informally, this is known as “can you hear the sound of a drum?” and was made famous by Kac [Kac66]. In [Mil64], Milnor constructed two 16-dimensional tori which are isospectral but not isometric, providing the first counterexample to 1. Gordon–Webb–Wolpert [GWW92] constructed a simpler counterexample using polygons in \mathbb{R}^2 . Hersch ([Her70]) showed that the round metric on S^2 is isospectrally unique by proving an isoperimetric inequality for the first eigenvalue.

Given the analogy between $\{\omega_p\}$ and $\{\lambda_p\}$, It is natural to pose the p -width isospectral problem:

Problem 2. *Is a closed manifold, (M^n, g) , uniquely determined by the values $\omega_p(g)$?*

We show that the answer is “no”:

Theorem 1.1. *For any $p > 0$, ω_p is constant on any fixed connected component of the space of Zoll metrics on S^2 .*

As a corollary, we can compute the p -widths explicitly for new metrics.

Corollary 1.1.1. *Let g be any Zoll metric on S^2 which lies in the connected component of g_{round} in the space of Zoll metrics on S^2 . Then $\omega_p(g) = \omega_p(g_{\text{round}}) = 2\pi\lfloor\sqrt{p}\rfloor$ for all p .*

The above gives rise to a continuous deformation of p -width isospectral metrics (compare [GW84]). In particular, any odd function $f : S^2 \rightarrow \mathbb{R}$ gives rise to a 1-parameter family of geometrically distinct Zoll metrics near the round metric (see [Gui76]). The p -widths of (S^2, g_{round}) were computed explicitly in [CM23], and this was the first example of the p -width spectrum being known for all p . Corollary 1.1.1 now gives uncountably more manifolds where the full width spectrum is known.

The proof of theorem 1.1 uses the defining properties of zoll metrics on S^2 , which are metrics for which all geodesics (with multiplicity one) are simple and length 2π (note this choice of length, as opposed to any constant c , agrees with [Gui76]). These were first constructed by Zoll [Zol01], and we refer the reader to [Gui76] [Fun11] [GG81] [Bes12] for a by no means a complete list of background sources. The proof also utilizes a recent result of Chodosh–Mantoulidis [CM23], which states that each $\omega_p(M^2, g)$ is achieved by a union of closed, potentially immersed geodesics.

This paper is organized as follows: in §2, we briefly define the p -widths, and recall the results of Chodosh–Mantoulidis [CM23]. We also define Zoll metrics and list some relevant properties. In §3 we prove the main theorem, and in §4 we pose some open questions. The author thanks Otis Chodosh for his conversations on p -widths and Lucas Ambrosio for his conversations on Zoll metrics.

2 Background

Let $X \subseteq I^k$ denote a cubical subcomplex.

Definition 2.1 ([MN17], Defn 4.1). A map $\Phi : X \rightarrow \mathcal{Z}_1(M; \mathbb{Z}_2)$ is a *p-sweepout* if it is continuous (with the standard flat norm topology on $\mathcal{Z}_1(M; \mathbb{Z}_2)$) and $\Phi^*(\bar{\lambda}^p) \neq 0$.

Definition 2.2 ([MN17] §3.3). A map $\Phi : X \rightarrow \mathcal{Z}_1(M; \mathbb{Z}_2)$ is said to have **no concentration of mass** if

$$\limsup_{r \rightarrow 0} \{\|\Phi(x)\|(B_r(p)) : x \in X, p \in M\} = 0.$$

Definition 2.3 ([Gro02], [Gut09]). We define $\mathcal{P}_p = \mathcal{P}_p(M)$ to be the set of all *p-sweepouts*, out of any cubical subcomplex X , with no concentration of mass. The *p-width* of (M, g) is

$$\omega_p(M, g) := \inf_{\Phi \in \mathcal{P}_p} \sup\{\mathbf{M}(\Phi(x)) : x \in \text{domain}(\Phi)\}.$$

To see the analogy between ω_p and λ_p , consider the following characterization of $\lambda_p(M)$ as

$$\lambda_p(M) := \inf_{\substack{V \subseteq C^\infty(M) \\ \dim V = p+1}} \sup_{f \in V \setminus \{0\}} \frac{\int_M |\nabla f|^2}{\int_M f^2}$$

Note the parallels between the min-max definitions of ω_p, λ_p , with $\Phi^*(\bar{\lambda}^p) \neq 0$ being analogous to $\dim(V) = p + 1$.

Having defined the *p-widths*, note that when $n + 1 = 2$, the widths may be achieved by geodesic networks a priori. In [CM23], Chodosh–Mantoulidis used the Allen–Cahn equation to prove that the *p-widths* can always be achieved by a union of closed geodesics:

Theorem 2.4 ([CM23], Thm 1.2). Let (M^2, g) be a closed Riemannian manifold. For every $p \in \mathbb{Z}^+$, there exists a collection of primitive, closed geodesics $\{\sigma_{p,i}\}$ such that

$$\omega_p(M, g) = \sum_{j=1}^{N(p)} m_j \cdot L_g(\sigma_{p,j}), \quad m_j \in \mathbb{Z}^+$$

Here, “primitive” means that the geodesic is connected and transversed with multiplicity one. They also showed:

Theorem 2.5 ([CM23], Thm 1.4). For g_0 the unit round metric on S^2

$$\forall p \in \mathbb{N}, \quad \omega_p(S^2, g_0) = 2\pi \lfloor \sqrt{p} \rfloor$$

We now collect relevant facts about Zoll metrics on S^2 .

Definition 2.6. A metric, g , on S^2 is called a *Zoll metric* if all of the primitive geodesics are closed and length 2π .

Zoll [Zol01], Funk [Fun11], Guillemin [Gui76], and Gromoll–Groß [GG81] each contributed significantly to the theory of Zoll metrics. In particular, Gromoll–Groß showed that all of the primitive geodesics on a Zoll surface are simple. Furthermore, the combined work of Guillemin and Funk demonstrated that the tangent space of Zoll metrics at the round metric is given by odd functions $f : S^2 \rightarrow \mathbb{R}$. We also mention a higher dimensional analogue of these metrics constructed by Ambrozio–Marques–Neves [AMN21].

Let $\mathcal{Z}(S^2)$ denote the space of Zoll metrics on S^2 , \mathcal{Z}_α , an arbitrary connected component of $\mathcal{Z}(S^2)$, and \mathcal{Z}_0 , the connected component containing g_0 , the unit round metric.

3 Proof of Main theorem

Let $\mathcal{M}_\infty(S^2)$ denote the set of all smooth metrics on S^2 , and let g_0 denote the round metric. For fixed p , $\omega_p : \mathcal{M}_\infty(S^2) \rightarrow \mathbb{R}$ is continuous due to the following lemma of Marques–Neves–Song:

Lemma 3.1 ([MNS19], Lemma 1). Let $C_1 < C_2$. There exists a $K(C_1, C_2)$, such that

$$|\omega_p(g) - \omega_p(g')| \leq K(C_1, C_2) p^{1/2} \|g - g'\|_{C^0(g_0)}$$

for any g, g' such that $C_1 g_0 \leq g, g' \leq C_2 g_0$.

Take any Zoll metric, g , on S^2 . Applying theorem 2.4, we have

$$\begin{aligned}\omega_p(S^2, g) &= \sum_{j=1}^{N(p)} m_j \cdot L_g(\sigma_{p,j}) \\ &= 2\pi \sum_{j=1}^{N(p)} m_j \in 2\pi\mathbb{Z}^+\end{aligned}$$

since each $\sigma_{p,j}$ is primitive and $L_g(\sigma_{p,j}) = 2\pi$ by definition of Zoll metric. Let \mathcal{Z}_α a fixed, connected component of \mathcal{Z} . Consider the restriction of $\omega_p : \mathcal{Z}_\alpha \rightarrow \mathbb{R}^\infty$. Its image is discrete, lying inside $2\pi\mathbb{N}$, but ω_p is continuous in each component due to lemma 3.1, thus the map is constant on connected components, proving theorem 1.1.

4 Open Questions

Having shown a counterexample to the isospectral p -width problem, it would be interesting to know any of the following

- Do there exist counterexamples to the isospectral problem in dimensions $3 \leq n + 1 \leq 7$? The p -widths behave differently in these dimensions compared to $n + 1 = 2$, as demonstrated by theorem 2.4. See [Aie16] for further differences relating to multiplicity and index bounds.
- Do there exist counterexamples to the isospectral problem on other surfaces (Σ^2, g) ?
- Do there exist other reasonable geometric conditions that one can impose on a manifold so that isospectral rigidity holds?

References

- [Aie16] Nicolau Sarquis Aieix, *The width of ellipsoids*, arXiv preprint arXiv:1601.01032 (2016).
- [AJ62] Frederick Justin Almgren Jr, *The homotopy groups of the integral cycle groups*, *Topology* **1** (1962), no. 4, 257–299.
- [AMN21] Lucas Ambrozio, Fernando C Marques, and André Neves, *Riemannian metrics on the sphere with Zoll families of minimal hypersurfaces*, arXiv preprint arXiv:2112.01448 (2021).
- [Bes12] Arthur L Besse, *Manifolds all of whose geodesics are closed*, vol. 93, Springer Science & Business Media, 2012.
- [CM20] Otis Chodosh and Christos Mantoulidis, *Minimal surfaces and the Allen–Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates*, *Annals of Mathematics* **191** (2020), no. 1, 213–328.
- [CM23] ———, *The p -widths of a surface*, *Publications mathématiques de l’IHÉS* **137** (2023), no. 1, 245–342.
- [FNT⁺20] Ailana Fraser, André Neves, Peter M Topping, Paul C Yang, Fernando C Marques, and André Neves, *Applications of min–max methods to geometry*, *Geometric Analysis: Cetraro, Italy 2018* (2020), 41–77.
- [Fun11] Paul Funk, *Über flächen mit lauter geschlossenen geodätischen linien*, W. Fr. Kaestner, 1911.
- [GG81] Detlef Gromoll and Karsten Grove, *On metrics on S^2 all of whose geodesics are closed*, *Inventiones mathematicae* **65** (1981), no. 1, 175–177.
- [Gro02] Mikhail Gromov, *Isoperimetry of waists and concentration of maps*, Tech. report, SIS-2003-246, 2002.
- [Gro06] Misha Gromov, *Dimension, non-linear spectra and width*, *Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1986–87*, Springer, 2006, pp. 132–184.
- [Gro10] Mikhail Gromov, *Singularities, expanders and topology of maps. part 2: From combinatorics to topology via algebraic isoperimetry*, *Geometric and Functional Analysis* **20** (2010), no. 2, 416–526.

- [Gui76] Victor Guillemin, *The radon transform on Zoll surfaces*, Advances in Mathematics **22** (1976), no. 1, 85–119.
- [Gut09] Larry Guth, *Minimax problems related to cup powers and Steenrod squares*, Geometric And Functional Analysis **18** (2009), 1917–1987.
- [GW84] Carolyn S Gordon and Edward N Wilson, *Isospectral deformations of compact solvmanifolds*, Journal of Differential Geometry **19** (1984), no. 1.
- [GWW92] Carolyn Gordon, David Webb, and Scott Wolpert, *Isospectral plane domains and surfaces via Riemannian orbifolds*, Inventiones mathematicae **110** (1992), no. 1, 1–22.
- [Her70] Joseph Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, CR Acad. Sci. Paris S\`{e}r. AB **270** (1970), A1645–A1648.
- [IMN18] Kei Irie, Fernando Marques, and André Neves, *Density of minimal hypersurfaces for generic metrics*, Annals of Mathematics **187** (2018), no. 3, 963–972.
- [Kac66] Mark Kac, *Can one hear the shape of a drum?*, The American Mathematical Monthly **73** (1966), no. 4P2, 1–23.
- [Li23] Yangyang Li, *Existence of infinitely many minimal hypersurfaces in higher-dimensional closed manifolds with generic metrics*, Journal of Differential Geometry **124** (2023), no. 2, 381–395.
- [LMN18] Yevgeny Liokumovich, Fernando Marques, and André Neves, *Weyl law for the volume spectrum*, Annals of Mathematics **187** (2018), no. 3, 933–961.
- [Mil64] John Milnor, *Eigenvalues of the Laplace operator on certain manifolds*, Proceedings of the National Academy of Sciences **51** (1964), no. 4, 542–542.
- [MMN23] Fernando C Marques, Rafael Montezuma, and André Neves, *Morse inequalities for the area functional*, Journal of Differential Geometry **124** (2023), no. 1, 81–111.
- [MN14] Fernando C Marques and André Neves, *Min-max theory and the Willmore conjecture*, Annals of Mathematics (2014), 683–782.
- [MN15] ———, *Morse index and multiplicity of min-max minimal hypersurfaces*, arXiv preprint arXiv:1512.06460 (2015).
- [MN17] ———, *Existence of infinitely many minimal hypersurfaces in positive Ricci curvature*, Inventiones mathematicae **209** (2017), no. 2, 577–616.
- [MN21] ———, *Morse index of multiplicity one min-max minimal hypersurfaces*, Advances in Mathematics **378** (2021), 107527.
- [MNS19] Fernando C Marques, André Neves, and Antoine Song, *Equidistribution of minimal hypersurfaces for generic metrics*, Inventiones mathematicae **216** (2019), no. 2, 421–443.
- [Son18] Antoine Song, *Existence of infinitely many minimal hypersurfaces in closed manifolds*, arXiv preprint arXiv:1806.08816 (2018).
- [Zho20] Xin Zhou, *On the multiplicity one conjecture in min-max theory*, Annals of Mathematics **192** (2020), no. 3, 767–820.
- [Zol01] Otto Zoll, *Ueber flächen mit scharen von geschlossenen geodätischen linien*, Dieterich, 1901.