

Equivalence between non-commutative harmonic oscillators and two-photon quantum Rabi models

Ryosuke Nakahama^{*†}

*NTT Institute for Fundamental Mathematics,
NTT Communication Science Laboratories,
Nippon Telegraph and Telephone Corporation,
3-9-11 Midori-cho, Musashino-shi, Tokyo 180-8585, Japan*

May 31, 2024

Abstract

We prove that the non-commutative harmonic oscillator on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$ introduced by Parmeggiani and Wakayama is equivalent to the two-photon quantum Rabi model, and they are also equivalent to a holomorphic differential equation on the unit disk. The confluence process of this differential equation and the relation with the one-photon quantum Rabi model are also discussed.

2020 Mathematics Subject Classification: 81Q10; 35P05; 34M46.

1 Introduction

The *quantum Rabi model* (QRM) is a light-matter interaction model introduced by Jaynes and Cummings [6]. Its Hamiltonian is given by the formally self-adjoint operator on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$,

$$H_{\text{QRM}}^{(g,\Delta)} := \mathbf{I}a^\dagger a + g\boldsymbol{\sigma}_1(a + a^\dagger) + \Delta\boldsymbol{\sigma}_3,$$

where $g, \Delta \in \mathbb{R}$, a^\dagger, a are the creation and annihilation operators on $L^2(\mathbb{R})$, and $\boldsymbol{\sigma}_1, \boldsymbol{\sigma}_3 \in M(2, \mathbb{C})$ are the Pauli matrices. The integrability of this model is proved by Braak [1, 2]. This Hamiltonian has discrete spectra. Then as a next step, the *two-photon quantum Rabi model* (or a special case of the *spin-boson model*)

$$H_{\text{2QRM}}^{(g,\Delta)} := \mathbf{I}a^\dagger a + g\boldsymbol{\sigma}_1(a^2 + (a^\dagger)^2) + \Delta\boldsymbol{\sigma}_3,$$

is considered (see e.g. [3, 9, 20] and references therein). This Hamiltonian has discrete spectra for $|g| < \frac{1}{2}$, but may have both discrete and continuous spectra for $|g| = \frac{1}{2}$.

Independently, as a purely mathematical model, Parmeggiani and Wakayama [16, 17, 18] introduced the *non-commutative harmonic oscillator* (NCHO), with the Hamiltonian operator on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$,

$$H_{\text{NCHO}}^{(\alpha,\beta)} := \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} \left(-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 \right) + \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \left(x \frac{d}{dx} + \frac{1}{2} \right),$$

^{*}This work was supported by JST CREST Grant Number JPMJCR2113, Japan.

[†]Email: ryosuke.nakahama@ntt.com

where $\alpha, \beta \in \mathbb{R}$. If $\alpha\beta > 1$, then this has discrete spectra, and if $\alpha\beta = 1$, then this has continuous spectra [14, 15]. The spectra of NCHO is studied mathematically also via the spectral zeta functions (see [5, 8, 13] and references therein). By [11, 12, 19, 21], the restriction of NCHO on the space of even or odd functions is equivalent to a Heun differential equation. On the other hand, by [2, 7], the (one-photon) QRM is equivalent to a confluent Heun differential equation, and NCHO is considered as a covering model of QRM via the confluence process of the Heun equation [19, 21]. However, this confluence process seems complicated and difficult to explain reasonably.

In this article, we prove that NCHO is equivalent to the two-photon QRM, and that their restriction to the space of even or odd functions is also equivalent to a holomorphic differential equation on the unit disk. We also discuss the confluence process of this holomorphic differential equation. Then we find that considering the two-photon QRM is more reasonable as a covering model of the one-photon QRM than considering NCHO. We note that the equivalence of NCHO and a holomorphic differential equation on the unit disk holds in a more general setting. See [10].

2 Preliminaries

First, let a, a^\dagger, H be the unbounded operators on $L^2(\mathbb{R})$ given by

$$a := \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right), \quad a^\dagger := \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right), \quad H := a^\dagger a + \frac{1}{2} = \frac{1}{2} \left(x^2 - \frac{d^2}{dx^2} \right).$$

Then for $\theta \in \mathbb{R}$, the unitary operator $e^{i\theta H}$ on $L^2(\mathbb{R})$ is given by

$$\begin{aligned} e^{i\theta H} f(x) &= \frac{1}{\sqrt{2\pi \cos \theta}} \int_{\mathbb{R}} e^{i(\tan \theta)(x^2 + \xi^2)/2 + i(\sec \theta)x\xi} \widehat{f}(\xi) d\xi & (\cos \theta \neq 0) \\ &= \frac{\sqrt{i}}{\sqrt{2\pi \sin \theta}} \int_{\mathbb{R}} e^{-i(\cot \theta)(x^2 + \xi^2)/2 + i(\csc \theta)x\xi} f(\xi) d\xi & (\sin \theta \neq 0), \end{aligned}$$

where $\widehat{f}(\xi) := \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-ix\xi} dx$ is the Fourier transform, if we choose the branches of $\sqrt{\cos \theta}, \sqrt{\sin \theta}$ suitably. This has the periodicity 4π (see e.g. [4, Corollary 4.55]).

Next, let $\mathcal{F}(\mathbb{C}) \subset \mathcal{O}(\mathbb{C})$ be the Fock space,

$$\mathcal{F}(\mathbb{C}) := \left\{ f \in \mathcal{O}(\mathbb{C}) \mid \frac{1}{\pi} \int_{\mathbb{C}} |f(w)|^2 e^{-|w|^2} dw < \infty \right\}.$$

Then $L^2(\mathbb{R})$ and $\mathcal{F}(\mathbb{C})$ are unitarily isomorphic by the Bargmann transform

$$\mathcal{B}: L^2(\mathbb{R}) \longrightarrow \mathcal{F}(\mathbb{C}), \quad (\mathcal{B}f)(w) := \pi^{-1/4} \int_{\mathbb{R}} f(x) e^{-(x^2 + w^2)/2 + \sqrt{2}xw} dx.$$

Then this satisfies

$$(\mathcal{B}af)(w) = \frac{\partial}{\partial w} (\mathcal{B}f)(w), \quad (\mathcal{B}a^\dagger f)(w) = w(\mathcal{B}f)(w), \quad (\mathcal{B}e^{i\theta H} f)(w) = e^{i\theta/2} (\mathcal{B}f)(e^{i\theta} w).$$

The orthonormal basis $\left\{ \frac{\pi^{-1/4}}{\sqrt{m!}} (a^\dagger)^m e^{-x^2/2} \right\} \subset L^2(\mathbb{R})$ is mapped to $\left\{ \frac{w^m}{\sqrt{m!}} \right\} \subset \mathcal{F}(\mathbb{C})$ by \mathcal{B} (see e.g. [4, Sections 1.6, 1.7]).

Next, let $\mathbf{H}, \mathbf{D} \subset \mathbb{C}$ be the upper half plane and the unit disk,

$$\mathbf{H} := \{y \in \mathbb{C} \mid \operatorname{Im} y > 0\}, \quad \mathbf{D} := \{z \in \mathbb{C} \mid |z| < 1\}.$$

For $\nu > 0$, let $\langle \cdot, \cdot \rangle_\nu$ be the weighted Bergman inner product on the space $\mathcal{O}(\mathbf{D})$ of holomorphic functions on \mathbf{D} given by, for $F(z) = \sum_{m=0}^{\infty} a_m z^m$, $G(z) = \sum_{m=0}^{\infty} b_m z^m$,

$$\begin{aligned} \langle F, G \rangle_\nu &:= \sum_{m=0}^{\infty} \frac{m!}{(\nu)_m} a_m \overline{b_m} & (\nu > 0) \\ &= \frac{\nu-1}{\pi} \int_{\mathbf{D}} F(z) \overline{G(z)} (1-|z|^2)^{\nu-2} dz & (\nu > 1), \end{aligned}$$

where $(\nu)_m := \nu(\nu+1)(\nu+2) \cdots (\nu+m-1)$, and let

$$\mathcal{H}_\nu(\mathbf{D}) := \{F(z) \in \mathcal{O}(\mathbf{D}) \mid \langle F, F \rangle_\nu < \infty\}.$$

Next, let $\mathcal{L}_{\text{even}} = \mathcal{L}_{1/2}$, $\mathcal{L}_{\text{odd}} = \mathcal{L}_{3/2}$ be the Laplace transforms

$$\begin{aligned} \mathcal{L}_{\text{even}} = \mathcal{L}_{1/2}: L^2(\mathbb{R})_{\text{even}} &\longrightarrow \mathcal{O}(\mathbf{H}), & (\mathcal{L}_{\text{even}} f)(y) &:= \pi^{-1/4} \int_{\mathbb{R}} f(x) e^{x^2 y i/2} dx, \\ \mathcal{L}_{\text{odd}} = \mathcal{L}_{3/2}: L^2(\mathbb{R})_{\text{odd}} &\longrightarrow \mathcal{O}(\mathbf{H}), & (\mathcal{L}_{\text{odd}} f)(y) &:= \sqrt{2} \pi^{-1/4} \int_{\mathbb{R}} f(x) x e^{x^2 y i/2} dx, \end{aligned}$$

let $\Phi_{\text{even}} = \Phi_{1/2}$, $\Phi_{\text{odd}} = \Phi_{3/2}$ be the integral transforms

$$\begin{aligned} \Phi_{\text{even}} = \Phi_{1/2}: \mathcal{F}(\mathbb{C})_{\text{even}} &\longrightarrow \mathcal{O}(\mathbf{D}), & (\Phi_{\text{even}} f)(z) &:= \frac{1}{\pi} \int_{\mathbb{C}} f(w) e^{z \overline{w}^2/2} e^{-|w|^2} dw, \\ \Phi_{\text{odd}} = \Phi_{3/2}: \mathcal{F}(\mathbb{C})_{\text{odd}} &\longrightarrow \mathcal{O}(\mathbf{D}), & (\Phi_{\text{odd}} f)(z) &:= \frac{1}{\pi} \int_{\mathbb{C}} f(w) \overline{w} e^{z \overline{w}^2/2} e^{-|w|^2} dw, \end{aligned}$$

and for $\nu \in \mathbb{C}$, let \mathcal{C}_ν be the weighted Cayley transform,

$$\mathcal{C}_\nu: \mathcal{O}(\mathbf{H}) \longrightarrow \mathcal{O}(\mathbf{D}), \quad (\mathcal{C}_\nu F)(z) := (1+z)^{-\nu} F\left(i \frac{1-z}{1+z}\right),$$

so that the following diagrams commute.

$$\begin{array}{ccc} L^2(\mathbb{R})_{\text{even}} & \xrightarrow{\mathcal{B}} & \mathcal{F}(\mathbb{C})_{\text{even}} \\ \mathcal{L}_{\text{even}} = \mathcal{L}_{1/2} \downarrow & & \downarrow \Phi_{\text{even}} = \Phi_{1/2} \\ \mathcal{O}(\mathbf{H}) & \xrightarrow{\mathcal{C}_{1/2}} & \mathcal{O}(\mathbf{D}), \end{array} \quad \begin{array}{ccc} L^2(\mathbb{R})_{\text{odd}} & \xrightarrow{\mathcal{B}} & \mathcal{F}(\mathbb{C})_{\text{odd}} \\ \mathcal{L}_{\text{odd}} = \mathcal{L}_{3/2} \downarrow & & \downarrow \Phi_{\text{odd}} = \Phi_{3/2} \\ \mathcal{O}(\mathbf{H}) & \xrightarrow{\mathcal{C}_{3/2}} & \mathcal{O}(\mathbf{D}). \end{array}$$

Then the compositions

$$\begin{aligned} \mathcal{C}_{1/2} \circ \mathcal{L}_{1/2} &= \Phi_{1/2} \circ \mathcal{B}: L^2(\mathbb{R})_{\text{even}} \longrightarrow \mathcal{H}_{1/2}(\mathbf{D}), \\ \mathcal{C}_{3/2} \circ \mathcal{L}_{3/2} &= \Phi_{3/2} \circ \mathcal{B}: L^2(\mathbb{R})_{\text{odd}} \longrightarrow \mathcal{H}_{3/2}(\mathbf{D}) \end{aligned}$$

are the unitary isomorphisms, so that

$$L^2(\mathbb{R}) = L^2(\mathbb{R})_{\text{even}} \oplus L^2(\mathbb{R})_{\text{odd}} \simeq \mathcal{H}_{1/2}(\mathbf{D}) \oplus \mathcal{H}_{3/2}(\mathbf{D}).$$

These isomorphisms satisfy, for $\nu = 1/2, 3/2$,

$$\begin{aligned} \mathcal{C}_\nu \circ \mathcal{L}_\nu \circ \left(a^\dagger a + \frac{1}{2} \right) &= \left(2z \frac{d}{dz} + \nu \right) \circ \mathcal{C}_\nu \circ \mathcal{L}_\nu, \\ \mathcal{C}_\nu \circ \mathcal{L}_\nu \circ \frac{1}{2} a^2 &= \frac{d}{dz} \circ \mathcal{C}_\nu \circ \mathcal{L}_\nu, \quad \mathcal{C}_\nu \circ \mathcal{L}_\nu \circ \frac{1}{2} (a^\dagger)^2 = \left(z^2 \frac{d}{dz} + \nu z \right) \circ \mathcal{C}_\nu \circ \mathcal{L}_\nu, \end{aligned}$$

and especially, for $f \in L^2(\mathbb{R})_{\text{even}}$ ($\nu = 1/2$) or $f \in L^2(\mathbb{R})_{\text{odd}}$ ($\nu = 3/2$),

$$(\mathcal{C}_\nu \mathcal{L}_\nu e^{i\theta H} f)(z) = e^{i\theta \nu} (\mathcal{C}_\nu \mathcal{L}_\nu f)(e^{2i\theta} z).$$

The orthonormal basis $\left\{ \frac{\pi^{-1/4}}{\sqrt{(2m)!}} (a^\dagger)^{2m} e^{-x^2/2} \right\} \subset L^2(\mathbb{R})_{\text{even}}$ is mapped to $\left\{ \sqrt{\frac{(1/2)_m}{m!}} z^m \right\} \subset \mathcal{H}_{1/2}(\mathbf{D})$ by $\mathcal{C}_{1/2} \circ \mathcal{L}_{1/2}$, and $\left\{ \frac{\pi^{-1/4}}{\sqrt{(2m+1)!}} (a^\dagger)^{2m+1} e^{-x^2/2} \right\} \subset L^2(\mathbb{R})_{\text{odd}}$ is mapped to $\left\{ \sqrt{\frac{(3/2)_m}{m!}} z^m \right\} \subset \mathcal{H}_{3/2}(\mathbf{D})$ by $\mathcal{C}_{3/2} \circ \mathcal{L}_{3/2}$ (see e.g. [4, Section 4.6]).

In summary, the correspondence of each operator is given as follows.

$$\begin{array}{c} L^2(\mathbb{R}) \\ \mathcal{F}(\mathbb{C}) \\ \mathcal{H}_\nu(\mathbf{D}) \ (\nu = \frac{1}{2}, \frac{3}{2}) \end{array} \quad \left\| \begin{array}{c} H = a^\dagger a + \frac{1}{2} \\ w \frac{d}{dw} + \frac{1}{2} \\ 2z \frac{d}{dz} + \nu \end{array} \right\| \left\| \begin{array}{c} \frac{1}{2} a^2 \\ \frac{1}{2} \frac{d^2}{dw^2} \\ \frac{d}{dz} \end{array} \right\| \left\| \begin{array}{c} \frac{1}{2} (a^\dagger)^2 \\ \frac{1}{2} w^2 \\ z^2 \frac{d}{dz} + \nu z \end{array} \right\| \begin{array}{c} e^{i\theta H} f(x) \\ e^{i\theta/2} f(e^{i\theta} w) \\ e^{i\theta \nu} F(e^{2i\theta} z) \end{array}$$

3 Equivalence between NCHO and two-photon QRM

For $\alpha, \beta \in \mathbb{R}$, $\eta\sqrt{\alpha\beta-1} \in \mathbb{R}$, in [19] the η -shifted non-commutative harmonic oscillator $H_{\text{NCHO}}^{(\alpha, \beta, \eta)}$ is defined as the Hamiltonian operator on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$,

$$H_{\text{NCHO}}^{(\alpha, \beta, \eta)} := \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} \left(-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 \right) + \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \left(x \frac{d}{dx} + \frac{1}{2} \right) + 2\eta\sqrt{\alpha\beta-1}i \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

$\eta = 0$ case is the original non-commutative harmonic oscillator introduced in [16, 17, 18].

Next, for $g, \Delta, \varepsilon \in \mathbb{R}$, let $\tilde{H}_{2\text{QRM}}^{(g, \Delta, \varepsilon)}$ be the operator on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$ given by

$$\tilde{H}_{2\text{QRM}}^{(g, \Delta, \varepsilon)} := \mathbf{I} \left(a^\dagger a + \frac{1}{2} \right) + g\sigma_1 (a^2 + (a^\dagger)^2) + \Delta\sigma_3 + \varepsilon\sigma_1,$$

where $\mathbf{I} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, and $\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $\sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$, $\sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ are the Pauli matrices. Then $H_{2\text{QRM}}^{(g, \Delta, \varepsilon)} := \tilde{H}_{2\text{QRM}}^{(g, \Delta, \varepsilon)} - \frac{1}{2}\mathbf{I}$ is a special case of the spin-boson model (e.g. [20]), and especially if $\varepsilon = 0$, then this is the two-photon quantum Rabi model (e.g. [3, 9]). In addition, for $\nu > 0$, let $\tilde{H}_{(\nu)}^{(g, \Delta, \varepsilon)}$ be the formally self-adjoint operator on $\mathcal{H}_\nu(\mathbf{D}) \otimes \mathbb{C}^2$ given by

$$\tilde{H}_{(\nu)}^{(g, \Delta, \varepsilon)} := \mathbf{I} \left(2z \frac{d}{dz} + \nu \right) + 2g\sigma_1 \left((1+z^2) \frac{d}{dz} + \nu z \right) + \Delta\sigma_3 + \varepsilon\sigma_1.$$

Then these are related as follows.

Theorem. Let $\alpha, \beta \in \mathbb{R}_{>0}$, $\eta\sqrt{\alpha\beta-1}, \lambda \in \mathbb{R}$, and let

$$\begin{aligned} \frac{1}{2\sqrt{\alpha\beta}} &=: g, & -\frac{2\eta\sqrt{\alpha\beta-1}}{\sqrt{\alpha\beta}} &=: \varepsilon, \\ \frac{\lambda}{2}(\alpha^{-1} + \beta^{-1}) &=: \mu, & -\frac{\lambda}{2}(\alpha^{-1} - \beta^{-1}) &=: \Delta, & \begin{bmatrix} \sqrt{\alpha}e^{-\pi i/4} & 0 \\ 0 & \sqrt{\beta}e^{\pi i/4} \end{bmatrix} &=: l. \end{aligned}$$

(1) Let $f(x) \in L^2(\mathbb{R}) \otimes \mathbb{C}^2$. Then the following (a) and (b) are equivalent.

- (a) $H_{\text{NCHO}}^{(\alpha, \beta, \eta)} f = \lambda f$.
- (b) $\tilde{H}_{2\text{QRM}}^{(g, \Delta, \varepsilon)} e^{\frac{\pi i}{4} H} l f = \mu e^{\frac{\pi i}{4} H} l f$.

(2) Suppose $f(x) \in L^2(\mathbb{R})_{\text{even}} \otimes \mathbb{C}^2$, and let

$$F(z) := e^{\frac{\pi i}{8}} (\mathcal{C}_{1/2} \mathcal{L}_{\text{even}} l f)(iz) = (\mathcal{C}_{1/2} \mathcal{L}_{\text{even}} e^{\frac{\pi i}{4} \mathbf{H}} l f)(z) \in \mathcal{H}_{1/2}(\mathbf{D}).$$

Then (a), (b) are equivalent to

$$(c) \quad \widetilde{\mathbf{H}}_{(1/2)}^{(g, \Delta, \varepsilon)} F = \mu F.$$

(3) Suppose $f(x) \in L^2(\mathbb{R})_{\text{odd}} \otimes \mathbb{C}^2$, and let

$$F(z) := e^{\frac{3\pi i}{8}} (\mathcal{C}_{3/2} \mathcal{L}_{\text{odd}} l f)(iz) = (\mathcal{C}_{3/2} \mathcal{L}_{\text{odd}} e^{\frac{\pi i}{4} \mathbf{H}} l f)(z) \in \mathcal{H}_{3/2}(\mathbf{D}).$$

Then (a), (b) are equivalent to

$$(d) \quad \widetilde{\mathbf{H}}_{(3/2)}^{(g, \Delta, \varepsilon)} F = \mu F.$$

Indeed, the formula (a) is rewritten as

$$\left[\begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} \left(\mathbf{a}^\dagger \mathbf{a} + \frac{1}{2} \right) + \frac{1}{2} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} (a^2 - (a^\dagger)^2) + 2\eta\sqrt{\alpha\beta - 1}i \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right] f = \lambda f.$$

Then by the definition of l and $g, \varepsilon, \mu, \Delta$, we have $l^\dagger l = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}$ and

$$\begin{aligned} \frac{1}{2} l^{\dagger-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} l^{-1} &= \frac{i}{2\sqrt{\alpha\beta}} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = g i \boldsymbol{\sigma}_1, \\ 2\eta\sqrt{\alpha\beta - 1} i l^{\dagger-1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} l^{-1} &= -\frac{2\eta\sqrt{\alpha\beta - 1}}{\sqrt{\alpha\beta}} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \varepsilon \boldsymbol{\sigma}_1, \\ \lambda l^{\dagger-1} l^{-1} &= \begin{bmatrix} \lambda\alpha^{-1} & 0 \\ 0 & \lambda\beta^{-1} \end{bmatrix} = \begin{bmatrix} \mu - \Delta & 0 \\ 0 & \mu + \Delta \end{bmatrix} = \mu \mathbf{I} - \Delta \boldsymbol{\sigma}_3. \end{aligned}$$

Hence the above equation is equivalent to

$$\left[\mathbf{I} \left(\mathbf{a}^\dagger \mathbf{a} + \frac{1}{2} \right) + g i \boldsymbol{\sigma}_1 (a^2 - (a^\dagger)^2) + \Delta \boldsymbol{\sigma}_3 + \varepsilon \boldsymbol{\sigma}_1 \right] l f = \mu l f. \quad (*)$$

Then by

$$\mathbf{a} e^{i\theta \mathbf{H}} = e^{i\theta} e^{i\theta \mathbf{H}} \mathbf{a}, \quad \mathbf{a}^\dagger e^{i\theta \mathbf{H}} = e^{-i\theta} e^{i\theta \mathbf{H}} \mathbf{a}^\dagger,$$

this is equivalent to

$$\left[\mathbf{I} \left(\mathbf{a}^\dagger \mathbf{a} + \frac{1}{2} \right) + g \boldsymbol{\sigma}_1 (a^2 + (a^\dagger)^2) + \Delta \boldsymbol{\sigma}_3 + \varepsilon \boldsymbol{\sigma}_1 \right] e^{\frac{\pi i}{4} \mathbf{H}} l f = \mu e^{\frac{\pi i}{4} \mathbf{H}} l f,$$

and we get (1). Next, when $f \in L^2(\mathbb{R})_{\text{even}}$ ($\nu = 1/2$) or $f \in L^2(\mathbb{R})_{\text{odd}}$ ($\nu = 3/2$), the formula (*) is transformed by $\mathcal{C}_\nu \circ \mathcal{L}_\nu$ to

$$\left[\mathbf{I} \left(2z \frac{d}{dz} + \nu \right) + 2g i \boldsymbol{\sigma}_1 \left(\frac{d}{dz} - z^2 \frac{d}{dz} - \nu z \right) + \Delta \boldsymbol{\sigma}_3 + \varepsilon \boldsymbol{\sigma}_1 \right] (\mathcal{C}_\nu \mathcal{L}_\nu l f)(z) = \mu l (\mathcal{C}_\nu \mathcal{L}_\nu f)(z),$$

and this is equivalent to

$$\left[\mathbf{I} \left(2z \frac{d}{dz} + \nu \right) + 2g \boldsymbol{\sigma}_1 \left((1 + z^2) \frac{d}{dz} + \nu z \right) + \Delta \boldsymbol{\sigma}_3 + \varepsilon \boldsymbol{\sigma}_1 \right] (\mathcal{C}_\nu \mathcal{L}_\nu l f)(iz) = \mu l (\mathcal{C}_\nu \mathcal{L}_\nu f)(iz).$$

Since we have $(\mathcal{C}_\nu \mathcal{L}_\nu e^{\frac{\pi i}{4} \mathbf{H}} l f)(z) = e^{\frac{\pi i}{4} \nu} (\mathcal{C}_\nu \mathcal{L}_\nu l f)(e^{\frac{\pi i}{2}} z) = e^{\frac{\pi i}{4} \nu} (\mathcal{C}_\nu \mathcal{L}_\nu l f)(iz)$, we get (2), (3).

Remark. By the definitions of μ, Δ , a spectrum μ of $\widetilde{\mathbf{H}}_{2\text{QRM}}^{(g, \Delta, \varepsilon)}$ concerns that of $\mathbf{H}_{\text{NCHO}}^{(\alpha, \beta, \eta)}$ only when $|\mu| > |\Delta|$. Especially, the non-existence of discrete spectra for $\mathbf{H}_{\text{NCHO}}^{(\alpha, \beta, 0)}$ with $\alpha\beta = 1$ [14, 15] and the existence of discrete spectra for $\widetilde{\mathbf{H}}_{2\text{QRM}}^{(g, \Delta, 0)}$ with $g = \frac{1}{2}$ [3, 9] do not contradict.

4 Covering of one-photon QRM

In this section, we consider the relation of $\tilde{H}_{(\nu)}^{(g,\Delta,\varepsilon)}$ on $\mathcal{H}_\nu(\mathbf{D}) \otimes \mathbb{C}^2$ ($\nu > 0$) and the one-photon asymmetric quantum Rabi model [1, 2, 22]

$$H_{1\text{QRM}}^{(g,\Delta,\varepsilon)} := \mathbf{I}a^\dagger a + g\sigma_1(a + a^\dagger) + \Delta\sigma_3 + \varepsilon\sigma_1$$

on $L^2(\mathbb{R}) \otimes \mathbb{C}^2$, according to [10, Remark 4.2]. First, the equation $\tilde{H}_{(\nu)}^{(g,\Delta,\varepsilon)}F = \mu F$,

$$\left[\mathbf{I} \left(2z \frac{d}{dz} + \nu \right) + 2g\sigma_1 \left((1+z^2) \frac{d}{dz} + \nu z \right) + \Delta\sigma_3 + \varepsilon\sigma_1 \right] F = \mu F,$$

that is,

$$2 \begin{bmatrix} z & g(1+z^2) \\ g(1+z^2) & z \end{bmatrix} \frac{dF}{dz} = \begin{bmatrix} \mu - \nu - \Delta & -\varepsilon - 2g\nu z \\ -\varepsilon - 2g\nu z & \mu - \nu + \Delta \end{bmatrix} F$$

is a Fuchsian differential equation with five regular singularities at $\left\{ \frac{\pm 1 \pm \sqrt{1-4g^2}}{2g}, \infty \right\}$ for $g \neq 0, \pm \frac{1}{2}$. We put

$$g' := \sqrt{\nu}g, \quad \Delta' := \frac{\Delta}{2}, \quad \varepsilon' := \frac{\varepsilon}{2}, \quad \mu' := \frac{\mu - \nu}{2}, \quad w := \sqrt{\nu}z.$$

Then the above becomes an equation

$$\left[\mathbf{I}w \frac{d}{dw} + g'\sigma_1 \left(\left(1 + \frac{w^2}{\nu} \right) \frac{d}{dw} + w \right) + \Delta'\sigma_3 + \varepsilon'\sigma_1 \right] F = \mu' F$$

on

$$\left\{ F(w) \in \mathcal{O}(\sqrt{\nu}\mathbf{D}) \mid \frac{\nu-1}{\nu\pi} \int_{\sqrt{\nu}\mathbf{D}} |F(w)|^2 \left(1 - \frac{|w|^2}{\nu} \right)^{\nu-2} dw < \infty \right\} \otimes \mathbb{C}^2.$$

Now we take the limit $\nu \rightarrow \infty$. Then this becomes an equation

$$\left[\mathbf{I}w \frac{d}{dw} + g'\sigma_1 \left(\frac{d}{dw} + w \right) + \Delta'\sigma_3 + \varepsilon'\sigma_1 \right] F = \mu' F$$

on

$$\mathcal{F}(\mathbb{C}) \otimes \mathbb{C}^2 = \left\{ F(w) \in \mathcal{O}(\mathbb{C}) \mid \frac{1}{\pi} \int_{\mathbb{C}} |F(w)|^2 e^{-|w|^2} dw < \infty \right\} \otimes \mathbb{C}^2,$$

or equivalently,

$$\begin{bmatrix} w & g' \\ g' & w \end{bmatrix} \frac{dF}{dw} = \begin{bmatrix} \mu' - \Delta' & -\varepsilon' - g'w \\ -\varepsilon' - g'w & \mu' + \Delta' \end{bmatrix} F.$$

This has two regular singularities at $\pm g'$ and one irregular singularity at ∞ , and hence this limit gives a confluence process of the Fuchsian equation. Also, this equation is equivalent to $H_{1\text{QRM}}^{(g',\Delta',\varepsilon')}f = \mu'f$ via the Bargmann transform $\mathcal{B}: L^2(\mathbb{R}) \rightarrow \mathcal{F}(\mathbb{C})$, and hence $\tilde{H}_{(\nu)}^{(g,\Delta,\varepsilon)}$ is regarded as a covering model of the one-photon quantum Rabi model.

Acknowledgements The author appreciates Professor M. Wakayama for a lot of helpful comments on this topic.

References

- [1] D. Braak, *Integrability of the Rabi Model*. Phys. Rev. Lett. **107** (2011), 100401.
- [2] D. Braak, *A generalized G-function for the quantum Rabi model*. Ann. Phys. **525** (2013), no. 3, L23–L28.
- [3] L. Duan, Y.F. Xie, D. Braak and Q.H. Chen, *Two-photon Rabi model: analytic solutions and spectral collapse*. J. Phys. A **49** (2016), no. 46, 464002, 13 pp.
- [4] G.B. Folland, *Harmonic analysis in phase space*. Ann. of Math. Stud., 122, Princeton University Press, Princeton, NJ, 1989. x+277 pp.
- [5] T. Ichinose and M. Wakayama, *Zeta functions for the spectrum of the non-commutative harmonic oscillators*. Comm. Math. Phys. **258** (2005), no. 3, 697–739.
- [6] E.T. Jaynes and F.W. Cummings, *Comparison of quantum and semiclassical radiation theories with application to the beam maser*. Proc. IEEE **51** (1963), 89–109.
- [7] K. Kimoto, C. Reyes-Bustos and M. Wakayama, *Determinant expressions of constraint polynomials and the spectrum of the asymmetric quantum Rabi model*. Int. Math. Res. Not. IMRN (2021), no. 12, 9458–9544.
- [8] K. Kimoto and M. Wakayama, *Apery-like numbers for non-commutative harmonic oscillators and automorphic integrals*. Ann. Inst. Henri Poincaré D **10** (2023), no. 2, 205–275.
- [9] C.F. Lo, *Demystifying the spectral collapse in two-photon Rabi model*. Scientific Reports **10**, Article number: 14792 (2020), 5 pp.
- [10] R. Nakahama, *Representation theory of $\mathfrak{sl}(2, \mathbb{R}) \simeq \mathfrak{su}(1, 1)$ and a generalization of non-commutative harmonic oscillators*. in Mathematical Foundation for Post-Quantum Cryptography, (edited by T. Takagi, et al.) “Mathematics for Industry” Springer, 2024 (To appear). arXiv:2310.17118.
- [11] H. Ochiai, *Non-commutative harmonic oscillators and Fuchsian ordinary differential operators*. Comm. Math. Phys. **217** (2001), no. 2, 357–373.
- [12] H. Ochiai, *Non-commutative harmonic oscillators and the connection problem for the Heun differential equation*. Lett. Math. Phys. **70** (2004), no. 2, 133–139.
- [13] H. Ochiai, *A special value of the spectral zeta function of the non-commutative harmonic oscillators*. Ramanujan J. **15** (2008), no. 1, 31–36.
- [14] A. Parmeggiani, *Non-commutative harmonic oscillators and related problems*. Milan J. Math. **82** (2014), 343–387.
- [15] A. Parmeggiani and A. Venni, *On the essential spectrum of certain non-commutative oscillators*. J. Math. Phys. **54** (2013), no. 12, 121507, 10 pp.
- [16] A. Parmeggiani and M. Wakayama, *Oscillator representations and systems of ordinary differential equations*. Proc. Natl. Acad. Sci. USA **98** (2001), 26–30.
- [17] A. Parmeggiani and M. Wakayama, *Non-commutative harmonic oscillators. I*. Forum Math. **14** (2002), no. 4, 539–604.

- [18] A. Parmeggiani and M. Wakayama, *Non-commutative harmonic oscillators. II*. Forum Math. **14** (2002), no. 5, 669–690.
- [19] C. Reyes-Bustos and M. Wakayama, *Covering models of the asymmetric quantum Rabi model: η -shifted non-commutative harmonic oscillators*. Comm. Math. Phys. **403** (2023), no. 3, 1429–1476.
- [20] N.H. Tong and Y.H. Hou, *Scaling analysis in the numerical renormalization group study of the sub-Ohmic spin-boson model*. Phys. Rev. B **85** (2012), 144425.
- [21] M. Wakayama, *Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun differential equations, eigenstates degeneration, and the Rabi model*. Int. Math. Res. Not. IMRN (2016), no. 3, 759–794.
- [22] Q. Xie, H. Zhong, M.T. Batchelor and C. Lee, *The quantum Rabi model: solution and dynamics*. J. Phys. A **50** (2017), no. 11, 113001, 40 pp.