

***E*-DISJUNCTIVE INVERSE SEMIGROUPS**

LUNA ELLIOTT, ALEX LEVINE AND JAMES MITCHELL

ABSTRACT. In this paper we provide an overview of the class of inverse semigroups S such that every congruence on S relates at least one idempotent to a non-idempotent; such inverse semigroups are called *E-disjunctive*. This overview includes the study of the inverse semigroup theoretic structure of *E*-disjunctive semigroups; a large number of natural examples; some asymptotic results establishing the rarity of such inverse semigroups; and a general structure theorem for all inverse semigroups where the building blocks are *E*-disjunctive.

Part 1. In the beginning

1. INTRODUCTION

In this paper we are concerned with a natural type of inverse semigroups. Recall that a *semigroup* is just a set with an associative binary operation, and an *inverse semigroup* is a semigroup S where for every $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. Inverse semigroups have been extensively studied in the literature since their inception. Roughly speaking, the class of inverse semigroups lies somewhere between the classes of semigroups and groups, having more structure, in general, than semigroups, and somewhat less structure than groups. If S is a semigroup, then $\rho \subseteq S \times S$ is a *congruence* if whenever $(x, y) \in \rho$ and $s \in S$, it follows that $(xs, ys), (sx, sy) \in \rho$ also. Congruences are to semigroups what normal subgroups are to groups. In this paper we are interested in the class of inverse semigroups S such that every congruence on S relates at least one idempotent to a non-idempotent element of S . Such inverse semigroups are called *E-disjunctive*; see [34, III.4] for further information.

In this paper we study the inverse semigroup theoretic structure of *E*-disjunctive semigroups; give a large number of natural examples; give some asymptotic results establishing the rarity of such inverse semigroups; and prove a general structure theorem for all inverse semigroups which maybe built from *E*-disjunctive inverse semigroups.

A congruence is called *idempotent-pure* if it never relates an idempotent to a non-idempotent. Idempotent-pure congruences have received significant attention since the study of inverse semigroups commenced; see, for example, [1, 3, 17, 26, 32, 33, 35]. Introduced by Green [13], they preserve much of the important structure of inverse semigroups and have multiple equivalent definitions of different flavours. In the kernel-trace representation (see [15, Section 5.3]) of inverse semigroup congruences, idempotent-pure congruences are those with trivial kernel, and thus are entirely determined by their restriction to the idempotents of a given inverse semigroup.

E-disjunctive inverse semigroups are those with no non-trivial idempotent-pure congruences. Every inverse semigroup has an *E*-disjunctive quotient by its syntactic congruence on its idempotents. It is not difficult to show that the symmetric inverse monoid is *E*-disjunctive, and so every inverse semigroup embeds into an *E*-disjunctive inverse semigroup. Slightly more non-trivial is the proof that every inverse semigroup occurs as the homomorphic image of an *E*-disjunctive inverse semigroup (Corollary 4.4), thus showing that this class captures a large variety of inverse semigroups.

2020 *Mathematics Subject Classification*. 20M18.

Key words and phrases. inverse semigroup, *E*-disjunctive, idempotent-pure.

There are a modest number of papers in the literature about E -disjunctive inverse semigroups. The first use of the term that we are aware of is in Petrich [34] from 1984. Shortly afterwards in 1985, Yoshida [40] published a short note on E -disjunctive inverse semigroups, where it is shown that the class of E -disjunctive inverse semigroups is closed under passing to full inverse subsemigroups; and alternative definition of E -disjunctivity was given. Yoshida also noted that an earlier work of Alimpić and Krgović [1] fully classifies when a Clifford inverse semigroup is E -disjunctive through the description of idempotent-pure homomorphisms. Additional classifications of E -disjunctivity were provided by Li and Zhang [20]. Petrich and Reilly [36], and Gigon [12] have also studied E -disjunctivity in the non-inverse case.

This paper has three parts. In the first we cover the basic properties of E -disjunctive inverse semigroups, and their interactions with the standard notions related to inverse semigroups (Section 2). These standard notions include: the natural partial order; adjoining identities and zeros; and basic closure properties such as direct products (Section 3). In Section 4 we describe some circumstances under which wreath products are E -disjunctive (Proposition 4.2, Theorem 4.3).

In the second part, we consider a compendium of examples of naturally occurring E -disjunctive semigroups. These include the symmetric inverse monoids I_X on any set X with at least 2 elements (see [15, Section 5.1] or the start of Section 5 for the definition, and Example 5.1 for the proof of E -disjunctivity); the dual symmetric inverse monoids (Section 5 and Example 5.2); some minimal examples of E -disjunctive semigroups with certain properties (Example 5.3); an infinite finitely generated Thompson's group-like E -disjunctive inverse monoid ([6] and Example 5.5); a proof that the arithmetic inverse monoid from [14] is E -disjunctive in Example 5.7. Graph inverse semigroups arise naturally from the study of Leavitt path algebras. Such semigroups have been studied extensively in the literature in recent years, see for example [2, 16, 21, 22, 28–30, 39]. In Section 6, we characterise the idempotent-pure congruences on graph inverse semigroups in Theorem 6.1, and characterise graph inverse semigroups that are E -disjunctive in terms of the underlying graphs in Theorem 6.2. In the final section of this part of the paper, we characterise the finite monogenic E -disjunctive inverse semigroups (Section 11.1); and use this to show that the number of monogenic E -disjunctive inverse semigroups as a proportion of all monogenic inverse semigroups of order n is asymptotically 0 in (Corollary 7.7).

In the third, and final, part of the paper we consider various structural properties of E -disjunctive semigroups. In Section 8, we show that there are fairly restrictive bounds on the number of idempotents and non-idempotent elements in finite E -disjunctive semigroups (Theorem 8.1). We explore the extent to which information about an arbitrary inverse semigroup can be recovered from its maximal E -disjunctive image in Section 9. In Section 11, we re-prove a theorem from [32]¹ which provides a means of constructing any inverse semigroup from an E -disjunctive semigroup acting on a partially ordered set (Theorem 8.1). This theorem implies similar in McAlister's famous P -theorem from [27] which characterises the E -unitary inverse semigroups via groups acting on partially ordered sets.

2. BASIC PROPERTIES

In this section we give some of the basic properties of E -disjunctive inverse semigroups. We will be using $E(S)$ to denote the set of idempotents in an inverse semigroup S .

We also show how to construct new examples from old: via ideals (Lemma 2.8); full subsemigroups (Lemma 2.9); direct products (Proposition 2.10); adjoining a zero or identity (Corollary 3.3 and Corollary 3.6); zero direct unions (Proposition 3.5); and wreath products (Theorem 4.3).

A congruence ρ on a semigroup S is called *idempotent-pure* if $(s, e) \in \rho$ and $e \in E(S)$ implies that $s \in E(S)$.

¹The authors of the present paper only discovered [32] at a late stage of the preparation of this paper, and proved the characterisation independently. The theorem and its proof is included for the sake of completeness.

| n | inverse semigroups [38] | E -unitary (non-semilattice) | E -disjunctive inverse semigroups | E -disjunctive inverse monoids |
|-----|-------------------------|--------------------------------|-------------------------------------|----------------------------------|
| 0 | 1 | 0 | 1 | 0 |
| 1 | 1 | 0 | 1 | 1 |
| 2 | 2 | 1 | 1 | 1 |
| 3 | 5 | 2 | 2 | 2 |
| 4 | 16 | 6 | 4 | 4 |
| 5 | 52 | 12 | 8 | 6 |
| 6 | 208 | 39 | 18 | 15 |
| 7 | 911 | 120 | 40 | 28 |
| 8 | 4,637 | 483 | 101 | 68 |
| 9 | 26,422 | 2,153 | 276 | 165 |
| 10 | 169,163 | 11,325 | 761 | 414 |
| 11 | 1,198,651 | 67,570 | 2,422 | 1,202 |
| 12 | 9,324,047 | 453,698 | 7,630 | 3,458 |

TABLE 1. Numbers of isomorphisms types of inverse semigroups of order n with certain properties; computed using the GAP package Semigroups [31], and [7, 23–25].

Definition 2.1 (*E -disjunctive*). An inverse semigroup S is called *E -disjunctive* if the only idempotent-pure congruence on S is the trivial congruence Δ_S .

The numbers of E -disjunctive inverse semigroups of size n for some small values of n are shown in Table 1.

Example 2.2. Every group is an E -disjunctive inverse semigroup and symmetric inverse monoids in 3 or more points are E -disjunctive; see Section 5. The free inverse monoids and the bicyclic monoid defined by the presentation $\langle b, c \mid bc = 1 \rangle$ are not E -disjunctive; see Section 6 for more details.

A useful tool when studying E -disjunctive inverse semigroups is the syntactic congruence with respect to the set of idempotents. This is the maximum idempotent-pure congruence on any inverse semigroup, and so will be trivial if and only if the semigroup is E -disjunctive.

Definition 2.3 (**Syntactic congruence**). Let S be an inverse semigroup. The *syntactic congruence* (with respect to $E(S)$) ρ on S is defined by $(s, t) \in \rho$ if and only if

$$\alpha s \beta \in E(S) \quad \text{if and only if} \quad \alpha t \beta \in E(S),$$

for all $\alpha, \beta \in S^1$ where S^1 is the monoid obtained by adjoining an identity to S .

Since the syntactic congruence with respect to $E(S)$ is the only syntactic congruence we will be using, we will use the term “syntactic congruence” to mean this exclusively.

The following lemma is well-known, we include a proof for completeness.

Lemma 2.4. *If S is an inverse semigroup, then the syntactic congruence ρ of S is the largest idempotent-pure congruence on S with respect to containment.*

Proof. We first check that ρ is an idempotent-pure congruence. It is immediate from the definition that ρ is both a right and left congruence, hence ρ is a congruence. Suppose that $e \in E(S)$ and $(e, s) \in \rho$. Then $1e1 \in E(S)$, so by the definition of ρ , $s = 1s1 \in E(S)$.

Let τ be an idempotent-pure congruence on S , and suppose that $(s, t) \in \tau$. Let $\alpha, \beta \in S^1$. Then

$$(\alpha s \beta, \alpha t \beta) \in \tau$$

so as τ is idempotent-pure, $\alpha s \beta \in E(S)$ if and only if $\alpha t \beta \in E(S)$. Hence $(s, t) \in \rho$. □

The next lemma relates E -disjunctivity and the syntactic congruence.

Lemma 2.5 (cf. Remark III.4.15(δ)). *Let S be an inverse semigroup. Then S is E -disjunctive if and only if the syntactic congruence is equality.*

The next result establishes that every inverse semigroup has an E -disjunctive quotient.

Lemma 2.6. *If S is any inverse semigroup, then the quotient of S by the syntactic congruence (which is idempotent-pure) is E -disjunctive.*

The following lemma provides an alternative means to showing inverse semigroups are E -disjunctive to computing the syntactic congruence.

Lemma 2.7. *Let S be an inverse semigroup. If every idempotent-pure congruence ρ is trivial on $E(S)$; that is, for all $e \in E(S)$ the congruence class of e is $\{e\}$, then S is E -disjunctive.*

Proof. Let ρ be an idempotent-pure congruence on S . As ρ is idempotent-pure, the kernel of ρ is $E(S)$. In addition, as the congruence is the trivial congruence when restricted to the idempotents, the trace of ρ is $\Delta_{E(S)}$. Thus the kernel-trace method (see for example [15, Theorem 5.3.3]) tells us that ρ must be the trivial congruence. We have thus shown that every idempotent-pure congruence on S is trivial, and so S is E -disjunctive. \square

We now consider various closure properties of the class of E -disjunctive inverse semigroups. This class is not closed under taking inverse subsemigroups. For example, every inverse semigroup is isomorphic to an inverse subsemigroup of some symmetric inverse monoid (by the Vagner-Preston Representation Theorem [15, Theorem 5.1.7]), which is E -disjunctive (see Example 5.1).

The class of E -disjunctive inverse semigroups is closed under passing to ideals.

Lemma 2.8. *Let I be an ideal of an E -disjunctive inverse semigroup S . Then I is E -disjunctive.*

Proof. Suppose that ρ is an idempotent-pure congruence on I . Define the binary relation $\bar{\rho}$ on S by

$$\bar{\rho} = \rho \cup \Delta_S.$$

We will show that $\bar{\rho}$ is an idempotent-pure congruence. It is immediate that $\bar{\rho}$ is an equivalence relation and idempotent-pure. We will show $\bar{\rho}$ is a congruence. Let $s, t, x \in S$ and suppose $s\bar{\rho}t$. Then $s = t$, in which case $sx = tx$ and so $sx\rho tx$, otherwise $s\rho t$, and $s, t \in I$. As I is an ideal, $sx, tx \in I$ and so $sx\rho tx$. Thus $\bar{\rho}$ is an idempotent-pure congruence on S . Since S is E -disjunctive, $\bar{\rho}$ is trivial, and so ρ is trivial, and I is E -disjunctive. \square

In the other direction, E -disjunctivity is not closed under passing to inverse supersemigroups, because adjoining an identity and then another identity will result in a non- E -disjunctive semigroup. However, a semigroup will be E -disjunctive if it has an E -disjunctive full inverse subsemigroup. Recall that an inverse subsemigroup T of an inverse semigroup S is *full* if $E(S) = E(T)$. We include the proof for completeness.

Lemma 2.9 ([40], Lemma 1). *If S is an inverse semigroup with a full E -disjunctive subsemigroup T , then S is E -disjunctive.*

Proof. Any non-trivial idempotent-pure congruence on S identifies two idempotents of S . Thus every congruence on S identifies two elements of T . It follows that any non-trivial idempotent-pure congruence on S , induces a non-trivial idempotent-pure congruence on T , so none can exist. \square

Another construction under which E -disjunctivity is preserved is taking finite direct products.

Proposition 2.10. *Let S_1 and S_2 be non-empty inverse semigroups. Then S_1 and S_2 are E -disjunctive if and only if $S_1 \times S_2$ is E -disjunctive.*

Proof. (\Rightarrow) Note that $E(S_1) \times E(S_2) = E(S_1 \times S_2)$. Let $(e_1, e_2), (f_1, f_2) \in E(S_1 \times S_2)$ be arbitrary.

We show that $\bar{\alpha}(e_1, e_2)\bar{\beta} \in E(S_1) \iff \bar{\alpha}(f_1, f_2)\bar{\beta} \in E(S_1)$ for all $\bar{\alpha}, \bar{\beta} \in (S_1 \times S_2)^1$ implies that $(e_1, e_2) = (f_1, f_2)$. This shows that the trace of the syntactic congruence on $S_1 \times S_2$ is just equality on the idempotents. The kernel of the syntactic congruence is $E(S_1 \times S_2)$, and congruences of inverse semigroups are determined by their kernel and trace (see, for example Theorem 5.3.3 of [15]), this implies that the syntactic congruence is equality (as it has the same kernel and trace).

Suppose that the following statement holds for all $\bar{\alpha}, \bar{\beta} \in (S_1 \times S_2)^1$: $\bar{\alpha}(e_1, e_2)\bar{\beta} \in E(S_1) \iff \bar{\alpha}(f_1, f_2)\bar{\beta} \in E(S_1)$. We show that $e_1 = f_1$, the other coordinate follows by symmetry. Let $\alpha, \beta \in S_1^1$ be arbitrary.

By the previous statement, when $\alpha, \beta \in S_1$

$$(\alpha, e_2 f_2)(e_1, e_2)(\beta, e_2 f_2) \in E(S_1 \times S_2) \iff (\alpha, e_2 f_2)(f_1, f_2)(\beta, e_2 f_2) \in E(S_1 \times S_2).$$

In the case that α or β is the identity 1 the above equivalence still holds as $(\alpha, e_2 f_2)$ and/or $(\beta, e_2 f_2)$ can be replaced with $1 \in (S_1 \times S_2)^1$ and the resulting statements are equivalent. So

$$(\alpha e_1 \beta, e_2 f_2) \in E(S_1 \times S_2) \iff (\alpha f_1 \beta, e_2 f_2) \in E(S_1 \times S_2)$$

and hence $\alpha e_1 \beta \in E(S_1) \iff \alpha f_1 \beta$. Since α and β were arbitrary and S_1 is E-disjunctive, it follows that $e_1 = f_1$.

(\Leftarrow) Suppose that $S_1 \times S_2$ is E-disjunctive. Let ρ be an idempotent-pure congruence on S_1 . We show that ρ is trivial. Let ρ' be the congruence on $S_1 \times S_2$ defined by $(s_1, s_2)\rho'(t_1, t_2)$ if and only if $s_1 \rho t_1$ and $s_2 = t_2$. As ρ' is idempotent-pure, it follows that ρ' is equality. Hence if S_2 is not empty, it follows that ρ is also equality. \square

Finally, we will mention the following result in Section 11.

Proposition 2.11 ([19, Proposition 2.4.5]). *A congruence ρ on an inverse semigroup S is idempotent-pure if and only if ρ is contained in the compatibility relation $\{(a, b) \in S^2 \mid ab^{-1}, a^{-1}b \in E(S)\}$.*

3. THE NATURAL PARTIAL ORDER, IDENTITIES AND ZEROS

In this section we consider the interaction of the notion of E-disjunctivity and the natural partial order on any inverse semigroup, and some applications. Recall that *the natural partial order* \leq on an inverse semigroup S is defined by $s \leq t$ if there exists $e \in E(S)$ such that $s = et$.

We define another partial order \preceq on an E-disjunctive inverse semigroup S so that $s \preceq t$ if

$$\alpha s \beta \in E(S) \iff \alpha t \beta \in E(S)$$

for all $\alpha, \beta \in S^1$.

Proposition 3.1. *The partial order \preceq on an E-disjunctive inverse semigroup is equal to the natural partial order.*

Proof. Let S be an E-disjunctive inverse semigroup, and let $s, t \in S$. Suppose $s \leq t$. Then $s = te$ for some $e \in E(S)$. Let $\alpha, \beta \in S^1$ be such that $\alpha t \beta \in E(S)$. Then $\alpha s \beta = \alpha te \beta \in E(S)$, and $s \preceq t$.

Suppose that $s \preceq t$. Since $t^{-1}t \in E(S)$, $t^{-1}s \in E(S)$, and so $s^{-1}t \in E(S)$. If $\alpha, \beta \in E(S)$ are such that $\alpha s \beta \in E(S)$, then $\alpha s s^{-1}t \beta \in E(S)$, and so $s s^{-1}t \preceq s$. If $\alpha, \beta \in S$ are such that $\alpha s s^{-1}t \beta \in E(S)$, then using the fact that $s \preceq t$, it follows that $\alpha s \beta = (\alpha s s^{-1})s \beta \in E(S)$, and so $s \preceq s s^{-1}t$. It follows that s is related to $s s^{-1}t$ by the syntactic congruence. Since S is E-disjunctive, $s = s s^{-1}t$, and so $s \leq t$. \square

The next lemma provide a characterisation of the identity element of an E-disjunctive inverse semigroup in terms of the natural partial order \leq .

Lemma 3.2. *Let S be an E-disjunctive inverse semigroup, and let $e \in S$ be such that $\alpha e \beta \in E(S)$ if and only if $\alpha \beta \in E(S^1)$ for all $\alpha, \beta \in S^1$. Then e is an identity.*

Proof. Using Proposition 3.1, $e \geq x$ for all $x \in E(S)$. It follows that $xe = x = ex$ for all $x \in E(S)$. If $x \in S \setminus E(S)$, then $xe = xx^{-1}xe = x(x^{-1}xe) = x(x^{-1}x) = x$. Similarly $ex = exx^{-1}x = xx^{-1}x = x$. Thus e is an identity for S . \square

A corollary of the previous lemma characterises when an E -disjunctive inverse semigroup S with identity adjoined S^1 is also E -disjunctive.

Corollary 3.3. *Let S be an E -disjunctive inverse semigroup. Then S^1 is E -disjunctive if and only if S does not contain an identity.*

Proof. (\Rightarrow) We prove the contrapositive; that is that if S contains an identity, then S^1 is not E -disjunctive. Suppose S contains an identity e . Then for all $x, y \in S^1$, $xey \in E(S^1)$ if and only if $xy \in E(S^1)$. Similarly, if 1 is the adjoined identity, $x1y \in E(S^1)$ if and only if $xy \in E(S^1)$ for all $x, y \in S^1$. Thus 1 and e are related by the syntactic congruence in S^1 , and so S^1 is not E -disjunctive.

(\Leftarrow) Suppose that S does not contain an identity. Since S is E -disjunctive, Lemma 3.2 tells us there is no element $e \in S$ such that for all $x, y \in S^1$, we have $xey \in E(S^1)$ if and only if $xy \in E(S)$. However, $x1y \in E(S^1)$ if and only if $xy \in E(S^1)$ for all $x, y \in S^1$. It follows that 1 is not related to any other element of S^1 by the syntactic congruence. As no two elements inside S are related by the syntactic congruence of S^1 , we have that the syntactic congruence of S^1 is equality, and so S^1 is E -disjunctive. \square

The next lemma is an analogue of Lemma 3.2 where “identity” is replaced by “zero”.

Lemma 3.4. *Let S be an E -disjunctive inverse semigroup, and let $e \in S$ be such that $\alpha e \beta \in E(S)$ for all $\alpha, \beta \in S^1$. Then e is a zero.*

Proof. Using Proposition 3.1, $e \leq x$ for all $x \in S$. If $x \in E(S)$, then $xe \leq e$ and $ex \leq e$. As $e \leq xe$ and $e \leq ex$, it follows that $xe = e = ex$. If $x \in S \setminus E(S)$, then $xe, ex \in E(S)$. Thus $xe = xe^2 = e$ and $ex = e^2x = e$, because we have shown that e acts as a zero when multiplied by elements of $E(S)$. Thus e is a (the) zero in S . \square

To obtain the analogue of Corollary 3.3 we prove a more general result.

Proposition 3.5. *A 0-direct union of E -disjunctive inverse semigroups $(S_i)_{i \in I}$ is E -disjunctive if and only if none of the semigroups S_i has a zero.*

Proof. (\Rightarrow) We prove the contrapositive. Suppose that S_i has a zero element 0_i for some $i \in I$. Then the congruence on the zero direct union generated by the pair $(0_i, 0)$ identifies only these two elements and is thus non-trivial and idempotent-pure.

(\Leftarrow) Let σ be an idempotent-pure congruence on the zero direct union. Suppose for a contradiction that there is $(a, b) \in \sigma$ with $a \neq b$. Since σ restricts to idempotent-pure congruences on each S_i , it follows that σ is trivial on each S_i . In particular, a and b do not belong to the same semigroup S_i for any $i \in I$.

Suppose without loss of generality that $(a, b) \in \sigma$, $i \in I$, $a \in S_i$ and $b \notin S_i$. Then $(a, 0) = (aa^{-1}a, ba^{-1}a) \in \sigma$. So a is the unique element of S_i related to 0 by σ . Since σ is a congruence, it follows that the set $\{a\}$ is an ideal of the semigroup S_i . This is a contradiction as S_i does not contain a zero. \square

Corollary 3.6. *Let S be an E -disjunctive inverse semigroup. Then S^0 is E -disjunctive if and only if S does not contain a zero.*

4. WREATH PRODUCTS AND QUOTIENTS

In this section we consider when wreath products of inverse semigroups are E -disjunctive and use this to show that every inverse semigroup is a homomorphic image of an E -disjunctive inverse

semigroup. We think of wreath products in terms of matrices. Recall that an element of a wreath product of groups $G \wr_X H$ where $H \leq S_X$ consists of a pair $((g_x)_{x \in X}, h) \in (G^X, H)$. We think of the elements of $G \wr_X H$ as an $X \times X$ matrix M such that the entry indexed by $(x, y) \in X \times X$ in M is g_x whenever $(x)h = y$ and 0 otherwise. We will also think of such matrices as functions $M : X \times X \rightarrow G \cup \{0\}$ where $M(x, y)$ is just the (x, y) -entry of the matrix. It is routine to verify that the group $G \wr_X H$ is isomorphic to the group consisting of the corresponding matrices, as just defined, where $0 + g = g + 0 = g$ for all $g \in G$.

We extend this definition of wreath products to inverse semigroups S and subsemigroups T of the symmetric inverse monoid I_X as follows. To do this nicely we introduce a semiring which contains S but only use the $S \cup \{0\}$ part of it.

Recall that $(R, +, \cdot)$ is a *semiring* if the following hold:

- (1) $(R, +)$ is a commutative monoid whose identity we call 0;
- (2) (R, \cdot) is a semigroup;
- (3) the operation \cdot distributes over $+$; and
- (4) if $r \cdot 0 = 0 \cdot r = 0$ for all $r \in R$.

If S is a semigroup, then we define $\mathbb{N}[S]$ to be the quotient of the free semiring over the set S by the relations $s \cdot t = st$, $s, t \in S$. That is, $\mathbb{N}[S]$ consists of finite formal sums of the form

$$\sum_{s \in S} n_s s$$

where $n_s \in \mathbb{N}$ for all $s \in S$ and only finitely many n_s are non-zero with the natural multiplication.

If R is a semiring and X is a set, then we define the (*row finite*) *matrix semiring* over R by

$$\begin{aligned} M_X(R) &= \{f : X \times X \rightarrow R \mid \text{all but finitely many entries of each row of } f \text{ are } 0\} \\ &= \{f : X \times X \rightarrow R \mid \text{for all } x \in X, |(R \setminus \{0\})f^{-1} \cap (\{x\} \times X)| < \infty\} \end{aligned}$$

with operation defined by

$$(x, y)fg = \sum_{i \in X} (x, i)f \cdot (i, y)g$$

for $f, g \in M_X(R)$.

If S is a semigroup, P_X is the partial transformation monoid on the set X , and $T \leq P_X$, then we define $S \wr T$ to be the following submonoid of $M_X(\mathbb{N}[S])$:

$$S \wr T = \{f \in M_X(\mathbb{N}[S]) \mid \text{im}(f) \subseteq S \cup \{0\} \text{ and } (S)f^{-1} \in T\}.$$

The condition $(S)f^{-1} \in T$ makes sense since $(S)f^{-1} \subseteq X \times X$, and so this condition simply asserts that the relation $(S)f^{-1}$ is a partial transformation that belongs to T .

We define $\phi : S \wr T \rightarrow T$ by $(f)\phi = (S)f^{-1}$. It is routine to verify that ϕ is a surjective homomorphism. As such the multiplication in $S \wr T$ can alternatively be defined as follows

$$(x, y)fg = \begin{cases} (x, z)f \cdot (z, y)g & \text{if } (x, z) \in (f)\phi, (z, y) \in (g)\phi \\ 0 & \text{if } (x, y) \notin (fg)\phi, \end{cases}$$

(since the sum only ever has one non-zero summand).

Lemma 4.1. *If S and $T \leq I_X \leq P_X$ are inverse semigroups, then $S \wr T$ is an inverse semigroup.*

Proof. If $f \in S \wr T$, then it is routine to verify that f is an idempotent if and only if

- (1) the preimage of S under f is an idempotent of T ;
- (2) the image of f contains only idempotents.

Since S and T are inverse semigroups, $f \in S \wr T$ is a diagonal matrix and so the idempotents of $S \wr T$ commute. Also if $f \in S \wr T$, then we define f^{-1} to be

$$(a, b)f^{-1} = \begin{cases} ((b, a)f)^{-1} & \text{if } (b, a)f \neq 0 \\ 0 & \text{if } (b, a)f = 0 \end{cases}.$$

In other words, f^{-1} is obtained from f by transposing f and inverting its entries. It is straightforward to show that f^{-1} is a semigroup theoretic inverse of f , and so $S \wr T$ is an inverse semigroup. \square

The following proposition is a special case of Theorem 4.3, however we include the proof below, as it is more straightforward, and helps exhibit the ideas behind the proof of Theorem 4.3.

Proposition 4.2. *Let G be a non-trivial group and $T \leq I_X$ be an inverse semigroup. Then $G \wr T$ is E -disjunctive.*

Proof. Seeking a contradiction suppose that ρ is a non-trivial idempotent-pure congruence on $G \wr T$. Then by Lemma 2.7 there exist $f \neq g \in E(G \wr T)$ such that $(f, g) \in \rho$. If $fg = f$, then $f < g$. If $fg \neq f$, then $f < fg$ are idempotents such that $(f, fg) \in \rho$. Thus we may assume without loss of generality that $f < g$. Note that f, g are both matrices whose entries are all 0 except for some idempotents on the diagonal. Since $f < g$, there exists $x \in X$ such that $(x, x)f < (x, x)g$. The entries of f and g belong to $G \cup \{0\}$ (whose only idempotents are the identity 1_G and 0). Thus $(x, x)g = 1_G$ and $(x, x)f = 0$. Let $h \in G \setminus \{1_G\}$ and let $g' \in S \wr T$ be the matrix with the same entries as g except that $(x, x)g' = h$. Thus $g'g = g'$ and $g'f = f$. It follows that

$$(f, g) \in \rho \Rightarrow (g'f, g'g) \in \rho \Rightarrow (f, g') \in \rho.$$

But g' is not an idempotent, and ρ is idempotent-pure. This is a contradiction. \square

The next theorem establishes that every inverse semigroup is a quotient of some E -disjunctive semigroup.

Theorem 4.3. *Let S be an E -disjunctive inverse semigroup without a zero and $T \leq I_X$ be a inverse semigroup. Then $S \wr T$ is E -disjunctive and has T as a quotient.*

Moreover, if $T \leq S_X$, then the assumption that S has no zero can be dropped.

Proof. If T is empty, then $S \wr T$ is empty, and so $S \wr T$ is E -disjunctive, as required. We therefore assume that T is non-empty.

Let D be equal to the set of diagonal matrices of $S \wr T$. Then D is a full inverse subsemigroup of $S \wr T$, i.e. D contains all of the idempotents of $S \wr T$. By Lemma 2.9, it suffices to show that D is E -disjunctive.

Let ρ be an idempotent-pure congruence on D . We show that ρ is trivial. It suffices by Lemma 2.7 to show that ρ is trivial on the idempotents. Suppose that $(f, g) \in \rho$ and f, g are idempotents. Let $h = fg \leq f$. We show that $f = h$, then by symmetry we will have $g = h$ and hence $f = g$ as required.

Let $x \in X$ be arbitrary. We need only show that $(x, x)f = (x, x)h$ (as they are idempotents they agree on their non-diagonal entries). There are two cases to consider.

Case 1: $(x, x)f \geq (x, x)h = 0$. We define

$$I = \{s \in S \mid \text{the matrix obtained from } f \text{ by replacing } (x, x)f \text{ with } s \text{ belongs to } h/\rho \}.$$

For all $s \in S \cup \{0\}$, let f_s be the matrix obtained from f by replacing $(x, x)f$ with s . It follows that

$$f_S := \{f_s \mid s \in S \cup \{0\}\}$$

is a semigroup isomorphic to $S \cup \{0\}$. The restriction of h/ρ to f_S is I . The natural map from S to f_S embeds I into a congruence class of f_S containing the zero element of f_S . Thus I is an ideal of S containing $(x, x)f$. Since ρ is idempotent-pure, it follows that I consists of idempotents. Since S

is E -disjunctive, it follows that I must therefore be a singleton (otherwise S/I would be a proper idempotent-pure quotient). Hence, since I is a singleton ideal, the unique element of I is a zero for S , a contradiction. So in fact Case 1 never occurs.

Showing that Case 1 does not occur is the only point in the proof requiring that S has no zero element. When $T \leq S_X$, this case does not occur because the only idempotent of T is the identity function on X so h has no zeros on the diagonal. Hence why the assumption is no longer needed.

Case 2: $(x, x)f \geq (x, x)h > 0$. For all $s \in S$ let $h_s \in D$ be the element which agrees with h on all entries except $(x, x)h_s = s$. Then $S_x := \{h_s \mid s \in S\}$ is a subsemigroup of D isomorphic to S . Hence, since ρ is idempotent-pure, the restriction of ρ to S_x is trivial. In particular, to show that $(x, x)h = (x, x)f$, we need only show that $(h, h_{(x,x)f}) = (h_{(x,x)h}, h_{(x,x)f}) \in \rho$.

We denote $h_{(x,x)f}$ by f' . Since $h \leq f$, $f'f = f'$ and $f'h = h$. It follows that

$$\begin{aligned} (f, g) \in \rho &\Rightarrow (ff, fg) \in \rho \\ &\Rightarrow (f, h) \in \rho \\ &\Rightarrow (f'f, f'h) \in \rho \\ &\Rightarrow (f', h) \in \rho \end{aligned}$$

as required. □

Corollary 4.4. *Every inverse semigroup is a quotient of an E -disjunctive inverse semigroup.*

Part 2. Some classes of E -disjunctive inverse semigroups

In this part of the paper we provide a number of examples of E -disjunctive inverse semigroups Section 5, and we characterise E -disjunctive inverse semigroups belonging to the classes of: graph inverse semigroups Section 6 (in terms of the underlying graphs); and monogenic inverse semigroups Section 7. As mentioned above, the Clifford E -disjunctive semigroup were characterised in [1, Theorem 6].

5. A COMPENDIUM OF EXAMPLES

In this section we give various examples of E -disjunctive inverse semigroups. These serve as counterexamples to various natural questions about E -disjunctive inverse semigroups.

Recall that if X is any set, then the *symmetric inverse monoid* I_X on X consists of the bijections between subsets of X and the operation is the usual composition of binary relations. By the Vagner-Preston Theorem [15, Theorem 5.1.7], every inverse semigroup is isomorphic to an inverse subsemigroup of some symmetric inverse monoid.

Example 5.1. The symmetric inverse monoid I_X on a set X with $|X| \geq 2$ is E -disjunctive.

Proof. Suppose that $s, t \in I_X$ and (s, t) belongs to the syntactic congruence on I_X . Then $\alpha s \beta \in E(I_X)$ if and only if $\alpha t \beta \in E(I_X)$ for all $\alpha, \beta \in I_X$. If $s \in I_X$, then we consider s as the subset of $X \times X$ consisting of the pairs $(x, (x)s)$. Let $x, y \in X$, and let $z \in X \setminus \{x\}$. Then $\{(x, x)\} \circ s \circ \{(y, z)\} \notin E(I_X)$ if and only if $(x, y) \in s$; and similarly for t . Since $\{(x, x)\} \circ s \circ \{(y, z)\} \in E(I_X)$ if and only if $\{(x, x)\} \cdot t \cdot \{(y, z)\} \in E(I_X)$, and so $(x, y) \in s$ if and only if $(x, y) \in t$. Thus $s = t$, and so the syntactic congruence on I_X is equality. □

If $|X| = 0$, then I_X is the trivial semigroup, and hence is E -disjunctive. If $|X| = 1$, then I_X is a semilattice of size 2, and as such is not E -disjunctive by Example 2.2.

Recall, from [10], for example, that the *dual symmetric inverse monoid* I_X^* is defined as follows. The underlying set of I_X^* is the set of partitions of $X \times \{0, 1\}$ such that each part intersects both $X \times \{0\}$ and $X \times \{1\}$. In other words, elements of I_X^* correspond to bijections between the parts of a partition of $X \times \{0\}$ and a partition of $X \times \{1\}$. We will use partitions and the corresponding

equivalence relations interchangeably. We define $\text{Diag}(s, t)$ to be the least equivalence relation on $X \times \{0, 1, 2\}$ containing

$$\{((x, a), (y, b)) \in (X \times \{0, 1, 2\})^2 : ((x, a), (y, b)) \in s \text{ or } ((x, a - 1), (y, b - 1)) \in t\}.$$

The product of s and t is defined to be

$$\{((x, a), (y, b)) \in (X \times \{0, 1\})^2 : ((x, 2a), (y, 2b)) \in \text{Diag}(s, t)\}$$

and is denoted st . Note that $e \in I_X^*$ is an idempotent whenever $((x, 0), (y, 0)) \in e$ if and only if $((x, 1), (y, 1)) \in e$ for all $x, y \in X$ (i.e. the partitions of $X \times \{0\}$ and $X \times \{1\}$ are the “same” and the corresponding function is the identity).

A congruence ρ is *idempotent-separating* if ρ never relates two distinct idempotents. Thus idempotent-separating is a dual property to idempotent-pure and inverse semigroups with no non-trivial idempotent-separating congruences, called *fundamental* inverse semigroups are a dual class of inverse semigroups to E -disjunctive inverse semigroups. It is not difficult to show that the symmetric inverse monoid is fundamental, thus providing a non-congruence free example of a semigroup in both of these classes.

Example 5.2. The dual symmetric inverse monoid I_X^* , where X is any set, is E -disjunctive.

Proof. If I_X^* has at most one idempotent, then I_X^* is either a group or the empty semigroup. In either case, I_X^* is E -disjunctive. By Lemma 2.7, it suffices to show that every idempotent-pure congruence is trivial on $E(S)$. Let ρ be an idempotent-pure congruence on I_X^* . Suppose for contradiction that there exist distinct idempotents $e, f \in E(I_X^*)$ such that $(e, f) \in \rho$. Since $e \neq f$, at most one of e and f equals ef ; assume without loss of generality that $e \neq ef$. Then there is a part of ef which is a union of at least two parts of e . Let $s \in I_X^*$ be an element which swaps two of these parts of e and fixes the others. If $(e, f) \in \rho$, then $(e, ef) = (e^2, ef) \in \rho$ and so $(se, sef) \in \rho$. But $se = s$, as e acts as the identity function on the image of s , which is not an idempotent. On the other hand, $sef = ef \in E(I_X^*)$, and so $(s, ef) \in \rho$. Since ef is an idempotent and s is not, this contradicts ρ being idempotent-pure, and so I_X^* is E -disjunctive. \square

The next example shows that “congruence” cannot be replaced by “right congruence” in the definition of E -disjunctive inverse semigroups. In fact, the next example is the unique inverse semigroup of smallest size up to isomorphism, showing this.

Example 5.3. Let S be the inverse semigroup defined by the inverse semigroup presentation:

$$\langle x, y \mid xy = xy^{-1} = yx = yx^{-1} = y^{-1}x^{-1} = x^2, yy^{-1} = xx^{-1} \rangle.$$

It can be shown, for example, using GAP [11, 31], that:

$$S = \{x, y, x^{-1}, y^{-1}, x^2, xx^{-1}, x^{-1}x, x^{-1}y, y^{-1}x, y^{-1}y, x^3\},$$

that S is E -disjunctive, the least right congruence ρ on S containing the pair (x^3, x) has non-trivial classes:

$$\{x, y, x^3\}, \{x^2, xx^{-1}\},$$

and that the idempotents of S are:

$$x^{-1}x, y^{-1}y, xx^{-1}, x^2.$$

Hence ρ is idempotent-pure, using the obvious definition of this notion for right congruences. It is also possible to show using [23, 24], based on [25], that there is no smaller E -disjunctive inverse semigroup admitting such a right congruence, that S is the unique inverse semigroup (up to isomorphism) of size 11 admitting such a right congruence, and even that ρ is the only such right congruence on S .

For our next example, we require the definition of Thompson’s group V , which we define below. For a more comprehensive introduction to Thompson’s group V , we refer the reader to [6].




FIGURE 1. Egg-box diagram of an E -disjunctive inverse semigroup with a non-trivial idempotent-pure right congruence.

Definition 5.4 (Thompson’s group V). Let \mathfrak{C} denote the Cantor space, with underlying set $\{0, 1\}^\omega$ (that is, infinite sequences of 0s and 1s), and using the product topology induced from the discrete topology of $\{0, 1\}$. We denote the free monoid on $\{0, 1\}$ by $\{0, 1\}^*$ (i.e. the monoid of finite sequences of 0s and 1s with concatenation as the operation). If $w \in \{0, 1\}^*$, then we define

$$w\mathfrak{C} = \{x \in \mathfrak{C} \mid w \text{ is a prefix of } x\}.$$

Note that these sets are clopen, and the collection of all such sets is a basis for \mathfrak{C} .

Let F_1 and F_2 be finite subsets of $\{0, 1\}^*$ such that $|F_1| = |F_2|$, and

$$\{w\mathfrak{C} \mid w \in F_1\} \quad \text{and} \quad \{w\mathfrak{C} \mid w \in F_2\}$$

are partitions of \mathfrak{C} . We call such subsets of $\{0, 1\}^*$ *complete antichains*. If $u \in \mathfrak{C}$, then since F_1 partitions \mathfrak{C} there exists $w_u \in F_1$ that is a prefix of u . In this case, we write $u = w_u v_u$ where $v_u \in \mathfrak{C}$ is just the suffix of u following w_u . The *prefix exchange map* $f: \mathfrak{C} \rightarrow \mathfrak{C}$ between F_1 and F_2 induced by a bijection $\phi: F_1 \rightarrow F_2$ is defined by

$$(u)f = (w_u \phi)v_u.$$

Every such prefix exchange map is a homeomorphism of \mathfrak{C} . The group of all prefix exchange maps between any pair of complete antichains is called *Thompson’s group V* .

The following example is a slight modification of the Thompson inverse monoid $\text{Inv}_{2,1}$ introduced in [4] (we modify it as that monoid does not have infinitely many \mathcal{J} -classes).

Example 5.5. We give an example of a Thompson’s group-like E -disjunctive inverse monoid that is finitely generated and has infinitely many \mathcal{J} -classes.

Let M be the inverse submonoid of the inverse monoid of partial permutations on \mathfrak{C} generated by Thompson’s group V and the identity functions on $1\mathfrak{C}$ and $\{1^n 0^\omega : n \in \mathbb{N} \cup \{0\}\}$. We denote the second of these identity functions by e . As Thompson’s group V is 2-generated (see for example [5]), M is 4-generated.

We next show that M has infinitely many \mathcal{J} -classes. We do so by showing that M contains the identity function on a set of size n for all $n \in \mathbb{N} \cup \{0\}$. For different n , these elements are not \mathcal{J} -related in $I_{\mathfrak{C}}$ and hence M has infinitely many \mathcal{J} -classes. It suffices to show that every identity f_n on the set

$$\{1^i 0^\omega : i < n\}$$

belongs to M . We denote the identity function on the set $\bigcup_{i < n} 1^i 0^\mathfrak{C}$ by g_n . It is straightforward to verify that $f_n = g_n e$, and so it suffices to show that $g_n \in M$. Note that

$$1\mathfrak{C} = 1^n \mathfrak{C} \cup \bigcup_{1 \leq i < n} 1^i 0^\mathfrak{C}$$

for any $n \geq 1$. If F is any complete antichain in $\{0, 1\}^*$ containing 0 and 1^n , and $\phi: F \rightarrow F$ is the bijection swapping 0 and 1^n . Then the corresponding prefix exchange map $\psi \in V$ maps $1\mathfrak{C}$ to $\text{dom}(g_n)$. Hence the conjugate of the identity function on $1\mathfrak{C}$ by ψ is f_n . In particular, since the identity on $1\mathfrak{C}$ is a generator of M , f_n belongs to M .

We now show that M is E -disjunctive. If $m \in M$ is arbitrary and the domain of m is uncountable, then m is a product of elements of V and the second generator. Since the domain of the second generator is clopen and the elements of V are prefix exchange maps, it follows that the image of m is clopen. Hence every element m of M has a domain which is either clopen or countable.

If J is the \mathcal{J} -class of f_1 , then $J = Vf_1V$ and so J consists of functions with domain of size 1 which map an element with an infinite tail of zeros to another such element.

Claim 5.6. If $a, b \in M \setminus \{\emptyset\}$ are distinct, there is an idempotent $e \in J$ such that $ea, eb \in J \cup \{\emptyset\}$ and $ea \neq eb$.

Proof. Suppose that a and b have the same domain. If $\text{dom}(a) = \text{dom}(b)$ is countable, then there is an element $a' \in J$ with a' less than a in the usual partial order of inverse semigroups. Since $a \neq b$, there exists $u \in \mathfrak{C}$ such that $(u)a \neq (u)b$. In particular, we may choose $e \in J$ such that e is the identity on u . In this case, $(u)ea = (u)a \neq (u)b = (u)eb$ and $ea, eb \in J$.

If $\text{dom}(a) = \text{dom}(b)$ is clopen, then the set $\{x \in \text{dom}(a) \mid (x)a \neq (x)b\}$ is open and non-empty. Thus it contains a set $w\mathfrak{C}$ for some $w \in \{0, 1\}^*$. By prefix replacement via Thompson's group V (using the prefix w) we can find $m' \in J$ such that $\text{im}(m') \subseteq w\mathfrak{C}$. Hence $m'^{-1}m'a, m'^{-1}m'b \in J$ are distinct and $m'^{-1}m'$ is the required idempotent in this case.

If $\text{dom}(a) \neq \text{dom}(b)$, then suppose without loss of generality that there is some $u \in \text{dom}(a) \setminus \text{dom}(b)$ such that u ends with an infinite tail of zeros. If we set e to be the identity function on $\{u\}$, then $e \in J$ since J comprises all functions with a domain of size 1 which map an element with an infinite tail of zeros to another such element. \square

As J contains both idempotents and non-idempotents, it suffices to show that every non-trivial congruence on M identifies the \mathcal{J} -class J with the \mathcal{J} -class of $f_0 = \emptyset$.

Let ρ be a non-trivial congruence on M . Let $a, b \in M$ be such that $a \neq b$ and $(a, b) \in \rho$. By the claim above, there is $e \in J$ such that $ea, eb \in J$ and $ea \neq eb$. Thus $ea(ea)^{-1}$ and $eb(ea)^{-1}$ are related by ρ . But one of these is zero and the other is not, so all elements of J are related to zero by ρ and we are done.

Example 5.7. The *arithmetic inverse monoid* \mathcal{A} , from [14], is the inverse submonoid of the symmetric inverse monoid $I_{(\mathbb{Z}_{\geq 0})}$ generated by the set $\{R_{a,b} \mid a, b \in \mathbb{Z}_{\geq 0}, a > b\}$, where $R_{a,b} \in I_{(\mathbb{Z}_{\geq 0})}$ is defined by

$$(n)R_{a,b} = \begin{cases} \frac{n-b}{a} & n \equiv b \pmod{a} \\ \text{undefined} & \text{otherwise.} \end{cases}$$

By [14, Theorem 12], every non-zero element of \mathcal{A} may be written uniquely in the form $R_{a,b}R_{c,d}^{-1}$, where $c > d$ and $a > b$ (note $(n)R_{c,d}^{-1} = nc + d$). Since idempotents of $I_{(\mathbb{Z}_{\geq 0})}$ are partial identities, it follows that the idempotents of \mathcal{A} are just $R_{a,b}R_{a,b}^{-1}$ or \emptyset (where \emptyset is the empty map), for all $a > b$ or the empty map \emptyset . By [14, Theorem 24], if $(R_{a,b}R_{c,d}^{-1}), (R_{e,f}R_{g,h}^{-1}) \in \mathcal{A} \setminus \{\emptyset\}$, then

$$(1) \quad (R_{a,b}R_{c,d}^{-1}) \cdot (R_{e,f}R_{g,h}^{-1}) = \begin{cases} R_{\frac{ae}{\gcd(c,e)}, \frac{a(r-d)+b}{c}} R_{\frac{ge}{\gcd(c,e)}, \frac{g(r-f)+h}{e}}^{-1} & \text{if } \gcd(c, e) \text{ divides } (d-f) \\ \emptyset & \text{otherwise,} \end{cases}$$

where r is minimal such that $r \equiv d \pmod{c}$ and $r \equiv f \pmod{e}$.

We will show that \mathcal{A} is E -disjunctive by showing that the syntactic congruence is trivial. Recall that the right syntactic congruence of a semigroup S is $\{(s, t) \in S \mid sx \in E(S) \iff tx \in E(S)\}$.

$E(S)$ for all $x \in S$. The right syntactic congruence is the maximum idempotent-pure right congruence on S . Since the syntactic congruence is idempotent-pure and a right congruence, the syntactic congruence is contained in the right syntactic congruence. Since the syntactic congruence ρ is idempotent-pure, the kernel of ρ is $E(\mathcal{A})$, and so ρ is trivial if and only if ρ has a trivial trace. Hence it suffices, since the trace of ρ is contained in the trace of the right syntactic congruence, to show that the trace of the syntactic right congruence is trivial. By definition, the right syntactic congruence has a trivial trace if and only if each idempotent e of \mathcal{A} defines a unique set $Y_e = \{s \in \mathcal{A} \mid es \in E(\mathcal{A})\}$.

Let $R_{a,b}R_{a,b}^{-1} \in E(\mathcal{A}) \setminus \{\emptyset\}$. We will describe the set $Y_{R_{a,b}R_{a,b}^{-1}}$. Suppose that $R_{e,f}R_{g,h}^{-1} \in \mathcal{A} \setminus \{\emptyset\}$ is such that $(R_{a,b}R_{a,b}^{-1}) \cdot (R_{e,f}R_{g,h}^{-1}) \in E(\mathcal{A})$. Then $(R_{a,b}R_{a,b}^{-1}) \cdot (R_{e,f}R_{g,h}^{-1}) = \emptyset$ or $= R_{x,y}R_{x,y}^{-1}$ for some x and y . Hence by (1) precisely one of the following holds:

- (1) $ae = ga$ and $r = \frac{g(r-f)}{e} + h$, where r is minimal such that $r \equiv b \pmod{a}$ and $r \equiv f \pmod{e}$;
- (2) $\gcd(a, e)$ does not divide $b - f$.

If (1) holds, then $ae = ga$ implies $e = g$, and so

$$r = \frac{g(r-f)}{e} + h = r - f + h.$$

Hence $f = h$. So $R_{e,f}R_{g,h}^{-1} = R_{e,f}R_{e,f}^{-1} \in E(\mathcal{A})$. If (2) holds, then immediately from (1) the product $(R_{a,b}R_{a,b}^{-1}) \cdot (R_{e,f}R_{g,h}^{-1}) = \emptyset$. Let

$$X_{a,b} = \{(e, f) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \mid e > f \text{ and } \gcd(a, e) \text{ does not divide } b - f\}.$$

Suppose $R_{a',b'}R_{a',b'}^{-1} \in E(\mathcal{A})$ is such that $X_{a',b'} = X_{a,b}$. Then for all $e > f \in \mathbb{Z}_{\geq 0}$, we have that $\gcd(a, e)$ divides $b - f$ if and only if $\gcd(a', e)$ divides $b' - f$. We will show that $a = a'$ and $b = b'$.

Let p be a prime greater than a and a' (and thus coprime to both). Then $\gcd(a, pa')$ divides $0 = b - b$, and so $a' = \gcd(a', pa')$ divides $b' - b$. By symmetry, a divides $b' - b$. If $f < pa'$, then $\gcd(a, a') = \gcd(a, pa')$ divides $b - (b' + f)$ if and only if $\gcd(a', pa')$ divides f . Since $b - b'$ is a multiple of a' , and hence a multiple of $\gcd(a, a')$ also, $\gcd(a, a')$ divides $b - (b' + f)$ if and only if $\gcd(a, a')$ divides f . Thus for all $f < pa'$, $\gcd(a, a')$ divides f if and only if $a' = \gcd(a', a')$ divides f . In particular, $\gcd(a, a')$ divides a , and so a' divides a . By symmetry, a divides a' , and since $a, a' \in \mathbb{Z}_{>0}$, it follows that $a = a'$. It remains to show that $b = b'$. We have already shown that a divides $b' - b$. Since $b' < a$ and $b < a$, $|b' - b| < a$ and so $|b' - b| = 0$, as required.

It follows that if $R_{a,b}R_{a,b}^{-1} \in E(\mathcal{A}) \setminus \{\emptyset\}$, then

$$Y_{R_{a,b}R_{a,b}^{-1}} \setminus E(\mathcal{A}) = \{R_{e,f}R_{g,h}^{-1} \mid (e, f) \in X_{a,b}\} \setminus E(\mathcal{A}).$$

If $a' \neq a$ or $b' \neq b$, then $X_{a,b} \neq X_{a',b'}$ and so $Y_{R_{a,b}R_{a,b}^{-1}} \neq Y_{R_{a',b'}R_{a',b'}^{-1}}$, as required. Finally, $R_{a,b} \in Y_{\emptyset}$, but $R_{a,b} \notin Y_{R_{a,b}R_{a,b}^{-1}}$, $Y_{\emptyset} \neq Y_{R_{a,b}R_{a,b}^{-1}}$ for any non-zero idempotent $R_{a,b}R_{a,b}^{-1}$.

6. GRAPH INVERSE SEMIGROUPS

In this section we give a full characterisation of when an arbitrary graph inverse semigroup is E -disjunctive.

For this purpose, we define a *graph* $\Gamma = (\Gamma^0, \Gamma^1, \mathbf{s}, \mathbf{r})$ to be a quadruple consisting of two sets, Γ^0 and Γ^1 , and two functions $\mathbf{s}, \mathbf{r} : \Gamma^1 \rightarrow \Gamma^0$, called the *source* and *range*, respectively. The elements of Γ^0 and Γ^1 are called *vertices* and *edges*, respectively. A sequence $p = e_1e_2 \cdots e_k$ of (not necessarily distinct) edges $e_i \in \Gamma^1$, such that $(e_i)\mathbf{r} = (e_{i+1})\mathbf{s}$ for $1 \leq i \leq k-1$, is a *path* from $(e_1)\mathbf{s}$ to $(e_k)\mathbf{r}$. We define $(p)\mathbf{s} = (e_1)\mathbf{s}$ and $(p)\mathbf{r} = (e_k)\mathbf{r}$, and refer to k as the *length* of p . The elements of Γ^0 are paths of length 0, and we denote by $\text{Path}(\Gamma)$ the set of all paths in Γ .

Define the *graph inverse semigroup* $S(\Gamma)$ of a graph Γ to be the inverse semigroup with zero $0 \notin \Gamma^0 \cup \Gamma^1$, generated by Γ^0 and Γ^1 , together with a set of elements $\Gamma^{-1} = \{e^{-1} \mid e \in \Gamma^1\}$, that satisfies the following four axioms, for all $u, v \in \Gamma_0$ and $e, f \in \Gamma^1$:

- (V): $vv = v$ and $vu = 0$ if $v \neq u$,
- (E1): $(e)s \ e = e \ (e)r = e$,
- (E2): $(e)r \ e^{-1} = e^{-1} \ (e)s = e^{-1}$,
- (CK1): $f^{-1}f = (f)r$ and $e^{-1}f = 0$ if $e \neq f$.

For every $v \in \Gamma^0$ we define $v^{-1} = v$, and for every $q = e_1 \cdots e_k \in \text{Path}(\Gamma)$ we define $q^{-1} = e_k^{-1} \cdots e_1^{-1}$. It follows directly, by repeated application of (CK1), that every non-zero element in $S(\Gamma)$ can be written in the form pq^{-1} for some $p, q \in \text{Path}(\Gamma)$. It is routine to show that $S(\Gamma)$ is an inverse semigroup, with $(pq^{-1})^{-1} = qp^{-1}$ for every non-zero $pq^{-1} \in S(\Gamma)$.

The congruences of a graph inverse semigroup were characterised in [39].

A *Wang triple* (H, W, f) on Γ consists of a hereditary set $H \subseteq \Gamma^0$, a set $W \subseteq \{v \in \Gamma^0 \setminus H \mid |(v)s_{\Gamma \setminus H}^{-1}| = 1\}$, and a *cycle function* $f : C(\Gamma^0) \rightarrow \mathbb{Z}^+ \cup \{\infty\}$ (i.e., $(c)f = 1$ for all $c \in C(H)$, $(c)f = \infty$ for all $c \notin C(H \cup W)$, and the restriction of f to $C(W)$ is invariant under cyclic permutations). In [39], the term ‘‘congruence triple’’ is used for this concept.

Given a Wang triple (H, W, f) on a graph Γ , we define the corresponding congruence to be the least congruence on $S(\Gamma)$ containing the following set:

$$(2) \quad (H \times \{0\}) \cup \{(w, ee^{-1}) \mid w \in W, (e)s = w, (e)r \notin H\} \\ \cup \{(c^{(c)}f, (c)s) \mid c \in C(W), (c)f \in \mathbb{Z}^+\}.$$

Henceforth, and forever, we write identify the Wang triples and the congruences they represent.

An *isolated vertex* in a graph Γ is a vertex $v \in \Gamma^0$ such that $v \neq (e)s$ and $v \neq (e)r$ for all $e \in \Gamma^1$. An *out-edge* of a vertex $v \in \Gamma^0$ is an edge $e \in \Gamma^1$ such that $(v)s = e$.

Theorem 6.1. *A Wang triple (H, W, f) of a graph inverse semigroup $S(\Gamma)$ is an idempotent-pure congruence if and only if H is a set of isolated vertices and $(c)f = \infty$ for all $c \in C(W)$.*

Proof. (\Rightarrow): We prove the contrapositive. If H contains a vertex v that is not isolated, then there is $e \in \Gamma^1$ such that $(e)r = v$ or $(e)s = v$. Thus $(e, 0) \in (H, W, f)$. But e is not an idempotent, and 0 is idempotent, and so (H, W, f) is not idempotent-pure.

If $(c)f = x \in \mathbb{Z}^+$ for some $c \in C(W)$, then $(c^x, (c)s) \in (H, W, f)$. Since c^x is not an idempotent and $(c)s$ is an idempotent, (H, W, f) is not idempotent-pure.

(\Leftarrow): We define $A = (\{0\} \cup H)^2$ and we define B to consist of the pairs $((xp_1)(yp_1)^{-1}, (xp_2)(yp_2)^{-1}) \in S(\Gamma)$ such that $x, y, p_1, p_2 \in \text{Path}(\Gamma)$, $(x)r = (p_1)s = (p_2)s = (y)r$; $(e)s \in W$, for all e in p_1 or p_2 . Let $\rho = A \cup B \cup \{(x, x) : x \in S(\Gamma)\}$. Note that ρ never relates an idempotent to a non-idempotent. It is thus sufficient to show that $\rho = (H, W, f)$. If $(x, y) \in A$, then x and y both belong to $H \cup \{0\}$, and so $(x, y) \in (H, W, f)$ also.

If $((xp_1)(yp_1)^{-1}, (xp_2)(yp_2)^{-1}) \in B$, where $p_1 = e_1 \cdots e_n$ for some $e_1, \dots, e_n \in \Gamma^1$, then

$$(xp_1(yp_1)^{-1}, xy^{-1}) = (xp_1p_1^{-1}y^{-1}, xy^{-1}) = (xe_1 \cdots e_n e_n^{-1} \cdots e_1^{-1}y^{-1}, xy^{-1}).$$

Since $(s(e_i), e_i e_i^{-1}) \in (H, W, f)$, it follows that for every i

$$(xe_1 \cdots e_i e_i^{-1} \cdots e_1^{-1}y^{-1}, xe_1 \cdots e_{i-1} (e_i) s e_{i-1}^{-1} \cdots e_1^{-1}y^{-1}) \\ = (xe_1 \cdots e_i e_i^{-1} \cdots e_1^{-1}y^{-1}, xe_1 \cdots e_{i-1} e_{i-1}^{-1} \cdots e_1^{-1}y^{-1}) \in (H, W, f).$$

Hence, by transitivity,

$$(xe_1 \cdots e_n e_n^{-1} \cdots e_1^{-1}y^{-1}, xy^{-1}) \in (H, W, f).$$

So $(xp_1(yp_1)^{-1}, xy^{-1}) \in (H, W, f)$ and hence by symmetry $(xp_2(yp_2)^{-1}, xy^{-1}) \in (H, W, f)$. It follows that $\rho \subseteq (H, W, f)$.

For the converse, it suffices to show that ρ is a congruence, and that ρ contains the pairs in (2).

To show that ρ is transitive, we will check individually if $A \circ B$, $A \circ A$ and $B \circ B$ are all contained in ρ . For $A \circ A$, if $(x, y), (y, z) \in A$, then $x, z \in H \cup \{0\}$ and so $(x, z) \in A$. For $A \circ B$, if $(z, xp_1(y_{p_1})^{-1}) \in A$ and $(xp_1(y_{p_1})^{-1}, xp_2(y_{p_2})^{-1}) \in B$. As it lies in a pair in A , $xp_1(y_{p_1})^{-1}$ is a vertex in H or 0 . As 0 can never occur in a pair in B , we can assume that $xp_1(y_{p_1})^{-1}$ is a vertex in H . However, as the first entry in a pair in B , $xp_1(y_{p_1})^{-1}$ either contains a vertex in W , or p_1 is the empty path. The first case cannot happen as there is no path from $\Gamma \setminus H$ to H , and $W \cap H = \emptyset$. In the second case, $xp_1(y_{p_1})^{-1} = xy^{-1}$, which lies in A . Thus xy^{-1} is a vertex in H , and so $x = y \in H$. As p_2 is either empty or a path starting in H which intersects a vertex in W , which never happens, we have that p_2 is empty. Therefore $(z, xp_2(y_{p_2})^{-1}) = (z, xy^{-1}) = (z, xp_1(y_{p_1})^{-1}) \in A$.

For $B \circ B$, suppose $\mathbf{q} = (xp_1(y_{p_1})^{-1}, xp_2(y_{p_2})^{-1}) \in B$ and $\mathbf{r} = (wp_3(z_{p_3})^{-1}, wp_4(z_{p_4})^{-1}) \in B$, where $xp_2(y_{p_2})^{-1} = wp_3(z_{p_3})^{-1}$. Either x is a prefix of w or w is a prefix of x ; without loss of generality suppose w is a prefix of x . So $x = wu$, for some path u . It follows that $p_3 = up_2$. So $xp_1 = wup_1$ and $\mathbf{q} = (wup_1(y_{p_1})^{-1}, wup_2(y_{p_2})^{-1})$. Additionally, as $p_3 = up_2$, $\mathbf{r} = (wup_2(z_{p_2})^{-1}, wp_4(z_{p_4})^{-1})$. We also have that $y = zu$, as $wup_2(y_{p_2})^{-1} = wup_2(z_{p_2})^{-1}$. Thus $\mathbf{q} = (wup_1(z_{p_1})^{-1}, wup_2(z_{p_2})^{-1})$ and $\mathbf{r} = (wp_3(z_{p_3})^{-1}, wp_4(z_{p_4})^{-1})$. Thus $(wup_1(z_{p_1})^{-1}, wp_4(z_{p_4})^{-1}) \in B$, as required.

We now show that ρ is a congruence. If $(x, y) \in \rho$, then $(x^{-1}, y^{-1}) \in \rho$, and so it suffices to show that ρ is a right congruence. If $(x, y) \in A$ and $s \in \Gamma^0 \cup \Gamma^1 \cup (\Gamma^1)^{-1}$, then as H consists of isolated vertices, $xs, ys \in H \cup \{0\}$, and so $(xs, ys) \in A$. If $(xp_1(y_{p_1})^{-1}, xp_2(y_{p_2})^{-1}) \in B$ and $s \in \Gamma^0 \cup \Gamma^1 \cup (\Gamma^1)^{-1}$, then at least one of the following cases applies:

- (1) $s \in \Gamma^0$ and $s = (y)\mathbf{s}$. In this case, $(xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (xp_1(y_{p_1})^{-1}, xp_2(y_{p_2})^{-1}) \in B$.
- (2) $s \in \Gamma^0$ and $s \neq (y)\mathbf{s}$. In this case $((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (0, 0) \in A$.
- (3) $s \in \Gamma^1$ and $(s)\mathbf{s} \neq (y)\mathbf{s}$. Again, $((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (0, 0) \in A$.
- (4) $s \in \Gamma^1$ and s is the first edge in y . Then $y = sy'$, for some path y' in Γ , and so $((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = ((xp_1(y'_{p_1})^{-1}, xp_2(y'_{p_2})^{-1}) \in B$.
- (5) $s \in \Gamma^1$, $y \notin \Gamma^0$, and $(s)\mathbf{s} = (y)\mathbf{s}$, s is not the first edge in y . Again,

$$((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (0, 0) \in A.$$

- (6) $s \in \Gamma^1$, $y \in \Gamma^0 \setminus W$ and $(s)\mathbf{s} = (y)\mathbf{s}$. From the definition of B , the source of each edge in p_1 lies in W , and $(p_1)\mathbf{s} = (y)\mathbf{r} = y \notin W$. In particular, p_1 contains no edges, i.e. $p_1 \in \Gamma^0$. Similarly, $p_2 \in \Gamma^0$. Thus $p_1 = p_2 = y$. So $((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = ((xy(y)^{-1}s, xy(y)^{-1}s) = (xs, xs) \in \rho$.
- (7) $s \in \Gamma^1$, $y \in W$, $(s)\mathbf{s} = y$, and $p_1, p_2 \notin \Gamma^0$. Then by the choice of W , s is the unique edge with source y , and so $p_1 = sp'_1$ and $p_2 = sp'_2$ for some paths p'_1, p'_2 . Hence

$$(xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (xsp'_1p'^{-1}_1, xsp'_2p'^{-1}_2) \in B.$$

- (8) $s \in \Gamma^1$, $y \in W$, $(s)\mathbf{s} = y$, and $p_1, p_2 \in \Gamma^0$. As $s(p_1) = \mathbf{s}(p_2) = \mathbf{r}(y) = y$, we have $p_1 = y = p_2$, and so $(xp_1(y_{p_1})^{-1}, xp_2(y_{p_2})^{-1}) = (xp_1(y_{p_1})^{-1}, xp_1(y_{p_1})^{-1}) \in \rho$.
- (9) $s \in \Gamma^1$ and $y \in W$, $(s)\mathbf{s} = y$ and precisely one of p_1 and p_2 lies in Γ^0 . Assume without loss of generality, that $p_1 \in \Gamma^0$. Then as W is part of a Wang triple, every vertex in W has a unique out-edge, and so s is the unique edge with source y , and so $p_2 = sp'_2$ for some path p'_2 . Hence

$$(xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (xs, xsp'_2p'^{-1}_2) \in B.$$

- (10) $s^{-1} \in \Gamma^1$ and $(s)\mathbf{s} = (y)\mathbf{s}$. Then let $z = s^{-1}y$. Note $(z)\mathbf{r} = (y)\mathbf{r}$. In addition,

$$((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = ((xp_1(s^{-1}y_{p_1})^{-1}, xp_2(s^{-1}y_{p_2})^{-1}) = ((xp_1(z_{p_1})^{-1}, xp_2(z_{p_2})^{-1}) \in B.$$

- (11) $s^{-1} \in \Gamma^1$ and $(s)\mathbf{s} \neq (y)\mathbf{s}$. Again, $((xp_1(y_{p_1})^{-1}s, xp_2(y_{p_2})^{-1}s) = (0, 0) \in A$.

Hence ρ is a congruence, as required. \square

Next, we state and prove the main theorem of this section.

Theorem 6.2. *A graph inverse semigroup defined using a graph Γ is E -disjunctive if and only if Γ has no isolated vertices, and every vertex in Γ has either 0 or at least 2 out-edges.*

Proof. By Theorem 6.1, $S(\Gamma)$ admits a non-trivial idempotent-pure congruence if and only if it has a (potentially empty) set H of isolated vertices and a set $W \subseteq \{v \in \Gamma^0 \setminus H \mid |(v)\mathbf{s}_{\Gamma \setminus H}^{-1}| = 1\}$ such that there is a cycle function f on Γ such that $(c)f = \infty$ for all $c \in C(W)$.

(\Rightarrow): Suppose that Γ has isolated vertices or there exists a vertex $v \in \Gamma^0$ with $|(v)\mathbf{s}^{-1}| = 1$. In the former case, taking H to be the non-empty set of isolated vertices, and $W = \emptyset$ gives a non-trivial idempotent-pure congruence. In the latter case, $H = \emptyset$ and $W = \{v \in \Gamma^0 : |(v)\mathbf{s}^{-1}| = 1\} \neq \emptyset$ gives a non-trivial idempotent-pure congruence. Thus $S(\Gamma)$ is not E -disjunctive.

(\Leftarrow): Suppose that Γ has no isolated vertices and that every vertex has either 0 or 2 out-edges. For Γ to admit a non-trivial idempotent-pure congruence, we would have $H = \emptyset$ and $W = \emptyset$, as these are the only possibilities for H and W . However, the congruence defined by this pair is equality, and so $S(\Gamma)$ is E -disjunctive. \square

7. FINITE MONOGENIC INVERSE MONOIDS

In this section we characterise those finite monogenic inverse monoids that are E -disjunctive. In order to do this, we require the following characterisation of finite monogenic inverse monoids, and some related results, mostly arising from [37].

Although we concentrate on monoids rather than semigroups in this section, analogues of the main results hold for monogenic inverse semigroups, and these can be concluded from the results in this section together with Corollary 3.3.

Lemma 7.1 (Lemma 8 in [9]). *If M is a finite monogenic inverse submonoid of I_n , then there exist $a, b \in \mathbb{N}$ such that $a + b = n$ and M is isomorphic to the inverse submonoid of I_n generated by*

$$x = [1, \dots, a] \cup p.$$

where p is some permutation on the set $\{a + 1, \dots, b + 1\}$. Moreover, the monoid is isomorphic to the submonoid of $I_{a+|p|}$ generated by $[1, \dots, a] \cup (a + 1, \dots, a + |p|)$.

Let $n, k \in \mathbb{N}$ be such that $n \geq 0$ and $k \geq 1$ and let $S_{n,k}$ be defined by the inverse monoid presentation

$$\text{Inv}\langle x \mid x^n x^{-n} = x^{n+1} x^{-(n+1)}, x^n x^{-n} = x^n x^{-n} x^k \rangle.$$

Theorem 7.2 (Theorem 10 in [9]). *The inverse submonoid of I_{n+k} generated by a partial permutation*

$$[1, 2, \dots, n] \cup (n + 1, n + 2, \dots, n + k)$$

is isomorphic to $S_{n,k}$ for all $n \geq 0$ and $k \geq 1$. Moreover, if $m \geq 0$ and $l \geq 1$, then $S_{n,k} \cong S_{m,l}$ if and only if $n = m$ and $k = l$.

Lemma 7.3. *Let $n \geq 0$ and $k \geq 1$. Then every element of $S_{n,k}$ can be uniquely expressed in the form $x^{-a} x^b x^{-b} x^c$ where $a, c \leq b < n$ or $x^n x^{-n} x^a$ where $0 \leq a < k$; and $E(S_{n,k}) = \{x^{-a} x^b x^{-b} x^a \mid a \leq b < n\} \cup \{x^n x^{-n}\}$.*

Proof. By [9, Lemma 6] (or [8, Proposition 4]) the set of words of the form $x^{-a} x^b x^{-b} x^c$ with $b < n$ contains representatives for every element of $S_{n,k}$. In particular, $x^{-a} x^b x^{-b} x^c$ when restricted to the set $\{1, \dots, n\}$ is the partial permutation $\{(a + 1, c + 1), (a + 2, c + 2), \dots, (a + (n - b), c + (n - b))\}$. Hence distinct words of the form $x^{-a} x^b x^{-b} x^c$ represent distinct elements of $S_{n,k}$. The remaining elements of the monoid are those in the ideal generated by $x^n x^{-n}$. Since $S_{n,k}$ and the inverse monoid generated by $x = [1, \dots, n](n + 1, \dots, n + k)$ coincide (by Theorem 7.2), $x^n x^{-n}$ is the

identity on $\{n+1, \dots, n+k\}$, and so $x^n x^{-n} x = (n+1, \dots, n+k)$ generates a cyclic group of order k .

The claim about idempotents is immediate from Theorem 7.2. \square

Lemma 7.4. [Theorem 2 in [37]] *If $n \geq 1$ and $k \geq 1$, then*

$$|S_{n,k}| = \frac{n(n+1)(2n+1)}{6} + k.$$

Note that $S_{0,k} = S_{1,k}$ and so $|S_{0,k}| = k+1$.

The next theorem is the main result of this section, characterising the idempotent-pure congruences on $S_{n,k}$ when $n \geq 0$ and $k \geq 1$.

Theorem 7.5. *If ρ is a non-trivial congruence on $S_{n,k}$, then $S_{n,k}/\rho \cong S_{n',k'}$, where $1 \leq n' \leq n$ and $k'|k$. Moreover, ρ is idempotent-pure if and only if $k' = k$ and $n \leq k$.*

Proof. Every homomorphic image of an inverse monoid is an inverse monoid, and since $S_{n,k}$ is monogenic, $S_{n,k}/\rho$ is monogenic also. In particular, $S_{n,k}/\rho$ is isomorphic to $S_{n',k'}$, for some $n' \leq n$ and $k' \leq k$ (by Lemma 7.1, and Theorem 7.2). Since $S_{n',k'}$ contains a cyclic subgroup of order k' (by Theorem 7.2) it follows that $k'|k$.

We now show that if ρ is idempotent-pure, then $k' = k$ and $n \leq k$. Suppose ρ is idempotent-pure. Since $k'|k$ it suffices to show that $k' \geq k$. Seeking a contradiction, suppose that $k' < k$. Then

$$[x^n x^{-n}]_\rho = [x^{n'} x^{-n'}]_\rho = [x^{n'} x^{-n'} x^{k'}]_\rho = [x^n x^{-n} x^{k'}]_\rho.$$

But $x^n x^{-n} x^{k'} \notin E(S_{n,k})$ by Lemma 7.3, a contradiction. It follows that $k' = k$.

It remains to show that $n \leq k$. If $n > k = k'$, then since $n' \leq n$, we can assume $n' < n$, as otherwise ρ would be trivial. So

$$[x^{n-1} x^{-(n-1)}]_\rho = [x^{n'} x^{-n'}]_\rho = [x^{n'} x^{-n'} x^k]_\rho = [x^{n'} x^{-n'} x^{k'}]_\rho = [x^{(n-1)} x^{-(n-1)} x^k]_\rho,$$

where $x^{(n-1)} x^{-(n-1)} x^k \notin E(S_{n,k})$ by Lemma 7.3. Thus ρ is not idempotent-pure, a contradiction. Hence $k' = k$ and $n \leq k$ as required.

For the converse, suppose that $k' = k$ and $n \leq k$. Then for all $0 \leq a, c \leq b < n$, $1 \leq f \leq n$, and $e, g < \max(n, k)$

$$\begin{aligned} [x^{-a} x^b x^{-b} x^a]_\rho = [x^{-e} x^f x^{-f} x^g]_\rho &\implies (b = f \text{ and } e = g = a) \text{ or } (b, f \geq n' \text{ and } -a + a = -e + g) \\ &\implies (b = f \text{ and } e = g = a) \text{ or } (b, f \geq n' \text{ and } e = g) \\ &\implies e = g. \end{aligned}$$

Hence $x^{-e} x^f x^{-f} x^g$ is an idempotent by Lemma 7.3. Also

$$[x^n x^{-n}]_\rho = [x^{-e} x^f x^{-f} x^g]_\rho \implies f \geq n' \text{ and } k|(g-e) \implies k|(g-e) \implies g-e=0 \implies e=g$$

so $x^{-e} x^f x^{-f} x^g$ is again an idempotent, and ρ is idempotent-pure. \square

Next, we use Theorem 7.5 to characterise the E -disjunctive finite monogenic inverse monoids.

Corollary 7.6. *A finite monogenic inverse monoid is E -disjunctive if and only if it is isomorphic to $S_{n,k}$ for some k, n with $n > k$ or $n = 1$.*

Proof. By Theorem 7.5, a congruence ρ on $S_{n,k}$ is idempotent-pure if and only if $k' = k$ and $n \leq k$ (where $S_{n,k}/\rho \cong S_{n',k'}$). Let $P(n, k)$ be the statement: for all $(n', k') \neq (n, k)$ with $1 \leq n' \leq n$, $1 \leq k'|k$, either $k' \neq k$ or $n > k$. So $S_{n,k}$ is E -disjunctive if and only if $P(n, k)$ is true. We show that $P(n, k)$ holds if and only if $n > k$ or $n = 1$.

(\Rightarrow): Suppose $P(n, k)$ is true. If $n = 1$, then the proof is complete. Otherwise set $n' = 1 < n$ and $k' = k$. Since $P(n, k)$ holds, either $k' \neq k$ or $n > k$. Hence, since $k' = k$, $n > k$, as required.

(\Leftarrow): If $n > 1$, then $n > k$, and so $P(n, k)$ holds immediately. Otherwise if $n = 1$, then for all $n' \leq n$ and $k' | k$ with $(n', k') \neq (n, k)$, $n' = n = 1$ and so $k' \neq k$. Hence in either case $P(n, k)$ holds. \square

We conclude this section by showing that, asymptotically, almost none of the monogenic inverse monoids are E -disjunctive.

Corollary 7.7. *The proportion of isomorphism classes of monogenic inverse monoids of size at most m which are E -disjunctive tends to 0 as m tends to infinity.*

Proof. Let $m \in \mathbb{Z}_{\geq 1}$ be given. By Theorem 7.2, the number of monogenic inverse monoids up to isomorphism with size at most m equals the number of $S_{n,k}$ such that $|S_{n,k}| \leq m$. Similarly, by Corollary 7.6, the number of E -disjunctive monogenic inverse monoids up to isomorphism with size at most m equals the number of $S_{n,k}$ such that $|S_{n,k}| \leq m$ where $n > k$ or $n = 1$. In particular, it suffices to find the proportion of pairs $\{(n, k) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1} \mid |S_{n,k}| \leq m\}$ such that $n > k$ or $n = 1$. By Lemma 7.4,

$$|S_{n,k}| = \frac{n(n+1)(2n+1)}{6} + k \leq m \quad \text{if and only if} \quad k \leq m - \sum_{i=0}^n i^2 = m - \frac{n(n+1)(2n+1)}{6}.$$

For all $j \in \mathbb{Z}_{\geq 1}$, there exists $m_j \in \mathbb{Z}_{\geq 1}$ such that $m_j \geq j$ and the number of (n, k) , such that $|S_{n,k}| \leq m_j$ is greater than jm_j . For example, if $j = 6$, then for $m \geq 105$, the number of (n, k) with $|S_{n,k}| \leq m$ is

$$\begin{aligned} \sum_{n=1}^{\infty} \max\left(m - \sum_{i=0}^n i^2, 0\right) &\geq \sum_{n=1}^6 \left(m - \sum_{i=0}^n i^2\right) \\ &= m + (m-1) + (m-5) + (m-14) + (m-30) + (m-55) \\ &= 7m - 105 \geq 6m. \end{aligned}$$

We next find an upper bound for the number of (n, k) such that $S_{n,k}$ is E -disjunctive and $|S_{n,k}| \leq m$. It suffices to give an upper bound for the number of pairs in $\{(n, k) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1} \mid |S_{n,k}| \leq m\}$ such that $n > k$ or $n = 1$:

$$\begin{aligned} &\sum_{n=1}^{\infty} |\{k \in \mathbb{Z}_{\geq 1} \mid (n > k \text{ or } n = 1) \text{ and } |S_{n,k}| \leq m\}| \\ &= |\{k \in \mathbb{Z}_{\geq 1} \mid |S_{1,k}| \leq m\}| + \sum_{n=2}^{\infty} |\{k \in \mathbb{Z}_{\geq 1} \mid n > k \text{ and } |S_{n,k}| \leq m\}| \\ &= |\{k \in \mathbb{Z}_{\geq 1} \mid k + 1 \leq m\}| + \sum_{n=2}^{\infty} |\{k \in \mathbb{Z}_{\geq 1} \mid n > k \text{ and } n(n+1)(2n+1)/6 + k \leq m\}| \\ &= m - 1 + \sum_{n=2}^{\infty} |\{k \in \mathbb{Z}_{\geq 1} \mid n > k \text{ and } n(n+1)(2n+1)/6 + k \leq m\}| \\ &= m - 1 + \sum_{n=2}^{\infty} |\{k \in \mathbb{Z}_{\geq 1} \mid n > k \text{ and } k \leq m - n(n+1)(2n+1)/6\}| \\ &\leq m - 1 + \sum_{n=2}^{\infty} \min(\max(m - n(n+1)(2n+1)/6, 0), n - 1) \\ &\leq m - 1 + \sum_{n=2}^{\infty} \min(\max(m - n^3/6, 0), n - 1) \end{aligned}$$

$$\begin{aligned}
 &\leq m - 1 + \sum_{n=2}^{\lfloor \sqrt[3]{6m} \rfloor} \min(m - n^3/6, n - 1) \\
 &\leq m - 1 + \sum_{n=2}^{\lfloor \sqrt[3]{6m} \rfloor} n - 1 \\
 &\leq m - 1 - \sqrt[3]{6m} + \frac{\sqrt[3]{6m}(\sqrt[3]{6m} + 1)}{2} \\
 &\leq m + \sqrt[3]{6m}\sqrt[3]{6m} + 1 \\
 &\leq 7m.
 \end{aligned}$$

Thus

$$\lim_{m \rightarrow \infty} \frac{\# \text{ monogenic } E\text{-disjunctive inverse semigroups of size at most } m}{\# \text{ monogenic inverse semigroups of size at most } m} \leq \lim_{n \rightarrow \infty} \frac{7m_n}{nm_n} = \lim_{n \rightarrow \infty} \frac{7}{n} = 0.$$

□

Part 3. A structure theory for E -disjunctive inverse semigroups

In this part of the paper we consider various structural properties of E -disjunctive semigroups. In Section 8 we find a bound on the ration of idempotent to non-idempotent elements in an E -disjunctive semigroup; in Section 9 we consider the maximum E -disjunctive homomorphic images of an inverse semigroup; in Section 10 we define a notion we refer to as preactions which is used extensively in the final section Section 11; where we prove that every inverse semigroup can be defined in terms of a semilattice and an E -disjunctive inverse semigroup.

8. RATIO OF IDEMPOTENTS TO NON-IDEMPOTENTS

Roughly speaking semilattices are as far from being E -disjunctive as possible. More specifically, every E -disjunctive homomorphic image of a semilattice is trivial. In this section, we precisely formalise this notion by showing that inverse semigroups with too many idempotents are not E -disjunctive. In particular, we will prove the following theorem.

Theorem 8.1. *Let S be an E -disjunctive inverse semigroup and $\kappa = |S \setminus E(S)|$. Then $|S| \leq 2^\kappa + \kappa$.*

If S is a finite inverse semigroup, then we define $\mathbf{h}: S \rightarrow \mathbb{N}_{\geq 0}$ so that $(s)\mathbf{h}$ is the largest value in $\mathbb{N}_{\geq 0}$ such that there is a chain with maximum element s of this length in the natural partial order of S .

Lemma 8.2. *If S is a finite inverse semigroup and $s, t \in S$ are such that $s \leq_{\mathcal{R}} t$ or $s \leq_{\mathcal{L}} t$, then $(s)\mathbf{h} \leq (t)\mathbf{h}$.*

Proof. We prove the lemma in the case that $s \leq_{\mathcal{R}} t$, the proof in the other case is similar. Let $s' \in S$ be such that $ts' = s$. Suppose that $n = (t)\mathbf{h}$ and suppose that $t_1 := t, t_2, \dots, t_n \in S$ are such that $t_i > t_{i+1}$ for all i . We set $s_i = t_i s'$ for all i . Since $t_{i+1} \leq t_i$ implies $t_{i+1} = (t_{i+1} t_{i+1}^{-1}) t_i$ (by [15, Proposition 5.2.1]) it follows that

$$s_{i+1} = t_{i+1} s' = (t_{i+1} t_{i+1}^{-1}) t_i s' = (t_{i+1} t_{i+1}^{-1}) s_i \leq s_i$$

and so $(s)\mathbf{h} \geq (t)\mathbf{h}$. □

Lemma 8.3. *Let S be a finite inverse semigroup. If $s, t \in S$ are such that $s \mathcal{S} t$, then $(s)\mathbf{h} = (t)\mathbf{h}$.*

Proof. Since $s \mathcal{S} t$, there exists $u \in S$ such that $s \mathcal{R} u \mathcal{L} t$ and so $(s)\mathbf{h} = (u)\mathbf{h} = (t)\mathbf{h}$, by Lemma 8.2. □

Lemma 8.4. *Let S be a finite inverse semigroup and let $e, f \in S$ be distinct idempotents. If $(f)\mathbf{h} \leq (e)\mathbf{h}$, then $(ef)\mathbf{h} < (e)\mathbf{h}$.*

Proof. If e and f are incomparable, then $ef < e$ or $ef < f$. In either case, $(e)\mathbf{h} \geq (f)\mathbf{h} > (ef)\mathbf{h}$. Otherwise, $f < e$, and so $ef < e$ and $(ef)\mathbf{h} < (e)\mathbf{h}$. \square

If S is a finite inverse semigroup, then we define $N(S)$ to be the set of idempotents in $e \in S$ such that there exists a non-idempotent $u \in S$ such that $uu^{-1} = e$. We also define

$$\phi_S: E(S) \rightarrow \mathcal{P}(N(S))$$

by

$$(f)\phi_S = \{e \in N(S) \mid e \leq f\}.$$

Lemma 8.5. *If S is a finite inverse semigroup, then $\phi_S: E(S) \rightarrow \mathcal{P}(N(S))$ defined by*

$$(f)\phi_S = \{e \in N(S) \mid e \leq f\}$$

is a homomorphism where the operation on $\mathcal{P}(N(S))$ is \cap .

Proof. Let $e, f \in E(S)$, and $g \in N(S)$. Then

$$\begin{aligned} g \in (e)\phi_S \cap (f)\phi_S &\iff g \leq e \text{ and } g \leq f \\ &\iff ge = g = gf \\ &\iff gef = g \\ &\iff g \leq ef \\ &\iff g \in (ef)\phi_S. \end{aligned} \quad \square$$

If I is an ideal of an inverse semigroup S , then the natural partial order on I is just the intersection of the natural partial order of S with $I \times I$.

Lemma 8.6. *If S is a finite E -disjunctive inverse semigroup, then $\phi_S: E(S) \rightarrow \mathcal{P}(N(S))$ is an embedding.*

Proof. By Lemma 8.5, we need only show that ϕ_S is injective. We proceed by induction on $(S)\mathbf{h} := \max\{(s)\mathbf{h} \mid s \in S\}$. If $(S)\mathbf{h} = 0$, then $S = \emptyset$ and so ϕ_S is an embedding.

Suppose that $(S)\mathbf{h} = k$ and the result holds for all finite E -disjunctive inverse semigroups T with $(T)\mathbf{h} < k$. Then the set $I := \{s \in S \mid (s)\mathbf{h} < k\}$ is an ideal of S , and $(I)\mathbf{h} = k - 1$, and I is E -disjunctive by Lemma 2.8. Thus by induction ϕ_I is an embedding.

Suppose that there exist $e, f \in E(S)$ such that $(e)\phi_I = (f)\phi_S$. If $e, f \in I$, then, since ϕ_I is just the restriction of ϕ_S to I ,

$$(e)\phi_I = (e)\phi_S = (f)\phi_S = (f)\phi_I$$

and so, since ϕ_I is injective, $e = f$, as required.

Hence it remains to prove the lemma in the case that $e \notin I$ or $f \notin I$. Suppose without loss of generality that $e \notin I$ and seeking a contradiction that $e \neq f$. By assumption,

$$(3) \quad (e)\phi_S = (e)\phi_S \cap (e)\phi_S = (e)\phi_S \cap (f)\phi_S = (ef)\phi_S.$$

If there exists a non-idempotent $u \in S$ such that $e = uu^{-1}$, then $e \in (e)\phi_S$. Since $e \neq f$ and $(e)\phi_S = (f)\phi_S$ by assumption, $e < f$. It follows that $(f)\mathbf{h} > (e)\mathbf{h} = k = (s)\mathbf{h}$, which is a contradiction.

Suppose that $uu^{-1} \neq e$ for all non-idempotents $u \in S$, and suppose that $\rho = \{(e, ef)\} \cup \Delta_S$. To reach our final contradiction it suffices to show that ρ is a congruence. We show ρ is a right congruence; the proof that ρ is a left congruence is symmetric.

Let $u \in S$ be arbitrary. If $u = e$, then $eu = ee = epef = efe = efu$ and so $eupefu$. If $u \neq e$, then we will also show that $eu = efu$. By assumption, $uu^{-1} \neq e$. Since $(e)\mathbf{h} = k$, it follows from Lemma 8.4, that $euu^{-1}, efuu^{-1} \in I$. So

$$\begin{aligned} (euu^{-1})\phi_I &= (euu^{-1})\phi_S \\ &= (e)\phi_S \cap (uu^{-1})\phi_S \\ &= (ef)\phi_S \cap (uu^{-1})\phi_S && \text{by (3)} \\ &= (efuu^{-1})\phi_S \\ &= (efuu^{-1})\phi_I. \end{aligned}$$

Thus $euu^{-1} = efuu^{-1}$, since ϕ_I is an embedding and so $eu = efu$ also. Therefore ρ is a non-trivial idempotent-pure congruence on S , and so S is not E -disjunctive, a contradiction. \square

The converse of Lemma 8.6 is not true, for example, if S is the strong semilattice of groups defined by an identity map from the cyclic group C_2 of order 2 to C_2 , then ϕ_S is injective, but S is not E -disjunctive.

Let S be an inverse semigroup and let $e \in E(S)$. Then *syntactic readout* of e is the function $\phi_e: ((S \setminus E(S)) \cup \{1_S\}) \times ((S \setminus E(S)) \cup \{1_S\}) \rightarrow \{0, 1\}$ defined by

$$(\alpha, \beta)\phi_e = \begin{cases} 0 & \text{if } \alpha\beta \in E(S) \\ 1 & \text{if } \alpha\beta \notin E(S) \end{cases}$$

for all $\alpha, \beta \in S \setminus E(S) \cup \{1\}$.

Lemma 8.7. *Let S be an E -disjunctive inverse semigroup and let $e, f \in E(S)$. Then $\phi_e = \phi_f$ (i.e. e and f have the same syntactic readout) if and only if $e = f$.*

Proof. The converse implication is trivial.

For the forward implication, since S is E -disjunctive, it follows from Lemma 2.5 that the syntactic congruence on S is Δ_S . Suppose that $e, f \in E(S)$ have the same syntactic readout. To show $e = f$, it suffices to show that (e, f) belongs to the syntactic congruence of S . In other words, to show that

$$(4) \quad \alpha e \beta \in E(S) \iff \alpha f \beta \in E(S)$$

for all $\alpha, \beta \in S^1$. By assumption, (4) holds for all $\alpha, \beta \in S^1 \setminus E(S)$.

Suppose that $\alpha \in E(S)$ or $\beta \in E(S)$. We may suppose without loss of generality that $\alpha \in E(S)$ and that $\alpha e \beta \in E(S)$. Since $\alpha \in E(S)$, $\alpha e \beta = e\alpha\beta$ and $\alpha f \beta = f\alpha\beta$. If $\alpha\beta \in E(S)$, then $f\alpha\beta = \alpha f \beta \in E(S)$. Otherwise, $\alpha\beta \notin E(S)$, and since $(1_S, \alpha\beta)\phi_e = (1_S, \alpha\beta)\phi_f$ it follows that $e\alpha\beta \in E(S)$ implies $f\alpha\beta \in E(S)$. Hence in both cases $\alpha e \beta \in E(S)$ implies $\alpha f \beta \in E(S)$, and the converse implication follows by symmetry. \square

Proof of Theorem 8.1. We consider the cases when S is finite and infinite separately.

Suppose that S is finite. Clearly, $|S| = |E(S)| + n$. By Lemma 8.6, it follows that

$$|S| = |E(S)| + n = |(E(S))\phi_S| + n \leq |\mathcal{P}(N(S))| + n.$$

The map sending a non-idempotent s of S to ss^{-1} , is surjective with image set $N(S)$. Thus $|N(S)| \leq n$. In particular

$$|S| \leq |\mathcal{P}(N(S))| + n = 2^{|N(S)|} + n \leq 2^n + n.$$

Suppose that S is infinite, and that $\kappa = |S \setminus E(S)|$. By Lemma 8.7, idempotents of S are uniquely determined by their syntactic readouts. There are at most $(2^\kappa)(2^\kappa) = 2^{2\kappa}$ syntactic readouts and so $|S| \leq 2^{2\kappa} + \kappa$. \square




FIGURE 2. Egg-box diagram of a semigroup that attains the bound in Theorem 8.1 when $\kappa = 3$

Corollary 8.8. *Let S be an infinite E -disjunctive inverse semigroup. Then S has infinitely many non-idempotents.*

The next example shows that the bound in Theorem 8.1 is sharp.

Example 8.9. Let κ be a finite or infinite cardinal. We define a Clifford semigroup S with κ non-idempotents and 2^κ idempotents by defining a strong semilattice of groups. The semilattice Y is the power set of κ under intersection; the groups are defined by:

$$G_y = \begin{cases} 1 & \text{if } |y| \neq 1 \\ C_2 & \text{if } |y| = 1 \end{cases}$$

for all $y \in Y$ where 1 denotes the trivial group and C_2 the cyclic group of order 2; and every homomorphism $\psi_{y,z} : G_y \rightarrow G_z$ with $y, z \in Y$ is constant; see Fig. 2 for an example. Clearly there is an idempotent in S for every subset of κ , and there is a non-idempotent for every element of κ . Hence $|S| = 2^\kappa + \kappa$. Since S is a semilattice of abelian groups, S is also commutative.

It remains to show that S is E -disjunctive. Let $e, f \in E(S)$ be such that $e < f$. If e and f are the idempotents belonging to G_y and G_z , respectively, then we abuse our notation by writing $\psi_{e,f}$ instead of $\psi_{y,z}$. By [1, Theorem 6], it suffices to show that there exists $g \in E(S)$ such that the map $\psi_{eg,fg}$ is not injective. Since $e < f$, f is non-zero. For this semigroup S , $N(S) = \{h \in E(S) \mid h \in G_y \text{ for some } y \in Y \text{ with } |y| = 1\}$. If $f \in N(S)$, then we set $g = f > e$ and so $g \not\leq e$. If $f \notin N(S)$, then there exists $g \in N(S)$ such that $g \leq f$ and $g \not\leq e$ by the construction of S . In either case, $g \in N(S)$, $g \not\leq e$, and $g \leq f$. Hence $eg < g$ and $fg = g$. The former implies that $eg = 0$, and so $\psi_{fg,eg}$ is a constant function from C_2 to a trivial group, and is not injective, as required.

9. MAXIMUM E -DISJUNCTIVE IMAGES

Every inverse semigroup S has an E -disjunctive quotient: the quotient of S by its syntactic congruence ρ . By Lemma 2.4, if T is any E -disjunctive semigroup such that T is a homomorphic image of S , then T is a quotient of S/ρ . As such we refer to S/ρ as the *maximum E -disjunctive quotient* of the inverse semigroup S . Since quotients and homomorphic images are interchangeable, we may also refer to S/ρ as the *maximum E -disjunctive image* of S . We will show that many properties can be exchanged between an inverse semigroup and its maximum E -disjunctive image. The situation is somewhat similar to the relationship between an E -unitary inverse semigroup and its maximum group homomorphic image. Of course, unlike the case for maximum group images,

maximum E -disjunctive images are not always groups, and so it is not clear to what extent this can be used to study inverse semigroups in general. In this section we will explore the relationship between an inverse semigroup and its maximum E -disjunctive image. In Section 11, we show that every inverse semigroup is described by its maximum E -disjunctive image and semilattice of idempotents, somewhat analogous to the description of E -unitary inverse semigroup via McAlister triples.

An inverse semigroup is E -unitary if and only if its minimum group congruence is idempotent-pure. Since every group is E -disjunctive, it follows that the maximum E -disjunctive image of an E -unitary inverse semigroup is a group. Conversely, if S is an inverse semigroup whose maximum E -disjunctive homomorphic image is a group, then the maximum group and E -disjunctive images coincide. In other words, we have the following result.

Proposition 9.1. *The maximum E -disjunctive image of an inverse semigroup S is a group if and only if S is E -unitary.*

Next, we use a theorem of Kambites [17] to show that a finitely generated inverse semigroup is finite if and only its maximum E -disjunctive image is finite. To do this, we first define the idempotent problem of a finitely generated inverse semigroup S as follows. Let Σ be a finite generating set for S . Then the *idempotent problem* of S , with respect to Σ , denoted $\text{IP}(S, \Sigma)$ is the language consisting of all words over $\Sigma \cup \Sigma^{-1}$ that represent idempotents in S .

Theorem 9.2 ([17]). *Let S be an inverse semigroup with a finite generating set Σ . Then S is finite if and only if $\text{IP}(S, \Sigma)$ is a regular language.*

Theorem 9.2 allows us to characterise the inverse semigroups with finite maximum E -disjunctive image as follows.

Theorem 9.3. *Let S be a finitely generated inverse semigroup. Then S has a finite maximum E -disjunctive image if and only if S is finite.*

Proof. (\Leftarrow): Since quotients of finite semigroups are finite, if S is finite, then so is its maximum E -disjunctive image.

(\Rightarrow): Suppose that S has a finite maximum E -disjunctive image T , that Σ is a finite generating set for S , and that $\phi: S \rightarrow T$ is an idempotent-pure epimorphism. Since ϕ is idempotent-pure and surjective, ϕ induces a bijection between $E(S)$ and $E(T)$ and an isomorphism from $(\Sigma \cup \Sigma^{-1})^*$ and $((\Sigma \cup \Sigma^{-1})\phi)^*$ that maps $\text{IP}(S, \Sigma)$ to $\text{IP}(T, (\Sigma)\phi)$. By Theorem 9.2, since T is finite, $\text{IP}(T, (\Sigma)\phi)$ is a regular language. Thus $\text{IP}(S, \Sigma)$, as the image under an isomorphism of a regular language, is itself regular. Applying Theorem 9.2 again yields that S is finite. \square

Every semilattice has trivial maximum E -disjunctive image. Since every finitely generated semilattice is finite, if S is an infinite semilattice, then S is not finitely generated, but its maximum E -disjunctive image is finite. In other words, the finitely generated hypothesis in Theorem 9.3 cannot be removed.

Lemma 9.4. *Let $\phi: S \rightarrow T$ be an idempotent-pure homomorphism. Then $\phi|_R$ is injective for every \mathcal{R} -class R of S .*

Proof. Suppose that $a, b \in S$ are such that $a\mathcal{R}b$ and suppose that $(a)\phi = (b)\phi$. By Green's Lemma the map $\lambda: R_a \rightarrow R_{a^{-1}a}$ defined by left multiplying by a^{-1} is a bijection. Hence

$$(a^{-1}a)\phi = (a^{-1})\phi \cdot (a)\phi = (a^{-1})\phi \cdot (b)\phi = (a^{-1}b)\phi$$

and $a^{-1}a\mathcal{R}a^{-1}b$ (since \mathcal{R} is a left congruence). Hence we may assume without loss of generality that a is an idempotent.

Since ϕ is idempotent-pure and $(a)\phi = (b)\phi$, it follows that b is an idempotent. Hence since $a\mathcal{R}b$ this implies $a = b$. \square

10. PREACTIONS

In this section we define a notion that is a weakening of the notion of an inverse semigroup action. This idea is somewhat analogous to the notion of partial actions introduced in [18]. We require this somewhat technical section in order to prove a generalisation of McAlister's P -Theorem [27] in Section 11. This generalisation describes every inverse semigroup in terms of an E -disjunctive inverse semigroup and a semilattice.

If \mathcal{Y} is a subset of a poset \mathcal{X} , then we write $\mathcal{Y}\downarrow = \{x \in \mathcal{X} \mid \exists y \in \mathcal{Y}, x \leq y\}$ for the order ideal of \mathcal{X} generated by \mathcal{Y} .

Recall that a *partial function* from X to Y is a function from a subset of X to a subset of Y . We will generalise the notation $f: X \rightarrow Y$ to denote a partial function from X to Y .

Definition 10.1 (Action.). Suppose that S is an inverse semigroup, that \mathcal{Y} is a poset (we view an unordered set as a poset in which all elements are incomparable when needed), and that $\alpha: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ is a partial function. If $s \in S$, then we define $\underline{s}_\alpha: \mathcal{Y} \rightarrow \mathcal{Y}$ to be the partial function defined by $(y)\underline{s}_\alpha = (y, s)\alpha$. We write \underline{s}_α rather than s_α , to avoid the having to write parentheses, for example, we write \underline{st}_α instead of $(st)_\alpha$. We say that α is an *action of S on \mathcal{Y}* if the following hold for all $s, t \in S$:

- (1) the partial function \underline{s}_α is an order isomorphism between subsets of \mathcal{Y} ;
- (2) $\underline{st}_\alpha = \underline{s}_\alpha \underline{t}_\alpha$ and $\underline{s}_\alpha^{-1} = \underline{s}^{-1}_\alpha$.

Definition 10.2 (Praction.). Suppose that S is an inverse semigroup, that \mathcal{Y} is a poset, and that $\alpha: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ is a partial function. If $s \in S^1$, then we define $\underline{s}_q: \mathcal{Y} \rightarrow \mathcal{Y}$ to be the partial function defined by $(y)\underline{s}_q = (y, s)q$ (using the identity function if s is the adjoined identity in S^1). We say that q is a *praction of S on \mathcal{Y}* if the following hold for all $s, t, u \in S^1$:

- (1) the partial function \underline{s}_q is an order isomorphism between subsets of \mathcal{Y} ;
- (2) if $s \leq t$, then $\underline{s}_q \subseteq \underline{t}_q$;
- (3) if $(x, y) \in \underline{s}_q$ and $(y, z) \in \underline{t}_q$, then $(x, z) \in \underline{st}_q$, and if $(x, y) \in \underline{s}_q$ then $(y, x) \in \underline{s}^{-1}_q$;
- (4) $\text{dom}(\underline{s}_q)\downarrow = \text{dom}(\underline{s}_q)$;
- (5) for all $y \in \mathcal{Y}$, there is $e \in E(S)$ such that $y \in \text{dom}(\underline{e}_q)$.

Lemma 10.3. *Suppose that S is an inverse semigroup, that \mathcal{Y} is a poset, and that $q: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ is a praction. If $s, t \in S^1$ and $e \in E(S)$, then the following hold:*

- (6) $\underline{s}_q \underline{t}_q \subseteq \underline{st}_q$;
- (7) *If $y, z \in \mathcal{Y}$, $s, t \in S$, and any two of $(x, y) \in \underline{s}_q$, $(y, z) \in \underline{t}_q$, $(x, z) \in \underline{st}_q$ hold, then so does the third;*
- (8) *If $e \in E(S)$ is an idempotent, then \underline{e}_q is a partial identity function.*

Proof. (6) This is an immediate consequence of (3).

(7) If $y, z \in \mathcal{Y}$, $s, t \in S$, $(x, y) \in \underline{s}_q$, and $(y, z) \in \underline{t}_q$, then we have $(x, z) \in \underline{st}_q$ by (3). If $(x, y) \in \underline{s}_q$ and $(x, z) \in \underline{st}_q$, then by (3), $(y, x) \in \underline{s}^{-1}_q$ and so $(y, z) \in \underline{s}^{-1}_q \underline{st}_q$. Thus by (2), $(y, z) \in \underline{t}_q$. If $(y, z) \in \underline{t}_q$ and $(x, z) \in \underline{st}_q$, then by (3), $(z, y) \in \underline{t}^{-1}_q$ and so $(x, y) \in \underline{st} \underline{t}^{-1}_q$. Thus by (2), $(x, y) \in \underline{s}_q$.

(8) From (6), we have that $\underline{e}_q \underline{e}_q \subseteq \underline{e}_q$, and so $(x)\underline{e}_q = x$ for all $x \in \text{im}(\underline{e}_q)$. By (3), $\text{dom } \underline{e}_q = \text{im}(\underline{e}^{-1}_q) = \text{im}(\underline{e}_q)$, and so \underline{e}_q is a partial identity function. \square

If $q: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ in Definition 10.2 is an inverse semigroup action, then q satisfies Definition 10.2(1), (2), and (3) and all conditions in Lemma 10.3.

Example 10.4. Suppose that $\min \mathbb{N} = 0$. We define a praction q of the additive group \mathbb{Z} on the set $\mathcal{Y} := -\mathbb{N}$ as follows:

$$\text{dom}(q) = \{(n, z) \in (-\mathbb{N}) \times \mathbb{Z} \mid n + z \in (-\mathbb{N})\}, \quad \text{and} \quad (n, z)q = n + z.$$

In this example, if $z \in \mathbb{Z}$, then $(n)\underline{z}_q = n + z$ and $\text{dom}(\underline{z}_q) = \{x \in \mathbb{Z} \mid x \leq -z\}$. Since we are using additive notation for \mathbb{Z} , the conditions in Definition 10.2 become additive; for example, (3) becomes “if $(x, y) \in \underline{s}_q$ and $(y, z) \in \underline{t}_q$, then $(x, z) \in \underline{s+t}_q$ ”. It is routine to verify that satisfies Definition 10.2. However, q is clearly not an action in the usual sense, however it naturally extends to one.

The main result in this section is the following lemma, which, roughly speaking, states that every preaction can be extended to an inverse semigroup action, albeit on a larger set.

Theorem 10.5. *Let \mathcal{Y} be a poset and let S be an inverse semigroup. If $q: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ is a preaction, then there is a poset $\mathcal{X}_q \supseteq \mathcal{Y}$ and an action (by partial order isomorphisms) $\alpha_q: \mathcal{X}_q \times S \rightarrow \mathcal{X}_q$ such that*

- (1) \mathcal{Y} is an order ideal of \mathcal{X}_q ;
- (2) the restriction of α_q to $(\mathcal{Y} \times S) \cap (\mathcal{Y})\alpha_q^{-1}$ equals q .
- (3) if $a, b \in \mathcal{X}_q$, then $a \leq b$ if and only if there is $s \in S^1$ such that $(a, s)\alpha_q, (b, s)\alpha_q \in \mathcal{Y}$ and $(a, s)\alpha_q \leq (b, s)\alpha_q$;

Proof. Roughly speaking, we define a set \mathcal{X}'_q to consist of the pairs in $\mathcal{Y} \times S$ we want to lie in the domain of α_q . In particular, if we can act on a point using an element s via α_q , we should be able to act on it with every left divisor of s first in order for composition to work properly. The proof has the following steps: we define \mathcal{X}'_q ; then we equip \mathcal{X}'_q with an action α'_q and a preorder \preceq ; and show that α'_q preserves \preceq . The poset \mathcal{X}_q is then defined to be the quotient of \mathcal{X}'_q by its strongly connected components and α_q is the action on \mathcal{X}_q induced by this quotient.

We define:

$$\mathcal{X}'_q = \{(y, s) \in \mathcal{Y} \times S \mid \text{there exists } s' \in sS \text{ with } (y, s') \in \text{dom}(q)\}$$

and $\alpha'_q: \mathcal{X}'_q \times S \rightarrow \mathcal{X}'_q$ by

$$((y, s), t)\alpha'_q = (y, st)$$

if and only if $(y, s) \in \mathcal{X}'_q$ satisfies $s \in St^{-1}$ and $(y, st) \in \mathcal{X}'_q$. As we did in Definition 10.2, if $t \in S$, then we define $\underline{t}\alpha'_q: \mathcal{X}'_q \rightarrow \mathcal{X}'_q$ by $(y, s)\underline{t}\alpha'_q = ((y, s), t)\alpha'_q$ for all $(y, s) \in \mathcal{X}'_q$. With this notation

$$\text{dom}(\underline{t}\alpha'_q) = \{(y, s) \in \mathcal{X}'_q \mid s \in St^{-1} \text{ and } (y, st) \in \mathcal{X}'_q\}.$$

To show that α'_q is an inverse semigroup action, it suffices to verify the domains:

$$\begin{aligned} \text{dom}(\underline{s}\alpha'_q) \cap (\text{dom}(\underline{t}\alpha'_q), s^{-1})\alpha'_q &= \{(y, v) \in \mathcal{X}'_q \mid v \in Ss^{-1}, (y, vs) \in \mathcal{X}'_q, v \in St^{-1}s^{-1} \text{ and } (y, vst) \in \mathcal{X}'_q\} \\ &= \{(y, v) \in \mathcal{X}'_q \mid v \in St^{-1}s^{-1} \text{ and } (y, vst) \in \mathcal{X}'_q\} \\ &= \text{dom}(\underline{st}\alpha'_q). \end{aligned}$$

Claim 10.6. If we define \preceq on \mathcal{X}'_q by $(y_1, s_1) \preceq (y_2, s_2)$ if there exists $s_3 \in S^1$ with

$$(y_1, s_1s_3), (y_2, s_2s_3) \in \text{dom}(q) \text{ and } (y_1, s_1s_3)q \leq (y_2, s_2s_3)q,$$

then \preceq is a preorder.

Proof. By the definition of \mathcal{X}'_q , \preceq is reflexive. It remains to show that \preceq is transitive. Suppose that $(y_1, s_1), (y_2, s_2), (y_3, s_3) \in \mathcal{X}'_q$ and there are $s_4, s_5 \in S$ such that

$$(5) \quad (y_1, s_1s_4), (y_2, s_2s_4) \in \text{dom}(q) \quad \text{and} \quad (y_1, s_1s_4)q \leq (y_2, s_2s_4)q$$

$$(6) \quad (y_2, s_2s_5), (y_3, s_3s_5) \in \text{dom}(q) \quad \text{and} \quad (y_2, s_2s_5)q \leq (y_3, s_3s_5)q.$$

It suffices to show that

$$(y_1, s_1s_5), (y_3, s_3s_5) \in \text{dom}(q) \text{ and } (y_1, s_1s_5)q \leq (y_3, s_3s_5)q.$$

By (6), it is thus sufficient to show that

$$(y_1, s_1 s_5) \in \text{dom}(q) \text{ and } (y_1, s_1 s_5)q \leq (y_2, s_2 s_5)q.$$

As $(y_2, s_2 s_4) \in \text{dom}(q)$ and $(y_2, s_2 s_5) \in \text{dom}(q)$, it follows from Definition 10.2(3) that $(y_2, s_2 s_4)q \in \text{dom}(\underline{s_4^{-1} s_5}_q)$ and hence by Definition 10.2(2) and Lemma 10.3(6):

$$((y_2, s_2 s_4)q, \underline{s_4^{-1} s_5}_q) = (y_2, s_2 s_4 s_4^{-1} s_5)q = (y_2, s_2 s_5)q.$$

Moreover as the domain of $\underline{s_4^{-1} s_5}_q$ is an order ideal by Definition 10.2(4), it follows that $((y_1, s_1 s_4)q, \underline{s_4^{-1} s_5}_q) \in \text{dom}(q)$ and hence

$$((y_1, s_1 s_4)q, \underline{s_4^{-1} s_5}_q) = (y_1, s_1 s_5)q.$$

Since $(y_1, s_1 s_4)q \leq (y_2, s_2 s_4)q$ and $\underline{s_4^{-1} s_5}_q$ is \leq -preserving (by Definition 10.2(1)), it follows that

$$(y_1, s_1 s_5)q = ((y_1, s_1 s_4)q, \underline{s_4^{-1} s_5}_q) \leq ((y_2, s_2 s_4)q, \underline{s_4^{-1} s_5}_q) = (y_2, s_2 s_5)q,$$

as required. \square

We now show that the action α'_q is \preceq -preserving. If $t \in S$ and $(y_1, s_1), (y_2, s_2) \in \text{dom}(\underline{t}_{\alpha'_q})$ and there is $s_3 \in S^1$ with $(y_1, s_1 s_3)q \leq (y_2, s_2 s_3)q$, then $s_1, s_2 \in St^{-1}$. Hence

$$(y_1, s_1 t t^{-1} s_3)q = (y_1, s_1 s_3)q \leq (y_2, s_2 s_3)q = (y_2, s_2 t t^{-1} s_3)q$$

so $((y_1, s_1), t)\alpha'_q = (y_1, s_1 t) \preceq (y_2, s_2 t) = ((y_2, s_2), t)\alpha'_q$, and the action α'_q is \preceq -preserving.

We write $(y_1, s_1) \sim (y_2, s_2)$ if $(y_1, s_1) \preceq (y_2, s_2)$ and $(y_2, s_2) \preceq (y_1, s_1)$, and denote by $[(y, s)]_{\sim}$ the \sim -equivalence class of $(y, s) \in \mathcal{X}'_q$. Let \mathcal{X}_q be the quotient of \mathcal{X}'_q by the equivalence relation \sim . Then \mathcal{X}_q is partially ordered by $[(y_1, s_1)]_{\sim} \leq [(y_2, s_2)]_{\sim}$ if $(y_1, s_1) \preceq (y_2, s_2)$.

We define $\phi: \mathcal{Y} \rightarrow \mathcal{X}_q$ by

$$(7) \quad (y)\phi = [(z, u)]_{\sim} \text{ if } (y)q^{-1} \subseteq [(z, u)]_{\sim}.$$

We will show that ϕ is a well-defined order-embedding and $\text{dom}(\phi) = \mathcal{Y}$.

If $(y)\phi = [(z, u)]_{\sim}$ and $(y)\phi = [(z', u')]_{\sim}$, then without loss of generality $(z, u)q = y = (z', u')q$ and so $(z, u) \sim (z', u')$, meaning that ϕ is well-defined. We show that there is $(z, u) \in \mathcal{X}'_q$ such that $(z, u)q = y$. By Definition 10.2(5) we can pick $z = y$ and $u = e$ for some idempotent e such that $y \in \text{dom}(e_q)$. This implies that the domain of ϕ is \mathcal{Y} . If $y_1, y_2 \in \mathcal{Y}$, and $(z_1, u_1) \in (y_1)q^{-1}$ and $(z_2, u_2) \in (y_2)q^{-1}$ for some z_1, z_2, u_1, u_2 , then

$$\begin{aligned} y_1 \leq y_2 &\iff (z_1, u_1)q \leq (z_2, u_2)q \\ &\implies (z_1, u_1) \preceq (z_2, u_2) \\ &\iff [(z_1, u_1)]_{\sim} \leq [(z_2, u_2)]_{\sim} \\ &\iff y_1 \phi \leq y_2 \phi. \end{aligned}$$

Thus to conclude both that ϕ is injective and order-preserving it suffices to show that $(z_1, u_1) \preceq (z_2, u_2)$ implies that $(z_1, u_1)q \leq (z_2, u_2)q$. By the definition of \preceq , there exists $s_3 \in S$ such that $(z_1, u_1 s_3)q \leq (z_2, u_2 s_3)q$. By assumption, for $i \in \{1, 2\}$, $z_i \in \text{dom}(\underline{u_i}_q)$ and $z_i \in \text{dom}(\underline{u_i s_3}_q)$. In other words, $(z_i, (z_i)\underline{u_i}_q) \in \underline{u_i}_q$ and $(z_i, (z_i)\underline{u_i s_3}_q) \in \underline{u_i s_3}_q$. Then Lemma 10.3(7) tells us $((z_i)\underline{u_i}_q, (z_i)\underline{u_i s_3}_q) \in \underline{s_3}_q$. In particular, $z_i \in \text{dom}(\underline{u_i s_3}_q)$. Hence

$$\begin{aligned} (z_1, u_1)q &= (z_1)\underline{u_1}_q && \text{definition of } \underline{u_1}_q \\ &= (z_1)\underline{u_1}_q \underline{s_3}_q \underline{s_3}_q^{-1} && \underline{s_3}_q \underline{s_3}_q^{-1} \text{ is the identity on } \text{dom}(\underline{s_3}_q) \\ &= (z_1)\underline{u_1 s_3}_q \underline{s_3}_q^{-1} \\ &= (z_1, u_1 s_3)q \cdot \underline{s_3}_q^{-1} \end{aligned}$$

$$\begin{aligned} &\leq (z_2, u_2 s_3)q \cdot \underline{s_3}_q^{-1} && \underline{s_3}_q \text{ is a partial order-isomorphism of } \mathcal{Y} \\ &= (z_2, u_2)q. \end{aligned}$$

Claim 10.7. If $(y_1, s_1) \in \text{dom}(q)$, and $(y_2, s_2) \preceq (y_1, s_1)$, then $(y_2, s_2) \in \text{dom}(q)$.

Proof. Let $(y_1, s_1) \in \text{dom}(q)$ and $(y_2, s_2) \preceq (y_1, s_1)$. Then by the definition of \preceq there exists $s_3 \in S^1$ such that

$$(y_1, s_1 s_3), (y_2, s_2 s_3) \in \text{dom}(q), \quad (y_2, s_2 s_3)q \leq (y_1, s_1 s_3)q.$$

Since $(y_1, (y_1)\underline{s_1}_q) \in \underline{s_1}_q$ and $(y_1, (y_1)\underline{s_1 s_3}_q) \in \underline{s_1 s_3}_q$, it follows by (3) that $((y_1)\underline{s_1}_q, (y_1)\underline{s_1 s_3}_q) \in \underline{s_3}_q$ $(y_1)\underline{s_1}_q \underline{s_3}_q = (y_1)\underline{s_1 s_3}_q$; see Fig. 3. In particular, $((y_1)\underline{s_1}_q, s_3) \in \text{dom}(q)$. Moreover $((y_1)\underline{s_1}_q, s_3)q = (y_1, s_1 s_3)q \geq (y_2, s_2 s_3)q$. By (3), $\text{im}(\underline{s_3}_q) = \text{dom}(\underline{s_3}_q^{-1})$ which is an order ideal by (4) and so $((y_2, s_2 s_3)q) \in \text{dom}(\underline{s_3}_q^{-1})$. Applying (3) to $(y_2, (y_2, s_2 s_3)q) \in \underline{s_2 s_3}_q$ and $((y_2, s_2 s_3)q, ((y_2, s_2 s_3)q, s_3^{-1})q) \in \underline{s_3}_q^{-1}$, we obtain $(y_2, ((y_2, s_2 s_3)q, s_3^{-1})q) \in \underline{s_2 s_3 s_3^{-1}}_q$.

Thus $(y_2, s_2 s_3 s_3^{-1}) \in \text{dom}(q)$ and, by (2), $\underline{s_2 s_3 s_3^{-1}}_q \subseteq \underline{s_2}_q$. In particular, $(y_2, s_2 s_3 s_3^{-1})q = (y_2, s_2)q$ and so $(y_2, s_2) \in \text{dom}(q)$. \square

A particular case of Claim 10.7 is the following.

Claim 10.8. If $(y, s) \in \text{dom}(q)$, then $[(y, s)]_\sim \subseteq \text{dom}(q)$ and $(y, s)q\phi = [(y, s)]_\sim = (y, s)qq^{-1}$. In other words, $[(y, s)]_\sim \phi^{-1} = (z, t)q$ for any $(z, t) \in [(y, s)]_\sim$.

The next claim establishes part (1) of the theorem.

Claim 10.9. If $\phi: \mathcal{Y} \rightarrow \mathcal{X}_q$ is defined as in (7), then $(\mathcal{Y})\phi$ is an order-ideal of \mathcal{X}_q .

Proof. Let $[(y_2, s_2)]_\sim \leq [(y_1, s_1)]_\sim \in (\mathcal{Y})\phi$. Then by the definition of ϕ , there is $(y, s) \in \text{dom}(q)$ with $[(y_1, s_1)]_\sim = [(y, s)]_\sim$. Then by Claim 10.8, $(y_1, s_1) \in \text{dom}(q)$. The assumption that $[(y_2, s_2)]_\sim \leq [(y_1, s_1)]_\sim$ implies that $(y_2, s_2) \preceq (y_1, s_1)$. Hence by Claim 10.7, $(y_2, s_2) \in \text{dom}(q)$ and so $[(y_2, s_2)]_\sim = ((y_2, s_2)q)\phi \in (\mathcal{Y})\phi$ as required. \square

Define the partial function $\alpha_q: \mathcal{X}_q \times S \rightarrow \mathcal{X}_q$ by

$$([(y, u)]_\sim, s)\alpha_q = [((y, u), s)\alpha'_q]_\sim = [(y, us)]_\sim$$

where

$$\begin{aligned} \text{dom}(\alpha_q) &= \{([(y, u)]_\sim, s) \in \mathcal{X}_q \times S \mid (y, u) \in \text{dom}(s\alpha'_q)\} \\ &= \{([(y, u)]_\sim, s) \in \mathcal{X}_q \times S \mid u \in Ss^{-1} \text{ there is } v \in S \text{ with } (y, usv) \in \text{dom}(q)\}. \end{aligned}$$

It remains to check that the induced action of S on the copy $(\mathcal{Y})\phi$ of \mathcal{Y} contained in \mathcal{X}_q is isomorphic to the action of S on \mathcal{Y} . To this end we define $\phi \oplus \text{id}_S: \mathcal{Y} \times S \rightarrow \mathcal{X}_q \times S$ by

$$(y, s)\phi \oplus \text{id}_S = ((y)\phi, s),$$

and

$$\mathcal{Z} = ((\mathcal{Y})\phi \times S) \cap ((\mathcal{Y})\phi)\alpha_q^{-1}.$$

That is we will show:

$$(\phi \oplus \text{id}_S) \circ \alpha_q|_{\mathcal{Z}} \circ \phi^{-1} = q.$$

We define the function on the left hand side by Q . Suppose that $(y, s) \in \text{dom}(Q)$. It follows that $y \in \text{dom}(\phi)$. From Definition 10.2(5), there exists $e_y \in E(S)$ be such that $(y, e_y) \in \text{dom}(q)$. Then

$$\begin{aligned} (y, s)Q &= (y, s)(\phi \oplus \text{id}_S) \circ \alpha_q|_{\mathcal{Z}} \circ \phi^{-1} \\ &= [([(y, e_y)]_\sim, s)\alpha_q|_{\mathcal{Z}} \circ \phi^{-1} \end{aligned}$$

$$\begin{array}{ccc}
& & y_1 \\
& \swarrow \xrightarrow{s_1 s_3 q} & \downarrow \xrightarrow{s_1 q} \\
(y_1) \underline{s_1 s_3 q} & \xleftarrow{\text{---} \underline{s_3 q} \text{---}} & (y_1) \underline{s_1 q}
\end{array}$$

∨

$$\begin{array}{ccc}
& & y_2 \\
& \swarrow \xrightarrow{s_2 s_3 q} & \downarrow \xrightarrow{s_2} \\
(y_2) \underline{(s_2 s_3) q} & \xleftarrow{\text{---} \underline{s_3^{-1} q} \text{---}} & (y_2) \underline{s_2 s_3 q s_3^{-1} q}
\end{array}$$

FIGURE 3. Diagram demonstrating the arguments in the second claim within the proof of Theorem 10.5. The dashed lines indicate an application of (3) starting with the arrow labelled $\underline{s_3 q}$ and proceeding anti-clockwise.

$$\begin{aligned}
&= ((y, e_y s)]_{\sim} \phi^{-1} && \text{by Claim 10.8} \\
&= (y, e_y s) q \\
&= (y, s) q && \text{by Definition 10.2(2).}
\end{aligned}$$

Thus $q|_{\text{dom}(Q)} = Q$. If $(y, s) \in \text{dom}(q)$, then, by the sequence of equalities above (in reverse order), $(y, s) \in \text{dom}(Q)$. Hence $q = Q$. The map α_q in the statement of the lemma can now be chosen by redefining $\mathcal{X}_q := (\mathcal{X}_q \setminus \text{im}(\phi)) \cup \mathcal{Y}$ and the map α_q by

$$(y, s) \alpha_q = \begin{cases} (y, s) \alpha_q & y \notin \mathcal{Y} \\ ((y) \phi, s) \alpha_q & y \in \mathcal{Y}. \end{cases}$$

As we previously showed that $Q = q$, it follows that now $q = \alpha_q|_{(\mathcal{Y} \times S) \cap (\mathcal{Y}) \alpha_q^{-1}}$ and so part (2) of the theorem holds.

Since part (3) of the theorem implies that α_q acts by partial order-isomorphisms on \mathcal{X}_q , the proof will be concluded by showing that part (3) holds. Suppose that $[(y_1, s_1)]_{\sim}, [(y_2, s_2)]_{\sim} \in \mathcal{X}_q$. We must show that $[(y_1, s_1)]_{\sim} \leq [(y_2, s_2)]_{\sim}$ if and only if there exists $t \in S^1$ such that $([(y_1, s_1)]_{\sim}, t) \alpha_q \leq ((y_2, s_2)]_{\sim}, t) \alpha_q \in (\mathcal{Y}) \phi$.

By the definition \leq , $[(y_1, s_1)]_{\sim} \leq [(y_2, s_2)]_{\sim}$ if and only if $(y_1, s_1) \preceq (y_2, s_2)$ if and only if (from the definition of \preceq) there exists $t \in S^1$ such that $(y_1, s_1 t), (y_2, s_2 t) \in \text{dom}(q)$ and $(y_1, s_1 t) q \leq (y_2, s_2 t) q \in \mathcal{Y}$. This holds if and only if there exists $t \in S^1$ such that $([(y_1, s_1)]_{\sim}, t) \alpha_q = [(y_1, s_1 t)]_{\sim} \in (\mathcal{Y}) \phi$, $([(y_2, s_2)]_{\sim}, t) \alpha_q = [(y_2, s_2 t)]_{\sim} \in (\mathcal{Y}) \phi$, and $([(y_1, s_1)]_{\sim}, t) \alpha_q \leq ((y_2, s_2)]_{\sim}, t) \alpha_q$, as required. Hence part (3) of the theorem holds, and the proof is complete at last. \square

The next corollary follows immediately from Theorem 10.5 and essentially states that preactions are precisely certain restrictions of certain actions.

Corollary 10.10. *If S is an inverse monoid, \mathcal{Y} is a poset, and $q: \mathcal{Y} \times S \rightarrow \mathcal{Y}$ is a partial function, then q is a preaction if and only if q satisfies Definition 10.2(4) and (5) and there is an action α_q of S on a poset $\mathcal{X} \supseteq \mathcal{Y}$ such that $q = \alpha_q|_{(\mathcal{Y} \times S) \cap (\mathcal{Y}) \alpha_q^{-1}}$.*

11. THE Q -THEOREM

The goal of this section is to introduce a means of defining an inverse semigroup in terms of a natural action of an E -disjunctive semigroup on a poset, and also to show that every inverse

semigroup can be defined this way. This construction generalises that of McAlister triples for E -unitary inverse semigroups [27]. This theorem was already known and proved in [32]. The authors of the present paper only discovered [32] at a late stage of the preparation of this paper, and proved the characterisation independently. The theorem and its proof is included for the sake of completeness.

Recall that if \mathcal{Y} is a subset of a poset \mathcal{X} , then we write $\mathcal{Y} \downarrow = \{x \in \mathcal{X} \mid \exists y \in \mathcal{Y}, x \leq y\}$. If $\mathcal{Y} = \mathcal{Y} \downarrow$, then we say that \mathcal{Y} is an *order ideal* in \mathcal{X} .

Definition 11.1 (*Q -semigroup*). Suppose that T is an inverse semigroup acting on a poset \mathcal{X} by partial order isomorphisms and \mathcal{Y} is a meet subsemilattice and order ideal of \mathcal{X} such that the following conditions hold:

- (1) For all $t \in T$, $\text{dom } t = (\text{dom } t) \downarrow$;
- (2) For all $y \in \mathcal{Y}$ there exists $t \in T$ with

$$\text{dom}(t|_{\mathcal{Y}}) = \bigcap \{\text{dom}(s|_{\mathcal{Y}}) \mid s \in T \text{ such that } y \in \text{dom}(s|_{\mathcal{Y}})\}.$$

We denote the set $\text{dom}(t|_{\mathcal{Y}})$ by $\delta(y)$.

- (3) For all $x \in \mathcal{X}$, there is $t \in T$ such that $(x)t \in \mathcal{Y}$.

Then we define $Q(T, \mathcal{Y}, \mathcal{X})$ to be

$$Q(T, \mathcal{Y}, \mathcal{X}) = \{(y, t) \in \mathcal{Y} \times T \mid \text{dom}(t) = \delta(y), (y)t \in \mathcal{Y}\},$$

with multiplication defined by $(y_1, t_1) \cdot (y_2, t_2) = (((y_1)t_1 \wedge y_2)t_1^{-1}, t_1 t_2)$.

We note that part (2) in Definition 11.1, can be reformulated as:

- (2)* for all $y \in \mathcal{Y}$, there exists $t \in T$ such that $\text{dom}(t|_{\mathcal{Y}})$ contains y and is the least possible with respect to containment.

Note that $\delta: \mathcal{Y} \rightarrow \mathcal{P}(\mathcal{Y})$ as defined in Definition 11.1 is a homomorphism (of semilattices) where the operation on $\mathcal{P}(\mathcal{Y})$ is \cap .

If $(G, \mathcal{Y}, \mathcal{X})$ is a McAlister triple, then it will turn out that the inverse semigroups $P(G, \mathcal{Y}, \mathcal{X})$ and $Q(G, \mathcal{Y}, \mathcal{X})$ coincide.

The main theorems of this section are the following; which we prove in Section 11.1 and Section 11.2, respectively.

Theorem 11.2. *If $(T, \mathcal{Y}, \mathcal{X})$ satisfy the conditions in Definition 11.1, then $Q(T, \mathcal{Y}, \mathcal{X})$ is an inverse semigroup.*

A converse of Theorem 11.2 also holds.

Theorem 11.3. *Every inverse semigroup S is isomorphic to some $Q(T, \mathcal{Y}, \mathcal{X})$ from Definition 11.1, where T is the maximum E -disjunctive homomorphic image of S , and \mathcal{Y} is the semilattice of idempotents of S .*

11.1. Q -semigroups are inverse semigroups. In this section we give the proof of Theorem 11.2.

Proof of Theorem 11.2. Let $Q = Q(T, \mathcal{Y}, \mathcal{X})$. We begin by showing that the multiplication defined in Definition 11.1 is well-defined. Let $(y_1, t_1), (y_2, t_2) \in Q$. Then we must show $((y_1)t_1 \wedge y_2)t_1^{-1}, t_1 t_2 \in Q$; that is $\text{dom}(t_1 t_2) = \delta(((y_1)t_1 \wedge y_2)t_1^{-1})$ and $((y_1)t_1 \wedge y_2)t_1^{-1} t_1 t_2 \in \mathcal{Y}$. Since $(y_1)t_1 \wedge y_2 \leq y_1 t_1 \in \text{dom}(t_1^{-1})$. Since $\text{dom}(t_1^{-1})$ is an order ideal, it follows that $(y_1)t_1 \wedge y_2 \in \text{dom}(t_1^{-1})$. Similarly, $(y_1)t_1 \wedge y_2 \leq y_2 \in \text{dom}(t_2)$ and so $(y_1)t_1 \wedge y_2 \in \text{dom}(t_2)$. Hence $((y_1)t_1 \wedge y_2)t_1^{-1} t_1 t_2 = ((y_1)t_1 \wedge y_2)t_2 \leq (y_2)t_2 \in \mathcal{Y}$.

It remains to show that

$$\delta(((y_1)t_1 \wedge y_2)t_1^{-1}) = \text{dom}(t_1 t_2).$$

First note that for all $t \in T$ and $y \in \text{dom}(t)$ we have

$$(\delta(y))t = \left(\bigcap_{\substack{s \in T \\ y \in \text{dom}(s)}} \text{dom}(s) \right) t \subseteq \bigcap_{\substack{s \in T \\ y \in \text{dom}(s)}} \text{dom}(s)t = \bigcap_{\substack{s \in T \\ y \in \text{dom}(s)}} \text{dom}(t^{-1}s) \subseteq \bigcap_{\substack{s \in T \\ yt \in \text{dom}(s)}} \text{dom}(s) = \delta((y)t).$$

Similarly, $\delta((y)t)t^{-1} \subseteq \delta(y)$ and $\delta((y)t) = \delta((y)t)t^{-1}t \subseteq (\delta(y))t$. It follows that

$$(8) \quad (\delta(y))t = \delta(yt).$$

Hence

$$\begin{aligned} \delta(((y_1)t_1 \wedge y_2)t_1^{-1}) &= \delta((y_1)t_1 \wedge y_2)t_1^{-1} \\ &= (\delta((y_1)t_1) \cap \delta(y_2))t_1^{-1} \\ &= (\delta(y_1)t_1 \cap \delta(y_2))t_1^{-1} \\ &= (\text{dom}(t_1)t_1 \cap \text{dom}(t_2))t_1^{-1} \\ &= \text{dom}(t_1) \cap \text{dom}(t_2)t_1^{-1} \\ &= \text{dom}(t_1 t_2). \end{aligned}$$

Next we prove that Q is a semigroup. It suffices to show that the multiplication is associative. Let $(y_1, t_1), (y_2, t_2), (y_3, t_3) \in Q$. Let

$$u = (((y_1)t_1 \wedge y_2)t_2 \wedge y_3)t_2^{-1} \text{ and } v = (y_1)t_1 \wedge ((y_2)t_2 \wedge y_3)t_2^{-1}.$$

We must first show that $u = v$, which we do by showing $u \leq v$ and $v \leq u$. To this end

$$u = (((y_1)t_1 \wedge y_2)t_2 \wedge y_3)t_2^{-1} \leq ((y_2)t_2 \wedge y_3)t_2^{-1}$$

and

$$u = (((y_1)t_1 \wedge y_2)t_2 \wedge y_3)t_2^{-1} \leq ((y_1)t_1 \wedge y_2)t_2 t_2^{-1} = (y_1)t_1 \wedge y_2 \leq (y_1)t_1.$$

Thus $u \leq v$. To show $v \leq u$, it is sufficient to show that $(v)t_2 \leq (u)t_2$. We have

$$(v)t_2 = ((y_1)t_1 \wedge ((y_2)t_2 \wedge y_3)t_2^{-1})t_2 \leq ((y_1)t_1 \wedge (y_2)t_2 t_2^{-1})t_2 = ((y_1)t_1 \wedge y_2)t_2$$

and

$$(v)t_2 = ((y_1)t_1 \wedge ((y_2)t_2 \wedge y_3)t_2^{-1})t_2 \leq ((y_2)t_2 \wedge y_3)t_2^{-1}t_2 = (y_2)t_2 \wedge y_3.$$

Hence $v \leq u$ and $u = v$, as required. From the definition of the multiplication,

$$\begin{aligned} ((y_1, t_1) \cdot (y_2, t_2)) \cdot (y_3, t_3) &= (((y_1)t_1 \wedge y_2)t_1^{-1}, t_1 t_2) \cdot (y_3, t_3) \\ &= (((y_1)t_1 \wedge y_2)t_1^{-1} t_1 t_2 \wedge y_3)t_2^{-1} t_1^{-1}, t_1 t_2 t_3) \\ &= (((y_1)t_1 \wedge y_2)t_2 \wedge y_3)t_2^{-1} t_1^{-1}, t_1 t_2 t_3) \quad ((y_1)t_1 \wedge y_2)t_1^{-1} t_1 = (y_1)t_1 \wedge y_2 \\ &= ((u)t_1^{-1}, t_1 t_2 t_3) \\ &= ((v)t_1^{-1}, t_1 t_2 t_3) \\ &= (((y_1)t_1 \wedge ((y_2)t_2 \wedge y_3)t_2^{-1})t_1^{-1}, t_1 t_2 t_3) \\ &= (y_1, t_1) \cdot ((y_2)t_2 \wedge y_3)t_2^{-1}, t_2 t_3) \\ &= (y_1, t_1) \cdot ((y_2, t_2) \cdot (y_3, t_3)). \end{aligned}$$

We conclude the proof by showing that Q is an inverse semigroup. We first show that Q is regular. Let $(y, t) \in Q$. We will show that $(yt, t^{-1}) \in Q$, and that this is an inverse for $(y, t) \in Q$. Since $(\delta(y))t = \delta(yt)$ by (8),

$$\text{dom } t^{-1} = (\text{dom } t)t = (\delta(y))t = \delta(yt).$$

Therefore $(yt, t^{-1}) \in Q$. In addition,

$$\begin{aligned} (y, t) \cdot ((y)t, t^{-1}) \cdot (y, t) &= (((y)t \wedge (y)t)t^{-1}, tt^{-1}) \cdot (y, t) \\ &= ((y)tt^{-1}, tt^{-1}) \cdot (y, t) \\ &= (y, tt^{-1}) \cdot (y, t) \\ &= (((y)tt^{-1} \wedge y)tt^{-1}, tt^{-1}t) \\ &= (y, t), \end{aligned}$$

and so Q is regular. It now suffices to show that the idempotents commute. If $(y, t) \in Q$ is an idempotent, then $t \in E(T)$. Conversely if $e \in E(T)$ then $(y, e) \cdot (y, e) = (((y)e \wedge y)e^{-1}, e^2) = (y, e)$. So $E(Q) = \{(y, e) \in Q \mid y \in \mathcal{Y}, e \in E(T)\}$. These elements commute:

$$(y_1, e_1) \cdot (y_2, e_2) = (((y_1)e_1 \wedge y_2)e_1^{-1}, e_1e_2) = (y_1 \wedge y_2, e_1e_2) = (y_2, e_2) \cdot (y_1, e_1). \quad \square$$

11.2. The proof of Theorem 11.3. For the remainder of this section, we suppose that S is a fixed inverse semigroup. The idea behind the proof of Theorem 11.3 is as follows. The semigroup S has a quotient by an idempotent-pure congruence that is an E -disjunctive inverse semigroup; see Section 9. By Lemma 9.4, an element $s \in S$ is determined by its image in this quotient together with the idempotent ss^{-1} . The set \mathcal{Y} in the definition of $Q(T, \mathcal{Y}, \mathcal{X})$ is the set of idempotents $E(S)$ of S , and T is the quotient of S by its maximum idempotent-pure congruence ρ . We will prove that the function $S \rightarrow Q(T, \mathcal{Y}, \mathcal{X})$ defined by

$$s \mapsto (ss^{-1}, s/\rho)$$

is an isomorphism for the correct choice of \mathcal{X} .

Roughly speaking, in order to capture the multiplication of S in the definition of $Q(T, \mathcal{Y}, \mathcal{X})$, we need to be able to recover the idempotents $(st)(st)^{-1}$ from the idempotents ss^{-1} and tt^{-1} . This is where the action comes in. The action we define comes from the conjugation (inverse semigroup) action of S on $E(S)$ which is defined as follows. We define $\alpha: E(S) \times S \rightarrow E(S)$ by

$$(9) \quad (e, s)\alpha = s^{-1}es \quad \text{and} \quad \text{dom}(\alpha) = \{(e, s) \in E(S) \times S \mid e \leq ss^{-1}\}.$$

It is routine to verify that this is an inverse semigroup action. We also define $\phi_\alpha: S \rightarrow I_{E(S)}$ to be the homomorphism associated to α . In Lemma 11.7 we will show that $T = S/\rho$ has a preaction (Definition 10.2) on $\mathcal{Y} = E(S)$ (the is essentially the same idea as Proposition 2.11). Hence by Theorem 10.5 there will exist an inverse semigroup action of T on a poset \mathcal{X} containing \mathcal{Y} .

Definition 11.4. Let α be the action given in (9). We define a multiplication on the set $\text{dom}(\alpha)$ by

$$(e, s)(f, t) = (((e, s)\alpha \wedge f, s^{-1})\alpha, st).$$

This multiplication is well-defined as $((e, s)\alpha \wedge f, s^{-1})\alpha \leq sfs^{-1} \leq stt^{-1}s^{-1} = (st)(st)^{-1}$. We do not assert that this multiplication is associative.

The natural magma homomorphism

$$(10) \quad \pi: \text{dom}(\alpha) \rightarrow S \text{ is defined by } (e, s)\pi = s.$$

We will show that a subset of $\text{dom}(\alpha)$ with the multiplication given in Definition 11.4 is a semigroup, by showing that the subset is (magma) isomorphic to a semigroup.

Lemma 11.5. *Let $\psi: S \rightarrow \text{dom}(\alpha)$ be the map defined by*

$$(s)\psi = (ss^{-1}, s),$$

Then ψ is an injective magma homomorphism and $\text{im}(\psi) \cong S$.

Proof. Note that ψ is well-defined by the definition of the set $\text{dom}(\alpha)$.

Let $s, t \in S$. Then

$$\begin{aligned} (s)\psi (t)\psi &= (ss^{-1}, s)(tt^{-1}, t) \\ &= ((s^{-1}ss^{-1}s \wedge tt^{-1}, s^{-1})\alpha, st) \\ &= ((s^{-1}s \wedge tt^{-1}, s^{-1})\alpha, st) \\ &= (ss^{-1}stt^{-1}s^{-1}, st) \\ &= (stt^{-1}s^{-1}, st) \\ &= (st)\psi. \end{aligned}$$

Since the homomorphism $\psi \circ \pi$ is the identity map on S , the restriction $\pi|_{\text{im}(\psi)} = \psi^{-1}$ is an isomorphism from a subsemigroup of $\text{dom}(\alpha)$ to S . \square

We have not yet defined the poset \mathcal{X} which we will be using to define our Q -semigroup. However, since the set of elements of the Q -semigroup do not depend on \mathcal{X} , only the multiplication within the Q -semigroup. The next lemma shows that S is contained in the set of elements in the Q -semigroup we are in the process of defining.

Lemma 11.6. *Let ρ be the syntactic congruence on S , and let $\psi_\rho: S \rightarrow E(S) \times S/\rho$ by*

$$(s)\psi_\rho = (ss^{-1}, s/\rho).$$

Then ψ_ρ is injective.

Proof. Let $s, t \in S$ be such that $(s)\psi_\rho = (ss^{-1}, s/\rho) = (tt^{-1}, t/\rho) = (t)\psi_\rho$. In particular, $ss^{-1} = tt^{-1}$ and so $s\mathcal{R}t$. Since the quotient homomorphism from S to S/ρ is idempotent-pure, Lemma 9.4 implies that this homomorphism is injective on the \mathcal{R} -classes of S . Hence $s/\rho = t/\rho$ implies that $s = t$. \square

We define

$$(11) \quad M = \text{im}(\psi_\rho) = \{(ss^{-1}, s/\rho) : s \in S\} \subseteq E(S) \times S/\rho.$$

and define multiplication on M such that $\psi_\rho: S \rightarrow M$ is an isomorphism. If $\psi: S \rightarrow \text{im}(\psi) \subseteq \text{dom}(\alpha)$ is the (semigroup) isomorphism from Lemma 11.5 and $\pi: \text{dom}(\alpha) \rightarrow S$ is from (10), then $\pi|_{\text{im}(\psi)}\psi_\rho = \psi^{-1}\psi_\rho: \text{im}(\psi) \rightarrow M$ is an isomorphism. If $(ss^{-1}, s) \in \text{im}(\psi)$, then

$$(ss^{-1}, s)\psi^{-1}\psi_\rho = (s)\psi_\rho = (ss^{-1}, s/\rho).$$

If $s, t \in S$ and $e, f \in E(S)$ are such that $(e, s/\rho), (f, t/\rho) \in M$, then there is $s_0 \in s/\rho$ such that $(s_0)\psi_\rho = (s_0s_0^{-1}, s_0/\rho) = (e, s/\rho)$. Since

$$\begin{aligned} (e, s/\rho)(f, t/\rho) &= ((e, s)(f, t))\psi^{-1}\psi_\rho \\ &= (((e, s_0)\alpha \wedge f), s_0^{-1})\alpha, st)\psi^{-1}\psi_\rho \\ &= (((e, s_0)\alpha \wedge f), s_0^{-1})\alpha, st/\rho), \end{aligned}$$

it follows that

$$(12) \quad (e, s/\rho)(f, t/\rho) = (((e, s_0)\alpha \wedge f), s_0^{-1})\alpha, st/\rho).$$

We will prove Theorem 11.3 by describing the multiplication of the given inverse semigroup S using only the E -disjunctive inverse semigroup S/ρ , the idempotents $E(S)$, and an action of S/ρ on a poset. Since M is isomorphic to S , the above equation almost does this. The problem is that α is defined in terms of S , and not only in terms of S/ρ and $E(S)$. We will show that S/ρ is sufficient to capture the needed information from this action.

Since the particular choice of representative of the classes in S/ρ is not important later, we denote S/ρ by T so that we may refer to the elements of T rather than choosing a representative for an element of S/ρ .

Lemma 11.7. *If $\alpha: E(S) \times S \rightarrow E(S)$ is the action defined in (9), then the partial function $q: E(S) \times T \rightarrow E(S)$ defined by*

$$(e, t)q = (e, s)\alpha$$

for all $(e, t) \in E(S) \times T$ such that there exists $t' \leq t$ with $s \in t'$ is a preaction. In particular,

$$\text{dom}(q) = \{(e, t) \in E(S) \times S/\rho \mid \exists s \in t' \leq t \text{ with } (e, s) \in \text{dom}(\alpha)\}.$$

Proof. We first show that q is well-defined. Let $s_1 \in t_1 \leq t \in T$, let $s_2 \in t_2 \leq t \in T$, and let $(e, s_1), (f, s_2) \in \text{dom}(\alpha)$. We will show that $(e, s_1)\alpha \leq (f, s_2)\alpha$ if and only if $e \leq f$. This will not only show that q is well-defined (by considering the case when $e = f$) but will also show it satisfies Definition 10.2(1).

Since $s_1 \in t_1 \leq t$ and $s_2 \in t_2 \leq t$, it follows that $(s_1^{-1}s_2/\rho) \leq t^{-1}t$ and $s_1s_2^{-1}/\rho \leq tt^{-1}$. Hence $s_1^{-1}s_2/\rho, s_1s_2^{-1}/\rho \in E(T)$, and so, by Lallement's Lemma, both $s_1^{-1}s_2/\rho$ and $s_1s_2^{-1}/\rho$ contain an idempotent. But ρ is idempotent-pure and so $s_1^{-1}s_2, s_1s_2^{-1} \in E(S)$. Thus $s_1^{-1}s_2$ and $s_1s_2^{-1}$ equal their inverses, that is,

$$(13) \quad s_1^{-1}s_2 = s_2^{-1}s_1 \quad \text{and} \quad s_1s_2^{-1} = s_2s_1^{-1}$$

Similarly $s_1s_2^{-1} = s_2s_1^{-1}$ is an idempotent. If $s \in S$ we define $\underline{s}_\alpha: E(S) \rightarrow E(S)$, by $(g)\underline{s}_\alpha = (g, s)\alpha = s^{-1}gs$ for all $g \in E(S)$, then $\text{dom}(\underline{s}_\alpha)$ is an order ideal. This notation coincides with the notation in Definition 10.2 although we have not shown that α is a preaction. We assumed at the start of the proof that $(e, s_1), (f, s_2) \in \text{dom}(\alpha)$ and so $(e, s_1)\alpha \in \text{im}(\underline{s}_{1\alpha})$ and $e \in \text{dom}(\underline{s}_{2\alpha})$ since $s_1 \leq s_2$ and $e \leq f$. In particular, $(e, s_1)\alpha \in \text{dom}(\underline{s}_{1\alpha}^{-1})$, and so $(e, s_1)\alpha \underline{s}_{1\alpha}^{-1} = (e)\underline{s}_{1\alpha} \underline{s}_{1\alpha}^{-1} = e$. Since $e \in \text{dom}(\underline{s}_{2\alpha})$, it follows that $(e, s_1)\alpha \underline{s}_{1\alpha}^{-1} \in \text{dom}(\underline{s}_{2\alpha})$ and so $(e, s_1)\alpha \in \text{dom}(\underline{s}_{1^{-1}s_2\alpha})$. Since $s_1^{-1}s_2$ is an idempotent, $(s_1^{-1}s_2)\alpha$ acts as the identity on every point in its domain, including $(e, s_1)\alpha$. In other words,

$$(14) \quad (e, s_1)\alpha = (e, s_1s_1^{-1}s_2)\alpha.$$

By definition, $(e, s_1s_1^{-1}s_2)\alpha = (e, s_1s_1^{-1})\alpha \cdot \underline{s}_{2\alpha}$. Since $\underline{s}_{1s_1^{-1}\alpha}$ is the identity $\text{dom}(\underline{s}_{1\alpha})$ and $e \in \text{dom}(\underline{s}_{1\alpha})$, it follows that

$$(15) \quad (e, s_1s_1^{-1}s_2)\alpha = (e)\underline{s}_{1s_1^{-1}\alpha} \underline{s}_{2\alpha} = (e)\underline{s}_{1s_1^{-1}\alpha} \circ \underline{s}_{2\alpha} = (e)\underline{s}_{2\alpha} = (e, s_2)\alpha.$$

Therefore if $s_1 \in t_1 \leq t \in T$, $s_2 \in t_2 \leq t \in T$, and $(e, s_1), (f, s_2) \in \text{dom}(\alpha)$, then

$$e \leq f \Rightarrow e \leq f \text{ and } (e, s_1)\alpha = (e, s_1s_1^{-1}s_2)\alpha = (e, s_2)\alpha \quad \text{by (14) and (15)}$$

$$\Rightarrow (e, s_1)\alpha = (e, s_2)\alpha \leq (f, s_2)\alpha \quad \underline{s}_{2\alpha} \text{ is an order isomorphism and } e \leq f$$

$$\Rightarrow (e, s_1)\alpha \leq (f, s_2)\alpha$$

$$\Rightarrow ((e, s_1)\alpha, s_2^{-1})\alpha \leq ((f, s_2)\alpha, s_2^{-1})\alpha$$

$$\Rightarrow e = (e, s_1s_2^{-1})\alpha \leq (f, s_2s_2^{-1})\alpha = f \quad \text{since } s_1s_2^{-1} \in E(S) \text{ by (13) and } e \in \text{dom}(\underline{s}_{1s_2^{-1}\alpha})$$

$$\Rightarrow e \leq f.$$

That Definition 10.2(2), (4) and (5) hold is clear. The remaining condition is condition (3). Suppose that $t_1, t_2 \in T$ and $(e)\underline{t}_{1q} = f$ and $(f)\underline{t}_{2q} = g$. We must show that $(e)\underline{t}_{1t_2q} = g$ and $(f)\underline{t}_{1q}^{-1} = e$.

We first want to show that $e \in \text{dom}(\underline{t}_{1t_2q})$. Since $e \in \text{dom}(\underline{t}_{1q})$, from the definition of q there exists $s_1 \in t'_1 \leq t_1$ such that $(e, s_1) \in \text{dom}(\alpha)$. Similarly, there exists $s_2 \in t'_2 \leq t_2$ with $(f, s_2) \in \text{dom}(\alpha)$. We want to show that there exists $s_3 \in t'_3 \leq t_1t_2$ with $(e, s_3) \in \text{dom}(\alpha)$.

Set $s_3 = s_1s_2$. By assumption $(e, s_1), (f, s_2) \in \text{dom}(\alpha)$, and so $(e, s_1)\alpha = (e)\underline{s_1}\alpha = (e, s_1)\alpha = (e, t_1)q = f$ and, similarly, $(f, s_2)\alpha = g$. Thus, since α is an action, $g = (f, s_2)\alpha = ((e, s_1)\alpha, s_2)\alpha = (e, s_1s_2)\alpha$. In particular, $(e, s_1s_2) \in \text{dom}(\alpha)$. Since $s_1 \in t'_1 \leq t_1$ and $s_2 \in t'_2 \leq t_2$, it follows that $s_1s_2 \in t'_1t'_2 \leq t_1t_2$ and so

$$(e)\underline{t_1t_2}q = (e, t_1t_2)q = (e, s_1s_2)\alpha = (f, s_2)\alpha = g,$$

as required.

It remains to show that $(f)\underline{t_1}^{-1}q = e$. Again, we begin by showing that $f \in \text{dom}(\underline{t_1}^{-1}q)$. Since $(e, s_1) \in \text{dom}(\alpha)$, there exists $s_1 \in t'_1 \leq t_1$ such that $(e, s_1) \in \text{dom}(\alpha)$. Since α is an inverse semigroup action, $((e)\underline{s_1}\alpha, s_1^{-1}) \in \text{dom}(\alpha)$. As we showed earlier, $(e)\underline{s_1}\alpha = f$, and so $(f, s_1^{-1}) \in \text{dom}(\alpha)$ and $(f, t_1^{-1}) \in \text{dom}(q)$. Thus $(f)\underline{t_1}^{-1}q = e$, as required. \square

We can now prove Theorem 11.3.

Theorem 11.3. *Every inverse semigroup S is isomorphic to some $Q(T, \mathcal{Y}, \mathcal{X})$ from Definition 11.1, where T is the maximum E -disjunctive homomorphic image of S , and \mathcal{Y} is the semilattice of idempotents of S .*

Proof. Let S be any inverse semigroup; let $\mathcal{Y} = E(S)$; let ρ be the syntactic congruence on S ; let $T = S/\rho$; and let $\phi_\rho: S \rightarrow T$ be the natural homomorphism defined by $(s)\phi_\rho = s/\rho$. We also recall the following:

- let $\alpha: E(S) \times S \rightarrow E(S)$ be the inverse semigroup action defined by $(e, s)\alpha = s^{-1}es$ (see (9)) for all $(e, s) \in E(S) \times S$ such that $e \leq ss^{-1}$;
- let $\pi: \text{dom}(\alpha) \rightarrow S$ be defined by $(e, s)\pi = s$ (see (10));
- let $\psi: S \rightarrow \text{dom}(\alpha)$ be defined by $(s)\psi = (ss^{-1}, s)$ (Lemma 11.5);
- let $M = \{(ss^{-1}, s/\rho) \in E(S) \times T \mid s \in S\}$ as defined in (11);
- let $q: E(S) \times T \rightarrow T$ be the preaction defined in Lemma 11.7;
- let \mathcal{X} be the poset defined in Theorem 10.5 (with respect to the preaction q) that contains \mathcal{Y} ;
- let $\beta: \mathcal{X} \times S \rightarrow \mathcal{X}$ be the action given in Theorem 10.5 such that β restricted to $(\mathcal{Y} \times S) \cap (\mathcal{Y})\beta^{-1}$ equals q .

We will verify that T , \mathcal{Y} , and \mathcal{X} satisfy the conditions in Definition 11.1. Firstly, by Theorem 10.5, β is an inverse semigroup action of T on \mathcal{X} by partial order isomorphisms, \mathcal{Y} is a meet subsemilattice, and order ideal, of \mathcal{X} .

- (1) Let $t \in T$ be arbitrary. We must show that $\text{dom } \underline{t}_\beta$ is an order ideal of \mathcal{X} . We can assume without loss of generality that t is an idempotent because $\text{dom}(\underline{t}_\beta) = \text{dom}(\underline{tt^{-1}}_\beta)$ for all $t \in T$. Let $a \in \mathcal{X}$ and $b \in \text{dom } \underline{t}_\beta \subseteq \mathcal{X}$ be such that $a \leq b$. By Theorem 10.5(3), there exists $s \in T^1$ such that $(a, s)\beta, (b, s)\beta \in \mathcal{Y}$ and $(a, s)\beta \leq (b, s)\beta$. In other words, $(a)\underline{s}_\beta \leq (b)\underline{s}_\beta$. Since $\text{dom}(\underline{s^{-1}ts}_q)$ is an order ideal in \mathcal{Y} (by Definition 10.2(1)) and \mathcal{Y} is an order ideal in \mathcal{X} , $\text{dom}(\underline{s^{-1}ts}_q) \cap \mathcal{Y}$ is an order ideal in \mathcal{X} . By assumption t is an idempotent, and so $s^{-1}ts$ is an idempotent also. It follows that $\text{dom}(\underline{s^{-1}ts}_q) \cap \mathcal{Y} = \text{dom}(\underline{s^{-1}ts}_\beta) \cap \mathcal{Y} = \text{dom}(\underline{s^{-1}ts}_q) \cap \mathcal{Y}$ is an order ideal of \mathcal{X} .

By assumption $b \in \text{dom } \underline{t}_\beta$ and so $(b)\underline{s}_\beta \in \text{dom}(\underline{s^{-1}ts}_\beta) \cap \mathcal{Y}$. Finally, since $\text{dom}(\underline{s^{-1}ts}_\beta) \cap \mathcal{Y}$ is an order ideal, $(a)\underline{s}_\beta \in \text{dom}(\underline{s^{-1}ts}_\beta)$, and so $a \in \text{dom}(\underline{t}_\beta)$, as required.

- (2) Firstly, for all $(y, t) \in \mathcal{Y} \times T$, the following hold:

$$\begin{aligned}
 (16) \quad y \in \text{dom}(\underline{t}_\beta) &\iff y \in \text{dom}(\underline{tt^{-1}}_\beta) \\
 &\iff y \in \text{dom}(\underline{tt^{-1}}_q) \\
 &\iff \text{there exists } e \in S \text{ such that } e/\rho \leq tt^{-1} \text{ and } y \leq e \\
 &\iff y/\rho \leq tt^{-1}.
 \end{aligned}$$

So if $y \in \mathcal{Y}$, then setting $t = y/\rho \in T$ and repeatedly applying (16) we obtain

$$\begin{aligned} \delta(y) &= \text{dom}(\underline{t}_\beta|_{\mathcal{Y}}) = \{z \in \mathcal{Y} \mid z/\rho \leq tt^{-1}\} \\ &= \bigcap \{\text{dom}(\underline{t}_{1\beta}|_{\mathcal{Y}}) \mid t_1 \in T \text{ such that } y/\rho \leq t_1 t_1^{-1}\} \\ &= \bigcap \{\text{dom}(\underline{t}_{1\beta}|_{\mathcal{Y}}) \mid t_1 \in T \text{ such that } y \in \text{dom}(\underline{t}_{1\beta}|_{\mathcal{Y}})\}. \end{aligned}$$

(3) We must show that for all $x \in \mathcal{X}$, there is $t \in T$ such that $(x, t)\beta \in \mathcal{Y}$. This is implied by Theorem 10.5(3).

Since S and M are isomorphic, by the definition of the multiplication of M , it suffices to show that we which was M and $Q(T, \mathcal{Y}, \mathcal{X})$ coincide (as semigroups).

The following holds:

$$\begin{aligned} Q(T, \mathcal{Y}, \mathcal{X}) &= \left\{ (y, t) \in \mathcal{Y} \times T \mid \text{dom}(\underline{t}_\beta) = \delta(y) = \text{dom}(\underline{y/\rho}_\beta|_{\mathcal{Y}}), (y, t)\beta \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid \text{dom}(\underline{s/\rho}_\beta) \cap \mathcal{Y} = \text{dom}(\underline{y/\rho}_\beta) \cap \mathcal{Y}, (y, s/\rho)\beta \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid \text{dom}(\underline{ss^{-1}/\rho}_\beta) \cap \mathcal{Y} = \text{dom}(\underline{y/\rho}_\beta) \cap \mathcal{Y}, (y, s/\rho)\beta \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid \bigcup (ss^{-1}/\rho) \downarrow = \bigcup (y/\rho) \downarrow \subseteq \mathcal{Y}, (y, s/\rho)\beta \in \mathcal{Y} \right\} \quad (\text{by (16)}) \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}/\rho) \downarrow = (y/\rho) \downarrow, (y, s/\rho)\beta \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, (y, s/\rho)\beta \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, (y, s/\rho)q \in \mathcal{Y} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, (y, s/\rho) \in \text{dom}(q) \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, \exists a \in S \text{ with } a/\rho \leq s/\rho \text{ and } y \leq aa^{-1} \right\} \quad (\text{by the definition of } q) \\ &= Q_1. \end{aligned}$$

If $(y, s'/\rho) \in Q_1$, then there exists $s \in s'/\rho$ such that $(ss^{-1}, y) \in \rho$ and there exists $a \in S$ with $a/\rho \leq s/\rho$ and $y \leq aa^{-1}$. Thus $ss^{-1}/\rho = y/\rho \leq aa^{-1}/\rho$. Moreover

$$ss^{-1}/\rho \leq aa^{-1}/\rho \iff ss^{-1}a/\rho \leq aa^{-1}a/\rho \iff s/\rho \leq a/\rho \iff s/\rho = a/\rho.$$

Thus

$$\begin{aligned} Q_1 &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, \exists a \in S \text{ with } a/\rho = s/\rho \text{ and } y \leq aa^{-1} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, \text{ and } y \leq ss^{-1} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid (ss^{-1}, y) \in \rho, y \leq ss^{-1} \right\} \\ &= \left\{ (y, s/\rho) \in \mathcal{Y} \times T \mid y = ss^{-1} \right\} \\ &= \left\{ (ss^{-1}, s/\rho) \in \mathcal{Y} \times T \mid s \in S \right\} = M. \end{aligned}$$

We have shown that M and $Q(T, \mathcal{Y}, \mathcal{X})$ are equal as sets. That their multiplications also coincide is precisely (12), and so the proof is complete. \square

ACKNOWLEDGEMENTS

The authors were supported by a Heilbronn Institute for Mathematical Research Small Grant during part of this work. The second named author was supported by a London Mathematical Society Early Career Fellowship and the Heilbronn Institute for Mathematical Research during this work.

REFERENCES

- [1] Branka P. Alimpić and Dragica N. Krgović, *Idempotent pure congruences on Clifford semigroups*, Algebraic Conference (Novi Sad, 1981), 1982, pp. 13–18. MR684674
- [2] Marina Anagnostopoulou-Merkouri, Zachary Mesyan, and James D. Mitchell, *Properties of congruence lattices of graph inverse semigroups*, International Journal of Algebra and Computation (April 2024), 1–26.
- [3] Bernd Billhardt, *On a wreath product embedding and idempotent pure congruences on inverse semigroups*, Semigroup Forum **45** (1992), no. 1, 45–54. MR1161418
- [4] Jean-Camille Birget, *Monoid generalizations of the Richard Thompson groups*, J. Pure Appl. Algebra **213** (2009), no. 2, 264–278. MR2468044
- [5] Collin Bleak and Martyn Quick, *The infinite simple group V of Richard J. Thompson: presentations by permutations*, Groups Geom. Dyn. **11** (2017), no. 4, 1401–1436. MR3737287
- [6] J. W. Cannon, W. J. Floyd, and W. R. Parry, *Introductory notes on Richard Thompson’s groups*, Enseign. Math. (2) **42** (1996), no. 3-4, 215–256. MR1426438
- [7] A. Distler and J. D. Mitchell, 2024.
- [8] G. G. Dyadchenko, *Structure of monogenic inverse semigroups*, Journal of Soviet Mathematics **24** (February 1984), no. 4, 428–434.
- [9] L. Elliott, A. Levine, and J. D. Mitchell, *Counting monogenic monoids and inverse monoids*, Comm. Algebra **51** (2023), no. 11, 4654–4661. MR4641433
- [10] D. G. Fitzgerald and Jonathan Leech, *Dual symmetric inverse monoids and representation theory*, Journal of the Australian Mathematical Society **64** (1998), no. 3, 345–367.
- [11] *GAP – Groups, Algorithms, and Programming, Version 4.12.2*, The GAP Group, 2022.
- [12] Roman S. Gigoń, *E -disjunctive semigroups and idempotent pure congruences*, Quasigroups Related Systems **21** (2013), no. 2, 229–238. MR3203149
- [13] David G. Green, *The lattice of congruences on an inverse semigroup*, Pacific J. Math. **57** (1975), no. 1, 141–152. MR390093
- [14] Peter Hines, *The inverse semigroup theory of elementary arithmetic*, arXiv e-prints (2022). <https://arxiv.org/abs/2206.07412>.
- [15] John M. Howie, *Fundamentals of semigroup theory*, London Mathematical Society Monographs. New Series, vol. 12, The Clarendon Press, Oxford University Press, New York, 1995. Oxford Science Publications. MR1455373
- [16] David G. Jones and Mark V. Lawson, *Graph inverse semigroups: their characterization and completion*, J. Algebra **409** (2014), 444–473. MR3198850
- [17] Mark Kambites, *Anisimov’s theorem for inverse semigroups*, Internat. J. Algebra Comput. **25** (2015), no. 1-2, 41–49. MR3325876
- [18] J. Kellendonk and Mark V. Lawson, *Partial actions of groups*, Internat. J. Algebra Comput. **14** (2004), no. 1, 87–114. MR2041539
- [19] Mark V. Lawson, *Inverse semigroups*, World Scientific Publishing Co., Inc., River Edge, NJ, 1998. The theory of partial symmetries. MR1694900
- [20] Yonghua Li and Moucheng Zhang, *On the E -disjunctive inverse semigroups*, Southeast Asian Bull. Math. **22** (1998), no. 2, 157–160. MR1684227
- [21] Yongle Luo and Zhengpan Wang, *Semimodularity in congruence lattices of graph inverse semigroups*, Comm. Algebra **49** (2021), no. 6, 2623–2632. MR4255032
- [22] Yongle Luo, Zhengpan Wang, and Jiaqun Wei, *Distributivity in congruence lattices of graph inverse semigroups*, Comm. Algebra **51** (2023), no. 12, 5046–5053. MR4652643
- [23] M. Malandro, *The inverse semigroups of order $j=15$* , 2024.
- [24] ———, *The unlabeled lattices on $j=15$ nodes*, 2024.
- [25] Martin E. Malandro, *Enumeration of finite inverse semigroups*, Semigroup Forum **99** (2019), no. 3, 679–723. MR4036868
- [26] Stuart W. Margolis and John C. Meakin, *Inverse monoids, trees and context-free languages*, Trans. Amer. Math. Soc. **335** (1993), no. 1, 259–276. MR1073775
- [27] D. B. McAlister, *Groups, semilattices and inverse semigroups. I, II*, Trans. Amer. Math. Soc. **192** (1974), 227–244; *ibid.* **196** (1974), 351–370. MR357660
- [28] John Meakin and Zhengpan Wang, *On graph inverse semigroups*, Semigroup Forum **102** (2021), no. 1, 217–234. MR4214502
- [29] Z. Mesyan, J. D. Mitchell, M. Morayne, and Y. H. Péresse, *Topological graph inverse semigroups*, Topology Appl. **208** (2016), 106–126. MR3506973
- [30] Zachary Mesyan and J. D. Mitchell, *The structure of a graph inverse semigroup*, Semigroup Forum **93** (2016), no. 1, 111–130. MR3528431
- [31] J. D. Mitchell et al., *Semigroups - GAP package, version 5.3.7*, 2024.

- [32] Liam O'Carroll, *Inverse semigroups as extensions of semilattices*, Glasgow Math. J. **16** (1975), no. 1, 12–21. MR407179
- [33] ———, *Idempotent determined congruences on inverse semigroups*, Semigroup Forum **12** (1976), no. 3, 233–243. MR404496
- [34] Mario Petrich, *Inverse semigroups*, Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, 1984. A Wiley-Interscience Publication. MR752899
- [35] Mario Petrich and Norman R. Reilly, *A network of congruences on an inverse semigroup*, Trans. Amer. Math. Soc. **270** (1982), no. 1, 309–325. MR642343
- [36] ———, *Operators related to E-disjunctive and fundamental completely regular semigroups*, J. Algebra **134** (1990), no. 1, 1–27.
- [37] G. B. Preston, *Monogenic inverse semigroups*, J. Austral. Math. Soc. Ser. A **40** (1986), no. 3, 321–342. MR832973
- [38] N. J. A. Sloane, *A001428*, The On-Line Encyclopedia of Integer Sequences (2023). OEIS Foundation Inc.
- [39] Zheng-Pan Wang, *Congruences on graph inverse semigroups*, J. Algebra **534** (2019), 51–64. MR3975713
- [40] Reikichi Yoshida, *On E-disjunctive inverse semigroups*, Acta Sci. Math. (Szeged) **49** (1985), no. 1-4, 49–52. MR839923