

Several classes of BCH codes of length $n = \frac{q^m-1}{2}$ Mengchen Lian^a, Shixin Zhu^{a,*}^a*School of Mathematics, HeFei University of Technology, Hefei 230601, China***Abstract**

BCH codes are an important class of linear codes and find extensive utilization in communication and disk storage systems. This paper mainly analyzes the negacyclic BCH code and cyclic BCH code of length $\frac{q^m-1}{2}$. For negacyclic BCH code, we give the dimensions of $\mathcal{C}_{(n,-1, \lceil \frac{\delta+1}{2} \rceil, 0)}$ for $\delta = a\frac{q^m-1}{q-1}, aq^{m-1} - 1 (1 \leq a < \frac{q-1}{2})$ and $\delta = a\frac{q^m-1}{q-1} + b\frac{q^m-1}{q^2-1}, aq^{m-1} + (a+b)q^{m-2} - 1 (2 \mid m, 1 \leq a+b \leq q-1, \lceil \frac{q-a-2}{2} \rceil \geq 1)$. Furthermore, the dimensions of negacyclic BCH codes $\mathcal{C}_{(n,-1,\delta,0)}$ with few nonzeros and $\mathcal{C}_{(n,-1,\delta,b)}$ with $b \neq 0$ are settled. For cyclic BCH code, we give the weight distribution of extended code $\bar{\mathcal{C}}_{(n,1,\delta,1)}$ and the parameters of dual code $\mathcal{C}_{(n,1,\delta,1)}^\perp$, where $\delta_2 \leq \delta \leq \delta_1$.

Keywords: BCH code, Negacyclic code, Dual code, Coset leader, Dually-BCH

1. Introduction

Let \mathcal{C} be an $[n, k, d]$ linear code over finite field \mathbb{F}_q where q is a prime power, having dimension k and minimum distance d , its dual code is

$$\mathcal{C}^\perp := \{\mathbf{b} \in \mathbb{F}_q^n : \mathbf{b} \cdot \mathbf{a}^T = 0 \text{ for any } \mathbf{a} \in \mathcal{C}\}$$

and \cdot denotes the standard inner product. The code \mathcal{C} is called λ -constacyclic if $(a_0, a_1, \dots, a_{n-1}) \in \mathcal{C}$ implies $(\lambda a_{n-1}, a_0, a_1, \dots, a_{n-2}) \in \mathcal{C}$, \mathcal{C} is cyclic if $\lambda = 1$ and \mathcal{C} is negacyclic if $\lambda = -1$. Representing $(a_0, a_1, \dots, a_{n-1}) \in \mathcal{C}$ as $a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1} \in \mathbb{F}_q[x]/\langle x^n - \lambda \rangle$, any λ -constacyclic code \mathcal{C} of length n is equivalent to an ideal of $\mathbb{F}_q[x]/\langle x^n - \lambda \rangle$, then there is a monic $g(x)$ such that $\mathcal{C} = \langle g(x) \rangle$. If $g(x)$ is the smallest degree, then $g(x)$ and $h(x) = \frac{x^n - \lambda}{g(x)}$ are referred to as the generator and parity-check polynomial of \mathcal{C} . Let \mathcal{C} have s nonzeros, i.e., $h(x)$ has s irreducible factors over \mathbb{F}_q . The extended linear code $\bar{\mathcal{C}}$ is

$$\bar{\mathcal{C}} = \{(a_0, a_1, \dots, a_{n-1}, a_n) : (a_0, a_1, \dots, a_{n-1}) \in \mathcal{C}, \sum_{i=0}^n a_i = 0\}.$$

In this paper, let $n, q \in \mathbb{Z}^+$ and $\gcd(n, q) = 1$. For $b, \delta \in \mathbb{Z}$. Let $m = \text{ord}_n(q)$, $\mathbb{F}_{q^m}^* = \langle \alpha \rangle$ and $\beta = \alpha^{\frac{q^m-1}{n}}$. We define $m_i(x)$ as the minimal polynomial corresponding to β^i with $i \in [0, n-1]$. Define

$$g_{(n,1,\delta,b)}(x) := \text{lcm}(m_b(x), m_{b+1}(x), \dots, m_{b+\delta-2}(x)) \text{ and } \delta \in [2, n],$$

*This research work is supported by the National Natural Science Foundation of China under Grant Nos. U21A20428 and 12171134.

*Corresponding author

Email addresses: lianmengchen0209@163.com (Mengchen Lian), zhushixinmath@hfut.edu.cn (Shixin Zhu)

lcm represents the lowest common multiple. Let $l = \text{ord}_{2n}(q)$, $\mathbb{F}_{q^l}^* = \langle \alpha \rangle$ and $\gamma = \alpha^{\frac{q^l-1}{2m}}$. We define $M_{1+2i}(x)$ as the minimal polynomial corresponding to γ^{1+2i} with $i \in [0, n-1]$. Define

$$g_{(n,-1,\delta,b)}(x) := \text{lcm}(M_{1+2b}(x), M_{1+2(b+1)}(x), \dots, M_{1+2(b+\delta-2)}(x)) \text{ and } \delta \in [2, n].$$

Let $\mathcal{C}_{(n,1,\delta,b)} = \langle g_{(n,1,\delta,b)}(x) \rangle$ and $\mathcal{C}_{(n,-1,\delta,b)} = \langle g_{(n,-1,\delta,b)}(x) \rangle$. Then $\mathcal{C}_{(n,1,\delta,b)}$ is a cyclic BCH code with the defining set $T = \bigcup_{i=b}^{b+\delta-2} C_i^n$, $\mathcal{C}_{(n,-1,\delta,b)}$ is a negacyclic BCH code with the defining set $T = \bigcup_{i=b}^{b+\delta-2} C_{1+2i}^{2n}$, where C_i^n is the q -cyclotomic coset of i (modulo n). Clearly, the dimension of $\mathcal{C}_{(n,\lambda,\delta,b)}$ is $n - |T|$, where $\lambda \in \{1, -1\}$. Let $\tilde{g}_{(n,1,\delta,b)}(x) = (x-1)g_{(n,1,\delta,b)}(x)$ and $\tilde{\mathcal{C}}_{(n,1,\delta,b)} = \langle \tilde{g}_{(n,1,\delta,b)}(x) \rangle$.

BCH codes studied for a long time and had many results in [1], [3], [6], [8], [15]-[21], [22]-[31]. In [10], [27], [33], [34], the authors examined the negacyclic codes of length $n = \frac{q^{2m}-1}{2}$, $\frac{q^m+1}{2}$, $\frac{q^m-1}{4}$ and $\frac{q^m-1}{2\lambda}$ respectively. In [12], [18], [21], the authors provided some research about weight distributions. There are many conclusions on the parameters in [4], [5], [14], [32]. Next, we list some known results for BCH codes of length $n = \frac{q^m-1}{2}$.

For negacyclic BCH code, Wang et al. [27] presented δ_i' (where $i = 1, 2, 3$ and δ_i' be defined in Section 2) and analysed the parameters of $\mathcal{C}_{(n,-1,\delta,0)}$ for δ in some ranges. Sun et al. [24] gave a class of ternary negacyclic code with parameters $[\frac{3^m-1}{2}, 2m, d \geq \frac{3^{m-1}-1}{2}]$ and its dual code with parameters $[\frac{3^m-1}{2}, \frac{3^m-1}{2} - 2m, 5]$.

For cyclic BCH code, Zhu et al. [35] determined the dimension of $\mathcal{C}_{(n,1,\delta,1)}$ with designed distance $2 \leq \delta \leq \frac{q^{\lfloor \frac{m+1}{2} \rfloor + 1}}{2}$, and gave δ_i (where $i = 1, 2, 3$ and δ_i be defined in Section 2) and the weight distributions of $\mathcal{C}_{(n,1,\delta_1,1)}$ and $\mathcal{C}_{(n,1,\delta_2,1)}$. Ling et al. [19] presented the minimum distances and weight enumerators of $\mathcal{C}_{(n,1,\delta,1)}$ for $\delta = \frac{q^m - q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i - 1}}{2}$ ($i \in [1, \lfloor \frac{m+11}{6} \rfloor]$) using the trace representation. Wang et al. [26] gave the parameters of $\mathcal{C}_{(n,1,\delta,1)}$ for $\delta = \frac{q^m - q^{m-1} - q^{\lfloor \frac{m-3}{2} \rfloor + i - 1}}{2}$ and $\frac{q^m - q^{m-1} - q^{\frac{3m-1}{4}} - q^{\frac{m+1}{2}} - q^{\frac{m-11}{4} + t} - 1}{2}$, where $1 \leq i \leq \lfloor \frac{m+6}{4} \rfloor$ and $2 \leq t \leq 4$.

In this paper, let $n = \frac{q^m-1}{2}$. The paper is segmented in the following sections: Section 2 gives some preliminaries. Section 3 presents the dimension of negacyclic codes $\mathcal{C}_{(n,-1,\delta,0)}$ with few nonzeros and δ takes some special forms. Section 4 gives the weight distribution of extended code and the parameters of dual code of cyclic BCH code. Section 5 serves as the conclusions of this paper.

2. Preliminaries

In this section, we introduce some notations and results about BCH codes.

- T^\perp is the defining set of \mathcal{C}^\perp .
- $CL(t) = \min\{i : i \in C_t^n\}$ is called the coset leader of C_t^n .
- $\Gamma_n = \{CL(t) : t \in Z_n\}$ is the set of all coset leaders.
- $\delta_i \in \Gamma_n$ is the i -th largest.
- $\delta_i' \in \Gamma_{2n}$ is odd and is the i -th largest.

- $\lceil a \rceil = \max\{a' : a' \leq a, a' \in Z\}$.
- $\lfloor a \rfloor = \min\{a' : a' \geq a, a' \in Z\}$.

Let $Tr_q^{q^r}$ be the trace mapping from \mathbb{F}_{q^r} to \mathbb{F}_q . Let $t = \sum_{s=0}^{m-1} t_s q^s$ be the q -adic expansion with $t \in Z$ and $0 < t < q^m - 1$. Denote $t = (t_{m-1}, t_{m-2}, \dots, t_0)$ is the sequence of t , the sequence of $tq^j \bmod n$ is $[tq^j]_n = (t_{m-1-j}, t_{m-2-j}, \dots, t_{m-j+1}, t_{m-j})$ which is the circular j -left-shift of t .

Lemma 1. ([27]) Let $i = (i_{m-1}, i_{m-2}, \dots, i_0)$ for $i \in [1, q^m - 1]$. Then $i \in \Gamma_{q^m-1}$ iff $[iq^j]_{q^m-1} \geq i$ for $j \in [0, m-1]$.

Lemma 2. ([13]) Let \mathcal{C} be a negacyclic code of length n and γ be a primitive $2n$ -th root of unity in \mathbb{F}_{q^m} . If $\{\gamma^{1+2^s} : t \leq s \leq t + \delta - 2\} \subseteq T$, then $d(\mathcal{C}) \geq \delta$.

Lemma 3. ([16]) Let $m = \text{ord}_n(q)$, $\mathbb{F}_{q^m}^* = \langle \alpha \rangle$ and $\beta = \alpha^{\frac{q^m-1}{n}}$. Let \mathcal{C} be a cyclic code of length n over \mathbb{F}_q and has s nonzeros. Let $\beta^{i_1}, \beta^{i_2}, \dots, \beta^{i_s}$ be the s roots of $h(x)$, they are not mutually conjugate. Denote $|C_{i_t}^n| = m_t$ for $1 \leq t \leq s$. Then the trace representation of \mathcal{C} is

$$\mathcal{C} = \{c(a_1, a_2, \dots, a_s) : a_t \in \mathbb{F}_{q^{m_t}}, 1 \leq t \leq s\},$$

where $c(a_1, a_2, \dots, a_s) = (\sum_{t=1}^s Tr_q^{q^{m_t}}(a_t \beta^{-li_t}))_{l=0}^{n-1}$.

Lemma 4. ([7], [11]) Let \mathcal{C} be an $[n, k, d]$ code over \mathbb{F}_q . Then

$$\begin{cases} \sum_{i=0}^{\lfloor \frac{d-1}{2} \rfloor} (q-1)^i \binom{n}{i} \leq q^{n-k}, \\ \sum_{i=0}^{\frac{d-2}{2}} (q-1)^i \binom{n-1}{i} \leq q^{n-k-1}, \quad \text{if } 2 \mid d. \end{cases}$$

Lemma 5. ([20]) Let $m \geq 3$, $2 \nmid m$ and $1 \leq i \leq q^{\frac{m+1}{2}} - 1$. Then $i \in \Gamma_{2n}$ iff $q \nmid i$. Moreover, $|C_i^{2n}| = m$.

Lemma 6. ([20]) Let $2 \mid m$ and $1 \leq i \leq 2q^{\frac{m}{2}} - 1$. Then $i \in \Gamma_{2n}$ iff $q \nmid i$. Moreover, $|C_{\frac{2n}{q^{\frac{m}{2}}+1}}^{2n}| = \frac{m}{2}$ and $|C_i^{2n}| = m$ for $i \neq q^{\frac{m}{2}} + 1$.

Lemma 7. ([27]) Let $2 \leq \delta \leq q^{\frac{m}{2}} + 1$ with $2 \mid m$ and $2 \leq \delta \leq \frac{q^{\frac{m+1}{2}} + 1}{2}$ with $2 \nmid m$. Then $\mathcal{C}_{(n, -1, \delta, 0)}$ is an $[n, n - m \lceil \frac{(2\delta-3)(q-1)}{2q} \rceil, d]$ negacyclic BCH code, where

$$d \geq \begin{cases} \delta + 1, & \text{if } q \mid (\delta - \frac{q+1}{2}); \\ \delta, & \text{otherwise.} \end{cases}$$

3. The parameters of negacyclic BCH code of length $n = \frac{q^m-1}{2}$

Let $n = \frac{q^m-1}{2}$. We will give $\dim(\mathcal{C}_{(n, -1, \lceil \frac{\delta+1}{2} \rceil, 0)})$ for $\tilde{\delta} = a \frac{q^m-1}{q-1}$, $a \frac{q^m-1}{q-1} + b \frac{q^m-1}{q^2-1}$ and $\frac{\delta_i'+1}{2}$. Furthermore, we discuss $\dim(\mathcal{C}_{(n, -1, \delta, b)})$ and the range of $d(\mathcal{C}_{(n, -1, \delta, 1)})$ for $b \neq 0$.

3.1. The dimension of $\mathcal{C}_{(n,-1,\delta,0)}$ with four specific designed distances

Let $1 \leq a < \frac{q-1}{2}$, $q > 3$ and $m > 2$ in Lemmas 8 and 10.

Lemma 8. Let $\delta = \left\lceil \frac{\tilde{\delta}+1}{2} \right\rceil$ and $\tilde{\delta} = a \frac{q^m-1}{q-1}$. Then $\mathcal{C}_{(n,-1,\delta,0)}$ is an $[n, k, d \geq \delta]$ negacyclic BCH code, where

$$k = \sum_{j=0}^{\lfloor \frac{m-1}{2} \rfloor} \left\lceil \frac{q-a-1}{2} \right\rceil^{2j+1} \left\lfloor \frac{q-a+1}{2} \right\rfloor^{m-2j-1} \binom{m}{2j+1}.$$

Proof. Note that

$$\tilde{\delta} = (a, a, \dots, a).$$

For code $\mathcal{C}_{(n,-1,\delta,0)}$, we have $T = \bigcup_{i=0}^{\delta-2} C_{1+2i}^{2n}$ and $\dim(\mathcal{C}_{(n,-1,\delta,0)}) = n - |T|$. To determine $\dim(\mathcal{C}_{(n,-1,\delta,0)})$, we need to determine the number of $s \notin T$ and $s \in \mathbb{Z}_{2n}$ with $2 \nmid s$. Let $A = \{s : s \in \mathbb{Z}_{2n}, 2 \nmid s, s \notin \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n}\}$. Hence, we can get $\dim(\mathcal{C}_{(n,-1,\delta,0)}) = |A|$.

Let $s' = \sum_{t=0}^{m-1} s_t q^t$ be the q -adic expansion. Let $s' \notin \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n}$ and $s' \in \mathbb{Z}_{2n}$. Clearly, $s' \notin \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n} \Leftrightarrow a \leq s_t \leq q-1$ for any $0 \leq t \leq m-1$ by Theorem 3.2 of [2]. Let $B = \{s_t : 2 \nmid s_t, 0 \leq t \leq m-1\}$. Since q^t is odd, then $2 \nmid s' \Leftrightarrow 2 \nmid |B|$.

• **Case 1:** $2 \mid m$.

If $2 \nmid a$, then $|\{t : 2 \nmid s_t\}| = \frac{q-a}{2}$ and $|\{t : 2 \mid s_t\}| = \frac{q-a}{2}$ for any $s_t \in [a, q-1]$. Since $2 \nmid s' \Leftrightarrow 2 \nmid |B|$, then $|A| = \sum_{j=0}^{\frac{m}{2}-1} \binom{m}{2j+1} \left(\frac{q-a}{2}\right)^m$. If $2 \mid a$, then $|\{t : 2 \nmid s_t\}| = \frac{q-a-1}{2}$ and $|\{t : 2 \mid s_t\}| = \frac{q-a+1}{2}$ for any $s_t \in [a, q-1]$. Since $2 \nmid s' \Leftrightarrow 2 \nmid |B|$, then $|A| = \sum_{j=0}^{\frac{m}{2}-1} \binom{m}{2j+1} \left(\frac{q-a-1}{2}\right)^{2j+1} \left(\frac{q-a+1}{2}\right)^{m-2j-1}$.

• **Case 2:** $2 \nmid m$.

If $2 \nmid a$, since $|\{t : 2 \nmid s_t\}| = |\{t : 2 \mid s_t\}| = \frac{q-a}{2}$ and $2 \nmid s' \Leftrightarrow 2 \nmid |B|$, then $|A| = \sum_{j=0}^{\frac{m-1}{2}} \binom{m}{2j+1} \left(\frac{q-a}{2}\right)^m$. By the same way, we have $|A| = \sum_{j=0}^{\frac{m-1}{2}} \binom{m}{2j+1} \left(\frac{q-a-1}{2}\right)^{2j+1} \left(\frac{q-a+1}{2}\right)^{m-2j-1}$ for $2 \mid a$.

Summing up all the discussions above, we complete our proof. \square

Example 9. Let $(q, m) = (5, 3)$ and $a = 1$, one has $n = 62$, $\tilde{\delta} = 31$ and $\delta = 16$. Then $\dim(\mathcal{C}_{(62,-1,16,0)}) = 32$. Let $(q, m) = (7, 3)$ and $a = 2$, one has $n = 171$, $\tilde{\delta} = 114$ and $\delta = 58$. Then $\dim(\mathcal{C}_{(171,-1,58,0)}) = 62$.

Lemma 10. Let $\delta = \left\lceil \frac{\tilde{\delta}'+1}{2} \right\rceil$ and $\tilde{\delta}' = aq^{m-1} - 1$. Then $\mathcal{C}_{(n,-1,\delta,0)}$ is an $[n, k, d \geq \delta]$ negacyclic BCH code, where

$$k = \sum_{j=0}^{\lfloor \frac{m-1}{2} \rfloor} \left\lceil \frac{q-a-1}{2} \right\rceil^{2j+1} \left\lfloor \frac{q-a+1}{2} \right\rfloor^{m-2j-1} \binom{m}{2j+1} + \kappa m,$$

$\kappa = 1$ for $2 \mid a$ and $\kappa = 0$ for $2 \nmid a$.

Proof. From Theorem 3.5 of [2], we know that $\bigcup_{i=1}^{\tilde{\delta}'-1} C_i^{2n} = \bigcup_{i=1}^{\tilde{\delta}'-1} C_i^{2n}$, $\tilde{\delta}' \in \Gamma_{2n}$ and $|C_{\tilde{\delta}'}^{2n}| = m$, where $\tilde{\delta}' = a \frac{q^m-1}{q-1}$. Similar to Lemma 8, we can get

$$\begin{aligned} \dim(\mathcal{C}_{(n,-1,\delta,0)}) &= |\{s : s \in \mathbb{Z}_{2n}, 2 \nmid s, s \notin \bigcup_{i=1}^{\tilde{\delta}'-1} C_i^{2n}\}| \\ &= |\{s : s \in \mathbb{Z}_{2n}, 2 \nmid s, s \notin \bigcup_{i=1}^{\tilde{\delta}'} C_i^{2n}\}| + |\{s : 2 \nmid s, s \in C_{\tilde{\delta}'}^{2n}\}|. \end{aligned}$$

Since $2 \mid \tilde{\delta}$ for $2 \nmid a$ and $2 \nmid \tilde{\delta}$ for $2 \mid a$, then

$$|\{s : 2 \nmid s, s \in C_{\tilde{\delta}'}^{2n}\}| = \begin{cases} m & \text{if } 2 \mid a; \\ 0 & \text{if } 2 \nmid a. \end{cases}$$

Combining Lemma 8, we get $\dim(\mathcal{C}_{(n,-1,\delta,0)})$. □

We assume that $1 \leq a + b \leq q - 1$, $m > 2$ is even and $\lceil \frac{q-a-2}{2} \rceil \geq 1$ in Lemmas 11 and 13.

Lemma 11. *Let $\delta = \lceil \frac{\tilde{\delta}+1}{2} \rceil$ and $\tilde{\delta} = a \frac{q^m-1}{q-1} + b \frac{q^m-1}{q^2-1}$. Then $\mathcal{C}_{(n,-1,\delta,0)}$ is an $[n, k, d \geq \delta]$ negacyclic BCH code, where*

$$k = \sum_{t=0}^{\frac{m}{2}} \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left\lfloor \frac{q-a-b}{2} \right\rfloor^j \left\lceil \frac{q-a-b}{2} \right\rceil^{t-j} \phi,$$

when $2 \nmid a$,

$$\phi = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left\lceil \frac{q-a-2}{2} \right\rceil^{2i} \left\lfloor \frac{q-a}{2} \right\rfloor^{m-2t-2i}, & \text{if } 2 \nmid t \text{ and } 2 \mid j, \text{ or } 2 \mid t \text{ and } 2 \nmid j; \\ \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left\lceil \frac{q-a-2}{2} \right\rceil^{2i+1} \left\lfloor \frac{q-a}{2} \right\rfloor^{m-2t-2i-1}, & \text{if } 2 \nmid t \text{ and } 2 \nmid j, \text{ or } 2 \mid t \text{ and } 2 \mid j; \end{cases}$$

when $2 \mid a$,

$$\phi = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left\lceil \frac{q-a-2}{2} \right\rceil^{2i} \left\lfloor \frac{q-a}{2} \right\rfloor^{m-2t-2i}, & \text{if } 2 \nmid j; \\ \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left\lceil \frac{q-a-2}{2} \right\rceil^{2i+1} \left\lfloor \frac{q-a}{2} \right\rfloor^{m-2t-2i-1}, & \text{if } 2 \mid j. \end{cases}$$

Proof. Note that

$$\tilde{\delta} = (a, a+b, a, a+b, \dots, a, a+b).$$

Let $A = \{s : s \in \mathbb{Z}_{2n}, 2 \nmid s, s \notin \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n}\}$. By the same way as Lemma 8, we have $\dim(\mathcal{C}_{(n,-1,\delta,0)}) = |A|$. Let $s' = \sum_{u=0}^{m-1} s_u q^u \notin \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n}$ and $s' \in \mathbb{Z}_{2n}$. This is equivalent to $a+b \leq s_{(u-1)} \leq q-1$ if $s_u = a$ and $a \leq s_{(u-1)} \leq q-1$ if $s_u > a$ by Theorem 3.1 of [22], where “ $s_{(u-1)}$ ” means “ $s_{(u-1) \pmod{m}}$ ”. Let $B = \{u : s_u = a, s_{(u-1)} \in [a+b, q-1], 0 \leq u \leq m-1\}$ and $|B| = t$. It is clear that $0 \leq t \leq \frac{m}{2}$ and there exist $\frac{m}{m-t} \binom{m-t}{t}$ alternatives for $|B| = t$. Since $2 \nmid s' \Leftrightarrow s' \in A \Leftrightarrow 2 \nmid (|\{u : 2 \nmid s_u\}| + |\{u-1 : 2 \nmid s_{(u-1)}\}| + |\{i : 2 \nmid s_i\}|)$ with $u \in B$ and $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$, then we will discuss the parity of a, b and t .

• **Case 1:** $2 \nmid a, b$.

In this case, $s_{(u-1)} \in [a+b, q-1]$ and $s_i \in [a+1, q-1]$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$. Clearly, $|\{u-1 : 2 \nmid s_{(u-1)}\}| = \frac{q-a-b-1}{2}$ and $|\{u-1 : 2 \mid s_{(u-1)}\}| = \frac{q-a-b+1}{2}$ for $u \in B$ and $s_{(u-1)} \in [a+b, q-1]$. Furthermore, $|\{i : s_i > a, 2 \nmid s_i\}| = \frac{q-a-2}{2}$ and $|\{i : s_i > a, 2 \mid s_i\}| = \frac{q-a}{2}$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$.

If $2 \nmid t$, then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b-1}{2}\right)^j \left(\frac{q-a-b+1}{2}\right)^{t-j} \phi_1$, where

$$\phi_1 = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left(\frac{q-a-2}{2}\right)^{2i} \left(\frac{q-a}{2}\right)^{m-2t-2i}, & \text{if } 2 \mid j; \\ \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left(\frac{q-a-2}{2}\right)^{2i+1} \left(\frac{q-a}{2}\right)^{m-2t-2i-1}, & \text{if } 2 \nmid j. \end{cases}$$

If $2 \mid t$, then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b-1}{2}\right)^j \left(\frac{q-a-b+1}{2}\right)^{t-j} \phi_2$, where

$$\phi_2 = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left(\frac{q-a-2}{2}\right)^{2i+1} \left(\frac{q-a}{2}\right)^{m-2t-2i-1}, & \text{if } 2 \mid j; \\ \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left(\frac{q-a-2}{2}\right)^{2i} \left(\frac{q-a}{2}\right)^{m-2t-2i}, & \text{if } 2 \nmid j. \end{cases}$$

- **Case 2:** $2 \nmid a$ and $2 \mid b$.

In this case, $s_{(u-1)} \in [a+b, q-1]$ for $u \in B$ and $s_i \in [a+1, q-1]$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$. Clearly, $|\{u-1 : 2 \nmid s_{(u-1)}\}| = \frac{q-a-b}{2}$ and $|\{u-1 : 2 \mid s_{(u-1)}\}| = \frac{q-a-b}{2}$ for $s_{(u-1)} \in [a+b, q-1]$. Furthermore, $|\{i : s_i > a, 2 \nmid s_i\}| = \frac{q-a-2}{2}$ and $|\{i : s_i > a, 2 \mid s_i\}| = \frac{q-a}{2}$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$.

If $2 \nmid t$, then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b}{2}\right)^j \left(\frac{q-a-b}{2}\right)^{t-j} \phi_3$, where $\phi_3 = \phi_1$.

If $2 \mid t$, then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b}{2}\right)^j \left(\frac{q-a-b}{2}\right)^{t-j} \phi_4$, where $\phi_4 = \phi_2$.

- **Case 3:** $2 \mid a, b$.

In this case, $|\{u : 2 \nmid s_u, u \in B\}| = 0$. By the same way as above, $|\{u-1 : 2 \nmid s_{(u-1)}\}| = \frac{q-a-b-1}{2}$ and $|\{u-1 : 2 \mid s_{(u-1)}\}| = \frac{q-a-b+1}{2}$ for $s_{(u-1)} \in [a+b, q-1]$. Furthermore, $|\{i : s_i > a, 2 \nmid s_i\}| = \frac{q-a-1}{2}$ and $|\{i : s_i > a, 2 \mid s_i\}| = \frac{q-a-1}{2}$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$.

Then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b-1}{2}\right)^j \left(\frac{q-a-b+1}{2}\right)^{t-j} \phi_5$, where

$$\phi_5 = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left(\frac{q-a-1}{2}\right)^{2i} \left(\frac{q-a-1}{2}\right)^{m-2t-2i}, & \text{if } 2 \nmid j; \\ \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left(\frac{q-a-1}{2}\right)^{2i+1} \left(\frac{q-a-1}{2}\right)^{m-2t-2i-1}, & \text{if } 2 \mid j. \end{cases}$$

- **Case 4:** $2 \mid a$ and $2 \nmid b$.

In this case, $|\{u : 2 \nmid s_u, u \in B\}| = 0$. Clearly, $|\{u-1 : 2 \nmid s_{(u-1)}\}| = \frac{q-a-b}{2}$ and $|\{u-1 : 2 \mid s_{(u-1)}\}| = \frac{q-a-b}{2}$ for $s_{(u-1)} \in [a+b, q-1]$. Furthermore, $|\{i : s_i > a, 2 \nmid s_i\}| = \frac{q-a-1}{2}$ and $|\{i : s_i > a, 2 \mid s_i\}| = \frac{q-a-1}{2}$ for $i \in \mathbb{Z}_m \setminus \{u, (u-1) \pmod{m} : u \in B\}$.

Then $|A| = \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left(\frac{q-a-b}{2}\right)^j \left(\frac{q-a-b}{2}\right)^{t-j} \phi_6$, where $\phi_6 = \phi_5$.

Summarizing the above, we arrive at the results. \square

Example 12. Let $(q, m) = (5, 4)$, one has $n = 312$. When $(a, b) = (1, 2)$, we have $\tilde{\delta} = 208$ and $\delta = 105$, then $\dim(\mathcal{C}_{(312, -1, 105, 0)}) = 80$. When $(a, b) = (2, 2)$, we have $\tilde{\delta} = 364$ and $\delta = 183$, then $\dim(\mathcal{C}_{(312, -1, 183, 0)}) = 16$.

Lemma 13. Let $\delta = \left\lceil \frac{\tilde{\delta}+1}{2} \right\rceil$ and $\tilde{\delta}' = a(q^{m-1} + q^{m-2}) + bq^{m-2} - 1$. Then $\mathcal{C}_{(n, -1, \delta, 0)}$ is an $[n, k, d \geq \delta]$ negacyclic BCH code, where

$$k = \sum_{t=0}^{\frac{m}{2}} \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left\lfloor \frac{q-a-b}{2} \right\rfloor^j \left\lceil \frac{q-a-b}{2} \right\rceil^{t-j} \phi + \kappa m,$$

ϕ is given in Lemma 11, $\kappa = 1$ for $2 \mid b$ and $\kappa = 0$ for $2 \nmid b$.

Proof. From the Theorem 3.4 of [22], $r \notin \Gamma_{2n}$ for any $\tilde{\delta}' < r < \tilde{\delta}$, where $\tilde{\delta} = a\frac{q^m-1}{q-1} + b\frac{q^m-1}{q^2-1}$, i.e., $\bigcup_{i=1}^{\tilde{\delta}'} C_i^{2n} = \bigcup_{i=1}^{\tilde{\delta}-1} C_i^{2n}$. Similar to Lemma 11, we can get

$$\dim(\mathcal{C}_{(n, -1, \delta, 0)}) = |\{s : s \in \mathbb{Z}_{2n}, 2 \nmid s, s \notin \bigcup_{i=1}^{\tilde{\delta}'} C_i^{2n}\}| + |\{s : 2 \nmid s, s \in C_{\tilde{\delta}'}^{2n}\}|.$$

Note that $\tilde{\delta}' \in \Gamma_{2n}$ and $|C_{\tilde{\delta}'}^{2n}| = m$, combining $2 \nmid \tilde{\delta}'$ for $2 \nmid b$ and $2 \nmid \tilde{\delta}'$ for $2 \mid b$, we have

$$|\{s : 2 \nmid s, s \in C_{\tilde{\delta}'}^{2n}\}| = \begin{cases} m & \text{if } 2 \nmid b; \\ 0 & \text{if } 2 \mid b. \end{cases}$$

Hence, we get $\dim(\mathcal{C}_{(n,-1,\delta,0)})$. □

We immediately get the following lemma for $a = 0$ and $1 \leq b \leq q - 1$ in Lemma 11.

Lemma 14. *Let $\delta = \left\lceil \frac{\delta''+1}{2} \right\rceil$ and $\delta'' = b \frac{q^m-1}{q^2-1}$. Then $\mathcal{C}_{(n,-1,\delta,0)}$ is an $[n, k, d \geq \delta]$ negacyclic BCH code, where*

$$k = \sum_{t=0}^{\frac{m}{2}} \frac{m}{m-t} \binom{m-t}{t} \sum_{j=0}^t \binom{t}{j} \left\lfloor \frac{q-b}{2} \right\rfloor^j \left\lceil \frac{q-b}{2} \right\rceil^{t-j} \phi,$$

where

$$\phi = \begin{cases} \sum_{i=0}^{\frac{m}{2}-t} \binom{m-2t}{2i} \left\lceil \frac{q-2}{2} \right\rceil^{2i} \left\lfloor \frac{q}{2} \right\rfloor^{m-2t-2i}, & \text{if } 2 \nmid j; \\ \sum_{i=0}^{\frac{m}{2}-t-1} \binom{m-2t}{2i+1} \left\lceil \frac{q-2}{2} \right\rceil^{2i+1} \left\lfloor \frac{q}{2} \right\rfloor^{m-2t-2i-1}, & \text{if } 2 \mid j. \end{cases}$$

3.2. The dimensions of some codes with few nonzeros

Lemma 15. ([27]) *Let $n = \frac{q^m-1}{2}$. Then*

$$\delta_1' = q^m - q^{m-1} - 1, \quad \delta_2' = q^m - q^{m-1} - q^{\lfloor \frac{2m-1}{3} \rfloor} - q^{\lfloor \frac{m-1}{3} \rfloor} - 1$$

and $|C_{\delta_1'}^{2n}| = m$, $|C_{\delta_2'}^{2n}| = \begin{cases} m, & \text{if } 3 \nmid m, \\ \frac{m}{3}, & \text{if } 3 \mid m \end{cases}$. If $q^m \geq 25$, then

$$\delta_3' = \begin{cases} q^m - q^{m-1} - q^{\lfloor \frac{2m-1}{3} \rfloor} - q^{\lfloor \frac{m-1}{3} \rfloor} - 1, & \text{if } 3 \nmid (m+1), \\ q^m - q^{m-1} - q^{\frac{2m-1}{3}} - q^{\frac{m+1}{3}} - 1, & \text{if } 3 \mid (m+1) \end{cases}$$

and $|C_{\delta_3'}^{2n}| = m$.

Lemma 16. *Let $m \equiv 2 \pmod{3}$, $m \geq 8$ and $2 \leq i \leq 7$. Then*

$$\delta_i' = q^m - q^{m-1} - q^{\frac{2m-1}{3} + \varepsilon_1} - q^{\frac{m+1}{3} + \zeta_1} - 1,$$

where

$$\varepsilon_1 = \begin{cases} 0, & \text{if } 2 \leq i \leq 3; \\ 1, & \text{if } 4 \leq i \leq 7, \end{cases} \quad \zeta_1 = \begin{cases} -1, & \text{if } i = 2; \\ 1, & \text{if } i = 3; \\ i - 6, & \text{if } 4 \leq i \leq 7. \end{cases}$$

And $|C_{\delta_i'}^{2n}| = m$ for $4 \leq i \leq 7$.

Proof. When $q \geq 3$ and $q^m \geq 25$, δ_1' , δ_2' and δ_3' were determined in [27]. Now we first prove that $\delta_i' = q^m - q^{m-1} - q^{\frac{2m-1}{3} + 1} - q^{\frac{m+1}{3} + i - 6} - 1$ is an odd coset leader for $4 \leq i \leq 7$, i.e., $q^t \delta_i' \geq \delta_i'$ for any $0 \leq t \leq m - 1$. Let Q_j be $\underbrace{q-1, \dots, q-1}_j$. Clearly,

$$\delta_i' = (q-2, Q_{\frac{m+1}{3}-3}, q-2, Q_{\frac{m-2}{3}+6-i}, q-2, Q_{\frac{m+1}{3}+i-6}).$$

(1) If $0 \leq t \leq \frac{m+1}{3} - 3$, it's obvious that $q^t \delta_i' > \delta_i'$.

(2) If $t = \frac{m+1}{3} - 2$, then

$$q^t \delta_i' = (q-2, Q_{\frac{m-2}{3}+6-i}, q-2, Q_{\frac{m+1}{3}+i-6}, q-2, Q_{\frac{m+1}{3}-3}).$$

Note that $\frac{m-2}{3} + 6 - i > \frac{m+1}{3} - 3$, thus $q^t \delta_i' > \delta_i'$.

(3) If $\frac{m+1}{3} - 1 \leq t \leq \frac{2m-1}{3} + 4 - i$, we have

$$[q^t \delta_i']_{2n} = q^m - q^{1+t-\frac{m+1}{3}} - q^{\frac{m+1}{3}+i-6+t} - q^{t-1} - 1 = [q^t \delta_i']_n > \delta_i',$$

since $\frac{m+1}{3} + i - 6 + t < m - 1$ and $t - 1 < \frac{2m-1}{3} + 1$.

(4) If $t = \frac{2m-1}{3} + 5 - i$, then

$$q^t \delta_i' = (q-2, Q_{\frac{m+1}{3}+i-6}, q-2, Q_{\frac{m+1}{3}-3}, q-2, Q_{\frac{m-2}{3}+6-i}),$$

since $\frac{m+1}{3} + i - 6 > \frac{m+1}{3} - 3$, thus $q^t \delta_i' > \delta_i'$.

(5) If $\frac{2m-1}{3} + 6 - i \leq t \leq m - 1$, then

$$[q^t \delta_i']_{2n} = q^m - q^{1+t-\frac{m+1}{3}} - q^{i-6+t-\frac{2m-1}{3}} - q^{t-1} - 1 = [q^t \delta_i']_n > \delta_i',$$

by $t < m$, $1 + t - \frac{m+1}{3} < \frac{2m-1}{3} + 1$ and $i - 6 + t - \frac{2m-1}{3} < \frac{m+1}{3} + i - 6$.

Hence, we have $\delta_i' \in \Gamma_{2n}$ and $2 \nmid \delta_i'$ for $4 \leq i \leq 7$. Since $q^t \delta_i' > \delta_i'$ for any $0 \leq t \leq m - 1$, then $|C_{\delta_i'}^{2n}| = m$. Next we prove that no integer $\delta' \in [\delta'_{i+1} + 1, \delta_i' - 1]$ satisfies $\delta' \in \Gamma_{2n}$ and $2 \nmid \delta'$. It is easy seen that

$$\delta'_{i+1} = (q-2, Q_{\frac{m+1}{3}-3}, q-2, Q_{\frac{m-2}{3}+5-i}, q-2, Q_{\frac{m+1}{3}+i-5}).$$

Let $\delta' \in [\delta'_{i+1} + 1, \delta_i' - 1]$. According to the Lemma 1. Then the form of δ is

$$\delta' = (q-2, Q_{\frac{m+1}{3}-3}, q-2, Q_{\frac{m-2}{3}+6-i}, \delta''_{\frac{m+1}{3}+i-6}, \delta''_{\frac{m+1}{3}+i-7}, \dots, \delta''_0).$$

Supposing δ' is an odd coset leader, then $q-2 \leq \delta''_t \leq q-1$ for $0 \leq t \leq \frac{m+1}{3} + i - 7$ and $\delta''_{\frac{m+1}{3}+i-6} = q-2$.

Let $A = \{\delta''_t : \delta''_t = q-2, 0 \leq t \leq \frac{m+1}{3} + i - 7\}$. If $|A| = 0$, then $\delta' = \delta_i'$, this contradicts $\delta' \in [\delta'_{i+1} + 1, \delta_i' - 1]$. If $|A| = 1$, then $2 \mid \delta'$, that is impossible. If $|A| = 2$, then

$$(\delta''_{\frac{m+1}{3}+i-7}, \dots, \delta''_0) = (Q_{a_1}, q-2, Q_{a_2}, q-2, Q_{a_3}),$$

where $\sum_{j=1}^3 a_j = \frac{m+1}{3} + i - 8$ and $a_j \geq 0$ for $4 \leq i \leq 7$ and $1 \leq j \leq 3$. There must exist an a_j that does not satisfy $a_j \geq \frac{m+1}{3} - 3$ for $1 \leq j \leq 3$, i.e., $\delta' \notin \Gamma_{2n}$. If $|A| > 2$, then δ' does not satisfy $\delta' \in \Gamma_{2n}$ and $2 \nmid \delta'$. \square

Theorem 17. *Let $m \equiv 2 \pmod{3}$, $m \geq 8$. Then $\dim(\mathcal{C}_{(n,-1,\frac{\delta_i'+1}{2},0)}) = im$, where δ_i' is given in Lemmas 15 and 16.*

Proof. Using lemmas 15 and 16, we can directly prove this conclusion. \square

Using the same approach as above, the following results can be established.

Lemma 18. Let $m \equiv 0 \pmod{3}$, $m \geq 9$ and $2 \leq i \leq 10$. Then

$$\delta_i' = q^m - q^{m-1} - q^{\frac{2m}{3} + \varepsilon_2} - q^{\frac{m}{3} + \zeta_2} - 1,$$

where

$$\varepsilon_2 = \begin{cases} -1, & \text{if } i = 2; \\ 0, & \text{if } 3 \leq i \leq 5; \\ 1, & \text{if } 6 \leq i \leq 10, \end{cases} \quad \zeta_2 = \begin{cases} -1, & \text{if } 2 \leq i \leq 3; \\ i - 4, & \text{if } 4 \leq i \leq 5; \\ i - 8, & \text{if } 6 \leq i \leq 10. \end{cases}$$

And $|C_{\delta_i'}^{2n}| = m$ for $4 \leq i \leq 10$.

Theorem 19. Let $m \equiv 0 \pmod{3}$, $m \geq 9$. Then $\dim(C_{(n, -1, \frac{\delta_i'+1}{2}, 0)}) = (i - \frac{2}{3})m$, where δ_i' is given in Lemma 18 with $2 \leq i \leq 10$.

Lemma 20. Let $m \equiv 1 \pmod{3}$, $m \geq 7$ and $2 \leq i \leq 6$. Then

$$\delta_i' = q^m - q^{m-1} - q^{\frac{2m+1}{3} + \varepsilon_3} - q^{\frac{m-1}{3} + \zeta_3} - 1,$$

where

$$\varepsilon_3 = \begin{cases} -1, & \text{if } i = 2; \\ 0, & \text{if } 3 \leq i \leq 6, \end{cases} \quad \zeta_3 = \begin{cases} 0, & \text{if } i = 2; \\ i - 4, & \text{if } 3 \leq i \leq 6. \end{cases}$$

And $|C_{\delta_i'}^{2n}| = m$ for $4 \leq i \leq 6$.

Theorem 21. Let $m \equiv 1 \pmod{3}$, $m \geq 7$. Then $\dim(C_{(n, -1, \frac{\delta_i'+1}{2}, 0)}) = im$, where δ_i' is given in Lemma 20 with $2 \leq i \leq 6$.

We discuss the parameters of negacyclic code C' with length $n = \frac{q^m-1}{2}$ and check polynomial $\text{lcm}(M_1(x), M_{\delta_1'}(x))$.

Theorem 22. Let $n = \frac{q^m-1}{2}$ and $m \geq 3$. Then C' is an $[n, 2m, d(C') \geq \frac{(q-2)q^{m-1}-1}{2}]$ code and C'^{\perp} is an $[n, n-2m, 3 \leq d(C'^{\perp}) \leq 5]$ dual code.

Proof. Since $1 \in \Gamma_{2n}$ and $\delta_1' \in \Gamma_{2n}$, then $\delta_1' \notin C_1^{2n} \Rightarrow \text{lcm}(M_1(x), M_{\delta_1'}(x)) = M_1(x)M_{\delta_1'}(x)$. From Lemma 15, we have $|C_1^{2n}| = |C_{\delta_1'}^{2n}| = m \Rightarrow \dim(C') = 2m$ and $\dim(C'^{\perp}) = n - 2m$.

Let

$$H = \{1 + 2i : \frac{q^{m-1} + 1}{2} \leq i \leq \frac{q^m - q^{m-1} - 4}{2}\}.$$

It is easy to get $H \subset T = \{1 + 2s : s \in [0, n-1]\} \setminus (C_1^{2n} \cup C_{\delta_1'}^{2n})$, and β^i is a root of C' for any $i \in H$. Hence, $d(C') \geq \frac{(q-2)q^{m-1}-1}{2}$ by Lemma 2.

Clearly, $-1, 1 \in C_1^{2n} \cup T^{\perp} = C_{\delta_1'}^{2n}$, then $d(C'^{\perp}) \geq 3$ by Lemma 2. If $d(C'^{\perp}) = 7$, by Lemma 4, $\sum_{i=0}^3 (q-1)^i \binom{n}{i} > q^{2m}$, this is infeasible, i.e., $d(C'^{\perp}) \leq 6$. If $d(C'^{\perp}) = 6$, we have $\sum_{i=0}^2 (q-1)^i \binom{n-1}{i} > q^{2m-1}$, this is infeasible, i.e., $3 \leq d(C'^{\perp}) \leq 5$. Thus we complete the proof. \square

3.3. The dimension of $\mathcal{C}_{(n,-1,\delta,b)}$ with $b \neq 0$

Let $CL(a_1, a_2) = \{1 + 2i \in [a_1, a_2] : 1 + 2i \in \Gamma_{2n}\}$, we will give some results about $\mathcal{C}_{(n,-1,\delta,b)}$ for $b \geq 1$.

Theorem 23. Let $m \geq 2$, $1 \leq 1 + 2(b + \delta - 2) \leq q^{\frac{m+1}{2}} - 1$ for $2 \nmid m$ and $1 \leq 1 + 2(b + \delta - 2) \leq 2q^{\frac{m}{2}} - 1$ for $2 \mid m$. Then $\mathcal{C}_{(n,-1,\delta,b)} = \mathcal{C}_{(n,-1,b+\delta,0)}$ for $1 + 2b \leq \left\lfloor \frac{b+\delta-2}{q} \right\rfloor$ and

$$\dim(\mathcal{C}_{(n,-1,\delta,b)}) = n - m \left\lceil \frac{(2\delta + 2b - 3)(q - 1)}{2q} \right\rceil.$$

Proof. We have $T = \bigcup_{i=b}^{b+\delta-2} C_{1+2i}^{2n}$ for code $\mathcal{C}_{(n,-1,\delta,b)}$. If $2 \nmid m$, then $|C_{1+2i}^{2n}| = m$ for $1 \leq 1 + 2i \leq q^{\frac{m+1}{2}} - 1$ from Lemma 5. If $2 \mid m$, since $q^{\frac{m}{2}} + 1$ is even, then $|C_{1+2i}^{2n}| = m$ for $1 \leq 1 + 2i \leq 2q^{\frac{m}{2}} - 1$ from Lemma 6. Hence, for any m we have

$$\dim(\mathcal{C}_{(n,-1,\delta,b)}) = n - \left| \bigcup_{i=b}^{b+\delta-2} C_{1+2i}^{2n} \right| = n - m |CL(1 + 2b, 1 + 2(b + \delta - 2))|.$$

Next we prove that there exists $t \in Z$ such that $1 + 2b \leq (1 + 2a)q^t \leq 1 + 2(b + \delta - 2)$ for any $1 + 2a$ with $1 \leq 1 + 2a \leq 1 + 2b$.

Let $t \in Z$ satisfy $(1 + 2a)q^{t-1} < 1 + 2b \leq (1 + 2a)q^t$, since $1 + 2a \leq 1 + 2b$. It follows that

$$1 + 2b \leq (1 + 2a)q^t < (1 + 2b)q \leq \left\lfloor \frac{b + \delta - 2}{q} \right\rfloor q \leq b + \delta - 2.$$

This means $CL(1, 1 + 2(b + \delta - 2)) = CL(1 + 2b, 1 + 2(b + \delta - 2))$, then $\mathcal{C}_{(n,-1,\delta,b)} = \mathcal{C}_{(n,-1,b+\delta,0)}$.

According to Lemma 7, we can get $|CL(1, 1 + 2(b + \delta - 2))| = \left\lceil \frac{(2\delta + 2b - 3)(q - 1)}{2q} \right\rceil$, where $1 \leq 1 + 2(b + \delta - 2) \leq q^{\frac{m+1}{2}} - 1$ for $2 \nmid m$ and $1 \leq 1 + 2(b + \delta - 2) \leq 2q^{\frac{m}{2}} - 1$ for $2 \mid m$. Thus we complete the proof. \square

Next we give a theorem about the bound of $d(\mathcal{C}_{(n,-1,\delta,1)})$.

Theorem 24. Let $(n, q) = 1$, $q - 1 \mid n$, $k \in [1, q - 1]$, $\delta_a \in \mathbb{Z}^+$ and $\delta_a \mid \frac{n}{q-1}$. Then

- (1) If $2 \nmid k$ and $\delta = \frac{k\delta_a - 1}{2}$, then $\frac{k\delta_a - 1}{2} \leq d(\mathcal{C}_{(n,-1,\delta,1)}) \leq \frac{k+1}{2}\delta_a$.
- (2) If $2 \mid k$ and $\delta = \frac{k\delta_a}{2}$, then $\frac{k\delta_a}{2} \leq d(\mathcal{C}_{(n,-1,\delta,1)}) \leq (\frac{k}{2} + 1)\delta_a$.

Proof. Define β to be the $2n$ -th root of unity in \mathbb{F}_{q^l} that is primitive, where $l = \text{ord}_{2n}(q)$. Clearly, $T = \bigcup_{i=1}^{\delta-1} C_{1+2i}^{2n}$ for code $\mathcal{C}_{(n,-1,\delta,1)}$.

- (1) If $2 \nmid k$ and $2\delta + 1 = k\delta_a$, we denote

$$f(x) = \frac{x^n + 1}{x^{\frac{n}{\delta_a}} + 1} \times \prod_{t=1}^{\frac{k-1}{2}} (x^{\frac{n}{\delta_a(q-1)}} - \beta^{\frac{n(2t-1)}{q-1}}).$$

Since $g_{(n,-1,\delta,1)}(\beta^{1+2i}) = 0$ for $i \in [1, \delta - 1]$, then $g_{(n,-1,\delta,1)}(\beta^{(2t-1)\delta_a}) = 0$ for $t \in [1, \frac{k-1}{2}]$. We can get $f(\beta^{1+2i}) = 0$ for $i \in [1, \delta - 1]$. Then $f(x) \in \mathcal{C}_{(n,-1,\delta,1)}$.

We know that polynomial $\frac{x^n + 1}{x^{\frac{n}{\delta_a}} + 1}$ has δ_a terms in expansion and polynomial $\prod_{t=1}^{\frac{k-1}{2}} (x^{\frac{n}{\delta_a(q-1)}} - \beta^{\frac{n(2t-1)}{q-1}})$ has at most $\frac{k+1}{2}$ terms. Hence, $w_H(f) \leq \frac{k+1}{2}\delta_a$. We know that $d(\mathcal{C}_{(n,-1,\delta,1)}) \geq \delta = \frac{k\delta_a - 1}{2}$ from Lemma 2. Hence, $\frac{k\delta_a - 1}{2} \leq d(\mathcal{C}_{(n,-1,\delta,1)}) \leq \frac{k+1}{2}\delta_a$.

(2) If $2 \mid k$ and $2\delta = k\delta_a$, denote

$$f(x) = \frac{x^n + 1}{x^{\frac{n}{\delta_a}} + 1} \times \prod_{t=1}^{\frac{k}{2}} (x^{\frac{n}{\delta_a(q-1)}} - \beta^{\frac{n(2t-1)}{q-1}}).$$

Similar to (1), then $f(x) \in \mathcal{C}_{(n,-1,\delta,1)}$ and $w_H(f) \leq (\frac{k}{2} + 1)\delta_a$. We know that $d(\mathcal{C}_{(n,-1,\delta,1)}) \geq \delta = \frac{k\delta_a}{2}$ from Lemma 2. Hence, $\frac{k\delta_a}{2} \leq d(\mathcal{C}_{(n,-1,\delta,1)}) \leq (\frac{k}{2} + 1)\delta_a$. \square

Example 25. Let $(q, m) = (3, 4)$, $n = \frac{q^m - 1}{2}$, $k = 1$ and $\delta_a = 10$. Then $\delta = 5$ and the code $\mathcal{C}_{(40,-1,5,1)}$ has parameters $[40, 28, 6]$, this is a best known linear code for these parameters in the Database [9].

4. The parameter of cyclic BCH code of length $n = \frac{q^m - 1}{2}$

Let $n = \frac{q^m - 1}{2}$ with $m \geq 2$. We will study the extended code and dual code of cyclic code.

4.1. Extended code $\bar{\mathcal{C}}$ and dual code \mathcal{C}^\perp

For $\delta_2 \leq \delta \leq \delta_1$, we discuss the weight distribution of $\bar{\mathcal{C}}_{(n,1,\delta,1)}$ and the parameters of $\mathcal{C}_{(n,1,\delta,1)}^\perp$.

Lemma 26. [35] Let $n = \frac{q^m - 1}{2}$. Then

$$\delta_1 = \frac{q^m - 1 - q^{m-1} - q^{\lfloor \frac{m-1}{2} \rfloor}}{2} \quad \text{and} \quad \delta_2 = \frac{q^m - 1 - q^{m-1} - q^{\lfloor \frac{m+1}{2} \rfloor}}{2}.$$

Moreover, $|C_{\delta_1}^n| = \begin{cases} m, & \text{if } 2 \nmid m; \\ \frac{m}{2}, & \text{if } 2 \mid m, \end{cases}$ and $|C_{\delta_2}^n| = m$.

Theorem 27. Let $2 \mid m$ and $\delta_2 + 1 \leq \delta \leq \delta_1$. Then $\bar{\mathcal{C}}_{(n,1,\delta,1)}$ is an $[\frac{q^m - 1}{2} + 1, \frac{m}{2} + 1, \delta_1 + 1]$ extended code and $\mathcal{C}_{(n,1,\delta,1)}^\perp$ is an $[\frac{q^m - 1}{2}, \frac{q^m - 1}{2} - \frac{m}{2} - 1, d]$ dual code, where

$$d = \begin{cases} 3, & \text{if } q = 3 \text{ and } m = 2; \\ 2, & \text{others.} \end{cases}$$

Moreover, Table 1 gives the weight distribution of $\bar{\mathcal{C}}_{(n,1,\delta,1)}$.

Proof. Since 1 and β^{δ_1} are zeroes of parity-check polynomial of \mathcal{C} and are non-conjugate roots of its. From Lemmas 3 and 26, we have

$$\mathcal{C}_{(n,1,\delta,1)} = \{(Tr_q^{q^{\frac{m}{2}}}(a\beta^{-\delta_1 i}) + b)_{i=0}^{n-1} : a \in \mathbb{F}_{q^{\frac{m}{2}}}, b \in \mathbb{F}_q\},$$

its weight distribution is equal to

$$\{\mathbf{c}(a, b) = (Tr_q^{q^{\frac{m}{2}}}(a\alpha^{(q^{\frac{m}{2}} + 1)i}) + b)_{i=0}^{n-1} : a \in \mathbb{F}_{q^{\frac{m}{2}}}, b \in \mathbb{F}_q\}. \quad (1)$$

Note that $\tilde{\mathcal{C}}_{(n,1,\delta,1)} = \langle \tilde{g}_{(n,1,\delta,1)}(x) \rangle$, where $\tilde{g}_{(n,1,\delta,1)}(x) = (x - 1)g_{(n,1,\delta,1)}(x)$ and $\tilde{g}_{(n,1,\delta,1)}(1) = 0$, i.e.,

$$\sum_{i=0}^{n-1} Tr_q^{q^{\frac{m}{2}}}(a\alpha^{(q^{\frac{m}{2}} + 1)i}) = 0.$$

Table 1:

Weight	Frequency
0	1
$\frac{q^m - q^{m-1} - q^{\frac{m}{2}-1} + 1}{2}$	$(q-1)(q^{\frac{m}{2}} - 1)$
$\frac{(q-1)(q^{m-1} + q^{\frac{m}{2}-1})}{2}$	$q^{\frac{m}{2}} - 1$
$\frac{q^m + 1}{2}$	$q - 1$

We have

$$\sum_{i=0}^{n-1} (Tr_q^{q^{\frac{m}{2}}} (a\alpha^{(q^{\frac{m}{2}}+1)^i}) + b) = \sum_{i=0}^{n-1} Tr_q^{q^{\frac{m}{2}}} (a\alpha^{(q^{\frac{m}{2}}+1)^i}) + nb = nb$$

for any codeword $\mathbf{c}(a, b)$ of (1). Since $b \in \mathbb{F}_q$, then $nb = (\frac{q^{m-1}-1}{2}q + \frac{q-1}{2})b = \frac{q-1}{2}b$. By the definition of extended code, we know $\bar{\mathbf{c}}(a, b) = (\mathbf{c}(a, b), -\frac{q-1}{2}b)$ is the codeword of $\bar{\mathcal{C}}_{(n,1,\delta_1,1)}$, where $a \in \mathbb{F}_{q^{\frac{m}{2}}}$ and $b \in \mathbb{F}_q$. Clearly, $wt(\bar{\mathbf{c}}(a, b)) = wt(\mathbf{c}(a, b)) + \varepsilon$ where

$$\varepsilon = \begin{cases} 0, & \text{if } b = 0; \\ 1, & \text{if } b \neq 0. \end{cases}$$

Thus, we give the weight distribution in Table 1.

For code $\mathcal{C}_{(n,1,\delta_1,1)}^\perp$ with $\delta_2 + 1 \leq \delta \leq \delta_1$, we have $T^\perp = (\mathbb{Z}_n \setminus T)^{-1} = C_0^n \cup C_{CL(n-\delta_1)}^n$. Then $d(\mathcal{C}_{(n,1,\delta_1,1)}^\perp) \geq 2$.

Since $n - \delta_1 = \frac{q^{m-1} + q^{\frac{m}{2}-1}}{2}$, then $CL(n - \delta_1) = \frac{q^{\frac{m}{2}} + 1}{2}$ and

$$C_{\frac{q^{\frac{m}{2}}+1}{2}}^n = \left\{ \frac{(1 + q^{\frac{m}{2}})q^i}{2} : 0 \leq i \leq \frac{m}{2} - 1 \right\}.$$

Thus $|C_{\frac{q^{\frac{m}{2}}+1}{2}}^n| = \frac{m}{2}$, and $\dim(\mathcal{C}_{(n,1,\delta_1,1)}^\perp) = n - \frac{m}{2} - 1$.

For $q > 3$, if $d(\mathcal{C}_{(n,1,\delta_1,1)}^\perp) = 3$, we have

$$\sum_{i=0}^1 (q-1)^i \binom{n}{i} > q^{\frac{m}{2}+1}$$

by Lemma 4, this is infeasible. Hence, $d(\mathcal{C}_{(n,1,\delta_1,1)}^\perp) = 2$. For the case of $q = 3$, we can get the result as above. \square

Theorem 28. Let $2 \nmid m$ and $\delta_2 + 1 \leq \delta \leq \delta_1$. Then $\bar{\mathcal{C}}_{(n,1,\delta_1,1)}$ is an $[\frac{q^m-1}{2} + 1, m + 1, \delta_1 + 1]$ extended code and $\mathcal{C}_{(n,1,\delta_1,1)}^\perp$ is an $[\frac{q^m-1}{2}, \frac{q^m-1}{2} - m - 1, d]$ dual code, where

$$\begin{cases} 3 \leq d \leq 4, & \text{if } q = 3; \\ d = 3, & \text{if } q > 3. \end{cases}$$

Moreover, Table 2 gives the weight distribution of $\bar{\mathcal{C}}_{(n,1,\delta_1,1)}$.

Table 2:

Weight	Frequency
0	1
$\frac{q^m - q^{m-1} - q^{\frac{m-1}{2}} + 1}{2}$	$\frac{(q-1)(q^m-1)}{2}$
$\frac{q^m - q^{m-1}}{2}$	$q^m - 1$
$\frac{q^m - q^{m-1} + q^{\frac{m-1}{2}} + 1}{2}$	$\frac{(q-1)(q^m-1)}{2}$
$\frac{q^{m+1}}{2}$	$q - 1$

Proof. If $2 \nmid m$, using the same approach as Theorem 27 for code $\mathcal{C}_{(n,1,\delta,1)}$, its weight distribution is equal to

$$\{\mathbf{c}(a, b) = (Tr_q^{q^m}(a\alpha^{(q^{\frac{m+1}{2}}+1)i}) + b)_{i=0}^{n-1} : a \in \mathbb{F}_{q^m}, b \in \mathbb{F}_q\} \quad (2)$$

and

$$\sum_{i=0}^{n-1} Tr_q^{q^m}(a\alpha^{(q^{\frac{m+1}{2}}+1)i}) = 0.$$

We have

$$\sum_{i=0}^{n-1} (Tr_q^{q^m}(a\alpha^{(q^{\frac{m+1}{2}}+1)i}) + b) = \sum_{i=0}^{n-1} Tr_q^{q^m}(a\alpha^{(q^{\frac{m+1}{2}}+1)i}) + nb = \frac{q-1}{2}b$$

for any codeword $\mathbf{c}(a, b)$ of (2). Similar to Theorem 27, we directly give the weight distribution in Table 2.

For code $\mathcal{C}_{(n,1,\delta,1)}^\perp$ with $\delta_2 + 1 \leq \delta \leq \delta_1$, we have $T^\perp = (\mathbb{Z}_n \setminus T)^{-1} = C_0^n \cup C_{CL(n-\delta_1)}^n$. Thus $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) \geq 2$.

Since $n - \delta_1 = \frac{q^{m-1} + q^{\frac{m-1}{2}}}{2}$, then $CL(n - \delta_1) = \frac{q^{\frac{m-1}{2}} + 1}{2}$ and

$$C_{\frac{q^{\frac{m-1}{2}} + 1}{2}}^n = \left\{ \frac{q^l + q^{\frac{m-1}{2}+l}}{2} : 1 \leq l \leq \frac{m-1}{2} \right\} \cup \left\{ \frac{q^l + q^{l-\frac{m+1}{2}}}{2} : \frac{m+1}{2} \leq l \leq m \right\}.$$

Set $m_1 = \frac{q^{\frac{m-1}{2}} + 1}{2}$, thus $|C_{m_1}^n| = m$. Then $\dim(\mathcal{C}_{(n,1,\delta,1)}^\perp) = n - m - 1$.

Suppose $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) = 2$, which means that there exist $a(x) = a_0 + a_1x + a_2x^2 + \cdots + a_ix^i + \cdots + a_jx^j + \cdots + a_{n-1}x^{n-1} \in \mathcal{C}_{(n,1,\delta,1)}^\perp$ and $b(x) = b_0 + b_1x + b_2x^2 + \cdots + b_ix^i + \cdots + b_jx^j + \cdots + b_{n-1}x^{n-1} \in \mathcal{C}_{(n,1,\delta,1)}^\perp$, where $a_t = b_t$ for any $t \in [0, n-1] \setminus \{i, j\}$ and $0 \leq i < j \leq n-1$. Since $a(1) = b(1) = 0$ and $a(\beta^{m_1}) = b(\beta^{m_1}) = 0$, i.e.,

$$\begin{cases} (a_i - b_i) + (a_j - b_j) = 0, \\ (a_i - b_i)\beta^{im_1} + (a_j - b_j)\beta^{jm_1} = 0, \end{cases}$$

which equals to

$$\beta^{(i-j)m_1} = 1 \iff (i-j)\left(\frac{q^{\frac{m-1}{2}} + 1}{2}\right) \equiv 0 \pmod{\frac{q^m - 1}{2}}. \quad (3)$$

Since $2 \nmid m$ and $\gcd(\frac{m-1}{2}, m) = 1$, then $\gcd(q^{\frac{m-1}{2}} + 1, q^m - 1) = 2$. Thus Eq.3 holds $\Leftrightarrow i = j$, a contradiction. Thus $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) \geq 3$.

By Lemma 4, if $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) = 5$, then $\sum_{i=0}^2 (q-1)^i \binom{n}{i} > q^{m+1}$, this is infeasible, i.e., $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) \leq 4$. For $q = 3$, if $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) = 4$, we have $\sum_{i=0}^1 (3-1)^i \binom{n-1}{i} \leq 3^m$ satisfying Lemma 4. Thus $3 \leq d(\mathcal{C}_{(n,1,\delta,1)}^\perp) \leq 4$. For $q > 3$, if $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) = 4$, then $\sum_{i=0}^1 (q-1)^i \binom{n-1}{i} > q^m$, this is infeasible, i.e., $d(\mathcal{C}_{(n,1,\delta,1)}^\perp) = 3$. \square

Example 29. Let $(q, m) = (3, 3)$. Then $\bar{\mathcal{C}}_{(n,1,\delta_1,1)}$ is an $[14, 4, 8]$ code and weight enumerator is $1 + 26z^8 + 26z^9 + 26z^{11} + 2z^{14}$, which has the best known parameters for linear code in Database [9]. Let $(q, m) = (5, 3)$. Then $\bar{\mathcal{C}}_{(n,1,\delta_1,1)}$ is an $[63, 4, 48]$ code and weight enumerator is $1 + 248z^{48} + 124z^{50} + 248z^{53} + 4z^{63}$, in Database [9], the best known linear code over \mathbb{F}_5 , having length 63 and dimension 4, holds a minimum distance of 49.

Theorem 30. Let $2 \nmid m$ and $\delta = \delta_2$. Then $\bar{\mathcal{C}}_{(n,1,\delta_2,1)}$ is an $[\frac{q^m-1}{2} + 1, 2m + 1, \delta_2 + 1]$ extended code and $\mathcal{C}_{(n,1,\delta_2,1)}^\perp$ is an $[\frac{q^m-1}{2}, \frac{q^m-1}{2} - 2m - 1, d]$ dual code, where

$$\begin{cases} 3 \leq d \leq 6, & \text{if } q = 3; \\ 3 \leq d \leq 5, & \text{if } 5 \leq q \leq 9; \\ 3 \leq d \leq 4, & \text{if } q \geq 11. \end{cases}$$

Moreover, Table 3 gives the weight distribution of $\bar{\mathcal{C}}_{(n,1,\delta_2,1)}$.

Proof. According to the Theorem 8 in [35], we have

$$\mathcal{C}_{(n,1,\delta_2,1)} = \{(Tr_{q^m/q}(a\beta^{-\delta_1 i} + b\beta^{-\delta_2 i}) + c)_{i=0}^{n-1} : a, b \in \mathbb{F}_{q^m}, c \in \mathbb{F}_q\},$$

its weight distribution is equal to

$$\left\{ \mathbf{v}_4(a, b, c) = Tr_q^{q^m} (a\alpha^{(q^{\frac{m-1}{2}+1}i)} + b\alpha^{(q^{\frac{m-3}{2}+1}i)} + c)_{i=0}^{n-1} : a, b \in \mathbb{F}_{q^m}, c \in \mathbb{F}_q \right\}. \quad (4)$$

Similar to Theorem 27, we have

$$\sum_{i=0}^{n-1} (Tr_q^{q^m} (a\alpha^{(q^{\frac{m-1}{2}+1}i)} + b\alpha^{(q^{\frac{m-3}{2}+1}i)} + c) = \frac{q-1}{2}c$$

for any codeword $\mathbf{v}_4(a, b, c)$ of (4). And $\bar{\mathbf{v}}_4(a, b, c) = (\mathbf{v}_4(a, b, c), c)$ is the codeword of $\bar{\mathcal{C}}_{(n,1,\delta_2,1)}$, where $a, b \in \mathbb{F}_{q^m}, c \in \mathbb{F}_q$. Then $wt(\bar{\mathbf{v}}_4(a, b, c)) = wt(\mathbf{v}_4(a, b, c)) + \varepsilon$ where

$$\varepsilon = \begin{cases} 0, & \text{if } c = 0; \\ 1, & \text{if } c \neq 0. \end{cases}$$

Thus, we give the weight distribution in Table 3.

For code $\mathcal{C}_{(n,1,\delta_2,1)}^\perp$, we have $T^\perp = (\mathbb{Z}_n \setminus T)^{-1} = C_0^n \cup C_{CL(n-\delta_1)}^n \cup C_{CL(n-\delta_2)}^n$. Then $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) \geq 2$.

Note that $n - \delta_2 = \frac{q^{m-1} + q^{\frac{m+1}{2}}}{2}$, then $CL(n - \delta_2) = \frac{q^{\frac{m-3}{2}+1}}{2}$ and

$$C_{\frac{q^{\frac{m-3}{2}+1}}{2}}^m = \left\{ \frac{q^l + q^{\frac{m-3}{2}+l}}{2} : 1 \leq l \leq \frac{m+3}{2} \right\} \cup \left\{ \frac{q^l + q^{l-\frac{m+3}{2}}}{2} : \frac{m+5}{2} \leq l \leq m \right\}.$$

Table 3:

Weight	Frequency
0	1
$\frac{q^m - q^{m-1} - q^{\frac{m+1}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{m-1} - 1)}{2(q+1)}$
$\frac{(q-1)(q^{m-1} - q^{\frac{m-1}{2}})}{2}$	$\frac{(q^m - 1)(q^{m-1} + q^{\frac{m-1}{2}})}{2}$
$\frac{q^m - q^{m-1} - q^{\frac{m-1}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{m+2} - q^m - q^{m-1} - q^{\frac{m+3}{2}} + q^{\frac{m-1}{2}} + q^2)}{2(q+1)}$
$\frac{q^m - q^{m-1}}{2}$	$(q^m - 1)(q^m - q^{m-1} + 1)$
$\frac{q^m - q^{m-1} + q^{\frac{m-1}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{m+2} - q^m - q^{m-1} + q^{\frac{m+3}{2}} - q^{\frac{m-1}{2}} + q^2)}{2(q+1)}$
$\frac{(q-1)(q^{m-1} + q^{\frac{m-1}{2}})}{2}$	$\frac{(q^m - 1)(q^{m-1} - q^{\frac{m-1}{2}})}{2}$
$\frac{q^m - q^{m-1} + q^{\frac{m+1}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{m-1} - 1)}{2(q+1)}$
$\frac{q^m + 1}{2}$	$q - 1$

Set $m_2 = \frac{q^{\frac{m-3}{2}} + 1}{2}$, thus $|C_{m_2}^\perp| = m$, i.e., $\dim(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = n - 2m - 1$.

Suppose $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = 2$, then we have $a(x) = a_0 + a_1x + a_2x^2 + \cdots + a_ix^i + \cdots + a_jx^j + \cdots + a_{n-1}x^{n-1} \in \mathcal{C}_{(n,1,\delta_2,1)}^\perp$ and $b(x) = b_0 + b_1x + b_2x^2 + \cdots + b_ix^i + \cdots + b_jx^j + \cdots + b_{n-1}x^{n-1} \in \mathcal{C}_{(n,1,\delta_2,1)}^\perp$, where $a_t = b_t$ for any $t \in [0, n-1] \setminus \{i, j\}$ and $0 \leq i < j \leq n-1$. Since $a(1) = b(1) = 0$, $a(\beta^{m_1}) = b(\beta^{m_1}) = 0$ and $a(\beta^{m_2}) = b(\beta^{m_2}) = 0$, i.e.

$$\begin{cases} (a_i - b_i) + (a_j - b_j) = 0, \\ (a_i - b_i)\beta^{im_1} + (a_j - b_j)\beta^{jm_1} = 0, \\ (a_i - b_i)\beta^{im_2} + (a_j - b_j)\beta^{jm_2} = 0, \end{cases}$$

which equals to

$$\beta^{(i-j)m_l} = 1 \Leftrightarrow (i-j)m_l \equiv 0 \pmod{\frac{q^m - 1}{2}}$$

for $l = 1, 2$. Obviously, we can get $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) \geq 3$.

For $3 \leq q \leq 9$, if $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = 7$, then $\sum_{i=0}^3 (q-1)^i \binom{n}{i} > q^{2m+1}$ by Lemma 4, this is infeasible, i.e., $3 \leq d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) \leq 6$. For $q \geq 11$, if $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = 5$, we have $\sum_{i=0}^2 (q-1)^i \binom{n}{i} > q^{2m+1}$ by Lemma 4, this is infeasible, i.e., $3 \leq d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) \leq 4$. The cases of $5 \leq q \leq 9$ and $q = 3$ are similar and we omit it here. \square

Theorem 31. *Let $2 \mid m$ and $\delta = \delta_2$. Then $\overline{\mathcal{C}}_{(n,1,\delta_2,1)}$ is an $[\frac{q^m-1}{2} + 1, \frac{3m}{2} + 1, \delta_2 + 1]$ extended code and $\mathcal{C}_{(n,1,\delta,1)}^\perp$ is an $[\frac{q^m-1}{2}, \frac{q^m-1}{2} - \frac{3m}{2} - 1, 2 \leq d \leq 4]$ dual code except for $(q, m) = (3, 2)$. Moreover, Table 4 gives the weight distribution of $\overline{\mathcal{C}}_{(n,1,\delta_2,1)}$.*

Proof. If $2 \mid m$, according to the Theorem 8 in [35] for $\mathcal{C}_{(n,1,\delta_2,1)}$, its weight distribution is equal to

$$\left\{ \mathbf{v}_5(a, b, c) = \text{Tr}_q^{\frac{m}{2}} (\alpha \alpha^{(q^{\frac{m}{2}} + 1)^i} + b \alpha^{(q^{\frac{m-2}{2}} + 1)^i} + c)_{i=0}^{n-1} : a \in \mathbb{F}_{q^{\frac{m}{2}}}, b \in \mathbb{F}_{q^m}, c \in \mathbb{F}_q \right\}. \quad (5)$$

With the same way as Theorem 30, we directly give the weight distribution in Table 4.

Table 4:

Weight	Frequency
0	1
$\frac{q^m - q^{m-1} - q^{\frac{m}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{\frac{m+2}{2}} + q^{\frac{m-2}{2}} - 2)}{2(q+1)}$
$\frac{(q-1)(q^{m-1} - q^{\frac{m}{2}-1})}{2}$	$\frac{(q^m - 1)(q^{\frac{m+2}{2}} + q)}{2(q+1)}$
$\frac{q^m - q^{m-1} - q^{\frac{m-2}{2}} + 1}{2}$	$\frac{(q^{\frac{m}{2}} - 1)(q^{m+1} - 2q^m + q)}{2}$
$\frac{q^m - q^{m-1}}{2}$	$(q^m - 1)q^{\frac{m-2}{2}}$
$\frac{q^m - q^{m-1} + q^{\frac{m-2}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{\frac{m+2}{2}} + q)(q-1)}{2(q+1)}$
$\frac{(q-1)(q^{m-1} + q^{\frac{m}{2}-1})}{2}$	$\frac{(q^{\frac{m+2}{2}} - q)(q^m - 2q^{m-1} + 1)}{2(q-1)}$
$\frac{q^m - q^{m-1} + q^{\frac{m}{2}} + 1}{2}$	$\frac{(q^m - 1)(q^{\frac{m}{2}} - q^{\frac{m-2}{2}})}{2}$
$\frac{(q-1)(q^{m-1} + q^{\frac{m}{2}})}{2}$	$\frac{(q^m - 1)(q^{\frac{m-2}{2}} - 1)}{(q^2 - 1)}$
$\frac{q^m + 1}{2}$	$q - 1$

Note that $n - \delta_2 = \frac{q^{m-1} + q^{\frac{m}{2}}}{2}$, then $CL(n - \delta_2) = \frac{q^{\frac{m}{2}-1} + 1}{2}$ and

$$C_{\frac{q^{\frac{m}{2}-1} + 1}{2}}^n = \left\{ \frac{q^l + q^{\frac{m}{2} + l - 1}}{2} : 1 \leq l \leq \frac{m}{2} + 1 \right\} \cup \left\{ \frac{q^l + q^{l - \frac{m}{2} - 1}}{2} : \frac{m}{2} + 2 \leq l \leq m \right\}.$$

Thus $|C_{\frac{q^{\frac{m}{2}-1} + 1}{2}}^n| = m$. Then $\dim(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = n - |T^\perp| = n - \frac{3m}{2} - 1$.

For $q = 3$ and $m = 2$, we can get $\dim(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = 0$. Otherwise, if $d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) = 5$, then $\sum_{i=0}^2 (q-1)^i \binom{n}{i} > q^{\frac{3m}{2}+1}$ by Lemma 4, this is impossible, i.e., $2 \leq d(\mathcal{C}_{(n,1,\delta_2,1)}^\perp) \leq 4$. \square

Example 32. Let $q = 3$, $m = 4$. Then $\bar{\mathcal{C}}_{(n,1,\delta_2,1)}$ is an $[41, 7, 23]$ linear code and weight enumerator is $1 + 280z^{23} + 300z^{24} + 336z^{26} + 240z^{27} + 600z^{29} + 168z^{30} + 240z^{32} + 20z^{36} + 2z^{40}$, which has the best known parameters for linear code in Database [9]. Let $q = 3$, $m = 3$. Then $\bar{\mathcal{C}}_{(n,1,\delta_2,1)}$ is an $[14, 7, 5]$ code and weight enumerator is $1 + 26z^5 + 156z^6 + 624z^8 + 494z^9 + 780z^{11} + 78z^{12} + 26z^{13} + 2z^{14}$, in Database [9], the best known linear code over \mathbb{F}_3 , having length 14 and dimension 7, holds a minimum distance of 6.

4.2. Dually-BCH code

If \mathcal{C} and \mathcal{C}^\perp are BCH codes with respect to β , then \mathcal{C} is a dually-BCH code [8]. For $b = 2$, we present the necessary and sufficient condition for $\mathcal{C}_{(n,1,\delta,b)}$ to be recognized as a dually-BCH code.

Theorem 33. Let $q \geq 3$ and $m \geq 3$ be odd. Then $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code iff $\delta_1 - 1 < \delta \leq n - 1$.

Proof. For $\mathcal{C}_{(n,1,\delta,2)}$ with $\delta \in [2, n - 1]$, we have $T = C_2^n \cup C_3^n \cup \dots \cup C_\delta^n$. Since $0 \notin T$, then $C_0^n \subset T^\perp$. Then $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code \Leftrightarrow there exists $t \geq 1$ such that $T^\perp = C_0^n \cup C_1^n \cup \dots \cup C_{t-1}^n$.

For $q \leq \delta \leq n - 1$, we have $C_1^n = C_q^n$, then $T = C_1^n \cup C_2^n \cup \dots \cup C_\delta^n$. According to Theorem 30 of [29], we deduce that $\mathcal{C}_{(n,1,\delta,2)}$ will be a dually-BCH code iff $\delta_1 - 1 < \delta \leq n - 1$.

For $2 \leq \delta \leq q-1$, we know that $|C_i^n| = m$ for any $1 \leq i \leq q-1$ and $q \nmid i$ by Lemma 5. Since $C_1^n \notin T$ and $C_2^n \in T$, then $\dim(\mathcal{C}_{(n,1,\delta,2)}) \in [m, n-m]$. Note that $\frac{q^{\frac{m-1}{2}}+1}{2} \in \Gamma_n$. We consider two cases.

(1) If $m \geq 5$, we have $\frac{q^{\frac{m-1}{2}}+1}{2} > q-1 > \delta$, then $\frac{q^{\frac{m-1}{2}}+1}{2} \notin T$. Since

$$CL(n - \frac{q^{\frac{m-1}{2}}+1}{2}) = \delta_1,$$

then $C_{\delta_1}^n \subset T^\perp$. If $T^\perp = C_0^n \cup C_1^n \cup \dots \cup C_{\delta_1}^n$, a contradiction.

(2) If $m = 3$, we discuss this in two parts.

For $q > 3$. If $2 \leq \delta \leq \frac{q-1}{2}$, then $\frac{q^{\frac{m-1}{2}}+1}{2} > \frac{q-1}{2} > \delta$, i.e., $\frac{q^{\frac{m-1}{2}}+1}{2} \notin T$. By the same way as (1), $\mathcal{C}_{(n,1,\delta,2)}^\perp$ is not a BCH code. If $\frac{q+1}{2} \leq \delta \leq q-1$, then $\frac{q+1}{2} \in T$. Note that $C_1^n \notin T$ and $T^\perp = (\mathbb{Z}_n \setminus T)^{-1}$, then $CL(n-1) = CL(\frac{q^3-3}{2}) = \frac{q^3-2q^2-1}{2} = \delta_2 \in T^\perp$. Thus, $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code $\Rightarrow C_0^n \cup C_1^n \cup \dots \cup C_{\delta_2}^n \in T^\perp$, i.e., $\dim(\mathcal{C}_{(n,1,\delta,2)}^\perp) \leq m$. Since $q > 3$, and $2, \frac{q+1}{2} \in T$ are coset leader by Lemma 5, then $\dim(\mathcal{C}_{(n,1,\delta,2)}) \leq n-2m$, a contradiction.

For $q = 3$, then $\delta = 2$. We can easily conclude that $\mathcal{C}_{(13,1,2,2)}$ does not satisfy the conditions for being a dually-BCH code. \square

Theorem 34. *Let $q \geq 3$ and $m \geq 2$ be even. Then $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code iff $\delta_1 - 1 < \delta \leq n-1$.*

Proof. For $\mathcal{C}_{(n,1,\delta,2)}$ with $\delta \in [2, n-1]$, we have $T = C_2^n \cup C_3^n \cup \dots \cup C_\delta^n$. Since $0 \notin T$, then $C_0^n \subset T^\perp$. Then $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code \Leftrightarrow there exists $t \geq 1$ such that $T^\perp = C_0^n \cup C_1^n \cup \dots \cup C_{t-1}^n$.

For $q \leq \delta \leq n-1$, we have $C_1^n = C_q^n$, then $T = C_1^n \cup C_2^n \cup \dots \cup C_\delta^n$. According to Theorem 30 of [29], we deduce that $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code iff $\delta_1 - 1 < \delta \leq n-1$.

For $2 \leq \delta \leq q-1$, we know that $|C_i^n| = m$ for any $1 \leq i \leq q-1$ and $q \nmid i$ by Lemma 6. Since $C_1^n \notin T$ and $C_2^n \in T$, then $\dim(\mathcal{C}_{(n,1,\delta,2)}) \in [m, n-m]$. Note that $\frac{q^{\frac{m}{2}}+1}{2} \in \Gamma_n$. We consider two cases:

(1) If $m \geq 4$, we have $\frac{q^{\frac{m}{2}}+1}{2} > q-1 > \delta$, then $\frac{q^{\frac{m}{2}}+1}{2} \notin T$. Since

$$CL(n - \frac{q^{\frac{m}{2}}+1}{2}) = \delta_1,$$

then $C_{\delta_1}^n \subset T^\perp$. If $T^\perp = C_0^n \cup C_1^n \cup \dots \cup C_{\delta_1}^n$, a contradiction.

(2) If $m = 2$, we discuss this in two parts.

For $q > 3$. If $2 \leq \delta \leq \frac{q-1}{2}$, then $\frac{q^{\frac{m}{2}}+1}{2} > \frac{q-1}{2} > \delta$, i.e., $\frac{q^{\frac{m}{2}}+1}{2} \notin T$. By the same way as (1), $\mathcal{C}_{(n,1,\delta,2)}^\perp$ is not a BCH code. If $\frac{q+1}{2} \leq \delta \leq q-1$, then $\frac{q+1}{2} \in T$. Note that $C_1^n \notin T$ and $T^\perp = (\mathbb{Z}_n \setminus T)^{-1}$, then $CL(n-1) = CL(\frac{q^2-3}{2}) = \frac{q^2-2q-1}{2} = \delta_2 \in T^\perp$. Thus, $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code $\Rightarrow C_0^n \cup C_1^n \cup \dots \cup C_{\delta_2}^n \in T^\perp$, i.e., $\dim(\mathcal{C}_{(n,1,\delta,2)}^\perp) \leq m$. Since $q > 3$, and $2, \frac{q+1}{2} \in T$ are coset leader by Lemma 6, then $\dim(\mathcal{C}_{(n,1,\delta,2)}) \leq n-2m$, a contradiction.

For $q = 3$, then $\delta = 2$. Thus we get a code $\mathcal{C}_{(4,1,2,2)}$, which is not a dually-BCH code obviously. \square

5. Conclusions

The principal contributions outlined in this paper are:

- For the negacyclic BCH code of length $n = \frac{q^m-1}{2}$, we get $\dim(\mathcal{C}_{(n,-1, \lceil \frac{\delta+1}{2} \rceil, 0)})$ for $\delta = a\frac{q^m-1}{q-1}$, $aq^{m-1} - 1 (1 \leq a < \frac{q-1}{2})$ (see Lemmas 8, 10) and $\delta = a\frac{q^m-1}{q-1} + b\frac{q^m-1}{q^2-1}$, $aq^{m-1} + (a+b)q^{m-2} - 1$ ($2 \mid m, 1 \leq a+b \leq q-1, \lceil \frac{q-a-2}{2} \rceil \geq 1$) (see Lemmas 11, 13). And we get $\dim(\mathcal{C}_{(n,-1,\delta,0)})$ with few nonzeros (see Theorems 17, 19, 21) and $\dim(\mathcal{C}_{(n,-1,\delta,b)})$ with $b \neq 0$ are settled (see Theorem 23).
- For the cyclic BCH code of length $n = \frac{q^m-1}{2}$, we give the weight distribution of extended code $\bar{\mathcal{C}}_{(n,1,\delta,1)}$ and the parameters of dual code $\mathcal{C}_{(n,1,\delta,1)}^\perp$ for $\delta_2 \leq \delta \leq \delta_1$ (see Theorems 27, 28, 30, 31). Moreover, we present the necessary and sufficient condition for $\mathcal{C}_{(n,1,\delta,2)}$ is a dually-BCH code (see Theorems 33, 34).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (No.U21A20428 and 12171134).

References

- [1] Aly, S.A., Klappenecker, A., Sarvepalli, P.K.: On quantum and classical BCH codes. *IEEE Trans. Inf. Theory* 53(3), 1183–1188 (2007)
- [2] Cherchem, A., Jamous, A., Liu, H., Maouche, Y.: Some new results on dimension and Bose distance for various classes of BCH codes. *Finite Fields Appl* 65, 101673 (2020)
- [3] Ding, C.: Parameters of several classes of BCH codes. *IEEE Trans. Inf. Theory* 61(10), 5322–5330 (2015)
- [4] Ding, C., Du, X., Zhou, Z.: The Bose and minimum distance of a class of BCH codes. *IEEE Trans. Inf. Theory* 61(5), 2351–2356 (2015)
- [5] Ding, C., Fan, C., Zhou, Z.: The dimension and minimum distance of two classes of primitive BCH codes. *Finite Fields Appl* 45, 237–263 (2017)
- [6] Ding, C., Li, C.: BCH cyclic codes. *Discrete Math* 347, 113918 (2024)

- [7] Fang, W., Wen, J., Fu, F.: A q -polynomial approach to constacyclic codes. *Finite Fields Appl* 47, 161–182 (2017)
- [8] Gong, B., Ding, C., Li, C.: The dual codes of several classes of BCH codes. *IEEE Trans. Inform. Theory* 68(2), 953–964 (2021)
- [9] Grassl, M.: Bounds on the minimum distance of linear codes and quantum codes. Online available at <http://www.codetables.de>
- [10] Guo, G., Li, R., Liu, Y., Wang, J.: A family of negacyclic BCH codes of length $n = \frac{q^{2m}-1}{2}$. *Cryptogr. Commun* 12, 187–203 (2020)
- [11] Huffman, W.C., Pless, V.: *Fundamentals of error-correcting codes*. Cambridge University Press, Cambridge (2010)
- [12] Krasikov, I., Litsyn, S.: On the distance distributions of BCH codes and their duals. *Des. Codes Cryptogr* 23(2), 223–232 (2001)
- [13] Krishna, A., Sarwate, D.V.: Pseudocyclic maximum-distance-separable codes. *IEEE Trans. Inf. Theory* 36(4), 880–884 (1990)
- [14] Li, S.: The minimum distance of some narrow-sense primitive BCH codes. *SIAM J. Discrete Math* 31(4), 2530–2569 (2017)
- [15] Li, C., Ding, C., Li, S.: LCD cyclic codes over finite fields. *IEEE Trans. Inf. Theory* 63(7), 4344–4356 (2017)
- [16] Li, S., Ding, C., Xiong, M., Ge, G.: Narrow-sense BCH codes over $\text{GF}(q)$ with length $n = \frac{q^m-1}{q-1}$. *IEEE Trans. Inf. Theory* 63(11), 7219–7236 (2017)
- [17] Li, S., Li, C., Ding, C., Liu, H.: Two families of LCD BCH codes. *IEEE Trans. Inf. Theory* 63(9), 5699–5717 (2017)
- [18] Li, F., Yue, Q., Liu, F.: The weight distribution of constacyclic codes. *Adv. Math. Commun* 11(3), 471–480 (2017)
- [19] Ling, X., Mesnager, S., Qi, Y., Tang, C.: A class of narrow-sense BCH codes over \mathbb{F}_q of length $\frac{q^m-1}{2}$. *Des. Codes Cryptogr* 88(2), 413–427 (2020)
- [20] Liu, H., Ding, C., Li, C.: Dimensions of three types of BCH codes over $\text{GF}(q)$. *Discrete Math* 340(8), 1910–1927 (2017)
- [21] Luo, J., Feng, K.: On the weight distributions of two classes of cyclic codes. *IEEE Trans. Inf. Theory* 54(12), 5332–5344 (2008)
- [22] Pang, B., Zhu, S., Kai, X.: Five families of the narrow-sense primitive BCH codes over finite fields. *Des. Codes Cryptogr* 89(12), 2679–2696 (2021)

- [23] Su, Y., Kai, X.: A family of cyclic codes with two zeros. *J. Appl. Math. Comput* 69, 3231–3243 (2023)
- [24] Sun, Z., Ding, C.: Several families of ternary negacyclic codes and their duals. *IEEE Trans. Inf. Theory* 70(5), 3226–3241 (2024)
- [25] Sun, Z., Zhu, S., Wang, L.: A class of constacyclic BCH codes. *Cryptography and Communications* 12, 265–284 (2020)
- [26] Wang, L., Lu, D., Zhu, S.: Parameters of some BCH codes over \mathbb{F}_q of length $\frac{q^m-1}{2}$. *Adv. Math. Commun* (2023)
- [27] Wang, X., Sun, Z., Ding, C.: Two families of negacyclic BCH codes. *J. Des. Codes Crypt* 91, 2395–2420 (2023)
- [28] Wang, X., Wang, J., Li, C., Wu, Y.: Two classes of narrow-sense BCH codes and their duals. *IEEE Trans. Inf. Theory* 70(1), 131–144 (2024)
- [29] Wang, X., Xiao, C., Zheng, D.: The duals of narrow-sense BCH codes with length $\frac{q^m-1}{\lambda}$. [arXiv:2312.05474 \[cs.IT\]](https://arxiv.org/abs/2312.05474).
- [30] Wu, P., Li, C., Peng, W.: On some cyclic codes of length $\frac{q^{2m}-1}{q+1}$. *Finite Fields Appl* 60, 101581 (2019)
- [31] Yue, D., Feng, Z.: Minimum cyclotomic coset representatives and their applications to BCH codes and Goppa codes. *IEEE Trans. Inf. Theory* 46(7), 2625–2628 (2000)
- [32] Yue, D., Hu, Z.: On the dimension and minimum distance of BCH codes over $\text{GF}(q)$. *J. Electronics (China)* 13(3), 216–221 (1996)
- [33] Zhang, Y., Liu, L., Xie, X.: Two types of negacyclic BCH codes. [arXiv:2307.12762 \[cs.IT\]](https://arxiv.org/abs/2307.12762)
- [34] Zhu, H., Li, J., Zhu, S.: Parameters of two classes of negacyclic BCH codes. *J. Appl. Math. Comput* 69(6), 4353–4380 (2023)
- [35] Zhu, S., Sun, Z., Kai, X.: A class of narrow-sense BCH codes. *IEEE Trans. Inf. Theory* 65(8), 4699–4714 (2019)