

Towards a characterization of the inverse systems of complete intersections. ^{*}

J. Elias [†]

May 31, 2024

Abstract

In this paper we give conditions on a homogeneous polynomial for which the associated graded Artin algebra is a complete intersection.

1 Introduction

We denote by $S = \mathbf{k}[x_0, \dots, x_n]$ the polynomial ring on the variables x_0, \dots, x_n over an algebraically closed field \mathbf{k} of characteristic zero. We denote the homogeneous maximal ideal of S by $\mathfrak{m} = (x_0, \dots, x_n)$.

Macaulay's duality, instance of Matlis' duality, establish a reversing one-to-one correspondence between the Artin Gorenstein quotients $A = S/I$ and their inverse system $I^\perp = \langle H \rangle$, see Proposition 1.1. In this paper we address the following problem: characterize the polynomials H such that its Macaulay's dual $I = \text{Ann}_R \langle H \rangle$ is a complete intersection, see [8, Chapter 9, L]. Very few instances of this problem are known. For example, if H is a monomial then $\text{Ann}_R \langle H \rangle$ is a monomial complete intersection. In [7] the case of quadratic complete intersections is considered; see [7] and its reference list for some other known cases.

In this paper we give explicit conditions on a homogeneous polynomial H for which $I = \text{Ann}_R \langle H \rangle$ is a complete intersection, Proposition 2.2. The key idea of this paper is to consider the resultant of a family of homogeneous forms $\underline{f} = (f_0, \dots, f_n)$ and the determinant of its jacobian matrix. Recall that the resultant of a family of homogeneous forms is the generator of a closed projective hypersurface, [4]. In the case considered in this paper, we get a locally constructible set, Proposition 2.4.

In the second section we characterize the polynomials H such that its Macaulay's dual $I = \text{Ann}_R \langle H \rangle$ is a complete intersection, Proposition 2.2. In the third section we apply this result to short algebras.

^{*}2010 *Mathematics Subject Classification*. Primary 13H10; Secondary 13H15; 14C05

Key words and Phrases: Number of generators, Artin Gorenstein local rings, Inverse system.

[†]Partially supported by PID2022-137283NB-C22.

PRELIMINARIES. Next, we fix some notations and we recall the basic results on Macaulay inverse systems, see [8] for more results on inverse systems.

Let $\Gamma = \mathbf{k}[y_0, \dots, y_n]$ be the polynomial ring on the variables y_0, \dots, y_n . This polynomial ring can be considered a S -module by derivation that we denote by \circ , defined by

$$\begin{aligned} \circ : S \times \Gamma &\longrightarrow \Gamma \\ (f, g) &\longrightarrow f \circ g = f(\partial_{y_1}, \dots, \partial_{y_n})(g) \end{aligned}$$

where ∂_{y_i} denotes the partial derivative with respect to y_i ; S is the injective hull of the residue field \mathbf{k} as S -module, [5].

Given a polynomial $H \in \Gamma$ we denote by $\langle H \rangle$ the sub- S -module of Γ generated by H . Notice that $\langle H \rangle$ is the \mathbf{k} -vector subspace of Γ generated by H and its derivatives of any order. On the other hand, we denote by $\langle H \rangle_{\mathbf{k}}$ the one-dimensional \mathbf{k} -vector space generated by H .

Proposition 1.1 (Macaulay-Matlis duality, [8]). *There is a order-reversing bijection \perp between the set of finitely generated (graded) sub- S -submodules of Γ and the set of (homogeneous) \mathfrak{m} -primary ideals of S given by: if M is a submodule of Γ then $M^\perp = (0 :_S M) \subset S$, and $I^\perp = (0 :_\Gamma I) \subset \Gamma$ for an ideal $I \subset S$. An ideal I is Gorenstein if and only if I^\perp is generated by a single polynomial $F \in \Gamma$.*

I^\perp is the so-called Macaulay's inverse system of I .

Given an Artinian graded quotient $A = S/I$ we denote by $\text{soc}(A) = (0 :_A \mathfrak{m})$ the socle of A . The ring A is Gorenstein if and only if $\dim_{\mathbf{k}}(\text{soc}(A)) = 1$, if this is the case then $\text{soc}(A)$ is a principal ideal of A generated by a homogeneous element of degree s . This integer is the socle degree of A . By Macaulay's duality there exists a homogeneous element F of degree s such that $I^\perp = \langle F \rangle$.

The examples of this paper are done by using the Singular library [2], [1].

2 Characterization of complete intersections

The following result is a consequence of [9], but in the Artinian case we can give an easier proof by using Macaulay's duality.

Lemma 2.1. *Let $A = S/K$ be a graded Gorenstein ring of socle degree s , and let J be a homogeneous ideal such that $K \subset J$. If $\text{soc}(R/J)_s \neq 0$ then $K = J$.*

Proof. We have $J^\perp \subset K^\perp = \langle F \rangle$ with F a homogeneous form of degree s . Since $\text{soc}(R/J)_s \neq 0$ there exist a homogeneous polynomial $G \in J^\perp \setminus \{0\}$ with $\deg(G) = s$. Hence $0 \neq G \in K_s^\perp = \langle F \rangle_{\mathbf{k}}$. Since $\dim_{\mathbf{k}} \langle F \rangle_{\mathbf{k}} = 1$ there exist $\lambda \in \mathbf{k} \setminus \{0\}$ such that $G = \lambda F$. From this we get that

$$K^\perp = \langle F \rangle = \langle G \rangle \subset J^\perp.$$

Hence $K^\perp = J^\perp$ and by Macaulay's duality $K = J$. □

We say that a homogeneous ideal $I \subset S$ is a complete intersection of type $\underline{d} = (d_0, \dots, d_n) \in \mathbb{N}^{d+1}$, with $2 \leq d_0 \leq \dots \leq d_n$, if there exists homogeneous regular sequence $\underline{f} = \{f_0, \dots, f_n\}$ minimally generating I such that $\deg(f_i) = d_i$, $i = 0, \dots, n$.

Given an integer $d \geq 1$ we denote by $\mathbb{P}_{\mathbf{k}}^{N(d)}$ the projective space parametrizing the forms of degree d on x_0, \dots, x_n , where $N(d) = \binom{n+d}{n} - 1$. Given integers $2 \leq d_0 \leq \dots \leq d_n$ we consider the multi-projective space $\mathbb{V} = \prod_{i=0}^n \mathbb{P}_{\mathbf{k}}^{N(d_i)}$ parameterizing the $(n+1)$ -pla of homogeneous forms $\underline{F} = \{F_0, \dots, F_n\}$, where F_i is a homogeneous form of degree d_i , $i = 0, \dots, n$. Recall that the resultant $\text{Res}(\underline{F})$ defines an irreducible hypersurface of \mathbb{V} , [4, Chapitre 5, Section 5.3]. In fact, $\text{Res}(\underline{F})$ is the generator of the defining ideal of the variety of \mathbb{V} obtained by projection on the first component of the incidence variety $\{(\underline{F}, p) \in \mathbb{V} \times \mathbb{P}_{\mathbf{k}}^n \mid F_i(p) = 0, i = 0, \dots, n\}$. The resultant $\text{Res}(\underline{F})$ is a polynomial of degree $\prod_{i \neq j} d_j$ on the coefficients of F_i , $i = 0, \dots, n$, [4, Proposition 5.21]. The resultant can be computed effectively, [4, Algorithme 5.22], and can be a dense polynomial, see [4, Example 5.20].

Given a $n+1$ -pla $\underline{f} = (f_0, \dots, f_n)$ of homogeneous elements of S of degree d_0, \dots, d_n the ideal (\underline{f}) is \mathfrak{m} -primary if and only if $\text{Res}(\underline{f}) \neq 0$, [4, Proposition 5.17]. These conditions are equivalent to $\dim_{\mathbf{k}}(S/(\underline{f})) < \infty$, or that \underline{f} is a maximal regular sequence of S .

The main result of this paper is the following:

Proposition 2.2. *Let $H \in S \setminus \{0\}$ be a homogeneous polynomial. Then the ideal $\text{Ann}_{\mathbf{R}}\langle H \rangle$ is a complete intersection of type $\underline{d} = (d_0, \dots, d_n)$ if and only if there exists homogeneous forms $\underline{f} = \{f_0, \dots, f_n\} \subset S$ such that $\deg(f_i) = d_i$, $i = 0, \dots, n$, and*

(i) $\text{Res}(\underline{f}) \neq 0$,

(ii) $\underline{f} \circ H = 0$, and

(iii) $\det(\text{Jac}(\underline{f})) \circ H \neq 0$.

If these equivalent conditions hold then $\text{Ann}_{\mathbf{R}}\langle H \rangle = (f_0, \dots, f_n)$ and $\deg(H) = \sum_{i=0}^n d_i - (n+1)$.

Proof. Assume that $K = \text{Ann}_{\mathbf{R}}\langle H \rangle$ is a complete intersection ideal generated by a homogeneous regular sequence $\underline{f} = \{f_0, \dots, f_n\} \in S$ of type (d_0, \dots, d_n) . Since $(f_0, \dots, f_n) \subset K$, by Macaulay duality we get (ii). Being $\{f_0, \dots, f_n\}$ a regular sequence we have $\text{Res}(\underline{f}) \neq 0$, [11] or [4, Proposition 5.17]. By hypothesis K is a complete intersection, so from [10], see also [12, Theorem 2.4], we have that the coset of $\det(\text{Jac}(\underline{f}))$ in S/K is a generator of its socle, in particular $\det(\text{Jac}(\underline{f})) \circ H \neq 0$.

Let now $\underline{f} = \{f_1, \dots, f_n\} \in S$ be a sequence of homogeneous elements of S satisfying the conditions (i), (ii) and (iii). From (i) we get that \underline{f} is a regular sequence, [11] or [4, Proposition 5.17]. From (ii) and Macaulay's duality we get that $K = (f_0, \dots, f_n) \subset J = \text{Ann}_{\mathbf{R}}\langle H \rangle$, so we can consider the natural projection

$$\frac{S}{K} \longrightarrow \frac{S}{J} \tag{1}$$

Being K a complete intersection of height $n + 1$ the socle of S/K is concentrated in a single degree $s = \sum_{i=0}^n d_i - (n + 1)$, and it is generated by the coset of $\det(\text{Jac}(\underline{f}))$, [10], see also [12, Theorem 2.4]. By (1) the coset of $\det(\text{Jac}(\underline{f}))$ in S/J belongs to its socle.

From (iii) we get that the ideal of S/J generated by $\det(\text{Jac}(\underline{f}))$ is not zero and of degree s . Since the coset of $\det(\text{Jac}(\underline{f}))$ in S/J is non-zero and belongs to $\text{soc}(R/J)$, from Lemma 2.1 we deduce that $K = J$, i.e. $(f_0, \dots, f_n) = \text{Ann}_R\langle H \rangle$.

We know that the coset of $\det(\text{Jac}(\underline{f}))$ in S/K is a generator of its socle, so $\deg(H) = \deg(\det(\text{Jac}(\underline{f}))) = \sum_{i=0}^n \deg(f_i) - (n + 1)$. \square

Remark 2.3. Let $H \neq 0$ be a homogeneous polynomial. Since $\text{Ann}_R\langle H \rangle$ is of maximal height there always exists a sequence \underline{f} satisfying (i) and (ii).

Given integers $1 \leq d_0 \leq \dots \leq d_n$ we consider the multi-projective space $\mathbb{E} = \mathbb{V} \times \mathbb{P}_{\mathbf{k}}^{N(s)}$ parameterizing the pairs (\underline{f}, H) where $\underline{f} = \{f_0, \dots, f_n\}$ is a $(n + 1)$ -pla of homogeneous forms of degrees d_1, \dots, d_n , and H is a homogeneous polynomial on y_0, \dots, y_n of degree $s = \sum_{i=0}^n d_i - (n + 1)$. We denote by π the natural projection in the second component of \mathbb{E} : $\pi : \mathbb{E} \rightarrow \mathbb{P}_{\mathbf{k}}^{N(s)}$.

Let U be the open subset of \mathbb{E} defined by the $(n + 1)$ -plas \underline{f} such that $\text{Res}(\underline{f}) \neq 0$. We consider the locally closed subset Z_1 (resp. closed subset Z_2) of \mathbb{E} defined by the condition $\underline{f} \circ H = 0$ and $\underline{f} \in U$ (resp. $\det(\text{Jac}(\underline{f})) \circ H = 0$). For all $[H] \in \mathbb{P}_{\mathbf{k}}^{N(s)}$, we denote the fiber of $[H]$ in Z_i by $\pi, i = 1, 2$,

$$(Z_i)_H = Z_i \cap \pi^{-1}([H]) = \{(\underline{f}, [H]) \in Z_i\}.$$

Proposition 2.4. *Let H be a degree s homogeneous form. Then $\text{Ann}_R\langle H \rangle$ is a complete intersection of type (d_0, \dots, d_n) if and only if*

$$(Z_1)_H \not\subseteq (Z_2)_H.$$

The set of homogeneous polynomials $[H] \in \mathbb{P}_{\mathbf{k}}^{N(s)}$ such that $(Z_1)_H \not\subseteq (Z_2)_H$ is a locally constructible set.

Proof. Assume that $\text{Ann}_R\langle H \rangle$ is not a complete intersection. Let (\underline{f}, H) be an element of $(Z_1)_H$. From Proposition 2.2 we get $\det(\text{Jac}(\underline{f})) \circ H = 0$, so $(\underline{f}, H) \in (Z_2)_H$.

Let H be an homogeneous form of degree s such that $(Z_1)_H \subset (Z_2)_H$. Assume that $\text{Ann}_R\langle H \rangle$ is a complete intersection of type d_0, \dots, d_n . Then there exists a family $\underline{f} = (f_0, \dots, f_n)$ satisfying the conditions of Proposition 2.2. In particular, condition (i) and (ii) is equivalent to $(\underline{f}, H) \in Z_1$. By hypothesis $(\underline{f}, H) \in Z_2$ so $\det(\text{Jac}(\underline{f})) = 0$. This a contradiction with the condition (iii) of Proposition 2.2.

From [6, Corollaire 9.5.2] we get that the set of homogeneous polynomials $[H] \in \mathbb{P}_{\mathbf{k}}^{N(s)}$ such that $(Z_1)_H \not\subseteq (Z_2)_H$ is a locally constructible set. \square

The computation of the defining equations of the locally constructible set of the last result, in contrast the case of the resultant, could be difficult. In the next section we will study a concrete example where Proposition 2.2 is applied for proving that some Gorenstein ideal $\text{Ann}_{\mathbf{R}}\langle H \rangle$ is a complete intersection. First we review the example 3.5 of [7].

Example 2.5. In this example we follow the notations of [7, Example 3.5]. We consider the ring of polynomials $S = \mathbf{k}[v, w, x, y, z]$.

(1) of [7, Example 3.5]. Given $H = vwxyz + wxyz^2$, $\text{Ann}_{\mathbf{R}}\langle H \rangle$ contains the sequence $\underline{f} = \{v^2, w^2, x^2, y^2z^2 - 2vz\}$, i.e. satisfies the condition (ii) of Proposition 2.2. Since $\overline{S}/(\underline{f})$ is Artinian we get condition (i) of Proposition 2.2. On the other hand, we get $\det(\text{Jac}(\underline{f})) = 2^5vwxy(z - v)$, so $\det(\text{Jac}(\underline{f})) \circ H = 2^5 \neq 0$. Hence $\text{Ann}_{\mathbf{R}}\langle H \rangle = (\underline{f})$ is a complete intersection.

(2) of [7, Example 3.5]. We proceed as the previous case.

(3) of [7, Example 3.5]. Let us consider $H = vwxyz + yz^4$. If $\text{Ann}_{\mathbf{R}}\langle H \rangle$ is a complete intersection, since $5 = d_0 + \dots + d_4 - 5$, then we get that $d_0 = \dots = d_4 = 2$. The \mathbf{k} -vector space of forms of degree two of $\text{Ann}_{\mathbf{R}}\langle H \rangle$ admits as \mathbf{k} -base the monomials: v^2, w^2, x^2, y^2 , i.e. a \mathbf{k} -base of \underline{f} satisfying (ii) of Proposition 2.2. Hence, any 5-pla $\underline{f} = (f_0, \dots, f_5)$ formed by 5 forms of degree two are linearly dependent, so $\det(\text{Jac}(\underline{f})) = 0$ and then $\text{Ann}_{\mathbf{R}}\langle H \rangle$ is not a complete intersection.

3 Case study: short algebras

In this section we consider short Gorenstein algebras: local Gorenstein algebras $A = \mathbf{k}[[x_1, \dots, x_r]]/I$ with Hilbert function $\{1, r, r, 1\}$. Recall that in the main result of [3] we prove that these algebras are isomorphic to their associated graded ring, i.e. we can consider that $A \cong \mathbf{k}[x_1, \dots, x_r]/K$ with K a homogeneous ideal. On the other hand if K is a complete intersection then $r \leq 3$. The cases $r \leq 2$ are computed directly, see [3]. In this example we study the case $r = 3$, following the general notations of this paper we have that $n = 2$. Hence $K = \text{Ann}_{\mathbf{R}}\langle H \rangle$ is an ideal of $S = \mathbf{k}[x_0, x_1, x_2]$ with H a degree 3 homogeneous polynomial.

In [3] we computed the isomorphism classes of these algebras in terms of the moduli of the degree three projective plane curve defined by H . Recall that a plane elliptic cubic curve $C \subset \mathbb{P}_{\mathbf{k}}^2$ is defined, in a suitable system of coordinates, by a Legendre's equation

$$H_{\lambda} = y_1^2y_2 - y_0(y_0 - y_2)(y_0 - \lambda y_2)$$

with $\lambda \neq 0, 1$. The j -invariant of C is:

$$j(\lambda) = 2^8 \frac{(\lambda^2 - \lambda + 1)^3}{\lambda^2(\lambda - 1)^2}.$$

It is well known that two plane elliptic cubic curves $C_i = V(H_{\lambda_i}) \subset \mathbb{P}_{\mathbf{k}}^2 = \mathbf{Spec}(S)$, $i = 1, 2$, are projectively isomorphic if and only if $j(\lambda_1) = j(\lambda_2)$. In [3, Proposition 3.7] we proved that:

Proposition 3.1. [3, Proposition 3.7] *Let A be an Artinian Gorenstein local \mathbf{k} -algebra with Hilbert function $\text{HF}_A = \{1, 3, 3, 1\}$. Then A is isomorphic to one and only one of the following graded quotients of $S = \mathbf{k}[x_0, x_1, x_2]$:*

<i>Model $A = S/K$</i>	<i>Inverse system</i>	<i>Geometry of $C = V(K) \subset \mathbb{P}_{\mathbf{k}}^2$</i>
(x_0^2, x_1^2, x_2^2)	$y_0 y_1 y_2$	<i>Three independent lines</i>
$(x_0^2, x_0 x_3^2, x_2 x_1^2, x_1^3, x_2^2 + x_0 x_1)$	$y_1(y_0 y_1 - y_2^2)$	<i>Conic and a tangent line</i>
$(x_0^2, x_1^2, x_2^2 + 6x_0 x_1)$	$y_3(y_1 y_2 - y_3^2)$	<i>Conic and a non-tangent line</i>
$(x_1^2, x_0 x_1, x_0^2 + x_1^2 - 3x_0 x_2)$	$y_1^2 y_2 - y_0^2(y_0 + y_2)$	<i>Irreducible nodal cubic</i>
$(x_2^2, x_0 x_1, x_0 x_2, x_1^3, x_0^3 + 3x_1^2 x_2)$	$y_1^2 y_2 - y_0^3$	<i>Irreducible cuspidal cubic</i>
$(x_1 x_2, x_0 x_2, x_0 x_1, x_1^3 - x_2^3, x_0^3 - x_2^3)$	$y_0^3 + y_1^3 + y_2^3$	<i>Elliptic Fermat curve</i>
$K_\lambda = (x_0 x_1, F_1, F_2)$	$H_\lambda, j(\lambda) \neq 0$	<i>Elliptic non Fermat curve</i>

with $F_1 = \lambda^2 x_0^2 + \lambda(1 + \lambda)x_0 x_2 + (\lambda^2 - \lambda + 1)x_2^2$, $F_2 = \lambda^2 x_1^2 + \lambda x_0 x_2 + (1 + \lambda)x_2^2$, and $K_{\lambda_1} \cong K_{\lambda_2}$ if and only if $j(\lambda_1) = j(\lambda_2)$.

Notice that the minimal number of generators of the height three Gorenstein ideal K of the last result are odd, as the main theorem of [13] prescribes. All cases in Proposition 3.1 can be easily computed by using Singular but the last one, [2]. In this case a parameter is involved: the j -invariant of the elliptic curve. In [3] we settle this case by showing first that $J = (x_0 x_1, F_1, F_2) \subset K_\lambda$ and then proving that $\dim_{\mathbf{k}}(S/J) = 8$. We can prove this fact by using Proposition 2.2. We know that $J = (x_0 x_1, F_1, F_2) \subset K_\lambda = \text{Ann}_{\mathbf{R}}\langle H_\lambda \rangle$, so $\underline{f} = \{x_0 x_1, F_1, F_2\}$ satisfies condition (ii) of Proposition 2.2. On the other hand it is easy to prove that J is \mathfrak{m} -primary, so we have condition (i) of Proposition 2.2. Since $j(\lambda) \neq 0$

$$\det(\text{Jac}(x_0 x_1, F_1, F_2)) \circ H_\lambda = -16\lambda^2(\lambda^2 - \lambda + 1) \neq 0.$$

Hence we get condition (iii) of Proposition 2.2 and then $(x_0 x_1, F_1, F_2) = K_\lambda$.

References

- [1] W. Decker, G.-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 4-3-0-A computer algebra system for polynomial computations, <https://www.singular.uni-kl.de>, 2022.
- [2] J. Elias, INVERSE-SYST–Singular library for computing Macaulay’s inverse systems, arXiv: 1501.01786, <https://www.ub.edu/C3A/elias/inverse-syst-v.5.2.lib>, 2015.
- [3] J. Elias and M. E. Rossi, *Isomorphism classes of short Gorenstein local rings via Macaulay’s inverse system*, Trans. Am. Math. Soc., **364** (2012), no. 9, 4589–4604.
- [4] M. Elkadi and B. Mourrain, *Introduction à la résolution des systèmes polynomiaux*, Mathématiques and Applications (Berlin) 59. Springer, 2007.
- [5] P. Gabriel, *Objects injectifs dans les catégories abéliennes*, Séminaire Pierre Dubriel 1958/59, Numdam, 1959, pp. 17–01, 32.

- [6] A. Grothendieck, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. (1966), no. 28, 255.
- [7] T. Harima, A. Wachi, and J. Watanabe, *A characterization of the Macaulay dual generators for quadratic complete intersections*, Illinois J. Math. **61** (2017), no. 3-4, 371–383.
- [8] A. Iarrobino and V. Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999, Appendix C by Iarrobino and Steven L. Kleiman.
- [9] A. R. Kustin and B. Ulrich, *If the socle fits*, J. Algebra **147** (1992), no. 1, 63–80.
- [10] G. Scheja and U. Storch, *Über Spurfunktionen bei vollständigen Durchschnitten*, J. Reine Angew. Math. **278/279** (1975), 174–190.
- [11] G. Scheja and U. Storch, *Regular sequences and resultants*, Research Notes in Mathematics, vol. 8, A K Peters, Ltd., Natick, MA, 2001.
- [12] W. V. Vasconcelos, *Jacobian matrices and constructions in algebra*, Applied algebra, algebraic algorithms and error-correcting codes (New Orleans, LA, 1991), Lecture Notes in Comput. Sci., vol. 539, Springer, Berlin, 1991, pp. 48–64.
- [13] J. Watanabe, *A note on Gorenstein rings of embedding codimension three*, Nagoya Math. J. **50** (1973), 227–232.

Joan Elias
 Departament de Matemàtiques i Informàtica
 Universitat de Barcelona
 Gran Via 585, 08007 Barcelona, Spain
 e-mail: elias@ub.edu