

# Totally geodesic submanifolds in the manifold SPD of symmetric positive-definite real matrices

Alice Barbara Tumpach\*, Gabriel Larotonda†

## Abstract

This paper is a self-contained exposition of the geometry of symmetric positive-definite real  $n \times n$  matrices  $\text{SPD}(n)$ , including necessary and sufficient conditions for a submanifold  $\mathcal{N} \subset \text{SPD}(n)$  to be totally geodesic for the affine-invariant Riemannian metric. A non-linear projection  $x \mapsto \pi(x)$  on a totally geodesic submanifold is defined. This projection has the minimizing property with respect to the Riemannian metric: it maps an arbitrary point  $x \in \text{SPD}(n)$  to the unique closest element  $\pi(x)$  in the totally geodesic submanifold for the distance defined by the affine-invariant Riemannian metric. Decompositions of the space  $\text{SPD}(n)$  follow, as well as variants of the polar decomposition of non-singular matrices known as Mostow's decompositions. Applications to decompositions of covariant matrices are mentioned.

*Keywords:* covariance matrices, reductive symmetric spaces, decompositions of Lie groups, symmetric positive-definite matrices.

*Subject Classification:* Riemannian Geometry.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Statement of Mostow's Theorem . . . . .	2
1.2	Examples of Applications . . . . .	3
1.2.1	Geodesic projection on the space of diagonal matrices . . . . .	3
1.2.2	Geodesic projection on the space of block diagonal matrices and on the space of block-anti-diagonal matrices . . . . .	3
1.2.3	Decomposition of $SL(2, \mathbb{R})$ . . . . .	5
1.3	Organization of the paper . . . . .	5
<b>2</b>	<b>The manifold <math>\text{SPD}(n)</math> of symmetric positive-definite real matrices</b>	<b>5</b>
<b>3</b>	<b>Totally geodesic submanifolds of the manifold <math>\text{SPD}(n)</math></b>	<b>8</b>
<b>4</b>	<b>Orthogonal projection on a totally geodesic submanifold</b>	<b>11</b>
<b>5</b>	<b>Mostow's decomposition Theorem for <math>GL(n, \mathbb{R})</math></b>	<b>14</b>
<b>A</b>	<b>The differential of the exponential map of a Lie group <math>G</math></b>	<b>15</b>
A.1	First expression of the differential of the exponential map . . . . .	15
A.2	Second expression of the differential of the exponential map . . . . .	17
<b>B</b>	<b>Geodesics in locally symmetric homogeneous spaces</b>	<b>18</b>

---

\*alice-barbora.tumpach@univ-lille.fr, Institut CNRS Pauli, Vienna, Austria.

†glaroton@dm.uba.ar, UBA & CONICET, Buenos Aires, Argentina

# 1 Introduction

Mostow's decomposition theorem is a generalization of polar decomposition and goes back to 1955 with the original proof by Mostow ([10]). Different proofs were given in [15, 8, 5, 16] in the context of complex infinite-dimensional Lie groups, in [1] in the context of von Neumann algebras, and [9] in the context of Finsler Lie groups. In this paper, we consider the real group  $\mathrm{GL}(n, \mathbb{R})$ , which is not semi-simple, but appears in a lot of applications. The aim of this paper is to provide a self-contained exposition of the geometry needed to understand this Theorem, and advertise some possible applications, which should be important in the fields of Probability and Statistics. For a general exposition of the geometry of the manifold of symmetric positive-definite real matrices we refer to [4].

## 1.1 Statement of Mostow's Theorem

Let us first state the theorem.

**Theorem 1.1** *Let  $E$  be a real subspace of the vector space  $\mathrm{Sym}(n)$  of symmetric  $n \times n$  real matrices, and denote by  $\mathcal{E} := \exp(E)$  the image of  $E$  by the exponential of matrices*

$$\mathcal{E} = \{y = \exp u := 1 + u + \frac{u^2}{2} + \frac{u^3}{3!} + \dots, u \in E\}. \quad (1)$$

*We endow the manifold  $\mathrm{SPD}(n)$  of symmetric positive-definite real  $n \times n$  matrices with the Riemannian metric (sometimes called *affined Riemannian metric*) given at  $x \in \mathrm{SPD}(n)$  by:*

$$\langle X, Y \rangle_x := \mathrm{Tr} (x^{-1} X x^{-1} Y) = \mathrm{Tr} (x^{-\frac{1}{2}} X x^{-\frac{1}{2}} \cdot x^{-\frac{1}{2}} Y x^{-\frac{1}{2}}), \quad (2)$$

where the dot denotes the product of matrices.

1. Then  $\mathcal{E} := \exp(E)$  is a totally geodesic submanifold of  $\mathrm{SPD}(n)$  and geodesically convex in  $\mathrm{SPD}(n)$ , if and only if the vector space  $E \subset \mathrm{Sym}(n)$  satisfies

$$[X, [X, Y]] \in E, \quad \text{for all } X, Y \in E, \quad (3)$$

where for any matrices  $A, B$ , the bracket is defined as  $[A, B] := A \cdot B - B \cdot A$ .

2. In this case, any  $x \in \mathrm{SPD}(n)$  can be decomposed uniquely as a product  $e \cdot f \cdot e$ , where  $e \in \mathcal{E} := \exp(E)$  and  $f \in \mathcal{F} := \exp(E^\perp)$ . The matrix  $x$  admits a unique projection  $\pi(x)$  onto  $\mathcal{E} := \exp(E)$  which minimises the geodesic distance from  $x$  to  $\mathcal{E} := \exp(E)$ , i.e.

$$\mathrm{dist}(x, \mathcal{E}) = \mathrm{dist}(x, \pi(x)).$$

The matrix  $e$  is related to  $\pi(x)$  by

$$e = \pi(x)^{\frac{1}{2}}. \quad (4)$$

3. If (3) is satisfied, any  $g \in \mathrm{GL}(n, \mathbb{R})$  can be decomposed uniquely as a product  $k \cdot \exp V \cdot \exp W$ , where  $k$  is an orthogonal matrix,  $W$  is in  $E$  and  $V$  is in the orthogonal space  $F := E^\perp$ . The Lie group  $\mathrm{GL}(n, \mathbb{R})$  is diffeomorphic to the product

$$\mathrm{GL}(n, \mathbb{R}) = \mathrm{O}(n) \times \exp F \times \exp E = \mathrm{O}(n) \times \mathcal{F} \times \mathcal{E}, \quad (5)$$

where

- $\mathrm{O}(n)$  is the orthogonal group

$$\mathrm{O}(n) := \{M \in \mathrm{GL}(n, \mathbb{R}), M^T M = M M^T = \mathrm{Id}\} \quad (6)$$

(here  $\mathrm{Id}$  is the  $n \times n$  identity matrix).

- $F$  denotes the orthogonal of  $E$  in  $\mathrm{Sym}(n)$ ,

$$F := E^\perp = \{Y \in \mathrm{Sym}(n), \mathrm{Tr} XY = 0 \quad \forall X \in E\}, \quad (7)$$

- $\mathcal{F} := \exp F$ .

**Corollary 1.2** *Any geodesically complete convex submanifold  $\mathcal{N}$  of  $\mathrm{SPD}(n)$  is of the form*

$$\mathcal{N} = x \cdot \exp E \cdot x$$

for some  $x \in \mathrm{SPD}(n)$ , and some subspace  $E \subset \mathrm{Sym}(n)$  satisfying  $[X, [X, Y]] \in E$  for all  $X, Y \in E$ .

## 1.2 Examples of Applications

### 1.2.1 Geodesic projection on the space of diagonal matrices

Let  $\mathbf{X} := (X_1, X_2, \dots, X_n)$  be a real random vector of dimension  $n$  and consider its *covariance matrix*

$$\text{cov}(\mathbf{X}, \mathbf{X}) = \mathbb{E} \left[ (\mathbf{X} - \mathbb{E}(\mathbf{X})) (\mathbf{X} - \mathbb{E}(\mathbf{X}))^T \right] \quad (8)$$

where the operator  $\mathbb{E}$  denotes the expected value (mean) of its argument.

Let  $E_1 \subset \text{Sym}(n)$  be the space of real diagonal matrices. Then condition (3) is satisfied, hence by Theorem 1.1 (see Theorem 4.2 for more details), any symmetric positive-definite matrix  $x \in \text{SPD}(n)$  has a unique projection  $\pi(x)$  in  $\mathcal{E}_1 := \exp(E_1)$  that minimises (for the Riemannian metric (2)) the distance between  $x$  and the manifold of real positive-definite diagonal matrices  $\mathcal{E}_1$ . Moreover

$$x = \pi(x)^{\frac{1}{2}} \cdot f \cdot \pi(x)^{\frac{1}{2}}, \quad (9)$$

for a unique element  $f$  in  $\mathcal{F}_1 := \exp(E_1^\perp)$  where

$$E_1^\perp := \{Y = (Y_{ij}) \in \text{Sym}(n), Y_{ii} = 0\}.$$

**Remark 1.3** The *correlation matrix* of  $\mathbf{X}$  is defined as

$$\text{corr}(\mathbf{X}, \mathbf{X}) = (\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{-\frac{1}{2}} \cdot \text{cov}(\mathbf{X}, \mathbf{X}) \cdot (\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{-\frac{1}{2}}), \quad (10)$$

where  $\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))$  is the matrix of diagonal elements of  $\text{cov}(\mathbf{X}, \mathbf{X})$ , i.e. the diagonal  $n \times n$  matrix containing the variances of  $X_i$ ,  $i = 1, \dots, n$ . Then

$$\text{cov}(\mathbf{X}, \mathbf{X}) = (\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{\frac{1}{2}} \cdot \text{corr}(\mathbf{X}, \mathbf{X}) \cdot (\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{\frac{1}{2}}).$$

In this decomposition, we remark that  $\text{cov}(\mathbf{X}, \mathbf{X})$  is the image of  $\text{corr}(\mathbf{X}, \mathbf{X})$  by the isometry of  $\text{SPD}(n)$  which sends  $y$  to

$$(\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{\frac{1}{2}} \cdot y \cdot (\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))^{\frac{1}{2}}).$$

However, in general,  $\text{diag}(x)$  differs from the projection  $\pi(x)$  onto the manifold of real positive-definite diagonal matrices for the Riemannian metric (2). In fact,  $\text{diag}(\text{cov}(\mathbf{X}, \mathbf{X}))$  is the projection of  $\text{cov}(\mathbf{X}, \mathbf{X})$  onto the vector space  $E_1 \subset \text{Sym}(n)$  of diagonal matrices for the Euclidian scalar product given by

$$\begin{aligned} (\cdot, \cdot) : \text{Sym}(n) \times \text{Sym}(n) &\longrightarrow \mathbb{R} \\ (X, Y) &\longmapsto \text{Tr}(XY). \end{aligned} \quad (11)$$

**Lemma 1.4** *Let  $x = \pi(x)^{1/2} e^v \pi(x)^{1/2}$  be Mostow's decomposition for  $x$ , with  $v^* = v \in E_1^\perp$  a matrix with zeros on the diagonal. Then  $\text{diag}(x) = \pi(x)$  if and only if  $\text{diag}(e^v) = 1$ .*

**Proof:** Write  $x = Pe^vP$  for short, and note that the assumption is  $P^2 = \text{diag}(x)$ . Then  $P^2 = \text{diag}(Pe^vP)$ , thus  $1 = P^{-1} \text{diag}(Pe^vP) P^{-1}$ . But taking diagonal has the bi-module property, i.e.  $\text{diag}(d_1 a d_2) = d_1 \text{diag}(a) d_2$  for any matrix  $a$  and any diagonal matrices  $d_1, d_2$ . Hence  $1 = \text{diag}(e^v)$ .  $\square$

In particular for  $n = 2$  note that  $v$  must be of the form  $v = \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix}$ , thus a straightforward computation shows that

$$e^v = \begin{pmatrix} \cosh(a) & \sinh(a) \\ \sinh(a) & \cosh(a) \end{pmatrix},$$

and  $\text{diag}(e^v) = 1$  is only possible if  $\cosh(a) = 1$ , or equivalently if  $a = 0$ . Thus it must be  $v = 0$  thus  $x = \pi(x)$  was already a diagonal matrix.

### 1.2.2 Geodesic projection on the space of block diagonal matrices and on the space of block-anti-diagonal matrices

Consider now two different random vectors  $\mathbf{X}$  of dimension  $p$  and  $\mathbf{Y}$  of dimension  $p$ , and the random vector

$$\mathbf{z} := \begin{pmatrix} \mathbf{X} \\ \mathbf{Y} \end{pmatrix}$$

of dimension  $p + q =: n$ . The *joint covariance matrix*  $\Sigma$  of  $\mathbf{X}$  and  $\mathbf{Y}$  is defined as

$$\Sigma = \mathbb{E} \left[ (\mathbf{Z} - \mathbb{E}(\mathbf{Z})) (\mathbf{Z} - \mathbb{E}(\mathbf{Z}))^T \right] = \begin{pmatrix} \text{cov}(\mathbf{X}, \mathbf{X}) & \text{cov}(\mathbf{X}, \mathbf{Y}) \\ \text{cov}(\mathbf{X}, \mathbf{Y}) & \text{cov}(\mathbf{Y}, \mathbf{Y}) \end{pmatrix}, \quad (12)$$

where  $\text{cov}(\mathbf{X}, \mathbf{Y})$  denotes the *cross-covariance matrix* between  $\mathbf{X}$  and  $\mathbf{Y}$ :

$$\text{cov}(\mathbf{X}, \mathbf{Y}) := \mathbb{E} \left[ (\mathbf{X} - \mathbb{E}(\mathbf{X})) (\mathbf{Y} - \mathbb{E}(\mathbf{Y}))^T \right]. \quad (13)$$

Let  $\mathbb{E}_2 \subset \text{Sym}(n)$  be the vector space of block-diagonal matrices and  $\mathbb{E}_3 \subset \text{Sym}(n)$  be the vector space of block-anti-diagonal matrices:

$$\begin{aligned} \mathbb{E}_2 &= \left\{ Z \in \text{Sym}(n), Z = \begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}, X \in \text{Sym}(p), Y \in \text{Sym}(q) \right\} \\ \mathbb{E}_3 &= \left\{ Z \in \text{Sym}(n), Z = \begin{pmatrix} 0 & C \\ C^T & 0 \end{pmatrix}, C \in \text{Mat}(p, q) \right\}. \end{aligned}$$

Then  $\mathbb{E}_2$  and  $\mathbb{E}_3$  both satisfy condition (3) and are orthogonal to each other for the Euclidian scalar product on  $\text{Sym}(n)$ . Note that  $\mathbb{E}_2$  is a Lie algebra, whereas  $\mathbb{E}_3$  is not. Therefore the cross-covariance matrix  $\Sigma$  has a unique expression of the form

$$\Sigma = \exp D \cdot \exp A \cdot \exp D, \quad (14)$$

where  $D$  is block-diagonal and  $A$  is block-anti-diagonal, as well as a unique expression of the form

$$\Sigma = \exp A' \cdot \exp D' \cdot \exp A', \quad (15)$$

where  $D'$  is block-diagonal and  $A'$  is block-anti-diagonal.

Moreover since the even powers of a block-anti-diagonal matrix are diagonal, one has the decomposition

$$\exp A = \cosh A + \sinh A,$$

where

$$\begin{aligned} \cosh A &:= \frac{(\exp(A) + \exp(-A))}{2} = \sum_{i=0}^n \frac{A^{2i}}{(2i)!} \in \mathbb{E}_2 \\ \sinh A &:= \frac{(\exp(A) - \exp(-A))}{2} = \sum_{i=1}^n \frac{A^{(2i-1)}}{(2i-1)!} \in \mathbb{E}_3 \end{aligned} \quad (16)$$

Therefore equation (14) leads to a decomposition

$$\Sigma = \exp D \cosh A \exp D + \exp D \sinh A \exp D, \quad (17)$$

where  $\exp D \cosh A \exp D \in \mathbb{E}_2$  is block-diagonal and  $\exp D \sinh A \exp D \in \mathbb{E}_3$  is block-anti-diagonal. Similarly equation (15) leads to a decomposition of  $\Sigma$  in a sum

$$(\cosh A' \cdot \exp D' \cdot \cosh A' + \sinh A' \cdot \exp D' \sinh A') + (\sinh A' \exp D' \cosh A' + \cosh A' \exp D' \sinh A') \quad (18)$$

where

$$(\cosh A' \cdot \exp D' \cdot \cosh A' + \sinh A' \cdot \exp D' \sinh A') \in \mathbb{E}_2$$

is block-diagonal and

$$(\sinh A' \exp D' \cosh A' + \cosh A' \exp D' \sinh A') \in \mathbb{E}_3$$

is block-anti-diagonal.

**Remark 1.5** For more than two blocks, the following happens: the space of block diagonal matrices still satisfies condition (3), hence there exists an orthogonal projection on the space of positive definite block diagonal matrices with an arbitrary (fixed) number of blocks, and the corresponding decomposition similar to (14). However the space of matrices with zero diagonal blocks do not satisfy condition (3) when there are more than two blocks, hence one can not deduce a decomposition of the type (15) in that case.

### 1.2.3 Decomposition of $SL(2, \mathbb{R})$

The group  $SL(2, \mathbb{R})$  is the group of isometries of the hyperbolic upper half plane  $\mathbb{H}^2$ :

$$\mathbb{H}^2 := \{z = x + hi, x \in \mathbb{R}, h \in \mathbb{R}^{+*}\} \subset \mathbb{C},$$

where the action is by homographies. Mostow's decomposition theorem applied to

$$E = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha \end{pmatrix}, \alpha \in \mathbb{R} \right\}$$

leads to a decomposition of isometries

$$SL(2, \mathbb{R}) = SO(2) \times \exp F \times \exp E$$

where

- $SO(2)$  is the stabilizer of  $i \in \mathbb{C}$ ;
- $\mathcal{E} = \exp E$  is the space of dilatations  $q \mapsto \alpha^2 q$ ,  $\alpha \in \mathbb{R}$ ;
- $\mathcal{F} = \exp F = \left\{ \begin{pmatrix} \cosh \beta & \sinh \beta \\ \sinh \beta & \cosh \beta \end{pmatrix}, \beta \in \mathbb{R} \right\}$  is the space of hyperbolic transformations with fixed points  $-1$  and  $1$  in  $\mathbb{R} \in \partial\mathbb{H}^2$  and preserving the unit half-circle containing  $i$ .

### 1.3 Organization of the paper

This paper is organized as follows. First we recall the geometry of the space  $SPD(n)$  of symmetric positive-definite  $n \times n$  matrices. We show in particular that it is a symmetric space of non-positive sectional curvature, homogeneous under the group  $GL(n, \mathbb{R})$  and that the exponential map defined by the usual power series is a diffeomorphism from the space  $\text{Sym}(n)$  of symmetric  $n \times n$  real matrices onto  $SPD(n)$ . Moreover we show that the exponential map is the Riemannian exponential map at the identity with respect to the affine Riemannian metric on  $SPD(n)$ . This is implied by a general result on the geodesics in locally symmetric spaces that we recall (Proposition B.1). This study implies the usual Al-Kashi inequality on the sides of a geodesic triangle in the non-positively curved space  $SPD(n)$ , and the convexity property of the distance between two geodesics. In the second subsection, the characterization of the geodesic subspaces of  $SPD(n)$  given by equation (3) is proved, mainly following arguments from [10]. In subsection 4, the key-step for the proof of Mostow's decomposition (5) is given by the construction of a non-linear projection from  $SPD(n)$  onto every closed geodesic subspace. The arguments given here for the existence of such projection are simpler and more direct than the ones given in the original paper [10], and apply to arbitrary dimension. In the last subsection, we use this projection to prove the theorem stated above.

## 2 The manifold $SPD(n)$ of symmetric positive-definite real matrices

Let  $M(n, \mathbb{R})$  be the vector space of real  $n \times n$  matrices. The group  $GL(n, \mathbb{R})$  is the group of invertible real  $n \times n$  matrices, or equivalently the space of real matrices with non-zero determinant:

$$GL(n, \mathbb{R}) := \{g \in M(n, \mathbb{R}), \det(g) \neq 0\},$$

The group law given by the multiplication of matrices makes the open set  $GL(n, \mathbb{R}) \subset M(n, \mathbb{R})$  into a Lie-group with Lie-algebra  $M(n, \mathbb{R})$ , where the Lie-bracket is given by the commutator of matrices:

$$[A, B] = AB - BA, \forall A, B \in M(n, \mathbb{R}). \quad (19)$$

The orthogonal group  $O(n)$  and its Lie algebra  $\mathfrak{so}(n)$  are defined by

$$O(n) := \{M \in GL(n, \mathbb{R}), M^T M = M M^T = \text{Id}\}$$

$$\mathfrak{so}(n) = \{A \in M(n, \mathbb{R}), A^T + A = 0\}$$

The vector space  $M(n, \mathbb{R})$  splits into the direct sum of  $\mathfrak{so}(n)$  and the linear subspace  $\text{Sym}(n)$  of symmetric elements in  $M(n, \mathbb{R})$

$$\text{Sym}(n) = \{M \in M(n, \mathbb{R}), A^T = A\}.$$

The exponential of matrices defined for all  $A$  in  $M(n, \mathbb{R})$  as

$$\exp(A) := \sum_{n=0}^{+\infty} \frac{A^n}{n!} \quad (20)$$

takes  $\text{Sym}(n)$  to the submanifold  $\text{SPD}(n)$  of  $\text{GL}(n, \mathbb{R})$  consisting of symmetric positive-definite matrices

$$\exp : \text{Sym}(n) = \{A \in M(n, \mathbb{R}), A^T = A\} \longrightarrow \text{SPD}(n) = \{A \in \text{Sym}(n), A^T A > 0\}.$$

Note that for  $A \in M(n, \mathbb{R})$ , the curve  $\gamma(t) := \exp(tA)$  satisfies

$$\dot{\gamma}(t) = (L_{\gamma(t)})_*(A) = \gamma(t) \cdot A. \quad (21)$$

where  $L_{\gamma(t)}$  denotes left translation by  $\gamma(t)$ , i.e. multiplication on the left by the matrix  $\gamma(t)$ , and where the dot denotes the multiplication of matrices. Hence the exponential map defined by (20) is the usual exponential map on the Lie group  $\text{GL}(n, \mathbb{R})$ . Let us endow  $\text{SPD}(n)$  with the following Riemannian metric :

$$\langle X, Y \rangle_x := \text{Tr} (x^{-1} X x^{-1} Y) = \text{Tr} \left( x^{-\frac{1}{2}} X x^{-\frac{1}{2}} \cdot x^{-\frac{1}{2}} Y x^{-\frac{1}{2}} \right) \quad (22)$$

where  $x \in \text{SPD}(n)$ ,  $X, Y \in T_x \text{SPD}(n)$ . The identity matrix  $\text{Id}$  belongs to  $\text{SPD}(n)$  and will be our reference point in the following.

**Proposition 2.1** *The left action of  $\text{GL}(n, \mathbb{R})$  on  $\text{SPD}(n)$  defined by*

$$\begin{aligned} \text{GL}(n, \mathbb{R}) \times \text{SPD}(n) &\rightarrow \text{SPD}(n) \\ (g, x) &\mapsto g \cdot x \cdot g^T, \end{aligned} \quad (23)$$

*is a transitive action by isometries (here the dot  $\cdot$  denotes the multiplication of matrices).*

□ **Proof of Proposition 2.1:**

For every  $x$  in  $\text{SPD}(n)$ , the square root of  $x$  is well-defined and belongs to  $\text{SPD}(n)$ . In other words there exists  $y$  in  $\text{SPD}(n)$  such that  $x = y^2$ . Since  $y^T = y$ , one has  $x = y \cdot y^T = y \cdot \text{Id} \cdot y^T$ , and the transitivity follows. To show that  $\text{GL}(n, \mathbb{R})$  acts by isometries, one has to show that the scalar product  $\langle \cdot, \cdot \rangle_x$  on the tangent space to the manifold  $\text{SPD}(n)$  at  $x = y \cdot \text{Id} \cdot y^T$  is related to the scalar product  $\langle \cdot, \cdot \rangle_{\text{Id}}$  at  $\text{Id}$  by

$$\langle X, Y \rangle_{\text{Id}} = \langle y_* X, y_* Y \rangle_{y \cdot \text{Id} \cdot y^T}, \quad (24)$$

where  $y_* : T_{\text{Id}} \text{SPD}(n) \rightarrow T_{y \cdot \text{Id} \cdot y^T} \text{SPD}(n)$  denotes the infinitesimal action of  $y \in \text{GL}(n, \mathbb{R})$  on tangent vectors to the manifold  $\text{SPD}(n)$ . To compute this infinitesimal action, consider a curve  $x(t) \in \text{SPD}(n)$  such that  $x(0) = x$  and  $\dot{x}(t) = X$ , and, for a fixed element  $g \in \text{GL}(n, \mathbb{R})$ , differentiate action (23) to get

$$g_* X = \frac{d}{dt} \Big|_{t=0} g \cdot x(t) \cdot g^T = g \cdot X \cdot g^T.$$

It follows that

$$\begin{aligned} \langle y_* X, y_* Y \rangle_{y \cdot \text{Id} \cdot y^T} &= \langle y_* X, y_* Y \rangle_{y y^T} = \text{Tr} (y y^T)^{-1} (y \cdot X \cdot y^T) (y y^T)^{-1} (y \cdot Y \cdot y^T) \\ &= \text{Tr} (y^T)^{-1} X Y y^T = \text{Tr} (X Y) = \langle X, Y \rangle_{\text{Id}}. \end{aligned}$$

In particular, the scalar product  $\langle \cdot, \cdot \rangle_{\text{Id}}$  is invariant by the action of the isotropy group of the reference point  $\text{Id}$ , which is exactly the orthogonal group  $\text{O}(n)$ . □

**Corollary 2.2** *The manifold  $\text{SPD}(n)$  of symmetric positive-definite real matrices is a homogeneous Riemannian manifold*

$$\text{SPD}(n) = \text{GL}(n, \mathbb{R}) / \text{O}(n).$$

**Theorem 2.3** *The exponential map is a diffeomorphism from  $\text{Sym}(n)$  onto  $\text{SPD}(n)$ .*

■ **Proof of Theorem 2.3:**

Since every  $x$  in  $\text{SPD}(n)$  admits an orthogonal basis of eigenvectors with positive eigenvalues, the exponential map from  $\text{Sym}(n)$  to  $\text{SPD}(n)$  is injective and onto, the inverse mapping being given by the logarithm. The differential of the exponential map is recalled in the appendix (Proposition A.2, equation (33)). By Lemma A.3, equation (39)), it can be written as

$$d_X \exp(Y) = R_{\exp(\frac{X}{2})} L_{\exp(\frac{X}{2})} \tau_X(Y),$$

where  $\tau_X(Y)$  is the linear isomorphism of  $\text{Sym}(n)$  given by equation (37). Hence  $d_X \exp$  is continuous and invertible. By the inverse function Theorem, it follows that  $\exp$  is a diffeomorphism. □

**Definition 2.4** Let  $G$  be a Lie group. An homogeneous space  $M = G/K$  is called *reductive* if the Lie algebra  $\mathfrak{g}$  of  $G$  splits into a direct sum  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ , where  $\mathfrak{k}$  is the Lie algebra of  $K$ , and  $\mathfrak{m}$  an  $\text{Ad}(K)$ -invariant complement. A reductive homogeneous space is called *locally symmetric* if the commutation relation  $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}$  holds.

A locally symmetric space is a particular case of a *naturally reductive* space (see Definition 7.84 page 196 in [3], Definition 23 page 312 in [14], or Proposition 5.2 page 125 in [2] and the definition that follows). In the finite-dimensional setting, the geodesics of a naturally reductive space are orbits of one-parameter subgroups of  $G$  (see Proposition 25 page 313 in [14] for a proof of this fact). The symmetric case is also treated in Theorem 3.3 page 173 in [7]. Its infinite-dimensional version has been given in Example 3.9 in [13]. For the sake of completeness, we give a proof in the appendix (see Proposition B.1) based on the fact that in this case the Levi-Civita connection is the homogeneous connection (see [17] for the infinite-dimensional case). We deduce the following theorem.

**Theorem 2.5** *The manifold  $\text{SPD}(n)$  is a locally symmetric homogeneous space. The curve  $\gamma(t) := \exp(t \log(x))$ , ( $0 \leq t \leq 1$ ) is the unique geodesic in  $\text{SPD}(n)$  joining the identity  $o = \text{Id}$  to the element  $x \in \text{SPD}(n)$ . More generally, the geodesic  $\gamma_{x,y}(t)$  between any two points  $x, y$  in  $\text{SPD}(n)$  exists and is unique, and is given by*

$$\gamma_{x,y}(t) := x^{\frac{1}{2}} \left( \exp t \log \left( x^{-\frac{1}{2}} y x^{-\frac{1}{2}} \right) \right) x^{\frac{1}{2}}. \quad (25)$$

■ **Proof of Theorem 2.5:**

This follows from the same arguments as in [10], or by the general result stated in Proposition B.1. Indeed the commutation relation  $[\text{Sym}(n), \text{Sym}(n)] \subset \mathfrak{so}(n)$  implies that  $\text{SPD}(n) = \text{GL}(n, \mathbb{R}) / \text{O}(n)$  is locally symmetric. It follows that

$$\gamma(t) := \exp(t \log(x)) = \left( \exp \left( \frac{t}{2} \log(x) \right) \right) \cdot \text{Id} \cdot \left( \exp \left( \frac{t}{2} \log(x) \right) \right)^T, \quad (0 \leq t \leq 1),$$

is a geodesic joining  $o = \text{Id}$  to  $x$ . The uniqueness of this geodesic follows from Proposition B.1 and Theorem 2.3, since any other geodesic  $\gamma_2$  joining  $o = \text{Id}$  to  $x$  is necessarily of the form

$$\gamma_2(t) = \exp t \dot{\gamma}_2(0) = \left( \exp \left( \frac{t}{2} \dot{\gamma}_2(0) \right) \right) \cdot \text{Id} \cdot \left( \exp \left( \frac{t}{2} \dot{\gamma}_2(0) \right) \right)^T$$

with velocity  $\dot{\gamma}_2(0) \in \text{Sym}(n)$ . Since  $\text{GL}(n)$  acts by isometries on  $\text{SPD}(n)$ , for  $x, y \in \text{SPD}(n)$ , the image of the geodesic  $t \mapsto \exp t \log \left( x^{-\frac{1}{2}} y x^{-\frac{1}{2}} \right)$  by the isometry  $u \mapsto x^{\frac{1}{2}} u x^{\frac{1}{2}} = x^{\frac{1}{2}} \cdot u \cdot (x^{\frac{1}{2}})^T$  is itself a geodesic. It follows that the unique geodesic  $\gamma_{x,y}$  joining two points  $x$  and  $y$  in  $\text{SPD}(n)$  is given by

$$\gamma_{x,y}(t) := x^{\frac{1}{2}} \left( \exp t \log \left( x^{-\frac{1}{2}} y x^{-\frac{1}{2}} \right) \right) x^{\frac{1}{2}}. \quad \square$$

**Proposition 2.6** *The manifold  $\text{SPD}(n)$  of symmetric positive-definite real matrices is a Riemannian manifold of non-positive sectional curvature.*

□ **Proof of Proposition 2.6:**

The Levi-Civita connection is defined by Koszul formula (see for instance Theorem 3.1 page 54 in [2] where this formula is recalled) and coincides with the homogeneous connection since  $\text{SPD}(n)$  is locally symmetric and the Riemannian metric is  $G$ -invariant. The sectional curvature  $K_{\text{Id}}$  at the reference point  $\text{Id}$  is given by :

$$K_{\text{Id}}(X, Y) := \frac{\langle R_{X,Y} X, Y \rangle_{\text{Id}}}{\langle X, X \rangle_{\text{Id}} \langle Y, Y \rangle_{\text{Id}} - \langle X, Y \rangle_{\text{Id}}^2} = \frac{\langle [[X, Y], X], Y \rangle_{\text{Id}}}{\langle X, X \rangle_{\text{Id}} \langle Y, Y \rangle_{\text{Id}} - \langle X, Y \rangle_{\text{Id}}^2},$$

for all  $X, Y$  in  $T_{\text{Id}} \text{SPD}(n) = \text{Sym}(n)$ , where we have used formula (44) for the curvature (see also Proposition 7.72 page 193 in [3], or Proposition 6.5 page 92 in [2]). Now the sign of the sectional curvature of the 2-plane generated by  $X$  and  $Y$  is the sign of  $\langle [[X, Y], X], Y \rangle_{\text{Id}}$ . One has :

$$\begin{aligned} \langle [[X, Y], X], Y \rangle_{\text{Id}} &= \text{Tr} [[X, Y], X] Y = \text{Tr} ([X, Y] X Y - X [X, Y] Y) \\ &= \text{Tr} [X, Y] [X, Y] = -\text{Tr} [X, Y]^T [X, Y] \leq 0, \end{aligned}$$

where the last identity following from the fact that  $[X, Y]$  belongs to  $[\text{Sym}(n), \text{Sym}(n)] \subset \mathfrak{so}(n)$ . □

The following two Lemmas are standard results in the geometry of non-positively curved spaces.

**Lemma 2.7** *The Riemannian angle between two paths  $f$  and  $g$  intersecting at  $o = Id$  is equal to the Euclidian angle between the two paths  $\log(f)$  and  $\log(g)$  at 0. Moreover, in any geodesic triangle  $ABC$  in  $SPD(n)$ ,*

$$c^2 \geq a^2 + b^2 - 2ab \cos \widehat{ACB}, \quad (26)$$

where  $a, b, c$  are the lengths of the sides opposite to  $A, B, C$  respectively, and  $\widehat{ACB}$  the angle at  $C$ .

$\Delta$  **Proof of Lemma 2.7:**

This follows from the same arguments as in [10]. The Al-Kashi inequality (26) is also a direct consequence of Corollary 13.2 in [7] page 73, since Lemma 2.5 implies that  $SPD(n)$  is a minimizing convex normal ball.  $\Delta$

**Lemma 2.8** *Let  $\gamma_1(t)$  and  $\gamma_2(t)$  be two constant speed geodesics in  $SPD(n)$ . Then the distance in  $SPD(n)$  between  $\gamma_1(t)$  and  $\gamma_2(t)$  is a convex function of  $t$ .*

$\Delta$

$\Delta$  **Proof of Lemma 2.8:**

This follows from the fact that a non-positively curved Riemannian manifold is a CAT(0) space.

Let us recall the following definition.

**Definition 2.9** A Riemannian manifold  $\mathcal{P}$  is called *symmetric* if for every  $p$  in  $\mathcal{P}$ , there exists a globally defined isometry  $s_p$  which fixes  $p$  and such that the differential of  $s_p$  at  $p$  is  $-\text{id}$ . The transformation  $s_p$  is called a *global symmetry with respect to  $p$* .

**Proposition 2.10** *The manifold  $SPD(n)$  is a globally symmetric homogeneous Riemannian manifold. The globally defined symmetry with respect to  $x \in SPD(n)$  is*

$$s_x(y) = x^{\frac{1}{2}}(x^{-\frac{1}{2}}yx^{-\frac{1}{2}})^{-1}x^{\frac{1}{2}} = xy^{-1}x. \quad (27)$$

$\square$  **Proof of Proposition 2.10:**

Consider the inversion  $\text{Inv} : SPD(n) \rightarrow SPD(n)$  defined by  $\text{Inv}(x) = x^{-1}$ . An easy computation shows that the differential  $T_x \text{Inv}$  of  $\text{Inv}$  at  $x \in SPD(n)$  takes a tangent vector  $X \in T_x SPD(n)$  to  $T_x \text{Inv}(X) = -x^{-1}Xx^{-1}$ . One therefore has :

$$\begin{aligned} \langle T_x \text{Inv}(X), T_x \text{Inv}(Y) \rangle_{\text{Inv}(x)} &= \langle -x^{-1}Xx^{-1}, -x^{-1}Yx^{-1} \rangle_{x^{-1}} = \text{Tr } x(-x^{-1}Xx^{-1})x(-x^{-1}Yx^{-1}) \\ &= \text{Tr } Xx^{-1}Yx^{-1} = \text{Tr } x^{-1}Xx^{-1}Y = \langle X, Y \rangle_x, \end{aligned}$$

hence  $\text{Inv}$  is an isometry of  $SPD(n)$ . Since  $\text{Inv}$  fixes the reference point  $\text{Id} \in SPD(n)$  and its differential at the reference point is  $-\text{Id}$  on  $T_{\text{Id}} SPD(n) = \text{Sym}(n)$ , it follows that  $\text{Inv}$  is a global symmetry with respect to  $\text{Id}$ . From the transitive action of  $\text{GL}(n, \mathbb{R})$  one can define a globally defined symmetry with respect to any  $x \in SPD(n)$ . Indeed note that  $s_x$  is the composition of the isometry  $\iota_x : y \mapsto x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}}$ , with the inversion  $\text{Inv}$  followed by the isometry  $(\iota_x)^{-1} : y \mapsto x^{\frac{1}{2}} \cdot y \cdot x^{\frac{1}{2}}$ . Hence  $s_x$  is an isometry, with fixed point  $x$ . Its differential at  $x$  is given by

$$T(\iota_x)^{-1} \circ T_{\text{Id}} \text{Inv} \circ T_x \iota_x = T_{\text{Id}}(\iota_x)^{-1} \circ (-\text{Id}) \circ T_x \iota_x = -\text{Id}.$$

Whence  $SPD(n)$  is a (globally) symmetric homogeneous Riemannian manifold.  $\square$

### 3 Totally geodesic submanifolds of the manifold $SPD(n)$

**Definition 3.1** A submanifold  $\mathcal{N}$  of a Riemannian manifold  $\mathcal{M}$  is called *totally geodesic* if any geodesic on  $\mathcal{N}$  for the induced Riemannian metric is also a geodesic on  $\mathcal{M}$ . In particular, the property of being totally geodesic is a local property of the submanifold  $\mathcal{N}$  inside the ambient manifold  $\mathcal{M}$ .

**Definition 3.2** A submanifold  $\mathcal{N}$  of a Riemannian manifold  $\mathcal{M}$  is called *geodesically convex* if there exists a unique minimizing geodesic of  $\mathcal{M}$  connecting two points in  $\mathcal{N}$  and entirely contained in  $\mathcal{N}$ . In particular, the property of being geodesically convex depends on global behavior of geodesics.

The following Theorem is based on [10]. We give below a self-contained proof based on the following Lemma.



**Theorem 3.3 ([10])** Let  $E$  be a linear subspace of  $\text{Sym}(n)$ . The following assertions are equivalent :

1.  $[X, [X, Y]] \in E$  for all  $X, Y \in E$ ,
2.  $e \cdot f \cdot e \in \mathcal{E} := \exp(E)$  for all  $e, f \in \mathcal{E} = \exp(E)$ ,
3.  $\mathcal{E} := \exp(E)$  is a geodesically convex submanifold of  $\text{SPD}(n)$ .
4.  $\mathcal{E} := \exp(E)$  is a totally geodesic submanifold of  $\text{SPD}(n)$ .
5.  $E$  is a Lie triple system, i.e. for all  $X, Y, Z \in E$ ,  $[X, [Y, Z]] \in E$ .

**Lemma 3.4 :**

Let  $f \in \mathcal{E} = \exp(E)$ ,  $Y \in E$ , and consider the differentiable curve in  $\text{Sym}(n)$  defined by

$$X(t) := \log(\exp tY \cdot f \cdot \exp tY).$$

Then

$$\dot{X}(t) = \text{ad}(X(t)) \coth(\text{ad}(X(t)/2))(Y). \quad (28)$$

**△ Proof of Lemma 3.4:**

One has

$$\frac{d}{dt}|_{t=t_0} \exp tY = Y \cdot \exp t_0Y = \exp t_0Y \cdot Y,$$

hence the differential of the right hand side of equation (29) is

$$\frac{d}{dt}|_{t=t_0} \exp tY \cdot f \cdot \exp tY = Y \exp X(t_0) + \exp X(t_0)Y.$$

On the other hand, the differential of the left hand side of equation (29) is

$$\frac{d}{dt}|_{t=t_0} \exp X(t) = (d_{X(t_0)} \exp)(\dot{X}(t_0)).$$

Hence  $X(t)$  satisfies the following differential equation :

$$\dot{X}(t) = (d_{X(t)} \exp)^{-1}(Y \cdot \exp X(t) + \exp X(t) \cdot Y).$$

By Lemma A.3, equation (41),

$$(d_X \exp)^{-1}(Z) = \frac{\text{ad}(X/2)}{\sinh(\text{ad}(X/2))} \left( \exp\left(-\frac{X}{2}\right) \cdot Z \cdot \exp\left(-\frac{X}{2}\right) \right)$$

It follows that

$$\begin{aligned} \dot{X}(t) &= \frac{\text{ad}(X(t)/2)}{\sinh(\text{ad}(X(t)/2))} \left( \exp\left(-\frac{X(t)}{2}\right) \cdot (Y \cdot \exp X(t) + \exp X(t) \cdot Y) \cdot \exp\left(-\frac{X(t)}{2}\right) \right) \\ &= \frac{\text{ad}(X(t)/2)}{\sinh(\text{ad}(X(t)/2))} \left( \exp\left(-\frac{X(t)}{2}\right) \cdot Y \cdot \exp\left(\frac{X(t)}{2}\right) + \exp\left(\frac{X(t)}{2}\right) \cdot Y \cdot \exp\left(-\frac{X(t)}{2}\right) \right) \\ &= \frac{\text{ad}(X(t)/2)}{\sinh(\text{ad}(X(t)/2))} \left( \text{Ad}\left(\exp\left(-\frac{X(t)}{2}\right)\right)(Y) + \text{Ad}\left(\exp\left(\frac{X(t)}{2}\right)\right)(Y) \right) \\ &= \frac{\text{ad}(X(t)/2)}{\sinh(\text{ad}(X(t)/2))} \left( \exp\left(\text{ad}\left(-\frac{X(t)}{2}\right)\right)(Y) + \exp\left(\text{ad}\left(\frac{X(t)}{2}\right)\right)(Y) \right) \\ &= \frac{\text{ad}(X(t))}{\sinh(\text{ad}(X(t)/2))} (\cosh(\text{ad}(X(t)/2)))(Y) \\ &= \text{ad}(X(t)) \coth(\text{ad}(X(t)/2))(Y) \end{aligned}$$

△

**■ Proof of Theorem 3.3:**

$1 \Rightarrow 2$  : Let  $f \in \mathcal{E}$  and  $Y \in E$ . We will show that

$$\exp X(t) = \exp tY \cdot f \cdot \exp tY \quad (29)$$

belongs to  $\mathcal{E} = \exp(\mathbb{E})$  for all  $t \in \mathbb{R}$ . By Lemma 3.4,

$$\dot{X}(t) = \text{ad}(X(t)) \coth(\text{ad}(X(t)/2))(Y).$$

Consider the vector field  $W$  on  $\text{Sym}(n)$  defined by

$$W(X) := \text{ad}(X) \coth(\text{ad}(X/2))(Y), X \in \text{Sym}(n).$$

Note that only even powers of  $\text{ad}(X)$  are involved in the operator  $\text{ad}(X) \coth(\text{ad}(X/2))$ . Since  $\text{ad}(X) \circ \text{ad}(X)(Y) \in \mathbb{E}$  for  $X, Y \in \mathbb{E}$ , the vector field  $W(X)$  is tangent to  $\mathbb{E}$  for every  $X \in \mathbb{E}$ . It follows that an integral curve  $X(t)$  of  $W$  starting at  $X(0) \in \mathbb{E}$  stays in  $\mathbb{E}$ . Moreover the flow of this vector field is defined for all  $t \in \mathbb{R}$ . Thus setting  $t = 1$  and  $Y = \log e$  with  $e \in \mathcal{E} = \exp \mathbb{E}$ , give the result  $e \cdot f \cdot e \in \mathcal{E}$ .

2  $\Rightarrow$  3 : Suppose that for all  $e$  and  $f$  in  $\mathcal{E} = \exp \mathbb{E}$ , the product  $e \cdot f \cdot e$  belongs to  $\mathcal{E} = \exp \mathbb{E}$ . Consider  $f \in \mathcal{E} = \exp \mathbb{E}$ . By Lemma 2.5, the geodesic of  $\text{SPD}(n)$  joining  $\text{Id} \in \mathcal{E} = \exp \mathbb{E}$  to  $f$  is  $t \mapsto \exp(t \log(f))$  hence lies entirely in the manifold  $\mathcal{E} = \exp \mathbb{E}$ . In particular, its mid-point  $\exp(\frac{1}{2} \log(f)) =: f^{\frac{1}{2}}$  belongs to  $\mathcal{E} = \exp \mathbb{E}$ . In other words, the square roots of elements in  $\mathcal{E} = \exp \mathbb{E}$  belong to  $\mathcal{E} = \exp \mathbb{E}$ . Consider now  $x$  and  $y$  two elements in  $\mathcal{E} = \exp \mathbb{E}$ . By hypothesis,  $x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}}$  belongs to  $\mathcal{E} = \exp \mathbb{E}$ , hence  $\log x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}}$  belongs to  $\mathbb{E}$  and the geodesic

$$t \mapsto \exp(t \log x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}})$$

joining  $\text{Id}$  to  $x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}}$  lies entirely in  $\mathcal{E} = \exp \mathbb{E}$ . Since  $x \in \mathcal{E} = \exp \mathbb{E}$ , by hypothesis the isometry  $z \mapsto x^{\frac{1}{2}} \cdot z \cdot x^{\frac{1}{2}}$  preserves  $\mathcal{E} = \exp \mathbb{E}$  and sends the previous geodesic to the unique geodesic of  $\text{SPD}(n)$  joining  $x$  to  $y$ , which therefore lies completely in  $\mathcal{E} = \exp \mathbb{E}$ . Consequently, the space  $\mathcal{E} = \exp \mathbb{E}$  is geodesically convex in  $\text{SPD}(n)$ .

3  $\Rightarrow$  4 : Suppose that  $\mathcal{E} = \exp \mathbb{E}$  is geodesically convex in  $\text{SPD}(n)$ . By the uniqueness of the geodesic joining two points in  $\text{SPD}(n)$  proved in Theorem 2.5, the geodesic of  $\text{SPD}(n)$  connecting two points  $x$  and  $y$  belonging to  $\mathcal{E} = \exp \mathbb{E}$  minimises the distance between  $x$  and  $y$  in  $\text{SPD}(n)$  and by hypothesis lies entirely in  $\mathcal{E} = \exp \mathbb{E}$ . Therefore it is also a geodesic of  $\mathcal{E} = \exp \mathbb{E}$ . Now consider an arbitrary geodesic  $t \mapsto \gamma(t)$  in  $\mathcal{E} = \exp \mathbb{E}$ , with  $t$  in some open interval  $I$ , and let us show that it is a geodesic in  $\text{SPD}(n)$ . For every  $t_0 \in I$ , there exists an  $\varepsilon > 0$  such that  $(t_0 - \varepsilon, t_0 + \varepsilon) \ni t \mapsto \gamma(t)$  is the unique geodesic in  $\mathcal{E} = \exp \mathbb{E}$  between  $\gamma(t_0 - \varepsilon)$  and  $\gamma(t_0 + \varepsilon)$ . By uniqueness, it is the geodesic segment in  $\text{SPD}(n)$  connecting  $\gamma(t_0 - \varepsilon)$  and  $\gamma(t_0 + \varepsilon)$ . Therefore  $t \mapsto \gamma(t)$  is a geodesic in  $\text{SPD}(n)$ .

4  $\Rightarrow$  2 : Suppose that  $\mathcal{E} = \exp \mathbb{E}$  is a totally geodesic submanifold of  $\text{SPD}(n)$ . Let us consider the symmetry  $s_x$  with respect to  $x \in \text{SPD}(n)$  defined from  $\text{SPD}(n)$  to  $\text{SPD}(n)$  by  $s_x : y \mapsto xy^{-1}x$  (see Proposition 2.10). Every geodesic of the form  $t \mapsto x^{\frac{1}{2}} \exp(tY)x^{\frac{1}{2}}$  is mapped to  $t \mapsto x^{\frac{1}{2}} \exp(-tY)x^{\frac{1}{2}}$  by  $s_x$ . It follows that every geodesic containing  $x$  is stable under  $s_x$ . Consequently  $s_x(\exp \mathbb{E}) \subset \exp \mathbb{E}$  for every  $x \in \exp \mathbb{E}$ . In particular, for  $e$  and  $f$  in  $\mathcal{E} = \exp \mathbb{E}$ ,

$$e \cdot f \cdot e = e^2 \cdot (e \cdot f^{-1} \cdot e)^{-1} \cdot e^2 = s_{e^2}(s_e(f))$$

belongs to  $\mathcal{E} = \exp \mathbb{E}$ .

2  $\Rightarrow$  1: Suppose that  $e \cdot f \cdot e \in \mathcal{E} := \exp(\mathbb{E})$  for all  $e, f \in \mathcal{E} = \exp(\mathbb{E})$ . Given  $X$  and  $Y$  in  $\mathbb{E}$ , consider the smooth function in  $\text{Sym}(n)$  defined by

$$Z(s, t) = \log(\exp tY \cdot \exp sX \cdot \exp tY).$$

In particular,

$$Z(s, 0) = \log(\exp sX) = sX.$$

By hypothesis,  $Z(s, t)$  belongs to  $\mathbb{E}$  for all  $(s, t) \in \mathbb{R}^2$ . Since  $\mathbb{E}$  is a vector space, all the derivatives of the function  $(s, t) \mapsto Z(s, t)$  belong to  $\mathbb{E}$  for all  $(s, t) \in \mathbb{R}^2$ . By Lemma 3.4,

$$\frac{\partial}{\partial t} Z(s, t) = \text{ad}(Z(s, t)) \coth(\text{ad}(Z(s, t)/2))(Y).$$

In particular

$$\frac{\partial}{\partial t} Z(s, t) \Big|_{t=0} = \text{ad}(Z(s, 0)) \coth(\text{ad}(Z(s, 0)/2))(Y) = \text{ad}(sX) \coth(\text{ad}(sX/2))(Y)$$

belongs to  $\mathbb{E}$  for all  $s \in \mathbb{R}$ . Consider the Taylor series of  $\frac{u}{2} \coth\left(\frac{u}{2}\right)$  in the neighborhood of  $u = 0$ :

$$\frac{u}{2} \coth\left(\frac{u}{2}\right) = 1 + \frac{u^2}{12} + u^2 \varepsilon(u), \text{ with } \lim_{u \rightarrow 0} \varepsilon(u) = 0.$$

It follows that

$$\frac{\partial}{\partial t}\Big|_{t=0} Z(s, t) = 2 \left( Y + \frac{1}{12} \text{ad}(sX) \circ \text{ad}(sX)(Y) + s^2 W \right) \in \mathbb{E} \quad \text{with} \quad \lim_{s \rightarrow 0} W = 0.$$

Therefore

$$\lim_{s \rightarrow 0} \frac{\frac{\partial}{\partial t}\Big|_{t=0} Z(s, t) - 2Y}{s^2} = \frac{1}{6} [X, [X, Y]] \in \mathbb{E}.$$

1  $\Rightarrow$  5: Consider  $X, Y, Z$  in  $\mathbb{E}$ . Developping  $[X + Y, [X + Y, Z]] \in \mathbb{E}$  we get

$$[X, [Y, Z]] + [Y, [X, Z]] \in \mathbb{E}.$$

By Jacobi identity,

$$[Y, [X, Z]] = [[Y, X], Z] + [X, [Y, Z]] \in \mathbb{E},$$

hence

$$A := [[Y, X], Z] + 2[X, [Y, Z]] \in \mathbb{E}.$$

Interchanging  $X$  and  $Z$  we also have

$$B := [[Y, Z], X] + 2[Z, [Y, X]] \in \mathbb{E}.$$

It follows that

$$2A + B = 2[[Y, X], Z] + 4[X, [Y, Z]] + [[Y, Z], X] + 2[Z, [Y, X]] = 3[X, [Y, Z]] \in \mathbb{E}.$$

5  $\Rightarrow$  1 Obvious by taking  $X = Y$ . ■

## 4 Orthogonal projection on a totally geodesic submanifold

The proof of Mostow's decomposition theorem given in [10] is based on the existence of an orthogonal projection from  $\text{SPD}(n)$  onto a totally geodesic submanifold  $\mathcal{E} := \exp \mathbb{E}$  which follows from compactness arguments. Here we use the completeness of  $\mathcal{E} := \exp \mathbb{E}$  to obtain this projection. Recall from [4] the following Proposition (we sketch the proof for the convenience of the reader).

**Proposition 4.1** *The manifold  $\text{SPD}(n)$  endowed with the distance induced from the Riemannian metric (22) is a complete metric space. For any linear subspace  $\mathbb{E}$  of  $\text{Sym}(n)$ , the manifold  $\mathcal{E} := \exp \mathbb{E}$ , endowed with the induced Riemannian metric, is closed in  $\text{SPD}(n)$  hence also a complete metric space.*

□ **Proof of Proposition 4.1** By (37), the exponential map  $\exp : \text{Sym}(n) \rightarrow \text{SPD}(n)$  increases distances, hence a Cauchy sequence in  $\text{SPD}(n)$  is the image by the exponential map of a Cauchy sequence in  $\text{Sym}(n)$  which is complete. By Theorem 2.3, it follows that  $\text{SPD}(n)$  is complete. The remainder follows from the fact that a linear subspace  $\mathbb{E}$  is closed in  $\text{Sym}(n)$ . □

**Theorem 4.2** *Let  $\mathbb{E}$  be a linear subspace of  $\text{Sym}(n)$  such that  $[X, [X, Y]] \in \mathbb{E}$ , for all  $X, Y \in \mathbb{E}$ . Then there exists a continuous orthogonal projection from  $\text{SPD}(n)$  onto the totally geodesic submanifold  $\mathcal{E} := \exp \mathbb{E}$ , i.e. a continuous map  $\pi : \text{SPD}(n) \rightarrow \mathcal{E}$  satisfying*

$$\text{dist}(x, \exp \mathbb{E}) = \text{dist}(x, \pi(x))$$

*and such that the geodesic joining  $x$  to  $\pi(x)$  is orthogonal to every geodesic starting at  $\pi(x)$  and included in  $\mathcal{E} := \exp \mathbb{E}$ .*

■ **Proof of Theorem 4.2:**

Let  $x$  be a element of  $\text{SPD}(n)$ . Denote by  $\delta$  the distance between  $x$  and  $\mathcal{E} := \exp \mathbb{E}$  in  $\text{SPD}(n)$  and let  $\{e_n\}_{n \in \mathbb{N}}$  be a minimizing sequence in  $\mathcal{E} := \exp \mathbb{E}$  thus that

$$\text{dist}(x, e_n)^2 \leq \delta^2 + \frac{1}{n}.$$

Let us show that  $\{e_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathcal{E} := \exp \mathbb{E}$ . For this purpose, consider for  $k > n$  the geodesic  $\gamma(t)$  joining  $e_n =: \gamma(0) \in \mathcal{E}$  to  $e_k =: \gamma(1) \in \mathcal{E}$ . This geodesic lies in  $\mathcal{E}$  since  $\mathcal{E}$  is a totally geodesic submanifold of  $\text{SPD}(n)$ , and is of the form:

$$\gamma(t) = e_n^{\frac{1}{2}} \exp(tH) e_n^{\frac{1}{2}},$$




Figure 1: Illustration of the orthogonal projection  $\pi(p)$  of a point  $p \in SPD(n)$  onto a totally geodesic submanifold  $\exp E$  and the translation of the fiber over  $\pi(p)$  by the action of  $\pi(p)^{-\frac{1}{2}}$ .




Figure 2: Illustration of geodesic triangles used in the proof of Theorem 4.2.

where  $H$  belongs to  $E$ . Denote by  $e_{n,k}$  the middle of the geodesic joining  $e_n$  to  $e_k$ , i.e.

$$e_{n,k} = e_n^{\frac{1}{2}} \exp\left(\frac{H}{2}\right) e_n^{\frac{1}{2}}.$$

By lemma 2.7 applied to the geodesic triangle joining  $x$ ,  $e_n$  and  $e_{n,k}$ , we have:

$$\text{dist}(x, e_n)^2 \geq \text{dist}(e_n, e_{n,k})^2 + \text{dist}(e_{n,k}, x)^2 - 2\text{dist}(e_n, e_{n,k})\text{dist}(e_{n,k}, x) \cos \widehat{e_n e_{n,k} x}. \quad (30)$$

On the other hand, lemma 2.7 applied to the geodesic triangle joining  $x$ ,  $e_k$  and  $e_{n,k}$  gives:

$$\text{dist}(x, e_k)^2 \geq \text{dist}(e_k, e_{n,k})^2 + \text{dist}(e_{n,k}, x)^2 - 2\text{dist}(e_k, e_{n,k})\text{dist}(e_{n,k}, x) \cos \widehat{e_k e_{n,k} x}. \quad (31)$$

By definition of  $e_{n,k}$  we have:  $\text{dist}(e_k, e_{n,k}) = \text{dist}(e_n, e_{n,k})$ . Moreover since the geodesic  $\gamma$  is a smooth curve, the sum of the following angles is the flat one:

$$\widehat{e_k e_{n,k} x} + \widehat{e_n e_{n,k} x} = \pi,$$

and  $\cos \widehat{e_k e_{n,k} x} = -\cos \widehat{e_n e_{n,k} x}$ . Summing inequalities (30) and (31), we obtain:

$$\text{dist}(x, e_n)^2 + \text{dist}(x, e_k)^2 \geq 2\text{dist}(e_k, e_{n,k})^2 + 2\text{dist}(e_{n,k}, x)^2.$$

It follows that:

$$\begin{aligned} \text{dist}(e_k, e_{n,k})^2 &\leq \frac{1}{2} (\text{dist}(x, e_n)^2 + \text{dist}(x, e_k)^2) - \text{dist}(e_{n,k}, x)^2 \\ &\leq \frac{1}{2} \left( \delta^2 + \frac{1}{n} + \delta^2 + \frac{1}{k} \right) - \delta^2 \\ &\leq \frac{1}{2} \left( \frac{1}{n} + \frac{1}{k} \right). \end{aligned}$$

This yields that

$$\text{dist}(e_n, e_k) \leq \sqrt{2} \left( \frac{1}{n} + \frac{1}{k} \right)^{\frac{1}{2}}.$$

Consequently  $\{e_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathcal{E} = \exp E$ . By Proposition 4.1,  $\mathcal{E}$  is a complete metric space hence the sequence  $\{e_n\}_{n \in \mathbb{N}}$  converges to a element  $\pi(x)$  in  $\mathcal{E}$  satisfying:

$$\text{dist}(x, \pi(x)) = \text{dist}(x, \mathcal{E}).$$

Denote by  $\alpha(t)$  the constant speed geodesic which satisfies  $\alpha(0) = \pi(x)$  and  $\alpha(1) = x$ . By uniqueness of the geodesic joining two points in  $\text{SPD}(n)$ , it follows that the length of  $\alpha$  is  $\text{dist}(x, \mathcal{E})$ . The map

$$y \mapsto (\pi(x))^{-\frac{1}{2}} y (\pi(x))^{-\frac{1}{2}}$$

being an isometry of  $\text{SPD}(n)$  which preserves  $\mathcal{E}$ , the curve  $t \mapsto (\pi(x))^{-\frac{1}{2}} \alpha(t) (\pi(x))^{-\frac{1}{2}}$  is a geodesic whose length is the distance between

$$(\pi(x))^{-\frac{1}{2}} \alpha(1) (\pi(x))^{-\frac{1}{2}} = (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$$

and  $\mathcal{E}$ . Therefore the projection of  $(\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$  onto  $\mathcal{E}$  is  $(\pi(x))^{-\frac{1}{2}} \alpha(0) (\pi(x))^{-\frac{1}{2}} = \text{Id}$ . From lemma 2.5 it follows that:

$$(\pi(x))^{-\frac{1}{2}} \alpha(t) (\pi(x))^{-\frac{1}{2}} = \exp tV,$$

for some  $V$  in  $\text{Sym}(n)$ . Since the length of  $t \mapsto \exp tV$  is  $\|V\|$ ,  $V$  is in  $F$  and  $(\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$  is in  $\exp F$ . Since  $E \perp F$ , by lemma 2.7,  $(\pi(x))^{-\frac{1}{2}} \alpha(\pi(x))^{-\frac{1}{2}}$  is orthogonal at the identity to every curve starting at the identity and contained in  $\exp E$ . Therefore  $\alpha$  is orthogonal at  $\pi(x)$  to every curve starting at  $\pi(x)$  and contained in  $\exp E$ .

To show that  $\pi$  is continuous, denote by  $\gamma(t)$  (resp.  $\alpha(t)$ ) the geodesic joining a point  $x_1$  (resp.  $x_2$ ) in  $\text{SPD}(n)$  to its projection onto  $\exp E$ , with  $\gamma(0) = \pi(x_1)$  (resp.  $\alpha(0) = \pi(x_2)$ ) and  $\gamma(1) = x_1$  (resp.  $\alpha(1) = x_2$ ). By the negative curvature property stated in Lemma 2.8, the map  $t \mapsto \text{dist}(\gamma(t), \alpha(t))$  is convex. Since, for  $t = 0$ ,  $\gamma(t)$  and  $\alpha(t)$  are orthogonal to the geodesic joining  $\pi(x_1)$  and  $\pi(x_2)$  (which is contained in  $\mathcal{E}$ ), the minimum of the distance between  $\gamma(t)$  and  $\alpha(t)$  is reached for  $t = 0$ , and  $\text{dist}(x_1, x_2) \geq \text{dist}(\pi(x_1), \pi(x_2))$ .  $\blacksquare$

## 5 Mostow's decomposition Theorem for $GL(n, \mathbb{R})$

**Theorem 5.1** *Let  $E$  be a linear subspace of  $\text{Sym}(n)$  such that:*

$$[X, [X, Y]] \in E, \quad \text{for all } X, Y \in E,$$

*and let  $F$  be its orthogonal in  $\text{Sym}(n)$ :*

$$F := E^\perp = \{ X \in \text{Sym}(n) \mid \text{Tr } XY = 0, \quad \forall Y \in E \}.$$

*Then for all symmetric positive-definite operator  $x$  in  $\text{SPD}(n)$ , there exist a unique element  $e \in \mathcal{E} := \exp E$  and a unique element  $f \in \mathcal{F} := \exp F$  such that  $x = efe$ . More precisely,*

$$\begin{aligned} e &:= (\pi(x))^{-\frac{1}{2}} \in \mathcal{E} = \exp E \\ f &:= (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}} \in \mathcal{F} := \exp F, \end{aligned}$$

*where  $\pi$  denotes the projection onto the totally geodesic submanifold  $\exp E$ . Moreover the map defined from  $\text{SPD}(n)$  to  $\exp E \times \exp F$  taking  $A$  to  $(e, f)$  is a homeomorphism.*

■ **Proof of Theorem 5.1:**

Let us introduce the map

$$\begin{aligned} \Upsilon : \exp E \times \exp F &\longrightarrow \text{SPD}(n) \\ (e, f) &\longmapsto efe \end{aligned}$$

Let us show that  $\Upsilon$  is one-one. Suppose that  $(e_1, f_1)$  and  $(e_2, f_2)$  are elements of  $\exp E \times \exp F$  such that  $e_1 f_1 e_1 = e_2 f_2 e_2$ . Consider the geodesic triangle joining  $e_1 f_1 e_1$ ,  $e_1^2$  and  $e_2^2$ . By Theorem 3.3,  $\exp E$  is a geodesic subspace of  $\text{SPD}(n)$ . Thus the geodesic joining  $e_1^2$  to  $e_2^2$  lies in  $\exp E$ . On the other hand the geodesic joining  $e_1 f_1 e_1$  to  $e_1^2$  lies in  $e_1 \exp F e_1$ . Since  $E$  is perpendicular to  $F$  at zero,  $\exp E$  is perpendicular to  $\exp F$  at the identity by lemma 2.7. Since the map taking any  $x \in \text{SPD}(n)$  to  $e_1 x e_1 \in \text{SPD}(n)$  is an isometry, the manifold  $e_1 \exp F e_1$  is perpendicular to the manifold  $e_1 \exp E e_1 = \exp E$  at  $e_1^2$ . Hence the angle at  $e_1^2$  of the above geodesic triangle is  $90^\circ$ . Similarly, the angle at  $e_2^2$  is  $90^\circ$  since it is formed by the geodesic joining  $e_2^2$  to  $e_2 f_2 e_2 = e_1 f_1 e_1$  which lies in  $e_2 \exp F e_2$  and the geodesic joining  $e_2^2$  to  $e_1^2$  which lies in  $\exp E$ . Denoting by  $a$  the length of the side of the geodesic triangle joining  $e_1^2$  to  $e_2^2$ ,  $b$  the length of the side joining  $e_1 f_1 e_1$  to  $e_1^2$  and  $c$  the length of the side joining  $e_1 f_1 e_1$  to  $e_2^2$ , one has  $c^2 \geq b^2 + a^2$  and  $b^2 \geq c^2 + a^2$  by lemma 2.7. This implies that  $a = 0$  and  $e_1^2 = e_2^2$ . It follows that  $e_1 = e_2$  and  $f_1 = f_2$ .

Let us show that  $\Upsilon$  is onto. Consider  $x$  in  $\text{SPD}(n)$ . By Theorem 4.2, the geodesic joining  $x$  to  $\pi(x) \in \exp E$  is orthogonal to every geodesic starting at  $\pi(x)$  and contained in  $\exp E$ . Denote by  $\gamma$  the geodesic satisfying  $\gamma(0) = \text{Id} \in \text{SPD}(n)$  and  $\gamma(1) = (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$ . Since  $y \mapsto (\pi(x))^{-\frac{1}{2}} y (\pi(x))^{-\frac{1}{2}}$  is an isometry,  $\gamma$  is orthogonal to every geodesic starting at the identity and contained in

$$(\pi(x))^{-\frac{1}{2}} \cdot \exp E \cdot (\pi(x))^{-\frac{1}{2}} = \exp E.$$

By lemma 2.7,  $\gamma$  is tangent to  $F = E^\perp$  at the identity and since it is of the form  $t \mapsto \exp tH$  by lemma 2.5, we have  $H$  in  $F$ . It follows that  $\gamma(1) = \exp H = (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$  is in  $\exp F$ . Therefore  $x = efe$  with  $e := (\pi(x))^{-\frac{1}{2}}$  in  $\exp E$  and  $f := (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$  in  $\exp F$ . Consequently  $\Upsilon$  is onto.

The continuity of the map that takes  $x$  to  $(e, f) \in \exp E \times \exp F$  with  $x = efe$ , i.e.  $e := (\pi(x))^{-\frac{1}{2}}$  and  $f := (\pi(x))^{-\frac{1}{2}} x (\pi(x))^{-\frac{1}{2}}$  follows directly from the continuity of the projection  $\pi$ . ■

**Theorem 5.2 (Mostow's Decomposition)** *Let  $E$  and  $F$  be as in Theorem 5.1. Then  $GL(n, \mathbb{R})$  is homeomorphic to the product  $O(n) \times \exp F \times \exp E$ .*

■ **Proof of Theorem 5.2:**

Denote by  $\Theta$  the map from  $O(n) \times \exp E \times \exp F$  to  $GL(n, \mathbb{R})$  that takes  $(k, f, e)$  to the product of matrices  $k \cdot f \cdot e$ .

Let us show that  $\Theta$  is one-one. Suppose that  $a = k_1 f_1 e_1 = k_2 f_2 e_2$  with  $(k_1, f_1, e_1)$  and  $(k_2, f_2, e_2)$  in  $O(n) \times \exp E \times \exp F$ . We have

$$a^T a = e_1 f_1^2 e_1 = e_2 f_2^2 e_2.$$

Since  $f_1^2$  and  $f_2^2$  are in  $\exp F$ , by Theorem 5.1, it follows that  $e_1 = e_2$  and  $f_1^2 = f_2^2$ . Thus  $f_1 = f_2$  and  $k_1 = k_2$ .

Let us show that  $\Theta$  is onto. Consider  $x$  in  $\text{GL}(n, \mathbb{R})$ . By Theorem 5.1, since  $x^T x$  is an element of  $\text{SPD}(n)$ , there exist  $e \in \exp E$  and  $f \in \exp F$  such that  $x^T x = ef^2e$ . Let  $k$  be  $x(fe)^{-1}$ . We have:

$$k^T k = ((fe)^{-1})^T x^T x (fe)^{-1} = (e^{-1}f^{-1})^T x^T x e^{-1}f^{-1} = f^{-1}e^{-1}(ef^2e)e^{-1}f^{-1} = \text{id}.$$

Thus  $k$  is in  $\text{O}(n)$  and  $x = kfe$ .

The continuity of the map that takes  $x$  in  $\text{GL}(n, \mathbb{R})$  to  $(k, f, e)$  in  $\text{O}(n) \times \exp F \times \exp E$  follows from the continuity of the map that takes  $x$  to  $x^T x$  and from Theorem 5.1.  $\blacksquare$

**■ Proof of Theorem 1.1:**

The Theorem of the Introduction is now a summary of previous considerations.  $\blacksquare$

$\square$  **Proof of Corollary 1.2:**

Consider  $N$  a geodesically complete and convex submanifold in  $\text{SPD}(n)$ . Let  $x$  be an arbitrary element in  $N$ . Then the image of  $N$  by the isometry  $y \mapsto x^{-\frac{1}{2}} \cdot y \cdot x^{-\frac{1}{2}}$  is a geodesically complete and convex submanifold in  $\text{SPD}(n)$  containing  $\text{Id}$ . By Theorem 1.1 it is of the form  $\exp E$  with  $E$  satisfying  $[X, [X, Y]] \in E$ .  $\square$

## A The differential of the exponential map of a Lie group $G$

### A.1 First expression of the differential of the exponential map

The following Proposition explicits the differential of the exponential map and is well-known in the theory of finite-dimensional linear group. For the sake of completeness, we give below a proof using the canonical connection on the tangent bundle of a Lie group  $G$ , which was explained to us by P. Gauduchon. The reader will find the computation of the differential of the exponential map using powers series as a consequence of Lemma 1 in [10] (this computation works as well in the infinite-dimensional setting, see for instance Proposition 2.5.3 page 116 in [16]). See also [6].

Let us introduce the connection on the tangent bundle  $TG$  of  $G$  for which the left-invariant vector fields are parallel. It is a flat connection since a trivialization of the tangent bundle is given by the left-invariant vector fields associated to an arbitrary basis of the Lie algebra  $\mathfrak{g}$  of  $G$ . We give in Proposition A.1 the corresponding covariant derivative  $\nabla$  in terms of the (left) Maurer-Cartan  $\mathfrak{g}$ -valued 1-form  $\theta : TG \rightarrow \mathfrak{g}$  defined by

$$\theta_g(W) = (L_{g^{-1}})_*(W),$$

where  $g \in G$ ,  $W \in T_g G$ , and where  $(L_{g^{-1}})_*$  denotes the differential of the left multiplication by  $g^{-1}$ . For a matrix group  $G$ ,  $(L_{g^{-1}})_*(W) = g^{-1}W$ .

**Proposition A.1** *Let  $W$  be a vector field on  $G$ , i.e. a section of the tangent bundle  $TG$ . For a tangent vector  $Z$  in  $T_g G$ , define*

$$(\nabla_Z W)(g) := (L_g)_* \left( \frac{d}{dt} \Big|_{t=0} \theta_{\gamma_Z(t)} (W(\gamma_Z(t))) \right) \in T_g G \quad (32)$$

where  $\gamma_Z$  is any smooth curve in  $G$  with  $\gamma_Z(0) = g$  and  $\frac{d}{dt} \Big|_{t=0} \gamma_Z(t) = Z$ . Then  $\nabla$  defines a connection for which left-invariant vector fields are parallel.

$\square$  **Proof of Proposition** Note that  $t \mapsto \theta_{\gamma_Z(t)} (W(\gamma_Z(t)))$  is a smooth curve in  $\mathfrak{g}$ , hence its derivative belongs to  $\mathfrak{g}$ . For a left-invariant vector field  $W$  this curve is constant, hence its derivative vanishes. In order to verify that formula (32) defines indeed a connection, we have to show that

$$\nabla_Z(fW) = df(Z)W + f\nabla_Z W,$$

for any function  $f$  on  $G$ , and any vector field  $W$ . One has

$$\begin{aligned}
(\nabla_Z(fW))(g) &= (L_g)_* \left( \frac{d}{dt} \Big|_{t=0} \theta_{\gamma_Z(t)} (f(\gamma_Z(t)) W(\gamma_Z(t))) \right) \\
&= (L_g)_* \left( \frac{d}{dt} \Big|_{t=0} (L_{\gamma_Z(t)^{-1}})_* (f(\gamma_Z(t)) W(\gamma_Z(t))) \right) \\
&= f(g) (L_g)_* \left( \frac{d}{dt} \Big|_{t=0} (L_{\gamma_Z(t)^{-1}})_* (W(\gamma_Z(t))) \right) \\
&\quad + (L_g)_* \left( \frac{d}{dt} \Big|_{t=0} (L_{g^{-1}})_* (f(\gamma_Z(t)) W(g)) \right) \\
&= f(g) (\nabla_Z W)(g) + df(Z)W(g)
\end{aligned}$$

□

**Proposition A.2** For every Lie group  $G$ , with Lie algebra  $\mathfrak{g}$ , the differential of the exponential map  $\exp : \mathfrak{g} \rightarrow G$  is given at  $X \in \mathfrak{g}$  by :

$$(d_X \exp)(Y) = L_{\exp(X)} \left( \frac{1 - e^{-\text{ad}(X)}}{\text{ad}(X)} \right) (Y). \quad (33)$$

for all  $Y$  in  $\mathfrak{g}$ .

□ **Proof of Proposition A.2:**

Let us define the following map :

$$\begin{aligned}
\Phi : \mathbb{R}^2 &\longrightarrow G \\
(t, s) &\longmapsto \exp(t(X + sY)).
\end{aligned}$$

Consider the push-forward  $U$  and  $V$  of the vector fields  $\frac{\partial}{\partial t}$  and  $\frac{\partial}{\partial s}$  on  $\mathbb{R}^2$  :

$$U(\Phi(t, s)) := \Phi_* \left( \frac{\partial}{\partial t} \right) \quad \text{and} \quad V(\Phi(t, s)) = \Phi_* \left( \frac{\partial}{\partial s} \right).$$

Denote by  $[\cdot, \cdot]_{\mathfrak{X}}$  the bracket of vector fields. One has:

$$[U, V]_{\mathfrak{X}} = \left[ \Phi_* \left( \frac{\partial}{\partial t} \right), \Phi_* \left( \frac{\partial}{\partial s} \right) \right]_{\mathfrak{X}} = \Phi_* \left[ \frac{\partial}{\partial t}, \frac{\partial}{\partial s} \right]_{\mathfrak{X}} = 0. \quad (34)$$

Note that

$$V(\Phi(t, s)) = \frac{\partial \Phi}{\partial s}(t, s) = \left( d \exp_{(tX + sY)} \right) (tY) \quad \text{and} \quad V(\Phi(1, 0)) = (d_X \exp)(Y).$$

The idea of this proof is to explicit the differential equation satisfied by the  $\mathfrak{g}$ -valued function

$$v(t) := \left( L_{\Phi(t, 0)}^{-1} \right)_* V(\Phi(t, 0)) = \left( L_{\exp(tX)}^{-1} \right)_* V(\exp(tX)).$$

For this purpose we will use the connection  $\nabla$  on the tangent bundle of  $G$  defined by (32). Note that by the very definition of the exponential map on a Lie group,  $t \mapsto \exp(tX) = \Phi(t, 0)$  is a geodesic for this connection, hence

$$\nabla_U U = 0 \quad (35)$$

along  $\Phi(t, 0)$ . Let us denote by  $T$  and  $R$  the torsion and the curvature of  $\nabla$ . By definition :

$$T(U, V) := \nabla_U V - \nabla_V U - [U, V]_{\mathfrak{X}}$$

$$\text{and} \quad R_{U, V} U := \nabla_V \nabla_U U - \nabla_U \nabla_V U - \nabla_{[U, V]_{\mathfrak{X}}} U.$$

By (34), one has

$$\nabla_U V = \nabla_V U + T(U, V),$$

hence

$$\nabla_U (\nabla_U V) = \nabla_U \nabla_V U + \nabla_U T(U, V).$$



But the curvature tensor vanishes, hence (34) and (35) imply

$$\nabla_U \nabla_V U = \nabla_V \nabla_U U - \nabla_{[V,U]_x} U = 0.$$

Consequently one has

$$\nabla_U (\nabla_U V) = \nabla_U T(U, V).$$

By the expression (32) of the connection, one has

$$T(U, V)(\Phi(t, 0)) = (\nabla_U V - \nabla_V U)(\Phi(t, 0)) = (L_{\Phi(t, 0)})_* (U \cdot \theta(V) - V \cdot \theta(U)).$$

Let us recall that the torsion is a tensor, hence  $T(U, V)(\Phi(t, 0))$  does not depend on the extensions of the vectors  $U(\Phi(t, 0))$  and  $V(\Phi(t, 0))$  into vector fields. Using the left-invariant extensions of these two vectors one see easily that by the very definition of the bracket in the Lie algebra  $\mathfrak{g}$  one has

$$T(U, V)(\Phi(t, 0)) = - (L_{\Phi(t, 0)})_* [\theta(U), \theta(V)].$$

Whence

$$\nabla_U (\nabla_U V) = -\nabla_U (L_{\Phi(t, 0)})_* [\theta(U), \theta(V)] = (L_{\Phi(t, 0)})_* \frac{d}{dt} [\theta(U), \theta(V)].$$

Now, along  $\Phi(t, 0)$ , the vector  $\theta(U)$  is the constant vector  $X$ , and  $\theta(V) = v(t)$ . It follows that

$$\frac{d^2 v(t)}{dt^2} = (L_{\Phi(t, 0)}^{-1})_* \nabla_U (\nabla_U V) = - (L_{\Phi(t, 0)}^{-1})_* \nabla_U (L_{\Phi(t, 0)})_* [X, \theta(V)] = -\nabla_U [X, \theta(V)].$$

This leads to the following differential equation

$$\frac{d^2 v(t)}{dt^2} = - \left[ X, \frac{dv}{dt} \right]$$

with initial conditions  $v(0) = 0$  and  $\frac{dv}{dt}|_{t=0} = Y$ . A first integration leads to

$$\frac{dv}{dt} = e^{-t \operatorname{ad}(X)}(Y)$$

and a second to

$$v(t) = \left( \frac{1 - e^{-t \operatorname{ad}(X)}}{\operatorname{ad}(X)} \right) (Y).$$

So the result follows from the identity  $v(1) = (L_{\exp(X)}^{-1})_* (d_X \exp)(Y)$ .  $\square$

## A.2 Second expression of the differential of the exponential map

**Lemma A.3** For  $X$  in  $\operatorname{Sym}(n)$ , define

$$\begin{aligned} \tau_X : \operatorname{Sym}(n) &\longrightarrow \operatorname{Sym}(n) \\ Y &\longmapsto \tau_X(Y) := L_{\exp(-\frac{X}{2})} R_{\exp(-\frac{X}{2})} d_X \exp(Y). \end{aligned} \quad (36)$$

Then

$$\tau_X = \frac{\sinh(\operatorname{ad}(X/2))}{\operatorname{ad}(X/2)} = \sum_{n=0}^{+\infty} \frac{(\operatorname{ad}(X/2))^{2n}}{(2n+1)!}. \quad (37)$$

Moreover  $\tau_X$  is a linear isomorphism of  $\operatorname{Sym}(n)$ , that depends smoothly on  $X \in \operatorname{Sym}(n)$ , and whose inverse is

$$\begin{aligned} \tau_X^{-1} : \operatorname{Sym}(n) &\longrightarrow \operatorname{Sym}(n) \\ Y &\longmapsto \frac{\operatorname{ad}(X/2)}{\sinh(\operatorname{ad}(X/2))}(Y), \end{aligned} \quad (38)$$

In particular,

$$d_X \exp(Y) = R_{\exp(\frac{X}{2})} L_{\exp(\frac{X}{2})} \tau_X(Y) = \exp\left(\frac{X}{2}\right) \cdot \left( \frac{\sinh(\operatorname{ad}(X/2))}{\operatorname{ad}(X/2)}(Y) \right) \cdot \exp\left(\frac{X}{2}\right) \quad (39)$$

$$(40)$$

and

$$(d_X \exp)^{-1}(Z) = \tau_X^{-1} L_{\exp(-\frac{X}{2})} R_{\exp(-\frac{X}{2})}(Z) = \frac{\operatorname{ad}(X/2)}{\sinh(\operatorname{ad}(X/2))} \left( \exp\left(-\frac{X}{2}\right) \cdot Z \cdot \exp\left(-\frac{X}{2}\right) \right) \quad (41)$$

△ **Proof of Lemma A.3:**

A direct consequence of formula (33) in Proposition A.2 is that, for all  $Y$  in  $\text{Sym}(n)$ , we have

$$\begin{aligned}\tau_X(Y) &= L_{\exp(-\frac{X}{2})} R_{\exp(-\frac{X}{2})} d_X \exp(Y) = L_{\exp(-\frac{X}{2})} R_{\exp(-\frac{X}{2})} L_{\exp(X)} \left( \frac{1 - e^{-\text{ad}(X)}}{\text{ad}(X)} \right) (Y) \\ &= L_{\exp(\frac{X}{2})} R_{\exp(-\frac{X}{2})} \left( \frac{1 - e^{-\text{ad}(X)}}{\text{ad}(X)} \right) (Y) = \text{Ad}_{\exp(\frac{X}{2})} \left( \left( \frac{1 - e^{-\text{ad}(X)}}{\text{ad}(X)} \right) (Y) \right) \\ &= \exp \left( \text{ad} \left( \frac{X}{2} \right) \right) \left( \left( \frac{1 - e^{-\text{ad}(X)}}{\text{ad}(X)} \right) (Y) \right) = \left( \frac{e^{\text{ad}(X/2)} - e^{-\text{ad}(X/2)}}{\text{ad}(X)} \right) (Y) \\ &= \frac{\sinh(\text{ad}(X/2))}{\text{ad}(X/2)} (Y),\end{aligned}$$

where

$$\frac{\sinh(\text{ad}(X/2))}{\text{ad}(X/2)} = \frac{\exp(\text{ad}(X/2)) - \exp(-\text{ad}(X/2))}{\text{ad}(X)} = \sum_{n=0}^{+\infty} \frac{(\text{ad}(X)/2)^{2n}}{(2n+1)!}. \quad (42)$$

Every  $X$  in  $\text{Sym}(n)$  is diagonalisable in an orthonormal basis  $\{e_i, 1 \leq i \leq n\}$  with real eigenvalues  $\lambda_i$ ,  $1 \leq i \leq n$  (counted with multiplicity). The eigenvalues of  $\text{ad}(X)$  acting on  $M(n, \mathbb{R})$  are the real numbers  $(\lambda_i - \lambda_j)$ ,  $1 \leq i, j \leq n$ , the corresponding eigenvectors being  $E_{ij} := e_i \otimes e_j$  (this follows immediately from the fact that  $x$  is similar to a diagonal matrix with eigenvalues  $\lambda_i$ ). The eigenvalues of  $\tau_X$  are

$$\frac{\sinh\left(\frac{\lambda_i - \lambda_j}{2}\right)}{\frac{(\lambda_i - \lambda_j)}{2}},$$

$1 \leq i, j \leq n$ , with  $E_{ij}$  as eigenvectors (the operator should be interpreted as the identity map when  $i = j$ ). Since for any real number  $u \in \mathbb{R}$ ,  $\frac{\sinh u}{u} \geq 1$ ,  $\tau_X$  is injective on  $\text{Sym}(n)$ . Since  $\tau_X$  is linear, it is also surjective. Moreover  $\tau_X$  depends smoothly on  $X$  because  $\frac{\sinh u}{u}$  is smooth and  $\text{ad}(X)$  depends smoothly on  $X$ . The inverse of  $\tau_X$  is the linear map on  $\text{Sym}(n)$  with  $E_{ij}$  as eigenvectors and corresponding eigenvalues

$$\frac{\frac{(\lambda_i - \lambda_j)}{2}}{\sinh\left(\frac{\lambda_i - \lambda_j}{2}\right)},$$

$1 \leq i, j \leq n$ . Therefore it can be written as

$$\begin{aligned}\tau_X^{-1} : \text{Sym}(n) &\rightarrow \text{Sym}(n) \\ Y &\mapsto \frac{\text{ad}(X/2)}{\sinh \text{ad}(X/2)} (Y),\end{aligned}$$

△

## B Geodesics in locally symmetric homogeneous spaces

**Proposition B.1** *Let  $M = G/K$  be a locally symmetric homogeneous space of a Lie group  $G$  with unit element  $e$  and Lie algebra  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ , where  $\mathfrak{k}$  is the Lie algebra of  $K$  and  $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}$ . Then, for any  $G$ -invariant Riemannian metric on  $M$ , the geodesics starting at  $o = eK$  are given by the action of one-parameter subgroups of  $G$  generated by elements in  $\mathfrak{m}$ , i.e. are of the form*

$$\gamma(t) = (\exp t\mathfrak{a}) \cdot o \in M,$$

where  $\mathfrak{a}$  belongs to  $\mathfrak{m}$  (here the dot denotes the action of  $G$  on  $M$ ).

The proof of previous Proposition is based on the fact that for locally symmetric homogeneous manifolds, the homogeneous connection is the Levi-Cevita connection of any  $G$ -invariant Riemannian metric. The homogeneous connection  $\hat{\nabla}$  on the tangent space of  $M$  is defined as follows. For every element  $\mathfrak{a}$  in  $\mathfrak{m}_x$  and every vector field  $X$  on  $M$ , one has

$$\hat{\nabla}_{X^\mathfrak{a}(x)} X = (\mathcal{L}_{X^\mathfrak{a}} X)(x) = [X^\mathfrak{a}, X]_{\mathfrak{x}} \quad (43)$$

where  $\mathcal{L}$  denotes the Lie derivative and  $[\cdot, \cdot]_{\mathfrak{x}}$  the bracket of vector fields. For  $\mathfrak{a}$  in  $\mathfrak{m}_x$  and  $\mathfrak{b}$  in  $\mathfrak{g}$ , one has :

$$\hat{\nabla}_{X^\mathfrak{a}(x)} X^\mathfrak{b} = [X^\mathfrak{a}, X^\mathfrak{b}]_{\mathfrak{x}}(x) = -X^{[\mathfrak{a}, \mathfrak{b}]}(x).$$

The torsion of the connection  $\hat{\nabla}$  is given by

$$T^{\hat{\nabla}}(X^a, X^b) = \hat{\nabla}_{X^a} X^b - \hat{\nabla}_{X^b} X^a - [X^a, X^b]_{\mathfrak{X}} = -X^{[a,b]}.$$

It follows that for a locally symmetric homogeneous space, the homogeneous connection is torsion free since for  $\mathfrak{a}$  and  $\mathfrak{b}$  in  $T_x M = \mathfrak{m}_x$ ,  $[\mathfrak{a}, \mathfrak{b}]$  belongs to the isotropy  $\mathfrak{k}_x$  thus  $X^{[\mathfrak{a}, \mathfrak{b}]}$  vanishes. On the other hand, it follows from definition (43) that the covariant derivative of any tensor field  $\Phi$  along  $Y \in T_x M$  is the Lie derivative of  $\Phi$  along the vector field  $X^a$  where  $\mathfrak{a} \in \mathfrak{m}_x$  is such that  $Y = X^a(x)$ . Thus the homogeneous connection preserves every  $G$ -invariant Riemannian metric. Consequently  $\hat{\nabla}$  is the Levi-Civita connection of every  $G$ -invariant Riemannian metric on the locally symmetric space  $M = G/H$ .

□ **Proof of Proposition B.1:**

Every element  $\mathfrak{a}$  in  $\mathfrak{g}$  generates a vector field  $X^a$  on the homogeneous space  $M = G/K$ . For every  $x = g \cdot o$ ,  $g \in G$ , the Lie algebra  $\mathfrak{g}$  splits into  $\mathfrak{g} = \mathfrak{k}_x \oplus \mathfrak{m}_x$ , where  $\mathfrak{k}_x := \text{Ad}(g)(\mathfrak{k})$  is the Lie algebra of the isotropy group at  $x$  and where  $\mathfrak{m}_x := \text{Ad}(g)(\mathfrak{m})$  can be identified with the tangent space  $T_x M$  of  $M$  at  $x$  by the application  $\mathfrak{a} \mapsto X^a(x)$ . To see that for  $\mathfrak{a} \in \mathfrak{m}$ , the curve

$$\gamma(t) = (\exp t\mathfrak{a}) \cdot o, \quad \mathfrak{a} \in \mathfrak{m}.$$

is a geodesic, note that the equality  $\mathfrak{a} = \text{Ad}(\exp t\mathfrak{a})(\mathfrak{a})$  implies that  $\mathfrak{a}$  belongs to the space  $\mathfrak{m}_{\gamma(t)}$  for all  $t$ . Hence from  $\dot{\gamma}(t) = X^a(\gamma(t))$  it follows that  $\hat{\nabla}_{\dot{\gamma}(t)} \dot{\gamma}(t) = \mathcal{L}_{X^a} X^a(\gamma(t)) = 0$ . In other words  $\gamma$  is a geodesic of  $M$ . □

**Remark B.2 (Curvature)** For a locally symmetric homogeneous space, the curvature tensor  $R$  of  $\hat{\nabla}$  defined by

$$R_{X,Y}Z := \hat{\nabla}_X \hat{\nabla}_Y Z - \hat{\nabla}_Y \hat{\nabla}_X Z - \hat{\nabla}_{[X,Y]} Z.$$

has the following simple formula

$$R_{X,Y}Z = [[X, Y], Z]. \quad (44)$$

Indeed, for  $\mathfrak{a}, \mathfrak{b}, \mathfrak{c} \in \mathfrak{m}_x$ , one has

$$\begin{aligned} R_{X^a, X^b} X^c &= \hat{\nabla}_{X^a}([X^b, X^c]_{\mathfrak{X}}) - \hat{\nabla}_{X^b}([X^a, X^c]_{\mathfrak{X}}) - \hat{\nabla}_{[X^a, X^b]_{\mathfrak{X}}} X^c \\ &= [X^a, [X^b, X^c]_{\mathfrak{X}}]_{\mathfrak{X}} - [X^b, [X^a, X^c]_{\mathfrak{X}}]_{\mathfrak{X}} + \hat{\nabla}_{X^{[\mathfrak{a}, \mathfrak{b}]}} X^c \\ &= [[X^a, X^b]_{\mathfrak{X}}, X^c]_{\mathfrak{X}}, \end{aligned}$$

where in the last equality we have used  $X^{[\mathfrak{a}, \mathfrak{b}]} = 0$  for  $\mathfrak{a}, \mathfrak{b} \in \mathfrak{m}_x$ , since  $[\mathfrak{m}_x, \mathfrak{m}_x] \subset \mathfrak{k}_x$  and  $\mathfrak{k}_x$  acts trivially on  $T_x M$ .

## References

- [1] Andruchow, E.; Larotonda, G. *Nonpositively curved metric in the positive cone of a finite von Neumann algebra*, Journal of the London Mathematical Society, Volume 74, Issue 1, 2006, 205–218.
- [2] A. Arvanitoyeorgos, *An Introduction to Lie Groups and the Geometry of Homogeneous Spaces*, Student Math. Library no. 22, American Math. Society, Providence, R.I., (2003).
- [3] A.L. Besse, *Einstein Manifolds*, Ergebnisse der Mathematik und ihre Grenzgebiete, Springer-Verlag, (1986).
- [4] R. Bhatia, *Positive Definite Matrices*, Princeton University Press, Princeton Series in Applied Mathematics, 2007.
- [5] C. Conde, G. Larotonda, *Manifolds of semi-negative curvature*, Proc. Lond. Math. Soc. (3) 100 (3) (2010) 670–704.
- [6] Godement, R. *Introduction a la theorie des groupes de Lie*, first edition : Publications Mathematiques de l'Universite Paris VII (Numeros 11-12), (1982), second edition : Springer (2004).
- [7] Helgason, S., *Differential Geometry and Symmetric Spaces*, Academic Press, New York, (1962).
- [8] Larotonda, G. *Geodesic Convexity, Symmetric Spaces and Hilbert-Schmidt Operators*, PhD Thesis, Universidad Nacional de General Sarmiento, Buenos Aires, Argentina (2005).

- [9] M. Miglioli, *Decompositions and complexifications of some infinite-dimensional homogeneous spaces*, Journal of Functional Analysis 266 (2014) 6599–6618.
- [10] Mostow, G.D. *Some new decomposition theorems for semi-simple groups*, Mem. Amer. Math. Soc. 1955 (1955), n° 14, 31-54.
- [11] Mumford, D. ; Fogarty, J. ; Kirwan, F. *Geometric Invariant Theory*, (3rd. ed.), Springer, (1994).
- [12] Neeb, K.-H. *Highest weight representations and infinite-dimensional Kähler manifolds*, Recent advances in Lie theory (Vigo, 2000), 367-392, Res. Exp. Math., 25, Heldermann, Lemgo, (2002).
- [13] Neeb, K.-H. *A Cartan-Hadamard theorem for Banach-Finsler manifolds*, Geom. Dedicata 95 (2002), 115–156.
- [14] B. O'Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Academic Press, United Kingdom, (1983).
- [15] H. Porta, L. Recht. *Conditional expectations and operator decompositions*. Ann. Global Anal. Geom. 12 (1994), no. 4, 335–339.
- [16] Tumpach, A.B. *Variétés kählériennes et hyperkähleriennes de dimension infinie*, Ph.D Thesis, École Polytechnique, Palaiseau, France, (july 2005), electronic version available at <https://theses.hal.science/tel-00012012v1>
- [17] Tumpach, A.B. *Mostow's decomposition theorem for  $L^*$ -groups and applications to affine coadjoint orbits and stable manifolds*, Journal of Geometry and Physics, Volume 191, 2023, 104881, ISSN 0393-0440, <https://doi.org/10.1016/j.geomphys.2023.104881s>

Alice Barbara TUMPACH  
 Institut CNRS Pauli  
 Oskar-Morgenstern-Platz 1  
 1090 Vienna, Austria

*E-mail address* : [barbara.tumpach@math.cnrs.fr](mailto:barbara.tumpach@math.cnrs.fr)

Gabriel Larotonda  
 Departamento de Matemática, FCEyN, UBA  
 Avenida Cantilo S/N, Ciudad Universitaria  
 1428 Buenos Aires, Argentina

[glaroton@dm.uba.ar](mailto:glaroton@dm.uba.ar)