A simple model of magnetic universe without singularity associated with a quadratic equation of state

Pierre-Henri Chavanis¹

¹Laboratoire de Physique Théorique, Université de Toulouse, CNRS, UPS, France

A model of magnetic universe based on nonlinear electrodynamics has been introduced by Kruglov [Phys. Rev. D 92, 123523 (2015)]. This model describes an early inflation era followed by a radiation era. We show that this model is related to the model of universe based on a quadratic equation of state introduced in our previous paper [P.H. Chavanis, Universe 1, 357 (2015)]. This correspondence may provide a more fundamental justification of our equation of state. It may arise from quantum corrections to linear electrodynamics when the electromagnetic field becomes very high. We discuss two quantitatively different models of early universe. In Model I, the primordial density of the universe is identified with the Planck density. At t = 0, the universe had the characteristics of a Planck black hole ("planckion" particle). During the inflation, which takes place on a Planck timescale, the size of the universe evolves from the Planck length to a size comparable to the Compton wavelength of the neutrino. If we interpret the radius of the universe at the end of the inflation (neutrino's Compton wavelength) as a minimum length related to quantum gravity and use Zeldovich's first formula of the vacuum energy, we obtain the correct value of the cosmological constant. In Model II, the primordial density of the universe is identified with the electron density as a consequence of nonlinear electrodynamics. At t = 0, the universe had the characteristics of an electron. This can be viewed as a refinement of the "primeval atom" of Lemaître. During the inflation, which takes place on a gravitoelectronic timescale, the size of the universe evolves from the electron's classical radius to a size comparable to the size of a dark energy star of the stellar mass. If we interpret the radius of the universe at the beginning of the inflation (electron's classical radius) as a minimum length related to quantum gravity and use Zeldovich's second formula of the vacuum energy, we obtain the correct value of the cosmological constant. This provides an accurate form of Eddington's relation between the cosmological constant and the mass of the electron. We use these arguments to show that the present universe contains about 10^{80} protons (Eddington's number). We also introduce a nonlinear electromagnetic Lagrangian that describes simultaneously the early inflation, the radiation era, and the dark energy era. It may account for a form of "generalized radiation" that is present from the beginning to the end of the cosmic evolution. Baryonic and dark matter are treated as independent noninteracting species. The dark energy era is due to the electromagnetic vacuum energy (zero-point radiation). In this approach, both the early inflation and the late acceleration of the universe (dark energy) arise as a consequence of nonlinear electrodynamics. This leads to a simple model of magnetic universe without singularity (aioniotic universe).

I. INTRODUCTION

In a series of papers [1-10], we have developed a simple model of universe based on a generalized equation of state of the form

$$P/c^2 = \alpha \rho + k \rho^{1+1/n},\tag{1}$$

where P is the pressure and ρc^2 is the energy density. This is the sum of a linear equation of state and a polytropic equation of state. When n > 0 this equation of state describes the early universe. When k < 0 the energy density interpolates between a phase of primordial inflation (de Sitter) where the scale factor increases exponentially rapidly with time and an α -era where the scale factor increases algebraically. For $\alpha = 1/3$, this period corresponds to the radiation era. When n < 0 this equation of state describes the late universe. When k < 0 the energy density interpolates between an α -era where the scale factor increases algebraically with time and a phase of late acceleration (de Sitter) where the scale factor increases exponentially rapidly.¹

We have also introduced the quadratic equation of state [1-10]

$$P = -(\alpha + 1)\frac{\rho^2}{\rho_P}c^2 + \alpha\rho c^2 - (\alpha + 1)\rho_{\Lambda}c^2,$$
(2)

¹ The case k > 0 has been treated in [3, 4] and leads to singular or peculiar models of universe that we do not consider here. The case of a phantom evolution where the density of the universe increases as the universe expands has been treated in [5]. We do not consider this case here neither.

which provides a simple model of nonsingular universe presenting an early inflation, an α -era, and a late acceleration (dark energy era). The density starts at early times with the Planck density $\rho_P = c^5/G^2\hbar = 5.16 \times 10^{99} \,\mathrm{g m}^{-3}$, decreases during the α -era, and tends to the cosmological density $\rho_A = \Lambda c^2/8\pi G = 5.96 \times 10^{-24} \,\mathrm{g m}^{-3}$ at late times (see Fig. 16 in [6]).² Accordingly, the universe experiences a first period of exponential acceleration (early inflation), decelerates (when $\alpha > -1/3$), and finally experiences a second period of exponential acceleration (late inflation), the one that we observe at present. The two de Sitter eras are bridged by an α -era. Therefore, this equation of state unifies the inflation in the early universe and the dark energy in the late universe. We argued that the equation of state (2) with $\alpha = 1/3$ describes a form of "generalized radiation". In order to obtain the complete evolution of the universe we have to introduce, in addition to this generalized radiation, baryonic and dark matter, and possibly other components, treated as independent noninteracting species. The resulting cosmological model turns out to be equivalent to the Λ CDM model except that it replaces the big bang singularity by a non singular (de Sitter) inflation era with a "graceful" exit. A summary of our main results is given in the introduction of [10].

A weakness of our model, however, is that the equation of state (2) is introduced in a rather *ad hoc* manner, without a "microscopic" derivation. Recently, we came accross the very interesting papers of Kruglov [11–13] who introduced a similar model (valid in the early universe) from the viewpoint of "magnetic cosmology" based on nonlinear electrodynamics. Following previous authors [14–20] he argued that electromagnetic fields play an important role in cosmology and that the evolution of the early universe is fueled by a stochastic magnetic field due to plasma fluctuations. In this sense, electromagnetic fields are the source of gravitational fields. When electromagnetic fields are very strong during the early evolution of the universe one must use nonlinear electrodynamics. This may take into account quantum gravity corrections to linear electrodynamics.³ The use of nonlinear electrodynamics can remove the big bang singularity. Instead of prescribing an equation of state as we did, Kruglov [11-13] (see also [22-24]) introduced a nonlinear electromagnetic Lagrangian – a sort of generalized Born-Infeld [25, 26] Lagrangian – that produces a phase of early inflation followed by a radiation era.⁴ In the present paper, we show that his nonlinear electromagnetic Lagrangian leads precisely to the equation of state (1) introduced in [1-10]. We also consider the inverse problem and explain how one can construct more general electromagnetic Lagrangians by prescribing an equation of state such as Eq. (1). We stress, however, that some electromagnetic constraints reduce the possible type of equations of state and select the indices $n = \pm 1$ among the whole polytropic family, leading in a natural manner to Eq. (2).

We discuss two quantitatively different models regarding the early inflation.

In Model I, the density of the primordial universe is identified with the Planck density $\rho_P = 5.16 \times 10^{99} \text{ g m}^{-3}$ (or a fraction of it). During the inflation, which takes place on a Planck timescale $t_P = 5.39 \times 10^{-44}$ s, the size of the universe increases by 30 orders of magnitude. It evolves from the Planck length $l_P = 1.62 \times 10^{-35}$ m to a size equal to the Compton wavelength of the neutrino $r_{\nu} = 3.91 \times 10^{-5}$ m with mass $m_{\nu} = 5.04 \times 10^{-3} \text{ eV/c}^2$. This creates 10^{90} particles of the Planck mass $M_P = (\hbar c/G)^{1/2} = 2.18 \times 10^{-5} \text{ g} = 1.22 \times 10^{19} \text{ GeV/c}^2$ during the inflation. This implies that, after the radiation era, the mass of the universe is equal to $10^{62}M_P$. If we interpret the radius of the universe at the end of the inflation (neutrino's Compton wavelength) as a minimum length related to quantum gravity in the sense of Amelino-Camelia [27] and use Zeldovich's first formula of the vacuum energy [28, 29], we obtain the correct value of the cosmological constant [30]:

$$\Lambda = \frac{G\hbar}{r_{\nu}^4 c^3} = \frac{Gcm_{\nu}^4}{\hbar^3} = 1.11 \times 10^{-52} \,\mathrm{m}^{-2}.$$
(3)

In Model II, the density of the primordial universe is identified with the electron density $\rho_e = 4.07 \times 10^{16} \text{ g m}^{-3}$ as a consequence of nonlinear electrodynamics. During the inflation, which takes place on a gravitoelectronic timescale $t_e^* = 0.0192 \text{ s}$, the size of the universe increase by 30 orders of magnitude. It evolves from the electron's classical radius $r_e = 2.82 \times 10^{-15} \text{ m}$ to a size of the order of the radius $\tilde{R}_2 = 7.07 \times 10^{15} \text{ m}$ of a dark energy star of the stellar mass. This creates 10^{90} particles of the electron mass $m_e = 9.11 \times 10^{-28} \text{ g} = 0.511 \text{ MeV/c}^2$ during the inflation. This implies that, after the radiation era, the universe is made of 10^{83} electrons or 10^{80} protons (Eddington's number). If we interpret the radius of the universe at the begining of the inflation (electron's classical radius) as a minimum length related to quantum gravity in the sense of Karolyhazy [31] and use Zeldovich's second formula of the vacuum

 $^{^2}$ The empirical value of the cosmological constant is $\Lambda = 1.11 \times 10^{-52} \, {\rm m}^{-2}.$

³ In addition, quantum mechanics could be a "classical" process arising from the stochastic fluctuations of the electromagnetic field (zero-point radiation) as suggested by the theory of stochastic electrodynamics [21].

⁴ This Lagrangian is also introduced in an *ad hoc* manner. However, since it is connected to a physical mechanism – nonlinear electrodynamics – this gives the hope to derive this model (or more general ones) from "microscopic" considerations. This is, however, beyond the scope of the present paper.

energy [28, 29], we obtain the correct value of the cosmological constant [30]:

$$\Lambda = \frac{G^2 m_e^6}{\alpha^6 \hbar^4} = \frac{G^2 \hbar^2}{r_e^6 c^6} = 1.36 \times 10^{-52} \,\mathrm{m}^{-2},\tag{4}$$

where $\alpha = e^2/(\hbar c) \simeq 1/137$ is the fine-structure constant. This provides a justification of Eddington's accurate relation [30, 32]. We note that there is no free parameter in this relation.

At the primordial time t = 0 the universe had the characteristics of a Planck black hole ("planckion" particle) in Model I and the characteristics of an electron in Model II. However, it is unstable and explodes, leading to a phase of cosmological expansion. This can be viewed as a refinement of the "primeval atom" of Lemaître [33].

Finally, we introduce a nonlinear electromagnetic Lagrangian associated with the quadratic equation of state (2) that describes simultaneously the early inflation, the radiation era, and the dark energy era. The dark energy era is due to the electromagnetic vacuum energy (zero-point radiation). We suggest that nonlinear electrodynamics must also be used when the magnetic field is very low. Therefore, in our approach, both the early inflation and the late acceleration of the universe (dark energy) arise as a consequence of nonlinear electrodynamics. It may account for a form of "generalized radiation" that is present from the begining to the end of the cosmic evolution. This leads to a simple model of magnetic universe without singularity (aioniotic universe).

The paper is organized as follows. In Sec. II, we recall the basic equations of nonlinear electrodynamics and gravitation (general relativity). In Sec. III, we recall the basic equations of cosmology and consider the case where the evolution of the universe is due to a stochastic electromagnetic field. In Sec. IV, we recall the basic equations of nonlinear electrostatics. In Sec. V, we consider the case of a purely magnetic universe. In Sec. VI, we consider the nonlinear electrodynamics associated with a generalized polytropic equation of state and explain why the polytropic indices $n = \pm 1$ and the equation of state parameter $\alpha = 1/3$ are naturally selected by electrodynamics. In Sec. VII and VIII, we discuss the evolution of the early and late universe in the context of nonlinear electrodynamics. In Sec. IX, we introduce a nonlinear Lagrangian associated with a generalized form of radiation that accounts both for the early inflation and for the late acceleration of the universe (dark energy). In Sec. XI, we study the electric field produced by a pointlike charge (electron) in the context of nonlinear electrodynamics. In Sec. XII, we point out interesting heuristic connections between nonlinear electrodynamics, quantum gravity, vacuum energy and dark energy.

II. NONLINEAR ELECTRODYNAMICS AND GRAVITATION

In this section, we recall the basic equations of (nonlinear) electrodynamics and general relativity [34, 35].

A. The total action

The geometry of spacetime in general relativity is specified by the metric tensor $g_{\mu\nu}$ which gives the spacetime interval ds between two infinitesimally separated events, that is⁵

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu. \tag{5}$$

Here and below the Greek indices μ, ν etc run over the spacetime coordinates (ranging from 0 to 3) while the Latin indices i, j etc run only over the space coordinates (ranging from 1 to 3). We assume summation over repeated indices.

We first discuss the Maxwell equations in a general curved spacetime and then focus on Friedmann-Lemaître-Roberston-Walker (FLRW) models of cosmology. Electrodynamics in curved spacetime is most conveniently formulated by giving the action for the electromagnetic fields and their interaction with charged particles. For the sake of generality, we also include the contribution of a perfect fluid (or matter field). The total action of the system, which is the sum of the Einstein-Hilbert action of general relativity + the action of the perfect fluid + the action of the electromagnetic field can be written as

$$S = \frac{c^4}{16\pi G} \int R\sqrt{-g} \, d^4x + \int \mathcal{L}_m \sqrt{-g} \, d^4x + \int \mathcal{L}(\mathcal{F}) \sqrt{-g} \, d^4x - \int A_\mu J^\mu \sqrt{-g} \, d^4x, \tag{6}$$

⁵ The $g_{\mu\nu}$ may also be viewed as gravitational potentials since the effect of gravity is to modify the curvature of spacetime.

where R is the Ricci curvature scalar, $g = \det(g_{\mu\nu})$ is the determinant of the metric tensor, $A^{\mu} = (U/c, \mathbf{A})$ is the electromagnetic quadripotential, and J^{μ} is the quadricurrent density. We have assumed that the Lagrangian density $\mathcal{L}(\mathcal{F})$ of the electromagnetic field is an arbitrary function of the electromagnetic invariant

$$\mathcal{F} = \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu},\tag{7}$$

where $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ is the Faraday or electromagnetic field strength tensor (one can replace the covariant derivative D_{μ} by the partial derivative ∂_{μ} in this equation). In the above equation, μ_0 denotes the magnetic permeability in vacuum. The electric permittivity in vacuum is denoted by ϵ_0 . They are related by $\mu_0\epsilon_0 = 1/c^2$. Ordinary (linear) Maxwell electrodynamics corresponds to the Lagrangian

$$\mathcal{L}_{\text{Maxwell}} = -\mathcal{F}.$$
(8)

B. The Einstein equations

The Einstein equations of the gravitational field can be derived from the principle of least action $\delta S = \delta(S_g + S_M) = 0$, where S_g and S_M are the actions of the gravitational field and all sources of mass-energy (including the electromagnetic field), by performing variations with respect to the metric $g_{\mu\nu}$. This yields

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}T_{\mu\nu},$$
(9)

where $R_{\mu\nu}$ is the Ricci tensor measuring the curvature of spacetime, $R = R^{\mu}_{\mu} = g^{\mu\nu}R_{\mu\nu}$ is the scalar curvature of spacetime, and $T_{\mu\nu}$ is the energy-momentum tensor of the matter (or electromagnetic field) given by

$$T_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\partial(\sqrt{-g}\mathcal{L})}{\partial g^{\mu\nu}} = 2 \frac{\partial \mathcal{L}}{\partial g^{\mu\nu}} - g_{\mu\nu}\mathcal{L}.$$
 (10)

From this formulation, the energy-momentum tensor (10) is automatically symmetric. For a macroscopic body, the energy-momentum tensor can be written as

$$T_{\mu\nu} = (\rho c^2 + P) u_{\mu} u_{\nu} - P g_{\mu\nu}, \tag{11}$$

where ρc^2 is the energy density, P is the pressure and u^{μ} is the quadrivelocity such that $u_{\mu}u^{\mu} = 1$ (for a fluid at rest $T_{00} = \rho c^2$ and $T_{xx} = T_{yy} = T_{zz} = P$).

The conservation of energy and momentum is expressed by

$$D_{\nu}T^{\mu\nu} = 0, \tag{12}$$

where $D_{\nu} = \frac{1}{\sqrt{-g}} \partial_{\nu} \sqrt{-g}$ is the covariant derivative in a curved spacetime. According to this equation, the divergence of the left hand side of Eq. (9) must be zero. This is actually the case because of the contracted Bianchi identities. As a result, the conservation of energy and momentum is contained in the Einstein equations (9). The equations of the gravitational field contain the equations for the matter which produces this field. However, for a complete determination of the distribution and motion of the matter (or electromagnetic field) one must still supplement to the Einstein equations the equation of state of the matter (or electromagnetic field), i.e., an equation relating the pressure P to the energy density ρc^2 . This equation must be given along with the gravitational field equations.

C. The nonlinear Maxwell equations

The fundamental equations of electromagnetism are the Maxwell equations. In a curved spacetime the first pair of Maxwell equations reads (one can replace the covariant derivative D_{μ} by the partial derivative ∂_{μ} in this equation)

$$\partial_{\lambda}F_{\mu\nu} + \partial_{\nu}F_{\lambda\mu} + \partial_{\mu}F_{\nu\lambda} = 0. \tag{13}$$

The field equations determining the second pair of Maxwell equations can be obtained from the principe of least action $\delta S = 0$ by varying the electromagnetic potential A_{μ} . This yields

$$D_{\nu}\left[\mathcal{L}'(\mathcal{F})F^{\mu\nu}\right] = \mu_0 J^{\mu}.\tag{14}$$

Equations (13) and (14) form the (nonlinear) Maxwell equations. We see that only the second pair of Maxwell equations is affected by a possible nonlinearity of the Lagrangian.

The conservation of charge is expressed by the equation of continuity

$$D_{\mu}J^{\mu} = 0.$$
 (15)

According to this equation, the divergence of the left hand side of Eq. (14) must be zero. This is actually the case because of the antisymmetry of the Faraday tensor. The conservation of charge is therefore included in the Maxwell equations.

D. Electromagnetic energy-momentum tensor

The electromagnetic energy-momentum tensor associated with the Lagrangian $\mathcal{L}(\mathcal{F})$ is

$$T_{\mu\nu} = -\frac{1}{\mu_0} \mathcal{L}'(\mathcal{F}) F_{\mu\alpha} F^{\alpha}_{\ \nu} - \mathcal{L}(\mathcal{F}) g_{\mu\nu}.$$
(16)

We note that

$$T^{\mu}_{\mu} = 4\mathcal{F}\mathcal{L}'(\mathcal{F}) - 4\mathcal{L}(\mathcal{F}). \tag{17}$$

In linear electrodynamics ($\mathcal{L}_{\text{Maxwell}} = -\mathcal{F}$) the energy-momentum tensor of the electromagnetic field has the property that $T = T^{\mu}_{\mu} = 0$. With the identity $R = -(8\pi G/c^4)T$ obtained from the Einstein equations (9) it follows that in the presence of an electromagnetic field without any masses the scalar curvature of spacetime is zero: R = 0. On the other hand, the identity $T = \rho c^2 - 3P$ obtained from Eq. (11) implies that $P = \rho c^2/3$. These relations are no more true for a nonlinear electrodynamics.

III. BASIC EQUATIONS OF COSMOLOGY

In this section, we recapitulate the basic equations of cosmology [35] and consider the case where the universe is filled with a stochastic electromagnetic field.

A. Friedmann equations

If we consider an expanding homogeneous and isotropic cosmological spacetime (background) with a uniform curvature, the line element is given by the Friedmann-Lemaître-Roberston-Walker (FLRW) metric

$$ds^{2} = c^{2}dt^{2} - a(t)^{2} \left[\frac{dr^{2}}{1 - kr^{2}} + r^{2} \left(d\theta^{2} + \sin^{2}\theta \, d\phi^{2} \right) \right],$$
(18)

where a(t) represents the radius of curvature of the 3-dimensional space, or the scale factor. By an abuse of language, we shall sometimes call it the "radius of the universe". On the other hand, k determines the curvature of space. The universe is closed if k > 0, flat if k = 0, or open if k < 0.

If the universe is isotropic and homogeneous at all points in conformity with the line element (18), and contains a uniform perfect fluid of energy density $\rho(t)c^2$ and isotropic pressure P(t), the energy-momentum tensor $T_{\mu\nu}$ is given by Eq. (11) and we have

$$T_0^0 = \rho c^2, \qquad T_1^1 = T_2^2 = T_3^3 = -P.$$
 (19)

The Einstein equations

$$R^{\mu}_{\nu} - \frac{1}{2}g^{\mu}_{\nu}R - \Lambda g^{\mu}_{\nu} = -\frac{8\pi G}{c^2}T^{\mu}_{\nu}$$
⁽²⁰⁾

relate the geometrical structure of spacetime $(g_{\mu\nu})$ to the material content of the universe $(T_{\mu\nu})$ including the electromagnetic field. For the sake of generality, we have included the cosmological constant Λ in the Einstein equations (20). Given Eqs. (18) and (19), the Einstein equations reduce to

$$8\pi G\rho + \Lambda = 3\frac{\dot{a}^2 + kc^2}{a^2}, \qquad \frac{8\pi G}{c^2}P - \Lambda = -\frac{2a\ddot{a} + \dot{a}^2 + kc^2}{a^2},$$
(21)

where dots denote differentiation with respect to time. These are the well-known Friedmann [36, 37] cosmological equations. The Friedmann equations are usually written under the form

$$H^{2} = \frac{8\pi G}{3}\rho - \frac{kc^{2}}{a^{2}} + \frac{\Lambda}{3},$$
(22)

$$2\dot{H} + 3H^2 = -\frac{8\pi G}{c^2}P - \frac{kc^2}{a^2} + \Lambda,$$
(23)

where $H = \dot{a}/a$ is the Hubble parameter. From these equations, one can derive the acceleration equation

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3P}{c^2}\right) + \frac{\Lambda}{3}.$$
(24)

The deceleration parameter is defined by

$$q = -\frac{\ddot{a}a}{\dot{a}^2}.$$
(25)

The universe is decelerating when q > 0 and accelerating when q < 0.

B. Energy conservation equation

Combining Eqs. (22) and (23), we obtain the energy conservation equation

$$\frac{d\rho}{dt} + 3H\left(\rho + \frac{P}{c^2}\right) = 0. \tag{26}$$

This equation can be directly derived from the relation (12) which is included in the Einstein equations through the contracted Bianchi identities. It can be written under the form

$$d\rho + 3\frac{da}{a}(\rho + P/c^2) = 0.$$
 (27)

For a given barotropic equation of state $P = P(\rho)$, we can solve Eq. (27) to obtain

$$\ln a = -\frac{1}{3} \int \frac{d\rho}{\rho + P(\rho)/c^2}.$$
(28)

This equation determines the relation $\rho(a)$ between the energy density and the scale factor. We can then solve the Friedmann equation (22) with $\rho = \rho(a)$ to obtain the temporal evolution of the scale factor a(t).

The energy conservation equation (27) can be rewritten as

$$d(\rho c^2 a^3) = -Pd(a^3).$$
(29)

Introducing the volume $V \propto a^3$ and the energy $E = \rho c^2 V$, Eq. (29) becomes dE = -PdV. It can be interpreted as the first principle of thermodynamics for an adiabatic evolution of the universe dS = 0 [3, 38, 39].

The equation of state parameter is defined by

$$w = \frac{P}{\rho c^2}.$$
(30)

According to Eq. (26) the energy density decreases with the scale factor when w > -1 (null dominant energy condition) and increases with the scale factor when w < -1. When w = -1 the energy density is constant. The case where the energy density increases with the scale factor corresponds to a phantom universe [5, 40].

Remark: It is possible to develop a useful mechanical analogy to study the Friedmann equation (22). Indeed, it can be cast in the suggestive form

$$E = \frac{1}{2}\dot{a}^2 + V(a), \tag{31}$$

where

$$E = -\frac{1}{2}kc^2, \qquad V(a) = -\frac{4\pi G}{3}\rho(a)a^2 - \frac{\Lambda}{6}a^2.$$
(32)

Eq. (31) has the structure of the first integral of motion for a particle of unit mass in a potential V(a). In that case, E represents its conserved energy. The Friedmann equation (22) then has the solution

$$t = \int \frac{da}{\sqrt{2[E - V(a)]}} \tag{33}$$

determining a(t) in reversed form.

C. Flat universe

In this paper, we consider a flat universe (k = 0) in agreement with the inflation paradigm [41] and the observations of the cosmic microwave background (CMB) [42, 43]. On the other hand, we set the cosmological constant equal to zero $(\Lambda = 0)$ because dark energy will be taken into account in the nonlinear electrodynamics. The Friedmann equations (22) and (23) then reduce to the form

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho,\tag{34}$$

$$2\dot{H} + 3H^2 = -\frac{8\pi G}{c^2}P.$$
(35)

The acceleration equation (24) becomes

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3P}{c^2}\right). \tag{36}$$

Using Eqs. (34) and (36) we see that the deceleration parameter (25) is related to the equation of state parameter (30) by

$$q = \frac{1+3w}{2}.\tag{37}$$

The universe is decelerating when w > -1/3 (strong energy condition) and accelerating when w < -1/3; when w = -1/3 the scale factor increases linearly with time [3].

D. Tolman-Ehrenfest averaging procedure

Let us study some general properties of nonlinear electrodynamics in cosmology [11–20]. We assume that the universe is filled with electromagnetic radiation. The electromagnetic field that is of cosmological interest is the cosmic microwave background (CMB). It can be considered as a random field of short coherent radiation wavelength as compared to the cosmological horizon scales. Due to the isotropy of the spatial sections of the FLRW model, an average procedure is needed for compatibility reason if the electromagnetic field is to act as a source of gravity. Using the usual Tolman-Ehrenfest [44] procedure, we assume that the averaged electromagnetic field obeys the equations

$$\overline{E}_i = 0, \quad \overline{B}_i = 0, \quad \overline{E_i B_j} = 0,$$
(38)

$$\overline{E_i E_j} = \frac{1}{3} E^2 \delta_{ij}, \quad \overline{B_i B_j} = \frac{1}{3} B^2 \delta_{ij}, \tag{39}$$

where \overline{X} denotes an average over a volume that is large compared to the radiation wavelength but small compared to the curvature of spacetime. In the following, we omit the averaging bars for notational simplicity.

With these conditions, the average value of the energy-momentum tensor of the electromagnetic field associated with the Lagrangian density $\mathcal{L}(\mathcal{F})$ [see Eq. (16)] can be written in the form of the energy-momentum tensor for a perfect fluid [see Eq. (11)]. The energy density and the pressure of the radiation are given by [15, 17]

$$\rho c^2 = -\mathcal{L}(\mathcal{F}) - E^2 \mathcal{L}'(\mathcal{F}), \tag{40}$$

$$P = \mathcal{L}(\mathcal{F}) + \frac{E^2 - 2B^2}{3} \mathcal{L}'(\mathcal{F}), \tag{41}$$

where E^2 and B^2 are the averaged electric and magnetic fields squared, respectively. For convenience we have rescaled **E** by $1/(\epsilon_0)^{1/2}$ and **B** by $(\mu_0)^{1/2}$. The electromagnetic invariant takes the form

$$\mathcal{F} = \frac{1}{4} F_{\mu\nu} F^{\mu\nu} = \frac{B^2 - E^2}{2}.$$
(42)

For Maxwell's linear electrodynamics described by the Lagrangian (8) the foregoing equations reduce to

$$\rho c^2 = \frac{1}{2}(E^2 + B^2), \qquad P = \frac{1}{6}(E^2 + B^2),$$
(43)

returning the usual equation of state of radiation

$$P = \frac{1}{3}\rho c^2. \tag{44}$$

IV. ELECTROSTATICS

In electrostatics, in the absence of magnetic field $(\mathbf{B} = \mathbf{0})$ and if we can neglect the expansion of the universe (a = 1), the nonlinear Maxwell equations can be written as

$$\nabla \times \mathbf{E} = 0 \tag{45}$$

and

$$\nabla \cdot [\mathcal{L}'(\mathcal{F})\mathbf{E}] = -\rho_e,\tag{46}$$

where ρ_e is the charge density and

$$\mathcal{F} = -\frac{1}{2}E^2. \tag{47}$$

The electric field \mathbf{E} is expressed in terms of the scalar potential U by the relation

$$\mathbf{E} = -\nabla U. \tag{48}$$

Substituting Eq. (48) into Eq. (46), we get

$$\nabla \cdot [\mathcal{L}'(\mathcal{F})\nabla U] = \rho_e. \tag{49}$$

In particular, in vacuum ($\rho_e = 0$), the electric potential satisfies the equation

$$\nabla \cdot \left[\mathcal{L}'(\mathcal{F}) \nabla U \right] = 0. \tag{50}$$

For Maxwell's linear electrodynamics, Eqs. (49) and (50) reduce to the usual Poisson and Laplace equations $\Delta U = -\rho_e$ and $\Delta U = 0$, respectively.

Using the Gauss law (46), the electric field produced by a point charge $[\rho_e = \sqrt{4\pi}e\delta(\mathbf{r})]$ satisfies the equation

$$-\nabla \cdot [\mathcal{L}'(\mathcal{F})\mathbf{E}] = \sqrt{4\pi}e\delta(\mathbf{r}). \tag{51}$$

Its solution is

$$-\mathcal{L}'(\mathcal{F})E(r) = \frac{e}{\sqrt{4\pi}r^2}.$$
(52)

For Maxwell's linear electrodynamics, we recover the Coulomb law $E(r) = e/(\sqrt{4\pi r^2})$.

Finally, using Eq. (47), the electric energy density and the pressure [see Eqs. (40) and (41)] can be written as

$$\rho c^2 = -\mathcal{L}(\mathcal{F}) + 2\mathcal{F}\mathcal{L}'(\mathcal{F}), \tag{53}$$

$$P = \mathcal{L}(\mathcal{F}) - \frac{2}{3}\mathcal{F}\mathcal{L}'(\mathcal{F}).$$
(54)

For a given Lagrangian $\mathcal{L}(\mathcal{F})$, these equations define the equation of state $P(\rho)$ in parametric form with parameter $\mathcal{F} = -E^2/2 < 0$. For Maxwell's linear electrodynamics, we get $\rho c^2 = E^2/2$ and $P = E^2/6$ leading to $P = \rho c^2/3$.

V. MAGNETIC UNIVERSE

We assume that the universe is filled with a magnetic fluid. We suppose that the magnetic field of nonlinear electrodynamics is the main source of gravity. We consider the case E = 0 and $B \neq 0$ because only the magnetic field is important in cosmology. Indeed, the electric field is screened by the charged primordial plasma, while the magnetic field lines are frozen [45]. This leads to the concept of "magnetic universe" [17, 19]. In this model, the cosmic dynamics is fueled by the magnetic fluid alone.

In a purely magnetic universe (E = 0), the energy density and the pressure are given by [see Eqs. (40) and (41)]

$$\rho c^2 = -\mathcal{L}(\mathcal{F}),\tag{55}$$

$$P = \mathcal{L}(\mathcal{F}) - \frac{4}{3}\mathcal{F}\mathcal{L}'(\mathcal{F})$$
(56)

with

$$\mathcal{F} = \frac{B^2}{2}.\tag{57}$$

For a given Lagrangian $\mathcal{L}(\mathcal{F})$, these equations define the equation of state $P(\rho)$ in parametric form with parameter $\mathcal{F} = B^2/2 > 0$. Conversely, for a given equation of state $P(\rho)$, we can obtain the Lagrangian $\mathcal{L}(\mathcal{F})$ as follows. Combining Eqs. (55) and (56), we find that

$$P/c^2 = -\rho + \frac{4}{3} \frac{d\rho}{d\ln \mathcal{F}}.$$
(58)

For a given equation of state $P = P(\rho)$, Eq. (58) is a just first order differential equation. Its solution is

$$\ln \mathcal{F} = \frac{4}{3} \int \frac{d\rho}{P(\rho)/c^2 + \rho},\tag{59}$$

which determines $\mathcal{F} = \mathcal{F}(\rho)$. If this relation can be inverted, the electromagnetic Lagrangian is given by $\mathcal{L}(\mathcal{F}) = -\rho(\mathcal{F})c^2$.

It is also possible to derive the electromagnetic Lagrangian from the equation of state (or the converse) in a more direct manner by using a simple identity and exploiting the results obtained in cosmology. Combining Eq. (58) with the energy conservation equation (27) we get

$$\frac{d\mathcal{F}}{\mathcal{F}} = -4\frac{da}{a}.\tag{60}$$

Integrating this relation, we obtain the important identity (see, e.g. [46])

$$\mathcal{F} = \frac{\mathcal{F}_0}{a^4} \quad \text{i.e.} \quad B = \frac{B_0}{a^2},\tag{61}$$

where B_0 is the present value of the magnetic field and $\mathcal{F}_0 = B_0^2/2$. Therefore, there exists a simple general relation [Eq. (61)] between the electromagnetic invariant \mathcal{F} and the scale factor a. This identity is useful because, in cosmology, we are used to prescribing an equation of state $P = P(\rho)$ and, from the energy conservation equation (27), derive the relation $\rho(a)c^2$ between the energy density and the scale factor a. Many equations of state have been introduced and

studied in cosmology. To each of these models, if we know $\rho(a)c^2$ explicitly, then using Eqs. (55) and (61), we can immediately associate an electromagnetic Lagrangian by simply writing

$$\mathcal{L}(\mathcal{F}) = -\rho[(\mathcal{F}_0/\mathcal{F})^{1/4}]c^2. \tag{62}$$

Therefore, we can produce a great number of electromagnetic Lagrangians associated with cosmological models. Some examples will be given below. Conversely, for a given electromagnetic Lagrangian $\mathcal{L}(\mathcal{F})$, we can immediately write down the relation $\rho(a)c^2$ between the energy density and the scale factor a without any calculation. Indeed, using Eqs. (55) and (61), we have

$$\rho(a)c^2 = -\mathcal{L}(\mathcal{F}_0/a^4). \tag{63}$$

Similarly, using Eqs. (56) and (61), the evolution of the pressure with the scale factor is immediately given by

$$P(a) = \mathcal{L}(\mathcal{F}_0/a^4) - \frac{4}{3}(\mathcal{F}_0/a^4)\mathcal{L}'(\mathcal{F}_0/a^4).$$
(64)

Equations (63) and (64) define the equation of state $P(\rho)$ in parametric form with parameter a.

Remark: For Maxwell's linear electrodynamics, representing normal radiation, 6 we have

$$\mathcal{L} = -\mathcal{F}, \quad \rho_{\rm rad}c^2 = \mathcal{F}, \quad P_{\rm rad} = \frac{1}{3}\mathcal{F}, \quad P_{\rm rad} = \frac{1}{3}\rho_{\rm rad}c^2. \tag{65}$$

Since

$$\rho_{\rm rad}c^2 = \mathcal{F} = \frac{\mathcal{F}_0}{a^4},\tag{66}$$

we find that

$$\mathcal{F}_0 = \rho_{\mathrm{rad},0} c^2 = \Omega_{\mathrm{rad},0} \rho_0 c^2, \tag{67}$$

where $\rho_{\rm rad,0}c^2$ is the present energy density of normal radiation and $\Omega_{\rm rad,0} = \rho_{\rm rad,0}/\rho_0$ is the present proportion of normal radiation ($\rho_0 c^2$ is the present energy density of the universe). Therefore, \mathcal{F}_0 represents the present energy density of normal radiation and $B_0 = \sqrt{2\mathcal{F}_0}$ the present magnetic field. The dimensional magnetic field is

$$B_{\rm dim} = \sqrt{\mu_0} B. \tag{68}$$

The vacuum permeability has the value $\mu_0 = 1.25663706212 \times 10^{-9} \,\mathrm{m \, s^2 \, T^2 \, g^{-1}}$. From the observations we have $\Omega_{\mathrm{rad},0} = 9.23765 \times 10^{-5}$ and $\rho_0 = 8.62 \times 10^{-24} \mathrm{g \, m^{-3}}$. This gives $B_0 = 4.23 \times 10^{-10} \,\mathrm{T}$. More generally, we have the relations

$$\rho_{\rm rad}c^2 = \frac{B^2}{2} = \mathcal{F} = \frac{\rho_{\rm rad,0}c^2}{a^4}.$$
 (69)

These relations have been established during the normal radiation era but, because of the a^{-4} law from Eq. (61) which coincides with the a^{-4} law of normal radiation [3], they are actually valid for all times, even when $\rho \neq \rho_{\rm rad}$. In other words, $\mathcal{F} = B^2/2$ always represents the energy density of the normal radiation, even when the normal radiation is subdominant.

VI. NONLINEAR ELECTRODYNAMICS CORRESPONDING TO A GENERALIZED POLYTROPIC EQUATION OF STATE $P/c^2 = \alpha \rho + k \rho^{1+1/n}$ IN COSMOLOGY

A. Generalized polytropic equation of state and the corresponding Lagrangian

We consider an equation of state of the form

$$P/c^2 = \alpha \rho + k \rho^{1+1/n},\tag{70}$$

Ì

⁶ We call it normal radiation in order to distinguish it from the generalized radiation introduced in Sec. IX.

where ρc^2 is the energy density. This is the sum of a linear equation of state and a polytropic equation of state. This equation of state has been used in cosmology to describe the evolution of the early and late universe [3–5]. Solving the energy conservation equation (27) with the equation of state (70), we obtain⁷

$$\rho = \frac{\rho_*}{\left[\left(\frac{a}{a_*}\right)^{[3(1+\alpha)]/n} \mp 1\right]^n},\tag{71}$$

where a_* is an integration constant and $\rho_* = [(\alpha + 1)/|k|]^n$. The upper sign corresponds to k > 0 and the lower sign corresponds to k < 0. The equation of state (70) can be rewritten as

$$P/c^{2} = \alpha \rho \pm (\alpha + 1)\rho \left(\frac{\rho}{\rho_{*}}\right)^{1/n}.$$
(72)

Using $\rho c^2 = -\mathcal{L}$ [see Eq. (55)] and $\mathcal{F} = \mathcal{F}_0/a^4$ [see Eq. (61)] yielding $\mathcal{F}/\mathcal{F}_* = (a_*/a)^4$ with $\mathcal{F}_* = \mathcal{F}_0/a_*^4$, we find that the Lagrangian associated with the equation of state (72) in cosmology reads

$$\mathcal{L} = \frac{-\rho_* c^2}{\left[\left(\frac{\mathcal{F}_*}{\mathcal{F}}\right)^{[3(1+\alpha)]/(4n)} \mp 1\right]^n}.$$
(73)

B. Conditions of validity

It is important to note that the electromagnetic invariant \mathcal{F} has not a constant sign depending whether the electric or magnetic field dominates. If the magnetic field dominates, we have $\mathcal{F} = B^2/2 > 0$ while $\mathcal{F} = -E^2/2 < 0$ when the electric field dominates. The Lagrangian (73) has been derived in a cosmological context where the magnetic field dominates. In that case, \mathcal{F} and \mathcal{F}_* are positive. Now, the same Lagrangian should also describe the case where the electric field dominates. But in that case \mathcal{F} is negative while \mathcal{F}_* has the same value as before, being a positive constant. Therefore, $\mathcal{F}/\mathcal{F}_* > 0$ in the magnetic regime while $\mathcal{F}/\mathcal{F}_* < 0$ in the electric regime. If we want to describe the two situations with the same Lagrangian, and since the Lagrangian (73) involves a power-law $(\mathcal{F}_*/\mathcal{F})^{[3(1+\alpha)]/(4n)}$, it is necessary that $[3(1 + \alpha)]/4n$ be an integer.⁸ For simplicity, we shall require that

$$\frac{3(1+\alpha)}{4n} = \pm 1.$$
 (74)

On the other hand, in order to recover the Maxwell Lagrangian $\mathcal{L}_{\text{Maxwell}} = -\mathcal{F}$ when $(\mathcal{F}/\mathcal{F}_*)^{[3(1+\alpha)]/(4n)} \ll 1$ (correspondence principle), we need

$$\alpha = \frac{1}{3} \quad \text{and} \quad \mathcal{F}_* = \rho_* c^2. \tag{75}$$

The two conditions (74) and (75) imply

$$n = \pm 1. \tag{76}$$

Interestingly, we recover the two canonical models considered in [3-5]:

(i) The model $\alpha = 1/3$ and n = 1 corresponds to

$$P/c^2 = \frac{1}{3}\rho \pm \frac{4}{3}\frac{\rho^2}{\rho_*}, \qquad \mathcal{L} = \frac{-\mathcal{F}}{1 \mp \frac{\mathcal{F}}{\mathcal{F}_*}}.$$
 (77)

It describes the early universe. In practice we shall consider the lower sign (k < 0) corresponding to a nonsingular (inflationary) early universe [3]. Interestingly, the Lagrangian from Eq. (77) coincides with the one introduced by Kruglov [11, 12] from other arguments.

⁷ Following [3, 4] we consider the case where the energy density decreases with the scale factor. The case of a phantom evolution where the energy density increases with the scale factor is considered in [5].

⁸ Of course, we could put an absolute value $|\mathcal{F}_*/\mathcal{F}|^{[3(1+\alpha)]/(4n)}$ in Eq. (73) but we find this procedure a bit artificial and prefer selecting models where no absolute value arises.

(ii) The model $\alpha = 1/3$ and n = -1 corresponds to

$$P/c^2 = \frac{1}{3}\rho \pm \frac{4}{3}\rho_*, \qquad \mathcal{L} = -\mathcal{F} \pm \mathcal{F}_*.$$
 (78)

It describes the late universe. In practice we shall consider the lower sign (k < 0) corresponding to a nonsingular (inflationary) late universe [4].

Remark: More generally, in the foregoing discussion, 1/n could be any positive or negative integer. However, by considering a polytropic equation of state with an arbitrary index n, we have shown in [10] that the values $n = \pm 1$ are selected by an extremum principle: they turn out to minimize the mass of the real SF associated with the generalized polytropic equation of state (70). This may give them a special status, in addition to the argument of simplicity invoked above.

VII. NONLINEAR ELECTRODYNAMICS BASED ON THE LAGRANGIAN $\mathcal{L} = -\mathcal{F}/(1 + \mathcal{F}/\mathcal{F}_I)$: COSMOLOGY IN THE EARLY UNIVERSE

In the early universe, we consider a nonlinear electrodynamics based on the Lagrangian

$$\mathcal{L} = \frac{-\mathcal{F}}{1 + \frac{\mathcal{F}}{\mathcal{F}_I}},\tag{79}$$

where $\mathcal{F}_I = \rho_I c^2$. We shall discuss later the value of the characteristic density ρ_I appearing in Eq. (79). From Eqs. (55) and (56) we find that the energy density and the pressure are given by

$$\rho c^2 = \frac{\mathcal{F}}{1 + \frac{\mathcal{F}}{\rho_I c^2}},\tag{80}$$

$$P = \frac{\mathcal{F}\left(\frac{1}{3} - \frac{\mathcal{F}}{\rho_I c^2}\right)}{\left(1 + \frac{\mathcal{F}}{\rho_I c^2}\right)^2}.$$
(81)

Eliminating \mathcal{F} between these two expressions we obtain the quadratic equation of state

$$P = -\frac{4\rho^2 c^2}{3\rho_I} + \frac{1}{3}\rho c^2.$$
(82)

This equation of state has been studied in [3, 6] to model the early universe. Below, we recall its main properties.

A. Generalized polytropic equation of state

For the sake of generality, we consider a generalized polytropic equation of state of the form

$$P = -(\alpha + 1)\rho_I c^2 \left(\frac{\rho}{\rho_I}\right)^{1+1/n} + \alpha \rho c^2$$
(83)

with n > 0 and $-1 < \alpha \le 1$ [3, 6], even if we shall finally take $\alpha = 1/3$ and n = 1 for the reasons explained in Sec. VIB. For $\rho \ll \rho_I$, we obtain the linear equation of state $P \sim \alpha \rho c^2$. For $\rho \to \rho_I$, we get $P \to -\rho_I c^2$ corresponding to the equation of state of vacuum energy.

B. Evolution of the density, pressure and scale factor

Solving the energy conservation equation (27) with the equation of state (83) we find that the energy density evolves with the scale factor as

$$\rho = \frac{\rho_I}{\left[1 + (a/a_1)^{\frac{3(1+\alpha)}{n}}\right]^n},$$
(84)

where a_1 is a constant of integration. The pressure depends on the scale factor as

$$\frac{P}{\rho_I c^2} = \frac{\alpha (a/a_1)^{\frac{3(\alpha+1)}{n}} - 1}{\left[(a/a_1)^{\frac{3(\alpha+1)}{n}} + 1 \right]^{n+1}}.$$
(85)

For $a \ll a_1$, the energy density is approximately constant

$$\rho \simeq \rho_I, \tag{86}$$

and the pressure tends to $P \rightarrow -\rho_I c^2$ corresponding to vacuum energy. The Hubble parameter is constant, with value $H = (8\pi/3)^{1/2} t_I^{-1}$, where $t_I = 1/(G\rho_I)^{1/2}$ is a characteristic time associated with ρ_I . This leads to a phase of early inflation during which the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_I}$ (early de Sitter era). For $a \gg a_1$, the energy density decreases algebraically as

$$\rho \sim \frac{\rho_I a_1^{3(1+\alpha)}}{a^{3(1+\alpha)}}.$$
(87)

In that case, it behaves as an α -fluid with a linear equation of state $P \sim \alpha \rho c^2$. This leads to an α -era during which the scale factor increases algebraically rapidly with time as $t^{2/[3(1+\alpha)]}$ and the density decreases as t^{-2} . The expansion of the universe is decelerating if $\alpha > -1/3$ and accelerating if $\alpha < -1/3$. We can write the energy density of the α -fluid as

$$\rho_{\alpha} = \frac{\Omega_{\alpha,0}\rho_0}{a^{3(1+\alpha)}},\tag{88}$$

where $\rho_0 c^2$ is the present energy density of the universe and $\Omega_{\alpha,0}$ is the present fraction of the α -fluid (e.g. radiation when $\alpha = 1/3$). Comparing Eq. (87) with Eq. (88) and introducing the convenient notation $\Omega_{I,0} = \rho_I/\rho_0$, we get

$$a_1 = \left(\frac{\Omega_{\alpha,0}}{\Omega_{I,0}}\right)^{\frac{1}{3(1+\alpha)}}.$$
(89)

This relation determines the constant a_1 (we note that its value is independent of n). We can then rewrite Eq. (84) as

$$\frac{\rho}{\rho_0} = \frac{\Omega_{\alpha,0}}{\left[a^{\frac{3(\alpha+1)}{n}} + \left(\frac{\Omega_{\alpha,0}}{\Omega_{I,0}}\right)^{1/n}\right]^n}.$$
(90)

The equation of state (83) thus describes the smooth transition between a phase of inflation and an α -era in the early universe. The characteristic scale a_1 marks the transition between the vacuum energy (de Sitter) era and the α -era. At $a = a_1$, we have $\rho_1 = \rho_I/2^n$ and $P_1 = [(\alpha - 1)/2^{n+1}]\rho_I c^2$. The equation of state (83) is studied in detail in [3, 6]. The energy density decreases monotonically from ρ_I to 0. When $\alpha \ge 0$, the pressure increases from $P = -\rho_I c^2$ to a maximum positive value $P_e/(\rho_I c^2) = \alpha^{n+1} n^n / [(\alpha + 1)^n (n+1)^{n+1}]$ at $a_e/a_1 = [(\alpha + n + 1)/(n\alpha)]^{n/[3(\alpha+1)]}$ and $\rho_e/\rho_I = [\alpha n/(\alpha + 1)(n+1)]^n$ then decreases to zero. The pressure vanishes when $a_w/a_1 = (1/\alpha)^{n/[3(\alpha+1)]}$ and $\rho_w/\rho_I = [\alpha/(\alpha+1)]^n$. When $\alpha < 0$, the pressure monotonically increases from $P = -\rho_I c^2$ to zero. It is always negative. In this model, there is no initial singularity (no big bang). The universe exists from the infinite past and the scale factor tends to zero when $t \to -\infty$. The temporal evolution of the scale factor a(t) can be obtained analytically (in reversed form) in terms of hypergeometric functions [see Eq. (61) in [3]]. The evolution of the temperature is discussed in [3]. We refer to Figs. 2, 4, 8 and 10 of [3] for an illustration of the previous results.

Equation of state parameter, deceleration parameter and squared speed of sound

The equation of state parameter $w = P/(\rho c^2)$ is given by

$$w = -(\alpha + 1) \left(\frac{\rho}{\rho_I}\right)^{1/n} + \alpha.$$
(91)

Using Eq. (84) we get

$$w = \frac{\alpha (a/a_1)^{\frac{3(\alpha+1)}{n}} - 1}{(a/a_1)^{\frac{3(\alpha+1)}{n}} + 1}.$$
(92)

The pressure vanishes (w = 0) when $a_w/a_1 = (1/\alpha)^{n/[3(\alpha+1)]}$ and $\rho_w/\rho_I = [\alpha/(\alpha+1)]^n$ (when $\alpha \ge 0$). The deceleration parameter [see Eq. (37)] is given by

$$q = \frac{1+3\alpha}{2} - \frac{3}{2}(\alpha+1)\left(\frac{\rho}{\rho_I}\right)^{1/n}.$$
(93)

Using Eq. (84) we get

$$q = \frac{(1+3\alpha)(a/a_1)^{\frac{3(\alpha+1)}{n}} - 2}{2\left[(a/a_1)^{\frac{3(\alpha+1)}{n}} + 1\right]}.$$
(94)

The universe is accelerating (q < 0) when $a < a_c$ and decelerating (q > 0) when $a > a_c$ with $a_c/a_1 = [2/(1 + 3\alpha)]^{n/[3(\alpha+1)]}$ and $\rho_c/\rho_I = (1 + 3\alpha)^n/[3(\alpha+1)]^n$ (provided that $\alpha > -1/3$). Therefore, a_c marks the end of the early inflation.

The squared speed of sound $c_s^2 = dP/d\rho$ is given by

$$\frac{c_s^2}{c^2} = -(\alpha+1)\left(1+\frac{1}{n}\right)\left(\frac{\rho}{\rho_I}\right)^{1/n} + \alpha.$$
(95)

Using Eq. (84) we get

$$\frac{c_s^2}{c^2} = \frac{\alpha(a/a_1)^{\frac{3(\alpha+1)}{n}} - \frac{\alpha+n+1}{n}}{(a/a_1)^{\frac{3(\alpha+1)}{n}} + 1}.$$
(96)

When $\alpha \ge 0$, the speed of sound is imaginary $(c_s^2 < 0)$ when $a < a_e$ and real $(c_s^2 > 0)$ when $a > a_e$ with $a_e/a_1 = [(\alpha + n + 1)/(n\alpha)]^{n/[3(\alpha+1)]}$ and $\rho_e/\rho_I = [\alpha n/(\alpha+1)(n+1)]^n$ (this is the point where the pressure reaches its maximum value P_e so that $c_s^2 = dP/d\rho = 0$). When $\alpha < 0$, the speed of sound is always imaginary $(c_s^2 < 0)$. When it is real, the speed of sound is always less than the speed of light.

As the universe expands from a = 0 to $a = +\infty$, the equation of state parameter w increases from -1 to α , the deceleration parameter q increases from -1 to $(1 + 3\alpha)/2$ and the ratio $(c_s/c)^2$ increases from $-(\alpha + n + 1)/n$ to α (see Fig. 6 in [3]).

D. Application to the radiation

In this section, we specifically apply the preceding results to the case n = 1 and $\alpha = 1/3$ (radiation). This corresponds to the equation of state (82). For $\rho \ll \rho_I$, we recover the equation of state of radiation $P \sim \rho c^2/3$. For $\rho \to \rho_I$, we get $P \to -\rho_I c^2$ corresponding to vacuum energy. The energy density and the pressure evolve with the scale factor as

$$\rho = \frac{\rho_I}{1 + (a/a_1)^4}, \qquad \frac{P}{\rho_I c^2} = \frac{\frac{1}{3}(a/a_1)^4 - 1}{\left[(a/a_1)^4 + 1\right]^2}.$$
(97)

For $a \ll a_1$, the energy density is approximately constant with value $\rho \simeq \rho_I$, and the pressure tends to $P \rightarrow -\rho_I c^2$ (vacuum energy). This leads to a phase of early inflation during which the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_I}$ (early de Sitter era). For $a \gg a_1$, the energy density decreases algebraically as

$$\rho \sim \frac{\rho_I a_1^4}{a^4},\tag{98}$$

corresponding to the radiation with a linear equation of state $P \sim \rho c^2/3$. During the radiation era, the scale factor increases algebraically rapidly with time as $t^{1/2}$ and the density decreases as t^{-2} . The expansion of the universe is decelerating. We can write the energy density of radiation as

$$\rho_{\rm rad} = \frac{\Omega_{\rm rad,0}\rho_0}{a^4},\tag{99}$$

where $\rho_0 c^2$ is the present energy density of the universe and $\Omega_{rad,0}$ is the present fraction of radiation.

The equation of state (82) describes the smooth transition between a phase of inflation and the radiation era in the early universe. The transition between the vacuum energy (de Sitter) era and the radiation era takes place at

$$a_1 = \left(\frac{\Omega_{\text{rad},0}}{\Omega_{I,0}}\right)^{1/4}.$$
(100)

At $a = a_1$, we have $\rho_1 = \rho_I/2$ and $P_1 = -\rho_I c^2/6$. The equation of state (82) is studied in detail in [3, 6]. The energy density decreases monotonically from ρ_I to 0. The pressure increases from $P = -\rho_I c^2$ to a maximum positive value $P_e/(\rho_I c^2) = 1/48$ at $a_e/a_1 = 7^{1/4}$ and $\rho_e/\rho_I = 1/8$ then decreases to zero. The pressure vanishes when $a_w/a_1 = 3^{1/4}$ and $\rho_w/\rho_I = 1/4$. There is no initial singularity (no big bang). The universe exists from the infinite past and the scale factor tends to zero when $t \to -\infty$. The temporal evolution of the scale factor a(t) is given analytically (in reversed form) by [1–3, 6, 9, 10]:

$$\sqrt{(a/a_1)^4 + 1} - \ln\left(\frac{1 + \sqrt{(a/a_1)^4 + 1}}{(a/a_1)^2}\right) = 2\left(\frac{8\pi}{3}\right)^{1/2}\frac{t}{t_I} + C.$$
(101)

The constant is given by $C \simeq 1 - \ln 2 + 2 \ln \epsilon$, where $\epsilon \equiv a(t=0)/a_1 \ll 1$. The value of ϵ will be determined in Sec. VII E. The evolution of the temperature is discussed in [1–3, 9]. We refer to Figs. 11-17 of [3] for an illustration of these results.

The equation of state parameter $w = P/(\rho c^2)$ is given by

$$w = -\frac{4\rho}{3\rho_I} + \frac{1}{3} = \frac{\frac{1}{3}(a/a_1)^4 - 1}{(a/a_1)^4 + 1}.$$
(102)

The pressure vanishes when $a_w/a_1 = 3^{1/4}$ and $\rho_w/\rho_I = 1/4$.

The deceleration parameter is given by

$$q = 1 - \frac{2\rho}{\rho_I} = \frac{(a/a_1)^4 - 1}{(a/a_1)^4 + 1}.$$
(103)

The universe is accelerating when $a < a_c$ and decelerating when $a > a_c$ with $a_c/a_1 = 1$ and $\rho_c/\rho_I = 1/2$. Therefore, a_c marks the end of the early inflation. We note that $a_c = a_1$.

The squared speed of sound $c_s^2 = dP/d\rho$ is given by

$$\frac{c_s^2}{c^2} = -\frac{8\rho}{3\rho_I} + \frac{1}{3} = \frac{\frac{1}{3}(a/a_1)^4 - \frac{7}{3}}{(a/a_1)^4 + 1}.$$
(104)

The speed of sound is imaginary $(c_s^2 < 0)$ when $a < a_e$ and real $(c_s^2 > 0)$ when $a > a_e$ with $a_e/a_1 = 7^{1/4}$ and $\rho_e/\rho_I = 1/8$ (this is the point where the pressure reaches its maximum value P_e so that $c_s^2 = dP/d\rho = 0$). When it is real, the speed of sound is always less than the speed of light.

As the universe expands from a = 0 to $a = +\infty$, the equation of state parameter w increases from -1 to 1/3, the deceleration parameter q increases from -1 to 1 and the ratio $(c_s/c)^2$ increases from -7/3 to 1/3 (see Fig. 6 of [3]).

Remark: The foregoing results can be directly obtained from the Lagrangian (79). We have the relation

$$\mathcal{F} = \frac{B^2}{2} = \frac{\rho_I c^2}{(a/a_1)^4},\tag{105}$$

which can be obtained by comparing Eq. (97) with Eqs. (80) and (81) or by comparing Eq. (98) with Eq. (69) in the radiation era. The results can be therefore expressed in terms of $\mathcal{F}/\rho_I c^2$ or $B^2/2\rho_I c^2$ instead of a/a_1 by using Eq. (105). Applying Eq. (105) at the present epoch (a = 1) we find that

$$a_1 = \left(\frac{\mathcal{F}_0}{\rho_I c^2}\right)^{1/4},\tag{106}$$

which is the same as Eq. (100) according to Eq. (67).

E. e-folding number and duration of the inflation

The *e*-folding number before inflation ends is defined by

$$N = \ln\left(\frac{a_c}{a}\right),\tag{107}$$

where a_c is the scale factor at the end of the inflation. It corresponds to q = 0 or w = -1/3. In this section, we take n = 1 and $\alpha = 1/3$ (radiation) for the reasons given in Sec. VIB. In that case, using the results of Sec. VIIC, we have $\rho_c = \rho_I/2$ and $a_c = a_1$, where a_1 is given by Eq. (100). The scale factor a_c at the end of the inflation coincides with the scale factor a_1 marking the transition between the inflation era and the radiation era. To determine a_1 we must specify the value of ρ_I . At that point we can have two positions:

(i) Model I: If we follow the arguments given in our previous papers [3, 6], we would identify ρ_I with the Planck density $\rho_P = c^5/G^2\hbar = 5.16 \times 10^{99} \,\mathrm{g\,m^{-3}}$ (or a fraction of the Planck density). In other words, we assume that the maximum density of the universe (corresponding to the early inflation era) is the Planck density. In that case, taking $\rho_0 = 8.62 \times 10^{-24} \,\mathrm{g\,m^{-3}}$ for the present density of the universe, we get $\Omega_{P,0} = 5.98 \times 10^{122}$ leading to

$$a_1 = 1.98 \times 10^{-32}.\tag{108}$$

On the other hand, we have argued in [3, 6] (see also Appendix E) that

$$\epsilon \equiv \frac{a(t=0)}{a_1} = 1.71 \times 10^{-30}.$$
(109)

Therefore, $a(t=0) = 3.39 \times 10^{-62}$. From the above results, we obtain [10]

$$N_0 = \ln\left[\frac{a_c}{a(t=0)}\right] = \ln\left[\frac{a_1}{a(t=0)}\right] = \ln\left(\frac{1}{\epsilon}\right) = 68.5.$$
(110)

This value is consistent with the value $N \simeq 65$ deduced from the observations in order to solve the horizon and flatness problems [47]. On the other hand, the duration t_c of the inflation can be evaluated by substituting $a = a_c = a_1$ into Eq. (101) giving

$$\sqrt{2} - \ln(1 + \sqrt{2}) = 2\left(\frac{8\pi}{3}\right)^{1/2} \frac{t_c}{t_P} + C,$$
(111)

where $C = 1 - \ln(2) + 2 \ln \epsilon = -137$ and $t_P = 1/(G\rho_P)^{1/2} = (\hbar G/c^5)^{1/2} = 5.39 \times 10^{-44}$ s is the Planck time. This gives $t_c = 23.8 t_P = 1.28 \times 10^{-42}$ s. Therefore, in model I, the duration of the early inflation is of the order of the Planck time t_P .

(ii) Model II: If we follow the arguments based on nonlinear electrodynamics given by Kruglov [11] and in Sec. XI, we should identify ρ_I with the electron density $\rho_e = m_e/r_e^3 = 4.07 \times 10^{16} \text{ g m}^{-3}$ (a more precise value would be $\rho_* = 0.0307\rho_e = 1.25 \times 10^{15} \text{ g m}^{-3}$ but it is convenient for the discussion to take ρ_e). In other words, the maximum density of the universe (corresponding to the early inflation era) is connected to the electron density, not to the Planck density. In that case, we get $\Omega_{e,0} = 4.72 \times 10^{39}$ leading to

$$a_1 = 1.18 \times 10^{-11}.\tag{112}$$

On the other hand, Kruglov [13] takes $\epsilon \equiv a(t=0)/a_1 = 6.5 \times 10^{-28}$. This value is consistent with Eq. (109). As discussed in Appendix E, it is convenient to determine a(t=0) in model II such that ϵ has the value given by Eq. (109). This gives $a(t=0) = 2.10 \times 10^{-41}$. In this manner, N_0 has the same value $N_0 = 68.5$ in the two models and this simplifies the comparison. Repeating the arguments given after Eq. (110), we find that the duration t_c of the early inflation in model II is given by $t_c = 23.8 t_e^* = 0.457$ s, where $t_e^* = 1/\sqrt{G\rho_e} = 0.0192$ s is a timescale constructed with the density of the electron (instead of the Planck density).⁹ It corresponds to the dynamical time of a self-gravitating system of density ρ_e [48]. We shall call it the gravitoelectronic time. In model II, the duration of the early inflation is of the order of t_e^* .

⁹ With the value $\epsilon \equiv a(t=0)/a_1 = 6.5 \times 10^{-28}$, Kruglov [13] finds that the inflation lasts approximately 2 s with the reasonable *e*-folding number $N \simeq 63$.

In conclusion, in model II based on nonlinear electrodynamics where $\rho_I \sim \rho_e$, the inflation is much longer (~ 1 s) than in model I where $\rho_I \sim \rho_P$ giving $t_c \sim 1.28 \times 10^{-42}$ s. This is because $\rho_e \sim 10^{-80} \rho_P \ll \rho_P$ [30]. It would be interesting to know if cosmological constraints on the duration of the inflation (or on the density of the primordial universe) are able to discriminate between the two models.

Remark: We can have a more direct estimate of the duration of the inflation by applying the very accurate approximate formula $a \simeq a(t=0)e^{(8\pi/3)^{1/2}t/t_I}$ at $a = a_1 = a_c$ [3, 6], giving

$$t_c = \left(\frac{3}{8\pi}\right)^{1/2} \ln\left(\frac{1}{\epsilon}\right) t_I \simeq 23.7 t_I.$$
(113)

VIII. NONLINEAR ELECTRODYNAMICS BASED ON THE LAGRANGIAN $\mathcal{L} = -\mathcal{F} - \mathcal{F}_{\Lambda}$: COSMOLOGY IN THE LATE UNIVERSE

In the late universe, we consider a nonlinear electrodynamics based on the Lagrangian

$$\mathcal{L} = -\mathcal{F} - \mathcal{F}_{\Lambda},\tag{114}$$

where $\mathcal{F}_{\Lambda} = \rho_{\Lambda}c^2$. We suggest that DE corresponds to the zero-point radiation energy that manifests itself as a constant term in the electromagnetic Lagrangian. We identify this constant with the cosmological density $\rho_{\Lambda} = \Lambda c^2/8\pi G = 5.96 \times 10^{-24} \,\mathrm{g \, m^{-3}}$. In this model, the late acceleration of the universe is due to the electromagnetic energy of point zero. From Eqs. (55) and (56) we find that the energy density and the pressure are given by

$$\rho c^2 = \mathcal{F} + \rho_\Lambda c^2, \tag{115}$$

$$P = \frac{1}{3}\mathcal{F} - \rho_{\Lambda}c^2. \tag{116}$$

Eliminating \mathcal{F} between these two expressions we obtain the affine equation of state

$$P = \frac{1}{3}\rho c^2 - \frac{4}{3}\rho_{\Lambda}c^2.$$
 (117)

This equation of state has been studied in [4, 6] to model the late universe. Below, we recall its main properties.

A. Generalized polytropic equation of state

For the sake of generality, we consider a generalized polytropic equation of state of the form

$$P = \alpha \rho c^2 - (\alpha + 1)\rho_{\Lambda} c^2 \left(\frac{\rho}{\rho_I}\right)^{1 - 1/|n|}$$
(118)

with n < 0 and $-1 < \alpha \le 1$ [4, 6], even if we shall finally take $\alpha = 1/3$ and n = -1 for the reasons explained in Sec. VIB. For $\rho \gg \rho_{\Lambda}$, we obtain the linear equation of state $P \sim \alpha \rho c^2$. For $\rho \rightarrow \rho_{\Lambda}$, we get $P \rightarrow -\rho_{\Lambda} c^2$ corresponding to the equation of state of dark energy.

Remark: The equation of state (118) in the late universe can be viewed as the "symmetric" version of the equation of state (83) in the early universe. The symmetrical structure of the equation of state in the early (n > 0) and late (n < 0) universe is developed in [1, 3, 4].

B. Evolution of the density, pressure and scale factor

Solving the energy conservation equation (27) with the equation of state (118) we find that the energy density evolves with the scale factor as

$$\rho = \rho_{\Lambda} \left[1 + \frac{1}{(a/a_2)^{\frac{3(1+\alpha)}{|n|}}} \right]^{|n|}, \tag{119}$$

where a_2 is a constant of integration. The pressure depends on the scale factor as

$$\frac{P}{\rho_{\Lambda}c^2} = \left[\frac{\alpha}{(a/a_2)^{\frac{3(\alpha+1)}{|n|}}} - 1\right] \left[\frac{1}{(a/a_2)^{\frac{3(\alpha+1)}{|n|}}} + 1\right]^{|n|-1}.$$
(120)

For $a \gg a_2$, the energy density is approximately constant

$$\rho \simeq \rho_{\Lambda},$$
(121)

and the pressure tends to $P \to -\rho_{\Lambda}c^2$ corresponding to DE. Here, ρ_{Λ} is unambiguously associated with the cosmological density ρ_{Λ} . The Hubble parameter is constant, with value $H = (8\pi/3)^{1/2}t_{\Lambda}^{-1}$, where $t_{\Lambda} = 1/(G\rho_{\Lambda})^{1/2} = (8\pi/\Lambda c^2)^{1/2} = 1.59 \times 10^{18}$ s is a characteristic time (cosmological time) associated with ρ_{Λ} . This leads to a phase of late accelerating expansion (or late inflation) during which the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_{\Lambda}}$ (late de Sitter era).

For $a \ll a_2$, the energy density decreases algebraically as

$$\rho \sim \frac{\rho_\Lambda a_2^{3(1+\alpha)}}{a^{3(1+\alpha)}}.\tag{122}$$

In that case, it behaves as an α -fluid with a linear equation of state $P \sim \alpha \rho c^2$. This leads to an α -era during which the scale factor increases algebraically rapidly with time as $t^{2/[3(1+\alpha)]}$ and the density decreases as t^{-2} . The expansion of the universe is decelerating if $\alpha > -1/3$ and accelerating if $\alpha < -1/3$. We can write the energy density of the α -fluid as

$$\rho_{\alpha} = \frac{\Omega_{\alpha,0}\rho_0}{a^{3(1+\alpha)}},\tag{123}$$

where $\rho_0 c^2$ is the present energy density of the universe and $\Omega_{\alpha,0}$ is the present fraction of the α -fluid (e.g. radiation when $\alpha = 1/3$). Comparing Eq. (122) with Eq. (123) and introducing the present fraction of DE $\Omega_{\Lambda,0} = \rho_{\Lambda}/\rho_0$, we get

$$a_2 = \left(\frac{\Omega_{\alpha,0}}{\Omega_{\Lambda,0}}\right)^{\frac{1}{3(1+\alpha)}}.$$
(124)

This relation determines the constant a_2 (we note that its value is independent of n). We have the relation $a_2 = a_1 \left(\rho_I / \rho_\Lambda\right)^{\frac{1}{3(1+\alpha)}}$. We can then rewrite Eq. (119) as

$$\frac{\rho}{\rho_0} = \Omega_{\alpha,0} \left[\frac{1}{a^{\frac{3(\alpha+1)}{|n|}}} + \left(\frac{\Omega_{\Lambda,0}}{\Omega_{\alpha,0}} \right)^{1/|n|} \right]^{|n|}.$$
(125)

The equation of state (118) thus describes the smooth transition between an α -era and a phase of accelerating expansion (DE or inflation) in the late universe. The characteristic scale a_2 marks the transition between the α -era and the dark energy (de Sitter) era. At $a = a_2$, we have $\rho_2 = 2^{|n|} \rho_{\Lambda}$ and $P_2 = (\alpha - 1)2^{|n|-1} \rho_{\Lambda} c^2$. The equation of state (118) is studied in detail in [4, 6]. The energy density decreases monotonically from $+\infty$ to ρ_{Λ} . The evolution of the pressure depends on the sign of α . When $\alpha > 0$, the pressure decreases from $+\infty$ to $P = -\rho_{\Lambda}c^2$. It vanishes when $a'_w/a_2 = \alpha^{|n|/[3(\alpha+1)]}$ and $\rho'_w/\rho_{\Lambda} = [(\alpha+1)/\alpha]^{|n|}$. When $\alpha < 0$, the pressure increases from $-\infty$ to $P = -\rho_{\Lambda}c^2$. When $\alpha = 0$, the evolution of the pressure depends on the value of |n|. When |n| > 1 the pressure increases from $-\infty$ to $P = -\rho_{\Lambda}c^2$. The pressure is constant $P = -\rho_{\Lambda}c^2$, and when |n| < 1 the pressure decreases from $-\infty$ to $P = -\rho_{\Lambda}c^2$. The pressure may present an extremum $P'_e/(\rho_{\Lambda}c^2) = \alpha^{-|n|+1}(-|n|)^{-|n|}/[(\alpha+1)^{-|n|}(-|n|+1)^{-|n|+1}]$ at the point $a'_e/a_2 = [-(\alpha - |n| + 1)/(|n|\alpha)]^{-|n|/[3(\alpha+1)]}$ and $\rho'_e/\rho_{\Lambda} = [-\alpha|n|/((\alpha+1)(-|n|+1))]^{-|n|}$. Its conditions of existence are detailed in [4]. We refer to Figs. 1 and 2 of [4] for an illustration of the previous results.

Remark: The temporal evolution of the scale factor a(t) is given in [4] assuming that the late universe is described by a single fluid with the equation of state (118). However, in general, there are other fluids that also contribute to the density of the universe and therefore change the temporal evolution of the scale factor (see below). This is why we have not given its expression here.

C. Equation of state parameter and squared speed of sound

The equation of state parameter $w = P/(\rho c^2)$ is given by

$$w = \alpha - (\alpha + 1) \left(\frac{\rho_{\Lambda}}{\rho}\right)^{1/|n|}.$$
(126)

Using Eq. (119) we get

$$w = \frac{\alpha (a_2/a)^{\frac{3(\alpha+1)}{|n|}} - 1}{(a_2/a)^{\frac{3(\alpha+1)}{|n|}} + 1}.$$
(127)

The pressure vanishes (w = 0) when $a'_w/a_2 = \alpha^{|n|/[3(\alpha+1)]}$ and $\rho'_w/\rho_{\Lambda} = [(\alpha+1)/\alpha]^{|n|}$ (assuming $\alpha > 0$). The squared speed of sound $c_s^2 = dP/d\rho$ is given by

$$\frac{c_s^2}{c^2} = -(\alpha+1)\left(1-\frac{1}{|n|}\right)\left(\frac{\rho_\Lambda}{\rho}\right)^{1/|n|} + \alpha.$$
(128)

Using Eq. (119) we get

$$\frac{c_s^2}{c^2} = \frac{\alpha (a_2/a)^{\frac{3(\alpha+1)}{|n|}} + \frac{\alpha - |n| + 1}{|n|}}{(a_2/a)^{\frac{3(\alpha+1)}{|n|}} + 1}.$$
(129)

The speed of sound may vanish at the point $a'_e/a_2 = [-(\alpha - |n| + 1)/(|n|\alpha)]^{-|n|/[3(\alpha+1)]}$ and $\rho'_e/\rho_{\Lambda} = [-\alpha|n|/((\alpha + 1)(-|n|+1))]^{-|n|}$. This is the point where the pressure is extremum. The speed of sound may equal the speed of light at the point $a'_s/a_2 = [(\alpha - 2|n| + 1)/(|n|(1 - \alpha))]^{-|n|/[3(\alpha+1)]}$ and $\rho'_s/\rho_{\Lambda} = [(1 - \alpha)|n|/((\alpha + 1)(-|n| + 1))]^{-|n|}$. The conditions of existence of these two points are detailed in [4].

As the universe expands from a = 0 to $a = +\infty$, the equation of state parameter w evolves from α to -1 and the squared speed of sound c_s^2/c^2 evolves from α to $(\alpha - |n| + 1)/|n|$ (see Fig. 5 of [4] for an illustration).

Remark: For the reason explained previously we have not given the deceleration parameter because its value may be affected by other fluids.

D. Application to the radiation

In this section, we specifically apply the preceding results to the case n = -1 and $\alpha = 1/3$ (radiation). This corresponds to the equation of state (117). It can be viewed as the "symmetric" version of the equation of state (82) in the early universe [1, 3, 4]. For $\rho \gg \rho_{\Lambda}$, we recover the linear equation of state of radiation $P \sim \rho c^2/3$. For $\rho \to \rho_{\Lambda}$, we get $P \to -\rho_{\Lambda}c^2$ corresponding to dark energy. The energy density and the pressure evolve with the scale factor as

$$\rho = \rho_{\Lambda} \left[1 + \frac{1}{(a/a_2)^4} \right], \qquad \frac{P}{\rho_{\Lambda} c^2} = \frac{1}{3(a/a_2)^4} - 1.$$
(130)

For $a \gg a_2$, the energy density is approximately constant with value $\rho \simeq \rho_{\Lambda}$, and the pressure tends to $P \rightarrow -\rho_{\Lambda}c^2$ corresponding to DE. This leads to a phase of late accelerating expansion (or late inflation) during which the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_{\Lambda}}$ (late de Sitter era). For $a \ll a_2$, the energy density decreases algebraically as

$$\rho \sim \frac{\rho_{\Lambda} a_2^4}{a^4},\tag{131}$$

corresponding to the radiation with a linear equation of state $P \sim \rho c^2/3$. During the radiation era, the scale factor increases algebraically rapidly with time as $t^{1/2}$ and the density decreases as t^{-2} . The expansion of the universe is decelerating. We can write the energy density of the radiation as

$$\rho_{\rm rad} = \frac{\Omega_{\rm rad,0}\rho_0}{a^4},\tag{132}$$

20

where $\rho_0 c^2$ is the present energy density of the universe and $\Omega_{rad,0}$ is the present fraction of radiation.

The equation of state (117) describes the smooth transition between the radiation era and a phase of accelerating expansion (DE or inflation) in the late universe. The transition between the radiation era and the the dark energy (de Sitter) era takes place at

$$a_2 = \left(\frac{\Omega_{\rm rad,0}}{\Omega_{\Lambda,0}}\right)^{1/4}.$$
(133)

At $a = a_2$, we have $\rho_2 = 2\rho_{\Lambda}$ and $P_2 = -(2/3)\rho_{\Lambda}c^2$. The equation of state (118) is studied in detail in [4, 6]. The energy density decreases monotonically from $+\infty$ to ρ_{Λ} . The pressure decreases monotonically from $+\infty$ to $P = -\rho_{\Lambda}c^2$. It vanishes when $a'_w/a_2 = 1/81$ and $\rho'_w/\rho_{\Lambda} = 4$.

The equation of state parameter $w = P/(\rho c^2)$ is given by

$$w = \frac{1}{3} - \frac{4\rho_{\Lambda}}{3\rho} = \frac{\frac{1}{3}(a_2/a)^4 - 1}{(a_2/a)^4 + 1}.$$
(134)

The pressure vanishes when $a'_w/a_2 = 1/81$ and $\rho'_w/\rho_{\Lambda} = 4$. As the universe expands from a = 0 to $a = +\infty$, the equation of state parameter w decreases from 1/3 to -1.

The squared speed of sound $c_s^2 = dP/d\rho$ is given by

$$\frac{c_s^2}{c^2} = \frac{1}{3}.$$
(135)

The speed of sound is constant and equal to $c_s/c = 1/\sqrt{3}$. The speed of sound is less than the speed of light.

We will see in Sec. IX that ρ given by Eq. (130) represents the density of generalized radiation in the late universe. For $a \ll a_2$ it corresponds to the ordinary radiation. For $a_{eq} \ll a \ll a_2$ (where a_{eq} is the value of the scale factor at radiation-matter equality) it is subdominant with respect to baryonic and dark matter viewed as different species. This is why we have not given the temporal evolution of the scale factor a(t) nor the deceleration parameter q(t) in this period because we have to take into account the contribution of matter. This is done in Sec. X where we present the complete model. By contrast, for $a_1 \ll a \ll a_{eq}$ we are in the radiation era and for $a \gg a'_2$ we are in the dark energy era where the asymptotic results for a(t) given above are valid. In these limits, the deceleration parameter is given by Eq. (37).

Remark: The foregoing results can be directly obtained from the Lagrangian (114). We have the relation

$$\mathcal{F} = \frac{B^2}{2} = \frac{\rho_\Lambda c^2}{(a/a_2)^4},\tag{136}$$

which can be obtained by comparing Eq. (130) with Eqs. (115) and (116) or by comparing Eq. (131) with Eq. (69) in the radiation era. The results can be therefore expressed in terms of $\mathcal{F}/\rho_{\Lambda}c^2$ or $B^2/2\rho_{\Lambda}c^2$ instead of a/a_2 by using Eq. (136). Applying Eq. (136) at the present epoch (a = 1) we find that

$$a_2 = \left(\frac{\mathcal{F}_0}{\rho_\Lambda c^2}\right)^{1/4},\tag{137}$$

which is the same as Eq. (133) according to Eq. (67).

IX. THE LAGRANGIAN OF THE GENERALIZED RADIATION

In Sec. VII we have considered a nonlinear electrodynamics based on a Lagrangian of the form

$$\mathcal{L} = \frac{-\mathcal{F}}{1 + \frac{\mathcal{F}}{\rho_I c^2}} \tag{138}$$

with $\rho_I = \rho_P$ (model I) or $\rho_I = \rho_e$ (model II). We have shown that this Lagrangian could describe the evolution of the early universe. In Sec. VIII we have considered a nonlinear electrodynamics based on a Lagrangian of the form

$$\mathcal{L} = -\mathcal{F} - \rho_{\Lambda} c^2. \tag{139}$$

We have shown that this Lagrangian could describe the evolution of the late universe. We now want to connect these two Lagrangians in order to describe the complete evolution of the universe (see Sec. X). We see that there is a common period corresponding to $\mathcal{F} \ll \mathcal{F}_I = \rho_I c^2$ in the early universe and $\mathcal{F} \gg \mathcal{F}_{\Lambda} = \rho_{\Lambda} c^2$ in the late universe. In this common period, corresponding to the normal radiation era, the Lagrangians (138) and (139) reduce to Maxwell's linear electrodynamics characterized by the Lagrangian

$$\mathcal{L} = -\mathcal{F}.\tag{140}$$

The Lagrangian of the nonlinear electrodynamics valid during the whole evolution of the universe is therefore

$$\mathcal{L} = \frac{-\mathcal{F}}{1 + \frac{\mathcal{F}}{\rho_I c^2}} - \rho_\Lambda c^2. \tag{141}$$

In the early universe, it reduces to Eq. (138) and in the late universe it reduces to Eq. (139). In the intermediate period (radiation era), it returns the ordinary Maxwell electrodynamics (140).

From Eqs. (55) and (56) we find that the energy density and the pressure are given by

$$\rho c^2 = \frac{\mathcal{F}}{1 + \frac{\mathcal{F}}{\rho_I c^2}} + \rho_\Lambda c^2, \tag{142}$$

$$P = \frac{\mathcal{F}\left(\frac{1}{3} - \frac{\mathcal{F}}{\rho_I c^2}\right)}{\left(1 + \frac{\mathcal{F}}{\rho_I c^2}\right)^2} - \rho_\Lambda c^2.$$
(143)

Eliminating \mathcal{F} between these two expressions we obtain in excellent approximation the quadratic equation of state¹⁰

$$P = -\frac{4\rho^2}{3\rho_I}c^2 + \frac{1}{3}\rho c^2 - \frac{4}{3}\rho_\Lambda c^2.$$
 (144)

This equation of state (see Fig. 1) was introduced and studied in [1–10]. In the early universe, we recover the quadratic equation of state (82) associated with the Lagrangian (138) and in the late universe we recover the affine equation of state (117) associated with the Lagrangian (139). In the intermediate period, we recover the linear equation of state $P = \frac{1}{3}\rho c^2$ of the ordinary radiation associated with Maxwell's electrodynamics. Following the interpretation given in our previous works [1–10], the equation of state (144) describes a form of *generalized radiation* which is responsible for the early inflation, the normal radiation era and the present and late acceleration (dark energy) of the universe. In the present paper, we have connected this "generalized radiation" to a form of nonlinear electrodynamics. Solving the energy conservation equation (27) with the equation of state (144) we obtain in excellent approximation (see footnote 10) the following relation between the density of the generalized radiation and the scale factor [1–10]:

$$\rho_{\rm Rad} = \frac{\rho_I}{1 + (a/a_1)^4} + \rho_{\Lambda}.$$
(145)

This is also the exact relation obtained from the Lagrangian (141) with Eq. (105) and (142). In the early universe, we recover the results of Sec. VII D. In the late universe we recover the results of Sec. VIII D. We stress that $\rho_{\text{Rad}} \neq \rho_{\text{rad}}$ in general. It is only in the intermediate period corresponding to Maxwell's linear electrodynamics that the energy density of the generalized radiation ρ_{Rad} reduces to the energy density of the normal radiation ρ_{rad} .

Remark: It is possible to find the exact relation $\rho_{\text{Rad}}(a)$ determined by the equation of state (144). It is given in Appendix C of [6]. Then, using Eqs. (55) and (105) we can obtain the exact Lagrangian $\mathcal{L}(\mathcal{F})$ corresponding to the equation of state (144). Conversely, eliminating \mathcal{F} between Eqs. (142) and (143) it is possible to obtain the exact equation of state corresponding to the Lagrangian (141). This equation of state exactly leads to the relation $\rho_{\text{Rad}}(a)$ from Eq. (145). However, the approximate expressions given above are so accurate (see footnote 10) that this refinement is not necessary.

¹⁰ To obtain this equation we have used the fact that $\rho_P/\rho_\Lambda \sim 10^{120} \gg 1$ and $\rho_e/\rho_\Lambda \sim 10^{40} \gg 1$. Since these dimensionless numbers are huge [30], the approximation is quasi perfect.



22



FIG. 1: Equation of state of the generalized radiation [see Eq. (144)] associated with the Lagrangian (141). The pressure P and the energy density ρc^2 are normalized by $\rho_I c^2$.

X. COMPLETE EVOLUTION OF THE UNIVERSE

The Lagrangian (141) describes the generalized radiation. It accounts for the early inflation, the radiation era and the late acceleration of the universe. Baryonic matter and dark matter must be treated independently, as additional components. As a result, the total Lagrangian describing the mass-energy content of the universe is

$$\mathcal{L} = \mathcal{L}_{\text{Rad}} + \mathcal{L}_{\text{matter}},\tag{146}$$

where \mathcal{L}_{Rad} is the Lagrangian of the generalized radiation¹¹ and $\mathcal{L}_{\text{matter}} = \mathcal{L}_{\text{b}} + \mathcal{L}_{\text{dm}}$ is the Lagrangian of the baryonic and dark matter. The energy density of the generalized radiation is given by Eq. (145) with $a_1 = (\rho_{\text{rad},0}/\rho_I)^{1/4}$ [see Eq. (100)]. If we assume for simplicity that baryonic matter and dark matter are pressureless $(P_{\text{b}} = P_{\text{dm}} = 0)$,¹² we find that their densities evolve with the scale factor as

$$\rho_{\rm b} = \frac{\rho_{\rm b,0}}{a^3}, \qquad \rho_{\rm dm} = \frac{\rho_{\rm dm,0}}{a^3}.$$
(147)

Therefore, the total energy density of the universe is

$$\rho = \frac{\rho_I}{1 + (a/a_1)^4} + \frac{\rho_{b,0}}{a^3} + \frac{\rho_{dm,0}}{a^3} + \rho_\Lambda.$$
(148)

It can be rewritten as

$$\rho = \frac{\rho_I}{1 + \frac{\rho_I a^4}{\rho_{\rm rad,0}}} + \frac{\rho_{\rm b,0}}{a^3} + \frac{\rho_{\rm dm,0}}{a^3} + \rho_\Lambda.$$
(149)

This model can account for the whole evolution of the universe from the early inflation to its late accelerating expansion. It exhibits two de Sitter eras connected by a radiation era and a matter era. We detail these different periods below.

Remark: In this model there is no past singularity (no big bang) nor future singularity (no little or big rip).¹³ The universe exists eternally in the past and in the future. The scale factor tends to zero when $t \to -\infty$ and to infinity when $t \to +\infty$. It has been called the "aioniotic" universe [3, 4].

¹¹ It is possible that there exist different forms of radiation. For example, a complex SF with a repulsive $|\varphi|^4$ self-interaction can behave like radiation [49]. In that case, we should describe each form of radiation by a specific Lagrangian. However, for simplicity, we shall regroup all forms of radiation in the Lagrangian (141).

¹² We could also consider a more general equation of state of the form $P = \alpha \rho c^2$ with $\alpha \simeq 0$ [4, 6, 10].

¹³ We have assumed a non-phantom evolution. See Ref. [5] for cosmological models presenting a singular or peculiar late evolution (phantom models).

A. Early universe: inflation + radiation

In the inflation + radiation era (early universe), the energy density of the universe is given by

$$\rho = \frac{\rho_I}{1 + (a/a_1)^4} \tag{150}$$

with $a_1 = (\Omega_{\text{rad},0}/\Omega_{P,0})^{1/4} = 1.98 \times 10^{-32}$ if $\rho_I = \rho_P$ and $a_1 = (\Omega_{\text{rad},0}/\Omega_{e,0})^{1/4} = 1.18 \times 10^{-11}$ if $\rho_I = \rho_e$. It can be rewritten as

$$\rho = \frac{\rho_I}{1 + \frac{\rho_I a^4}{\rho_{\rm rad,0}}}.$$
(151)

We recover the results detailed in Sec. VII. When $a \ll a_1$, we are in the inflation era. The density is approximately constant $\rho \simeq \rho_I$. This leads to a phase of early accelerating expansion (or early inflation) where the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_I}$ (early de Sitter era). The universe is accelerating. The transition between the radiation era and the matter era takes place at $a_{eq} = \Omega_{rad,0}/\Omega_{m,0} = 3.00 \times 10^{-4}$ (see Sec. XB). When $a_1 \ll a \ll a_{eq}$, we are in the ordinary radiation era described by the linear equation of state $P_{rad} = \rho c^2/3$. The density decreases algebraically as $\rho = \rho_{rad,0}/a^4$. The scale factor increases algebraically with time as $t^{1/2}$ and the density decreases as t^{-2} . The universe is decelerating. We thus have a transition between a phase of accelerating expansion (vacuum energy/de Sitter) in the early universe and a phase of decelerating expansion in the radiation era. The transition takes place at $a \simeq a_1$. This transition between the inflation era and the radiation era is studied in detail in [3, 6]. The temporal evolution of the scale factor a(t) is given analytically (in reversed form) by

$$\sqrt{(a/a_1)^4 + 1} - \ln\left(\frac{1 + \sqrt{(a/a_1)^4 + 1}}{(a/a_1)^2}\right) = 2\left(\frac{8\pi}{3}\right)^{1/2}\frac{t}{t_I} + C \tag{152}$$

with $C \simeq 1 - \ln 2 + 2 \ln \epsilon$, where $\epsilon = a(t=0)/a_1 = 1.71 \times 10^{-30}$ (see Sec. VII E). In the inflation era $(a \ll a_1)$:

$$a \sim a(t=0)e^{(8\pi/3)^{1/2}t/t_I}, \qquad \rho \simeq \rho_I.$$
 (153)

In the radiation era $(a_1 \ll a \ll a_{eq})$:

$$a \sim \left(2\sqrt{\Omega_{\text{rad},0}}H_0t\right)^{1/2}, \qquad \frac{\rho}{\rho_0} \sim \frac{1}{\left(2H_0t\right)^2},$$
(154)

where $H_0 = (8\pi G \rho_0/3)^{1/2} = 2.195 \times 10^{-18} \,\text{s}^{-1}$ is the present value of the Hubble parameter.

B. Intermediate universe: radiation + matter

In the radiation + matter era (intermediate universe), the energy density of the universe evolves with the scale factor as

$$\rho = \frac{\rho_{\rm rad,0}}{a^4} + \frac{\rho_{\rm m,0}}{a^3}.$$
(155)

If we normalize the density by its present value, we get

$$\frac{\rho}{\rho_0} = \frac{\Omega_{\rm rad,0}}{a^4} + \frac{\Omega_{\rm m,0}}{a^3}.$$
(156)

When $a \ll a_{eq}$, we are in the ordinary radiation era described by a linear equation of state $P_{rad} = \rho c^2/3$. The density decreases algebraically as $\rho = \rho_{rad,0}/a^4$. The scale factor increases algebraically with time as $t^{1/2}$ and the density decreases as t^{-2} . The universe is decelerating. When $a \gg a_{eq}$, we are in the matter era (Einstein-de Sitter) with a vanishing pressure $P_m = 0$. The density decreases algebraically as $\rho = \rho_{m,0}/a^3$. The scale factor increases algebraically with time as $t^{2/3}$ and the density decreases as t^{-2} . The universe is decelerating. We note that $a \gg a_{eq}$ we are in the matter era (Einstein-de Sitter) with a vanishing pressure $P_m = 0$. The density decreases algebraically as $\rho = \rho_{m,0}/a^3$. The scale factor increases algebraically with time as $t^{2/3}$ and the density decreases as t^{-2} . The universe is decelerating. We thus have a transition between two phases of decelerating expansion (radiation and matter). The transition takes place at $a \simeq a_{eq}$ with

$$a_{\rm eq} = \frac{\Omega_{\rm rad,0}}{\Omega_{\rm m,0}} = 3.00 \times 10^{-4}.$$
 (157)

We have taken $\Omega_{rad,0} = 9.23765 \times 10^{-5}$ and $\Omega_{m,0} = 0.3075$. This transition between the radiation era and the matter era is studied in detail in [7]. The temporal evolution of the scale factor is given analytically (in reversed form) by

$$H_0 t = -\frac{2}{3} \frac{1}{(\Omega_{\rm m,0})^{1/2}} \left(\frac{2\Omega_{\rm rad,0}}{\Omega_{\rm m,0}} - a\right) \sqrt{\frac{\Omega_{\rm rad,0}}{\Omega_{\rm m,0}} + a} + \frac{4}{3} \frac{(\Omega_{\rm rad,0})^{3/2}}{(\Omega_{\rm m,0})^2}.$$
(158)

It can also be written as

$$a^{3} - 3\frac{\Omega_{\rm rad,0}}{\Omega_{\rm m,0}}a^{2} = \frac{9}{4}\Omega_{\rm m,0}H_{0}^{2}t^{2} - 6\frac{\Omega_{\rm rad,0}^{3/2}}{\Omega_{\rm m,0}}H_{0}t.$$
(159)

This is a cubic equation for a which can be solved by standard methods. However, in order to plot the curve a(t), it is simpler to compute t(a) and represent a versus t. In the radiation era we recover Eq. (154) and in the matter era we recover Eq. (165) given below.

C. Late universe: matter + dark energy (Λ CDM)

In the matter + dark energy era (late universe), the energy density is given by

$$\rho = \frac{\rho_{\mathrm{m},0}}{a^3} + \rho_{\Lambda}.$$
(160)

If we normalize the density by its present value, we get

$$\frac{\rho}{\rho_0} = \frac{\Omega_{\mathrm{m},0}}{a^3} + \Omega_{\Lambda,0}.$$
(161)

We recover the Λ CDM model. We note that, in the present model, dark energy comes from the generalized radiation associated with nonlinear electrodynamics. It arises from the constant term in the Lagrangian (141) taking into account the electromagnetic energy of point zero. Introducing $a'_2 = (\Omega_{m,0}/\Omega_{\Lambda,0})^{1/3} = 0.7634$ (we have taken $\Omega_{m,0} = 0.3075$ and $\Omega_{\Lambda,0} = 0.6911$), the relation between the energy density and the scale factor can be rewritten as

$$\frac{\rho}{\rho_{\Lambda}} = \frac{1}{(a/a_2')^3} + 1. \tag{162}$$

When $a_{eq} \ll a \ll a'_2$, we are in the matter era (Einstein-de Sitter) with a vanishing pressure $P_m = 0$. The density decreases algebraically as $\rho = \rho_{m,0}/a^3$. The scale factor increases algebraically with time as $t^{2/3}$ and the density decreases as t^{-2} . The universe is decelerating. When $a \gg a'_2$, we are in the dark energy era. The density is approximately constant $\rho \simeq \rho_{\Lambda}$, equal to the cosmological density. This leads to a phase of late accelerating expansion (or late inflation) where the scale factor increases exponentially rapidly with time as $e^{(8\pi/3)^{1/2}t/t_{\Lambda}}$ (late de Sitter era). The universe is accelerating. We thus have a transition between a phase of decelerating expansion in the matter era (Einstein-de Sitter) and a phase of accelerating expansion (vacuum energy/de Sitter) in the late universe. The transition takes place at $a \simeq a'_2$. This transition between the matter era and the dark energy era is studied in detail in [4, 6]. The temporal evolution of the scale factor a(t) and density $\rho(t)$ is given analytically by

$$\frac{a}{a_2'} = \sinh^{2/3} \left[\frac{3}{2} \left(\frac{8\pi}{3} \right)^{1/2} \frac{t}{t_\Lambda} \right], \qquad \frac{\rho}{\rho_\Lambda} = \frac{1}{\tanh^2 \left[\frac{3}{2} \left(\frac{8\pi}{3} \right)^{1/2} \frac{t}{t_\Lambda} \right]},\tag{163}$$

with $t_{\Lambda} = 1/\sqrt{G\rho_{\Lambda}} = (8\pi/\Lambda c^2)^{1/2} = 1.59 \times 10^{18} \,\mathrm{s}$ (cosmological time). It can also be written as

$$a = \left(\frac{\Omega_{m,0}}{\Omega_{\Lambda,0}}\right)^{1/3} \sinh^{2/3} \left(\frac{3}{2}\sqrt{\Omega_{\Lambda,0}}H_0t\right), \qquad \frac{\rho}{\rho_0} = \frac{\Omega_{\Lambda,0}}{\tanh^2\left(\frac{3}{2}\sqrt{\Omega_{\Lambda,0}}H_0t\right)}.$$
(164)

In the matter era $(a_{eq} \ll a \ll a'_2)$:

$$a \sim \left(\frac{3}{2}\sqrt{\Omega_{m,0}}H_0t\right)^{2/3}, \qquad \frac{\rho}{\rho_0} \sim \frac{1}{\left(\frac{3}{2}H_0t\right)^2}.$$
 (165)

In the dark energy era $(a \gg a'_2)$:

$$a \sim \left(\frac{\Omega_{m,0}}{4\Omega_{\Lambda,0}}\right)^{1/3} e^{\sqrt{\Omega_{\Lambda,0}}H_0 t}, \qquad \rho \simeq \rho_{\Lambda}.$$
 (166)

D. Numerical applications

In order to describe quantitatively the physical evolution of the universe from the early inflation to the late acceleration that we observe today we use the "radius of the universe" $R(t) = a(t)R_{\Lambda}$ with $R_{\Lambda} = ct_{\Lambda} = (8\pi/\Lambda)^{1/2} = 4.77 \times 10^{26}$ m defined in Appendix B. We also use the results of Appendix E. We consider the two models of Sec. VIIE which differ from each other only in the early universe. The temporal evolution of the radius and density of the universe in the two models are represented in Figs. 2 and 3.

Model I: In the first model, the density of the primordial universe (maximum density) is equal to the Planck density $\rho_I = \rho_P = 5.16 \times 10^{99} \,\mathrm{g\,m^{-3}}$. In that model, the universe starts at t = 0 (begining of the inflation) with a size equal to the Planck length $R(t = 0) = l_P = 1.62 \times 10^{-35} \,\mathrm{m}$ (i.e. $a(t = 0) = 3.39 \times 10^{-62}$) and reaches a size $R_c = R_1 \sim \lambda_A^* = 3.91 \times 10^{-5} \,\mathrm{m}$ of the order of the neutrino's Compton wavelength (i.e. $a_1 = a_c = 1.98 \times 10^{-32}$) at the end of the inflation which occurs on a timescale $t_1 = t_c = 23.8 \,t_P$ of the order of a few Planck times $t_P = 5.39 \times 10^{-44} \,\mathrm{s}$. The *e*-folding number is $N_0 = 68.5$ (the size of the universe increases by a factor $1/\epsilon \sim 10^{30}$ during the inflation). At t = 0 the density and the magnetic field are $\rho(t = 0) \simeq \rho_P$ and $B(t = 0) = 3.69 \times 10^{113} \,\mathrm{T}$. At $t = t_1 = t_c$ the density and the magnetic field are $\rho_1 = \rho_c = \rho_P/2$ and $B_1 = B_c = 1.08 \times 10^{54} \,\mathrm{T}$.¹⁴ We also note that the mass of the universe as a Planck black hole or a planckion particle) while its mass at the end of the inflation is $\rho_P R_c^3 \sim 10^{90} M_P$. This suggests that 10^{90} particles of mass M_P have been created during the inflation, implying that, after the radiation era, the mass of the universe is equal to $10^{62} M_P$ (see Appendix B).



FIG. 2: Temporal evolution of the radius of the universe in logarithmic scales in Model I (black) and Model II (red). The universe exists at all times in the past and in the future. There is no singularity (aioniotic universe). The early universe undergoes a phase of exponential inflation. During the early inflation, the scale factor increases by ~ 30 orders of magnitude in ~ 10^{-42} s (Model I) or in ~ 1 s (Model II). This is followed by the radiation era ($a \propto t^{1/2}$), by the matter era ($a \propto t^{2/3}$), and by the dark energy era responsible for the accelerated expansion of the universe observed at present. The universe exhibits two types of inflation corresponding to the Planck density ρ_P (Model I) or to the electron density (Model II) and a late inflation corresponding to the cosmological density ρ_A (dark energy or cosmological constant). The evolution of the early and late universe is remarkably symmetric. In our model it is described by two polytropic equations of state with index n = 1 and n = -1, respectively. The dashed line corresponds to the Λ CDM model which presents a big bang singularity at t = 0. We have also represented the location of the present universe that is just at the transition between the matter era and the dark energy era (cosmic coincidence).

Model II: In the second model, the density of the primordial universe (maximum density) is equal to the electron density $\rho_I = \rho_e = 4.07 \times 10^{16} \,\mathrm{g\,m^{-3}}$. This value is justified by applying the nonlinear electrodynamics with the Lagrangian from Eq. (79) to the electron (see Sec. XI). In that model, the universe starts at t = 0 (begining of the inflation) with a size of the order of the electron radius $R(t=0) \sim r_e = 2.82 \times 10^{-15} \,\mathrm{m}$ (i.e. $a(t=0) \sim 5.91 \times 10^{-42}$) and reaches a size $R_c = R_1 \sim \tilde{R}_2 = 7.07 \times 10^{15} \,\mathrm{m}$ of the order of the radius of a dark energy star of the stellar mass (i.e. $a_1 = a_c = 1.18 \times 10^{-11}$) at the end of the inflation which occurs on a timescale $t_1 = t_c = 23.8 \, t_e^*$ of the order of a few gravitoelectronic times $t_e^* = 0.0192 \,\mathrm{s}$. The *e*-folding number is $N_0 = 68.5$ (the size of the universe increases by a factor $1/\epsilon \sim 10^{30}$ during the inflation). At t = 0 the density and the magnetic field are

¹⁴ We have used Eqs. (68), (69) and (105) to compute the magnetic field.



FIG. 3: Temporal evolution of the density of the universe in logarithmic scales in Model I (black) and Model II (red). The density goes from a maximum value ρ_{max} equal to the Planck density ρ_P (Model I) or to the electron density ρ_e (Model II) to a minimum value $\rho_{\text{min}} = \rho_{\Lambda}$ equal to the cosmological density. These two bounds, which are fixed by fundamental constants of physics, are responsible for the early and late inflation of the universe. In between, the density decreases as t^{-2} . The dashed line corresponds to the Λ CDM model with a big bang singularity at t = 0.

 $\rho(t=0) \simeq \rho_e$ and $B(t=0) = 1.04 \times 10^{72} \text{ T}$. At $t_1 = t_c$ the density and the magnetic field are $\rho_1 = \rho_c = \rho_e/2$ and $B_c = 3.03 \times 10^{12} \text{ T}$ (see footnote 14). We also note that the mass of the universe at t=0 is equal to the electron mass $m_e = \rho_e r_e^3 = 9.11 \times 10^{-28} \text{ g}$ (the primordial universe has the same characteristics as an electron) while its mass at the end of the inflation is $\rho_e R_c^3 \sim 10^{90} m_e$. This suggests that 10^{90} particles of mass m_e have been created during the inflation, implying that, after the radiation era, the universe is made of 10^{83} electrons or 10^{80} protons, which is Eddington's number (see Appendix B).

After the inflation, the evolution of the universe is the same in the two models. The universe undergoes a radiation era, then enters in the matter era at $t_{eq} = 5.25 \times 10^4$ yrs (i.e. $a_{eq} = 3.00 \times 10^{-4}$) and in the dark energy era at $t'_2 = 10.2$ Gyrs (i.e. $a'_2 = 0.7634$). The age of the universe is $t_0 = 0.956 H_0^{-1} = 13.8$ Gyrs (i.e. $a_0 = 1$). Its size and density are $R_0 = R_{\Lambda} = 4.77 \times 10^{26}$ m and $\rho_0 = 8.62 \times 10^{-24}$ g m⁻³. The present magnetic field is $B_0 = 4.23 \times 10^{-10}$ T (see Sec. V).

XI. NONLINEAR ELECTRODYNAMICS BASED ON THE LAGRANGIAN $\mathcal{L} = -\mathcal{F}/(1 + \mathcal{F}/\mathcal{F}_*)$: ELECTROSTATICS

In this section, we apply the nonlinear electrodynamics considered previously to the electron in the spirit of the Born-Infeld [25, 26] theory. A similar approach has been developed by Kruglov [11]. We consider a purely electrostatic situation. We compute the electric field created by a pointlike charge $\sqrt{4\pi e}$ and determine the total electric energy that it carries. We then identify this electric energy with the mass-energy of the electron and obtain an estimate of its classical radius and density.

We consider a nonlinear electrodynamics based on the Lagrangian

$$\mathcal{L} = \frac{-\mathcal{F}}{1 + \frac{\mathcal{F}}{\mathcal{F}_*}}.$$
(167)

This is the Lagrangian (141) of the generalized radiation without the constant term (vacuum energy) that yields an infinite total energy. We leave the constant $\mathcal{F}_* = \rho_* c^2$ undetermined for the moment. It will be determined at the end by applying this model to the electron. The linear (Maxwell) electrodynamics is recovered in the limit $\mathcal{F}_* \to +\infty$. According to Eqs. (40), (41) and (167), the general expressions of the energy density and pressure are

$$\rho c^2 = \frac{\mathcal{F}}{1 + \frac{\mathcal{F}}{\mathcal{F}_*}} + \frac{E^2}{\left(1 + \frac{\mathcal{F}}{\mathcal{F}_*}\right)^2},\tag{168}$$

$$P = -\frac{\mathcal{F}}{1 + \frac{\mathcal{F}}{\mathcal{F}_*}} - \frac{E^2 - 2B^2}{3} \frac{1}{\left(1 + \frac{\mathcal{F}}{\mathcal{F}_*}\right)^2}.$$
(169)

A. Electric field

In electrostatics, using Eq. (47), the Lagrangian (167) takes the form

$$\mathcal{L} = \frac{E^2}{2\left(1 - \frac{E^2}{2\mathcal{F}_*}\right)}.$$
(170)

According to Eqs. (52) and Eq. (167) the electric field created by a pointlike charge $\sqrt{4\pi e}$ is given by

$$\frac{E(r)}{\left[1 - \frac{E^2(r)}{2\mathcal{F}_*}\right]^2} = \frac{e}{\sqrt{4\pi r^2}}.$$
(171)

In this nonlinear electrodynamics, the electric field is finite at the origin r = 0 (unlike in Maxwell's electrodynamics) with the maximum value $E(0) = E_* = \sqrt{2F_*}$. It decreases monotonically with the distance. At small distances $(r \ll r_*)$ it decreases as $E(r)/\sqrt{2F_*} = 1 - r/2r_* + ...,^{15}$ where

$$r_* = \left(\frac{e}{\sqrt{8\pi\mathcal{F}_*}}\right)^{1/2} \tag{172}$$

is a characteristic radius determined by the finite value of \mathcal{F}_* in the nonlinear electrodynamics based on Eq. (167). It can be interpreted as the radius of the electron in this model. It reduces to zero in Maxwell's electrodynamics $\mathcal{F}_* \to +\infty$. At large distances $(r \gg r_*)$ we recover the Coulomb law $E(r) \sim e/(\sqrt{4\pi}r^2)$. The electric field is plotted as a function of r in Fig. 4.



FIG. 4: Electric field E (normalized by $\sqrt{2\mathcal{F}_*}$) as a function of the distance r (normalized by r_*).

Remark: Conversely, Eq. (172) can be written as

$$\mathcal{F}_* = \rho_* c^2 = \frac{e^2}{8\pi r_*^4}.$$
(173)

Remark: Let us call ρ_e^* the extended charge density in the usual (Maxwell) electrodynamics that produces the same electric field as above. It can be determined by the usual Gauss law

$$\nabla \cdot \mathbf{E} = \rho_e^*,\tag{174}$$

where E(r) is given by Eq. (171). Using the Gauss theorem, we have $E(r) = Q(r)/(4\pi r^2)$ where $Q(r) = \int_0^r \rho_e^*(r') 4\pi r'^2 dr'$ is the charge contained within the sphere of radius r. Since $E(r) \sim e/(\sqrt{4\pi}r^2)$ for $r \to +\infty$, we have $Q = \lim_{r \to +\infty} E(r) 4\pi r^2 = \sqrt{4\pi e}$. Therefore, the total charge Q associated with the extended distribution ρ_e^* in linear electrodynamics coincides with the charge $\sqrt{4\pi e}$ of the singular point-charge in nonlinear electrodynamics. This is a general result that was first made in connection to the Born-Infeld model [25, 26].

¹⁵ By contrast, in the Born-Infeld [25, 26] model the electric field decreases as $E(r)/\sqrt{2\mathcal{F}_*} = 1 - \frac{1}{2}(r/r_*)^4 + \dots$

B. Energy density and pressure

The energy density and the pressure are given by Eqs. (53) and (54) with Eq. (167) yielding

$$\rho c^{2} = \frac{E^{2}(r) \left[1 + \frac{E^{2}(r)}{2\mathcal{F}_{*}} \right]}{2 \left[1 - \frac{E^{2}(r)}{2\mathcal{F}_{*}} \right]^{2}},$$
(175)

$$P = \frac{E^2(r) \left[1 - \frac{3E^2(r)}{2\mathcal{F}_*}\right]}{6 \left[1 - \frac{E^2(r)}{2\mathcal{F}_*}\right]^2}.$$
(176)

The energy density and the pressure diverge at the origin r = 0 as $\rho/\rho_* \sim 2(r_*/r)^2$ and $P/\rho_*c^2 \sim -(2/3)(r_*/r)^2$ respectively (this is the same behavior as in the Born-Infeld [25, 26] model). The energy density starts from $+\infty$ at r = 0 and decreases to 0^+ as $\rho/\rho_* \sim (r_*/r)^4$ for $r \to +\infty$. The pressure starts from $-\infty$ at r = 0, increases, vanishes at $r_0/r_* = 2/3^{3/4}$ (corresponding to $E_0/\sqrt{2\mathcal{F}_*} = 1/\sqrt{3}$ and $\rho_0/\rho_* = 1$), becomes positive, reaches a maximum $P_{\rm max}/\rho_*c^2 = 1/24$ at $r_m/r_* = 4/5^{3/4}$ (corresponding to $E_m/\sqrt{2\mathcal{F}_*} = 1/\sqrt{5}$ and $\rho_m/\rho_* = 3/8$) and decreases to 0^+ as $P/\rho_*c^2 \sim (1/3)(r_*/r)^4$ for $r \to +\infty$. The energy density and the pressure are plotted as a function of r in Fig. 5.



FIG. 5: Energy density ρc^2 (normalized by $\rho_* c^2$) and pressure P (normalized by $\rho_* c^2$) as a function of the distance r (normalized by r_*).

C. Equation of state

Eliminating the electric field E(r) between Eqs. (175) and (176) we obtain the equation of state

$$\frac{P}{\rho_* c^2} = \frac{1}{6} \left(-1 - 2\frac{\rho}{\rho_*} + \sqrt{1 + 8\frac{\rho}{\rho_*}} \right).$$
(177)

For small energy densities $\rho c^2 \rightarrow 0$ (large distances) we recover the usual equation of state of the radiation

$$P \sim \frac{1}{3}\rho c^2,\tag{178}$$

corresponding to Maxwell's linear electrodynamics. For large energy densities $\rho c^2 \rightarrow +\infty$ (short distances) we obtain the equation of state

$$P \sim -\frac{1}{3}\rho c^2. \tag{179}$$

This is a linear equation of state of the form $P = \alpha \rho c^2$ with $\alpha = -1/3$. The same equation of state is obtained in the Born-Infeld [25, 26] model at high densities (both for the electric field of the electron and for the magnetic universe).¹⁶ We note that the pressure in the core $(r < r_0)$ is negative. If we view the electron as an extended charge of typical radius r_* (see below), a negative pressure is necessary to counteract the ordinary electrostatic repulsion and ensure its cohesion. This is similar to the Poincaré stress [51, 52] introduced in the Abraham-Lorentz [53, 54] electromagnetic model of the electron to stabilize the particle (see Appendix F of [55]). The equation of state (177) is plotted in Fig. 6. We note that this "electric" equation of state differs from the "magnetic" equation of state (82) even though the electromagnetic Lagrangian is the same.



FIG. 6: Pressure P (normalized by ρ_*c^2) as a function of the energy density ρ (normalized by ρ_*). This is the electrostatic equation of state associated with the Lagrangian (167).

D. Classical radius of the electron and value of $\mathcal{F}_* = \rho_* c^2$

The electric energy density $\rho(r)c^2$ diverges at the origin r=0 as r^{-2} . However, the total electric energy

$$\mathcal{E} = \int_0^{+\infty} \rho(r) c^2 4\pi r^2 \, dr \tag{180}$$

is finite. This is like in the Born-Infeld [25, 26] model. Following Born [25] we shall identify the electrostatic energy with the mass of the electron via the relation¹⁷

$$\mathcal{E} = m_e c^2. \tag{181}$$

This relation gives an electromagnetic origin to the mass of the electron and allows one to determine \mathcal{F}_* (or ρ_* or r_*). Using Eq. (171), (172), (175) and (180) we find that the electrostatic energy of the electron is

$$\mathcal{E} = \frac{e^2}{r_*} \int_0^1 \frac{(1+y^2)(3y^2+1)}{4\sqrt{y}} \, dy = \frac{16}{15} \frac{e^2}{r_*},\tag{182}$$

where $y = E/\sqrt{2\mathcal{F}_*}$. Together with Eq. (181) this gives

$$m_e c^2 = \frac{16}{15} \frac{e^2}{r_*}.$$
(183)

¹⁶ Interestingly, this equation of state occurs in cosmology in relation to the Milne model of the universe [50]. This is also the equation of state of a gas of cosmic string. Therefore, it may describe a cosmic stringlike era (see, e.g., [49]).

¹⁷ This relation has a long history in physics even before Einstein's theory of relativity (see Appendix F of [55]). It appeared in relation to the Abraham-Lorentz model of the electron where it was believed that the mass of a particle had an electromagnetic origin [56]. The Born-Infeld [25, 26] theory of the electron can be considered as a revival of the old idea of the electromagnetic origin of mass; namely, that the electron is a singularity in the electromagnetic field and that its mass is purely electromagnetic. By contrast, it is not possible to identify $\mathcal{E} = m_e c^2$ in Maxwell's electrodynamics because the electromagnetic energy \mathcal{E} is infinite whereas m_e is finite.

Comparing Eq. (183) with Eq. (C1) we get

$$r_* = \frac{16}{15} r_e = 3.00 \times 10^{-15} \,\mathrm{m},\tag{184}$$

which may be interpreted as the electron radius in the present model (it turns out that r_* is very close to r_e since $16/15 \simeq 1.07$). In this sense, the present model (similarly to the Born-Infeld [25, 26] model) justifies the relation from Eq. (C1) defining the classical radius of the electron. In the present model (as in the Born-Infeld [25, 26] electrodynamics), the electron has a finite effective radius because the electric field and the electrostatic energy of a point charge are finite. This is basically due to the finite value of \mathcal{F}_* in the Lagrangian. According to Eqs. (C1), (173) and (184) we have

$$\mathcal{F}_* = \frac{e^2}{8\pi r_*^4} = \frac{1}{8\pi (\frac{16}{15})^4} \frac{e^2}{r_e^4} = \frac{1}{8\pi (\frac{16}{15})^4} \frac{m_e c^2}{r_e^3}.$$
(185)

Using Eqs. (75) and (C5), this gives

$$\rho_* = \frac{1}{8\pi (\frac{16}{15})^4} \frac{m_e}{r_e^3} = \frac{1}{8\pi (\frac{16}{15})^4} \rho_e = 1.25 \times 10^{15} \,\mathrm{g \, m^{-3}}.$$
(186)

Therefore, as could have been expected, we find that ρ_* is of the order of the density of the electron ρ_e (up to a factor 0.0307). This is much smaller than the Planck density ρ_P . These densities differ by 80 orders of magnitude (see Appendix C).

Remark: We must, however, point out a difficulty with this model. If we assume that \mathcal{F}_* is a universal constant and if we apply the same argument to the proton (which has a charge opposite to that of the electron) we would find the same radius and the same mass as the electron, which is obviously incorrect. This suggests that the mass of the proton has not an electromagnetic origin, even in a context of nonlinear electrodynamics.

E. Heisenberg-Euler Lagrangian

Following Kruglov [13] we can compare the previous Lagrangian model at the weak field limit with the Heisenberg-Euler [57] Lagrangian, which is the QED Lagrangian with one loop correction. The Heisenberg-Euler Lagrangian reads

$$\mathcal{L}_{\rm HE} \simeq -\mathcal{F} + \lambda \mathcal{F}^2 + \dots \tag{187}$$

with

$$\lambda = \frac{8\alpha^2}{45} \frac{\hbar^3}{m_e^4 c^5},\tag{188}$$

where $\alpha = e^2/\hbar c$ is the fine-structure constant. When $\mathcal{F}/\mathcal{F}_* \ll 1$, we can expand Eq. (167) to first order and we obtain

$$\mathcal{L} \simeq -\mathcal{F} + \frac{\mathcal{F}^2}{\mathcal{F}_*}.\tag{189}$$

Comparing this expression with the Heisenberg-Euler Lagrangian (187) we get

$$\mathcal{F}_* = \frac{45}{8\alpha^2} \frac{m_e^4 c^5}{\hbar^3}.$$
 (190)

Recalling that $\mathcal{F}_* = \rho_* c^2$ [see Eq. (75)] we find that

$$\rho_* = \frac{45}{8\alpha^2} \frac{m_e^4 c^3}{\hbar^3} = \frac{45}{8} \alpha \rho_e = 1.67 \times 10^{15} \,\mathrm{g \, m^{-3}},\tag{191}$$

where ρ_e is the electron density from Eq. (C6). Therefore, this argument confirms that ρ_* is of the order of the electronic density (up to a factor 0.0410). The comparison between Eqs. (186) and (191) gives $\alpha = 1/[45\pi(16/15)^4] = 1/183$.

F. Fundamental length

In the present model of nonlinear electrodynamics, the electric field created by a point like charge (electron) is nonsingular at the origin and the electrostatic energy is finite even though the central energy density diverges. On the other hand, the mass of a charged particle like the electron has a purely electromagnetic nature as in the Abraham-Lorentz model [53, 54]. If we identify the electromagnetic energy \mathcal{E} with the mass-energy $m_e c^2$ of the electron we can define the classical radius r_e of the electron and its density ρ_e .

These results are similar to the Born-Infeld [25, 26] electrodynamics and very different from Maxwell's electrodynamics where the electric field created by a point like charge is singular at the origin and the electrostatic energy is infinite. In that case, the electron has a vanishing radius and an infinite density. Therefore, its mass (which is finite) cannot have an electromagnetic origin.

The regularization of the divergences (infinities) is due to the finite value of \mathcal{F}_* which plays a role similar to the speed of light c in the theory of relativity (this analogy is at the basis of the Born-Infeld [25, 26] model who adopted a Lagrangian of nonlinear electrodynamics similar to the Lagrangian of a relativistic particle). The finiteness of \mathcal{F}_* prevents the electric field to be larger than $E(0) = E_* = \sqrt{2\mathcal{F}_*}$. It gives an upper bound on the possible electric field. Similarly, the finiteness of the speed of light c imposes v < c in relativistic mechanics. When $\mathcal{F}_* \to +\infty$ we recover Maxwell's electrodynamics. Similarly, when $c \to +\infty$, we recover Newton's mechanics.

If we assume that the nonlinear Lagrangian (167) applies both to the magnetic universe like in Sec. VII and to the electron like in the present section we come to the conclusion that ρ_I in Sec. VII should be identified to the electron density ρ_e , not to the Planck density ρ_P . This argument selects Model II with respect to Model I in Sec. VII E. Therefore, in this model, the primordial density of the universe (maximum density) is equal to the electron density, not to the Planck density. This implies (see Sec. VII E) that the duration of the inflation (~ 1 s) is much longer than usually believed (~ 10^{-42} s).

Following Kruglov [11], one can interpret the quantity $r_* \sim r_e$ as a fundamental length due to quantum gravity effects. Indeed, for strong electromagnetic fields, nonlinear electrodynamics may arise from possible quantum gravity corrections to linear electrodynamics. This is how a new parameter r_* with the dimension of a length is introduced in the model [see Eq. (172)]. We can substantiate this claim with the following arguments. We have already mentioned that r_e represents in Model II the initial size of the universe at time t = 0 (see Appendix E and Sec. X D). On the other hand, we will show in Sec. XII that the electron classical radius determines the correct value of the cosmological constant (vacuum energy) through Zeldovich's second formula [28, 29]. This justifies the Eddington formula relating the cosmological constant to the radius (or to the mass) of the electron and the other fundamental constants of physics [30, 32]. The electron radius can also be interpreted as a minimum scale in quantum gravity in the sense of Karolyhazy [31]. All these arguments suggest that r_e should be interpreted as a fundamental minimum length. Since this minimum length is much larger than the Planck length ($r_e \gg l_P$) this shows that quantum gravity is not a Planck scale phenomenon. This is very different from the results obtained in Model I where the initial size of the universe at t = 0 is the Planck length l_P .

XII. HEURISTIC CONNECTIONS BETWEEN NONLINEAR ELECTRODYNAMICS, QUANTUM GRAVITY, VACUUM ENERGY AND DARK ENERGY

In this section we relate the two cosmological models discussed in Sec. VIIE to the two models of vacuum energy introduced by Zeldovich [28, 29] and to the two models of minimum length introduced in quantum gravity by Karolyhazy [31] and Amelino-Camelia [27]. We show striking heuristic connections between these apparently disconnected topics.

A. Model I

Lemaître [58] was the first to understand that the effect of the cosmological constant Λ is equivalent to that of a fluid with a constant density $\rho_{\Lambda} = \Lambda c^2/8\pi G$ described by an equation of state $P = -\rho c^2$. He interpreted $\rho_{\Lambda} c^2$ as the vacuum energy density. However, he did not connect his interpretation with the zero-point energy, nor relate it to quantum mechanics. The origin of the vacuum energy was first discussed by Zeldovich [28, 29] and Sakharov [59] in relation to quantum field theory. When one tries to compute the vacuum energy density ρ_{Λ} from first principles, one encounters a severe problem of divergence at small scales (UV divergence). However, following the seminal study of Zeldovich [29], several procedures have been devised to deal with these divergences (see [60] for a review). Zeldovich [28, 29] introduced two models of vacuum energy, each depending on a fundamental mass scale m. In his first model, the regularized, finite, energy density of the vacuum which leads to the equation of state of vacuum $P = -\rho c^2$ as a consequence of relativistic Lorentz invariance reads¹⁸

$$\rho_{\Lambda}c^2 \sim \frac{m^4c^5}{\hbar^3} \sim \frac{\hbar c}{\lambda_C^4},\tag{192}$$

where we have introduced the Compton wavelength $\lambda_C = \hbar/mc$ of the elementary particle of mass m. It represents the relevant small scale cut-off λ_{\min} in Zeldovich's model I. The expression (192) of the vacuum energy density is commonly adopted in the literature [60]. We note that this expression of the vacuum energy density does not involve the gravitational constant G. This may unveil a problem with this approach if vacuum energy arises from quantum gravity (see below). In his original paper, Zeldovich [29] used Eq. (192) with the proton mass and obtained a discrepency of 40 orders of magnitude with the empirical cosmological density. If we use the Planck mass and the Planck length in Eq. (192), we find that $\rho_{\Lambda} \sim \rho_P$, yielding a discrepency of 120 orders of magnitude with the empirical value. This is the usual formulation of the cosmological constant problem [62]. Alternatively, if we adopt the measured value of the cosmological constant and reverse the relation from Eq. (192) we get

$$m_{\Lambda}^* = \left(\frac{\Lambda \hbar^3}{Gc}\right)^{1/4} = \sqrt{m_{\Lambda} M_P} = 5.04 \times 10^{-3} \,\mathrm{eV/c^2},$$
 (193)

$$\lambda_{\Lambda}^* = \left(\frac{G\hbar}{\Lambda c^3}\right)^{1/4} = \sqrt{R_{\Lambda}^* l_P} = 3.91 \times 10^{-5} \,\mathrm{m},\tag{194}$$

where $m_{\Lambda} = \hbar \sqrt{\Lambda}/c = 2.08 \times 10^{-33} \text{ eV/c}^2$ is the cosmon mass and $R_{\Lambda}^* = 1/\sqrt{\Lambda} = 9.49 \times 10^{25} \text{ m}$ is the cosmological length [30]. In Ref. [30] we arrived at the mass and length scales (193) and (194) by different considerations and we identified them with the mass and Compton wavelength of the neutrino. We conclude therefore that the mass m and the length λ_C leading to the observed value of the cosmological constant in Zeldovich's model I correspond to the mass and Compton wavelength of the neutrino.

Some authors have tried to determine the limit on the measurability of spacetime distances in quantum gravity or the minimum uncertainty of spacetime geodesics [27]. It arises when taking into account the quantum properties of the devices used for measurement. According to Amelino-Camelia [27] the minimum uncertainty of the measure of the length of an object of size l due to quantum fluctuations is given by

$$\delta l \sim (ll_P)^{1/2}.$$
 (195)

When applied to the Universe as a whole $(l \sim R_{\Lambda}^*)$ this gives

$$\lambda_{\min} = \sqrt{R_{\Lambda}^* l_P} = 3.91 \times 10^{-5} \,\mathrm{m},$$
(196)

which corresponds to Eq. (194). This minimum length is larger than the Planck length $l_P = 1.62 \times 10^{-35}$ m showing that quantum gravity is not a Planck scale phenomenon.

In Model I of Sec. VII E we have identified the primordial (maximum) density of the universe with the Planck density ρ_P . Then, we have shown that the universe starts with a size $R(0) = l_P$ equal to the Planck length at t = 0 and reaches a size $R_1 = \lambda_{\Lambda}^*$ equal to the neutrino's Compton wavelength at the end of the inflation of duration $t_c = 23.8 t_P$ (see Sec. VII E, Sec. X D and Appendix E).

Combining the above results, we conclude that the minimum length λ_{\min} in Zeldovich's model I leading to the correct value of the cosmological constant corresponds to:

(i) the Compton wavelength of the neutrino [Eq. (194)].

(ii) the limit on the measurability of spacetime distances [Eq. (196)] in quantum gravity according to Amelino-Camelia [27].¹⁹

(iii) The radius of the universe at the end of the inflation in model I of Sec. VII E [Eq. (E4)]. This radius defines an effective or physical "minimum" length $\lambda_{\Lambda}^* = 3.91 \times 10^{-5}$ m which could replace the Planck length $l_P = 1.62 \times 10^{-35}$ m and yield the correct value of the vacuum energy density when substituted into Eq. (192).

In conclusion, if we identify λ_{\min} to the Compton wavelength of the neutrino in Zeldovich's model I, we can express the Compton wavelength λ_{Λ}^* and the mass m_{Λ}^* of the neutrino as a function of the cosmological constant [see Eqs. (193) and (194)]. Conversely, assuming that the neutrino has these characteristic parameters, we can explain the measured value of Λ [see Eq. (3)].

¹⁸ The energy density of vacuum fluctuations can be related to the Casimir effect [61] and is given qualitatively by $\rho_{\Lambda}c^2 \sim (e^2/r)/r^3 \sim \hbar c/r^4$ where e^2/r is the electrostatic energy and we have used $e^2 \sim \hbar c$ in order of magnitude [30].

¹⁹ This lengthscale can also be related to the radius of a fifth extra dimension [63].

B. Model II

By considering the gravitational interaction energy between virtual pairs of the quantum electrodynamic vacuum, Zeldovich [28, 29] obtained another formula for the vacuum energy density²⁰

$$\rho_{\Lambda}c^2 \sim \frac{Gm^6c^4}{\hbar^4} \sim \frac{G\hbar^2}{c^2\lambda_C^6},\tag{198}$$

where we have introduced the Compton wavelength $\lambda_C = \hbar/mc$ of the particle of mass m. It represents the relevant minimum length λ_{\min} in Zeldovich's model II. We note that this expression of the vacuum density explicitly involves the gravitational constant G. Therefore, it may be related to a theory of quantum gravity. In his original papers, Zeldovich [28, 29] used Eq. (198) with the proton mass and obtained a discrepency of 7 orders of magnitude with the empirical cosmological density. If we use the Planck mass and the Planck length in Eq. (198), we find that $\rho_{\Lambda} \sim \rho_P$, yielding a discrepency of 120 orders of magnitude with the empirical value (cosmological constant problem). Alternatively, if we adopt the measured value of the cosmological constant and reverse the relation from Eq. (198) we get

$$m_e \sim \left(\frac{\Lambda \hbar^4}{G^2}\right)^{1/6} \sim (m_\Lambda M_P^2)^{1/3} = 6.77 \times 10^7 \,\mathrm{eV/c^2},$$
 (199)

$$\lambda_e \sim \left(\frac{G^2\hbar^2}{\Lambda c^6}\right)^{1/6} \sim (R_\Lambda^* l_P^2)^{1/3} = 2.92 \times 10^{-15} \,\mathrm{m.}$$
 (200)

In Ref. [30] we arrived at the mass and length scales (199) and (200) by different considerations and we identified them with the mass and Compton wavelength of the electron (in order of magnitude). Actually, we obtained accurate expressions of the electron mass and electron Compton wavelength under the form [30, 64, 65]

$$m_e \simeq \alpha \left(\frac{\Lambda \hbar^4}{G^2}\right)^{1/6}, \qquad \lambda_e = \frac{r_e}{\alpha} \simeq \frac{1}{\alpha} \left(\frac{G^2 \hbar^2}{\Lambda c^6}\right)^{1/6},$$
 (201)

where $\alpha = e^2/(\hbar c) \simeq 1/137$ is the fine-structure constant (see Appendix C). We see that Eq. (200) corresponds more closely to the classical electron radius r_e than to the electron Compton wavelength λ_e (we note that the classical electron radius does not depend explicitly on α). The formula (201) provides an accurate form of the Eddington formula [30, 32] relating the mass of the electron to the cosmological constant (or the converse).²¹ It can be written $m_e \simeq \alpha (m_\Lambda M_P^2)^{1/3}$ and $r_e = \alpha \lambda_e \simeq (R_\Lambda^* l_P^2)^{1/3}$. We conclude therefore that the mass m and the length λ_C leading to the observed value of the cosmological constant in Zeldovich's model II correspond to the typical mass and typical Compton wavelength of the electron (more precisely to m_e/α and $\alpha \lambda_e = r_e$).

Another formula has been obtained for the minimum uncertainty in the measure of the length of an object of size l due to quantum fluctuations. According to Karolyhazy [31] it is given by

$$\delta l \sim (ll_P^2)^{1/3}.$$
 (202)

This is the condition that the device used to make the measurement does not turn into a black hole. This expression was later related to the holographic principle and to the theory of quantum information. The minimum total uncertainty in the measurement of a length equal to the size of the universe $(l \sim R_{\Lambda}^*)$, which is a consequence of combining the principles of quantum mechanics and general relativity, is given by

$$\lambda_{\min} = (R_{\Lambda}^* l_P^2)^{1/3} = 2.92 \times 10^{-15} \,\mathrm{m},\tag{203}$$

$$\rho_{\Lambda}c^2 \sim \frac{Gm^2}{\lambda_C} \times \frac{1}{\lambda_C^3}.$$
(197)

 $^{^{20}}$ He wrote the vacuum energy density under the form

This expression assumes that the vacuum contains virtual pairs of particles with effective density $n \sim 1/\lambda_C^3$ and that these pairs have a gravitational energy of interaction Gm^2/λ_C .

²¹ It would be desirable to know if this formula is *exact* or just a good approximate relation (possibly the leading term in an expansion in powers of α).

which corresponds to Eq. (200). This minimum length is larger than the Planck length $l_P = 1.62 \times 10^{-35}$ m showing that quantum gravity is not a Planck scale phenomenon.

In Model II of Sec. VII E we have identified the primordial (maximum) density of the universe with the electron density ρ_e . This was justified by applying the same Lagrangian (79) of nonlinear electrodynamics both to the magnetic universe (see Sec. VII E) and to the electron (see Sec. XI). Then, we have shown that the universe starts with a size $R(0) = r_e$ equal to the classical radius of the electron at t = 0 and reaches a size $R_1 = \tilde{R}_2$ of the order of the radius of a dark energy star of the stellar mass at the end of the inflation of duration $t_c = 23.8 t_e^*$ (see Sec. VII E).

Combining the above results, we conclude that the minimum length λ_{\min} in Zeldovich's model II leading to the correct value of the cosmological constant corresponds to:

(i) the classical radius of the electron [Eqs. (200) and (201)].

(ii) the limit on the measurability of spacetime distances [Eq. (203)] in quantum gravity according to Karolyhazy [31].

(iii) The radius of the universe at the beginning of the inflation in model II of Sec. VII E [Eq. (E18)]. This radius defines an absolute "minimum" length $r_e = 2.82 \times 10^{-15}$ m which could replace the Planck length $l_P = 1.62 \times 10^{-35}$ m and yield the correct value of the vacuum energy density when substituted into Eq. (198).

In conclusion, if we identify λ_{\min} to the classical radius of the electron in Zeldovich's model II, we can express the radius r_e , the Compton wavelength λ_e and the mass m_e of the electron as a function of the cosmological constant [see Eqs. (199)-(201)] and we justify the mysterious Eddington relation [30, 32]. Conversely, we can express the cosmological constant in terms of r_e , λ_e or m_e and, by using the empirical value of the mass of the electron, we can explain the measured value of Λ [see Eq. (4)]. The physical reason to identify λ_{\min} to the classical radius of the electron is that it corresponds to the initial radius of the universe at t = 0 according to the model of nonlinear electrodynamics based on the Lagrangian (79) provided that this model applies both to the early magnetic universe and to the electron.

XIII. CONCLUSION

In this paper we have discussed the connection between the model of magnetic universe based on nonlinear electrodynamics introduced by Kruglov [11-13] and our model of universe based on a quadratic (or polytropic) equation of state [1-10]. These models both predict a period of early inflation followed by a radiation era. These models are essentially equivalent since the equation of state introduced in [1-10] can be deduced from the nonlinear electromagnetic Lagrangian introduced in [11-13] and *vice versa*. However, nonlinear electrodynamics might give a physical interpretation to our quadratic equation of state. It may arise from possible quantum gravity corrections to linear electrodynamics.

We have also generalized the nonlinear electromagnetic Lagrangian (79) by including a zero-point energy [see Eq. (141)] so that it describes a form of "generalized radiation" [1–10] that accounts simultaneously for the early inflation, the ordinary radiation, and the dark energy. Baryonic matter and dark matter are added as independent species. This leads to a complete model of universe (see Sec. X). This model essentially coincides with the Λ CDM model but it includes a period of early inflation. The density decreases from a maximum density ρ_I equal to the Planck density ρ_P or to the electron density ρ_e up to a minimum density ρ_Λ equal to the cosmological density (see Fig. 3). In this sense, it connects two de Sitter eras of accelerating expansion separated by a radiation era and a matter era of decelerated expansion (see Fig. 2).²²

By applying the nonlinear electrodynamics to the electron in the spirit of the Born-Infeld model [25, 26], following Kruglov [11], we have obtained an extended model of electron and justified the fact that the electron may have a finite radius and a finite density like in the Abraham-Lorentz model [53, 54]. Indeed, in this model, the electric field is nonsingular at the origin and the electric energy is finite.²³ By identifying the electrostatic energy with the mass of the electron we can obtain the electron radius and the electron density. This determines the fundamental density ρ_* , or the fundamental length r_* , in the nonlinear electromagnetic Lagrangian (see Sec. XI). Then, by applying the same nonlinear electromagnetic Lagrangian to cosmology we have shown that the initial density of the universe is

²² In this paper, we have considered a model similar to the Λ CDM model where the dark energy density is constant (n = -1) but, following [4, 5], we could consider more general models where the dark energy density varies with time (see Secs. VI and VIII with n < 0 and $n \neq -1$). There is, however, a difficulty with such models when interpreted in terms of nonlinear electrodynamics as pointed out in Sec. VI B.

²³ When the Lagrangian (167) is applied to the electron, we find that the electric field E(r) is finite at r = 0. By contrast, in cosmology, the magnetic field $B \sim 1/a^2$ diverges when $a \to 0$.

equal to the electron density ρ_e , its initial radius is equal to the electron radius r_e and its initial mass is equal to the electron mass m_e . We have then interpreted the radius of the electron as a fundamental minimum length and we have mentioned the connection with the result of Karolyhazy [31] in quantum gravity. By introducing this minimum length in the second Zeldovich [28, 29] formula of vacuum energy [see Eq. (198)] we have obtained a refined Eddington relation between the cosmological constant and the mass of the electron [see Eq. (4)]. This relation provides the exact (or almost exact) value of the cosmological constant. We have thus justified the Eddington relation [30, 32] and the value of the cosmological constant from nonlinear electrodynamics. This is a true prediction of the cosmological constant without free parameter since the mass of the electron is experimentally known. We have also shown that this cosmological model produces 10^{83} electrons or 10^{80} protons in the present universe, which is precisely the Eddington number [30, 32].

We have considered another model in which the initial density of the universe is equal to the Planck density ρ_P , its initial radius is equal to the Planck length l_P and its initial mass is equal to the Planck mass M_P . We have shown that the radius of the universe at the end of the inflation is equal to the Compton wavelength of the neutrino r_{ν} introduced in [30]. We have then interpreted this radius as an effective fundamental minimum length and we have mentioned the connection with the result of Amelino-Camelia [27] in quantum gravity. Then, by introducing this minimum length in the first Zeldovich [28, 29] formula of vacuum energy [see Eq. (192)] we have obtained the correct value of the cosmological constant [see Eq. (3)]. However, this is not a true prediction of Λ since the mass of the neutrino is not firmly known. Indeed, in [30] it has been precisely determined in terms of the cosmological constant.

In the two models of inflation described above the e-folding number is the same, $N_0 = 68.5$, corresponding to an increase of the size of the universe by 30 orders of magnitude between the begining and the end of the inflation. However, the duration of the inflation is very different. In Model I (where the initial density is equal to the Planck density $\rho_P = 5.16 \times 10^{99} \,\mathrm{g\,m^{-3}}$) the size of the universe increases from the Planck length $l_P = 1.62 \times 10^{-35} \,\mathrm{m}$ to the Compton wavelength of the neutrino $r_{\nu} = 3.91 \times 10^{-5} \,\mathrm{m}$ on a timescale $t_c = 23.8 \,t_P = 1.28 \times 10^{-42} \,\mathrm{s}$. In Model II (where the initial density is equal to the electron density $\rho_e = 4.07 \times 10^{16} \,\mathrm{g\,m^{-3}}$) the size of the universe increases from the classical electron radius $r_e = 2.82 \times 10^{-15} \,\mathrm{m}$ to the radius $\tilde{R}_2 = 7.07 \times 10^{15} \,\mathrm{m}$ of a dark energy star of the stellar mass on a timescale $t_c = 23.8 \,t_e^* = 0.457 \,\mathrm{s}$. We also note that the fundamental minimum length, which gives the correct value of the cosmological constant when introduced in the Zeldovich formula of vacuum energy, is different in the two models. In Model I it corresponds to the Compton wavelength of the neutrino which is equal to the size of the universe at the end of the inflation and in Model II it corresponds to the classical radius of the electron which is equal to the size of the universe at the begining of the inflation. In the two cases, the fundamental minimum length is much larger than the Planck length suggesting that quantum gravity is not a Planck scale phenomenon. It would be interesting to know if observations favor one model over the other.

Appendix A: The logotropic model

In a series of papers [64–70] we have developed a model of unified dark matter and dark energy based on the logotropic equation of state

$$P = A \ln \left(\frac{\rho}{\rho_P}\right),\tag{A1}$$

where A is a new fundamental constant of physics and $\rho_P = c^5/G^2\hbar = 5.16 \times 10^{99} \,\mathrm{g \, m^{-3}}$ is the Planck density. The constant A can be interpreted as a sort of "logotropic temperature" in a generalized thermodynamical framework [66–68]. This model leads to dark matter halos with a density profile that is flat at the center (thereby solving the core-cusp problem of the CDM model) and that decreases at large distances as

$$\rho \sim \left(\frac{A}{8\pi G}\right)^{1/2} \frac{1}{r}.\tag{A2}$$

At even larger distances the density falls like r^{-2} (isothermal profile) or like r^{-3} (NFW and Burkert profiles), or even more rapidly in order to have a finite mass. This confinement may result from an incomplete relaxation [65, 70]. The logotropic model implies that the dark matter halos ("small" scales) have a universal surface density given by

$$\Sigma_0 = \rho_0 r_h = 5.85 \left(\frac{A}{4\pi G}\right)^{1/2},\tag{A3}$$

where the prefactor is deduced from the theory (here ρ_0 is the central density and r_h is the halo radius where the central density is divided by 4). By applying the logotropic equation of state (A1) to the universe as a whole

36

("large" scales) we found that the universal constant A is related to the present value of the dark energy density $\rho_{\Lambda} \equiv \rho_{de,0} = 5.96 \times 10^{-24} \,\mathrm{g \, m^{-3}}$ by

$$A/c^{2} = \frac{\rho_{\Lambda}}{\ln\left(\frac{\rho_{P}}{\rho_{\Lambda}}\right)} = 2.10 \times 10^{-26} \,\mathrm{g \, m^{-3}}.$$
 (A4)

Using $\rho_{\Lambda} = \Lambda c^2 / 8\pi G$ with $\Lambda = 1.11 \times 10^{-52} \,\mathrm{m}^{-2}$, we can rewrite the foregoing equation as

$$A = 1.40 \times 10^{-4} \, \frac{c^4 \Lambda}{G}.$$
 (A5)

This allows us to express the universal surface density of dark matter halos in terms of the cosmological constant Λ of the Λ CDM model by

$$\Sigma_0 = 0.01955 \frac{c^2 \sqrt{\Lambda}}{G} = 133 \, M_{\odot} / \text{pc}^2. \tag{A6}$$

This value turns out to be in good agreement with the observational value $\Sigma_0^{\text{obs}} = 141^{+83}_{-52} M_{\odot}/\text{pc}^2$ [71]. The average surface density of dark matter halos is $\langle \Sigma \rangle = M_h/(\pi r_h^2) = 0.474 \Sigma_0 = 63.1 M_{\odot}/\text{pc}^2$ where the prefactor in the second equality is deduced from the theory. It is in good agreement with the observational value $\langle \Sigma \rangle_{\text{obs}} = 0.51 \Sigma_0^{\text{obs}} = 72^{+42}_{-27} M_{\odot}/\text{pc}^2$ [72]. The logotropic model therefore implies a universal gravitational acceleration

$$g = \frac{GM_h}{r_h^2} = \pi G \langle \Sigma \rangle = 1.49 \, G\Sigma_0. \tag{A7}$$

Using Eq. (A6) the universal gravitational acceleration can be expressed in terms of the cosmological constant as

$$g = 0.0291 \, c^2 \sqrt{\Lambda} = 2.76 \times 10^{-11} \,\mathrm{m \, s^{-2}}.$$
 (A8)

This value is in good agreement with the observational value $g_{\rm obs} = \pi G \langle \Sigma \rangle_{\rm obs} = 3.2^{+1.8}_{-1.2} \times 10^{-11} \,\mathrm{m/s^2}$ [72]. Using the foregoing relations, the asymptotic behavior of the logotropic density can be rewritten as

$$\rho \sim 0.121 \frac{\Sigma_0}{r} \sim 0.0811 \frac{g}{Gr} \sim 0.00236 \frac{c^2 \sqrt{\Lambda}}{Gr}.$$
(A9)

As noted in [70] this r^{-1} behavior is similar to the density cusp in the NFW model $\rho = \rho_s r_s / [r(1 + r/r_s)^2] \sim \rho_s r_s / r$. We find that

$$\rho_s r_s = 0.121 \,\Sigma_0 = 0.0811 \,\frac{g}{G} = 0.00236 \,\frac{c^2 \sqrt{\Lambda}}{G} = 16.0 \,M_{\odot}/\mathrm{pc}^2. \tag{A10}$$

We also noticed [64, 65] that the surface density $\Sigma_0 = 133 M_{\odot}/\text{pc}^2$ of dark matter halos (and the surface density of the universe $\Sigma_{\Lambda} = c^2 \sqrt{\Lambda}/G = 6800 M_{\odot}/\text{pc}^2$) is of the same order as the surface density of the electron

$$\Sigma_e = \frac{m_e}{r_e^2} = 54.9 \, M_{\odot} / \text{pc}^2. \tag{A11}$$

Using the accurate Eddington relation (C9) we find that

$$\Sigma_e \simeq \alpha \frac{c^2 \sqrt{\Lambda}}{G} \simeq \alpha \Sigma_\Lambda \simeq 0.373 \, \Sigma_0.$$
 (A12)

This relation may be interpreted in terms of the holographic principle [30, 64].

The circular velocity at the halo radius is

$$v_h^2 = \frac{GM_h}{r_h}.$$
(A13)

Combining the foregoing relations, we find that

$$\frac{v_h^4}{M_h} = Gg = \pi \langle \Sigma \rangle G^2 = 1.49 \,\Sigma_0 G^2. \tag{A14}$$

This relation is connected to the Tully-Fisher relation [73, 74] which involves the baryon mass M_b instead of the DM halo mass M_h . Introducing the baryon fraction $f_b = M_b/M_h \sim 0.17$, we obtain $(M_b/v_h^4)^{\text{th}} = f_b/(1.49 \Sigma_0 G^2) = 46.4 M_{\odot} \text{km}^{-4} \text{s}^4$ which is close to the observed value $(M_b/v_h^4)^{\text{obs}} = 47 \pm 6 M_{\odot} \text{km}^{-4} \text{s}^4$ [75].

More generally, the rotation curve is given by

$$v^2(r) = \frac{GM(r)}{r}.$$
(A15)

For $r \to +\infty$, we have

$$M(r) \sim 2\pi \left(\frac{A}{8\pi G}\right)^{1/2} r^2 \sim 0.760 \,\Sigma_0 r^2 \sim 0.510 \,\frac{gr^2}{G} \sim 0.0148 \,\frac{c^2 \sqrt{\Lambda} r^2}{G} \tag{A16}$$

and

$$v^2(r) \sim 2\pi G \left(\frac{A}{8\pi G}\right)^{1/2} r \sim 0.760 \, G\Sigma_0 r \sim 0.510 \, gr \sim 0.0148 \, c^2 \sqrt{\Lambda} r.$$
 (A17)

Asymptotically, the gravitational acceleration (the gravitational force F(r) = mg(r) by unit of mass) produced by the logotropic distribution tends to a constant

$$g(r) = \frac{d\Phi}{dr} = \frac{GM(r)}{r^2} \to 0.510 \, g = g_{\infty}.$$
 (A18)

The gravitational potential behaves as $\Phi(r) \sim g_{\infty} r$. We find $M(r)/M_{\odot} \sim 101 (r/\text{pc})^2$, $\log(\frac{v}{\text{km s}^{-1}}) = 1.32 + \frac{1}{2} \log(\frac{r}{\text{kpc}})$ and $g_{\infty} = 1.41 \times 10^{-11} \text{ m s}^{-2}$ to be compared with the observational expressions $M(r)^{\text{obs}}/M_{\odot} = 200^{+200}_{-120} (r/\text{pc})^2$, $\log(\frac{v_{\text{obs}}}{\text{km s}^{-1}}) = 1.47^{+0.15}_{-0.19} + 0.5 \log(\frac{r}{\text{kpc}})$ and $g_{\infty}^{\text{obs}} = 3^{+3}_{-2} \times 10^{-11} \text{ m s}^{-2}$ [76].

When the logotropic model is applied in a cosmological framework, the evolution of the density of the generalized radiation is given by [64–70]

$$\rho_{\text{Rad}} = \frac{\rho_I}{1 + \left(\frac{a}{a_1}\right)^4} + \rho_\Lambda (1 + 3B\ln a), \tag{A19}$$

where $B = A/(\rho_{\Lambda}c^2) = 1/\ln(\rho_P/\rho_{\Lambda}) = 3.53 \times 10^{-3}$. In the logotropic model, the density of dark energy increases slowly (logarithmically) with the scale factor. This corresponds to a phantom [5, 40] behavior leading to a little rip [77] (the energy density and the scale factor become infinite in infinite time). The model of Sec. IX, where the dark energy density is constant, is recovered for B = 0. Using Eqs. (55) and (61) the nonlinear electromagnetic Lagrangian inspired by the logotropic model is

$$\mathcal{L} = \frac{-\mathcal{F}}{1 + \frac{\mathcal{F}}{\rho_I c^2}} - \rho_\Lambda c^2 - \frac{3}{4} B \rho_\Lambda c^2 \ln\left(\frac{\mathcal{F}_*}{\mathcal{F}}\right). \tag{A20}$$

Remark: The above results can be expressed in terms of the present value of the Hubble parameter $H_0 = (8\pi G\rho_0/3)^{1/2} = 2.195 \times 10^{-18} \,\mathrm{s}^{-1}$ instead of the cosmological constant by using the relation $\rho_{\Lambda} = \Omega_{\Lambda,0}\rho_0$ with $\Omega_{\Lambda,0} = 0.6911$ giving

$$\Lambda c^2 = 3\Omega_{\Lambda,0} H_0^2 = 2.07 H_0^2. \tag{A21}$$

For example, the universal surface density can be written as $\Sigma_0 = 0.02815 H_0 c/G$ and the universal gravitational acceleration can be written as $g = 0.0419 H_0 c$. This relation explains why the fundamental constant $a_0 = g/f_b$ that appears in the MOND (modification of Newtonian dynamics) theory [78] is of order $a_0 \simeq H_0 c/4 = 1.65 \times 10^{-10} \text{ m/s}^2$ in good agreement with the observational value $a_0^{\text{obs}} = (1.3 \pm 0.3) \times 10^{-10} \text{ m/s}^2$. Note, however, that our model is completely different from the MOND theory.

Appendix B: The mass of the universe

1. Cosmological scales

The empirical value of the cosmological constant deduced from the observation of the accelerated expansion of the universe is $\Lambda = 1.11 \times 10^{-52} \,\mathrm{m}^{-2}$. By using general arguments based on physical considerations and dimensional

analysis, we can introduce cosmological scales. The cosmological density $\rho_{\Lambda}^* = \Lambda c^2/G = 1.50 \times 10^{-22} \,\mathrm{g\,m^{-3}}$ is of the order of the density of the universe, the cosmological time $t_{\Lambda}^* = 1/(G\rho_{\Lambda}^*)^{1/2} = 1/(c\sqrt{\Lambda}) = 3.16 \times 10^{17} \,\mathrm{s}$ is of the order of the age of the universe, the cosmological length $R_{\Lambda}^* = ct_{\Lambda}^* = 1/\sqrt{\Lambda} = 9.49 \times 10^{25} \,\mathrm{m}$ is of the order of the size of the visible universe (the distance travelled by a photon on a timescale t_{Λ}^*), and the cosmological mass $M_{\Lambda}^* = \rho_{\Lambda}^* R_{\Lambda}^{*3} = c^2/(G\sqrt{\Lambda}) = 1.28 \times 10^{56} \,\mathrm{g}$ is of the order of the mass of the universe.²⁴ In astronomical units, $t_{\Lambda}^* = 10.0 \,\mathrm{Gyrs}, R_{\Lambda}^* = 3.07 \,\mathrm{Gpc}$ and $M_{\Lambda}^* = 6.42 \times 10^{22} \,M_{\odot}$. The typical number of electrons in the universe is $N_e \sim M_{\Lambda}^*/m_e \sim 10^{83}$ and the typical number of protons (Eddington's number) is $N_p \sim M_{\Lambda}^*/m_p \sim 10^{80}.^{25}$ These quantities play an important role in the theory of large numbers [30].

2. Evolution of the mass of the universe during the different epochs

We define the radius and the mass of the of the universe at time t by

$$R(t) = a(t)R_{\Lambda}, \qquad M(t) = \rho(t)R(t)^3. \tag{B1}$$

During the inflation era, the density of the universe is constant ($\rho = \rho_I$) and the mass of the universe increases as

$$M(t) = \rho_I R(t)^3. \tag{B2}$$

During the radiation era, the density of the universe decreases as $\rho \sim R^{-4}$ and the mass of the universe decreases as

$$M(t) \sim \frac{1}{R(t)}.$$
(B3)

During the matter era, the density of the universe decreases as $\rho \sim R^{-3}$ and the mass of the universe is constant

$$M(t) = \text{cst.} \tag{B4}$$

During the dark energy era, the density of the universe is constant ($\rho = \rho_{\Lambda}$) and the mass of the universe increases as

$$M(t) = \rho_{\Lambda} R(t)^3. \tag{B5}$$

The evolution of the mass of the universe as a function of the radius in Models I and II of Sec. VIIE is plotted in Fig. 7.



FIG. 7: Evolution of the mass of the universe as a function of the radius in Model I (black) and Model II (red).

²⁴ These quantities are just orders of magnitude. They are given without any prefactor, and this is why they have been written with the symbol * (we have $\rho_{\Lambda}^* = 8\pi\rho_{\Lambda}, t_{\Lambda}^* = t_{\Lambda}/\sqrt{8\pi}, R_{\Lambda}^* = R_{\Lambda}/\sqrt{8\pi}$, and $M_{\Lambda}^* = M_{\Lambda}/\sqrt{8\pi}$). These relations can be derived from the Friedmann equations by using the fact that the present density of the universe is of the order of the cosmological density on account of the cosmic coincidence [30].

²⁵ These estimates assume that the Universe is made only of electrons or protons, which is of course not correct. We should also take into account dark matter and dark energy. But, because of the cosmic coincidence, the density of baryonic matter, dark matter and dark energy is of the same order of magnitude at present so our estimates are meaningful.

3. Model II

In Model II, the initial density of the universe is equal to the density of the electron: $\rho_I = \rho_e$. The initial radius of the universe is equal to the radius of the electron $(R(0) = r_e)$ and its initial mass is equal to the mass m_e of the electron (see Appendix E). At t = 0, the primordial universe has the same characteristics as the electron but it is "unstable" and "explodes". This picture can be viewed as a refinement of the "primeval atom" of Lemaître [33]. Let us follow its expansion accross the ages. Between the begining and the end of the inflation, its radius increases by 30 orders of magnitude (see Appendix E). Therefore, at the end of the inflation, its radius is $R_1 \sim 10^{30} r_e$ and its volume is $R_1^3 \sim 10^{90} r_e^3$. Since its density is constant ($\rho \sim \rho_e$) its mass at the end of the inflation (or at the begining of the radiation era) is $M_1 \sim 10^{90} m_e$.

Between the end of the inflation (or the beginning of the radiation era) and the time of radiation-matter equality, the radius of the universe passes from R_1 to $R_{eq} = a_{eq}R_{\Lambda}$. Using Eqs. (112) and (157), we find that $R_{eq}/R_1 = a_{eq}/a_1 \sim 10^7$. Therefore, its radius increases by 7 orders of magnitude. Using Eq. (B3), we find that its mass at the time of radiation-matter equality is $M_{eq} \sim 10^{-7}M_1 \sim 10^{83}m_e$.

This mass remains constant during the matter era (see Eq. (B4)) and may be identified with the present mass of the universe: $M_{\Lambda} \sim 10^{83} m_e$. This shows that the mass of the universe is equal to $N_e \sim 10^{83}$ electrons of mass m_e . Since $m_p/m_e = 1836$, we find that the mass of the universe is equal to $N_p \sim 10^{80}$ protons of mass m_p . This justifies the Eddington number [30, 32]. These results are in good agreement with the observations.

4. Model I

In Model I, the initial density of the universe is equal to the Planck density: $\rho_I = \rho_P$. The initial radius of the universe is equal to the Planck length $(R(0) = l_P)$ and its initial mass is equal to the Planck mass M_P (see Appendix E). At t = 0, the primordial universe has the same characteristics as a Planck black hole (or a "planckion" particle) but it is "unstable" and "explodes". Let us follow its expansion accross the ages. Between the begining and the end of the inflation, its radius increases by 30 orders of magnitude (see Appendix E). Therefore, at the end of the inflation, its radius is $R_1 \sim 10^{30} l_P$, which is of the order of the Compton wavelength of the neutrino, and its volume is $R_1^3 \sim 10^{90} l_P^3$. Since its density is constant ($\rho \sim \rho_P$) its mass at the end of the inflation (or at the begining of the radiation era) is $M_1 \sim 10^{90} M_P$.

Between the end of the inflation (or the beginning of the radiation era) and the time of radiation-matter equality, the radius of the universe passes from R_1 to $R_{\rm eq} = a_{\rm eq}R_{\Lambda}$. Using Eqs. (108) and (157), we find that $R_{\rm eq}/R_1 = a_{\rm eq}/a_1 \sim 10^{28}$. Therefore, its radius increases by 28 orders of magnitude. Using Eq. (B3), we find that its mass at the time of radiation-matter equality is $M_{\rm eq} \sim 10^{-28}M_1 \sim 10^{62}M_P$. This mass remains constant during the matter era (see Eq. (B4)) and may be identified with the present mass

This mass remains constant during the matter era (see Eq. (B4)) and may be identified with the present mass of the universe: $M_{\Lambda} \sim 10^{62} M_P$. This shows that the mass of the universe is equal to $N_P \sim 10^{62}$ particles of mass M_P . Although this result is quantitatively correct, no such particles of mass M_P exist in abundance in the universe. Therefore, this picture is not in agreement with the observations. This suggests that Model II may be more relevant than Model I.

Appendix C: The electron

The classical radius r_e of the electron is defined through the relation

$$E = m_e c^2 = \frac{e^2}{r_e}.$$
(C1)

This equation expresses the equality (in order of magnitude) between the rest-mass energy of the electron and its electrostatic energy, assuming that the electron has a certain size. This is a convenient manner to define the "radius" of the electron. This relation first appeared in the Abraham-Lorentz [53, 54] model of the extended electron with an electromagnetic mass and later in the Born-Infeld [25, 26] theory of nonlinear electrodynamics (see Appendix F of [55] for a short review of these old theories). Recalling the value of the charge of the electron $e = 4.80 \times 10^{-13} \text{ g}^{1/2} \text{ m}^{3/2} \text{ s}^{-1}$ and its mass $m_e = 9.11 \times 10^{-28} \text{ g} = 0.511 \text{ MeV/c}^2$, we obtain

$$r_e = \frac{e^2}{m_e c^2} = 2.82 \times 10^{-15} \,\mathrm{m.}$$
 (C2)

The Compton wavelength of the electron is $\lambda_e = \hbar/(m_e c) = 3.86 \times 10^{-13}$ m. It is related to the classical radius of the electron by

$$\lambda_e = \frac{r_e}{\alpha} \simeq 137 \, r_e,\tag{C3}$$

where

$$\alpha = \frac{e^2}{\hbar c} \simeq \frac{1}{137} \simeq 7.30 \times 10^{-3} \tag{C4}$$

is Sommerfeld's fine-structure constant.²⁶ The typical electron density is

$$\rho_e = \frac{m_e}{r_e^3} = 4.07 \times 10^{16} \,\mathrm{g}\,\mathrm{m}^{-3}.$$
(C5)

It can also be written as

$$\rho_e = \frac{m_e^4 c^6}{e^6} = \frac{m_e^4 c^3}{\alpha^3 \hbar^3} \tag{C6}$$

or as

$$\rho_e = \frac{\alpha \hbar}{c r_e^4}.\tag{C7}$$

The characteristic (dynamical) time associated with the electron is

$$t_e = \left(\frac{m_e r_e^3}{e^2}\right)^{1/2} = \frac{e^2}{m_e c^3} = \frac{r_e}{c} = 9.40 \times 10^{-24} \,\mathrm{s.}$$
(C8)

This is the time it takes for a light wave to travel accross the "size" of an electron. This timescale first appeared in the Abraham-Lorentz [53, 54] theory of the extended electron when they tried to calculate the recoil force on an accelerated charged particle caused by the particle emitting electromagnetic radiation. It can be written as $t_e = \alpha \hbar/(m_e c^2)$. It is connected to the flight time in the relativistic extension of Nelson's stochastic quantum mechanics [79] developed by Lehr and Park [80]. This is what Caldirola [81] called the "chronon", which is a sort of "quantum of time". This is also the unit of time provided by the atomic constants that Dirac used in his cosmological theory based on a large number hypothesis [30, 82, 83].

In [30] we have obtained an accurate formula that relates the mass of the electron to the cosmological constant:

$$m_e \simeq \alpha \left(\frac{\Lambda \hbar^4}{G^2}\right)^{1/6}.$$
 (C9)

This equation can be viewed as an accurate form of Eddington formula [30, 32]. The classical radius of the electron is then given in good approximation by

$$r_e \simeq \left(\frac{G^2 \hbar^2}{\Lambda c^6}\right)^{1/6}.$$
 (C10)

With the empirical value $\Lambda = 1.11 \times 10^{-52} \,\mathrm{m}^{-2}$ of the cosmological constant deduced from the observations of the accelerated expansion of the universe we obtain the approximate value $8.80 \times 10^{-28} \,\mathrm{g}$ for the mass of the electron which is very close to the measured value $m_e = 9.11 \times 10^{-28} \,\mathrm{g}$ (similarly we get $2.92 \times 10^{-15} \,\mathrm{m}$ for the classical radius of the electron which is very close to $r_e = 2.82 \times 10^{-15} \,\mathrm{m}$). Using Eq. (C9) we find that

$$\lambda_e \simeq \frac{1}{\alpha} \left(\frac{G^2 \hbar^2}{\Lambda c^6}\right)^{1/6}, \qquad \rho_e \simeq \alpha \left(\frac{\Lambda^2 c^9}{G^4 \hbar}\right)^{1/3}, \qquad t_e \simeq \left(\frac{G^2 \hbar^2}{\Lambda c^{12}}\right)^{1/6}.$$
 (C11)

²⁶ Since quantum effects enter at a distance of the order λ_e which is much larger than r_e , a purely classical electromagnetic model of the electron is not relevant.

In [30] we have developed the theory of large numbers pioneered by Weyl, Eddington and Dirac. We have introduced the "largest large number" [30]

$$\chi = \frac{\rho_P}{\rho_{\Lambda}^*} = \frac{c^3}{G\hbar\Lambda} = 3.44 \times 10^{121} \sim 10^{120},\tag{C12}$$

which is the ratio between the Planck density and the cosmological density. In terms of this number, we have

$$\frac{m_e}{M_P} \sim \chi^{-1/6} \sim 10^{-20}, \quad \frac{r_e}{l_P} \sim \chi^{1/6} \sim 10^{20}, \quad \frac{\rho_e}{\rho_P} \sim \chi^{-2/3} \sim 10^{-80}, \quad \frac{t_e}{t_P} \sim \chi^{1/6} \sim 10^{20}.$$
 (C13)

Remark: If we assume that Eqs. (C9) and (C10) are *exact* (see footnote 21), then the "theoretical" values of the cosmological constant and of the "largest large number" are

$$\Lambda_{\rm th} = \frac{G^2 m_e^6}{\alpha^6 \hbar^4} = 1.36 \times 10^{-52} \,\mathrm{m}^{-2}, \qquad \chi_{\rm th} = \left(\frac{c\hbar\alpha^2}{Gm_e^2}\right)^3 = 2.81 \times 10^{121}. \tag{C14}$$

Appendix D: The neutrino

In [30] we have suggested that the mass of the neutrino is related to the cosmological constant by the relation

$$m_{\Lambda}^* = \left(\frac{\Lambda \hbar^3}{Gc}\right)^{1/4} = 5.04 \times 10^{-3} \,\mathrm{eV/c^2}.$$
 (D1)

The Compton wavelength $\lambda_C = \hbar/mc$ of the neutrino is then given by

$$\lambda_{\Lambda}^* = \left(\frac{G\hbar}{\Lambda c^3}\right)^{1/4} = 3.91 \times 10^{-5} \,\mathrm{m}.\tag{D2}$$

To make the numerical applications we have used the empirical value $\Lambda = 1.11 \times 10^{-52} \,\mathrm{m}^{-2}$ of the cosmological constant. In terms of the "largest large number" defined by Eq. (C12) we have

$$\frac{m_{\Lambda}^*}{M_P} \sim \chi^{-1/4} \sim 10^{-30}, \quad \frac{\lambda_{\Lambda}^*}{l_P} \sim \chi^{1/4} \sim 10^{30}.$$
 (D3)

Remark: Eliminating the cosmological constant between Eqs. (C9) and (D1) we obtain the following relation

$$\frac{m_e^6}{(m_\Lambda^*)^4} = \alpha^6 M_P^2 \tag{D4}$$

between the mass of the neutrino and the mass of the electron. This allows us to determine the "exact" value of the neutrino mass, independently of the uncertainty on the value of the cosmological constant. We find $(m_{\Lambda}^*)_{\rm th} = 5.30 \times 10^{-3} \, {\rm eV/c^2}$.

Appendix E: Transition between the inflation era and the radiation era

As discussed in Sec. VII the transition between the inflation era and the radiation era corresponds to a value of the scale factor a_1 given by Eq. (100).²⁷ We want to determine the quantity

$$\epsilon = \frac{a(t=0)}{a_1},\tag{E1}$$

²⁷ Basically, this relation can be obtained as follows. The density of radiation evolves with the scale factor as $\rho_{\rm rad} = \rho_{\rm rad,0}/a^4$. Owing to the fact that $\rho_{\rm rad} \sim \rho_I$ at the end of the inflation, we get $a_1 \sim (\rho_{\rm rad,0}/\rho_I)^{1/4}$. If we make the additional rough approximation $\rho_{\rm rad,0} \sim \rho_{\Lambda}$ on account of the cosmic coincidence we obtain $a_1 \sim (\rho_{\Lambda}/\rho_I)^{1/4}$ in order of magnitude.

(i) Model I: We first assume that the initial density of the universe (upper bound) is of the order of the Planck density: $\rho_I = \rho_P = c^5/G^2\hbar = 5.16 \times 10^{99} \,\mathrm{g \, m^{-3}}$. According to Eq. (100) we have

$$a_1 = \left(\frac{\Omega_{\rm rad,0}}{\Omega_{\Lambda,0}}\right)^{1/4} \left(\frac{\rho_{\Lambda}}{\rho_P}\right)^{1/4} = 1.98 \times 10^{-32} \sim \chi^{-1/4} \sim 10^{-30},\tag{E2}$$

where χ is the "largest large number" defined by Eq. (C12). Therefore, the radius of the universe at the end of the inflation (i.e. at the transition between the inflation era and the radiation era) is

$$R_1 = a_1 R_{\Lambda} = \left(\frac{\Omega_{\text{rad},0}}{\Omega_{\Lambda,0}}\right)^{1/4} \left(\frac{8\pi G\hbar}{\Lambda c^3}\right)^{1/4} = 9.42 \times 10^{-6} \,\text{m.}$$
(E3)

Comparing Eq. (E3) with Eq. (D2) we see that the radius of the universe at the end of the inflation is of the order of the Compton wavelength of the neutrino:

$$R_1 \sim \lambda_{\Lambda}^*. \tag{E4}$$

On the other hand, it is natural to identify R(0) with the Planck length:

$$R(0) = l_P = ct_P = \left(\frac{G\hbar}{c^3}\right)^{1/2} = 1.62 \times 10^{-35} \,\mathrm{m.}$$
(E5)

Therefore

$$a(t=0) = \frac{R(0)}{R_{\Lambda}} = \frac{l_P}{R_{\Lambda}} = \left(\frac{G\hbar\Lambda}{8\pi c^3}\right)^{1/2} = 3.39 \times 10^{-62} \sim \chi^{-1/2},\tag{E6}$$

where we have used

$$\frac{l_P}{R_\Lambda} = \frac{t_P}{t_\Lambda} = \left(\frac{\rho_\Lambda}{\rho_P}\right)^{1/2} = \left(\frac{1}{8\pi\chi}\right)^{1/2} \sim \chi^{-1/2} \sim 10^{-60}$$
(E7)

to get the last estimates. We then get

$$\epsilon = \frac{R(0)}{R_1} = \frac{l_P}{R_1} = \frac{l_P}{a_1 R_\Lambda}.$$
 (E8)

Using Eq. (E2), we can write Eq. (E8) as

$$\epsilon = \left(\frac{\Omega_{\Lambda,0}}{\Omega_{\rm rad,0}}\right)^{1/4} \left(\frac{\rho_P}{\rho_\Lambda}\right)^{1/4} \frac{l_P}{R_\Lambda}.$$
(E9)

Substituting Eqs. (C12) and (E7) into Eq. (E9) we obtain

$$\epsilon = \left(\frac{\Omega_{\Lambda,0}}{\Omega_{\rm rad,0}}\right)^{1/4} (8\pi\chi)^{-1/4} = 1.71 \times 10^{-30} \sim \chi^{-1/4} \sim 10^{-30}.$$
 (E10)

The increase in size of the universe during the inflation is

$$\frac{R_1}{l_P} = \frac{1}{\epsilon} = \left(\frac{\Omega_{\text{rad},0}}{\Omega_{\Lambda,0}}\right)^{1/4} (8\pi\chi)^{1/4} = 5.83 \times 10^{29} \sim \chi^{1/4} \sim 10^{30}.$$
(E11)

²⁸ This time corresponds to the moment of the big bang singularity ($\rho = \infty$ and a = 0) in the case of a purely radiative early universe.

Therefore, during the inflation, the size of the universe increases by about 30 orders of magnitude. We note that the value of ϵ gives an *e*-folding number $N_0 = 68.5$ which is fully consistent with the observations (see Sec. VII E). This value has been obtained by assuming $R(0) = l_P$. Conversely, assuming $N \simeq 60 - 70$ from the observations implies that $R(0) \sim l_P$.

(ii) *Model II:* We now assume that the initial density of the universe is of the order of the density of the electron: $\rho_I = \rho_e = 4.07 \times 10^{16} \,\mathrm{g \, m^{-3}}$. According to Eq. (100) we have

$$a_1 = \left(\frac{\Omega_{\rm rad,0}}{\Omega_{\Lambda,0}}\right)^{1/4} \left(\frac{\rho_{\Lambda}}{\rho_e}\right)^{1/4} = 1.18 \times 10^{-11} \sim \chi^{-1/12} \sim 10^{-10},\tag{E12}$$

where we have used Eqs. (C12) and (C13) to get the last estimates. Therefore, the radius of the universe at the end of the inflation (i.e. at the transition between the inflation era and the radiation era) is

$$R_1 = a_1 R_{\Lambda} \simeq \frac{1}{\alpha^{1/4}} \left(\frac{\Omega_{\rm rad,0}}{\Omega_{\Lambda,0}}\right)^{1/4} \left(\frac{512\pi^3 G\hbar}{c^3 \Lambda^5}\right)^{1/12} = 5.82 \times 10^{15} \,\mathrm{m},\tag{E13}$$

where we have used Eq. (C11). In Ref. [30] we have introduced the lengthscale

$$\tilde{R}_2 = \left(\frac{\hbar G}{c^3 \Lambda^5}\right)^{1/12} = \chi^{5/12} \, l_P = 7.07 \times 10^{15} \,\mathrm{m} \tag{E14}$$

that we have interpreted as being the radius of a dark energy star of the stellar mass. Therefore, we see that the radius of the universe at the end of the inflation is of the order of the radius of a dark energy star of the stellar mass:

1

$$R_1 \sim \tilde{R}_2.$$
 (E15)

We note that this radius is gigantic as compared to the Compton wavelength of the neutrino obtained when we take $\rho_I = \rho_P$ [see Eqs. (D2) and (E4)]. They differ by 20 orders of magnitude ($\chi^{1/6} \sim 10^{20}$). On the other hand, we shall determine R(0) in order to get the same value of ϵ as above (see Sec. VII E). First, we note that

$$\epsilon = \frac{R(0)}{R_1} = \frac{R(0)}{a_1 R_\Lambda} = \left(\frac{\Omega_{\Lambda,0}}{\Omega_{\rm rad,0}}\right)^{1/4} \left(\frac{\rho_e}{\rho_\Lambda}\right)^{1/4} \frac{R(0)}{R_\Lambda},\tag{E16}$$

where we have used Eq. (E12). Equating Eqs. (E16) and (E9) we find that

$$R(0) = \left(\frac{\rho_P}{\rho_e}\right)^{1/4} l_P.$$
(E17)

Using Eq. (C7), we get

$$R(0) = \frac{r_e}{\alpha^{1/4}} = 9.65 \times 10^{-15} \,\mathrm{m.}$$
(E18)

In this model, the initial radius of the universe is of the order of the size of the electron instead of being of the order of the Planck length [see Eq. (E5)]. This ensures that ϵ is given by Eq. (E10) in agreement with the observations (the size of the universe increases by about 30 orders of magnitude during the inflation). Conversely, assuming that the initial radius of the universe is of the order of the size of the electron, we find that ϵ is given by Eq. (E10). Finally, we find that

$$a(t=0) = \frac{R(0)}{R_{\Lambda}} = \frac{1}{\alpha^{1/4}} \frac{r_e}{R_{\Lambda}} = 2.10 \times 10^{-41} \sim \chi^{-1/3} \sim 10^{-40},$$
(E19)

where we have used Eqs. (C13) and (E7) to get the last estimates. The increase in size of the universe during the inflation is

$$\frac{R_1}{R(0)} = \frac{\alpha^{1/4} R_1}{r_e} = \frac{1}{\epsilon} = \left(\frac{\Omega_{\rm rad,0}}{\Omega_{\Lambda,0}}\right)^{1/4} (8\pi\chi)^{1/4} = 5.83 \times 10^{29} \sim \chi^{1/4} \sim 10^{30}.$$
(E20)

- [1] P.H. Chavanis, AIP Conference Proceedings 1548, 75 (2013)
- [2] P.H. Chavanis, Journal of Gravity **2013**, 682451 (2013)
- [3] P.H. Chavanis, Eur. Phys. J. Plus **129**, 38 (2014)
- [4] P.H. Chavanis, Eur. Phys. J. Plus 129, 222 (2014)
- [5] P.H. Chavanis, arXiv:1208.1185
- [6] P.H. Chavanis, Universe 1, 357 (2015)
- [7] P.H. Chavanis, Phys. Rev. D 92, 103004 (2015)
- [8] P.H. Chavanis, Eur. Phys. J. Plus 130, 181 (2015)
- [9] P.H. Chavanis, J. Phys.: Conf. Ser. 1030, 012009 (2018)
- [10] P.H. Chavanis, Universe 8, 92 (2022)
- [11] S.I. Kruglov, Ann. Phys. 353, 299 (2015)
- [12] S.I. Kruglov, Phys. Rev. D 92, 123523 (2015)
- [13] S.I. Kruglov, Int. J. Mod. Phys. A 35, 2050168 (2020)
- [14] V.A. De Lorenci, R. Klippert, M. Novello, J.M. Salim, Phys. Rev. D 65, 063501 (2002)
- [15] M. Novello, S.E. Perez Bergliaffa, J.M. Salim, Phys. Rev. D 69, 127301 (2004)
- [16] C.S. Camara, M.R. de Garcia Maia, J.C. Carvalho, J.A.S. Lima, Phys. Rev. D 69, 123504 (2004)
- [17] M. Novello, E. Goulart, J.M. Salim, S.E. Perez Bergliaffa, Class. Quantum Grav. 24, 3021 (2007)
- [18] D.N. Vollick, Phys. Rev. D 78, 063524 (2008)
- [19] M. Novello, J.M. Salim, A.N. Araújo, Phys. Rev. D 85, 023528 (2012)
- [20] R. García-Salcedo, T. Gonzalez, I. Quiros, Phys. Rev. D 89, 084047 (2014)
- [21] L. de la Peña, A.M. Cetto, A.V. Hernández, The Emerging Quantum. The Physics Behind Quantum Mechanics (Springer, 2015)
- [22] A. Övgün, G. Leon, J. Magaña, K. Jusufi, Eur. Phys. J. C 78, 462 (2018)
- [23] H.B. Benaoum, A. Övgün, Class. Quantum Grav. 38 135019 (2021)
- [24] H.B. Benaoum, G. Leon, A. Övgün, H. Quevedo, Eur. Phys. J. C 83, 367 (2023)
- [25] M. Born, Proc. Roy. Soc. **143**, 410 (1933)
- [26] M. Born, L. Infeld, Proc. Roy. Soc. 144, 425 (1934)
- [27] G. Amelino-Camelia, Nature **398**, 216 (1999)
- [28] Ya. B. Zel'dovich, J. Exp. Theor. Phys. Lett. 6, 316 (1967)
- [29] Ya. B. Zel'dovich, Sov. Phys. Uspek. 11, 381 (1968)
- [30] P.H. Chavanis, Phys. Dark Univ. 44, 101420 (2024)
- [31] F. Karolyhazy, Nuovo Cimento A **42**, 390 (1966)
- [32] A.S. Eddington, Proc. Roy. Soc. A 133, 605 (1931)
- [33] G. Lemaître, L'expansion de l'espace, Revue des questions scientifiques 20, 391 (1931)
- [34] L.D. Landau, E.M. Lifshitz The Classical Theory of Fields, Pergamon Press (1951)
- [35] S. Weinberg, Gravitation and Cosmology (John Wiley, 2002)
- [36] Friedmann A. 1922, Z. Physik, 10, 377
- [37] Friedmann A. 1924, Z. Physik, 21, 326
- [38] G. Lemaître, Ann. Soc. Sci. Bruxelle 47, 49 (1927)
- [39] G. Lemaître, Monthly Not. Roy. Astron. Soc. 91, 483 (1931)
- [40] R.R. Caldwell, Phys. Lett. B 545, 23 (2002)
- [41] A.H. Guth, Phys. Rev. D 23, 347 (1981)
- [42] Planck Collaboration, Astron. Astrophys. 571, 66 (2014)
- [43] Planck Collaboration, Astron. Astrophys. 594, A13 (2016)
- [44] R. Tolman, P. Ehrenfest, Phys. Rev. **36**, 1791 (1930)
- [45] D. Lemoine, M. Lemoine, Phys. Rev. D 52, 1955 (1995)
- [46] E.J. Kim, A.V. Olinto, R. Rosner, Astrophys. J. 468, 28 (1996)
- [47] A.R. Liddle, H. Lyth, Cosmological Inflation and Large-Scale Structure (Cambridge University Press, 2000)
- [48] J. Binney, S. Tremaine, *Galactic Dynamics* (Princeton Series in Astrophysics, 2008)
- [49] A. Suárez, P.H. Chavanis, Phys. Rev. D 95, 063515 (2017)
- [50] E.A. Milne, Zeit. für Astrophy. 6, 1 (1933)
- [51] H. Poincaré, Comptes Rendus 140, 1504 (1905)
- [52] H. Poincaré, Rend. Circ. Matem. Palermo 21 129 (1906)
- [53] M. Abraham, Theorie der Elektrizitat (Teubner, Leipzig, 1905)
- [54] H.A. Lorentz The Theory of Electrons (Teubner, Leipzig, 1909)
- [55] P.H. Chavanis, Phys. Rev. D **107**, 103503 (2023)
- [56] J.J. Thomson, Phil. Mag. 11, 229 (1881)

- [57] W. Heisenberg, E. Euler, Z. Physik 98, 714 (1936)
- [58] G. Lemaître, Proc. Nat. Acad. Sci. 20, 12 (1934)
- [59] A.D. Sakharov, Dokl. Akad. Nauk SSSR 177, 70 (1967)
- [60] J. Martin, Comptes Rendus Physique 13, 566 (2012)
- [61] H.B.G. Casimir, Proc. Kon. Ned. Akad. Wet. 51, 793 (1948)
- [62] S. Weinberg, Rev. Mod. Phys. 61, 1 (1989)
- [63] A. Dupays, B. Lamine, A. Blanchard, Astron. Astrophys. 554, A60 (2013)
- [64] P.H. Chavanis, Phys. Dark Univ. 24, 100271 (2019)
- [65] P.H. Chavanis, Phys. Dark Univ. 37, 101098 (2022)
- [66] P.H. Chavanis, Eur. Phys. J. Plus 130, 130 (2015)
- [67] P.H. Chavanis, Phys. Lett. B **758**, 59 (2016)
- [68] P.H. Chavanis, S. Kumar, J. Cosmol. Astropart. Phys. 5, 018 (2017)
- [69] P.H. Chavanis, Astronomy 1, 126 (2022)
- [70] P.H. Chavanis, Phys. Rev. D 106, 063525 (2022)
- [71] F. Donato et al., Mon. Not. R. Astron. Soc. 397, 1169 (2009)
- [72] G. Gentile, B. Famaey, H. Zhao, P. Salucci, Nature 461, 627 (2009)
- [73] R.B. Tully, J.R. Fisher, Astron. Astrophys. 54, 661 (1977)
- [74] S.S. McGaugh, J.M. Schombert, G.D. Bothun, W.J.G. de Blok, Astrophys. J. 533, L99 (2000)
- [75] S.S. McGaugh, Astron. J. **143**, 40 (2012)
- [76] M.G. Walker, S.S. McGaugh, M. Mateo, E.W. Olszewski, R. Kuzio de Naray, Astrophys. J. 717, L87 (2010)
- [77] P.H. Frampton, K.J. Ludwick, R.J. Scherrer, Phys. Rev. D 84, 063003 (2011)
- [78] M. Milgrom, Astrophys. J. **270**, 365 (1983)
- [79] E. Nelson, Phys. Rev. **150**, 1079 (1966)
- [80] W.J. Lehr, J.L. Park, J. Math. Phys. 18, 1235 (1977)
- [81] P. Caldirola, Nuovo Cimento 45, 549 (1978)
- [82] P.A.M. Dirac, Nature 139, 323 (1937)
- [83] P.A.M. Dirac, Proc. Roy. Soc. A 165, 199 (1938)