

Group-theoretical classification of orientable objects and particle phenomenology

D. M. Gitman^{1,2*} and A. L. Shelepin^{3†}

¹ P.N. Lebedev Physical Institute,

53 Leninskiy ave., 119991 Moscow, Russia.

² Institute of Physics, University of São Paulo,

Rua do Matão, 1371, CEP 05508-090, São Paulo,, Brazil

³MIREA – Russian Technological University,

Prospect Vernadskogo, 78, 117454, Moscow, Russia

June 4, 2024

Abstract

In our previous works, we have proposed a quantum description of relativistic orientable objects by a scalar field on the Poincaré group. This description is, in a sense, a generalization of ideas used by Wigner, Casimir and Eckart back in the 1930's in constructing a non-relativistic theory of a rigid rotator. The present work is a continuation and development of the above mentioned our works. The position of the relativistic orientable object in Minkowski space is completely determined by the position of a body-fixed reference frame with respect to the space-fixed reference frame, and can be specified by elements q of the motion group of the Minkowski space - the Poincaré group $M(3,1)$. Quantum states of relativistic orientable objects are described by scalar wave functions $f(q)$ where the arguments $q = (x, z)$ consist of Minkowski space-time points x , and of orientation variables z given by elements of the matrix $Z \in SL(2, C)$. Technically, we introduce and study the so-called double-sided representation $\mathbf{T}(\mathbf{g})f(q) = f(g_l^{-1}qg_r)$, $\mathbf{g} = (g_l, g_r) \in \mathbf{M}$, of the group \mathbf{M} , in the space of the scalar functions $f(q)$. Here the left multiplication by g_l^{-1} corresponds to a change of space-fixed reference frame, whereas the right multiplication by g_r corresponds to a change of body-fixed reference frame. On this basis, we develop a classification of the orientable objects and draw the attention to a possibility of connecting these results with the particle phenomenology. In particular, we demonstrate how one may identify fields described by linear and quadratic functions of z with known elementary particles of spins $0, \frac{1}{2}$, and 1 . The developed classification does not contradict the phenomenology of elementary particles and, moreover, in some cases give its group-theoretic explanation.

1 Introduction

It is known that the first and most known example of a theory in which wave functions depend on some orientation variables is the non-relativistic theory of a rigid rotator, constructed by Wigner, Casimir and Eckart back in the 1930's (see Ref. [6] for references and

*dmitrygitman@hotmail.com

†alex@shelepin.msk.ru

historical remarks). In their construction, one reference frame (laboratory, space-fixed, *s.r.f.* in what follows) is connected with surrounding objects, while another one (body-fixed, *b.r.f.* in what follows) is connected with a rotating body. Correspondingly, there appear two sets of operators of the angular momentum: one related to the *s.r.f.* (left generators of the rotation group \hat{J}_k^L), and another related to the *b.r.f.* (right generators of the rotation group \hat{J}_k^R). These two sets of generators commute with each other, and the rotator Hamiltonian is constructed from the right generators \hat{J}_k^R . The interpretation of the generators \hat{J}_k^R as projections of the momentum on the *b.r.f.* belongs to Wigner and Casimir (1931), and underlies the theory of molecular spectra. In fact, the non-relativistic theory of the rigid rotator is formulated as a theory of a scalar field on the group $SO(3) \sim SU(2)$.

It should be also noted that a description of relativistic spinning particles can be reformulated in terms of one scalar field depending on the Minkowski space-time coordinates as well as on some auxiliary continuous variables that describe spin. Such a reformulation has a long history, starting with the work [7] of Ginzburg and Tamm. At the end of 1940's and the beginning of 1950's, independently by various authors, see Refs. [7, 8, 9, 10] (mainly in connection constructing of relativistic wave equations (RWE)), were introduced some fields depending not only on the Minkowski space-time coordinates but also on a certain set of spinning variables. A systematic treatment of these fields as fields on homogeneous spaces of the Poincaré group was presented by Finkelstein in 1955; see Ref. [11]. He also considered a classification and explicit constructions of homogeneous spaces of the Poincaré group that contains the Minkowski space which is a homogeneous space of the latter group. In 1964, Lurçat suggested to construct a quantum theory on the whole Poincaré group, instead on the Minkowski space, see Ref. [12], but this idea did not receive immediate development.

In 70-90s of the last century, ideas of considering fields on homogeneous spaces of the Poincaré group, gained a certain development, in particular, in the Refs [13, 14, 15, 16, 17, 18, 19, 20, 21]. Properties of different spaces were studied, as well as some possibilities of introducing interactions in the spin phase-space and possibilities of constructing corresponding Lagrangian formulations. It was found that the choice of a homogeneous space generates restrictions on the corresponding scalar fields. Thus, authors of the Ref. [13] had arrived at the conclusion that the minimal dimension of a homogeneous space suitable for a description of integer and semi-integer spins is equal to eight.

In our previous works [1, 3, 4, 5], following the ideas of the pioneer works cited above, we have developed a quantum description of relativistic orientable objects by a scalar field $f(q)$, $q = (x, z)$ that depends on the Minkowski space-time coordinates $x = (x^\mu) = (x^0, \mathbf{x})$, $\mathbf{x} = (x^k)$, and on the orientation variables $z = (z_{\alpha\beta})$ given by elements of the matrix $Z \in SL(2, C)$, in fact, on the Poincaré group $M(3, 1)$, $q \in M(3, 1)$.

At the beginning, we have proposed an approach to constructing fields on groups of motions in Euclidean and pseudo-Euclidean spaces, elaborating in detail the cases of 2, 3- and 4 dimensions; see Ref. [1]. In that construction, a scalar field appears as a generating function of usual multicomponent fields. In particular, it has been demonstrated that, as distinct from the case of scalar fields on homogeneous spaces, the field on a group as a whole is closed with respect to discrete transformations. The discrete transformations (C, P, T) correspond to involutive automorphisms of the group and are reduced to a complex conjugation and to a change of the arguments of scalar functions $f(q)$; see Ref. [2]. Most types of RWE were written in terms of scalar functions $f(q)$.

By analogy with the non-relativistic theory of the rigid rotator, in the relativistic case, it is supposed that one reference frame (laboratory, space-fixed, *s.r.f.*) is connected with surrounding objects, while another one (body-fixed, *b.r.f.*) is connected with an

orientable object. The position of the latter is completely determined by the position of the corresponding *b.r.f.* with respect to some *s.r.f.*, such that all the positions can be given by elements of the Poincaré group. It implies that quantum states of the relativistic orientable objects are described (as was already said above) by scalar wave functions $f(q)$ on the Poincaré group.

On this basis, in the Ref. [3] two types of transformations in the space of scalar functions $f(q)$ (see just below) were defined and studied. In the Refs. [4, 5] a method of classifying such objects with respect to the corresponding transformation properties of the functions $f(q)$ was proposed, as well as possible ways of introducing interactions of orientable objects.

By the present consideration we develop results of the latter works. As an important part of this development we represent a more complete classification of the orientable objects and draw the attention of readers to the possibility of connecting these results with the particle phenomenology. In particular, we observe the possibility of identifying fields described by linear and quadratic functions of z with known elementary particles of spin $0, \frac{1}{2}$, and 1.

Now we pass to technical details of the mentioned above transformations. In this regard, it is useful once again to stress that the position of an orientable object is completely determined by the position of the corresponding *b.r.f.* with respect to some *s.r.f.*. Symmetries of the space containing the orientable object (space-time or external symmetries) determine the behavior of wave functions with respect to the left transformations $T_L(g_l)$, $g_l \in M(3, 1)_{ext}$, whereas the symmetries of the orientable object itself (internal symmetries) determine the behavior of wave functions with respect to the right transformations $T_R(g_r)$, $g_r \in M(3, 1)_{int}$,

$$T_L(g_l) f(q) = f(g_l^{-1}q), \quad T_R(g_r) f(q) = f(qg_r). \quad (1)$$

As it is known, operations (1) define left and right regular representations of the Poincaré group. Studying these representations is equivalent to studying representation $\mathbf{T}(\mathbf{g})$, of the group $\mathbf{M} = M(3, 1)_{ext} \times M(3, 1)_{int}$, $\mathbf{g} \in \mathbf{M}$, in the space of scalar functions $f(q)$,

$$\mathbf{T}(\mathbf{g})f(q) = f(g_l^{-1}qg_r), \quad \mathbf{g} = (g_l, g_r) \in \mathbf{M}. \quad (2)$$

Here the left multiplication by g_l^{-1} corresponds to a change of the *s.r.f.*, whereas the right multiplication by g_r corresponds to a change of the *b.r.f.*. It is easy to verify that Eq. (2) specifies a representation $\mathbf{T}(\mathbf{g})$ of the group \mathbf{M} in the space of the scalar functions $f(q)$. In what follows we call such a representation the *double-sided* representation.

A classification of functions $f(q)$ on a group is the subject of the harmonic analysis on Lie groups and is carried out using the maximal sets of commuting operators of the group; see e.g. [24]. The number of the operators in the maximal set is equal to the number of the group parameters, and the set consists of the Casimir operators and of the (equal in number) functions of the generators in left and right regular representations (left and right generators).

It is important to emphasize that due to the fact that the groups $M(3, 1)_{ext}$ and $M(3, 1)_{int}$ act in the same space of function $f(q)$, Casimir operators constructed from right and left generators coincide. This means that states transformed by a fixed representation of $M(3, 1)_{ext}$ are transformed by the same representation of $M(3, 1)_{int}$.

In 3 + 1 dimensions, according to the above said, and as a consequence of Eq. (2), there appear two sets of 10 generators, the right and the left ones. Having in hands these generators one can construct a maximal set of commuting operators acting in the space of the functions $f(q)$. The maximal set of commuting operators consists of 10

operators – two Casimir operators, four functions of left generators and four functions of right generators. The last four functions of generators $M(3, 1)_{int}$, which are not expressed in terms of generators $M(3, 1)_{ext}$, commute with all the generators $M(3, 1)_{ext}$, i.e., they correspond to the internal quantum numbers of the orientable object. The question arises about their physical meaning and (due to the coincidence of the Casimir operators) the possible connection of their eigenvalues with the eigenvalues of the generators of external symmetries $M(3, 1)_{int}$.

Thus, considering the two-sided representation of the group $\mathbf{M} = M(3, 1)_{ext} \times M(3, 1)_{int}$, we naturally come to the question of the relation between external and internal symmetries and corresponding quantum numbers. In this regards, one ought to mention that in the 1960s, attempts were made to unite internal and external symmetries in the framework of a one group. Soon, however, the so-called no-go theorem [22] was proved (under some very general assumptions), stating that the symmetry group of the S -matrix is locally isomorphic to a direct product of the Poincaré group and the group of internal symmetries. However, on this basis, one sometimes makes too strong conclusion that a non-trivial relation between internal and external symmetries is impossible.

As was already said, transformations (2) form the direct product of the groups of internal and external symmetries, which removes the case under consideration from the prohibitions of this theorem. Moreover, as is demonstrated below, a non-trivial relation between internal and external quantum numbers is possible. Both transformation groups, corresponding to changing the *s.r.f.* and the *b.r.f.*, are acting in the same space of scalar functions $f(q)$ and have the same Casimir operators which define the mass and the spin. By fixing eigenvalues of the Casimir operators, and therefore fixing a representation, we obviously impose some conditions on the spectra of both left and right operators that enter the maximal set. Thus, in spite of the fact that the left and the right generators commute, their spectra are not independent. Below we are observed such a relationship on some concrete examples.

We note that the maximal sets of commuting operators are different for the massless and the massive cases, that is why, in what follows, we consider them separately.

By orientable objects, we understand states that are transformed according to irrep of the group $M(3, 1)_{ext}$. By fixing irrep of $M(3, 1)_{ext}$, we, as a consequence of the law of transformation (2), fix irrep of $\mathbf{M} = M(3, 1)_{ext} \times M(3, 1)_{int}$. This means that irreps (describing multiplets of particles) of $\mathbf{M} = M(3, 1)_{ext} \times M(3, 1)_{int}$ decompose into a sum of $M(3, 1)_{ext}$ irreps, differing by eigenvalues of generators $M(3, 1)_{int}$ (internal quantum numbers). In other words, different particles are states belonging to irrep \mathbf{M} , not connected by transformations of $M(3, 1)_{ext}$. Note that, unlike the external symmetries the internal ones can be violated, which is quite similar to the situation observed in the case of the rigid rotator, for which internal symmetries are the symmetries of the rotating body itself.

The work is organized as follows. In Sections 2 and 3, we construct and study maximal sets of commuting operators in the space of functions $f(q)$ on the Poincaré group for the cases of the mass $m = 0$ and $m \neq 0$. In Sections 4 and 5, we represent a classification of massless orientable objects of chirality $0, \pm 1/2, \pm 1, \pm 3/2$ and formulate corresponding RWE. In Section 6 we give a classification of orientable objects with mass $m \neq 0$ and spin $0, 1/2, 1, 3/2$. In Sections 7 and 8 we consider the ultra-relativistic limit and formulate corresponding RWE. A comparison of the obtained classification of the orientable objects with the particle phenomenology is given in Sect. 9; some concluding remarks are summarized in the same Section. In the Appendix (Section 9) we construct generators of the Lorentz group in the space of the functions $f(z)$.

2 Field on the Poincaré group and maximal sets of commuting operators

The functions $f(q)$ on the Poincaré group $M(3, 1)$ depend on 10 parameters,

$$q = X, Z, \quad X = \{x^\mu\}, \quad Z = \{z_{\underline{a}}^\alpha\} = \begin{pmatrix} z_{\underline{1}}^1 & z_{\underline{2}}^1 \\ z_{\underline{1}}^2 & z_{\underline{2}}^2 \end{pmatrix} = \begin{pmatrix} z^1 & \underline{z}^1 \\ z^2 & \underline{z}^2 \end{pmatrix} \in SL(2, C), \quad (3)$$

where $\det Z = 1$. Action of the groups $M(3, 1)_{ext}$ and $M(3, 1)_{int}$ in the space of functions $f(q)$

$$M(3, 1)_{ext} : \quad T_L(g_l)f(q) = f(g_l^{-1}q), \quad (4)$$

$$M(3, 1)_{int} : \quad T_R(g_r)f(q) = f(qg_r), \quad (5)$$

corresponds to the replacements of s.r.f. and b.r.f., respectively. Representations (4), (5) provide the most general consideration, since regular representations contain all (up to equivalence) irreducible representations (irreps) of the group [23]. Transformations from $M(3, 1)_{ext}$ correspond to the symmetries of the containing space, or external ones, and transformations from $M(3, 1)_{int}$ corresponds to symmetries of the orientable object, or internal symmetries.

The actions on the left and the right correspond to two sets of generators of the Poincaré group. There are ten generators each in the left and right regular representations, which for brevity we will call the left and right generators.

The generators that correspond to translations and rotations have the form

$$M(3, 1)_{ext} : \quad \hat{p}_\mu = i\partial/\partial x^\mu, \quad \hat{J}_{\mu\nu} = \hat{L}_{\mu\nu} + \hat{S}_{\mu\nu}, \quad \mu, \nu = 0, 1, 2, 3, \quad (6)$$

$$M(3, 1)_{int} : \quad \hat{p}_{\underline{m}}^R = -v^\nu_{\underline{m}} \hat{p}_\nu, \quad \hat{J}_{\underline{mn}}^R = \hat{S}_{\underline{mn}}^R, \quad \underline{m}, \underline{n} = 0, 1, 2, 3, \quad (7)$$

where $\hat{L}_{\mu\nu} = i(x_\mu \partial_\nu - x_\nu \partial_\mu)$ are the operators of orbital momentum projections and $\hat{S}_{\mu\nu}$ are the operators of spin projections. Here $v^\mu_{\underline{n}} = \frac{1}{2}(\sigma^\mu)_{\dot{\beta}\alpha}(\bar{\sigma}_{\underline{n}})^{\underline{a}\underline{b}}z_{\underline{a}}^\alpha z_{\underline{b}}^{\dot{\beta}}$, the operators $\hat{S}_{\mu\nu}$ and $\hat{S}_{\underline{mn}}^R$ depend only on z and $\partial/\partial z$. Their explicit form is given in the Appendix. It is also convenient to use a three-dimensional notations: $\hat{S}^k = \frac{1}{2}\epsilon_{ijk}\hat{S}^{ij}$, $\hat{B}^k = \hat{S}^{0k}$. Generators from different sets commute with each other.

In consequence of the unimodularity of 2×2 matrices Z there exist invariant antisymmetric tensors $\varepsilon^{\alpha\beta} = -\varepsilon^{\beta\alpha}$, $\varepsilon^{\dot{\alpha}\dot{\beta}} = -\varepsilon^{\dot{\beta}\dot{\alpha}}$, $\varepsilon^{12} = \varepsilon^{\dot{1}\dot{2}} = 1$, $\varepsilon_{12} = \varepsilon_{\dot{1}\dot{2}} = -1$. Now spinor indices are lowered and raised according to the rules

$$z_\alpha = \varepsilon_{\alpha\beta} z^\beta, \quad z^\alpha = \varepsilon^{\alpha\beta} z_\beta. \quad (8)$$

Discrete transformations correspond to the automorphisms of the Poincaré group, their action is reduced to replacing arguments and complex conjugation of functions $f(x, z)$ [2, 3].

Classification of functions on the group is carried out using maximal sets of commuting operators. A maximal set of commuting operators in the space of the functions on the group contains Casimir operators and an equal number operators that are some functions of left and right generators [24]. In total, in our case, the set contains 10 operators, according to the number of group parameters.

Casimir operators, that label irreps, can be composed of both left and right generators, so that the “left” and “right” mass and spin are the same. For the Poincaré group $M(3, 1)$,

we have:

$$\hat{p}^2 = \eta^{\mu\nu} \hat{p}_\mu \hat{p}_\nu = \eta^{\underline{mn}} \hat{p}_{\underline{m}}^R \hat{p}_{\underline{n}}^R, \quad (9)$$

$$\hat{W}^2 = \eta^{\mu\nu} \hat{W}_\mu \hat{W}_\nu = \eta^{\underline{mn}} \hat{W}_{\underline{m}}^R \hat{W}_{\underline{n}}^R, \quad (10)$$

$$\text{where } \hat{W}_\mu = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \hat{p}^\nu \hat{J}^{\rho\sigma} = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \hat{p}^\nu \hat{S}^{\rho\sigma}, \quad \hat{W}_{\underline{m}}^R = \frac{1}{2} \epsilon_{\underline{mnr}s} \hat{p}_{\underline{n}}^R \hat{S}_{\underline{r}s}^R. \quad (11)$$

As four operators, composed of left generators (generators of the group $M(3, 1)_{\text{ext}}$), one can choose the Casimir operator $\hat{\mathbf{p}}\hat{\mathbf{J}} = \hat{\mathbf{p}}\hat{\mathbf{S}}$ and the generators \hat{p}_k of the subgroup $M(3)_{\text{ext}}$. The latter correspond to additive quantum numbers. Eigenfunctions of these operators correspond to definite values of the helicity and the momentum.

Functions of right generators can be chosen in different ways. We choose sets, corresponding to reduction $M(3, 1)_{\text{int}} \supset SL(2, C)_{\text{int}}$. Two Casimir operators of the ‘‘right’’ Lorentz group $SL(2, C)_{\text{int}}$ have the form

$$\hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2 = \frac{1}{2} \hat{S}_{\underline{mn}}^R \hat{S}_{\underline{R}}^{\underline{mn}} = \frac{1}{2} \hat{S}_{\mu\nu} \hat{S}^{\mu\nu}, \quad \hat{\mathbf{S}}\hat{\mathbf{B}} = \frac{1}{16} \epsilon^{\underline{mnr}s} \hat{S}_{\underline{mn}}^R \hat{S}_{\underline{r}s}^R = \frac{1}{16} \epsilon^{\mu\nu\rho\sigma} \hat{S}_{\mu\nu} \hat{S}_{\rho\sigma}. \quad (12)$$

where \hat{S}^i and \hat{B}^i are operators of spin and boost projections, see Appendix. In contrast to $\hat{S}_{\underline{mn}}^R$, being the generators of the group $M(3, 1)_{\text{int}}$, the operators of spin projections $\hat{S}_{\mu\nu}$ are not generators of the group $M(3, 1)_{\text{ext}}$, see (7) and (6), and therefore operators (12) are functions of right (but not left) generators of the Poincaré group.

For $SL(2, C)_{\text{int}}$ we use two sets of commuting operators corresponding to the reduction schemes

$$\begin{aligned} SL(2, C)_{\text{int}} &\supset U(1) \times U(1), \\ SL(2, C)_{\text{int}} &\supset SU(2) \supset U(1). \end{aligned}$$

In the first case two generators \hat{S}_R^3 and \hat{B}_R^3 of the maximal commutative (Cartan) subgroup of $SL(2, C)_{\text{int}}$ correspond to additive quantum numbers. In the second case the set includes \hat{S}_R^3 and Casimir operator $\hat{\mathbf{S}}_R^2$ of $SU(2)_{\text{int}}$ with eigenvalues $S_R(S_R + 1)$.

Therefore, we will consider two sets of 10 commuting operators on the group $M(3, 1)$:

$$\hat{W}^2, \hat{p}_\mu, \hat{\mathbf{p}}\hat{\mathbf{S}} (\hat{S}^3 \text{ in the rest frame}), \hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2, \hat{\mathbf{S}}\hat{\mathbf{B}}, \hat{S}_R^3, \hat{B}_R^3, \quad (13)$$

$$\hat{W}^2, \hat{p}_\mu, \hat{\mathbf{p}}\hat{\mathbf{S}} (\hat{S}^3 \text{ in the rest frame}), \hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2, \hat{\mathbf{S}}\hat{\mathbf{B}}, \hat{\mathbf{S}}_R^2, \hat{S}_R^3, \quad (14)$$

including the Lubanski–Pauli operator \hat{W}^2 , four left generators \hat{p}_μ (the eigenvalue of the Casimir operator \hat{p}^2 , is evidently expressed through their eigenvalues), and helicity $\hat{\mathbf{p}}\hat{\mathbf{S}}$, expressed through the left generators. The Casimir operators $\hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2$ and $\hat{\mathbf{S}}\hat{\mathbf{B}}$ of the subgroup $SL(2, C)_{\text{int}}$ determine characteristics j_1, j_2 of the irrep $T_{[j_1 j_2]}$ of the Lorentz group (see Appendix).

2.1 Case of non-zero mass

As is known, particles with non-zero mass are characterized by intrinsic parity η – eigenvalue of the space reflection operator P (which defines the outer automorphism of the Poincaré group).

The action of space reflection P on orientation variables is given by the formula

$$P : Z \rightarrow (Z^\dagger)^{-1}, \text{ or } \begin{pmatrix} z^1 & \underline{z}^1 \\ z^2 & \underline{z}^2 \end{pmatrix} \rightarrow \begin{pmatrix} * & * \\ -\underline{z}_1 & z_1 \\ * & * \\ -\underline{z}_2 & z_2 \end{pmatrix}. \quad (15)$$

The operator of the space reflection \hat{P} anticommutes with the operators $\hat{\mathbf{S}}\hat{\mathbf{B}}$ and \hat{B}_R^3 . Therefore, eigenfunctions of the latter three operators change their sign under the action of \hat{P} . The set (14) is more convenient to describe states of definite internal parity, because only one operator $\hat{\mathbf{S}}\hat{\mathbf{B}}$ from this set doesn't commute with \hat{P} . Eigenvalues of $\hat{\mathbf{S}}\hat{\mathbf{B}}$ are proportional to $(j_1 - j_2)(j_1 + j_2 + 1)$, see (63). Thus, one can use quantum numbers, corresponding to the set (14), to characterize eigenstates of \hat{P} , with only one change (replacement the sign of $j_1 - j_2$ by intrinsic parity η).

Taking into account the above said, for the massive fields on the Poincaré group, we will consider the following set of quantum numbers:

$$p_\mu, m^2 = p_\mu p^\mu \neq 0, s, \eta, \mathbf{p}\hat{\mathbf{S}} (S^3 \text{ in the rest frame}), j_1 + j_2, |j_1 - j_2|, S_R, S_R^3. \quad (16)$$

Along with the eigenvalues of the two Casimir operators, intrinsic parity η labels irreps of the improper Poincaré group.

2.2 Case of zero mass

The massless case corresponds to the zero eigenvalue of the Casimir operator, $\hat{p}^2 f(x, z) = 0$. The small group is a three-parameter group of motions of a two-dimensional Euclidean space $E(2)$. This is a non-compact group, its unitary irreps are either infinite-dimensional (the so-called representations with continuous spin) or one-dimensional. In the latter case for representations of zero mass (classified by irreps of $SO(2) \subset E(2)$) we have [25, 26]

$$\hat{W}_\mu f(x, z) = \lambda \hat{p}_\mu f(x, z). \quad (17)$$

In this case, irreps of the Poincaré group are labelled by the chirality $\lambda = j_1 - j_2$ [24]. Taking into account the identity $\hat{W}\hat{p} \equiv 0$, we obtain for square of Lubanski-Pauli operator

$$\hat{W}^2 f(x, z) = (\hat{W}^2 + \lambda^2 \hat{p}^2) f(x, z) = (\hat{W} - \lambda \hat{p})^2 f(x, z) = 0.$$

Let's write out the first equation of the system (17) for the component $\hat{W}_0 = \frac{1}{2} \varepsilon_{0\nu\rho\sigma} \hat{p}^\nu \hat{S}^{\rho\sigma} = \mathbf{p}\hat{\mathbf{S}}$:

$$\mathbf{p}\hat{\mathbf{S}} f(x, z) = (j_1 - j_2) \hat{p}_0 f(x, z). \quad (18)$$

This equation establishes the equality of helicity and chirality, which significantly limits the number of possible states of massless particles.

For spatial components $\hat{W}^k = \hat{p}^0 \hat{S}^k - \varepsilon^{kli} \hat{p}^i \hat{B}^l$ we get

$$(\hat{p}^0 \hat{S}^k - \varepsilon^{kji} \hat{p}^i \hat{B}^j) f(x, z) = (j_1 - j_2) \hat{p}^k f(x, z). \quad (19)$$

In the case of representations $T_{[j_0]}$ and $T_{[0j]}$, the equations (19) can be simplified by excluding boost operators \hat{B}^k . Consider the operators $\hat{M}^k = \frac{1}{2}(\hat{S}^k - i\hat{B}^k)$, $\hat{M}^k = -\frac{1}{2}(\hat{S}^k + i\hat{B}^k)$, see (62). We have $\hat{M}^k f(x, z) = \frac{1}{2}(\hat{S}^k - i\hat{B}^k) f(x, z) = 0$ for $T_{[0j]}$ and $\hat{M}^k f(x, z) = -\frac{1}{2}(\hat{S}^k + i\hat{B}^k) f(x, z) = 0$ for $T_{[j_0]}$, see (62), hence $\hat{B}^k = i \text{sign} \lambda \hat{S}^k$,

$$\left(\hat{p}^0 \hat{S}^k - i \text{sign}(j_1 - j_2) \varepsilon_{kji} \hat{p}^i \hat{S}^j + (j_1 - j_2) \hat{p}^k \right) f(x, z) = 0. \quad (20)$$

Using (19), one can exclude \hat{p}^0 ,

$$\left((j_1 - j_2)^{-1} \hat{p}^i \hat{S}^i \hat{S}^k - i \text{sign}(j_1 - j_2) \varepsilon_{kji} \hat{p}^i \hat{S}^j + (j_1 - j_2) \hat{p}^k \right) f(x, z) = 0. \quad (21)$$

If the spin for a massive particle marks irreps of the Poincaré group and is determined through the eigenvalue of the Casimir operator \hat{W}^2 , then for massless particles irreps

are marked by the chirality, and the spin in this case is often defined as the modulus of chirality. However, spin 1 particles can have chirality 1 or 0 in the massless limit; in this sense the identification of spin with the modulus of chirality is not always correct.

For massless particles we arrive at a set of operators

$$\hat{p}^2 f = 0, \hat{W} = \lambda \hat{p} (\hat{\mathbf{p}} \hat{\mathbf{S}} f = \lambda \hat{p}_0 f), \hat{p}_\mu, \hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2, \hat{\mathbf{S}} \hat{\mathbf{B}}, \hat{B}_R^3, \hat{S}_R^3. \quad (22)$$

The eigenvalues of the operators \hat{p}^2 and \hat{W}^2 are fixed and equal to 0, the helicity (function of the left generators) is equal to the chirality (function of the right generators) and labels the irreps of the Poincaré group.

Thus, in contrast to the massive case, instead of 10 we have only 7 different quantum numbers – helicity (equal to the chirality $j_1 - j_2$), three functions of the left generators (momentum components), three functions of the right generators ($j_1 + j_2$ and two internal quantum numbers B_R^3, S_R^3 or S_R, S_R^3).

In contrast to the massive case, described by the Dirac and Duffin-Kemmer equations, associated with operators that do not commute with some generators of $M(3, 1)_{int}$ (right generators), and, therefore, associated with broken inner symmetries, the massless case does not a priori imply such a violation.

The case $m^2 = 0$ includes also functions on the group $f(z)$ that depend only on the orientation variables and do not depend on the spatial variables x^μ . They are classified according to the irreps of the Lorentz group (the maximal set in this case includes 6 operators – two Casimir operators $\hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2$ and $\hat{\mathbf{S}} \hat{\mathbf{B}}$, the left operators $\hat{\mathbf{S}}^2$ and \hat{S}^3 , the right operators $\hat{\mathbf{S}}_R^2$ and \hat{S}_R^3). However, these states cannot be localized in any region of the space, and, therefore, they cannot be interpreted as particle states, so we will not consider them here.

The case $m^2 < 0$ corresponds to tachyons, the small group $SU(1, 1)$ is non-compact, its unitary irreps are either one-dimensional or infinite-dimensional. As in the case of zero mass, spin-finite-dimensional representations are classified according to the irreps of the group $SO(2) \subset SU(1, 1)$.

3 S_R^3 charge

The physical meaning of the left generators included in the maximal set of operators is well known; two of the four functions of the right generators define the irrep of the Lorentz group $SL(2, C)_{int}$. Let's try to understand the physical meaning of the remaining two operators $\hat{\mathbf{S}}_R^2$ and \hat{S}_R^3 .

One can attempt to associate the charge corresponding to the right generator \hat{S}_R^3 of the Poincaré group to observable characteristics of physical particles. Notice that although the internal quantum numbers corresponding to the right generators do not change under the left transformations, the discrete transformations (automorphisms of the Poincaré group) act on both right and left generators (see [3] for details). The known behavior under the discrete transformations may help one to identify right characteristics with properties of physical particles.

In this regard, let's list the properties of the operator \hat{S}_R^3 :

1. Its quantum number is additive.
2. It has integer eigenvalues for particles of integer spin and half-integer eigenvalues for particles of half-integer spin.
3. It does not change the sign under the space reflection.
4. It changes the sign under the charge conjugation.

Then it follows from item 4 that this number equals to zero for real neutral particle (i.e., particle that coincides its own antiparticle); at the same time, real neutral particles with definite S_R^3 must to have an integer spin.

If one considers S_R^3 -charge of particles described by finite-dimensional representations of the Lorentz group $T_{[j_1, j_2]}$, $j_1 + j_2 = s$, then for particles of spin one-half two values are possible: $1/2$ and $-1/2$; for particles of spin one three values are possible: $1, 0$ and -1 . In particular, for a photon and Z^0 -boson, as for real neutral particles, we have $S_R^3 = 0$.

Let us see which eigenvalues S_R^3 can be associated with particles described by eigenfunctions $f(x, z)$ of the operator \hat{S}_R^3 ,

$$\hat{S}_R^3 f(x, z) = S_R^3 f(x, z) .$$

Since the sign of \hat{S}_R^3 changes under the charge conjugation, particles and antiparticles must have S_R^3 -charges of opposite signs. For definiteness, let an electron e^- has the S_R^3 -charge $-\frac{1}{2}$; then a positron e^+ has $S_R^3 = \frac{1}{2}$. For W^- , as a charged particle, it is natural to expect $S_R^3 = \pm 1$. Next, since $\tilde{\nu}_e$ admits only the values $\pm\frac{1}{2}$, the reaction $W^- \rightarrow e^- + \tilde{\nu}_e$ implies $S_R^3 = -\frac{1}{2}$ for $\tilde{\nu}_e$ and $S_R^3 = -1$ for W^- . Therefore, we have

$$\begin{aligned} S_R^3 = \frac{1}{2} : & \quad \nu_e, e^+; & S_R^3 = 1 : & \quad W^+; \\ S_R^3 = -\frac{1}{2} : & \quad e^-, \tilde{\nu}_e; & S_R^3 = -1 : & \quad W^- \quad S_R^3 = 0 : \gamma, Z^0, H; \end{aligned} \quad (23)$$

Applying similar reasons to other families of fundamental fermions, we find the following classification with respect to the sign of the quantum number S_R^3 :

$$\begin{aligned} S_R^3 = \frac{1}{2} : & \quad \nu_e \quad \nu_\mu \quad \nu_\tau \quad u \quad c \quad t \\ S_R^3 = -\frac{1}{2} : & \quad e^- \quad \mu \quad \tau \quad d \quad s \quad b \end{aligned} \quad (24)$$

Therefore, the S_R^3 -charge, whose sign changes under the charge conjugation C , distinguishes not only particles and antiparticles but also the ‘‘up-down’’ components in doublets of elementary fermions. This charge is conserved in any interactions, since the carriers of electromagnetic and strong interactions are characterized by $S_R^3 = 0$, whereas we have already considered weak interaction case.

As a consequence of Eqs. (23) and (24), we may write the following relation between S_R^3 and some other charges:

$$S_R^3 = \frac{L - B}{2} + Q, \quad (25)$$

where L, B , and Q are the lepton, baryon and electric charges, respectively. Equation (25) relates the ‘‘right’’ charge S_R^3 with observable characteristics of the particles.

For the above-mentioned fundamental particles of spin one, the charge S_R^3 coincides with the electric charge. Taking into account that $(B - L)/2 = \bar{Q}$, where \bar{Q} – average electric charge of corresponding lepton (ν, e^-) or quark (d, u) doublet one can rewrite Eq. (25) as $S_R^3 = Q - \bar{Q}$.

Formula (25) is closely related to the concept of R -parity. In 1961 Michel and Lurçat [27] observed that for all the known particles with integer B there holds the relation $(-1)^{B+L} = (-1)^{2S}$, in other words, the number $B + L + 2S$ is always even. Later, this observation resulted in the concept of R -parity, being positive for all the known particles, and defined as

$$R = (-1)^{3(B-L)+2S}, \quad (26)$$

where the multiplier 3 is introduced to include quarks in the consideration.

We have already noted that the spectra of right and left spin operators are not independent, in particular (see item 2 above), S_R^3 -charge and spin S must both be either integers or half-integers,

$$(-1)^{2S_R^3} = (-1)^{2S}. \quad (27)$$

Due to (25) we have $(-1)^{L-B+2Q} = (-1)^{2S}$, therefore, the value

$$R_1 = (-1)^{2S_R^3+2S} = (-1)^{L-B+2Q+2S} \quad (28)$$

is always positive. For integer Q we have $R_1 = R$, for fractional Q (quarks), it is not difficult to verify that $(-1)^{L-B+2Q} = (-1)^{3(B-L)}$. We see, that for all known particles $R_1 = R$.

Thus, consideration of the field on the Poincaré group together with the empirical formula (25) leads to the connection of spin with internal quantum numbers, expressed in the concept of R -parity.

We also note that S_R^3 coincides with the projection of the weak isospin T_3 for left components of particles and right of antiparticles. In fact, in the massless limit, the formula (25) turns into the well-known relation $T_3 = -Y/2 + Q$, where Y is a weak hypercharge. The possible connection of $SU(2)_{int}$ with the weak gauge group is considered in Ref. [5].

In the next section we describe functions $f(x, z)$, polynomial in z . Considering polynomials of fixed degree n in z , we will actually consider the irreps of the Lorentz group $T_{[j_1 j_2]}$ with $j_1 + j_2 = n/2$. According to the reduction formula (64), the values j_1 and j_2 define a possible range of spin values S , S_R and projections S^3 and S_R^3 .

Irreps of $SL(2, C)_{int}$	S, S_R	projections S^3, S_R^3
$T_{[1/2 0]}, T_{[0 1/2]}$	1/2	$\pm 1/2$
$T_{[1 0]}, T_{[0 1]}$	1	$0, \pm 1$
$T_{[1/2 1/2]}$	0, 1	$0, \pm 1$
$T_{[3/2 0]}, T_{[0 3/2]}$	3/2	$\pm 1/2; \pm 3/2$
$T_{[1 1/2]}, T_{[1/2 1]}$	1/2, 3/2	$\pm 1/2; \pm 3/2$

(29)

4 Massless case

4.1 $j_1 + j_2 = 0$, chirality 0


The case of ordinary scalar functions $f(x) = e^{-ipx^0 + ipx^3}$, which do not depend on orientation variables z . Corresponds to a scalar particle.

4.2 $j_1 + j_2 = 1/2$, chirality $\pm 1/2$

Let us consider functions linear in z . Irreps enter into the decomposition of the left (right) regular representation with a multiplicity equal to the dimension of the representation; see [24]. This can be clearly illustrated on the Fig. 1. It shows weight diagrams of the representations $T_{[1/2 0]}$ (chirality $\frac{1}{2}$) and $T_{[0 1/2]}$ (chirality $-\frac{1}{2}$) of the dimension 2. When acting from the left, the spinors (z_1, z_2) and $(\underline{z}_1, \underline{z}_2)$ are transformed identically and correspond to equivalent representations; when acting from the right they are transformed identically (z_1, \underline{z}_1) and (z_2, \underline{z}_2) . If we consider simultaneously the action on the left and the right (external and internal transformations), then all four states are mixed.


In total we have 8 states (orientation variables $z^\alpha, \underline{z}^\alpha, \overset{*}{z}^\alpha, \overset{*}{\underline{z}}^\alpha$), corresponding to four different particles (in other words, states that cannot be translated into each other by transformations of the laboratory reference system). If states inside quadruplets are connected

Figure 1: The weight diagrams of the representations $T_{[1/2\ 0]}$ and $T_{[0\ 1/2]}$ of $SL(2, C)_{\text{ext}} \times SL(2, C)_{\text{int}}$. Vertical lines join states related by transformations of $SL(2, C)_{\text{ext}}$, horizontal lines join states related by transformations of $SL(2, C)_{\text{int}}$.



by left or right transformations of the Poincaré group, then quadruplets are connected to each other by discrete transformations (space reflection, complex (charge) conjugation), which are external automorphisms of the group. Particle-antiparticle pairs correspond to complex conjugate pairs $z^\alpha, z^{\alpha*}$ and $\underline{z}^\alpha, \underline{z}^{\alpha*}$; inside the pairs the signs of the chirality $j_1 - j_2$ and the charge S_R^3 are opposite.

Figure 2: The weight diagram of the representation $T_{[1/2\ 0]} \oplus T_{[0\ 1/2]}$ of $SL(2, C)_{\text{int}}$. The dotted line joins states related by transformations of $SL(2, C)_{\text{int}}$. a) States with definite chirality, functions (30). b) States with definite parity η , functions (41).



The decomposition of the representation of the Poincaré group in the space of linear in z functions $f(x, z)$ is carried out using a set of operators (22). Consider the equation $\hat{\mathbf{p}}\hat{\mathbf{S}}f(x, z) = (j_1 - j_2)\hat{p}_0f(x, z)$, establishing the equality of the chirality and the helicity, from the set. For a massless particle with a certain momentum $p^\mu = (p^0, 0, 0, p^3)$, $p^3 = \pm p^0$, we have $p^3 S^3 = (j_1 - j_2)p^0$; for a positive p^0 signs of p^3 and S^3 must coincide for particles, and be opposite for antiparticles (positive and negative helicity, right-handed particles and left-handed antiparticles). Thus, the classification of functions on the Poincaré group gives two left-handed particles and two right-handed antiparticles, each having one chiral state,

$$\begin{aligned} R (j_1 - j_2 = 1/2) : & \quad e^{-ipx^0 + ipx^3} z^1 & \quad e^{-ipx^0 + ipx^3} \underline{z}^1 \\ L (j_1 - j_2 = -1/2) : & \quad e^{-ipx^0 - ipx^3} \underline{z}^1 & \quad e^{-ipx^0 - ipx^3} z^1 \end{aligned} \quad (30)$$

These two particle/antiparticle pairs differ in signs of the charge S_R^3 , states with the same chirality are connected by a group transformation $SU(2)_{\text{int}}$, see Fig. 2a.

We see that in the space of the functions on the Poincaré group, particles of spin one half appear not in pairs, but in quadruplets.


4.3 $j_1 + j_2 = 1$, chirality ± 1

Let us consider functions on the group that are quadratic in z .

From the variables z, \underline{z} as well as from z^*, \underline{z}^* , 10 different quadratic combinations can be constructed. 9 of them correspond to the irreps $T_{[1\ 0]}$ and $T_{[0\ 1]}$ (chirality $j_1 - j_2 = \pm 1$), two combinations are invariants, since, due to unimodularity, $\det Z = z^1 \underline{z}^2 - z^2 \underline{z}^1 = 1$ is a Lorentz scalar.

The representations $T_{[1\ 0]}$ and $T_{[0\ 1]}$ correspond to figures similar to the one Fig. 1, but each of them has not 4, but 9 states.

Figure 3: The weight diagrams of the representation $T_{[1\ 0]} \oplus T_{[0\ 1]}$ of $SL(2, C)_{int}$, $j_1 - j_2 = \pm 1$ (a), $\eta = \pm 1$ (b). The dotted lines join states related by the transformations $SL(2, C)_{int}$.



By restoring the laboratory indices, one can easily observe that each point of the weight diagram (Fig.3) corresponds to three states (according to the number of possible spin projections), that transform under 3-dimensional irreps of $SL(2, C)_{ext}$. In particular, for states with $S_R^3 = 0$ among four pairwise products 3 functions correspond to the chirality one, namely $z^1 \underline{z}^1$, $z^2 \underline{z}^2$, and $z^1 \underline{z}^2 + z^2 \underline{z}^1$.

Making a reduction to the compact group $SU(2)_{int}$, we also obtain two triplets: left and right, corresponding to the diagonals on Fig.3a, or triplets with a fixed parity, Fig.3b.

For a massless particle with a certain momentum $p^\mu = (p^0, 0, 0, p^3)$, $p^3 = \pm p^0$, we have $p^3 S^3 = (j_1 - j_2) p^0$; for a positive p^0 signs of p^3 and S^3 must coincide for particles, and be opposite for antiparticles (positive and negative helicity, right-handed particles and left-handed antiparticles). Thus, we have 6 states (particles) $W_L^\pm, W_L^0, W_R^\pm, W_R^0$, corresponding to three values of S_R^3 and two values of the chirality; for the states with $p^\mu = (p, 0, 0, p)$ we have

$$\begin{array}{l}
 R (j_1 - j_2 = 1) : \\
 L (j_1 - j_2 = -1) :
 \end{array}
 \begin{array}{ccc}
 S_R^3 = -1 & S_R^3 = 0 & S_R^3 = 1 \\
 e^{-ipx^0 + ipx^3} z^1 z^1 & e^{-ipx^0 + ipx^3} z^1 \underline{z}^1 & e^{-ipx^0 + ipx^3} \underline{z}^1 \underline{z}^1 \\
 e^{-ipx^0 - ipx^3} z_i^* z_i^* & e^{-ipx^0 - ipx^3} z_i^* \underline{z}_i^* & e^{-ipx^0 - ipx^3} \underline{z}_i^* \underline{z}_i^*
 \end{array}
 \quad (31)$$

By direct substitution, it is not difficult to make sure that the states (31) satisfies not only (18), but also (20).

In this case, particles with the same S_R^3 differ only by signs of the chirality/helicity and can be considered as different polarization states of the same particle.

4.4 $j_1 + j_2 = 1$, chirality 0

Let us now consider the remaining quadratic combinations of z , which are products of z, \underline{z} to the complex conjugates z^*, \underline{z}^* . In total, there are 16 such combinations and they belong to the $T_{[\frac{1}{2}, \frac{1}{2}]}$ representation of the Lorentz group. For this representation, all the states are characterized by the chirality $j_1 - j_2 = 0$. During a reduction to the compact group $SU(2)_{int}$ (or $SU(2)_{ext}$) we obtain a triplet and a singlet.

Each point on the Fig. 4a corresponds to 4 states, namely 3 vector components and a scalar with respect to the rotation group $SU(2)_{ext}$. States with a fixed charge S_R^3 can be distinguished either by B_R^3 , or by S_R and the parity η .

For the states with a certain momentum $p^\mu = (p^0, 0, 0, p^3)$, $p^3 = \pm p^0$, the equation (18) gives $\hat{p}^3 \hat{S}^3 f(x, z) = 0$. Thus, the only one components remains with $S^3 = 0$,

$$f(x, z) = e^{-ipx^0 \pm ipx^3} \varphi(z), \quad \hat{S}^3 \varphi(z) = 0.$$

These states must also satisfy the equations (19) for spatial components W_i , $W^i f(x, z) = p^0 \hat{S}^i - \varepsilon_{ijk} p^j \hat{B}^k f(x, z) = 0$. The third equation is fulfilled identically, the first two equations give $p^3 = -p^0$ for functions of the form $z^1 z^1$ and $p^3 = p^0$ for functions of the form $z^2 z^2$. Since these functions are connected by rotation from $SU(2)_{ext}$, then we have the following 4 states corresponding to 4 particles,

$$\begin{array}{ccc} S_R^3 = -1 & S_R^3 = 0 & S_R^3 = 1 \\ e^{-ipx^0 - ipx^3} z^1 \underline{z}^1 & e^{-ipx^0 - ipx^3} \underline{z}^1 \underline{z}^1 & e^{-ipx^0 - ipx^3} \underline{z}^1 z^1 \\ & e^{-ipx^0 - ipx^3} z^1 \underline{z}^1 & \end{array} \quad (32)$$

The two states with $S_R^3 = 0$ differ in the values of B_R^3 .

Let us consider scalars with respect to the group $SU(2)_{ext}$:

$$H^- : z^1 \underline{z}^1 + z^2 \underline{z}^2, \quad H^+ : \underline{z}^1 z^1 + \underline{z}^2 z^2, \quad (33)$$

$$H_-^0 : z^1 \underline{z}^1 + z^2 \underline{z}^2, \quad H_+^0 : \underline{z}^1 z^1 + \underline{z}^2 z^2, \quad (34)$$

– eigenfunctions of spin projection operators \hat{S}_k with zero eigenvalues. Their linear combinations are the triplet H^-, H^0, H^+ ($S_R = 1, \eta = 1$) and the singlet H_0^0 ($S_R = 0, \eta = -1$) with respect to $SU(2)_{int}$,

$$H_0^0 : z^1 \underline{z}^1 + z^2 \underline{z}^2 - \underline{z}^1 z^1 - \underline{z}^2 z^2, \quad H_0^0 : z^1 \underline{z}^1 + z^2 \underline{z}^2 + \underline{z}^1 z^1 + \underline{z}^2 z^2, \quad (35)$$

see Fig.4b. All this functions satisfy the equation (18) identically, for them both spin projections and chirality are equal to zero. However, the set of these states is not invariant with respect to the boost operators contained in the equations (19), which convert these states into vectors with respect to $SU(2)_{ext}$.

The results of the consideration of massless case can be presented in the form of a Table 1.

For fixed j_1 and j_2 , the number of particles (states not connected by transformations $SL(2, C)_{ext}$) is given by the dimension of the irrep of the $SU(2)_{int}$ group. Each of the listed massless particles can be in one chiral state, for which the projection of the spin onto the direction of motion coincides with $j_1 - j_2$.

For the case $j_1 + j_2 > 1$ the consideration scheme is similar. For example, let us consider polynomials of degree 3 in z , namely, $j_1 + j_2 = 3/2$, and chirality is $\pm 1/2$ and $\pm 3/2$.

Figure 4: The weight diagrams of the representation $T_{[\frac{1}{2} \frac{1}{2}]}$ of $SL(2, C)_{int}$, chirality $j_1 - j_2 = 0$. On Fig.a the dotted line joins states related by transformations of $SL(2, C)_{int}$, on Fig.b the dotted line joins states related by transformations of $SU(2) \in SL(2, C)_{int}$.




Table 1: Classification of scalar functions and its interpretation, massless case.

chirality $j_1 - j_2$	$j_1 j_2$	irreps of $SU(2)_{int} \times SU(2)_{ext}$ and its dimension	dimension of $SU(2)_{int}$ irreps	particles
$\frac{1}{2}$	$\frac{1}{2} 0$	$T_{\frac{1}{2}} \times T_{\frac{1}{2}}$ 4	2	$e_R^+, \bar{\nu}_R$
$-\frac{1}{2}$	$0 \frac{1}{2}$	$T_{\frac{1}{2}} \times T_{\frac{1}{2}}$ 4	2	e_L^-, ν_L
1	1 0	$T_1 \times T_1$ 9	3	W_R^+, W_R^-, W_R^0
-1	0 1	$T_1 \times T_1$ 9	3	W_L^+, W_L^-, W_L^0
0	$\frac{1}{2} \frac{1}{2}$	$(T_1 + T_0) \times T_1$ 12 $(T_1 + T_0) \times T_0$ 4	3 1	HL^+, HL^0, HL^- HL_0^0

4.5 $j_1 + j_2 = 3/2$, chirality $\pm 3/2$

The representations $T_{[\frac{3}{2} 0]}$ and $T_{[0 \frac{3}{2}]}$ of the Lorentz group have the dimension 4 and are associated with 16 monomials of the third degree in z, \underline{z} . In total, there are 8 states (particles) not connected by transformations $SL(2, C)_{int}$, differing by the values S_R^3 ($-3/2, -1/2, 1/2, 3/2$) and by the chirality sign ($-3/2, 3/2$), corresponding to 8 points on the Fig. 5a. Each point of the weight diagram (Fig. 5) corresponds to four states (according to the number of possible spin projections).

The equation (18) for a fixed momentum directed along the x^3 axis has the form $p_3 \hat{S}^3 f(x, z) = (j_1 - j_2) p_0 f(x, z)$; accordingly, for a fixed chirality and the charge S_R^3 , one component remains, characterized by $S^3 = j_1 - j_2$. These states can be interpreted as 8 particles or as 4 particles in 2 polarization states.

4.6 $j_1 + j_2 = 3/2$, chirality $\pm 1/2$

The representations $T_{[1 \frac{1}{2}]}$ and $T_{[\frac{1}{2} 1]}$ of the Lorentz group have the dimension 6 and each of them corresponds to 36 monomials of third degree. Being reduced to a compact subgroup $SU(2)$, they split into 4-dimensional $T_{3/2}$ and 2-dimensional $T_{1/2}$ representations. These states can be interpreted as 24 particles or 12 particles in two polarization states.





Figure 5: The weight diagram of the representation $T_{[3/2 0]} \oplus T_{[0 3/2]}$ of $SL(2, C)_{\text{int}}$. The dotted line joins states related by transformations of $SL(2, C)_{\text{int}}$. a) States with definite chirality $j_1 - j_2 = \pm 3/2$. b) States with definite parity η .

Figure 6: The weight diagram of the representation $T_{[1 1/2]} \oplus T_{[1/2 1]}$ of $SL(2, C)_{\text{int}}$. The dotted line joins states related by transformations of $SL(2, C)_{\text{int}}$. a) States with definite chirality $j_1 - j_2 = \pm 1/2$. b) States with definite parity η .



5 Relativistic wave equations for massless particles

The operators \hat{W}_μ and \hat{S}_k in equations (17)–(20), acting in the space of functions $f(x, z)$ on the group, are differential in z and for all representations have the same form. By writing $f(x, z)$ as polynomials of z with coefficients $\psi_n(x)$ and comparing the coefficients at powers of z , we can write equations (17)–(20) as matrix equations for $\psi_n(x)$, and their form will be different in different representations.

1. $j_1 + j_2 = 1/2$, the chirality $\pm 1/2$, $T_{[1/2 0]}$ and $T_{[0 1/2]}$. Let us write down the functions corresponding to $S_R^3 = 1/2$:

$$\begin{aligned} f(x, \underline{z}) &= \psi_1(x) \underline{z}^1 + \psi_2(x) \underline{z}^2, & \psi_R(x) &= (\psi_1(x) \ \psi_2(x))^T, \\ f(x, \underline{z}^*) &= \psi^1(x) \underline{z}_1^* + \psi^2(x) \underline{z}_2^*, & \psi_L(x) &= (\psi^1(x) \ \psi^2(x))^T \end{aligned}$$

Substituting these forms into Eq. (18) and comparing coefficients at \underline{z}_i and at \underline{z}_i^* , we obtain a pair of Weyl equations,

$$(\hat{p}_0 - \boldsymbol{\sigma} \hat{\mathbf{p}}) \psi_R = 0, \quad (\hat{p}_0 + \boldsymbol{\sigma} \hat{\mathbf{p}}) \psi_L = 0. \quad (36)$$

The same equations we obtain for the case $S_R^3 = -1/2$. For chiralities $\pm 1/2$ the equations (21) for spatial components of \hat{W}_μ will be fulfilled automatically if (18) is fulfilled. To see this, it is enough to note that in the case under consideration $\mathbf{S} = \frac{1}{2} \boldsymbol{\sigma}$.

2. $j_1 + j_2 = 1$, the chirality ± 1 , $T_{[10]}$ and $T_{[01]}$. The representations $T_{[10]}$ and $T_{[01]}$ of the Lorentz group are three-dimensional complex conjugate representations by the matrices $SO(3, C)$. For the complex functions $\psi_k = E_k + iH_k$ transforming under $T_{[10]}$, the combination $\sum \psi_k^2 = \sum (E_k + iH_k)^2 = \mathbf{E}^2 - \mathbf{H}^2 + 2i\mathbf{E}\mathbf{H}$ is conserved. Let us write functions $f(x, z, \underline{z})$ and $f(x, \underline{z}, \underline{z}^*)$, corresponding to $S_R^3 = 0$ and connected by space reflection:

$$\begin{aligned} f(x, z, \underline{z}) &= \psi_1(x)i(z^2\underline{z}^2 - z^1\underline{z}^1) + \psi_2(x)(-z^1\underline{z}^1 - z^2\underline{z}^2) + \psi_3(x)i(z^1\underline{z}^2 + z^2\underline{z}^1), \\ \xi &= \psi_R(x) = (\psi_1 \psi_2 \psi_3)^T; \\ f(x, \underline{z}, \underline{z}^*) &= \psi^1(x)i(\underline{z}_2^*\underline{z}_2^* - \underline{z}_1^*\underline{z}_1^*) + \psi^2(x)(-\underline{z}_1^*\underline{z}_1^* - \underline{z}_2^*\underline{z}_2^*) + \psi^3(x)i(\underline{z}_1^*\underline{z}_2^* + \underline{z}_2^*\underline{z}_1^*), \\ \eta &= \psi_L(x) = (\psi^1 \psi^2 \psi^3)^T. \end{aligned}$$

Substituting these forms into Eqs. (18) and (21) and comparing expressions at the orientation variables, we obtain Maxwell equations in the Majorana form:

$$i\frac{\partial \xi}{\partial t} = \hat{\mathbf{p}}\mathbf{S}\xi, \quad \hat{\mathbf{p}}\xi = 0, \quad i\frac{\partial \eta}{\partial t} = -\hat{\mathbf{p}}\mathbf{S}\eta, \quad \hat{\mathbf{p}}\eta = 0, \quad (37)$$

where $(S^i)_{kl} = \varepsilon_{ikl}$,

$$S^1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad S^2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad S^3 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The column $(\xi \ \eta)^T$ is considered as the photon wave function [28, 29].

3. $j_1 + j_2 = 1$, the chirality 0, $T_{[\frac{1}{2}\frac{1}{2}]}$. Let us write down the functions corresponding to $S_R^3 = 1/2$:

$$\begin{aligned} f(x, z) &= \psi_\mu(x)\sigma^\mu_{\alpha\dot{\beta}}\underline{z}^\alpha\underline{z}^{\dot{\beta}} \\ &= \psi_0(x)(\underline{z}^1\underline{z}^1 + \underline{z}^2\underline{z}^2) + \psi_1(x)(\underline{z}^1\underline{z}^2 + \underline{z}^2\underline{z}^1) + \psi_2(x)i(\underline{z}^2\underline{z}^1 - \underline{z}^1\underline{z}^2) + \psi_3(x)(\underline{z}^1\underline{z}^1 - \underline{z}^2\underline{z}^2). \end{aligned}$$

Substituting these forms into Eq. (18) and comparing coefficients at the orientation variables, we obtain

$$\hat{\mathbf{p}}\mathbf{S}\psi = 0, \quad \psi(x) = (\psi_1(x), \psi_2(x), \psi_3(x))^T, \quad (38)$$

where $(S^i)_{kl} = \varepsilon_{ikl}$, or $\nabla \times \psi = 0$ in vector notations. Functions $\psi_0(x)$, corresponding to scalar H^+ , are not included in the equation, because $\hat{S}_k(\underline{z}^1\underline{z}^1 + \underline{z}^2\underline{z}^2) = 0$. Equation (19) can be written in matrix form,

$$(\hat{p}^0\tilde{S}^i - \varepsilon_{ijk}\hat{p}^j B^k)\Psi(x) = 0, \quad \Psi(x) = (\psi_0(x), \psi(x))^T, \quad (39)$$

where matrices B^k have two nonzero elements $(B^k)_{0k} = (B^k)_{k0} = i$, and \tilde{S}^i are obtained from S^i by adding the first zero row and column. The latter equation is equivalent to the system of equation (38) and three equations $\frac{\partial \psi_k}{\partial t} - \frac{\partial \psi_0}{\partial x^k} = 0$.

6 Massive case

Here we consider eigenfunctions of the maximal set of 10 commuting operators (12). Since the representations $T_{[j_1 j_2]}$ and $T_{[j_2 j_1]}$ are connected by a space reflection, then, as was mentioned above, to construct states with a certain internal parity instead of chirality $j_1 - j_2$ one has to use the modulus $|j_1 - j_2|$ and the internal parity given by the set of numbers (16): the mass m , the spin S , the parity η , the momentum \mathbf{p} , the spin projection S_3 , parameters of irrep of the Lorentz group $j_1 + j_2$, $|j_1 - j_2|$, S_R and its projection S_R^3 .

We will consider the rest frame ($p_0 = m$, $p_k = 0$); in this frame, the operator \hat{W}^2 reads:

$$\hat{W}^2 f(x, z) = -m^2 S(S+1) f(x, z), \quad (40)$$

where $S(S+1)$ is an eigenvalue of operator $\hat{\mathbf{S}}^2$.

The irreps $T_{[j_0]}$ and $T_{[0j]}$ of the group $SL(2, C)$ when reduced to the compact subgroup $SU(2)$ remain irreducible T_j of the dimension $2j+1$. For such representations $S = j$. For other irreps of $SL(2, C)$, upon the reduction, the representation becomes reducible and, for a given j_1, j_2 , we obtain different values of the spin and irreps of the Poincaré group.

6.1 $j_1 + j_2 = 0$, spin 0

The case of ordinary scalar functions $f(x) = e^{-imx^0}$, which do not depend on orientation variables z . Corresponds to a scalar particle with $\eta = 1$.

6.2 $j_1 + j_2 = 1/2$, spin 1/2

Consider the representations $T_{[\frac{1}{2}0]}$ and $T_{[0\frac{1}{2}]}$. The spin $S = 1/2$, in the rest frame

$$\begin{array}{ccc} S_R^3 = -1/2 & S_R^3 = 1/2 & \\ \eta = 1 & e^{-imx^0} (z^\alpha - \underline{z}_{\dot{\alpha}}^*) & e^{-imx^0} (\underline{z}^\alpha + \underline{z}_{\dot{\alpha}}^*) \\ \eta = -1 & e^{-imx^0} (z^\alpha + \underline{z}_{\dot{\alpha}}^*) & e^{-imx^0} (\underline{z}^\alpha - \underline{z}_{\dot{\alpha}}^*) \end{array} \quad (41)$$

Each of 4 functions corresponds to a point on the Fig. 2b. Different signs of the spin projection $S^3 = \pm 1/2$ correspond to two possible values of the index α in Eq. (41).

There are 8 states, 4 particles; the particle states in the rest frame differ in two spin projection.

6.3 $j_1 + j_2 = 1$, spin 1

The representations $T_{[10]}$ and $T_{[01]}$ are connected by space reflection, $S = j_1 + j_2 = 1$. Each point in Fig. 3b corresponds to 3 states with different spin projections.

$$\begin{array}{ccc} S_R^3 = -1 & S_R^3 = 0 & S_R^3 = 1 \\ \eta = \pm 1 & e^{-imx^0} (z^\alpha z^\beta \pm \underline{z}_{\dot{\alpha}}^* \underline{z}_{\dot{\beta}}^*) & e^{-imx^0} (z^{(\alpha} \underline{z}^{\beta)} \mp \underline{z}_{(\dot{\beta}}^* \underline{z}_{\dot{\alpha}}^*)} & e^{-imx^0} (\underline{z}_\alpha \underline{z}_\beta \pm \underline{z}_{\dot{\alpha}}^* \underline{z}_{\dot{\beta}}^*) \end{array} \quad (42)$$

Here, the parentheses of the indexes indicate symmetrization.

There are 18 states, 3 particles; the particle states differ in parity $\eta = \pm 1$ and 3 spin projection values in the rest frame.

6.4 $j_1 + j_2 = 1$, spin 0 and 1

The representation $T_{[\frac{1}{2}\frac{1}{2}]}$, 16 states in total. The four-dimensional representation, when reduced to a compact subgroup, decomposes into a scalar ($S = 0$) and a vector ($S = 1$).

For $S = 0$, we have 4 particles, a singlet H_0^0 and a triplet H^-, H^0, H^+ with respect to $SU(2)_{int}$. In the rest frame

$$\begin{array}{ccc} S_R^3 = -1 & S_R^3 = 0 & S_R^3 = 1 \\ S^R = 1 & e^{-imx^0} (z^1 \underline{z}^{*1} + z^2 \underline{z}^{*2}) & e^{-imx^0} (z^1 \underline{z}^{*1} + z^2 \underline{z}^{*2} - \underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2}) & e^{-imx^0} (\underline{z}^1 \underline{z}^{*1} + \underline{z}^2 \underline{z}^{*2}) \\ S^R = 0 & & e^{-imx^0} (z^1 \underline{z}^{*1} + z^2 \underline{z}^{*2} + \underline{z}^1 \underline{z}^{*1} + \underline{z}^2 \underline{z}^{*2}) & \end{array} \quad (43)$$

Scalars represent 4-plet of the $SL(2, C)_{int}$. Being reduced to a compact subgroup $SU(2)_{int}$, they split into the triplet H^-, H^0, H^+ and the singlet H_0^0 (see Fig. 4b). In the triplet $S_R^3 = -1, 0, +1$, and, according to the empirical formula (25), the same values of electrical charge. With respect to the operator M_k algebra (see 62) the scalars form two doublets H^-, H_-^0 and H_+^0, H^+ (see Fig. 4a).

We note, that quantum numbers of the H_0^0 state (zero electric charge and the parity $\eta = 1$) coincide with those of the Higgs boson.

For $S = 1$, we also have 4 particles, a singlet L_0^0 and a triplet L^-, L^0, L^+ with respect to $SU(2)_{int}$.

$$\begin{array}{llll}
& S^3 = -1 & S^3 = 0 & S^3 = 1 \\
L^-, S_R^3 = -1 & e^{-imx^0} \underline{z}^1 \underline{z}^{*2} & e^{-imx^0} (\underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2}) & e^{-imx^0} \underline{z}^2 \underline{z}^{*1} \\
L^+, S_R^3 = 1 & e^{-imx^0} \underline{z}^1 \underline{z}^{*2} & e^{-imx^0} (\underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2}) & e^{-imx^0} \underline{z}^2 \underline{z}^{*1} \\
L^0, S_R^3 = 0 & e^{-imx^0} (\underline{z}^1 \underline{z}^{*2} - \underline{z}^1 \underline{z}^{*2}) & e^{-imx^0} (\underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2} + \underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2}) & e^{-imx^0} (\underline{z}^2 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*1}) \\
L_0^0, S_R^3 = 0 & e^{-imx^0} (\underline{z}^1 \underline{z}^{*2} + \underline{z}^1 \underline{z}^{*2}) & e^{-imx^0} (\underline{z}^1 \underline{z}^{*1} - \underline{z}^2 \underline{z}^{*2} - \underline{z}^1 \underline{z}^{*1} + \underline{z}^2 \underline{z}^{*2}) & e^{-imx^0} (\underline{z}^2 \underline{z}^{*1} + \underline{z}^2 \underline{z}^{*1})
\end{array} \tag{44}$$

The case $S_R^3 = 0$ (fields L^0, L_0^0) corresponds to a truly neutral hypothetical particle, which is called the notoph in the literature [30, 31, 32] (in this case, the corresponding potentials are tensors transforming according to the representations $T_{[10]}$ and $T_{[01]}$).

Thus, just 8 particles (states not connected by transformations $SL(2, C)_{ext}$) are associated with the representation $T_{\frac{1}{2}} \times T_{\frac{1}{2}}$.

The above can be summarized in the Table 2. Each of the listed particles can be in several spin states, their number coincides with the dimension of the $SU(2)_{ext}$ representation.

Table 2: Classification of functions and their interpretation, $j_1 + j_2 \leq 1$, $m \neq 0$.

Spin	η	$ j_1 - j_2 $	Irreps of $SL(2, C)_{int}$	Irreps of $SU(2)_{int} \times SU(2)_{ext}$ and its dimension	dimension of $SU(2)_{int}$ irreps	particles	
$\frac{1}{2}$	1	$\frac{1}{2}$	$T_{[\frac{1}{2} 0]}, T_{[0, \frac{1}{2}]}$	$T_{\frac{1}{2}} \times T_{\frac{1}{2}}$	4	2	e^-, ν
$\frac{1}{2}$	-1	$\frac{1}{2}$	$T_{[\frac{1}{2} 0]}, T_{[0, \frac{1}{2}]}$	$T_{\frac{1}{2}} \times T_{\frac{1}{2}}$	4	2	$e^+, \bar{\nu}$
1	± 1	1	$T_{[1 0]}, T_{[0, 1]}$	$T_1 \times T_1, T_1 \times T_1$	18	3, 3	W^+, W^-, W^0
		0	$T_{[\frac{1}{2} \frac{1}{2}]}$		16		
1	1	0		$T_1 \times T_1$	9	3	L^+, L^-, L^0
1	-1	0		$T_1 \times T_0$	3	1	L^{00}
0	-1	0		$T_0 \times T_1$	3	3	H^+, H^0, H^-
0	1	0		$T_0 \times T_0$	1	1	H^{00}

6.5 $j_1 + j_2 = 3/2$, spin 1/2 and 3/2

The representations $T_{[\frac{3}{2} 0]}$ and $T_{[0 \frac{3}{2}]}$ are related by space reflection, $S = j_1 + j_2 = 3/2$. There are 32 states in total and 8 particles (states that are not connected by transformations $SL(2, C)_{int}$, differing by the values S_R^3 ($\pm 1/2, \pm 3/2$) and by the parity, corresponding to 8 points on the Fig. 5b. Each point of the weight diagram (Fig. 5b) corresponds to four states (according to the number of possible spin projections).

The representations $T_{[1\frac{1}{2}]}$ and $T_{[\frac{1}{2}1]}$ are related by a space reflection, $S = j_1 + j_2 = 1$. There are 72 states in total. These are 12 spin 3/2 particles, differing by the parity, by the values of S_R (1/2, 3/2) and S_R^3 ($\pm 1/2, \pm 3/2$), and 12 spin 1/2 particles, differing by the parity, by the values of S_R (1/2, 3/2) and S_R^3 ($\pm 1/2, \pm 3/2$).

Table 3: Classification of scalar functions, $j_1 + j_2 = 3/2$, $m \neq 0$.

Spin	η	$ j_1 - j_2 $	Irreps of $SL(2, C)_{int}$	Irreps of $SU(2)_{int} \times SU(2)_{ext}$ and its dimension	dimension of $SU(2)_{int}$ irreps	
3/2	± 1	$\frac{3}{2}$	$T_{[\frac{3}{2}0]}, T_{[0, \frac{3}{2}]}$	$T_{\frac{3}{2}} \times T_{\frac{3}{2}}, T_{\frac{3}{2}} \times T_{\frac{3}{2}}$	32	4, 4
3/2	1	$\frac{1}{2}$	$T_{[\frac{3}{2}\frac{1}{2}]}, T_{[\frac{1}{2}\frac{3}{2}]}$	$T_{\frac{3}{2}} \times T_{\frac{3}{2}}, T_{\frac{3}{2}} \times T_{\frac{3}{2}}$	36	4, 4
3/2	-1	$T_{\frac{3}{2}} \times T_{\frac{1}{2}}, T_{\frac{3}{2}} \times T_{\frac{1}{2}}$		16	2, 2	
1/2	-1	$T_{\frac{1}{2}} \times T_{\frac{3}{2}}, T_{\frac{1}{2}} \times T_{\frac{3}{2}}$		16	4, 4	
1/2	1	$T_{\frac{1}{2}} \times T_{\frac{1}{2}}, T_{\frac{1}{2}} \times T_{\frac{1}{2}}$		8	2, 2	

7 The ultra-relativistic limit

Let us now consider functions corresponding to a massive particle, moving along the axis x^3 . They can be obtained from functions in the rest frame, which are characterized by a certain internal parity with the help of a Lorentz transformation $U \in SL(2, C)$,

$$P = UP_0U^\dagger, \quad Z = UZ_0, \quad \text{where } P_0 = \text{diag}\{m, m\}, \quad U = \text{diag}\{e^a, e^{-a}\},$$

$$p_0 = m \cosh 2a, \quad p_3 = m \sinh 2a, \quad e^{\pm a} = \sqrt{(p_0 \pm p_3)/m}. \quad (45)$$

For the orientation variables we have:

$$(z^1, \underline{z}^1, \overset{*}{z}_2, \overset{*}{\underline{z}}_2) \rightarrow e^a(z^1, \underline{z}^1, \overset{*}{z}_2, \overset{*}{\underline{z}}_2), \quad (z^2, \underline{z}^2, \overset{*}{z}_1, \overset{*}{\underline{z}}_1) \rightarrow e^{-a}(z^2, \underline{z}^2, \overset{*}{z}_1, \overset{*}{\underline{z}}_1).$$

If we apply these transformations to functions (41)–(44) and consider the ultra-relativistic limit, then only states for which helicity is equal to chirality will remain in it.

For example, applying these transformations to the state with $S_R^3 = -1/2$, $\eta = 1$ from (41), we find

$$f(x, z) \rightarrow f^{i(k_0x^0 + k_3x^3)} \left[C_1(z^1 e^a - \overset{*}{\underline{z}}_1 e^{-a}) + C_2(z^2 e^{-a} - \overset{*}{\underline{z}}_2 e^a) \right], \quad (46)$$

where the first term in the square brackets corresponds to $S^3 = 1/2$, and the second term corresponds to $S^3 = -1/2$. In the ultra-relativistic case with a positive a (i.e. with $k_3 > 0$) there remain only two components,

$$f'(x, z) \approx e^{i(k_0x^0 + k_3x^3)} \left(C_1 z^1 e^a - C_2 \overset{*}{\underline{z}}_2 e^a \right). \quad (47)$$

The first term corresponds to $S^3 > 0$ and chirality $\lambda = 1/2$, the second term corresponds to $S^3 < 0$ and chirality $\lambda = -1/2$. We see, that in the ultra-relativistic limit with $p_0 > 0$ the signs of chirality and helicity $\hat{p}\hat{S}$ are the same.

As should be expected, the results of this analysis of the ultra-relativistic case coincide with the conclusions on the chirality and helicity of a particle with spin 1/2, obtained on a basis of the Dirac equation.

For states (42) of spin 1 (representations of $T_{[10]}$ and $T_{[01]}$) in the ultra-relativistic limit also remain only 2 terms with the same signs of helicity and chirality.

For states (44) of spin 1 (representation $T_{[\frac{1}{2}, \frac{1}{2}]}$), in the ultra-relativistic limit there remains 1 term with $S^3 = 0$ and zero chirality.

8 First order left-invariant equations

If in the case of massless particles, first-order equations in \hat{p}^μ appear within the framework of the classification of functions with respect to the irreps of the Poincaré group, then for massive particles we have two second-order eigenvalue equations for the Casimir operators \hat{p}^2 and \hat{W}^2 . For massive particles, it is impossible to construct first-order equations using only the generators of the Poincaré group. The desired equations must be invariant under the transformations $M(3, 1)_{ext}$, and for particles with a certain parity, also under the space reflection. In this case, the invariance with respect to $M(3, 1)_{int}$ is not required, the internal symmetries can be broken.

From the orientation variables z one can construct operators that are transformed as vectors with respect to $M(3, 1)_{ext}$ and $M(3, 1)_{int}$, preserving the degree of polynomials in z ; see Ref. [3],

$$\hat{\Gamma}^{\mu n} = \frac{1}{2} \left(\bar{\sigma}^{\mu\dot{\alpha}\beta} \sigma_{\underline{b}\dot{\alpha}}^n z_{\dot{\alpha}}^{\dot{a}} \partial_{\underline{\beta}}^{\underline{b}} + \sigma^\mu_{\beta\dot{\alpha}} \bar{\sigma}^{n\dot{a}\underline{b}} z_{\underline{b}}^{\dot{\beta}} \partial_{\dot{a}}^{\dot{\alpha}} \right), \quad (48)$$

$$\underline{\hat{\Gamma}}^{\mu n} = \frac{1}{2} \left(\bar{\sigma}^{\mu\dot{\alpha}\beta} \sigma_{\underline{b}\dot{\alpha}}^n z_{\dot{\alpha}}^{\dot{a}} \partial_{\underline{\beta}}^{\underline{b}} - \sigma^\mu_{\beta\dot{\alpha}} \bar{\sigma}^{n\dot{a}\underline{b}} z_{\underline{b}}^{\dot{\beta}} \partial_{\dot{a}}^{\dot{\alpha}} \right), \quad (49)$$

with one external (left) and one internal (right) index. It is only the operators $\hat{p}_\mu \hat{\Gamma}^{\mu 0}$ and $\hat{p}_\mu \underline{\hat{\Gamma}}^{\mu k}$, $k = 1, 2, 3$, that are invariant under the space reflection, and the associated 4 equations possess solutions with a definite inner parity.

Let us consider the following equation:

$$\hat{p}^\mu \hat{\Gamma}^{\mu 0} f(x, z) = \mathcal{M} f(x, z), \quad (50)$$

where

$$\begin{aligned} 2\hat{\Gamma}^{\mu 0} &= \bar{\sigma}^{\mu\dot{\alpha}\beta} (z_{\dot{\alpha}}^{\dot{1}} \partial_{\underline{\beta}}^{\underline{1}} + z_{\dot{\alpha}}^{\dot{2}} \partial_{\underline{\beta}}^{\underline{2}}) + \sigma^\mu_{\beta\dot{\alpha}} (z_{\underline{1}}^{\dot{\beta}} \partial_{\dot{1}}^{\dot{\alpha}} + z_{\underline{2}}^{\dot{\beta}} \partial_{\dot{2}}^{\dot{\alpha}}) \\ &= (\bar{\sigma}^{\mu\dot{\alpha}\beta} z_{\dot{\alpha}}^{\dot{1}} \partial / \partial z^{\underline{\beta}} + \sigma^\mu_{\beta\dot{\alpha}} z_{\underline{\beta}}^{\dot{\beta}} \partial / \partial z_{\dot{\alpha}}^{\dot{\beta}}) + (\bar{\sigma}^{\mu\dot{\alpha}\beta} z_{\dot{\alpha}}^{\dot{2}} \partial / \partial z^{\underline{\beta}} + \sigma^\mu_{\beta\dot{\alpha}} z_{\underline{\beta}}^{\dot{\beta}} \partial / \partial z_{\dot{\alpha}}^{\dot{\beta}}). \end{aligned} \quad (51)$$

The set of the operators $\hat{S}^{\mu\nu}$ and $\hat{\Gamma}^{\mu 0}$ satisfies the commutation relations of the group $SO(3, 2)$.

Obviously, the equation (50) is not invariant under the right boosts, since the operator $\hat{p}_\mu \hat{\Gamma}^{\mu 0}$ does not commute with a part of the right generators, namely with $\hat{\mathbf{S}}\hat{\mathbf{B}}$, \hat{B}_R^3 and p_μ^R . However, it commutes with $\hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2$ and with generator S_R^3 of $SU(2)_{int}$, that is why particles described by the equation (50), may have certain values $j_1 + j_2$ and S_R^3 . And if the parity operator connects only states with different signs of $j_1 - j_2$, then the operator $\hat{p}_\mu \hat{\Gamma}^\mu$ connects states with all possible values j_1 and j_2 for a fixed sum $j_1 + j_2$.

Thus, the invariant subspaces of the operators $\hat{\Gamma}^{\mu 0}$ are subspaces of polynomials of degree $2(j_1 + j_2)$ by z with fixed values S_R^3 . In particular, such subspaces are the spaces of polynomials $f(x, z, \underline{z})$ and $f(x, \underline{z}, \dot{z})$ of degree $2(j_1 + j_2)$, characterized by $S_R^3 = \mp(j_1 + j_2)$.

Let's take a closer look at the case of spin 1/2. Linear functions from the subspace $f(x, z, \underline{z})$ depend on 4 variables $z^\alpha, z_{\dot{\alpha}}^*$, $S_R^3 = -1/2$. For them, we have 2 equations of the form (50) with different signs of the mass term. Each of these two equations combines two

irreps of the improper Poincaré group, characterized by the same sign of $p_0\eta$. One can see that in the rest frame

$$\varkappa^0 = \varepsilon m s, \quad \varepsilon = \eta \operatorname{sign} p_0 \operatorname{sign} S_R^3. \quad (52)$$

In total, for functions linear in orientation variables, we have 4 equations, their solutions in the rest frame at $p_0 > 0$ are given by formulas (41). Although the particle and its antiparticle belong to different subspaces with $S_R^3 = \pm 1/2$, they are described by equations with the same sign before the mass term.

Let us introduce into the consideration the generator of the group $U(1)$ of phase transformations of orientation variables z ,

$$\hat{\Gamma}^5 = \frac{1}{2} \left(z_{\underline{a}}^\alpha \partial_\alpha^{\underline{a}} - \overset{*}{z}_{\underline{b}}^{\dot{\beta}} \partial_{\dot{\beta}}^{\underline{b}} \right). \quad (53)$$

These transformations, along with external and internal automorphisms of the Poincaré group, can be considered as field symmetry transformations on the Poincaré group; see Ref. [3]. Polynomials of degree n in z, \underline{z} and of degree m in $\overset{*}{z}, \overset{*}{\underline{z}}$ are eigenfunctions of the operator $\hat{\Gamma}^5$ with the eigenvalue $n - m$. In the subspaces of the functions $f(x, z, \underline{z})$ and $f(x, \underline{z}, \overset{*}{z})$ its eigenvalues coincide with the chirality, defined as $j_1 - j_2$. The chirality operator Γ^5 commutes with the right and the left generators of the Poincaré group, but does not commute with $\Gamma^{\mu 0}$.

The matrix representation of the Dirac equation can be obtained from Eq. (50) for the functions $f(x, z, \underline{z})$ and $f(x, \underline{z}, \overset{*}{z})$, which are polynomials of the first order in z .

For example, substituting linear in z functions corresponding to the spin 1/2 from the subspace $f(x, z, \underline{z})$,

$$f_D(x, z) = \chi_\alpha(x) z^\alpha + \psi^{\dot{\alpha}}(x) \overset{*}{z}_{\dot{\alpha}} = Z_D \Psi_D(x), \quad Z_D = (z^\alpha \overset{*}{z}_{\dot{\alpha}}), \quad \Psi_D(x) = \begin{pmatrix} \chi_\alpha(x) \\ \psi^{\dot{\alpha}}(x) \end{pmatrix}, \quad (54)$$

into equation (50) and making a comparison of the coefficients at z^α and $\overset{*}{z}_{\dot{\alpha}}$ in the left- and right-hand sides, we obtain the Dirac equation

$$(\hat{p}_\mu \gamma^\mu - \varkappa) \Psi_D(x) = 0, \quad (55)$$

where $\varkappa = 2\varkappa^0$. The action of the chirality operator (53) amounts to the multiplication by $\gamma^5/2$, $\gamma^5 = \operatorname{diag}\{\sigma^0, -\sigma^0\}$,

$$\hat{\Gamma}^5 f_D(x, z) = \frac{1}{2} Z_D \gamma^5 \Psi_D(x).$$

Similarly, considering polynomials of the second degree leads to the 10-component Duffin-Kemmer equations, and higher degrees to the Bhabha equations. Note that starting with 2nd-order polynomials in z , states of the form (42) transforming according to representations of $T_{[j \ 0]}$ or $T_{[0 \ j]}$ are not, generally speaking, eigenfunctions of $\hat{p}^\mu \hat{\Gamma}^{\mu 0}$, and in the rest frame, solutions of these equations contain components that transform according to other representations.

9 Concluding remarks

A classification of relativistic orientable objects and its connection to the particle phenomenology is studied. This study is based on the possibility to describe these objects by a scalar field on the Poincaré group. Scalar functions $f(q)$, $q = (x, z)$, on the group

depend on the Minkowski space coordinates $x = (x^\mu) = (x^0, \mathbf{x})$, $\mathbf{x} = (x^k)$, as well as on the orientation variables $z = (z_{\alpha\beta})$ given by elements of the matrix $Z \in SL(2, C)$. The functions have certain transformation properties with respect to both groups $M(3, 1)_{ext}$ and $M(3, 1)_{int}$, see Sec. 1. We classify them with the help of maximal sets of commuting operators constructed from generators of the groups $M(3, 1)_{ext}$ and $M(3, 1)_{int}$.

We suppose that different particles correspond to states described by functions $f(x, z)$ that are not connected by transformations from $M(3, 1)_{ext}$ (i.e., by a change of space-fixed reference frame). Such states are classified by eigenvalues of Casimir operators and some functions of the right generators of the group (i.e., generators of the group $M(3, 1)_{int}$).

This way of the classification, applied to linear and quadratic functions of z allows one to identify some of them with known elementary particles of spin $0, \frac{1}{2}$, and 1 . Below, we list results of such an identification (the enumeration is given by values of spin and chirality in the massless limit).

Spin 1/2, chirality $\pm 1/2$. Such particles appear in quadruplets, but not in particle/antiparticle pairs.

In the massless case, we have 4 states and 4 particles – two $SL(2, C)_{int}$ doublets of chirality $\lambda = -1/2$ (particles) and $\lambda = 1/2$ (their antiparticles), interconnected by both space reflection and charge conjugation. The particle and the antiparticle differ in charge signs S_R^3 , each of them appear in one chiral state only. When reduced to $SU(2)_{int} \subset SL(2, C)_{int}$, these two-dimensional representations remain irreducible.

In the real world, the four fundamental fermions (in massless limit) have identical properties, for example, (e_L^-, ν_L) and $(e_R^+, \tilde{\nu}_R)$ or (d_L, u_L) and $(\tilde{d}_L, \tilde{u}_L)$.

In the massive case, we have 8 states and 4 particles, differing in sign S_R^3 and the parity ($\eta = 1$ corresponds to particles and $\eta = -1$ corresponds to antiparticles). In the rest frame, states of one particle differ by two spin projections $S_3 = \pm 1/2$. The four (e^-, ν) and $(e^+, \tilde{\nu})$ or (d, u) and (\tilde{d}, \tilde{u}) of fundamental fermions have analogous properties, excluding the equality of masses.

Spin 1, chirality ± 1 .

In the massless case, there appear 6 states and 6 particles: two $SO(3, C)_{int} \sim SL(2, C)_{int}$ triplets of chirality $\lambda = -1$ (particles) and $\lambda = 1$ (antiparticles), interconnected by both space reflection and charge conjugation. Components of the triplets differ in charge $S_R^3 = 0, \pm 1$, each particle is in only one chiral state. When reduced to $SU(2)_{int} \subset SL(2, C)_{int}$, these three-dimensional representations remain irreducible.

In the massive case, we have 18 states and 6 particles, differing in quantum numbers $S_R^3 = 0, \pm 1$ and in the parity, $\eta = 1$ (three particles) and $\eta = -1$ (three antiparticles). In the rest frame, states of one particle differ by three projections of spin, $S_3 = 0, \pm 1$.

These two triples differ only in signs of chirality ($m = 0$) or parity ($m \neq 0$). Therefore, if it turns out that the other charges of such particles are the same, then they can be considered as different polarization states of one particle. With this interpretation, we have one triplet, each component of which has 2 ($m = 0$) or 6 states ($m \neq 0$).

The triplet (W^-, W^0, W^+) of vector bosons (in massless limit) has identical properties with the one considered above. In the massive case the triplet (W^-, Z^0, W^+) has similar properties, excluding the equality of masses.

Spin 1, chirality 0.

In the case $m \neq 0$, we have $SO(3, 1)_{int} \sim SL(2, C)_{int}$ quadruplet of particles, containing two truly neutral particles ($\eta = \pm 1, S_R^3 = 0$) and two charged particles ($\eta = 1, S_R^3 = \pm 1$). Each particle from the quadruplet can be in 3 states, which differ in the rest frame by three projections of the spin $S_3 = 0, \pm 1$. In the ultra-relativistic limit, remains only one state with zero projection of the spin on the direction of motion.

The case $S_R^3 = 0$ corresponds to a truly neutral hypothetical particle, which is called

the notoph in the literature.

Chirality 0. In the massless case $SO(3, 1)_{int} \sim SL(2, C)_{int}$ quadruplet of particles contain two truly neutral particles ($S_R^3 = 0$), differing by the values of B_3^R , and two charged particles ($S_R^3 = \pm 1$). Each particle from the quadruplet has only one chiral state with zero projection of the spin on the direction of motion.

Spin 0. Scalar particles are represented by a singlet (independent of z functions $f(x)$) and a quadruplet (2nd order polynomials in z). The quadruplet consists of two truly neutral particles ($\eta = \pm 1$, $S_R^3 = 0$) and two charged particles ($\eta = -1$, $S_R^3 = \pm 1$). The characteristics of the Higgs boson correspond to both the singlet and to the component of the quadruplet with $\eta = 1$.

This list exhausts orientable objects described by polynomials of the 0, 1st and 2nd orders in orientation variables.

Among the states discussed above there exist states with the same S_R^3 that differ only by signs of the chirality (or the parity for $m \neq 0$). The question is: to consider these states describing different particles, or they are different polarization states of one and the same particle. If we start only from the representation theory of the Poincaré group, we do not get any answer to such a question. But, using the empirical formula (25), we see that in the case of spin 1/2, these states differ in the electric charge and, therefore, are states of different particles (electron and neutrino). At the same time, in the case of the spin 1, Eq. (25) does not prohibit interpreting them as different polarization states of one and the same particle (photon, W bosons).

When considering relativistic wave equations and particles described by them, only 8 out of 10 from the maximal set of commuting operators of the Poincaré group are usually used (6 left generators and 2 of 4 functions of the right generators, the latter define the irrep of the Lorentz group $SL(2, C)_{int}$). Here, for the classification of the orientable objects we have used the all 10 operators from the maximal set.

We stress that the results of developed classification, do not contradict the phenomenology of elementary particles, and, moreover, in some cases, present the corresponding group-theoretic explanation. Besides, the relationship of the spectra of the left and right generators allows us to give a possible explanation of the relationship between spin and the internal quantum numbers (charges) of particles, expressed in the positivity of R -parity.

In addition, it seems important to note that the composition and properties of particles predicted by the presented classification follow exclusively from the properties of the space-time and are, in a sense, model independent.

Appendix. Generators of the Lorentz group in the space of functions $f(z)$

The Lorentz group $SL(2, C)$ is a group of complex 2×2 matrices with a unit determinant,

$$Z = \begin{pmatrix} z^1_{\underline{1}} & z^1_{\underline{2}} \\ z^2_{\underline{1}} & z^2_{\underline{2}} \end{pmatrix} = \begin{pmatrix} z^1 & \underline{z}^1 \\ z^2 & \underline{z}^2 \end{pmatrix}, \quad \det Z = z^1 \underline{z}^2 - z^2 \underline{z}^1 = 1. \quad (56)$$

For the sake of brevity, we have used the notation that we applied in [3, 1] and $z^\alpha = z^\alpha_{\underline{1}}$, $z_{\dot{\alpha}}^* = z_{\dot{\alpha}}^{\underline{2}}$, $z^\alpha = z^\alpha_{\underline{2}}$, $z_{\dot{\alpha}}^* = z_{\dot{\alpha}}^{\underline{1}}$. In consequence of the unimodularity of 2×2 matrices Z there exist invariant antisymmetric tensors $\varepsilon^{\alpha\beta} = -\varepsilon^{\beta\alpha}$, $\varepsilon^{\dot{\alpha}\dot{\beta}} = -\varepsilon^{\dot{\beta}\dot{\alpha}}$, $\varepsilon^{12} = \varepsilon^{\dot{1}\dot{2}} = 1$, $\varepsilon_{12} = \varepsilon_{\dot{1}\dot{2}} = -1$. Now spinor indices are lowered and raised according to the rules

$$z_\alpha = \varepsilon_{\alpha\beta} z^\beta, \quad z^\alpha = \varepsilon^{\alpha\beta} z_\beta, \quad (57)$$

and, in particular

$$\hat{Z} = \begin{pmatrix} \overset{*}{z}_1 & \overset{*}{z}_2 \\ \overset{*}{z}_1 & \overset{*}{z}_2 \end{pmatrix} = \begin{pmatrix} -\overset{*}{z}_2 & \overset{*}{z}_1 \\ \overset{*}{z}_1 & -\overset{*}{z}_2 \end{pmatrix} = \begin{pmatrix} -\overset{*}{z}_2 & \overset{*}{z}_2 \\ \overset{*}{z}_1 & -\overset{*}{z}_1 \end{pmatrix}. \quad (58)$$

Besides the four-dimensional vector notation for spin operators, it is also convenient to use a three-dimensional notation: $\hat{S}^k = \frac{1}{2}\epsilon_{ijk}\hat{S}^{ij}$, $\hat{B}^k = \hat{S}_{0k}$. In the space of functions on the group $f(z)$ (functions of the elements of the matrix $SL(2, C)$ (56) and their complex conjugates) a direct calculation yields for left and right generators [1]:

$$\hat{S}^k = \frac{1}{2}(z\sigma^k\partial_z - \overset{*}{z}\overset{*}{\sigma}^k\partial_{\overset{*}{z}}) + \dots, \quad \hat{B}^k = \frac{i}{2}(z\sigma^k\partial_z + \overset{*}{z}\overset{*}{\sigma}^k\partial_{\overset{*}{z}}) + \dots, \quad z = (z^1 \ z^2), \quad \partial_z = (\partial/\partial z^1 \ \partial/\partial z^2)^T; \quad (59)$$

$$\hat{S}_R^k = -\frac{1}{2}(\chi\overset{*}{\sigma}^k\partial_\chi - \overset{*}{\chi}\sigma^k\partial_{\overset{*}{\chi}}) + \dots, \quad \hat{B}_R^k = -\frac{i}{2}(\chi\overset{*}{\sigma}^k\partial_\chi + \overset{*}{\chi}\sigma^k\partial_{\overset{*}{\chi}}) + \dots, \quad \chi = (z^1 \ \underline{z}^1), \quad \partial_\chi = (\partial/\partial z^1 \ \partial/\partial \underline{z}^1)^T. \quad (60)$$

The dots here stand for analogous expressions obtained by the change $z \rightarrow z' = (\underline{z}^1 \ \underline{z}^2)$, $\chi \rightarrow \chi' = (z^2 \ \underline{z}^2)$. In particular,

$$\hat{S}_R^3 = \frac{1}{2}(-z\partial_z + \underline{z}\partial_{\underline{z}} + \overset{*}{z}\partial_{\overset{*}{z}} - \overset{*}{\underline{z}}\partial_{\overset{*}{\underline{z}}}), \quad \hat{B}_R^3 = \frac{i}{2}(-z\partial_z + \underline{z}\partial_{\underline{z}} - \overset{*}{z}\partial_{\overset{*}{z}} + \overset{*}{\underline{z}}\partial_{\overset{*}{\underline{z}}}). \quad (61)$$

It is known that from \hat{S}^k and \hat{B}^k one can construct such linear combinations \hat{M}^k and $\hat{\bar{M}}^k$,

$$\hat{M}^k = \frac{1}{2}(\hat{S}^k - i\hat{B}^k) = z\sigma^k\partial_z + \underline{z}\sigma^k\partial_{\underline{z}}, \quad \hat{M}_+ = z^1\partial/\partial z^2, \quad \hat{M}_- = z^2\partial/\partial z^1, \quad \hat{\bar{M}}^k = -\frac{1}{2}(\hat{S}^k + i\hat{B}^k) = \overset{*}{z}\overset{*}{\sigma}^k\partial_{\overset{*}{z}} + \overset{*}{\underline{z}}\overset{*}{\sigma}^k\partial_{\overset{*}{\underline{z}}}, \quad \hat{\bar{M}}_+ = \overset{*}{z}^1\partial/\partial \overset{*}{z}^2, \quad \hat{\bar{M}}_- = \overset{*}{z}^2\partial/\partial \overset{*}{z}^1, \quad (62)$$

that $[\hat{M}^i, \hat{\bar{M}}^k] = 0$; in addition, for unitary representations of the Lorentz group, as it follows from the condition $\hat{S}^{k\dagger} = \hat{S}^k$, $\hat{B}^{k\dagger} = \hat{B}^k$, the relation $\hat{M}^{k\dagger} = \hat{\bar{M}}^k$ must be fulfilled (for finite-dimensional non-unitary representations $\hat{S}^{k\dagger} = \hat{S}^k$, $\hat{B}^{k\dagger} = -\hat{B}^k$ and $\hat{M}^{k\dagger} = -\hat{\bar{M}}^k$).

Taking into account the fact that the operators \hat{M}^k and $\hat{\bar{M}}^k$ satisfy the commutation relations of the algebra $su(2)$, we find the following relations for the spectra of the Casimir operators of the Lorentz subgroup:

$$\hat{\mathbf{S}}^2 - \hat{\mathbf{B}}^2 = 2(\hat{\mathbf{M}}^2 + \hat{\bar{\mathbf{M}}}^2) = 2(j_1(j_1 + 1) + j_2(j_2 + 1)) = 2(j_1 + j_2 + 1)(j_1 + j_2), \quad \hat{\mathbf{S}}\hat{\mathbf{B}} = -i(\hat{\mathbf{M}}^2 - \hat{\bar{\mathbf{M}}}^2) = -i(j_1(j_1 + 1) - j_2(j_2 + 1)) = -i(j_1 + j_2 + 1)(j_1 - j_2). \quad (63)$$

That is, the irreps of the Lorentz group $SL(2, C)$ are labeled by a pair of numbers $[j_1, j_2]$.

The difference $j_1 - j_2$ (the difference between the number of dotted and undotted indices) can also be obtained as an eigenvalue of the chirality operator $\hat{\Gamma}^5$ (53).

For finite-dimensional irreps of the group $SL(2, C)$, the formula of reduction to the compact $SU(2)$ -subgroup have the form

$$T_{[j_1, j_2]} = \sum_{j=|j_1-j_2|}^{j_1+j_2} T_j. \quad (64)$$

References

- [1] D.M. Gitman and A.L. Shelepin. Fields on the Poincaré group: Arbitrary spin description and relativistic wave equations. *Int. J. Theor. Phys.*, 40:603–684, 2001. arXiv:hep-th/0003146.
- [2] I.L. Buchbinder, D.M. Gitman, and A.L. Shelepin. Discrete symmetries as automorphisms of the proper Poincaré group. *Int. J. Theor. Phys.*, 41(4):753–790, 2002. arXiv:hep-th/0010035.
- [3] D.M. Gitman and A.L. Shelepin. Fields on the Poincaré group and quantum description of orientable objects. *Eur. Phys. J. C*, 61(1):111–139, 2009. arXiv:hep-th/0901.2537.
- [4] D.M. Gitman and A.L. Shelepin. Classification of quantum relativistic orientable objects. *Phys. Scr.*, 83:015103 (14pp), 2011. arXiv:1001.5290.
- [5] D.M. Gitman and A.L. Shelepin. Interaction of orientable object fields with gauge fields. *Phys. Scr.*, 84:055101 (13pp), 2011.
- [6] L.S. Biedenharn and J.D. Louck. *Angular Momentum in Quantum Physics*. Addison-Wesley, Reading, Massachusetts, 1981.
- [7] V.L. Ginzburg and I.E. Tamm. On the theory of spin. *Zh. Ehksp. Teor. Fiz.*, 17:227–237, 1947.
- [8] V. Bargmann and E.P. Wigner. Group theoretical discussion of relativistic wave equations. *Proc. Nat. Acad. USA*, 34:211–223, 1948.
- [9] H. Yukawa. Quantum theory of non-local fields. I. Free fields. *Phys. Rev.*, 77(2):219–226, 1950.
- [10] Yu.M. Shirokov. Relativistic theory of spin. *Zh. Ehksp. Teor. Fiz.*, 21(6):748–760, 1951.
- [11] D. Finkelstein. Internal structure of spinning particles. *Phys. Rev.*, 100(3):924–931, 1955.
- [12] F. Lurçat. Quantum field theory and the dynamical role of spin. *Physics*, 1:95, 1964.
- [13] H. Bacry and A. Kihlberg. Wavefunctions on homogeneous spaces. *J. Math. Phys.*, 10(12):2132–2141, 1969.
- [14] A. Kihlberg. Fields on a homogeneous space of the Poincaré group. *Ann. Inst. Henri Poincaré*, 13(1):57–76, 1970.
- [15] C.P. Boyer and G.N. Fleming. Quantum field theory on a seven-dimensional homogeneous space of the Poincaré group. *J. Math. Phys.*, 15(7):1007–1024, 1974.
- [16] H. Arodź. Metric tensors, Lagrangian formalism and Abelian gauge field on the Poincaré group. *Acta Phys. Pol., Ser. B*, 7(3):177–190, 1976.
- [17] M. Toller. Classical field theory in the space of reference frames. *Nuovo Cimento B*, 44(1):67–98, 1978.

- [18] M. Toller. Free quantum fields on the Poincaré group. *J. Math. Phys.*, 37(6):2694–2730, 1996.
- [19] W. Drechsler. Geometro-stochastically quantized fields with internal spin variables. *J. Math. Phys.*, 38(11):5531–5558, 1997.
- [20] L. Hannibal. Relativistic spin on the Poincaré group. *Found. Phys.*, 27(1):43–56, 1997.
- [21] V.V. Varlamov. Towards the quantum electrodynamics on the Poincare group. arXiv: hep-th/0403070.
- [22] S. Coleman and J. Mandula. All possible symmetries of the S -matrix. *Phys. Rev.*, 159(5):1251–1256, 1967.
- [23] N.Ya. Vilenkin. *Special Functions and the Theory of Group Representations*. AMS, Providence, 1968.
- [24] A.O. Barut and R. Raczká. *Theory of Group Representations and Applications*. PWN, Warszawa, 1977.
- [25] N.N. Bogolyubov, A.A. Logunov, A.I. Oksak, and I.T. Todorov. *General Principles of Quantum Field Theory*. Kluwer Acad. Pub., Dordrecht, 1990.
- [26] J.P. Elliott and P.O. Dawber. *Symmetry in Physics*, volume 2. The Macmillan Press LTD, London, 1979.
- [27] F. Lurçat and L. Michel. Sur les relations entre charges et spin. *Nuovo Cimento*, 21(3):574–576, 1961.
- [28] V.B. Berestetskii, E.M. Lifshitz, and L.P. Pitaevskii. *Relativistic Quantum Theory*. Pergamon, New York, 1971.
- [29] I. Białynicki-Birula. On the wave function of the photon. *Acta Phys. Pol., Ser. B*, 86(1-2):97–116, 1994.
- [30] V.I. Ogievetsky and I.V. Polubarinov. Notoph and its possible interactions. *Soviet J. Nucl. Phys.*, 4:156–161, 1967.
- [31] M. Kalb and P. Ramond. Classical direct interstring action. *Phys. Rev. D*, 9:2273–2284, 1974.
- [32] M. V. Chizhov. Theory and phenomenology of spin-1 chiral particles. *Phys. Part. Nucl.*, 42:93–183, 2011.