

There and Back Again: Mapping and Factorising Cosmological Observables

David Stefanyszyn*

*School of Mathematical Sciences & School of Physics and Astronomy,
University of Nottingham,
University Park, Nottingham, NG7 2RD, UK*

Xi Tong†

*Department of Applied Mathematics and Theoretical Physics,
University of Cambridge,
Wilberforce Road, Cambridge, CB3 0WA, UK*

Yuhang Zhu‡

*Cosmology, Gravity and Astroparticle Physics Group, Center for Theoretical Physics of the Universe,
Institute for Basic Science, Daejeon, 34126, Korea*

(Dated: June 4, 2024)

Cosmological correlators encode invaluable information about the wavefunction that characterises the state of the primordial universe. In this *letter* we present a simple yet novel duality between correlators and wavefunction coefficients that is valid to all orders in the loop expansion. The duality manifests itself as a \mathbb{Z}_4 symmetry in the correspondence between correlators and the physical parts of wavefunction coefficients. To demonstrate the power of the duality already at tree-level, we derive a correlator-to-correlator factorisation (CCF) formula for the parity-odd part of cosmological correlators that relates n -point observables to lower-point ones via a series of diagrammatic cuts. These relations are in principle testable as they involve observables defined for arbitrary physical kinematics (i.e. without performing any analytic continuation or taking any soft limits). For $n = 4$, the CCF formula provides a relation between the parity-odd part of the trispectrum of curvature perturbations, the bispectra involving two curvature perturbations and an additional state with a complementary series mass and integer spin, and the power spectrum of this additional state.

I. INTRODUCTION

One of the ultimate goals of physics is to understand the laws of nature at the beginning of time. A cosmologist’s approach to this problem is to measure spatial correlations in the Cosmic Microwave Background radiation and the Large Scale Structure of the universe. These correlations are seeded by primordial *cosmological correlators* of quantum fields evolving during inflation, with their momentum dependence encoding the secrets of the underlying physics at play during the universe’s first moments. For example, the soft limit of correlators probes the cosmic expansion history [1, 2] and the inflationary particle spectrum [3–30], the equilateral limit probes higher-dimensional self-interactions of the inflaton [31–35], while the collinear limit probes the initial state [32, 36–40] and environmental effects [41]. Understanding the structure of cosmological correlators is therefore of upmost importance in our quest to understand the early universe and therefore physics at extreme energy scales.

In recent years, however, much attention has been paid to more primitive objects, namely *wavefunction coefficients* that encode the inflationary dynamics in the per-

turbative expansion of the wavefunction of the universe (given that we work on a fixed background geometry, “field-theoretic wavefunction” might be a more appropriate name) [42, 43]. Although these objects are not directly observable, cosmological correlators can be extracted from them by applying the Born rule, and their somewhat simpler kinematic dependence means that constraints from cherished physical principles such as symmetries, locality and unitarity turn out to be more transparent [44–64]. They also play an important role in defining cosmological amplitudes [58, 65–68], can be used to understand the origin of IR-divergences in de Sitter space [69–71], contain (boost-breaking) flat-space amplitudes in a certain singular kinematic limit [72–74], and have neat connections to geometry [49, 75–77].

In this *letter*, we derive a duality between cosmological correlators B_n and the “physical” part of wavefunction coefficients $\rho_n \equiv \psi_n + \psi_n^\sharp$, where \sharp stands for complex conjugation and momentum reversal,¹ that is valid to all orders in perturbation theory (i.e. to all orders in the loop expansion). We show that in the dictionary that translates the $\{\rho_n\}$ to the $\{B_n\}$, there exists a \mathbb{Z}_4 symmetry that *syntactically* swaps $\rho_n \leftrightarrow B_n$ and maps any valid

* david.stefanyszyn@nottingham.ac.uk

† xt246@cam.ac.uk

‡ yhzhu@ibs.re.kr

¹ It is this combination that appears in the square modulus of the wavefunction of the universe and therefore contributes to the Born rule. In momentum space, it reads $\rho_n(\{\mathbf{k}\}) = \psi_n(\{\mathbf{k}\}) + \psi_n^*(\{-\mathbf{k}\})$ where $\{\mathbf{k}\}$ collectively denotes the external momenta.

equation to another valid equation within the dictionary. The duality therefore takes us from wavefunction coefficients to cosmological correlators, *and back again*.

We use this duality, in combination with the results of [78] where we showed that parity-odd correlators are factorised, to prove that under a set of mild assumptions, parity-odd correlators of inflatons and gravitons factorise into a *structured sum of products of lower-point correlators*. This factorisation holds for physical kinematics (no need for analytic continuation) and for generic momentum dependence (no need to take specific kinematic limits) and is in principle a relation that can be tested with observations. We show that it also has a neat interpretation in terms of diagrammatic cuts. The first non-trivial example of this correlator-to-correlator factorisation (CCF) relates the trispectrum of primordial perturbations to the bispectra involving two curvature perturbations and one additional state with integer spin and a complementary series mass, and the power spectrum of this state. The power of this relation lies in the fact that it maps an observable to a combination of other observables.

Notations and conventions. For conciseness, we use the DeWitt notation [79], where both field indices and spatial coordinates are abbreviated as a single Latin index as $\varphi^A(\mathbf{x}) \equiv \varphi_i$. Contractions are interpreted as $\varphi_i \chi^i \equiv \sum_A \int d^3x \varphi_A(\mathbf{x}) \chi^A(\mathbf{x})$. We adopt the following diagrammatic notations:

$$B_n \equiv \text{blob with } n \text{ legs} , \quad \rho_n \equiv \text{blob with } n \text{ legs} . \quad (1)$$

These blobs represent the abstract notion of correlators and wavefunction coefficients respectively, without specific perturbation theory structures inside. They should be distinguished from the typical Schwinger-Keldysh/Witten diagrams that compute B_n and ρ_n . For conciseness, all momentum and tensor indices have been suppressed.

II. CORRELATOR-WAVEFUNCTION DUALITY

We start with a lightning review of the wavefunction approach to primordial perturbations. Consider a set of weakly interacting quantum fields collectively denoted by $\Phi^i(\eta)$ evolving in a classical inflationary spacetime $g_{\mu\nu} = a^2(\eta)\eta_{\mu\nu}$, $a(\eta) = -1/(H\eta)$. In the Schrödinger picture, we define the wavefunction of the universe by projecting the time-evolved Bunch-Davies vacuum onto a field-value eigenstate:

$$\Psi[\varphi] = \langle \varphi | U(\eta_0, -\infty) | \text{BD} \rangle = \int_{\text{BD}}^{\Phi(\eta_0)=\varphi} \mathcal{D}\Phi e^{iS[\Phi]} . \quad (2)$$

In practice, the Bunch-Davies vacuum choice means that we integrate over fields that vanish in the far past (with a suitable contour deformation). In perturbation theory,

the wavefunction is conventionally parametrised by

$$\Psi[\varphi] = \exp \left[+ \sum_{n=1}^{\infty} \frac{1}{n!} (\psi_n)^{i_1 \dots i_n} \varphi_{i_1} \dots \varphi_{i_n} \right] , \quad (3)$$

with the coefficients ψ_n computed via Witten diagrammatics. Given our notation, there is an implicit sum over different fields in (3) thereby ensuring that each wavefunction coefficient has the appropriate normalisation e.g. for ψ_3 with only two identical fields, we have an overall factor of $\frac{3}{3!} = \frac{1}{2!}$. We unconventionally start the summation with $n = 1$. This tadpole term starts out at loop-level and is necessary for the cancellation of the monopole moment in observables.

The full n -point correlators of quantum fields are computed by the Born rule as²

$$\langle \varphi_{i_1} \dots \varphi_{i_n} \rangle = \frac{\int \mathcal{D}\varphi |\Psi[\varphi]|^2 \varphi_{i_1} \dots \varphi_{i_n}}{\int \mathcal{D}\varphi |\Psi[\varphi]|^2} . \quad (4)$$

However, the quantities of more observational relevance are their connected part, which we will denote as B_n , in which only one momentum-conserving δ -function is present in momentum space. These are computed from the generating functional

$$Z[J] = \int \mathcal{D}\varphi |\Psi[\varphi]|^2 e^{iJ^i \varphi_i} , \quad (5)$$

by taking derivatives with respect to the auxiliary current i.e.

$$(B_n)_{i_1 \dots i_n} = \frac{\partial}{i \partial J^{i_1}} \dots \frac{\partial}{i \partial J^{i_n}} \ln Z[J] \Big|_{J=0} . \quad (6)$$

This yields the correct normalisation since the denominator in (4) is simply $Z[0]$, and we emphasise that the derivatives act on $\ln Z[J]$ rather than $Z[J]$ thereby ensuring we only extract connected correlators. Alternatively, we can integrate (6) to obtain

$$\begin{aligned} & \exp \left[+ \sum_{n=1}^{\infty} \frac{i^n}{n!} (B_n)_{i_1 \dots i_n} J^{i_1} \dots J^{i_n} \right] \\ &= \int \mathcal{D}\varphi \exp \left[+ \sum_{n=1}^{\infty} \frac{1}{n!} (\rho_n)^{i_1 \dots i_n} \varphi_{i_1} \dots \varphi_{i_n} \right] e^{iJ^i \varphi_i} , \end{aligned} \quad (7)$$

where $\rho_n \equiv \psi_n + \psi_n^\dagger$ is the physical part of a wavefunction coefficient. Therefore, the *connected correlators* can be viewed as a Fourier transformation of the physical wavefunction coefficients. Now notice that the two sides of

² Note that in perturbation theory, the path integral is always computed by expanding and truncating the non-Gaussian part of $|\Psi[\varphi]|^2$, with convergence guaranteed by the Gaussian part with $\rho_2 < 0$.

(7) take completely analogous forms. We can therefore perform an inverse Fourier transformation to rewrite (7) as

$$\begin{aligned} & \exp \left[+ \sum_{n=1}^{\infty} \frac{1}{n!} (\rho_n)^{i_1 \dots i_n} \phi_{i_1} \dots \phi_{i_n} \right] \\ &= \int \mathcal{D}J \exp \left[+ \sum_{n=1}^{\infty} \frac{i^n}{n!} (B_n)_{i_1 \dots i_n} J^{i_1} \dots J^{i_n} \right] e^{-iJ^i \phi_i}. \end{aligned} \quad (8)$$

After a change of the dummy variable $J \rightarrow -J$, we obtain

$$\begin{aligned} & \exp \left[+ \sum_{n=1}^{\infty} \frac{1}{n!} (\rho_n)^{i_1 \dots i_n} \phi_{i_1} \dots \phi_{i_n} \right] \\ &= \int \mathcal{D}J \exp \left[+ \sum_{n=1}^{\infty} \frac{(-i)^n}{n!} (B_n)_{i_1 \dots i_n} J^{i_1} \dots J^{i_n} \right] e^{iJ^i \phi_i}. \end{aligned} \quad (9)$$

Comparing (7) and (9), we see that a syntactic replacement³

$$g : \begin{pmatrix} \rho_n \\ B_n \end{pmatrix} \mapsto (-i)^n \begin{pmatrix} B_n \\ \rho_n \end{pmatrix}, \quad (10)$$

maps them to each other (after some labelling). Since both (7) and (9) are equivalent statements of the correlator-wavefunction dictionary, we deduce that g is an exact symmetry of the dictionary. Since $g^4 = 1$, the duality mapping generates a \mathbb{Z}_4 group that maps the dictionary to itself. The formal proof above shows that such a \mathbb{Z}_4 symmetry is valid to *arbitrary* finite orders in perturbation theory, since the Gaussian integrals in both (7) and (9) are well-defined as long as $\rho_2 < 0 < B_2$. Note that in this derivation we have dropped an integration constant since ultimately the relationship between the ρ_n and B_n comes from taking derivatives with respect to an auxiliary current and therefore an overall constant in (9) is inconsequential.

To demonstrate the duality, let us inspect some simple examples. Up to $n = 5$ at tree-level, we have

$$B_2 = -\frac{1}{\rho_2}, \quad (11a)$$

$$B_3 = -\frac{1}{\rho_2^3} \rho_3, \quad (11b)$$

$$B_4 = \frac{1}{\rho_2^4} \left[\rho_4 - \left(\rho_3 \frac{1}{\rho_2} \rho_3 + 2 \text{ perms} \right) \right], \quad (11c)$$

$$\begin{aligned} B_5 = & -\frac{1}{\rho_2^5} \left[\rho_5 - \left(\rho_4 \frac{1}{\rho_2} \rho_3 + 9 \text{ perms} \right) \right. \\ & \left. + \left(\rho_3 \frac{1}{\rho_2} \rho_3 \frac{1}{\rho_2} \rho_3 + 14 \text{ perms} \right) \right], \end{aligned} \quad (11d)$$

where the internal DeWitt indices are understood as contracted. Note that, up to minus signs, the coefficient of each term is unity thanks to the normalisation of the wavefunction. Under the duality mapping g , equations (11) become

$$-\rho_2 = -\frac{1}{-B_2}, \quad (12a)$$

$$i\rho_3 = -\frac{1}{-B_2^3} iB_3, \quad (12b)$$

$$\rho_4 = \frac{1}{B_2^4} \left[B_4 - \left(iB_3 \frac{1}{-B_2} iB_3 + 2 \text{ perms} \right) \right], \quad (12c)$$

$$\begin{aligned} -i\rho_5 = & -\frac{1}{-B_2^5} \left[-iB_5 - \left(B_4 \frac{1}{-B_2} iB_3 + 9 \text{ perms} \right) \right. \\ & \left. + \left(iB_3 \frac{1}{-B_2} iB_3 \frac{1}{-B_2} iB_3 + 14 \text{ perms} \right) \right], \end{aligned} \quad (12d)$$

which are equivalent to solving the original equations (11) for the ρ 's. Note also that since the $n = 1$ entry starts out at loop-level, we have neglected it here. In practice, it is convenient to remove the tree-level tadpoles, so that the Gaussian term dominates the typical field fluctuations. However, we note that the duality (10) works for any values of ρ_1 and B_1 , since convergence is always guaranteed by the Gaussian term at non-typically large field fluctuations. In Appendix A, we explicitly verify the duality including the $n = 1$ tadpole terms up to 4-point 1-loop order, and show that consistently keeping tadpole terms is essential for the duality to work. We have further successfully confirmed the validity of the duality with a channel-insensitive check at 9-point 4-loop order using a computer algorithm.⁴

In general, the tree-level dictionary translating physical wavefunction coefficients to correlators (in the absence of linear mixings which we assume throughout) reads

$$B_n = \frac{1}{(-\rho_2)^n} \sum_{k=0}^{n-3} (-1)^k (k\text{-cuts})_{\rho}, \quad (13)$$

with

$$\begin{aligned} (k\text{-cuts})_{\rho} \equiv & \sum_{n-k \geq n_1 \dots n_{k+1} \geq 3} \left[\rho_{n_1} \frac{1}{\rho_2} \rho_{n_2} \dots \rho_{n_k} \frac{1}{\rho_2} \rho_{n_{k+1}} \right. \\ & \left. + (\pi_{n_1 \dots n_{k+1}} - 1)\text{-perms} \right]. \end{aligned} \quad (14)$$

The correlator-wavefunction duality then implies the reciprocal formula

$$\rho_n = \frac{1}{B_2^n} \sum_{k=0}^{n-3} (-1)^k (k\text{-cuts})_B, \quad (15)$$

³ With an alternative wavefunction sign convention $\hat{\rho}_2 \equiv -\rho_2 > 0$, the duality mapping reads $g : \hat{\rho}_2 \mapsto B_2$ while the $n \neq 2$ mapping rules remain the same.

⁴ The Mathematica notebook that verifies the duality is available [here](#).

where $\left(k\text{-cuts}\right)_B$ is obtained from (14) via a *syntactic* substitution $\rho \rightarrow B$. Notice that here we have applied the tree-level topology to write $(-i)^{n_1+\dots+n_{k+1}-n} = (-1)^k$.

In summary, the power of the duality allows us to directly invert the dictionary without the need of ever performing the algebraic inversion in practice, and all the combinatorics are automatically left intact.

III. FACTORISATION OF PARITY-ODD CORRELATORS

Let's now see how the duality we derived above can be put to good use. In [78], we derived a factorisation theorem for cosmological correlators of the inflaton and graviton which states that n -point functions of these states are factorised into lower-point objects if these observables are parity-odd (PO).⁵ This theorem allows for correlators arising from the exchange of additional states of *any* mass and integer spin (in addition to contact diagrams), and relies on the following small set of mild assumptions:

- Unitarity and locality
- The tree-level approximation
- Bunch-Davies vacuum conditions
- IR-convergence of the nested time integrals that compute cosmological correlators
- Scale invariance of the interactions

An immediate consequence of the theorem is the absence of total-energy singularities in the PO sector of primordial perturbations which leads to a very nice distinction between the PO and parity-even (PE) sectors.⁶ The theorem does not, however, state what objects such correlators factorise into. This is where the duality we have derived here comes to fruition. Indeed, the basis of the factorisation theorem of [78] is a proof that for the PO sector we have $\rho_n^{\text{PO}} = 0$ (as long as the external states are the inflaton and/or the helicity-summed graviton, and any internal states that are produced during inflation and decay into these massless states are in the *complementary series* of dS representations or the $SO(3)$ representations of [26]). This follows from the fact that for the PO sector we have $\rho_n^{\text{PO}} = \psi_n(\mathbf{k}) - \psi_n^*(\mathbf{k})$ i.e. it is the imaginary part of wavefunction coefficients that contribute to cosmological observables, yet under the above assumptions, wavefunction coefficients are purely real. This can be proven on very general grounds by performing Wick rotations of the time variables and using the fact that the

time-ordered part of the bulk-bulk propagator is purely real after this rotation,⁷ as are the vertices and the bulk-boundary propagator [78]. The result holds for exchanging fields of arbitrary integer spin.

Turning our attention to (15), in the PO sector the factorisation theorem of [78] therefore implies

$$\frac{1}{B_2^n} \sum_{k=0}^{n-3} (-1)^k \left[\left(k\text{-cuts}\right)_B \right]^{\text{PO}} = \rho_n^{\text{PO}} = 0, \quad (16)$$

which can be rearranged to yield a formula for B_n^{PO} in terms of lower-point *correlators*:

$$B_n^{\text{PO}} = \sum_{k=1}^{n-3} (-1)^{k-1} \left[\left(k\text{-cuts}\right)_B \right]^{\text{PO}}, \quad \forall n \geq 4. \quad (17)$$

In these expressions $[\dots]^{\text{PO}}$ indices that we are projecting correlators onto their PO part, which for correlators of the inflaton means that we take the imaginary part.⁸ Note that this formula holds regardless of how the parity violation arises which could be due to parity-violating vertices or due to the exchange of a spinning field with a parity-violating two-point function. We therefore see that the correlator-wavefunction duality has enabled us to derive, using the factorisation theorem of [78], a correlator-to-correlator factorisation (CCF) formula for the PO sector of primordial perturbations which has a neat interpretation in terms of correlator cuts. As an example, for $n = 4$ there is only one possibility where an exchange diagram is cut into two cubic diagrams and (17) can be diagrammatically represented by

$$\left(\text{tree} \right)^{\text{PO}} = 3 \left(\text{tree} \right)^{\text{PO}}, \quad (18)$$

with the factor of 3 symbolising the three different channels. The absence of a contact diagram contribution was already noted in [83, 84]. For $n = 5$ there is more structure with single and double cuts possible. We have

$$\left(\text{tree} \right)^{\text{PO}} = 10 \left(\text{tree} \right)^{\text{PO}} - 15 \left(\text{tree} \right)^{\text{PO}}, \quad (19)$$

where again the numerical factors are counting distinct channels, and the different colours allow for the exchange of different fields.

⁵ For the graviton, the reality and factorisation theorems of [78] hold once we sum over the two helicities.

⁶ This is a tree-level statement: total-energy singularities can arise at loop-level [80].

⁷ Note that with this rotation we are still computing dS wavefunction coefficients rather than EAdS correlators. For connections between dS and AdS observables see [81, 82].

⁸ This follows from the fact that in momentum space the inflaton is a parity transformation away from being real.

In the above derivation we took additional states to be in the complementary series as this allowed us to set $\rho_n^{\text{PO}} = 0$ [78]. If there are also principle series fields then this is not possible since then $\rho_n^{\text{PO}} \neq 0$. Correlators are indeed still factorised, but they do not factorise into other correlators. Our CCF formula is still useful for principle series fields, however, since if we have an explicit expression for B_n^{PO} due to the exchange of complementary series fields (which can be computed using the CCF formula), we can analytically continue the mass parameter to derive corresponding expressions for the exchange of principle series fields.⁹

The most relevant case for phenomenology is $n = 4$ (corresponding to the trispectrum) and with curvature perturbations as the external states. This observable has received much attention recently due to the purported detection of parity-violation in the galaxy four-point function [85, 86] (see also [87, 88], however). In this case our CCF formula reads

$$B_4^{\text{PO}}(\{\mathbf{k}\}) = \left[B_3(\mathbf{k}_1, \mathbf{k}_2, -\mathbf{s}) \cdot \frac{1}{B_2(\mathbf{s})} \cdot B_3(\mathbf{s}, \mathbf{k}_3, \mathbf{k}_4) \right]^{\text{PO}} + (\mathbf{t} + \mathbf{u})\text{-channels}, \quad (20)$$

which provides a neat connection between distinct observables, namely between the PO part of the trispectrum of curvature perturbations, the bispectrum consisting of two curvature perturbations and one additional state with integer spin and mass in the complementary series, and the power spectrum of this new state. This relation is in principle testable, and any violation of this relation would imply one of the above listed assumptions is violated. One case of particular interest is where the exchanged field is the graviton. Parity-violation in such a set-up can come from a number of sources e.g. as a dynamical Chern-Simons correction to the graviton propagator [89] or due to a mixing between the graviton and an $SU(2)$ gauge field [90, 91]. It would be interesting to study these cases in more detail given our CCF formula.

Finally, let us point out that although the CCF relation (17) is almost completely transparent in the wavefunction language thanks the correlator-wavefunction duality, it seems rather mysterious from the traditional in-in/Schwinger-Keldysh (SK) diagrammatics of [92]. One can indeed derive CCF relations using SK diagrammatics for specific lower-point examples, as we demonstrate for $n = 4$ in Appendix B, yet the general proof seems to be hidden from sight. It remains an intriguing question as to why boundary correlators should factorise into other

correlators even away from specific kinematic limits (e.g. the OPE limit).

IV. SUMMARY

In this work we have derived a duality in the dictionary between cosmological correlators $\{B_n\}$ and the physical part of wavefunction coefficients $\{\rho_n\}$ that is valid to all orders in perturbation theory. This duality allows us to derive a reciprocal formula that reconstructs $\{\rho_n\}$ from $\{B_n\}$ in a syntactic fashion. When combined with the results of [78] which states that $\rho_n^{\text{PO}} = 0$ for massless scalar/graviton external states, and complementary series internal states, we obtain an infinite set of correlator-to-correlator factorisation (CCF) formulae. These relations state that n -point PO correlators at tree-level are factorised into structured combinations of lower-point correlators. These CCF relations involve observables defined for *physical* kinematics and can therefore in principle be tested observationally. Any violation of these relations would directly point to the failure of the tree-level assumption, unitarity, locality, scale invariance or the Bunch-Davies vacuum. We showed how our CCF formulae can be understood in terms of diagrammatic cuts, and since taking these cuts does not require any analytical continuation, they serve as a way to understand the general structure of cosmological observables in a physically accessible manner.

Our work certainly opens up many avenues for future exploration. For instance, the \mathbb{Z}_4 symmetry goes beyond the context of cosmology all the way to connected Green functions and EFT Wilson coefficients in general QFTs. It would be interesting to see if one can make general statements there too. In addition, it would be interesting to extend our CCF formula (or something akin to CCF) to loop-level. Finally, it would be neat to find a full proof of the CCF formula directly using the Schwinger-Keldysh formalism, where only observables are involved from the get-go.

Acknowledgements We thank Paolo Benincasa, Trevor Cheung, Zongzhe Du, Sadra Jazayeri, Eiichiro Komatsu, Hayden Lee, Ciaran McCulloch, Enrico Pajer, Ayngaran Thavanesan, Dong-Gang Wang, Yi Wang, Zhong-Zhi Xianyu and Masahide Yamaguchi for helpful discussions. D.S. is supported by a UKRI Stephen Hawking Fellowship [grant number EP/W005441/1] and a Nottingham Research Fellowship from the University of Nottingham. X.T. is supported by STFC consolidated grants ST/T000694/1 and ST/X000664/1. Y.Z. is supported by the IBS under the project code, IBS-R018-D3. X.T. thanks the University of Nottingham for kind hospitality. For the purpose of open access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising.

Data access statement No new data were created or analysed during this study.

⁹ We need to take the $\eta_0 \rightarrow 0$ limit before performing any analytical continuation. We emphasise that the continuation is only possible at the level of the correlators and not wavefunction coefficients, since only the Schwinger-Keldysh propagators are analytic in the mass parameters. Indeed, the future boundary condition placed on the wavefunction bulk-bulk propagator makes a distinction between complementary and principle series fields.

Appendix A: The Duality at 4-point, 1-loop

In this section, we present an explicit check of the correlator-wavefunction duality at the 4-point 1-loop-level. To establish a consistent power-counting scheme in perturbation theory, we have in mind an underlying interacting theory of the form $\lambda\Phi^3$, with $\lambda \ll 1$ being an expansion parameter. We use ξ to count the number of loops, namely the number of unknown momenta that are integrated over. These can come from *quantum loop* corrections to the wavefunction coefficients, or from *classical loops* that arise when we perform the path integral to take us from wavefunction coefficients to correlators. Without loss of generality, we organise the perturbation series expansion in powers of λ and ξ . As long as the duality works at order $\mathcal{O}(\lambda^p \xi^q)$ for *any* $p, q \in \mathbb{N}$, we can claim the validity of the duality is independent of this choice of power-counting scheme. Notice also that B_n and ρ_n are always of the same order in λ and ξ . To illustrate this counting and to explain our diagrammatic notation, consider the following diagrams:

$$\text{---}\bullet\text{---} = \mathcal{O}(1), \quad (\text{A1a}) \quad \text{---}\bullet\text{---}\text{---}\text{---} = \mathcal{O}(\lambda\xi), \quad (\text{A1d})$$

$$\text{---}\bullet\text{---}\text{---}\text{---} = \mathcal{O}(\lambda), \quad (\text{A1b}) \quad \text{---}\bullet\text{---}\text{---}\text{---} = \mathcal{O}(\lambda^2\xi), \quad (\text{A1e})$$

$$\text{---}\bullet\text{---} = \mathcal{O}(\lambda\xi), \quad (\text{A1c}) \quad \text{---}\bullet\text{---}\text{---}\text{---} = \mathcal{O}(\lambda^2), \quad (\text{A1f})$$

where we have removed the tree-level contribution to tadpoles and regarded both ρ_1 and B_1 as small loop-order perturbations. (A1a) simply represents ρ_2 computed to any order in perturbation theory. There is a tree-level contribution and therefore the leading contribution is independent of λ and ξ . (A1b) and (A1c) similarly represent ρ_3 and ρ_1 computed to any order in perturbation theory. The former is $\mathcal{O}(\lambda)$ since the leading contribution comes from a tree-level contact diagram with a single cubic vertex, while for ρ_1 there is no tree-level contribution so the leading term is at one-loop and comes from a single cubic vertex and so it is $\mathcal{O}(\lambda\xi)$. (A1d) represents ρ_3 computed to any order in perturbation theory but with two of the external momenta fixed to be equal and opposite and integrated over thereby producing a contribution to the 1-point function. Since ρ_3 starts out at tree-level and is $\mathcal{O}(\lambda)$, once we perform the momentum integration we have $\mathcal{O}(\lambda\xi)$. Note that if ρ_3 is computed at tree-level, (A1d) represents a ‘‘classical loop’’ in the language of [69, 80]. (A1e) represents a product of ρ_3 and ρ_1 for which we still only have a single overall momentum-conserving delta function. Such contributions to correlation functions once we perform the Born rule are usually referred to as ‘‘factorised contributions’’ which should be distinguished from disconnected contributions that carry more than one momentum-conserving delta function. Since ρ_3 starts at tree-level and is $\mathcal{O}(\lambda)$, while ρ_1 starts out at loop level and is $\mathcal{O}(\lambda\xi)$, this dia-

gram is $\mathcal{O}(\lambda^2\xi)$. Finally, (A1f) represents ρ_4 computed to any order in perturbation theory. The leading contribution comes from a tree-level exchange diagram so it is $\mathcal{O}(\lambda^2)$.

As an explicit example we will work up to $\mathcal{O}(\lambda^2\xi)$, in which case the Born rule yields the following relations between the B_n and ρ_n :

$$\text{---}\text{---}\text{---}\text{---} = - \frac{\text{---}\bullet\text{---}}{\text{---}\bullet\text{---}} + \frac{1}{2} \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^2}, \quad (\text{A2a})$$

$$\text{---}\text{---}\text{---}\text{---} = - \frac{1}{\text{---}\bullet\text{---}} - \frac{1}{2} \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^3} - \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^3} + \frac{1}{2} \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^4} + \frac{1}{2} \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^4}, \quad (\text{A2b})$$

$$\text{---}\text{---}\text{---}\text{---} = - \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^3}, \quad (\text{A2c})$$

$$\text{---}\text{---}\text{---}\text{---} = \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^4} - 3 \frac{\text{---}\bullet\text{---}\text{---}\text{---}}{(\text{---}\bullet\text{---})^5}. \quad (\text{A2d})$$

Note that the factor of 3 in the second term on the RHS of (A2d) is counting the three distinct channels whereas all other factors in other expressions are just numerical factors that arise from doing the path integral. Any relations that involve B_n or ρ_n with $n \geq 5$ are higher-order in λ so we don’t consider them here. As an example, to derive (A2a), we first expand the generating functional (5) to $\mathcal{O}(J\lambda^2\xi)$,

$$Z[J] = \text{Det} \left(\frac{-\rho_2}{2\pi} \right)^{-1/2} \left\{ 1 + i J^i (\rho_2^{-1})_{ij} \left[-(\rho_1)^j + \frac{1}{2} (\rho_3)^{jkl} (\rho_2^{-1})_{kl} \right] + \dots \right\}, \quad (\text{A3})$$

and then differentiate to obtain the 1-point correlator

$$(B_1)_i = \frac{\partial}{i \partial J^i} \ln Z[J] \Big|_{J=0} = -(\rho_2^{-1})_{ij} (\rho_1)^j + \frac{1}{2} (\rho_2^{-1})_{ij} (\rho_3)^{jkl} (\rho_2^{-1})_{kl}, \quad (\text{A4})$$

which corresponds to (A2a). In momentum space for a single-field theory, this reads

$$B_1(\mathbf{k}) = -\frac{\rho_1(\mathbf{k})}{\rho_2(\mathbf{k})} + \frac{1}{2} \int \frac{d^3q}{(2\pi)^3} \frac{\rho_3(\mathbf{k}, \mathbf{q}, -\mathbf{q})}{\rho_2(\mathbf{k})\rho_2(\mathbf{q})}, \quad (\text{A5})$$

where all momentum-conserving δ -functions are stripped. The second contribution on the RHS of this expression is the ‘‘classical loop’’.

To check the duality relation, we need to invert equations (A2a)-(A2d) and express ρ in terms of B before expanding and truncating at order $\mathcal{O}(\lambda^2\xi)$. The expressions for ρ_3 and ρ_4 can be easily worked out as

$$\rho_3 = \frac{\text{diagram}}{(\text{diagram})^3}, \quad (\text{A6})$$

$$\rho_4 = \frac{\text{diagram}}{(\text{diagram})^4} - 3 \frac{\text{diagram}}{(\text{diagram})^5}. \quad (\text{A7})$$

Up to $\mathcal{O}(\lambda^2\xi)$, all contributions in (A6) and (A7) are in fact tree-level ones. With the expression (A6) for ρ_3 , we can then invert (A2a) to get

$$\rho_2 = \frac{\text{diagram}}{\text{diagram}} - \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^2}. \quad (\text{A8})$$

The ρ_2 case turns out to be more complicated. After solving for ρ_2 and substituting each ρ_n by their expression in terms of B_n , $n = 1, 3, 4$, we obtain

$$\begin{aligned} \rho_2 &= - \frac{1}{\text{diagram}} \\ &- \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^3} + \frac{2}{2} \frac{\text{diagram}}{(\text{diagram})^4} + \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4} \\ &- \frac{\text{diagram}}{(\text{diagram})^3} + \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4} \\ &- \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4} \\ &- \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4}, \end{aligned} \quad (\text{A9})$$

where the last four lines correspond to the last four terms in (A2b), respectively. Note that among the three channels of $B_3 \cdot B_3$, two of them contract into a bubble diagram, and the remaining channel becomes a 1-loop tadpole diagram (the last term in the second line here). We have highlighted identical terms with the same colour to indicate the intricate cancellation of combinatorics at

play here. Finally, we arrive at

$$\begin{aligned} \rho_1 &= - \frac{1}{\text{diagram}} - \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^3} - \frac{\text{diagram}}{(\text{diagram})^3} \\ &+ \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4} + \frac{1}{2} \frac{\text{diagram}}{(\text{diagram})^4}. \end{aligned} \quad (\text{A10})$$

Now it is straightforward to check the duality. Applying $g : (\rho_n, B_n) \rightarrow (-i)^n (B_n, \rho_n)$ to (A2), we indeed land on the desired equations (A8),(A10),(A6) and (A7).

As we can see from above, the inclusion of tadpole terms ρ_1, B_1 is essential in maintaining the duality. In practice, insisting tadpole cancellation $\rho_1 = 0$ or $B_1 = 0$ would then spontaneously break the duality at loop-level. However, the \mathbb{Z}_4 symmetry is still secretly at play, meaning the structure of the inverted dictionary is not arbitrary: it must be consistent with the \mathbb{Z}_4 symmetry if one reinserts the tadpole terms.

Before ending this section, we point out that the inversion procedure above can be automated by a computer algorithm. We have successfully confirmed the validity of the duality with a channel-insensitive check at 9-point 4-loop order.¹⁰

Appendix B: Correlator-to-Correlator Factorisation from Schwinger-Keldysh Diagrams at 4pt

Even though we have used the duality and the result that $\rho_n^{\text{PO}} = 0$ to prove that PO correlators are factorised into combinations of lower-point correlators for any n , it is instructive to see how we could draw the same conclusion directly at the level of Schwinger-Keldysh diagrams. We will illustrate this for the simplest case of $n = 4$, and to be concrete we will focus on the PO trispectrum of curvature perturbations generated via the exchange of a massive spin-1 field (the generalisation to other theories is straightforward). In principle, B_4^{PO} is computed as a sum over contact and exchange diagrams. However, it is now very well-known that the contact diagram contributions at tree-level yield a vanishing correlator once both sides of the in-in contour are added together [83, 84] (as is also the case for parity-odd Weyl³ graviton correlator [72, 93] and the same cancellation explains the simplicity and rarity of other parity-odd graviton correlators [94–97]), so let's instead focus solely on exchange diagrams. We assume that there are two cubic vertices each of the form $\pi\pi\sigma$ where π is the inflaton and σ is the massive spin-1 field with its index suppressed. Such vertices can

¹⁰ See [here](#) for the Mathematica codes verifying the duality at any finite orders.

be constructed within the effective field theory of inflation (EFToI), as in [7]. The precise form of these interactions will play little role for us, however we take one vertex to be PO and the other to be PE such that we realise a PO trispectrum. Since cubic interactions of $SO(3)$ scalars cannot violate parity, only the transverse modes of the massive field will contribute. Diagrammatically, we therefore have [92]

$$B_4 = \text{[Diagram 1]} + \text{[Diagram 2]} + \text{[Diagram 3]} + \text{[Diagram 4]} \quad (\text{B1})$$

The external propagators are given by

$$K_+^\pi = \pi(\eta_0, k)\pi^*(\eta, k), \quad (\text{B2})$$

$$K_-^\pi = (K_+^\pi)^*, \quad (\text{B3})$$

while the internal ones are given by

$$G_{++}^\sigma = \sigma(\eta, k)\sigma^*(\eta', k)\theta(\eta - \eta') + (\eta \leftrightarrow \eta'), \quad (\text{B4})$$

$$G_{+-}^\sigma = \sigma^*(\eta, k)\sigma(\eta', k), \quad (\text{B5})$$

$$G_{--}^\sigma = (G_{++}^\sigma)^*, \quad (\text{B6})$$

$$G_{-+}^\sigma = (G_{+-}^\sigma)^*, \quad (\text{B7})$$

where a $+/-$ indicates that a propagator is connected to a black/white vertex, and η_0 indicates the time at which the correlator is evaluated and we will take $\eta_0 \rightarrow 0$.¹¹ Concentrating on the transverse modes only, for the massive spin-1 field we write $\sigma_i^\pm = \sigma\epsilon_i^\pm$ where ϵ_i^\pm are the transverse polarisation vectors and σ is the mode function which is given by [7]

$$\sigma(k, \eta) = \sqrt{\frac{\pi}{2}} \frac{1}{\sqrt{2k}} e^{\frac{i\pi}{4} + \frac{i\pi\nu}{2}} (-k\eta)^{1/2} H_\nu^{(1)}(-k\eta), \quad (\text{B8})$$

with

$$\nu = \sqrt{\frac{1}{4} - \frac{m^2}{H^2}}. \quad (\text{B9})$$

Here $H_\nu^{(1)}$ is the Hankel function of the first kind. As always, in (B1) we sum over the two helicity modes in each exchange diagram. Now, recall that black and white vertices are related by complex conjugation *and* a flip of all momenta at that vertex. Since we are interested in the PO sector, flipping the sign of all momenta yields a minus sign so we can write

$$B_4^{\text{PO}} = \left(\text{[Diagram 1]} + \text{[Diagram 2]} \right) - \text{c.c.}, \quad (\text{B10})$$

which manifestly shows that PO correlators are purely imaginary. In [78], the Feynman propagator G was written as the sum of two new propagators $G = C + F$ with all time-orderings contained within C (the connected part indicated by double lines), and with the role of F (the factorised part indicated by dashed lines) to ensure that C is *purely real* once both time variables are rotated clockwise by 90° in the complex plane. For fields in the complementary series, which will be our focus here and requires $m^2 < H^2/4$, C in fact becomes the bulk-bulk propagator of the wavefunction formalism (for principle series fields the situation is different [78]). We therefore have

$$C_{++}^\sigma = G_{++}^\sigma - \frac{\sigma(\eta_0, k)}{\sigma^*(\eta_0, k)} \sigma^*(\eta, k)\sigma^*(\eta', k), \quad (\text{B11})$$

$$F_{++}^\sigma = \frac{\sigma(\eta_0, k)}{\sigma^*(\eta_0, k)} \sigma^*(\eta, k)\sigma^*(\eta', k). \quad (\text{B12})$$

With this decomposition, (B10) becomes

$$B_4^{\text{PO}} = \left(\text{[Diagram 1]} + \text{[Diagram 2]} + \text{[Diagram 3]} \right) - \text{c.c.} \quad (\text{B13})$$

By construction, C_{++} is purely real after a Wick rotation of both time variables to Euclidean time, while the other components that enter in the Feynman rules that compute the first diagram in (B13) are also real after the same rotation [78]:

$$\text{Integration measures : } i \int \frac{d\eta}{\eta^4}, \quad i \int \frac{d\eta'}{\eta'^4}, \quad (\text{B14})$$

$$\text{Derivatives : } \eta \partial_\eta, \quad i\eta \mathbf{k}, \quad (\text{B15})$$

$$\text{External propagator : } K_+^\pi = \frac{H^2}{2k^3} (1 - ik\eta) e^{ik\eta}, \quad (\text{B16})$$

$$\text{Polarisations : } \sum_{h=\pm 1} \epsilon_i^{(h)}(\mathbf{k}) \epsilon_j^{(h)*}(\mathbf{k}) = \delta_{ij} - \hat{k}_i \hat{k}_j. \quad (\text{B17})$$

Note that the derivatives take this form due to scale invariance. If the time integrals are convergent,¹² we can close the contour, drop the arc at infinity thanks to the Bunch-Davies vacuum choice, and use the Wick rotation to compute the necessary integrals. Since everything is manifestly real in Euclidean time, the diagrams with C propagators cancel. This is a summary of one of the main results of [78]. We therefore have

$$B_4^{\text{PO}} = \left(\text{[Diagram 1]} + \text{[Diagram 2]} \right) - \text{c.c.} \quad (\text{B18})$$

¹¹ Note that the form of these propagators holds for any field. For example, if the massive field goes to the boundary we will have $K_+^\pi = \sigma(\eta_0, k)\sigma^*(\eta, k)$.

¹² This is guaranteed if we work within the EFToI since the non-linear symmetries of the inflaton dictate a minimal number of derivatives that is sufficient to cancel the would-be divergences arising from the integration measure [78].

We now note that the F_{++}^σ and G_{+-}^σ propagators can be written as

$$F_{++}^\sigma = \frac{1}{B_2^\sigma(k)} K_+^\sigma(\eta, k) K_+^\sigma(\eta', k), \quad (\text{B19})$$

$$G_{+-}^\sigma = \frac{1}{B_2^\sigma(k)} K_+^\sigma(\eta, k) K_-^\sigma(\eta', k), \quad (\text{B20})$$

where B_2^σ is the power spectrum of σ . It follows that (B18) can be expressed as

$$\begin{aligned} B_4^{\text{PO}} &= \frac{1}{\text{diagram}} \left(\text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{diagram 4} \right) \\ &= \left(\text{diagram 5} + \text{diagram 6} \right) \frac{1}{\text{diagram 7}} \left(\text{diagram 8} + \text{diagram 9} \right) \\ &= \left[B_3 \cdot \frac{1}{B_2} \cdot B_3 \right]^{\text{PO}}, \\ &= \sum_{h=\pm 1} \left[B_3^h \frac{1}{B_2^h} B_3^h \right]^{\text{PO}}, \end{aligned} \quad (\text{B21})$$

where in the final line we have converted to the helicity basis with B_3^h the three-point function with two curvature perturbations and the helicity- h mode of the spin-1 field. We have also used that in the SK formalism three-point functions for $\pi\pi\sigma$ are computed via [92]

$$B_3 = \text{diagram 10} + \text{diagram 11}. \quad (\text{B22})$$

We have therefore recovered the result we derived in the main text (there is no factor of 3 here since we have worked with only one of the three channels). For principle series fields we can arrive at an equation similar to (B18), but the two diagrams do not have the same factors and therefore the RHS is not simply a combination of correlators.

-
- [1] X. Chen, M. H. Namjoo, and Y. Wang, Quantum Primordial Standard Clocks, *JCAP* **02**, 013, [arXiv:1509.03930 \[astro-ph.CO\]](#).
- [2] Y. Wang, Z. Wang, and Y. Zhu, Non-standard primordial clocks from induced mass in alternative to inflation scenarios, *JCAP* **11**, 026, [arXiv:2007.09677 \[hep-th\]](#).
- [3] X. Chen and Y. Wang, Quasi-Single Field Inflation and Non-Gaussianities, *JCAP* **04**, 027, [arXiv:0911.3380 \[hep-th\]](#).
- [4] D. Baumann and D. Green, Signatures of Supersymmetry from the Early Universe, *Phys. Rev. D* **85**, 103520 (2012), [arXiv:1109.0292 \[hep-th\]](#).
- [5] T. Noumi, M. Yamaguchi, and D. Yokoyama, Effective field theory approach to quasi-single field inflation and effects of heavy fields, *JHEP* **06**, 051, [arXiv:1211.1624 \[hep-th\]](#).
- [6] N. Arkani-Hamed and J. Maldacena, Cosmological Collider Physics, (2015), [arXiv:1503.08043 \[hep-th\]](#).
- [7] H. Lee, D. Baumann, and G. L. Pimentel, Non-Gaussianity as a Particle Detector, *JHEP* **12**, 040, [arXiv:1607.03735 \[hep-th\]](#).
- [8] S. Jazayeri and S. Renaux-Petel, Cosmological bootstrap in slow motion, *JHEP* **12**, 137, [arXiv:2205.10340 \[hep-th\]](#).
- [9] G. L. Pimentel and D.-G. Wang, Boostless cosmological collider bootstrap, *JHEP* **10**, 177, [arXiv:2205.00013 \[hep-th\]](#).
- [10] G. Cabass, O. H. E. Philcox, M. M. Ivanov, K. Akitsu, S.-F. Chen, M. Simonović, and M. Zaldarriaga, BOSS Constraints on Massive Particles during Inflation: The Cosmological Collider in Action, (2024), [arXiv:2404.01894 \[astro-ph.CO\]](#).
- [11] P. Chakraborty and J. Stout, Light scalars at the cosmological collider, *JHEP* **02**, 021, [arXiv:2310.01494 \[hep-th\]](#).
- [12] X. Chen, R. Ebadi, and S. Kumar, Classical cosmological collider physics and primordial features, *JCAP* **08**, 083, [arXiv:2205.01107 \[hep-ph\]](#).
- [13] X. Tong and Z.-Z. Xianyu, Large spin-2 signals at the cosmological collider, *JHEP* **10**, 194, [arXiv:2203.06349 \[hep-ph\]](#).
- [14] M. Reece, L.-T. Wang, and Z.-Z. Xianyu, Large-field inflation and the cosmological collider, *Phys. Rev. D* **107**, L101304 (2023), [arXiv:2204.11869 \[hep-ph\]](#).
- [15] L.-T. Wang and Z.-Z. Xianyu, Gauge Boson Signals at the Cosmological Collider, *JHEP* **11**, 082, [arXiv:2004.02887 \[hep-ph\]](#).
- [16] S. Kumar and R. Sundrum, Cosmological Collider Physics and the Curvaton, *JHEP* **04**, 077, [arXiv:1908.11378 \[hep-ph\]](#).
- [17] W. Sohn, D.-G. Wang, J. R. Fergusson, and E. P. S. Shellard, Searching for Cosmological Collider in the Planck CMB Data, (2024), [arXiv:2404.07203 \[astro-ph.CO\]](#).
- [18] C. McCulloch, E. Pajer, and X. Tong, A Cosmological Tachyon Collider: Enhancing the Long-Short Scale Coupling, (2024), [arXiv:2401.11009 \[hep-th\]](#).
- [19] N. Craig, S. Kumar, and A. McCune, An Effective Cosmological Collider, (2024), [arXiv:2401.10976 \[hep-ph\]](#).
- [20] S. Jazayeri, S. Renaux-Petel, and D. Werth, Shapes of the cosmological low-speed collider, *JCAP* **12**, 035, [arXiv:2307.01751 \[hep-th\]](#).

- [21] Z. Qin and Z.-Z. Xianyu, Phase information in cosmological collider signals, *JHEP* **10**, 192, [arXiv:2205.01692 \[hep-th\]](#).
- [22] A. Bodas, S. Kumar, and R. Sundrum, The Scalar Chemical Potential in Cosmological Collider Physics, *JHEP* **02**, 079, [arXiv:2010.04727 \[hep-ph\]](#).
- [23] L.-T. Wang and Z.-Z. Xianyu, In Search of Large Signals at the Cosmological Collider, *JHEP* **02**, 044, [arXiv:1910.12876 \[hep-ph\]](#).
- [24] P. D. Meerburg, M. Münchmeyer, J. B. Muñoz, and X. Chen, Prospects for Cosmological Collider Physics, *JCAP* **03**, 050, [arXiv:1610.06559 \[astro-ph.CO\]](#).
- [25] X. Chen, Y. Wang, and Z.-Z. Xianyu, Standard Model Background of the Cosmological Collider, *Phys. Rev. Lett.* **118**, 261302 (2017), [arXiv:1610.06597 \[hep-th\]](#).
- [26] L. Bordin, P. Creminelli, A. Khmelnitsky, and L. Senatore, Light Particles with Spin in Inflation, *JCAP* **10**, 013, [arXiv:1806.10587 \[hep-th\]](#).
- [27] Z.-Z. Xianyu and J. Zang, Inflation correlators with multiple massive exchanges, *JHEP* **03**, 070, [arXiv:2309.10849 \[hep-th\]](#).
- [28] S. Aoki, L. Pinol, F. Sano, M. Yamaguchi, and Y. Zhu, Cosmological Correlators with Double Massive Exchanges: Bootstrap Equation and Phenomenology, (2024), [arXiv:2404.09547 \[hep-th\]](#).
- [29] P. Chakraborty and J. Stout, Compact scalars at the cosmological collider, *JHEP* **03**, 149, [arXiv:2311.09219 \[hep-th\]](#).
- [30] X. Chen, J. Fan, and L. Li, New inflationary probes of axion dark matter, *JHEP* **12**, 197, [arXiv:2303.03406 \[hep-ph\]](#).
- [31] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan, and L. Senatore, The Effective Field Theory of Inflation, *JHEP* **03**, 014, [arXiv:0709.0293 \[hep-th\]](#).
- [32] X. Chen, M.-x. Huang, S. Kachru, and G. Shiu, Observational signatures and non-Gaussianities of general single field inflation, *JCAP* **01**, 002, [arXiv:hep-th/0605045](#).
- [33] D. Baumann and D. Green, Equilateral Non-Gaussianity and New Physics on the Horizon, *JCAP* **09**, 014, [arXiv:1102.5343 \[hep-th\]](#).
- [34] P. Creminelli, On non-Gaussianities in single-field inflation, *JCAP* **10**, 003, [arXiv:astro-ph/0306122](#).
- [35] D. Seery and J. E. Lidsey, Primordial non-Gaussianities in single field inflation, *JCAP* **06**, 003, [arXiv:astro-ph/0503692](#).
- [36] R. Holman and A. J. Tolley, Enhanced Non-Gaussianity from Excited Initial States, *JCAP* **05**, 001, [arXiv:0710.1302 \[hep-th\]](#).
- [37] P. D. Meerburg, J. P. van der Schaar, and P. S. Corasaniti, Signatures of Initial State Modifications on Bispectrum Statistics, *JCAP* **05**, 018, [arXiv:0901.4044 \[hep-th\]](#).
- [38] J. Ganc, Calculating the local-type fNL for slow-roll inflation with a non-vacuum initial state, *Phys. Rev. D* **84**, 063514 (2011), [arXiv:1104.0244 \[astro-ph.CO\]](#).
- [39] R. Flauger, D. Green, and R. A. Porto, On squeezed limits in single-field inflation. Part I, *JCAP* **08**, 032, [arXiv:1303.1430 \[hep-th\]](#).
- [40] D. Green and R. A. Porto, Signals of a Quantum Universe, *Phys. Rev. Lett.* **124**, 251302 (2020), [arXiv:2001.09149 \[hep-th\]](#).
- [41] S. A. Salcedo, T. Colas, and E. Pajer, The Open Effective Field Theory of Inflation, (2024), [arXiv:2404.15416 \[hep-th\]](#).
- [42] J. M. Maldacena, Non-Gaussian features of primordial fluctuations in single field inflationary models, *JHEP* **05**, 013, [arXiv:astro-ph/0210603](#).
- [43] D. Anninos, T. Anous, D. Z. Freedman, and G. Konstantinidis, Late-time Structure of the Bunch-Davies De Sitter Wavefunction, *JCAP* **11**, 048, [arXiv:1406.5490 \[hep-th\]](#).
- [44] N. Arkani-Hamed, D. Baumann, H. Lee, and G. L. Pimentel, The Cosmological Bootstrap: Inflationary Correlators from Symmetries and Singularities, *JHEP* **04**, 105, [arXiv:1811.00024 \[hep-th\]](#).
- [45] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee, and G. L. Pimentel, The cosmological bootstrap: weight-shifting operators and scalar seeds, *JHEP* **12**, 204, [arXiv:1910.14051 \[hep-th\]](#).
- [46] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee, and G. L. Pimentel, The Cosmological Bootstrap: Spinning Correlators from Symmetries and Factorization, *SciPost Phys.* **11**, 071 (2021), [arXiv:2005.04234 \[hep-th\]](#).
- [47] D. Baumann, W.-M. Chen, C. Duaso Pueyo, A. Joyce, H. Lee, and G. L. Pimentel, Linking the singularities of cosmological correlators, *JHEP* **09**, 010, [arXiv:2106.05294 \[hep-th\]](#).
- [48] S. Albayrak, P. Benincasa, and C. Duaso Pueyo, Perturbative Unitarity and the Wavefunction of the Universe, (2023), [arXiv:2305.19686 \[hep-th\]](#).
- [49] N. Arkani-Hamed, P. Benincasa, and A. Postnikov, Cosmological Polytopes and the Wavefunction of the Universe, (2017), [arXiv:1709.02813 \[hep-th\]](#).
- [50] P. Benincasa, Wavefunctionals/S-matrix techniques in de Sitter, *PoS CORFU2021*, 358 (2022), [arXiv:2203.16378 \[hep-th\]](#).
- [51] H. Goodhew, S. Jazayeri, and E. Pajer, The Cosmological Optical Theorem, *JCAP* **04**, 021, [arXiv:2009.02898 \[hep-th\]](#).
- [52] S. Melville and E. Pajer, Cosmological Cutting Rules, *JHEP* **05**, 249, [arXiv:2103.09832 \[hep-th\]](#).
- [53] H. Goodhew, S. Jazayeri, M. H. G. Lee, and E. Pajer, Cutting cosmological correlators, *JCAP* **08**, 003, [arXiv:2104.06587 \[hep-th\]](#).
- [54] E. Pajer, Building a Boostless Bootstrap for the Bispectrum, *JCAP* **01**, 023, [arXiv:2010.12818 \[hep-th\]](#).
- [55] S. Jazayeri, E. Pajer, and D. Stefanyzyn, From locality and unitarity to cosmological correlators, *JHEP* **10**, 065, [arXiv:2103.08649 \[hep-th\]](#).
- [56] S. A. Salcedo, M. H. G. Lee, S. Melville, and E. Pajer, The Analytic Wavefunction, *JHEP* **06**, 020, [arXiv:2212.08009 \[hep-th\]](#).
- [57] D. Meltzer, The inflationary wavefunction from analyticity and factorization, *JCAP* **12** (12), 018, [arXiv:2107.10266 \[hep-th\]](#).
- [58] Y. Donath and E. Pajer, The In-Out Formalism for In-In Correlators, (2024), [arXiv:2402.05999 \[hep-th\]](#).
- [59] M. H. G. Lee, From amplitudes to analytic wavefunctions, *JHEP* **03**, 058, [arXiv:2310.01525 \[hep-th\]](#).
- [60] S. Agui-Salcedo and S. Melville, The cosmological tree theorem, *JHEP* **12**, 076, [arXiv:2308.00680 \[hep-th\]](#).
- [61] S. Céspedes, A.-C. Davis, and S. Melville, On the time evolution of cosmological correlators, *JHEP* **02**, 012, [arXiv:2009.07874 \[hep-th\]](#).
- [62] D. Baumann, D. Green, A. Joyce, E. Pajer, G. L. Pimentel, C. Sleight, and M. Taronna, Snowmass White Paper: The Cosmological Bootstrap, in *Snowmass 2021* (2022) [arXiv:2203.08121 \[hep-th\]](#).
- [63] L. Di Pietro, V. Gorbenko, and S. Komatsu, Analyticity

- and unitarity for cosmological correlators, *JHEP* **03**, 023, [arXiv:2108.01695 \[hep-th\]](#).
- [64] M. Hogervorst, J. a. Penedones, and K. S. Vaziri, Towards the non-perturbative cosmological bootstrap, *JHEP* **02**, 162, [arXiv:2107.13871 \[hep-th\]](#).
- [65] S. Melville and G. L. Pimentel, A de Sitter S -matrix for the masses, (2023), [arXiv:2309.07092 \[hep-th\]](#).
- [66] S. Melville and G. L. Pimentel, A de Sitter S -matrix from amputated cosmological correlators, (2024), [arXiv:2404.05712 \[hep-th\]](#).
- [67] B. Fan and Z.-Z. Xianyu, Cosmological Amplitudes in Power-Law FRW Universe, (2024), [arXiv:2403.07050 \[hep-th\]](#).
- [68] H. Gomez, R. Lipinski Jusinkas, and A. Lipstein, Cosmological scattering equations at tree-level and one-loop, *JHEP* **07**, 004, [arXiv:2112.12695 \[hep-th\]](#).
- [69] S. Céspedes, A.-C. Davis, and D.-G. Wang, On the IR divergences in de Sitter space: loops, resummation and the semi-classical wavefunction, *JHEP* **04**, 004, [arXiv:2311.17990 \[hep-th\]](#).
- [70] A. Bzowski, P. McFadden, and K. Skenderis, Renormalisation of IR divergences and holography in de Sitter, (2023), [arXiv:2312.17316 \[hep-th\]](#).
- [71] V. Gorbenko and L. Senatore, $\lambda\phi^4$ in dS, (2019), [arXiv:1911.00022 \[hep-th\]](#).
- [72] J. M. Maldacena and G. L. Pimentel, On graviton non-Gaussianities during inflation, *JHEP* **09**, 045, [arXiv:1104.2846 \[hep-th\]](#).
- [73] S. Raju, New Recursion Relations and a Flat Space Limit for AdS/CFT Correlators, *Phys. Rev. D* **85**, 126009 (2012), [arXiv:1201.6449 \[hep-th\]](#).
- [74] E. Pajer, D. Stefnyszyn, and J. Supel, The Boostless Bootstrap: Amplitudes without Lorentz boosts, *JHEP* **12**, 198, [Erratum: *JHEP* **04**, 023 (2022)], [arXiv:2007.00027 \[hep-th\]](#).
- [75] P. Benincasa and F. Vazão, The Asymptotic Structure of Cosmological Integrals, (2024), [arXiv:2402.06558 \[hep-th\]](#).
- [76] P. Benincasa and G. Dian, The Geometry of Cosmological Correlators, (2024), [arXiv:2401.05207 \[hep-th\]](#).
- [77] N. Arkani-Hamed, D. Baumann, A. Hillman, A. Joyce, H. Lee, and G. L. Pimentel, Differential Equations for Cosmological Correlators, (2023), [arXiv:2312.05303 \[hep-th\]](#).
- [78] D. Stefnyszyn, X. Tong, and Y. Zhu, Cosmological Correlators Through the Looking Glass: Reality, Parity, and Factorisation, (2023), [arXiv:2309.07769 \[hep-th\]](#).
- [79] B. S. DeWitt, Quantum Theory of Gravity. 2. The Manifestly Covariant Theory, *Phys. Rev.* **162**, 1195 (1967).
- [80] M. H. G. Lee, C. McCulloch, and E. Pajer, Leading loops in cosmological correlators, *JHEP* **11**, 038, [arXiv:2305.11228 \[hep-th\]](#).
- [81] C. Sleight and M. Taronna, From AdS to dS exchanges: Spectral representation, Mellin amplitudes, and crossing, *Phys. Rev. D* **104**, L081902 (2021), [arXiv:2007.09993 \[hep-th\]](#).
- [82] C. Sleight and M. Taronna, From dS to AdS and back, *JHEP* **12**, 074, [arXiv:2109.02725 \[hep-th\]](#).
- [83] T. Liu, X. Tong, Y. Wang, and Z.-Z. Xianyu, Probing P and CP Violations on the Cosmological Collider, *JHEP* **04**, 189, [arXiv:1909.01819 \[hep-ph\]](#).
- [84] G. Cabass, S. Jazayeri, E. Pajer, and D. Stefnyszyn, Parity violation in the scalar trispectrum: no-go theorems and yes-go examples, *JHEP* **02**, 021, [arXiv:2210.02907 \[hep-th\]](#).
- [85] J. Hou, Z. Slepian, and R. N. Cahn, Measurement of parity-odd modes in the large-scale 4-point correlation function of Sloan Digital Sky Survey Baryon Oscillation Spectroscopic Survey twelfth data release CMASS and LOWZ galaxies, *Mon. Not. Roy. Astron. Soc.* **522**, 5701 (2023), [arXiv:2206.03625 \[astro-ph.CO\]](#).
- [86] O. H. E. Philcox, Probing parity violation with the four-point correlation function of BOSS galaxies, *Phys. Rev. D* **106**, 063501 (2022), [arXiv:2206.04227 \[astro-ph.CO\]](#).
- [87] O. H. E. Philcox and M. Shiraishi, Testing Parity Symmetry with the Polarized Cosmic Microwave Background, (2023), [arXiv:2308.03831 \[astro-ph.CO\]](#).
- [88] P. Paul, C. Clarkson, and R. Maartens, Parity violation in the observed galaxy trispectrum, (2024), [arXiv:2402.16478 \[astro-ph.CO\]](#).
- [89] C. Creque-Sarbinowski, S. Alexander, M. Kamionkowski, and O. Philcox, Parity-violating trispectrum from Chern-Simons gravity, *JCAP* **11**, 029, [arXiv:2303.04815 \[astro-ph.CO\]](#).
- [90] A. Maleknejad, Axion Inflation with an $SU(2)$ Gauge Field: Detectable Chiral Gravity Waves, *JHEP* **07**, 104, [arXiv:1604.03327 \[hep-ph\]](#).
- [91] A. Maleknejad and E. Komatsu, Production and Backreaction of Spin-2 Particles of $SU(2)$ Gauge Field during Inflation, *JHEP* **05**, 174, [arXiv:1808.09076 \[hep-ph\]](#).
- [92] X. Chen, Y. Wang, and Z.-Z. Xianyu, Schwinger-Keldysh Diagrammatics for Primordial Perturbations, *JCAP* **12**, 006, [arXiv:1703.10166 \[hep-th\]](#).
- [93] M. Shiraishi, D. Nitta, and S. Yokoyama, Parity Violation of Gravitons in the CMB Bispectrum, *Prog. Theor. Phys.* **126**, 937 (2011), [arXiv:1108.0175 \[astro-ph.CO\]](#).
- [94] P. Creminelli, J. Gleyzes, J. Noreña, and F. Vernizzi, Resilience of the standard predictions for primordial tensor modes, *Phys. Rev. Lett.* **113**, 231301 (2014), [arXiv:1407.8439 \[astro-ph.CO\]](#).
- [95] G. Cabass, E. Pajer, D. Stefnyszyn, and J. Supel, Bootstrapping large graviton non-Gaussianities, *JHEP* **05**, 077, [arXiv:2109.10189 \[hep-th\]](#).
- [96] L. Bordin and G. Cabass, Graviton non-Gaussianities and Parity Violation in the EFT of Inflation, *JCAP* **07**, 014, [arXiv:2004.00619 \[astro-ph.CO\]](#).
- [97] N. Bartolo and G. Orlando, Parity breaking signatures from a Chern-Simons coupling during inflation: the case of non-Gaussian gravitational waves, *JCAP* **07**, 034, [arXiv:1706.04627 \[astro-ph.CO\]](#).