

COMBINATORIAL PROOFS OF INEQUALITIES INVOLVING THE NUMBER OF PARTITIONS WITH PARTS SEPARATED BY PARITY

CRISTINA BALLANTINE AND AMANDA WELCH

ABSTRACT. We consider the number of various partitions of n with parts separated by parity and prove combinatorially several inequalities between these numbers. For example, we show that for $n \geq 5$ we have $p_{\text{od}}^{\text{eu}}(n) < p_{\text{ed}}^{\text{ou}}(n)$, where $p_{\text{od}}^{\text{eu}}(n)$ is the number of partitions of n with odd parts distinct and even parts unrestricted and all odd parts less than all even parts and $p_{\text{ed}}^{\text{ou}}(n)$ is the number of partitions of n with even parts distinct and odd parts unrestricted and all even parts less than all odd parts. We also prove a conjectural inequality of Fu and Tang involving partitions with parts separated by parity with restrictions on the multiplicity of parts.

Keywords: partitions with parts separated by parity, combinatorial injections.
MSC 2020: 05A17, 05A20, 11P81

1. INTRODUCTION

A partition of a non-negative integer n is a non-increasing sequence of positive integers that sum to n . We write a partition of n as $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell$ and $\lambda_1 + \lambda_2 + \dots + \lambda_\ell = n$. We refer to the terms λ_i as the parts of λ .

In this article, we consider the number of partitions of n with parts separated by parity. These partitions were introduced by Andrews in [3, 2]. Because the numbers of partitions of n with parts separated by parity (in different combinations) display modular properties, several authors have studied different aspects of their behavior, see for example [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]. Following the notation in [3], we denote by $\mathcal{P}_{yz}^{wx}(n)$ the set of partitions of n with parts of type w satisfying condition x or of type y satisfying condition z and all parts of type w satisfying condition x are larger than all parts of type y satisfying condition z . Here, the symbols w and y satisfy $w \neq y$ and are e (for even) or o (for odd). The symbols x and y are u (for unrestricted) or d (for distinct). For example, $\mathcal{P}_{\text{ou}}^{\text{ed}}(n)$ is the set of partitions of n in which all even parts are greater than all odd parts, odd parts are unrestricted and even parts are distinct. We set $p_{yz}^{wx}(n) := |\mathcal{P}_{yz}^{wx}(n)|$.

In this article, we allow partitions in $\mathcal{P}_{yz}^{wx}(n)$ to have no parts of type y or no parts of type w .

Example 1. Let $n = 6$. Then

$$\mathcal{P}_{\text{ou}}^{\text{ed}}(6) = \{(6), (5, 1), (4, 2), (4, 1^2), (3^2), (3, 1^3), (2, 1^5), (1^7)\}.$$

Thus, $p_{\text{ou}}^{\text{ed}}(6) = 8$.

Bringmann, Craig and Nazaroglu [4] studied the asymptotic behavior of the eight sequences $p_{yz}^{wx}(n)$ (for all possible combinations of w, x, y, z). At the end of their

article they conclude that for sufficiently large n ,

$$(1) \quad p_{\text{ed}}^{\text{od}}(n) < p_{\text{od}}^{\text{ed}}(n) < p_{\text{od}}^{\text{eu}}(n) < p_{\text{eu}}^{\text{od}}(n) < p_{\text{ed}}^{\text{ou}}(n) < p_{\text{eu}}^{\text{ou}}(n) < p_{\text{ou}}^{\text{ed}}(n) < p_{\text{ou}}^{\text{eu}}(n).$$

The second, fifth, and last inequalities are immediate since $\mathcal{P}_{\text{od}}^{\text{ed}}(n) \subseteq \mathcal{P}_{\text{od}}^{\text{eu}}(n)$, $\mathcal{P}_{\text{ed}}^{\text{ou}}(n) \subseteq \mathcal{P}_{\text{eu}}^{\text{ou}}(n)$ and $\mathcal{P}_{\text{ou}}^{\text{ed}}(n) \subseteq \mathcal{P}_{\text{ou}}^{\text{eu}}(n)$. Bringmann, Craig and Nazarov asked for combinatorial proofs of the other inequalities.

In section 3, we give combinatorial injections proving the following inequalities:

$$(2) \quad p_{\text{ed}}^{\text{od}}(n) < p_{\text{od}}^{\text{ed}}(n), \quad \text{for } n \geq 11;$$

$$(3) \quad p_{\text{eu}}^{\text{ou}}(n) < p_{\text{ou}}^{\text{eu}}(n), \quad \text{for } n \geq 3;$$

$$(4) \quad p_{\text{od}}^{\text{eu}}(n) < p_{\text{ed}}^{\text{ou}}(n), \quad \text{for } n \geq 5;$$

$$(5) \quad p_{\text{eu}}^{\text{od}}(n) < p_{\text{ou}}^{\text{ed}}(n), \quad \text{for } n \geq 2;$$

$$(6) \quad p_{\text{od}}^{\text{ed}}(n) < p_{\text{eu}}^{\text{od}}(n), \quad \text{for } n \geq 8.$$

In [11], Fu and Tang considered partitions with parts separated by parity with additional conditions on the multiplicity of parts. We adapt their notation to match the notation of this article. Let $\overline{\mathcal{P}}_{\text{eu}}^{\text{ou}}(n)$ be the subset of partitions in $\mathcal{P}_{\text{eu}}^{\text{ou}}(n)$ in which the largest even part appears with odd multiplicity and all other parts appear with even multiplicity. Note that a partition of n with only odd parts is considered to be in $\overline{\mathcal{P}}_{\text{eu}}^{\text{ou}}(n)$ with a single part equal to 0 (and multiplicity one). We set $\overline{p}_{\text{eu}}^{\text{ou}}(n) := |\overline{\mathcal{P}}_{\text{eu}}^{\text{ou}}(n)|$. Similarly, let $\overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(n)$ be the subset of partitions in $\mathcal{P}_{\text{ou}}^{\text{eu}}(n)$ in which both even and odd parts occur, the largest even part and the largest odd part each appear with odd multiplicity and all other parts appear with even multiplicity. We set $\overline{p}_{\text{ou}}^{\text{eu}}(n) := |\overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(n)|$. Fu and Tang made the following conjecture.

Conjecture 1. *Let $n \geq 3$. Then,*

$$\overline{p}_{\text{eu}}^{\text{ou}}(2n) < \overline{p}_{\text{ou}}^{\text{eu}}(2n + 1).$$

We prove the conjecture in section 4.

2. DEFINITIONS AND NOTATION

Throughout the article we use the following notation. Given a partition λ , the length of λ , denoted $\ell(\lambda)$, is the number of parts in λ . We denote by λ^o , respectively λ^e , the partition whose parts are precisely the odd, respectively even, parts of λ . We denote by $\ell_o(\lambda)$, respectively $\ell_e(\lambda)$, the number of odd, respectively even, parts in λ , and when λ contains both even and odd parts, $\ell_m(\lambda)$ denotes the minimum of $\ell_o(\lambda)$ and $\ell_e(\lambda)$. Moreover, λ_s^o , respectively λ_s^e , denotes the last part in λ^o , respectively λ^e . By λ_1^o , respectively λ_1^e , we mean the first part in λ^o , respectively λ^e . If j is a part of λ , we denote by $a_{>j}$ the smallest part in λ that is larger than j , if such a part exists. We write $m_\lambda(j)$ for the multiplicity of j in λ , i.e., the number of times j occurs as a part in λ . We also write $\ell_{>j}(\lambda)$, respectively $\ell_{\geq j}(\lambda)$, for the number of parts greater than, respectively greater than or equal to, j in λ . Occasionally, we use the frequency notation for partitions and write $\lambda = (i_1^{m_1}, i_2^{m_2}, \dots, i_k^{m_k})$, where $m_j = m_\lambda(i_j)$. For convenience, if $j > \ell(\lambda)$, we set $\lambda_j = 0$. The set of partitions of n with distinct parts is denoted by $\mathcal{D}(n)$.


If $\mathcal{A}(n)$ is the set of partitions of n satisfying certain properties, we define

$$\mathcal{A} := \bigcup_{n \geq 0} \mathcal{A}(n).$$

We identify partitions with their multiset of parts. If a is a positive integer, we write $a \in \lambda$ to mean that a is a part of λ . If λ and μ are partitions, by $\lambda \cup \mu$ and $\lambda \setminus \mu$ we mean the obvious operations on the multisets of parts of λ and μ . Moreover, $\lambda \setminus \mu$ is only defined if $\mu \subseteq \lambda$ as multisets.

The Ferrers diagram of a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)})$ is an array of left justified boxes such that the i th row from the top contains λ_i boxes. We abuse notation and use λ to mean a partition or its Ferrers diagram.

Example 2. The Ferrers diagram of $\lambda = (5, 4, 3, 3, 2)$ is shown below.



3. COMBINATORIAL PROOFS OF INEQUALITIES (2) – (6)

3.1. The inequality $p_{\text{ed}}^{\text{od}}(n) < p_{\text{od}}^{\text{ed}}(n)$. Let $n \geq 0$. We create an injection

$$\varphi_1 : \mathcal{P}_{\text{ed}}^{\text{od}}(n) \rightarrow \mathcal{P}_{\text{od}}^{\text{ed}}(n).$$

Let $\lambda \in \mathcal{P}_{\text{ed}}^{\text{od}}(n)$. If $\lambda^o = \emptyset$ or $\lambda^e = \emptyset$, define $\varphi_1(\lambda) := \lambda$. If $\lambda^e, \lambda^o \neq \emptyset$, define $\varphi_1(\lambda)$ to be the partition obtained from λ by subtracting one from each of the last $\ell_m(\lambda)$ parts of λ and adding one to each of the first $\ell_m(\lambda)$ parts of λ . In each case, the partition $\varphi_1(\lambda) \in \mathcal{P}_{\text{od}}^{\text{ed}}(n)$. Moreover, if $\lambda^o, \lambda^e \neq \emptyset$, the smallest even part of $\varphi_1(\lambda)$ differs by at least three from the largest odd part of $\varphi_1(\lambda)$.

If we denote by $\tilde{\mathcal{P}}_{\text{od}}^{\text{ed}}(n)$ the subset of partitions $\mu \in \mathcal{P}_{\text{od}}^{\text{ed}}(n)$ with $\mu^o = \emptyset$ or $\mu^e = \emptyset$, or $\mu^o, \mu^e \neq \emptyset$ and $\mu_s^e - \mu_1^o \geq 3$, then $\varphi_1 : \mathcal{P}_{\text{ed}}^{\text{od}}(n) \rightarrow \tilde{\mathcal{P}}_{\text{od}}^{\text{ed}}(n)$ is a bijection. To see that φ_1 is invertible, let $\mu \in \tilde{\mathcal{P}}_{\text{od}}^{\text{ed}}(n)$. If $\mu^o = \emptyset$ or $\mu^e = \emptyset$, then $\varphi_1^{-1}(\mu) = \mu$. If $\mu^o, \mu^e \neq \emptyset$, then $\varphi_1^{-1}(\mu)$ is the partition obtained from μ by subtracting one from each of the first $\ell_m(\mu)$ parts of μ and adding one to each of the last $\ell_m(\mu)$ parts of μ .

Example 3. Let $n = 35$ and $\lambda = (17, 12, 6)$. Then $\ell_m(\lambda) = \ell_o(\lambda) = 1$. Thus, $\varphi_1(\lambda) = (18, 12, 5)$.

We have the following interpretation of $p_{\text{od}}^{\text{ed}}(n) - p_{\text{ed}}^{\text{od}}(n)$.

Proposition 1. *Let $n \geq 0$. Then, $p_{\text{od}}^{\text{ed}}(n) - p_{\text{ed}}^{\text{od}}(n)$ equals the number of partitions $\lambda \in \mathcal{P}_{\text{od}}^{\text{ed}}(n)$ with $\lambda^o, \lambda^e \neq \emptyset$ and $\lambda_s^e - \lambda_1^o = 1$.*

One can verify directly that there is no partition $\lambda \in \mathcal{P}_{\text{od}}^{\text{ed}}(10)$ with both even and odd parts such that $\lambda_s^e - \lambda_1^o = 1$. If $n \geq 11$, there is a partition $\lambda \in \mathcal{P}_{\text{od}}^{\text{ed}}(n)$ with both even and odd parts such that $\lambda_s^e - \lambda_1^o = 1$. For example,

- (i) if $n = 4k$, $k \geq 3$, then $\lambda = (2k, 2k - 1, 1)$;
- (ii) if $n = 13$, then $\lambda = (6, 4, 3)$;
if $n = 17$, then $\lambda = (10, 4, 3)$;
if $n = 4k + 1$, $k \geq 5$, then $\lambda = (2k - 2, 2k - 3, 5, 1)$;
- (iii) if $n = 4k + 2$, $k \geq 3$, then $\lambda = (2k, 2k - 1, 3)$;
- (iv) if $n = 4k + 3$, $k \geq 2$, then $\lambda = (2k + 2, 2k + 1)$.

The combinatorial injection φ_1 together with the discussion above proves the following theorem.

Theorem 3.1. *The inequality $p_{ed}^{od}(n) \leq p_{od}^{ed}(n)$ holds for all $n \geq 0$. Moreover, $p_{ed}^{od}(n) < p_{od}^{ed}(n)$ for $n \geq 11$.*

Inclusion shows that $p_{od}^{ed}(n) \leq p_{ou}^{ed}(n)$ for $n \geq 0$. Since for $n \geq 2$ we have $(1^n) \in \mathcal{P}_{ou}^{ed}(n) \setminus \mathcal{P}_{od}^{ed}(n)$, we have $p_{od}^{ed}(n) < p_{ou}^{ed}(n)$ for $n \geq 2$. Hence, Theorem 3.1 gives the following inequality.

Corollary 3.2. *If $n \geq 2$, then $p_{ed}^{od}(n) < p_{ou}^{ed}(n)$.*

3.2. The inequality $p_{eu}^{ou}(n) < p_{ou}^{eu}(n)$. Let $n \geq 0$. The mapping φ_1 defined in the previous section can be extended to $\mathcal{P}_{eu}^{ou}(n)$. It becomes an injection

$$\varphi_1 : \mathcal{P}_{eu}^{ou}(n) \rightarrow \mathcal{P}_{ou}^{eu}(n).$$

We denote by $\tilde{\mathcal{P}}_{ou}^{eu}(n)$ the subset of partitions $\mu \in \mathcal{P}_{ou}^{eu}(n)$ satisfying one of the following conditions

- (i) $\mu^o = \emptyset$;
- (ii) $\mu^e = \emptyset$;
- (iii) $\mu^o, \mu^e \neq \emptyset$, $\ell_o(\mu) \leq \ell_e(\mu)$ and $\mu_{\ell_o(\mu)} - \mu_{\ell_o(\mu)+1} \geq 2$;
- (iv) $\mu^o, \mu^e \neq \emptyset$, $\ell_e(\mu) < \ell_o(\mu)$ and $\mu_{\ell(\mu)-\ell_e(\mu)} - \mu_{\ell(\mu)-\ell_e(\mu)+1} \geq 2$.

Then $\varphi_1 : \mathcal{P}_{eu}^{ou}(n) \rightarrow \tilde{\mathcal{P}}_{ou}^{eu}(n)$ is a bijection. To see that φ_1 is invertible, let $\mu \in \tilde{\mathcal{P}}_{ou}^{eu}(n)$. If $\mu^o = \emptyset$ or $\mu^e = \emptyset$, then $\varphi_1^{-1}(\mu) = \mu$. If $\mu^o, \mu^e \neq \emptyset$, then $\varphi_1^{-1}(\mu)$ is the partition obtained from μ by subtracting one from each of the first $\ell_m(\mu)$ parts of μ and adding one to each of the last $\ell_m(\mu)$ parts of μ .

Example 4. Let $n = 43$ and $\lambda = (11, 9^2, 6, 4, 2^2)$. Then $\ell_m(\lambda) = \ell_o(\lambda) = 3$. Thus, $\varphi_1(\lambda) = \mu = (12, 10^2, 6, 3, 1^2)$. We note that this falls under Case (iii) as $\ell_e(\mu) > \ell_o(\mu)$ and $\mu_{\ell_o(\mu)} - \mu_{\ell_o(\mu)+1} = \mu_3 - \mu_4 = 10 - 6 = 4$.

Similarly, let $n = 44$ and $\lambda = (11, 9^2, 7, 4, 2^2)$. Then $\ell_m(\lambda) = \ell_e(\lambda) = 3$. Thus, $\varphi_1(\lambda) = \mu = (12, 10^2, 7, 3, 1^2)$. We note that this falls under Case (iv) as $\ell_e(\mu) < \ell_o(\mu)$ and $\mu_{\ell(\mu)-\ell_e(\mu)} - \mu_{\ell(\mu)-\ell_e(\mu)+1} = \mu_4 - \mu_5 = 7 - 3 = 4$.

Remark 1. We could interpret the excess $p_{ou}^{eu}(n) - p_{eu}^{ou}(n)$ in the manner of Proposition 1, but the description is less elegant.

One can verify directly that $p_{ou}^{eu}(2) = p_{eu}^{ou}(2) = 2$. If $n \geq 3$, the partition $(2, 1^{n-2}) \in \mathcal{P}_{ou}^{eu}(n)$ is not in $\tilde{\mathcal{P}}_{ou}^{eu}(n)$. Together with the combinatorial injection φ_1 , this proves the following theorem.

Theorem 3.3. *The inequality $p_{eu}^{ou}(n) \leq p_{ou}^{eu}(n)$ holds for all $n \geq 0$. Moreover, $p_{eu}^{ou}(n) < p_{ou}^{eu}(n)$ for $n \geq 3$.*

3.3. The inequality $p_{od}^{eu}(n) < p_{ed}^{ou}(n)$. In this section, given a partition $\lambda \notin \mathcal{D}$, we denote by λ_r the smallest part size in λ with multiplicity greater than one; we also denote by λ_{rr} the second smallest part size of λ with multiplicity greater than one, if such a part exists.

Let $n \geq 0$. We create an injection

$$\varphi_2 : \mathcal{P}_{od}^{eu}(n) \rightarrow \mathcal{P}_{ed}^{ou}(n).$$

Let $\lambda \in \mathcal{P}_{od}^{eu}(n)$. We consider five cases. For $1 \leq i \leq 5$, we denote by $\mathcal{B}_i(n)$ the image of the set of all partitions in Case i under φ_2 . At the end of this section, we give examples for each case.

Case 1: If $\lambda^e = \emptyset$, define $\varphi_2(\lambda) := \lambda$. Then,

$$\mathcal{B}_1(n) = \{\mu \in \mathcal{P}_{\text{ed}}^{\text{ou}}(n) \mid \mu = \mu^o, \mu \in \mathcal{D}\}.$$

Case 2: (n even) If $\lambda^o = \emptyset$ and $\ell_e(\lambda) = 1$, define $\varphi_2(\lambda) := \lambda$. Then,

$$\mathcal{B}_2(n) = \{(n)\}.$$

Case 3: If $\ell_o(\lambda) \geq 2$ and $\ell_e(\lambda) = 1$, define

$$\varphi_2(\lambda) := (\lambda_1^e - 1, \lambda_1^o, \dots, \lambda_{s-1}^o, \lambda_s^o + 1),$$

the partition obtained from λ by subtracting one from the even part of λ and adding one to the last part of λ . Then,

$$\mathcal{B}_3(n) = \{\mu \in \mathcal{P}_{\text{ed}}^{\text{ou}}(n) \mid \ell_e(\mu) = 1, \ell_o(\mu) \geq 2, \mu_1^o \geq \mu_2^o > \mu_3^o > \dots > \mu_s^o > \mu_1^e\}.$$

Case 4: (n even) If $\ell_e(\lambda) \geq 2$ and $\lambda^o = (\lambda_s^e - 1, 1)$, define

$$\varphi_2(\lambda) := (\lambda_1^e + 1, \lambda_2^e - 1, \dots, \lambda_s^e - 1, \lambda_1^o = \lambda_s^e - 1) \cup (1^{\ell_e(\lambda)-1}),$$

the partition obtained from λ by removing a part equal to 1, adding one to the largest part of λ , subtracting one from every even part of λ other than λ_1^e and inserting $\ell_e(\lambda) - 1$ parts equal to 1. Then,

$$\mathcal{B}_4(n) = \left\{ \mu \in \mathcal{P}_{\text{ed}}^{\text{ou}}(n) \mid \begin{array}{l} \mu = \mu^o, \mu \notin \mathcal{D}, \ell(\mu) \geq 4, \ell(\mu) \text{ even,} \\ \mu_1 \neq \mu_2, m_\mu(1) = \ell_{>1}(\mu) - 2, m_\mu(a_{>1}) > 1 \end{array} \right\}.$$

Case 5: If $\ell_e(\lambda) = \ell_o(\lambda) = 1$, or $\ell_e(\lambda) \geq 2$ and $\lambda^o \neq (\lambda_s^e - 1, 1)$, let

$$k := \begin{cases} \min\{\lambda_1^o, \lambda_s^e - \lambda_1^o\} & \text{if } \lambda^o \neq \emptyset, \\ 1 & \text{if } \lambda^o = \emptyset. \end{cases}$$

and define

$$\varphi_2(\lambda) := (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k) \cup \lambda^o \cup (k^{\ell_e(\lambda)}),$$

the partition obtained from λ by subtracting k from each part of λ^e and inserting $\ell_e(\lambda)$ parts equal to k .

Then, $\mathcal{B}_5(n)$ is the subset of partitions $\mu \in \mathcal{P}_{\text{ed}}^{\text{ou}}(n)$ with $\mu = \mu^o$, $\mu \notin \mathcal{D}$, $\ell(\mu) \geq 3$, and if $\ell(\mu) > 3$ then

- (I) If $\mu_r = 1$ and $m_\mu(1) < \ell_{>1}(\mu)$, then μ_{rr} exists and $m_\mu(1) \in \{\ell_{\geq \mu_{rr}}(\mu) - 1, \ell_{\geq \mu_{rr}}(\mu)\}$.
- (II) If $\mu_r > 1$, then $\ell(\mu) \geq 5$ and
 - (i) if $m_\mu(\mu_r) < \ell_{>\mu_r}(\mu)$, then μ_{rr} exists and $m_\mu(\mu_r) \in \{\ell_{\geq \mu_{rr}}(\mu) - 1, \ell_{\geq \mu_{rr}}(\mu)\}$;
 - (ii) if $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu)$, then $m_\mu(a_{>\mu_r}) > 1$;
 - (iii) if $m_\mu(\mu_r) > \ell_{>\mu_r}(\mu)$, then $\ell_{\geq \mu_r}(\mu)$ is odd.

Note: If $\mu_r = 1$ and $m_\mu(1) \geq \ell_{>1}(\mu)$, there are no additional conditions.

Examples 6. – 15. below illustrate different partitions that belong to Case 5 and the discussion before the various examples clarifies the description of $\mathcal{B}_5(n)$.

The sets $\mathcal{B}_i(n)$, $1 \leq i \leq 5$, are mutually disjoint. To see that $\mathcal{B}_4(n) \cap \mathcal{B}_5(n) = \emptyset$, let $\mu \in \mathcal{B}_5(n)$ and consider the following cases:

- if $\mu_r > 1$, then $\ell(\mu) \geq 5$ and $m_\mu(1) = 1$ and thus $m_\mu(1) \neq \ell_{>1}(\mu) - 2$;

- if $\mu_r = 1$ and $m_\mu(1) = \ell_{>1}(\mu) - 2$, then the condition $m_\mu(1) \in \{\ell_{\geq \mu_{rr}}(\mu) - 1, \ell_{\geq \mu_{rr}}(\mu)\}$ implies that $m_\mu(a_{>1}) = 1$.

Note: In fact, if $\mu \in \mathcal{B}_5(n)$ with $\mu_r = 1$ and $m_\mu(1) \leq \ell_{>1}(\mu) - 2$, then $m_\mu(a_{>1}) = 1$.

The mapping

$$\varphi_2 : \mathcal{P}_{\text{od}}^{\text{eu}}(n) \rightarrow \bigcup_{i=1}^5 \mathcal{B}_i(n)$$

is a bijection. To see that φ_2 is invertible, let $\mu \in \bigcup_{i=1}^5 \mathcal{B}_i(n)$.

If $\mu \in \mathcal{B}_1(n) \cup \mathcal{B}_2(n)$, then

$$\varphi_2^{-1}(\mu) = \mu.$$

If $\mu \in \mathcal{B}_3(n)$, then μ has one even part and at least two odd parts and only the first two parts may be equal. We have

$$\varphi_2^{-1}(\mu) = (\mu_1 + 1, \mu_2, \dots, \mu_{\ell(\mu)-1}, \mu_{\ell(\mu)} - 1).$$

If n is even and $\mu \in \mathcal{B}_4(n)$, then all parts of μ are odd and $\mu_1 \neq \mu_2$. If we set $m = m_\mu(1)$, we have $\ell(\mu) = 2m + 2$ and $\mu_{m+1} = \mu_{m+2} > 1$. Then,

$$\varphi_2^{-1}(\mu) = (\mu_1 - 1, \mu_2 + 1, \dots, \mu_{m+1} + 1, \mu_{m+2}, 1).$$

If $\mu \in \mathcal{B}_5(n)$, then we have the following cases.

If $\ell(\mu) = 3$, we have $\varphi_2^{-1}(\mu) = (\mu_1 + \mu_3, \mu_2)$.

If $\ell(\mu) > 3$, then

- if $m_\mu(\mu_r) < \ell_{>\mu_r}(\mu)$, or $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu)$ and $\mu_r \neq 1$, $\varphi_2^{-1}(\mu)$ is obtained from μ by adding μ_r to each of the first $\ell_{\geq \mu_{rr}}(\mu) - 1$ parts and removing $\ell_{\geq \mu_{rr}}(\mu) - 1$ parts equal to μ_r .
- if $m_\mu(\mu_r) > \ell_{>\mu_r}(\mu)$, or $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu)$ and $\mu_r = 1$, $\varphi_2^{-1}(\mu)$ is obtained from μ by adding μ_r to each of the first $\lfloor \ell_{\geq \mu_r}(\mu)/2 \rfloor$ parts of μ and removing $\lfloor \ell_{\geq \mu_r}(\mu)/2 \rfloor$ parts equal to μ_r .

One can verify directly that $p_{\text{od}}^{\text{eu}}(4) = p_{\text{ed}}^{\text{ou}}(4) = 3$. If $n \geq 5$, the partition $(n-2, 2) \in \mathcal{P}_{\text{ed}}^{\text{ou}}$ is not in $\bigcup_{i=1}^5 \mathcal{B}_i(n)$. Together with the combinatorial injection φ_2 , this proves the following theorem.

Theorem 3.4. *The inequality $p_{\text{od}}^{\text{eu}}(n) \leq p_{\text{ed}}^{\text{ou}}(n)$ holds for all $n \geq 0$. Moreover, $p_{\text{od}}^{\text{eu}}(n) < p_{\text{ed}}^{\text{ou}}(n)$ for $n \geq 5$.*

3.3.1. *Examples.* To help clarify the map φ_2 and its image, we provide examples for all the cases. We also discuss Case 5 in more detail.

Example 5. Let $n = 16$. We give examples for partitions in $\mathcal{P}_{\text{od}}^{\text{eu}}(16)$ in each of the Cases 1 – 4.

Case 1: If $\lambda = (15, 1)$, then $\varphi_2(\lambda) = (15, 1) \in \mathcal{B}_1(16)$.

Case 2: If $\lambda = (16)$, then $\varphi_2(\lambda) = (16) \in \mathcal{B}_2(16)$.

Case 3: If $\lambda = (8, 5, 3)$, then $\varphi_2(\lambda) = (7, 5, 4) \in \mathcal{B}_3(16)$.

Case 4: If $\lambda = (8, 4, 3, 1)$, then $\varphi_2(\lambda) = (9, 3, 3, 1) \in \mathcal{B}_4(16)$.

First, we consider partitions λ with $\ell_e(\lambda) = \ell_o(\lambda) = 1$. If $\lambda = (\lambda_1^e, \lambda_1^o)$, then $\varphi_2(\lambda) = (\lambda_1^e - k, \lambda_1^o) \cup (k) \in \mathcal{B}_5(n)$, as there are exactly three parts, they are all odd, and either $\lambda_1^e - k = \lambda_1^o$ or $k = \lambda_1^o$ (or both if $\lambda_1^e = 2\lambda_1^o$). This leads to all partitions $\mu \in \mathcal{B}_5(n)$ with $\ell(\mu) = 3$. Below, we consider one such example.

Example 6. Let $n = 13$ and $\lambda = (8, 5)$. Then $k = 3$ and $\varphi_2(\lambda) = (5, 5, 3) \in \mathcal{B}_5(13)$.

Next, we restrict our attention to partitions with $k = 1$. We start with partitions λ with $\lambda = \lambda^e = (\lambda_1^e, \lambda_2^e, \dots, \lambda_s^e)$ and $\ell(\lambda) > 1$. Then $k = 1$ and $\mu = \varphi_2(\lambda) = (\lambda_1^e - 1, \lambda_2^e - 1, \dots, \lambda_s^e - 1) \cup (1^{\ell_e(\lambda)}) \in \mathcal{B}_5(n)$, as $\mu_r = 1$ and $m_\mu(1) \geq \ell_{>1}(\mu)$. Note that in this case $\ell(\mu)$ is even and $m_\mu(1) = \ell_{>1}(\lambda) + j$ with $j \geq 0$ even. Below, we give a specific example for a partition with all parts even and length greater than one.

Example 7. Let $n = 16$ and $\lambda = (8, 6, 2)$. Then $\varphi_2(\lambda) = (7, 5, 1^4) \in \mathcal{B}_5(16)$.

Still restricting to partitions with $k = 1$, it remains to consider partitions λ in Case 5 satisfying $\ell_e(\lambda) \geq 2$, $\ell_o(\lambda) \geq 1$, and $\lambda^o \neq (\lambda_s^e - 1, 1)$ for which $k = 1$. We have two subcases.

First, if $\lambda = \lambda^e \cup (1)$, then $k = 1$ and

$$\mu = \varphi_2(\lambda) = (\lambda_1^e - 1, \lambda_2^e - 1, \dots, \lambda_s^e - 1) \cup (1^{\ell_e(\lambda)+1}) \in \mathcal{B}_5(n),$$

as $\mu_r = 1$ and $m_\mu(1) > \ell_{>1}(\mu)$. Note that in this case $\ell(\mu)$ is odd, and $m_\mu(1) = \ell_{>1}(\mu) + j$ with $j \geq 1$ odd. Below, we give a specific example.

Example 8. Let $n = 27$ and $\lambda = (8, 6, 6, 4, 2, 1)$. Then $\varphi_2(\lambda) = (7, 5, 5, 3, 1^7) \in \mathcal{B}_5(27)$.

Second, if $\lambda_1^o = \lambda_s^e - 1$, $\lambda^o \neq (1)$, and $\lambda^o \neq (\lambda_s^e - 1, 1)$, then $k = 1$ and

$$\mu = \varphi_2(\lambda) = (\lambda_1^e - 1, \lambda_2^e - 1, \dots, \lambda_s^e - 1) \cup (\lambda_s^e - 1, \lambda_2^o, \dots, \lambda_s^o) \cup (1^{\ell_e(\lambda)}) \in \mathcal{B}_5(n),$$

as $\mu_r = 1$, $m_\mu(1) < \ell_{>1}(\mu)$, $\lambda_s^e - 1$ is repeated, and if $\lambda_s^o > 1$, then $m_\mu(1) = \ell_{\geq \mu_{rr}}(\mu) - 1$ and if $\lambda_s^o = 1$, then $m_\mu(1) = \ell_{\geq \mu_{rr}}(\mu)$. This falls under restriction (I). Below, we give a specific example.

Example 9. Let $n = 29$ and $\lambda = (8, 6, 6, 5, 3, 1)$. Then $\varphi_2(\lambda) = (7, 5, 5, 5, 3, 1^4) \in \mathcal{B}_5(29)$.

Next, we consider partitions λ in Case 5 satisfying $\ell_e(\lambda) \geq 2$, $\ell_o(\lambda) \geq 1$, and $\lambda^o \neq (\lambda_s^e - 1, 1)$ for which $k > 1$. The images of these partitions under φ_2 all fall under restriction (II).


• First, assume $3 \leq k = \lambda_s^e - \lambda_1^o < \lambda_1^o$. In addition, assume $\lambda_2^o = k$. Then

$$\mu = \varphi_2(\lambda) = (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k = \lambda_1^o) \cup (\lambda_1^o) \cup (k^{\ell_e(\lambda)+1}) \cup (\lambda_3^o, \lambda_4^o, \dots, \lambda_s^o).$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu)$ and $m_\mu(a_{>\mu_r}) > 1$. We obtain partitions in $\mathcal{B}_5(n)$ under restriction (II)(ii). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 10. Let $n = 35$ and $\lambda = (10, 8, 8, 5, 3, 1)$. Then $k = 3$ and

$$\varphi_2(\lambda) = (7, 5, 5, 5, 3, 3, 3, 3, 1) \in \mathcal{B}_5(35).$$




- If $3 \leq k = \lambda_s^e - \lambda_1^o < \lambda_1^o$ with the additional assumption $\lambda_1^o > k > \lambda_2^o$, then

$$\mu = \varphi_2(\lambda) = (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k = \lambda_1^o) \cup (\lambda_1^o) \cup (k^{\ell_e(\lambda)}) \cup (\lambda_2^o, \lambda_3^o, \dots, \lambda_s^o).$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu) - 1$, $m_\mu(a_{>\mu_r}) > 1$, so $\mu_{rr} = a_{>\mu_r}$ and $m_\mu(\mu_r) = \ell_{\geq\mu_{rr}}(\mu) - 1$. We obtain partitions in $\mathcal{B}_5(n)$ falling under restriction (II)(i). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 11. Let $n = 32$ and $\lambda = (10, 8, 8, 5, 1)$. Then $k = 3$ and

$$\varphi_2(\lambda) = (7, 5, 5, 5, 3, 3, 3, 1) \in \mathcal{B}_5(32).$$




- If $3 \leq k = \lambda_s^e - \lambda_1^o < \lambda_1^o$ with the additional assumption $\lambda_i^o = k$ for some $i > 2$, then

$$\begin{aligned} \mu = \varphi_2(\lambda) = & (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k = \lambda_1^o) \cup (\lambda_1^o, \lambda_2^o, \dots, \lambda_{i-1}^o) \\ & \cup (k^{\ell_e(\lambda)+1}) \cup (\lambda_{i+1}^o, \lambda_{i+2}^o, \dots, \lambda_s^o). \end{aligned}$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) < \ell_{>\mu_r}(\mu)$, μ_{rr} exists but is not $a_{>\mu_r}$, and $m_\mu(\mu_r) = \ell_{\geq\mu_{rr}}(\mu)$. We obtain partitions in $\mathcal{B}_5(n)$ falling under condition (II)(i). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 12. Let $n = 66$ and $\lambda = (16, 14, 10, 10, 7, 5, 3, 1)$. Then $k = 3$ and

$$\varphi_2(\lambda) = (13, 11, 7, 7, 7, 5, 3, 3, 3, 3, 1) \in \mathcal{B}_5(66).$$




- If $3 \leq k = \lambda_s^e - \lambda_1^o < \lambda_1^o$ with the additional assumption $\lambda_i^o > k > \lambda_{i+1}^o$ for some $i \geq 2$ (recall that if $j > \ell(\lambda)$, we set $\lambda_j = 0$ for convenience), then

$$\begin{aligned} \varphi_2(\lambda) = & (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k = \lambda_1^o) \cup (\lambda_1^o, \lambda_2^o, \dots, \lambda_i^o) \\ & \cup (k^{\ell(\lambda^e)}) \cup (\lambda_{i+1}^o, \lambda_{i+2}^o, \dots, \lambda_s^o). \end{aligned}$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) < \ell_{>\mu_r}(\mu)$, μ_{rr} exists but is not $a_{>\mu_r}$, and $m_\mu(\mu_r) = \ell_{\geq \mu_{rr}}(\mu) - 1$. We obtain partitions in $\mathcal{B}_5(n)$ falling under restriction (II) CB: (II)(i). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 13. Let $n = 63$ and $\lambda = (16, 14, 10, 10, 7, 5, 1)$. Then $k = 3$ and

$$\varphi_2(\lambda) = (13, 11, 7, 7, 7, 5, 3, 3, 3, 3, 1) \in \mathcal{B}_5(63).$$



Lastly, we consider partitions λ in Case 5, satisfying $\ell_e(\lambda) \geq 2$, $\ell_o(\lambda) \geq 1$, and $\lambda^o \neq (\lambda_s^e - 1, 1)$, and such that $\lambda_s^e - \lambda_1^o \geq \lambda_1^o = k \geq 3$.


- If $\lambda_s^e - \lambda_1^o > \lambda_1^o$, then

$$\begin{aligned} \mu = \varphi_2(\lambda) = & (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k \neq \lambda_1^o) \cup (\lambda_1^o = k) \\ & \cup (k^{\ell(\lambda^e)}) \cup (\lambda_2^o, \lambda_3^o, \dots, \lambda_s^o). \end{aligned}$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) = \ell_{>\mu_r}(\mu) + 1$ and $\ell_{\geq \mu_r}(\mu)$ is odd. We obtain partitions in $\mathcal{B}_5(n)$ falling under restriction (II)(iii). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 14. Let $n = 54$ and $\lambda = (16, 14, 10, 10, 3, 1)$. Then $k = 3$ and

$$\varphi_2(\lambda) = (13, 11, 7, 7, 3, 3, 3, 3, 3, 1) \in \mathcal{B}_5(63).$$




- If $\lambda_s^e - \lambda_1^o = \lambda_1^o$, then

$$\begin{aligned} \mu = \varphi_2(\lambda) = & (\lambda_1^e - k, \lambda_2^e - k, \dots, \lambda_s^e - k = \lambda_1^o) \cup (\lambda_1^o = k) \\ & \cup (k^{\ell(\lambda^e)}) \cup (\lambda_2^o, \lambda_3^o, \dots, \lambda_s^o). \end{aligned}$$

We have $\mu \in \mathcal{B}_5(n)$, as $m_\mu(\mu_r) = \ell_{\geq \mu_r}(\mu) + j$ with $j > 1$ odd and $\ell_{\geq \mu_r}(\mu)$ is odd. We obtain partitions in $\mathcal{B}_5(n)$ falling under restriction (II)(iii). Below, we give a specific example which we illustrate with Ferrers diagrams.

Example 15. Let $n = 59$ and $\lambda = (16, 14, 10, 10, 5, 3, 1)$. Then $k = 5$ and

$$\varphi_2(\lambda) = (11, 9, 5, 5, 5, 5, 5, 5, 3, 1) \in \mathcal{B}_5(59).$$



3.4. The inequality $p_{\text{eu}}^{\text{od}}(n) < p_{\text{ou}}^{\text{ed}}(n)$. Let $n \geq 0$. We define a bijection

$$\varphi_3 : \mathcal{P}_{\text{eu}}^{\text{od}}(n) \rightarrow \mathcal{P}_{\text{ou}}^{\text{ed}}(n).$$

Let $\lambda \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$. We consider three cases.

Case 1: If $\lambda^e = \emptyset$, define $\varphi_3(\lambda) := \lambda$.

Case 2: If $\lambda^o = \emptyset$ and $\lambda \in \mathcal{D}(n)$, define $\varphi_3(\lambda) := \lambda$.

Case 3: If $\lambda^o = \emptyset$ and $\lambda \notin \mathcal{D}(n)$ or $\lambda^o, \lambda^e \neq \emptyset$, define

$$\varphi_3(\lambda) := (\lambda_1 - 1, \lambda_2 - 1, \dots, \lambda_{\ell(\lambda)} - 1) \cup (1^{\ell(\lambda)}),$$

the partition obtained from λ by subtracting one from each of its parts and inserting $\ell(\lambda)$ parts equal to 1.

Let $\tilde{\mathcal{P}}_{\text{ou}}^{\text{ed}}(n)$ be the subset of partitions $\mu \in \mathcal{P}_{\text{ou}}^{\text{ed}}(n)$ satisfying one of the following conditions.

- (i) all parts of μ are odd and distinct;
- (ii) all parts of μ are even and distinct;
- (iii) $\mu^o \neq \emptyset$, $\ell(\mu) \geq 4$, $\ell(\mu)$ even, $m_\mu(1) \geq 2$, $m_\mu(1) \geq \ell_{>1}(\mu)$, and if $\mu^e = \emptyset$ and 1 is the only repeated part, then either $m_\mu(1) = \ell(\mu)$ or $m_\mu(1) \geq \ell_{>1}(\mu) + 4$.

Then, $\varphi_3 : \mathcal{P}_{\text{eu}}^{\text{od}}(n) \rightarrow \tilde{\mathcal{P}}_{\text{ou}}^{\text{ed}}(n)$ is a bijection. To see that φ_3 is invertible, let $\mu \in \tilde{\mathcal{P}}_{\text{ou}}^{\text{ed}}(n)$. If $\mu = \mu^o$ or $\mu = \mu^e \in \mathcal{D}(n)$, then $\varphi_3^{-1}(\mu) = \mu$. Otherwise, μ satisfies all conditions in (iii) above and $\varphi_3^{-1}(\mu)$ is the partition obtained from μ by removing $\ell(\mu)/2$ parts equal to 1 and adding one to each of the first $\ell(\mu)/2$ parts of μ .

Example 16. Let $n = 22$ and $\lambda = (10, 8, 2, 2)$. Then

$$\varphi_3(\lambda) = (9, 7, 1^6) \in \tilde{\mathcal{P}}_{\text{ou}}^{\text{ed}}(22).$$

- μ has a single repeated part and the repeated part is even, of size at least 4, and it is either the smallest or second smallest part of μ ;
- the partition $\tilde{\mu}$ consisting of the parts of μ that are larger than the repeated part is in $\mathcal{C}_1 \cup \mathcal{C}_2$;
- if $\tilde{\mu} \neq \emptyset$ and the repeated part of μ is $2k+2$, then $|\mu| - |\tilde{\mu}| - (2k+2) > \mu_1$.

Case 4: (n even) Suppose $\lambda^o, \lambda^e \neq \emptyset$, $1 \in \lambda$, $\ell_o(\lambda) = 2$, and $\lambda_s^e - \lambda_1^o = 1$. If $\lambda_1^o = 2k+1$ for some $k \geq 1$, then

$$\varphi_4(2k+2, 2k+1, 1) := (2k+2, 2k+2),$$

and if $\ell(\lambda) \geq 4$, then

$$\varphi_4(\lambda) := (\lambda_2^e, \dots, \lambda_{\ell_e(\lambda)-1}^e, \lambda_s^e = 2k+2, 2k+2) \cup (2^{\lambda_1/2}),$$

the partition obtained from λ by removing the first part and the last two parts of λ and inserting a part equal to $2k+2$ and $\lambda_1/2$ parts equal to 2. Note that $\lambda_1 \geq 6$.

Then, $\mathcal{C}_4(n)$ is the subset of partitions $\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ satisfying all of the following conditions:

- $\mu = \mu^e \notin \mathcal{D}(n)$;
- If $\ell(\mu) > 2$, then $2 \in \mu$, the two smallest part sizes in μ are repeated and the repeated part not equal to 2 has multiplicity exactly two. Moreover, $2m_\mu(2) > \mu_1$.

Case 5: (n odd, $n \geq 9$). Suppose $\lambda^e \neq \emptyset$, $\lambda^o = (1)$. We have several subcases.

(i) If $\ell(\lambda) = 2$, then $\varphi_4(\lambda) = \varphi_4(n-1, 1) := (n-4, 2, 2)$. Then

$$\mathcal{C}_{5,(i)}(n) = \{\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n) \mid \ell(\mu) = 3, \ell_0(\mu) = 1, m_\mu(2) = 2\}.$$

(ii) If $\ell(\lambda) = 3$, then $\varphi_4(\lambda) := \varphi_4(\lambda_1, \lambda_2, 1) = (\lambda_2+1, 2^{\lambda_1/2})$. Then, since $n \geq 9$,

$$\mathcal{C}_{5,(ii)}(n) = \{\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n) \mid \ell_o(\mu) = 1, \mu = (\mu_1, 2^{m_\mu(2)}), m_\mu(2) \geq 3, 2m_\mu(2) > \mu_1\}.$$

(iii) If $\ell(\lambda) = 4$ and $2 \in \lambda$, then $\varphi_4(\lambda) := \varphi_4(\lambda_1, \lambda_2, 2, 1) = (\lambda_1+1, 2^{\lambda_2/2+1})$.

Then,

$$\mathcal{C}_{5,(iii)}(n) = \{\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n) \mid \ell_o(\mu) = 1, \mu = (\mu_1^o, 2^{m_\mu(2)}), m_\mu(2) \geq 3, 2m_\mu(2) < \mu_1\}.$$

(iv) If $\ell(\lambda) \geq 4$ and $2 \notin \lambda$, then $\varphi_4(\lambda) := (\lambda_1+1, \lambda_2, \dots, \lambda_{\ell(\lambda)-2}, 2^{\lambda_{\ell(\lambda)-1}/2})$, the partition obtained from λ by adding one to the first part of λ , removing the last two parts of λ , and inserting $\lambda_{\ell(\lambda)-1}/2$ parts equal to 2. Then, $\mathcal{C}_{5,(iv)}(n)$ is the subset of partitions $\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ satisfying all of the following conditions:

- $\ell_o(\mu) = 1$;
- μ^e has at least one part greater than 2;
- the only repeated part of μ is 2 and $2m_\mu(2) < a_{>2}$.

Note that if $\mu \in \mathcal{C}_{5,(iv)}(n)$, then $2m_\mu(2) < \mu_1$.

(v) If $\ell(\lambda) \geq 5$ and $2 \in \lambda$, then $\varphi_4(\lambda) := (\lambda_2+1, \lambda_3, \dots, \lambda_{\ell(\lambda)-2}, 2^{\lambda_1/2+1})$, the partition obtained from λ by removing the first and last part of λ , adding one to second part of λ , and inserting $\lambda_1/2$ parts equal to 2. Then, $\mathcal{C}_{5,(v)}(n)$ is the subset of partitions $\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ satisfying all of the following conditions:

- $\ell_o(\mu) = 1$;
- μ^e has at least two parts greater than 2;
- the only repeated part of μ is 2 and $2m_\mu(2) > \mu_1 + 2$.

Let $\mathcal{C}_5(n) = \mathcal{C}_{5,(i)} \cup \mathcal{C}_{5,(ii)} \cup \mathcal{C}_{5,(iii)} \cup \mathcal{C}_{5,(iv)} \cup \mathcal{C}_{5,(v)}$. Then, $\mathcal{C}_5(n)$ is the subset of partitions $\mu \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ satisfying all of the following conditions:

- $\ell_o(\mu) = 1$;
- the only repeated part of μ is 2;
- if μ has exactly one even part greater than 2, then $2m_\mu(2) < a_{>2}$;
- if μ has at least two even parts greater than 2, then $2m_\mu(2) < a_{>2}$ or $2m_\mu(2) > \mu_1 + 2$.

Clearly, the sets $\mathcal{C}_i(n)$, $1 \leq i \leq 5$, are mutually disjoint.

The mapping

$$\varphi_4 : \mathcal{P}_{\text{od}}^{\text{ed}}(n) \rightarrow \bigcup_{i=1}^5 \mathcal{C}_i(n)$$

is a bijection. To see that φ_2 is invertible, let $\mu \in \bigcup_{i=1}^5 \mathcal{C}_i(n)$.

If $\mu \in \mathcal{C}_1(n)$, then $\varphi_4^{-1}(\mu) = \mu$.

If $\mu \in \mathcal{C}_2(n)$, then $\varphi_4^{-1}(\mu)$ is the partition obtained from μ by subtracting one from the first $\ell_m(\mu)$ parts of μ and adding one to the last $\ell_m(\mu)$ parts of μ .

If $\mu \in \mathcal{C}_3(n)$, let $\tilde{\mu}$ be as in the description of $\mathcal{C}_3(n)$. Then

$$\varphi_4^{-1}(\mu) = \varphi_4^{-1}(\tilde{\mu}) \cup (|\mu| - |\tilde{\mu}| - (2k + 2), 2k + 1, 1).$$

If $\mu \in \mathcal{C}_4(n)$, then

- if $\ell(\mu) = 2$, then $\varphi_4^{-1}(\mu) = (\mu_1, \mu_2 - 1, 1)$;
- if $\ell(\mu) > 2$ and the repeated part greater than 2 is $2k + 2$ for some $k \geq 1$, then

$$\varphi_4^{-1}(\mu) = (\mu \setminus (2k + 2, 2^{m_\mu(2)})) \cup (2m_\mu(2), 2k + 1, 1).$$

If $\mu \in \mathcal{C}_5(n)$, then

- if $\ell(\mu) = 3$, then $\varphi_4^{-1}(\mu) = \varphi_4^{-1}(\mu_1, 2, 2) = (\mu_1 + 3, 1)$;
- If $\ell(\mu) \geq 4$ and μ has no even part greater than 2, then

$$\varphi_4^{-1}(\mu) = \begin{cases} (2m_\mu(2), \mu_1 - 1, 1), & \text{if } 2m_\mu(2) > \mu_1, \\ (\mu_1 - 1, 2m_\mu(2) - 2, 2, 1), & \text{if } 2m_\mu(2) < \mu_1; \end{cases}$$

- If $\ell(\mu) \geq 4$ and μ has an even part greater than 2, then

$$\varphi_4^{-1}(\mu) = \begin{cases} (\mu \setminus (\mu_1, 2^{m_\mu(2)})) \cup (\mu_1 - 1, 2m_\mu(2), 1), & \text{if } 2m_\mu(2) < a_{>2}, \\ (\mu \setminus (\mu_1, 2^{m_\mu(2)-1})) \cup (2m_\mu(2) - 2, \mu_1 - 1, 1), & \text{if } 2m_\mu(2) > \mu_1 + 2. \end{cases}$$

If $n \geq 8$ is even, then the partition $(n - 4, 2, 2) \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ is not in $\bigcup_{i=1}^5 \mathcal{C}_i(n)$. If $n \geq 9$ is odd, then for $n = 4k + 1$ for some $k \geq 2$, the partition $(2k + 1, 2k) \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ is not in $\bigcup_{i=1}^5 \mathcal{C}_i(n)$, and for $n = 4k + 3$ for some $k \geq 2$, the partition $(2k + 1, 2k, 2) \in \mathcal{P}_{\text{eu}}^{\text{od}}(n)$ is not in $\bigcup_{i=1}^5 \mathcal{C}_i(n)$. Together with the combinatorial injection φ_4 , this proves the following theorem.

Theorem 3.6. *The inequality $p_{\text{od}}^{\text{ed}}(n) < p_{\text{eu}}^{\text{od}}(n)$ holds for all $n \geq 8$.*

Case 1: If $\lambda^e = \emptyset$, define

$$\varphi_5(\lambda) = \varphi_5(\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}) := (\lambda_1 + 1, \lambda_2, \dots, \lambda_{\ell(\lambda)}).$$

Then

$$\mathcal{E}_1(2n+1) = \{\mu \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \mid \ell_e(\mu) = 1, \mu_1 - \mu_2 = 1\}.$$

Case 2: If $\lambda^o = \emptyset$, define

$$\varphi_5(\lambda) := \lambda \cup (1).$$

Then

$$\mathcal{E}_2(2n+1) = \{\mu \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \mid \mu^o = (1)\}.$$

Note that if $\mathcal{E}_1(2n+1) \cap \mathcal{E}_2(2n+1) \neq \emptyset$, then $n = 1$.

Case 3: If $\lambda^e, \lambda^o \neq \emptyset$, define

$$\varphi_5(\lambda) = \varphi_5(\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}) := (\lambda_1 + \lambda_2, \lambda_3 - 1, \dots, \lambda_{\ell(\lambda)} - 1) \cup (1^{\ell(\lambda)-1}),$$

the partition obtained from λ by merging the first two parts of λ , subtracting one from each of the remaining parts, and inserting $\ell(\lambda) - 1$ parts of size 1. Then

$$\mathcal{E}_3(2n+1) = \left\{ \mu \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \left| \begin{array}{l} \mu_1 \equiv 2 \pmod{4}, \ell_o(\mu) \geq 3, \\ \mu_1 - \mu_2 \geq \mu_2 + 2 \geq 3, m_\mu(1) \geq \ell_{>1}(\mu) \end{array} \right. \right\}.$$

Clearly, the sets $\mathcal{E}_i(2n+1)$, $1 \leq i \leq 3$, are mutually disjoint.

The mapping

$$\varphi_5 : \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n) \rightarrow \bigcup_{i=1}^3 \mathcal{E}_i(2n+1)$$

is a bijection. To see that φ_5 is invertible, let $\mu \in \bigcup_{i=1}^3 \mathcal{E}_i(2n+1)$.

If $\mu \in \mathcal{E}_1(2n+1)$, then $\varphi_5^{-1}(\mu) = (\mu_1 - 1, \mu_2, \dots, \mu_{\ell(\mu)})$.

If $\mu \in \mathcal{E}_2(2n+1)$, then $\varphi_5^{-1}(\mu) = \mu \setminus (1)$.

If $\mu \in \mathcal{E}_3(2n+1)$, then $\varphi_5^{-1}(\mu) = (\mu_1/2, \mu_1/2, \mu_2 + 1, \dots, \mu_{\ell(\mu)/2} + 1)$, the partition obtained from μ by removing $\ell(\mu)/2$ parts equal to 1, splitting the first part into two equal parts, and adding one to each of the remaining parts.

Finally, we show that if $n \geq 3$, then $\overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \setminus \bigcup_{i=1}^3 \mathcal{E}_i(2n+1) \neq \emptyset$.

Let $n \geq 3$. If $2n+1 = 4k+1$ for $k \geq 2$, then

$$(4(k-1), 1^5) \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \setminus \bigcup_{i=1}^3 \mathcal{E}_i(2n+1).$$

If $2n+1 = 4k+3$ for $k \geq 1$, then

$$(4k, 1^3) \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(2n+1) \setminus \bigcup_{i=1}^3 \mathcal{E}_i(2n+1).$$

□

4.1. **Examples.** To help clarify the map φ_5 , we provide examples for the three cases.

Example 25. Let $n = 12$. We give examples for partitions in $\overline{\mathcal{P}}_{\text{eu}}^{\text{ou}}(24)$ in Cases 1 and 2.

Case 1: If $\lambda = (7^2, 5^2)$, then $\varphi_5(\lambda) = (8, 7, 5^2) \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(25)$.

Case 2: If $\lambda = (12, 6^2)$, then $\varphi_5(\lambda) = (12, 6^2, 1) \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(25)$.

Example 26. Let $n = 26$ and $\lambda = (11^2, 9^2, 8, 2^2)$. Then, $\lambda \in \overline{\mathcal{P}}_{\text{eu}}^{\text{ou}}(52)$ and $\varphi_5(\lambda) = (22, 8^2, 7, 1^8) \in \overline{\mathcal{P}}_{\text{ou}}^{\text{eu}}(53)$.

5. CONCLUDING REMARKS

We defined combinatorial injections to prove several inequalities between the numbers of different partitions of n with parts separated by parity. However, we were unable to find combinatorial arguments for the following inequalities in (1):

$$(7) \quad p_{\text{od}}^{\text{eu}}(n) < p_{\text{eu}}^{\text{od}}(n);$$

$$(8) \quad p_{\text{eu}}^{\text{od}}(n) < p_{\text{ed}}^{\text{ou}}(n);$$

$$(9) \quad p_{\text{eu}}^{\text{ou}}(n) < p_{\text{ou}}^{\text{ed}}(n).$$

We make some observations about inequality (9). In particular, we show that it suffices to prove the inequality for even n .

First, note that in [4] the authors show that the injection $\lambda \mapsto \lambda \cup (1)$ proves combinatorially that $p_{\text{ou}}^{\text{ed}}(n) \leq p_{\text{ou}}^{\text{ed}}(n+1)$ for all $n \geq 0$. In addition, for $n \geq 2$, the partition (n) is in $\mathcal{P}_{\text{ou}}^{\text{ed}}(n+1)$ but not in the image of the injection above. Thus, for $n \geq 2$, we have the strict inequality $p_{\text{ou}}^{\text{ed}}(n) < p_{\text{ou}}^{\text{ed}}(n+1)$. This inequality together with the result of the next proposition shows that it suffices to prove inequality (9) for even n .

Proposition 2. *Let $k \geq 0$. Then*

$$p_{\text{eu}}^{\text{ou}}(2k) = p_{\text{eu}}^{\text{ou}}(2k+1).$$

Proof. Let $k \geq 0$. We define a bijection $\psi : \mathcal{P}_{\text{eu}}^{\text{ou}}(2k) \rightarrow \mathcal{P}_{\text{eu}}^{\text{ou}}(2k+1)$.

Let $\lambda \in \mathcal{P}_{\text{eu}}^{\text{ou}}(2k)$.

If $\lambda^e = \emptyset$, define $\psi(\lambda) = \lambda \cup (1)$.

If $\lambda^e \neq \emptyset$, define $\psi(\lambda)$ to be the partition obtained from λ by adding 1 to the largest even part of λ .

To see that ψ is invertible, let $\mu \in \mathcal{P}_{\text{eu}}^{\text{ou}}(2k+1)$. Clearly, $\mu^o \neq \emptyset$.

If $1 \in \mu$ (which implies that $\mu^e = \emptyset$), then $\psi^{-1}(\mu) = \mu \setminus (1)$ is the partition obtained from μ by removing a part equal to 1.

If $1 \notin \mu$, then $\psi^{-1}(\mu)$ is the partition obtained from μ by subtracting 1 from the smallest odd part of μ . \square

While not needed for the argument that proving inequality (9) for even n is sufficient, we also have the following strict inequality for $p_{\text{eu}}^{\text{ou}}(n)$.

Proposition 3. *Let $k \geq 1$. Then*

$$(10) \quad p_{\text{eu}}^{\text{ou}}(2k-1) < p_{\text{eu}}^{\text{ou}}(2k).$$

Proof. As noted in [4], the injection

$$f(\eta) = f(\eta_1, \eta_2, \dots, \eta_{\ell(\eta)}) := (\eta_1 + 2, \eta_2, \dots, \eta_{\ell(\eta)})$$

shows that, for $n \geq 1$, we have $p_{\text{eu}}^{\text{ou}}(n) \leq p_{\text{eu}}^{\text{ou}}(n+2)$. Moreover, if $n \geq 1$, then the partition (1^{n+2}) is in $\mathcal{P}_{\text{eu}}^{\text{ou}}(n+2)$ but not in the image of $\mathcal{P}_{\text{eu}}^{\text{ou}}(n)$ under f . Hence, if $n \geq 1$, we have $p_{\text{eu}}^{\text{ou}}(n) < p_{\text{eu}}^{\text{ou}}(n+2)$.

Inequality (10) is clearly true for $k = 1$ since $p_{\text{eu}}^{\text{ou}}(1) = 1$ and $p_{\text{eu}}^{\text{ou}}(2) = 2$. If $k \geq 2$, and ψ is the bijection of Proposition 2, the mapping $f \circ \psi^{-1}$ is an injection from $\mathcal{P}_{\text{eu}}^{\text{ou}}(2k-1)$ to $\mathcal{P}_{\text{eu}}^{\text{ou}}(2k)$. Since f is not surjective, we have $p_{\text{eu}}^{\text{ou}}(2k-1) < p_{\text{eu}}^{\text{ou}}(2k)$. \square

We end this section by discussing further directions of study for partitions with parts separated by parity. In addition to the modifiers u (unrestricted) and d (distinct), we could also include the modifier nd to mean that the relevant parts (even or odd) are not all distinct. For example, $\mathcal{P}_{\text{ou}}^{\text{end}}(n)$ is the set of partitions of n with all even parts larger than all odd parts, there is at least one even part with multiplicity larger than 1, and odd parts unrestricted. From the inequalities (1), we can find several inequalities involving the modifier nd (all holding for sufficiently large n).

First, consider the inequalities $p_{\text{od}}^{\text{eu}}(n) < p_{\text{eu}}^{\text{od}}(n) < p_{\text{ed}}^{\text{ou}}(n) < p_{\text{eu}}^{\text{ou}}(n) < p_{\text{ou}}^{\text{eu}}(n)$ and calculate several differences:

- $p_{\text{ou}}^{\text{eu}}(n) - p_{\text{od}}^{\text{eu}}(n) = p_{\text{ond}}^{\text{eu}}(n)$,
- $p_{\text{eu}}^{\text{ou}}(n) - p_{\text{eu}}^{\text{od}}(n) = p_{\text{eu}}^{\text{ond}}(n)$,
- $p_{\text{eu}}^{\text{ou}}(n) - p_{\text{ed}}^{\text{ou}}(n) = p_{\text{end}}^{\text{ou}}(n)$,
- $p_{\text{od}}^{\text{eu}}(n) - p_{\text{od}}^{\text{ed}}(n) = p_{\text{od}}^{\text{end}}(n)$.

Therefore,

$$p_{\text{eu}}^{\text{ond}}(n) - p_{\text{end}}^{\text{ou}}(n) = p_{\text{eu}}^{\text{ou}}(n) - p_{\text{eu}}^{\text{od}}(n) - (p_{\text{eu}}^{\text{ou}}(n) - p_{\text{ed}}^{\text{ou}}(n)) = p_{\text{ed}}^{\text{ou}}(n) - p_{\text{eu}}^{\text{od}}(n) > 0$$

and

$$\begin{aligned} p_{\text{ond}}^{\text{eu}}(n) - p_{\text{eu}}^{\text{ond}}(n) &= p_{\text{ou}}^{\text{eu}}(n) - p_{\text{od}}^{\text{eu}}(n) - (p_{\text{eu}}^{\text{ou}}(n) - p_{\text{eu}}^{\text{od}}(n)) \\ &= (p_{\text{ou}}^{\text{eu}}(n) - p_{\text{eu}}^{\text{ou}}(n)) + (p_{\text{eu}}^{\text{od}}(n) - p_{\text{od}}^{\text{eu}}(n)) > 0. \end{aligned}$$

This leads to the following inequalities.

Proposition 4. *For n sufficiently large,*

$$p_{\text{end}}^{\text{ou}}(n) < p_{\text{eu}}^{\text{ond}}(n) < p_{\text{ond}}^{\text{eu}}(n).$$

Next, consider the inequalities $p_{\text{ed}}^{\text{od}}(n) < p_{\text{od}}^{\text{ed}}(n) < p_{\text{od}}^{\text{eu}}(n) < p_{\text{eu}}^{\text{od}}(n) < p_{\text{ed}}^{\text{ou}}(n)$ and calculate several differences:

- $p_{\text{ed}}^{\text{ou}}(n) - p_{\text{ed}}^{\text{od}}(n) = p_{\text{ed}}^{\text{ond}}(n)$,
- $p_{\text{eu}}^{\text{od}}(n) - p_{\text{ed}}^{\text{od}}(n) = p_{\text{end}}^{\text{od}}(n)$,
- $p_{\text{od}}^{\text{eu}}(n) - p_{\text{od}}^{\text{ed}}(n) = p_{\text{od}}^{\text{end}}(n)$,
- $p_{\text{eu}}^{\text{ou}}(n) - p_{\text{ed}}^{\text{ou}}(n) = p_{\text{end}}^{\text{ou}}(n)$.

Therefore,

$$\begin{aligned} p_{\text{end}}^{\text{od}}(n) - p_{\text{od}}^{\text{end}}(n) &= p_{\text{ed}}^{\text{od}}(n) - p_{\text{ed}}^{\text{od}}(n) - (p_{\text{od}}^{\text{eu}}(n) - p_{\text{od}}^{\text{ed}}(n)) \\ &= (p_{\text{ed}}^{\text{od}}(n) - p_{\text{ed}}^{\text{od}}(n)) + (p_{\text{eu}}^{\text{od}}(n) - p_{\text{od}}^{\text{eu}}(n)) > 0 \end{aligned}$$

and

$$p_{\text{ed}}^{\text{od}}(n) - p_{\text{end}}^{\text{od}}(n) = p_{\text{ed}}^{\text{ou}}(n) - p_{\text{ed}}^{\text{od}}(n) - (p_{\text{eu}}^{\text{od}}(n) - p_{\text{ed}}^{\text{od}}(n)) = p_{\text{ed}}^{\text{ou}}(n) - p_{\text{eu}}^{\text{od}}(n) > 0$$

This leads to the following inequalities.

Proposition 5. *For n sufficiently large,*

$$p_{od}^{end}(n) < p_{end}^{od}(n) < p_{ed}^{ond}(n).$$

Moreover, to prove (8) combinatorially it suffices to find an injection from $\mathcal{P}_{end}^{od}(n)$ into $\mathcal{P}_{ed}^{ond}(n)$.

Lastly, consider the inequalities $p_{od}^{eu}(n) < p_{eu}^{od}(n) < p_{ed}^{ou}(n) < p_{eu}^{ou}(n) < p_{ou}^{eu}(n)$ and calculate several differences:

- $p_{ou}^{eu}(n) - p_{od}^{eu}(n) = p_{ond}^{eu}(n)$,
- $p_{eu}^{ou}(n) - p_{eu}^{od}(n) = p_{eu}^{ond}(n)$,
- $p_{eu}^{ou}(n) - p_{ed}^{ou}(n) = p_{end}^{ou}(n)$,
- $p_{ou}^{eu}(n) - p_{ou}^{ed}(n) = p_{ou}^{end}(n)$.

Therefore

$$p_{eu}^{ond}(n) - p_{end}^{ou}(n) = p_{eu}^{ou}(n) - p_{eu}^{od}(n) - (p_{eu}^{ou}(n) - p_{ed}^{ou}(n)) = p_{ed}^{ou}(n) - p_{eu}^{od}(n) > 0$$

and

$$\begin{aligned} p_{ond}^{eu}(n) - p_{eu}^{ond}(n) &= p_{ou}^{eu}(n) - p_{od}^{eu}(n) - (p_{eu}^{ou}(n) - p_{eu}^{od}(n)) \\ &= (p_{ou}^{eu}(n) - p_{eu}^{ou}(n)) + (p_{ed}^{ou}(n) - p_{od}^{eu}(n)) > 0 \end{aligned}$$

This leads to the following inequalities.

Proposition 6. *For n sufficiently large,*

$$p_{end}^{ou}(n) < p_{eu}^{ond}(n) < p_{ond}^{eu}(n).$$

It would be interesting to find combinatorial proofs for the inequalities in Propositions 4–6.

REFERENCES

- [1] G. E. Andrews, *The Theory of Partitions*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1998. Reprint of the 1976 original.
- [2] G. E. Andrews, *Integer partitions with even parts below odd parts and the mock theta functions*, Ann. Comb. **22** (2018), no. 3, 433–445.
- [3] G. E. Andrews, *Partitions with parts separated by parity*, Ann. Comb. **23** (2019), no. 2, 241–248.
- [4] K. Bringmann, W. Craig and C. Nazaroglu, *On the asymptotic behavior for partitions separated by parity*, arXiv:2403.00635 [math.CO].
- [5] K. Banerjee and M. G. Dastidar, *Ramanujan’s theta functions and parity of parts and cranks of partitions*, Annals of Combinatorics **27.3** (2023): 565–578.
- [6] H. E. Burson and D. Eichhorn, *On the positivity of infinite products connected to partitions with even parts below odd parts and copartitions*, arXiv preprint arXiv:2209.13784 (2022).
- [7] K. Bringmann and C. Jennings-Shaffer, *A Note on Andrews’ Partitions with Parts Separated by Parity*, Ann. Comb. **23**, 573–578 (2019). <https://doi.org/10.1007/s00026-019-00440-z>
- [8] D. Chen and R. Chen, *Congruence modulo 4 for Andrews’ even parts below odd parts partition function*, Annals of Combinatorics **27.2** (2023): 269–279.
- [9] S. Chern, *On a problem of George Andrews concerning partitions with even parts below odd parts*, Afrika Matematika **30.5** (2019): 691–695.
- [10] S. Chern, *Note on partitions with even parts below odd parts*, Mathematical Notes **110.3** (2021): 454–457.
- [11] S. Fu and D. Tang, *Partitions with parts separated by parity: conjugation, congruences and the mock theta functions*, Proceedings of the Royal Society of Edinburgh Section A: Mathematics (2023): 1–21.

- [12] A. Goswami and A. K. Jha, *On the Parity of Some Partition Functions*, Technical report no. 20-06 in RISC Report Series, Research Institute for Symbolic Computation (RISC), Johannes Kepler University Linz, Austria. ISSN 2791-4267 (online). 2019.
- [13] P. J. Mahanta, M. P. Saikia, and A. Sarma. *Biases in non-unitary partitions*, arXiv preprint arXiv:2301.09845 (2023).
- [14] D. Passary, *Studies of partition functions with conditions on parts and parity*, The Pennsylvania State University, 2019.
- [15] C. Ray and R. Barman, *On Andrews' integer partitions with even parts below odd parts*, Journal of Number Theory, Volume 215, 2020, Pages 321-338, ISSN 0022-314X, <https://doi.org/10.1016/j.jnt.2020.02.001>.
- [16] R. Rahman and N. Saikia. *Congruences for Partition Functions $\overline{\mathcal{EO}}(n)$ and $\mathcal{EO}_e(n)$* , Kyungpook Mathematical Journal 63.2 (2023): 155-166.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, COLLEGE OF THE HOLY CROSS,
WORCESTER, MA 01610, USA,

Email address: `cballant@holycross.edu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EASTERN ILLINOIS UNIVERSITY, CHARLESTON,
IL 61920, USA,

Email address: `arwelch@eiu.edu`