

# GRAPHS AND THEIR VIETORIS-RIPS COMPLEXES HAVE THE SAME PSEUDOTOPOLOGICAL WEAK HOMOTOPY TYPE

JONATHAN TREVIÑO-MARROQUÍN

ABSTRACT. In this document, we propose a bridge between the graphs and the geometric realizations of their Vietoris Rips complexes, i.e. Graphs, with their canonical Čech closure structure, have the same homotopy type that the realization of their Vietoris Rips complex.

## 1. INTRODUCTION

The homotopical study of (reflexive) graphs has recently seen increased activity and interest from a number of different perspectives. For example, in [5] the authors develop  $A$ -theory, a cubical-type homotopy is defined in graphs and uses the inductive product and the discrete interval to generate their homotopies; in [9, 11]  $\times$ -homotopy is introduced for studying not-necessarily reflexive graphs; there are also homotopy theories for digraphs [12] and the non-topological homotopy for metric spaces considered in [20]; restricting to reflexive graphs on cubical lattices whose edges are determined by their position in the lattice, we obtain digital homotopy [16, 17], finally, Antonio Rieser proposed that Čech closure spaces [22] and pseudotopological spaces [23] could be used as categories in which to simultaneously develop the homotopy theory of topological spaces, graphs, digraphs, and point clouds (the latter for topological data analysis [6, 7]), extending earlier work of Demaria [10] on graphs.

In this article, we continue the development of homotopy theory in pseudotopological spaces begun in [22, 23], and we prove that every graph  $X$  has the same weak homotopy type as its Vietoris-Rips complex  $|VR(X)|$ . To achieve this, we define an intermediate pretopological space  $R(X)$  (Definition 2.6), which is a simplicial complex with a modified closure structure such that (1) there is a continuous inclusion of the vertices of  $X$  into  $R(X)$ , (2)  $R(X)$  is strong homotopy equivalent to the graph, and (3)  $R(X)$  is weak homotopy equivalent to the geometric realization  $|VR(X)|$  of the Vietoris-Rips complex of  $X$ . The construction of the weak homotopy equivalence further requires that we construct three transformations of functions  $f : Y \rightarrow R(X)$ , which are the key to the proof:

- **the discrete modification** (Definition 2.9), which sends a map  $f$  with codomain  $R(X)$  to one whose image lies only in  $X \subset R(X)$ ;
- **the flood** (Definition 2.14), which expands the preimage of a vertex  $v \in X \subset R(X)$  in order to create a neighborhood  $U$  of  $f^{-1}(v) \subset Y$  small enough

---

This work was supported by the CONACYT postgraduate studies scholarship number 839062, and the results in this article form a part of the author's PhD thesis project supervised by Antonio Rieser.

so that the image of every element in  $U$  is contained in the closure of the image of any of its points;

- **the convex transformation** (Definition 2.19), which converts maps  $f : Y \rightarrow R(X)$  whose image is in  $X \subset R(X)$  into a simplicial map from a triangulation of  $Y$  to  $R(X)$ .

We study the properties of each of these transformations in turn, which we then use in combination for the proof of our main result. The critical property of each is that these transformations, when applied to a continuous maps from a triangulizable compact metric space, produce a continuous map.

Applying the flood transformation after the discrete modification uses the same fundamental idea as that studied in digital homotopy to find the second homotopy group for the digital sphere [15]. In contrast to that paper, whose domain is the lattice  $\mathbb{Z}^2$  and the distances are discrete, in this paper, the domain will be a triangulizable compact metric space, in particular  $S^n$  and  $S^n \times I$ .

The results in this paper, when combined with well-known results about the Vietoris-Rips complex [13], we observe that for a “big enough” cloud of points sampled from a manifold, the pseudotopological homotopy groups (defined in [23]) from certain closure spaces built on those points are isomorphic to the classical homotopy groups of the manifold being sampled. In addition, the Vietoris-Rips complex is difficult to understand and only a few complete calculations are known for metric spaces at any scale, for example [1, 2, 4] and is used to bound Gromov-Hausdorff distance [3, 14, 18]; thus, with this paper and growing wave of studies in these categories mentioned above, we may hope for new ways to approach the Vietoris-Rips complex as well.

The isomorphism between the pseudotopological homotopy groups of finite graphs and those of its clique complex has been also established independently by Milicevic and Scoville in [19]. Their proof is very different from this one, particularly as it does not pass through the space  $R(X)$ , and it only applies to finite graphs, whereas the technique we study here is also valid for arbitrary, even uncountably infinite, reflexive graphs.

## 2. WEAK HOMOTOPY EQUIVALENCE

Closure spaces have two definitions which are equivalent categories: one defined through the closure operation, which is a map from the power set to the power set, and another defined through convergence of filters, which is a function from the space to the collection of all of the filters. For this paper, it is enough to know that we can use both definition for the same object. By completeness, we briefly introduce these both categories. (To go deeper with why these are equivalent, we invite the reader to check [21], proposition 2.3.1.8)

**Definition 2.1** ([21], 2.3.1.6). A *closure space* is a pair  $(X, c)$ , where  $X$  is a set and  $c : 2^X \rightarrow 2^X$  a function, called a closure operation on  $X$ , satisfying the following conditions:

- (1)  $c(\emptyset) = \emptyset$ .
- (2)  $A \subset c(A)$  for each  $A \in 2^X$ .
- (3)  $c(A \cup B) = c(A) \cup c(B)$  for each  $A, B \in 2^X$ .

A map  $f : X \rightarrow Y$  is continuous from  $(X, c)$  to  $(Y, c')$ , closure spaces, denoted as  $f : (X, c) \rightarrow (Y, c')$ , provided that  $f(c(A)) \subset c'(f(A))$  for every  $A \in 2^X$ .

**Definition 2.2** ([8], 14 B.1). Let  $(X, c)$  a closure space. We say that  $U \subset X$  is *neighborhood* of a point  $x \in X$  if  $x \in X \setminus c(X \setminus U)$ . The neighborhood system of a point  $x$ , denoted  $\mathcal{U}(x)$ , is the collection of neighborhoods of  $x$ .

**Proposition 2.3** ([21], 2.3.1.6). *Let  $(X, c)$  and  $(Y, c')$  closure spaces. Then  $f : X \rightarrow Y$  is continuous if and only if  $f(\mathcal{U}(x)) \supset \mathcal{U}(f(x))$  for every  $x \in X$ .*

**Example.** Let  $S^1$  be the circle (of diameter one) and  $C_4$  be 4-cyclic graph (or the digital circle), i.e. the set  $\{0, 1, 2, 3\}$  such that  $cl(y) = \{y - 1, y, y + 1\}$  module 4. We have that the map  $f : S^1 \rightarrow C_4$  such that

$$f(x) = \begin{cases} 0 & x \in [0, 1/4) \\ 1 & x \in [1/4, 2/4) \\ 2 & x \in [2/4, 3/4) \\ 3 & x \in [3/4, 1) \end{cases}$$

is continuous.

**Definition 2.4.** Let  $f, g : X \rightarrow Y$  be continuous maps between closure spaces.  $f$  and  $g$  are said to be *homotopic*, denoted by  $f \simeq g$ , if there exists a continuous map  $H : X \times I \rightarrow Y$  such that  $H(x, 0) = f(x)$  and  $H(x, 1) = g(x)$  for every  $x \in X$ .  $X$  and  $Y$  are called *strong homotopy equivalent* if there exist continuous maps  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  such that  $gf \simeq 1_X$  and  $fg \simeq 1_Y$ .

We define the *homotopy classes* for a closure space  $X$  for every  $n \geq 0$ , denoted by  $\pi_n(X, *)$ , as the set of equivalence classes of continuous maps  $(S^n, *) \rightarrow (X, *)$ . Using the standard  $\star$  operation in those maps and standard arguments, we can prove that  $\pi_n(X, *)$  is a group for all  $n \geq 1$  and an abelian group for every  $n > 1$ . Two closure spaces  $X$  and  $Y$  are *weak homotopy equivalent*, and  $f$  (respectively  $g$ ) is a *weak homotopy equivalence* if there exists a continuous map  $f : X \rightarrow Y$  (or  $g : Y \rightarrow X$ ) such that  $\pi_n(f) : \pi_n(X, *) \rightarrow \pi_n(Y, *)$  (or  $\pi_n(g) : \pi_n(Y, *) \rightarrow \pi_n(X, *)$ ) is an isomorphism for every  $n \geq 0$ . We say that  $X$  and  $Y$  have the same weak homotopy type if there exists a zig-zag of weak homotopy equivalences from  $X$  to  $Y$

$$X \rightarrow X_1 \leftarrow X_2 \rightarrow \dots \leftarrow X_{n-1} \rightarrow X_n \leftarrow Y.$$

**Remark.** A Čech closure space is a particular case of a pseudotopological space. The pseudotopological weak homotopy studied in [23] agrees with the one defined for closure spaces in Definition 2.4.

Every undirected (also valid for directed) simple graph has a natural Čech closure structure and induces a simplicial complex. We recall these structures in the following definition.

**Definition 2.5.** Let  $(X, E)$  be a simple, reflexive graph (i.e. the graph does not have multiple edges and every loop  $(x, x)$  is an edge in the graph). We say that  $(X, E)$  is *locally finite* if for every  $x \in X$  there exists a finite number of vertices  $(x, y) \in E$ . We define the *canonical closure structure for a graph  $(X, E)$* , denoted by  $c_X$ , such that

$$c_X(x) := \{y \in X \mid (x, y) \in E\} \text{ for every } x \in X$$

$$c_X(A) := \bigcup_{x \in A} c_X(x) \text{ for every } A \subset X$$

Observe that this closure operation induces the neighborhood system at  $x$  in  $X$  as  $\mathcal{U}_X(x) = [c_X(x)]_X$ .

When the context is clear, we omit to write the set of edges, i.e. we write just  $X$  instead of  $(X, E)$  when we refer to a graph. We can (and we implicitly do) set an arbitrary order  $<$  over the vertices of  $X$ ; this is helpful in a couple of construction during the document.

The following definition introduce two realizations of a graph. The first realization is the classical simplicial complex induced by a graph (the clique complex); on the other hand, the second realization is a hybrid between the graph and the clique complex. This second realization is actually finer than the first one and allows a strong homotopy equivalence between itself and the graph.

**Definition 2.6.** Let  $X$  be a graph.

- (1)  $|VR(X)|$  is the geometric realization of the simplicial complex with  $n$ -simplices  $\{v_0, \dots, v_n\}$  such that  $v_i \in cl(v_j)$  for every  $i, j$ . This space is a topological, and then it has a natural topological closure operation which we denote by  $c_{|VR(X)|}$ .
- (2)  $R(X)$  is defined over the same set of  $|VR(X)|$ , but the neighborhood at  $v \in X$  in  $R(X)$  is defined as

$$\mathcal{U}_v := \left[ \bigcup_{v \in \sigma} |\sigma| \right]_{R(X)}$$

which is a neighborhood system at  $v$ . The neighborhood system of every point in  $|VR(X)| - X$  is the equal to the neighborhood filter in  $|VR(X)|$ . We denote the closure defined by the neighborhood system as  $c_{R(X)}$ .

It is straightforward to see that  $(|VR(X)|, c_{|VR(X)|})$  is a finer pretopological space than  $(|VR(X)|, c_{R(X)})$ , they are over the same set and by construction there are fewer sets in the neighborhood at  $v$  in  $R(X)$  than in  $|VR(X)|$  (i.e. the identity is continuous as PsTop spaces, and then as closure spaces). In the same sense, we construct  $R(X)$  satisfying that  $X$  is a (pretopological) subspaces of it.

**Definition 2.7** (The Barycentric Cover of a Simplex). Let  $\Delta^n := \{v_0, \dots, v_n\}$  an  $n$ -simplex. Define

$$b_{\{v_{i_0}, \dots, v_{i_k}\}} \text{ the barycenter of the } k\text{-simplex } \{v_{i_0}, \dots, v_{i_k}\}$$

for every  $i \in [n]$ , let's define

$$A_i := \bigcup_{\alpha_j \in [n]} |\{v_i, b_{\{v_i, v_{\alpha_1}\}}, \dots, b_{\{v_i, v_{\alpha_1}, \dots, v_{\alpha_n}\}}\}|$$

The collection  $\{A_0, \dots, A_n\}$  is said to be *the barycentric cover of  $|\Delta^n|$*

**Proposition 2.8.** Let  $\Delta^n := \{v_0, \dots, v_n\}$  be an  $n$ -simplex and  $\{A_0, \dots, A_n\}$  the barycentric cover of  $\Delta^n$ . Then  $|\Delta^n| = \bigcup_{i=0}^n |A_i|$ .

*Proof.* Let  $x \in |\Delta^n|$ , then for every  $i \in [n]$  there exists  $t_i \geq 0$  such that  $\sum_{i=0}^n t_i = 1$  and

$$x = \sum_{i=0}^n t_i v_i$$

Since we have a finite number of elements, there exists an index in  $[n]$ , say  $\beta_0$  such that  $t_{\beta_0} = \min_{i \in [n]} \{t_i\}$ , thus

$$x = \sum_{i \in [n] \setminus \{\beta_0\}} (t_i - t_{\beta_0})x_i + \frac{t_{\beta_0}}{n+1} \left( \frac{1}{n+1} \sum_{i=0}^n x_i \right)$$

With the same argument we can find the index  $\beta_1$  such that  $t_{\beta_1} = \min_{i \in [n] \setminus \beta_0}$  and

$$\begin{aligned} x &= \sum_{i \in [n] \setminus \{\beta_0, \beta_1\}} (t_i - t_{\beta_1})x_i + (t_{\beta_1} - t_{\beta_0})n \left( \frac{1}{n} \sum_{i \in [n] \setminus \{\beta_0\}} x_i \right) \\ &\quad + t_{\beta_0}(n+1) \left( \frac{1}{n+1} \sum_{i=0}^n x_i \right) \end{aligned}$$

Following this procedure until we cover all of the vertexes, we obtain that

$$x = \sum_{k=0}^n \left( (t_{\beta_{n-k}} - t_{\beta_{n-k-1}})(k+1) \left( \frac{1}{k+1} \sum_{i \in [n] \setminus \{\beta_j | j \in [k]\}} x_i \right) \right)$$

defining  $t_{\beta_{-1}} = 0$ . Observe that, by definition of barycenter,

$$b_{\{v_{\beta_j} | j \in [k]\}} = \frac{1}{k+1} \sum_{i \in [n] \setminus \{\beta_j | j \in [k]\}} x_i$$

Concluding that  $x \in A_{\beta_n}$ .  $\square$

Take a  $n$ -simplex  $\Delta^n := \{v_0, \dots, v_n\}$  and a  $m$ -simplex  $\Delta^m := \{w_0, \dots, w_m\}$  which share a  $k$ -simplex, say  $\Delta^k := \{v_0, \dots, v_k\}$  with  $v_i = w_i$  for every  $i \in [k]$ ; suppose that  $\{A_0, \dots, A_n\}$ ,  $\{B_0, \dots, B_m\}$  and  $\{C_0, \dots, C_k\}$  the barycentric cover of  $\Delta^n$ ,  $\Delta^m$  and  $\Delta^k$ , respectively. Observe that  $C_i = A_i \cap \Delta^n = B_i \cap \Delta^m$  for every  $i \in [k]$  given that the barycentric cover of  $C_i$  only depends on the barycenters in  $\Delta^k$ .

With this in mind, we can extend the definition below (Definition 2.9) for all  $R(X)$  if we can define it in the simplexes.

**Definition 2.9** (Discrete Modification). Let  $X$  be a graph, we define *the discrete modification* as  $\Theta_X : R(X) \rightarrow X \subset R(X)$  as follows: Let  $\Delta^n := \{v_0, \dots, v_n\}$  be an  $n$ -simplex and define

$$b_{\{v_{i_0}, \dots, v_{i_k}\}} \text{ the barycenter of the } k\text{-simplex } \{v_{i_0}, \dots, v_{i_k}\}$$

for every  $i \in [n]$ , recall that

$$A_i := \bigcup_{\alpha_j \in [n]} |\{v_i, b_{\{v_i, v_{\alpha_1}\}}, \dots, b_{\{v_i, v_{\alpha_1}, \dots, v_{\alpha_n}\}}\}|$$

Observe that if we have some repeat elements, then they are just simplexes contained in the one which every element is different. Then we define:

$$\Theta_X(x) := \begin{cases} v_0 & x \in A_0 \\ v_1 & x \in A_1 - A_0 \\ \vdots & \\ v_n & x \in A_n - \bigcup_{i=0}^{n-1} A_i \end{cases}$$




FIGURE 1. Illustration of the discrete modification for the geometric realization of a 2-simplex

**Lemma 2.10.** *Let  $X$  be a graph, the discrete modification  $\Theta_X : R(X) \rightarrow X \subset R(X)$  is continuous.*

*Proof.* Let  $x \in R(X)$ . In the first case, let  $x \in X$ , by definition we have that  $\mathcal{U}(x) = [\cup_{x \in \sigma} |\sigma|]_{R(X)}$ . By the definition of the discrete modification,  $\Theta_X |\sigma| = \{v \mid v \in \sigma\} \subset |\sigma|$ . Then  $\Theta_X(\cup_{x \in \sigma} |\sigma|) = \cup_{x \in \sigma} \{v \mid v \in \sigma\} \subset \cup_{x \in \sigma} |\sigma|$ . Thus  $\Theta_X[\cup_{x \in \sigma} |\sigma|]_{R(X)} \rightarrow x = \Theta_X(x)$ .

Let  $x \notin X$ , by the definition of  $VR(X)$ , there exists a unique simplex  $\sigma$  containing  $x$  in its relative interior, then  $\Theta_X(x) \in \{v \mid v \in \sigma\}$ . A filter  $\mathcal{F} \rightarrow x$  in  $VR(X)$  (and then in  $R(X)$  because  $x \notin X$ ) if it converges in the restriction of every simplex which contains  $x$ , then  $\cup_{\{v \mid v \in \sigma\} \subset \sigma'} |\sigma'|$ . Then  $\Theta_X(\mathcal{U}(x)) \supset \Theta_X[\cup_{\{v \mid v \in \sigma\} \subset \sigma'} |\sigma'|]_{R(X)} \supset [\cup_{\Theta_X(x) \in \sigma} |\sigma|]_{R(X)}$ . Thus  $\Theta_X(\mathcal{U}(x)) \rightarrow \Theta_X(x)$ .  $\square$

**Remark.** By construction, we have that  $Im(\Theta_X) \subset X$ ,  $\Theta_X \Theta_X = \Theta_X$  and  $\Theta_X(v) = v$  for every  $v \in X \subset R(X)$ .

When  $(X, d)$  is a metric space, we write  $B_r(x) := \{y \in X \mid d(x, y) < r\}$ .

**Proposition 2.11.** *Let  $X$  be a graph.  $X$  and  $R(X)$  are strong homotopy equivalent.*

*Proof.* Let  $g : X \rightarrow R(X)$  be the map which sends the points in  $X$  to the corresponding vertices of the simplices in  $R(X)$ .

It is straightforward to observe that  $\Theta_X g = 1_X$ , so we only need to express a homotopy from  $g \Theta_X$  to  $1_{R(X)}$ .

First we restrict for a simplex  $\Delta \in VR(X)$ . Let  $H|_{\Delta} : \Delta \times I \rightarrow \Delta \subset R(X)$  be defined by

$$H|_{\Delta}(x, t) := \begin{cases} 1_{\Delta}(x) & \text{if } t \in [0, 1) \\ \Theta_X(x) & \text{if } t = 1. \end{cases}$$

We claim that  $H|_{\Delta}$  is continuous: For every point  $(x, 1)$ , its image contained in  $\Delta$ , getting that  $H$  is continuous in that point; for every point  $(x, t)$  with  $t < 1$ , there exist  $r > 0$  such that  $B_r(x, t) \cap (\Delta \times \{1\}) = \emptyset$ , concluding that  $H(B_r(x, t)) = B_r(x)$  and  $H$  is continuous in that point.

Denote the neighborhood system of  $t$  in  $[0, 1]$  by  $\mathcal{I}(t)$ . For the whole domain  $R(X)$ , we observe that for all  $t < 1$  there exists  $\varepsilon > 0$  such that for every  $U \in \mathcal{U}$  satisfies that  $U \times (t - \varepsilon, t + \varepsilon) \cap U \times \{1\} = \emptyset$ , then  $H(\mathcal{U}(x) \times (t - \varepsilon, t + \varepsilon)) = \mathcal{U}(x)$ , and then  $H(\mathcal{U}(x) \times \mathcal{I}(t)) \rightarrow x$ .

On the other hand,  $H(\mathcal{U} \times \mathcal{I}(1)) = [\{H(A \times [1 - 1/n, 1]) \mid A \in \mathcal{U}(x), n \in \mathbb{N}\}]_{R(X)} = \{\Theta_X(A) \cup U(A) \mid A \in \mathcal{U}(x)\}_{R(X)}$  by definition of  $H$ . As we see in Lemma 2.10, there exists an element  $B \in \mathcal{U}(x)$  such that  $\Theta_X(B) \subset \cup_{\Theta(x) \in \sigma} |\sigma|$  and  $B \subset \cup_{\Theta(x) \in \sigma} |\sigma|$  for every  $B \in \mathcal{U}(x)$ . Thus,  $H(\mathcal{U}(x) \times \mathcal{I}(1)) \subset [\cup_{\Theta(x) \in \sigma} |\sigma|]$  concluding that  $H(\mathcal{U}(x) \times \mathcal{I}(1)) \rightarrow \Theta_X(x)$ .  $\square$

**Lemma 2.12.** *Let  $X$  be a graph,  $Y$  be a compact metric space with the topology induced by the metric and suppose that  $f : Y \rightarrow R(X)$  is continuous. Then  $f_d := \Theta_X f : Y \rightarrow R(X)$  is a continuous map such that*

- (1)  $Im(f_d) \subset X \subset R(X)$ ,
- (2) If  $f(y) \in X \subset R(X)$ , then  $f_d(y) = f(y)$ ,
- (3)  $f_d \simeq f$ .

*In addition, if  $X$  is locally finite, then  $f$  satisfies that*

- (4)  $\#(Im(f_d)) < \infty$

*Proof.* First suppose that  $X$  is any graph. Let  $f : Y \rightarrow R(X)$  a morphism in PreTop, and define  $f_d := \Theta_X f$ . Then by definition  $Im(f_d) \subset X$  and  $f_d(y) = f(y)$  if  $f(y) \in X$ .

To observe that  $f_d \simeq f$ , Proposition 2.11 establishes that  $\Theta_X$  is homotopic to the identity  $1_{R(X)} : R(X) \rightarrow R(X)$ . Thus,  $f_d = \Theta_X f \simeq 1_{R(X)} f = f$ .

In the last part of the proof, suppose that  $X$  is locally finite. Since  $f_d$  is continuous, being the composition of continuous maps, and  $cl(f_d(y))$  is the minimal neighborhood, then for every  $y \in Y$  there exists  $r_y > 0$  such that  $f_d(B_{r_y}(y)) \subset cl(f_d(y))$  since  $cl(f_d(y))$  is a neighborhood of  $f_d(y)$  in  $R(X)$ . Given that  $X$  is locally finite, then  $\#f_d(B_{r_y}(y)) \leq \#(cl(f_d(y)) \cap X) < \infty$ . Since  $Y$  is compact, there exists a finite number of points of  $Y$ , say  $\{y_1, \dots, y_k\}$ , such that  $Y = \cup_{i=1}^k B_{r_{y_i}}(y_i)$ . Thus  $\#f_d(Y) \leq \sum_{i=1}^k \#f_d(B_{r_{y_i}}(y_i))$ , i.e.  $\#f_d(Y)$  is finite.  $\square$

The following lemma removes the sufficient condition in Lemma 2.12 that  $X$  be a locally finite graph.

**Lemma 2.13.** *Let  $X$  be a graph,  $Y$  be a compact metric space with the topology induced by the metric,  $A \subset Y$  a compact subset,  $v \in X$ , and suppose that  $f : (Y, A) \rightarrow (R(X), v)$  be continuous. Then there exists  $f'_d : (Y, A) \rightarrow (X, v)$  a continuous map such that*

- (1)  $Im(f'_d) \subset X \subset R(X)$ .
- (2)  $\#(Im(f'_d)) < \infty$ .
- (3)  $f'_d(A) = v$ .
- (4)  $f'_d \simeq f$ .

*Proof.* Define  $f_d := \Theta_X f : Y \rightarrow R(X)$ . By Lemma 2.12, we obtain that

- $Im(f_d) \subset X \subset R(X)$ .
- $f_d(A) = v$ , because the one hypothesis is that  $f(A) = v$ .
- $f_d \simeq f$ .

The following construction is very similar to the subsequent proof of Lemma 2.16 but with some nuances: Since for every  $x \in X$ ,  $cl(x)$  is the minimal neighborhood of  $x$ , so for all  $y \in Y$  there exists  $r_y > 0$  such that  $f_d(B_{r_y}(y)) \subset cl(f_d(y))$ . In addition, since  $A$  is a compact subset of a topological Hausdorff space by hypothesis,  $A$  is closed, so we can choose  $r_y$  sufficiently small so that  $B_{r_y}(y) \cap A = \emptyset$  for every  $y \in Y \setminus A$ . Then  $\{B_{r_{y_i}/2}(y_i) \mid y_i \in Y\}$  is an open cover of the compact space  $Y$ , and there exists a finite number of points  $\{y_1, \dots, y_n\}$  such that

$$Y = \bigcup_{i=1}^n B_{r_{y_i}/2}(y_i)$$

We need two additional observations before we can construct a candidate for  $f'_d$ . First, observe that if  $y_i \notin A$ , then  $B_{r_{y_i}/2}(y_i) \subset B_{r_{y_i}}(y_i) \subset Y \setminus A$ . Second, consider  $y_i$  and  $y_j$  such that  $B_{r_{y_i}/2}(y_i) \cap B_{r_{y_j}/2}(y_j) \neq \emptyset$ , then

$$d(y_i, y_j) < (r_{y_i} + r_{y_j})/2 \leq \max(r_{y_i}, r_{y_j})$$

and thus  $y_i \in B_{r_{y_j}}(y_j)$  or  $y_j \in B_{r_{y_i}}(y_i)$ ; both imply that  $f_d(y_i) \in cl(f_d(y_j))$ , recalling that the closure structure on  $X \subset R(X)$  is symmetric (i.e.  $y_i, y_j \in X \subset R(X)$  and  $y_i \in cl(y_j) \iff y_j \in cl(y_i)$ ), since the closure structure on  $X$  is induced by a graph.

For every  $y_i \in \{y_1, \dots, y_n\}$ , define  $f_i : Y \rightarrow X \subset R(X)$  by

$$f_i(y) := \begin{cases} f_d(y_i) & y \in B_{r_{y_i}/2}(y_i), \\ f_{i-1}(y) & \text{otherwise,} \end{cases}$$

where  $f_0 = f_d$ . We first note that, by definition, since  $Im(f_d) \subset X \subset R(X)$ , it follows that  $Im(f_i) \subset X \subset R(X)$  for every  $i \in \{1, \dots, n\}$ . We now claim that, for every  $i \in \{1, \dots, n\}$ , each  $f_i$  is continuous,  $f_i \simeq f_{i-1}$ , and  $f_i(A) = v$ .

By the observations above,  $B_{r_{y_i}/2}(y_i) \subset Y \setminus A$  for any  $y_i \in \{y_1, \dots, y_n\}$  with  $y_i \in Y \setminus A$ , and therefore for every such  $y_i \in Y \setminus A$ ,  $f_i(y) = f_{i-1}(y)$  for any point  $y \in A$ . Also note that for any  $y_i \in A$ ,  $f_i(y_i) = v$ . Since, by construction,  $A$  is covered by the  $B_{r_{y_i}}(y_i)$  with  $y_i \in A$ , we conclude that  $f_i(A) = v$ .

We now prove that  $f_i$  is continuous, by proceeding by induction over  $i$ . Observe first that  $f_0$  is continuous by Lemma 2.12, we obtain the base case. Suppose that  $f_{i-1}$  is continuous and consider the following three cases:

Case 1:  $y \in B_{r_{y_i}/2}(y_i)$ . Then there exists  $r > 0$  such that  $B_r(y) \subset B_{r_{y_i}/2}(y_i)$ , and then  $f_i(y) = f_i(y_i) = f_d(y_i)$ , and, in particular,  $f_i(B_r(y)) = f_i(B_{r_{y_i}/2}(y_i)) = \{f_i(y)\} \subset cl(f_i(y_i)) = cl(f_i(y))$ .

Case 2:  $y \in \partial B_{r_{y_i}/2}(y_i)$ . By the induction hypothesis,  $f_{i-1}$  is continuous, and therefore there exists  $r > 0$  such that  $B_r(y) \subset B_{r_{y_i}}(y_i)$  and  $f_{i-1}(B_r(y)) \subset cl(f_{i-1}(y))$ .

Now define  $\Lambda := \{k \mid y \in B_{r_{y_k}}(y_k)\}$  and observe that  $f_{i-1}(y) \in \{f_d(y)\} \cup \{f_d(y_k) \mid k \in \Lambda\}$  and  $y \in B_{r_{y_k}}(y_k) \cap B_{r_{y_i}}(y_i) \neq \emptyset$ .

In addition,  $y \in B_{r_{y_i}}(y_i)$ , and therefore  $f_d(y) \in cl(f_d(y_i))$  by the hypothesis on  $r_{y_i}$ . Thus, given that the closure structure on  $X \subset R(X)$  is symmetric, it follows that  $f_d(y_i) \in cl(f_{i-1}(y))$ . Putting this together and observing that  $f_{i-1}(y) = f_i(y)$  by definition, we have

$$f_i(B_r(y)) \subset f_{i-1}(B_r(y)) \cup \{f_d(y_i)\} \subset cl(f_{i-1}(y)) \cup \{f_d(y_i)\} \subset cl(f_{i-1}(y)) = cl(f_i(y)).$$

Case 3:  $y \notin \overline{B_{r_{y_i}/2}(y_i)}$ . Then there exists  $r > 0$  such that  $B_r(y) \subset Y \setminus \overline{B_{r_{y_i}/2}(y_i)}$ . Thus  $f_i(B_r(y)) = f_{i-1}(B_r(y)) \subset cl(f_{i-1}(y)) = cl(f_i(y))$ .

Finally, we prove the existence of a homotopy between  $f_i$  and  $f_{i-1}$ . Define  $H_i : Y \times I \rightarrow X \subset R(X)$  such that

$$H_i(y, t) := \begin{cases} f_i(y) & t = 1 \\ f_{i-1}(y) & t \in [0, 1) \end{cases}$$

for  $i \in \{1, \dots, n\}$ . Observe that for  $t \in [0, 1)$  we can find  $r > 0$  small enough such  $H_i(B_r(y, t)) = f_{i-1}(B_r(y, t))$ , and therefore  $H$  is continuous on  $Y \times [0, 1)$ .

We now show that  $H$  is continuous at any point of the form  $(y, 1) \in Y \times [0, 1]$ , considering the same three cases as above:



Case 1:  $y \in B_{r_{y_i}/2}(y_i)$ . Define  $\Lambda := \{k \mid y \in B_{r_{y_k}/2}(y_k)\}$ , then

$$f_d(y) \in f(B_{r_{y_k}/2}(y_k)) \subset cl(f_d(y_k))$$

for every  $k \in \Lambda$  and there exists  $r > 0$  such that  $f_d(B_r(y)) \subset cl(f_d(y))$  and  $B_r(y) \subset B_{r_{y_k}/2}(y_k)$  for every  $k \in \Lambda$ . Recall that,  $\forall y \in Y$ ,  $r_y > 0$  was chosen so that  $f_d(B_{r_y}(y)) \subset cl(f_d(y))$ . Therefore,  $B_r(y) \subset B_{r_{y_i}/2}$  implies that  $f_d(B_r(y)) \subset cl(f_d(y_i))$ . Also, by construction,  $B_{r_{y_k}}(y_k) \cap B_{r_{y_i}}(y_i) \neq \emptyset$  for all  $k \in \Lambda$ , and from the observations above, it follows that  $f_d(y_k) \in cl(f_d(y_i))$  for all  $k \in \Lambda$ . This implies that

$$\begin{aligned} f_{i-1}(B_r(y)) &\subset f_d(B_r(y)) \cup \{f_d(y_j) \mid j \in \Lambda\} \subset \\ &cl(f_d(y_i)) \cup \{f_d(y_j) \mid j \in \Lambda\} \subset cl(f_d(y_i)). \end{aligned}$$

Furthermore,  $f_i(y) = f_d(y_i)$  and  $f_i(B_r(y)) = \{f_d(y_i)\}$  by the definition of  $f_i$  and the choice of  $r > 0$ . We conclude that

$$H_i(B_r(y, 1)) \subset f_i(B_r(y)) \cup f_{i-1}(B_r(y)) \subset cl(f_d(y_i)) = cl(f_i(y)) = cl(H_i(y, 1)),$$

as desired.

Case 2:  $y \in \partial B_{r_{y_i}/2}(y_i)$ . Then there exists  $r > 0$  such that  $f_i(B_r(y)) \subset cl(f_i(y))$  and  $f_{i-1}(B_r(y)) \subset cl(f_{i-1}(y)) = cl(f_i(y))$ . Concluding  $H_i(B_r(y, 1)) \subset f_i(B_r(y)) \cup f_{i-1}(B_r(y)) \subset cl(f_i(y))$ .

Case 3:  $y \notin \overline{B_{r_{y_i}/2}(y_i)}$ . Then there exists  $r > 0$  such that

$$\begin{aligned} H_i(B_r(y, 1)) &= f_i(B_r(y)) = f_{i-1}(B_r(y)) \subset \\ &cl(f_{i-1}(B_r(y))) = cl(f_i(B_r(y))) = cl(H_i(B_r(y, 1))). \end{aligned}$$

Since we replace the values in the image of  $B_{r_{y_i}/2}(y_i)$  by  $y_i$ , then  $f_n$  has just a finite number of value in its image. It now follows that  $f'_d := f_n$  satisfies the conclusion of the lemma.  $\square$

In the consecutive,  $\Theta_X f$  is going to denote  $f'_d$ .

In Proposition 3.6 and Theorem 3.2 below, we have that the domain is compact and connected so we can apply Lemma 2.13, which is all we will use in the remainder of the article; however, we keep Lemma 2.12 above to provide a different sufficient condition for the image of the discrete modification to be finite which may be useful in the future.

The maps  $f_i$  in the proof of Lemma 2.13 are an example of the following definition: The  $v$ -flood; this map is an adaption from the  $v$ -flood in digital topology [15].

**Definition 2.14** (The  $v$ -flood). Given a continuous map  $f : Y \rightarrow R(X)$  and  $v \in X \subset R(X)$ , and let  $B \subset Y$ . We define the not-necessarily continuous function  $f_B^v : Y \rightarrow R(X)$

$$f_B^v(y) := \begin{cases} v & y \in B \\ f(y) & \text{otherwise} \end{cases}$$

which we call *the flood of  $f$  on  $B$  with  $v$*  or *the  $v$ -flood of  $f$  on  $B$* . (See Figure 2.)

**Lemma 2.15.** *Let  $Y$  be a metric space with the topology induced by the metric. Let  $f : Y \rightarrow R(X)$  be a function such that  $Im(f) \subset X \subset R(X)$  and let  $v \in X \subset R(X)$ .*




FIGURE 2. Illustration of a flood. The pictures represent the image in a region of  $Y$  with blue, green and red representing point in  $X$ ; in particular  $v$  is red. The first picture represents the image of  $f$ . In the second picture the gray transparent zone is  $B$  in that region. The last picture represents the image of  $f_B^v$ .

If for every  $y \in Y$  such that  $f(y) = v$ , there exists an  $r_y > 0$  such that  $f(B_{r_y}(y)) \subset cl(v)$ , then  $f_B^v$  is continuous in  $X \subset R(X)$  and  $f \simeq f_B^v$  for

$$B := \bigcup_{y \in f^{-1}(v)} B_{r_y/2}(y)$$

and any  $r'_y \leq r_y$ .

*Proof.* First we show continuity of  $f_B^v$ . We consider two cases:

- (1) Let  $y \in B \subset Y$ . Since  $B$  is open in  $Y$ , being the union of open sets, there exists  $r > 0$ , in particular  $r_y$ , such that  $B_r(y) \subset B$  and therefore  $f_B^v(B_r(y)) = v \in cl(v)$ . In particular,  $f_B^v(B_r(y)) \subset cl(v)$ .
- (2) Now let  $y \in Y \setminus B$ . Then by continuity of  $f$ , and being  $cl(f(y))$  the minimal neighborhood of  $f(y)$  in  $R(X)$ , there exists  $r > 0$  such that  $f(B_r(y)) \subset cl(f(y))$ . Suppose that:

- (a)  $v \in cl(f(y))$ . Then  $f_B^v(B_r(y)) \subset f(B_r(y)) \cup \{v\} \subset cl(f(y))$ .
- (b)  $v \notin cl(f(y))$ . Since  $cl(f(y))$  is the minimal neighborhood, there exists  $r' > 0$  such that  $f(B_{r'}(y))$  does not contain which is not neighbor of  $f(y)$ , in particular  $v \notin f(B_{r'}(y))$ . That  $r'$  has to be less or equal to the distance from  $y$  to any point with image  $v$  under  $f$ ; thus  $B_{\min\{r,r'\}/2}(y) \subset Y \setminus B$  and  $f^v(B_{\min\{r,r'\}/2}(y)) = f(B_{\min\{r,r'\}/2}(y))$ . Therefore  $f_B^v(B_{\min\{r,r'\}/2}(y)) = f(B_{\min\{r,r'\}/2}(y)) \subset cl(f(y)) = cl(f^v(y))$

The next step is to define the homotopy. We define  $H : Y \times I \rightarrow X \subset R(X)$  by

$$H(y, t) := \begin{cases} f(y) & \text{if } t \in [0, 1) \\ f_B^v(y) & \text{if } t = 1 \end{cases}$$

Analogous to Proposition 2.11, observe that  $H$  is continuous in  $Y \times [0, 1)$ . Now, consider a point  $(y, 1)$ . We have two cases:

- (1) If  $f(y) = f^v(y)$ , then there exist  $r, r' > 0$  such that  $f(B_r(y)) \subset cl(f(y)) = cl(f^v(y))$  and  $f_B^v(B_{r'}(y)) \subset cl(f^v(y))$ , thus  $H(B_{\min\{r,r'\}}(y), 1) \subset f(B_r(y)) \cup f_B^v(B_{r'}(y)) \subset cl(f^v(y)) = cl(H(y, 1))$ .
- (2) If  $f(y) \neq f^v(y)$ , then  $f_B^v(y) = v$  and there exists  $r > 0$  and  $y_0 \in f^{-1}(v)$  such that  $f(B_r(y)) \subset f(B_{r_{y_0}}(y_0)) \subset cl(v)$ . For the same reason, there exists  $r' > 0$  such that  $f_B^v(B_{r'}(y)) = \{v\}$ , then  $f_B^v(B_{\min\{r,r'\}}(y)) = f(B_{\min\{r,r'\}}(y)) \cup \{v\} \subset cl(v)$ , and  $H$  continuous.  $\square$

Later when we work with homotopy groups, then we need a version of *Lemma 2.15* which conserve a point. We prove the extension of this result for preserving a compact set.

**Lemma 2.16.** *Let  $Y$  be a metric space with topology induced by the metric and  $A \subset Y$  compact, and let  $v, v' \in X \subset R(X)$ . Let  $f : (Y, A) \rightarrow (R(X), v')$  be a function such that  $\text{Im}(f) \subset X \subset R(X)$ . If for every  $y \in Y$  such that  $f(y) = v$ , there exists  $r_y > 0$  such that*

- $f(B_{r_y}(y)) \subset \text{cl}(v)$  and
- if  $v \neq v'$ ,  $A \cap B_{r_y}(y) = \emptyset$ ,

then  $f_B^v(A) = v'$ ,  $f_B^v$  is continuous in  $X \subset R(X)$  and  $f \simeq f_B^v \text{ rel}(A)$  in the closure space  $R(X)$  for

$$B := \bigcup_{y \in f^{-1}(v)} B_{r'_y/2}(y)$$

with  $r'_y \leq r_y$ .

*Proof.* First we prove that  $f_B^v(A) = v'$ . If  $v = v'$  there is nothing to do, by hypothesis  $f_B^v(f^{-1}(v)) = v$ . On the other hand, if  $v \neq v'$ , then  $B \subset Y \setminus A$ . Thus  $f_B^v(A) = f(A) = v'$  by definition.

Observe that  $f_B^v$  is continuous and  $f \simeq f_B^v$  for Lemma 2.15. Also in Lemma 2.15, we define the homotopy such that  $H(-, 1) = f_B^v$  and  $H(-, t) = f$  for every  $t \in [0, 1]$ . Thus  $H(A, t) = v'$ , and then the homotopy is relative to  $A$ .  $\square$

Given a compact metric space  $Y$  and  $A \subset Y$  compact, for any map  $f : Y \rightarrow R(X)$  such that  $f(A) = v' \in X$ , we have that in the image of  $\Theta_X f : Y \rightarrow X \subset R(X)$  there is just a finite bunch and  $\Theta_X f(A) = v'$  by Lemma 2.12. In addition, for every  $y \in Y$  such that  $f(y) = v_0$ , by continuity and  $A$  being a compact, we can find  $r_y > 0$  such that  $f(B_{r_y}(y)) \subset \text{cl}(v_0)$  and  $A \cap B_{r_y}(y) = \emptyset$ . Thus, by Lemma 2.16, we conclude that  $(\Theta_X f)_{B_0}^{v_0} \simeq \Theta_X f \simeq f$  and  $(\Theta_X f)_{B_0}^{v_0}(A) = v'$  with

$$B_0 = \bigcup_{y \in (\Theta_X f)^{-1}(v_0)} B_{r_y/2}(y)$$

Observe we can repeat this process for  $(\Theta_X f)_{B_0}^{v_0}$ ,  $v_1$  and building in a similar fashion  $B_1 \subset Y$ ; obtaining  $((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1}$  which is homotopy to  $(\Theta_X f)_{B_0}^{v_0}$  and preserves the image in  $A$ .

Then we repeat this process a finite number of times for all of the vertices in  $\text{Im}(\Theta_X f)$ , obtaining

$$\begin{aligned} f &\simeq (\Theta_X f)_{B_0}^{v_0} \simeq ((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1} \simeq \dots \simeq (\dots ((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1}) \dots)_{B_n}^{v_n} \\ v' &= (\dots ((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1}) \dots)_{B_n}^{v_n}(A). \end{aligned}$$

For simplification, we denote  $(\dots ((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1}) \dots)_{B_n}^{v_n}$  by  $f^{v_n}$ .

**Proposition 2.17.** *Suppose that we have the sequences  $\{y_n\} \rightarrow x$  and  $\{z_n\} \rightarrow x$  in a metric space  $X$  and we have a function  $f : X \rightarrow Y$  (not necessarily continuous) such that  $\#Y < \infty$ . Then we can take a subsequence of  $\{(y_n, z_n)\}$  such that  $f(y_n) = f(y_k)$  and  $f(z_n) = f(z_k)$  for every  $k, n$ .*

*Proof.* Observe that the image of the sequence contains at most  $(\#Y)^2$  points. If we suppose that we cannot make that construction, then for every  $(v, w) \in Y \times Y$  there exists a  $r_{v,w}$  such that through that point the image cannot be  $(v, w)$ . If we

take  $r := \min_{v,w} r_{v,w}$  ( $\rightarrow\leftarrow$ ). From that  $r$ , we cannot take more point and then their images are out of  $Y$ .  $\square$

**Lemma 2.18.** *Let  $Y$  be a compact metric space and  $X$  a graph. Given a continuous map  $f : Y \rightarrow R(X)$  with image  $\{v_0, \dots, v_m\}$ , for every  $x \in Y$  there exists  $r_x > 0$  such that if  $v, w \in f^{v_m}(B_{r_x}(x))$ , then  $v \in cl(w)$  (equivalently  $w \in cl(v)$ ).*

*Proof.* We proceed by contradiction. Suppose that there exists  $x \in Y$  such that for every  $r > 0$  there exists a pair  $w_r, w'_r \in f^{v_m}(B_r(x))$  we have that  $w_r \notin cl(w'_r)$ . We can restrict just to the elements in  $1/k$  for every  $k \in \mathbb{N}$ ; in consequence we have elements  $y_k, z_k \in X$  such that  $y_k, z_k \in B_{1/k}(x)$ ,  $f^{v_m}(y_k) = w_{1/k}$  and  $f^{v_m}(z_k) = w'_{1/k}$ . Thus we have the sequences  $\{y_k\}$  and  $\{z_k\}$  converge to  $x$ ; therefore, by Proposition 2.17, we can take a subsequence of both such that the image of all of the elements of the subsequence  $\{y_k\}$  (resp.  $\{z_k\}$ ) is a fixed element  $w$  (resp.  $w'$ ).

We name  $v := f^{v_m}(x)$ , and divide this proof in two cases:

(1) Suppose that  $w < v$  or  $w' < v$ , we obtain a direct contradiction. The elements only change their values to  $w$  (resp.  $w'$ ) only appears at  $f^w$  (resp.  $f^{w'}$ ), and in  $f^v(B_r(x)) = v$  for some  $r > 0$ . (See Figure 3.)




FIGURE 3. Illustration of  $w < v$  or  $w' < v$ . The pictures represent the image in a region of  $Y$  with blue, green and red representing  $w$ ,  $w'$  and  $v$  respectively. The white point with black contour is  $x$ . From left to right: The first image is before  $f^v$ . Second image represents a point with image  $v$  and its neighborhood. The last one represents  $f^v$ . (We just ignored all of the rest of points in the image.)

(2) Suppose that  $w > v$  and  $w' > v$ . Without loss of generality, we can assume that  $w < w'$ . Observe once again that in the map  $f^v$  we have a  $r > 0$  such that the image of  $B_r(x)$  is  $v$ . Then in  $f^v$  the image of  $\{y_k\}$  must become  $w$ . Take any element  $z \in S^m$  such that  $f^w(z) = w'$ , then

$$\lim_n d(y_k, z) \rightarrow d(x, z),$$

obtaining that there are no points generated in  $f^{w'}$  with image  $w'$  outside of  $B_{d(x,z)/2}(z)$ , which implies that there are no points with image generated in  $f^{w'}$  with image  $w'$  inside of  $B_{r/2}(x)$ . ( $\rightarrow\leftarrow$ ) (See Figure 4.)

Observe that  $v = w$  or  $v = w'$  is also an automatic contradiction.  $\square$

The discrete modification (Definition 2.9) is built to convert all of the point in the image into vertexes. The next definition seeks to perform an inverse process, moving from a function with only image at the vertexes to one with image at  $|VR(X)|$ .




FIGURE 4. Illustration of  $v < w < w'$ . The pictures represent the image in a region of  $Y$  with blue, green and red representing  $w$ ,  $w'$  and  $v$  respectively. The white point with black contour is  $x$ . From left to right: The first image represent  $f^w$ , and then we must have a sequence with image  $w$ ; we have a green point in the contour of the red circle. In the second image, the circles the distance and the half of the distance between the green and the white point  $x$ . The last one represents  $f^{w'}$  and shows the space without green points. (We just ignored all of the rest of points in the image.)

**Definition 2.19** (The Convex Transformation). Let  $X$  be a graph, be  $Y$  a triangulizable compact space with a triangulation (with a finite number of triangles)  $K$ ,  $\mathcal{U}$  an open cover of  $Y$  and  $f : Y \rightarrow X \subset R(X)$  such that:

- (1) For every  $U \in \mathcal{U}$ , all  $y, y' \in U$  satisfies that  $f(y) \in c_X(f(y'))$ .
- (2) Every simplex in  $Y$  is totally contained in some  $U \in \mathcal{U}$ .

We define the *convex transformation of  $f$* , denoted by  $f_K^{\text{conv}} : Y \rightarrow |VR(X)|$ , such that for every simplex in the triangulation of  $Y$  we look at its vertices, say  $\{y_0, \dots, y_n\}$ , and define

$$f_K^{\text{conv}} \left( \sum_{i=0}^n t_i y_i \right) = \sum_{i=0}^n t_i f(y_i)$$

such that  $t_i \geq 0$  and  $\sum_{i=0}^n t_i = 1$

the convex sums  $\sum t_i y_i$  and  $\sum t_i f(y_i)$  are an abuse of notation for convenience;  $\sum t_i y_i$  means the image of the point in the triangulation in  $Y$ , and  $\sum t_i f(y_i)$  means the convex sum of the vertices  $f(y_i)$  seen in  $|VR(X)|$ , which has the same set as  $R(X)$ .

When the context is clear, we write  $f^{\text{conv}}$  omitting the triangulation of  $Y$  in the notation.

**Lemma 2.20.** Consider  $X$  a graph,  $Y$  a triangulizable compact space with a triangulation  $K$ ,  $\mathcal{U}$  and  $f$  as in Definition 2.19.  $f_K^{\text{conv}} : Y \rightarrow (|VR(X)|, c_{|VR(X)|})$  is well-defined.

*Proof.* To show that  $f_K^{\text{conv}}$  is well-defined, it is enough to show that if of two simplexes, say  $\Delta^n = \{v_0, \dots, v_n\}$  and  $\Delta^m = \{w_0, \dots, w_m\}$ , share a face, say  $\Delta^k = \{x_0, \dots, x_n\}$  such that  $x_i = v_i = w_i$  for  $i \in [k]$ , then the value  $f_K^{\text{conv}}$  is exactly the same in shared face. For  $t_i, s_j, r_a \in [0, 1]$  for every  $i \in [n], j \in [m], a \in [k]$  such that  $\sum_{i=0}^n t_i = \sum_{j=0}^m s_j = \sum_{a=0}^k r_a = 1$  and  $t_i = s_j = 0$  for  $i \in [n] \setminus [k], j \in [m] \setminus [k]$  we

that

$$\begin{aligned} f_K^{\text{conv}} \Big|_{\Delta^k \subset \Delta^n} \left( \sum_{i=0}^n t_i x_i \right) &= \sum_{i=0}^n t_i f^{v_m}(v_i) = \sum_{a=0}^k r_a f^{v_m}(x_a) = f_K^{\text{conv}} \Big|_{\Delta^k} \left( \sum_{a=0}^k r_a x_a \right) \\ &= \sum_{j=0}^m s_j f^{v_m}(w_j) = f_K^{\text{conv}} \Big|_{\Delta^k \subset \Delta^m} \left( \sum_{j=0}^m s_j w_j \right). \end{aligned}$$

Obtaining that  $f_K^{\text{conv}}$  is well-defined.  $\square$

**Lemma 2.21.** *Consider  $X, Y, \mathcal{U}$  and  $f$  as in Definition 2.19.  $f^{\text{conv}} : Y \rightarrow (|VR(X)|, c_{|VR(X)|})$  is continuous and  $f^{\text{conv}} \simeq f$  in  $R(X)$ .*

*Proof.* By construction, it follows that  $f^{\text{conv}}$  is continuous in geometric realization of the simplexes because that map is just the geometric realization from an  $n$ -simplex. Now, since we have a locally finite closed cover, observe that  $f^{\text{conv}}$  is continuous by 17 A.18 in [8].

The next step is to build the homotopy. Let  $H : S^n \times I \rightarrow R(X)$  be the map

$$H(x, t) := \begin{cases} f^{\text{conv}}(x) & \text{if } t \in [0, 1) \\ f(x) & \text{if } t = 1. \end{cases}$$

As we have seen before, continuity in  $S^n \times [0, 1)$  is straightforward: we can take a radius small enough such that  $B_r(x, t) \cap S^n \times \{1\} = \emptyset$ , and then  $H(B_r(x, t)) = B_r(x, t)$ ; thus, by continuity in  $f^{\text{conv}}$ , we obtain the continuity in that set. Consider a point  $(x, 1)$ ; by construction, the image of every simplex  $\Delta$  in  $Y$  is inside an open set  $U$  such that for all  $y, y' \in U$  satisfies that  $f(y) \in cl(f(y'))$ . Thus, every point  $z \in f(\delta)$  is neighbor of the vertexes as well as the elements of the geometric realization of that simplex (a.k.a  $f^{\text{conv}}$  of them). Therefore, remembering that  $f(x) \in X \subset R(X)$ , for every simplex  $\{\Delta_1, \dots, \Delta_a\}$  such that  $x \in \Delta_i$  we have that  $\cup_{i=1}^a f^{\text{conv}}(\Delta_i) \subset cl(f(x))$ , and then for  $r > 0$  such that  $B_r(f(x)) \subset \cup_{i=1}^a \Delta_i$ , we obtain that  $H(B_r(x, 1)) \subset f^{\text{conv}}(x) \cup f(x) \subset cl(f(x))$ .  $\square$

### 3. WEAK HOMOTOPY EQUIVALENCE OF $(X, \lambda)$ AND $|VR(X, \lambda)|$

In this section, we prove that the homomorphism  $\pi_n(1_{|VR(X)|}) : \pi_n(|VR(X)|) \rightarrow \pi_n(R(X))$  is an isomorphism, with  $1_{|VR(X)|} : |VR(X)| \rightarrow R(X)$  the inclusion map. The core of the proof is showing that functions  $S^n \rightarrow R(X)$  and  $S^n \times I \rightarrow R(X)$  are homotopic to maps whose image is in  $X \subset R(X)$  and which have the property that, for every point  $x$  in the domain, there exists a neighborhood  $U$  of  $x$  such that every pair of points in  $U$  are neighbors. We can then prove that there is a triangulation of the domain such that the latter map is homotopic to some realization of a simplicial map from this triangulation to  $R(X)$ .

We define  $S^n := \{x \in \mathbb{R}^{n+1} \mid |x| = 1\}$  with its subspace topological structure. We will also use a metric  $d : S^n \times S^n \rightarrow \mathbb{R}$  which we define to be the geodesic distance using the standard Riemmanian metric.

We will need the following classical result before starting the proof that  $\pi_n(1_{|VR(X)|})$  is an epimorphism.

**Lemma 3.1** (Lebesgue's number lemma; [25], 22.5). *Let  $X$  be a compact metric space and  $\mathcal{U}$  an open cover of  $X$ , then there exists a number  $\delta > 0$  such that every subset of  $X$  having diameter less than  $\delta$  is contained in some member of the cover.*

We now show the first main result of this section.

**Theorem 3.2.** *Let  $X$  be a graph. Every map  $f : S^n \rightarrow R(X)$  is homotopic to the composition of the identity  $1_{|VR(X)|} : |VR(X)| \rightarrow R(X)$  and some map  $g : S^n \rightarrow |VR(X)|$ . This implies that  $\pi_n(1_{|VR(X)|})$  is an epimorphism.*

*Proof.* Let  $f : (S^n, *) \rightarrow (R(X), v_\otimes)$  a continuous based map with basepoints  $* \in S^n, v_\otimes \in X$ . Since  $S^n$  is compact and connected, by Lemma 2.13  $\Theta_X f \simeq f$  in  $R(X)$  and  $Im(\Theta_X f)$  is a finite number of vertices, say  $\{v_0, \dots, v_m\}$ .

Since  $\Theta_X f$  is continuous and the set  $\{*\}$  is compact, then since  $c(v_0)$  is the minimal neighborhood of  $v_0 \in R(X)$ , for every  $x \in S^n$  such that  $f(x) = v_0$  there exists  $r_x > 0$  such that  $\Theta_X f(B_{r_x}(x)) \subset cl(v_0)$ . If, furthermore,  $v_\otimes \neq v_0$ , then we may choose  $r_x$  such that  $* \notin B_{r_x}(x)$ . Thus  $\Theta_X f \simeq (\Theta_X f)_{B_0}^{v_0}$  by Lemma 2.16, where

$$B_0 := \bigcup_{x \in (\Theta_X f)^{-1}(v_0)} B_{r_x}(x).$$

With similar definitions, for every  $i \in \{0, \dots, m\}$  we can build  $B_i$ , and we define  $f^{v_i}$  as in Lemma 2.16 by

$$f^{v_i} := (\dots((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1} \dots)_{B_i}^{v_i},$$

and we observe that

$$f \simeq (\Theta_X f)_{B_0}^{v_0} \simeq ((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1} \simeq \dots \simeq (\dots((\Theta_X f)_{B_0}^{v_0})_{B_1}^{v_1} \dots)_{B_m}^{v_m} =: f^{v_m}$$

$$v_\otimes = f^{v_m}(*).$$

Thus, by Lemma 2.18, for every  $x \in S^n$ , there exists  $r > 0$  such that for every  $w, w' \in f^{v_m}(B_r(x))$  we have that  $w \in cl(w')$ ; we denote this open cover of  $S^n$  as  $\mathcal{U}$ .

Take the classical triangulation of  $S^n$  such that  $*$  is one of the vertexes of its simplices and apply the barycentric subdivision until we have that every simplex has diameter less than the Lebesgue's number of the cover  $\mathcal{U}$ . Then, by Lemma 3.1 and Lemma 2.18, the triangulation satisfies the hypothesis of Lemma 2.21, and therefore  $(f^{v_m})^{\mathbf{conv}} \simeq f^{v_m}$ , and, slightly abusing notation,  $(f^{v_m})^{\mathbf{conv}} : S^n \rightarrow |VR(X)|$  is also continuous with the closure operation  $c_{|VR(X)|}$ .

In conclusion,  $f \simeq f^{v_m} \simeq (f^{v_m})^{\mathbf{conv}}$ , and the latter may be lifted to a continuous map  $S^n \rightarrow |VR(X)|$ .

$$\begin{array}{ccc} & & (|VR(X)|, v_\otimes) \\ & \nearrow & \downarrow 1_{|VR(X)|} \\ (S^n, *) & \xrightarrow{(f^{v_m})^{\mathbf{conv}}} & (R(X), v_\otimes) \end{array}$$

□

To prove  $\pi_n(1_{|VR(X)|})$  is a monomorphism, we are going to need two more technical results. Following the notation of [24], we denote by  $sd$  the barycentric subdivision of a simplicial complex and we write  $sd^n$  when we apply the barycentric subdivision  $n$ -times.

**Definition 3.3.** Let  $K_1$  and  $K_2$  be simplicial complexes and let  $f : |K_1| \rightarrow |K_2|$  be a continuous map. A simplicial map  $\varphi : K_1 \rightarrow K_2$  is called a *simplicial approximation* to  $f$  if  $f(\alpha) \in |s|$  implies that  $|\varphi(\alpha)| \in |s|$  for  $\alpha \in |K_1|$  and  $s_2 \in K_2$ .

**Theorem 3.4** (Simplicial-approximation Theorem; [24], 3.4.8). *Let  $(K_1, L_1)$  be a finite simplicial pair and let  $f : (|K_1|, |L_1|) \rightarrow (|K_2|, |L_2|)$  be a map. There exists an integer  $N$  such that if  $n \geq N$  there is a simplicial approximation  $(sd^n K_1, sd^n L_1) \rightarrow (K_2, L_2)$  to  $f$ .*

Thus, considering for example the canonical triangulation  $K_n$  of  $S^n$ , for each homotopy class  $\beta \in \pi_k(|VR(X)|, v)$ , there is a non-negative integer  $m$  and a simplicial map  $sd^m K_n \rightarrow VR(X)$  whose geometric realization represents  $\beta$ .

The following results show that the convex modification is homotopic to the convex modification of the barycentric subdivision (with respect to the closure operation on  $c_{|VR(X)|}$ ). This result is essential in the proof of having a monomorphism and, as we will see, imposes some technical conditions in the selection of balls when we apply the floods in the proof of Lemma 2.16.

**Proposition 3.5.** *Let  $Y$  a triangulizable compact metric space with finite triangulation  $K$  and let  $f : Y = |K| \rightarrow X \subset R(X)$  be such that the image of every simplex in the triangulation  $K$  is a simplex in  $R(X)$ . Then  $f_K^{\text{conv}}$  and  $f_{sd K}^{\text{conv}} : Y = |sd K| \rightarrow |VR(X)|$  are homotopic.*

*Proof.* Since the image of every simplex in  $K$  is a simplex of  $R(X)$ , then every simplex in  $sd K$  is contained in a simplex of  $R(X)$ , then  $f_{sd K}^{\text{conv}}$  is well-defined. Consider an  $n$ -simplex  $\Delta = \{x_0, \dots, x_n\}$  in the triangulation  $sd K$  such that  $x_0 < x_1 < \dots < x_n$  and describe the interval  $I$  as the geometric realization of  $\{0, 1\}$ . Recall that  $|\Delta| \times I$  is a triangulizable metric space with  $(n+1)$ -simplexes such that after any  $(w_i, w'_i) \in \Delta \times \{0, 1\}$  such that either  $w_i < w_{i+1}$  and  $w'_i = w'_{i+1}$  or  $w_i = w_{i+1}$  and  $w'_i < w'_{i+1}$ .

Consider the restriction  $f_K^{\text{conv}}|_{|\Delta|} : |\Delta| \rightarrow |VR(X)|$  and note that  $f_{sd K}^{\text{conv}}|_{|\Delta|} : |\Delta| \rightarrow |VR(X)|$  is, by definition,

$$f_{sd K}^{\text{conv}}|_{|\Delta|} \left( \sum_{i=0}^n t_i x_i \right) := \sum_{i=0}^n t_i f(x_i).$$

Now, triangulate  $|\Delta| \times I$  as described above and define  $H_\Delta : |\Delta| \times I \rightarrow |VR(X)|$  such that for every  $(n+1)$ -simplex  $\{(w_0, w'_0), \dots, (w_{n+1}, w'_{n+1})\}$

$$H_\Delta \left( \sum_{i=0}^{n+1} t_i (w_i, w'_i) \right) := \sum_{i=0}^{n+1} t_i H_\Delta(w_i, w'_i)$$

with  $H_\Delta(w_i, 0) = f_K^{\text{conv}}(w_i)$  and  $H_\Delta(w_i, 1) = f_{sd K}^{\text{conv}}(w_i)|_{|\Delta|}$ .  $H_\Delta$  is continuous because is continuous in every  $(n+1)$ -simplex, being a lineal map when we restrict them to those  $(n+1)$ -simplex.

Define  $H$  such that  $H|_{|\Delta| \times I} = H_\Delta$ .  $H$  is continuous because is continuous in  $|\Delta| \times I$  and by 17 A.18 in [8], and, by the same result, the map  $f_K^{\text{conv}} : |K| \rightarrow |VR(X)|$  is continuous; by the definition of  $H$  we obtain that  $H(y, 0) = f_K^{\text{conv}}(y)$  and  $H(y, 1) = f_{sd K}^{\text{conv}}(y)$ ; concluding that both maps are homotopic.  $\square$

**Proposition 3.6.** *Let  $X$  be a graph.  $\pi_n(1_{|VR(X)|}) : \pi_n(|VR(X)|) \rightarrow \pi_n(R(X))$  is a monomorphism.*

*Proof.* Let  $f : (S^n, *) \rightarrow (|VR(X)|, v_\otimes)$  be such that  $1_{|VR(X)|} \circ f : (S^n, *) \rightarrow (R(X), v_\otimes)$  is homotopic to the constant map  $\hat{v}_\otimes : (S^n, *) \rightarrow (R(X), v_\otimes)$  such that  $\hat{v}_\otimes(x) = v_\otimes$  for every  $x \in S^n$ , i.e. there exists  $H : S^n \times I \rightarrow R(X)$  such that



$H(\{*\} \times I) = v_{\otimes}$ ,  $H(x, 0) = 1_{|VR(X)|} \circ f(x)$  and  $H(x, 1) = \hat{v}_{\otimes}(x) = v_{\otimes}$ . We want to build a continuous homotopy  $\hat{H} : S^n \times I \rightarrow |VR(X)|$  from  $f$  to  $\hat{v}_{\otimes}$ .

Without loss of generality, we can consider  $f$  to be a simplicial-approximation to itself by Theorem 3.4. Let  $K$  denote the triangulation on  $S^n$  used in this simplicial approximation. Since we only need a homotopy from  $f$  to  $\hat{v}_{\otimes}$ , so it is enough to consider a map which we know to be homotopic (in  $|VR(X)|$ ) to the original  $f$ .

By Lemma 2.13, because  $S^n \times I$  is compact, then  $\Theta_X H : S^n \times I \rightarrow X \subset R(X)$  satisfies that  $\Theta_X H \simeq H$  and  $\#Im(\Theta_X H) < \infty$ , say  $\{v_0, \dots, v_m\}$ .

For convenience, we will say that:

- $(x, t)$  is in the bottom (of  $S^n \times I$ ) if  $t = 0$ .
- $(x, t)$  is in the top (of  $S^n \times I$ ) if  $t = 1$ .
- $(x, t)$  is in the middle (of  $S^n \times I$ ) if  $t \in (0, 1)$ .

Since  $\Theta_X H$  is continuous in  $R(X)$  and  $\{*\} \times I$  is compact, and because  $H(x, t) = v_0 \implies \Theta_X(H)(x, t) = v_0$  and  $cl(v_0)$  is the minimal neighborhood of  $v_0$  in  $R(X)$ , we have that for every  $(x, t) \in S^n \times I$  such that  $H(x, t) = v_0$  there exists  $r_{(x,t)} > 0$  such that  $\Theta_X H(B_{r_{(x,t)}}(x, t)) \subset cl(H(x, t))$ . Furthermore, if  $v \neq v_0$  then we may choose  $r(x, y)$  such that  $B_{r_{(x,t)}}(x, t) \cap (\{*\} \times I) = \emptyset$ , and

- if  $(x, t)$  is in the top,  $B_{r_{(x,t)}}(x, t) \cap S^n \times \{0\} = \emptyset$ .
- if  $(x, t)$  is in the middle,  $B_{r_{(x,t)}}(x, t) \cap S^n \times \{0, 1\} = \emptyset$ .
- if  $(x, t)$  is in the bottom,  $B_{r_{(x,t)}}(x, t) \cap S^n \times \{1\} = \emptyset$  and  $B_{r_{(x,t)}}(x, t)$  does not touch any vertex of the triangulation of  $S^n$  except for  $(x, t)$ , which is possible because we have just a finite number of simplexes. We define  $B_0$  as the union of those balls with center  $(x, t) \in (\Theta_X H)^{-1}(v)$ .

Thus, by Lemma 2.16, we obtain that  $(\Theta_X H)_{B_0}^{v_0} \simeq \Theta_X H$ ,  $(\Theta_X H)_{B_0}^{v_0}(\{*\} \times I) = v_{\otimes}$ ,  $(\Theta_X H)_{B_0}^{v_0}(x, 1) = v_{\otimes}$  and the  $v_0$ -flood does not change the value of the vertices on the triangulation of  $S^n$ .

With similar definitions, we can successively build  $B_1, \dots, B_m$  from the  $v_i$ -floods to obtain  $H^{v_m}$ , observing that

$$\begin{aligned} H &\simeq (\Theta_X H)_{B_0}^{v_0} \simeq ((\Theta_X H)_{B_0}^{v_0})_{B_1}^{v_1} \simeq \dots \simeq (\dots ((\Theta_X H)_{B_0}^{v_0})_{B_1}^{v_1}) \dots)_{B_m}^{v_m} =: H^{v_m} \\ v_{\otimes} &= H^{v_m}(\{*\} \times I) \end{aligned}$$

Thus, by Lemma 2.18, for every  $(x, t) \in S^n \times I$  there exists  $r > 0$  such that for every  $w, w' \in H^{v_m}(B_r(x, t))$  we have that  $w \in cl(w')$ ; we denote this open cover of  $S^n \times I$  as  $\mathcal{U}$ .

Take the triangulation  $K$  and let  $I$  be the geometric realization of  $\{0, 1\}$ . Now take the barycentric subdivision in both  $K$  and  $I$  until the diameter of the simplices in  $sd^k K \times sd^k \{0, 1\}$  is less than the Lebesgue number of  $\mathcal{U}$ . Then, by Lemma 3.1 and Lemma 2.18, these objects satisfy the hypothesis of Lemma 2.21. Thus  $(H^{v_m})_{sd^k K \times sd^k \{0, 1\}}^{\text{conv}}$  is continuous in  $|VR(X)|$ ,  $(H^{v_m})_{sd^k K \times sd^k \{0, 1\}}^{\text{conv}}(x, 1) = \hat{v}_{\otimes}(x) = v_{\otimes}$ , and  $(H^{v_m})_{sd^k K \times sd^k \{0, 1\}}^{\text{conv}}(x, 0) = (f^{v_m})_{sd^k K}^{\text{conv}}(x)$ .

Finally, since the points of the vertexes of the triangulation were not changed, we have that

$$(f^{v_m})_{sd^k K}^{\text{conv}} \simeq (f^{v_m})_{sd^{k-1} K}^{\text{conv}} \simeq \dots \simeq (f^{v_m})_{sd K}^{\text{conv}} \simeq (f^{v_m})_K^{\text{conv}} = f$$

in  $|VR(X)|$  by Proposition 3.5. We conclude that  $f \simeq \hat{v}_{\otimes}$  in  $|VR(X)|$ , as desired.  $\square$

**Theorem 3.7.** *Let  $G = (X, E)$  be a graph, and let  $(X, c_G)$  be the induced Čech closure space. Then  $|VR(X)|$  has the same weak homotopy type as  $(X, c_G)$ .*

*Proof.* Given the strong homotopy equivalence in Proposition 2.11 between the maps  $1_X : X \rightarrow R(X)$  and  $\Theta_X : R(X) \rightarrow X$ , we obtain that  $\pi_n(\Theta_X) : \pi_n(R(X)) \rightarrow \pi_n(X)$  is an isomorphism.

In addition, by Theorem 3.2 and Proposition 3.6, we have that  $\pi_n(1_{|VR(X)|}) : \pi_n(|VR(X)|) \rightarrow \pi_n(R(X))$  is also an isomorphism. Thus, we obtain that

$$\pi_n(\Theta_X) \circ \pi_n(1_{|VR(X)|}) : \pi_n(|VR(X)|) \rightarrow \pi_n(X)$$

is an isomorphism. □

#### ACKNOWLEDGMENTS

I would like to thank my thesis advisor, Antonio Rieser, who has helped with several suggestions about the order of the ideas of these paper to improve the delivery and to reduce the number of particular lemmas for specific cases. I am deeply grateful to the AIM Workshop on Discrete and Combinatorial Homotopy Theory, its organizers: Helen Barcelo, Antonio Rieser and Volkmar Welker, and certainly with my one-week teammates: Greg Lupton, Oleg Musin, Nicholas Scoville and Chris Staecker; I learned a great deal from them about how to decompose a problem, discuss the ideas and try to simplify them in order to improve the exposition. I am grateful to Antonio Rieser for telling me about the existence of [19] after I showed him the first versions of this work.

#### REFERENCES

- [1] Michał Adamaszek and Henry Adams, *On Vietoris-Rips complexes of hypercube graphs*, J. Appl. Comput. Topol. **6** (2022), no. 2, 177–192. MR4423676
- [2] Michał Adamaszek, Henry Adams, Ellen Gasparovic, Maria Gommel, Emilie Purvine, Radmila Sazdanovic, Bei Wang, Yusu Wang, and Lori Ziegelmeier, *On homotopy types of Vietoris-Rips complexes of metric gluings*, J. Appl. Comput. Topol. **4** (2020), no. 3, 425–454. MR4130978
- [3] Henry Adams, Johnathan Bush, Nate Clause, Florian Frick, Mario Gómez, Michael Harrison, R. Amzi Jeffs, Evgeniya Lagoda, Sunhyuk Lim, Facundo Mémoli, Michael Moy, Nikola Sadovek, Matt Superdock, Daniel Vargas, Qingsong Wang, and Ling Zhou, *Gromov-hausdorff distances, borsuk-ulam theorems, and vietoris-rips complexes*, 2022.
- [4] Henry Adams and Žiga Virk, *Lower bounds on the homology of Vietoris-Rips complexes of hypercube graphs*, Bull. Malays. Math. Sci. Soc. **47** (2024), no. 3, Paper No. 72, 32. MR4712847
- [5] Eric Babson, Hélène Barcelo, Mark de Longueville, and Reinhard Laubenbacher, *Homotopy theory of graphs*, J. Algebraic Combin. **24** (2006), no. 1, 31–44. MR2245779
- [6] Gunnar Carlsson, *Topology and data*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 2, 255–308. MR2476414
- [7] Gunnar Carlsson, Vin de Silva, Tigran Ishkhanov, and Afra Zomorodian, *On the local behavior of spaces of natural images*, International Journal of Computer Vision **76** (2008), 1–12.
- [8] Eduard Čech, *Topological spaces*, Revised edition by Zdeněk Frolíc and Miroslav Katětov. Scientific editor, Vlastimil Pták. Editor of the English translation, Charles O. Junge, Publishing House of the Czechoslovak Academy of Sciences, Prague; Interscience Publishers John Wiley & Sons, London-New York-Sydney, 1966. MR0211373
- [9] Tien Chih and Laura Scull, *A homotopy category for graphs*, J. Algebraic Combin. **53** (2021), no. 4, 1231–1251. MR4263649
- [10] Davide Carlo Demaria, *On some applications of homotopy to directed graphs*, Proceedings of the Geometry Conference (Milan and Gargnano, 1987), 1987, pp. 183–202 (1989). MR1017928
- [11] Anton Dochtermann, *Hom complexes and homotopy theory in the category of graphs*, European J. Combin. **30** (2009), no. 2, 490–509. MR2489282

- [12] Alexander Grigor'yan, Yong Lin, Yuri Muranov, and Shing-Tung Yau, *Homotopy theory for digraphs*, Pure Appl. Math. Q. **10** (2014), no. 4, 619–674. MR3324763
- [13] Janko Latschev, *Vietoris-Rips complexes of metric spaces near a closed Riemannian manifold*, Arch. Math. (Basel) **77** (2001), no. 6, 522–528. MR1879057
- [14] Sunhyuk Lim, Facundo Mémoli, and Zane Smith, *The Gromov-Hausdorff distance between spheres*, Geom. Topol. **27** (2023), no. 9, 3733–3800. MR4674839
- [15] Gregory Lupton, Oleg Musin, Nicholas A. Scoville, P. Christopher Staecker, and Jonathan Treviño-Marroquín, *A second homotopy group for digital images*, 2023.
- [16] Gregory Lupton, John Oprea, and Nicholas A. Scoville, *Homotopy theory in digital topology*, Discrete Comput. Geom. **67** (2022), no. 1, 112–165. MR4356246
- [17] Gregory Lupton and Nicholas A. Scoville, *Digital fundamental groups and edge groups of clique complexes*, J. Appl. Comput. Topol. **6** (2022), no. 4, 529–558. MR4496690
- [18] Facundo Mémoli, Zane Smith, and Zhengchao Wan, *The Gromov-Hausdorff distance between ultrametric spaces: its structure and computation*, J. Comput. Geom. **14** (2023), no. 1, 78–143. MR4643242
- [19] Nikola Milicevic and Nicholas A. Scoville, *Homotopy and singular homology groups of finite graphs from the closure space perspective*, 2024. <https://meetings.ams.org/math/jmm2024/meetingapp.cgi/Paper/31808> [Accessed: (May 17th, 2024)].
- [20] Conrad Plaut and Jay Wilkins, *Discrete homotopies and the fundamental group*, Adv. Math. **232** (2013), 271–294. MR2989983
- [21] Gerhard Preuss, *Foundations of topology*, Kluwer Academic Publishers, Dordrecht, 2002. An approach to convenient topology. MR2033142
- [22] Antonio Rieser, *Čech closure spaces: A unified framework for discrete and continuous homotopy*, Topology and its Applications **296** (2021), 107613.
- [23] ———, *Cofibration and model category structures for discrete and continuous homotopy*, arXiv, 2022.
- [24] Edwin H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York-Toronto, Ont.-London, 1966. MR0210112
- [25] Stephen Willard, *General topology*, Dover Publications, Inc., Mineola, NY, 2004. Reprint of the 1970 original [Addison-Wesley, Reading, MA; MR0264581]. MR2048350