MAP MONOIDALES AND DUOIDAL ∞ -CATEGORIES

TAKESHI TORII

ABSTRACT. In this paper we give an example of duoidal ∞ -categories. We introduce map \mathcal{O} -monoidales in an \mathcal{O} -monoidal ($\infty, 2$)-category for an ∞ -operad \mathcal{O}^{\otimes} . We show that the endomorphism mapping ∞ -category of a map \mathcal{O} -monoidale is a coCartesian ($\Delta^{\mathrm{op}}, \mathcal{O}$)-duoidal ∞ -category. After that, we introduce a convolution product on the mapping ∞ -category from an \mathcal{O} -comonoidale to an \mathcal{O} -monoidale. We show that the \mathcal{O} -monoidal structure on the duoidal endomorphism mapping ∞ -category of a map \mathcal{O} -monoidale is equivalent to the convolution product on the mapping ∞ -category from the dual \mathcal{O} -comonoidale to the map \mathcal{O} -monoidale.

1. INTRODUCTION

Duoidal categories are categories equipped with two monoidal products in which one is (op)lax monoidal with respect to the other. Duoidal categories are generalizations of braided monoidal categories. Braided monoidal categories can be regarded as pseudomonoids in the monoidal 2-category of monoidal categories and strong monoidal functors. On the other hand, duoidal categories can be regarded as pseudomonoids in the monoidal 2-category of monoidal categories and lax monoidal functors. The notion of duoidal category was introduced by Aguiar-Mahajan by name of 2-monoidal category in [2]. The terminology of duoidal category was proposed by Street in [21].

Examples and applications of duoidal categories are presented in [2] and [21]. We can make sense of monoidal categories enriched over duoidal categories. In [4, 5] duoidal categories play fundamental role in their theory of centers and homotopy centers of monoids in monoidal categories enriched over duoidal categories, and in developing of a generalization of Deligne's conjecture. A general theory of duoidal categories have further been developed in [8]. We can consider bimonoids in duoidal categories, which are generalizations of bialgebras. There are several equivalent conditions for bialgebras to be Hopf algebras. The fundamental theorem of Hopf modules for bimonoids in duoidal categories was studied in [6]. The notion of bimonoids in duoidal categories have been generalized to that of weak bimonoids in [9]. Hopf conditions have further been studied for bimonoids in duoidal endomorphism hom-categories of map monoidales in monoidal bicategories in [7].

As explained in [7], the endomorphism hom-category of a map monoidale in a monoidal bicategory admits the structure of a duoidal category. One of the monoidal products is given by the composition of morphisms and the other is by convolution product. In this paper we generalize this construction in the setting of higher categories.

In higher category theory we replace the category of sets in ordinary category theory to the homotopy theory of spaces based on the homotopy hypothesis [17]. In other words, the theory of $(\infty, 0)$ -categories is equivalent to the homotopy theory of spaces. An (∞, n) -category is an ∞ -category whose k-morphisms are invertible for k > n. On way to view an (∞, n) -category is to regard it as an ∞ -category enriched over $(\infty, n-1)$ -categories. In particular, an $(\infty, 1)$ -category

Date: June 1, 2024 (version 1.0).

²⁰²⁰ Mathematics Subject Classification. 18N60 (primary), 18N70, 18M50, 55U40 (secondary).

Key words and phrases. Monoidale, duoidal ∞ -category, convolution product, ∞ -bicategory, ∞ -operad.

is an ∞ -category enriched over spaces. Another way to view an (∞, n) -category is to regard it as an *n*-category up to coherent homotopy. By taking the set of components of mapping spaces, we can associate to an $(\infty, 1)$ -category its homotopy category which is an ordinary category. An $(\infty, 2)$ -category is regarded as an ∞ -category enriched over $(\infty, 1)$ -categories. By taking homotopy categories of mapping $(\infty, 1)$ -categories, we obtain the homotopy bicategory of an $(\infty, 2)$ -category.

In [22] we introduced the notion of duoidal ∞ -categories, which are counterparts of duoidal categories in the setting of higher categories. In [23] we generalized duoidal ∞ -categories to higher monoidal ∞ -categories and studied their duality property. Higher monoidal categories are generalizations of duoidal categories, which are also introduced in [2]. Higher monoidal categories have multiple monoidal products together with multiple interchange laws. Higher monoidal ∞ -categories are counterparts of higher monoidal categories in higher category theory. In [25] we gave an example of duoidal ∞ -categories of operadic modules. Lurie [18] showed that the $(\infty, 1)$ -category of \mathcal{O} -A-modules in \mathcal{C} admits the structure of an \mathcal{O} -monoidal ∞ -category, where \mathcal{O}^{\otimes} is a small coherent symmetric ∞ -operad, \mathcal{C} is an \mathcal{O} -monoidal $(\infty, 1)$ -category with a sufficient supply of colimits, and A is an \mathcal{O} -algebra in \mathcal{C} . By generalizing this construction, we showed that if \mathcal{P}^{\otimes} is a symmetric ∞ -operad, \mathcal{O}^{\otimes} is a small coherent symmetric ∞ -operad, \mathcal{O} is a small coherent symmetric ∞ -operad, \mathcal{O}^{\otimes} is a small coherent symmetric ∞ -operad, \mathcal{O} is a small coherent symmetric ∞ -operad, \mathcal{O}^{\otimes} is a small coherent symmetric ∞ -operad

The purpose of this paper is to give another example of duoidal ∞ -categories. We show that we can construct a duoidal ∞ -category by the same procedure as in [7] in the setting of higher categories. We consider an \mathcal{O} -monoidal ($\infty, 2$)-category \mathcal{Z} for an ∞ -operad \mathcal{O}^{\otimes} over a perfect operator category. We introduce an \mathcal{O} -monoidale as an \mathcal{O} -monoid object in the underlying ($\infty, 1$)category of \mathcal{Z} . We can consider the 1-full sub ($\infty, 2$)-category \mathcal{Z}^L spanned by left adjoint 1morphisms, which inherits \mathcal{O} -monoidal structure from \mathcal{Z} . A map \mathcal{O} -monoidale is defined to be an \mathcal{O} -monoidale in \mathcal{Z}^L .

For an object A in \mathbb{Z} , the endomorphism mapping ∞ -category $\mathbb{Z}(A, A)$ is a monoidal $(\infty, 1)$ category by composition of morphisms. Let Δ^{op} be the opposite category of nonempty totally ordered sets. We regard it as an ∞ -operad over the perfect operator category \mathbf{O} of finite ordered sets. We formulate the composition monoidal structure on $\mathbb{Z}(A, A)$ by using the ∞ -operad Δ^{op} .

The first main theorem of this paper is as follows.

Theorem 1.1 (cf. Theorem 6.3). Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category and let A be a map \mathcal{O} -monoidale in \mathcal{Z} . The endomorphism mapping ∞ -category $\mathcal{Z}(A, A)$ admits the structure of a coCartesian $(\Delta^{\mathrm{op}}, \mathcal{O})$ -duoidal ∞ -category,

Next, we compare the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)$ with convolution product. We call an \mathcal{O} comonoid object in the underlying $(\infty, 1)$ -category of \mathcal{Z} an \mathcal{O} -comonoidale. For an \mathcal{O} -monoidale A and an \mathcal{O} -comonoidale C, we can define a convolution product on the mapping ∞ -category $\mathcal{Z}(C, A)$. Informally speaking, when \mathcal{O}^{\otimes} is the associative ∞ -operad, the convolution product is given by

$$C \stackrel{c}{\longrightarrow} C \otimes C \stackrel{f \otimes g}{\longrightarrow} A \otimes A \stackrel{m}{\longrightarrow} A$$

for $f, g \in \mathcal{Z}(C, A)$, where *m* is the product on *A* and *c* is the coproduct on *C*. The convolution product gives $\mathcal{Z}(C, A)$ the structure of an \mathcal{O} -monoidal $(\infty, 1)$ -category.

We denote by \mathcal{Z}^R the 1-full sub $(\infty, 2)$ -category of \mathcal{Z} spanned by right adjoint 1-morphisms. By taking adjoint morphisms, there is a dual equivalence between the $(\infty, 1)$ -category of \mathcal{O} -monoidales in \mathcal{Z}^L and the $(\infty, 1)$ -category of \mathcal{O} -comonoidales in \mathcal{Z}^R . We denote by A^* the \mathcal{O} -comonoidale in \mathcal{Z}^R corresponding to a map \mathcal{O} -monoidale A under this equivalence. Since the underlying object of

 A^* is equivalent to that of A, the underlying $(\infty, 1)$ -category of $\mathcal{Z}(A^*, A)$ is equivalent to that of $\mathcal{Z}(A, A)$.

The second main theorem of this paper is as follows.

Theorem 1.2. Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category and let A be a map \mathcal{O} -monoidale in \mathcal{Z} . The \mathcal{O} -monoidal structure on the coCartesian $(\Delta^{\operatorname{op}}, \mathcal{O})$ -duoidal ∞ -category $\mathcal{Z}(A, A)$ is equivalent to the convolution product on $\mathcal{Z}(A^*, A)$.

The organization of this paper is as follows: The paper is divided into two parts. In Part 1 we introduce a notion of map monoidales and construct a duoidal structure on the endomorphism mapping ∞ -category of a map monoidale. In §2 we review duoidal categories in the setting of higher categories. We recall the definition of (virtual) coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -categories, where \mathcal{O}^{\otimes} and \mathcal{P}^{\otimes} are ∞ -operads over perfect operator categories. In §3 we review scaled simplicial sets which are models of $(\infty, 2)$ -categories. We recall the model structure on the category of scaled simplicial sets whose underlying ∞ -category is equivalent to the theory of $(\infty, 2)$ -categories. In §4 we study adjoint morphisms in $(\infty, 2)$ -categories. We construct 1-full sub $(\infty, 2)$ -categories spanned by left and right adjoint 1-morphisms, respectively. We show that there is a natural dual equivalence between the underlying $(\infty, 1)$ -categories of them. In §5 we introduce a map \mathcal{O} -monoidale in an \mathcal{O} -monoidal ($\infty, 2$)-category, where \mathcal{O}^{\otimes} is an ∞ -operad over a perfect operator category. We define a map \mathcal{O} -monoidale to be an \mathcal{O} -monoidale in \mathcal{Z}^L . We show that there is a dual equivalence between the $(\infty, 1)$ -category of \mathcal{O} -monoidales in \mathcal{Z}^L and the $(\infty, 1)$ -category of \mathcal{O} -comonoidales in \mathcal{Z}^R by taking adjoint morphisms. In §6 we prove Theorem 1.1. First, we show that the endomorphism mapping ∞ -category $\mathcal{Z}(A, A)$ of an \mathcal{O} -monoidale A is a virtual coCartesian $(\Delta^{\mathrm{op}}, \mathcal{O})$ -duoidal ∞ -category. After that, we show that it is in fact a coCartesian $(\Delta^{\mathrm{op}}, \mathcal{O})$ -duoidal ∞ -category when A is a map \mathcal{O} -monoidale.

In Part 2 we compare the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)$ with convolution product. In §7 we introduce a convolution product on the mapping ∞ -category. For an \mathcal{O} -monoidale A and an \mathcal{O} -comonoidale C, we show that the mapping ∞ -category $\mathcal{Z}(C, A)$ has the structure of an \mathcal{O} monoidal $(\infty, 1)$ -category by convolution product. In order to compare the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)$ with convolution product when A is a map \mathcal{O} -monoidale, we need to establish an equivalence between a wide subcategory $\operatorname{Tw}_{sR}^l(\mathcal{X})$ of the twisted arrow ∞ -category and a wide subcategory $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X})$ of the oplax arrow ∞ -category of an $(\infty, 2)$ -category \mathcal{X} . For this purpose, in §8 we introduce a twisted square ∞ -category $\operatorname{TS}(\mathcal{X})$, which is a generalization of twisted arrow ∞ category. For a fibrant scaled simplicial set X which represents \mathcal{X} , we construct a simplicial space $\operatorname{TS}(X)_{\bullet}$ which represents $\operatorname{TS}(\mathcal{X})$. We defer to §11 a proof of $\operatorname{TS}(X)_{\bullet}$ being a complete Segal space. In §9 we show that there is an equivalence of $(\infty, 1)$ -categories between $\operatorname{Tw}_{sR}^l(\mathcal{X})$ and $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X})$. For this purpose, we construct a perfect pairing between them by taking a suitable subcategory of $\operatorname{TS}(\mathcal{X})$. Finally, in §10 we show that the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)$ is equivalent to the convolution product on $\mathcal{Z}(A^*, A)$, where A^* is the dual \mathcal{O} -comonoidale associated to the map \mathcal{O} -monoidale A.

Notation 1.3. We denote by $\operatorname{Set}_{\Delta}$ the category of simplicial sets. We write $\operatorname{Set}_{\Delta}^{\operatorname{Kan}}$ and $\operatorname{Set}_{\Delta}^{\operatorname{Joy}}$ for the category $\operatorname{Set}_{\Delta}$ equipped with the Kan and Joyal model structures, respectively. In this paper we mean that an ∞ -groupoid is a Kan complex, that an ∞ -category is a quasi-category, and that an ∞ -bicategory is a fibrant scaled simplicial set. We let rev : $\Delta^{\operatorname{op}} \to \Delta^{\operatorname{op}}$ be a functor given by assigning to a nonempty finite ordered set its reverse ordered set. For a simplicial object S_{\bullet} , we denote by $S_{\bullet}^{\operatorname{rev}}$ the composite functor $S_{\bullet} \circ \operatorname{rev}$.

We denote by S the ∞ -category of (small) spaces, by $\operatorname{Cat}_{(\infty,1)}$ the ∞ -category of (small) $(\infty, 1)$ categories, and by $\operatorname{Cat}_{(\infty,2)}$ the ∞ -category of (small) $(\infty, 2)$ -categories. We denote by $(-)^{\simeq}$: $\operatorname{Cat}_{(\infty,1)} \to S$ the right adjoint to the inclusion functor $S \to \operatorname{Cat}_{(\infty,1)}$, and by $u_1(-): \operatorname{Cat}_{(\infty,2)} \to$

 $\operatorname{Cat}_{(\infty,1)}$ the right adjoint to the inclusion functor $\operatorname{Cat}_{(\infty,1)} \to \operatorname{Cat}_{(\infty,2)}$. For an ∞ -category \mathcal{C} which admits finite products, we denote by \mathcal{C}^{\times} the associated Cartesian symmetric monoidal ∞ -category.

For an $(\infty, 1)$ -category \mathcal{C} with objects c and d, we denote by $\operatorname{Map}_{\mathcal{C}}(c, d)$ the mapping space from c to d. For an $(\infty, 2)$ -category \mathcal{X} with objects x and y, we denote by $\mathcal{X}(x, y)$ the mapping $(\infty, 1)$ -category from x to y.

For a nonempty finite totally ordered set S, we denote by Δ^S the (|S|-1)-dimensional simplex with S as the set of vertices. For $s \in S$, $\Delta^{S-\{s\}}$ is the codimension 1 face of Δ^S opposite to the vertex s. For a subset $\emptyset \neq N \subsetneq S$, we set

$$\Lambda_N^S = \bigcup_{s \in S-N} \Delta^{S-\{s\}}.$$

For $[r] = \{0 < 1 < \cdots < r\}$, we write Δ^r and Λ^r_i for $\Delta^{[r]}$ and $\Lambda^{[r]}_{\{i\}}$, respectively.

Part 1. Duoidal endomorphism mapping ∞ -categories of map monoidales

In ordinary category theory a map monoidale is a pseudomonoid in a monoidal bicategory whose unit and multiplication morphisms are left adjoint. The endomorphism category of a map monoidale admits the structure of a duoidal category. In Part 1 we consider counterparts of this construction in higher category theory.

In §5 we introduce a notion of a map \mathcal{O} -monoidale in an \mathcal{O} -monoidal (∞ , 2)-category \mathcal{Z} , where \mathcal{O} is an ∞ -operad over a perfect operator category. In §6 we show that the endomorphism mapping ∞ -category of a map \mathcal{O} -monoidale in \mathcal{Z} admits the structure of a coCartesian ($\Delta^{\text{op}}, \mathcal{O}$)-duoidal ∞ -category.

For this purpose, in §2 we recall the notion of duoidal ∞ -categories introduced in [22, 23]. In §3 we recall scaled simplicial sets which are models of $(\infty, 2)$ -categories. In §4 we study adjoint morphisms in $(\infty, 2)$ -categories.

2. Duoidal ∞ -categories

In ordinary category theory a duoidal category has two monoidal products in which one is (op)lax monoidal with respect to the other. In this section we review the notion of duoidal category in the setting of higher category theory (cf. [22, 23]).

Let Φ be a prefect operator category in the sense of [3, Definitions 1.2 and 4.6]. Associated to Φ , we have the Leinster category $\Lambda(\Phi)$ equipped with collections of inert morphisms and active morphisms. An ∞ -operad over Φ is a functor $p: \mathcal{O}^{\otimes} \to \Lambda(\Phi)$ of ∞ -categories satisfying certain conditions (see [3, Definition 7.8] for precise definition). By definition, a perfect operator category Φ has a final object *. We denote by \mathcal{O} the fiber $\mathcal{O}^{\otimes}_{*}$ at $* \in \Lambda(\Phi)$ and say that it is the underlying ∞ -category of \mathcal{O}^{\otimes} . A morphism of \mathcal{O}^{\otimes} is said to be inert if it is a *p*-coCartesian morphism over an inert morphism, and active if it covers an active morphism of $\Lambda(\Phi)$. A morphism of ∞ -operads over Φ between $\mathcal{O}^{\otimes} \to \Lambda(\Phi)$ and $\mathcal{P}^{\otimes} \to \Lambda(\Phi)$ is a functor $f: \mathcal{O}^{\otimes} \to \mathcal{P}^{\otimes}$ over $\Lambda(\Phi)$ that preserves inert morphisms. We denote by $O_{\infty/\Lambda(\Phi)}$ the ∞ -category of ∞ -operads over Φ . We notice that $O_{\mathbb{P}_{\infty/\Lambda(\Phi)}}$ admits finite products.

Example 2.1. Let **F** be the category of finite sets. By [3, Example 4.9.3], **F** is a perfect operator category. By [3, Example 6.5], $\Lambda(\mathbf{F})$ is equivalent to the category Fin_{*} of pointed finite sets. By [3, Example 7.9], the notion of ∞ -operads over **F** coincides with that of Lurie's ∞ -operads in [18, Chapter 2].

Example 2.2. Let **O** be the category of ordered finite sets. By [3, Example 4.9.2], **O** is a perfect operator category. By [3, Example 6.6], $\Lambda(\mathbf{O})$ is equivalent to Δ^{op} . The notion of ∞ -operads over **O** coincides with that of non-symmetric ∞ -operads in [11, §3].

Let $C^{\otimes} \to \mathcal{O}^{\otimes}$ be a map of ∞ -operads over Φ . We say that it is an \mathcal{O} -monoidal ∞ -category if it is a coCartesian fibration. In this case we have a multiplication map

$$\otimes_{\phi}: \prod_{i \in |I|} C_{a_i}^{\otimes} \xleftarrow{\simeq} C_a^{\otimes} \xrightarrow{\phi_*} C_b^{\otimes}$$

for an active morphism $\phi : a \to b$ in \mathcal{O}^{\otimes} , where p(a) = I, $a \simeq (a_i)_{i \in |I|}$, and $a_i, b \in \mathcal{O}$.

Let C and D be \mathcal{O} -monoidal ∞ -categories, and let $g: C^{\otimes} \to D^{\otimes}$ be a map of ∞ -categories over \mathcal{O}^{\otimes} . We say that g is a lax \mathcal{O} -monoidal functor if it is a map of ∞ -operads over \mathcal{O}^{\otimes} , that is, it preserves inert morphisms. We say that g is a strong \mathcal{O} -monoidal functor if it preserves coCartesian morphisms. We denote by $\mathsf{Mon}_{\mathcal{O}}^{\mathrm{lax}}(\mathsf{Cat}_{(\infty,1)})$ the subcategory of $\mathsf{Cat}_{(\infty,1)/\mathcal{O}^{\otimes}}$ spanned by \mathcal{O} -monoidal ∞ -category and lax \mathcal{O} -monoidal functors. We denote by $\mathsf{Mon}_{\mathcal{O}}(\mathsf{Cat}_{(\infty,1)})$ the wide subcategory of $\mathsf{Mon}_{\mathcal{O}}^{\mathrm{lax}}(\mathsf{Cat}_{(\infty,1)})$ spanned by strong \mathcal{O} -monoidal functors. We notice that $\mathsf{Mon}_{\mathcal{O}}^{\mathrm{lax}}(\mathsf{Cat}_{(\infty,1)})$ and $\mathsf{Mon}_{\mathcal{O}}(\mathsf{Cat}_{(\infty,1)})$ admit finite products.

As in [18, §2.4.2], we can identify an \mathcal{O} -monoidal ∞ -category with a functor from \mathcal{O}^{\otimes} to $\operatorname{Cat}_{(\infty,1)}$. Let Y be an ∞ -category with finite products, and let M be a functor $\mathcal{O}^{\otimes} \to Y$ of ∞ -categories. For any $a \in \mathcal{O}^{\otimes}$ with p(a) = I, we can take a family of p-coCartesian morphisms $\{a \to a_i | i \in |I|\}$ lying over the inert morphisms $\{\rho_i : I \to \{i\} | i \in |I|\}$. The family $\{a \to a_i | i \in |I|\}$ induces a morphism $M(a) \to \prod_{i \in |I|} M(a_i)$ in Y, which we call a Segal map. We say that M is an \mathcal{O} -monoid object in Y if the Segal map is an equivalence for each $a \in \mathcal{O}^{\otimes}$. We denote by $\operatorname{Mon}_{\mathcal{O}}(Y)$ the full subcategory of $\operatorname{Fun}(\mathcal{O}^{\otimes}, Y)$ spanned by \mathcal{O} -monoid objects. There is an equivalence of ∞ -categories between $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$ and $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$ (cf. [18, Remark 2.4.2.6]).

Now, we define a notion of virtual coCartesian duoidal ∞ -category. Let $p_{\mathcal{O}} : \mathcal{O}^{\otimes} \to \Lambda(\Phi)$ and $p_{\mathcal{P}} : \mathcal{P}^{\otimes} \to \Lambda(\Psi)$ be ∞ -operads over perfect operator categories Φ and Ψ , respectively. We denote by $\pi_{\mathcal{O}}$ and $\pi_{\mathcal{P}}$ the projections from $\mathcal{O}^{\otimes} \times \mathcal{P}^{\otimes}$ to the factors, respectively.

Definition 2.3 (Definition of virtual coCartesian Duoidal ∞ -categories). Let

$$f: X \to \mathcal{O}^{\otimes} \times \mathcal{P}^{\otimes}.$$

be a categorical fibration of ∞ -categories. We say that f is a virtual coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category if it satisfies the following conditions:

- (1) The composite $\pi_{\mathcal{O}} \circ f : X \to \mathcal{O}^{\otimes}$ is a coCartesian fibration, and f carries $(\pi_{\mathcal{O}} \circ f)$ coCartesian morphisms to $\pi_{\mathcal{O}}$ -coCartesian morphisms.
- (2) For each $a \in \mathcal{O}^{\otimes}$, the restriction $p_a : X_a \to \mathcal{P}^{\otimes}$ is a morphism of ∞ -operads over $\Lambda(\Psi)$.
- (3) For each morphism $a \to a'$ in \mathcal{O}^{\otimes} , the induced map $X_a \to X_{a'}$ over \mathcal{P}^{\otimes} is a morphism of ∞ -operads over \mathcal{P}^{\otimes} .
- (4) For each $a \in \mathcal{O}^{\otimes}$ with $p_{\mathcal{O}}(a) = I$, the Segal morphism

$$X_a \longrightarrow \prod_{i \in |I|} \mathcal{P}^{\otimes} X_{a_i}$$

is an equivalence in the ∞ -category $\operatorname{Op}_{\infty/\mathcal{P}^{\otimes}}$ of ∞ -operads over \mathcal{P}^{\otimes} , where $\{a \to a_i | i \in |I|\}$ is a family of $p_{\mathcal{O}}$ -coCartesian morphisms lying over the inert morphisms $\{\rho_i : I \to \{i\} | i \in |I|\}$, and the right hand side is a product in $\operatorname{Op}_{\infty/\mathcal{P}^{\otimes}}$.

Remark 2.4. A virtual coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category corresponds to an \mathcal{O} -monoid object of the Cartesian symmetric monoidal ∞ -category $\operatorname{Op}_{\infty/\mathcal{P}^{\otimes}}$ under the straightening along the map $\pi_{\mathcal{O}} \circ f : X \to \mathcal{O}^{\otimes}$.

Next, we recall the definition of coCartesian duoidal ∞ -categories.

Definition 2.5 (Definition of coCartesian Duoidal ∞ -categories (cf. [22, Definition 4.23] and [23, Definition 3.14])). Let

$$f: X \to \mathcal{O}^{\otimes} \times \mathcal{P}^{\otimes}.$$

be a categorical fibration of ∞ -categories. We say that f is a coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category if it satisfies the following conditions:

- (1) The map f is a virtual coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category.
- (2) For each $a \in \mathcal{O}^{\otimes}$, the restriction $p_a : X_a \to \mathcal{P}^{\otimes}$ is a \mathcal{P} -monoidal ∞ -category.

Remark 2.6. A coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category corresponds to an \mathcal{O} -monoid object of the Cartesian symmetric monoidal ∞ -category $\mathsf{Mon}_{\mathcal{P}}^{\mathsf{lax}}(\mathsf{Cat}_{(\infty,1)})$ under the straightening along the map $\pi_{\mathcal{O}} \circ f : X \to \mathcal{O}^{\otimes}$.

Informally speaking, a coCartesian $(\mathcal{O}, \mathcal{P})$ -duoidal ∞ -category has \mathcal{O} -monoidal and \mathcal{P} -monoidal products in which the \mathcal{O} -monoidal structure is lax monoidal with respect to the \mathcal{P} -monoidal structure.

3. Scaled simplicial sets and $(\infty, 2)$ -categories

In this section we review scaled simplicial sets which are models of $(\infty, 2)$ -categories. We recall the model structure on the category of scaled simplicial sets introduced in [19] whose underlying ∞ -category is equivalent to the ∞ -category $\operatorname{Cat}_{(\infty,2)}$ of $(\infty, 2)$ -categories.

First, we recall the definition of scaled simplicial sets.

Definition 3.1 ([19, Definition 3.1.1])). A scaled simplicial set is a pair (\overline{X}, T_X) in which \overline{X} is a simplicial set and T_X is a set of 2-simplices of \overline{X} that contains all the degenerate 2-simplices. We say that a 2-simplex is thin if it is contained in T_X . For scaled simplicial sets $X = (\overline{X}, T_X)$ and $Y = (\overline{Y}, T_Y)$, a morphism $f : X \to Y$ of scaled simplicial sets is a map $\overline{f} : \overline{X} \to \overline{Y}$ of simplicial sets that carries T_X into T_Y . We denote by $\operatorname{Set}_{\Delta}^{\mathrm{sc}}$ the category of scaled simplicial sets.

For a simplicial set S, we have two canonical scaled simplicial sets S_{\sharp} and S_{\flat} . The scaled simplicial set S_{\sharp} has the underlying simplicial set S equipped with all 2-simplices as thin triangles. On the other hand, S_{\flat} has the underlying simplicial set S equipped with all degenerate 2-simplices as thin triangles.

Now, we recall scaled anodyne maps of scaled simplicial sets which characterize fibrant objects in $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$.

Definition 3.2 ([19, Definition 3.1.3]). The collection of scaled anodyne maps is the weakly saturated class of morphisms of $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ generated by the following maps:

(An1) For each 0 < i < n, the inclusion map

$$(\Lambda_i^n, \{\Delta^{\{i-1,i,i+1\}}\}|_{\Lambda_i^n} \cup \{\text{degenerate}\}) \longrightarrow (\Delta^n, \{\Delta^{\{i-1,i,i+1\}}\} \cup \{\text{degenerate}\}).$$

(An2) The inclusion map

$$(\Delta^4, T) \longrightarrow (\Delta^4, T \cup \{\Delta^{\{0,3,4\}}, \Delta^{0,1,4}\}),$$

where

$$T = \{\Delta^{\{0,2,4\}}, \Delta^{\{1,2,3\}}, \Delta^{\{0,1,3\}}, \Delta^{\{1,3,4\}}, \Delta^{\{0,1,2\}}\} \cup \{\text{degenerate}\}$$

(An3) For n > 2, the inclusion map

$$(\Lambda_0^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\} \cup \{\text{degenerate}\}) \longrightarrow (\Delta^n \coprod_{\Delta^{\{0,1\}}} \Delta^0, \{\Delta^{\{0,1,n\}}\} \cup \{\text{degenerate}\}).$$

A model structure on $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ was introduced in [19] and showed that the underlying ∞ -category of $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ is equivalent to the ∞ -category $\operatorname{Cat}_{(\infty,2)}$ of $(\infty,2)$ -categories.

Theorem 3.3 (cf. [19, Theorem 4.2.7], [14, Theorem 1.29], and [13, Theorem 1.12]). There exists a model structure on $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ whose cofibrations are the monomorphisms and whose fibrant objects are those objects that have the right lifting properties with respect to the scaled anodyne maps.

Definition 3.4. We call a fibrant scaled simplicial set an ∞ -bicategory.

Remark 3.5. In [19, Definition 4.1.1] scaled simplicial sets satisfying the right lifting property with respect to the scaled anodyne maps are referred to as weak ∞ -bicategories, and fibrant objects in Set^{sc}_{Δ} are referred to as ∞ -bicategories. However, it is shown that these two notations coincide in [14, Theorem 5.1].

We recall that $\operatorname{Set}_{\Delta}^{\operatorname{Joy}}$ is the category of simplicial sets equipped with the Joyal model structure. By assigning the scaled simplicial set S_{\sharp} to a simplicial set S, we obtain a left Quillen functor $(-)_{\sharp} : \operatorname{Set}_{\Delta}^{\operatorname{Joy}} \to \operatorname{Set}_{\Delta}^{\operatorname{sc}}$. We denote by

$$u_1 : \operatorname{Set}_{\Delta}^{\operatorname{sc}} \longrightarrow \operatorname{Set}_{\Delta}^{\operatorname{Joy}}$$

the right adjoint to $(-)_{\sharp}$, which assigns to a scaled simplicial set $X = (\overline{X}, T_X)$ the subcomplex of \overline{X} spanned by those simplices whose 2-dimensional faces are all thin. Since u_1 is a right Quillen functor, u_1X is an ∞ -category for an ∞ -bicategory X. The functor u_1 induces the functor $u_1 : \operatorname{Cat}_{(\infty,2)} \to \operatorname{Cat}_{(\infty,1)}$ of the underlying ∞ -categories, which is a right adjoint to the inclusion functor $\operatorname{Cat}_{(\infty,1)} \to \operatorname{Cat}_{(\infty,2)}$.

The model structure on $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ is cartesian closed by [19, Proposition 3.1.8 and Lemma 4.2.6]. (see, also, [14, Remark 1.31] or [13, the paragraph before Lemma 1.24]). Thus, we have a function object FUN(X, Y) in $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$ for scaled simplicial sets X and Y. When Y is an ∞ -bicategory, FUN(X, Y) is also an ∞ -bicategory. We denote by

$$\operatorname{Fun}(X,Y)$$

its underlying ∞ -category u_1 FUN(X, Y), and by

$$\operatorname{Map}^{\operatorname{sc}}(X,Y)$$

its underlying ∞ -groupoid $u_1 \text{FUN}(X, Y)^{\simeq}$.

The Gray tensor product of scaled simplicial sets was introduced in [13]. This construction determines a functor

$$(-)\otimes(-):\operatorname{Set}_{\Delta}^{\operatorname{sc}}\times\operatorname{Set}_{\Delta}^{\operatorname{sc}}\longrightarrow\operatorname{Set}_{\Delta}^{\operatorname{sc}},$$

which is a left Quillen bifunctor by [13, Theorem 2.14]. Hence, it induces a functor $(-) \otimes (-)$: Cat $_{(\infty,2)} \times$ Cat $_{(\infty,2)} \to$ Cat $_{(\infty,2)}$ of ∞ -categories, which preserves colimits separately in each variable. For example, the Gray tensor product $\Delta^{1}_{\sharp} \otimes \Delta^{1}_{\sharp}$ is a scaled simplicial set whose underlying simplicial set is $\Delta^{1} \times \Delta^{1}$ equipped with the 2-simplex $\Delta^{\{(0,0),(1,0),(1,1)\}}$ together with degenerate 2-simplices as thin. It represents an oplax square depicted by



The Gray tensor product on scaled simplicial sets has adjoints $FUN^{lax}(-,-)$ and $FUN^{opl}(-,-)$ which satisfy

$$\operatorname{Hom}_{\operatorname{Set}_{\Delta}^{\operatorname{sc}}}(Z, \operatorname{FUN}^{\operatorname{iax}}(X, Y)) \cong \operatorname{Hom}_{\operatorname{Set}_{\Delta}^{\operatorname{sc}}}(Z \otimes X, Y),$$

$$\operatorname{Hom}_{\operatorname{Set}_{\Delta}^{\operatorname{sc}}}(Z, \operatorname{FUN}^{\operatorname{opl}}(X, Y)) \cong \operatorname{Hom}_{\operatorname{Set}_{\Delta}^{\operatorname{sc}}}(X \otimes Z, Y).$$

If Y is an ∞ -bicategory, then $\mathrm{FUN}^{\mathrm{lax}}(X,Y)$ and $\mathrm{FUN}^{\mathrm{opl}}(X,Y)$ are also ∞ -bicategories. We write

$$\operatorname{Fun}^{\operatorname{lax}}(X,Y), \quad \operatorname{Fun}^{\operatorname{opl}}(X,Y)$$

for the underlying ∞ -categories $u_1 \text{FUN}^{\text{lax}}(X, Y)$ and $u_1 \text{FUN}^{\text{opl}}(X, Y)$, respectively.

4. Adjoint morphisms in $(\infty, 2)$ -categories

In this section we study adjoint morphisms in $(\infty, 2)$ -categories. For an $(\infty, 2)$ -category \mathcal{X} , we construct 1-full sub $(\infty, 2)$ -categories \mathcal{X}^L and \mathcal{X}^R spanned by left and right adjoint 1-morphisms, respectively. We show that there is a natural dual equivalence between the underlying $(\infty, 1)$ -categories of \mathcal{X}^L and \mathcal{X}^R .

First, we observe a property of adjoint 1-morphisms in an $(\infty, 2)$ -category. Let (m^L, m^R, η) be an adjunction in an $(\infty, 2)$ -category \mathcal{X} , where $m^L : x \to x'$ is a left adjoint 1-morphism to $m^R : x' \to x$, and $\eta : \mathrm{id}_x \Rightarrow m^R m^L$ is its unit 2-morphism. For 1-morphisms $f : x \to y$ and $g : x' \to y$ of \mathcal{X} , we consider a map of mapping spaces

$$\operatorname{Map}_{\mathcal{X}(x',y)}(fm^R,g) \longrightarrow \operatorname{Map}_{\mathcal{X}(x,y)}(f,gm^L)$$

given by the composite

 $\operatorname{Map}_{\mathcal{X}(x',y)}(fm^{R},g) \xrightarrow{(-)m^{L}} \operatorname{Map}_{\mathcal{X}(x,y)}(fm^{R}m^{L},gm^{L}) \xrightarrow{(f\eta)^{*}} \operatorname{Map}_{\mathcal{X}(x,y)}(f,gm^{L}).$

The following lemma might be well-known but we give a proof of it for convenience of readers.

Lemma 4.1. The map $\operatorname{Map}_{\mathcal{X}(x',y)}(fm^R,g) \to \operatorname{Map}_{\mathcal{X}(x,y)}(f,gm^L)$ is an equivalence.

 $\mathit{Proof.}$ We shall construct a homotopy inverse of the map. We consider a map given by the composite

$$\operatorname{Map}_{\mathcal{X}(x,y)}(f,gm^{L}) \xrightarrow{(-)m^{R}} \operatorname{Map}_{\mathcal{X}(x',y)}(fm^{R},gm^{L}m^{R}) \xrightarrow{(g\epsilon)_{*}} \operatorname{Map}_{\mathcal{X}(x,y)}(fm^{R},g),$$

where $\epsilon : m^L m^R \Rightarrow id_{x'}$ is a counit of the adjunction. By the triangle identity, we see that it is the desired homotopy inverse.

For an $(\infty, 2)$ -category \mathcal{X} , we construct 1-full sub $(\infty, 2)$ -categories \mathcal{X}^L and \mathcal{X}^R of \mathcal{X} spanned by left and right adjoint 1-morphisms, respectively. We give a construction of ∞ -bicategories X^L and X^R which represent \mathcal{X}^L and \mathcal{X}^R , respectively, when \mathcal{X} is represented by an ∞ -bicategory X.

Let $X = (\overline{X}, T_X)$ be a fibrant scaled simplicial set. We write \overline{X}^L for the subcomplex of \overline{X} spanned by those simplices whose 1-dimensional faces are all left adjoint. We define X^L to be the scaled simplicial set $(\overline{X}^L, T_{X^L})$, where T_{X^L} is the restriction of T_X to \overline{X}^L . We define a scaled simplicial set X^R in a similar manner by using right adjoints instead of left adjoints. We show that X^L and X^R are fibrant scaled simplicial sets and hence they represent $(\infty, 2)$ -categories.

Lemma 4.2. The scaled simplicial sets X^L and X^R are fibrant.

Proof. We shall prove the case for X^L . The case for X^R can be proven in a similar manner. We consider an extension problem of scaled anodyne maps with respect to X^L . Since X is fibrant, we can find an extension in X. If a scaled anodyne map is of type (An1) for $n \ge 3$, (An2), or (An3), then the extension lands in X^L since the source of scaled anodyne maps contains all 1-simplices of

the target. If a scaled anodyne map is of type (An1) for n = 2, then the image of $\Delta^{\{0,2\}}$ in X is equivalent to a composite of left adjoints, and hence the extension also lands in X^L .

Definition 4.3. Let \mathcal{X} be an $(\infty, 2)$ -category represented by a fibrant scaled simplicial set X. We denote by \mathcal{X}^L and \mathcal{X}^R the $(\infty, 2)$ -categories represented by fibrant scaled simplicial sets X^L and X^R , respectively.

We recall that the inclusion functor $\operatorname{Cat}_{(\infty,1)} \to \operatorname{Cat}_{(\infty,2)}$ has the right adjoint $u_1 : \operatorname{Cat}_{(\infty,2)} \to \operatorname{Cat}_{(\infty,1)}$. For an $(\infty,2)$ -category \mathcal{X} , $u_1\mathcal{X}$ is the underlying $(\infty,1)$ -category of \mathcal{X} obtained by discarding all non-invertible 2-morphisms. Next, we show that there is a dual equivalence between $u_1\mathcal{X}^L$ and $u_1\mathcal{X}^R$ by taking adjoint 1-morphisms. For this purpose, we construct a perfect pairing between $u_1\mathcal{X}^L$ and $u_1\mathcal{X}^R$.

First, we construct a pairing of $u_1 \mathcal{X}^L$ and $u_1 \mathcal{X}^R$. For this purpose, we recall the oplax arrow ∞ -category

$$\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$$

which is represented by the ∞ -category Fun^{opl}(Δ^1_{\sharp}, X) (cf. [16, Definition 7.5]). An object of $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$ is a 1-morphism of \mathcal{X} . A morphism of $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$ from $f: x \to y$ to $f': x' \to y'$ is an oplax square depicted by



The inclusion map $\partial \Delta^{1}_{\sharp} \to \Delta^{1}_{\sharp}$ induces a map $\mathrm{FUN}^{\mathrm{opl}}(\Delta^{1}_{\sharp}, X) \to X \times X$, which is a fibration between fibrant objects in $\mathrm{Set}^{\mathrm{sc}}_{\Delta}$. Since $u_{1} : \mathrm{Set}^{\mathrm{sc}}_{\Delta} \to \mathrm{Set}^{\mathrm{Joy}}_{\Delta}$ is a right Quillen functor, we obtain a map $\mathrm{Ar}^{\mathrm{opl}}(X) \to u_{1}X \times u_{1}X$, which is a fibration of fibrant objects in $\mathrm{Set}^{\mathrm{Joy}}_{\Delta}$. We write

 $(s,t):\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})\longrightarrow u_1\mathcal{X}\times u_1\mathcal{X}$

for the corresponding map of $(\infty, 1)$ -categories.

Remark 4.4. The map (s,t): $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X}) \to u_1 \mathcal{X} \times u_1 \mathcal{X}$ is an orthofibration by [16, Proposition 7.9].

Now, we give a sufficient condition for 1-morphisms in $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$ to be (s,t)-coCartesian.

Lemma 4.5. Let σ be a morphism in $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$ depicted by



where g^L is a left adjoint 1-morphism. Then σ is an (s,t)-coCartesian morphism if f' is equivalent to hfg^R , and the 2-morphism u is equivalent to the composite $hf \stackrel{hf\eta}{\Longrightarrow} hfg^Rg^L \simeq f'g^L$, where (g^L, g^R, η) is an adjunction in \mathcal{X} that is an extension of g^L .

Proof. Let $f'': x'' \to y''$ be an object of $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})$. We would like to show that the following commutative diagram

$$\begin{array}{cccc} \operatorname{Map}_{\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})}(f',f'') & \longrightarrow & \operatorname{Map}_{\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})}(f,f'') \\ & & & \downarrow \\ & & & \downarrow \\ \operatorname{Map}_{u_{1}\mathcal{X}\times u_{1}\mathcal{X}}((x',y'),(x'',y'')) & \longrightarrow & \operatorname{Map}_{u_{1}\mathcal{X}\times u_{1}\mathcal{X}}((x,y),(x'',y'')) \end{array}$$

is a pullback in the $(\infty, 1)$ -category of spaces. We take a 1-morphism $(k, l) : (x', y') \to (x'', y'')$ in $u_1 \mathcal{X} \times u_1 \mathcal{X}$. The induced map on fibers

$$\operatorname{Map}_{\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})}(f', f'')_{(k,l)} \to \operatorname{Map}_{\operatorname{Ar}^{\operatorname{opl}}(\mathcal{X})}(f, f'')_{(kg^L, lh)}$$

is identified with the map

$$\operatorname{Map}_{\mathcal{X}(x',y'')}(lf',f''k) \longrightarrow \operatorname{Map}_{\mathcal{X}(x,y'')}(lhf,f''kg^L)$$

given by the composition with σ , which is an equivalence by Lemma 4.1.

We define maps

$$\begin{array}{rccc} (s,t)_{sL} : & \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}) & \longrightarrow & u_1 \mathcal{X}^L \times u_1 \mathcal{X}, \\ (s,t)_{L,R} : & \operatorname{Ar}_{L,R}^{\operatorname{opl}}(\mathcal{X}) & \longrightarrow & u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R \end{array}$$

by the following pullback diagrams

$$\begin{array}{cccc} \operatorname{Ar}_{L,R}^{\operatorname{opl}}(\mathcal{X}) & \longrightarrow & \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}) & \longrightarrow & \operatorname{Ar}^{\operatorname{opl}}(\mathcal{X}) \\ (s,t)_{L,R} & & & & & & \\ (s,t)_{sL} & & & & & & \\ u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R & \longrightarrow & u_1 \mathcal{X}^L \times u_1 \mathcal{X} & \longrightarrow & u_1 \mathcal{X} \times u_1 \mathcal{X} \end{array}$$

By the criterion of (s, t)-coCartesian morphisms in Lemma 4.5, we obtain the following corollary.

Corollary 4.6. The maps $(s,t)_{sL}$: $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}) \to u_1 \mathcal{X}^L \times u_1 \mathcal{X}$ and $(s,t)_{L,R}$: $\operatorname{Ar}_{L,R}^{\operatorname{opl}}(\mathcal{X}) \to u_1 \mathcal{X}^L \times u_1 \mathcal{X}$ are coCartesian fibrations.

Next, we construct the desired pairing from the map $(s,t)_{L,R}$. We define $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X})$ to be the full subcategory of $\operatorname{Ar}_{L,R}^{\operatorname{opl}}(\mathcal{X})$ spanned by those objects that are right adjoint 1-morphisms in \mathcal{X} . By restricting the map $(s,t)_{L,R}$ to $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X})$, we obtain a map

$$\overline{(s,t)}_{L,R}: \overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X}) \longrightarrow u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R.$$

Lemma 4.7. The map $\overline{(s,t)}_{L,R}$: $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X}) \to u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R$ is a coCartesian fibration.

Proof. Let $f: x \to y$ be an object of $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X})$ and let $(g^L, h^R): (x, y) \to (x', y')$ be a morphism of $u_1\mathcal{X}^L \times u_1\mathcal{X}^R$. We take a $(s, t)_{L,R}$ -coCartesian morphism $f \to f'$ in $\operatorname{Ar}_{L,R}^{\operatorname{opl}}(\mathcal{X})$ covering (g^L, h^R) . It suffices to show that f' belongs to $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X})$. This follows from the fact that $h^R f g^R$ is a right adjoint 1-morphism.

We define $\operatorname{Pair}(\mathcal{X})$ to be the wide subcategory of $\overline{\operatorname{Ar}}_{L,R}^{\operatorname{opl}}(\mathcal{X})$ spanned by $\overline{(s,t)}_{L,R}$ -coCartesian morphisms. We denote by

$$(s,t)^{\operatorname{pair}} : \operatorname{Pair}(\mathcal{X}) \longrightarrow u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R$$

the restriction of $\overline{(s,t)}_{L,R}$ to Pair(\mathcal{X}). By Lemma 4.7, the map $(s,t)^{\text{pair}}$ is a left fibration, and hence it defines a pairing of $(\infty, 1)$ -categories between $u_1 \mathcal{X}^L$ and $u_1 \mathcal{X}^R$.

Theorem 4.8. The pairing $(s,t)^{\text{pair}}$: $\operatorname{Pair}(\mathcal{X}) \to u_1 \mathcal{X}^L \times u_1 \mathcal{X}^R$ is perfect.

Proof. By [18, Corollary 5.2.1.22], it suffices to show that the pairing $(s, t)^{\text{pair}}$ is both left and right representable and that an object of $\text{Pair}(\mathcal{X})$ is left universal if and only if it is right universal.

First, we show that $(s,t)^{\text{pair}}$ is left representable. Let x be an object of $u_1 \mathcal{X}^L$. We set $S(x) = \{x\} \times_{u_1 \mathcal{X}^L} \text{Pair}(\mathcal{X})$ for simplicity. We take the identity morphism $\text{id}_x : x \to x$, which we regard as

an object of S(x). We have to show that id_x is an initial object. For any object $f : x \to y$ in S(x), we can identify $Map_{S(x)}(id_x, f)$ with $Map_{u_1\chi^R}(x, y)_{/f}$, which is contractible.

Next, we shall show that $(s, t)^{\text{pair}}$ is right representable. Let y be an object of $u_1 \mathcal{X}^R$. We set $T(y) = \text{Pair}(\mathcal{X}) \times_{u_1 \mathcal{X}^R} \{y\}$ for simplicity. We take the identity morphism $\text{id}_y : y \to y$, which we regard as an object of T(y). We have to show that id_y is an initial object. For any object $f : x \to y$ in T(y), we can identify $\text{Map}_{T(y)}(\text{id}_y, f)$ with the space of units (g, f, u) that are extensions of f, which is contractible by [20, Theorem 4.4.18].

Finally, we shall show that an object of $\operatorname{Pair}(\mathcal{X})$ is left universal if and only if it is right universal. By the above argument, we see that an object $f : x \to y$ of $\operatorname{Pair}(\mathcal{X})$ is left universal if and only if f is equivalent to id_x in S(x). This means that f is an equivalence in \mathcal{X} . In a similar manner we see that f is right universal if and only if f is an equivalence in \mathcal{X} . This completes the proof. \Box

Corollary 4.9. There is a dual equivalence of $(\infty, 1)$ -categories between $u_1 \mathcal{X}^L$ and $u_1 \mathcal{X}^R$.

Remark 4.10. The dual equivalence in Corollary 4.9 is the identity on objects and assigns to a left adjoint 1-morphism its right adjoint 1-morphism.

Finally, we show that the dual equivalence in Corollary 4.9 is natural with respect to \mathcal{X} . We let $u_1(-)^L$ and $u_1(-)^R$ be functors from $\operatorname{Cat}_{(\infty,2)}$ to $\operatorname{Cat}_{(\infty,1)}$ which assigns $u_1\mathcal{Y}^L$ and $u_1\mathcal{Y}^R$ to an $(\infty,2)$ -category \mathcal{Y} , respectively.

Corollary 4.11. There is a natural dual equivalence of functors between $u_1(-)^L$ and $u_1(-)^R$.

Proof. Let $f: \mathcal{Y} \to \mathcal{Z}$ be a functor of $(\infty, 2)$ -categories. This induces a morphism of pairings

(4.1)
$$\begin{array}{ccc} \operatorname{Pair}(\mathcal{Y}) & \xrightarrow{\operatorname{Pair}(f)} & \operatorname{Pair}(\mathcal{Z}) \\ & \downarrow & & \downarrow \\ & u_1 \mathcal{Y}^L \times u_1 \mathcal{Y}^R & \xrightarrow{u_1 f^L \times u_1 f^R} & u_1 \mathcal{Z}^L \times u_1 \mathcal{Z}^R \end{array}$$

By the characterization of right universal objects in $\operatorname{Pair}(\mathcal{X})$ in the proof of Theorem 4.8, we see that (4.1) is right representable in the sense of [18, Variant 5.2.1.16]. By this observation, we obtain a functor

$$\operatorname{Cat}_{(\infty,2)} \longrightarrow \operatorname{CPair}^{\operatorname{perf}}$$

of ∞ -categories by assigning to an $(\infty, 2)$ -category \mathcal{Y} the perfect pairing $\operatorname{Pair}(\mathcal{Y}) \to u_1 \mathcal{Y}^L \times u_1 \mathcal{Y}^R$, where CPair^{perf} is the ∞ -category of perfect pairings and right representable morphisms (see [18, Remark 5.2.1.20] for the definition of CPair^{perf}). This implies that the dual equivalence in Corollary 4.9 is natural with respect to \mathcal{X} .

Remark 4.12. We can prove an existence of a natural dual equivalence between $u_1 \mathcal{X}^L$ and $u_1 \mathcal{X}^R$ by using [15, Remark 4.11] as well. Let \mathfrak{adj} be the free 2-category containing an adjunction, and let \mathfrak{adj}^{\bullet} be the cosimplicial $(\infty, 2)$ -category whose *n*th object is the $(\infty, 2)$ -category of *n* composable adjunctions $\mathfrak{adj} \coprod_{[0]} \cdots \coprod_{[0]} \mathfrak{adj}$. By [15, Theorem 4.4 and Remark 4.11], there are equivalences of simplicial spaces

$$\operatorname{Map}_{\operatorname{Cat}_{(\infty,2)}}([\bullet]^{\operatorname{op}},\mathcal{X})^{\operatorname{ladj}} \xleftarrow{\simeq} \operatorname{Map}_{\operatorname{Cat}_{(\infty,2)}}(\mathfrak{adj}^{\bullet},\mathcal{X}) \xrightarrow{\simeq} \operatorname{Map}_{\operatorname{Cat}_{(\infty,2)}}([\bullet],\mathcal{X})^{\operatorname{radj}}$$

This implies that there is a natural dual equivalence of complete Segal spaces associated to $u_1 \mathcal{X}^L$ and $u_1 \mathcal{X}^R$.

5. Map monoidales in monoidal ∞ -bicategories

In this section we generalize map monoidales in monoidal bicategories in the setting of higher category theory. We introduce notions of left adjoint \mathcal{O} -monoidales and right adjoint \mathcal{O} -comonoidales in an \mathcal{O} -monoidal (∞ , 2)-category. We show that there is a dual equivalence between the (∞ , 1)category of left adjoint \mathcal{O} -monoidales and the (∞ , 1)-category of right adjoint \mathcal{O} -comonoidales by taking adjoint morphisms.

Since $\operatorname{Cat}_{(\infty,2)}$ has finite products, we can consider \mathcal{O} -monoid objects in $\operatorname{Cat}_{(\infty,2)}$. We call an \mathcal{O} -monoid object in $\operatorname{Cat}_{(\infty,2)}$ an \mathcal{O} -monoidal $(\infty, 2)$ -category. In the following of this section we let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category.

Since the functor $u_1 : \operatorname{Cat}_{(\infty,2)} \to \operatorname{Cat}_{(\infty,1)}$ is right adjoint to the inclusion functor, it preserves small limits. In particular, it preserves finite products and hence it induces a functor $u_1 : \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)}) \to \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$. From this observation, we see that $u_1\mathcal{Z}$ is an \mathcal{O} monoidal $(\infty, 1)$ -category.

Let \mathcal{W} be an \mathcal{O} -monoidal $(\infty, 1)$ -category. We identify \mathcal{W} with a coCartesian fibration $\mathcal{W}^{\otimes} \to \mathcal{O}^{\otimes}$ of ∞ -operads. An \mathcal{O} -algebra object A in \mathcal{W} is a map of ∞ -operads $\mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$ over \mathcal{O}^{\otimes} . We define $\operatorname{Alg}_{\mathcal{O}}(\mathcal{W})$ to be the full subcategory of $\operatorname{Fun}_{\mathcal{O}^{\otimes}}(\mathcal{O}^{\otimes}, \mathcal{W}^{\otimes})$ spanned by \mathcal{O} -algebra objects, and call it the ∞ -category of \mathcal{O} -algebra objects in \mathcal{W} .

The opposite $(\infty, 1)$ -category \mathcal{W}^{op} inherits an \mathcal{O} -monoidal structure from \mathcal{W} . An \mathcal{O} -coalgebra object in \mathcal{W} is an \mathcal{O} -algebra object in \mathcal{W}^{op} . We define $\operatorname{coAlg}_{\mathcal{O}}(\mathcal{W})$ to be $\operatorname{Alg}_{\mathcal{O}}(\mathcal{W}^{\text{op}})^{\text{op}}$, and call it the ∞ -category of \mathcal{O} -coalgebra objects in \mathcal{W} .

In ordinary category theory a monoidale is a pseudomonoid in a monoidal bicategory. We define a monoidale in the setting of higher category theory.

Definition 5.1. Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category. An \mathcal{O} -monoidale in \mathcal{Z} is an \mathcal{O} -algebra object in $u_1\mathcal{Z}$. An \mathcal{O} -comonoidale in \mathcal{Z} is an \mathcal{O} -coalgebra object in $u_1\mathcal{Z}$.

Next, we would like to introduce a notion of a map monoidale in the setting of higher category theory. For this purpose, we show that the $(\infty, 2)$ -categories \mathcal{Z}^L and \mathcal{Z}^R inherit \mathcal{O} -monoidal structure from \mathcal{Z} .

Lemma 5.2. If \mathcal{Z} is an \mathcal{O} -monoidal $(\infty, 2)$ -category, then \mathcal{Z}^L and \mathcal{Z}^R are also \mathcal{O} -monoidal $(\infty, 2)$ -categories.

Proof. We shall show that \mathcal{Z}^L is an \mathcal{O} -monoidal $(\infty, 2)$ -category. The case of \mathcal{Z}^R is similar. The construction $\mathcal{X} \mapsto \mathcal{X}^L$ determines a functor $(-)^L : \operatorname{Cat}_{(\infty,2)} \to \operatorname{Cat}_{(\infty,2)}$. This functor preserves finite products. Thus, it induces a functor $(-)^L : \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)}) \to \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)})$. \Box

By Lemma 5.2, we have \mathcal{O} -monoidal $(\infty, 1)$ -categories $u_1 \mathcal{Z}^L$ and $u_1 \mathcal{Z}^R$. In ordinary category theory a monoidale is said to be a map monoidale if the product and unit morphisms are left adjoint 1-morphisms. We define a notion of a map monoidale in an \mathcal{O} -monoidal $(\infty, 2)$ -category.

Definition 5.3. Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category. We define a map \mathcal{O} -monoidale in \mathcal{Z} to be an \mathcal{O} -monoidale in \mathcal{Z}^L . We also say that a map \mathcal{O} -monoidale is a left adjoint \mathcal{O} -monoidale. We define a right adjoint \mathcal{O} -comonoidale in \mathcal{Z} to be an \mathcal{O} -comonoidale in \mathcal{Z}^R .

Next, we study a duality of left adjoint \mathcal{O} -monoidales and right adjoint \mathcal{O} -comonoidales under the dual equivalence in Corollary 4.9. For this purpose, we promote the dual equivalence in Corollary 4.9 to a dual equivalence of monoidal (∞ , 1)-categories.

Lemma 5.4. When \mathcal{Z} is an \mathcal{O} -monoidal $(\infty, 2)$ -category, there is a natural dual equivalence of \mathcal{O} -monoidal $(\infty, 1)$ -categories between $u_1 \mathcal{Z}^L$ and $u_1 \mathcal{Z}^R$.

Proof. By the proof of Corollary 4.11, there is a functor $\operatorname{Cat}_{(\infty,2)} \to \operatorname{CPair}^{\operatorname{pair}}$ which assigns to an $(\infty, 2)$ -category \mathcal{X} the perfect pairing $\operatorname{Pair}(\mathcal{X}) \to u_1 \mathcal{X}^L \times \mathcal{X}^R$. We can verify that this functor preserves finite products. Hence we obtain a functor $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)}) \to \operatorname{Mon}_{\mathcal{O}}(\operatorname{Pair}^{\operatorname{perf}})$. This implies that the dual equivalence in Corollary 4.9 can be promoted to a natural dual equivalence of \mathcal{O} -monoidal $(\infty, 1)$ -categories. \Box

By Lemma 5.4, we obtain a duality between left adjoint \mathcal{O} -monoidales and right adjoint \mathcal{O} comonoidales.

Corollary 5.5. There is a natural dual equivalence between the ∞ -category of left adjoint \mathcal{O} -monoidales and the ∞ -category of right adjoint \mathcal{O} -comonoidales

$$\operatorname{Alg}_{\mathcal{O}}(u_1 \mathcal{Z}^L) \simeq \operatorname{coAlg}_{\mathcal{O}}(u_1 \mathcal{Z}^R)^{\operatorname{op}}$$

Definition 5.6. For a left adjoint \mathcal{O} -monoidale $A \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathbb{Z}^L)$, we let A^* be the corresponding object in $\operatorname{coAlg}_{\mathcal{O}}(u_1 \mathbb{Z}^R)$ under the equivalence in Corollary 5.5. We call A^* the right adjoint \mathcal{O} -comonoidale associated to A. Similarly, for a right adjoint \mathcal{O} -comonoidale $C \in \operatorname{coAlg}_{\mathcal{O}}(u_1 \mathbb{Z}^R)$, we denote by C^* the corresponding object in $\operatorname{Alg}_{\mathcal{O}}(u_1 \mathbb{Z}^L)$ under the equivalence in Corollary 5.5. We call C^* the left adjoint \mathcal{O} -monoidale associated to C.

6. Map monoidales and Duoidal ∞ -categories

Let A be a map \mathcal{O} -monoidale in an \mathcal{O} -monoidal $(\infty, 2)$ -category \mathcal{Z} . The goal of this section is to show that the endomorphism mapping ∞ -category $\mathcal{Z}(A, A)$ admits the structure of a coCartesian $(\Delta^{\mathrm{op}}, \mathcal{O})$ -duoidal ∞ -category.

First, we construct a simplicial \mathcal{O} -monoidal $(\infty, 1)$ -category $\operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes}$. For an $(\infty, 2)$ -category \mathcal{K} , the functor $\operatorname{Fun}^{\operatorname{opl}}(\mathcal{K}, -) : \operatorname{Cat}_{(\infty, 2)} \to \operatorname{Cat}_{(\infty, 1)}$ preserves finite products. Hence, it induces a functor

$$\operatorname{Fun}^{\operatorname{opl}}(\mathcal{K},-)^{\otimes}:\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)})\to\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}).$$

Thus, Fun^{opl}(\mathcal{K}, \mathcal{Z})^{\otimes} is an \mathcal{O} -monoidal ($\infty, 1$)-category for an \mathcal{O} -monoidal ($\infty, 2$)-category \mathcal{Z} . We notice that it is functorial with respect to \mathcal{K} . We define a simplicial \mathcal{O} -monoidal ($\infty, 1$)-category

$$\operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes} : \Delta^{\operatorname{op}} \longrightarrow \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty, 1)})$$

which is given by $[n] \mapsto \operatorname{Fun}^{\operatorname{opl}}([n], \mathcal{Z})^{\otimes}$.

Notice that the 0th \mathcal{O} -monoidal $(\infty, 1)$ -category $\operatorname{Fun}^{\operatorname{opl}}([0], \mathcal{Z})^{\otimes}$ is equivalent to $u_1 \mathcal{Z}^{\otimes}$. The inclusion map $i : \{[0]\} \hookrightarrow \Delta^{\operatorname{op}}$ induces an adjunction

$$i^* : \operatorname{Fun}(\Delta^{\operatorname{op}}, \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})) \rightleftharpoons \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}) : i_*,$$

where i^* is the restriction and i_* is its right adjoint given by the right Kan extension along i. Hence, the equivalence $i^* \operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes} \simeq u_1 \mathcal{Z}^{\otimes}$ gives rise to a map

$$\operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes} \longrightarrow i_* u_1 \mathcal{Z}^{\otimes}$$

in Fun($\Delta^{\operatorname{op}}, \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$). By using the composite $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}) \simeq \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}) \to \operatorname{Mon}_{\mathcal{O}}^{\operatorname{lax}}(\operatorname{Cat}_{(\infty,1)}) \to \operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$, we regard it as a map in Fun($\Delta^{\operatorname{op}}, \operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$).

Let $A \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathcal{Z})$ be an \mathcal{O} -monoidale in \mathcal{Z} . We regard A as a morphism of ∞ -operads $A : \mathcal{O}^{\otimes} \to u_1 \mathcal{Z}^{\otimes}$ over \mathcal{O}^{\otimes} . By the right Kan extension along i, we obtain a morphism

$$i_*A: i_*\mathcal{O}^\otimes \longrightarrow i_*u_1\mathcal{Z}^\otimes$$

in Fun($\Delta^{\mathrm{op}}, \mathrm{Op}_{\infty/\mathcal{O}^{\otimes}}$).

Definition 6.1. We define $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ by the following pullback diagram in Fun $(\Delta^{\mathrm{op}}, \mathrm{Op}_{\infty/\mathcal{O}^{\otimes}})$

(6.1)
$$\begin{array}{cccc} \mathcal{Z}(A,A)^{\otimes}_{\bullet} & \longrightarrow & \operatorname{Fun}^{\operatorname{opl}}([\bullet],\mathcal{Z})^{\otimes} \\ \downarrow & & \downarrow \\ i_*\mathcal{O}^{\otimes} & \xrightarrow{i_*A} & i_*u_1\mathcal{Z}^{\otimes}. \end{array}$$

We would like to show that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a Δ^{op} -monoid object in $\mathsf{Mon}^{\mathrm{lax}}_{\mathcal{O}}(\mathrm{Cat}_{(\infty,1)})$, which is a full subcategory of $\mathrm{Op}_{\infty/\mathcal{O}^{\otimes}}$, when A is a map \mathcal{O} -monoidale. First, we show that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a Δ^{op} -monoid object of $\mathrm{Op}_{\infty/\mathcal{O}^{\otimes}}$, and hence it corresponds to a virtual coCartesian ($\Delta^{\mathrm{op}}, \mathcal{O}$)-duoidal ∞ -category by Remark 2.4.

Proposition 6.2. The functor $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a Δ^{op} -monoid object of $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$.

Proof. For a simplicial object F in an $(\infty, 1)$ -category which admits finite limits, we say that F is a category object if the inert maps $[n] \to [i]$ in Δ^{op} for i = 0, 1 induce an equivalence

$$F([n]) \to F(\{0,1\}) \times_{F(\{1\})} \cdots \times_{F(\{n-1\})} F(\{n-1,n\})$$

for each $n \ge 0$. In order to prove the proposition, it suffices to show that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a category object and that $\mathcal{Z}(A, A)^{\otimes}_{0}$ is a final object in $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$.

First, we shall show that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a category object in $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$. Since category objects are closed under limits, it suffices to show that $i_*\mathcal{O}^{\otimes}$, $i_*u_1\mathcal{Z}^{\otimes}$, and $\operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes}$ are category objects. For an ∞ -operad \mathcal{P}^{\otimes} over \mathcal{O}^{\otimes} , the simplicial object $i_*\mathcal{P}^{\otimes}$ is given by

$$[n]\mapsto \overbrace{\mathcal{P}^{\otimes}\times_{\mathcal{O}^{\otimes}}\cdots\times_{\mathcal{O}^{\otimes}}\mathcal{P}^{\otimes}}^{n+1}.$$

By this description, we see that $i_*\mathcal{O}^{\otimes}$ and $i_*u_1\mathcal{Z}^{\otimes}$ are category objects. By using an equivalence $\{0,1\} \coprod_{\{1\}} \cdots \coprod_{\{n-1\}} \{n-1,n\} \to [n]$ of $(\infty, 1)$ -categories and the fact that the Gray tensor product preserves colimits in each variable, we obtain an equivalence

$$(\{0,1\}\otimes\mathcal{Y})\coprod_{\{1\}\otimes\mathcal{Y}}\cdots\coprod_{\{n-1\}\otimes\mathcal{Y}}(\{n-1,n\}\otimes\mathcal{Y})\xrightarrow{\simeq}[n]\otimes\mathcal{Y}$$

of $(\infty, 2)$ -categories for any $(\infty, 1)$ -category \mathcal{Y} . This equivalence implies that the simplicial object $\operatorname{Map}_{\operatorname{Cat}_{(\infty,1)}}(\mathcal{Y}, \operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes})$ is a category object in $\operatorname{Mon}_{\mathcal{O}}(\mathcal{S})$. By the Yoneda lemma and the fact that $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}) \to \operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$ preserves limits, we see that $\operatorname{Fun}^{\operatorname{opl}}([\bullet], \mathcal{Z})^{\otimes}$ is a category object of $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$.

Next, we shall show that $\mathcal{Z}(A, A,)_0^{\otimes}$ is a final object of $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$. Since the right vertical arrow in (6.1) induces an equivalence, so does the left vertical arrow. The claim follows from the equivalence $i_*\mathcal{O}^{\otimes}([0]) \simeq \mathcal{O}^{\otimes}$.

Finally, when A is a map \mathcal{O} -monoidale in \mathcal{Z} , we show that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is a Δ^{op} -monoid object of $\mathsf{Mon}^{\mathrm{lax}}_{\mathcal{O}}(\mathrm{Cat}_{(\infty,1)})$, which corresponds to a coCartesian ($\Delta^{\mathrm{op}}, \mathcal{O}$)-duoidal ∞ -category by Remark 2.6.

Theorem 6.3. We assume that A is a map \mathcal{O} -monoidale in an \mathcal{O} -monoidal $(\infty, 2)$ -category \mathcal{Z} . Then, the functor $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is an object of $\operatorname{Mon}_{\Delta^{\operatorname{op}}}(\operatorname{Mon}^{\operatorname{lax}}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)}))$, and hence it corresponds to a coCartesian $(\Delta^{\operatorname{op}}, \mathcal{O})$ -duoidal ∞ -category.

Proof. By Proposition 6.2, it suffices to show that $\mathcal{Z}(A, A)_1^{\otimes}$ is an \mathcal{O} -monoidal ∞ -category. We write $\operatorname{Ar}^{\operatorname{opl}}(\mathcal{Z})^{\otimes}$ for $\operatorname{Fun}^{\operatorname{opl}}([1], \mathcal{Z})^{\otimes}$. Since A is a map \mathcal{O} -monoidale, the map $A : \mathcal{O}^{\otimes} \to u_1 \mathcal{Z}^{\otimes}$

factors through $(u_1 \mathcal{Z}^L)^{\otimes}$. We consider the following pullback diagram

$$\begin{array}{cccc} \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z}) & \longrightarrow & \operatorname{Ar}^{\operatorname{opl}}(\mathcal{Z})^{\otimes} \\ & & & & \downarrow \\ (s,t)_{sL}^{\otimes} & & & \downarrow \\ (u_1\mathcal{Z}^L)^{\otimes} \times_{\mathcal{O}^{\otimes}} u_1\mathcal{Z}^{\otimes} & \longrightarrow & u_1\mathcal{Z}^{\otimes} \times_{\mathcal{O}^{\otimes}} u_1\mathcal{Z}^{\otimes} \end{array}$$

in $\operatorname{Op}_{\infty/\mathcal{O}^{\otimes}}$. Since $\mathcal{Z}(A, A)_1^{\otimes} \to \mathcal{O}^{\otimes}$ is obtained by taking pullback of $(s, t)_{sL}^{\otimes}$ along the map $(A, A) : \mathcal{O}^{\otimes} \to (u_1 \mathcal{Z}^L)^{\otimes} \times_{\mathcal{O}^{\otimes}} u_1 \mathcal{Z}^{\otimes}$, it suffices to show that $(s, t)_{sL}^{\otimes}$ is a coCartesian fibration.

We have a commutative diagram



where p and q are coCartesian fibrations. First, we shall show that $(s,t)_{sL}^{\otimes}$ is a locally coCartesian fibration. For this, by the dual of [17, Proposition 2.4.2.11], it suffices to show that the following two conditions hold: (1) The map $(s,t)_{sL}^{\otimes}$ carries p-coCartesian morphisms to q-coCartesian morphisms. (2) For any $a \in \mathcal{O}^{\otimes}$, the induced map $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_a \to (u_1 \mathcal{Z}^L)_a^{\otimes} \times u_1 \mathcal{Z}_a^{\otimes}$ on fibers is a coCartesian fibration.

For (1), we let $f: x \to x'$ be an object of $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})$ over $a \in \mathcal{O}^{\otimes}$, and let $\phi: a \to b$ be a morphism in \mathcal{O}^{\otimes} . We may assume that $\phi: a \to b$ is active, where $a \simeq \{a_i\}_{i \in |I|}$, and $a_i, b \in \mathcal{O}$. We can write $f \simeq \{f_i\}_{i \in |I|}$, where $f_i: x_i \to x'_i$. A *p*-coCartesian morphism $f \to \phi_* f$ is identified with $\{f_i\}_{i \in |I|} \to \otimes_{\phi} f_i$. Then $(s, t)_{sL}^{\otimes}(f \to \phi_* f)$ is identified with $\{(x_i, x'_i)\}_{i \in |I|} \to (\otimes_{\phi} x_i, \otimes_{\phi} x'_i)$. Thus, $(s, t)_{sL}^{\otimes}$ carries *p*-coCartesian morphisms to *q*-coCartesian morphisms. For (2), we may assume that $a \in \mathcal{O}$. Then the map $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_a \to (u_1 \mathcal{Z}_a)^{\otimes} \times u_1 \mathcal{Z}_a^{\otimes}$ is a coCartesian fibration by Corollary 4.6.

Finally, we would like to show that $(s,t)_{sL}^{\otimes}$ is in fact a coCartesian fibration. By the dual of [17, Proposition 2.4.2.8] together with the characterization of locally $(s,t)_{sL}^{\otimes}$ -coCartesian morphisms in the dual of [17, Proposition 2.4.2.11], it suffices to show that the following condition holds: (3) For any morphism $\phi : a \to b$ in \mathcal{O}^{\otimes} , the induced functor $\phi_* : \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_a \to \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_b$ carries $(s,t)_{sL,a}^{\otimes}$ -coCartesian morphisms to $(s,t)_{sL,b}^{\otimes}$ -coCartesian morphisms.

For (3), we may assume that $\phi : a \to b$ is active, where $a \simeq \{a_i\}_{i \in |I|}$, and $a_i, b \in \mathcal{O}$. We let $\{f_i\}_{i \in |I|} \to \{g_i\}_{i \in |I|}$ be a morphism of $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_a \simeq \prod_{i \in |I|} \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})_{a_i}$. Then we have an oplax square σ_i :



for each $i \in |I|$, where m_i^L is a left adjoint 1-morphism in \mathcal{Z}_{a_i} . By Lemma 4.5, the morphism $\{f_i\}_{i\in |I|} \to \{g_i\}_{i\in |I|}$ is $(s,t)_{sL,a}^{\otimes}$ -coCartesian if and only if the map $v_i : n_i f_i m_i^R \Longrightarrow g_i$ is invertible for each i, where v_i is a 2-morphism corresponding to u_i under the equivalence in Lemma 4.1. We

can identify the morphism $\phi_*(\{f_i\}_{i\in |I|} \to \{g_i\}_{i\in |I|})$ with an oplax square σ :



By Lemma 4.5, σ is an $(s,t)_{sL,b}^{\otimes}$ -coCartesian morphism if and only if $v : (\otimes_{\phi} n_i)(\otimes_{\phi} f_i)(\otimes_{\phi} m_i)^R \Rightarrow \otimes_{\phi} g_i$ is invertible, where $(\otimes_{\phi} m_i)^R$ is a right adjoint to $\otimes_{\phi} m_i^L$. Since there is an equivalence $(\otimes_{\phi} m_i)^R \simeq \otimes_{\phi} (m_i^R)$, we see that $v \simeq \otimes_{\phi} v_i$ is invertible if v_i is invertible for each i.

Part 2. Convolution product in monoidal $(\infty, 2)$ -categories

In ordinary category theory the endomorphism category of a map monoidale has the structure of a duoidal category. One of the two monoidal products is given by composition of morphisms, and the other is by convolution product.

Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category and let A be a map \mathcal{O} -monoidale in \mathcal{Z} . In §6 we showed that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ corresponds to a coCartesian $(\Delta^{\operatorname{op}}, \mathcal{O})$ -duoidal ∞ -category. The $\Delta^{\operatorname{op}}$ -monoidal structure is given by composition of morphisms in \mathcal{Z} . In Part 2 we show that the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ is given by convolution product.

In §7 we introduce convolution product on the mapping ∞ -category $\mathcal{Z}(C, A)$ from an \mathcal{O} comonoidale C to an \mathcal{O} -monoidale A in an \mathcal{O} -monoidal $(\infty, 2)$ -category \mathcal{Z} . We construct an \mathcal{O} -monoidal ∞ -category $\mathcal{Z}(C, A)_{\text{conv}}^{\otimes}$ whose underlying ∞ -category is $\mathcal{Z}(C, A)$. In §10 we prove
an equivalence of \mathcal{O} -monoidal ∞ -categories between $\mathcal{Z}(A, A)_1^{\otimes}$ and $\mathcal{Z}(A^*, A)_{\text{conv}}^{\otimes}$, where A is a left
adjoint \mathcal{O} -monoidale and A^* is the right adjoint \mathcal{O} -comonoidale associated to A.

In order to prove this, we need to establish an equivalence between some wide subcategories of the twisted arrow ∞ -category and of the oplax arrow ∞ -category. For this purpose, in §8 we introduce a twisted square ∞ -category, which is a generalization of twisted arrow ∞ -category. In §9, we show that there is such an equivalence by constructing a perfect pairing between them by taking a suitable subcategory of the twisted square ∞ -category.

7. Convolution product

Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category over a perfect operator category. We take an \mathcal{O} monoidale A and an \mathcal{O} -comonoidale C in \mathcal{Z} . In this section we introduce a convolution product on the mapping ∞ -category $\mathcal{Z}(C, A)$, which gives $\mathcal{Z}(C, A)$ the structure of an \mathcal{O} -monoidal $(\infty, 1)$ category.

Definition 7.1 (cf. [16, Definition 7.15]). For $n \ge 0$, we denote by $[n]^{\text{op}} \star_{\text{opl}} [n]$ a strict 2-category informally depicted as

$$\overline{0} \xleftarrow{\overline{1}} \overbrace{\overline{2}} \xleftarrow{\overline{2}} \xleftarrow{\overline{n}} \overbrace{\overline{n}}$$

$$\downarrow \Longrightarrow \downarrow \Longrightarrow \downarrow \Longrightarrow \Longrightarrow \downarrow$$

$$0 \longrightarrow 1 \longrightarrow 2 \longrightarrow \cdots \longrightarrow n.$$

The functor $[n] \mapsto [n]^{\text{op}} \star_{\text{opl}} [n]$ forms a cosimplicial object in the category of strict 2-categories.

Definition 7.2. For an $(\infty, 2)$ -category \mathcal{X} , we define a simplicial ∞ -groupoid $\operatorname{Tw}^{l}(\mathcal{X})_{\bullet}$ by

$$\operatorname{Tw}^{l}(\mathcal{X})_{n} = \operatorname{Map}_{\operatorname{Cat}_{(\infty,2)}}([n]^{\operatorname{op}} \star_{\operatorname{opl}} [n], \mathcal{X}).$$

As in the proof of [16, Corollary 7.17], we see that $\operatorname{Tw}^{l}(\mathcal{X})_{\bullet}$ is a complete Segal ∞ -groupoid. We denote by $\operatorname{Tw}^{l}(\mathcal{X})$ the $(\infty, 1)$ -category corresponding to $\operatorname{Tw}^{l}(\mathcal{X})_{\bullet}$.

The inclusion functors $[n]^{\mathrm{op}} \hookrightarrow [n]^{\mathrm{op}} \star_{\mathrm{opl}} [n]$ and $[n] \hookrightarrow [n]^{\mathrm{op}} \star_{\mathrm{opl}} [n]$ induce a functor

$$(s,t): \operatorname{Tw}^{l}(\mathcal{X}) \longrightarrow u_{1}\mathcal{X}^{\operatorname{op}} \times u_{1}\mathcal{X}.$$

We understand that an object of $\operatorname{Tw}^{l}(\mathcal{X})$ is a 1-morphism $f: x \to y$ in \mathcal{X} and that a morphism of $\operatorname{Tw}^{l}(\mathcal{X})$ from $f: x \to y$ to $f': x' \to y'$ is a diagram σ :

$$\begin{array}{cccc} x & \xleftarrow{g} & x' \\ f & \xrightarrow{\alpha} & \downarrow f' \\ y & \xrightarrow{h} & y' \end{array}$$

in \mathcal{X} . By [1] (see also [16, §7]), the map $(s, t) : \operatorname{Tw}^{l}(\mathcal{X}) \longrightarrow u_{1}\mathcal{X}^{\operatorname{op}} \times u_{1}\mathcal{X}$ is a coCartesian fibration classified by the restricted mapping ∞ -category functor of \mathcal{X} :

$$\mathcal{X}(-,-): u_1\mathcal{X}^{\mathrm{op}} \times u_1\mathcal{X} \longrightarrow \mathrm{Cat}_{(\infty,1)}.$$

We note that a morphism $\sigma : f \to f'$ in $\operatorname{Tw}^{l}(\mathcal{X})$ is (s, t)-coCartesian if and only if the 2-morphism α is invertible in \mathcal{X} .

Now, we let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category, where \mathcal{O}^{\otimes} is an ∞ -operad over a perfect operator category. First, we show that $\operatorname{Tw}^{l}(\mathcal{Z})$ is an \mathcal{O} -monoidal $(\infty, 1)$ -category. The construction $\mathcal{X} \mapsto \operatorname{Tw}^{l}(\mathcal{X})$ determines a functor $\operatorname{Tw}^{l} : \operatorname{Cat}_{(\infty,2)} \to \operatorname{Cat}_{(\infty,1)}$. Since the functor Tw^{l} preserves finite products, it induces a functor $\operatorname{Tw}^{l} : \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)}) \to \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$. Hence, we obtain the following lemma.

Lemma 7.3. If \mathcal{Z} is an \mathcal{O} -monoidal $(\infty, 2)$ -category, then $\operatorname{Tw}^{l}(\mathcal{Z})$ has an \mathcal{O} -monoidal structure.

We have a natural transformation (s,t): $\operatorname{Tw}^{l}(-) \to u_{1}(-)^{\operatorname{op}} \times u_{1}(-)$. Since the functors $\operatorname{Tw}^{l}(-)$ and $u_{1}(-)^{\operatorname{op}} \times u_{1}(-)$ preserve finite products, we obtain the following lemma.

Lemma 7.4. The map $(s,t): \operatorname{Tw}^{l}(\mathcal{Z}) \to (u_{1}\mathcal{Z})^{\operatorname{op}} \times u_{1}\mathcal{Z}$ is a strong \mathcal{O} -monoidal functor.

By Lemma 7.4, we have a commutative diagram in the ∞ -category of ∞ -operads



where p and q are coCartesian fibrations, and the top horizontal arrow $(s, t)^{\otimes}$ preserves coCartesian morphisms. Next, we show that the map $(s, t)^{\otimes} : \operatorname{Tw}^{l}(\mathcal{Z})^{\otimes} \to (u_{1}\mathcal{Z}^{\operatorname{op}})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes}$ is a coCartesian fibration.

Lemma 7.5. The map $(s,t)^{\otimes} : \operatorname{Tw}^{l}(\mathcal{Z})^{\otimes} \to (u_{1}\mathcal{Z}^{\operatorname{op}})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes}$ is a coCartesian fibration.

Proof. First, we show that $(s,t)^{\otimes}$ is a locally coCartesian fibration. The maps p and q are coCartesian fibrations and the map $(s,t)^{\otimes}$ carries p-coCartesian morphisms to q-coCartesian morphisms. Since the map (s,t): $\operatorname{Tw}^{l}(\mathcal{Z}) \to u_{1}\mathcal{Z}^{\operatorname{op}} \times u_{1}\mathcal{Z}$ is a coCartesian fibration, we see that the map $\operatorname{Tw}^{l}(\mathcal{Z})_{a}^{\otimes} \to (u_{1}\mathcal{Z}^{\operatorname{op}})_{a}^{\otimes} \times u_{1}\mathcal{Z}_{a}^{\otimes}$ induced on fibers is also a coCartesian fibration for any $a \in \mathcal{O}^{\otimes}$. Thus, $(s,t)^{\otimes}$ is a locally coCartesian fibration by the dual of [17, Proposition 2.4.2.11].

Next, we show that $(s,t)^{\otimes}$ is in fact a coCartesian fibration. By the dual of [17, Proposition 2.4.2.8], it suffices to show that the induced functor $\phi_* : \operatorname{Tw}^l(\mathcal{Z})_a^{\otimes} \to \operatorname{Tw}^l(\mathcal{Z})_b^{\otimes}$ sends $(s,t)_a^{\otimes} - \operatorname{coCartesian}$ morphisms to $(s,t)_b^{\otimes}$ -coCartesian morphisms for any morphism $\phi : a \to b$ in \mathcal{O}^{\otimes} . We may assume that $\phi : a \to b$ is active, p(a) = I, $a \simeq (a_i)_{i \in |I|}$, and $a_i, b \in \mathcal{O}$. We can identify a morphism σ of $\operatorname{Tw}^l(\mathcal{Z})_a^{\otimes}$ with a family $(\sigma_i)_{i \in |I|}$, where σ_i is a diagram

$$egin{array}{cccc} x_i & \xleftarrow{y_i} & x'_i \ f_i & \xrightarrow{lpha_i} & igcup_{f'_i} \ y_i & \xrightarrow{h_i} & y'_i \end{array}$$

in $\mathcal{Z}_{a_i}^{\otimes}$. We see that the morphism $\phi_* \sigma \simeq \otimes_{\phi} \sigma_i$ is a diagram

$$\begin{array}{cccc} \otimes_{\phi} x_i & \xleftarrow{\otimes_{\phi} g_i} & \otimes_{\phi} x'_i \\ \otimes_{\phi} f_i & \Longrightarrow & & \downarrow \otimes_{\phi} f'_i \\ y_i & \xleftarrow{\otimes_{\phi} h_i} & \otimes_{\phi} y'_i \end{array}$$

in \mathcal{Z}_b^{\otimes} . If σ is an $(s,t)_a^{\otimes}$ -coCartesian morphism, then α_i is invertible for any $i \in |I|$. This implies that $\otimes_{\phi} \alpha_i$ is also invertible. By the characterization of coCartesian morphisms of twisted arrow ∞ -categories, we see that $\phi_*\sigma$ is an $(s,t)_b^{\otimes}$ -coCartesian morphism.

Now, we introduce a convolution product on the mapping ∞ -category $\mathcal{Z}(C, A)$, where $C \in \operatorname{coAlg}_{\mathcal{O}}(u_1 \mathcal{Z})$ is an \mathcal{O} -comonoidale and $A \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathcal{Z})$ is an \mathcal{O} -monoidale in \mathcal{Z} . The objects C and A determine morphisms $C : \mathcal{O}^{\otimes} \to (u_1 \mathcal{Z}^{\operatorname{op}})^{\otimes}$ and $A : \mathcal{O}^{\otimes} \to u_1 \mathcal{Z}^{\otimes}$ of ∞ -operads over \mathcal{O}^{\otimes} , respectively. We consider the following pullback diagram

(7.2)
$$\begin{array}{cccc} \mathcal{Z}(C,A)_{\mathrm{conv}}^{\otimes} & \longrightarrow & \mathrm{Tw}^{l}(\mathcal{Z})^{\otimes} \\ & & & & \downarrow \\ & & & \downarrow \\ \mathcal{O}^{\otimes} & \xrightarrow{(C,A)} & (u_{1}\mathcal{Z}^{\mathrm{op}})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes} \end{array}$$

We can regard (7.2) as a pullback diagram in the ∞ -category of ∞ -operads over \mathcal{O}^{\otimes} . Hence the left vertical arrow $\mathcal{Z}(C, A)_{\text{conv}}^{\otimes} \to \mathcal{O}^{\otimes}$ is a map of ∞ -operads.

Theorem 7.6. Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category. For an \mathcal{O} -monoidale $A \in \operatorname{Alg}_{\mathcal{O}}(u_1\mathcal{Z})$ and an \mathcal{O} -comonoidale $C \in \operatorname{coAlg}_{\mathcal{O}}(u_1\mathcal{Z})$ in \mathcal{Z} , the mapping $(\infty, 1)$ -category $\mathcal{Z}(C, A)$ has an \mathcal{O} monoidal structure.

Proof. It suffices to show that the map $\mathcal{Z}(C, A)_{\text{conv}}^{\otimes} \to \mathcal{O}^{\otimes}$ is a coCartesian fibration. Since the right vertical arrow in (7.2) is a coCartesian fibration by Lemma 7.5, so is the left vertical arrow. \Box

Definition 7.7. We call the \mathcal{O} -monoidal structure on $\mathcal{Z}(C, A)$ in Theorem 7.6 a convolution product.

Remark 7.8. We notice that the construction of convolution product is functorial. We consider a functor $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)}) \to \operatorname{Cat}_{(\infty,1)}$ which assigns to an \mathcal{O} -monoidal $(\infty,2)$ -category \mathcal{Z} the $(\infty,1)$ -category $\operatorname{coAlg}_{\mathcal{O}}(u_1\mathcal{Z})^{\operatorname{op}} \times \operatorname{Alg}_{\mathcal{O}}(u_1\mathcal{Z})$. We denote by $\operatorname{coAlg}_{\mathcal{O}}(u_1(-))^{\operatorname{op}} \times \operatorname{Alg}_{\mathcal{O}}(u_1(-)) \to$ $\operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,2)})$ a coCartesian fibration obtained by unstraightening the functor. Informally speaking, an object of $\operatorname{coAlg}_{\mathcal{O}}(u_1(-))^{\operatorname{op}} \times \operatorname{Alg}_{\mathcal{O}}(u_1(-))$ is a triple (\mathcal{Z}, C, A) in which \mathcal{Z} is an

18

 \mathcal{O} -monoidal $(\infty, 2)$ -category, C is an \mathcal{O} -comonoidale, and A is an \mathcal{O} -monoidale in \mathcal{Z} . We can construct a functor

$$\operatorname{coAlg}_{\mathcal{O}}(u_1(-)^{\operatorname{op}}) \times \operatorname{Alg}_{\mathcal{O}}(u_1(-)) \longrightarrow \operatorname{Mon}_{\mathcal{O}}(\operatorname{Cat}_{(\infty,1)})$$

which assigns to (\mathcal{Z}, C, A) the mapping ∞ -category $\mathcal{Z}(C, A)$ equipped with the convolution \mathcal{O} monoidal structure.

8. Twisted square ∞ -categories

In this section we introduce a twisted square ∞ -category of an $(\infty, 2)$ -category, which is a generalization of twisted arrow ∞ -category. We construct a simplicial space $\mathrm{TS}(X)_{\bullet}$ for an ∞ -bicategory X, which is equipped with a map to $\mathrm{Tw}^{r}(X)_{\bullet} \times \mathrm{Ar}^{\mathrm{lax}}(X)_{\bullet}$. We defer a proof of $\mathrm{TS}(X)_{\bullet}$ being in fact a complete Segal space to §11 below.

First, in order to introduce a twisted square ∞ -category, we define a cosimplicial object ts[•] of scaled simplicial sets.

Definition 8.1. We define a cosimplicial object ts_{UR}^{\bullet} of scaled simplicial sets. For $n \geq 0$, the underlying simplicial set of ts_{UR}^{\bullet} is $\Delta^2 \times \Delta^n$. We depict it by the following grid:



The set $T_{ts_{UR}^n}$ of thin 2-simplices is given by

$$\begin{split} T_{\mathrm{ts}_{\mathrm{UR}}^n} &= \{\Delta^{\{ijk,ijk',ijk''\}} | \ (i,j) = (0,0), (0,1), (1,1), \ 0 \leq k < k' < k'' \leq n \} \\ &\cup \{\Delta^{\{00k,01k',01k''\}} | \ 0 \leq k \leq k' < k'' \leq n \} \\ &\cup \{\Delta^{\{01k,01k',11k''\}} | \ 0 \leq k < k' \leq k'' \leq n \} \\ &\cup \{\Delta^{\{00k,01k',11k''\}} | \ 0 \leq k \leq k' \leq k'' \leq n \} \\ &\cup \{\Delta^{\{00k,01k',11k''\}} | \ 0 \leq k \leq k' \leq k'' \leq n \} \\ &\cup \{\mathrm{degenerate}\}. \end{split}$$

We can verify that $ts_{UR}^{\bullet} = \{ts_{UR}^n\}_{n\geq 0}$ forms a cosimplicial object.

Definition 8.2. We define a cosimplicial object ts_{LL}^{\bullet} of scaled simplicial sets. For $n \ge 0$, we consider the simplicial set $\Delta^n \star \Delta^{n, \text{op}} \star \Delta^n$. We depict it by the following grid:



For a vertex v = (i, k) of $\Delta^2 \times \Delta^n$, we set $\Omega(v) = 00k$ if i = 0, $\Omega(v) = 10k$ if i = 1, $\Omega(v) = 11k$ if i = 2. For a simplex σ of $\Delta^2 \times \Delta^n$, we denote by $\Omega(\sigma)$ the simplex of $\Delta^n \star \Delta^{n, \text{op}} \star \Delta^n$ spanned by $\Omega(v)$ for vertices v of σ . For a subcomplex K of $\Delta^2 \times \Delta^n$, we denote by $\Omega(K)$ the subcomplex of $\Delta^n \star \Delta^{n, \text{op}} \star \Delta^n$ spanned by $\Omega(\sigma)$ for simplices σ of K.

The underlying simplicial set of ts_{LL}^n is $\Omega(\Delta^2 \times \Delta^n)$. In other words, it is spanned by (n+2)-dimensional simplices

$$\Lambda$$
 {000,...,00k,10k,...10k',11k',...,11n}

for $0 \le k \le k' \le n$. The set $T_{\mathrm{ts}_{\mathrm{LL}}^n}$ of thin 2-simplices is given by

$$\begin{split} T_{\mathrm{ts}_{\mathrm{LL}}^n} &= \{ \Delta^{\{ijk,ijk',ijk''\}} | \ (i,j) = (0,0), (1,0), (1,1), \ 0 \le k < k' \le n \} \\ &\cup \{ \Delta^{\{00k,00k',10k''\}} | \ 0 \le k < k' \le k'' \le n \} \\ &\cup \{ \Delta^{\{10k,11k',11k''\}} | \ 0 \le k \le k' < k'' \le n \} \\ &\cup \{ \mathrm{degenerate} \}. \end{split}$$

We can verify that $ts_{LL}^{\bullet} = {ts_{LL}^n}_{n \ge 0}$ forms a cosimplicial object.

Definition 8.3. We define a cosimplicial object ts[•] of scaled simplicial sets. For $n \ge 0$, we have monomorphisms from $(\Delta^1 \times \Delta^n)_{\flat}$ to $\operatorname{ts}_{\operatorname{UR}}^n$ and $\operatorname{ts}_{\operatorname{LL}}^n$, respectively, given by $(i,k) \mapsto iik$. We define a scaled simplicial set ts^n as a pushout

$$\mathrm{ts}^n = \mathrm{ts}_{\mathrm{UR}}^n \coprod_{(\Delta^1 \times \Delta^n)_\flat} \mathrm{ts}_{\mathrm{LL}}^n.$$

We can verify that $ts^{\bullet} = \{ts^n\}_{n>0}$ forms a cosimplicial object.

Notation 8.4. For a subcomplex K of the underlying simplicial set of ts^n , we denote by K_{\dagger} the scaled simplicial set whose underlying simplicial set is K equipped with the induced scaling from ts^n .

For scaled simplicial sets A and B with B fibrant, we recall that $\operatorname{Map}^{\operatorname{sc}}(A, B)$ is the underlying ∞ -groupoid of the ∞ -bicategory FUN(A, B).

Definition 8.5. For an ∞ -bicategory X, we define a simplicial space $TS(X)_{\bullet}$ by

$$TS(X)_n = Map^{sc}(ts^n, X).$$

We show that $TS(X)_{\bullet}$ is a complete Segal space and hence it represents an $(\infty, 1)$ -category.

Proposition 8.6. For any ∞ -bicategory X, $TS(X)_{\bullet}$ is a complete Segal space.

We defer the proof of Proposition 8.6 to §11 below.

Remark 8.7. By the assignment $X \mapsto TS(X)_{\bullet}$, we obtain a functor $TS : (Set_{\Delta}^{sc})^{\circ} \to CSS^{\circ}$, where CSS is the category of simplicial spaces equipped with the complete Segal space model structure. We can verify that TS is an enriched functor between $(Set_{\Delta})^{\circ}$ -enriched categories. By taking simplicial nerves, we obtain a functor $TS : Cat_{(\infty,2)} \to Cat_{(\infty,1)}$ of ∞ -categories.

By restrictions to faces of ts^{\bullet} , we construct maps out of $TS(X)_{\bullet}$. For this purpose, we set

$$\partial^{\mathrm{T}} \mathrm{ts}^{n} = (\Delta^{\{0,1\}} \times \Delta^{n})_{\dagger},$$

$$\partial^{\mathrm{F}} \mathrm{ts}^{n} = (\Delta^{\{1,2\}} \times \Delta^{n})_{\dagger},$$

$$\partial^{\mathrm{R}} \mathrm{ts}^{n} = \Omega(\Delta^{\{0,1\}} \times \Delta^{n})_{\dagger},$$

$$\partial^{\mathrm{B}} \mathrm{ts}^{n} = \Omega(\Delta^{\{1,2\}} \times \Delta^{n})_{\dagger},$$

We can verify that $\partial^{\mathrm{F}} \mathrm{ts}^{\bullet}$, $\partial^{\mathrm{R}} \mathrm{ts}^{\bullet}$, $\partial^{\mathrm{T}} \mathrm{ts}^{\bullet}$, $\partial^{\mathrm{B}} \mathrm{ts}^{\bullet}$ form cosimplicial objects.

Definition 8.8. For f = T, F, R, B, we define simplicial spaces $\partial^f TS(X)_{\bullet}$ by

 $\partial^f \mathrm{TS}(X)_{\bullet} = \mathrm{Map}^{\mathrm{sc}}(\partial^f \mathrm{ts}^{\bullet}, X).$

Remark 8.9. For each $n \ge 0$, there are natural isomorphisms $\partial^{\mathrm{T}}\mathrm{TS}(X)_n \cong \mathrm{Fun}^{\mathrm{opl}}([n], X)^{\simeq}$ and $\partial^{\mathrm{F}}\mathrm{TS}(X)_n \cong \mathrm{Fun}^{\mathrm{lax}}([n], X)^{\simeq}$ of simplicial sets. Thus, the simplicial spaces $\partial^{\mathrm{T}}\mathrm{TS}(X)_{\bullet}$ and $\partial^{\mathrm{F}}\mathrm{TS}(X)_{\bullet}$ are complete Segal spaces.

As for $\partial^{\mathrm{R}}\mathrm{TS}(X)_{\bullet}$ and $\partial^{\mathrm{B}}\mathrm{TS}(X)_{\bullet}$, we show that they are also complete Segal spaces by comparing with twisted arrow ∞ -categories. We recall that rev : $\Delta^{\mathrm{op}} \to \Delta^{\mathrm{op}}$ is a functor given by assigning to a nonempty finite ordered set its reverse ordered set. For a simplicial object S_{\bullet} , we denote by $S_{\bullet}^{\mathrm{rev}}$ a simplicial object given by the composite functor $S_{\bullet} \circ \mathrm{rev}$. We notice that $\partial^{\mathrm{B}}\mathrm{TS}(X)_{\bullet}$ is isomorphic to $\partial^{\mathrm{R}}\mathrm{TS}(X)_{\bullet}^{\mathrm{rev}}$ as simplicial spaces. Thus, it suffices to show that $\partial^{\mathrm{R}}\mathrm{TS}(X)_{\bullet}$ is a complete Segal space.

We recall the twisted arrow ∞ -category of an $(\infty, 2)$ -category. Let $Q(\bullet)$ be a cosimplicial scaled simplicial set, in which Q(n) is given as in [1, Definition 2.2]. For a ∞ -bicategory X, we define a simplicial space

$$\operatorname{Tw}^{r}(X)_{\bullet} = \operatorname{Map}^{\operatorname{sc}}(Q(\bullet), X),$$

which is a complete Segal space by [25, Theorem 4.9].

We have the following lemma.

Lemma 8.10. The simplicial spaces $\partial^{\mathrm{R}}\mathrm{TS}(X)_{\bullet}$ and $\partial^{\mathrm{B}}\mathrm{TS}(X)_{\bullet}$ are complete Segal spaces. There are equivalences of complete Segal spaces $\partial^{\mathrm{R}}\mathrm{TS}(X)_{\bullet} \simeq \mathrm{Tw}^{r}(X)_{\bullet}$ and $\partial^{\mathrm{B}}\mathrm{TS}(X)_{\bullet} \simeq \mathrm{Tw}^{r}(X)_{\bullet}^{\mathrm{rev}}$.

Proof. Since we have an isomorphism $\partial^{B}TS(X)_{\bullet} \cong \partial^{R}TS(X)_{\bullet}^{rev}$ of simplicial spaces, it suffices to show that $\partial^{R}TS(X)_{\bullet}$ is a complete Segal space and that there is an equivalence of complete Segal spaces between $\partial^{R}TS(X)_{\bullet}$ and $Tw^{r}(X)_{\bullet}$. We notice that the cosimplicial scaled simplicial set $\partial^{R}ts^{\bullet}$ is isomorphic to $T(\bullet)$ defined in [25, Definition 5.1]. The claim follows from [25, Proposition 5.8 and Theorem 5.9].

Definition 8.11. For f = T, F, R, B, we define maps of complete Segal spaces by restrictions

$$\partial^f : \mathrm{TS}(X)_{\bullet} \longrightarrow \partial^f \mathrm{TS}(X)_{\bullet}$$

9. An equivalence between $\mathrm{Tw}^l_{sR}(\mathcal{X})$ and $\mathrm{Ar}^{\mathrm{opl}}_{sL}(\mathcal{X})$

For an $(\infty, 2)$ -category \mathcal{X} , we set

$$\operatorname{Tw}_{sR}^{l}(\mathcal{X}) = \operatorname{Tw}^{l}(\mathcal{X}) \times_{(u_{1}\mathcal{X} \times u_{1}\mathcal{X})} (u_{1}\mathcal{X}^{R} \times u_{1}\mathcal{X}).$$

We recall that $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}) = \operatorname{Ar}^{\operatorname{opl}}(\mathcal{X}) \times_{(u_1\mathcal{X}\times u_1\mathcal{X})} (u_1\mathcal{X}^L \times u_1\mathcal{X})$. We would like to show that there is an equivalence of $(\infty, 1)$ -categories between $\operatorname{Tw}_{sR}^l(\mathcal{X})$ and $\operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X})$, which is compatible with the dual equivalence of $u_1\mathcal{X}^R$ and $u_1\mathcal{X}^L$ in Corollary 4.9. We set

$$\operatorname{Tw}_{sL}^{r}(\mathcal{X}) \simeq \operatorname{Tw}_{sR}^{l}(\mathcal{X}^{2\text{-op}})^{\operatorname{op}}, \qquad \operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X}) \simeq \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}^{2\text{-op}}).$$

In this section we show that there is a dual equivalence of $(\infty, 1)$ -categories between $\operatorname{Tw}_{sL}^r(\mathcal{X})$ and $\operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X})$. For this purpose, we construct a perfect pairing between them by taking a suitable subcategory of the $(\infty, 1)$ -category of twisted squares in \mathcal{X} .

First, by using Proposition 8.6, we define an $(\infty, 1)$ -category of twisted squares in an $(\infty, 2)$ -category \mathcal{X} .

Definition 9.1. Let \mathcal{X} be an $(\infty, 2)$ -category represented by an ∞ -bicategory X. We denote by $TS(\mathcal{X})$ the $(\infty, 1)$ -category corresponding to the complete Segal space $TS(X)_{\bullet}$. We call $TS(\mathcal{X})$ the $(\infty, 1)$ -category of twisted squares in \mathcal{X} .

Informally speaking, the objects of $TS(\mathcal{X})$ are lax squares



and morphisms are diagrams



with 2-morphisms

in \mathcal{X} equipped with an equivalence

$$\alpha_{01} : (B_1 b_{001} m_{001}) \cdot (S_1 m_{001}) \cdot (b_{011} T_1) \xrightarrow{\simeq} (m_{111} a_{100} R_1) \cdot (m_{111} S_0) \cdot (F_1 a_{000})$$

in $\operatorname{Map}_{\mathcal{X}(x(000),x(111))}(f_{01},g_{01})$, where $f_{01} = b_{011}m_{011}a_{000}$ and $g_{01} = m_{111}a_{100}n_{101}b_{001}m_{001}$. The maps $\partial^f : \operatorname{TS}(X)_{\bullet} \to \partial^f \operatorname{TS}(X)_{\bullet}$ of complete Segal spaces for $f = \mathbb{R}$, \mathbb{F} induce a map of $(\infty, 1)$ -categories

$$(\partial^{\mathrm{R}}, \partial^{\mathrm{F}}) : \mathrm{TS}(\mathcal{X}) \longrightarrow \mathrm{Tw}^{r}(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X}),$$

where $\operatorname{Ar}^{\operatorname{lax}}(\mathcal{X})$ is the lax arrow ∞ -category represented by $\operatorname{Fun}^{\operatorname{lax}}(\Delta^1_{\sharp}, X)$ (cf. [16, Remark 7.7]).

Now, we study conditions in which diagram (9.2) is an $(\partial^{R}, \partial^{F})$ -coCartesian morphism. For this purpose, we consider a diagram



in \mathcal{X} . We set

$$\begin{aligned} f_{01} &= b_{011}m_{011}a_{000}, \\ f_{12} &= b_{012}m_{012}a_{001}, \\ f_{02} &= b_{012}m_{012}m_{011}a_{000}. \end{aligned}$$

and

 $\begin{array}{rcl} g_{01} & = & m_{111}a_{100}n_{101}b_{001}m_{001}, \\ g_{12} & = & m_{112}a_{101}n_{102}b_{002}m_{002}, \\ g_{02} & = & m_{112}m_{111}a_{100}n_{101}n_{102}b_{002}m_{002}m_{001}. \end{array}$

By composition with T_1 and B_1 , we obtain a map of spaces

$$B_1(-)T_1: \operatorname{Map}_{\mathcal{X}(x(001),x(112))}(f_{12},g_{12}) \longrightarrow \operatorname{Map}_{\mathcal{X}(x(000),x(112))}(f_{02},g_{02}).$$

Lemma 9.2. We assume that m_{001} admits a right adjoint $m_{001}^R : x(001) \to x(000)$, that a mate $m_{011}a_{000}m_{001}^R \Rightarrow a_{001}$ of T_1 is invertible and that the 2-morphism B_1 is invertible. Then the map $B_1(-)T_1$ is an equivalence.

Proof. By the assumption that B_1 is invertible, the induced map

$$(B_1)_* : \operatorname{Map}_{\mathcal{X}(x(000), x(112))}(f_{02}, g_{12}m_{001}) \xrightarrow{\simeq} \operatorname{Map}_{\mathcal{X}(x(000), x(112))}(f_{02}, g_{02})$$

is an equivalence. Since m_{001} admits a right adjoint m_{001}^R , we have an equivalence

$$\operatorname{Map}_{\mathcal{X}(x(001),x(112))}(f_{02}m_{001}^{R},g_{12}) \simeq \operatorname{Map}_{\mathcal{X}(x(000),x(112))}(f_{02},g_{12}m_{001})$$

by Lemma 4.1. The lemma follows from the fact that there is an equivalence $f_{12} \simeq f_{02} m_{001}^R$ since a mate of T_1 is invertible. \square

For simplicity, we write

$$\mathcal{X}(f_{01}, g_{01}), \quad \mathcal{X}(f_{12}, g_{12}), \quad \mathcal{X}(f_{02}, g_{02})$$

for $\operatorname{Map}_{\mathcal{X}(x(000),x(111))}(f_{01},g_{01}), \operatorname{Map}_{\mathcal{X}(x(001),x(112))}(f_{12},g_{12}), \operatorname{Map}_{\mathcal{X}(x(000),x(112))}(f_{02},g_{02}),$ respectively. We set ות

Φ_{01}	=	$B_1 b_{001} m_{001} \cdot S_1 m_{001} \cdot b_{011} T_1$
Φ_{12}	=	$B_2 b_{002} m_{002} \cdot S_2 m_{002} \cdot b_{012} T_2$
Φ_{02}	=	$B_{12}b_{002}m_{002}m_{001} \cdot S_2m_{002}m_{001} \cdot b_{012}T_{12}$

and

Ψ_{01}	=	$m_{111}a_{100}R_1 \cdot m_{111}S_0 \cdot F_1a_{000}$
Ψ_{12}	=	$m_{112}a_{101}R_2 \cdot m_{112}S_1 \cdot F_2a_{001}$
Ψ_{02}	=	$m_{112}m_{111}a_{100}R_{12} \cdot m_{112}m_{111}S_0 \cdot F_{12}a_{000}$

where

$$\begin{array}{rclcrcrc} B_{12} & = & m_{112}B_1n_{102}\cdot B_2,\\ T_{12} & = & T_2m_{001}\cdot m_{012}T_1,\\ R_{12} & = & n_{101}R_2m_{001}\cdot R_1,\\ F_{12} & = & m_{112}F_1\cdot F_2m_{011}. \end{array}$$

Note that $\Phi_{01}, \Psi_{01} \in \mathcal{X}(f_{01}, g_{01}), \Phi_{12}, \Psi_{12} \in \mathcal{X}(f_{12}, g_{12}), \text{ and } \Phi_{02}, \Psi_{02} \in \mathcal{X}(f_{02}, g_{02}).$

р

Suppose that we have morphism (9.2). This determines an equivalence $\alpha_{01}: \Phi_{01} \xrightarrow{\simeq} \Psi_{01}$ in $\mathcal{X}(f_{01}, g_{01})$. Using this, we construct a map of spaces

$$\operatorname{Map}_{\mathcal{X}(f_{12},g_{12})}(\Phi_{12},\Psi_{12}) \longrightarrow \operatorname{Map}_{\mathcal{X}(f_{02},g_{02})}(\Phi_{02},\Psi_{02})$$

as follows: An equivalence $\alpha_{12}: \Phi_{12} \xrightarrow{\simeq} \Psi_{12}$ induces an equivalence $\Phi_{02} = B_1 \cdot \Phi_{12} \cdot T_1 \xrightarrow{\simeq} B_1 \cdot \Psi_{12} \cdot T_1$. We obtain an equivalence $B_1 \cdot \Psi_{12} \cdot T_1 \xrightarrow{\simeq} R_2 \cdot \Phi_{01} \cdot F_2$ by using the canonical equivalences $B_1 \cdot R_2 \simeq$ $R_2 \cdot B_1$ and $F_2 \cdot T_1 \simeq T_1 \cdot F_2$. Composing with the equivalence $\alpha_{01} : \Phi_{01} \xrightarrow{\simeq} \Psi_{01}$, we obtain an equivalence

$$\Phi_{02} = B_1 \cdot \Phi_{12} \cdot T_1 \xrightarrow{\alpha_{12}} B_1 \cdot \Psi_{12} \cdot T_1 \xrightarrow{\simeq} R_2 \cdot \Phi_{01} \cdot F_2 \xrightarrow{\alpha_{01}} R_2 \cdot \Psi_{01} \cdot F_2 = \Psi_{02}.$$

Lemma 9.3. The map $\operatorname{Map}_{\mathcal{X}(f_{12}, q_{12})}(\Phi_{12}, \Psi_{12}) \to \operatorname{Map}_{\mathcal{X}(f_{02}, q_{02})}(\Phi_{02}, \Psi_{02})$ is an equivalence.

Proof. By Lemma 9.2, there is an equivalence

$$\operatorname{Map}_{\mathcal{X}(f_{12},g_{12})}(\Phi_{12},\Psi_{12}) \xrightarrow{\simeq} \operatorname{Map}_{\mathcal{X}(f_{02},g_{02})}(B_1 \cdot \Phi_{12} \cdot T_1, B_1 \cdot \Psi_{12} \cdot T_1).$$

By composing with the equivalence $B_1 \cdot \Psi_{12} \cdot T_1 \xrightarrow{\simeq} R_2 \cdot \Phi_{01} \cdot F_2 \xrightarrow{\alpha_{01}} R_2 \cdot \Psi_{01} \cdot F_2 = \Psi_{02}$, we obtain the desired equivalence.

Proposition 9.4. If the 1-morphism m_{001} admits a right adjoint $m_{001}^R : x(001) \to x(000)$, a mate $m_{011}a_{000}m_{001}^R \Rightarrow a_{001}$ of T_1 is invertible, and the 2-morphism B_1 is invertible, then diagram (9.2) is an $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})$ -coCartesian morphism.

24

Proof. For simplicity, we set $TS = TS(\mathcal{X})$, $Tw = Tw^{r}(\mathcal{X})$ and $Ar = Ar^{lax}(\mathcal{X})$. We have to show that the following commutative diagram

$$\begin{array}{cccc} \operatorname{Map}_{\mathrm{TS}}(S_1, S_2) & \longrightarrow & \operatorname{Map}_{\mathrm{TS}}(S_0, S_2) \\ & & & \downarrow \\ & & & \downarrow \\ \operatorname{Map}_{\mathrm{Tw} \times \mathrm{Ar}}((b_{001}, b_{011}), (b_{002}, b_{012})) & \longrightarrow & \operatorname{Map}_{\mathrm{Tw} \times \mathrm{Ar}}((b_{000}, b_{010}), (b_{002}, b_{012})) \end{array}$$

is pullback. We take fibers of the vertical arrows at (R_2, F_2) and (R_{12}, F_{12}) , respectively, where $R_{12} = n_{101}R_2m_{001} \cdot R_1$ and $F_{12} = m_{112}F_1 \cdot F_2m_{011}$. It suffices to show that the induced map

$$\operatorname{Map}_{\mathrm{TS}}(S_1, S_2)_{(R_2, F_2)} \longrightarrow \operatorname{Map}_{\mathrm{TS}}(S_0, S_2)_{(R_{12}, F_{12})}$$

is an equivalence.

Taking top and bottom faces of diagram (9.3), we obtain a commutative diagram

where

and

$$btm_{12} = Map_{\mathcal{X}(x(102),x(112))}(m_{112}a_{101}n_{102},a_{102}),btm_{02} = Map_{\mathcal{X}(x(102),x(112))}(m_{112}m_{111}a_{100}n_{101}n_{102},a_{102}).$$

The map $top_{12} \rightarrow top_{02}$ is an equivalence by the assumptions that m_{001} admits a right adjoint and that a mate of T_1 is invertible. The map $btm_{12} \rightarrow btm_{02}$ is also an equivalence by the assumption that B_1 is invertible. Thus, it suffices to show that diagram (9.4) is pullback.

We take $(T_2, B_2) \in \text{top}_{12} \times \text{btm}_{12}$ and $(T_{12}, B_{12}) \in \text{top}_{02} \times \text{btm}_{02}$, where $T_{12} = T_2 m_{001} \cdot m_{012} T_1$ and $B_{12} = m_{112} B_1 n_{102} \cdot B_2$. It suffices to show that the induced map on fibers is an equivalence. We can identify it with the map $\text{Map}_{\mathcal{X}(f_{12},g_{12})}(\Phi_{12},\Psi_{12}) \to \text{Map}_{\mathcal{X}(f_{02},g_{02})}(\Phi_{02},\Psi_{02})$, which is an equivalence by Lemma 9.3.

We define $\text{TS}_L(\mathcal{X})$ to be the wide subcategory of $\text{TS}(\mathcal{X})$ spanned by those morphisms (9.2) in which m_{001} is a left adjoint 1-morphism in \mathcal{X} . Then, we obtain a map

$$(\partial^{\mathrm{R}}, \partial^{\mathrm{F}})_{L} : \mathrm{TS}_{L}(\mathcal{X}) \longrightarrow \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X})$$

by restricting $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})$ to $\mathrm{TS}_{L}(\mathcal{X})$. We notice that there is a pullback diagram

$$\begin{array}{ccc} \mathrm{TS}_{L}(\mathcal{X}) & \longrightarrow & \mathrm{TS}(\mathcal{X}) \\ & & & & & \\ \left(\partial^{\mathrm{R}}, \partial^{\mathrm{F}}\right)_{L} \\ & & & & \\ \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X}) & \longrightarrow & \mathrm{Tw}^{r}(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X}) \end{array}$$

Corollary 9.5. The map $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_L : \mathrm{TS}_L(\mathcal{X}) \to \mathrm{Tw}_{sL}^r(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X})$ is a coCartesian fibration. Diagram (9.2) is a $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_L$ -coCartesian morphism if and only if a mate $m_{011}a_{000}m_{001}^R \Rightarrow a_{001}$ of T_1 and the 2-morphism $B_1 : a_{101} \Rightarrow m_{111}a_{100}n_{101}$ are invertible. We define $TS_{L,R}(\mathcal{X})$ by the following pullback diagram

$$\begin{array}{cccc} \mathrm{TS}_{L,R}(\mathcal{X}) & \longrightarrow & \mathrm{TS}_{L}(\mathcal{X}) \\ (\partial^{\mathrm{R}}, \partial^{\mathrm{F}})_{L,R} & & & & & \\ \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}_{sR}^{\mathrm{lax}}(\mathcal{X}) & \longrightarrow & \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}^{\mathrm{lax}}(\mathcal{X}). \end{array}$$

By Corollary 9.5, the left vertical arrow $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})_{L,R}$ is a coCartesian fibration. We denote by $\mathrm{TS}_{L,R}^{\mathrm{cocart}}(\mathcal{X})$ the wide subcategory of $\mathrm{TS}_{L,R}(\mathcal{X})$ spanned by $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})_{L,R}$ -coCartesian morphisms. By the restriction of $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})_{L,R}$ to $\mathrm{TS}_{L,R}^{\mathrm{cocart}}(\mathcal{X})$, we obtain a left fibration

(9.5)
$$(\partial^{\mathrm{R}}, \partial^{\mathrm{F}})_{L,R}^{\mathrm{cocart}} : \mathrm{TS}_{L,R}^{\mathrm{cocart}}(\mathcal{X}) \to \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}_{sR}^{\mathrm{lax}}(\mathcal{X}).$$

Thus, we obtain a pairing between $\operatorname{Tw}_{sL}^{r}(\mathcal{X})$ and $\operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X})$.

Pairing (9.5) is not a perfect pairing in general. By restricting $\operatorname{TS}_{L,R}^{\operatorname{cocart}}(\mathcal{X})$ to a full subcategory, we construct a perfect pairing between $\operatorname{Tw}_{sL}^{r}(\mathcal{X})$ and $\operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X})$. We denote by $\operatorname{TS}_{L,R}^{\operatorname{pair}}(\mathcal{X})$ the full subcategory of $\operatorname{TS}_{L,R}^{\operatorname{cocart}}(\mathcal{X})$ spanned by those objects (9.1) in which the 1-morphism $a_{00}: x(00) \to x(01)$ is right adjoint. By restricting $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_{R,L}^{\operatorname{cocart}}$ to $\operatorname{TS}_{L,R}^{\operatorname{pair}}(\mathcal{X})$, we obtain a map

$$(\partial^{\mathrm{R}}, \partial^{\mathrm{F}})_{L,R}^{\mathrm{pair}} : \mathrm{TS}_{L,R}^{\mathrm{pair}}(\mathcal{X}) \to \mathrm{Tw}_{sL}^{r}(\mathcal{X}) \times \mathrm{Ar}_{sR}^{\mathrm{lax}}(\mathcal{X}).$$

Lemma 9.6. The map $(\partial^{\mathrm{R}}, \partial^{\mathrm{F}})_{L,R}^{\mathrm{pair}}$ is a left fibration.

Proof. Since $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_{L,R}^{\operatorname{coCart}}$ is a left fibration, it suffices to show that if the source of a morphism of $\operatorname{TS}_{L,R}^{\operatorname{cocart}}(\mathcal{X})$ is in $\operatorname{TS}_{L,R}^{\operatorname{pair}}(\mathcal{X})$, then the target is also in $\operatorname{TS}_{L,R}^{\operatorname{pair}}(\mathcal{X})$. We suppose that diagram (9.2) is an $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_{L,R}$ -coCartesian morphism and that a_{000} is a right adjoint 1-morphism. Then a_{001} is also a right adjoint 1-morphism since m_{001} is left adjoint and a mate $m_{001}a_{000}m_{001}^R \Rightarrow a_{001}$ of T_1 is invertible.

Theorem 9.7. The map $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_{L,R}^{\text{pair}} : \operatorname{TS}_{L,R}^{\text{pair}}(\mathcal{X}) \to \operatorname{Tw}_{sL}^{r}(\mathcal{X}) \times \operatorname{Ar}_{sR}^{\text{lax}}(\mathcal{X})$ is a perfect pairing.

Proof. By Lemma 9.6, the map $(\partial^{\mathbb{R}}, \partial^{\mathbb{F}})_{L,R}^{\text{pair}}$ is a pairing of ∞ -categories between $\operatorname{Tw}_{sL}^{r}(\mathcal{X})$ and $\operatorname{Ar}_{sR}^{\text{lax}}(\mathcal{X})$. In order to prove that it is perfect, by [18, Corollary 5.2.1.22], we need to show that it is both left and right representable, and that an object of $\operatorname{TS}_{L,R}^{\text{pair}}(\mathcal{X})$ is left universal if and only if it is right universal.

First, we shall show that $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})_{L,R}^{\text{pair}}$ is left representable. We regard a 1-morphism $b: x \to y$ in \mathcal{X} as an object of $\operatorname{Tw}_{sL}^r(\mathcal{X})$. We set $\mathcal{C}(b) = \{b\} \times_{\operatorname{Tw}_{sL}^r(\mathcal{X})} \operatorname{TS}_{L,R}^{\text{pair}}(\mathcal{X})$. We regard id_b as an object of $\mathcal{C}(b)$ and show that it is initial. We take an object σ of $\mathcal{C}(b)$ that is represented by a lax square



26

A morphism from id_b to σ in $\mathcal{C}(b)$ is represented by a cube



We note that the 2-morphisms $T_1 : m \Rightarrow a$ and $B_1 : a' \Rightarrow m'$ are invertible. By observing that there is an equivalence between $\operatorname{Map}_{\mathcal{C}(b)}(\operatorname{id}_b, \sigma)$ and $\operatorname{Map}_{\mathcal{X}(x,y')}(b'a, a'b)_{\sigma/}$, we see that $\operatorname{Map}_{\mathcal{C}(b)}(\operatorname{id}_b, \sigma)$ is contractible.

Next, we shall show that $(\partial^{\mathbf{R}}, \partial^{\mathbf{F}})_{L,R}^{\text{pair}}$ is right representable. We regard a 1-morphism $b: x \to y$ in \mathcal{X} as an object of $\operatorname{Ar}_{sR}^{\text{lax}}(\mathcal{X})$. We set $\mathcal{D}(b) = \{b\} \times_{\operatorname{Ar}_{sR}^{\text{lax}}(\mathcal{X})} \operatorname{TS}_{L,R}^{\text{pair}}(\mathcal{X})$. We regard id_{b} as an object of $\mathcal{D}(b)$ and show that it is initial. We take an object τ of $\mathcal{D}(b)$ that is represented by a lax square



A morphism from id_b to τ in $\mathcal{D}(b)$ is represented by a cube



We note that a mate $m^R \Rightarrow c$ of T_1 and the 2-morphism $B_1 : c' \Rightarrow n$ are invertible. By observing that there is an equivalence between $\operatorname{Map}_{\mathcal{D}(b)}(\operatorname{id}_b, \tau)$ and $\operatorname{Map}_{\mathcal{X}(x',y)}(bc,c'b')_{/\tau}$, we see that $\operatorname{Map}_{\mathcal{D}(b)}(\operatorname{id}_b, \tau)$ is contractible.

By the above argument, we see that an object



of $\operatorname{TS}_{L,R}^{\operatorname{pair}}(\mathcal{X})$ is left universal if and only if a and a' are equivalence 1-morphisms and S is an invertible 2-morphism in \mathcal{X} if and only if it is right universal.

Theorem 9.7 implies that there is a dual equivalence of $(\infty, 1)$ -categories between $\operatorname{Tw}_{sL}^r(\mathcal{X})$ and $\operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X})$. Next, we would like to show that this equivalence is compatible with the dual equivalence in Corollary 4.9.

Taking the top and bottom faces in (9.2), we obtain a map

$$(\partial^{\mathrm{T}}, \partial^{\mathrm{B}}) : \mathrm{TS}(\mathcal{X}) \longrightarrow \mathrm{Ar}^{\mathrm{opl}}(\mathcal{X}) \times \mathrm{Tw}^{l}(\mathcal{X}^{2\mathrm{-op}}).$$

By restriction, we obtain a map of pairings

Lemma 9.8. The map of pairings in (9.6) is left and right representable in the sense of [18, Variant 5.2.1.16].

Proof. Since the both pairings are perfect, it suffices to show that it is left representable. From the description of left representable objects in the pairings in the proofs of Theorems 4.8 and 9.7, we see that the map $(\partial^{\mathrm{T}}, \partial^{\mathrm{B}})$: $\mathrm{TS}_{L,R}^{\mathrm{pair}}(\mathcal{X}) \to \mathrm{Pair}(\mathcal{X}) \times \mathrm{Tw}^{l}(\mathcal{X}^{2-\mathrm{op}})$ carries left representable objects to left representable objects.

Making use of [18, Proposition 5.2.1.17], we obtain the following corollary by Theorems 4.8 and 9.7, and Lemma 9.8.

Corollary 9.9. There is a natural equivalence of $(\infty, 1)$ -categories

$$\operatorname{Tw}_{sL}^{r}(\mathcal{X})^{\operatorname{op}} \xrightarrow{\simeq} \operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{X})$$

which makes the following diagram commute

$$\begin{array}{cccc} \mathrm{Tw}_{sL}^{r}(\mathcal{X})^{\mathrm{op}} & \xrightarrow{\simeq} & \mathrm{Ar}_{sR}^{\mathrm{lax}}(\mathcal{X}) \\ (s,t)_{sL} & & & \downarrow \\ (u_{1}\mathcal{X}^{L})^{\mathrm{op}} \times u_{1}\mathcal{X} & \xrightarrow{\simeq} & u_{1}\mathcal{X}^{R} \times u_{1}\mathcal{X}. \end{array}$$

By applying Corollary 9.9 to the $(\infty, 2)$ -category $\mathcal{X}^{2\text{-op}}$, we obtain the following corollary.

Corollary 9.10. There is a natural equivalence of $(\infty, 1)$ -categories

$$\operatorname{Tw}_{sR}^{l}(\mathcal{X}) \xrightarrow{\simeq} \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X})$$

which makes the following diagram commute

$$\begin{array}{cccc} \operatorname{Tw}_{sR}^{l}(\mathcal{X}) & \xrightarrow{\simeq} & \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{X}) \\ & & & & \downarrow \\ (s,t)_{sR} & & & \downarrow \\ (u_{1}\mathcal{X}^{R})^{\operatorname{op}} \times u_{1}\mathcal{X} & \xrightarrow{\simeq} & u_{1}\mathcal{X}^{L} \times u_{1}\mathcal{X}. \end{array}$$

Let \mathcal{Z} be an \mathcal{O} -monoidal $(\infty, 2)$ -category and let A be a map \mathcal{O} -monoidale in \mathcal{Z} . In §6 we showed that $\mathcal{Z}(A, A)^{\otimes}_{\bullet}$ corresponds to a coCartesian $(\Delta^{\operatorname{op}}, \mathcal{O})$ -duoidal ∞ -category. In this section we show that the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, A)^{\otimes}_1$ given by Theorem 6.3 is equivalent to the convolution product $\mathcal{Z}(A^*, A)^{\otimes}_{\operatorname{conv}}$ defined in §7.

First, we promote the equivalences in Corollaries 9.9 and 9.10 to those of Cartesian symmetric monoidal ∞ -categories. We consider the functors $\mathrm{TS}_{L,R}^{\mathrm{pair}}(-)$, $\mathrm{Tw}_{sL}^{r}(-)$, $\mathrm{Ar}_{sR}^{\mathrm{lax}}(-)$, $\mathrm{Pair}(-) \times \mathrm{Tw}^{l}((-)^{2-\mathrm{op}})$, and $(u_{1}(-)^{L} \times u_{1}(-)^{R}) \times (u_{1}(-)^{\mathrm{op}} \times u_{1}(-))$ from $\mathrm{Cat}_{(\infty,2)}$ to $\mathrm{Cat}_{(\infty,1)}$. Since these functors preserve finite products, they extend to symmetric monoidal functors between the Cartesian symmetric monoidal ∞ -categories $\mathrm{Cat}_{(\infty,2)}^{\times}$ and $\mathrm{Cat}_{(\infty,1)}^{\times}$. Furthermore, the natural transformations appearing in Corollary 9.9 extend to natural transformations of symmetric monoidal functors. Hence, we obtain the following proposition.

Proposition 10.1. For an O-monoidal $(\infty, 2)$ -category \mathcal{Z} , there is a natural equivalence

$$\operatorname{Tw}_{sL}^r(\mathcal{Z})^{\otimes} \xrightarrow{\simeq} \operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{Z})^{\otimes}$$

of \mathcal{O} -monoidal $(\infty, 1)$ -categories, which makes the following diagram commute

$$\begin{array}{cccc} \operatorname{Tw}_{sL}^{r}(\mathcal{Z})^{\otimes} & \xrightarrow{\simeq} & \operatorname{Ar}_{sR}^{\operatorname{lax}}(\mathcal{Z})^{\otimes} \\ & & & \downarrow \\ (s,t)_{sL}^{\otimes} & & \downarrow \\ ((u_{1}\mathcal{Z}^{L})^{\operatorname{op}})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes} & \xrightarrow{\simeq} & (u_{1}\mathcal{Z}^{R})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes}. \end{array}$$

By applying Proposition 10.1 to the 2-opposite \mathcal{O} -monoidal $(\infty, 2)$ -category $(\mathcal{Z}^{2-\text{op}})^{\otimes}$, we obtain the following proposition.

Proposition 10.2. For an \mathcal{O} -monoidal $(\infty, 2)$ -category \mathcal{Z} , there is a natural equivalence

$$\operatorname{Tw}_{sR}^{l}(\mathcal{Z})^{\otimes} \xrightarrow{\simeq} \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})^{\otimes}$$

of \mathcal{O} -monoidal $(\infty, 1)$ -categories, which makes the following diagram commute

$$\begin{array}{cccc} \operatorname{Tw}_{sR}^{l}(\mathcal{Z})^{\otimes} & \xrightarrow{\simeq} & \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})^{\otimes} \\ & & & \downarrow \\ (s,t)_{sR}^{\otimes} \\ ((u_{1}\mathcal{Z}^{R})^{\operatorname{op}})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes} & \xrightarrow{\simeq} & (u_{1}\mathcal{Z}^{L})^{\otimes} \times u_{1}\mathcal{Z}^{\otimes} \end{array}$$

Next, we consider the \mathcal{O} -monoidal $(\infty, 1)$ -category $\mathcal{Z}(A, A)_1^{\otimes}$ for a map \mathcal{O} -monoidale A in \mathcal{Z} . We shall slightly generalize the construction of $\mathcal{Z}(A, A)_1^{\otimes}$. Let $B \in \text{Alg}_{\mathcal{O}}(u_1 \mathcal{Z})$ be an \mathcal{O} -monoidale in \mathcal{Z} . We define an ∞ -operad $\mathcal{Z}(A, B)^{\boxtimes}$ by the following pullback diagram

$$\begin{array}{cccc} \mathcal{Z}(A,B)^{\boxtimes} & \longrightarrow & \operatorname{Ar}_{sL}^{\operatorname{opl}}(\mathcal{Z})^{\otimes} \\ & & & & \downarrow \\ & & & \downarrow \\ \mathcal{O}^{\otimes} & \stackrel{(A,B)}{\longrightarrow} & (u_1\mathcal{Z}^L)^{\otimes} \times (u_1\mathcal{Z})^{\otimes} \end{array}$$

By Corollary 4.6, $\mathcal{Z}(A, B)^{\boxtimes}$ is an \mathcal{O} -monoidal ∞ -category. We notice that there is an equivalence $\mathcal{Z}(A, A)^{\boxtimes} \simeq \mathcal{Z}(A, A)_1^{\otimes}$ of \mathcal{O} -monoidal ∞ -categories.

By Proposition 10.2, we obtain the following theorem which means that the \mathcal{O} -monoidal structure on $\mathcal{Z}(A, B)^{\boxtimes}$ is equivalent to that given by convolution product. **Theorem 10.3.** Let $A \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathbb{Z}^L)$ be a map \mathcal{O} -monoidale and let $B \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathbb{Z})$ be an \mathcal{O} monoidale in \mathbb{Z} . We denote by $A^* \in \operatorname{coAlg}_{\mathcal{O}}(u_1 \mathbb{Z}^R)$ the right adjoint \mathcal{O} -comonoidale associated to A. There is an equivalence of \mathcal{O} -monoidal ∞ -categories

$$\mathcal{Z}(A^*, B)_{\operatorname{conv}}^{\otimes} \simeq \mathcal{Z}(A, B)^{\boxtimes}.$$

Proof. There is a commutative diagram

$$((u_1 \mathcal{Z}^R)^{\mathrm{op}})^{\otimes} \times u_1 \mathcal{Z}^{\otimes} \xrightarrow{\mathcal{O}^{\otimes}} (u_1 \mathcal{Z}^L)^{\otimes} \times u_1 \mathcal{Z}^{\otimes}$$

of ∞ -operads. The theorem follows by taking pullbacks along (A^*, B) and (A, B) in the commutative diagram in Proposition 10.2.

Corollary 10.4. Let $A \in \operatorname{Alg}_{\mathcal{O}}(u_1 \mathcal{Z}^L)$ be a map \mathcal{O} -monoidale in \mathcal{Z} . There is an equivalence of \mathcal{O} -monoidal ∞ -categories

$$\mathcal{Z}(A^*, A)_{\operatorname{conv}}^{\otimes} \simeq \mathcal{Z}(A, A)_1^{\otimes}.$$

11. Proof of Proposition 8.6

In this section we give a proof of Proposition 8.6, which says that $TS(X)_{\bullet}$ is a complete Segal space for any ∞ -bicategory X.

First, we show that $TS(X)_{\bullet}$ is a Segal space.

Proposition 11.1. For any ∞ -bicategory X, the simplicial object $TS(X)_{\bullet}$ is a Segal space.

In order to prove Proposition 11.1, we recall the following lemma.

Lemma 11.2 (cf. [25, Lemma 4.6]). Let A^{\bullet} be a cosimplicial object of $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$, and let X be an ∞ bicategory. If A^{\bullet} is Reedy cofibrant and satisfies the co-Segal condition, then the simplicial object Map^{sc} (A^{\bullet}, X) is a Segal space.

By Lemma 11.2, in order to prove Proposition 11.1, it suffices to show that the cosimplicial scaled simplicial set ts[•] is Reedy cofibrant and satisfies the co-Segal condition.

Lemma 11.3. The cosimplicial scaled simplicial set ts[•] is Reedy cofibrant.

Proof. The lemma follows by observing that the underlying simplicial set of the *n*th latching object of ts[•] is isomorphic to $(\Delta^2 \times \partial \Delta^n) \cup \Omega(\Delta^2 \times \partial \Delta^n)$, which is a subcomplex of the underlying simplicial set $(\Delta^2 \times \Delta^n) \cup \Omega(\Delta^2 \times \Delta^n)$ of tsⁿ.

Next, we would like to show that ts[•] satisfies the co-Segal condition. For this purpose, we make some preliminary results. For 0 < i < n, we set

$$\Lambda_i^n \text{ts} = ((\Delta^2 \times \Lambda_i^n) \cup \Omega(\Delta^2 \times \Lambda_i^n))_{\dagger}.$$

We show that the inclusion map $\Lambda_i^n \text{ts} \to \text{ts}^n$ is scaled anodyne. In order to prove this, we recall the following lemma, which is a special case of [1, Lemma 1.18]. Before that, we recall the notation. For a nonempty finite totally ordered set S and a subset $\emptyset \neq N \subsetneqq S$, we set

$$\Lambda_N^S = \bigcup_{s \in S-N} \Delta^{S-\{s\}}.$$

Lemma 11.4 ([25, Lemma 3.7], [1, Lemma 1.18]). Let $\Delta_{\dagger}^r = (\Delta^r, T)$ be a scaled simplicial set for $r \geq 3$, and let M be a nonempty subset of $\{0, 1, \ldots, r-1\}$. We assume that there exists an integer s with $0 \leq s < t$ such that $s \notin M$ and $i \in M$ for all $s < i \leq t$, where t is the largest number of M. Furthermore, we assume that $|M| \leq r-2$ and that the triangle $\Delta^{[r]-M}$ is not thin in Δ_{\dagger}^r when |M| = r-2. If triangles $\Delta^{\{i,t,t+1\}}$ are thin in Δ_{\dagger}^r for all $s \leq i < t$, then the inclusion map $\Lambda_{M,\dagger}^r \to \Delta_{\dagger}^r$ is scaled anodyne, where $\Lambda_{M,\dagger}^r$ is Λ_M^r equipped with the induced scaling from Δ_{\dagger}^r .

We decompose Λ_i^n into two parts. We set

$$\begin{split} \Lambda_i^n \mathbf{t}_{SUR} &= (\Delta^2 \times \Lambda_i^n)_{\dagger}, \\ \widehat{\Lambda}_i^n \mathbf{t}_{SLL} &= \Omega((\Delta^2 \times \Lambda_i^n) \cup (\Delta^{\{0,2\}} \times \Delta^n))_{\dagger}. \end{split}$$

Lemma 11.5. For any 0 < i < n, the inclusion map $\Lambda_i^n \operatorname{ts}_{\mathrm{UR}} \to \operatorname{ts}_{\mathrm{UR}}^n$ is scaled anodyne.

Proof. First, we shall show that the inclusion map $\Lambda_i^n \operatorname{tsu}_R \to \overline{\Lambda}_i^n \operatorname{tsu}_R$ is scaled anodyne, where $\overline{\Lambda}_i^n \operatorname{tsu}_R = \Lambda_i^n \operatorname{tsu}_R \cup (\Lambda_1^2 \times \Delta^n)_{\dagger}$. Since $\Lambda_{i,\sharp}^n \to \Delta_{\sharp}^n$ is a scaled anodyne map of type (An1), the inclusion map $\Lambda_i^n \operatorname{tsu}_R \to \Lambda_i^n \operatorname{tsu}_R \cup \bigcup_{i=0,1,2} (\Delta^{\{i\}} \times \Delta^n)_{\dagger}$ is scaled anodyne. By using [13, Proposition 2.16] and isomorphisms $(\Delta^{\{0,1\}} \times \Delta^n)_{\dagger} \cong \Delta_{\sharp}^1 \otimes \Delta_{\sharp}^n$ and $(\Delta^{\{1,2\}} \times \Delta^n)_{\dagger} \cong \Delta_{\sharp}^n \otimes \Delta_{\sharp}^1$, we see that the inclusion map $\Lambda_i^n \operatorname{tsu}_R \cup \bigcup_{i=0,1,2} (\Delta^{\{i\}} \times \Delta^n)_{\dagger} \to \overline{\Lambda}_i^n \operatorname{tsu}_R$ is scaled anodyne.

Next, we shall show that the inclusion map $\overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{UR}} \to \operatorname{ts}_{\mathrm{UR}}^n$ is scaled anodyne. For $0 \leq r \leq n$ and $0 \leq k \leq n-r$, we denote by $\sigma(r,k)$ the (n+2)-dimensional simplex

$$\Lambda$$
{000,...,00k,01k,...,01(k+r),11(k+r),...,11n}

of $\Delta^2 \times \Delta^n$. For $0 \le s \le n$, we set

$$X(s) = \overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{UR}} \cup \bigcup_{s \le r \le n} \bigcup_{0 \le k \le n-r} \sigma(r,k)_{\dagger}$$

Then, we obtain a filtration

$$\overline{\Lambda}_i^n \operatorname{ts}_{\operatorname{UR}} = X(n+1) \to X(n) \to X(n-1) \to \dots \to X(0) = \operatorname{ts}_{\operatorname{UR}}^n$$

It suffices to show that the inclusion map $X(s+1) \to X(s)$ is scaled anodyne for each $0 \le s \le n$.

We fix s with $0 \le s \le n$ and show that $X(s+1) \to X(s)$ is scaled anodyne. We identify $\sigma(s,k)_{\dagger}$ with a map $\sigma(s,k) : \Delta_{\dagger}^{n+2} \to X(s)$, where Δ_{\dagger}^{n+2} is Δ^{n+2} equipped with the induced scaling. There is a pushout diagram

where

$$M(k) = \begin{cases} \{00i, 01k, 01(k+s)\} & (0 < i < k), \\ \{01k, 01i, 01(k+s)\} & (k \le i \le k+s), \\ \{01k, 01(k+s), 11i\} & (k+s < i < n). \end{cases}$$

Thus, it suffices to show that the inclusion map $\Lambda_{M(k),\dagger}^{n+2} \to \Delta_{\dagger}^{n+2}$ is scaled anodyne. This follows from Lemma 11.4.

Lemma 11.6. For any 0 < i < n, the inclusion map $\widehat{\Lambda}_i^n \operatorname{ts}_{LL} \to \operatorname{ts}_{LL}^n$ is scaled anodyne.

Proof. First, we shall show that the inclusion map $\widehat{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \to \overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}}$ is scaled anodyne, where $\overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} = \widehat{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \cup \Omega(\Lambda_1^2 \times \Delta^n)_{\dagger}$. Since $\Lambda_{i,\sharp}^n \to \Delta_{\sharp}^n$ is a scaled anodyne map of type (An1), the inclusion map $\widehat{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \to \widehat{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \cup \Omega(\Delta^{\{1\}} \times \Delta^n)_{\dagger}$ is scaled anodyne. By the proof of [25, Lemma 5.5] and its dual, we see that the inclusion map $\widehat{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \cup \Omega(\Delta^{\{1\}} \times \Delta^n)_{\dagger} \to \Omega(\Delta^{\{1\}} \times \Delta^n)_{\dagger} \to \overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}}$ is scaled anodyne.

Next, we shall show that the inclusion map $\overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \to \operatorname{ts}_{\mathrm{LL}}^n$ is scaled anodyne. For $0 \leq r \leq n$ and $0 \leq k \leq n-r$, we denote by $\overline{\sigma}(r,k)$ the (n+2)-dimensional simplex

 $\Lambda \{000,...,00k,10k,...,10(k+r),11(k+r),...,11n\}$

of $\Delta^n \star \Delta^{n, \text{op}} \star \Delta^n$. For $0 \leq s \leq n$, we set

$$Y(s) = \overline{\Lambda}_i^n \operatorname{ts}_{\mathrm{LL}} \cup \bigcup_{0 \le r \le s} \bigcup_{0 \le k \le n-r} \overline{\sigma}(r,k)_{\dagger}.$$

Then, we obtain a filtration

$$\overline{\Lambda}_i^n \operatorname{ts}_{\operatorname{LL}} = Y(-1) \to Y(0) \to Y(1) \to \dots \to Y(n) = \operatorname{ts}_{\operatorname{LL}}^n$$

It suffices to show that the inclusion map $Y(s-1) \to Y(s)$ is scaled anodyne for each $0 \le s \le n$.

We fix s with $0 \le s \le n$ and show that $Y(s-1) \to Y(s)$ is scaled anodyne. We identify $\overline{\sigma}(s,k)_{\dagger}$ with a map $\overline{\sigma}(s,k) : \Delta_{\dagger}^{n+2} \to Y(s)$, where Δ_{\dagger}^{n+2} is Δ^{n+2} equipped with the induced scaling. There is a pushout diagram

where

$$M(k) = \begin{cases} \{10i, 11s\} & (0 = k < i < s < n), \\ \{10i\} & (0 = k < i < s = n), \\ \{11s, 11i\} & (0 = k \le s \le i < n), \\ \{00i, 00k, 11(k + s)\} & (0 < i \le k \le k + s < n), \\ \{00k, 10i, 11(k + s)\} & (0 < k < i < k + s < n), \\ \{00k, 11(k + s), 11i\} & (0 < k \le k + s \le i < n), \\ \{00i, 00k\} & (0 < i \le k \le k + s = n), \\ \{00k, 10i\} & (0 < k < i < k + s = n). \end{cases}$$

Thus, it suffices to show that the inclusion map $\Lambda_{M(k),\sharp}^{n+2} \to \Delta_{\dagger}^{n+2}$ is scaled anodyne. This follows from Lemma 11.4.

Lemma 11.7. For 0 < i < n, the inclusion map $\Lambda_i^n ts \to ts^n$ is scaled anodyne.

Proof. The map $\Lambda_i^n ts \to ts^n$ can be written as the composite $\Lambda_i^n ts \to \Lambda_i^n ts \cup ts_{\text{UR}}^n \to ts^n$. The first and second maps are scaled anodyne since they are obtained as pushouts of the maps in Lemmas 11.5 and 11.6, respectively.

Proof of Proposition 11.1. By Lemmas 11.2 and 11.3, it suffices to show that the cosimplicial object ts^{\bullet} satisfies the co-Segal condition. We can prove that the co-Segal map $ts^1 \coprod_{ts^0} \cdots \coprod_{ts^0} ts^1 \to ts^n$ is scaled anodyne by induction on n together with Lemma 11.7.

Next, we show that the Segal space $TS(X)_{\bullet}$ is complete. In order to prove this, we make some preliminary results.

By Definition 8.11, we have maps of Segal spaces

$$\partial^f : \mathrm{TS}(X)_{\bullet} \longrightarrow \partial^f \mathrm{TS}(X)_{\bullet}$$

for f = T, F, R, B. We set

$$\mathrm{Bd}(X)_{\bullet} = \prod_{f} \partial^{f} \mathrm{TS}(X)_{\bullet},$$

where the product is taken over f = T, F, R, B. Note that $Bd(X)_{\bullet}$ is a complete Segal space, that is, the map $s_0 : Bd(X)_1 \to Bd(X)_1^{eq}$ is an equivalence, where $Bd(X)_1^{eq}$ is the full subspace of $Bd(X)_1$ spanned by equivalences.

In order to prove that the Segal space $\mathrm{TS}(X)_{\bullet}$ is complete, it suffices to show that the map $s_0 : \mathrm{TS}(X)_0 \to \mathrm{TS}(X)_1^{\mathrm{eq}}$ is an equivalence. We denote by $\partial^{\mathrm{Bd}} : \mathrm{TS}(X)_{\bullet} \to \mathrm{Bd}(X)_{\bullet}$ the map of Segal spaces obtained as a product of ∂^f for $f = \mathrm{T}$, F, R, B. We define $\mathrm{TS}(X)_1^{\mathrm{bd,eq}}$ to be

$$\operatorname{TS}(X)_1 \times_{\operatorname{Bd}(X)_1} \operatorname{Bd}(X)_1^{\operatorname{eq}}$$

We notice that the map $s_0 : TS(X)_0 \to TS(X)_1$ factors through $TS(X)_1^{bd,eq}$.

Lemma 11.8. If $s_0 : TS(X)_0 \to TS(X)_1^{bd,eq}$ is an equivalence, then $s_0 : TS(X) \to TS(X)^{eq}$ is also an equivalence.

Proof. The lemma follows from the fact that $TS(X)_1^{eq}$ is a full subspace of $TS(X)_1^{bd,eq}$.

We consider the following commutative diagram

Since the bottom horizontal arrow is an equivalence, it suffices to show that diagram (11.1) is pullback in order to prove that the top horizontal arrow is an equivalence. For $P \in Bd(X)_0$, let $(s_0)_P : TS(X)_{0,P} \to TS(X)_{1,s_0(P)}^{bd,eq}$ be the induced map on fibers. We show that $(s_0)_P$ is an equivalence for any $P \in Bd(X)_0$. In fact, we show that $(d_i)_P : TS(X)_{1,s_0(P)}^{bd,eq} \to TS(X)_{0,P}$ is an equivalence for i = 0, 1.

We consider a scaled simplicial set

$$\mathrm{ts}^1_{\diamondsuit} = ((\Delta^2 \times \Delta^1) \cup \Omega(\Delta^2 \times \Delta^1), \widetilde{T})$$

where \widetilde{T} is a set of 2-simplices that consists of the thin triangles of ts^1 together with

$$\Delta^{\{000,001,011\}}, \quad \Delta^{\{010,110,111\}}, \quad \Delta^{\{000,100,101\}}, \\ \Delta^{\{100,101,111\}}, \quad \Delta^{\{000,001,111\}}, \quad \Delta^{\{000,110,111\}}.$$

Note that there is an inclusion map $ts^1 \to ts^1_{\Diamond}$ of scaled simplicial sets.

For a subcomplex K of (the underlying simplicial set of) ts^1 , we write K_{\diamond} for the scaled simplicial set K equipped with the induced scaling from ts^1_{\diamond} . We set

$$\mathrm{FSR}^{i} = (\partial^{\mathrm{F}} \mathrm{ts}^{1} \cup d^{i}(\mathrm{ts}^{0}) \cup \partial^{\mathrm{R}} \mathrm{ts}^{1})_{\Diamond}$$

for i = 0, 1. We consider inclusion maps

$$\begin{aligned} \theta_0 : & \operatorname{FSR}^0 \coprod_{\Delta_{\sharp}^{\{110,111\}}} \Delta_{\sharp}^{\{0\}} & \longrightarrow & \operatorname{ts}_{\diamondsuit}^1 \coprod_{\Delta_{\sharp}^{\{110,111\}}} \Delta_{\sharp}^{\{0\}}, \\ \theta_1 : & \operatorname{FSR}^1 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^{\{0\}} & \longrightarrow & \operatorname{ts}_{\diamondsuit}^1 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^{\{0\}}. \end{aligned}$$

Lemma 11.9. The inclusion maps θ_0 and θ_1 are trivial cofibrations in Set^{sc}_{Δ}.

Proof. We shall show that the map θ_1 is a trivial cofibration. The other case is similar. For simplicity, we set

$$E_{0} = \operatorname{FSR}^{1} \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^{\{0\}},$$

$$E_{1} = (\operatorname{FSR}^{1} \cup \operatorname{ts}_{\operatorname{LL}\Diamond}^{1}) \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^{\{0\}},$$

$$E_{2} = \operatorname{ts}_{\Diamond}^{1} \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^{\{0\}}.$$

It suffices to show that the inclusion maps $E_0 \to E_1$ and $E_1 \to E_2$ are trivial cofibrations. First, we shall show that $E_0 \to E_1$ is a trivial cofibration. We set

$$F_{0} = FSR^{1},$$

$$F_{1} = FSR^{1} \cup \Delta^{\{000,100,110,111\}}_{\Diamond},$$

$$F_{2} = FSR^{1} \cup (\Delta^{\{000,100,110,111\}} \cup \Delta^{\{000,100,101,111\}})_{\Diamond}.$$

We can see that the inclusion maps $F_0 \to F_1$ and $F_1 \to F_2$ are scaled anodyne since they are obtained by iterated pushouts along scaled anodyne maps of type (An1). By taking a pushout along the map $\Delta_{\sharp}^{\{000,001\}} \to \Delta_{\sharp}^{\{0\}}$, we see that $E_0 \to F_2 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^0$ is scaled anodyne. By [14, Remark 1.33], the map $\Lambda_{0,\sharp}^2 \coprod_{\Delta_{\sharp}^{\{0,1\}}} \Delta_{\sharp}^{\{0\}} \to \Delta_{\sharp}^2 \coprod_{\Delta_{\sharp}^{\{0,1\}}} \Delta_{\sharp}^{\{0\}}$ is a trivial cofibration. By taking a pushout along it, we obtain a trivial cofibration $F_2 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^0 \to E_1$. Hence, $E_0 \to E_1$ is a trivial cofibration by composition.

Next, we shall show that $E_1 \to E_2$ is a trivial cofibration. We set

We can see that the inclusion maps $G_0 \to G_1$ and $G_1 \to G_2$ are scaled anodyne since they are obtained by taking pushouts along scaled anodyne maps of type (An1). By taking a pushout along the map $\Delta_{\sharp}^{\{000,001\}} \to \Delta_{\sharp}^{\{0\}}$, we see that $E_1 \to G_2 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^0$ is scaled anodyne. By taking a pushout along the trivial cofibration $\Lambda_{0,\sharp}^2 \coprod_{\Delta_{\sharp}^{\{0,1\}}} \Delta_{\sharp}^{\{0\}} \to \Delta_{\sharp}^2 \coprod_{\Delta_{\sharp}^{\{0,1\}}} \Delta_{\sharp}^{\{0\}}$, we obtain a trivial cofibration $G_2 \coprod_{\Delta_{\sharp}^{\{000,001\}}} \Delta_{\sharp}^0 \to E_2$. Hence, $E_1 \to E_2$ is a trivial cofibration by composition. \Box

For i = 0, 1, Lemma 11.9 implies that the induced map

$$\Delta^{1}_{\sharp} \coprod_{\partial^{\mathrm{F}} \mathrm{ts}^{1}_{\Diamond}} \mathrm{FSR}^{i} \coprod_{\partial^{\mathrm{R}} \mathrm{ts}^{1}_{\Diamond}} \Delta^{1}_{\sharp} \longrightarrow \Delta^{1}_{\sharp} \coprod_{\partial^{\mathrm{F}} \mathrm{ts}^{1}_{\Diamond}} \mathrm{ts}^{1}_{\Diamond} \coprod_{\partial^{\mathrm{R}} \mathrm{ts}^{1}_{\Diamond}} \Delta^{1}_{\sharp}$$

is a trivial cofibration, where $\partial^{\mathrm{F}} \mathrm{ts}^{1}_{\Diamond} \to \Delta^{1}_{\sharp}$ is a map given by 010, 011 \mapsto 0 and 110, 111 \mapsto 1, and $\partial^{\mathrm{R}} \mathrm{ts}^{1}_{\Diamond} \to \Delta^{1}_{\sharp}$ is a map given by 000, 001 \mapsto 0 and 100, 101 \mapsto 1. Since the map $d^{i} : \mathrm{ts}^{0} \to \mathrm{ts}^{1}$ induces an isomorphism of scaled simplicial sets between ts^{0} and $\Delta^{1}_{\sharp} \coprod_{\partial^{\mathrm{F}} \mathrm{ts}^{1}_{\Diamond}} \mathrm{FSR}^{i} \coprod_{\partial^{\mathrm{R}} \mathrm{ts}^{1}_{\Diamond}} \Delta^{1}_{\sharp}$, it induces a trivial cofibration

$$\mathrm{ts}^{0} \longrightarrow \Delta^{1}_{\sharp} \coprod_{\partial^{\mathrm{F}} \mathrm{ts}^{1}_{\Diamond}} \mathrm{ts}^{1}_{\Diamond} \coprod_{\partial^{\mathrm{R}} \mathrm{ts}^{1}_{\Diamond}} \Delta^{1}_{\sharp}$$

in $\operatorname{Set}_{\Delta}^{\operatorname{sc}}$. This implies that the restriction map $d_i : \operatorname{TS}(X)_{1,(s_0(b_{000}),s_0(b_{010}))}^{\operatorname{eq}} \to \operatorname{TS}(X)_{0,(b_{000},b_{010})}$ is an equivalence. Hence, we obtain that $(d_i)_P : \operatorname{TS}(X)_{1,s_0(P)}^{\operatorname{bd,eq}} \to \operatorname{TS}(X)_{0,P}$ is an equivalence. Since the composite $(d_i)_P(s_0)_P$ is an identity, we obtain the following corollary.

Corollary 11.10. The map $(s_0)_P : TS(X)_{0,P} \to TS(X)_{1,s_0(P)}^{bd,eq}$ is an equivalence for any $P \in Bd(X)_0$.

Proof of Proposition 8.6. By Proposition 11.1 and Lemma 11.8, it suffices to show that the map $s_0 : TS(X)_0 \to TS(X)_1^{bd,eq}$ is an equivalence. By Corollary 11.10, (11.1) is a pullback diagram. Since the bottom horizontal arrow in (11.1) is an equivalence, the top horizontal arrow is also an equivalence.

References

- F. Abellán García, and W. H. Stern, Enhanced twisted arrow categories, Theory Appl. Categ. 39 (2023), Paper No. 4, 98–149.
- [2] M. Aguiar and S. Mahajan, Monoidal functors, species and Hopf algebras, CRM Monograph Series, 29. American Mathematical Society, Providence, RI, 2010.
- [3] C. Barwick, From operator categories to higher operads, Geom. Topol. 22 (2018), no. 4, 1893–1959.
- M. Batanin, and M. Markl, Centers and homotopy centers in enriched monoidal categories, Adv. Math. 230 (2012), no. 4–6, 1811–1858.
- [5] M. Batanin, and M. Markl, Operadic categories and duoidal Deligne's conjecture, Adv. Math.285(2015), 1630– 1687.
- [6] G. Böhm, Y. Chen, and L. Zhang, On Hopf monoids in duoidal categories, J. Algebra 394 (2013), 139–172.
- [7] G. Böhm, S. Lack, Hopf comonads on naturally Frobenius map-monoidales, J. Pure Appl. Algebra 220 (2016), no. 6, 2177–2213.
- [8] T. Booker and R. Street, Tannaka duality and convolution for duoidal categories, Theory Appl. Categ. 28 (2013), No. 6, 166–205.
- [9] Y. Chen, and G. Böhm, Weak bimonoids in duoidal categories, J. Pure Appl. Algebra218(2014), no.12, 2240– 2273.
- [10] H. Chu, and R. Haugseng, Homotopy-coherent algebra via Segal conditions, Adv. Math. 385 (2021), Paper No. 107733, 95 pp.
- [11] D. Gepner and R. Haugseng, Enriched ∞ -categories via non-symmetric ∞ -operads. Adv. Math. 279 (2015), 575–716.
- [12] D. Gepner, R. Haugseng, and T. Nikolaus, Lax colimits and free fibrations in ∞-categories. Doc. Math. 22 (2017), 1225–1266.
- [13] A. Gagna, Y. Harpaz, and E. Lanari, Gray tensor products and Lax functors of (∞, 2)-categories, Adv. Math.391(2021), Paper No. 107986, 32 pp.
- [14] A. Gagna, Y. Harpaz, and E. Lanari, On the equivalence of all models for $(\infty, 2)$ -categories, J. Lond. Math. Soc. (2)106(2022), no.3, 1920–1982.
- [15] R. Haugseng, On lax transformations, adjunctions, and monads in $(\infty, 2)$ -categories, High. Struct. 5 (2021), no. 1, 244–281.
- [16] R. Haugseng, F. Hebestreit, S. Linskens, and J. Nuiten, Two-variable fibrations, factorisation systems and ∞-categories of spans, Forum Math. Sigma 11 (2023), Paper No. e111, 70 pp.
- [17] J. Lurie, Higher topos theory, Annals of Mathematics Studies, 170. Princeton University Press, Princeton, NJ, 2009.
- [18] J. Lurie, Higher algebra, available at http://www.math.harvard.edu/~lurie/.
- [19] J. Lurie, $(\infty, 2)$ -categories and the Goodwillie calculus I, available at http://www.math.harvard.edu/~lurie/.
- [20] E. Riehl and D. Verity, Homotopy coherent adjunctions and the formal theory of monads, Adv. Math. 286 (2016), 802–888.
- [21] R. Street, Monoidal categories in, and linking, geometry and algebra, Bull. Belg. Math. Soc. Simon Stevin 19 (2012), no. 5, 769–821.
- [22] T. Torii, On duoidal ∞ -categories, preprint, arXiv:2106.14121.
- [23] T. Torii, On higher monoidal ∞ -categories, preprint, arXiv:2111.00158.
- [24] T. Torii, Duoidal ∞-categories of operadic modules, preprint, arXiv:2204.11152.
- [25] T. Torii, A simple model for twisted arrow ∞ -categories, preprint, arXiv:2405.06912.

DEPARTMENT OF MATHEMATICS, OKAYAMA UNIVERSITY, OKAYAMA 700-8530, JAPAN *Email address*: torii@math.okayama-u.ac.jp