

A Free-fermion Formulation of Two-dimensional Ising Models

De-Zhang Li,¹ Xin Wang,^{1,2,*} and Xiao-Bao Yang^{3,†}

¹Quantum Science Center of Guangdong-Hong Kong-Macao Greater Bay Area, Shenzhen 518045, China

²Department of Physics, City University of Hong Kong, Hong Kong SAR, China

³Department of Physics, South China University of Technology, Guangzhou 510640, China

Ising model is a famous toy model in condensed matter and statistical physics. In this work we present a free-fermion formulation of the two-dimensional classical Ising models on the honeycomb, triangular and Kagomé lattices. Each Ising model is studied in the case of a zero field and the case of an imaginary field $i(\pi/2)k_B T$. We employ the decorated lattice technique, star-triangle transformation and weak-graph expansion method to exactly map each Ising model in both cases into an eight-vertex model on the square lattice. The resulting vertex weights are shown to satisfy the free-fermion condition. In the zero field case, each Ising model is an even free-fermion model. In the case of the imaginary field, the Ising model on the honeycomb lattice is an even free-fermion model while those on the triangular and Kagomé lattices are odd free-fermion models. The exact solution of the Kagomé lattice Ising model in the imaginary field $i(\pi/2)k_B T$ is obtained, which has not been reported in previous works. The frustrated Ising models on the triangular and Kagomé lattices in the imaginary field still exhibit a non-zero residual entropy.

I. INTRODUCTION

Ising model plays a crucial role in the search of condensed matter and statistical physics of lattice systems, especially in understanding the phase transition via statistical mechanics. It has a long history since 1920s [1]. The one-dimensional Ising model was first solved by Ising himself [2]. Kramers and Wannier introduced the transfer matrix method to rederive the exact solution in one dimension and study the two-dimensional case [3, 4]. Onsager first used the transfer matrix method to present the exact solution of the two-dimensional Ising model on the square lattice in the case without a magnetic field [5]. This famous result has been obtained from various approaches [6–14]. Lee and Yang proposed the solution for the case in an imaginary field $i(\pi/2)k_B T$ in their work of Lee-Yang zeros [15]. This solution has also been given by a variety of different methods [16–22]. Onsager’s solution [5] has a great influence in the field of exactly solvable lattice models [23–27]. Since then the mathematical exact results for Ising models have shown their beauty and power, and attracted considerable attention. Two-dimensional Ising models on the honeycomb lattice [28–31], the triangular lattice [32–36], the Kagomé lattice [37, 38], and the checkerboard lattice [39, 40] were also exactly solved. Although the Ising model in a non-zero field is a well-known unsolved problem, the case of an imaginary field $i(\pi/2)k_B T$ originated in the work of Lee and Yang [15] is special and of significant interest.

In this work we focus on the exact solutions of isotropic Ising models on typical two-dimensional lattices, both in the zero field and in the imaginary field. The Hamilto-

nian of N spins $\{s_i = \pm 1, i = 1, \dots, N\}$ simply reads

$$H = \sum_{\langle ij \rangle} J s_i s_j - H_{\text{ex}} \sum_{i=1}^N s_i, \quad (1)$$

where the sum $\sum_{\langle ij \rangle}$ is over all nearest-neighbours and the field H_{ex} can be 0 or $i(\pi/2)k_B T$. The approach we employ is the mapping into the free-fermion model [41–45], which was proposed in the vertex model problem. Vertex model is also an important and intriguing issue in statistical mechanics. Studies of vertex models such as the six-vertex [46–57], eight-vertex [41–43, 58–66] and sixteen-vertex models [67–72] have a close relation with those of Ising models. The free-fermion models are the solvable eight-vertex models which satisfy the free-fermion condition. The exact solution of a free-fermion model can be obtained in a variety of ways, such as using the S -matrix [41], the Pfaffian [42] and the transfer matrix methods [73–78]. Usually an eight-vertex model on the square lattice is defined as either the even subcase or the odd subcase of the general sixteen-vertex model [72], as shown in Fig. 1. In this paper we adopt the notation of vertex configurations in Ref. [69] (see Fig. 1 there), and refer to the eight-vertex model of even/odd subcase satisfying the free-fermion condition as even/odd free-fermion model.

There have been lots of studies connecting the Ising model with the vertex model [21, 22, 79–82]. Ref. [21] introduced a rather simple and direct correspondence between the square lattice Ising spin states and the arrow configurations of the sixteen-vertex model. The value $+1/-1$ of an Ising spin is naturally mapped to the direction of the arrow settled on the corresponding site. By further performing a transformation from the sixteen vertex weights to those in the even subcase, the author showed that the Ising model is equivalent with an even free-fermion model, both for the zero field case and the imaginary field case. The present paper is inspired by the

* Corresponding author: x.wang@cityu.edu.hk

† Corresponding author: scxbyang@scut.edu.cn




FIG. 1. The vertex configurations of the sixteen-vertex model. The even subcase consists of vertices (1)–(8), while the odd subcase consists of vertices (9)–(16).

success of this work. The method of transforming the square lattice Ising model to the free-fermion model is extended and applied to the Ising models on the honeycomb, triangular and Kagomé lattices.

The structure of the paper is as follows. In Sec. II we briefly introduce the methods and techniques used for transforming the Ising models into the free-fermion models, specifically the decorated lattice technique [37, 83, 84], the star-triangle transformation [37, 83–86] and the weak-graph expansion [69, 87–89]. In Sec. III we present the exact results of the Ising models on the honeycomb, triangular and Kagomé lattices. Each of these models is shown to be equivalent with a free-fermion model, both in the zero field and in the imaginary field. The exact solutions are thereby obtained in the free-fermion formulation. Summary is outlined in Sec. IV.

II. METHODS AND TECHNIQUES

It is essential to introduce the methods and techniques for transformation of lattice systems and vertex weights, before we present the main results of the Ising models. The decorated lattice technique [37, 83, 84] and the star-triangle transformation [37, 83–86] help us to obtain a lattice structure, which is convenient for translating the Boltzmann factors to the vertex weights of a sixteen-vertex model. The resulting sixteen-vertex model is then transformed into an even or odd eight-vertex model by the weak-graph expansion [69, 87–89].

A. Decorated lattice technique

Decorated lattice technique is suitable for the case when we would like to add a spin site between a pair of nearest-neighbours, keeping the partition function invariant. As shown in Fig. 2, the interaction J is replaced by J_1 . In our consideration, the additional spin \tilde{s} is used to represent the direction of the arrow on the edge linking s_1 and s_2 . This may help us to build up an effective




FIG. 2. The decorated lattice technique.

vertex model formulation of the Ising problem. Consider the Boltzmann factors of this pair contributed to the partition function, which should be conserved after the decorated lattice transformation. That is

$$e^{-\beta J s_1 s_2} = A \sum_{\tilde{s}=\pm 1} e^{-\beta J_1 \tilde{s}(s_1 + s_2)}. \quad (2)$$

We can see clearly from Eq. (2) that

$$e^{-\beta J} = A (e^{-2\beta J_1} + e^{2\beta J_1}), \quad e^{\beta J} = 2A.$$

Thus the interaction J_1 and the multiplier A are determined

$$\cosh(2\beta J_1) = e^{-2\beta J}, \quad (3)$$

$$A = \frac{1}{2} e^{\beta J}. \quad (4)$$

B. Star-triangle transformation

Fig. 3 shows the star-triangle transformation. The triangle with internal interaction J is replaced by a “star” consisting of an additional spin site coupled to the three vertices with an interaction J' . The partition function should be invariant after the transformation, i.e., the Boltzmann factors of this unit should be conserved

$$e^{-\beta J (s_1 s_2 + s_2 s_3 + s_3 s_1)} = B \sum_{\tilde{s}=\pm 1} e^{-\beta J' \tilde{s}(s_1 + s_2 + s_3)}. \quad (5)$$

This leads to

$$e^{-3\beta J} = B (e^{-3\beta J'} + e^{3\beta J'}), \quad e^{\beta J} = B (e^{-\beta J'} + e^{\beta J'}).$$




FIG. 3. The star-triangle transformation.

The interaction J' and the multiplier B can be easily determined from the equations above

$$\frac{\cosh(3\beta J')}{\cosh(\beta J')} = 4\cosh^2(\beta J') - 3 = e^{-4\beta J}, \quad (6)$$

$$B = (e^{-4\beta J} + 3)^{-1/2} e^{\beta J}. \quad (7)$$

One can easily verify that the star-triangle transformation directly connects the partition function of the triangular lattice model with that of the honeycomb lattice model. This enables us to simplify the derivation of the results for the triangular lattice model.

C. Weak-graph expansion

Weak-graph expansion is a useful tool in lattice statistical problems [87, 89]. In this paper, we use the version of weak-graph expansion method proposed by Wu, which was originally designed for the even eight-vertex model [88] and later extended to the sixteen-vertex model [69]. Applying the method to the sixteen-vertex model enables us to obtain a new set of vertex weights, with the partition function invariant. The details of the method are shown in Ref. [88], and we only list the result for all sixteen rearranged weights

$$\begin{aligned} \tilde{\omega}_1 &= \frac{1}{4} (\omega_1 + \omega_2 + \omega_3 + \omega_4 + \omega_5 + \omega_6 + \omega_7 + \omega_8 + \omega_9 + \omega_{10} + \omega_{11} + \omega_{12} + \omega_{13} + \omega_{14} + \omega_{15} + \omega_{16}), \\ \tilde{\omega}_2 &= \frac{1}{4} (\omega_1 + \omega_2 + \omega_3 + \omega_4 + \omega_5 + \omega_6 + \omega_7 + \omega_8 - \omega_9 - \omega_{10} - \omega_{11} - \omega_{12} - \omega_{13} - \omega_{14} - \omega_{15} - \omega_{16}), \\ \tilde{\omega}_3 &= \frac{1}{4} (\omega_1 + \omega_2 + \omega_3 + \omega_4 - \omega_5 - \omega_6 - \omega_7 - \omega_8 - \omega_9 - \omega_{10} - \omega_{11} - \omega_{12} + \omega_{13} + \omega_{14} + \omega_{15} + \omega_{16}), \\ \tilde{\omega}_4 &= \frac{1}{4} (\omega_1 + \omega_2 + \omega_3 + \omega_4 - \omega_5 - \omega_6 - \omega_7 - \omega_8 + \omega_9 + \omega_{10} + \omega_{11} + \omega_{12} - \omega_{13} - \omega_{14} - \omega_{15} - \omega_{16}), \\ \tilde{\omega}_5 &= \frac{1}{4} (\omega_1 + \omega_2 - \omega_3 - \omega_4 + \omega_5 + \omega_6 - \omega_7 - \omega_8 - \omega_9 - \omega_{10} + \omega_{11} + \omega_{12} + \omega_{13} + \omega_{14} - \omega_{15} - \omega_{16}), \\ \tilde{\omega}_6 &= \frac{1}{4} (\omega_1 + \omega_2 - \omega_3 - \omega_4 + \omega_5 + \omega_6 - \omega_7 - \omega_8 + \omega_9 + \omega_{10} - \omega_{11} - \omega_{12} - \omega_{13} - \omega_{14} + \omega_{15} + \omega_{16}), \\ \tilde{\omega}_7 &= \frac{1}{4} (\omega_1 + \omega_2 - \omega_3 - \omega_4 - \omega_5 - \omega_6 + \omega_7 + \omega_8 - \omega_9 - \omega_{10} + \omega_{11} + \omega_{12} - \omega_{13} - \omega_{14} + \omega_{15} + \omega_{16}), \\ \tilde{\omega}_8 &= \frac{1}{4} (\omega_1 + \omega_2 - \omega_3 - \omega_4 - \omega_5 - \omega_6 + \omega_7 + \omega_8 + \omega_9 + \omega_{10} - \omega_{11} - \omega_{12} + \omega_{13} + \omega_{14} - \omega_{15} - \omega_{16}), \\ \tilde{\omega}_9 &= \frac{1}{4} (-\omega_1 + \omega_2 + \omega_3 - \omega_4 + \omega_5 - \omega_6 + \omega_7 - \omega_8 + \omega_9 - \omega_{10} + \omega_{11} - \omega_{12} - \omega_{13} + \omega_{14} + \omega_{15} - \omega_{16}), \\ \tilde{\omega}_{10} &= \frac{1}{4} (-\omega_1 + \omega_2 + \omega_3 - \omega_4 + \omega_5 - \omega_6 + \omega_7 - \omega_8 - \omega_9 + \omega_{10} - \omega_{11} + \omega_{12} + \omega_{13} - \omega_{14} - \omega_{15} + \omega_{16}), \\ \tilde{\omega}_{11} &= \frac{1}{4} (-\omega_1 + \omega_2 + \omega_3 - \omega_4 - \omega_5 + \omega_6 - \omega_7 + \omega_8 + \omega_9 - \omega_{10} + \omega_{11} - \omega_{12} + \omega_{13} - \omega_{14} - \omega_{15} + \omega_{16}), \\ \tilde{\omega}_{12} &= \frac{1}{4} (-\omega_1 + \omega_2 + \omega_3 - \omega_4 - \omega_5 + \omega_6 - \omega_7 + \omega_8 - \omega_9 + \omega_{10} - \omega_{11} + \omega_{12} - \omega_{13} + \omega_{14} + \omega_{15} - \omega_{16}), \\ \tilde{\omega}_{13} &= \frac{1}{4} (-\omega_1 + \omega_2 - \omega_3 + \omega_4 - \omega_5 + \omega_6 + \omega_7 - \omega_8 - \omega_9 + \omega_{10} + \omega_{11} - \omega_{12} + \omega_{13} - \omega_{14} + \omega_{15} - \omega_{16}), \\ \tilde{\omega}_{14} &= \frac{1}{4} (-\omega_1 + \omega_2 - \omega_3 + \omega_4 - \omega_5 + \omega_6 + \omega_7 - \omega_8 + \omega_9 - \omega_{10} - \omega_{11} + \omega_{12} - \omega_{13} + \omega_{14} - \omega_{15} + \omega_{16}), \\ \tilde{\omega}_{15} &= \frac{1}{4} (-\omega_1 + \omega_2 - \omega_3 + \omega_4 + \omega_5 - \omega_6 - \omega_7 + \omega_8 + \omega_9 - \omega_{10} - \omega_{11} + \omega_{12} + \omega_{13} - \omega_{14} + \omega_{15} - \omega_{16}), \\ \tilde{\omega}_{16} &= \frac{1}{4} (-\omega_1 + \omega_2 - \omega_3 + \omega_4 + \omega_5 - \omega_6 - \omega_7 + \omega_8 - \omega_9 + \omega_{10} + \omega_{11} - \omega_{12} - \omega_{13} + \omega_{14} - \omega_{15} + \omega_{16}). \end{aligned} \quad (8)$$

The sixteen-vertex model is symmetric, when the weight of each vertex configuration is invariant under the rever-

sal of arrows, i.e.

$$\omega_1 = \omega_2, \dots, \omega_{15} = \omega_{16}. \quad (9)$$

Similarly, the antisymmetric sixteen-vertex model is that satisfying

$$\omega_1 = -\omega_2, \dots, \omega_{15} = -\omega_{16}. \quad (10)$$

It is straightforward to see from Eq. (8) that, the symmetric sixteen-vertex model is transformed into an even eight-vertex model (Eq. (3) of Ref. [69] shows this case), while the antisymmetric case is transformed into an odd subcase.

We will show in the next section that each Ising model considered can be mapped to a symmetric/antisymmetric sixteen-vertex model, and further transformed into an even/odd eight-vertex model by Eq. (8). We can find that each case satisfies either the even free-fermion condition [41, 42]

$$\tilde{\omega}_1\tilde{\omega}_2 + \tilde{\omega}_3\tilde{\omega}_4 = \tilde{\omega}_5\tilde{\omega}_6 + \tilde{\omega}_7\tilde{\omega}_8 \quad (11)$$

or the odd free-fermion condition [43]

$$\tilde{\omega}_9\tilde{\omega}_{10} + \tilde{\omega}_{11}\tilde{\omega}_{12} = \tilde{\omega}_{13}\tilde{\omega}_{14} + \tilde{\omega}_{15}\tilde{\omega}_{16}. \quad (12)$$

III. RESULTS AND DISCUSSIONS

A. The honeycomb lattice

We apply the decorated lattice technique to the honeycomb lattice, as shown in Fig. 4(a). It is easy to verify that, the region bounded by dash lines can be regarded as a vertex site on the square lattice. Each vertex site consists of two honeycomb lattice Ising spins. Fig. 4(b) shows the details with J_1 determined by Eq. (3). It is natural to introduce a one-to-one mapping from the spin states of $\{\tilde{s}\}$ to the arrow configurations

$$-\cdot - + / - \mapsto \rightarrow / \leftarrow, \begin{array}{c} | \\ | \end{array} + / - \mapsto \uparrow / \downarrow. \quad (13)$$

Therefore, a sixteen-vertex model can be built up with an appropriate correspondence between the vertex weights and the Boltzmann factors of the Ising model.

1. Zero field

In the zero field, the partition function can be expressed by

$$Z = A^N \sum_{\{\tilde{s}\}=\pm 1} \omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4) \quad (14)$$

with

$$\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4) = \sum_{s_1=\pm 1, s_2=\pm 1} e^{-\beta J s_1 s_2} \times e^{-\beta J_1 [s_1(\tilde{s}_1+\tilde{s}_2)+s_2(\tilde{s}_3+\tilde{s}_4)]} \quad (15)$$

and A defined in Eq. (4). $\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4)$ is the Boltzmann factor of the vertex unit shown in Fig. 4(b).




FIG. 4. The diagrammatic representation of applying the decorated lattice technique to the honeycomb lattice model. The region bounded by dash lines in (a) forms a vertex site of the sixteen-vertex model. The details of the vertex unit are shown in (b).

Then the vertex weights can be determined directly from $\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4)$, according to the vertex configurations in Fig. 1 and the mapping in Eq. (13). That is

$$\begin{aligned} \omega_1 &= \omega(+, +, +, +) = 2e^{\beta J} + e^{-\beta J} (e^{-4\beta J_1} + e^{4\beta J_1}) \\ &= 2(e^{\beta J} - e^{-\beta J}) + 4e^{-5\beta J}, \\ \omega_2 &= \omega_1, \\ \omega_3 &= \omega(+, -, -, +) = 2(e^{\beta J} + e^{-\beta J}), \\ \omega_4 &= \omega_5 = \omega_6 = \omega_3, \\ \omega_7 &= \omega(-, -, +, +) = 2e^{-\beta J} + e^{\beta J} (e^{-4\beta J_1} + e^{4\beta J_1}) \\ &= 2(e^{-\beta J} - e^{\beta J}) + 4e^{-3\beta J}, \\ \omega_8 &= \omega_7, \\ \omega_9 &= \omega(+, -, +, +) = (e^{-\beta J} + e^{\beta J}) (e^{-2\beta J_1} + e^{2\beta J_1}) \\ &= 2(e^{-3\beta J} + e^{-\beta J}), \\ \omega_{10} &= \dots = \omega_{16} = \omega_9. \end{aligned} \quad (16)$$

We see that the resulting sixteen-vertex model is symmetric. As stated before, the vertex weights can be rearranged by performing the weak-graph expansion such that an even eight-vertex model is obtained. The new weights are straightforward to show from Eq. (8)

$$\begin{aligned} \tilde{\omega}_1 &= \frac{1}{2}\omega_1 + \omega_3 + \frac{1}{2}\omega_7 + 2\omega_9 = 2e^{\beta J} (e^{-2\beta J} + 1)^3, \\ \tilde{\omega}_2 &= \frac{1}{2}\omega_1 + \omega_3 + \frac{1}{2}\omega_7 - 2\omega_9 \\ &= 2e^{\beta J} (e^{-2\beta J} + 1) (e^{-2\beta J} - 1)^2, \\ \tilde{\omega}_3 &= \tilde{\omega}_4 = \tilde{\omega}_5 = \tilde{\omega}_6 = \tilde{\omega}_2, \\ \tilde{\omega}_7 &= \frac{1}{2}\omega_1 - \omega_3 + \frac{1}{2}\omega_7 = 2e^{\beta J} (e^{-2\beta J} + 1)^2 (e^{-2\beta J} - 1), \\ \tilde{\omega}_8 &= \tilde{\omega}_7. \end{aligned} \quad (17)$$

One can easily examine that the even free-fermion condition Eq. (11) is satisfied. Thus, the honeycomb lattice Ising model in the zero field is equivalent with an

even free-fermion model. Denote the number of vertex sites by N_v . The exact solution of the even free-fermion model $\lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{even}}$ had been given in Refs. [41, 42] (note that the ordering of vertices (5)–(8) in Fig. 1 differs slightly from that of Refs. [41, 42], but the expression for the solution is the same). We show the result quoting this well-known solution and noticing Eq. (14)

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z &= \ln A + \frac{1}{2} \lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{even}} \\ &= \ln \left(\frac{1}{2} e^{\beta J} \right) + \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln [a \\ &\quad + b \cos \theta + c \cos \phi + d \cos(\theta - \phi) + e(\theta + \phi)], \end{aligned} \quad (18)$$

with

$$\begin{aligned} a &= \tilde{\omega}_1^2 + \tilde{\omega}_2^2 + \tilde{\omega}_3^2 + \tilde{\omega}_4^2, \\ b &= 2(\tilde{\omega}_1 \tilde{\omega}_3 - \tilde{\omega}_2 \tilde{\omega}_4), \\ c &= 2(\tilde{\omega}_1 \tilde{\omega}_4 - \tilde{\omega}_2 \tilde{\omega}_3), \\ d &= 2(\tilde{\omega}_3 \tilde{\omega}_4 - \tilde{\omega}_7 \tilde{\omega}_8), \\ e &= 2(\tilde{\omega}_3 \tilde{\omega}_4 - \tilde{\omega}_5 \tilde{\omega}_6). \end{aligned} \quad (19)$$

A straightforward calculation of the coefficients in Eq. (19) from the vertex weights in Eq. (17) gives

$$\begin{aligned} a &= 2^7 e^{-4\beta J} [\cosh^3(2\beta J) + 1], \\ b &= c = 2^7 e^{-4\beta J} \sinh^2(2\beta J), \\ d &= -2^7 e^{-4\beta J} \sinh^2(2\beta J), \\ e &= 0. \end{aligned} \quad (20)$$

Inserting Eq. (20) into Eq. (18) and making some arrangements, the exact solution turns out to be

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z &= \frac{3}{4} \ln 2 + \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \\ &\quad \ln \{ \cosh^3(2\beta J) + 1 \\ &\quad + \sinh^2(2\beta J) [\cos \theta + \cos \phi + \cos(\theta - \phi)] \}. \end{aligned} \quad (21)$$

It can be easily verified that, this expression is consistent with the results published in the previous literatures [28–31].

2. Imaginary field

In the case of $H_{\text{ex}} = i(\pi/2)k_B T$, we notice that

$$e^{\beta H_{\text{ex}} \sum_{i=1}^N s_i} = i^{\sum_{i=1}^N s_i} = i^N \prod_{i=1}^N s_i \quad (22)$$

by using the identity $i^{s_i} = i \times s_i$ [22]. We can see that, i^N is a constant and the product $\prod_{i=1}^N s_i$ can be conveniently

included in the factors $\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4)$. We just need to rewrite the partition function as

$$Z = i^N A^N \sum_{\{\tilde{s}\}=\pm 1} \omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4) \quad (23)$$

with

$$\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4) = \sum_{s_1=\pm 1, s_2=\pm 1} (s_1 s_2) e^{-\beta J s_1 s_2} \times e^{-\beta J_1 [s_1(\tilde{s}_1 + \tilde{s}_2) + s_2(\tilde{s}_3 + \tilde{s}_4)]}. \quad (24)$$

The vertex weights are then determined in the similar way to those in Eq. (16)

$$\begin{aligned} \omega_1 &= \omega_2 = -2(e^{\beta J} + e^{-\beta J}) + 4e^{-5\beta J}, \\ \omega_3 &= \omega_4 = \omega_5 = \omega_6 = 2(e^{-\beta J} - e^{\beta J}), \\ \omega_7 &= \omega_8 = 2(e^{\beta J} + e^{-\beta J}) - 4e^{-3\beta J}, \\ \omega_9 &= \dots = \omega_{16} = 2(e^{-3\beta J} - e^{-\beta J}). \end{aligned} \quad (25)$$

Again, the Boltzmann factors are translated into the weights of a symmetric sixteen-vertex model. We still perform the weak-graph expansion via Eq. (8)

$$\begin{aligned} \tilde{\omega}_1 &= -16e^{-2\beta J} \cosh^2(\beta J) \sinh(\beta J), \\ \tilde{\omega}_2 &= -16e^{-2\beta J} \sinh^3(\beta J), \\ \tilde{\omega}_3 &= \tilde{\omega}_4 = \tilde{\omega}_5 = \tilde{\omega}_6 = \tilde{\omega}_1, \\ \tilde{\omega}_7 &= \tilde{\omega}_8 = 16e^{-2\beta J} \cosh(\beta J) \sinh^2(\beta J). \end{aligned} \quad (26)$$

Clearly, we have obtained an even free-fermion model again. Notice Eq. (23) and the partition function is

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z = i \frac{\pi}{2} + \ln A + \frac{1}{2} \lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{even}} \quad (27)$$

with the even free-fermion solution expressed in Eqs. (18) and (19). Omitting the detailed calculations, we show the final exact result

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z &= i \frac{\pi}{2} + \frac{3}{4} \ln 2 + \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \\ &\quad \ln \{ \cosh^3(2\beta J) - 1 \\ &\quad + \sinh^2(2\beta J) [\cos \theta + \cos \phi + \cos(\theta - \phi)] \}. \end{aligned} \quad (28)$$

One can examine that our result recovers the proposed solutions of this case [16, 22].

B. The triangular lattice

To deal with the triangular lattice Ising model, we employ the star-triangle transformation as shown in Fig. 5(a). The transformed spin system is on the honeycomb lattice, with the replacement of the nearest-neighbour interaction $J \rightarrow J'$. J' is determined by Eq. (6). Applying the decorated lattice technique in the same way as that in Fig. 4(a), we construct a sixteen-vertex model of which the vertex site is shown Fig. 5(b). The mapping from the spin states to the arrow configurations is still that in Eq. (13). The relation between J' and J_1 is the same with that in Eq. (3).




FIG. 5. The diagrammatic representation of applying the star-triangle transformation to the triangular lattice model. The transformed system, which is on the honeycomb lattice as shown by the dash lines in (a), can be mapped into the sixteen-vertex model in the same way as that in Fig. 4(a). The details of the vertex unit are shown in (b).

1. Zero field

In the zero field, the partition function is simply

$$Z = B^N Z(\text{honeycomb}, J \rightarrow J', H_{\text{ex}} = 0)$$

with B defined in Eq. (7). Obviously this case is also equivalent with an even free-fermion model. Since the number of spins on the transformed honeycomb lattice is double that on the triangular lattice, the partition function should be

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z = \ln B + 2 \times \text{Eq. (21)} (J \rightarrow J'). \quad (29)$$

Taking into account Eqs. (6), (7) and (21), the exact solution can be obtained

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z = \frac{1}{8\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln \{ e^{-6\beta J} + 3e^{2\beta J} - 4 \sinh(2\beta J) [\cos \theta + \cos \phi + \cos(\theta - \phi)] \}. \quad (30)$$

This expression agrees with the known results of this case [32–35].

2. Imaginary field

We still make use of Eq. (22) to deal with the effect of the imaginary field. The partition function of this case should be

$$Z = i^N B^N A'^{2N} \sum_{\{\tilde{s}\}=\pm 1} \omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4), \quad (31)$$

with $A' = \frac{1}{2} e^{\beta J'}$ and

$$\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4) = \sum_{s_1=\pm 1, s_2=\pm 1} (s_1) e^{-\beta J' s_1 s_2} \times e^{-\beta J'_1 [s_1(\tilde{s}_1 + \tilde{s}_2) + s_2(\tilde{s}_3 + \tilde{s}_4)]}. \quad (32)$$

Note that the factors $\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4)$ differ from that in Eq. (24) as the spin s_2 is not on the original triangular lattice. Similarly, the factors $\omega(\tilde{s}_1, \tilde{s}_2, \tilde{s}_3, \tilde{s}_4)$ are translated into the vertex weights of a sixteen-vertex model. We find that the vertex weights in this case are antisymmetric

$$\begin{aligned} \omega_1 &= 4e^{-3\beta J'} (e^{-4\beta J'} - 1)^{1/2}, \\ \omega_2 &= -\omega_1, \\ \omega_3 &= \omega_4 = \omega_5 = \omega_6 = 0, \\ \omega_7 &= -4e^{-\beta J'} (e^{-4\beta J'} - 1)^{1/2}, \\ \omega_8 &= -\omega_7, \\ \omega_9 &= \omega_{13} = 2 (e^{-\beta J'} - e^{\beta J'}) (e^{-4\beta J'} - 1)^{1/2}, \\ \omega_{10} &= \omega_{14} = -\omega_9, \\ \omega_{11} &= \omega_{15} = -2 (e^{-\beta J'} + e^{\beta J'}) (e^{-4\beta J'} - 1)^{1/2}, \\ \omega_{12} &= \omega_{16} = -\omega_{11}. \end{aligned} \quad (33)$$

From these weights, the weak-graph expansion produces an odd eight-vertex model by Eq. (8)

$$\begin{aligned} \tilde{\omega}_9 &= -2e^{\beta J'} (e^{-2\beta J'} + 1)^2 (e^{-4\beta J'} - 1)^{1/2}, \\ \tilde{\omega}_{10} &= -2e^{\beta J'} (e^{-4\beta J'} - 1)^{3/2}, \\ \tilde{\omega}_{11} &= -2e^{\beta J'} (e^{-2\beta J'} - 1)^2 (e^{-4\beta J'} - 1)^{1/2}, \\ \tilde{\omega}_{12} &= \tilde{\omega}_{14} = \tilde{\omega}_{16} = \tilde{\omega}_{10}, \\ \tilde{\omega}_{13} &= \tilde{\omega}_9, \\ \tilde{\omega}_{15} &= \tilde{\omega}_{11}. \end{aligned} \quad (34)$$

It is obvious that the odd free-fermion condition Eq. (12) is satisfied. Therefore, the triangular lattice Ising model in the imaginary field is equivalent with an odd free-fermion model. Now the result is

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z = i \frac{\pi}{2} + \ln B + 2 \ln A' + \lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{odd}}, \quad (35)$$

where the exact solution of the odd free-fermion model is given by Eq. (21) of Ref. [43]

$$\lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{odd}} = \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln [a + b \cos \theta + c \cos \phi + d \cos(\theta - \phi) + e(\theta + \phi)] \quad (36)$$

with

$$\begin{aligned}
a &= 2 \left[(\tilde{\omega}_9 \tilde{\omega}_{10} + \tilde{\omega}_{11} \tilde{\omega}_{12})^2 + (\tilde{\omega}_9 \tilde{\omega}_{12})^2 + (\tilde{\omega}_{10} \tilde{\omega}_{11})^2 \right. \\
&\quad \left. + (\tilde{\omega}_{13} \tilde{\omega}_{16})^2 + (\tilde{\omega}_{14} \tilde{\omega}_{15})^2 \right], \\
b &= 2 \left[-(\tilde{\omega}_9 \tilde{\omega}_{12})^2 - (\tilde{\omega}_{10} \tilde{\omega}_{11})^2 + 2\tilde{\omega}_{13} \tilde{\omega}_{14} \tilde{\omega}_{15} \tilde{\omega}_{16} \right], \\
c &= 2 \left[(\tilde{\omega}_{13} \tilde{\omega}_{16})^2 + (\tilde{\omega}_{14} \tilde{\omega}_{15})^2 - 2\tilde{\omega}_9 \tilde{\omega}_{10} \tilde{\omega}_{11} \tilde{\omega}_{12} \right], \\
d &= -2(\tilde{\omega}_9 \tilde{\omega}_{10} - \tilde{\omega}_{15} \tilde{\omega}_{16})^2, \\
e &= -2(\tilde{\omega}_{11} \tilde{\omega}_{12} - \tilde{\omega}_{15} \tilde{\omega}_{16})^2.
\end{aligned} \tag{37}$$

We should remark that the relation between our notation of weights and that of Ref. [43] is

$$\begin{aligned}
\tilde{\omega}_9 &= u_1, \quad \tilde{\omega}_{10} = u_2, \quad \tilde{\omega}_{11} = u_4, \quad \tilde{\omega}_{12} = u_3, \\
\tilde{\omega}_{13} &= u_7, \quad \tilde{\omega}_{14} = u_8, \quad \tilde{\omega}_{15} = u_6, \quad \tilde{\omega}_{16} = u_5.
\end{aligned}$$

Substituting $A' = \frac{1}{2}e^{\beta J'}$ and Eqs. (6), (7) and (34) into Eq. (35), we give the final expression

$$\begin{aligned}
\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z &= i\frac{\pi}{2} + \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln \left\{ e^{-12\beta J} - 4 \right. \\
&\quad \left. + 3e^{4\beta J} - 8 \sinh^2(2\beta J) [\cos \theta - \cos \phi + \cos(\theta - \phi)] \right\}.
\end{aligned} \tag{38}$$

Then we have rederived the exact solution of this case [16, 22], and indicated that the model in this case can be seen as an odd free-fermion model.

It is easy to find that this frustrated system (when $J > 0$) in the imaginary field has a finite residual entropy in the zero temperature limit

$$\begin{aligned}
S &= \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln \left\{ 3 - 2 [\cos \theta - \cos \phi \right. \\
&\quad \left. + \cos(\theta - \phi)] \right\} \\
&= 0.161533.
\end{aligned} \tag{39}$$

Here we omit the Boltzmann constant k_B . Interestingly, the residual entropy in the imaginary field is half that in the zero field [32] (we can also obtain the residual entropy in the zero field by Eq. (30), and compare the value with Eq. (39)).

C. The Kagomé lattice

Fig. 6 shows the Kagomé lattice. The region bounded by dash lines, which consists of two triangles, is chosen as the vertex site on the square lattice. The states of four spins (s_1, s_2, s_3, s_4) around each site can be mapped to the arrow configurations around this vertex by Eq. (13), thus the sixteen-vertex model is conveniently constructed.




FIG. 6. The diagrammatic representation of the Kagomé lattice. The region bounded by dash lines forms a vertex site of the sixteen-vertex model.

1. Zero field

The partition function is in the form of

$$Z = \sum_{\{(s_1, s_2, s_3, s_4)\} = \pm 1} \omega(s_1, s_2, s_3, s_4) \tag{40}$$

with

$$\omega(s_1, s_2, s_3, s_4) = \sum_{s_5 = \pm 1} e^{-\beta J [s_5(s_1 + s_2 + s_3 + s_4) + s_1 s_2 + s_3 s_4]}. \tag{41}$$

The translated vertex weights from the mapping in Eq. (13) are

$$\begin{aligned}
\omega_1 &= \omega_2 = 2e^{-2\beta J} \cosh(4\beta J), \\
\omega_3 &= \omega_4 = \omega_5 = \omega_6 = 2e^{2\beta J}, \\
\omega_7 &= \omega_8 = 2e^{-2\beta J}, \\
\omega_9 &= \dots = \omega_{16} = 2 \cosh(2\beta J).
\end{aligned} \tag{42}$$

Again, this symmetric sixteen-vertex model are transformed into an even eight-vertex model by Eq. (8) with the weights

$$\begin{aligned}
\tilde{\omega}_1 &= \frac{1}{2} (e^{-3\beta J} + 3e^{\beta J})^2, \\
\tilde{\omega}_2 &= \tilde{\omega}_3 = \tilde{\omega}_4 = \tilde{\omega}_5 = \tilde{\omega}_6 = \frac{1}{2} (e^{-3\beta J} - e^{\beta J})^2, \\
\tilde{\omega}_7 &= \tilde{\omega}_8 = \frac{1}{2} (e^{-3\beta J} + 3e^{\beta J}) (e^{-3\beta J} - e^{\beta J}).
\end{aligned} \tag{43}$$

One can see immediately that this case is an even free-fermion model. Making use of the even free-fermion solution as expressed in Eqs. (18) and (19), we give the exact

result

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z \\
&= \frac{1}{3} \lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{even}} \\
&= \frac{1}{24\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln \{ e^{-12\beta J} + 18e^{-4\beta J} + 24 \\
&\quad + 21e^{4\beta J} + 4(e^{-8\beta J} - e^{-4\beta J} - 1 + e^{4\beta J}) [\cos \theta \\
&\quad\quad + \cos \phi - \cos(\theta - \phi)] \} \quad (44)
\end{aligned}$$

This result agrees with the known expressions (see Eq. (37) of Ref. [38] for instance).

2. Imaginary field

Taking into account Eq. (22), the partition function in the presence of the imaginary field is

$$Z = i^N \sum_{\{(s_1, s_2, s_3, s_4)\} = \pm 1} \omega(s_1, s_2, s_3, s_4) \quad (45)$$

with

$$\begin{aligned}
\omega(s_1, s_2, s_3, s_4) = \sum_{s_5 = \pm 1} (s_1 s_2 s_5) e^{-\beta J s_5 (s_1 + s_2 + s_3 + s_4)} \times \\
e^{-\beta J (s_1 s_2 + s_3 s_4)}. \quad (46)
\end{aligned}$$

We find that the resulting sixteen-vertex model is anti-symmetric, like in the case of triangular lattice model in the imaginary field. The vertex weights are listed

$$\begin{aligned}
\omega_1 &= e^{-6\beta J} - e^{2\beta J}, \\
\omega_2 &= -\omega_1, \\
\omega_3 &= \omega_4 = \omega_5 = \omega_6 = \omega_7 = \omega_8 = 0, \\
\omega_9 &= \omega_{11} = \omega_{13} = \omega_{15} = e^{2\beta J} - e^{-2\beta J}, \\
\omega_{10} &= \omega_{12} = \omega_{14} = \omega_{16} = -\omega_9. \quad (47)
\end{aligned}$$

Then the transformed weights of the odd eight-vertex model are

$$\begin{aligned}
\tilde{\omega}_9 &= \frac{1}{2} e^{2\beta J} (3 + e^{-4\beta J}) (1 - e^{-4\beta J}), \\
\tilde{\omega}_{10} &= -\frac{1}{2} e^{2\beta J} (1 - e^{-4\beta J})^2, \\
\tilde{\omega}_{11} &= \tilde{\omega}_{13} = \tilde{\omega}_{15} = \tilde{\omega}_9, \\
\tilde{\omega}_{12} &= \tilde{\omega}_{14} = \tilde{\omega}_{16} = \tilde{\omega}_{10}. \quad (48)
\end{aligned}$$

It is trivial to see that this case is equivalent with an odd free-fermion model, the exact result is thereby given by the odd free-fermion solution in Eqs. (36) and (37)

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z \\
&= i \frac{\pi}{2} + \frac{1}{3} \lim_{N_v \rightarrow \infty} \frac{1}{N_v} \ln Z_{\text{odd}} \\
&= i \frac{\pi}{2} + \frac{1}{6} \ln \left[(1 + 3e^{4\beta J}) |1 - e^{-4\beta J}|^3 \right]. \quad (49)
\end{aligned}$$

The expression is surprisingly simple. To our knowledge, this result has not been reported previously. There have been research works studying the Kagomé lattice Ising model in a magnetic field by transformation into the honeycomb lattice model [90–92]. One can examine that, the case in the imaginary field was not included in these works. Our analysis demonstrates that this case is also exactly solvable.

As is well-known, most of the exact solutions of two-dimensional Ising models are of the form of a double integral. Onsager’s solution of the square lattice model [5] is a very early example. Each model studied in this paper using the free-fermion approach involves a double integral in the solution, except for the Kagomé lattice model in the imaginary field. It is interesting to compare the result Eq. (49) with Eq. (1) of Ref. [93], which is the solution of the dimer model on the Kagomé lattice and can also be derived by mapping to the odd free-fermion model. Both formulas take a very simply logarithmic form. Ref. [93] stated that the novel and unique expression “points to the special role played by the Kagomé lattice”. Here we provide new evidence for this statement.

When $J > 0$, the Kagomé lattice Ising model is geometrically frustrated. The residual entropy in the imaginary field is easily calculated from Eq. (49) by taking the zero temperature limit

$$S = \frac{1}{6} \ln 3 = 0.183102. \quad (50)$$

IV. SUMMARY

The exact solutions of Ising models on typical two-dimensional lattices, specifically the honeycomb, triangular and Kagomé lattices, are studied in the free-fermion formulation. For each Ising model, both the case of the zero field and the case of an imaginary field $i(\pi/2)k_B T$ are considered. Five known solutions are rederived and one new solution is found. In particular, the Kagomé lattice model in the imaginary field is shown to be exactly solvable, exhibiting a very simple form of solution. Hence, we have introduced a new member to the family of exactly solvable lattice models.

Each solution in this paper is achieved by mapping the Ising spin system into a free-fermion model. The mapping procedure is simple and easy to extend to other lattice systems. Applying the mapping method to the Ising model in a physical (real) and non-zero magnetic field would be interesting and likely very fruitful. We expect that our work provides new insights into the mathematical exact results of spin systems in a magnetic field. Furthermore, the free-fermion formulation may be suitable for other statistical lattice problems, such as the dimer covering problem. Study of the two-dimensional dimer models using the free-fermion approach is currently in progress.

ACKNOWLEDGMENTS

This work was supported by Guangdong Provincial Quantum Science Strategic Initiative (Grant No.

GDZX2200001), Guangdong Basic and Applied Basic Research Foundation (Grant No. 2021A1515010328) and National Natural Science Foundation of China (Grant No. 11874312).

-
- [1] M. Niss, History of the Lenz-Ising Model 1920–1950: From Ferromagnetic to Cooperative Phenomena, *Arch. Hist. Exact Sci.* **59**, 267 (2005).
- [2] E. Ising, Beitrag zur Theorie des Ferromagnetismus, *Z. Phys.* **31**, 253 (1925).
- [3] H. A. Kramers and G. H. Wannier, Statistics of the Two-Dimensional Ferromagnet. Part I, *Phys. Rev.* **60**, 252 (1941).
- [4] H. A. Kramers and G. H. Wannier, Statistics of the Two-Dimensional Ferromagnet. Part II, *Phys. Rev.* **60**, 263 (1941).
- [5] L. Onsager, Crystal Statistics. I. A Two-Dimensional Model with an Order-Disorder Transition, *Phys. Rev.* **65**, 117 (1944).
- [6] B. Kaufman, Crystal Statistics. II. Partition Function Evaluated by Spinor Analysis, *Phys. Rev.* **76**, 1232 (1949).
- [7] Y. Nambu, A Note on the Eigenvalue Problem in Crystal Statistics, *Prog. Theor. Phys.* **5**, 1 (1950).
- [8] M. Kac and J. C. Ward, A Combinatorial Solution of the Two-Dimensional Ising Model, *Phys. Rev.* **88**, 1332 (1952).
- [9] T. D. Schultz, D. C. Mattis, and E. H. Lieb, Two-Dimensional Ising Model as a Soluble Problem of Many Fermions, *Rev. Mod. Phys.* **36**, 856 (1964).
- [10] C. A. Hurst and H. S. Green, New Solution of the Ising Problem for a Rectangular Lattice, *J. Chem. Phys.* **33**, 1059 (1960).
- [11] C. A. Hurst, New Approach to the Ising Problem, *J. Math. Phys.* **7**, 305 (1966).
- [12] R. W. Gibberd and C. A. Hurst, New Approach to the Ising Model. II, *J. Math. Phys.* **8**, 1427 (1967).
- [13] M. E. Fisher, On the Dimer Solution of Planar Ising Models, *J. Math. Phys.* **7**, 1776 (1966).
- [14] S. Samuel, The use of anticommuting variable integrals in statistical mechanics. I. The computation of partition functions, *J. Math. Phys.* **21**, 2806 (1980).
- [15] T. D. Lee and C. N. Yang, Statistical Theory of Equations of State and Phase Transitions. II. Lattice Gas and Ising Model, *Phys. Rev.* **87**, 410 (1952).
- [16] G. Baxter, Weight Factors for the Two-Dimensional Ising Model, *J. Math. Phys.* **6**, 1015 (1965).
- [17] G. Baxter, Local Weights Which Determine Area, and the Ising Model, *J. Math. Phys.* **8**, 399 (1967).
- [18] B. M. McCoy and T. T. Wu, Theory of Toeplitz Determinants and the Spin Correlations of the Two-Dimensional Ising Model. II, *Phys. Rev.* **155**, 438 (1967).
- [19] T. W. Marshall, The Ising model in a magnetic field of $\frac{1}{2}\pi$, *Mol. Phys.* **21**, 847 (1971).
- [20] D. Merlini, On the solution of the two-dimensional Ising model with an imaginary magnetic field $\beta H = h = i\pi/2$, *Lett. Nuovo Cimento* **9**, 100 (1974).
- [21] A. Gaaff, The 2-d Ising model in a magnetic field $i\pi/2$ as a soluble case of the sixteen vertex model, *Phys. Lett. A* **49**, 103 (1974).
- [22] K. Y. Lin and F. Y. Wu, ISING MODEL IN THE MAGNETIC FIELD $i\pi kT/2$, *Int. J. Mod. Phys. B* **02**, 471 (1988).
- [23] G. F. Newell and E. W. Montroll, On the Theory of the Ising Model of Ferromagnetism, *Rev. Mod. Phys.* **25**, 353 (1953).
- [24] R. J. Baxter, *Exactly solved models in statistical mechanics* (Academic Press, London, 1982).
- [25] R. J. Baxter, Solvable models in statistical mechanics, from Onsager onward, *J. Stat. Phys.* **78**, 7 (1995).
- [26] E. H. Lieb, Some of the Early History of Exactly Soluble Models, *Int. J. Mod. Phys. B* **11**, 3 (1997).
- [27] R. J. Baxter, Some comments on developments in exact solutions in statistical mechanics since 1944, *J. Stat. Mech.* **2010**, P11037 (2010).
- [28] K. Husimi and I. Syôzi, The Statistics of Honeycomb and Triangular Lattice. I, *Prog. Theor. Phys.* **5**, 177 (1950).
- [29] I. Syôzi, The Statistics of Honeycomb and Triangular Lattice. II, *Prog. Theor. Phys.* **5**, 341 (1950).
- [30] R. M. F. Houtappel, Order-disorder in hexagonal lattices, *Physica* **16**, 425 (1950).
- [31] H. N. V. Temperley, Statistical mechanics of the two-dimensional assembly, *Proc. R. Soc. Lond. A* **202**, 202 (1950).
- [32] G. H. Wannier, Antiferromagnetism. The Triangular Ising Net, *Phys. Rev.* **79**, 357 (1950).
- [33] G. H. Wannier, Erratum - Antiferromagnetism. The Triangular Ising Net, *Phys. Rev. B* **7**, 5017 (1973).
- [34] G. F. Newell, Crystal Statistics of a Two-Dimensional Triangular Ising Lattice, *Phys. Rev.* **79**, 876 (1950).
- [35] R. B. Potts, Combinatorial Solution of the Triangular Ising Lattice, *Proc. Phys. Soc. A* **68**, 145 (1955).
- [36] A. Nourhani, V. H. Crespi, and P. E. Lammert, Classical triangular lattice antiferromagnetic Ising model as a free-fermion/superconductor system, *J. Phys. A: Math. Theor.* **56**, 205001 (2023).
- [37] I. Syôzi, Statistics of Kagomé Lattice, *Prog. Theor. Phys.* **6**, 306 (1951).
- [38] K. Kanô and S. Naya, Antiferromagnetism. The Kagomé Ising Net, *Prog. Theor. Phys.* **10**, 158 (1953).
- [39] H. J. Giacomini, Exact results for a checkerboard Ising model with crossing and four-spin interactions, *J. Phys. A: Math. Gen.* **18**, L1087 (1985).
- [40] F. Y. Wu, Two-dimensional Ising model with crossing and four-spin interactions and a magnetic field $i(\pi/2)kT$, *J. Stat. Phys.* **44**, 455 (1986).
- [41] C. Fan and F. Y. Wu, Ising Model with Second-Neighbor Interaction. I. Some Exact Results and an Approximate Solution, *Phys. Rev.* **179**, 560 (1969).
- [42] C. Fan and F. Y. Wu, General Lattice Model of Phase Transitions, *Phys. Rev. B* **2**, 723 (1970).
- [43] F. Y. Wu and H. Kunz, The Odd Eight-Vertex Model, *J. Stat. Phys.* **116**, 67 (2004).
- [44] R. J. Baxter, Free-fermion, checkerboard and Z-invariant lattice models in statistical mechanics, *Proc. R. Soc.*

- Lond. A **404**, 1 (1986).
- [45] P. Melotti, The Free-Fermion Eight-Vertex Model: Couplings, Bipartite Dimers and Z-Invariance, *Commun. Math. Phys.* **381**, 33 (2021).
- [46] J. C. Slater, Theory of the Transition in KH_2PO_4 , *J. Chem. Phys.* **9**, 16 (1941).
- [47] F. Rys, Über ein zweidimensionales klassisches Konfigurationsmodell, *Helv. Phys. Acta* **36**, 537 (1963).
- [48] E. H. Lieb, Residual Entropy of Square Ice, *Phys. Rev.* **162**, 162 (1967).
- [49] E. H. Lieb, Exact Solution of the Problem of the Entropy of Two-Dimensional Ice, *Phys. Rev. Lett.* **18**, 692 (1967).
- [50] F. Y. Wu, Exactly Soluble Model of the Ferroelectric Phase Transition in Two Dimensions, *Phys. Rev. Lett.* **18**, 605 (1967).
- [51] E. H. Lieb, Exact Solution of the F Model of An Antiferroelectric, *Phys. Rev. Lett.* **18**, 1046 (1967).
- [52] E. H. Lieb, Exact Solution of the Two-Dimensional Slater KDP Model of a Ferroelectric, *Phys. Rev. Lett.* **19**, 108 (1967).
- [53] B. Sutherland, Exact Solution of a Two-Dimensional Model for Hydrogen-Bonded Crystals, *Phys. Rev. Lett.* **19**, 103 (1967).
- [54] C. P. Yang, Exact Solution of a Model of Two-Dimensional Ferroelectrics in an Arbitrary External Electric Field, *Phys. Rev. Lett.* **19**, 586 (1967).
- [55] B. Sutherland, C. N. Yang, and C. P. Yang, Exact Solution of a Model of Two-Dimensional Ferroelectrics in an Arbitrary External Electric Field, *Phys. Rev. Lett.* **19**, 588 (1967).
- [56] F. Y. Wu, Remarks on the Modified Potassium Dihydrogen Phosphate Model of a Ferroelectric, *Phys. Rev.* **168**, 539 (1968).
- [57] M. L. Glasser, Evaluation of the Partition Functions for Some Two-Dimensional Ferroelectric Models, *Phys. Rev.* **184**, 539 (1969).
- [58] B. Sutherland, Two-Dimensional Hydrogen Bonded Crystals without the Ice Rule, *J. Math. Phys.* **11**, 3183 (1970).
- [59] R. J. Baxter, Eight-Vertex Model in Lattice Statistics, *Phys. Rev. Lett.* **26**, 832 (1971).
- [60] R. J. Baxter, Partition function of the Eight-Vertex lattice model, *Ann. Phys.* **70**, 193 (1972).
- [61] L. P. Kadanoff and F. J. Wegner, Some Critical Properties of the Eight-Vertex Model, *Phys. Rev. B* **4**, 3989 (1971).
- [62] R. Baxter, Eight-vertex model in lattice statistics and one-dimensional anisotropic heisenberg chain. I. Some fundamental eigenvectors, *Ann. Phys.* **76**, 1 (1973).
- [63] R. Baxter, Eight-vertex model in lattice statistics and one-dimensional anisotropic heisenberg chain. II. Equivalence to a generalized ice-type lattice model, *Ann. Phys.* **76**, 25 (1973).
- [64] R. Baxter, Eight-vertex model in lattice statistics and one-dimensional anisotropic heisenberg chain. III. Eigenvectors of the transfer matrix and hamiltonian, *Ann. Phys.* **76**, 48 (1973).
- [65] R. J. Baxter, Solvable eight-vertex model on an arbitrary planar lattice, *Philos. Trans. R. Soc. London, Ser. A* **289**, 315 (1978).
- [66] R. J. Baxter, The Six and Eight-Vertex Models Revisited, *J. Stat. Phys.* **116**, 43 (2004).
- [67] F. Y. Wu, Critical Behavior of Two-Dimensional Hydrogen-Bonded Antiferroelectrics, *Phys. Rev. Lett.* **22**, 1174 (1969).
- [68] F. Y. Wu, Critical Behavior of Hydrogen-Bonded Ferroelectrics, *Phys. Rev. Lett.* **24**, 1476 (1970).
- [69] F. Y. Wu, Exact results on a general lattice statistical model, *Solid State Commun.* **10**, 115 (1972).
- [70] F. Y. Wu, Phase Transition in a Sixteen-Vertex Lattice Model, *Phys. Rev. B* **6**, 1810 (1972).
- [71] A. Gaaff and J. Hijmans, Symmetry relations in the sixteen-vertex model, *Physica A* **80**, 149 (1975).
- [72] M. Assis, The 16-vertex model and its even and odd 8-vertex subcases on the square lattice, *J. Phys. A: Math. Theor.* **50**, 395001 (2017).
- [73] B. U. Felderhof, Direct diagonalization of the transfer matrix of the zero-field free-fermion model, *Physica* **65**, 421 (1973).
- [74] B. U. Felderhof, Diagonalization of the transfer matrix of the free-fermion model. II, *Physica* **66**, 279 (1973).
- [75] B. U. Felderhof, Diagonalization of the transfer matrix of the free-fermion model. III, *Physica* **66**, 509 (1973).
- [76] V. V. Bazhanov and Y. G. Stroganov, Hidden symmetry of free Fermion model, *Theor. Math. Phys.* **62**, 253 (1985).
- [77] V. V. Bazhanov and Y. G. Stroganov, Hidden symmetry of free fermion model. II. Partition function, *Theor. Math. Phys.* **63**, 519 (1985).
- [78] V. V. Bazhanov and Y. G. Stroganov, Hidden symmetry of free fermion model. III. Inversion relations, *Theor. Math. Phys.* **63**, 604 (1985).
- [79] M. Debauche, H. T. Diep, P. Azaria, and H. Giacomini, Exact phase diagram of a generalized Kagomé Ising lattice: Reentrance and disorder lines, *Phys. Rev. B* **44**, 2369 (1991).
- [80] A. Kooiman and J. M. J. van Leeuwen, Free fermion approximation for the Ising model with further-neighbor interactions on a triangular lattice, *J. Stat. Phys.* **69**, 247 (1992).
- [81] T. Morita and K. Tanaka, Diagrammatical Techniques for Two-Dimensional Ising Models. III. Ising Model to Vertex Model, *J. Phys. Soc. Jpn.* **62**, 873 (1993).
- [82] R. Krčmár, A. Gendiar, and L. Šamaj, Ising ferromagnets and antiferromagnets in an imaginary magnetic field, *Phys. Rev. E* **105**, 054112 (2022).
- [83] M. E. Fisher, Transformations of Ising Models, *Phys. Rev.* **113**, 969 (1959).
- [84] I. Syözi, Transformation of Ising Models, in *Phase Transitions and Critical Phenomena. Vol. 1: Exact results* (Academic Press, London, 1972).
- [85] L. Onsager, The Ising model in two dimensions, in *Critical phenomena in alloys, magnets and superconductors* (McGraw-Hill, New York, 1971).
- [86] H. Au-Yang and J. H. H. Perk, Onsager's Star-Triangle Equation: Master Key to Integrability, in *Integrable Sys Quantum Field Theory* (Academic Press, San Diego, 1989).
- [87] J. F. Nagle, Weak-Graph Method for Obtaining Formal Series Expansions for Lattice Statistical Problems, *J. Math. Phys.* **9**, 1007 (1968).
- [88] F. Y. Wu, Exact Solution of a Model of an Antiferroelectric Transition, *Phys. Rev.* **183**, 604 (1969).
- [89] J. F. Nagle, Ferroelectric Models, in *Phase Transitions and Critical Phenomena. Vol. 3: Series Expansions for Lattice Models* (Academic Press, London, 1974).
- [90] H. Giacomini, An exact result for the Kagome lattice Ising model with magnetic field, *J. Phys. A: Math. Gen.*

- [21, L31 \(1988\)](#).
- [91] H. J. Giacomini, An exact functional relation for the partition function of a 2D Ising model with magnetic field, *J. Phys. A: Math. Gen.* **21**, L599 (1988).
- [92] W. T. Lu and F. Y. Wu, Soluble kagome Ising model in a magnetic field, *Phys. Rev. E* **71**, 046120 (2005).
- [93] F. Wang and F. Y. Wu, Exact solution of close-packed dimers on the kagomé lattice, *Phys. Rev. E* **75**, 040105 (2007).