

GREEDY LEARNING TO OPTIMIZE WITH CONVERGENCE GUARANTEES

Patrick Fahy¹, Mohammad Golbabaee², Matthias Ehrhardt¹

¹University of Bath, ²University of Bristol

{pf341, me549}@bath.ac.uk, m.golbabaee@bristol.ac.uk

ABSTRACT

Learning to optimize is an approach that leverages training data to accelerate the solution of optimization problems. Many approaches use unrolling to parametrize the update step and learn optimal parameters. Although L2O has shown empirical advantages over classical optimization algorithms, memory restrictions often greatly limit the unroll length and learned algorithms usually do not provide convergence guarantees. In contrast, we introduce a novel method employing a greedy strategy that learns iteration-specific parameters by minimizing the function value at the next iteration. This enables training over significantly more iterations while maintaining constant memory usage. We parameterize the update such that parameter learning corresponds to solving a convex optimization problem at each iteration. In particular, we explore preconditioned gradient descent with multiple parametrizations including a novel convolutional preconditioner. With our learned algorithm, convergence in the training set is proven even when the preconditioner is neither symmetric nor positive definite. Convergence on a class of unseen functions is also obtained, ensuring robust performance and generalization beyond the training data. We test our learned algorithms on two inverse problems, image deblurring and Computed Tomography, on which learned convolutional preconditioner demonstrates improved empirical performance over classical optimization algorithms such as Nesterov’s Accelerated Gradient Method and the quasi-Newton method L-BFGS.

1 INTRODUCTION

We consider the optimization problem

$$\min_x f(x), \quad (1)$$

with the assumption that $f : \mathcal{X} \rightarrow \mathbb{R}$ is convex, L -smooth and bounded below, where \mathcal{X} is a Hilbert space. Classic optimization methods are built in a theoretically justified manner, with guarantees on their performance and convergence properties. For example, Nesterov’s Accelerated Gradient Method (NAG) (Nesterov, 1983) accelerates classical first-order algorithms using momentum. However, practitioners often concentrate on problems within a much smaller class. For example, in reconstructing images from blurred observations y generated by a blurring operator A , one might minimize a function from the class:

$$\mathcal{F} = \left\{ f : \mathcal{X} \rightarrow \mathbb{R} : f(x) = \frac{1}{2} \|Ax - y\|^2 + R(x), y \in \mathcal{Y} \right\}, \quad (2)$$

where $R : \mathcal{X} \rightarrow \mathbb{R}$ is a chosen regularizer. Learning to optimize (L2O) uses data to learn how to minimize functions $f \in \mathcal{F}$ in a small number of iterations. Typically, the solution at each iteration t is updated by a parametrized function $G_\theta : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X}$ (i.e. the update rule) as dependent on parameters θ_t at iteration t as

$$x_{t+1} = x_t - G_{\theta_t}(x_t, \nabla f(x_t)). \quad (3)$$

Unrolling algorithms e.g. Monga et al. (2021) directly parametrize the update step as a neural network, often taking the previous iterates of the solution updates and the gradients as input arguments to the

neural network. For some $T > 0$, the parameters $\theta = (\theta_0, \dots, \theta_{T-1})$ can be learned to minimise the loss

$$L(\theta) = \mathbb{E}_{f \in \mathcal{F}} \left[\sum_{t=1}^T f(x_t) \right]. \quad (4)$$

Learned optimization algorithms often lack convergence guarantees, including many that use RNNs (Andrychowicz et al., 2016; Metz et al., 2019) or Reinforcement learning (Li & Malik, 2016). Liu et al. (2023) consider methods of the form $x_{t+1} = x_t - G_t \nabla f(x_t) + b_t$, for $f : \mathbb{R}^n \rightarrow \mathbb{R}$, a diagonal matrix $G_t \in \mathbb{R}^{n \times n}$, and a vector $b_t \in \mathbb{R}^n$. The G_t and b_t are constructed using the outputs of neural networks. However, their method does not guarantee convergence to a minimizer.

Other approaches achieve provable convergence, which can be enforced with safeguarding (Heaton et al., 2023), or constructing convergent algorithms by learning parameters within a provably convergent set (Banert et al., 2020; 2024). Tan et al. (2023a;b) learn mirror maps using input-convex neural networks within the mirror descent optimization algorithm such that the algorithm is provably convergent. Lastly, Sucker et al. (2024) and Sambharya & Stellato (2024) consider applying the PAC-Bayes framework to L2O.

Unlike NAG, Newton’s method accelerates convergence by applying the inverse Hessian to the gradient, which can be costly in practice. Quasi-Newton methods like BFGS (Nocedal & Wright, 2006b) approximate the Hessian, and L-BFGS (Liu & Nocedal, 1989) is used when BFGS is too memory-intensive. Similarly, we aim to accelerate the optimization by learning a preconditioner G_t in the update $x_{t+1} = x_t - G_t \nabla f(x_t)$.

1.1 CONTRIBUTIONS

Our paper contributes in the following ways:

- A novel approach to L2O that learns parameters at each iteration sequentially, using a greedy approach by minimizing the function value at the next iteration. This enables training over significantly more iterations while maintaining constant memory usage: Section 3.
- Convergence in the training set is proven even when the preconditioner is neither symmetric nor positive definite: Section 4.
- Convergence is proven on a class of unseen functions under certain conditions: Section 4.
- Learning parameters is a convex optimization problem for ‘linear parametrizations’ of G_t , enabling training that is significantly faster, with closed-form solutions for least-squares functions: Section 5.
- A novel parametrization of G_t as a convolution operator. At iteration t we learn a convolutional kernel κ_t such that $G_t x = \kappa_t * x$. This parametrization is shown to outperform Nesterov’s Accelerated Gradient and L-BFGS on test data: Section 6.

In Section 6, we validate our learned algorithms on two inverse problems: image deblurring and Computed Tomography (CT). Inverse problems represent a crucial class of optimization problems that appear in important fields such as medical imaging and machine learning. Many such problems have an associated forward operator which is highly ill-conditioned, making them an ideal test for optimization algorithms.

2 NOTATION

We require the following notation. Let \mathcal{X} be a Hilbert space with its corresponding field given by \mathbb{R} and norm $\|\cdot\|$. A function $f : \mathcal{X} \rightarrow \mathbb{R}$ is convex if for all $x, y \in \mathcal{X}$ and for all $\alpha \in [0, 1]$:

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y). \quad (5)$$

A function $f : \mathcal{X} \rightarrow \mathbb{R}$ is L -smooth with parameter $L > 0$ if its gradient is Lipschitz continuous, i.e., if for all $x, y \in \mathcal{X}$, the following inequality holds:

$$\|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|. \quad (6)$$

A function $f : \mathcal{X} \rightarrow \mathbb{R}$ is bounded below if there exists some $M \in \mathbb{R}$ such that $f(x) \geq M$ for all $x \in \mathcal{X}$. We say that $f \in \mathcal{F}_L$ if f is convex, L -smooth, and bounded below. We assume that the Hilbert space \mathcal{X} has dimension $\dim(\mathcal{X}) = n$ and, therefore, admits a finite orthonormal basis $\{e_1, \dots, e_n\}$. For $x \in \mathcal{X}$ and $j \in \{1, \dots, n\}$, define $[x]_j := \langle x, e_j \rangle$. For $x, y \in \mathcal{X}$, define the pointwise product $x \odot y$ by

$$[x \odot y]_j = [x]_j [y]_j. \quad (7)$$

For Hilbert spaces \mathcal{X} and \mathcal{Y} , denote the space of linear operators from \mathcal{X} to \mathcal{Y} by $\mathcal{L}(\mathcal{X}, \mathcal{Y})$. If $\mathcal{Y} = \mathcal{X}$, we write $\mathcal{L}(\mathcal{X})$. For example, if $\mathcal{X} = \mathbb{R}^n$, $\mathcal{L}(\mathcal{X})$ is the space of $n \times n$ matrices. Denote the adjoint of $A \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ by A^* , meaning that for $x \in \mathcal{X}, y \in \mathcal{Y}$, $\langle Ax, y \rangle = \langle x, A^*y \rangle$. Denote by $I \in \mathcal{L}(\mathcal{X})$ the identity operator: $I(x) = x$ for all $x \in \mathcal{X}$.

3 GREEDY LEARNING TO OPTIMIZE OF PRECONDITIONED GRADIENT DESCENT

This section introduces the proposed method: greedy learning to optimize. Firstly, we introduce how we parametrize the optimization algorithm as preconditioned gradient descent. Next, we detail our training data and define a loss function with which we learn parameters. We then provide an algorithm of how parameters are learned sequentially using a greedy approach. Lastly, we show how our learned algorithm is applied to unseen functions.

At each iteration $t \in \{0, 1, 2, \dots\}$, we parametrize the linear operator G_t using a Hilbert space Θ and learn parameters $\theta_t \in \Theta$ in the update

$$x_{t+1} = x_t - G_{\theta_t} \nabla f(x_t). \quad (8)$$

The following propositions show that it is possible to obtain convergence after just one iteration of the update (8). Firstly, we show that it is possible to even when G is a pointwise operator, i.e. $Gx := p \odot x$ for some $p \in \mathcal{X}$.

Proposition 1. *Assume that $f : \mathcal{X} \rightarrow \mathbb{R}$ is convex, continuously differentiable, and has a global minimum, and take any initial point $x_0 \in \mathcal{X}$. Then there exists $p \in \mathcal{X}$ such that, $x_0 - p \odot \nabla f(x_0) \in \arg \min_x f(x)$.*

While the pointwise parametrization obtains convergence after one iteration for one function, for an arbitrary linear operator $G \in \mathcal{L}(\mathcal{X})$, under certain conditions, one can obtain convergence after one iteration for multiple functions.

Proposition 2. *For $k \in \{1, \dots, N\}$, assume that $f_k : \mathcal{X} \rightarrow \mathbb{R}$ is convex, continuously differentiable, and has a global minimum, with any initial point $x_k^0 \in \mathcal{X}$. Assume that the set of gradients $\{\nabla f_1(x_1^0), \dots, \nabla f_N(x_N^0)\}$ is linearly independent. Then if $N \leq n$, there exists an operator $P \in \mathcal{L}(\mathcal{X})$ such that $x_k^0 - P \nabla f_k(x_k^0) \in \arg \min_x f_k(x)$, for all $k \in \{1, \dots, N\}$.*

Propositions 1 and 2 motivate learning θ_t by considering the function values only at the next iteration. In order to learn the parameters θ_t for $t \in \{0, 1, 2, \dots\}$, we use a training dataset of functions $\mathcal{T} := \{f_1, \dots, f_N\}$, with $f_k \in \mathcal{F}_{L_k}$ for $k \in \{1, \dots, N\}$, with corresponding initial points $\mathcal{X}_0 := \{x_1^0, \dots, x_N^0\}$.

We consider learning parameters using a regularizer $R : \Theta \rightarrow \mathbb{R}$ so that undesirable properties are penalized. At iteration t , we solve the optimization problem

$$\theta_t \in \arg \min_{\theta} \left\{ g_{t, \lambda_t}(\theta) := \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - G_{\theta} \nabla f_k(x_k^t)) + \lambda_t R(\theta) \right\}, \quad (9)$$

for some regularization parameter $\lambda_t \geq 0$, which is used to balance the importance of the regularizer. Such a strategy is greedy, as learning refers to tuning the parameters θ_t considering only the function values at the next iteration, $f_k(x_k^{t+1})$. The sequential training procedure for parameter learning is detailed in Algorithm 1. Unlike unrolling, where memory requirements scale linearly with the number of training iterations, greedy training is a memory-friendly method, with memory that is constant with increasing training iterations.

Algorithm 1 Training algorithm for greedy parameter learning in preconditioned gradient descent

```

1: Input: Functions  $f_1, \dots, f_N$ , initial points  $x_1^0, \dots, x_N^0$ , final iteration  $T$ , regularization parameters  $\lambda_0, \dots, \lambda_{T-1} \geq 0$ .
2: for  $t = 0, 1, 2, \dots, T - 1$  do
3:    $\theta_t \in \arg \min_{\theta} g_{t, \lambda_t}(\theta)$ 
4:   for  $k = 1, 2, \dots, N$  do
5:      $x_k^{t+1} = x_k^t - G_{\theta_t} \nabla f_k(x_k^t)$ 
6:   end for
7: end for
8: Output: Learned parameters  $\theta_0, \dots, \theta_{T-1}$ .

```

Suppose that training is terminated after iteration T , having learned the parameters $\theta_0, \dots, \theta_{T-1}$. To minimise an unseen function f with initial point x_0 , we propose Algorithm 2.

Algorithm 2 Learned algorithm applied to a new function f

```

1: Input: Function  $f$  with initial point  $x_0$ .
2: for  $t = 0, 1, 2, \dots$  do
3:   if  $t < T$  then
4:      $x_{t+1} = x_t - G_{\theta_t} \nabla f(x_t)$ 
5:   else
6:      $x_{t+1} = x_t - G_{\theta_{T-1}} \nabla f(x_t)$ 
7:   end if
8: end for
9: Output:  $x_t$ .

```

4 CONVERGENCE RESULTS

This section contains convergence results for our learned Algorithm 2. Firstly, convergence is obtained on training functions as $T \rightarrow \infty$, without the need for the learned operators G_{θ_t} to have properties such as being symmetric or positive definite. Following this, we show convergence results and rates for a class of unseen functions if λ_t is asymptotically non-vanishing. Before we present the convergence results, we require the following definitions, the first of which provides a condition for which the update rule (8) generalizes gradient descent (GD): $x_{t+1} = x_t - \alpha_t \nabla f(x_t)$ for $\alpha_t > 0$.

Definition 1. We say that the family (G_{θ}) is GGD (generalizes gradient descent) if for all $\alpha > 0$, there exist parameters θ such that

$$G_{\theta} = \alpha I. \quad (10)$$

Parametrizations of (G_{θ}) that satisfy the GGD property are shown in section 5. Let $\tau = 1/L_{\text{train}}$, where $L_{\text{train}} = \max\{L_1, \dots, L_N\}$ is the largest smoothness coefficient in the training data set. This choice of step size in gradient descent ensures convergence for all functions $f_k \in \mathcal{T}$. From this point forward, we assume (G_{θ}) is GGD, meaning there exists some $\tilde{\theta}$ such that $G_{\tilde{\theta}} = \tau I$. Furthermore, the GGD property can be leveraged to establish provable convergence for a set of unseen functions by introducing a penalty when the parameters deviate significantly from $\tilde{\theta}$. With this purpose, we define $R(\theta)$ in (9) as

$$R(\theta) := \frac{1}{2} \|\theta - \tilde{\theta}\|_2^2. \quad (11)$$

The next definition is to ensure the parametrized algorithm adopts the convergence properties of gradient descent on the training data.

Definition 2. We say that θ_t is BGD (better than gradient descent) with regularization parameter λ_t if

$$g_{t, \lambda_t}(\theta_t) \leq g_{t, 0}(\tilde{\theta}) = \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - \tau \nabla f_k(x_k^t)). \quad (12)$$

4.1 CONVERGENCE ON TRAINING DATA

Theorem 1. Convergence on training data. *Suppose that $\lambda_t \geq 0$ and $(\theta_t)_{t=0}^\infty$ is a BGD sequence of parameters. Then with Algorithm (1), we have $\nabla f_k(x_k^t) \rightarrow 0$ as $t \rightarrow \infty$ for all $k \in \{1, \dots, N\}$.*

Convergence rates can also be obtained for training data, see Appendix Section 4.

4.2 CONVERGENCE ON UNSEEN DATA

We now show convergence on unseen data. Firstly, we show that if the regularization parameters λ_t are eventually non-vanishing, then the learned parameters tend towards $\tilde{\theta}$.

Lemma 1. *If $\liminf_{t \rightarrow \infty} \lambda_t > 0$ and $(\theta_t)_{t=0}^\infty$ is BGD, then $\theta_t \rightarrow \tilde{\theta}$ as $t \rightarrow \infty$.*

This result is useful to ensure convergence on unseen data as if $G : \Theta \rightarrow \mathcal{L}(\mathcal{X})$ is continuous, then under the same conditions, $G_{\theta_t} \rightarrow G_{\tilde{\theta}} = \tau I$ as $t \rightarrow \infty$, i.e. our learned algorithm gets close to GD for large t . The idea is that we start with a method that fits the data very well leading to quick initial convergence, but in the interest of safety, over time we become closer to an algorithm with proven convergence.

Theorem 2. Convergence on unseen data for regularized parameter learning

Assume that $G : \Theta \rightarrow \mathcal{L}(\mathcal{X})$ is continuous, θ_t is BGD and $\liminf_{t \rightarrow \infty} \lambda_t > 0$. Then, there exists a constant $\tilde{L} > 0$, and a final training iteration T such that for all $f \in \mathcal{F}_{\tilde{L}}$ and any starting point x_0 , using Algorithm 2 (which depends on T), we have $\nabla f(x_t) \rightarrow 0$ as $t \rightarrow \infty$.

In practice, provable convergence can be verified during training. At iteration t , the regularization parameter λ_t may be selected large enough such that $\|G_{\theta_t} - \tau I\| < \tau$, which guarantees convergence if training is then stopped with $T = t$. A proof is provided with Proposition 6 in the Appendix. The following theorem presents the convergence rate obtained for test functions.

Theorem 3. Convergence rates on unseen data for regularized parameter learning

Under the same assumptions as Theorem 2 and if $(x_t)_{t=1}^\infty$ is a bounded sequence then there exists a constant $C > 0$, such that

$$f(x_t) - f(x^*) \leq \frac{C}{t}. \quad (13)$$

This result gives the worst-case convergence rate of the learned algorithm. In Section 6 we will see that the empirical performance of the learned algorithms may exceed that of NAG and L-BFGS.

5 LINEAR PARAMETRIZATIONS

In this section, we consider 'linear parametrizations' of G , defined below.

Definition 3. *We call G a linear parameterization if $G : \Theta \rightarrow \mathcal{L}(\mathcal{X})$ is a linear map. This means there exists a linear operator $B_k^t \in \mathcal{L}(\Theta, \mathcal{X})$ such that*

$$G_\theta \nabla f_k(x_k^t) = B_k^t(\theta). \quad (14)$$

The motivation is that when G is a linear parametrization, each optimization problem (9) is convex (as it is the composition of a convex function with a linear function (Beck, 2014)). Therefore, learning comprises solving a sequence of convex optimization problems. In this case, there exist fast, provably convergent algorithms to find global solutions. Such problems also require fewer hyperparameters to tune. Due to the speed of training enabled by linear parametrizations and the constant memory of the greedy approach, we are able to learn algorithms up to significantly higher iterations. Four examples of linear parametrizations of G are provided in Table 1. These parametrizations are used for the numerical experiments in Section 6.

Lemma 2. *All parametrizations G_θ in Table 1 satisfy the GGD property (10), and are all continuous with respect to their parameters.*

Corollary 1. *We obtain the convergence results from Theorems 2 and 3 for linear parametrizations in Table 1, when θ_t is BGD and $\liminf_{t \rightarrow \infty} \lambda_t > 0$.*

Table 1: Examples of linear parametrizations

Label	Description	parametrization	# parameters
(PS)	Scalar step size	$G_{\alpha_t} = \alpha_t I, \alpha_t \in \mathbb{R}$	1
(PP)	Pointwise operator	$G_{p_t} x = p_t \odot x, p_t, x \in \mathcal{X}$	$\dim(\mathcal{X})$
(PC)	Image convolution	$G_{\kappa_t} x = \kappa_t * x, \kappa_t \in \mathbb{R}^{m_1 \times m_2}$	$m_1 m_2$
(PF)	Full linear operator	$G_{P_t} = P_t \in \mathcal{L}(\mathcal{X})$	$\dim(\mathcal{X})^2$

5.1 CLOSED-FORM SOLUTIONS

If each function $f_k \in \mathcal{T}$ can be written as a least-squares function, then the parameters θ_t at iteration t have a closed-form solution. For $k \in \{1, \dots, N\}$, let f_k be given by

$$f_k(x) = \frac{1}{2} \|A_k x - y_k\|^2, \quad (15)$$

with corresponding $y_k \in \mathcal{Y}$, for a Hilbert space \mathcal{Y} , and linear operator $A_k \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$.

Proposition 3. *For a linear parametrization G , let B_k^t be given as in (14), and f_k as in (15). Then θ_t given by*

$$\theta_t = \left(\lambda_t I_{\Theta} + \frac{1}{N} \sum_{k=1}^N (A_k B_k^t)^* (A_k B_k^t) \right)^\dagger \left(\lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla f_k(x_k^t) \right) \quad (16)$$

is a solution to (9), where M^\dagger represents the Moore–Penrose pseudoinverse of a linear operator M .

Note, for this class of functions, we also have a closed-form equation for exact line search for a scalar step size (Nocedal & Wright, 2006a), which we recover with $\lambda_t = 0, N = 1$ for the scalar parametrization (PS) in Table 1. Therefore the optimization problem (9) can be seen as an extension of exact line search to include linear operators. Calculations for the closed-form solutions for the four examples in Table 1 are detailed in Appendix Section 5. In general, we require optimization algorithms to approximate θ_t . Due to the optimization problem being convex, we provide gradient calculations and smoothness constants for these parametrizations in Appendix Section C.1.

6 NUMERICAL EXPERIMENTS

In this section, we test the four linear parametrizations in Table 1 on two inverse problems in imaging: image deblurring and CT. We consider linear inverse problems, defined by receiving an observation $y \in \mathcal{Y}$, generated from a ground-truth x_{true} via some linear forward operator $A : \mathcal{X} \rightarrow \mathcal{Y}$, such that $y = Ax_{\text{true}} + \varepsilon$, where $\varepsilon \in \mathcal{Y}$ is some random noise, and the goal is to recover x_{true} . For both experiments, $\mathcal{X} = \mathbb{R}^{h_1 \times h_2}$ for $h_1, h_2 \in \mathbb{N}$, and ε is noise sampled from a zero-mean Gaussian distribution. To approximate x_{true} from y , we solve

$$\min_x \left\{ f(x) := \frac{1}{2} \|Ax - y\|_2^2 + \alpha H_\epsilon(x) \right\}, \quad (17)$$

for a given regularization parameter α . The regularizer H_ϵ is the Huber Total Variation (Rudin et al., 1992; Huber, 1992) defined by

$$H_\epsilon(x) = \sum_{i,j=1}^{h_1, h_2} h_\epsilon \left(\sqrt{(Dx)_{i,j,1}^2 + (Dx)_{i,j,2}^2} \right), \quad h_\epsilon(s) = \begin{cases} \frac{1}{2\epsilon} s^2, & \text{if } |s| \leq \epsilon \\ |s| - \frac{\epsilon}{2}, & \text{otherwise,} \end{cases} \quad (18)$$

where finite difference operator $D : \mathbb{R}^{h_1 \times h_2} \rightarrow \mathbb{R}^{h_1 \times h_2 \times 2}$ is defined in Chambolle & Pock (2016). We take $\epsilon = 0.01$ and normalize the forward operator in both cases so that $\|A\| = 1$. Then, each function f is L -smooth, where $L = 1 + \frac{8\alpha}{\epsilon}$ (Chambolle & Pock, 2016).

For each parametrization in Table 1, to learn parameters θ_t we apply NAG for solving the optimization problem 9. The initialization is set as $\theta_t^0 = \tilde{\theta}$ for $t = 0$, and $\theta_t^0 = \theta_{t-1}^0$ for $t > 0$. NAG is stopped when $\|\nabla g_t(\theta_t^\ell)\|_2 / \|\nabla g_t(\theta_t^0)\|_2 < 10^{-3}$, or when $\ell = \ell_{\text{stop}} = 5000$. For both problems, we use a

training set of 100 functions for parametrizations (PS), (PP), and (PC). For the parametrization (PF), the model is trained using 1000 functions and is only implemented for the CT problem. Testing is performed on a separate set of 100 functions for all parametrizations. The learned convolutional kernel (PP) has dimensions $h_1 \times h_2$, matching the size of the images in \mathcal{X} .

Given a dataset of functions f_1, \dots, f_N , the mean value at iteration t is defined as $F(x_t) = \frac{1}{N} \sum_{k=1}^N f_k(x_k^t)$ and let $F^* := \min_{x \in \mathcal{X}^N} F(x)$. This function is used to visualize the performance of the learned methods. The learned algorithms are compared with NAG, L-BFGS, and GD with constant step size τ . All computations were carried out on an Nvidia RTX 3600 12GB GPU.

6.1 IMAGE DEBLURRING

In this example, the forward operator A in (17) is a Gaussian blur with a 5×5 kernel size and a standard deviation of $\sigma = 1.5$. We use the STL-10 dataset (Coates et al., 2011) with greyscale images of size 96×96 as \mathcal{X} . The noise ε is modeled with a standard deviation of 2.5×10^{-3} , and we set $\alpha = 10^{-5}$, resulting in $L = 1.008$. The initial point x_0 is taken equal to the observation y .

Training with the greedy method was performed up to iteration $T = 600$ with $\lambda_t = 0$ for all t for the parametrizations (PS), (PP), (PC). This resulted in learning 600 parameters for (PS) and 5529600 for both (PP) and (PC). For the convolutional parametrization (PC) at iterations $t > 50$ we set $\ell_{\text{stop}} = 500$. The total training time for (PS) and (PP) was approximately 10 minutes each, while (PC) required 19 hours.

Figure 1a shows that the learned scalar parameters (PS) eventually fluctuate above and below $2/L$, which is outside of the range of provable convergence of gradient descent with a constant step size. Despite this, the learned algorithm leads to convergence on training data as $t \rightarrow \infty$ by Theorem 1. In Figure 1b, Oscillatory behavior also occurs in the pointwise parametrization (PP), with most fluctuations occurring near the boundary. We also see values outside the interval $(0, 2/L)$, including negative values. The learned convolutional kernels (PC) in Figure 1c also contain positive and negative values. These kernels are predominantly weighted towards the center, suggesting that information from neighboring pixels is prioritized over more distant ones. As the number of iterations increases, the kernels exhibit increasing similarity, though no formal convergence result for θ_t has been established when $\lambda_t = 0$.




Figure 1: Learned parameters for the image deblurring problem with $\lambda_t = 0$ for all $t \in \{0, 1, \dots, T\}$. (a) Learned scalar parameters (PS) for $t \in \{0, 1, \dots, 100\}$ for the image deblurring problem compared to $2/L$. (b) Learned 96×96 pointwise operators (PP) restricted to the interval $[-10, 10]$, against iteration t for $t \in \{5, 6, 7, 8, 9\}$. (c) Learned 96×96 convolutional kernels (PC) restricted to the interval $[-10, 10]$, against iteration t for $t \in \{0, 2, 5, 15, 100\}$.

Figure 2 shows that the learned parametrizations (PS), (PP), and (PC) generalize well to unseen data due to the closeness of the train and test curves. The learned pointwise parametrization performs very similarly to the learned scalar parameterization for this example. This is despite (PP) having an equal number of parameters as the learned convolutional kernels, which use their parameters to capture global information across the image, rather than focusing on specific pixel-level details. Furthermore,

the learned convolutional algorithm significantly outperforms both NAG and L-BFGS on the test data, reaching a tolerance of 10^{-6} in just over 100 iterations, compared with about 400 and 450 for L-BFGS and NAG, respectively. More information on the number of iterations to reach specific tolerances can be found in Appendix Section D.3. Additionally, Appendix Section D.1 provides comparisons of the best and worst performance between the learned convolutional parametrization and NAG, while D.2 explores the impact of different kernel sizes on the performance of the (PC) parametrization.

We now consider the effect of regularization on the convergence of the learned Algorithm 2. We use regularization at iteration T for the parametrizations (PS) and (PP) to ensure convergence when applying Algorithm 2 to further iterations. The (PC) parametrization was not considered as it has already reached a suitable tolerance within the training iterations. As discussed in Section 4, we may select λ_T large enough such that the learned parameters lead to guaranteed convergence. For the (PS) parametrization, we find $\lambda_T = 2 \times 10^{-7}$ guarantees convergence of Algorithm 2, and for the (PP) parametrization, $\lambda_T = 3.90625 \times 10^{-9}$ guarantees convergence. Figure 2 shows that when learned without regularization, the learned algorithm quickly diverges. However, if the regularization parameter λ_T is chosen large enough, the algorithm converges as proved in Theorem 2.




Figure 2: Left: Performance comparison of the learned methods on training versus test data for the image deblurring problem within training iteration. Center: Test performance versus benchmark optimization algorithms within training iterations. Right: Performance beyond training iterations for the (PP) and (PS) parameterizations when learned with and without guaranteed convergence. The final training iteration T is indicated by the dotted vertical line.

Figure 3 illustrates the reduced number of iterations needed to obtain a deblurred image reconstruction using the learned convolution parameterization. It requires approximately four times more iterations to achieve a comparable reconstruction quality with NAG compared to the learned convolutional algorithm.




Figure 3: Comparisons of reconstructions for the image deblurring problem.

We now compare the time taken for training with the greedy learning approach versus the unrolling approach. In the unrolling approach, we fix the number of iterations to $T = 20$ and jointly learn the parameters $\theta_0, \dots, \theta_{T-1}$ in the update rule (8) with the (PC) parametrization. The same training dataset as the greedy method is used, using a batch size of 4, with the objective function defined in equation (4). Parameter updates are performed using Adam (Kingma, 2014), with the learning rate selected via grid search, while keeping all other parameters at their default values. The unrolling method was trained for 13000 epochs, taking approximately 41 hours in total. As shown in Figure 4, the performance of the learned parameters using the unrolled method is similar to that learned with

the greedy method over the first 20 iterations. However, during the 41-hour training period for the unrolling approach, the greedy approach (9) could learn parameters for at least 1864 iterations using NAG (with a stopping iteration of $\ell_{\text{stop}} = 500$ for $t > 50$), showing the benefit of solving a sequence of convex optimization problems.




Figure 4: Performance of the learned unrolled algorithm versus the greedy learned algorithm on test data.

6.2 COMPUTED TOMOGRAPHY

In this case, the forward operator A in (17) is the Radon transform in 2D. We simulate CT measurements using ODL (Adler et al., 2017) with a parallel-beam geometry employing 90 projection angles evenly distributed over a 180-degree range. For the dataset, we extract 40×40 pixel crops from the center of each ground-truth image in the SARS-CoV-2 CT-scan dataset (Soares et al., 2020). The noise ε is modeled with a standard deviation of 10^{-2} , and we set $\alpha = 10^{-4}$, resulting in $L = 1.08$. The initial point x_0 is taken equal to $0 \in \mathcal{X}$.

Greedy training was performed up to iteration $T = 200$ with $\lambda_t = 0$ for all iterations t for the (PS), (PP), and (PC) parametrizations. The total time for training (PS) was about 10 minutes, for (PP) was about 67 minutes, and (PC) took approximately 10 hours. For the (PF) parametrization, training was performed up to iteration $T = 11$ with $\lambda_t = 0$ for all t . Furthermore, the (PF) parametrization was trained with regularization such that $\lambda_t = 10^{-10}$ for $t < T = 101$ iterations. At iteration 101, the learned operator G_{θ_T} satisfied $\|G_{\theta_T} - \tau I\| < \tau$, guaranteeing convergence on iterations $t \geq T$. For each iteration t , solving the optimization problem (9) with the (PF) parametrization took one hour.

Figure 5 shows that the learned scalar values again eventually fluctuate above and below $2/L$, similar to the behavior observed for the deblurring problem. The learned pointwise operators also exhibit oscillations between consecutive iterations, with many values falling outside the interval $(0, 2/L)$. Likewise, the learned convolutional kernels for the CT problem contain both positive and negative values, are predominantly weighted toward the center of the kernel, and become increasingly similar as the iterations progress.




Figure 5: (a) Learned scalars for $t \in \{0, 1, \dots, 100\}$. (b) Learned pointwise operators for $t \in \{4, 5, 6, 7, 8\}$. (c) Learned kernels restricted to the interval $[-5, 5]$, for $t \in \{0, 1, 3, 5, 10\}$. (d) Example CT observation. (e) Reconstruction by minimizing 17.

Figure 6 shows that the learned parametrizations (PS), (PP), and (PC) generalize well to unseen data for the CT problem. In contrast, the full parametrization (PF) shows signs of overfitting, as it does not generalize well. The (PF) parametrization with regularization mitigates this issue, as the generalization performance improves. Similar to the deblurring problem, the learned (PC) parametrization outperforms NAG and L-BFGS on the CT test data, reaching a tolerance of 10^{-10} in approximately 30 iterations, compared with about 80 and more than 100 for L-BFGS and NAG, respectively. The (PF) parametrization with regularization initially converges quickly but its speed decreases later due to regularization. This is because, with increasing iterations, the learned update gets closer to gradient descent.




Figure 6: Left: Train versus test set performance of the learned parametrizations. Right: Test performance versus benchmark optimization algorithms.

Figure 7 demonstrates that the learned parametrization (PC) achieves high-quality reconstructions with significantly fewer iterations compared to NAG for the CT problem. Also, note that the regularized (PF) parametrization achieves a good visual reconstruction after only two iterations while also providing guaranteed convergence.




Figure 7: Left: An example reconstruction for the CT problem. Right: The absolute difference between the final reconstruction and the intermediate reconstruction for the full parametrization with regularization at iteration 2, the convolutional parametrization at iteration 10, and for NAG at iterations 10 and 70.

7 CONCLUSIONS

In this paper, we presented a novel L2O approach for minimizing unconstrained convex problems with differentiable objective functions. Our method employs a greedy strategy to learn a linear operator at each iteration of an optimization algorithm, meaning that memory requirements are constant with respect to the iterations. Parameter learning in our framework also corresponds to solving convex optimization problems, enabling the use of fast algorithms. Both factors allow training over a large number of training iterations, which would otherwise be prohibitively expensive. Furthermore, we obtain convergence results on the training set, even when the preconditioner is neither symmetric nor positive definite. Convergence is also proven for a class of unseen functions under certain conditions. The numerical results on imaging inverse problems demonstrate that our approach with a novel convolutional parametrization outperforms classical optimization algorithms.

REFERENCES

- Jonas Adler, Holger Kohr, and Ozan Öktem. Operator discretization library (odl). *Zenodo*, 2017.
- Marcin Andrychowicz, Misha Denil, Sergio Gomez, Matthew W Hoffman, David Pfau, Tom Schaul, Brendan Shillingford, and Nando De Freitas. Learning to learn by gradient descent by gradient descent. *Advances in neural information processing systems*, 29, 2016.
- Sebastian Banert, Axel Ringh, Jonas Adler, Johan Karlsson, and Ozan Öktem. Data-driven nonsmooth optimization. *SIAM Journal on Optimization*, 30(1):102–131, 2020.
- Sebastian Banert, Jevgenija Rudzusika, Ozan Öktem, and Jonas Adler. Accelerated forward-backward optimization using deep learning. *SIAM Journal on Optimization*, 34(2):1236–1263, 2024.
- Amir Beck. *Introduction to nonlinear optimization: Theory, algorithms, and applications with MATLAB*. SIAM, 2014.
- Antonin Chambolle and Thomas Pock. An introduction to continuous optimization for imaging. *Acta Numerica*, 25:161–319, 2016.
- Adam Coates, Andrew Ng, and Honglak Lee. An analysis of single-layer networks in unsupervised feature learning. In *Proceedings of the fourteenth international conference on artificial intelligence and statistics*, pp. 215–223. JMLR Workshop and Conference Proceedings, 2011.
- Gene H Golub and Charles F Van Loan. *Matrix computations*. JHU press, 2013.
- Robert M Gower, Mark Schmidt, Francis Bach, and Peter Richtárik. Variance-reduced methods for machine learning. *Proceedings of the IEEE*, 108(11):1968–1983, 2020.
- Howard Heaton, Xiaohan Chen, Zhangyang Wang, and Wotao Yin. Safeguarded learned convex optimization. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 37, pp. 7848–7855, 2023.
- Peter J Huber. Robust estimation of a location parameter. In *Breakthroughs in statistics: Methodology and distribution*, pp. 492–518. Springer, 1992.
- Diederik P Kingma. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- Ke Li and Jitendra Malik. Learning to optimize. *arXiv preprint arXiv:1606.01885*, 2016.
- Dong C Liu and Jorge Nocedal. On the limited memory bfgs method for large scale optimization. *Mathematical programming*, 45(1):503–528, 1989.
- Jialin Liu, Xiaohan Chen, Zhangyang Wang, Wotao Yin, and HanQin Cai. Towards constituting mathematical structures for learning to optimize. In *International Conference on Machine Learning*, pp. 21426–21449. PMLR, 2023.
- Luke Metz, Niru Maheswaranathan, Jeremy Nixon, Daniel Freeman, and Jascha Sohl-Dickstein. Understanding and correcting pathologies in the training of learned optimizers. In *International Conference on Machine Learning*, pp. 4556–4565. PMLR, 2019.
- Vishal Monga, Yuelong Li, and Yonina C Eldar. Algorithm unrolling: Interpretable, efficient deep learning for signal and image processing. *IEEE Signal Processing Magazine*, 38(2):18–44, 2021.
- Yurii Nesterov. A method for solving the convex programming problem with convergence rate $o(1/k^2)$. In *Dokl akad nauk Sssr*, volume 269, pp. 543, 1983.
- Yurii Nesterov et al. *Lectures on convex optimization*, volume 137. Springer, 2018.
- Jorge Nocedal and Stephen J Wright. Line search methods. *Numerical optimization*, pp. 30–65, 2006a.
- Jorge Nocedal and Stephen J Wright. Quasi-newton methods. *Numerical optimization*, pp. 135–163, 2006b.

Leonid I Rudin, Stanley Osher, and Emad Fatemi. Nonlinear total variation based noise removal algorithms. *Physica D: nonlinear phenomena*, 60(1-4):259–268, 1992.

Rajiv Sambharya and Bartolomeo Stellato. Data-driven performance guarantees for classical and learned optimizers. *arXiv preprint arXiv:2404.13831*, 2024.

Eduardo Soares, Plamen Angelov, Sarah Biaso, Michele Higa Froes, and Daniel Kanda Abe. Sars-cov-2 ct-scan dataset: A large dataset of real patients ct scans for sars-cov-2 identification. *MedRxiv*, pp. 2020–04, 2020.

Michael Sucker, Jalal Fadili, and Peter Ochs. Learning-to-optimize with pac-bayesian guarantees: Theoretical considerations and practical implementation. *arXiv preprint arXiv:2404.03290*, 2024.

Hong Ye Tan, Subhadip Mukherjee, Junqi Tang, and Carola-Bibiane Schönlieb. Boosting data-driven mirror descent with randomization, equivariance, and acceleration. *arXiv preprint arXiv:2308.05045*, 2023a.

Hong Ye Tan, Subhadip Mukherjee, Junqi Tang, and Carola-Bibiane Schönlieb. Data-driven mirror descent with input-convex neural networks. *SIAM Journal on Mathematics of Data Science*, 5(2): 558–587, 2023b.

A FURTHER NOTATION

The following notation is required in this section.

A function $f : \mathcal{X} \rightarrow \mathbb{R}$ is strongly convex with parameter $\mu > 0$ if $f - \frac{\mu}{2} \|\cdot\|^2$ is convex. We say $f \in \mathcal{F}_{L,\mu}$ if $f \in \mathcal{F}_L$ and f is μ -strongly convex.

If $\dim(\mathcal{X}) = n$ and $\dim(\mathcal{Y}) = m$, denote an orthonormal basis of \mathcal{X} by $\{e_1, \dots, e_n\}$, and an orthonormal basis of \mathcal{Y} by $\{\tilde{e}_1, \dots, \tilde{e}_m\}$, then A can be uniquely determined by mn scalars γ_{ij} for $i \in \{1, \dots, m\}, j \in \{1, \dots, n\}$: $A(e_j) = \sum_{i=1}^m \gamma_{ij} \tilde{e}_i$, and denote $[A]_{ij} = \gamma_{ij}$.

Denote $\mathbf{1} \in \mathcal{X}$ to be such that $[\mathbf{1}]_j = 1$ for $j \in \{1, \dots, n\}$. For operators $A, B \in \mathcal{L}(\mathcal{X})$, and elements $x, y, z \in \mathcal{X}$, define the linear operators $A \odot B$ and $x \otimes y$ by

$$[A \odot B]_{ij} := [A]_{ij}[B]_{ij}, \quad (19)$$

$$(x \otimes y)z := \langle y, z \rangle x, \quad (20)$$

with the property that

$$[x \otimes y]_{qi} = \langle y, e_i \rangle \langle x, e_q \rangle = [x]_q [y]_i. \quad (21)$$

For two linear operators $A, B \in \mathcal{L}(\mathcal{X})$, define $A \otimes B$ by

$$[A \otimes B]_{ij,kl} = [A]_{ik}[B]_{jl}. \quad (22)$$

For a linear operator $A \in \mathcal{L}(\mathcal{X})$ and $x \in \mathcal{X}$,

$$[Ax]_i = \sum_{j=1}^n [A]_{ij}[x]_j. \quad (23)$$

B PROOFS FOR SECTION 4

The following lemma is required to prove the convergence of our learned method.

Lemma 3. Define $F : \mathcal{X}^N \rightarrow \mathbb{R}$ by

$$F(x) = \frac{1}{N} \sum_{k=1}^N f_k(x_k), \quad x = (x_1, x_2, \dots, x_N) \in \mathcal{X}^N. \quad (24)$$

Then

1. Each $f_k \in \mathcal{F}_{L_k}$ implies $F \in \mathcal{F}_{L_F}$, with $L_F = \frac{1}{N} L_{\text{train}}$ where $L_{\text{train}} = \max\{L_1, \dots, L_N\}$.
2. Each $f_k \in \mathcal{F}_{L_k, \mu_k}$ implies $F \in \mathcal{F}_{L, \mu_F}$ with $\mu_F = \frac{1}{N} \mu_{\min}$ where $\mu_{\min} = \min\{\mu_1, \dots, \mu_N\}$.

Proof. We have

$$\nabla F(x) = \frac{1}{N} (\nabla f_1(x_1), \dots, \nabla f_N(x_N)), \quad (25)$$

and for any $y \in \mathcal{X}^N$,

$$\|x - y\| = \sqrt{\sum_{k=1}^N \|x_k - y_k\|^2}.$$

Then

$$\begin{aligned} \|\nabla F(x) - \nabla F(y)\| &= \frac{1}{N} \sqrt{\sum_{k=1}^N \|\nabla f_k(x_k) - \nabla f_k(y_k)\|^2} \\ &\leq \frac{1}{N} \sqrt{\sum_{k=1}^N L_k^2 \|x_k - y_k\|^2} \quad (L_k\text{-smoothness of } f_k) \\ &\leq \frac{\max\{L_1, \dots, L_N\}}{N} \|x - y\|, \end{aligned}$$

which proves 1.

For strong convexity, it is required to show that $x \mapsto (F(x) - \frac{\min\{\mu_1, \dots, \mu_N\}}{N} \|x\|^2)$ is convex. We have

$$F(x) - \frac{\min\{\mu_1, \dots, \mu_N\}}{N} \|x\|^2 = \frac{1}{N} \sum_{k=1}^N (f_k(x_k) - \min\{\mu_1, \dots, \mu_N\} \|x_k\|^2). \quad (26)$$

Notice that due to the strong convexity of f_k for all k , and that $\mu_k \geq \min\{\mu_1, \dots, \mu_N\}$,

$$x_k \mapsto (f_k(x_k) - \min\{\mu_1, \dots, \mu_N\} \|x_k\|^2) \quad (27)$$

is convex. Therefore the function $x \mapsto (F(x) - \frac{\min\{\mu_1, \dots, \mu_N\}}{N} \|x\|^2)$ is convex as it is the sum of convex functions, as required. \square

Theorem 1. Convergence on training data.

Suppose $\lambda_t \geq 0$. If θ_t is BGD then with Algorithm (1), we have

$$\nabla f_k(x_k^t) \rightarrow 0 \text{ as } t \rightarrow \infty, \quad (28)$$

for all $k \in \{1, \dots, N\}$.

Bonus: Convergence rates

Furthermore, if we denote

$$x_0 = (x_1^0, \dots, x_N^0), \quad x^* = (x_1^*, \dots, x_N^*), \quad (29)$$

then

$$F(x_t) - F(x^*) \leq \frac{\max\{L_1, \dots, L_N\}}{2tN} \|x_0 - x^*\|^2. \quad (30)$$

If, in addition, each f_k is μ_k -strongly convex, then we have linear convergence given by

$$F(x_t) - F(x^*) \leq \left(1 - \frac{\max\{L_1, \dots, L_N\}}{\min\{\mu_1, \dots, \mu_N\}}\right)^t (F(x_0) - F(x^*)). \quad (31)$$

Note that this result gives a worst-case convergence bound among train functions. However, provable convergence is still acquired. Also, note that this is not an issue for a function class with constant smoothness and strongly convex parameters.

Proof. As θ_t is BGD, we have that

$$\begin{aligned} F(x_{t+1}) &= g_{t, \lambda_t}(\theta_t) \leq g_{t, \lambda_t}(\tilde{\theta}) \\ &= \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - \tau \nabla f(x_k^t)) \\ &= F(x_t - \tau (\nabla f_1(x_1^t), \dots, \nabla f_N(x_N^t))) \\ &= F(x_t - \tau N \nabla F(x_t)) \\ &= F(x_t - \tau_F \nabla F(x_t)), \end{aligned}$$

where $\tau_F = \frac{1}{L_F}$.

F is L_F -smooth as each f_k is L_k -smooth and μ -strongly convex if each f_k is μ_k -strongly convex, where

$$\begin{aligned} L_F &= \frac{\max\{L_1, \dots, L_N\}}{N} \\ \mu_F &= \frac{\min\{\mu_1, \dots, \mu_N\}}{N}, \end{aligned}$$

and therefore, using standard convergence rate results of gradient descent (Nesterov et al., 2018), we have

$$F(x_t) - F(x^*) \leq \frac{L_F}{2t} \|x_0 - x^*\|^2, \quad (32)$$

as F is L_F -smooth, and if F is also μ_F -strongly convex we have

$$F(x_t) - F(x^*) \leq \left(1 - \frac{L_F}{\mu_F}\right)^t (F(x_0) - F(x^*)). \quad (33)$$

In both cases, we have that $\|\nabla F(x_t)\|^2 = \frac{1}{N^2} \sum_{k=1}^N \|\nabla f_k(x_k^t)\|^2 \rightarrow 0$ as $t \rightarrow \infty$, which implies that $\nabla f_k(x_k^t) \rightarrow 0$ as $t \rightarrow \infty$ for all $k \in \{1, \dots, N\}$. \square

We have proved convergence for the mean of our train functions. The following proposition proves the same convergence rate holds for each function in our training set.

Proposition 4. *Suppose we have a convergence rate for F of*

$$F(x_t) - F^* \leq C\rho(t), \quad (34)$$

for some constant $C > 0$. Then the convergence rate for all $f_k \in \mathcal{T}$ is given by

$$f_k(x_k^t) - f_k^* \leq c\rho(t), \quad (35)$$

for some constant $c > 0$.

Proof. Let $k \in \{1, \dots, N\}$. Note that by the definition of F , we may write

$$\begin{aligned} F(x_t) &= \frac{1}{N} f_k(x_k^t) + \frac{1}{N} \sum_{i=1, i \neq k}^N f_i(x_i^t), \\ \implies F^* &= \frac{1}{N} f_k^* + \frac{1}{N} \sum_{i=1, i \neq k}^N f_i^*. \end{aligned}$$

Therefore, using the convergence rate of F gives

$$\frac{1}{N} f_k(x_k^t) - \frac{1}{N} f_k^* + \frac{1}{N} \sum_{i=1, i \neq k}^N (f_i(x_i^t) - f_i^*) \leq C\rho(t) \quad (36)$$

which implies

$$\begin{aligned} f_k(x_k^t) - f_k^* + \sum_{i=1, i \neq k}^N (f_i(x_i^t) - f_i^*) &\leq NC\rho(t), \\ \implies f_k(x_k^t) - f_k^* &\leq NC\rho(t), \end{aligned}$$

as $f_i(x_i^t) \geq f_i^* 0$ for all i . \square

Proposition 5. *Assume that $f : \mathcal{X} \rightarrow \mathbb{R}$ is convex, continuously differentiable, and has a global minimum. Then for a point $z \in \mathcal{X}$ if there exists some $x^* \in \arg \min_x f(x)$ such that $[z]_i = [x^*]_i$, then $[\nabla f(x)]_i = 0$.*

Proof. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $g(t) := f(x + te_i)$, then g is convex as for $\alpha \in [0, 1], t_1, t_2 \in \mathbb{R}$, we have $g(\alpha t_1 + (1 - \alpha)t_2) = f(\alpha(x + t_1 e_i) + (1 - \alpha)(x + t_2 e_i)) \leq \alpha g(t_1) + (1 - \alpha)g(t_2)$. Note that $g'(0) = [\nabla f(x)]_i$. Assume there exists $\delta \neq 0$ such that $g(\delta) < g(0)$, then $g(\delta) < g(0) = f(x)$, which is a contradiction of x^* being an optimal point, as one can take $x = x^*$. Therefore g achieves a minimum at $t = 0$, then $[\nabla f(x)]_i = 0$. \square

Proof of Proposition 1. Choose the vector $p \in \mathcal{X}$ such that

$$[p]_i = \begin{cases} \frac{[x_0 - x^*]_i}{[\nabla f(x_0)]_i}, & \text{if } [\nabla f(x_0)]_i \neq 0, \\ 0, & \text{otherwise,} \end{cases} \quad (37)$$

and let $I = \{i : [\nabla f(x_0)]_i \neq 0\}$ Then for any $i \in I$, we have

$$\begin{aligned} [x_0 - p \odot \nabla f(x_0)]_i &= [x_0]_i - [p]_i [\nabla f(x_0)]_i \\ &= [x_0]_i - \frac{[x_0 - x^*]_i}{[\nabla f_k(x_0)]_i} [\nabla f(x_0)]_i \\ &= [x^*]_i. \end{aligned}$$

Thus, by proposition 5, $[\nabla f(x_0 - p \odot \nabla f(x_0))]_i = 0$, for all $i \in I$, and similarly $[\nabla f(x_0 - p \odot \nabla f(x_0))]_i = 0$, for all $i \notin I$, and therefore $\nabla f(x_0 - p \odot \nabla f(x_0)) = 0$, meaning that $x_0 - p \odot \nabla f(x_0) \in \arg \min_x f(x)$ as required. \square

Proof of Proposition 2. We require

$$\begin{cases} x_1^* &= x_1^0 - P \nabla f_1(x_1^0), \\ &\vdots \\ x_N^* &= x_N^0 - P \nabla f_N(x_N^0). \end{cases}$$

Each of these equations gives n linear equations in n^2 unknowns. There are N such equations and so we have nN linear equations in n^2 unknowns. Rewritten, these read

$$P [\nabla f_1(x_1^0) | \cdots | \nabla f_N(x_N^0)] = [x_1^0 - x_1^* | \cdots | x_N^0 - x_N^*]. \quad (38)$$

Such a P exists if $nN \leq n^2$, which is equivalent to $N \leq n$, and if the columns of $[\nabla f_1(x_1^0) | \cdots | \nabla f_N(x_N^0)]$ are linearly independent. \square

B.1 UNSEEN DATA

Proof of Lemma 1. Firstly,

$$g_t(\tilde{\theta}, \lambda_t) = \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - \tau \nabla f_k(x_k^t)) \rightarrow \frac{1}{N} \sum_{k=1}^N f_k^* \text{ as } t \rightarrow \infty. \quad (39)$$

Note also that as $(\theta_t)_{t=0}^\infty$ is a BGD sequence of parameters then $\frac{1}{N} \sum_{k=1}^N f_k^* \leq g_t(\theta_t, \lambda_t) \leq g_t(\tilde{\theta}, \lambda_t)$ and so $g_t(\theta_t, \lambda_t) \rightarrow \frac{1}{N} \sum_{k=1}^N f_k^*$ as $t \rightarrow \infty$ as $g_t(\theta_t, \lambda_t) \geq \frac{1}{N} \sum_{k=1}^N f_k^*$. Furthermore,

$$\begin{aligned} g_t(\tilde{\theta}, \lambda_t) - g_t(\theta_t, \lambda_t) &= -\frac{\lambda_t}{2} \|\theta_t - \tilde{\theta}\|_2^2 + \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - \tau \nabla f_k(x_k^t)) \\ &\quad - \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - G_{\theta_t} \nabla f_k(x_k^t)). \end{aligned}$$

Therefore,

$$0 = \lim_{t \rightarrow \infty} g_t(\tilde{\theta}, \lambda_t) - g_t(\theta_t, \lambda_t) = \lim_{t \rightarrow \infty} -\frac{\lambda_t}{2} \|\theta_t - \tilde{\theta}\|_2^2. \quad (40)$$

Now, $\liminf_{t \rightarrow \infty} \lambda_t > 0$ implies that

$$\frac{\lambda}{2} \|\theta_t - \tilde{\theta}\|_2^2 \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (41)$$

In particular, $\theta_t \rightarrow \tilde{\theta}$ as $t \rightarrow \infty$, as required. \square

Lemma 4. Suppose $\liminf_{t \rightarrow \infty} \lambda_t > 0$, $G : \Theta \rightarrow \mathcal{L}(\mathcal{X})$ is continuous and at each training iteration θ_t is BGT. Then for any ν such that $0 \leq \nu < \tau$, there exists an iteration T such that

$$\|G_{\theta_T} - \tau I\| \leq \nu < \tau. \quad (42)$$

Proof. By Lemma 1, $\theta_t \rightarrow \tilde{\theta}$ as $t \rightarrow \infty$, therefore as G_θ is continuous in θ we have $G_{\theta_t} \rightarrow \tau I$ as $t \rightarrow \infty$. Therefore for any $\nu > 0$ there exists some iteration $T > 0$ such that

$$\|G_{\theta_t} - \tau I\| \leq \nu \quad (43)$$

for all $t \geq T$, in particular for $t = T$. \square

Proof of Theorem 2. By Lemma 4, for any tolerance $\nu < \tau$ there exists an iteration T such that

$$\|G_{\theta_T} - \tau I\| \leq \nu. \quad (44)$$

Let

$$G_{\theta_T} = \tau I + M, \quad (45)$$

then

$$\|M\| \leq \nu. \quad (46)$$

Using L -smoothness of f , we have

$$\begin{aligned} f(x - G_{\theta_T} \nabla f(x)) &\leq f(x) - \langle G_{\theta_T} \nabla f(x), \nabla f(x) \rangle + \frac{L}{2} \|G_{\theta_T} \nabla f(x)\|^2 \\ &= f(x) - \left\langle G_{\theta_T} \nabla f(x), \nabla f(x) - \frac{L}{2} G_{\theta_T} \nabla f(x) \right\rangle \\ &= f(x) - \left\langle (\tau I + M) \nabla f(x), \nabla f(x) - \frac{L}{2} (\tau I + M) \nabla f(x) \right\rangle \\ &= f(x) - \left\langle \tau \nabla f(x) + M \nabla f(x), \nabla f(x) - \frac{L}{2} \tau \nabla f(x) - \frac{L}{2} M \nabla f(x) \right\rangle \\ &= f(x) - \left\langle \tau \nabla f(x) + M \nabla f(x), \left(1 - \frac{L\tau}{2}\right) \nabla f(x) - \frac{L}{2} M \nabla f(x) \right\rangle \\ &= f(x) - \tau \left(1 - \frac{L\tau}{2}\right) \|\nabla f(x)\|^2 + \frac{L}{2} \|M \nabla f(x)\|^2 \\ &\quad - (1 - \tau L) \langle \nabla f(x), M \nabla f(x) \rangle \\ &\leq f(x) - \tau \left(1 - \frac{L\tau}{2}\right) \|\nabla f(x)\|^2 + \frac{L\nu^2}{2} \|\nabla f(x)\|^2 + \nu |1 - \tau L| \|\nabla f(x)\|^2 \\ &= f(x) - \left(\tau \left(1 - \frac{\tau L}{2}\right) - \frac{L\nu^2}{2} - \nu |1 - \tau L| \right) \|\nabla f(x)\|^2 \\ &= f(x) - c(\nu, L, \tau) \|\nabla f(x)\|^2, \end{aligned}$$

where

$$\begin{aligned} c(\nu, L, \tau) &= \tau \left(1 - \frac{\tau L}{2}\right) - \frac{L\nu^2}{2} - \nu |1 - \tau L| \\ &= \frac{L}{2} \left(\frac{1}{L} - \nu - \left| \frac{1}{L} - \tau \right| \right) \left(\nu + \left| \frac{1}{L} - \tau \right| + \frac{1}{L} \right). \end{aligned}$$

Therefore

$$f(x_{t+1}) \leq f(x_t) - c(\nu, L, \tau) \|\nabla f(x)\|^2. \quad (47)$$

Note that $\nu + \left| \frac{1}{L} - \tau \right| + \frac{1}{L} > 0$, so for $c(\nu, L, \tau)$ to be positive, we require

$$\frac{1}{L} - \nu - \left| \frac{1}{L} - \tau \right| > 0. \quad (48)$$

Case 1 - $\frac{1}{L} \geq \tau$

Then we require

$$\tau - \nu > 0 \iff \nu < \tau, \quad (49)$$

which is true by assumption.

Case 2 - $\frac{1}{L} < \tau$

Then we require

$$\frac{2}{L} - \nu - \tau > 0 \iff \frac{1}{L} > \frac{\tau + \nu}{2}. \quad (50)$$

To conclude both cases, we have $\nu < \tau$ and therefore as $\frac{1}{\tau} < \frac{1}{\tau + \nu}$, we require only case 2 to be satisfied for $c(\nu, L, \tau) > 0$:

$$L < \frac{2}{\tau + \nu} \quad (51)$$

□

Proposition 6. Assume that $G : \Theta \rightarrow \mathcal{L}(\mathcal{X})$ is continuous. Then at any iteration t there exists $\lambda_t \geq 0$ and a constant $\tilde{L} > 0$ such that for all $f \in \mathcal{F}_{\tilde{L}}$ and any starting point x_0 , using Algorithm 2 gives $\nabla f(x_t) \rightarrow 0$ as $t \rightarrow \infty$.

Proof. Take $\nu \in (0, \tau)$. Define $h(\lambda) = \|G_{\arg \min_{\theta} g_{t,\lambda}(\theta)} - \tau I\| - \nu$ for $g_{t,\lambda}(\theta)$ as in (9). Note that $\lim_{\lambda \rightarrow \infty} h(\lambda) = -\nu < 0$. If $h(0) < 0$ then we are done as for $\lambda_t = 0$, the corresponding learned parameters θ_t satisfy $\|G_{\theta_t} - \tau I\| < \nu$, leading to a provably convergent algorithm for $f \in \mathcal{F}_{\tilde{L}}$ for some $\tilde{L} > 0$. Else, suppose that $h(0) > 0$. Then as h is continuous in λ , there exists some λ such that $h(\lambda) < 0$. \square

At iteration t , to find a λ_t that is large enough to ensure convergence, we start at an initial point $\lambda = 10^{-6}$ and find $\phi \in \arg \min_{\theta} g_{t,\lambda}(\theta)$. If $\|G_{\phi} - \tau I\| < \tau$, then increase λ by a multiple and re-evaluate. Repeat until this inequality no longer holds, and take λ_t to be the most recent λ such that $\|G_{\phi} - \tau I\| < \tau$. Else if $\lambda = 10^{-6}$ and $\phi \in \arg \min_{\theta} g_{t,\lambda}(\theta)$ satisfies $\|G_{\phi} - \tau I\| > \tau$ then reduce λ by a multiple and re-evaluate until $\|G_{\phi} - \tau I\| < \tau$, then take $\lambda_t = \lambda$. For the (PS) parametrization we take the multiple to be 5, and for the (PP) parametrization, we take this multiple to be 2.

Proof of Theorem 3. Define $D = \max_{t=0,1,\dots} \{\|x_t - x^*\|\}$, which is finite as (x_t) is bounded. Due to the convexity of f and the Cauchy-Schwarz inequality, we have that

$$\begin{aligned} f(x_t) - f(x^*) &\leq \langle \nabla f(x_t), x_t - x^* \rangle \\ &\leq \|\nabla f(x_t)\| \|x_t - x^*\| \\ &\leq D \|\nabla f(x_t)\|. \end{aligned}$$

Therefore

$$\|\nabla f(x_t)\|^2 \geq \frac{1}{D^2} (f(x_t) - f(x^*))^2, \quad (52)$$

and for $t \geq T$ we have

$$\begin{aligned} f(x_{t+1}) &\leq f(x_t) - c(\nu, L, \tau) \|\nabla f(x_t)\|^2 \\ &\leq f(x_t) - \frac{c(\nu, L, \tau)}{D^2} (f(x_t) - f(x^*))^2. \end{aligned}$$

Denote $\Delta_t = f(x_{t+T}) - f(x^*)$, then in the spirit of (Nesterov et al., 2018), we have for all $t \geq 0$

$$\begin{aligned} \Delta_{t+1} &\leq \Delta_t - \frac{c}{D^2} \Delta_t^2 \\ \implies \frac{1}{\Delta_t} &\leq \frac{1}{\Delta_{t+1}} - \frac{c}{D^2} \frac{\Delta_t}{\Delta_{t+1}} \leq \frac{1}{\Delta_{t+1}} - \frac{c}{D^2} \\ \implies \frac{c}{D^2} + \frac{1}{\Delta_t} &\leq \frac{1}{\Delta_{t+1}}. \end{aligned}$$

Taking a summation gives

$$\begin{aligned} \sum_{k=0}^{t-1} \frac{c}{D^2} &\leq \sum_{k=0}^{t-1} \left(\frac{1}{\Delta_{k+1}} - \frac{1}{\Delta_k} \right) \\ \implies \frac{c}{D^2} t &\leq \frac{1}{\Delta_t} - \frac{1}{\Delta_0}. \end{aligned}$$

Therefore

$$\Delta_t \leq \frac{1}{\frac{1}{\Delta_0} + \frac{c}{D^2} t} = \frac{D^2 \Delta_0}{D^2 + c \Delta_0 t} \leq \frac{D^2 \Delta_0}{c \Delta_0 t} = \frac{D^2}{t},$$

as required. \square

C PROOFS FOR SECTION 5

- Proof of Lemma 2.**
1. For scalar step sizes, $G_\theta = \theta I$, take $\tilde{\theta} = \tau$.
 2. For a pointwise parametrization, $G_\theta x = \theta \odot x$, take $\tilde{\theta} = \tau \mathbf{1}$.
 3. For full operator parametrization, $G_\theta = \theta \in \mathcal{L}(\mathcal{X})$, take $\tilde{\theta} = \tau I$.
 4. For the convolutional parametrization, $G_\theta x = \theta * x$, take

$$\theta(i, j) = \begin{cases} \tau, & \text{if } i = j = 0, \\ 0, & \text{otherwise.} \end{cases} \quad (53)$$

G_θ are clearly continuous in θ for all listed parametrizations. □

Proof of Corollary 1. Apply lemma 2. □

Proof of Proposition 3. Because this problem is convex, if a solution θ is found by differentiating the objective function and equating equal to zero, this is a global minimizer. First, note that

$$\begin{aligned} f_k(x_k^t - B_k^t \theta) &= \frac{1}{2} \|A_k(x_k^t - B_k^t \theta) - y_k\|^2 \\ &= \frac{1}{2} \|A_k x_k^t - y_k\|^2 + \frac{1}{2} \| -A_k B_k^t \theta \|^2 + \langle -A_k B_k^t \theta, A_k x_k^t - y_k \rangle \\ &= \frac{1}{2} \|A_k x_k^t - y_k\|^2 + \frac{1}{2} \|A_k B_k^t \theta\|^2 - \langle \theta, (B_k^t)^* \nabla f_k(x_k^t) \rangle. \end{aligned}$$

Now,

$$\begin{aligned} \nabla_\theta \left\{ \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - B_k^t \theta) + \frac{\lambda_t}{2} \|\theta - \tilde{\theta}\|_2^2 \right\} \\ = \frac{1}{N} \sum_{k=1}^N (A_k B_k^t)^* (A_k B_k^t \theta) - (B_k^t)^* \nabla f_k(x_k^t) + \lambda_t (\theta - \tilde{\theta}) \end{aligned}$$

is equal to zero if and only if

$$\left(\frac{1}{N} \sum_{k=1}^N (A_k B_k^t)^* (A_k B_k^t) + \lambda_t I_\Theta \right) \theta = \lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla f_k(x_k^t).$$

□

A bonus proposition regarding the uniqueness of optimal parameters:

Proposition 7. $g_{t, \lambda_t}(\theta)$ has a unique global minimizer θ_t^* if at least one of the following are satisfied:

- $\lambda_t > 0$,
- f_k is twice continuously differentiable for $k \in \{1, \dots, N\}$, and there exists some $j \in \{1, \dots, N\}$ for which both B_j^t is injective and also f_j is μ_j -strongly convex.

Proof. **Case 1 - $\lambda_t > 0$**

$\frac{1}{N} \sum_{k=1}^N (A_k B_k^t)^* (A_k B_k^t)$ is self-adjoint and positive semi-definite as it is the sum of self-adjoint operators, $\frac{1}{N} \sum_{k=1}^N (A_k B_k^t)^* (A_k B_k^t) + \lambda_t I$ is a self-adjoint, positive-definite operator and therefore invertible.

Case 2 - $\lambda_t = 0$

If each f_k is twice continuously differentiable; then g_{t, λ_t} is twice continuously differentiable. It is then sufficient to show there exists $m > 0$ such that

$$\nabla^2 g_{t, \lambda_t}(\theta) \succeq mI, \quad (54)$$

for all θ , as this implies that g_{t,λ_t} is strongly convex and has a unique global minimizer. Note that

$$\nabla^2 g_{t,\lambda_t}(\theta) = \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla^2 f_k(x_k^t - B_k^t \theta) B_k^t. \quad (55)$$

Note that

$$\langle v, \nabla^2 g_{t,\lambda_t}(\theta) v \rangle = \langle v, \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla^2 f_k(x_k^t - B_k^t \theta) B_k^t v \rangle \quad (56)$$

$$= \frac{1}{N} \sum_{k=1}^N \langle v, (B_k^t)^* \nabla^2 f_k(x_k^t - B_k^t \theta) B_k^t v \rangle \quad (57)$$

$$= \frac{1}{N} \sum_{k=1}^N \langle B_k^t v, \nabla^2 f_k(x_k^t - B_k^t \theta) B_k^t v \rangle. \quad (58)$$

$$(59)$$

Each f_k is convex and so for all $v \in \mathcal{X}$,

$$\langle v, \nabla^2 f_k(x_k^t - B_k^t \theta) v \rangle \geq 0, \quad (60)$$

and f_j is μ_j -strongly convex, therefore

$$\langle v, \nabla^2 f_j(v_j^t - B_j^t \theta) v \rangle \geq \mu_j \|v\|^2. \quad (61)$$

For $v \in \mathcal{X}$,

$$\begin{aligned} \langle v, \nabla^2 g_{t,\lambda_t}(\theta) v \rangle &\geq \frac{1}{N} \mu_j v^T (B_j^t)^* B_j^t v \\ &\geq \left(\frac{1}{N} \mu_j \rho_{\min}^j \right) \|v\|^2, \end{aligned}$$

where ρ_{\min}^j is the minimum eigenvalue of $M_j^t = (B_j^t)^* B_j^t$ (a symmetric linear operator). Due to the symmetry of M_j^t , $\rho_{\min}^j \geq 0$ and is greater than zero if and only if B_j^t is injective. As B_j^t is injective, then $\rho_{\min}^j > 0$ and therefore $g_{t,\lambda_t}(\theta)$ is strongly-convex. \square

Proposition 7 applied to least-square functions:

Corollary 2. Uniqueness of optimal parameters in the least-squares case

When our f_k can be written as least-squares functions (15), then $g_{t,\lambda_t}(\theta)$ has a unique global minimizer θ_t^* if at least one of the following are satisfied:

- $\lambda_t > 0$,
- there exists some $j \in \{1, \dots, N\}$ for which both B_j^t and A_j are injective.

Proof. If A_j is injective then $A_j^* A_j$ is invertible which means that $f_j(x) = \frac{1}{2} \|A_j x - y^j\|^2$ is strongly convex. \square

Proposition 8. p_t given by

$$p_t = \left(\lambda_t I_{\Theta} + \frac{1}{N} \sum_{k=1}^N (\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)) \odot (A_k^* A_k) \right)^{\dagger} \left(\lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t) \odot \nabla f_k(x_k^t) \right) \quad (62)$$

is a solution to (9) with the pointwise parametrization $G_{p_t} x = p_t \odot x$ for any $x \in \mathcal{X}$.

Proof. Define for $x \in \mathcal{X}$,

$$B_k^t x = \nabla f_k(x_k^t) \odot x$$

then, for $x \in \mathcal{X}$, $(B_k^t)^*(x) = B_k^t(x)$. Now,

$$\begin{aligned} (A_k B_k^t)^*(A_k B_k^t)p &= (B_k^t)^*(A_k^* A_k B_k^t p) \\ &= \nabla f_k(x_k^t) \odot (A_k^* A_k (\nabla f_k(x_k^t) \odot p)). \end{aligned}$$

Now,

$$\begin{aligned} &[\nabla f_k(x_k^t) \odot (A_k^* A_k (\nabla f_k(x_k^t) \odot p))]_j \\ &= [\nabla f_k(x_k^t)]_j [A_k^* A_k (\nabla f_k(x_k^t) \odot p)]_j, \text{ by (7)} \\ &= [\nabla f_k(x_k^t)]_j \sum_{i=1}^n [\nabla f_k(x_k^t) \odot p]_i [A_k^* A_k]_{ji}, \text{ by (23)} \\ &= \sum_{i=1}^n [\nabla f_k(x_k^t)]_i [p]_i [\nabla f_k(x_k^t)]_j [A_k^* A_k]_{ji}, \text{ by (7)}. \end{aligned}$$

Secondly,

$$\begin{aligned} &[(\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)) \odot (A_k^* A_k)]_j p]_j \\ &= \sum_{i=1}^n [p]_i [(\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)) \odot (A_k^* A_k)]_{ji}, \text{ by (23)} \\ &= \sum_{i=1}^n [p]_i [\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)]_{ji} [A_k^* A_k]_{ji}, \text{ by (19)} \\ &= \sum_{i=1}^n [p]_i [\nabla f_k(x_k^t)]_j [\nabla f_k(x_k^t)]_i [A_k^* A_k]_{ji} \text{ by (22)} \\ &= [\nabla f_k(x_k^t) \odot (A_k^* A_k (\nabla f_k(x_k^t) \odot p))]_j, \text{ by (21)}. \end{aligned}$$

Finally,

$$\lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla f_k(x_k^t) = \lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t) \odot \nabla f_k(x_k^t).$$

Then the result follows from proposition 3. \square

Proposition 9. Let $B_k^t : \mathcal{L}(\mathcal{X}) \rightarrow \mathcal{X}$ be such that for any linear operator $P \in \mathcal{L}(\mathcal{X})$, we have $B_k^t(P) = P \nabla f_k(x_k^t)$. Then its adjoint $(B_k^t)^* : \mathcal{X} \rightarrow \mathcal{L}(\mathcal{X})$ is given by

$$(B_k^t)^*(w) = \nabla f_k(x_k^t) \otimes w, \quad (63)$$

for any element $w \in \mathcal{X}$. Then

$$\theta_t = \left(\lambda_t I_{\Theta} + \frac{1}{N} \sum_{k=1}^N (A_k^* A_k) \otimes (\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)) \right)^{\dagger} \left(\lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t) \right), \quad (64)$$

(16) is a solution to (9) for the full operator parametrization.

Proof. $\theta_t \in \mathcal{L}(\mathcal{X})$ and we require $\theta_t \nabla f_k(x_k^t) = B_k^t(\theta_t)$, so can take $B_k^t(\theta_t) = \theta_t \nabla f_k(x_k^t)$. For the adjoint,

$$\langle B_k^t(P), w \rangle = \sum_{i=1}^n [P \nabla f_k(x_k^t)]_i [w]_i \quad (65)$$

$$= \sum_{i=1}^n \sum_{j=1}^n [P]_{ij} [\nabla f_k(x_k^t)]_j [w]_i \quad (66)$$

$$= \langle P, (B_k^t)^* w \rangle \quad (67)$$

$$= \sum_{i=1}^n \sum_{j=1}^n [P]_{ij} [(B_k^t)^* w]_{ij}, \quad (68)$$

and therefore $[(B_k^t)^* w]_{ij} = w_i [\nabla f_k(x_k^t)]_j$, which means $(B_k^t)^*(w) = w \otimes \nabla f_k(x_k^t)$. Now,

$$\begin{aligned} [(A_k B_k^t)^* (A_k B_k^t) \theta]_{ij} &= [(B_k^t)^* (A_k^* A_k B_k^t \theta)]_{ij} \\ &= [(A_k^* A_k B_k^t \theta) \otimes \nabla f_k(x_k^t)]_{ij} \\ &= [\nabla f_k(x_k^t)]_j [A_k^* A_k B_k^t \theta]_i, \text{ by (21)} \\ &= [\nabla f_k(x_k^t)]_j \sum_{q=1}^n [A_k^* A_k]_{iq} [B_k^t \theta]_q \\ &= [\nabla f_k(x_k^t)]_j \sum_{q=1}^n [A_k^* A_k]_{iq} \sum_{\ell=1}^n [\theta]_{q\ell} [\nabla f_k(x_k^t)]_\ell, \text{ (definition of } B_k^t). \end{aligned}$$

Similarly,

$$\begin{aligned} & [((A_k^* A_k) \otimes (\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t))) \theta]_{ij} \\ &= \sum_{q,\ell=1}^n [(A_k^* A_k) \otimes (\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t))]_{ij,q\ell} [\theta]_{q\ell} \\ &= \sum_{q,\ell=1}^n [A_k^* A_k]_{iq} [\nabla f_k(x_k^t) \otimes \nabla f_k(x_k^t)]_{j\ell} [\theta]_{q\ell}, \text{ by (22)} \\ &= \sum_{q,\ell=1}^n [A_k^* A_k]_{iq} [\nabla f_k(x_k^t)]_j [\nabla f_k(x_k^t)]_\ell [\theta]_{q\ell} \\ &= [(A_k B_k^t)^* (A_k B_k^t) \theta]_{ij}, \end{aligned}$$

as required, due to $[A_k^* A_k]_{jq} = [A_k^* A_k]_{qj}$. □

Proposition 10. *If each f_k can be written as a least-squares function (15), then α_t can be given as*

$$\alpha_t = \frac{\lambda_t \tilde{\theta} + \frac{1}{N} \sum_{k=1}^N \|\nabla f_k(x_k^t)\|^2}{\lambda_t + \frac{1}{N} \sum_{k=1}^N \|A_k \nabla f_k(x_k^t)\|^2}, \quad (69)$$

if $\lambda_t > 0$ or $A_j \nabla f_j(x_j^t) \neq \underline{0}$ for some $j \in \{1, \dots, N\}$.

Proof. Take $B_k^t : \mathbb{R} \rightarrow \mathcal{X}$ such that

$$B_k^t(\alpha) = \alpha \nabla f_k(x_k^t).$$

Then for $\alpha \in \mathbb{R}$

$$\langle B_k^t(\alpha), w \rangle = \langle \alpha \nabla f_k(x_k^t), w \rangle = \alpha \langle \nabla f_k(x_k^t), w \rangle.$$

Therefore

$$(B_k^t)^*(w) = \langle \nabla f_k(x_k^t), w \rangle. \quad (70)$$

then general formula 3 gives the desired result as

$$\begin{aligned} (B_k^t)^*(A_k^* A_k B_k^t(\alpha)) &= \alpha \langle \nabla f_k(x_k^t), A_k^* A_k \nabla f_k(x_k^t) \rangle = \alpha \|A_k \nabla f_k(x_k^t)\|^2, \\ (B_k^t)^*(\nabla f_k(x_k^t)) &= \langle \nabla f_k(x_k^t), \nabla f_k(x_k^t) \rangle = \|\nabla f_k(x_k^t)\|^2. \end{aligned}$$

Then the result follows from proposition 3. \square

Proposition 11. For $n_1, n_2 \in \mathbb{N}$, let $\mathcal{X} = \mathbb{R}^{n_1 \times n_2}$. Define $B_k^t : \mathcal{X} \rightarrow \mathcal{X}$ be such that for any convolutional kernel $\kappa \in \mathcal{X}$, we have $B_k^t(\kappa) = \kappa * \nabla f_k(x_k^t)$. Then its adjoint $(B_k^t)^* : \mathcal{X} \rightarrow \mathcal{X}$ is given by

$$(B_k^t)^*(w) = w * \overline{\nabla f_k(x_k^t)}, \quad (71)$$

where for $x \in \mathcal{X}$,

$$\bar{x}(k, l) = x(-k, -l). \quad (72)$$

Proof. For the adjoint of B_k^t , we have

$$\langle B_k^t(\kappa), w \rangle = \langle \kappa * \nabla f_k(x_k^t), w \rangle \quad (73)$$

$$= \sum_{i,j} [\kappa * \nabla f_k(x_k^t)](i, j) w(i, j) \quad (74)$$

$$= \sum_{i,j} \sum_{k,l} \kappa(k, l) [\nabla f_k(x_k^t)](i - k, j - l) w(i, j) \quad (75)$$

$$= \sum_{i,j} \sum_{k,l} \kappa(k, l) [\nabla f_k(x_k^t)](i, j) w(i + k, j + l) \quad (76)$$

$$= \sum_{i,j} \sum_{k,l} \kappa(k, l) [\nabla f_k(x_k^t)](i, j) w(i + k, j + l) \quad (77)$$

$$= \sum_{i,j} [\nabla f_k(x_k^t)](i, j) \left(\sum_{k,l} \kappa(k, l) w(i + k, j + l) \right) \quad (78)$$

$$= \sum_{i,j} [\nabla f_k(x_k^t)](i, j) \left(\sum_{k,l} \kappa(-k, -l) w(i - k, j - l) \right) \quad (79)$$

$$= \langle \nabla f_k(x_k^t), \bar{\kappa} * w \rangle, \quad (80)$$

where $\bar{\kappa}(k, l) = \kappa(-k, -l)$. \square

C.1 APPROXIMATING OPTIMAL LINEAR PARAMETERS

For general functions f_k , a closed-form solution does not exist for calculating linear parameters. Instead, we require an optimization algorithm to approximate these quantities. With information of $\nabla g_{t, \lambda_t}(\theta)$, and $L_{g_{t, \lambda_t}}$, the Lipschitz constant of $\nabla g_{t, \lambda_t}(\theta)$, one can use any first-order convex optimization algorithm, such as gradient descent, Nesterov accelerated gradient (Nesterov et al., 2018), or stochastic optimization methods such as SGD, and SVRG (Gower et al., 2020) (especially for large N , due to both speed and memory considerations) to approximate θ_t^* . For example, one can start at an initial point θ_t^0 at iteration t and update via gradient descent

$$\theta_t^{w+1} = \theta_t^w - \frac{1}{L_{g_{t, \lambda_t}}} \nabla g_{t, \lambda_t}(\theta_t^w). \quad (81)$$

The following result illustrates how $\nabla g_{t, \lambda_t}(\theta)$ and $L_{g_{t, \lambda_t}}$ can be calculated.

Proposition 12. For a general linear parametrization G , the gradient of g_{t,λ_t} with respect to θ and its associated Lipschitz constant can be calculated as

$$\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla f_k(x_k^t - G_\theta \nabla f_k(x_k^t)), \quad (82)$$

$$L_{g_{t,\lambda_t}} = \lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \|B_k^t\|^2. \quad (83)$$

Proof. As

$$\begin{aligned} g_{t,\lambda_t}(\theta) &= \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - G_\theta \nabla f_k(x_k^t)) + \frac{\lambda_t}{2} \|\theta - \tilde{\theta}\|^2 \\ &= \frac{1}{N} \sum_{k=1}^N f_k(x_k^t - B_k^t \theta) + \frac{\lambda_t}{2} \|\theta - \tilde{\theta}\|^2, \end{aligned}$$

then by the chain rule

$$\nabla g_{t,\lambda_t}(\theta) = -\frac{1}{N} \sum_{k=1}^N (B_k^t)^* \nabla f_k(x_k^t - B_k^t \theta) + \lambda_t(\theta - \tilde{\theta}), \quad (84)$$

as required. To calculate the smoothness constant, we have

$$\begin{aligned} &\|\nabla g_{t,\lambda_t}(\theta_1) - \nabla g_{t,\lambda_t}(\theta_2)\| \\ &= \left\| \lambda_t(\theta_1 - \theta_2) + \frac{1}{N} \sum_{k=1}^N (B_k^t)^* (\nabla f_k(x_k^t - B_k^t \theta_2) - \nabla f_k(x_k^t - B_k^t \theta_1)) \right\| \\ &\leq \lambda_t \|\theta_1 - \theta_2\| + \frac{1}{N} \sum_{k=1}^N \left\| (B_k^t)^* (\nabla f_k(x_k^t - B_k^t \theta_2) - \nabla f_k(x_k^t - B_k^t \theta_1)) \right\| \\ &\leq \lambda_t \|\theta_1 - \theta_2\| + \frac{1}{N} \sum_{k=1}^N \|B_k^t\| \|\nabla f_k(x_k^t - B_k^t \theta_2) - \nabla f_k(x_k^t - B_k^t \theta_1)\| \\ &\leq \lambda_t \|\theta_1 - \theta_2\| + \frac{1}{N} \sum_{k=1}^N L_k \|B_k^t\| \|B_k^t(\theta_1 - \theta_2)\| \\ &\leq \left(\lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \|B_k^t\|^2 \right) \|\theta_1 - \theta_2\| \end{aligned}$$

Due to the properties of the triangle inequality, the Cauchy-Schwarz inequality, and the operator norm, this bound is tight. Therefore the Lipschitz constant of $\nabla g_{t,\lambda_t}(\theta)$ is given by

$$\lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \|B_k^t\|^2 \quad (85)$$

as required. \square

Using this general result, we can calculate these values for specific parametrizations of G .

Corollary 3. Suppose each $f_k \in \mathcal{F}_{L_k}$.

Pointwise parametrization

For the pointwise parametrization, $\theta \in \mathcal{X}$, and

$$\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - \theta \odot \nabla f_k(x_k^t)) \odot \nabla f_k(x_k^t), \quad (86)$$

and an upper bound of the Lipschitz constant of $\nabla_{\theta}g$ is given by

$$L_{\nabla_{\theta}g} = \lambda_t + \frac{1}{N} \sum_{k=1}^N L_k (\max\{|\nabla f_k(x_k^t)|_1, \dots, |\nabla f_k(x_k^t)|_n\})^2. \quad (87)$$

Full operator parametrization

In this case we have $\theta \in \mathcal{L}(\mathcal{X})$. The gradient of $g_{t,\lambda_t}(\theta)$ is given by

$$\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - \theta \nabla f_k(x_k^t)) \otimes \nabla f_k(x_k^t), \quad (88)$$

and an upper bound of the Lipschitz constant of $\nabla g_{t,\lambda_t}(\theta)$ is given by

$$\lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \|\nabla f_k(x_k^t)\|^2. \quad (89)$$

Scalar step size

We now take $\theta \in \mathbb{R}$. The derivative of g_{t,λ_t} with respect to θ is given by

$$g'_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \langle \nabla f_k(x_k^t - \theta \nabla f_k(x_k^t)), \nabla f_k(x_k^t) \rangle, \quad (90)$$

and the Lipschitz constant of $g'(\theta)$ is given by

$$\lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \|\nabla f_k(x_k^t)\|^2. \quad (91)$$

Convolution

In this case we have $\theta \in \mathbb{R}^{n_1 \times n_2}$. The gradient of $g_{t,\lambda_t}(\theta)$ is given by

$$\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - G_{\theta} \nabla f_k(x_k^t)) * \overline{\nabla f_k(x_k^t)}. \quad (92)$$

Proof. **Pointwise parametrization**

In this case, we have $\theta \in \mathcal{X}$ and $B_k^t(x) = \nabla f_k(x_k^t) \odot x$ and $(B_k^t)^*(x) = B_k^t x$ for $x \in \mathcal{X}$. Furthermore,

$$\begin{aligned} \|B_k^t\| &= \max_{x \neq 0} \frac{\|x \odot \nabla f_k(x_k^t)\|}{\|x\|} = \max_{x \neq 0} \sqrt{\frac{\sum_{i=1}^n [x]_i^2 [\nabla f_k(x_k^t)]_i^2}{\sum_{i=1}^n [x]_i^2}} \\ &\leq \max_q \|\nabla f_k(x_k^t)\|_q \max_{x \neq 0} \sqrt{\frac{\sum_{i=1}^n [x]_i^2}{\sum_{i=1}^n [x]_i^2}} = \max\{|\nabla f_k(x_k^t)|_1, \dots, |\nabla f_k(x_k^t)|_n\}. \end{aligned}$$

Full operator parametrization

In the case of the full operator parametrization, we have $(B_k^t)^*(w) = w \otimes \nabla f_k(x_k^t)$. Therefore, using Proposition 12 gives (88). For the Lipschitz constant, note that

$$\|B_k^t(P)\| = \|P \nabla f_k(x_k^t)\| \leq \|P\| \|\nabla f_k(x_k^t)\|,$$

and therefore

$$\|B_k^t\| = \max_{P \neq 0} \frac{\|B_k^t(P)\|}{\|P\|} \leq \|\nabla f_k(x_k^t)\|.$$

Scalar step size

Let B_k^t be defined for any $\alpha \in \mathbb{R}$ by $B_k^t(\alpha) = \alpha \nabla f_k(x_k^t)$, then for an element $w \in \mathcal{X}$, $(B_k^t)^*(w) = \langle \nabla f_k(x_k^t), w \rangle$. Furthermore,

$$\begin{aligned} \|B_k^t(\alpha)\| &= \|\alpha \nabla f_k(x_k^t)\| \\ &= |\alpha| \|\nabla f_k(x_k^t)\|, \end{aligned}$$

and so

$$\|B_k^t\| = \max_{\alpha \neq 0} \frac{\|B_k^t(\alpha)\|}{|\alpha|} = \|\nabla f_k(x_k^t)\|.$$

Convolution

For the gradient,

$$(B_k^t)^*(\nabla f_k(x_k^t) - G_\theta \nabla f_k(x_k^t)) = \nabla f_k(x_k^t) - G_\theta \nabla f_k(x_k^t) * \overline{\nabla f_k(x_k^t)}.$$

□

For any chosen linear parametrization, one can approximate the operator norm of B_k^t using the power method (Golub & Van Loan, 2013). The following table summarises the previous propositions:

Table 2: Example parametrization properties

Parametrization	Equations
Pointwise	<ul style="list-style-type: none"> • $\tilde{\theta} = \tau \mathbf{1} \in \mathcal{X}$ • $B_k^t(x) = \nabla f_k(x_k^t) \odot x$, $(B_k^t)^*(x) = B_k^t(x)$ • $\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - \theta \odot \nabla f_k(x_k^t)) \odot \nabla f_k(x_k^t)$ • $L_{g_{t,\lambda_t}} \leq \lambda_t + \frac{1}{N} \sum_{k=1}^N L_k (\max\{ \nabla f_k(x_k^t)_1 , \dots, \nabla f_k(x_k^t)_n \})^2$
Full operator	<ul style="list-style-type: none"> • $\tilde{\theta} = \tau I \in \mathcal{L}(\mathcal{X})$ • $B_k^t(P) = P \nabla f_k(x_k^t)$, $(B_k^t)^*(w) = w \otimes \nabla f_k(x_k^t)$ • $\nabla g_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - \theta \nabla f_k(x_k^t)) \otimes \nabla f_k(x_k^t)$ • $L_{g_{t,\lambda_t}} \leq \lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \ \nabla f_k(x_k^t)\ ^2$
Scalar	<ul style="list-style-type: none"> • $\tilde{\theta} = \tau \in \mathbb{R}$ • $B_k^t(\alpha) = \alpha \nabla f_k(x_k^t)$, $(B_k^t)^*(w) = \langle w, \nabla f_k(x_k^t) \rangle$ • $g'_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \langle \nabla f_k(x_k^t - \theta \nabla f_k(x_k^t)), \nabla f_k(x_k^t) \rangle$ • $L_{g_{t,\lambda_t}} = \lambda_t + \frac{1}{N} \sum_{k=1}^N L_k \ \nabla f_k(x_k^t)\ ^2$
Convolution	<ul style="list-style-type: none"> • $\tilde{\theta}(i, j) = \begin{cases} \tau, & \text{if } i = j = 0, \\ 0, & \text{otherwise.} \end{cases}$ • $B_k^t(\kappa) = \kappa * \nabla f_k(x_k^t)$, $(B_k^t)^*(\kappa) = \kappa * \overline{\nabla f_k(x_k^t)}$ • $g'_{t,\lambda_t}(\theta) = \lambda_t(\theta - \tilde{\theta}) - \frac{1}{N} \sum_{k=1}^N \nabla f_k(x_k^t - \theta * \nabla f_k(x_k^t)) * \overline{\nabla f_k(x_k^t)}$

D ADDITIONAL NUMERICAL RESULTS**D.1 BEST AND WORST PERFORMANCE OF LEARNED CONVOLUTION**

In this section, we compare the performance of the learned algorithm on individual functions in the test dataset (of 100 functions). Figure 8 shows that for the deblurring problem best performance of the learned convolutional algorithm take less than one quarter of the iterations as NAG to reach a tolerance of 10^{-7} , and the worst performance takes approximately one third the iterations. Therefore, across all 100 functions in the test set the learned convolutional algorithm performs significantly better and quite similarly for different functions.




Figure 8: Best and worst case convergence of the learned convolutional algorithm compared to NAG for the deblurring problem, considering the ratio of number of iterations to reach a tolerance of 10^{-7} .

Figure 9 shows that for the CT problem, the worst performance of the learned convolutional algorithm takes less than one third of the iterations as NAG to reach a tolerance of 10^{-9} . The best case reaches a tolerance of 10^{-9} in about one fifth of the iterations. However, a tolerance of 10^{-5} is a sufficient tolerance for reconstruction quality and is achieved in only 4 iterations with the learned method, whereas NAG requires 28 iterations. Thus, for both of the problems, we achieve superior performance with the learned convolutional algorithm for every single function in both of the test sets. We also see similar speedups for both problems for all functions when using the learned algorithm with a convolutional parametrization.




Figure 9: Best and worst case convergence of the learned convolutional algorithm compared to NAG for the CT problem, considering the ratio of number of iterations to reach a tolerance of 10^{-9} .

D.2 ABLATION STUDY: SIZE OF LEARNED KERNELS

Figure 10 shows the many of the learned convolutional algorithms outperform NAG for the deblurring problem. The 3×3 learned kernels outperform NAG initially but are eventually overtaken before a tolerance of 10^{-7} . We see that the 5×5 kernels significantly outperform the NAG kernels, with then incremental improvements with larger kernel sizes as the 96×96 kernels achieve a tolerance of 10^{-8} is half the iterations as the 5×5 kernels, while the 3×3 kernels achieve of tolerance of 10^{-8} in approximately 900 iterations, while the 5×5 kernels achieve this tolerance in about 500 iterations, compared to just over 200 iterations for the 96×96 kernels.




Figure 10: Test performance of different kernel sizes in the convolutional parametrization, averaged over the test dataset in the deblurring problem. Tested kernel sizes are 3×3 , 5×5 , 7×7 , 11×11 , 3×3 , 96×96 .

D.3 TOLERANCE TABLES

Table 3: The first row shows error thresholds for the deblurring problem. The entries in the table show the number of required iterations to fall below the respective error threshold. "na" means that the threshold was not reached within 2000 iterations.

	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	10^{-7}
Convolution	1	2	14	40	72	115	176
L-BFGS	3	10	39	104	220	384	565
NAG	4	15	56	137	266	432	647
Backtracking Line Search	2	12	105	553	1970	na	na
Scalar	3	17	173	930	na	na	na
Pointwise	2	14	190	1368	na	na	na
GD	5	32	346	1758	na	na	na

Table 4: The first row shows error thresholds for the deblurring problem. The entries in the table show the number of required iterations to fall below the respective error threshold. "na" means that the threshold was not reached within 1000 iterations. The convolutional method clearly performs best

	10^{-3}	10^{-4}	10^{-5}	10^{-6}	10^{-7}	10^{-8}	10^{-9}	10^{-10}
Convolution	1	2	4	5	8	14	22	29
L-BFGS	4	8	15	26	39	55	71	83
NAG	5	11	19	33	49	71	110	141
Backtracking Line Search	5	9	21	47	87	141	201	na
Pointwise	2	8	22	56	113	189	na	na
Scalar	3	8	23	56	111	184	na	na
GD	6	19	52	129	na	na	na	na