

FINITE GROUPS WITH GEODETIC CAYLEY GRAPHS

MURRAY ELDER, ADAM PIGGOTT, FLORIAN STOBER, ALEXANDER THUMM, AND ARMIN WEISS

ABSTRACT. A connected undirected graph is called *geodetic* if for every pair of vertices there is a unique shortest path connecting them. It has been conjectured that for finite groups, the only geodetic Cayley graphs which occur are odd cycles and complete graphs. In this article we present a series of theoretical results which contribute to a computer search verifying this conjecture for all groups of size up to 1024. The conjecture is also verified theoretically for several infinite families of groups including dihedral and some families of nilpotent groups. Two key results which enable the computer search to reach as far as it does are: if the center of a group has even order, then the conjecture holds (this eliminates all 2-groups from our computer search); if a Cayley graph is geodetic then there are bounds relating the size of the group, generating set and center (which cuts down the number of generating sets which must be searched significantly).

1. INTRODUCTION

Cayley graphs of groups form an important subclass of the vertex-transitive and regular graphs. The undirected Cayley graph of a group G with respect to a generating set Σ is the connected graph on the set of vertices $V = G$ where $\{g, h\} \in E$ if and only if there is a generator $a \in \Sigma$ such that $ga = h$. Besides being an important tool in combinatorial group theory, there are also interesting graph-theoretic questions about Cayley graphs. One example which has been much studied is the longstanding conjecture that every finite undirected Cayley graph has a Hamiltonian cycle (see for example [25]).

A connected undirected graph $\Gamma = (V, E)$ is called *geodetic* if for any pair $u, v \in V$, the shortest path from u to v is unique. Research on geodetic graphs began in 1962, when Ore posed the problem of classifying all such graphs [24]. This goal has been achieved so far for planar geodetic graphs, and geodetic graphs of diameter two [33, 32]; yet, after decades of active research, a full classification of finite geodetic graphs has not been attained (for some recent developments, see for example [14, 17, 34]).

In 1997, Shapiro [31] asked whether each finitely generated group that admits a geodetic Cayley graph with respect to some finite generating set is *plain*, that is, isomorphic to the free product of finitely many finite groups and finitely many copies of \mathbb{Z} . It is well-known and not hard to see that the converse holds: each plain group admits a geodetic Cayley graph (with respect to the generating set consisting of each non-trivial element of each finite factor, and a cyclic generator for each \mathbb{Z} factor). Recently, significant progress has been made on this question [11] and variants of it [10, 12, 13] (see also the paragraph below on related work).

For a finite group, one may consider a generating set consisting of every non-trivial element. Then the corresponding Cayley graph is the complete graph, which is geodetic. For cyclic groups of odd order, there is a second possibility: taking an arbitrary single generator, the Cayley graph is an odd cycle, which is also geodetic. The question we study here is whether there is any other possibility. In a 2017 PhD thesis [15], Federici conjectured there is none.

Conjecture A ([15, Conjecture 6]). *Every finite geodetic Cayley graph is either a cycle of odd length or a complete graph.*

In this paper we report on a systematic computer experiment which confirms the conjecture for a significant number of groups.

Theorem B. *All groups up to size 1024 satisfy Conjecture A.*

In fact, we can go even further than that by showing that all groups of even order up to 2014, all simple groups of order below 5000, and the symmetric group S_7 satisfy Conjecture A. Given that there are

Date: June 4, 2024.

2020 Mathematics Subject Classification. 05C12, 05C25, 20F05.

Key words and phrases. geodetic graph, geodetic group, finite group, Cayley graph, group center.

approximately 49.5 billion groups of order at most 1024, and each has exponentially many different generating sets, a naive computer search could not possibly achieve the result in Theorem B. Instead, our computer search relies on a series of theoretical results concerning finite groups and when they can admit geodesic Cayley graphs. Some of these results confirm the conjecture for infinite families of groups, as summarized by the following.

Theorem C. *A finite group G satisfies Conjecture A if any one of the following conditions holds.*

- (1) *The center $Z(G)$ of G has even order (Theorem 3.14).*
- (2) *The group G contains an abelian subgroup of index two (Theorem 5.4).*
- (3) *The group G is nilpotent and its order does not have certain small divisors depending on the nilpotency class of G (see Theorem 5.9 for the precise statement).*
- (4) *The group G has large commutativity degree (Theorem 5.17).*

This cuts down the number of groups which need to be considered enormously, from 49.5 billion to 3197. In fact, just excluding abelian groups and groups with even-order center leaves 4734 groups of order at most 1024. We note that Conjecture A was shown to be satisfied by all abelian groups by Georgakopoulos as reported in [15, Proposition 10].

A second key theoretical result is the following, which provides an upper bound on the number of generating sets that the search must consider for each group.

Theorem D. *Let G be a finite group with generating set Σ such that the Cayley graph $\text{Cay}(G, \Sigma)$ is geodesic and neither complete nor a cycle. Then $|\Sigma| < 0.845\sqrt{|G|}$.*

Article organization. Theorems C and D are proved via a series of results presented in Sections 3 to 5. In Section 6 we give details of the implementation of our computer program. We use GAP [18] and its small group (SmallGrp) library [2] to enumerate the groups and check which of them are already covered by Theorem C. Then we apply an exhaustive search to check whether there is any geodesic generating set for any of the remaining groups. This exhaustive search crucially relies on several pruning methods based on variants of Theorem D and other theoretical results developed in Sections 3 and 4. The code is available at https://osf.io/9ay6s/?view_only=37e18301e4e74e12bfe4e07b90b924c0.

Related work. As noted above, the first mention of Conjecture A that we are aware of is in the 2017 thesis of Federici [15], where the conjecture is proved for abelian groups and where it is shown that the Cayley graph of a semidirect product $C_m \rtimes C_n$ with C_n acting faithfully on C_m with respect to the “standard” generating set is not geodesic. Notice that for both m and n sufficiently large, this class of groups turns out to be one of the most difficult cases for our computer search (see Section 6.5). Besides this result, Federici proves some helpful lemmas for general geodesic Cayley graphs and runs computer experiments (unfortunately undocumented) which do not find any geodesic Cayley graphs except the obvious ones.

In 2022 Che [5] programmed an exhaustive computer search as part of an undergraduate project supervised by Alexey Talambutsa establishing Conjecture A for subgroups of the symmetric group S_4 . Moreover, for

- all subgroups of S_6 with generating sets of size five,
- all subgroups of S_7 with generating sets of size four,
- all subgroups of S_9 with generating sets of size three,
- all subgroups of S_{10} with generating sets of size two,

Che showed that none of the corresponding Cayley graphs except complete ones and odd cycles are geodesic. For the source code, see <https://gitlab.com/andr0901/kayley-geodesics>. Be aware that restricting the number of generators to five or less is a strong restriction. (Note that by contrast, Theorem B verifies all generating sets for S_6 , and Theorem 6.6 below verifies all generating sets for S_7 . The groups S_9 and S_{10} are beyond our scope.)

For the case of infinite groups, Gardam, Spriano, Townsend and the first two authors have proved that if a quasi-transitive locally-finite graph is geodesic, then it is quasi-isometric to a tree [11]. This implies that the only groups that can have geodesic Cayley graphs are the virtually free groups. Moreover, they show that the class of groups admitting geodesic Cayley graphs are a strict subclass of virtually free ([11, Theorem B]), which almost resolves Shapiro’s question. To make further progress it will be useful to understand finite geodesic Cayley graphs as, if Conjecture A is false and some interesting (family of) counterexamples can be obtained, these might be used to construct non-trivial infinite geodesic Cayley graphs for virtually free groups that are not plain groups.

A crucial role for the study of infinite geodetic Cayley graphs is played by so-called *inverse-closed finite confluent length-reducing rewriting systems* (icfclrrs). Indeed, it is an easy exercise to show that a group presented by an icfclrrs has a geodetic Cayley graph. For finite groups the converse is also straightforward; hence, in this case the two properties are equivalent. Recently, as a corollary of the above-mentioned result, this equivalence has been shown to be true for all finitely generated groups [11, Theorem C] – yielding a connection between a rather geometric property (being geodetic) and another coming from formal languages (being presented by an icfclrrs). Earlier, Eisenberg and the second author [10] solved a variant of a conjecture of Madlener and Otto [22, p. 143] (known as Gilman’s conjecture) by showing that all groups presented by a monadic (all right-hand sides are of length one) finite confluent length-reducing rewriting system are plain. Furthermore, [12] shows that groups presented by icfclrrs with left-hand sides of length at most three are also plain. In [26, Corollary 3.13] it is shown that the finite cyclic groups of order at least three are the only finite groups admitting more than one “normalized” finite confluent monadic rewriting system, and those groups admit exactly two such rewriting systems.

In [13] Townsend and the first two authors generalize the concept of a geodetic graph to graphs which have at most k different geodesics between any pair of vertices for some constant k , which they call k -geodetic, and study properties of groups which admit k -geodetic Cayley graphs. While the main focus in [13] is on infinite groups, [13, Example 1.1] gives examples of Cayley graphs of finite groups that are k -geodetic but not $(k - 1)$ -geodetic or complete for $k \geq 2$. The easiest such examples are cyclic groups of even order with a non-complete 2-geodetic Cayley graph as well as a complete bipartite Cayley graph.

A graph is called *strongly geodetic* if for every pair of vertices there is at most *one* non-backtracking path connecting them that has length at most the diameter of the graph [3]. Clearly, all strongly geodetic graphs are geodetic. By [3, Theorem 1] the class of strongly geodetic regular graphs coincides with the class of so-called *Moore graphs*, which also have been thoroughly studied in graph theory (for a definition see [20] or the survey paper [23]). Interestingly, for diameter two the Moore graphs coincide exactly with the regular geodetic graphs [32]. Moreover, by [1, 7] the Moore graphs are completely classified: cycles of odd length, complete graphs, the Petersen graph, Hoffman-Singleton graph, and potentially hypothesized graphs (not known to exist) with 3250 vertices and degree 57. Of these, only the cycles and the complete graphs are Cayley graphs [4, 23] (we provide an alternative proof of this fact in Proposition 3.17 below). Hence, Conjecture A is true if we replace “geodetic” with “strongly geodetic”. On the other hand, Cayley graphs provide examples that are “close” to being Moore graphs [9].

2. PRELIMINARIES

2.1. Groups and words. Throughout this article, we only consider finite groups. We write these multiplicatively, mostly omitting the operator altogether, and denote their neutral element by 1. As usual, we write $[g, h] = g^{-1}h^{-1}gh$ for the *commutator* of two elements g, h of a group G . The *center* of G is the subgroup $Z(G) = \{g \in G \mid [g, h] = 1 \text{ for all } h \in G\}$. Two elements $g, g' \in G$ are *conjugate* if $g' = h^{-1}gh$ ($=: g^h$) for some $h \in G$. Given $g \in G$ and $H \subseteq G$, we write $g^H = \{g^h \mid h \in H\}$.

We denote by $\text{ord}(g)$ the *order* of $g \in G$, i.e., the smallest positive integer n with $g^n = 1$. The *order* (number of elements) of the group G is denoted by $|G|$, and the *exponent* of G by $\text{exp}(G)$, i.e., the smallest positive integer n such that $g^n = 1$ for all $g \in G$. Our notation is similar to [29].

For a subset $X \subseteq G$, we write $\langle X \rangle$ for the subgroup generated by X . It consists of those group elements that can be written as words over the alphabet $\Sigma = X \cup X^{-1}$. We denote the set of all such words by Σ^* . Given words $v, w \in \Sigma^*$, we write $v = w$ with the meaning that v and w evaluate to the same group element in G , whereas $v \equiv w$ denotes equality of words.

A subset $\Sigma \subseteq G$ with $\langle \Sigma \rangle = G$ is called a *generating set* of G . Throughout, we assume that all generating sets satisfy $1 \notin \Sigma$ and are *symmetric*, i.e., $a \in \Sigma$ implies $a^{-1} \in \Sigma$. We sometimes represent the inverse a^{-1} of a generator $a \in \Sigma$ by \bar{a} to emphasize that it is a single letter.

The length of a word $w = a_1 \cdots a_n \in \Sigma^*$ (with $a_i \in \Sigma$) is denoted by $|w| = n$. We denote the set of words over Σ of length n by Σ^n . A word $w \in \Sigma^*$ is called a *geodesic* for (or representing) a group element $g \in G$ if $w = g$ and w is shortest among all words with that property. The geodesic length of $g \in G$ is defined as the length of a geodesic word representing g . If g admits a unique geodesic, we denote its geodesic by $\text{geod}(g)$.

We use the following notation for specific groups. C_n is the cyclic group of order n , D_{2n} the dihedral group of order $2n$, and S_n (resp. A_n) the symmetric (resp. alternating) group on n elements. A direct product is

denoted by \times and a *semidirect product* by \rtimes , where $G = N \rtimes H$ means that N is a normal subgroup of G and H a subgroup of G such that $G = NH$ and $N \cap H = \{1\}$. Notice that $G/N = H$ in this case. Be aware that writing $G = N \rtimes H$ does not completely define G – instead we need to also specify a homomorphism from H to the automorphism group of N describing the action $(h, n) \mapsto hnh^{-1}$ of H on N (in other words H acts on N via automorphisms).

2.2. Graphs and Cayley graphs. We consider only *undirected finite simple graphs* $\Gamma = (V, E)$ where $E \subseteq \binom{V}{2}$. The *Cayley graph* $\text{Cay}(G, \Sigma) = (V, E)$ of a group G (with respect to a generating set $\Sigma \subseteq G$) is defined by $V = G$ and $E = \{\{g, ga\} \mid g \in G, a \in \Sigma\}$. In other literature sometimes the *directed* Cayley graph is considered; however, throughout this paper we only consider the undirected graph defined as above. Moreover, we always assume that $1 \notin \Sigma$; hence, $\text{Cay}(G, \Sigma)$ contains no loops.

The Cayley graph is connected because Σ generates G . Note also that G acts on $\text{Cay}(G, \Sigma)$ by left translations, i.e., $g.v = gv$ and $g.\{h, ha\} = \{gh, gha\}$. In particular, $\text{Cay}(G, \Sigma)$ is vertex-transitive.

Given a graph $\Gamma = (V, E)$, a *path* of length n from $v_0 \in V$ to $v_n \in V$ in Γ is a sequence v_0, \dots, v_n with $\{v_{i-1}, v_i\} \in E$ for all $1 \leq i \leq n$. A *cycle* is a path with $v_0 = v_n$. The *distance* $d(u, v)$ between vertices u and v is defined as the length, i.e., the number of edges, of a shortest path (*geodesic*) connecting u and v . The *diameter* of Γ is $\max\{d(u, v) \mid u, v \in V\}$. Moreover, given $v \in V$ and $H \subseteq V$, we define $d(v, H) = \min\{d(v, h) \mid h \in H\}$ to be the distance of v to H , $\mathcal{N}(v) = \{u \mid \{u, v\} \in E\}$ to be the neighborhood of v , and $\mathcal{N}(H) = \{u \mid \exists v \in H : \{u, v\} \in E\}$ to be the neighborhood of $H \subseteq V$.

A subset $H \subseteq V$ is called a *clique* (resp. *independent set*) if $\{u, v\} \in E$ (resp. $\{u, v\} \notin E$) for all $u, v \in H$ with $u \neq v$. A graph $\Gamma = (V, E)$ is called *complete* if V is a clique; it is called a *cycle* if $V = \{v_1, \dots, v_n\}$ where $|V| = n$ such that $E = \{\{v_i, v_{i+1}\} \mid i \in \{1, \dots, n\}\}$ (calculating modulo n).

2.3. Geodetic graphs.

Definition 2.1. *A graph is geodetic if each pair of vertices is connected by a unique geodesic.*

The following equivalent definition using even cycles is due to Stemple.

Lemma 2.2 ([33, Theorem 2]). *A connected graph Γ is geodetic if and only if Γ contains no even cycle $x_0, \dots, x_{2n} = x_0$ such that $d(x_i, x_{i+n}) = n$ for all $0 \leq i \leq n$.*

An important special case are cycles of length four or six, the conditions under which those can exist are described in [32]. Of these we recall some basic facts on 4-cycles.

Lemma 2.3 ([32, Theorem 3.3]). *Suppose that $\Gamma = (V, E)$ is a geodetic graph. Then the vertices of every cycle v_0, v_1, v_2, v_3, v_0 of length four in Γ induce a complete subgraph.*

Remark 2.4. Throughout, we use Lemma 2.3 in the following way without giving further reference. Suppose that Σ be a generating set of a group G such that the corresponding Cayley graph is geodetic. Then, given $a, b, c, d \in \Sigma$ with $a \neq c$ such that $ab = cd$, we conclude that also $ab, cd, c^{-1}a, bd^{-1} \in \Sigma$.

Lemma 2.5 ([32, Theorem 3.5]). *Let $\Gamma = (V, E)$ be a geodetic graph and $C \subseteq V$ be a clique in Γ . If $v \in V$ is adjacent to at least two distinct vertices in C , then $C \cup \{v\}$ is a clique.*

Lemma 2.6. *Let $\Gamma = (V, E)$ be a geodetic graph. Then the neighbors of any vertex $v \in V$ can be partitioned into a set of disjoint cliques.*

Proof. Assume for contradiction that $x, y, z \in \mathcal{N}(v)$ with $\{x, y\} \in E$, $\{y, z\} \in E$ but $\{x, z\} \notin E$. Then there are two geodesics from x to z (one via v and one via y), contradicting Γ being geodetic. \square

After fixing a starting point in a Cayley graph $\text{Cay}(G, \Sigma)$ (for instance the vertex corresponding to the neutral element 1) paths starting at 1 are in bijection with words $w \in \Sigma^*$ (recall that we always assume the generating set $\Sigma \subseteq G$ to be symmetric). This allows us to denote paths by words in Σ^* rather than vertex sequences and leads to the following observation.

Observation 2.7. *The Cayley graph $\text{Cay}(G, \Sigma)$ of a group G is geodetic if and only if each group element $g \in G$ is represented by a unique geodesic $w \in \Sigma^*$.*

3. STRUCTURE OF GEODETIC CAYLEY GRAPHS

3.1. Complete subgroups.

Definition 3.1. Let G be a group with generating set Σ . We call $H \leq G$ a complete subgroup (with respect to Σ) if $H \setminus 1 \subseteq \Sigma$ or, equivalently, if $H \subseteq G$ induces a complete subgraph of $\text{Cay}(G, \Sigma)$.

Lemma 3.2. Let $\Gamma = \text{Cay}(G, \Sigma)$ be geodesic and let $C \subseteq G$ be a maximal clique of Γ with $1 \in C$.

- (1) Then $g^{-1}C = C$ or $g^{-1}C \cap C = \{1\}$ for each $g \in C$.
- (2) If C is the only maximal clique of its size, then C is closed under inversion.

Proof. Notice that all left translations are isomorphisms of Γ . Since $1 \in g^{-1}C \cap C$ for each $g \in C$, the first statement follows from Lemma 2.6. For the second statement, note that $g^{-1}C = C$ holds for each $g \in C$ by uniqueness of C . As such, $g^{-1} \in g^{-1}C = C$. \square

Note that, in the second case of the previous lemma, where C is the only maximal clique of its size, C is not only closed under inversion, it is also a subgroup. We prove a more general statement.

Lemma 3.3. Let $\Gamma = \text{Cay}(G, \Sigma)$ be geodesic. If $X \subseteq G$ is a clique of Γ with $1 \in X$ and such that X is closed under inversion, then $\langle X \rangle$ is a complete subgroup.

Proof. Let $C \subseteq G$ be a maximal clique containing X . Then $g^2 \in C$ for each $g \in X$. If $g^2 = 1$ then this is obvious. Otherwise $\{1, g, g^2\} \subseteq gX$ is a clique of Γ and thus $g^2 \in C$ by maximality of C .

In particular, for each $g \in X$, we have $\{1, g\} \subseteq g^{-1}C = C$ and thus $g^{-1}C = C$ by Lemma 3.2. As X is closed under inversion, this proves that $X^2 = X \cdot X \subseteq X \cdot C = C$ is a clique of Γ . Moreover, X^2 is closed under inversion and $1 \in X^2 \subseteq C$. Hence $\langle X \rangle = X \cup X^2 \cup X^4 \cup \dots \subseteq C$ by induction. \square

We observe that the second part of Lemma 3.2 and Lemma 3.3 together imply [15, Lemma 14] of Federici.

Lemma 3.4. Let $\text{Cay}(G, \Sigma)$ be geodesic and let $H_1, H_2 \leq G$ be complete subgroups. If $H_1 \cap H_2 \neq 1$ then $\langle H_1, H_2 \rangle$ is also a complete subgroup of G .

Proof. Under these assumptions $H_1 \cup H_2$ is a clique (Lemma 2.5). Since $1 \in H_1 \cup H_2$ and $H_1 \cup H_2$ is closed under inversion, $\langle H_1, H_2 \rangle$ is a clique by Lemma 3.3. \square

Lemma 3.5. Let $\Gamma = (V, E)$ be a biconnected vertex-transitive non-complete geodesic graph and let $C \subseteq V$ be a clique of Γ of size k . Then there is an independent set $I \subseteq \mathcal{N}(C)$ of size $k^2 - k$.

Proof. By [19, Corollary 1] every biconnected non-complete geodesic graph containing a clique of size k also contains the star graph $K_{1,k}$ as an induced subgraph. In other words, there exists a vertex $v \in V$ with an independent set of size k in its neighborhood $\mathcal{N}(v)$. As Γ is vertex-transitive, for every vertex $v \in V$ there is an independent set $I(v) \subseteq \mathcal{N}(v)$ of size k .

Fix a maximal clique \tilde{C} of Γ with $C \subseteq \tilde{C}$ and set $\tilde{I}(v) := I(v) \setminus \tilde{C}$. Since an independent set can contain at most one vertex of any given clique, we have $|\tilde{I}(v)| \geq |I(v)| - 1 = k - 1$.

We claim that $\tilde{I}(u) \cap \tilde{I}(v) = \emptyset$ for distinct $u, v \in C$. If there were a vertex $x \in \tilde{I}(u) \cap \tilde{I}(v)$, then we would have $x \in \tilde{C}$ by Lemma 2.5, because x is adjacent to two vertices in \tilde{C} . That contradicts the definition of \tilde{I} which excludes vertices in \tilde{C} .

Let $I := \bigcup_{v \in C} \tilde{I}(v) \subseteq \mathcal{N}(C)$ and observe that $|I| \geq k(k - 1) = k^2 - k$, as the union is disjoint. It remains to show that I is indeed an independent set. Assume there are $x, y \in I$ with $\{x, y\} \in E$. By definition of I there are $u, v \in C$ such that $x \in \tilde{I}(u)$ and $y \in \tilde{I}(v)$. We must have $u \neq v$, as $\tilde{I}(u)$ is an independent set. Now we have a 4-cycle u, x, y, v . By Lemma 2.3 this implies the existence of the edge $\{u, y\}$. Now y is a neighbor of both u and v . That implies $y \in \tilde{C}$ by Lemma 2.5, a contradiction with definition of \tilde{I} . \square

Proposition 3.6. Let $\Gamma = \text{Cay}(G, \Sigma)$ be a geodesic but not complete Cayley graph. If $H \leq G$ is a complete subgroup with respect to Σ , then $|G| \geq |H|^3 - |H|^2 + |H|$.

Proof. By Lemma 3.5 there is an independent set of size $|H|^2 - |H|$ in the neighborhood of H . We now look at the left cosets of H . Each such coset is a clique of Γ . Thus, each coset can contain at most one point of the independent set. Since H itself does not contain any point of the independent set, the index of H is at least $|H|^2 - |H| + 1$. Hence, $|G| = |G : H| |H| \geq |H|^3 - |H|^2 + |H|$. \square

3.2. Conjugacy classes and elements of order two.

Lemma 3.7 (Conjugacy class of generators). *Let G be a group with geodetic Cayley graph $\text{Cay}(G, \Sigma)$ and $H \leq G$ such that $H = \langle H \cap \Sigma \rangle$. If $x^H \subseteq \Sigma$ for some $x \in \Sigma$, then $H \cap \Sigma = \{x^{\pm 1}\}$ or H is a complete subgroup.*

Proof. If $y \in H \cap \Sigma$ with $y \neq x^{\pm 1}$, then $x^y \in \Sigma$ and, therefore, $yx^y = xy \in \Sigma$ as this element cannot have length zero or two. We also have $x^{-1}y \in \Sigma$ and $yx^{\pm 1} \in \Sigma$ by symmetry.

If $H \cap \Sigma \neq \{x^{\pm 1}\}$, then there exists $y \in H \cap \Sigma$ with $y \neq x^{\pm 1}$. We will show that every $uv \in G$ with $u, v \in H \cap \Sigma$ has length at most one. If $u \notin \{x^{\pm 1}\}$ and $v \in \{x^{\pm 1}\}$, or vice versa, then this follows from the argument above. Otherwise, first consider the case that $u, v \notin \{x^{\pm 1}\}$ and define $u' := ux$ and $v' := \bar{x}v$. As $u' = ux = x\bar{u}u$, we have $u' \in \Sigma$ and likewise $v' \in \Sigma$. Since $uv = u'v'$, it follows that $uv \in \Sigma$. Finally, if $u, v \in \{x^{\pm 1}\}$, we set $u' := uy$ and $v' := y^{-1}v$ and by the same argument conclude that $uv \in \Sigma$. \square

The above lemma may seem technical but is extremely useful, for example, as demonstrated by the following consequences.

Corollary 3.8 (Commuting generators). *Let $\text{Cay}(G, \Sigma)$ be geodetic. If $ab = ba$ and $a \neq b^{\pm 1}$ for some $a, b \in \Sigma$, then the subgroup $\langle a, b \rangle$ is complete.*

Proof. Apply Lemma 3.7 with $H = \langle a, b \rangle$ and $x = a$. \square

Since in any non-cyclic abelian group we must have two commuting generators, we obtain the following result first shown by Georgakopoulos and presented in [15].

Corollary 3.9 ([15, Proposition 10]). *Let G be a non-cyclic abelian group with generating set Σ such that the Cayley graph is geodetic. Then $\text{Cay}(G, \Sigma)$ is complete.*

If in a cyclic group of order at least 3 we have $\Sigma = \{a, \bar{a}\}$ for some generator a of G , then the Cayley graph is a cycle; hence, it is geodetic if and only if $|G|$ is odd. (Note that the cyclic group of order 2 is a special (redundant) case: its only geodetic Cayley graph comes from the generating set consisting of the unique non-trivial element.) Otherwise, we can again apply Corollary 3.8 and obtain the following.

Corollary 3.10. *Let G be a cyclic group with generating set Σ such that the Cayley graph is geodetic. Then either the Cayley graph is a complete graph or a cycle of odd length (and, hence, $|G|$ is odd).*

Another immediate consequence of Lemma 3.7 is that if a geodetic generating set Σ contains any (necessarily non-trivial) conjugacy class x^G , then the associated Cayley graph $\text{Cay}(G, \Sigma)$ is complete or an odd cycle. As particular special cases of this observation, we obtain:

Corollary 3.11 (Generator in center). *Let G be a group with generating set Σ and $Z(G) \cap \Sigma \neq \emptyset$. Then the Cayley graph $\text{Cay}(G, \Sigma)$ is geodetic if and only if it is a complete graph or an odd cycle.*

Corollary 3.12 (Complete normal subgroup). *Let G be a group with generating set Σ and $N \trianglelefteq G$ a non-trivial normal subgroup which is complete with respect to Σ . Then $\text{Cay}(G, \Sigma)$ is geodetic if and only if it is a complete graph.*

If a group has an element of order 2 we have the following.

Lemma 3.13 (Order-two elements). *Let G be a group with generating set Σ such that the Cayley graph is geodetic. Let $g \in G$ be an element of order two. Then the geodesic for g is of the form*

$$(w_1 \cdots w_\ell) \cdot w_{\ell+1} \cdot (\bar{w}_\ell \cdots \bar{w}_1)$$

of length $2\ell + 1$ where $\ell \in \mathbb{N}$ and each $w_i \in \Sigma$. In particular, Σ contains an element of order two conjugate to g , namely $w_{\ell+1}$.

Proof. Let $w_1 \cdots w_k \in \Sigma^*$ be a geodesic for g . We have $g = g^{-1}$ and thus $w_1 \cdots w_k = \bar{w}_k \cdots \bar{w}_1$. Hence, we have two paths of length k that lead from 1 to g . Since $\text{Cay}(G, \Sigma)$ is geodetic, g must have a unique geodesic, i.e., the two paths must coincide; hence, $w_i = \bar{w}_{k-i+1}$ for $i \in \{1, \dots, k\}$.

If k is even, then $g = (w_1 \cdots w_{k/2}) \cdot (w_{k/2+1} \cdots w_k) \equiv (w_1 \cdots w_{k/2}) \cdot (\bar{w}_{k/2} \cdots \bar{w}_k) = 1$, contradicting the assumption that g is of order two. Thus, $k = 2\ell + 1$ must be odd and we obtain the equation

$$\begin{aligned} g &= (w_1 \cdots w_\ell) \cdot w_{\ell+1} \cdot (w_{\ell+2} \cdots w_k) \\ &\equiv (w_1 \cdots w_\ell) \cdot w_{\ell+1} \cdot (\bar{w}_\ell \cdots \bar{w}_1). \end{aligned}$$

Hence, g is conjugate to $w_{\ell+1} \in \Sigma$ and $w_{\ell+1} = \overline{w_{\ell+1}}$, proving the lemma. \square

The above results lead to the following observation which turns out to be extremely useful in any computer search, since it shows that every 2-group immediately satisfies Conjecture A.

Theorem 3.14 (Even order center). *Let G a group such that $Z(G)$ is of even order. Then the only geodesic Cayley graph of G is the complete graph.*

Proof. Assume we have a geodesic Cayley graph of G . By Lemma 3.13, the generating set Σ must contain one element of each conjugacy class of order two elements in G . Since $Z(G)$ is of even order it must contain an order two element g . As g is in the center of the group, it is not conjugate to any other elements. Hence, $g \in \Sigma$ and by Corollary 3.11, the Cayley graph is either complete, or an odd cycle. Since G contains an even order subgroup, G itself has even order. Therefore, the Cayley graph cannot be an odd cycle, so it must be complete graph. \square

Remark 3.15. All but 4734 groups of the approximately 49.5 billion groups of order at most 1024 are covered by the combination of Theorem 3.14 and Corollary 3.9.

3.3. Cayley graphs of diameter two.

Lemma 3.16. *If $\Gamma = \text{Cay}(G, \Sigma)$ is geodesic and has diameter 2, then Γ is the cycle C_5 of length 5.*

Proof. From Stemple [32] we know that every geodesic graph of diameter two falls into one of the following classes: block graphs joining all cliques in one vertex, biconnected graphs with exactly two different vertex degrees, and Moore graphs of diameter two.

Among these, the Moore graphs are the only regular graphs. The Moore graphs of diameter two are the C_5 , the Petersen graph, the Hoffman-Singleton graph and a hypothetical graph of degree 57 [1, 7]. A degree-57 Moore graph cannot be vertex-transitive [4, Theorem 1.13], and, the Petersen graph and Hoffman-Singleton graph are not Cayley graphs [28] (see also [23] or our own computer search). Thus C_5 is the only geodesic Cayley graph of diameter 2. \square

The combination of Lemma 3.13 and Lemma 3.7 affords a short elementary proof of the (well-known) fact that the Moore graphs of diameter 2, other than C_5 , are not Cayley graphs.

Proposition 3.17. *The Petersen graph, the Hoffman-Singleton graph, and all of the hypothetical Moore graphs of degree 57 and diameter 2 are not Cayley graphs.*

Proof. Assume that $\Gamma = \text{Cay}(G, \Sigma)$ for one of these graphs. Then $|G|$ is even and, as such, there exist elements of order two in G . Since Γ is geodesic and has diameter two, each such element is contained in Σ by Lemma 3.13. But then Σ contains an entire conjugacy class. By Lemma 3.7, the graph Γ would therefore have to be complete or a cycle, both of which is absurd. \square

3.4. Central elements.

Lemma 3.18. *Let $\text{Cay}(G, \Sigma)$ be geodesic and $b, t \in \Sigma$. Suppose that $b^2 \neq 1$, $t^2 = 1$, and $b^2t = tb^{\pm 2}$. Then the subgroup $\langle b, t \rangle \leq G$ is complete with respect to Σ .*

Proof. If $bt = tb^{\pm 1}$, then the statement follows from Lemma 3.7, so assume that $bt \neq tb^{\pm 1}$ holds. As $bbt = tbb$ or $bbt = t\bar{b}\bar{b}$, we have $s := bt\bar{b} = \bar{b}tb$ or $s := btb = \bar{b}t\bar{b}$, respectively. In particular, there are two words of length three representing s ; thus there must be a shorter one. Moreover, since s has order two, its geodesic must have odd length. Hence, $s \in \Sigma$. The situation is as shown in Figure 1.

Let $H := \langle b, t, s \rangle = \langle H \cap \Sigma \rangle \leq G$. One may verify by straightforward computations (conjugating st by b, \bar{b}, s, t) that $(st)^H = \{st, ts\} \subseteq \Sigma$. The statement then follows using Lemma 3.7. \square

Lemma 3.19. *Let $\text{Cay}(G, \Sigma)$ be geodesic. If there exists some $b \in \Sigma$ with $b^2 \neq 1$ and $(b^2)^G \subseteq \{b^{\pm 2}\}$, then $\text{Cay}(G, \Sigma)$ is an odd cycle or a complete graph.*

Proof. Suppose first, that $(b^2)^G = \{b^2\}$ and thus $b^2 \in Z(G)$. Then either $b \in Z(G)$ or $b \notin Z(G)$. In the first case, the statement follows immediately from Lemma 3.7. In the second case, $|G : Z(G)|$ is even; hence so is $|G|$. As such, there exists an element of order two in G . In fact, there even exists $t \in \Sigma$ with $t^2 = 1$ by Lemma 3.13. The statement now follows from Lemma 3.18 and Lemma 3.7.

Finally, suppose that $(b^2)^G = \{b^{\pm 2}\}$ has two elements. Then $|G : C_G(b^2)| = 2$ and, as before, there thus exists some $t \in \Sigma$ with $t^2 = 1$. The statement follows from Lemma 3.18 and Lemma 3.7. \square




FIGURE 1. The situation in the proof of Lemma 3.18.

Lemma 3.20 (Flat coset). *Let G be a group with generating set Σ such that the Cayley graph is geodesic. If $w \in \Sigma^m$ is a geodesic of length m and $a \in \Sigma$ such that $aw = wa$ and $w \neq a^{\pm m}$, then each $h \in w\langle a \rangle$ has a geodesic of length at most m .*

Moreover, if $w \notin \langle a \rangle$, then each $h \in w\langle a \rangle$ has a geodesic of length exactly m .


Proof. By w_i we denote the geodesic of wa^i . Let r be the order of a in G . Observe that $w_0 = w_r$.

First, we prove the lemma for the case $w \notin \langle a \rangle$. Now aw_i and $w_i a$ are two different words of length $|w_i| + 1$ representing wa^{i+1} . Thus, there must be a shorter word, implying that $|w_{i+1}| \leq |w_i|$. Therefore, $m = |w_r| \leq \dots \leq |w_i| \leq \dots \leq |w_0| = m$, and we conclude $|w_i| = m$ for all i .

Second, we consider the case $w = a^k$. Recall that w_i is the geodesic for $wa^i = a^{i+k}$. Observe that, because the geodesic of a^k has length m , we must have $m \leq k$. In fact, because $w \neq a^{\pm m}$, we have $m < k$. Using the same argument, we also have $m < r - k$. We proceed by induction on i that $|w_i| \leq m$. For $w_0 = w_r = w$ this obviously holds.

If $0 < i \leq r - k$, then we have two words, $w_{i-1}a$ and aw_{i-1} of length $|w_{i-1}| + 1$ for $a^{i+k} = w_i$. The two words are different, because $w_{i-1} \neq a^{|w_{i-1}|}$: If w_{i-1} were a power of a , then $w_{i-1} = a^{i+k-1}$. For $i = 1$ this contradicts one of the conditions of the lemma, for $i > 1$ we have $|w_{i-1}| \leq m < k < i + k - 1$. Thus, the geodesic w_i has length at most $|w_{i-1}| \leq m$.

For $r - k \leq i < r$, we also use induction on decreasing values of i , multiplying with \bar{a} . Now $w_r = w$ is the induction base. Assuming the statement holds for $i + 1$, the two words $w_{i+1}\bar{a}$ and $\bar{a}w_{i+1}$ both correspond to the group element $a^{i+k} = w_i$. For $i = r - 1$ they are different because of the conditions of the lemma, for $i < r - 1$ we show that $w_{i+1} \neq a^{|w_{i+1}|}$: If w_{i+1} were a power of \bar{a} , then $w_{i+1} = \bar{a}^{2r-k-i-1}$ leading to the contradiction $2r - k - i - 1 > r - k > m \geq |w_{i+1}|$. Therefore, we must have $|w_i| \leq |w_{i+1}| \leq m$. \square

FIGURE 2. The coset $w\langle a \rangle$ in the case $w = a^k$ in the proof of Lemma 3.20.

The second case of the proof, where $w = a^k$ is illustrated in Figure 2. One can observe the two inductions, starting at w , one multiplying with a , the other multiplying with \bar{a} covering the entire subgroup. Before

arriving at 1 we encounter small powers of a , respectively \bar{a} . In fact, there is an integer $d \leq |w|$, such that a^i and \bar{a}^i are geodesic for $0 \leq i \leq d$, but for $j > d$ the words a^j and \bar{a}^j are not geodesics. By applying Lemma 3.20 to the geodesic of a^{d+1} , which has length d , we obtain $d = m$.

Lemma 3.21 (Geodesics for central elements are powers of generators). *Let G be a group with geodesic Cayley graph $\text{Cay}(G, \Sigma)$, and let $z \in Z(G) \setminus 1$. Then the unique geodesic of z is a^k for some $a \in \Sigma$ and $k \geq 1$.*

Proof. Let $w_1 \cdots w_k \in \Sigma^*$ be a geodesic for z . As z is in the center we have

$$w_1 w_2 \cdots w_k = z = z^{w_1} = w_2 \cdots w_k w_1.$$

Now we have two words of length k representing z . There cannot be a shorter word, as we chose $w_1 \cdots w_k$ to be a geodesic. The Cayley graph is geodesic, thus the two words must coincide: we obtain $w_1 = w_2 = \cdots = w_k$. \square

Theorem 3.22 (Geodesics for central elements are powers with the same exponent). *Let G be a group with geodesic Cayley graph $\text{Cay}(G, \Sigma)$ and $\Sigma \cap Z(G) = \emptyset$. Then, there is an integer $k \geq 3$ such that for each $z \in Z(G) \setminus 1$ there exists an element $c_z \in \Sigma$ such that c_z^k is the geodesic for z .*

In light of this theorem, we can refer to the *length of central geodesics* as the integer k so that $z = c_z^k$ for any element $z \in Z(G)$. Note that k divides the order of $G/Z(G)$. We refer to the above elements c_z as *central roots*.

Proof. We know by Lemma 3.21 that the geodesic of every element in the center is a power of a generator. We first show that no two central elements can have a geodesic that is a power of the same generator (note that a generator and its inverse is allowed and will happen). Let b^k be a geodesic representing a central element and assume k is minimal so that b^k is central. Let g be a generator different to b^\pm (such a generator must exist, otherwise G would be a cyclic group, but then $b \in Z(G)$). By Lemma 3.20 the elements $b^k g$ and $b^k \bar{g}$ both have a geodesic of length at most k , which we write as $(b^k g)$ and $(b^k \bar{g})$. Now $(b^k g)(b^k \bar{g})$ is a word of length at most $2k$ representing the central element b^{2k} . Thus, b^{2k} is not a geodesic. It follows that no power of b other than k gives a geodesic for a central element, since k is minimal and b^{k+i} geodesic and central would imply b^i geodesic and central, so $i \geq k$.

Next we show that the geodesics of all central elements have the same length. For a central element and its inverse this is obvious. Assume there are $c, d \in \Sigma$ with $c \notin \{d, \bar{d}\}$ and $k, \ell \in \mathbb{N}$ such that c^k and d^ℓ both are geodesics representing different central elements. Note that $c^k \neq \bar{d}^\ell$ because both are geodesics. Now assume for a contradiction that $k \neq \ell$, w.l.o.g. $k < \ell$. Lemma 3.20 tells us that $c^k d^\ell \in c^k \langle d \rangle$ has a geodesic u of length at most k . But $c^k d^\ell$ is also in the center (and non-trivial), so $u = f^m$ for some $f \in \Sigma, m \leq k$. Applying Lemma 3.20 again to $d^\ell = f^m c^{-k} \in f^m \langle c \rangle$, we obtain a geodesic v of length at most $m \leq k < \ell$, contradicting d^ℓ being a geodesic.

It remains to show, that no central element has a geodesic of length two. In that case the Cayley graph would be cyclic or complete by Lemma 3.19, contradicting $\Sigma \cap Z(G) = \emptyset$. \square

Lemma 3.23. *Let $\text{Cay}(G, \Sigma)$ be geodesic. If there is an element $z \in Z(G)$, with $\text{ord}(z) \geq 4$, then the length of central geodesics must be odd.*

Proof. Let $m = \text{ord}(z)$. If m is even, then the statement follows from Corollary 3.11. Thus from here on $m \geq 5$.

Assume that b^{2k} is the geodesic for z . Let y be the geodesic for $z^{(m-1)/2}$. By Theorem 3.22 it has length $2k$. As $m \geq 5$, we have $y \neq b^{\pm 2k}$. By Lemma 3.20, the element $t = b^{km} = y b^k$ has length at most $2k$. It also has order two, thus by Lemma 3.13 it must have odd length, that is, the length of t is at most $2k - 1$. Applying Lemma 3.20 to $y = t \bar{c}^k$ we obtain the contradiction $|y| \leq 2k - 1$. \square

4. BOUNDS ON THE SIZE OF GENERATING SETS

In this section we establish bounds on the possible sizes of those generating sets Σ of a group G which result in a geodesic but neither complete nor cyclic Cayley graph $\Gamma = \text{Cay}(G, \Sigma)$.

Our method utilizes the following properties of the structure balls of radius one and two. Throughout, we denote the r -ball in Γ centered at a vertex $g \in G$ by $\mathcal{B}_r(g) := \{h \in G \mid d(g, h) \leq r\}$. As Cayley graphs are vertex-transitive, all balls of the same radius r are isomorphic subgraphs of Γ .

We begin by analyzing the structure of one-balls. By Lemma 2.6 the neighbors of any $g \in G$ can be partitioned into a set of disjoint cliques. We denote by m the number of these disjoint cliques and by $\delta_1 \leq \delta_2 \leq \dots \leq \delta_m$ their sizes. Clearly, $\delta := \sum_{i=1}^m \delta_i$ is the degree of Γ and thus $\delta = |\Sigma|$.

Lemma 4.1. *We have $m \geq 1 + \delta_m \geq 1 + \delta/m$. If Γ is not complete or a cycle, then $m \geq 3$.*

Proof. The Cayley graph Γ contains a clique of size $\delta_m + 1$. By [19, Corollary 1] the neighborhood of some and, hence, of every vertex contains an independent set of size $\delta_m + 1$. Thus $m \geq \delta_m + 1$ by definition of m . The second inequality follows from $\delta = \sum_{i=1}^m \delta_i \leq \sum_{i=1}^m \delta_m \leq m\delta_m$.

For the final inequality, note that $m \leq 1$ holds if and only if Γ is complete. If $m = 2$, then the above argument shows that $\delta_1 = \delta_2 = 1$ and thus $\delta = 2$. If both elements of Σ are of order two, then G is a dihedral group and Γ is an even cycle; hence, Γ is not geodetic. Otherwise, the two elements of Σ are inverses of one another and, as such, the group G is cyclic. But then Γ is complete, a cycle, or not geodetic by Corollary 3.10. The only remaining possibility is $m \geq 3$. \square

Lemma 4.2. *For every $g \in G$, the size of the two-ball centered at g satisfies*

$$|\mathcal{B}_2(g)| = 1 + \delta + \delta^2 - \sum_{i=1}^m \delta_i^2 \geq 1 + \delta + \frac{1}{2}\alpha\delta^2 \quad \text{where} \quad \alpha = \alpha(m) := \frac{3m-4}{2m-2}.$$

Proof. Every vertex at distance one from g which is contained in a clique of size δ_i has $\delta - \delta_i$ neighbors at distance two from g . As each vertex at distance two from g has a unique neighbor at distance one from g , we obtain the equality

$$|\mathcal{B}_2(g)| = 1 + \delta + \sum_{i=1}^m \delta_i(\delta - \delta_i) = 1 + \delta + \delta^2 - \sum_{i=1}^m \delta_i^2.$$

We then apply the reversed Cauchy-Schwarz inequality due to Pólya and Szegő [27, pp. 57, 213–214] with the bounds $1 \leq \delta_i \leq m - 1$ obtained in Lemma 4.1. This yields

$$m \sum_{i=1}^m \delta_i^2 = \left(\sum_{i=1}^m 1^2 \right) \cdot \left(\sum_{i=1}^m \delta_i^2 \right) \leq \frac{1}{4} \frac{m^2}{m-1} \left(\sum_{i=1}^m \delta_i \right)^2 = \frac{1}{4} \frac{m^2}{m-1} \delta^2,$$

from which the claimed inequality follows. \square

Lemma 4.3. *Let $g, h \in G$. If $d(g, h) \geq 3$, then $|\mathcal{B}_2(g) \cap \mathcal{B}_2(h)| \leq 2\delta - 1$.*

Proof. Let h' be the vertex preceding h on the geodesic path from g to h . Then $\mathcal{B}_2(g) \cap \mathcal{B}_1(h) \subseteq \{h'\}$ by uniqueness of geodesics. Hence, $|\mathcal{B}_2(g) \cap \mathcal{B}_1(h')| \leq \delta$. If $h'' \in \mathcal{N}(h) \setminus \{h'\}$, then $d(g, h'') \geq 3$ which implies $|\mathcal{B}_2(g) \cap \mathcal{B}_1(h'')| \leq 1$. In total we obtain $|\mathcal{B}_2(g) \cap \mathcal{B}_2(h)| \leq 2\delta - 1$. \square

We now have all the tools necessary to prove the main result of this section.

Theorem 4.4. *Let $\text{Cay}(G, \Sigma)$ be geodetic but neither complete nor a cycle. If*

$$\alpha_0 = \alpha(m_0) = \frac{3m_0 - 4}{2m_0 - 2}$$

where $m_0 \geq 3$ is any integer with $\frac{1}{2}(m_0 - 1)(3m_0 - 4)(m_0 - 2)^2 < |G|$, then $\alpha_0 |\Sigma|^2 < |G|$.

Remark 4.5. Before proceeding with the proof, we note that the factor $\alpha_0 = \alpha(m_0)$ is monotonically increasing in m_0 . The choice $m_0 = 3$ is always valid and yields $\alpha_0 = \frac{5}{4}$. On the other hand, $\alpha_0 \rightarrow \frac{3}{2}$ as $m_0 \rightarrow \infty$. Moreover, as a byproduct of our computer experiments, we obtain *a posteriori* that the conclusion of Theorem 4.4 holds with $\alpha_0 = \frac{7}{5}$, i.e., we obtain Theorem D: after verifying that Conjecture A holds every group of order at most 560 (see Theorem B), we set $m_0 = 6$.

We use a combination of the bound obtained in Theorem 4.4, as well as other ones discussed hereafter, in our computer experiments. This results in a massive reduction in the number of generating sets we have to consider. In order to make these bounds as tight as possible, we choose the maximal value m_0 that is permitted for the group currently under examination. For some groups, we also employ Theorem 3.22 to

establish an improved lower bound on the diameter which, in turn, allows us to replace Lemma 4.3 with a better estimate.

Proof. For the sake of deriving a contradiction, we assume that $\alpha_0 |\Sigma|^2 \geq |G|$. We also continue to employ the notation established above. In particular, the previous inequality becomes $\alpha_0 \delta^2 \geq |G|$, and Lemma 4.1 then yields $\alpha_0 (m^2 - m)^2 \geq \alpha_0 \delta^2 \geq |G|$. We cannot have $m < m_0$, for otherwise

$$\frac{1}{2}(m_0 - 1)(3m_0 - 4)(m_0 - 2)^2 = \alpha_0((m_0 - 1)^2 - (m_0 - 1)) \geq \alpha_0(m^2 - m)^2 \geq |G|$$

which contradicts the choice of m_0 . Hence $m \geq m_0$ and, therefore, also $\alpha = \alpha(m) \geq \alpha(m_0) = \alpha_0$.

Finally, observe that $\text{Cay}(G, \Sigma)$ has diameter at least three by Lemma 3.16. Hence there exist elements $g, h \in G$ with $d(g, h) = 3$. Using Lemma 4.2 and Lemma 4.3 we arrive at

$$|\mathcal{B}_2(g) \cup \mathcal{B}_2(h)| = \underbrace{|\mathcal{B}_2(g)| + |\mathcal{B}_2(h)|}_{\geq 2+2\delta+\alpha\delta^2} - \underbrace{|\mathcal{B}_2(g) \cap \mathcal{B}_2(h)|}_{\leq 2\delta-1} \geq 3 + \alpha\delta^2 \geq 3 + \alpha_0\delta^2 \geq 3 + |G|.$$

As such, $|G| \geq |\mathcal{B}_2(g) \cup \mathcal{B}_2(h)| \geq |G| + 3 > |G|$, which is the desired contradiction. \square

Corollary 4.6. *If $\text{Cay}(G, \Sigma)$ is geodetic but not complete or a cycle, then $\alpha_0(|Z(G)| - 1)^2 \leq |G|$.*

Proof. This follows from Theorem 4.4, as $|\Sigma| \geq |Z(G)| - 1$ by Theorem 3.22 and Corollary 3.11. \square

In Theorem 4.4 we have used two balls of radius two at distance three from each other to give a lower bound to the size of the group. In cases where the group has a non-trivial center, we can improve upon this. By Theorem 3.22, the distance between any two central elements is at least three. Using a subset of these as centers of balls of radius two, and Lemma 4.3 to bound the size of pairwise intersections, we arrive at the following.

Theorem 4.7. *Let $\text{Cay}(G, \Sigma)$ be geodetic but neither complete nor a cycle. Then the inequality*

$$\frac{1}{2}\alpha_0 z_0 |\Sigma|^2 - z_0(z_0 - 2) |\Sigma| + \frac{1}{2}z_0(z_0 - 1) \leq |G|,$$

where

$$\alpha_0 = \alpha(m_0) = \frac{3m_0 - 4}{2m_0 - 2},$$

holds for all integers z_0 with $1 \leq z_0 \leq |Z(G)|$ and all lower bounds m_0 on the number $m \geq m_0$ of maximal cliques in the neighborhood $\mathcal{N}(1)$ of $1 \in G$.

Proof. Choose a set of z_0 central elements, say $Z_0 \subseteq Z(G)$. By Lemma 4.2 and Lemma 4.3,

$$|G| \geq \left| \bigcup_{g \in Z_0} \mathcal{B}_2(g) \right| \geq \sum_{g \in Z_0} |\mathcal{B}_2(g)| - \sum_{\substack{g, h \in Z_0 \\ g \neq h}} |\mathcal{B}_2(g) \cap \mathcal{B}_2(h)| \geq z_0(1 + \delta + \frac{1}{2}\alpha\delta^2) - \binom{z_0}{2}(2\delta - 1)$$

with $\alpha \geq \alpha_0$. Upon expanding the rightmost expression one obtains the stated inequality. \square

Note that for groups with small center, the best lower bound is achieved by $z_0 = |Z(G)|$. However, if the center is large enough, then at some point it is no longer beneficial to place more balls, due to the way we approximate the intersection. The best lower bound is obtained for $z_0 \approx \frac{1}{4}\alpha_0\delta$. Based on this observation we obtain the following bound on the center using $|\Sigma| \geq |Z(G)| - 1$ and setting $z_0 = \lceil \frac{1}{4}\alpha_0(|Z(G)| - 1) \rceil + 1$.

Corollary 4.8. *If $\text{Cay}(G, \Sigma)$ is geodetic but not complete or a cycle, then $\frac{1}{16}\alpha_0^2(|Z(G)| - 1)^3 \leq |G|$.*

If the central geodesics are sufficiently long, then we can obtain an even better bound by placing balls along these central geodesics (as opposed to only their endpoints). We begin with the following estimate on the distance between two points on two different central geodesics.

Lemma 4.9. *Let $\text{Cay}(G, \Sigma)$ be geodetic but not complete or a cycle. Let $g = ya^i$ and $h = zb^j$, where $y, z \in Z(G)$ and $a, b \in \Sigma$, be two points on two different central geodesics. Then*

$$d(g, h) \geq \max\{\min\{i, k - i\}, \min\{j, k - j\}\},$$

where k is the length of the central geodesics.

Proof. If any of y , ya^k , z and zb^k coincide, then we may assume that $ya^k = zb^k$. This can be achieved by replacing y with ya^k and i with $k - i$ and/or replacing z and j suitably. This step is the origin of the minima appearing in the statement. By Lemma 3.20, the geodesic from y to h has length at least k , thus $d(g, h) \geq k - \min\{i, k - i\} = \min\{i, k - i\}$. Similarly, $d(g, h) \geq \min\{j, k - j\}$. \square

Multiple balls of radius two can be placed on each central geodesic if its length permits that. However, we will only give an explicit bound for the simplest case, placing a single such ball in the middle of each central geodesic.

Proposition 4.10. *If $\text{Cay}(G, \Sigma)$ is geodetic but not complete or a cycle, $3 \nmid |G|$, and $5, 7 \nmid |G : Z(G)|$, then $\frac{1}{4}\alpha_0(|Z(G)| - 1)^4 < |G|$.*

Proof. As $3 \nmid |G|$, the length k of central geodesics must be odd by Lemma 3.23. As $5, 7 \nmid |G : Z(G)|$ we have $k \geq 11$. We place a 2-ball on the middle of each central geodesic. The intersection between any pair of 2-balls is trivial, as the distance between any two of their center points is at least 5. Thus we obtain

$$\begin{aligned} |G| &\geq \frac{1}{2}|Z(G)|(|Z(G)| - 1)|\mathcal{B}_2(1)| \\ &\geq \frac{1}{2}(|Z(G)| - 1)^2\left(\frac{1}{2}\alpha\delta^2 + \delta + 1\right) \\ &\geq \frac{1}{4}\alpha(|Z(G)| - 1)^2\delta^2. \end{aligned}$$

Using the inequality $\delta \geq |Z(G)| - 1$ yields the desired bound. \square

5. FURTHER CASES: DIHEDRAL, NILPOTENT, GROUPS WITH LARGE COMMUTATIVITY DEGREE

In this section we show that further large families of groups satisfy Conjecture A by combining results from Section 3 with more detailed knowledge about finite group theory and insights gleaned from the computer search.

We first consider groups that have an abelian subgroup of index two (which is necessarily normal); these include the dihedral groups. Pushing this to index three presents more challenges; so for that case we are able to prove the conjecture only in the two important special cases when the subgroup is not normal or when the center is trivial. For nilpotent groups we can prove the conjecture holds in all groups of class two except for a particular subfamily (see Corollary 5.16), and in groups of any class provided certain numerical conditions are satisfied (see Theorem 5.9). Each of these families of groups in some sense is close to abelian, which is emphasized by the fact that these classes cover all groups with a high *commutativity degree* (see Theorem 5.17).

5.1. Abelian subgroups of index two.

Lemma 5.1. *Let G be a group and $N \leq G$ a non-trivial subgroup with $|G : N| = 2$. If $\text{Cay}(G, \Sigma)$ is geodetic then $N \cap \Sigma \neq \emptyset$.*

Proof. If $\Sigma \cap N = \emptyset$, then every word $w \in \Sigma^*$ representing $1 \in G$ has even length. Hence, all cycles in $\text{Cay}(G, \Sigma)$ must have even length. But then $\text{Cay}(G, \Sigma)$ cannot be geodetic (see Lemma 2.2). \square

Theorem 5.2. *Let $\phi : A \rightarrow A$ be an order-two automorphism of an abelian group. Let*

$$D_{A,\phi} := A \rtimes_{\phi} C_2 = \langle X \cup \{t\} \mid R \cup \{t^2, txt = \phi(x); x \in X\} \rangle$$

where $A = \langle X \mid R \rangle$ be the corresponding semidirect product. Then the only geodetic Cayley graph of the group $D_{A,\phi}$ is the complete graph.

Proof. Assume that $\text{Cay}(D_{A,\phi}, \Sigma)$ is geodetic but not complete (in particular, $Z(D_{A,\phi}) \cap \Sigma = \emptyset$ by Corollary 3.11). Since $\text{ord}(t) = 2$, the generating set Σ contains a conjugate of t by Lemma 3.13. Upon replacing Σ with a suitable conjugate if necessary, we may therefore assume that $t \in \Sigma$.

Since $|D_{A,\phi} : A| = 2$, there exists some $x \in A \cap \Sigma$ by Lemma 5.1. If $\phi(x) = x$, that is, $txt = x$ then $x \in Z(D_{A,\phi})$ which is not possible. If $\phi(x) = \bar{x}$ then $x^{D_{A,\phi}} = \{x^{\pm 1}\} \subseteq \Sigma$. Since $\Sigma \neq \{x^{\pm 1}\}$, Lemma 3.7 implies that $\text{Cay}(D_{A,\phi}, \Sigma)$ is complete. Thus $\phi(x) \notin \{x^{\pm 1}\}$. Note that $\phi(x)x$ commutes with t since $t\phi(x)x = ttxt = xttx = x\phi(x)t = \phi(x)xt$ as $x, \phi(x) \in A$. Hence $\phi(x)x \in Z(D_{A,\phi}) \setminus 1$.

By Theorem 3.22, there exists $y \in \Sigma$ such that $\phi(x)x = y^k$ with $k \geq 3$ and, therefore, with $k = 3$ by uniqueness of geodesics (since $\phi(x)x = txtx$). Thus, in particular, the length of central geodesics is three. Now, suppose that $z \in \Sigma$ with $z^3 \in Z(G)$. If $z \notin A$ then $z = at$ for some $a \in A$. But then $z^3 = atatat = a\phi(a)at$ and, since this is central, $(a\phi(a)at)t = t(a\phi(a)at) = \phi(a)a\phi(a)$. As such, $a^2\phi(a) = a\phi(a)^2$ which implies $a = \phi(a)$. Hence t and a commute, so a is central. Now $z^3 = a^3t$ implies $t = a^{-3}z^3 \in Z(G)$, which is a contradiction. It follows that every element of $Z(D_{A,\phi}) \setminus 1$ is of the form z^3 for some $z \in \Sigma \cap A$. In particular, this shows that $y \in \Sigma \cap A$.

We now argue that we can find an element $z \in \Sigma \cap A$ with $z \notin \{y^{\pm 1}\}$. If so, then y and z commute; hence $z^{\langle y \rangle} = \{z\} \subseteq \Sigma$ and then we can apply Lemma 3.7 to show that $\langle y \rangle$ is complete. However, this implies $y^3 \in \Sigma$ and thus $Z(G) \cap \Sigma \neq \emptyset$; contradicting Corollary 3.11.

Firstly, suppose that $y = x$, then $x^3 = y^3 = \phi(x)x$ and thus $\phi(x) = x^2$. We obtain the contradiction $y^3 = x^3 = \bar{x}x^4 = \bar{x}\phi(\phi(x)) = \bar{x}x = 1$. Next, suppose that $y = x^{-1}$ holds. Then $x^{-3} = y^3 = \phi(x)x$ and thus $\phi(x) = x^{-4}$, or equivalently $\phi(y) = y^{-4}$. Then $\phi(\phi(y)) = \phi(y^{-4}) = y^{16}$ and since ϕ is order two, this means $y = y^{16}$ so $y^{15} = 1$. This means the order of y is either 3, 5, or 15. It cannot be 3 since y^3 is a non-trivial element of the center. If $y^5 = 1$ then $y \in Z(G)$ and so $Z(G) \cap \Sigma \neq \emptyset$ and we are done. Thus $\text{ord}(y) = 15$. We then have $y^6 \in Z(D_{A,\phi})$ and $y^6 \notin \{y^{\pm 3}\}$. Thus $y^6 = z^3$ for some $z \in \Sigma \cap A$ with $z \notin \{y^{\pm 1}\}$. Lastly if $y \neq x^{\pm 1}$ then $z := x$ satisfies our requirements. Thus we have found an element $z \in \Sigma \cap A$ with $\{y^{\pm 1}\}$ which by the above paragraph shows that the Cayley graph is complete, contradicting our assumption. \square

Corollary 5.3. *The only geodetic Cayley graph of a dihedral group is the complete graph.*

We will now reduce the general case, of admitting an abelian subgroup of index two, to the situation discussed in Theorem 5.2. In other words, we will prove the following.

Theorem 5.4. *Let G be a group. If there exists an abelian subgroup $A \leq G$ such that $|G : A| = 2$, then the only geodetic Cayley graph of G is the complete graph.*

Our reduction will rely on a certain relationship between the structure of the group G and the parity of the order of A as well as that of $Z(G)$. Part of this relationship is captured by the following observation (with $p = 2$). It will be used again (with $p = 3$) in Section 5.2.

Lemma 5.5. *Let G be a group and $A \trianglelefteq G$ a normal abelian subgroup of prime index $|G : A| = p$. If $g \in G \setminus A$, then $g^p \in Z(G) \cap A$. Moreover, if $p \nmid |A|$, then there exists some $\tilde{g} \in G \setminus A$ with $\tilde{g}^p = 1$.*

Proof. The image of any given $g \in G \setminus A$ in the quotient $G/A \cong C_p$ has order p and, as such, $g^p \in A$. In particular, g^p commutes with every element $a \in A$ and, clearly, g^p also commutes with g . We have $\langle A, g \rangle = G$ since $A \leq G$ is a maximal subgroup (its index is prime) and $g \in G \setminus A$. Therefore, the element g^p commutes with all elements of G , i.e., $g^p \in Z(G)$ as claimed.

Now assume that $p \nmid |A|$ and choose $g \in G \setminus A$. By the above, we have $g^p \in Z(G) \cap A$ and thus the cyclic subgroup $H := \langle g^p \rangle$ is contained in $Z(G) \cap A$ and, in particular, $H \leq A$. From this we infer that $p \nmid |H|$ and, as p is prime, there exists $a \in H$ with $a^{-p} = g^p$. The element $\tilde{g} := ag$ has the desired properties. Indeed, $\tilde{g} = ag \in G \setminus A$ holds since $a \in A$ and $g \in G \setminus A$, and $\tilde{g}^p = (ag)^p = a^p g^p = 1$ holds since $a \in Z(G)$. \square

Proof of Theorem 5.4. If $|Z(G)|$ is even then Theorem 3.14 applies, so assume $|Z(G)|$ is odd. We claim that $|A|$ is odd as well. Suppose otherwise. Then there exists some $a \in A$ with $\text{ord}(a) = 2$. Note that we cannot have $a \in Z(G)$ for $|Z(G)|$ is odd. Let $g \in G \setminus A$ and consider $\tilde{a} := a^g a \in A$. Clearly, $a^g \neq a$ for otherwise we would have $a \in Z(G)$. Therefore, $\tilde{a} \neq 1$. Since $\tilde{a}^2 = 1$, we conclude that $\text{ord}(\tilde{a}) = 2$. Moreover $\tilde{a}^g = a^{g^2} a^g = a a^g = a^g a = \tilde{a}$ since $g^2 \in Z(G)$ by Lemma 5.5. But then we conclude that $\tilde{a} \in Z(G)$ and, therefore, $|Z(G)|$ would have to be even.

Finally, since $|A|$ is odd, there exists $t \in G \setminus A$ with $t^2 = 1$ by Lemma 5.5. Consider the automorphism $\phi : A \rightarrow A$ with $\phi(a) = a^t$. It satisfies $\phi^2 = \text{id}$ and $\phi \neq \text{id}$ since $t^2 = 1$ and $t \notin Z(G)$, respectively. As such, it has order two and we can apply Theorem 5.2. \square

5.2. Abelian subgroups of index three. For index 3 abelian we are able to show the conjecture holds for two cases: first if A is not normal, and second if the center of G is trivial.

The general intuition is that, if a group has a lot of commuting elements (as it does when it contains an abelian subgroup of small index), then all of its Cayley graphs will need to have a lot of squares. However, we can only manage to make this precise with particular hypothesis. In Section 5.4 we investigate this further.

Lemma 5.6. *Let G be a group. If there exists an abelian subgroup $A \leq G$ such that $|G : A| = 3$ and A is not normal in G , then the only geodetic Cayley graph of G is the complete graph.*

Proof. Suppose that $A \leq G$ is abelian with $|G : A| = 3$ and that A is not normal in G . Let $A^g \leq G$ be a conjugate subgroup with $A \neq A^g$. Then $\langle A, A^g \rangle = G$ since A is a maximal subgroup of G (as its index is prime). Moreover, we observe that $A \cap A^g \subseteq Z(G)$ as every element of $A \cap A^g$ commutes with every element of $\langle A, A^g \rangle$.

Now consider the action of G by left-multiplication on the set of cosets G/A . It gives rise to a homomorphism $\rho : G \rightarrow S_3$ with $\text{Ker}(\rho) = \{h \in G \mid hgA = gA \text{ for all } g \in G\} = \bigcap_{g \in G} A^g \leq Z(G)$. The ρ -preimage of $C_3 \leq S_3$ is an abelian subgroup $\rho^{-1}(C_3) \leq G$, as it is an extension of $Z(G)$ by a cyclic group. Since $|G : \rho^{-1}(C_3)| = 2$, the statement now follows from Theorem 5.4. \square

Remark 5.7. In the situation described in Lemma 5.6, we have $|G : Z(G)| = 6$. In Corollary 4.6 we have proved an inequality relating the size of the center and the size of the group, which in the case at hand gives us the bound $|G| \leq 36$; these remaining groups are covered by our computer search.

We note that this alternative proof can be adapted, so as to cover groups with an abelian but *not* normal subgroup of index 5 or 7 (the center of such a group has index at most 20 or 42 and, thus, by Corollary 4.8 we need only consider such groups with order at most 300 and 840, respectively).

The other case is that G contains a normal abelian subgroup of index three. For this we can make progress when we restrict to the case the center is trivial.

Theorem 5.8. *Let G be a group with $Z(G) = 1$. If there exists an abelian subgroup $A \leq G$ such that $|G : A| = 3$, then the only geodetic Cayley graph of G is the complete graph.*

Proof. Let G and A be as in the hypothesis and assume that $\text{Cay}(G, \Sigma)$ is geodetic but not complete. By Lemma 5.6 we may additionally assume that A is a normal subgroup of G . Note that we then have $\text{ord}(\psi) = 3$ for each $\psi \in \Sigma \setminus A$ by Lemma 5.5 and our assumption on $Z(G)$. In order to derive a contradiction, we establish two claims regarding the conjugacy classes of elements of A with specific geodesics, in particular, elements of the set $C := \{g \in A \mid \text{geod}(g) \equiv \psi_1\psi_2 \text{ with } \psi_1, \psi_2 \in \Sigma \setminus A\}$.

Claim (1). *If $x \in \Sigma \cap A$, then $x^G = \{x, \psi_1\psi_2, \psi_2\psi_1\}$ for some $\psi_1, \psi_2 \in \Sigma \setminus A$ with $\psi_1\psi_2, \psi_2\psi_1 \in C$.*

Since $A \leq G$ is a maximal subgroup, there exists some $\psi \in \Sigma \setminus A$. Because $G = A \cdot \{1, \psi, \psi^{-1}\}$, we know that $x^G = \{x, \psi^{-1}x\psi, \psi x\psi^{-1}\}$. We also note that $\psi x\psi x\psi x = (\psi x)^3 = 1$ as $\psi x \in G \setminus A$. Therefore, the element $\psi x\psi = x^{-1}\psi^{-1}x^{-1}$ has length at most two. In turn, we conclude that both of the elements $\psi^{-1}x\psi = \psi(\psi x\psi)$ and $\psi x\psi^{-1} = (\psi x\psi)\psi$ have length at most two.

If one of these elements, $\psi^{-1}x\psi$ say, had length one, then, so would $x\psi$ and $\psi^{-1}x$ by Lemma 2.3; hence, so would $\psi(x\psi) = x^{-1}(\psi^{-1}x^{-1})$ and $(\psi^{-1}x)\psi = (x^{-1}\psi)x^{-1}$; hence, so would ψx and $x\psi^{-1}$. It follows that the other element $(\psi x)\psi^{-1} = \psi(x\psi^{-1})$ would also have length one. But then all elements of the conjugacy class x^G would have length one, i.e., $x^G \subseteq \Sigma$. This contradicts Lemma 3.7.

Next, we assume that $\text{geod}(\psi^{-1}x\psi) \equiv y_1y_2$ with $y_1, y_2 \in \Sigma \cap A$. This implies $y_1, y_2 \in \{x^{\pm 1}\}$, for otherwise $\langle \Sigma \cap A \rangle \leq G$ is a complete subgroup by Lemma 3.7 and thus y_1y_2 has length one. If $\psi^{-1}x\psi = x^{\pm 2}$, then either $\psi x\psi^{-1} = \psi_1\psi_2 \in C$ or, for the same reason as above, $\psi x\psi^{-1} = x^{\mp 2}$.

In the second case, x is conjugate to x^{-1} by transitivity of conjugacy. Since x is the only element of length one in x^G , this implies $x = x^{-1}$. But then $\psi^{-1}x\psi = x^{\pm 2} = 1$, which is absurd.

In the first case, i.e., $x^G = \{x, x^{\pm 2}, \psi_1\psi_2\}$, we have $\psi^{-1}x\psi = x^{\pm 2}$ and, hence, $x = x^{\pm 8}$ as $\psi^3 = 1$. If $\psi^{-1}x\psi = x^{-2}$, then $x^9 = 1$ and thus $\psi^{-1}x^3\psi = x^{-6} = x^3 \in Z(G)$. But this implies $x^3 = 1$ and, therefore, $x^{-2} = x \in \Sigma$; a contradiction. In case $\psi^{-1}x\psi = x^2$, we first observe that $\psi_2\psi_1 = x$. Indeed, $\psi_2\psi_1$ is conjugate to x and cannot equal x^2 or $\psi_1\psi_2$ by uniqueness of geodesics. Using this, we obtain $\psi_2x\psi_2^{-1} = x^2$. Now $\psi_1x = \psi_2^{-1}\psi_2\psi_1x = \psi_2^{-1}x^2 = \psi_2^{-1}\psi_2x\psi_2^{-1} = x\psi_2^{-1} \in \Sigma$, since it has two expressions of length two. But then the same is true for $x^2 = \psi_2(\psi_1x)$; hence $x^2 \in \Sigma$, which contradicts our assumption that x^2 has length two.

The only remaining possibility is that $\text{geod}(\psi^{-1}x\psi) \equiv \psi_1\psi_2$ and, hence, $\text{geod}(\psi x\psi^{-1}) \equiv \psi_2\psi_1$ for some $\psi_1, \psi_2 \in \Sigma \setminus A$. This establishes Claim (1).

Claim (2). *If $\psi_1, \psi_2 \in \Sigma \setminus A$ with $g = \psi_1\psi_2 \in C$, then for some $x \in \Sigma \cap A$*

$$g^G = \{x, \psi_1\psi_2, \psi_2\psi_1\} \quad \text{or} \quad g^G = \{x^2, \psi_1\psi_2, \psi_2\psi_1\}.$$

Clearly, $\psi_1\psi_2$ and $\psi_2\psi_1$ are distinct elements of g^G . Moreover, $\psi_2\psi_1$ also has length two by Claim (1). The third element of g^G is $h := \psi_1^{-1}\psi_2\psi_1^{-1} = \psi_2^{-1}\psi_1\psi_2^{-1}$, which therefore has length at most two. The claim is trivial if h has length one. Suppose that h has length two and $\text{geod}(h) \equiv \psi'_1\psi'_2$ with $\psi'_1, \psi'_2 \in \Sigma \setminus A$. Clearly, $\psi'_2\psi'_1 \neq \psi'_1\psi'_2$ and $\psi'_2\psi'_1 \in g^G$. Moreover, $\psi'_2\psi'_1$ also has length two by what we have just shown. It follows that $\{\psi'_1, \psi'_2\} = \{\psi_1, \psi_2\}$, which is clearly impossible.

If $\text{geod}(h) \equiv y_1y_2$ with $y_1, y_2 \in \Sigma \cap A$, then $y_1 = y_2$ by uniqueness of geodesics and the fact that $y_1y_2 = y_2y_1$. Hence $h = x^2$ with $x = y_1 = y_2 \in \Sigma$. This completes our proof of the claim.

We finally derive the desired contradiction. To this end, recall that there exists some $\psi \in \Sigma \setminus A$. Furthermore, we cannot have $\Sigma = \{\psi^{\pm 1}\}$ for G is not cyclic. As such, we either have $\Sigma \cap A \neq \emptyset$ or there exists some $\psi' \in \Sigma \setminus A$ with $\psi' \notin \{\psi^{\pm 1}\}$. In the latter case, $\Sigma \cap A \neq \emptyset$ by Claim (2).

Now, choose some $x \in \Sigma \cap A$ and let $\psi_1, \psi_2 \in \Sigma \setminus A$ with $x^G = \{x, \psi_1\psi_2, \psi_2\psi_1\}$ as in Claim (1). We then have $x = \psi_1^{-1}\psi_2\psi_1^{-1} = \psi_2^{-1}\psi_1\psi_2^{-1}$. Let $\psi_3 := x\psi_1 = \psi_1^{-1}\psi_2 \in \Sigma$ and observe $\psi_3 \notin \{\psi_1^{\pm 1}, \psi_2\}$. In fact, we also have $\psi_3 = \psi_1^{-1}\psi_2 \neq \psi_2^{-1}$ for otherwise $\psi_2\psi_1 = 1$. Now note, that $\psi_3^{-1}\psi_2^{-1} = x$ and thus $\psi_2^{-1}\psi_3^{-1} \in x^G \setminus \{x\} = \{\psi_1\psi_2, \psi_2\psi_1\}$, which contradicts uniqueness of geodesics. \square

5.3. Nilpotent groups. In every nilpotent group, certain iterated commutators evaluate to central elements. This fact, together with our results concerning central elements developed in Section 3.4, imposes restrictions on the structure of a nilpotent group with an alleged non-complete geodesic Cayley graph. For groups of nilpotency class two, we obtain further restrictions based on a more detailed analysis of the involved commutator maps; see Corollary 5.16 below.

Recall that a group G is *nilpotent* if $G = 1$ or $G/Z(G)$ is nilpotent. If G is nilpotent, then there exists a number s such that $[g_1, \dots, g_{s+1}] := [[g_1, \dots, g_s], g_{s+1}] = 1$ for all $g_1, \dots, g_{s+1} \in G$. The smallest such number s is the *nilpotency class* of G . A group is nilpotent if and only if it is a direct product of p -groups; see [29, Theorem 5.2.4]. In particular, every nilpotent group of even order has even-order center.

Theorem 5.9. *Let G be a nilpotent but not cyclic group of nilpotency class s and suppose that*

$$p \nmid \frac{\exp(G)}{\exp(Z(G))}$$

for each odd prime $p < 3 \cdot 2^{s-1} - 2$. Then the only geodesic Cayley graph of G is the complete graph.

Proof. All finite nilpotent groups are direct products of p -groups. Thus, if 2 divides $|G|$, then there must be an order 2 element in the center and the statement follows from Theorem 3.14. If G is abelian, then the statement is a consequence of Corollary 3.9. We now assume that $s > 1$ and $2 \nmid |G|$, and that there exists a geodesic Cayley graph $\text{Cay}(G, \Sigma)$ which is not complete.

In every finite group of nilpotency class s , by [6, Lemma 2.6], there are generators $a_1, \dots, a_s \in \Sigma$ such that $z = [a_1, a_2, \dots, a_s] \neq 1$. Further, since G is nilpotent of class s , we have $z \in Z(G)$; hence, by Theorem 3.22, $z = a^q$ with $a \in \Sigma$ and $q \geq 3$ dividing the order of a in the quotient group $G/Z(G)$. Since a^q is a geodesic, its length is shorter than the length of the commutator, which is $3 \cdot 2^{s-1} - 2$.

Let p be a prime divisor of $q = pr$. Then p also divides $|G : Z(G)|$, $|G|$, and $|Z(G)|$. Furthermore, p is odd and $p \leq q < 3 \cdot 2^{s-1} - 2$. Let p^n be the largest power of p dividing $\exp(Z(G))$ (note that $n \geq 1$). Let $\tilde{z} \in Z(G)$ with $\text{ord}(\tilde{z}) = p^n$. Once more, $\tilde{z} = \tilde{a}^q$ for some $\tilde{a} \in \Sigma$ by Theorem 3.22. Now consider the element $b = \tilde{a}^r$ (where $q = pr$ as above). Clearly, $(b^p)^{p^n} = \tilde{z}^{p^n} = 1$. As such, $\text{ord}(b) = p^{n+1}$ and, thus, p^{n+1} divides $\exp(G)$. We conclude that p divides $\exp(G)/\exp(Z(G))$, which contradicts our assumption on G . \square

Remark 5.10. The condition in Theorem 5.9 is satisfied if no odd prime $p < 3 \cdot 2^{s-1} - 2$ divides the order of G or, more generally, if p^2 does not divide $\exp(G)$ for any such prime p . If, for example, G is a nilpotent group of nilpotency class two with $9 \nmid \exp(G)$, then G satisfies Conjecture A; if G is of nilpotency class three and neither 3, 5, nor 7 divide $|G|$, then G satisfies Conjecture A.

To further explore the applicability of Theorem 5.9, as well as the following Corollary 5.16, we have examined all non-abelian groups of order p^k with $p \in \{3, 5, 7, 11\}$ and $k \leq 7$ in GAP [18] using its SmallGrp library [2]. The results of this examination are summarized in Table 1.

We now turn to the case of nilpotency class two. In the following we always assume that $\text{Cay}(G, \Sigma)$ is non-complete geodesic Cayley graph of a nilpotent group G of nilpotency class 2. We will consider the

p	$s \leq 1$	$s = 2$	$s = 3$	$s = 4$	$s = 5$	$s = 6$
3	45	587 / 1,926	150 / 6,362	36 / 1,386	0 / 180	0 / 6
5	45	7,256	247 / 23,073	119 / 3,382	5 / 1,227	0 / 9
7	45	26,914	255 / 76,783	131 / 8,034	60 / 2,140	15 / 198
11	45	204,912	514,627	139 / 26,882	62 / 5,170	19 / 402

TABLE 1. The number of groups of order p^k with $k \leq 7$ and nilpotency class s covered by Theorem 5.9 and Corollary 5.16 / total number of such groups (if distinct).

following subsets of Σ :

$$\Sigma_i := \{x \in \Sigma \mid \text{ord}(x) = 3^i\} \quad \text{and} \quad \Delta := \{x \in \Sigma \mid x^3 \in Z(G) \setminus 1\}.$$

Since $[x, y] \in Z(G) \setminus 1$ for some $x, y \in \Sigma$ (see the proof of Theorem 5.9), the length of central geodesics is three according to Theorem 3.22. As such, the map $\alpha: G \rightarrow G$ given by $g \mapsto g^3$ induces a bijection from Δ onto $Z(G) \setminus 1$. In particular, we conclude that $\Delta \neq \emptyset$. As there exists an element of order three in $Z(G)$, we must also have $\Delta \cap \Sigma_2 \neq \emptyset$.

Lemma 5.11. *Let $x, y \in \Sigma$ with $x \in \Delta$ or $y \in \Delta$. Then $[x, y] = 1$ if and only if $\{x^{\pm 1}\} = \{y^{\pm 1}\}$.*

Proof. If $[x, y] = 1$ and $\{x^{\pm 1}\} \neq \{y^{\pm 1}\}$, then $\langle x, y \rangle \leq G$ is a complete subgroup by Lemma 3.7. If, furthermore, $x \in \Delta$, then $x^3 \in Z(G) \cap \Sigma$. But then $\text{Cay}(G, \Sigma)$ is complete by Corollary 3.11. \square

Lemma 5.12. *Let $x \in \Sigma$ and $y_1, y_2 \in \Sigma \setminus \{x^{\pm 1}\}$ with $y_1 \in \Delta$. Then $[x, y_1] = [x, y_2]$ implies $y_1 = y_2$.*

Proof. Let $z := [x, y_1] = [x, y_2] \in Z(G)$. Then $z \neq 1$ by Lemma 5.11 and, thus, $z = a^3$ with $a \in \Delta$. Assuming $y_1 \neq y_2$, the element $xz = x^{y_1} = x^{y_2}$ has two distinct representatives of length three; therefore, it has length at most two. But then $\bar{x} = a$ by uniqueness of geodesics, as $\bar{x}(xz) = z = a^3$. In particular, this shows that $x = \bar{a} \in \Delta$ and that $x^y = xz = x^{-2}$ for $y \in \{y_1, y_2\}$.

Let $\text{ord}(x) = 2k + 1$ and note that $k \geq 4$ as $x \in \Delta$. Then $(x^k)^y = x^{-2k} = x$ for $y \in \{y_1, y_2\}$. As such, $x^k = y_1 x y_1^{-1} = y_2 x y_2^{-1}$ has length at most two. Since $x^k \notin \{1, x^{\pm 1}, x^{\pm 2}\}$, we conclude that all powers of x , in particular $z = x^{-3}$, have length at most two by Lemma 3.20; a contradiction! \square

As usual, we denote the *commutator (or derived) subgroup* of G by $G' = \langle [a, b]; a, b \in G \rangle$, and the *Frattini subgroup* of G by $\Phi(G)$, i.e., $\Phi(G)$ is the intersection of all maximal subgroups of G .

Using the above, we will now show that a nilpotent group G of class two that violates Conjecture A would have to be *special*, i.e., it would satisfy $G' = \Phi(G) = Z(G)$. In some sense, such groups are as non-abelian as possible given the constraint $G' \leq Z(G)$ imposed by G being nilpotent of class two.

Proposition 5.13. *If G is a nilpotent group of nilpotency class 2 which is not a special 3-group with $\exp(G) = 9$, then G satisfies Conjecture A.*

Proof. Let $H_1 := \langle \Sigma \setminus \Delta \rangle$ and $H_2 := \langle \Delta \rangle$. Clearly $H_2' \leq G' \leq Z(G) \leq H_2 \leq G$. Furthermore, the quotient $H_2/Z(G)$ is an elementary abelian 3-group.

We first assume that there exists an element $u \in \Sigma \setminus \Delta$ and consider the homomorphism

$$\phi: G \rightarrow Z(G); \quad g \mapsto [u, g].$$

By Lemma 5.11 and Lemma 5.12, ϕ is injective on Δ . Since $|\Delta| = |Z(G) \setminus 1|$, we have $\phi(\Delta) = Z(G) \setminus 1$ and, therefore, $\phi(H_2) = \text{Im}(\phi) = Z(G)$. Moreover, $\phi(\Sigma \setminus \Delta) = 1$ by Lemma 5.12, i.e., $H_1 \leq \text{Ker}(\phi)$.

The homomorphism ϕ factors through the quotient $G \rightarrow G/Z(G)$ and the inclusion $G' \rightarrow Z(G)$. As such, $G' = Z(G)$ is isomorphic to a quotient of $H_2/Z(G)$; hence, $Z(G)$ is an elementary abelian 3-group. It follows that $\Delta \subseteq \Sigma_2$, i.e., $\text{ord}(x) = 9$ holds for all $x \in \Delta$. On the other hand, we have $u^3 \in Z(G)$ since $[u^3, x] = [u, x]^3 = \phi(x)^3 = 1$ for each $x \in \Sigma$. This implies $u^3 = 1$ since $u \in \Sigma \setminus \Delta$. Moreover, the same holds for all $u \in \Sigma \setminus \Delta$, that is $\Sigma \setminus \Delta = \Sigma_1$ and, thus, $\Delta = \Sigma_2$. This implies that $G/Z(G)$ is an elementary abelian 3-group. Hence, $\exp(G) = 9$ and $G' = \Phi(G) = Z(G)$ as claimed.

We now deal with the case $\Sigma \setminus \Delta = \emptyset$, i.e., $\Sigma = \Delta$ and $H_2 = G$. As such, $G/Z(G)$ is an elementary abelian 3-group and thus so is $Z(G)$. Therefore, $\Sigma_1 = \Sigma \setminus \Delta = \emptyset$ and $\Sigma_2 = \Delta = \Sigma$. It remains to show that $G' = Z(G)$. To this end, fix $x \in \Sigma$ and consider the homomorphism

$$\psi: G \rightarrow Z(G); \quad g \mapsto [x, g].$$

By Lemma 5.11 and Lemma 5.12, ψ is injective on $\Sigma \setminus \{x^{\pm 1}\}$. But then $Z(G) \setminus \psi(G)$ contains at most two elements. Since $\psi(G) \leq G' \leq Z(G)$ and $4 \leq |\Sigma| < |Z(G)|$, there are still enough elements in $\psi(G)$ to generate $Z(G)$. We therefore conclude that $\psi(G) = G' = Z(G)$. \square

Proposition 5.14. *The following hold.*

- (1) $\Sigma = \Sigma_1 \cup \Sigma_2$ where $\Sigma_i := \{x \in \Sigma \mid \text{ord}(x) = 3^i\}$ and $x^3 \in Z(G)$ for all $x \in \Sigma$.
- (2) The subgroup $H_1 := \langle \Sigma_1 \rangle \leq G$ is either trivial or $H_1 \cong C_3$.

Proof. The first statement is obtained directly as part of the proof of Proposition 5.13. It remains to justify the second statement. To this end, we assume that $H_1 \neq 1$ and $H_1 \not\cong C_3$. Then there exist elements $y_1, y_2 \in \Sigma_1$ with $\{y_1^{\pm 1}\} \neq \{y_2^{\pm 1}\}$ and $x \in \Sigma_2$ such that $z := [x, y_1] = [x, y_2] \neq 1$. Indeed, for any given $x \in \Sigma_2$, we must have $[x, \Sigma_1] \subseteq Z(G) \setminus [x, \Sigma_2]$ by Lemma 5.11 and Lemma 5.12 and the latter set has (at most) two elements. As in the proof of Lemma 5.12 we conclude that $z = x^3$.

Finally, we have $x^4 = x^{y_1^{-1}} = x^{y_2^{-1}}$, as $\text{ord}(x) = 9$, and this element has length at most two. By uniqueness of the geodesic representative of $x^3 = x^{-1}(x^4)$, we obtain $x = x^{-1}$; a contradiction! \square

Corollary 5.15. *If G is an extra-special 3-group, i.e., $G' = \Phi(G) = Z(G) \cong C_3$, then the only geodesic Cayley graph of G is the complete graph.*

Proof. Assume, for the sake of deriving a contradiction, that $Z(G) \cong C_3$. Then $|\Sigma| = |\Sigma_1| + |\Sigma_2| \leq 4$. Clearly, $|\Sigma| \geq 4$ as G is not cyclic. Therefore, $|\Sigma_1| = |\Sigma_2| = 2$. By Lemma 5.11, $[x, y] = x^{-3}$ for some $x \in \Sigma_2$ and $y \in \Sigma_1$. Then $yxxy = y^{-1}x^{-2}$ has length at most two; hence, $yxxy^{-1} = (yxxy)y = x^4$ has length at most two; hence, $x^3 = x^{-1}(x^4)$ has length at most two. This contradicts Theorem 3.22. \square

Corollary 5.16 (Nilpotent class 2). *Let G be a nilpotent group of nilpotency class 2 which is either*

- *not a special 3-group, i.e., $G' = \Phi(G) = Z(G)$, with $\exp(G) = 9$, or*
- *an extra-special 3-group, i.e., $G' = \Phi(G) = Z(G) \cong C_3$.*

Then G satisfies Conjecture A.

In other words, the case we cannot handle theoretically is nilpotent of class 2 with $\exp(G) = 9$ and $G' = \Phi(G) = Z(G) \not\cong C_3$.

5.4. Groups with large commutativity degree. Many of our results address Conjecture A in the case of groups that are, in some sense, close to being abelian. This property can be quantified by a group G 's *commutativity degree* $\mathbf{P}(G)$, which is the probability that two randomly chosen elements of G commute, i.e.,

$$\mathbf{P}(G) := \frac{|\{(g, h) \in G \times G \mid gh = hg\}|}{|G \times G|}.$$

The interested reader is referred to the survey by Das, Nath, and Pournaki [8] for further details and historical context. We will content ourselves here with the following observation.

Theorem 5.17. *Conjecture A holds for every group G with $\mathbf{P}(G) > \frac{11}{32}$.*

Proof. For abelian groups, this follows from Corollary 3.9 (see also [15]). If G is non-abelian and $\mathbf{P}(G) > \frac{11}{32}$, then according to Rusin [30, p. 246] the structure of G must be as indicated in Table 2.

In most of these cases, $Z(G)$ contains a subgroup isomorphic to C_2 ; hence, Theorem 3.14 applies. If $G/Z(G) \cong S_3$ or $G/Z(G) \cong D_{10}$, then G contains an abelian subgroup of index two corresponding to $C_3 \leq S_3$ or $C_5 \leq D_{10}$ in $G/Z(G)$, respectively. As such, we can apply Theorem 5.4. In the remaining case, i.e., if $C_3 \cong G' \leq Z(G)$, then G is covered by Corollary 5.16. \square

Group Structure			Applicable Result
G'	$G' \cap Z(G)$	$G/Z(G)$	
C_2	C_2	C_2^{2r}	Theorem 3.14
C_3	1	S_3	Theorem 5.4
C_2^2 or C_4	C_2	D_8	Theorem 3.14
C_2^2	C_2	C_2^3 or C_2^4	Theorem 3.14
C_3	C_3	C_3^2	Corollary 5.16
C_5	1	D_{10}	Theorem 5.4
C_2^2	C_2^2	C_2^3 or C_2^4	Theorem 3.14
C_6	C_2	$S_3 \times C_2$ or $C_3 \rtimes C_4$	Theorem 3.14

TABLE 2. The possible structures of a non-abelian group G with $\mathbf{P}(G) > \frac{11}{32}$.

6. EXPERIMENTS.

We now turn to the exhaustive computer search to check which groups of order up to 1024, as well as all even orders up to 2014 and all non-abelian finite simple groups up to order 5000, have a generating set that yields a geodetic Cayley graph. The aim of this experiment was to either verify Conjecture A for as many group orders as possible, or to find a group and a generating set that yields a non-trivial geodetic Cayley graph, i.e., a graph that is neither complete nor an odd cycle. For our code, see

https://osf.io/9ay6s/?view_only=37e18301e4e74e12bfe4e07b90b924c0.

Improving the computer search has been a major motivation for the theoretical work in the previous sections. An important result in this regard are the bounds on the generating set discussed in Section 4. Another are the results covering entire classes of groups, the most important of which is Theorem 3.14 excluding all groups with even-order center.

In turn, the results from the computer search also influenced our theoretical work. For example, at one point, groups which were semidirect products with C_2 had a long running time in our computer search. This directed our theoretical work to focus on such groups, leading up to Theorem 5.4 showing that Conjecture A holds for all groups with an abelian subgroup of index two.

6.1. Overview. Our approach consists of three stages. The first is the *filtering stage* in which we identify the relevant groups which are not covered by our theoretical results, and thus need to be considered in our computer search. We realized this stage using GAP [18].

The second stage is a *preprocessing stage*, also realized in GAP, during which we compute the information required for the search and store it in a JSON file. Besides the multiplication table describing the group, the most important information computed in this stage comprised a set of *forbidden elements* and a set of *required subsets*.

Forbidden elements are elements which cannot be part of any geodetic generating set except for the complete one. A required subset is a set of elements of which each geodetic generating set needs to contain at least one. We will give more details on how we compute and use these below.

The third and final stage is the actual search, which we have implemented in Rust. Obviously, enumerating all generating sets is infeasible even for relatively small groups. For example, already for the symmetric group S_5 , which has 120 elements, there are 2^{72} potential generating sets (symmetric subsets not containing the neutral element). To circumvent this problem, we discard generating sets based on the theoretical results described in the previous sections. We implement this using a binary counter for enumerating generating sets, which allows us to systematically skip the respective ranges. For each of the remaining generating sets, we test whether or not the resulting Cayley graph is geodetic and report those that are.

6.2. Filtering. We used GAP [18] and its SmallGrp library [2] to obtain a list of all finite groups up to order 1024 relevant to our search. (In a second run we repeated the experiment also filtering out all odd-ordered groups, which we report on below.) When generating this list of groups, we ignored the following.

- All abelian groups (Corollary 3.9).
- All groups with even-order center (Theorem 3.14).
- All groups with a large center (Corollary 4.8 and Proposition 4.10).

Total Number of Groups	1,206,579
Abelian Groups	2,034
Groups with Center of Even Order	1,200,151
Groups with Abelian Subgroup of Index 2	989
Nilpotent Groups as in Corollary 5.16	170
Nilpotent Groups as in Theorem 5.9	18
Groups with Abelian Subgroup of Index 3 as in Theorem 5.8	86
Groups with Large Center as in Corollary 4.8 and Proposition 4.10	274
Remaining Groups	3,197

TABLE 3. Number of groups of order up to 1024 and excluding 2-groups which are caught by the different filtering steps. The filtering is performed in the same order as in the table, and each group is only counted towards the first category it matches.

- All groups with abelian index-2 subgroups (Theorem 5.4).
- The groups with abelian index-3 subgroups covered by Theorem 5.8.
- The nilpotent groups covered by Theorem 5.9 and Corollary 5.16.

Note that, while there are approximately $50 \cdot 10^9$ groups up to order 1024, most of those are 2-groups, which all have an even-order center. Excluding the 2-groups there are only 1206579 groups of order at most 1024, and after excluding the groups in the above list only 3197 groups remain. We provide more details on the number of groups falling into each of the above categories in Table 3.

Remark 6.1. The smallest group with a center larger than the bound from Corollary 4.8 is $C_{13} \times A_4$. Its center has order 13 – just above the bound, which in this case is 12.

To find an example for Proposition 4.10, we have to look a bit further. The proposition excludes 3 as prime divisor in the order of the group, and, 5 and 7 are excluded as prime divisors in the index of the center. The smallest group covered by Proposition 4.10 that is not already covered by Corollary 4.8 is $C_7 \times (C_{13} \times C_4)$. Its center has order 7, just larger than the bound of 6.

During filtering we only verify those bounds on the size of the center based on Corollary 4.8 and Proposition 4.10 (with $m_0 = 3$). More precise bounds on the size of non-complete geodetic generating sets, and thus also on the size of the center, are computed within the initialization step of our search algorithm. Therein, we utilize most results of Section 4 (with optimal parameters). Due to tighter bounds, we were able to exclude an additional 240 groups from the search.

Remark 6.2. For filtering the groups of even order up to 2014, we use the same method. However, there is one special case: the groups of order $1536 = 3 \cdot 2^9$. There are more than $4 \cdot 10^8$ of them – too many for running through the entire filtering stage. However, most of them have a normal Sylow 3-subgroup. Because such groups have an even-ordered center, they can be safely ignored by Theorem 3.14. Therefore, we only run the filtering procedure for the remaining groups, indexed 408526598–408641062 in the SmallGrp library.

6.3. Preprocessing. The main objective of the preprocessing stage is to compute the forbidden elements and the required subsets of each group. As mentioned above, an element is forbidden if, whenever it is part of a geodetic generating set, the associated Cayley graph is necessarily complete. Note that, since we filter out abelian groups in the filtering stage, here we do not need to consider the possibility that the Cayley graph is an odd cycle. The set of forbidden elements comprises all elements $g \in G$ such that $h = g$ or $h = g^2$ is nontrivial and satisfies $h^G \subseteq \{h^{\pm 1}\}$; see Lemma 3.7 and Lemma 3.19. In particular, this includes central elements (Corollary 3.11) and their square roots (Theorem 3.22).

A required subset is a set of which each geodetic generating set needs to contain at least one element. In the preprocessing we compute the following sets, which we know to be required.

- Each conjugacy class of elements of order two (Lemma 3.13).
- Each normal subgroup of index two (Lemma 5.1).
- Each complement of a maximal subgroup (as we want a generating set).

There is one other family of required subsets: the potential roots of each central element. However, since these sets are smaller, and thus their inclusion is more effective, when we know the diameter of the center, we do not add these sets in the preprocessing stage, but later in the search algorithm.

At this stage we also take advantage of automorphisms to reduce the size of the required subsets and thus reduce the number of generating sets we need to look at. The procedure is to go through the required subsets one by one. For each, we compute the orbits of its elements under the automorphism group and select one element from each orbit. As we are interested only in symmetric generating sets, we consider the orbits of an element and its inverse as a single orbit. Continuing with the next required subset, we no longer use the full automorphism group, but only the point-wise stabilizer of the elements that were selected in the previous required subsets. This is justified by the following straight-forward observation:

Lemma 6.3. *Let $R \subseteq G$ and $Y \subseteq G$ and let $\tilde{R} \subseteq R$ be a system of representatives of $R/\text{Stab}(Y)$. Then for all $X \subseteq G$ with $R \cap X \neq \emptyset$, there exists some $\phi \in \text{Stab}(Y)$ such that $\phi(X) \cap \tilde{R} \neq \emptyset$.*

In cases where the required subset contains only part of an orbit under the action of the automorphism group, we take care to select representatives that are part of the original set. This way, for each generating set that contains an element of each of the original required subsets, there is a generating set which contains at least one element of each of the smaller required subsets obtained after applying the automorphism group.

Finally, we discard required subsets that are supersets of smaller ones.

Remark 6.4. For the finite simple groups $\text{PSL}(2, q)$ with $q \in \{17, 19, 16\}$, for parallelization, we split the computation of the search algorithm into several chunks. This is implemented by generating several instances during the preprocessing stage with different required and forbidden subsets.

6.4. The search algorithm. For a group G , a set of *candidates* is a subset $C \subseteq G$ such that C contains exactly one representative of a non-trivial group element and its inverse. Thus if $g \in G$ has order two, then $g \in C$, and otherwise we choose either g or g^{-1} to belong to C . Note that if more than half of the elements of G are of order two, then, according to Liebeck and MacHale [21], G has an abelian subgroup of index two or $Z(G)$ has even order. As such groups were excluded during the filtering stage, we may assume at most half the elements of G have order two; hence, $|C| \leq \frac{3}{4}|G|$.

In the following, we identify C with the numbers $\{0, \dots, |C| - 1\}$ (in particular, we fix an order on C). We enumerate the generating sets of a group by enumerating all subsets of C in the order of a binary counter from 0 to $2^{|C|}$ (where a binary number naturally corresponds to a subset of C). The main loop of our search algorithm is presented in Algorithm 1. Here we write the counter as a subset $X \subseteq \{0, \dots, |C| - 1\}$. Incrementing the binary counter is done in lines 6 to 14. In line 18 we check whether the resulting Cayley graph is geodesic (`TESTGEODETIC`, for details see below). We use a stack to keep track of the candidate set X , the corresponding increment I , and some additional information that is not displayed in Algorithm 1. During the execution of the algorithm, we maintain the following invariants of the stack (which we consider to grow upwards).

- (I₁) If (X, I) is above (X', I') , then $X \supseteq X'$ and $I < I'$. Moreover, $X \cap [I', |C|] = X' \cap [I', |C|]$.
- (I₂) If (X, I) is directly above (X', I') , then additionally $X \cap [I, |C|] = (X' \cap [I, |C|]) \cup \{I\}$.

We use several pruning methods to shortcut the counting process. The variable I_{next} in Algorithm 1 serves as the index of the bit to be increased next (meaning that $I_{\text{next}} = 0$ yields a usual increment by one); by increasing I_{next} , we can skip over certain values for the counter.

The pruning methods, described in detail below, rely upon

- bounds on the size of the generating set,
- the forbidden candidate array (for simplicity not included in the pseudocode Algorithm 1),
- the required subsets described above in Section 6.3 (handled in line 5 in Algorithm 1),
- saturating the generating set (`FILLIN`),
- handling of collisions at distance 3 discovered during the check whether the Cayley graph is geodesic (`HANDLECOLLISIONS`).

Finally, for groups of even order and groups with non-trivial center, we employ further pruning techniques during `TESTGEODETIC` and `HANDLECOLLISIONS`.

Bounds on the number of generators. In Section 4 we developed several bounds on the size of non-complete geodesic generating sets. The most general such bound is due to Theorem 4.4: it gives us an upper bound of

<pre> procedure CHECKGROUP(G) 1: $(X, I) \leftarrow (\emptyset, \infty)$ 2: Stack.PUSH(X, I) 3: $I_{\text{next}} \leftarrow 0$ 4: while Stack $\neq \emptyset$ do 5: $I_{\text{next}} \leftarrow \max\{I_{\text{next}}, \text{REQUIRED}(X)\}$ 6: while $I_{\text{next}} \in X$ do 7: $I_{\text{next}} \leftarrow I_{\text{next}} + 1$ 8: if $I_{\text{next}} \geq C$ then 9: return 10: while $I_{\text{next}} \geq I$ do 11: $(X, I) \leftarrow \text{Stack.POP}()$ 12: $I_{\text{last}} \leftarrow I$ 13: $(X, I) \leftarrow (X \cup \{I_{\text{next}}\}, I_{\text{next}})$ 14: Stack.PUSH(X, I) 15: $I_{\text{next}} \leftarrow \text{CHECKGENSET}(X, I, I_{\text{last}})$ </pre>	<pre> procedure CHECKGENSET(X, I, I_{last}) 16: if FILLIN(X, I, I_{last}) fails then 17: return I 18: $T \leftarrow \text{TESTGEODETIC}(X)$ 19: if $T = \text{GEODETIC}$ then 20: output X 21: return 0 22: if HANDLECOLLISIONS(T, X, I, I_{last}) fails then 23: return I 24: else 25: return 0 </pre>
---	---

ALGORITHM 1. Outline of our search algorithm.

$c \cdot \sqrt{n}$ where n is the order of the group. The constant factor c is at most $2/\sqrt{5}$ but, depending on the size of the group, the factor can be even smaller. In fact, we compute the optimal bound given by Theorem 4.4 in our program. If the group has non-trivial center, we use Theorem 4.7 to obtain an even smaller bound in the order of $\sqrt[3]{n}$. We use these bounds to prune the search tree as follows: whenever there are too many bits set to one in the counter (i.e., $|X|$ is too big – detected either in FILLIN or HANDLECOLLISIONS), we increment the least-significant bit currently set to one by setting $I_{\text{next}} \leftarrow I$ (see lines 17 and 23 of Algorithm 1).

In some cases, the bound computed at this stage is even impossible to satisfy and, therefore, the group possess no geodetic generating set other than the complete one. In this way, we exclude an additional 240 (of the remaining 3197) groups from the search.

Forbidden candidates. As a first improvement, we use a bit array indicating candidates that are forbidden given the other candidates already contained in X . For each entry on the stack (stack frame) we keep a separate forbidden candidate array. Initially, the forbidden candidate array of the first stack frame comprises the forbidden candidates computed during preprocessing (Section 6.3). When creating a new stack frame, the forbidden candidates of its predecessor are copied; afterwards additional candidates might be marked as forbidden on the new stack frame. An additional candidate can be marked as forbidden if its inclusion in X would

- lead to a generating set that is too large,
- require the inclusion of a candidate that is already forbidden, or
- violate the order in which we search through the generating sets.

These conditions are tested for during the FILLIN and HANDLECOLLISIONS procedures. For details we refer to the respective paragraphs below.

We use the forbidden candidate array in the FILLIN and HANDLECOLLISIONS procedures as well as in the counter logic. While omitted from the description in Algorithm 1 for simplicity, its incorporation is rather straightforward: whenever I_{next} is a forbidden candidate, it is incremented.

Required subsets. We incorporate the required subsets computed during the preprocessing step as detailed in Section 6.3. Whenever there is an unmet required subset, we increase I_{next} to the smallest candidate of this required subset (see line 5 of Algorithm 1). This dispenses with all candidate sets in between which do not contain an element of the required subset and, thus, need not be considered.

If multiple required subsets are unmet, we apply the following heuristic: choose a required subset to be satisfied first such that the smallest candidate contained in it is maximal.

It may happen that the smallest candidate of a required subset is forbidden. In this case we simply consider the smallest candidate I in the required subset that is not forbidden. If I is larger than I_{last} , the candidate of the previous stack frame, then we cannot satisfy the required subset by pushing a new stack frame without violating the order in which we search through the generating sets (Invariant (I₁)). Therefore,

we assign $I_{\text{next}} \leftarrow I_{\text{last}}$ in order to skip a number of candidate sets to which we cannot add any candidate from the required subset.

Finally, we remark that further required subsets are created during the `HANDLECOLLISIONS` procedure and at the beginning of the search for groups with non-trivial center.

Saturating the generating set (FILLIN). The most crucial improvement of our algorithm over a naïve search is based on Lemma 2.3. If $a, b, c, d \in \Sigma$ with $ab = cd$ for a geodesic generating set $\Sigma \subseteq G$, then $(ab), (c^{-1}a) \in \Sigma$. Therefore, whenever we add a new candidate to X , we test whether a product of the newly added element(s) and some other element of the generating set Σ associated with X equals a different product of two elements of Σ . If so, we also add this product to Σ by adding the corresponding candidate to X . By repeating this we obtain a candidate set X' . If the corresponding generating set Σ' is too large, or if X' contains a forbidden candidate, then the `FILLIN` procedure fails and marks the current increment I as forbidden on the previous stack frame.

In order to avoid considering the same generating set multiple times, we only want to add a candidate J to X during the `FILLIN` procedure if $J < I$; otherwise, the resulting candidate set would be considered again after incrementing the counter.

As such, if X' contains a candidate J with $I < J$, then the `FILLIN` procedure fails and, thus, in line 17 of Algorithm 1 we set $I_{\text{next}} \leftarrow I$ skipping a number of candidate sets that would violate the search order. Moreover, if X' contains a candidate J with $I_{\text{last}} < J$, then we also mark I as forbidden on the previous stack frame, which corresponds to I_{last} . This is because introducing I as a candidate in any stack frame above the previous one would violate the second part of Invariant (I₁).

Testing whether the Cayley graph is geodesic (TESTGEODETIC). As the Cayley graph is vertex-transitive, it suffices to check whether geodesics from the origin $1 \in G$ to each other vertex are unique. We implemented this using a breadth-first search. If during this breadth-first search we encounter an element with two different geodesics of length three, then we record this element for later handling. We collect a certain number of such elements depending on the order of the group and other parameters such as, e.g., the order of the center and the number of elements of order two. In the case of groups with non-trivial center we implemented some further checks as detailed below.

Handling collisions at distance three (HANDLECOLLISIONS). While testing whether the Cayley graph is geodesic, we compute a list of elements which have multiple geodesics of length three. We aim to extend the generating set to a geodesic generating set. To achieve this, we need to add new generators (by adding the respective candidates) such that each element in the list either becomes a generator or can be uniquely written as a product of two generators.

Since there are many options as for which generators to add, for each of the recorded collisions, we create a temporary required subset. It remains valid until the current candidate is removed from the candidate set, i.e., the lifetime of these new required subsets are tied to the current stack frame.

The temporary required subset associated with a collision is constructed as follows. Suppose that adding a single candidate J to X would resolve the collision. Then we insert J into the required subset, unless invoking `FILLIN` with $X' = X \cup \{J\}$ fails. For pairs of candidates whose addition would resolve the collision, we proceed similarly, but only add the larger of the two candidates to the required subset.

Finally, if all calls to `FILLIN` failed, and thus the required subset is empty, then the call to `HANDLECOLLISIONS` fails as we cannot resolve the collision. Crucially, if all calls to `FILLIN` forbade the corresponding inclusion, then we also forbid I on the stack frame corresponding to I_{last} .

Modifications for groups of even order. If during `TESTGEODETIC` we find an element of order two among the collisions at distance three, then the only choice is to add the element itself to the generating set. Recall that, by Lemma 3.13, such an element cannot have a geodesic of length two. As such, we do not construct a required subset in this case (by calling `HANDLECOLLISIONS`), but add the respective candidates directly to the candidate set instead (or possibly forbid the current increment I) and continue afterwards with `FILLIN`.

Based on the same observation, as a further improvement for groups of even order (more precisely, for groups whose order is divisible by six) we have implemented the following. During `FILLIN`, we check whether a generator is of order six, and thus generates a subgroup isomorphic to C_6 , or whether two generators together generate a subgroup isomorphic to S_3 . If so, we add all non-trivial elements of the corresponding

subgroup to the generating set Σ since such a subgroup is complete with respect to any geodetic generating set.

Modifications for groups with non-trivial center. For groups with non-trivial center, we compute all possible diameters of the center given by Theorem 3.22. Then we run the search repeatedly, once for each central diameter. We add the corresponding central roots as required subsets and add the central roots which are too short as forbidden elements. When testing whether the Cayley graph is geodetic, we additionally check whether there are central elements with geodesics shorter than the center diameter. If that happens, then the only geodetic graph the Cayley graph can be extended to by adding more generators is the complete graph or one with smaller central diameter.

6.5. Experimental results. Our experiments were conducted on a machine with an AMD Ryzen 9 5900X CPU (12 cores, 24 threads, 3.7 GHz) and 128 GB of RAM. Running the experiments for all groups up to order 1024 took 299 hours of total CPU time, and we were able to establish

Theorem B. *All groups up to size 1024 satisfy Conjecture A.*

With parallelization, the running time was dominated by a few “difficult” groups. See Table 4 for the running time of select groups and Figure 3 for a plot of the running time of every group of order up to 1024 compared to the group’s order.




FIGURE 3. Running time of our computer search for all groups of order below 1024; groups of even order are marked in black, those of odd order are marked in red.

As it can be seen from Table 4 and Figure 3, our search is substantially faster for groups of even order. Therefore, for groups of even order we extended our search up to order 2014. We did not go beyond that since larger groups are not completely listed in the GAP SmallGrp library [2]. Moreover, note that, as detailed in Remark 6.2, for groups of order 1536, we had to take some special care during the filtering stage.

Theorem 6.5. *Conjecture A holds for all groups of even order at most 2014.*

In the light of Corollary 3.8 and Theorem 5.9 it seems reasonable to search for counterexamples to Conjecture A within classes of groups that are far from commutative. Thus, it is natural to consider non-abelian finite simple groups. Nevertheless, in our experiments these groups turned out to be even easier to handle than many of the other groups of even order. (Note that by the famous Feit-Thompson theorem [16] all non-abelian simple groups have even order.) Indeed, for this special case we could go far beyond order 2014 and succeeded to show the following.

Theorem 6.6. *Conjecture A holds for the groups $\text{PSL}(2, 17)$, A_7 , $\text{PSL}(2, 19)$, $\text{PSL}(2, 16)$ and S_7 . In particular, it holds for all simple groups of order at most 5000.*

To give an indication which groups are “easy” and which are “difficult”, we give an overview on some selected groups in Table 4. The table contains the following classes of groups:

- alternating and symmetric groups,
- further finite simple groups of order exceeding 2000,
- the five groups with the longest running time,
- different groups of order $729 = 3^6$,
- groups of even order below 1024 with the longest running time.

Note that the order of the groups certainly plays a role for the running time, but there was also a huge variance in running time for different groups of roughly the same order. For instance this can be seen very prominently for groups of order 729 in Fig. 3.

There are various reasons for this difference. For example we observe an impact of the size of the center, which is to be expected given the results in Section 4. Moreover, we see the clear effect of the groups having even order: for these we have additional possibilities to generate small required subsets and we can use the improved FILLIN procedure. However, also among groups of odd order and with trivial center there is a huge variation in running time. We observe that the most difficult instances are groups $C_p \rtimes C_n$ with a faithful action, p prime, and n as large as possible.

Additional statistics on our experiments may be found at

https://osf.io/9ay6s/?view_only=37e18301e4e74e12bfe4e07b90b924c0.

7. DISCUSSION

We have shown that for several infinite classes of finite groups there are no geodesic Cayley graphs except the complete graphs. This includes all abelian groups (except cyclic groups of odd order), dihedral groups, and groups with even-order center, as well as many nilpotent groups. Moreover, we have verified by a computer search that Conjecture A holds for all groups up to order 1024, all groups of even order up to 2014, all simple groups of order below 5000, and the symmetric group S_7 .

The main open problem, of course, remains whether Conjecture A holds for all finite groups, i.e., that every geodesic Cayley graph of a finite group is either complete or a cycle of odd length. Our experiments suggest that it might be reasonable to aim for proving Conjecture A for all groups of even order.

8. ACKNOWLEDGMENTS

The authors sincerely thank Alexey Talambutsa for pointing out previous work. In completing this work, the first two authors were supported by Australian Research Council grant DP210100271 and the fifth author was partially supported by DFG grant WE 6835/1–2.

9. DISCLOSURE STATEMENT

The authors report there are no competing interests to declare.

REFERENCES

- [1] Eiichi Bannai and Tatsuro Ito. On finite Moore graphs. *Journal of the Faculty of Science, the University of Tokyo. Sect. 1 A, Mathematics*, 20:191–208, 1973. URL: <https://api.semanticscholar.org/CorpusID:118193462>.
- [2] H. U. Besche, B. Eick, E. O’Brien, and M. Horn. SmallGrp, the GAP small groups library, Version 1.5.3. <https://gap-packages.github.io/smallgrp/>, May 2023. Refereed GAP package.
- [3] Juraj Bosák, Anton Kotzig, and Štefan Znám. Strongly geodesic graphs. *J. Combin. Theory*, 5:170–176, 1968.

Group	Order	Index	Sets tested	Duration
A_5	60	5	31	0.3 ms
S_5	120	34	207	1.3 ms
A_6	360	118	6249	23 ms
S_6	720	763	27590	120 ms
A_7	2,520	—	74,946,283	52 min
S_7	5,040	—	1,059,510,737	197 min
$\text{PSL}(2, 17)^{\parallel}$	2,448	—	123,451,769	23 min
$\text{PSL}(2, 19)^{\parallel}$	3,420	—	2,696,472,513	21 h
$\text{PSL}(2, 16)^{\parallel}$	4,080	—	4,869,673,337	13 h
$C_{109} \times C_9$	981	3	5,683,264,056	14 h
$(C_7 \times C_3) \times (C_{13} \times C_3)$	819	6	8,728,959,134	19 h
$C_{89} \times C_{11}$	979	1	10,178,934,027	24 h
$C_{61} \times C_{15}$	915	1	43,174,839,011	87 h
$C_{43} \times C_{21}$	903	1	43,967,855,355	88 h
$(C_9 \times C_9) \times C_9$	729	75	97,944,803	52 min
$(C_{27} \times C_9) \times C_3$	729	390	175,898,535	87 min
$((C_9 \times C_3) \times C_3) \times C_3^2$	729	399	240,194,985	128 min
$C_{27} \times C_{27}^{\ddagger}$	729	22	10,776,997	255 s
$(C_3 \times (C_{27} \times C_3)) \times C_3^{\ddagger}$	729	90	11,276,468	208 s
C_9^2 by C_3^2	729	96	13,525	39 ms
$C_{31} \times C_{30}$	930	1	821,601	18 s
$C_2^5 \times C_{31}$	992	194	3,260,710	70 s
$C_{17} \times A_5$	1,020	9	1,030,890	18 s

TABLE 4. Experiments for selected groups including the five groups with the longest running time. The first column shows the group according to the GAP structure description, the second column displays the order of the group, the third column the index in the SmallGrp library, the fourth column the number of calls to the TESTGEODETIC procedure, and the final column contains the running time. The computation was parallelized for the groups $\text{PSL}(2, q)$ (marked with \parallel) according to Remark 6.4; the quantities displayed in the last two columns are cumulative. The center of the groups of order 729 is either of order three or nine (marked with \ddagger).

- [4] Peter J. Cameron. *Permutation Groups*. London Mathematical Society Student Texts. Cambridge University Press, 1999. doi:10.1017/CB09780511623677.
- [5] Andrey Che. Uniqueness of geodesics in Cayley graphs of finite groups. Technical report, 2022.
- [6] Anthony E. Clement, Stephen Majewicz, and Marcos Zyman. *The theory of nilpotent groups*. Birkhäuser/Springer, Cham, 2017.
- [7] R. M. Damerell. On Moore graphs. *Math. Proc. Cambridge Philos. Soc.*, 74(2):227–236, 1973. doi:10.1017/S0305004100048015.
- [8] A. K. Das, R. K. Nath, and M. R. Pournaki. A Survey on the Estimation of Commutativity in Finite Groups. *Southeast Asian Bull. Math.*, 37(2):161–180, 2013.
- [9] Michael J. Dinneen and Paul R. Hafner. New results for the degree/diameter problem. *Networks*, 24(7):359–367, 1994. doi:10.1002/net.3230240702.
- [10] Andy Eisenberg and Adam Piggott. Gilman’s conjecture. *J. Algebra*, 517:167–185, 2019. doi:10.1016/j.jalgebra.2018.09.022.
- [11] Murray Elder, Giles Gardam, Adam Piggott, Davide Spriano, and Kane Townsend. Graphs and groups with unique geodesics. *arXiv preprint*, 2023. arXiv:2311.03730.
- [12] Murray Elder and Adam Piggott. Rewriting systems, plain groups, and geodetic graphs. *Theoret. Comput. Sci.*, 903:134–144, 2022.
- [13] Murray Elder, Adam Piggott, and Kane Townsend. On k -geodetic graphs and groups. *Internat. J. Algebra Comput.*, 2023. doi:10.1142/S0218196723500534.

- [14] Asaf Etgar and Nati Linial. On the connectivity and diameter of geodetic graphs. *European J. Combin.*, 116:Paper No. 103886, 10, 2024. doi:10.1016/j.ejc.2023.103886.
- [15] Bruno Federici. *Interactions between large-scale invariants in infinite graphs*. PhD thesis, University of Warwick, 2017. URL: <https://wrap.warwick.ac.uk/108882/>.
- [16] Walter Feit and John G. Thompson. Solvability of groups of odd order. *Pacific J. Math.*, 13:775–1029, 1963. URL: <http://projecteuclid.org/euclid.pjm/1103053943>.
- [17] Carlos Frasser and George Vostrov. Geodetic Graphs Homeomorphic to a Given Geodetic Graph. *Int. J. Graph Theory Appl.*, 3:13–44, 12 2020.
- [18] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.12.2*, 2022. URL: <https://www.gap-system.org>.
- [19] Dmitriy Gorovoy and David Zmiaikou. On graphs with unique geodesics and antipodes. *Discrete Math.*, 347(4):Paper No. 113864, 17, 2024. doi:10.1016/j.disc.2023.113864.
- [20] A. J. Hoffman and R. R. Singleton. On Moore Graphs with Diameters 2 and 3. *IBM J. Res. Dev.*, 4(5):497–504, 1960. doi:10.1147/rd.45.0497.
- [21] Hans Liebeck and Desmond MacHale. Groups with Automorphisms Inverting most Elements. *Math. Z.*, 124(1):51–63, Mar 1972. doi:10.1007/BF01142582.
- [22] Klaus Madlener and Friedrich Otto. Groups presented by certain classes of finite length-reducing string-rewriting systems. In *Rewriting techniques and applications (Bordeaux, 1987)*, volume 256 of *Lecture Notes in Comput. Sci.*, pages 133–144. Springer, Berlin, 1987. doi:10.5555/30432.30444.
- [23] Mirka Miller and Jozef Širáň. Moore graphs and beyond: a survey of the degree/diameter problem. *Electron. J. Combin.*, DS14:61, 2005.
- [24] Oystein Ore. Theory of Graphs. In *Colloquium Publications*. American Mathematical Society, 1962.
- [25] Igor Pak and Radoš Radoičić. Hamiltonian paths in Cayley graphs. *Discrete Math.*, 309(17):5501–5508, 2009. doi:<https://doi.org/10.1016/j.disc.2009.02.018>.
- [26] Adam Piggott. On groups presented by monadic rewriting systems with generators of finite order. *Bull. Aust. Math. Soc.*, 91(3):426–434, 2015. doi:10.1017/S0004972715000015.
- [27] G. Pólya and G. Szegő. *Aufgaben und Lehrsätze aus der Analysis: Erster Band: Reihen · Integralrechnung Funktionentheorie*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 1925. doi:10.1007/978-3-662-38381-0.
- [28] Primož Potočnik, Pablo Spiga, and Gabriel Verret. Cubic vertex-transitive graphs on up to 1280 vertices. *J. Symbolic Comput.*, 50:465–477, 2013. doi:10.1016/j.jsc.2012.09.002.
- [29] Derek J. S. Robinson. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996. doi:10.1007/978-1-4419-8594-1.
- [30] David J. Rusin. What is the probability that two elements of a finite group commute? *Pacific J. Math.*, 82(1):237–247, 1979.
- [31] Michael Shapiro. Pascal’s triangles in Abelian and hyperbolic groups. *J. Aust. Math. Soc.*, 63(2):281–288, 1997.
- [32] Joel G. Stemple. Geodetic graphs of diameter two. *J. Combin. Theory Ser. B*, 17(3):266–280, 1974.
- [33] Joel G. Stemple and Mark E. Watkins. On planar geodetic graphs. *J. Combin. Theory*, 4(2):101–117, 1968.
- [34] Florian Stober and Armin Weiß. Geodetic graphs: experiments and new constructions. *arXiv preprint*, 2023. arXiv: 2308.08970.

SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES, UNIVERSITY OF TECHNOLOGY SYDNEY, BROADWAY NSW 2007, AUSTRALIA

Email address: murray.elder@uts.edu.au

MATHEMATICAL SCIENCES INSTITUTE, AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA ACT 2601, AUSTRALIA

Email address: adam.piggott@anu.edu.au

INSTITUT FÜR FORMALE METHODEN DER INFORMATIK, UNIVERSITÄT STUTTGART, UNIVERSITÄTSSTRASSE 38, 70569 STUTTGART, GERMANY

Email address: florian.stober@fmi.uni-stuttgart.de

INSTITUT FÜR FORMALE METHODEN DER INFORMATIK, UNIVERSITÄT STUTTGART, UNIVERSITÄTSSTRASSE 38, 70569 STUTTGART, GERMANY

Email address: alexander.thumm@fmi.uni-stuttgart.de

INSTITUT FÜR FORMALE METHODEN DER INFORMATIK, UNIVERSITÄT STUTTGART, UNIVERSITÄTSSTRASSE 38, 70569 STUTTGART, GERMANY

Email address: armin.weiss@fmi.uni-stuttgart.de