

A QUEST FOR CONVERGENCE: EXPLORING SERIES IN NON-LINEAR ENVIRONMENTS

GEIVISON RIBEIRO

ABSTRACT. This note presents an extension of a result within the concept of $[\mathcal{S}]$ -lineability, originally developed in 2019 by L. Bernal-González, J.A. Conejero, M. Murillo-Arcila, and J.B. Seoane-Sepúlveda (see [4]). Additionally, we provide a characterization in terms of lineability in the context of complements of unions of closed subspaces in F -spaces, and finally, we present a negative result in both normed spaces and p -Banach spaces. These findings contribute to the understanding of linearity in exotic settings in topological vector spaces.

1. INTRODUCTION AND BACKGROUND

In recent years, the exploration of linearity within unconventional mathematical contexts has emerged as a notable trend. This pursuit has been underscored by the introduction of terms like *lineability* and *spaceability*, originally coined by V.I. Gurariy, which gained prominence following their appearance in the seminal paper by Aron, Gurariy, and Seoane-Sepúlveda; we refer the reader to [2], (see also [11]). The inception of these concepts sparked interest among many mathematicians, catalyzing an increase in research efforts in this direction. According to these terminologies, a subset A of a vector space X is said to be lineable if $A \cup \{0\}$ encompasses an infinite dimensional vector subspace. Furthermore, in the context of X being a topological vector space, A is called spaceable if $A \cup \{0\}$ contains a closed infinite dimensional vector subspace. A detailed account of lineability can be found in [1, 3, 6, 10].

In 2004, during an Analysis Seminar at Kent State University, Gurariy introduced the following concept of $[\mathcal{S}]$ -lineability (see [4]): Given a Hausdorff topological vector space $X \neq \{0\}$ and a vector subspace \mathcal{S} of $\mathbb{K}^{\mathbb{N}}$ (where $\mathbb{K} = \mathbb{R}$ or \mathbb{C}), a subset A in X is defined as:

- $[(u_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable in X if, for each sequence $(c_n)_{n=1}^{\infty} \in \mathcal{S}$, the series $\sum_{n=1}^{\infty} c_n u_n$ converges in X to a vector of $A \cup \{0\}$.
- $[\mathcal{S}]$ -lineable in X if it is $[(u_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable for some sequence $(u_n)_{n=1}^{\infty}$ of linearly independent elements in X .

As far as we know, the notion of $[\mathcal{S}]$ -lineability has been developed only in [4].

To avoid trivial or undesirable situations, we shall assume throughout this paper that a vector space X never collapses to $\{0\}$. Throughout this note, we will use the term subspace instead of vector subspace.

The symbol $\text{card}(M)$ will denote the cardinality of the set M , and in particular, we will denote \aleph_0 as the cardinality of \mathbb{N} .

This note is organized as follows. In Section 2, we consider that X is an F -space and we establish that a set of the form $X \setminus \bigcup_{i \in I} Y_i$ is lineable if and only if it is $[\mathcal{S}]$ -lineable for every nontrivial subspace \mathcal{S} of ℓ_{∞} . Recall that an F -space is a complete metrizable topological vector space, and that every

2020 *Mathematics Subject Classification.* 15A03, 46B87, 46A16.

Key words and phrases. Lineability, spaceability, convergence, series, basic sequences, topological vector space, quotient space.

G. Ribeiro is supported by Grant 2022/1962, Paraíba State Research Foundation (FAPESQ).

F -space X supports an F -norm $\|\cdot\|_X$. Recall also that an F -norm on a vector space X is a function $\|\cdot\| : X \rightarrow [0, \infty)$ satisfying, for all $x, y \in X$ and $\lambda \in \mathbb{K}$ the following properties:

- (1) $\|x\|_X = 0$ if and only if $x = 0$.
- (2) $\|\lambda x\|_X \leq \|x\|_X$ if $|\lambda| \leq 1$.
- (3) $\|\lambda x\|_X \rightarrow 0$ if $|\lambda| \rightarrow 0$.
- (4) $\|x + y\|_X \leq \|x\|_X + \|y\|_X$.

Moving to Section 3, we extend a result in the realm of $[\mathcal{S}]$ -lineability, originally attributed to L. Bernal-González, J.A. Conejero, M. Murillo-Arcila, and J.B. Seoane-Sepúlveda. Lastly, in Section 4, we present negative results.

2. $[\mathcal{S}]$ -LINEABILITY IN THE CONTEXT OF COMPLEMENTS OF UNIONS

In this section, we aim to characterize the family of complements of unions of closed subspaces through the notion of $[\mathcal{S}]$ -lineability. To this end, let us begin with the following notion as appeared in [5].

Definition 2.1. *We say that a sequence $(u_n)_{n=1}^\infty$ of elements of a topological vector space X is topologically linearly independent, if for each sequence $(c_n)_{n=1}^\infty \in \mathbb{K}^\mathbb{N}$ with $\sum_{n=1}^\infty c_n u_n = 0$, we have $(c_n)_{n=1}^\infty = 0$.*

Based on this definition, if $\mathcal{S} \neq \{0\}$ is a subspace of $\mathbb{K}^\mathbb{N}$ then we will say that a sequence $(u_n)_{n=1}^\infty$ of elements of a topological vector space X is \mathcal{S} -topologically linearly independent in X (or \mathcal{S} -independent) if for each sequence $(c_n)_{n=1}^\infty \in \mathcal{S}$ with $\sum_{n=1}^\infty c_n u_n = 0$, we have $(c_n)_{n=1}^\infty = 0$.

Within this perspective, still in [5], Drewnowski et al. demonstrated the following result, which will be a crucial ingredient for the proof of the theorem 2.4. It is worth mentioning that this result establishes a connection between linear ℓ_∞ -independence and linear independence.

Proposition 2.2. *Assume that $(x_n)_{n=1}^\infty$ is a linearly independent sequence in a Hausdorff topological vector space X . Then there is $\lambda_n > 0$ such that $(\lambda_n x_n)_{n=1}^\infty$ is ℓ_∞ -independent.*

Em [8], the authors made a notable contribution with the following result:

Theorem 2.3. [8, Theorem 7.4.2] *Let Z_r ($r \in \mathbb{N}$) be Banach spaces and let X be a Fréchet space. Let $T_r : Z_r \rightarrow X$ be continuous linear mappings and $Y = \text{span}(\bigcup_{r=1}^\infty T_r(Z_r))$. If Y is not closed in X then the complement $X \setminus Y$ is spaceable.*

At this point, a characterization of spaceability for the complements of non-enumerable unions of subspaces in the context of F -spaces in terms of lineability has yet to be developed. It is not even known if these complements are $[\mathcal{S}]$ -lineable, and it is precisely for this purpose that we present the following result:

Theorem 2.4. *Let $X \neq \{0\}$ be an F -space and $\{Y_i\}_{i \in I}$ be a family of nontrivial closed subspaces of X . The set $X \setminus \bigcup_{i \in I} Y_i$ is lineable if and only if it is $[\mathcal{S}]$ -lineable for every subspace $\mathcal{S} \neq \{0\}$ of ℓ_∞ .*

Proof. Assume that $X \setminus \bigcup_{i \in I} Y_i$ is lineable and consider the quotient map $Q_i : X \rightarrow X/Y_i$ ($i \in I$). Let E be an infinite dimensional subspace of X such that

$$E \cap \left(\bigcup_{i \in I} Y_i \right) = \{0\}.$$

Let $(x_n)_{n=1}^\infty$ be a sequence of elements linearly independent in E such that

$$(2.1) \quad \sum_{n=1}^{\infty} \|x_n\|_X < \infty.$$

Fix $i \in I$. Due to the fact that $X \setminus \bigcup_{i \in I} Y_i$ is lineable, we can infer that the sequence $(Q_i(x_n))_{n=1}^\infty$ is also linearly independent in X/Y_i . Indeed, if for some $N \in \mathbb{N}$, we have

$$\sum_{n=1}^N a_n Q_i(x_n) = 0,$$

then

$$Q_i \left(\sum_{n=1}^N a_n x_n \right) = 0.$$

Hence

$$\sum_{n=1}^N a_n x_n \in Y_i \cap E = \{0\}.$$

That is,

$$a_n = 0,$$

which proves the desired linear independence in X/Y_i . Since X/Y_i is a Hausdorff topological vector space, we can invoke Proposition 2.2 to obtain a sequence of positive real numbers $(\lambda_n)_{n=1}^\infty$ such that the sequence $(Q_i(\lambda_n x_n))_{n=1}^\infty$ is ℓ_∞ -independent in X/Y_i . We also claim that the sequence $(\alpha_n)_{n=1}^\infty$, where $\alpha_n := \frac{\lambda_n}{1+\lambda_n}$ is such that $(Q_i(\alpha_n x_n))_{n=1}^\infty$ is ℓ_∞ -independent in X/Y_i . Indeed, let $(t_n)_{n=1}^\infty \in \ell_\infty$ be such that

$$\sum_{n=1}^\infty t_n Q_i(\alpha_n x_n) = 0.$$

Hence,

$$(2.2) \quad \sum_{n=1}^\infty \frac{t_n}{1+\lambda_n} Q(\lambda_n x_n) = 0.$$

Since

$$\left(\frac{t_n}{1+\lambda_n} \right)_{n=1}^\infty \in \ell_\infty,$$

and $(Q_i(\lambda_n x_n))_{n=1}^\infty$ is ℓ_∞ -independent in X/Y_i , we get

$$\frac{t_n}{1+\lambda_n} = 0 \text{ for each } n \in \mathbb{N},$$

and this entails that

$$t_n = 0 \text{ for each } n \in \mathbb{N}.$$

Therefore, $(Q_i(\alpha_n x_n))_{n=1}^\infty$ is really ℓ_∞ -independent in X/Y_i . Furthermore, since $|\alpha_n| \leq 1$ for all $n \in \mathbb{N}$, we have

$$\|\alpha_n x_n\|_X \leq \|x_n\|_X \text{ for all } n \in \mathbb{N}.$$

Since

$$\sum_{n=1}^\infty \|x_n\|_X < \infty,$$

we obtain

$$\sum_{n=1}^\infty \|\alpha_n x_n\|_X \leq \sum_{n=1}^\infty \|x_n\|_X < \infty.$$

Now, let $c = (c_n)_{n=1}^\infty \in \ell_\infty \setminus \{0\}$. We claim that the series $\sum_{n=1}^\infty c_n \alpha_n x_n$ converges in X . Indeed, due to the fact that

$$\left\| \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X \leq \|\alpha_n x_n\|_X < \infty,$$

if we take $\varepsilon > 0$, then there exists $n_0 \in \mathbb{N}$ such that

$$\sum_{n=n_0+1}^\infty \left\| \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X < \varepsilon.$$

Thus, for $r, s \in \mathbb{N}$ with $r, s \geq n_0$, we have

$$\begin{aligned} \left\| \sum_{n=1}^r \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n - \sum_{n=1}^s \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X &= \left\| \sum_{n=s+1}^r \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X \\ &\leq \sum_{n=s+1}^r \left\| \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X \\ &\leq \sum_{n=n_0+1}^\infty \left\| \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right\|_X \\ &< \varepsilon. \end{aligned}$$

Since X is complete and Hausdorff, we can conclude that sequence $\left(\sum_{n=1}^r \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \right)_{r=1}^\infty$ converges in X for the series

$$\sum_{n=1}^\infty \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n.$$

Hence,

$$\sum_{n=1}^\infty \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \in X,$$

and consequently,

$$\sum_{n=1}^\infty c_n \alpha_n x_n = \|c\|_{\ell_\infty} \sum_{n=1}^\infty \|c\|_{\ell_\infty}^{-1} c_n \alpha_n x_n \in X.$$

Our aim from this point forward will be to demonstrate that the series $\sum_{n=1}^\infty c_n \alpha_n x_n$ converges to a vector in $(X \setminus \bigcup_{i \in I} Y_i) \cup \{0\}$. To achieve this, we will assume, by way of contradiction, the existence of $y \in (\bigcup_{i \in I} Y_i) \setminus \{0\}$ such that

$$(2.3) \quad \sum_{n=1}^\infty c_n \alpha_n x_n = y.$$

Since $y \in Y_i$ for some $i \in I$, we have

$$Q_i \left(\sum_{n=1}^\infty c_n \alpha_n x_n \right) = Q_i(y) = 0.$$

That is,

$$\sum_{n=1}^\infty c_n Q_i(\alpha_n x_n) = 0.$$

Since $(Q_i(\alpha_n x_n))_{n=1}^\infty$ is ℓ_∞ -independent in X/Y_i and $(c_n)_{n=1}^\infty \in \ell_\infty$, we get

$$(c_n)_{n=1}^\infty = 0.$$

From (2.3), we obtain

$$y = 0,$$

which leads to a contradiction. Therefore, the series $\sum_{n=1}^{\infty} c_n \alpha_n x_n$ converges to a vector in $(X \setminus \bigcup_{i \in I} Y_i) \cup \{0\}$. This shows that $X \setminus \bigcup_{i \in I} Y_i$ is $[\ell_{\infty}]$ -lineable and by [4], we conclude that $X \setminus \bigcup_{i \in I} Y_i$ is $[\mathcal{S}]$ -lineable for each subspace \mathcal{S} of ℓ_{∞} . For the proof of the converse, it suffices to take $\mathcal{S} = c_{00}$ and utilize the fact that $X \setminus Y$ is $[c_{00}]$ -lineable if and only if it is lineable. \square

3. A KIND OF EXTENSION

A result due Bernal et al. in [4] states that if A is a subset $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable of a metrizable vector space X , where \mathcal{S} is an infinite dimensional subspace of $\mathbb{K}^{\mathbb{N}}$ and $(x_n)_{n=1}^{\infty}$ is a basic sequence, (recall that a sequence $(x_n)_{n=1}^{\infty}$ of a metrizable topological vector space X is said to be basic whenever every $x \in \text{span}\{x_n : n \in \mathbb{N}\}$, the closed linear span of the x_n 's, can be uniquely represented as a convergent series $x = \sum_{n=1}^{\infty} c_n x_n$), then A is lineable. More precisely, they showed the following:

Proposition 3.1. *Assume that $(x_n)_{n=1}^{\infty}$ is a basic sequence in X , where X is metrizable, and that A is a $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable subset of X , where \mathcal{S} is an infinite dimensional subspace of $\mathbb{K}^{\mathbb{N}}$. Then A is lineable.*

In this section, we complement this result by removing the assumption of metrizability and the requirement for the sequence to be basic.

Theorem 3.2. *Let $X \neq \{0\}$ be a Hausdorff topological vector space and A be a nonvoid subset of X . If $(x_n)_{n=1}^{\infty}$ is an ℓ_{∞} -independent sequence of elements of X and A is $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable, for some infinite dimensional subspace \mathcal{S} of ℓ_{∞} , then A is lineable.*

Proof. Let \mathcal{S} be an infinite dimensional subspace of ℓ_{∞} and assume that A is $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable. Since $(x_n)_{n=1}^{\infty}$ is ℓ_{∞} -independent in X , we can conclude that the operator $T: \mathcal{S} \rightarrow X$ given by $T((c_n)_{n=1}^{\infty}) = \sum_{n=1}^{\infty} c_n x_n$ is not only well-defined and linear, but also injective. Thus,

$$\dim T(\mathcal{S}) = \dim \mathcal{S} \geq \aleph_0.$$

Furthermore, due to the fact that X is Hausdorff, we have

$$T(\mathcal{S}) \subset A \cup \{0\}.$$

Hence, A is lineable and the proof is complete. \square

4. NEGATIVE RESULTS

In this section, we consider results that provide an enlightening insight into the $[\mathcal{S}]$ -lineability of subsets in some class of some classes of infinite dimensional vector spaces. The next proposition highlights the importance of sequence properties concerning their distance from the origin in normed vector spaces. Roughly speaking, the next statement shows that sequences away from zero do not generate $[\mathcal{S}]$ -lineability.

Proposition 4.1. *Let X be an infinite dimensional normed space. If $(x_n)_{n=1}^{\infty}$ is a linearly independent sequence in X such that $\inf_{n \in \mathbb{N}} \|x_n\|_X > 0$, then for any infinite dimensional subspace \mathcal{S} of ℓ_{∞} properly containing c_0 , there is no subset of X that is $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable.*

Proof. Assume that A is a subset of X which is $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable in X for some infinite dimensional subspace \mathcal{S} of ℓ_{∞} properly containing c_0 . Given $(c_n)_{n=1}^{\infty} \in \mathcal{S}$, we have

$$\|c_n x_n\|_X = |c_n| \|x_n\|_X \geq |c_n| \inf_{n \in \mathbb{N}} \|x_n\|_X \text{ for all } n \in \mathbb{N}.$$

That is,

$$(4.1) \quad \left(\inf_{n \in \mathbb{N}} \|x_n\|_X \right)^{-1} \cdot \|c_n x_n\|_X \geq |c_n| \text{ for all } n \in \mathbb{N}.$$

Due to (4.1), if we take $(c_n)_{n=1}^\infty \in \mathcal{S} \setminus c_0$, we can infer that the sequence $(\|c_n x_n\|_X)_{n=1}^\infty$ does not converge to zero. However, this leads to a contradiction, since the series $\sum_{n=1}^\infty c_n x_n$ converges in X for some vector of $A \cup \{0\}$. Therefore, the desired result follows. \square

The result above remains valid in the context of p -Banach spaces as well. Recall that if $p \in (0, 1]$, then a p -Banach space X is a vector space equipped with a non-negative function denoted by $\|\cdot\|$, also known as a p -norm, which satisfies the following conditions:

- (1) $\|x\|_p = 0$ if and only if $x = 0$.
- (2) $\|tx\|_p = |t|^p \|x\|_p$ for all $x \in X$ and $t \in \mathbb{R}$.
- (3) $\|x + y\|_p \leq \|x\|_p + \|y\|_p$ for all $x, y \in X$.

The space ℓ_p (with $p \in (0, 1)$) of all real sequences $(x_n)_{n=1}^\infty$ such that the series $\sum_{n=1}^\infty |x_n|^p$ converges is an example of p -Banach space. If we consider the sequence $(e_n)_{n=1}^\infty$ in ℓ_∞ , where $e_n = (0, 0, \dots, 0, 1, 0, 0, \dots)$ (with the 1 at the n th place), we obtain the following result:

Corollary 4.2. *For $p \in (0, 1)$ and any infinite dimensional subspace \mathcal{S} of ℓ_∞ properly containing c_0 , there is no subset of ℓ_p that is $[(e_n)_{n=1}^\infty, \mathcal{S}]$ -lineable.*

REFERENCES

- [1] R.M. Aron, L. Bernal-González, D. Pellegrino and J.B. Seoane-Sepúlveda, *Lineability: the search for linearity in mathematics*, Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2016. xix+308 pp.
- [2] R.M. Aron, V.I. Gurariy and J.B. Seoane-Sepúlveda, *Lineability and spaceability of sets of functions on \mathbb{R}* , Proc. Amer. Math. Soc. **133** (2004), 795–803.
- [3] L. Bernal-González and M.O. Cabrera, *Lineability criteria, with applications*, J. Funct. Anal. **266** (2014), 3997–4025.
- [4] L. Bernal-González, J. A. Conejero, M. Murillo-Arcila, J.B. Seoane-Sepúlveda, *[S]-linear and convex structures in function families*, Linear Algebra Appl. **579** (2019), 463–483.
- [5] L. Drewnowski, I. Labuda and Z. Lipecki, *Existence of quasi-bases for separable topological linear spaces*, Arch. Math. (Basel) **37** (1981), no. 5, 454–456.
- [6] P.H. Enflo, V.I. Gurariy, and J.B. Seoane-Sepúlveda, *Some Results and Open Questions on Spaceability in Function Spaces*, Trans. Amer. Math. Soc., **366** (2014), no. 2, 611–625.
- [7] V.I. Gurariy, *Subspaces and bases in spaces of continuous functions*, Dokl. Akad. Nauk SSSR **167** (1966), 971–973 (Russian).
- [8] D. Kitson and R.M. Timoney, *Operator ranges and spaceability*. J. Math. Anal. Appl. **378** (2011), 680–686.
- [9] B. Levine and D. Milman, *On linear sets in space C consisting of functions of bounded variation*, Comm. Inst. Sci. Math. Méc. Univ. Kharkoff [Zapiski Inst. Mat. Mech.] (4) **16** (1940), 102–105 (Russian, with English summary).
- [10] D.M. Pellegrino and E. V. Teixeira, *Norm optimization problem for linear operators in classical Banach spaces*, Bull. Braz. Math. Soc. (N.S.) **40** (2009), no. 3, 417–431.
- [11] J.B. Seoane-Sepúlveda, *Chaos and lineability of pathological phenomena in analysis*, Thesis (Ph.D.), Kent State University, ProQuest LLC, Ann Arbor, MI (2006), 139 pages.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.
 Email address: geivison.ribeiro@academico.ufpb.br