Removable edges in near-bipartite bricks

Yipei Zhang¹, Fuliang Lu², Xiumei Wang¹, Jinjiang Yuan¹

 1 School of Mathematics and Statistics, Zhengzhou University, Zhengzhou 450001, China

² School of Mathematics and Statistics, Minnan Normal University, Zhangzhou 363000, China

Abstract

An edge e of a matching covered graph G is removable if G - e is also matching covered. The notion of removable edge arises in connection with ear decompositions of matching covered graphs introduced by Lovász and Plummer. A nonbipartite matching covered graph G is a brick if it is free of nontrivial tight cuts. Carvalho, Lucchesi, and Murty proved that every brick other than K_4 and $\overline{C_6}$ has at least $\Delta - 2$ removable edges. A brick G is near-bipartite if it has a pair of edges $\{e_1, e_2\}$ such that $G - \{e_1, e_2\}$ is a bipartite matching covered graph. In this paper, we show that in a near-bipartite brick G with at least six vertices, every vertex of G, except at most six vertices of degree three contained in two disjoint triangles, is incident with at most two nonremovable edges; consequently, G has at least $\frac{|V(G)|-6}{2}$ removable edges. Moreover, all graphs attaining this lower bound are characterized.

Keywords near-bipartite graph; brick; removable edge; perfect matching

1 Introduction

Graphs considered in this paper may have multiple edges, but no loops. We follow [1] for undefined notation and terminology. A connected graph G is k-extendable if each set of kindependent edges extends to a perfect matching of G. An edge e of a graph G is admissible if G has a perfect matching that contains e, and nonadmissible otherwise. A connected nontrivial graph is matching covered if each of its edges is admissible. Matching covered graphs are also called 1-extendable [11]. For $X \subseteq V(G)$, by $\partial(X)$ we mean the edge cut of G, which is the set of edges of G with one end in x and the other in \overline{X} , where $\overline{X} = V(G) \setminus X$; by $G/X \to x$ or simply G/X we mean the graph obtained by contracting X to a single vertex x, the graph $G/\overline{X} \to \overline{x}$ or simply G/\overline{X} is defined analogously.

^{*}Corresponding author.

Email addresses: zyipei@163.com, flianglu@163.com, wangxiumei@zzu.edu.cn, yuanjj@zzu.edu.cn.

Let G be a matching covered graph. An edge cut $C = \partial(X)$ of G is tight if $|M \cap C| = 1$ for each perfect matching M of G and is separating if G/X and G/\overline{X} are matching covered. A matching covered graph that is free of nontrivial tight cuts is a *brace* if it is bipartite, and a brick if it is nonbipartite. Edmonds et al. [5] showed that a graph G is a brick if and only if G is 3-connected and for any two distinct vertices x and y of G, G - x - y has a perfect matching. A brick is *solid* if it is free of nontrivial separating cuts. We denote the number of vertices of G by n. An edge e of G is removable if G - e is also matching covered. A pair of edges $\{e_1, e_2\}$ is a removable doubleton of G if neither e_1 nor e_2 is removable in G but $G - \{e_1, e_2\}$ is matching covered. The notion of removable edge arises in connection with ear decompositions of matching covered graphs introduced by Lovász and Plummer. The existence of removable edges, especially of special types, plays an important role in the generation of some bricks, see [9, 10]. Lovász [12] first showed the existence of removable edges of bricks other than K_4 and $\overline{C_6}$. Carvalho, Lucchesi and Murty [2, 4] proved that every brick other than K_4 and $\overline{C_6}$ has at least $\Delta - 2$ removable edges; in every solid brick G with six or more vertices, each vertex is incident with at most two nonremovable edges, consequently, G has at least $\frac{n}{2}$ removable edges. Zhai, Lucchesi and Guo [17] showed that every matching covered graph G has at least m(G) removable classes (including removable edges and removable doubletons), where m(G) denotes the minimum number of perfect matchings needed to cover all edges of G. For bipartite matching covered graphs, He et al. [7] gave a characterization of graphs each of whose edges is removable.



Figure 1: The four near-bipartite tri-ladders

A nonbipartite matching covered graph G is near-bipartite if it has a pair of edges $\{e_1, e_2\}$ such that $G - \{e_1, e_2\}$ is a bipartite matching covered graph; such a pair $\{e_1, e_2\}$ is referred to as a removable doubleton of G. The graphs K_4 and $\overline{C_6}$ are the only simple near-bipartite bricks on four and six vertices, respectively, each of which has three removable doubletons but no removable edges. The significance of near-bipartite graphs arises from the theory of ear decompositions. Fischer and Little characterized Pfaffian near-bipartite graphs in [6]. Kothari [9], and Kothari and Carvalho [10] investigated generation procedures for near-bipartite bricks and simple near-bipartite bricks, respectively. Inspiring by the structure with respect to nonremovable edges in solid bricks, we consider near-bipartite bricks in this paper. The main results are stated as follows.

Theorem 1.1. Let G be a near-bipartite brick other than K_4 . Then every vertex of G, except at most six vertices of degree three contained in two disjoint triangles, is incident with at most two nonremovable edges.

Theorem 1.2. Every near-bipartite brick G other than K_4 has at least $\frac{n-6}{2}$ removable edges, and near-bipartite tri-ladders (see Section 4) are the only graphs attaining the lower bound.

This paper is organized as follows. In Section 2, we present some basic results. In Section 3 and 4, we give proofs of Theorems 1.1 and 1.2, respectively.

2 Preliminaries

We begin with some notations. Let G be a graph with the vertex set V(G) and the edge set E(G). By V(e) we mean the set of the two ends of the edge e. For $X \subseteq V(G)$, by $N_G(X)$, or simply N(X), we mean the set of vertices that are not in X but have neighbours in X; by G[X] we mean the subgraph of G induced by X. For $X, Y \subseteq V(G)$, by $E_G[X, Y]$, or simply E[X, Y], we mean the set of edges of G with one end in X and the other end in Y. If Y = X, we use E(X) instead of E[X, X]. Then the edge cut $\partial(X) = E[X, \overline{X}]$. An edge cut $\partial(X)$ is a k-cut if $|\partial(X)| = k$; is trivial if either |X| = 1 or $|\overline{X}| = 1$, and nontrivial otherwise.

2.1 Basic results

Tutte [16] proved that a graph G has a perfect matching if and only if $o(G-S) \leq |S|$ for every $S \subseteq V(G)$, where o(G-S) denotes the number of odd components of G-S. A nonempty subset S of V(G) is a *barrier* of G that has a perfect matching if o(G-S) = |S|. A graph G is *factor-critical* if, for each vertex v of G, G-v has a perfect matching. Using Tutte's theorem, Lemma 2.1 may be easily derived (see [4]).

Lemma 2.1. Let G be a graph with a perfect matching. Then

- (i) if G is a brick, then every barrier of G is trivial,
- (ii) an edge e of G is admissible if and only if G has no barriers containing V(e), and
- (iii) for each maximal barrier B of G, all components of G B are factor-critical.

Lemma 2.2 ([4]). Let G be a brick, and let f_1 and f_2 be two adjacent edges of G. If, for $i = 1, 2, S_i$ is a barrier of $G - f_i$, then $|S_1 \cap S_2| \le 1$.

Lemma 2.3. Let G be a graph with a perfect matching, let S and S' be two subsets of V(G) such that $N(S) \subseteq S'$, $S \cap S' = \emptyset$ and $|S'| \leq |S| + 1$. If S is an independent set of G, then S' is a barrier of G.

Proof. Since $N(S) \subseteq S'$, $S \cap S' = \emptyset$ and S is an independent set, we have $o(G - S') \ge |S|$. Furthermore, since G has a perfect matching, by Tutte's theorem and the assumption that $|S'| \le |S| + 1$, we have

$$|S'| - 1 \le |S| \le o(G - S') \le |S'|,$$

Thus o(G - S') = |S'| or |S'| - 1. Since |V(G)| is even, o(G - S') and |S'| have the same parity. Thus o(G - S') = |S'| and then S' is a barrier of G.

Note that it is possible that S' = N(S) in Lemma 2.3, i.e., if S is an independent set of a graph G with a perfect matching and $|N(S)| \leq |S| + 1$, then N(S) is a barrier of G.

Lemma 2.4 ([11]). Let H[U,W] be a bipartite graph with four or more vertices, where |U| = |W|. Then H is matching covered if and only if $|N(S)| \ge |S| + 1$ for every nonempty proper subset S of U.

Lemma 2.5 ([13]). Let H[U, W] be a bipartite matching covered graph, and u a vertex of G of degree d, where $u \in U$ and $d \geq 3$. If uw_1, uw_2, \ldots, uw_r , $0 < r \leq d$, are nonremovable edges of H incident with u, then there exist partitions (U_0, U_1, \ldots, U_r) of U and (W_0, W_1, \ldots, W_r) of W such that $u \in U_0$ and, for $i \in \{1, 2, \cdots, r\}$, (a) $|U_i| = |W_i|$, (b) $w_i \in W_i$ and (c) $N(W_i) = U_i \cup \{u\}$; in particular, uw_i is the only edge in $E[U_0, W_i]$.

From Lemma 2.5, we can obtain the following lemma.

Lemma 2.6. Let H be a bipartite matching covered graph, and u a vertex of H with degree three or more. If f_1 and f_2 are two edges incident with u that lie in a 4-cycle, then at least one of f_1 and f_2 is removable.

2.2 Two types of nonremovable edges

For the rest of Section 2 and Section 3, we assume that

- (i) G is a near-bipartite brick with a removable doubleton $\{e_1, e_2\}$, and
- (ii) (U, W) is the bipartition of H such that e_1 connects two vertices of U and e_2 connects two vertices of W, where $H = G \{e_1, e_2\}$.

Note that both U and W are stable set of H, e_1 is the only edge in $E_G(U)$ and e_2 is the only edge in $E_G(W)$. Unless otherwise specified, we use N(X), E(X), and E[X, Y] for $N_G(X)$, $E_G(X)$, and $E_G[X, Y]$, respectively, where $X, Y \subseteq V(G)$. Then $E(U) = \{e_1\}$ and $E(W) = \{e_2\}$.

Lemma 2.7. Let S be a subset of U (or W) that contains at most one end of e_1 (or e_2). If $N(S) \subseteq S'$ and $|S'| \ge 2$, then $|S'| \ge |S| + 2$.

Proof. If $S = \emptyset$, the assertion is trivial. Now suppose that $S \neq \emptyset$. Since H is a bipartite matching covered graph, Lemma 2.4 implies that $|N_H(S)| \ge |S| + 1$. If S contains exactly one end of e_1 , then $|N(S)| = |N_H(S)| + 1 \ge |S| + 2$. If S contains no ends of e_1 , then |N(S)| =

 $|N_H(S)| \ge |S| + 1$. When |N(S)| = |S| + 1, Lemma 2.3 implies that N(S) is a nontrivial barrier of G. Since G is a brick, Lemma 2.1(i) implies that G has no nontrivial barriers, a contradiction. Thus $|N(S)| \ge |S| + 2$. Since $N(S) \subseteq S'$, we have $|S'| \ge |N(S)| \ge |S| + 2$. \Box

Since $\{e_1, e_2\}$ is a removable doubleton of G, none of e_1 and e_2 is removable in G. Let e be a nonremovable edge of G such that $e \notin \{e_1, e_2\}$. We say that e is of type I if e is removable in H, and of type I otherwise.

Lemma 2.8. If e is an edge of type I, then

- (i) ([2]) there exist partitions (A_1, A_2) of U and (B_1, B_2) of W such that $|B_1| = |A_1| + 1$, $|A_2| = |B_2| + 1$, $\{e_1\} = E(A_2)$, $\{e_2\} = E(B_1)$ and $\{e\} = E[A_1, B_2]$, and
- (ii) B_1 and A_2 are barriers of G e.

Proof. (ii) If e is of type I, by (i), $N_{G-e}(A_1) \subseteq B_1$. Since $|B_1| = |A_1| + 1$, Lemma 2.3 implies that B_1 is a barrier of G - e. Similarly, A_2 is also a barrier of G - e.

Now assume that e is of type II, by the definition, e is nonremovable in both G and H. So both G-e and H-e have nonadmissible edges. Note that if h is a nonadmissible edge of H-e and is admissible in G-e, then the perfect matching of G-e containing h contains e_1 and e_2 . Therefore, there exists an edge e^* that is nonadmissible in both H-e and G-e. Otherwise, G-e is matching covered, a contradiction. By Lemma 2.1(ii), G-e has a barrier containing $V(e^*)$. Let B be a maximal such barrier. Lemma 2.1(ii) implies that each component of G-e-B is factor-critical. In particular, each bipartite component is trivial, so G-e-B has at most two nontrivial components, one contains e_1 and the other contains e_2 .

Let U_2 and W_1 be the sets of the vertices in trivial components of G - e - B that lie in U and W, respectively. Then U_2 and W_1 are independent set of G. Let u and w be the two ends of e such that $u \in U$ and $w \in W$. Since e^* is admissible in G, e connects two distinct components of G - e - B, i.e., u and w lie in distinct components of G - e - B. Let $U_1 = B \cap U$ and $W_2 = B \cap W$. Then $u \notin U_1, w \notin W_2$,

$$e^* \in E[U_1, W_2], N_{H-e}(W_1) \subseteq U_1 \text{ and } N_{H-e}(U_2) \subseteq W_2.$$
 (1)

Let ω be the number of nontrivial components of G - e - B. Then $\omega \leq 2$ and

$$|B| = |U_1| + |W_2| = |W_1| + |U_2| + \omega.$$
(2)

Assume that G_1 and G_2 are the two nontrivial components of G - e - B. When $\omega = 0$, both G_1 and G_2 are null. When $\omega = 1$, for convenience, we assume that G_1 is null if $|U_1| = |W_1|$ and G_2 is null otherwise. Let $U_{i+2} = V(G_i) \cap U$ and $W_{i+2} = V(G_i) \cap W$, i = 1, 2. Then (U_1, U_2, U_3, U_4) is a partition of U and (W_1, W_2, W_3, W_4) is a partition of W. Note that

$$N_{H-e}(U_i) \subseteq W_2 \cup W_i \text{ and } N_{H-e}(W_i) \subseteq U_1 \cup U_i \text{ for } i = 3, 4.$$
 (3)

Observation 2.9. When $\omega = 0$, $U_3 = W_3 = U_4 = W_4 = \emptyset$; when $\omega = 1$, $U_3 = W_3 = \emptyset$ if $|U_1| = |W_1|$ and $U_4 = W_4 = \emptyset$ otherwise.

2.3 Properties

Proposition 2.10. $U_1 \neq \emptyset$, $W_2 \neq \emptyset$, $|U_1| \ge |W_1|$ and $|W_2| \ge |U_2|$.

Proof. Since $e^* \in E[U_1, W_2]$, we have $U_1 \neq \emptyset$ and $W_2 \neq \emptyset$. To show $|U_1| \geq |W_1|$ and $|W_2| \geq |U_2|$, it suffices to consider the case $W_1 \neq \emptyset$ and $U_2 \neq \emptyset$. Note that e is incident with at most one vertex of W_1 and U_2 , respectively. Since H is a bipartite matching covered graph, Lemma 2.4 and (1) implies that $|U_1| + 1 \geq |N_H(W_1)| \geq |W_1| + 1$ and $|W_2| + 1 \geq |N_H(U_2)| \geq |U_2| + 1$, so $|U_1| \geq |W_1|$ and $|W_2| \geq |U_2|$.

Proposition 2.11. If $\omega = 0$, then (i) $W_1 \neq \emptyset$ and $U_2 \neq \emptyset$; (ii) $|U_1| = |W_1|$ and $|U_2| = |W_2|$; (iii) $\{e\} = E[W_1, U_2], \{e_1\} = E[U_1, U_2]$ and $\{e_2\} = E[W_1, W_2]$.

Proof. Since $\omega = 0$, we have $u \in U_2$ and $w \in W_1$, so $U_2 \neq \emptyset$ and $W_1 \neq \emptyset$. Since e is the only edge connecting two distinct components of G - e - B, $\{e\} = E[U_2, W_1]$ and $V(e_i) \cap B \neq \emptyset$ for i = 1, 2. By (2), $|U_1| + |W_2| = |W_1| + |U_2|$. By Proposition 2.10, $|U_1| = |W_1|$ and $|U_2| = |W_2|$. Then $U_1 \neq \emptyset$. If W_1 contains no ends of e_2 , then $N(W_1) \subseteq U_1 \cup \{u\}$. By Lemma 2.7, $|U_1| \geq |W_1| + 1$, a contradiction. Thus $\{e_2\} = E[W_1, W_2]$. Similarly, $\{e_1\} = E[U_1, U_2]$.

Proposition 2.12. Assume that $\omega = 1$. Then

- (i) $|U_1| = |W_1| + 1$ and $|U_2| = |W_2|$, or $|U_1| = |W_1|$ and $|W_2| = |U_2| + 1$;
- (ii) if $|U_1| \neq |W_1|$, then $U_2 \neq \emptyset$, $\{e_1\} = E[U_1, U_2]$, $\{e_2\} = E(W_3)$, $|W_3| = |U_3| + 1$, and $\{e\} = E[U_2, W_1]$ if $W_1 \neq \emptyset$ and $\{e\} = E[U_2, W_3]$ otherwise;
- (iii) if $|U_1| = |W_1|$, then $W_1 \neq \emptyset$, $\{e_1\} = E(U_4)$, $\{e_2\} = E[W_1, W_2]$, $|U_4| = |W_4| + 1$, and $\{e\} = E[W_1, U_2]$ if $U_2 \neq \emptyset$ and $\{e\} = E[W_1, U_4]$ otherwise.

Proof. By (2), $|U_1| + |W_2| = |W_1| + |U_2| + 1$. (i) follows from Proposition 2.10.

(ii) If $|U_1| \neq |W_1|$, by Observation 2.9 and (i), $U_4 = W_4 = \emptyset$ and $|U_2| = |W_2|$. By Proposition 2.10, $W_2 \neq \emptyset$, so $U_2 \neq \emptyset$. If at most one edge in $\{e, e_1\}$ has an end in U_2 , say e, then $N(U_2) \subseteq W_2 \cup \{w\}$. By Lemma 2.7, $|W_2| \geq |U_2| + 1$, a contradiction. Thus each edge of $\{e, e_1\}$ has exactly one end in U_2 . This implies that $\{e\} = E[U_2, W_1 \cup W_3]$ and $\{e_1\} = E[U_1, U_2]$, so $\{e_2\} = E(W_3)$. Recall that G_1 is factor-critical. Then $|W_3| = |U_3| + 1$. If $\{e\} = E[U_2, W_3]$, then $N(W_1) \subseteq U_1$. Since $|U_1| = |W_1| + 1$, Lemma 2.3 implies that U_1 is a barrier of G. By Lemma 2.1(i), U_1 is a singleton and $W_1 = \emptyset$. (ii) holds.

If $|U_1| = |W_1|$, then $|U_2| \neq |W_2|$. Analogously, (iii) holds.

Recall that when $\omega = 2$, e_1 and e_2 lie in two nontrivial components of G - e - B, respectively. Assume, without loss of generality, that $e_1 \in E(G_2)$ and $e_2 \in E(G_1)$. Recall that $\{e_1\} = E(U)$ and $\{e_2\} = E(W)$. Then

$$\{e_1\} = E(U_4) \text{ and } \{e_2\} = E(W_3).$$
 (4)

Proposition 2.13. Assume that $\omega = 2$. Then

(i) $|U_i| = |W_i| + 1$ for i = 1, 4 and $|W_i| = |U_i| + 1$ for i = 2, 3;

(ii) when $U_2 \neq \emptyset$, $\{e\} = E[U_2, W_1]$ if $W_1 \neq \emptyset$ and $\{e\} = E[U_2, W_3]$ otherwise; (iii) when $U_2 = \emptyset$, $\{e\} = E[U_4, W_1]$ if $W_1 \neq \emptyset$ and $\{e\} = E[U_4, W_3]$ otherwise.

Proof. (i) By (1), $N(W_1) \subseteq U_1 \cup \{u\}$. By Proposition 2.10, $U_1 \neq \emptyset$. By Lemma 2.7, $|U_1| \ge |W_1| + 1$. Similarly, $|W_2| \ge |U_2| + 1$. By (2), $|U_1| + |W_2| = |W_1| + |U_2| + 2$, so $|U_1| = |W_1| + 1$ and $|W_2| = |U_2| + 1$. Recall that both G_1 and G_2 are factor-critical. By (4), we have $|U_4| = |W_4| + 1$ and $|W_3| = |U_3| + 1$.

To show (ii) and (iii), we first claim that $V(e) \cap (U_3 \cup W_4) = \emptyset$. Let $B' = U_1 \cup W_2 \cup U_3 \cup W_4$. In fact, if $V(e) \cap (U_3 \cup W_4) \neq \emptyset$, then $\overline{B'}$ is an independent set of H. By (i), $|\overline{B'}| = |B'|$. Since $N_H(\overline{B'}) \subseteq B'$, by Lemma 2.3, B' is a barrier of H. By (1), $e^* \in E[U_1, W_2]$, so $e^* \in E(B')$. Lemma 2.1(ii) implies that e^* is nonadmissible in H, a contradiction. The claim holds.

(ii) When $U_2 \neq \emptyset$, by (i), $|W_2| = |U_2| + 1 \ge 2$. If $u \notin U_2$, then $N(U_2) \subseteq W_2$. By Lemma 2.3, W_2 is a nontrivial barrier of G, a contradiction. Thus $u \in U_2$. Likewise, if $W_1 \neq \emptyset$, then $|U_1| \ge 2$ and $w \in W_1$. Thus $\{e\} = E[U_2, W_1]$. If $W_1 = \emptyset$, then $w \in W_3 \cup W_4$. By the above claim, $w \in W_3$, so $\{e\} = E[U_2, W_3]$.

(iii) When $U_2 = \emptyset$, by the above claim again, $u \in U_4$. If $W_1 = \emptyset$, then $\{e\} = E[U_4, W_3]$. If $W_1 \neq \emptyset$, by the same reason as the case $U_2 \neq \emptyset$, then $w \in W_1$, so $\{e\} = E[U_4, W_1]$. (iii) holds.

Corollary 2.14. (i) If $U_2 \neq \emptyset$, then $u \in U_2$; otherwise, $u \in U_4$ and $|W_2| = 1$.

- (ii) If $W_1 \neq \emptyset$, then $w \in W_1$; otherwise, $w \in W_3$.
- (iii) If $U_4 \neq \emptyset$, then $\{e_1\} = E(U_4)$; otherwise, $\{e_1\} = E[U_1, U_2]$.
- (iv) If $W_3 \neq \emptyset$, then $|U_1| = |W_1| + 1$ and $\{e_2\} = E(W_3)$; otherwise, $|U_1| = |W_1|$, $\{e_2\} = E[W_1, W_2]$ and $U_3 = \emptyset$.
- (v) $|U_2 \cup U_4| = |W_2 \cup W_4|.$

Proof. Since e = uv with $u \in U$ and $w \in W$, (i) and (ii) follow from Proposition 2.11-2.13. (iii), (iv) and (v) follow from (4), Observation 2.9, and Proposition 2.11-2.13.

Lemma 2.15. For any $u^* \in V(G)$, we have

- (i) u^* is incident with at most one nonremovable edge of type I;
- (ii) u^* is incident with at most two nonremovable edges of type II.

Proof. Without loss of generality, we may assume that $u^* \in U$.

(i) Suppose to the contrary that u^* is incident with two nonremovable edges of type I, say u^*w_1 and u^*w_2 . By Lemma 2.8, there is a barrier S_i of $G - u^*w_i$ that contains $V(e_1)$, i = 1, 2. Thus $|S_1 \cap S_2| \ge 2$, contradicting Lemma 2.2. (i) holds.

(ii) Suppose to the contrary that u^* is incident with three nonremovable edges of type II, say u^*w_1, u^*w_2 and u^*w_3 . Then u^*w_i $(1 \le i \le 3)$ is nonremovable in both G and H. By Lemma 2.5, there exist partitions (U'_0, U'_1, U'_2, U'_3) of U and (W'_0, W'_1, W'_2, W'_3) of W, such that $u^* \in U'_0$, and for $i \in \{1, 2, 3\}$: (a) $|U'_i| = |W'_i|$, (b) $w_i \in W'_i$, and (c) $N_H(W'_i) = U'_i \cup \{u^*\}$; in particular, u^*w_i is the only edge between U'_0 and W'_i . Note that at least one of W'_1, W'_2 and W'_3 contain no ends of e_2 , say W'_3 . Then $N(W'_3) = U'_3 \cup \{u^*\}$. By Lemma 2.3, $U'_3 \cup \{u^*\}$ is a nontrivial barrier of G, a contradiction. (ii) holds.

3 Proof of Theorem 1.1

We first present a lemma, the proof of which will be given later.

Lemma 3.1. Let $u \in U \setminus V(e_1)$, and uy and uw be two nonremovable edges in G such that uy is of type I and uw is of type II. Let w' be a neighbour of u other than y and w. Then either uw' is removable in G, or uw' is of type II, $e_2 = ww'$ and d(u) = d(w) = d(w') = 3.

Proof of Theorem 1.1. Assume u is a vertex in $U \setminus V(e_1)$ that is incident with more than two nonremovable edges of G. Then, by Lemma 2.15, u is incident with exactly one edge of type I and two edges of type II. We adopt the notational conventions stated in Lemma 3.1. Then uy is of type I, and uw and uw' are of type II. By Lemma 3.1, we have $e_2 = ww'$ and d(u) = d(w) = d(w') = 3. Therefore, u lies in the triangle containing e_2 and each vertex in this triangle is of degree three.

If there exist two vertices u_1 and u_2 in $U \setminus V(e_1)$ such that u_1 and u_2 are incident with more than two nonremovable edges of G, then we get two triangles T_1 and T_2 that contain u_1 and u_2 , respectively, and have the edge e_2 in common. By Lemma 2.6, for $i = 1, 2, u_i$ is incident with a removable edge of H in T_i that is of type I in G, contradicting the fact that it is of type II. Thus $U \setminus V(e_1)$ contains at most one vertex that is incident with more than two nonremovable edges of G.

The same result is also true for vertices in $W \setminus V(e_2)$. So the theorem follows.

Now we turn to the proof of Lemma 3.1. By Corollary 2.14 (i) and (ii), $u \in U_2 \cup U_4$, $w \in W_1 \cup W_3$ and $\{uw\} = E[U_2 \cup U_4, W_1 \cup W_3]$. Thus $\{y, w'\} \subset W_2 \cup W_4$. Since uy is of type I, Lemma 2.8 implies that there exists partitions (A_1, A_2) of U and (B_1, B_2) of W such that $|B_1| = |A_1| + 1$, $|A_2| = |B_2| + 1$, $\{e_1\} = E(A_2)$, $\{e_2\} = E(B_1)$ and $\{uy\} = E[A_1, B_2]$. Furthermore, B_1 and A_2 are barriers of G - uy. Note that

$$u \in A_1 \cap (U_2 \cup U_4), w \in B_1 \cap (W_1 \cup W_3), y \in B_2 \cap (W_2 \cup W_4), w' \in B_1 \cap (W_2 \cup W_4), \quad (5)$$

$$N(A_1 \setminus \{u\}) \subseteq B_1 \text{ and } N(B_2 \setminus \{y\}) \subseteq A_2.$$
 (6)

Combining (3) and Corollary 2.14(iii), we have

$$N((U_i \cap A_1) \setminus \{u\}) \subseteq (W_i \cup W_2) \cap B_1, i = 3, 4.$$

$$\tag{7}$$

3.1 Properties

Lemma 3.2. If $E[w', U_1 \cup U_3] \neq \emptyset$, then uw' is removable in G.

Proof. Suppose to the contrary that uw' is nonremovable in G. Since uy is of type I, Lemma 2.15(i) implies that uw' is of type II, so uw' is nonremovable in H. Then, there exists an edge f of H - uw' such that each perfect matching of H containing f contains uw'. Let M be a perfect matching of H containing both f and uw'. Let M_1 be a perfect matching of H containing uy. Then uw is neither in M nor in M_1 . Recall that $\{uw\} = E[U_2 \cup U_4, W_1 \cup W_3]$ and $\{y, w'\} \subseteq W_2 \cup W_4$. By Corollary 2.14(v), $|U_2 \cup U_4| = |W_2 \cup W_4|$. Let $X = U_2 \cup U_4 \cup W_2 \cup W_4$. Then $M \cap \partial(X) = \emptyset$ and $M_1 \cap \partial(X) = \emptyset$. Since $f \in M$, $f \in E(X)$ or $f \in E(\overline{X})$. If $f \in E(\overline{X})$, then $(M \cap E(\overline{X})) \cup (M_1 \cap E(X))$ is a perfect matching of H that contains f but does not contain uw', a contradiction. So $f \in E(X)$. Recall that $E[w', U_1 \cup U_3] \neq \emptyset$. Let z' be a neighbour of w' in $U_1 \cup U_3$ and M_2 a perfect matching of H that contains w'z'. Then $uw \in M_2$, so $M_2 \cap \partial(X) = \{w'z', uw\}$. It follows that $((M \cap E(X)) \setminus \{uw'\}) \cup (M_2 \cap (E(\overline{X}) \cup \partial(X)))$ is a perfect matching of H that contains. The assertion follows.

Proposition 3.3. (i) $|W_2 \cap B_1| \le 1$, $|U_1 \cap A_2| \le 1$ and $W_1 \subseteq B_1$; (ii) If $\{e_1\} = E[U_1, U_2]$, then $|U_1 \cap A_2| = 1$; (iii) If $U_2 \ne \emptyset$, then $W_2 \cap B_1 = \{w'\}$, $U_2 \cap A_1 = \{u\}$ and d(u) = 3.

Proof. (i) Recall that $U_1 \cup W_2$ is a barrier of G - uw, and B_1 and A_2 are barriers of G - uy. By Lemma 2.2, $|W_2 \cap B_1| \leq 1$ and $|U_1 \cap A_2| \leq 1$. By (5), $w \notin B_2$ and $y \notin W_1$. Recall that $\{e_2\} = E(B_1)$. By (1) and (6), $N(W_1 \cap B_2) \subseteq U_1 \cap A_2$. Since G is 3-connected, $W_1 \cap B_2 = \emptyset$, so $W_1 \subseteq B_1$. Therefore, (i) holds.

(ii) The result follows from (i) and the fact that $\{e_1\} = E(A_2) \cap E[U_1, U_2]$.

(iii) If $U_2 \neq \emptyset$, by (5) and Corollary 2.14(i), $u \in U_2 \cap A_1$. By (1) and (5), the neighbour w' of u lies in $W_2 \cap B_1$. As $|W_2 \cap B_1| \leq 1$, we have $W_2 \cap B_1 = \{w'\}$, so $N(u) = \{y, w, w'\}$, i.e., d(u) = 3. By (1) and (6), $N((U_2 \cap A_1) \setminus \{u\}) \subseteq W_2 \cap B_1$. Since G is 3-connected, $(U_2 \cap A_1) \setminus \{u\} = \emptyset$, so $U_2 \cap A_1 = \{u\}$. Therefore, (iii) holds.

Proposition 3.4. Assume that $U_2 = \emptyset$ and uw' is nonremovable in G. Then $w' \in W_4 \cap B_1$, $|W_4 \cap B_1| \ge |U_4 \cap A_1|$, and $|(W_2 \cup W_3) \cap B_1| \ge |U_3 \cap A_1| + 2$ if $W_3 \ne \emptyset$.

Proof. By Lemma 3.2, $E[w', U_1 \cup U_3] = \emptyset$. By (1), $E[W_2, U_1] \neq \emptyset$. By Corollary 2.14(i), $|W_2| = 1$, so $w' \notin W_2$. By (5), $w' \in W_4 \cap B_1$. By Corollary 2.14(iv), if $W_3 \neq \emptyset$, then $\{e_2\} = E(W_3)$. Recall that $\{e_2\} = E(B_1)$. Then $|W_3 \cap B_1| \ge 2$. By (7) and Corollary 2.14(i), $N(U_3 \cap A_1) \subseteq (W_3 \cup W_2) \cap B_1$ and $N((U_4 \cap A_1) \setminus \{u\}) \subseteq (W_4 \cup W_2) \cap B_1$. By Lemma 2.7, $|(W_2 \cup W_3) \cap B_1| \ge |U_3 \cap A_1| + 2$ if $W_3 \neq \emptyset$, and $|W_4 \cap B_1| \ge |U_4 \cap A_1|$ if $W_2 \subseteq B_1$. If $W_2 \subseteq B_2$, when $(U_4 \cap A_1) \setminus \{u\} \neq \emptyset$, Lemma 2.4 implies that $|W_4 \cap B_1| \ge |U_4 \cap A_1|$, which is also true when $U_4 \cap A_1 = \{u\}$. The assertion follows.

By Proposition 3.3(i), we proceed to consider the following two cases.

3.2 The case $|U_1 \cap A_2| = 1$

Lemma 3.5. Assume that $|U_1 \cap A_2| = 1$.

- (i) If $U_2 = \emptyset$ or $W_3 \neq \emptyset$, then uw' is removable in G.
- (ii) If $U_2 \neq \emptyset$ and $W_3 = \emptyset$, then uw' is removable in G, or uw' is of type II, $e_2 = ww'$ and d(u) = d(w) = 3.

Proof. Since $|U_1 \cap A_2| = 1$, $|U_1 \cap A_1| = |U_1| - 1$. By Proposition 3.3(i), $W_1 = W_1 \cap B_1$. By Proposition 2.11-2.13, $|W_1| \le |U_1| \le |W_1| + 1$, so $|W_1| - 1 \le |U_1 \cap A_1| \le |W_1 \cap B_1|$. In particular, if $|W_1| = |U_1|$, then $|W_1 \cap B_1| = |U_1 \cap A_1| + 1$. We now show the following claim.

Claim. If $U_2 \neq \emptyset$ and uw' is nonremovable in G, then $|W_2 \cap B_1| = |U_2 \cap A_1| = 1$, $|W_4 \cap B_1| \ge |U_4 \cap A_1|$, and $|W_3 \cap B_1| \ge |U_3 \cap A_1| + 2$ if $W_3 \ne \emptyset$.

Since $U_2 \neq \emptyset$, by Proposition 3.3(iii), $U_2 \cap A_1 = \{u\}$ and $W_2 \cap B_1 = \{w'\}$. Then $u \notin U_3 \cup U_4$. By Lemma 3.2, $E[W_2 \cap B_1, U_1 \cup U_3] = \emptyset$. By (7), $N(U_3 \cap A_1) \subseteq W_3 \cap B_1$ and $N(U_4 \cap A_1) \subseteq (W_4 \cap B_1) \cup \{w'\}$. By Lemma 2.4, $|W_4 \cap B_1| \ge |U_4 \cap A_1|$. If $W_3 \neq \emptyset$, by Corollary 2.14(iv), $\{e_2\} = E(W_3)$. Recall that $\{e_2\} = E(B_1)$. Then $|W_3 \cap B_1| \ge 2$. By Lemma 2.7, $|W_3 \cap B_1| \ge |U_3 \cap A_1| + 2$. Claim holds.

(i) Suppose to the contrary that uw' is nonremovable in *G*. We will show $|B_1| \ge |A_1| + 2$, contradicting the fact that $|B_1| = |A_1| + 1$. Recall that $|W_1 \cap B_1| \ge |U_1 \cap A_1|$. If $U_2 \ne \emptyset$ and $W_3 \ne \emptyset$, by the above claim, we have $|B_1| = \sum_{i=1}^4 |W_i \cap B_1| \ge \sum_{i=1}^4 |U_i \cap A_1| + 2 = |A_1| + 2$. If $U_2 = \emptyset$, by Proposition 3.4, $|W_4 \cap B_1| \ge |U_4 \cap A_1|$, and $|(W_2 \cup W_3) \cap B_1| \ge |U_3 \cap A_1| + 2$ if $W_3 \ne \emptyset$. Then $|B_1| \ge |A_1| + 2$. If $W_3 = \emptyset$, by Corollary 2.14(iv), $U_3 = \emptyset$, $\{e_2\} = E[W_1, W_2]$ and $|U_1| = |W_1|$. Then $|W_1 \cap B_1| = |U_1 \cap A_1| + 1$. Recall that $\{e_2\} = E(B_1)$. By Corollary 2.14(i), $|W_2| = 1$, so $W_2 \subseteq B_1$, i.e., $|W_2 \cap B_1| = 1$. Consequently, $|B_1| \ge |A_1| + 2$. (i) holds.

(ii) Since $U_2 \neq \emptyset$ and $W_3 = \emptyset$, by Corollary 2.14, $u \in U_2$, $\{e_2\} = E[W_1, W_2]$, $|U_1| = |W_1|$, $U_3 = \emptyset$ and $w \in W_1$. By Proposition 3.3(iii), d(u) = 3 and $W_2 \cap B_1 = \{w'\}$, so $w' \in V(e_2)$. Assume that uw' is nonremovable in G. By Lemma 2.15, uw' is of type II. Suppose to the contrary that $e_2 \neq ww'$. Then W_1 contains at least two vertices, w and one end of e_2 , i.e., $|W_1| \ge 2$. Since $|U_1| = |W_1|$, we have $|W_1 \cap B_1| = |U_1 \cap A_1| + 1$. Since $|B_1| = |A_1| + 1$, by the above claim, we have $|W_4 \cap B_1| = |U_4 \cap A_1|$. Recall that $\{e_1\} = E(A_2)$. Since $u \notin U_4$, by (7), $N(U_4 \cap A_1) \subseteq (W_4 \cap B_1) \cup \{w'\}$. By Lemma 2.7, $W_4 \cap B_1 = \emptyset$. By Lemma 3.2, $E[W_2 \cap B_1, U_1 \cup U_3] = \emptyset$, so $N(U_1 \cap A_1) \subseteq W_1 \cap B_1 = W_1$. By Lemma 2.3, W_1 is a nontrivial barrier of G. This contradiction implies that $e_2 = ww'$. By (1), $N(W_1 \setminus \{w\}) \subseteq U_1$. Since $|U_1| = |W_1 \setminus \{w\}| + 1$, Lemma 2.3 implies that U_1 is a barrier of G. By Lemma 2.1(i), $|U_1| = 1$, so $W_1 = \{w\}$. Then w has exactly three neighbours u, w' and the vertex in U_1 , i.e., d(w) = 3. (ii) holds.

3.3 The case $U_1 \cap A_2 = \emptyset$

If $U_1 \cap A_2 = \emptyset$, then $U_1 \subseteq A_1$, i.e., $U_1 \cap A_1 = U_1$. Recall that $W_1 \cap B_1 = W_1$. By Corollary 2.14(i), $u \notin U_1$ and, by (6), $N(U_1) \subseteq N(A_1 \setminus \{u\}) \subseteq B_1$.

Lemma 3.6. If $U_1 \cap A_2 = \emptyset$ and $U_2 \neq \emptyset$, then uw' is removable in G.

Proof. Suppose to the contrary that uw' is nonremovable in G. Since $U_2 \neq \emptyset$, by Proposition 3.3(iii), $W_2 \cap B_1 = \{w'\}$. By (1), $E[U_1, W_2] \neq \emptyset$. Since $N(U_1) \subseteq B_1$, we have $E[U_1, W_2 \cap B_1] \neq \emptyset$, i.e., $E[U_1, w'] \neq \emptyset$, contradicting Lemma 3.2.

To deal with the case $U_1 \cap A_2 = \emptyset$ and $U_2 = \emptyset$, we need to consider the properties with respect to uw'. If uw' is of type II, assume that (U'_1, U'_2, U'_3, U'_4) and (W'_1, W'_2, W'_3, W'_4) are partitions of U and W, respectively, with respect to uw' defined as those with respect to uwin Section 2. Then all the properties with respect to uw are also true with respect to uw'. If $U_2 = \emptyset$, by Proposition 2.11(i), $\omega \ge 1$. We say the edge uw is of type 1 if $\omega = 1$, and of type 2 if $\omega = 2$. Analogously, we may define the type of the edge uw' when $U'_1 \cap A_2 = \emptyset$ and $U'_2 = \emptyset$.

Lemma 3.7. Assume that $U_1 \cap A_2 = \emptyset$, $U_2 = \emptyset$ and uw' is nonremovable in G. Then uw' is neither of type 1 nor of type 2.

The proof of Lemma 3.8 requires Lemma 3.7, whose proof is presented in the next subsection.

Lemma 3.8. If $U_1 \cap A_2 = \emptyset$ and $U_2 = \emptyset$, then uw' is removable in G.

Proof. Suppose to the contrary that uw' is nonremovable in G. By Proposition 3.4, $w' \in W_4$. Recall that $N(u) = \{w, w', y\}$, uy is of type I and uw is of type II. By Lemma 2.15(i), uw' is of type II. So $|U'_1 \cap A_2| \leq 1$. By Lemma 3.7, uw' is neither of type 1 nor of type 2. Then $|U'_1 \cap A_2| = 1$ or $U'_2 \neq \emptyset$. By Corollary 2.14(iv), $\{e_2\} = E(W_3) \cup E[W_1, W_2]$. Since $w' \in W_4$, $w' \notin V(e_2)$, so $e_2 \neq ww'$. If $|U'_1 \cap A_2| = 1$, Lemma 3.5 implies that uw is removable in G, a contradiction. If $U'_1 \cap A_2 = \emptyset$ and $U'_2 \neq \emptyset$, Lemma 3.6 implies that uw is removable in G, also a contradiction.

Proof of Lemma 3.1. By Lemmas 3.5, 3.6 and 3.8, we only need to show d(w') = 3 when uw' is of type II. In this case, noticing uw is of type II, with w' playing the role of w, we have d(w') = 3 by Lemmas 3.5, 3.6 and 3.8 again.

3.4 Proof of Lemma 3.7

In this subsection, we show Lemma 3.7 by contradiction. Suppose that uw' is of type 1 or type 2. Then uw' is of type II, $U'_1 \cap A_2 = \emptyset$, $U'_2 = \emptyset$ and $|W'_2| = 1$. We may let $W'_2 = \{w'_2\}$. We also have (a) $\{e_1\} = E(U'_4)$, (b) if $W'_1 \neq \emptyset$, then $\{uw'\} = E[U'_4, W'_1]$; otherwise, $\{uw'\} = E[U'_4, W'_3]$, and (c) if $W'_3 \neq \emptyset$, then $\{e_2\} = E(W'_3)$; otherwise, $\{e_2\} = E[W'_1, W'_2]$. Recall that uw is of type II. Since $U_1 \cap A_2 = \emptyset$ and $U_2 = \emptyset$, uw is also of type 1 or type 2, so $|W_2| = 1$. Let $W_2 = \{w_2\}$. Note that $u \in U_4 \cap U'_4 \cap A_1$ and $\{e_1\} = E(U_4) \cap E(U'_4)$. The following properties with respect to uw are also true with respect to uw'. **Proposition 3.9.** $|U_4 \cap A_1| = |W_4 \cap B_1|, A_2 \subseteq U_4 \text{ and } B_2 \subseteq W_4.$

Proof. Recall that $W_1 \subseteq B_1$, $U_1 \subseteq A_1$, $U_2 = \emptyset$ and $N(U_1) \subseteq B_1$. By (1), $E[U_1, w_2] \neq \emptyset$, so $w_2 \in B_1$, i.e., $|W_2 \cap B_1| = 1$. By Proposition 3.4, $|W_4 \cap B_1| \ge |U_4 \cap A_1|$, and $|(W_2 \cup W_3) \cap B_1| \ge |U_3 \cap A_1| + 2$ if $W_3 \neq \emptyset$. If uw is of type 1, by Proposition 2.12(ii), $|W_1| = |U_1|$, so $|W_1 \cap B_1| = |U_1 \cap A_1|$. By Observation 2.9, $U_3 = W_3 = \emptyset$, so $A_2 \subseteq U_4$ and $B_2 \subseteq W_4$. Since $|B_1| = |A_1| + 1$, we have $|W_4 \cap B_1| = |U_4 \cap A_1|$. If uw is of type 2, by (4) and Proposition 2.13, $W_3 \neq \emptyset$ and $|W_1| = |U_1| - 1$, so $|W_1 \cap B_1| = |U_1 \cap A_1| - 1$. Since $|B_1| = |A_1| + 1$, we have $|W_4 \cap B_1| = |U_3 \cap A_1| = |U_3 \cap A_1| + 1$. Note that $|W_3| = |U_3| + 1$. Then $|W_3 \cap B_2| = |U_3 \cap A_2|$. Recall that $w \in B_1$, $y \in B_2 \cap (W_2 \cup W_4)$ and $N(U_1) \subseteq B_1$. By (3) and (6), $N(W_3 \cap B_2) \subseteq U_3 \cap A_2$. By Lemma 2.4, $W_3 \cap B_2 = \emptyset$, so $U_3 \cap A_2 = \emptyset$, i.e., $W_3 \subseteq B_1$ and $U_3 \subseteq A_1$. Thus $A_2 \subseteq U_4$ and $B_2 \subseteq W_4$.

Recall that $\{w_2\} \cup U_1$ and $\{w'_2\} \cup U'_1$ are barriers of G - uw and G - uw', respectively. Lemma 2.2 implies that

$$|(\{w_2\} \cup U_1) \cap (\{w_2'\} \cup U_1')| \le 1.$$
(8)

Since $u \in U_4$ and $\{e_1\} = E(U_4)$, by (3),

$$N(U_3 \cap U'_j) \subseteq (W_3 \cup \{w_2\}) \cap (W'_j \cup \{w'_2\}), \text{ where } j = 3, 4.$$
(9)

Proposition 3.10. (i) $W_1 \cap W'_1 = \emptyset$; and (ii) if $w_2 \in W'_1 \cup W'_3$ and $w'_2 \in W_1 \cup W_3$, then $|W_4 \cap W'_4 \cap B_1| \ge |U_4 \cap U'_4 \cap A_1| + 1$.

Proof. By Proposition 3.4, $w' \in W_4$ and $w \in W'_4$.

(i) By (8), $|U_1 \cap U'_1| \leq 1$. Note that $w, w' \notin W_1 \cap W'_1$. By (1), $N_{G-e_2}(W_1 \cap W'_1) \subseteq U_1 \cap U'_1$. As G is 3-connected, (i) follows.

(ii) Note that $w_2 \neq w'$ and $w'_2 \neq w$. Since $w_2 \in W'_1 \cup W'_3$ and $w'_2 \in W_1 \cup W_3$, by (1) and (3), w_2 and w'_2 have no neighbours in $U_4 \cap U'_4$. Recall that $u \in U_4 \cap U'_4 \cap A_1$. By Proposition 3.9, $A_2 \subseteq (U_4 \cap U'_4) \setminus \{u\}$. By (3), $N(A_2) \subseteq N((U_4 \cap U'_4) \setminus \{u\}) \subseteq W_4 \cap W'_4$. By (6), $N(B_2 \setminus \{y\}) \subseteq A_2$. Then $N(A_2 \cup (B_2 \setminus \{y\})) \subseteq \{y\} \cup (W_4 \cap W'_4 \cap B_1)$. Since G is 3-connected, $|W_4 \cap W'_4 \cap B_1| \ge 2$. By (3) and (6), $N((U_4 \cap U'_4 \cap A_1) \setminus \{u\}) \subseteq W_4 \cap W'_4 \cap B_1$, (ii) follows from Lemma 2.7.

Proposition 3.11. Neither uw nor uw' is of type 2.

To complete the proof of Lemma 3.7, we need Proposition 3.11, the proof of that is a main burden. We present it later. From Proposition 3.11, we see that both uw and uw' are of type 1. By Proposition 2.12 and Observation 2.9, $\{e_2\} = E[W_1, W_2] \cap E[W'_1, W'_2]$ and $U_3 = W_3 = \emptyset$. Then each of w_2 and w'_2 is an end of e_2 . By Proposition 3.10(i), $W_1 \cap W'_1 = \emptyset$. If $w_2 = w'_2$, then the other end of e_2 lies in $W_1 \cap W'_1$, a contradiction. Thus $w_2 \neq w'_2$. Then $e_2 = w_2w'_2$, so $w'_2 \in W_1$ and $w_2 \in W'_1$. The former implies that $|W_1 \cap W'_2| = 1$. Proposition 3.10(ii) implies that $|U_4 \cap U'_4 \cap A_1| \leq |W_4 \cap W'_4 \cap B_1| - 1$. By Proposition 3.9, $|U'_4 \cap A_1| = |W'_4 \cap B_1|, U_1 \subseteq A_1$ and $W_1 \subseteq B_1$. Note that (U_1, U_4) is a partition of U and (W_1, W_2, W_4) is a partition of W. Since $w_2 \in W'_1$, we have $|U_1 \cap U'_4 \cap A_1| \ge |W_1 \cap W'_4 \cap B_1| + 1$, so $|U_1 \cap U'_4| \ge |W_1 \cap W'_4| + 1$. By Proposition 3.4, $w' \in W_4$, so $w_2 \ne w'$. By (1), $N(w_2) \subseteq U'_1 \cup \{w'_2\}$. Since $E[w_2, U_1] \ne \emptyset$ and $d(w_2) \ge 3$, we have $U_1 \cap U'_1 \ne \emptyset$, so $|U_1 \cap U'_1| = 1$. Consequently, $|U_1| \ge |W_1| + 1$, contradicting the fact that $|U_1| = |W_1|$. So Lemma 3.7 holds.

3.5 Proof of Proposition 3.11

Assume, without loss of generality, that uw is of type 2. We first present some basic properties.

Proposition 3.12. Assume that $|U_1| \ge 2$. Then $|U_1 \cap U'_1| + |U_1 \cap U'_4| = |W_1 \cap W'_4| + 1$ and, if $w'_2 \notin W_1$, then $U_1 \cap U'_3 = W_1 \cap W'_3 = \emptyset$; otherwise, $|U_1 \cap U'_1| = 1$, $|U_1 \cap U'_3| = |W_1 \cap W'_3| + 1$ and $w_2 \in W'_1 \cup W'_3$.

Proof. Since *uw* is of type 2, $|U_1| = |W_1| + 1$, so $W_1 \neq \emptyset$. By Corollary 2.14(ii) and Proposition 3.4, *w* ∈ $W_1 \cap W'_4$. Note that $\{e_2\} = E(W_3)$. By (1) and (3), $N(W_1 \cap W'_4) \subseteq (U_1 \cap (U'_4 \cup U'_1)) \cup \{u\}$ and $N(W_1 \cap W'_3) \subseteq U_1 \cap (U'_1 \cup U'_3)$. For the former case, since $d(w) \ge 3$, we have $|U_1 \cap (U'_4 \cup U'_1)| \ge 2$. By Lemma 2.7, $|U_1 \cap (U'_4 \cup U'_1)| \ge |W_1 \cap W'_4| + 1$. For the latter, if $W_1 \cap W'_3 \neq \emptyset$, Lemma 2.4 implies that $|U_1 \cap (U'_1 \cup U'_3)| \ge |W_1 \cap W'_3| + 1$. By (8), $|U_1 \cap U'_1| \le 1$, so $|U_1 \cap U'_3| \ge |W_1 \cap W'_3|$, which is also true when $W_1 \cap W'_3 = \emptyset$. Recall that $W_1 \cap W'_1 = \emptyset = U_2 = U'_2$.

Assume that $w'_2 \notin W_1$. Then $W_1 \cap W'_2 = \emptyset$. Since $|U_1| = |W_1| + 1$, we have $\sum_{i=1}^4 |U_1 \cap U'_i| = \sum_{i=1}^4 |W_1 \cap W'_i| + 1$. Then $|U_1 \cap (U'_4 \cup U'_1)| = |W_1 \cap W'_4| + 1$ and $|U_1 \cap U'_3| = |W_1 \cap W'_3|$, the latter implies that $|U_1 \cap (U'_1 \cup U'_3)| \le |W_1 \cap W'_3| + 1$. Since $W_1 \cap W'_3$ is an independent set, Lemma 2.3 implies that $U_1 \cap (U'_1 \cup U'_3)$ is a barrier of G. By Lemma 2.1(i), $U_1 \cap (U'_1 \cup U'_3)$ is a singleton. As G is 3-connected, $W_1 \cap W'_3 = \emptyset = U_1 \cap U'_3$.

Assume that $w'_2 \in W_1$. Then uw' is of type 2, otherwise, $w'_2 \in V(e_2)$, so $w'_2 \in W_3$, a contradiction. So $\{e_2\} = E(W'_3)$ and $|W'_3| = |U'_3| + 1$. If $E[w'_2, U'_3] = \emptyset$, then $N(U'_3) \subseteq W'_3$. By Lemma 2.3, W'_3 is a nontrivial barrier of G, a contradiction. Therefore, $E[w'_2, U'_3] \neq \emptyset$. Recall that $E[w'_2, U'_1] \neq \emptyset$ and $w'_2 \neq w$. Since $w'_2 \in W_1$ and $N(W_1 \setminus \{w\}) \subseteq U_1$, we have $N(w'_2) \subseteq U_1$. So $U_1 \cap U'_1$ and $U_1 \cap U'_3$ are nonempty, the former implies that $|U_1 \cap U'_1| = 1$. By Lemma 2.7, $|U_1 \cap (U'_1 \cup U'_3)| \ge |W_1 \cap W'_3| + 2$, so $|U_1 \cap U'_3| \ge |W_1 \cap W'_3| + 1$. Since $|U_1| = |W_1| + 1$ and $|W_1 \cap W'_2| = 1$, by the same reason as the case $w'_2 \notin W_1$, we have $|U_1 \cap (U'_4 \cup U'_1)| = |W_1 \cap W'_4| + 1$ and $|U_1 \cap U'_3| = |W_1 \cap W'_3| + 1$.

We now show that $w_2 \in W'_1 \cup W'_3$. If not, $w_2 \in \{w'_2\} \cup W'_4$. By (8), $w_2 \neq w'_2$, so $w_2 \in W'_4$. If $|U'_1| = 1$, then $W'_1 = \emptyset$. Combining $|U_1 \cap U'_1| = 1$, we have $W_3 \cap W'_1 = U_3 \cap U'_1 = \emptyset$, which is also true when $|U'_1| \ge 2$ (as above) since $w_2 \notin W'_1$. Recall that $w'_2 \in W_1$. By (9), $N(U_3 \cap U'_3) \subseteq W_3 \cap W'_3$ and $N(U_3 \cap U'_4) \subseteq \{w_2\} \cup (W_3 \cap W'_4)$. Because $\{e_2\} = E(W_3) \cap E(W'_3)$, Lemma 2.7 implies that $|W_3 \cap W'_3| \ge |U_3 \cap U'_3| + 2$. If $U_3 \cap U'_4 \ne \emptyset$, Lemma 2.4 implies that $|W_3 \cap W'_4| \ge |U_3 \cap U'_4|$, which is also true when $U_3 \cap U'_4 = \emptyset$. Note that $W_3 \cap W'_2 = \emptyset$. We have $|W_3| \ge |U_3| + 2$, contradicting the fact that $|W_3| = |U_3| + 1$. **Proposition 3.13.** Assume that $w'_2 \in W_1 \cup W_3$. If $|U_1| \ge 2$ and $|U_1 \cap U'_1| = 1$, or $|U_1| = 1$ and $U_1 \cap U'_4 = \emptyset$, then $w_2 \in W'_4$.

Proof. Suppose to the contrary that $w_2 \notin W'_4$. Then $W_2 \cap W'_4 = \emptyset$. Since $w'_2 \in W_1 \cup W_3$, we have $w_2 \neq w'_2$, so $w_2 \in W'_1 \cup W'_3$. By Proposition 3.10(ii), $|W_4 \cap W'_4 \cap B_1| \ge |U_4 \cap U'_4 \cap A_1| + 1$. By (9), $N(U_3 \cap U'_4) \subseteq W_3 \cap (W'_4 \cup \{w'_2\})$. Lemma 2.4 implies that $|W_3 \cap W'_4| \ge |U_3 \cap U'_4|$. We assert that $|W_1 \cap W'_4| = |U_1 \cap U'_4|$, which follows from Proposition 3.12 if $|U_1| \ge 2$ and $|U_1 \cap U'_1| = 1$, and from the fact that $U_1 \cap U'_4 = \emptyset$ and $W_1 = \emptyset$ if $|U_1| = 1$. By Proposition 3.9, $U_1, U_3 \subseteq A_1$ and $W_1, W_3 \subseteq B_1$. Then $|W'_4 \cap B_1| = \sum_{i=1}^4 |W'_4 \cap B_1 \cap W_i| \ge \sum_{i=1}^4 |U'_4 \cap A_1 \cap U_i| + 1 = |U'_4 \cap A_1| + 1$, contradicting the fact that $|W'_4 \cap B_1| = |U'_4 \cap A_1|$.

If uw' is of type 1, then $w' \in W'_1$, $\{e_2\} = E[w'_2, W'_1]$, and let k = 1; otherwise, $w' \in W'_1$ or $w' \in W'_3$, $\{e_2\} = E(W'_3)$, and let k = 3. Since uw is of type 2, we have $\{e_2\} = E(W_3)$. Thus, $W_3 \cap W'_k$ contains at least one end of e_2 , i.e., $W_3 \cap W'_k \neq \emptyset$. By Proposition 3.4, $w \in W'_4$ and $w' \in W_4$, so $w' \notin V(e_2)$ and $N_H(W_3 \cap W'_k) \subseteq (U_1 \cup U_3) \cap (U'_1 \cup U'_k)$. Lemma 2.4 implies that

$$|(U_1 \cup U_3) \cap (U_1' \cup U_k')| \ge |W_3 \cap W_k'| + 1.$$
(10)

By Lemma 3.2, $E[w', U_1 \cup U_3] = \emptyset$, so $N(w') \subseteq (U_4 \cap (U'_1 \cup U'_k)) \cup \{u\}$. Since $d(w') \geq 3$, $|U_4 \cap (U'_1 \cup U'_k)| \geq 2$. Since $N((W_4 \cap W'_k) \setminus \{w'\}) \subseteq (U_1 \cup U_4) \cap (U'_1 \cup U'_k)$, Lemma 2.7 implies that

$$|(U_1 \cup U_4) \cap (U_1' \cup U_k')| \ge |W_4 \cap W_k'| + 1.$$
(11)

If uw' is of type 1, then k = 1 and $|U_1'| = |W_1'|$. By (10) and (11), $|W_1'| = |(U_1 \cup U_4) \cap U_1'| + |(U_1 \cup U_3) \cap U_1'| - |U_1 \cap U_1'| \geq |W_4 \cap W_1'| + |W_3 \cap W_1'| - |U_1 \cap U_1'| + 2$. Since $W_1 \cap W_1' = \emptyset$, we have $|W_1'| = \sum_{i=2}^4 |W_i \cap W_1'|$, so $|W_2 \cap W_1'| + |U_1 \cap U_1'| \geq 2$. Since $|U_1 \cap U_1'| \leq 1$, we have $|U_1 \cap U_1'| = 1$ and $w_2 \in W_1'$. If $|U_1| = 1$, then $U_1 \cap U_4' = \emptyset$. Since $w_2' \in V(e_2)$, $w_2' \in W_3$. By Proposition 3.13, $w_2 \in W_4'$, a contradiction. Then uw' is of type 2, so $|W_3'| = |U_3'| + 1$ and k = 3. Recall that uw is of type 2. Then $\{e_2\} = E(W_3) \cap E(W_3')$. Let $Z_3 = (U_3 \cap U_3') \cup (W_3 \cap W_3')$. Then $|Z_3| \geq 2$. Since $w, w' \notin W_3 \cap W_3'$, $N(W_3 \cap W_3') \subseteq (U_1 \cup U_3) \cap (U_1' \cup U_3')$. Combining (9), we have

$$N(Z_3) \subseteq (\{w_2\} \cap (\{w_2'\} \cup W_3')) \cup (\{w_2'\} \cap W_3) \cup (U_1 \cap (U_1' \cup U_3')) \cup (U_1' \cap U_3).$$
(12)

If $|U_1| = 1$ or $|U'_1| = 1$, let $U_1 = \{u_1\}$ or $U'_1 = \{u'_1\}$, respectively.

Proposition 3.14. If $|U_1| \ge 2$ and $w'_2 \notin W_1$, or $|U_1| = 1$, then $|U'_1| \ge 2$. Analogously, if $|U'_1| \ge 2$ and $w_2 \notin W'_1$, or $|U'_1| = 1$, then $|U_1| \ge 2$.

Proof. Suppose to the contrary that $|U'_1| = 1$. Then $W'_1 = \emptyset$ and $|(U_1 \cup U_3 \cup U_4) \cap U'_1| = 1$. First consider the case $|U_1| \ge 2$ and $w'_2 \notin W_1$. By Proposition 3.12, $U_1 \cap U'_3 = W_1 \cap W'_3 = \emptyset$. By (10) and (11), $|U_1 \cap U'_1| + |U_4 \cap U'_3| + |U_3 \cap U'_3| \ge |W_4 \cap W'_3| + |W_3 \cap W'_3| + 1$. If $|U_1 \cap U'_1| = 0$, since $|W_2 \cap W'_3| \le 1$, we have $|U'_3| \ge |W'_3|$, a contradiction. If $|U_1 \cap U'_1| = 1$, by (8), $w'_2 \ne w_2$, so $w'_2 \in W_3 \cup W_4$. If $w'_2 \in W_4$, by (12) and the fact that $U_1 \cap U'_3 = \emptyset$, then $N(Z_3) \subseteq \{u'_1, w_2\}$, contradicting the fact that G is 3-connected. Thus $w'_2 \in W_3$. By Proposition 3.13, $w_2 \in W'_4$, i.e., $|W_2 \cap W'_3| = 0$. Therefore, $|U'_3| \ge |W'_3|$, a contradiction.

Now assume that $|U_1| = 1$. Then $W_1 = \emptyset$. If $w'_2 = w_2$, since G is 3-connected, by (12), we have $N(Z_3) = \{w_2, u_1, u'_1\}, u_1 \in U'_3 \text{ and } u'_1 \in U_3$. By (10) and (11), we have $|U_3 \cap U'_3| \ge |W_3 \cap W'_3| - 1$ and $|(U_1 \cup U_4) \cap U'_3| \ge |W_4 \cap W'_3| + 1$. Consequently, $|U'_3| \ge |W'_3|$, a contradiction. So $w'_2 \ne w_2$. Then $w'_2 \in W_3 \cup W_4$. Recall that $w'_2 u'_1 \in E(G)$. If $w'_2 \in W_4$, then $u'_1 \in U_4 \cup U_1$. By (12), $N(Z_3) \subseteq \{u_1, w_2\}$, a contradiction. Then $w'_2 \in W_3$, so $u'_1 \in U_1 \cup U_3$. Analogously, $w_2 \in W'_3$ and $u_1 \in U'_1 \cup U'_3$. Then $U_1 \cap U'_4 = \emptyset$. By Proposition 3.13, $w_2 \in W'_4$, a contradiction.

We are ready to complete the proof of Proposition 3.11 by distinguishing the following two cases to get contradictions. By Proposition 3.4, $w \in W'_4$, so $w'_2 \neq w$.

Case 1. $|U_1| \ge 2$ and $|U'_1| \ge 2$. By Proposition 3.12, we may claim that $U_1 \cap U'_3 = \emptyset$ if $w'_2 \notin W_1$ and $U'_1 \cap U_3 = \emptyset$ if $w_2 \notin W'_1$. First suppose that $U_1 \cap U'_1 = \emptyset$. We assert that $w'_2 \notin W_1$. Otherwise, $N(w'_2) \subseteq N(W_1 \setminus \{w\}) \subseteq U_1$. Since $E[w'_2, U'_1] \neq \emptyset$, $N(w'_2) \cap (U_1 \cap U'_1) \neq \emptyset$, a contradiction. Analogously, $w_2 \notin W'_1$. By the above claim, $U_1 \cap U'_3 = U'_1 \cap U_3 = \emptyset$. By (12), $N(Z_3) \subseteq \{w_2, w'_2\}$, contradicting the fact that G is 3-connected. Now suppose that $|U_1 \cap U'_1| = 1$. By (8), $w'_2 \neq w_2$. When $w'_2 \in W_1 \cup W_3$, Proposition 3.13 implies that $w_2 \in W'_4$, so $w_2 \notin W'_1$. If $w'_2 \notin W_1$, by (12), $N(Z_3) \subseteq \{w'_2\} \cup (U_1 \cap U'_1)$, a contradiction. Thus $w'_2 \in W_1$. By Proposition 3.12, $w_2 \in W'_1 \cup W'_3$, a contradiction. When $w'_2 \in W_4$, analogously, we deduce that $w_2 \in W'_1$, so $w'_2 \in W_1 \cup W_3$, a contradiction.

Case 2. At least one of U_1 and U'_1 is a singleton. Assume, without loss of generality, that $|U'_1| = 1$. By Proposition 3.14, $|U_1| \ge 2$ and $w'_2 \in W_1$. Proposition 3.12 implies that $w_2 \in W'_1 \cup W'_3$. Recall that $w'_2 \ne w$. Since $N(w'_2) \subseteq N(W_1 \setminus \{w\}) \subseteq U_1$ and $u'_1 w'_2 \in E(G)$, we have $u'_1 \in U_1$, so $|U'_1 \cap U_1| = 1$. By Proposition 3.13, $w_2 \in W'_4$, a contradiction.

4 Proof of Theorem 1.2

Let G be a graph and $C = \partial(X)$ an edge cut of G. We call G/X and G/\overline{X} the C-contractions of G, and we say that G is a splicing of G/X and G/\overline{X} (at x and \overline{x}), or G is obtained by splicing G/X and G/\overline{X} (at x and \overline{x}). A graph G is a tri-ladder if it is $\overline{C_6}$ or it may be obtained from $\overline{C_6}$ by iterative splicings K_4 at vertices in triangles. More precisely, there exists a sequence of graphs (G_0, G_1, \ldots, G_r) such that $G_0 = \overline{C_6}$, $G = G_r$, and, for $1 \leq i \leq r$, G_i is a splicing of G_{i-1} and K_4 at vertices in triangles. Note that each G_i is a tri-ladder and G_0 (= $\overline{C_6}$) has exactly two disjoint triangles, say T and T_0 . For convenience, we may assume that the vertices in T are never used to splice with K_4 . Let u, v and w be three vertices of T_0 . Let w' be a vertex of K_4 . Suppose that G_1 is a splicing of G_0 and K_4 at w and w'. Then $uv \in E(G_1)$, and uv is referred to as a rung of G_i $(1 \leq i \leq r)$. Moreover, G_1 has two disjoint triangles, one is T and the other is $T_1 = K_4 - w'$. Continuing in this way, we see that each G_i , $1 \leq i \leq r$, has one more rung than G_{i-1} , and has two disjoint triangles, one is T and the other is denoted by T_i . Then G_i has i rungs. For the rung e in G_i not in G_{i-1} , we refer to i as the rank of e, denoted by R(e). Note that the r rungs form a matching. Deleting them from G, the resulting graph has exactly three vertex-disjoint paths connecting vertices in T and T_r , which are referred to as *ridges* of G. The four graphs depicted in Figure 1 are near-bipartite tri-ladders whose rungs are illustrated by bold lines.

Since K_4 and $\overline{C_6}$ are cubic bricks, by the following lemma, a tri-ladder is a cubic brick.

Lemma 4.1 ([3]). Any splicing of two cubic bricks is a cubic brick.

Lemma 4.2 ([3]). Any splicing of two matching covered graphs is a matching covered graph.

The following lemma is an immediate consequence of Lemma 4.2, also see [15].

Lemma 4.3. Suppose that G is a splicing of two matching covered graphs G_1 and G_2 at u_1 and u_2 , where $u_i \in V(G_i)$, i = 1, 2. Then every removable edge in G_i that is not incident with u_i is removable in G.

For a bipartite graph with a perfect matching, by Hall's Theorem, we can obtain a characterization of nonadmissible edges of the graph. Using this characterization, the following lemma can be easily proved.

Lemma 4.4 ([15]). Let G be a 3-connected cubic nonbipartite graph. If G has a pair of edges e_1 and e_2 such that $G - \{e_1, e_2\}$ is bipartite, then G is a near-bipartite graph with removable doubleton $\{e_1, e_2\}$.

Lemma 4.5 ([2]). In a brace on six or more vertices, every edge is removable.

Lemma 4.6 ([8]). In a cubic matching covered graph, each 3-cut is a separating cut.

It is known that a cubic graph is 3-connected if and only if it is 3-edge-connected. A cubic graph is termed *essentially* 4-edge-connected if it is 2-edge-connected and free of nontrivial 3-cuts. Note that every nontrivial 3-cut is a matching in a 3-connected graph. An edge cut C of a matching covered G is good if it is separating but not tight.

Lemma 4.7 ([8]). Every essentially 4-edge-connected cubic graph is either a brick or a brace.

Lemma 4.8. Let G be a 3-connected cubic near-bipartite graph with removable doubleton $\{e_1, e_2\}$, and $H = G - \{e_1, e_2\}$ with a bipartition (U, W) such that $V(e_1) \subseteq U$. Suppose that $\partial(X)$ is a nontrivial 3-cut of G. Then |X| is odd, and G/\overline{X} and G/X are 3-connected cubic graphs. Moreover, if $|X \cap U| \ge |X \cap W|$, then the following statements hold.

(i) $|X \cap U| = |X \cap W| + 1$.

- (ii) ([9]) $\partial(X)$ is tight in G if and only if $E[X \cap W, \overline{X} \cap U] = \emptyset$, one of e_1 and e_2 has both ends in $X \cap U$ or $\overline{X} \cap W$ (adjust notation so that $V(e_1) \subseteq X \cap U$), and e_2 has at least one end in $X \cap W$. In addition, G/X is bipartite and G/\overline{X} is a near-bipartite graph with removable doubleton $\{e_1, e_2\}$.
- (iii) $\partial(X)$ is good in G if and only if $V(e_1) \subseteq X \cap U$, $V(e_2) \subseteq \overline{X} \cap W$, $|E[X \cap U, \overline{X} \cap W]| = 2$ and $E[X \cap W, \overline{X} \cap U]$ contains only one edge (say zw, where $w \in X \cap W$). In addition, zwis nonremovable in G, and both G/\overline{X} and G/X are near-bipartite graphs with removable doubleton $\{e_1, w\overline{x}\}$ and $\{e_2, zx\}$, respectively.

Proof. Since G is cubic and $\partial(X)$ is a 3-cut, G/\overline{X} and G/X are cubic, and $|\partial(X)|$ and |X| have the same parity, so |X| is odd. Since G is 3-connected, so are G/\overline{X} and G/X. To show (i) and (iii), let $U' = X \cap U$, $W' = X \cap W$, $U'' = \overline{X} \cap U$, $W'' = \overline{X} \cap W$, a = |U'| and b = |W'|.

(i) Suppose to the contrary that $a \ge b+2$. Since G[U'] has at most one edge e_1 , $|\partial(X)| \ge 3a-2-3b \ge 4$, a contradiction. (i) holds.

(iii) Suppose that $\partial(X)$ is good in G. Let $c = |E[U', \overline{X}]|$. Recall that $\partial(X)$ is a 3-cut. Then $c \leq 3$ and $|E[W', \overline{X}]| = 3 - c$. By counting the number of edges in E[U', W'] in two ways, we have 3a - 2|E(U')| - c = 3b - 2|E(W')| - (3 - c). Since a = b + 1, we have

$$c = |E(W')| - |E(U')| + 3.$$
(13)

If U' has at most one end of e_1 , then $E(U') = \emptyset$. Since $c \leq 3$, we have c = 3, $E(W') = \emptyset$ and $E[W', \overline{X}] = \emptyset$. Thus $e_2 \in E(W'')$. By (ii), $\partial(X)$ is tight in G, a contradiction. So $e_1 \in E(U')$. Analogously, $e_2 \in E(W'')$, so $E(W') = \emptyset$. By (13), we have c = 2, so |E[U', W'']| = 2 and E[W', U''] contains only one edge, say zw, where $w \in W'$. Because every perfect matching of G that contains e_1 contains zw, zw is nonremovable in G. Since both $G/\overline{X} - \{e_1, w\overline{x}\}$ and $G/X - \{e_2, zx\}$ are bipartite, Lemma 4.4 implies that both G/\overline{X} and G/X are near-bipartite graphs with removable double $\{e_1, w\overline{x}\}$ and $\{e_2, zx\}$, respectively.

Conversely, by Lemma 4.6, $\partial(X)$ is a separating cut of G. From (ii), we see that $\partial(X)$ is not tight in G, so it is good in G. (iii) holds.

Let G be a 3-connected cubic graph. If n = 4, then $G = K_4$. If $n \ge 6$ and G has a nontrivial 3-cut C, then each C-contraction of G is also a 3-connected cubic graph that has strictly fewer vertices than G. If either of the C-contractions has a nontrivial 3-cut, then the graph can be further decomposed into even smaller graphs. This procedure can be repeated until we obtain a list of 3-connected cubic graphs each of which is free of nontrivial 3-cuts. We refer to it as a 3-cut-decomposition of G. A 3-cut-decomposition is referred to as a K_4 decomposition if it results in a list of K_4 s. If a graph has a K_4 -decomposition, by Lemma 4.1, it is a cubic brick and can be obtained by sequentially splicing cubic bricks.

Lemma 4.9. Let G be a cubic near-bipartite brick with at least six vertices. Then G has a K_4 -decomposition if and only if G is a tri-ladder.

Proof. Use induction on n. Note that $\overline{C_6}$ is the only graph obtained by splicing two K_4 s, and it is a near-bipartite tri-ladder. Thus the result holds when n = 6. Now suppose that $n \ge 8$.

If G is a tri-ladder, then G has two vertex-disjoint triangles. Let Y be the set of the vertices of one triangle. Then $\partial(Y)$ is a nontrivial 3-cut of G and $G/\overline{Y} = K_4$. By Lemma 4.6 and the fact that G is a brick, $\partial(Y)$ is good in G. Let G' = G/Y. Then G' is a tri-ladder, which is also a cubic brick. By Lemma 4.8(iii), G' is near-bipartite. By the induction hypothesis, G' has a K_4 -decomposition, so does G.

If G has a K_4 -decomposition, then G has a nontrivial 3-cut, say $\partial(X)$. Let $G_1 = G/\overline{X}$ and $G_2 = G/X$. Then G_i , i = 1, 2, has a K_4 -decomposition, so it is a cubic brick. As above, $\partial(X)$ is good. Lemma 4.8(iii) implies that both G_1 and G_2 are near-bipartite, and \overline{x} and xare incident with edges in removable doubletons of G_1 and G_2 , respectively. If G_i , i = 1 or 2, has at least six vertices, by the induction hypothesis, it is a tri-ladder. Then the contracted vertex, \overline{x} or x, lies in a triangle of G_i . If one of G_1 and G_2 is K_4 , say G_1 , then G_2 has at least six vertices, so it is a tri-ladder. Consequently, G is a tri-ladder, which is a splicing of G_2 and K_4 at vertices in triangles. If each of G_1 and G_2 has at least six vertices, then both of them are tri-ladders. Since a splicing of two tri-ladders at vertices in triangles is still a tri-ladder, G is a tri-ladder. The result holds.

Lemma 4.10. Every near-bipartite tri-ladder has exactly $\frac{n-6}{2}$ removable edges.

Proof. Let G be a near-bipartite tri-ladder with r^* rungs. If n = 6, then $G = \overline{C_6}$, which has no removable edges. The result holds. Now assume that $n \ge 8$. Then $r^* \ge 1$. By Theorem 1.1, G has at least $\frac{n-6}{2}$ removable edges. Note that $2r^* = n - 6$. From the following claim, G has exactly $\frac{n-6}{2}$ removable edges. We are done.

Claim. Every edge e that is not a rung is nonremovable in G.

To show this claim, we present some notions and notations. Let T and T^* be the two disjoint triangles of G with vertex set $\{u_0, v_0, w_0\}$ and $\{u, v, w\}$, respectively. Assume that uand u_0 , v and v_0 , and w and w_0 are ends of three ridges of G, which are referred to as U-ridge, V-ridge and W-ridge, respectively. Let f_i be the rung of G with rank i, $1 \leq i \leq r^*$. We denote by u_i , v_i , or w_i the end of f_i that lies in U-ridge, V-ridge, or W-ridge, respectively. For convenience, we refer to the edges in T as rungs with rank zero. By $u_i u_j$ -subpath we mean the subpath of the U-ridge connecting u_i and u_j . We use similar terminology for subpaths in Vridge and W-ridge. Since G is near-bipartite, G has two edges e_1 and e_2 such that $G - \{e_1, e_2\}$ is bipartite. Thus one of e_1 and e_2 lies in T and the other lies in T^* . Let $H = G - \{e_1, e_2\}$. Colour the vertices in the two color classes of H white and black, respectively. We are now ready to prove the claim. We distinguish two cases according to whether e lies in triangles or not.

Case 1. $e \in E(T \cup T^*)$. Suppose, without loss of generality, that e = uv. Assume that r, s and t are the maximum rank of rungs that has no end in W-ridge, U-ridge and V-ridge, respectively. Thus these three rungs are $u_r v_r, v_s w_s$ and $u_t w_t$. For i = r, s, t, let $d_i = r^* - i$.

Then $d_i \geq 0$. Assume, without loss of generality, that $s \leq t$. Then we consider the neighbour of v in the V-ridge, say y. Thus y is either v_r or v_s . Assume, without loss of generality, that y is white. Then v is black. If u is black, then uv is either e_1 or e_2 , so uv is nonremovable in G. Suppose now that u is white. Recall that $r^* \geq 1$.

First consider the case $y = v_r$ $(r \ge 0)$. Recall that $0 \le d_r \le r^*$. Let U' and W' be the sets of all the black vertices in the uu_r -subpath and the $ww_{r'}$ -subpath, respectively, where r' = r + 1 when $0 < d_r < r^*$ and r' = 0 when $d_r = r^*$ (i.e., r = 0). If $d_r = 0$, let $U' = \{u_r\}$ and $W' = \{w\}$. Let $B' = U' \cup W' \cup \{v_r\}$. If $r \ge 1$, since $u_rv_r \in E(H)$, u_r is black. If r = 0 and d_r is even, since the U-ridge is a path of H, u_0 is black. In both cases, $u_r \in B'$. Let $e' = u_rv_r$. If r = 0 and d_r is odd, u_0 is white but w_0 is black, so $w_0 \in B'$. Let $e' = w_0v_0$. Consequently, B' is a barrier of G - uv, which contains V(e'). By Lemma 2.1(ii), e' is nonadmissible in G - uv, so uv is nonremovable in G.

We now consider the case $y = v_s$ and $s \ge 1$. Then w_s is black and $s \ne t$. Since $s \le t$, we have $r^* = t > s$, so $d_s \ge 1$. Then u_{s+1} is black. Therefore v_s and all the black vertices in the ww_s -subpath and the uu_{s+1} -subpath form a barrier of G - uv, which contains v_s and w_s . As above, uv is nonremovable in G.

Case 2. $e \notin E(T \cup T^*)$. Then e lies in a ridge, say the V-ridge. Let $v_{r^*+1} = v$. Suppose, without loss of generality, that $e = v_j v_k$ $(0 \le j < k \le r^* + 1)$, v_j is white, and the other end of the rung f_j lies in the U-ridge. Then $f_j = u_j v_j$. If j = 0, then $u_0 w_0$ is nonadmissible in $G - v_0 v_k$, so $v_0 v_k$ is nonremovable in G. Suppose now that $j \ge 1$. Then u_j is black. Let w_i be the vertex in the W-ridge with i < j and i as large as possible. Then $f_{i+1} = u_{i+1}v_{i+1}$.

Suppose that w_i is white. When j - i is odd, u_{i+1} is black and v_{i+1} is white. Then v_i is black, because $f_i = w_i v_i$ when $i \ge 1$, and $v_i v_{i+1} \in E(H)$ otherwise. Let B' be the set of the vertex w_i and all the black vertices in the $u_{i+1}u_j$ -subpath and the $v_i v_j$ -subpath. Analogously, when j - i is even, u_{i+1} is white and v_{i+1} is black. Then u_i is black, because $f_i = w_i u_i$ when $i \ge 1$, and $u_i u_{i+1} \in E(H)$ otherwise. Let B' be the set of the vertex w_i and all the black vertices in the $u_i u_j$ -subpath and the $v_{i+1}v_j$ -subpath. In both cases, B' is a barrier of $G - v_j v_k$ containing two ends of $w_i v_i$ or $w_i u_i$, so $v_j v_k$ is nonremovable in G.

Suppose that w_i is black. Let X be the set of all vertices in the u_0u_j -subpath, the v_0v_j subpath and the w_0w_i -subpath. Then $\partial(X)$ is a nontrivial 3-cut, which consists of the edge v_jv_k and two edges incident with u_j and w_i , respectively. Since G is a cubic brick, by Lemma 4.6, $\partial(X)$ is a separating cut but not tight, so it is good in G. Recall that v_j is white, and u_j and w_i are black. Let x_w and x_b be the numbers of white and black vertices in X, respectively, and x_{wb} be the number of edges of G each of which connects a white vertex and a black vertex in X. By counting in two ways, we have $3x_b - 2 \ge x_{wb} \ge 3x_w - 1 - 2$, so $x_b \ge x_w$. By Lemma 4.8(iii), v_jv_k is nonremovable in G. The result holds.

Next, we show that if G is a near-bipartite brick other than K_4 that has exactly $\frac{n-6}{2}$ removable edges, then G is a tri-ladder. Once this is proved. Theorem 1.2 holds. To do this,

we need the following four lemmas.

Lemma 4.11 ([8]). In an essentially 4-edge-connected cubic brick, each edge is either removable or lies in a removable doubleton.

Lemma 4.12 ([14]). Suppose that \mathscr{E} is a set of removable doubletons of an essentially 4edge-connected cubic brick G and $|\mathscr{E}| \geq 2$. Then G can be decomposed into balanced bipartite vertex-induced subgraphs G_i $(i = 1, 2, ..., |\mathscr{E}|)$ satisfying $E_G[V(G_j), V(G_k)]$ is a removable doubleton of G if $|j - k| \equiv 1 \pmod{|\mathscr{E}|}$, and $E_G[V(G_j), V(G_k)] = \emptyset$ otherwise.

Here, a bipartite graph G[A, B] is balanced if |A| = |B|.

Lemma 4.13 ([14]). Suppose that $\{e_1, e'_1\}$ and $\{e_2, e'_2\}$ are removable doubletons of a cubic brick G. If both e_1 and e_2 are incident with v_0 , then e'_1 and e'_2 are adjacent, and $v_0u_0 \in E(G)$, where u_0 is the common end of e'_1 and e'_2 .

Lemma 4.14. Let $\partial(X)$ be a good 3-cut of a cubic matching covered graph G. Assume that $H = G/X \to x$ is an essentially 4-edge-connected cubic near-bipartite brick other than K_4 , and $\{xu, yv\}$ is a removable doubleton of H. Let (A, B) be the bipartition of $H - \{xu, yv\}$ such that $x, u \in A$. Then each vertex of $B \setminus \{y, v\}$ is incident with at least two removable edges of G.

Proof. If $\{xu, yv\}$ is the only removable doubleton of H, by Lemma 4.11, each edge of $E(H) \setminus \{xu, yv\}$ is removable in H. By Lemma 4.3, the result holds. Now assume that H contains s removable doubletons, where $s \geq 2$.

By Lemma 4.12, H can be decomposed into balanced bipartite subgraphs H_1, H_2, \ldots, H_s . Let $A_i = V(H_i) \cap A$ and $B_i = V(H_i) \cap B$, $i = 1, 2, \ldots, s$. Let $x_1 = x$, $u_s = u$, $y_1 = y$, and $v_s = v$. Moreover, we may suppose that $\{u_i y_{i+1}, v_i x_{i+1}\}$, $i = 1, 2, \ldots, s - 1$, are s - 1removable doubletons of H other than $\{x_1 u_s, y_1 v_s\}$, where $\{u_i, x_i\} \subseteq A_i$ and $\{y_i, v_i\} \subseteq B_i$. Then $E_H[V(H_i), V(H_{i+1})] = \{u_i y_{i+1}, v_i x_{i+1}\}$, $i = 1, 2, \ldots, s - 1$. Let w_1 and w_2 be the two neighbors of x_1 other than u_s in H. Since H is a brick, $H - \{w_1, w_2\}$ has a perfect matching, say N_1 . Note that $x_1 u_s \in N_1$ because $d_H(x_1) = 3$. Since $\partial_G(X)$ is a good 3-cut of G, there exists a perfect matching of G, say M, containing each edge of $\partial_G(X)$. Let $M' = (N_1 \setminus \{x_1 u_s\} \cup (M \setminus E(G[\overline{X}]))$. Then M' is also a perfect matching of G. Note that $w_1, w_2 \in B_1 \cup B_2$ and at most one of w_1 and w_2 lies in B_2 . We now consider the following two cases.

Case 1. $w_1, w_2 \in B_1$. Since $|A_1 \setminus \{x_1\}| = |B_1 \setminus \{w_1, w_2\}| + 1$ and $E_H[A_1 \setminus \{x_1\}, V(H) \setminus V(H_1)] = \{u_1y_2\}$, we have $u_1y_2 \in N_1$ and $v_1x_2 \notin N_1$. So $u_1 \neq x_1$. Similarly, if $|s| \geq 3$, we have $u_ty_{t+1} \in N_1$ and $v_tx_{t+1} \notin N_1$, $t = 2, 3, \ldots, s - 1$. Then M' is a perfect matching of $G - (\bigcup_{i=1}^{s-1}\{v_ix_{i+1}\})$ and $\bigcup_{i=1}^{s-1}\{u_iy_{i+1}\} \subseteq M'$. Note that G/\overline{X} is matching covered because $\partial_G(X)$ is a good cut of G. For $i \in \{1, 2, \ldots, s - 1\}$, since $\{u_iy_{i+1}, v_ix_{i+1}\}$ is a removable doubleton of H, $H - \{u_iy_{i+1}, v_ix_{i+1}\}$ is matching covered. By Lemma 4.2, $G - \{u_iy_{i+1}, v_ix_{i+1}\}$ is matching covered. Recall that M' is a perfect matching of $G - v_ix_{i+1}$ and $u_iy_{i+1} \in M'$.

Therefore, $G - v_i x_{i+1}$ is matching covered. Then each edge of $\bigcup_{i=1}^{s-1} \{v_i x_{i+1}\}$ is removable in G. By Lemma 4.11, each edge of $E(H) \setminus (\{x_1 w_1, x_1 w_2, x_1 u_s, y_1 v_s\} \cup (\bigcup_{i=1}^{s-1} \{u_i y_{i+1}, v_i x_{i+1}\}))$ is removable in H, which is also removable in G by Lemma 4.3. This implies that each vertex of $B \setminus \{y_1, v_s\}$ is incident with at least two removable edges of G. The result holds.

Case 2. One of w_1 and w_2 lies in B_2 , say w_2 . Then $w_1 \in B_1$ and $x_1w_2 = u_1y_2$. So $x_1 = u_1$ and $w_2 = y_2$. Since $\{x_1u_s, y_1v_s\}$ and $\{x_1w_2, v_1x_2\}$ are removable doubletons of H, by Lemma 4.13, we have $v_1 = y_1$ and $x_1y_1 \in E(H)$. So $w_1 = y_1 = v_1$. It follows that $|V(H_1)| = 2$ because $E_H[V(H_1), V(H_2)] = \{x_1w_2, y_1x_2\}$ and H is a cubic brick. If |s| = 2, by Lemma 4.11, each edge of $E(H) \setminus \{x_1w_1, x_1w_2, x_1u_2, y_1v_2, y_1x_2\}$ is removable in H, which is also removable in Gby Lemma 4.3. So the result holds. We now consider $|s| \ge 3$. Since $|A_2| = |B_2 \setminus \{w_2\}| + 1$ and $E_H[A_2, V(H) \setminus V(H_2)] = \{x_2y_1, u_2y_3\}$, we have $u_2y_3 \in N_1$ and $v_2x_3 \notin N_1$. Similarly, if $|s| \ge 4$, we have $u_ty_{t+1} \in N_1$ and $v_tx_{t+1} \notin N_1$, $t = 3, 4, \ldots, s - 1$. By the same reason as the above case, we can show that each edge of $\bigcup_{i=2}^{s-1}\{v_ix_{i+1}\}$ is removable in G, and each edge of $E(H) \setminus (\{x_1w_1, x_1w_2, x_1u_s, y_1v_s, y_1x_2\} \cup (\bigcup_{i=2}^{s-1}\{u_iy_{i+1}, v_ix_{i+1}\}))$ is removable in G. So the result holds.

Lemma 4.15. If G is a near-bipartite brick other than K_4 that has exactly $\frac{n-6}{2}$ removable edges, then G is a tri-ladder.

Proof. Let $\{e_1, e_2\}$ be a removable doubleton of G. Since G is a brick, G is 3-connected. Thus, $\delta(G) \geq 3$. By Theorem 1.1, every vertex of G is incident with at most two nonremovable edges and so at least one removable edge, except at most six vertices of degree three contained in two disjoint triangles of G. From the proof of Theorem 1.1 (in the first part of Section 3), we see that these two triangles contain e_1 and e_2 , respectively. Since G has exactly $\frac{n-6}{2}$ removable edges, every vertex of G other than the six vertices of degree three is incident with exactly one removable edge and two nonremovable edges. Thus, G is cubic. Note that $n \geq 6$. If G has a K_4 -decomposition, by Lemma 4.9, G is a tri-ladder. We will show that G has a K_4 -decomposition in the following.

Since G has triangles and $G \neq K_4$, it has nontrivial 3-cuts. Let G^* be any graph obtained by a 3-cut-decomposition of G. Since G is a cubic brick, G^* is 3-connected and cubic. Thus, G^* is an essentially 4-edge-connected cubic graph. Let $\partial(X_1)$ be a nontrivial 3-cut of G such that x_1 is a contraction vertex of G^* . Since G is a brick, $\partial(X_1)$ is good in G. By Lemma 4.8(iii), only one of e_1 and e_2 has both ends in X_1 . This implies that G^* has at most two contraction vertices. By Lemma 4.7, G^* is a brick or a brace. For the latter case, $|V(G^*)| \ge 6$ because G^* is 3-connected and cubic. By Lemma 4.5, every edge of G^* is removable in G^* . Since G^* has at most two contraction vertices, there exists a vertex v of G^* that is incident with at most one contraction vertex of G^* . By Lemma 4.3, v is incident with at least two removable edges in G. This contradiction implies that G^* is a brick.

By Lemma 4.1, each graph generated in the procedure of the 3-cut-decomposition of G is a cubic brick. Thus all the nontrivial 3-cuts used in this procedure are good. By Lemma

4.8(iii), G^* is a near-bipartite brick whose contraction vertices are ends of the edges in its removable doubletons. Then G^* is an essentially 4-edge-connected cubic near-bipartite brick. Recall that x_1 is a contraction vertex of G^* . Let $\{x_1x'_1, x_2x'_2\}$ be a removable doubleton of G^* . Suppose that U' and W' are two color classes of $G^* - x_1x'_1 - x_2x'_2$ such that $x_1, x'_1 \in U'$. Suppose to the contrary that $G^* \neq K_4$. Then $|V(G^*)| \geq 6$. If G^* has exactly one contraction vertex x_1 , by Lemma 4.14, each vertex of $W' \setminus \{x_2, x'_2\}$ is incident with at least two removable edges in G, a contradiction. So G^* has exactly two contraction vertices, one is x_1 , by Lemma 4.8(iii), the other is x_2 or x'_2 , say x_2 . Let $G' = G/X_1$ and let X_2 be a subset of V(G) such that $G^* = G'/X_2$. Then $\partial(X_1)$ and $\partial(X_2)$ are good 3-cut in G and G', respectively, and G'is a cubic near-bipartite brick. By Lemma 4.14, each vertex of $U' \setminus \{x_1, x'_1\}$ is incident with at least two removable edges in G', which are also removable in G by Lemma 4.3, a contradiction. Thus $G^* = K_4$. The result holds.

By Theorem 1.1, Lemmas 4.10 and 4.15, Theorem 1.2 holds.

Acknowledgements

The authors would like to thank the anonymous referee for helpful comments on improving the representation of the paper. This work is supported by the National Natural Science Foundation of China (Nos. 12171440, 12271235 and 12371318), NSF of Fujian (Nos. 2021J01978, 2021J06029) and Fujian Alliance of Mathematics (No. 2023SXLMMS09).

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory, Springer-Verlag, Berlin 2008.
- [2] M.H. Carvalho, C.L. Lucchesi, and U.S.R. Murty, Ear decompositions of matching covered graphs, Combinatorica 19 (1999), 151-174.
- [3] M.H. Carvalho, C.L. Lucchesi, and U.S.R. Murty, Graphs with independent perfect matchings, J. Graph Theory 48 (2005), 19-50.
- [4] M.H. Carvalho, C.L. Lucchesi, and U.S.R. Murty, A generalization of Little's Theorem on Pfaffian orientations, J. Combin. Theory Ser. B 102 (2012), 1241-1266.
- [5] J. Edmonds, L. Lovász, and W.R. Pulleyblank, Brick decompositions and the matching rank of graphs, Combinatorica 2 (1982), 247-274.
- [6] I. Fischer and C.H.C. Little, A characterisation of Pfaffian near bipartite graphs, J. Combin. Theory Ser. B 82 (2001), 175-222.

- J.H. He, E.L. Wei, D. Ye, and S.H. Zhai, On perfect matchings in matching covered graphs, J. Graph Theory 90 (2019), 535-546.
- [8] N. Kothari, M.H. Carvalho, C.L. Lucchesi, and C.H.C. Little, On essentially 4-edgeconnected cubic bricks, Electron. J. Combin. 27 (2020), #P1.22.
- [9] N. Kothari, Generating near-bipartite bricks, J. Graph Theory 90 (2019), 565-590.
- [10] N. Kothari and M.H. Carvalho, Generating simple near-bipartite bricks, J. Graph Theory 95 (2020), 594-637.
- [11] L. Lovász and M.D. Plummer, Matching Theory, Annals of Discrete Mathematics, vol.29, Elsevier Science, 1986.
- [12] L. Lovász, Matching structure and the matching lattice, J. Combin. Theory Ser. B 43 (1987), 187-222.
- [13] L. Lovász and S. Vempala, On removable edges in matching covered graphs, Unpublished manuscript.
- [14] F.L. Lu, X. Feng, and Y. Wang, b-invariant edges in essentially 4-edge-connected nearbipartite cubic bricks, Electron. J. Combin., 27 (2020), #P1.55.
- [15] F.L. Lu and J.G. Qian, Removable edges in cubic matching covered graphs, https://arxiv.org/abs/2202.04279.
- [16] W.T. Tutte, The factorization of linear graphs, J. London Math. Soc. 22 (1947), 107-111.
- [17] S.H. Zhai, C.L. Lucchesi, and X.F. Guo, A lower bound on the number of removable ears of 1-extendable graphs, Discrete Math. 310 (2010), 1123-1126.