

# CONTINUOUS DEPENDENCE FOR MCKEAN-VLASOV SDES UNDER DISTRIBUTION-DEPENDENT LYAPUNOV CONDITIONS

ZHENXIN LIU AND JUN MA

**ABSTRACT.** In this paper, we consider the continuous dependence on initial values and parameters of solutions as well as invariant measures for McKean-Vlasov SDEs under distribution-dependent Lyapunov conditions. In contrast to the classical SDEs, the solutions for McKean-Vlasov SDEs do not converge in probability although the initial values converge in probability, which is due to the mismatch of the distances between measures. Finally, we give some examples to illustrate our theoretical results.

## 1. INTRODUCTION

A signature of structural stability in dynamical systems is the continuous dependence of solutions and invariant measures on initial values and parameters. Investigating this problem enables us to understand deeply the robustness and global bifurcation phenomenon of the dynamical systems in consideration, which, in addition, is of great significance in practical applications. In this paper, we prove the continuous dependence of solutions and invariant measures for McKean-Vlasov SDEs (MVSDEs):

$$(1.1) \quad dX_t = b(t, X_t, \mathcal{L}_{X_t})dt + \sigma(t, X_t, \mathcal{L}_{X_t})dW_t,$$

where  $\mathcal{L}_{X_t}$  denotes the law of  $X_t$ . Due to the dependence of coefficients on distribution, MVSDEs are also known as distribution-dependent SDEs. In addition, MVSDEs are called mean-field SDEs since they are the limit of  $N$ -particle systems if the coefficients satisfy some conditions:

$$dX_{i,t}^N = b(t, X_{i,t}^N, \mu_t^N)dt + \sigma(t, X_{i,t}^N, \mu_t^N)dW_t,$$

for each  $i = 1, 2, \dots, N$ , where  $\mu_t^N := \frac{1}{N} \sum_{j=1}^N \delta_{X_{j,t}^N}$ ; for details, see Sznitman [29, 30] for instance.

With the increasing demands on practical financial markets and social systems, MVSDEs have drawn much attention. For example, the change rate of prices in financial markets may depend on the macrocosmic distribution. Inspired by Kac's work on the Vlasov kinetic equation [19], MVSDEs were first studied by McKean [25] which showed the propagation of chaos in physical systems of  $N$ -interacting particles related to the Boltzmann equation. In order to study large population deterministic and stochastic differential games, Lasry, Lions [20, 21, 22] introduced mean-field games, which were independent of the work of Huang, Malhame and Caines [16, 17]. Owing to their work, MVSDEs are studied more extensively. Wang [32], Ren et al. [28] as well as Liu and Ma [23, 24] obtained the existence and uniqueness under different conditions. Butkovsky [5], Eberle et al. [12], Bogachev et al. [3], Wang [32] as well as Liu and Ma [23, 24] showed ergodicity under different conditions. Buckdahn et al. [4] investigated the relationship between the functional and the associated second-order PDE. Moreover, Cannarsa et al. [6] find solutions by weak KAM approach, which was also utilized by Iturriaga and Wang [18] to study the asymptotic behavior for first-order mean-field games.

---

2010 *Mathematics Subject Classification.* 37H30, 60H10.

*Key words and phrases.* McKean-Vlasov SDEs, distribution-dependent Lyapunov conditions, continuous dependence.

In this paper, we are interested in the continuous dependence of solutions and invariant measures on initial values and parameters for MVSDEs under distribution-dependent Lyapunov conditions. Until now, we find that there are few study about this issue. Bahlali et al. [1] proved the convergence of solutions in  $L^2$  uniformly in  $[0, T]$  under Lipschitz and linear growth conditions, where the linear growth condition is linear in space variable and bounded in distribution variable. Qiao [26] showed the convergence in  $L^2$  of the corresponding solutions for multivalued MVSDEs and the convergence in  $W_1$ -Wasserstein distance of the corresponding invariant measures under monotone and linear growth conditions. Wang [32] proved the continuous dependence of solutions on determinant initial values under monotone and linear growth conditions. Hammersley et al. [15] investigated the continuous dependence on initial values under a Lyapunov condition, which is different from ours. Wu et al. [33] obtained the continuous dependence on initial values for McKean-Vlasov stochastic functional differential equations.

Although there are few study about continuous dependence for MVSDEs, there exist some results for the classical SDEs. In general, there are three ways to illustrate the convergence of the corresponding solutions, i.e. convergence in  $L^2$ , probability and distribution. Friedman [13] showed that the corresponding solutions varied continuously in  $L^2$  if the coefficients converge on any compact set. Gihman and Skorohod [14] studied the convergence in  $L^2$  and probability. Da Prato and Tudor [11] proved the continuous dependence in  $L^2$  uniformly on  $[0, T]$ , probability and distribution for semi-linear stochastic partial differential equations (SPDEs) when the coefficients converge point-wise and initial values converges in the above three ways. Qiu and Wang [27] get the similar results of recurrent solutions. Cheng and Liu [9] illustrated similar results for SPDEs. In summary, the solutions inherit the convergence of initial values if the coefficients converge point-wise or on any compact set. Thus, we want to know whether these results also hold for MVSDEs.

In this paper, we mainly investigate the continuous dependence of solutions and invariant measures for MVSDEs under distribution-dependent Lyapunov conditions. To be specific, we study that  $X_{k,t}$  converges to  $X_t$  in  $W_2$ -Wasserstein distance uniformly on  $[0, T]$  under some suitable conditions (for details see Section 3), where  $X_{k,t}$  and  $X_t$  are solutions of the following MVSDEs, respectively:

$$(1.2) \quad \begin{cases} dX_{k,t} = b_k(t, X_{k,t}, \mathcal{L}_{X_{k,t}})dt + \sigma_k(t, X_{k,t}, \mathcal{L}_{X_{k,t}})dW_t, \\ X_{k,0} = \xi_k, \end{cases}$$

where  $k = 1, 2, \dots$ , and

$$(1.3) \quad \begin{cases} dX_t = b(t, X_t, \mathcal{L}_{X_t})dt + \sigma(t, X_t, \mathcal{L}_{X_t})dW_t, \\ X_0 = \xi. \end{cases}$$

In order to illustrate this result, we utilize Skorokhod's representation theorem and martingale representation theorem. Moreover, the invariant measure of equation (1.2) converges weakly to the invariant measure of equation (1.3) if the coefficients  $b, \sigma, b_k, \sigma_k$  are independent of  $t$ , which is obtained by the same method as above. However, in contrast to classical SDEs, we do not have the result about the convergence in probability when initial values converge in probability, since the coefficients depend on the distribution and are continuous with respect to the distribution variables in  $W_2$ -Wasserstein distance, which gives rise to the mismatch of distance between measures. To illustrate this situation, we give a counterexample in which the coefficients are Lipschitz. Meanwhile, for the completeness of this article, we also show that solutions are convergent in  $L^2$  uniformly on  $[0, T]$ , and invariant measures are convergent in  $W_2$  under Lipschitz conditions when the coefficients converge point-wise.

The rest of this paper is arranged as follows. In Section 2, we collect a number of preliminary results concerning the existence and uniqueness of solutions as well as invariant measures

for MVSDEs and so on. Section 3 presents the continuous dependence of solutions and invariant measures on initial values and parameters under distribution-dependent Lyapunov conditions, and give a counterexample which illustrates that the solution does not inherit its convergence if the initial values converge in probability. In Section 4, we provide some examples to illustrate our theoretical results. In Appendix, we give the proof of continuous dependence under Lipschitz conditions.

## 2. PRELIMINARIES

Throughout the paper, let  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$  be a filtered complete probability space. Assume that the filtration  $\{\mathcal{F}_t\}$  satisfies the usual condition, i.e. it is right continuous and  $\mathcal{F}_0$  contains all  $P$ -null sets. Let  $W$  be an  $n$ -dimensional Brownian motion defined on  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ . We denote by  $A^\top$  the transpose of matrix  $A \in \mathbb{R}^{n \times m}$  with  $n, m \geq 1$ ,  $tr(A)$  the trace of  $A$  and  $|A| := \sqrt{tr(A^\top A)}$  the norm of  $A$ . Assume that the coefficients  $b, b_k : [0, \infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d) \rightarrow \mathbb{R}^d$ ,  $\sigma, \sigma_k : [0, \infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d) \rightarrow \mathbb{R}^{d \times n}$ , and  $X_0, X_{k,0}$  are  $\mathcal{F}_0$ -measurable and satisfy some integrable condition to be specified below.

Let us introduce some basic notations as follows. Let  $\mathcal{P}(\mathbb{R}^d)$  be the space of probability measures on  $\mathbb{R}^d$ . We use the usual Wasserstein distance  $W_p$  on  $\mathcal{P}_p(\mathbb{R}^d)$  with  $p = 1, 2$  in what follows, i.e.

$$W_p(\mu, \nu) := \inf_{\pi \in \mathcal{C}(\mu, \nu)} \left[ \int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y|^p \pi(dx, dy) \right]^{1/p}$$

for  $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^d)$ , where  $\mathcal{P}_p(\mathbb{R}^d) := \{\mu \in \mathcal{P}(\mathbb{R}^d) : \int_{\mathbb{R}^d} |x|^p \mu(dx) < \infty\}$ , and  $\mathcal{C}(\mu, \nu)$  denotes the set of all coupling between  $\mu$  and  $\nu$ . We also define Wasserstein distance  $\bar{W}_2$  on  $\mathcal{P}(C[0, T])$  by

$$\bar{W}_2(\mu, \nu) := \inf_{\pi \in \mathcal{C}(\mu, \nu)} \left[ \int_{C[0, T] \times C[0, T]} |x - y|^2 \pi(dx, dy) \right]^{1/2},$$

for  $\mu, \nu \in \mathcal{P}(C[0, T])$ , where  $\mathcal{P}(C[0, T])$  denotes the set of probability measures on  $C[0, T]$ . As usual, we also denote  $\mu(f) := \int_{\mathbb{R}^d} f(x) \mu(dx)$  in what follows for any function  $f$  defined on  $\mathbb{R}^d$  and  $\mu \in \mathcal{P}(\mathbb{R}^d)$ .

We introduce two propositions which show the existence and uniqueness of solutions as well as invariant measures for MVSDEs under distribution-dependent Lyapunov conditions; for details see [24]. Before giving these results, we introduce some definitions about Lions derivative and differentiable space in Cardaliaguet [7] and Chassagneux et al. [8].

**Definition 2.1** (Lions derivative). A function  $f : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$  is called differentiable at  $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ , denoted by  $\partial_\mu f$ , if there exists a random variable  $X \in L^2(\Omega)$  with  $\mu = \mathcal{L}_X$  such that  $F(X) := f(\mathcal{L}_X)$  and  $F$  is Fréchet differentiable at  $X$ .  $f$  is called differentiable on  $\mathcal{P}_2(\mathbb{R}^d)$  if  $f$  is differentiable at any  $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ .

**Definition 2.2.** The space  $C^{(1,1)}(\mathcal{P}_2(\mathbb{R}^d))$  contains the functions  $f : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$  satisfying the following conditions: (i)  $f$  is differentiable and its derivative  $\partial_\mu f(\mu)(y)$  has a jointly continuous version in  $(\mu, y)$ , still denoted  $\partial_\mu f(\mu)(y)$ ; (ii)  $\partial_\mu f(\mu)(\cdot)$  is continuously differentiable for any  $\mu$ , and its derivative  $\partial_y \partial_\mu f(\mu)(y)$  is jointly continuous at any  $(\mu, y)$ .

(H1) For any  $N \geq 1$ , there exists a constant  $C_N \geq 0$  such that for any  $|x|, |y| \leq N$  and  $\text{supp} \mu, \text{supp} \nu \subset B(0, N)$  we have

$$|b(t, x, \mu)| + |\sigma(t, x, \mu)| \leq C_N,$$

$$|b(t, x, \mu) - b(t, y, \nu)| + |\sigma(t, x, \mu) - \sigma(t, y, \nu)| \leq C_N(|x - y| + W_2(\mu, \nu)).$$

Here  $B(0, N)$  denotes the closed ball in  $\mathbb{R}^d$  centered at the origin with radius  $N$ .

(H2) (Lyapunov condition) There exists a nonnegative function  $V \in C^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d))$  such that there exist nonnegative constants  $\lambda \in \mathbb{R}^+$  satisfying for all  $(t, x, \mu) \in \mathbb{R}^+ \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$

$$\begin{aligned} (\mathcal{L}V)(t, x, \mu) &\leq \lambda V(x, \mu), \\ V_R(\mu) &:= \inf_{|x| \geq R} V(x, \mu) \rightarrow \infty \text{ as } R \rightarrow \infty, \end{aligned}$$

where

$$\begin{aligned} C^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)) &:= \{f : \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \mid f(\cdot, \mu) \in C^2(\mathbb{R}^d) \text{ for } \mu \in \mathcal{P}_2(\mathbb{R}^d), \\ &\quad f(x, \cdot) \in C^{(1,1)}(\mathcal{P}_2(\mathbb{R}^d)) \text{ for } x \in \mathbb{R}^d\}, \end{aligned}$$

and

$$\begin{aligned} (\mathcal{L}V)(t, x, \mu) &:= b(s, x, \mu) \cdot \partial_x V(x, \mu) + \frac{1}{2} \text{tr}((\sigma \sigma^\top)(s, x, \mu) \cdot \partial_x^2 V(x, \mu)) \\ &\quad + \int [b(s, y, \mu) \cdot \partial_\mu V(x, \mu)(y) + \frac{1}{2} \text{tr}((\sigma \sigma^\top)(s, y, \mu) \cdot \partial_y \partial_\mu V(x, \mu)(y))] \mu(dy). \end{aligned}$$

(H2') (Lyapunov condition) There exists a nonnegative function  $V \in C^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d))$  such that there is a constant  $\gamma \geq 0$  satisfying

$$\begin{aligned} (\mathcal{L}V)(x, \mu) &\leq -\gamma, \\ V_R &:= \inf_{|x| \vee |\mu|_2 \geq R} V(x, \mu) \rightarrow \infty \text{ as } R \rightarrow \infty, \end{aligned}$$

for all  $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$ , where  $|\mu|_2^2 := \int_{\mathbb{R}^d} |x|^2 \mu(dx)$ .

(H3) (Continuity) For any bounded sequences  $\{x_n, \mu_n\} \in \mathbb{R}^d \times \mathcal{P}_V(\mathbb{R}^d)$  with  $x_n \rightarrow x$  and  $\mu_n \rightarrow \mu$  weakly in  $\mathcal{P}(\mathbb{R}^d)$  as  $n \rightarrow \infty$ , we have

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} |b(t, x_n, \mu_n) - b(t, x, \mu)| + |\sigma(t, x_n, \mu_n) - \sigma(t, x, \mu)| = 0.$$

where  $\mathcal{P}_V(\mathbb{R}^d) := \{\mu \in \mathcal{P}(\mathbb{R}^d) : \int_{\mathbb{R}^d} V(x, \mu) \mu(dx) < \infty\}$ .

(H4) There exist constants  $\ell > 1, K > 0$  such that

$$|b(t, x, \mu)|^{2\ell} + |\sigma(t, x, \mu)|^{2\ell} \leq K(1 + V(x, \mu)).$$

(H5) There exist constants  $M, \epsilon > 0$  and increasing unbounded function  $L : \mathbb{N} \rightarrow (0, \infty)$  such that for any  $n \geq 1, x, y \in C[0, T], |x(t)| \vee |y(t)| \leq n$  and  $\mu, \nu \in \mathcal{P}(C[0, T])$  satisfying

$$\begin{aligned} &|b(t, x_t, \mu_t) - b(t, y_t, \nu_t)| + |\sigma(t, x_t, \mu_t) - \sigma(t, y_t, \nu_t)| \\ &\leq L_n(|x_t - y_t| + W_{2,n}(\mu_t, \nu_t) + M e^{-\epsilon L_n} (1 \wedge W_2(\mu_t, \nu_t))), \end{aligned}$$

where

$$\begin{aligned} W_{2,n}^2(\mu_t, \nu_t) &:= \inf_{\pi \in \mathcal{C}(\mu, \nu)} \int_{C[0, T] \times C[0, T]} |x_{t \wedge \tau_x^n \wedge \tau_y^n} - y_{t \wedge \tau_x^n \wedge \tau_y^n}|^2 \pi(dx, dy), \\ \tau_x^n &:= \inf\{t \geq 0 : |x_t| \geq n\}, \tau_y^n := \inf\{t \geq 0 : |y_t| \geq n\}. \end{aligned}$$

**Proposition 2.3.** [24, Theorem 3.2] *Assume (H1)–(H4). Then for any  $T > 0, X_0 \in L^2(\Omega, \mathcal{F}, P)$ , equation (1.3) has a solution  $X$  which satisfies*

$$(2.1) \quad EV(X_t, \mathcal{L}_{X_t}) \leq e^{\lambda t} EV(X_0, \mathcal{L}_{X_0}), \quad \text{for } t \in [0, T].$$

Moreover, if (H5) holds, the solution is unique.

**Proposition 2.4.** [24, Theorem 4.3] *Assume that (H1), (H2'), (H3)-(H5) holds with time-independent coefficients, semigroup  $P_t$  is Feller, and that there is a function  $C : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  satisfying  $\sup_{t \in \mathbb{R}^+} C(t) < \infty$  such that*

$$\omega(P_t^* \mu, P_t^* \nu) \leq C(t) \omega(\mu, \nu) \quad \text{for } \mu, \nu \in \mathcal{P}_2(\mathbb{R}^d),$$

where  $\omega$  is Lévy-Prohorov distance and  $P_t^* \mu$  denotes the distribution at time  $t$  with initial distribution  $\mu$ . Then there exists at least one invariant measure. Moreover, if there exists a constant  $t_0 > 0$  such that  $C(t_0) < 1$ , there is a unique invariant measure.

### 3. CONTINUOUS DEPENDENCE

In this section, we consider continuous dependence of solutions and invariant measures on initial values and parameters for MVSDs under distribution-dependent Lyapunov conditions. And we divide this section into three parts.

**3.1. Continuous dependence of solutions.** In this subsection, we prove that solutions vary continuously with respect to initial values and parameters under distribution-dependent Lyapunov conditions, which is obtained by Skorokhod's representation theorem and martingale representation theorem.

**Theorem 3.1.** *Assume that the conditions in Proposition 2.3 hold with  $b, \sigma$  replaced by  $b_k, \sigma_k$ , and*

$$(3.1) \quad \lim_{k \rightarrow \infty} EV(X_{k,0}, \mathcal{L}_{X_{k,0}}) = EV(X_0, \mathcal{L}_{X_0}).$$

Assume further that

$$(3.2) \quad \lim_{k \rightarrow \infty} |b_k(t, x, \mu) - b(t, x, \mu)| + |\sigma_k(t, x, \mu) - \sigma(t, x, \mu)| = 0 \quad \text{point-wise.}$$

Then solutions vary continuously with respect to initial values and parameters. To be precise,

$$X_k \rightarrow X \text{ in } W_2 \text{ on } C[0, T].$$

*Proof.* By Hölder's inequality, BDG's inequality and (H4), there exists a constant  $C(\ell, K) > 0$  such that

$$(3.3) \quad \begin{aligned} & E\left( \sup_{t \in [s, (s+\epsilon) \wedge T]} |X_{k,t} - X_{k,s}|^{2\ell} \right) \\ &= E\left( \sup_{t \in [s, (s+\epsilon) \wedge T]} \left| \int_s^t b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) dr + \int_s^t \sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) dW_r \right|^{2\ell} \right) \\ &\leq C(\ell) E \int_s^{(s+\epsilon) \wedge T} |b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^{2\ell} dr \cdot \epsilon^{2\ell-1} \\ &\quad + C(\ell) E \left( \int_s^{(s+\epsilon) \wedge T} |\sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^2 dr \right)^\ell \\ &\leq C(\ell, K) E \int_s^{(s+\epsilon) \wedge T} 1 + V(X_{k,r}, \mathcal{L}_{X_{k,r}}) dr \cdot \epsilon^{2\ell-1} \\ &\quad + C(\ell) E \left( \int_s^{(s+\epsilon) \wedge T} |\sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^{2\ell} dr \right) \cdot \epsilon^{\ell-1} \\ &\leq C(\ell, K) (1 + e^{\lambda T} EV(X_{k,0}, \mathcal{L}_{X_{k,0}})) \cdot \epsilon^\ell. \end{aligned}$$

Let  $n = \lceil \frac{T}{\epsilon} \rceil + 1$  where  $[m]$  denotes the integer part of  $m \in \mathbb{R}$ . Then we get

$$E\left( \sup_{s, t \in [0, T], |t-s| \leq \epsilon} |X_{k,t} - X_{k,s}|^{2\ell} \right)$$

$$\begin{aligned}
&\leq C(\ell) \sum_{j=1}^n E\left( \sup_{t \in [(j-1)\epsilon, j\epsilon]} |X_{k,t} - X_{k,(j-1)\epsilon}|^{2\ell} \right) \\
&\leq C(\ell, K)(1 + e^{\lambda T} EV(X_{k,0}, \mathcal{L}_{X_{k,0}})) \cdot (T + \epsilon)\epsilon^{\ell-1}.
\end{aligned}$$

Thus, by Ascoli-Arzelà theorem,  $\{\mathcal{L}_{X_k}\}$  is tight on  $C[0, T]$ , i.e. there exists a subsequence, still denoted  $\{\mathcal{L}_{X_k}\}$ , which is weakly convergent to some measure  $\mu \in \mathcal{P}(C[0, T])$ . By Skorokhod's representation theorem, there are  $C[0, T]$ -valued random variables  $\bar{X}_k$  and  $\bar{X}$  on some probability space  $(\bar{\Omega}, \bar{\mathcal{F}}, \bar{P})$  such that  $\mathcal{L}_{\bar{X}_k} = \mathcal{L}_{X_k}$ ,  $\mathcal{L}_{\bar{X}} = \mu$ , and

$$(3.4) \quad \bar{X}_k \rightarrow \bar{X} \text{ a.s.}$$

Note that we have

$$\begin{aligned}
\bar{E}V(\bar{X}_{k,t}, \mathcal{L}_{\bar{X}_{k,t}}) &\leq e^{\lambda t} \bar{E}V(\bar{X}_{k,0}, \mathcal{L}_{\bar{X}_{k,0}}), \\
\bar{E}V(\bar{X}_t, \mathcal{L}_{\bar{X}_t}) &\leq e^{\lambda t} EV(\bar{X}_0, \mathcal{L}_{\bar{X}_0}).
\end{aligned}$$

And by Hölder's inequality, BDG's inequality and (H4), we obtain

$$\begin{aligned}
\bar{E} \sup_{t \in [0, T]} |\bar{X}_{k,t}|^2 &= E \sup_{t \in [0, T]} |X_{k,t}|^2 \\
&\leq 2E\left( \sup_{t \in [0, T]} |X_{k,t} - X_{k,0}|^2 \right) + 2E|X_{k,0}|^2 \\
&= 2E\left( \sup_{t \in [0, T]} \left| \int_0^t b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) dr \right. \right. \\
&\quad \left. \left. + \int_0^t \sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) dW_r \right|^2 \right) + 2E|X_{k,0}|^2 \\
&\leq 4E \int_0^T |b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^2 dr \cdot T \\
&\quad + 4E \left( \int_0^T |\sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^2 dr \right) + 2E|X_{k,0}|^2 \\
&\leq C(K)E \int_0^T 1 + V(X_{k,r}, \mathcal{L}_{X_{k,r}}) dr \cdot T \\
&\quad + 4E \left( \int_0^T |\sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}})|^2 dr \right) + 2E|X_{k,0}|^2 \\
&\leq C(K)(1 + e^{\lambda T} EV(X_{k,0}, \mathcal{L}_{X_{k,0}})) \cdot (T^2 + T) + 2E|X_{k,0}|^2 \\
&< \infty.
\end{aligned}$$

Thus, by limit (3.1) and Lebesgue dominated convergence theorem we get

$$(3.5) \quad \lim_{k \rightarrow \infty} \bar{E} \sup_{t \in [0, T]} |\bar{X}_{k,t} - \bar{X}_t|^2 = 0.$$

If  $\bar{X}$  is a weak solution of equation (1.3), by the weak uniqueness we have  $\mathcal{L}_X = \mathcal{L}_{\bar{X}}$  on  $C[0, T]$ . Therefore we get

$$\lim_{k \rightarrow \infty} \bar{W}_2(\mathcal{L}_{X_k}, \mathcal{L}_X) = \lim_{k \rightarrow \infty} \bar{W}_2(\mathcal{L}_{\bar{X}_k}, \mathcal{L}_{\bar{X}}) \leq \lim_{k \rightarrow \infty} \bar{E} \sup_{t \in [0, T]} |\bar{X}_{k,t} - \bar{X}_t|^2 = 0,$$

and the proof is complete.

Now we prove that  $\bar{X}$  is a weak solution for equation (1.3). Let  $\bar{M}_{k,t} := \bar{X}_{k,t} - \bar{X}_{k,0} - \int_0^t b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) dr$ . By the identical distribution between  $\bar{X}_k$  and  $X_k$ , we have

$$(3.6) \quad \bar{E}[(\bar{M}_{k,t} - \bar{M}_{k,s}) \cdot \Psi(\bar{X}_k|_{[0,s]})]$$

$$\begin{aligned}
&= \bar{E} \left[ \left( \bar{X}_{k,t} - \bar{X}_{k,s} - \int_s^t b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) dr \right) \cdot \Psi(\bar{X}_k|_{[0,s]}) \right] \\
&= 0,
\end{aligned}$$

and

$$\begin{aligned}
(3.7) \quad &\bar{E} \left[ \left( \bar{M}_{k,t} \cdot \bar{M}_{k,t} - \bar{M}_{k,s} \cdot \bar{M}_{k,s} \right. \right. \\
&\quad \left. \left. - \int_s^t (\sigma_k \cdot \sigma_k^\top)(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) dr \right) \cdot \Psi(\bar{X}_k|_{[0,s]}) \right] = 0,
\end{aligned}$$

for any bounded continuous function  $\Psi$  on  $C[0, T]$ . That is to say that  $\bar{M}_k$  is a square-integrable martingale, and its quadratic variation

$$[\bar{M}_k]_t = \int_0^t (\sigma_k \cdot \sigma_k^\top)(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) dr.$$

By the continuity of coefficient  $b_k, b$ , we get

$$\lim_{n \rightarrow \infty} |b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) - b(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r})| = 0 \quad \bar{P}\text{-a.s.}$$

and due to (H4), we have

$$\begin{aligned}
&\bar{E} \int_0^t |b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) - b(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r})|^{2\ell} dr \\
&\leq C(\ell) \bar{E} \int_0^t |b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}})|^{2\ell} + |b(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r})|^{2\ell} dr \\
&\leq C(\ell, K) \int_0^t 1 + \bar{E}V(\bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) + \bar{E}V(\bar{X}_r, \mathcal{L}_{\bar{X}_r}) dr \\
&< \infty.
\end{aligned}$$

Therefore, by Vitali convergence theorem, we obtain

$$\lim_{n \rightarrow \infty} \bar{E} \int_0^t |b_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) - b(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r})|^2 dr = 0.$$

In a similar way, we have

$$\lim_{n \rightarrow \infty} \bar{E} \int_0^t |\sigma_k(r, \bar{X}_{k,r}, \mathcal{L}_{\bar{X}_{k,r}}) - \sigma(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r})|^2 dr = 0.$$

Therefore, by limit (3.5) let  $k \rightarrow \infty$  in equality (3.6) as well as (3.7) and denote  $\bar{M} := \lim_{k \rightarrow \infty} \bar{M}_k$ , we get

$$\begin{aligned}
&\bar{E}[(\bar{M}_t - \bar{M}_s) \cdot \Psi(\bar{X}|_{[0,s]})] = 0, \\
&\bar{E}[(\bar{M}_t \cdot \bar{M}_t - \bar{M}_s \cdot \bar{M}_s - \int_s^t \sigma \cdot \sigma^\top(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r}) dr) \cdot \Psi(\bar{X}|_{[0,s]})] = 0.
\end{aligned}$$

By martingale limit theorem (see [10, Proposition 1.3] for instance),  $\bar{M}$  is a square integrable martingale, and its quadratic variation

$$[\bar{M}]_t = \int_0^t \sigma \cdot \sigma^\top(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r}) dr.$$

Therefore, by martingale representation theorem, there exists a Brownian motion  $\bar{W}$  on the probability space  $(\bar{\Omega}, \bar{\mathcal{F}}, \bar{P})$  such that

$$\bar{M}_t = \int_0^t \sigma(r, \bar{X}_r, \mathcal{L}_{\bar{X}_r}) d\bar{W}_r.$$

Thus, let  $k \rightarrow \infty$  in the following SDE:

$$(3.8) \quad \begin{cases} d\bar{X}_{k,t} = b_k(t, \bar{X}_{k,t}, \mathcal{L}_{\bar{X}_{k,t}})dt + d\bar{M}_{k,t} \\ \bar{X}_{k,0} = \bar{\xi}_k, \end{cases}$$

we obtain that  $(\bar{X}, \bar{W})$  is a weak solution for MVSDE (1.3).  $\square$

**3.2. Continuous dependence of invariant measures.** In this subsection, we investigate the continuous dependence of invariant measures on parameters under distribution-dependent Lyapunov conditions for the following MVSDEs:

$$(3.9) \quad dX_{k,t} = b_k(X_{k,t}, \mathcal{L}_{X_{k,t}})dt + \sigma_k(X_{k,t}, \mathcal{L}_{X_{k,t}})dW_t,$$

where  $k = 1, 2, \dots$ ,

$$(3.10) \quad dX_t = b(X_t, \mathcal{L}_{X_t})dt + \sigma(X_t, \mathcal{L}_{X_t})dW_t,$$

which is obtained by Skorokhod's representation theorem and martingale representation theorem. Moreover, in this subsection, we do not need condition ' $X_{k,0} \rightarrow X_0$  in some sense'.

**Theorem 3.2.** *Assume that the conditions of Proposition 2.4 hold with  $b, \sigma$  replaced by  $b_k, \sigma_k$ . Assume further that*

$$(3.11) \quad \lim_{k \rightarrow \infty} |b_k(x, \mu) - b(x, \mu)| + |\sigma_k(x, \mu) - \sigma(x, \mu)| = 0 \quad \text{point-wise.}$$

Then we have

$$\mu_k \rightarrow \mu$$

where  $\mu_k, \mu$  denote invariant measures for equations (3.9) and (3.10), respectively.

*Proof.* It is obvious that  $\mathcal{L}_{X_k}$  is tight on  $C[0, T]$  by the proof of Theorem 3.1, where  $X_k$  denotes the solution of the following MVSDE:

$$(3.12) \quad \begin{cases} dX_{k,t} = b_k(X_{k,t}, \mathcal{L}_{X_{k,t}})dt + \sigma_k(X_{k,t}, \mathcal{L}_{X_{k,t}})dW_t, \\ X_{k,0} = \xi_k. \end{cases}$$

That is to say that for any  $\epsilon > 0$ , there exists a compact set  $K_\epsilon \subset \mathbb{R}^d$  uniformly on  $[0, T]$ , such that  $\mathcal{L}_{X_{k,t}}(K_\epsilon) > 1 - \epsilon$ . And by Proposition 2.4, we obtain that  $\mathcal{L}_{X_{k,t}}$  weakly converges to  $\mu_k$  as  $t \rightarrow \infty$ . Applying [2, Theorem 2.1], we have

$$\mu_k(K_\epsilon) \geq \limsup_{t \rightarrow \infty} \mathcal{L}_{X_{k,t}}(K_\epsilon) > 1 - \epsilon.$$

Therefore,  $\{\mu_k\}$  is tight. So there is a subsequence, still denoted  $\{\mu_k\}$ , which is weakly convergent to some measure  $\bar{\mu}$ . And similar to the proof of Theorem 3.1, we have  $\bar{\mu} = \mu$ . Therefore, the proof is complete.  $\square$

*Remark 3.3.* By Theorem 3.1, it seems that we could obtain  $\lim_{k \rightarrow \infty} W_2(\mu_k, \mu) = 0$ . However, it may not hold since we can only get  $X_k \rightarrow X$  in  $W_2$  on  $C[0, T]$  for any  $T > 0$ . If we can get the  $X_k \rightarrow X$  in  $W_2$  on  $C[0, \infty)$ , this result directly follows.

**3.3. Counterexample.** For the classical SDEs, if the initial value is convergent in probability, the corresponding solution is convergent in probability uniformly on  $[0, T]$ . However, this result does not hold for MVSDEs since the metric between measures does not match. That is to say that the distance, under which the coefficients are continuous with respect to the distribution variables, does not match the distance between initial values. Therefore, we utilize the property, which is that the convergence in probability does not imply the convergence in  $L^2$ , to give a counterexample to illustrate this situation.



*Example 3.4.* For any  $k = 1, 2, \dots$ , let  $X_0, X_{k,0}$  be random variables satisfying  $X_{k,0} \geq X_0, X_{k,0} \neq X_0$  a.s., and  $X_{k,0} \rightarrow X_0$  in probability but not in  $L^p$  for any  $p > 0$ . And denote  $X_k, X$  by the solutions of the following MVSDs with initial values  $X_{k,0}, X_0$ , respectively:

$$(3.13) \quad dX_{k,t} = \int_{\mathbb{R}^d} y \mathcal{L}_{k,t}(dy) dt + \sqrt{2} dW_t,$$

and

$$(3.14) \quad dX_t = \int_{\mathbb{R}^d} y \mathcal{L}_{X_t}(dy) dt + \sqrt{2} dW_t.$$

Then  $X_k$  does not converge to  $X$  in probability.

*Proof.* Now we construct two sequences of stochastic processes  $\{X_k^{(n)}\}$  and  $\{X^{(n)}\}$  as follows:

$$X_{k,t}^{(0)} = X_{k,0}, X_t^{(0)} = X_0, t \geq 0,$$

for any  $n \geq 1$ ,

$$(3.15) \quad \begin{cases} dX_{k,t}^{(n)} = \int_{\mathbb{R}^d} y \mathcal{L}_{X_{k,t}^{(n-1)}}(dy) dt + \sqrt{2} dW_t, \\ X_{k,0}^{(n)} = X_{k,0}, \end{cases}$$

and

$$(3.16) \quad \begin{cases} dX_k^{(n)} = \int_{\mathbb{R}^d} y \mathcal{L}_{X_t^{(n-1)}}(dy) dt + \sqrt{2} dW_t, \\ X_0^{(n)} = X_0, \end{cases}$$

By Wang [32, Lemma 2.3], we have

$$(3.17) \quad \begin{aligned} \lim_{n \rightarrow \infty} E \sup_{0 \leq t \leq T} |X_{k,t} - X_{k,t}^{(n)}|^2 &= 0, \\ \lim_{n \rightarrow \infty} E \sup_{0 \leq t \leq T} |X_t - X_t^{(n)}|^2 &= 0. \end{aligned}$$

By the definition of  $\{X_k^{(n)}\}$  and  $\{X^{(n)}\}$ , we get

$$\begin{aligned} X_{k,t}^{(1)} - X_t^{(1)} &= X_{k,0} - X_0 + \int_0^t E(X_{k,s}^{(0)} - X_s^{(0)}) ds = X_{k,0} - X_0 + E(X_{k,0} - X_0)t, \\ X_{k,t}^{(2)} - X_t^{(2)} &= X_{k,0} - X_0 + \int_0^t E(X_{k,s}^{(1)} - X_s^{(1)}) ds \\ &= X_{k,0} - X_0 + \int_0^t (E(X_{k,0} - X_0) + E(X_{k,0} - X_0)s) ds \\ &= X_{k,0} - X_0 + E(X_{k,0} - X_0)t + \frac{1}{2} E(X_{k,0} - X_0)t^2, \\ &\dots \\ X_{k,t}^{(n)} - X_t^{(n)} &= X_{k,0} - X_0 + \int_0^t E(X_{k,s}^{(n-1)} - X_s^{(n-1)}) ds \\ &= X_{k,0} - X_0 + E(X_{k,0} - X_0)t + \frac{1}{2} E(X_{k,0} - X_0)t^2 + \dots + \frac{1}{n!} E(X_{k,0} - X_0)t^n. \end{aligned}$$

Note that we have

$$(3.18) \quad P(|X_k^{(n)} - X^{(n)}| > \epsilon) \leq P(|X_k^{(n)} - X_k| > \frac{\epsilon}{3}) + P(|X_k - X| > \frac{\epsilon}{3}) + P(|X - X^{(n)}| > \frac{\epsilon}{3}),$$

for any  $\epsilon > 0$ . Since the convergence in  $L^p$  implies the convergence in probability, by two equalities in the right side of (3.17) we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} P\left(\sup_{0 \leq t \leq T} |X_{k,t}^{(n)} - X_{k,t}| > \frac{\epsilon}{3}\right) &= 0, \\ \lim_{n \rightarrow \infty} P\left(\sup_{0 \leq t \leq T} |X_t - X_t^{(n)}| > \frac{\epsilon}{3}\right) &= 0. \end{aligned}$$

Let  $n \rightarrow \infty$  then  $k \rightarrow \infty$  in inequality (3.18), we get

$$\lim_{k \rightarrow \infty} P(|X_{k,t} - X_t| > \frac{\epsilon}{3}) \geq \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} P(|X_{k,t}^{(n)} - X_t^{(n)}| > \epsilon) = 1,$$

where the last equality holds for some sufficiently small constant  $\epsilon > 0$  and  $t \neq 0$ . The proof is complete.  $\square$

However convergence in probability for initial values does not imply that the corresponding solutions converge in probability under Lipschitz condition by the above counterexample, we obtain that solutions inherit the  $L^2$ -convergence of initial values, which is showed by the following proposition. And we think that the proof is simple, so we put it in Appendix.

**Proposition 3.5.** *Suppose that the coefficients  $b, \sigma, b_k, \sigma_k$  of equations (1.3) and (1.2) are Lipschitz with Lipschitz constant  $L$ , and satisfy the linear growth conditions with linear growth constant  $K$  for all  $k = 1, 2, \dots$ . Assume further that*

$$(3.19) \quad \lim_{k \rightarrow \infty} E(|X_{k,0} - X_0|^2) = 0,$$

and

$$(3.20) \quad \lim_{k \rightarrow \infty} |b_k(t, x, \mu) - b(t, x, \mu)| + |\sigma_k(t, x, \mu) - \sigma(t, x, \mu)| = 0 \quad \text{point-wise.}$$

Then we have

$$(3.21) \quad \lim_{k \rightarrow \infty} E \sup_{0 \leq t \leq T} |X_{k,t} - X_t|^2 = 0.$$

*Remark 3.6.* By the above Proposition, we directly obtain [26, Proposition 5.2] and [1, Theorem 5.1 and Theorem 6.1].

As a direct consequence of Proposition 3.5, we have the following corollary.

**Corollary 3.7.** *Assume that the conditions of Proposition 3.5 hold with (3.19) replaced by*

$$\lim_{k \rightarrow \infty} W_2(\mathcal{L}_{X_{k,0}}, \mathcal{L}_{X_0}) = 0.$$

Then we have

$$\lim_{k \rightarrow \infty} \bar{W}_2(\mathcal{L}_{X_k}, \mathcal{L}_X) = 0.$$

*Proof.* For each  $k \geq 1$ , there exist random variables  $\tilde{X}_{k,0}, \tilde{X}_0^{(k)}$  such that  $\mathcal{L}_{X_{k,0}} = \mathcal{L}_{\tilde{X}_{k,0}}$ ,  $\mathcal{L}_{X_0} = \mathcal{L}_{\tilde{X}_0^{(k)}}$  and

$$W_2(\mathcal{L}_{X_{k,0}}, \mathcal{L}_{X_0})^2 = E|\tilde{X}_{k,0} - \tilde{X}_0^{(k)}|^2.$$

Let  $\tilde{X}_{k,t}, \tilde{X}_t^{(k)}$  be the solutions of MVSEs (1.3) and (1.2) with initial value  $\tilde{X}_{k,0}, \tilde{X}_0^{(k)}$  respectively. By Proposition 3.5, we have

$$\lim_{k \rightarrow \infty} E \sup_{0 \leq s \leq T} |\tilde{X}_{k,s} - \tilde{X}_s^{(k)}|^2 = 0.$$

Therefore, by the weak uniqueness we obtain

$$\bar{W}_2(\mathcal{L}_{X_k}, \mathcal{L}_X)^2 = \bar{W}_2(\mathcal{L}_{\tilde{X}_k}, \mathcal{L}_{\tilde{X}^{(k)}})^2 \leq E \sup_{0 \leq t \leq T} |\tilde{X}_{k,t} - \tilde{X}_t^{(k)}|^2 \rightarrow 0.$$

The proof is complete.  $\square$

*Remark 3.8.* We know that the optimal coupling is attained, but the optimal one depend on  $k$  and we could not find a random variable  $\tilde{X}_0$  such that  $W_2(\mathcal{L}_{X_{k,0}}, \mathcal{L}_{X_0})^2 = E|\tilde{X}_{k,0} - \tilde{X}_0|^2$  holds for every  $k = 1, 2, \dots$

*Example 3.9.* (Small ‘distribution’ limit). In particular, the random trajectory of the MVSDE

$$\begin{cases} dX_t^\epsilon = b_\epsilon(X_t^\epsilon, \mathcal{L}_{X_t^\epsilon})dt + \sigma_\epsilon(X_t^\epsilon, \mathcal{L}_{X_t^\epsilon})dW_t \\ X_0^\epsilon = \xi \end{cases}$$

converge in  $L^2$  uniformly on  $[0, T]$  as  $\epsilon \rightarrow 0$  to the trajectory of

$$\begin{cases} dX_t = b(X_t)dt + \sigma(X_t)dW_t \\ X_0 = \xi, \end{cases}$$

where  $\lim_{\epsilon \rightarrow 0} b_\epsilon(x, \mu) = b(x)$ ,  $\lim_{\epsilon \rightarrow 0} \sigma_\epsilon(x, \mu) = \sigma(x)$  for each  $x \in \mathbb{R}^d$ ,  $\mu \in \mathcal{P}(\mathbb{R}^d)$ .

We next give the continuous dependence of invariant measures on parameters without the dependence on initial values. The proof is also put in Appendix.

**Proposition 3.10.** *Suppose that the conditions of Proposition 3.5 hold except for the condition (3.19), and for the Lipschitz conditions of coefficients  $b, b_k$  replaced by the following conditions: there exist nonnegative constants  $L_1, L_2$  with  $2L_1 - 2L_2 - 8L^2 > 0$  such that*

$$\langle x - y, b(x, \mu) - b(y, \nu) \rangle \leq -L_1|x - y|^2 + L_2|x - y|W_2(\mu, \nu),$$

for any  $x, y \in \mathbb{R}^d$ ,  $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ . Assume further that the coefficients  $b, b_k, \sigma, \sigma_k$  are independent of  $t$ . Then we have

$$(3.22) \quad \lim_{k \rightarrow \infty} W_2(\mu_k, \mu) = 0,$$

where  $\mu_k, \mu$  are the unique invariant measures of MVSDs (3.9) and (3.10), respectively.

#### 4. APPLICATION

In this section, we give some examples to illustrate our theoretical results.

*Example 4.1.* Consider the McKean-Vlasov SDEs

$$\begin{aligned} dX_{\lambda,t} &= -\lambda X_{\lambda,t} \int_{\mathbb{R}} y^2 \mathcal{L}_{X_{\lambda,t}}(dy)dt + (\sqrt{2} + \lambda)X_{\lambda,t}dW_t \\ &=: b_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dt + \sigma_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dW_t, \end{aligned}$$

where  $\lambda \in [0, 1]$  is a parameter and  $W$  is a one-dimensional Brownian motion. Then the solution  $X_{\lambda,t}$  with initial value  $X_{\lambda,0}$  for the above MVSDs converges to  $X_t$ , which is the solution of the following SDEs with initial value  $X_0$ , in  $W_2$  uniformly on  $[0, T]$

$$\begin{aligned} dX_t &= -X_t \int_{\mathbb{R}} y^2 \mathcal{L}_{X_t}(dy)dt + (\sqrt{2} + 1)X_t dW_t \\ &= b(X_t, \mathcal{L}_{X_t})dt + \sigma(X_t, \mathcal{L}_{X_t})dW_t. \end{aligned}$$

*Proof.* (H1), (H3) and (H5) obviously hold. Now we prove that (H2) and (H4) hold. For any  $x \in \mathbb{R}$ ,  $\mu \in \mathcal{P}(\mathbb{R})$ , let  $V(x, \mu) = x^4 + \int_{\mathbb{R}} y^6 \mu(dy)$ . Then (H4) holds and

$$\begin{aligned} \partial_x V(x, \mu) &= 4x^3, \quad \partial_x^2 V(x, \mu) = 12x^2, \\ \partial_\mu V(x, \mu)(z) &= 6z^5, \quad \partial_z \partial_\mu V(x, \mu)(z) = 30z^4. \end{aligned}$$

Thus, we have

$$\begin{aligned} (\mathcal{L}V)(x, \mu) &= -\lambda x \int y^2 \mu(dy) \cdot 4x^3 + \frac{1}{2} \cdot (\sqrt{2} + \lambda)^2 x^2 \cdot 12x^2 \\ &\quad + \int \left( (-\lambda z \int y^2 \mu(dy)) \cdot 6z^5 + \frac{1}{2} \cdot (\sqrt{2} + \lambda)^2 z^2 \cdot 30z^4 \right) \mu(dz) \\ &\leq 36x^4 + \int 90z^6 \mu(dz) \leq 90V(x, \mu), \end{aligned}$$

i.e. (H2) holds. Therefore, by Theorem 3.1, we get the desired result.  $\square$

*Example 4.2.* For any  $x \in \mathbb{R}, \mu \in \mathcal{P}(\mathbb{R})$ , let

$$\begin{aligned} b_\lambda(x, \mu) &= -3\lambda x + 3\lambda \int_{\mathbb{R}} y \mu(dy), \quad \sigma_\lambda(x, \mu) = x - \int_{\mathbb{R}} y \mu(dy), \\ b(x, \mu) &= -3x + 3 \int_{\mathbb{R}} y \mu(dy), \quad \sigma(x, \mu) = x - \int_{\mathbb{R}} y \mu(dy), \end{aligned}$$

where  $\lambda$  is a parameter in  $[1, 2]$ . Then  $\mu_\lambda \rightarrow \mu$  weakly as  $\lambda \rightarrow 1$ , where  $\mu_\lambda$  and  $\mu$  are invariant measures of following MVSDs:

$$\begin{aligned} dX_{\lambda,t} &= b_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dt + \sigma_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dW_t, \\ dX_t &= b(X_t, \mathcal{L}_{X_t})dt + \sigma(X_t, \mathcal{L}_{X_t})dW_t. \end{aligned}$$

*Proof.* Conditions (H1), (H3) and (H5) obviously hold. Let  $V(x, \mu) = \frac{1}{4}(x - \int_{\mathbb{R}} y \mu(dy))^4$ . Then it is immediately to see that (H4) holds. Note that we have

$$\begin{aligned} \partial_x V(x, \mu) &= (x - \int y \mu(dy))^3, \quad \partial_x^2 V(x, \mu) = 3(x - \int y \mu(dy))^2, \\ \partial_\mu V(x, \mu)(z) &= - (x - \int y \mu(dy))^3, \quad \partial_z \partial_\mu V(x, \mu)(z) = 0, \end{aligned}$$

and

$$\begin{aligned} (\mathcal{L}V)(x, \mu) &= -3\lambda(x - \int y \mu(dy))^4 + \frac{3}{2}(x - \int y \mu(dy))^4 \\ &\quad + \int 3(z - \int y \mu(dy))(x - \int y \mu(dy))^3 \mu(dz) \\ &= (-3\lambda + \frac{3}{2})(x - \int y \mu(dy))^4 \leq -\frac{3}{2}V(x, \mu). \end{aligned}$$

That is to say that (H2) holds. Therefore, by Theorem 3.2, we obtain the desired results.  $\square$

*Remark 4.3.* It is easy to verify that the coefficients  $b_\lambda, \sigma_\lambda, b, \sigma$  do not satisfy the conditions of Proposition 3.10, so here we apply Theorem 3.2 to illustrate the result.

*Example 4.4.* Consider the McKean-Vlasov SDE

$$\begin{aligned} dX_{\lambda,t} &= ((-6 + \lambda)X_{\lambda,t} + \int_{\mathbb{R}} y \mathcal{L}_{X_{\lambda,t}}(dy))dt + (X_{\lambda,t} + \lambda \int_{\mathbb{R}} y \mathcal{L}_{X_{\lambda,t}}(dy))dW(t) \\ &=: b_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dt + \sigma_\lambda(X_{\lambda,t}, \mathcal{L}_{X_{\lambda,t}})dW_t, \end{aligned}$$

where  $\lambda \in [0, 1)$  is a parameter and  $W$  is a one-dimensional Brownian motion. Clearly  $b_\lambda$  and  $\sigma_\lambda$  are Lipschitz and satisfy linear growth condition. We note that for any  $x \in \mathbb{R}, \mu \in \mathcal{P}(\mathbb{R})$ ,

$$\lim_{\lambda \rightarrow \lambda_0} |b_\lambda(x, \mu) - b_{\lambda_0}(x, \mu)| + |\sigma_\lambda(x, \mu) - \sigma_{\lambda_0}(x, \mu)| = 0.$$

Then, by Proposition 3.5 and Corollary 3.7, the unique solution  $X_\lambda$  converges to  $X_{\lambda_0}$  in  $L^2$  uniformly on  $[0, T]$ , and the corresponding measure  $\mathcal{L}_{X_\lambda}$  converges to  $\mathcal{L}_{X_{\lambda_0}}$  under  $\bar{W}_2$ .

Moreover, we have

$$\begin{aligned} \langle x - y, b_\lambda(x, \mu) - b_\lambda(y, \nu) \rangle &= \langle x - y, (-6 + \lambda)(x - y) + \int z\mu(dz) - \int z\nu(dz) \rangle \\ &\leq (-6 + \lambda)|x - y|^2 + |x - y|W_2(\mu, \nu), \\ |\sigma_\lambda(x, \mu) - \sigma_\lambda(y, \nu)| &= |x - y + \lambda \int z\mu(dz) - \lambda \int z\nu(dz)| \\ &\leq |x - y| + \lambda W_2(\mu, \nu). \end{aligned}$$

Therefore,  $L_1 = 6 - \lambda$ ,  $L_2 = L = 1$ , and  $2L_1 - 2L_2 - 8L^2 = 2(6 - \lambda) - 2 - 8 = 2(1 - \lambda) > 0$ . Applying Proposition 3.10, we obtain that the uniqueness invariant measure  $\mu_{\lambda, I}$  converges to  $\mu_{\lambda_0, I}$  under  $W_2$ .

#### APPENDIX

In this section, we give the proof of Proposition 3.5 and Proposition 3.10.

*Proof of Proposition 3.5.* Note that we have

$$\begin{aligned} X_{k,t} - X_t &= X_{k,0} - X_0 + \int_0^t b_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - b(s, X_s, \mathcal{L}_{X_s}) ds \\ &\quad + \int_0^t \sigma_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - \sigma(s, X_s, \mathcal{L}_{X_s}) dW_s \\ &= X_{k,0} - X_0 + \int_0^t b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s}) ds \\ &\quad + \int_0^t \sigma_k(s, X_s, \mathcal{L}_{X_s}) - \sigma(s, X_s, \mathcal{L}_{X_s}) dW_s \\ &\quad + \int_0^t b_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - b_k(s, X_s, \mathcal{L}_{X_s}) ds \\ &\quad + \int_0^t \sigma_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - \sigma_k(s, X_s, \mathcal{L}_{X_s}) dW_s \\ &=: \eta_{k,t} + \int_0^t b_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - b_k(s, X_s, \mathcal{L}_{X_s}) ds \\ &\quad + \int_0^t \sigma_k(s, X_{k,s}, \mathcal{L}_{X_{k,s}}) - \sigma_k(s, X_s, \mathcal{L}_{X_s}) dW_s, \end{aligned}$$

where

$$\begin{aligned} \eta_{k,t} &:= X_{k,0} - X_0 + \int_0^t b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s}) ds \\ &\quad + \int_0^t \sigma_k(s, X_s, \mathcal{L}_{X_s}) - \sigma(s, X_s, \mathcal{L}_{X_s}) dW_s. \end{aligned}$$

By Cauchy-Schwarz inequality and martingale inequality, we get

$$\begin{aligned} &E\left(\sup_{0 \leq s \leq t} |X_{k,s} - X_s|^2\right) \\ &\leq 3E\left(\sup_{0 \leq s \leq t} |\eta_{k,s}|^2\right) + 3E\left(\sup_{0 \leq s \leq t} \left| \int_0^s b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) - b_k(r, X_r, \mathcal{L}_{X_r}) dr \right|^2\right) \\ &\quad + 3E\left(\sup_{0 \leq s \leq t} \left| \int_0^s \sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) - \sigma_k(r, X_r, \mathcal{L}_{X_r}) dW_r \right|^2\right) \end{aligned}$$

$$\begin{aligned}
&\leq 3E\left(\sup_{0\leq s\leq t} |\eta_{k,s}|^2\right) + 3TE\left(\int_0^t |b_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) - b_k(r, X_r, \mathcal{L}_{X_r})|^2 dr\right) \\
&\quad + 3 \times 4E\left(\int_0^t |\sigma_k(r, X_{k,r}, \mathcal{L}_{X_{k,r}}) - \sigma_k(r, X_r, \mathcal{L}_{X_r})|^2 dr\right) \\
&\leq 3E\left(\sup_{0\leq s\leq t} |\eta_{k,s}|^2\right) + 3TE\int_0^t 2L^2(|X_{k,r} - X_r|^2 + W_2(\mathcal{L}_{X_{k,r}}, \mathcal{L}_{X_r})^2) dr \\
&\quad + 3 \times 4E\int_0^t 2L^2(|X_{k,r} - X_r|^2 + W_2(\mathcal{L}_{X_{k,r}}, \mathcal{L}_{k,r})^2) dr \\
&\leq 3E\left(\sup_{0\leq s\leq t} |\eta_{k,s}|^2\right) + 2(6TL^2 + 24L^2)\int_0^t E\left(\sup_{0\leq s\leq r} |X_{k,s} - X_s|^2\right) dr,
\end{aligned}$$

for any  $t \in [0, T]$ . Applying Gronwall's inequality, we obtain

$$E\sup_{0\leq s\leq t} |X_{k,s} - X_s|^2 \leq 3E\left(\sup_{0\leq s\leq t} |\eta_{k,s}|^2\right)e^{2(6TL^2+24L^2)t}.$$

So (3.21) holds if we can prove

$$\lim_{t\rightarrow\infty} E\left(\sup_{0\leq t\leq T} |\eta_{k,t}|^2\right) = 0.$$

By Cauchy-Schwarz inequality and martingale inequality, we obtain

$$\begin{aligned}
E\left(\sup_{0\leq t\leq T} |\eta_{k,t}|^2\right) &\leq 3E|X_{k,0} - X_0|^2 \\
&\quad + 3E\left(\sup_{0\leq t\leq T} \left|\int_0^t b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s}) ds\right|^2\right) \\
&\quad + 3E\left(\sup_{0\leq t\leq T} \left|\int_0^t \sigma_k(s, X_s, \mathcal{L}_{X_s}) - \sigma(s, X_s, \mathcal{L}_{X_s}) dW_s\right|^2\right) \\
&\leq 3E|X_{k,0} - X_0|^2 \\
&\quad + 3TE\int_0^T |b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s})|^2 ds \\
&\quad + 3 \times 4E\int_0^T |\sigma_k(s, X_s, \mathcal{L}_{X_s}) - \sigma(s, X_s, \mathcal{L}_{X_s})|^2 ds.
\end{aligned}$$

We note that

$$\begin{aligned}
|b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s})|^2 &\leq 2|b_k(s, X_s, \mathcal{L}_{X_s})|^2 + 2|b(s, X_s, \mathcal{L}_{X_s})|^2 \\
&\leq 12K^2(1 + |X_s|^2 + \mathcal{L}_{X_s}(|\cdot|^2)),
\end{aligned}$$

and  $E\sup_{0\leq s\leq T} |X_s|^2 < \infty$  by [32, Theorem 2.1] or [23, Theorem 3.1]. Applying Lebesgue dominated convergence theorem, we get

$$\lim_{k\rightarrow\infty} E\left|\int_0^T b_k(s, X_s, \mathcal{L}_{X_s}) - b(s, X_s, \mathcal{L}_{X_s}) ds\right|^2 = 0.$$

In the same way, we obtain

$$\lim_{k\rightarrow\infty} E\left|\int_0^T \sigma_k(s, X_s, \mathcal{L}_{X_s}) - \sigma(s, X_s, \mathcal{L}_{X_s}) dW_s\right|^2 = 0.$$

Now  $\lim_{k\rightarrow\infty} E\sup_{0\leq t\leq T} |\eta_{k,t}|^2 = 0$  holds and hence (3.21) is proved.  $\square$

*Proof of Proposition 3.10.* By [32, Theorem 3.1], there exist invariant measures for MVSDs (3.9) and (3.10), denoted by  $\mu_k, \mu$ , respectively. Let  $X_{k,0}, X_0$  be random variables such that

$$\lim_{k \rightarrow \infty} E|X_{k,0} - X_0|^2 = 0,$$

Denote  $X_{k,t}, X_t$  by the corresponding solutions with initial values  $X_{k,0}, X_0$ , respectively. By Itô's formula, we have

$$\begin{aligned} E|X_{k,t} - X_t|^2 &= E|X_{k,0} - X_0|^2 \\ &\quad + 2E \int_0^t \langle X_{k,s} - X_s, b_k(X_{k,s}, \mathcal{L}_{X_{k,s}}) - b(X_s, \mathcal{L}_{X_s}) \rangle ds \\ &\quad + E \int_0^t |\sigma_k(X_{k,s}, \mathcal{L}_{X_{k,s}}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq E|X_{k,0} - X_0|^2 \\ &\quad + 2E \int_0^t \langle X_{k,s} - X_s, b_k(X_{k,s}, \mathcal{L}_{X_{k,s}}) - b_k(X_s, \mathcal{L}_{X_s}) \rangle ds \\ &\quad + 2E \int_0^t \langle X_{k,s} - X_s, b_k(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s}) \rangle ds \\ &\quad + 2E \int_0^t |\sigma_k(X_{k,s}, \mathcal{L}_{X_{k,s}}) - \sigma_k(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\quad + 2E \int_0^t |\sigma_k(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq E|X_{k,0} - X_0|^2 \\ &\quad + E \int_0^t -2L_1|X_{k,s} - X_s|^2 + 2L_2|X_{k,s} - X_s|W_2(\mathcal{L}_{X_{k,s}}, \mathcal{L}_{X_s}) ds \\ &\quad + 4L^2 E \int_0^t |X_{k,s} - X_s|^2 + W_2(\mathcal{L}_{X_{k,s}}, \mathcal{L}_{X_s})^2 ds \\ &\quad + 2E \int_0^t |X_{k,s} - X_s| \cdot |b_k(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s})| ds \\ &\quad + 2E \int_0^t |\sigma_k(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq E|X_{k,0} - X_0|^2 + \int_0^t (-2L_1 + 2L_2 + 8L^2 + \epsilon) E|X_{k,s} - X_s|^2 ds \\ &\quad + \frac{1}{\epsilon} E \int_0^t |b_k(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\quad + 2E \int_0^t |\sigma_k(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds, \end{aligned}$$

where  $0 < \epsilon < 2L_1 - 2L_2 - 8L^2$ , and the last inequality holds owing to Cauchy-Schwarz inequality. By Gronwall's inequality, we have

$$(4.1) \quad E|X_{k,t} - X_t|^2 \leq e^{(-2L_1 + 2L_2 + 8L^2 + \epsilon)t} E\eta_{k,t},$$

where

$$\eta_{k,t} := |X_{k,0} - X_0|^2 + \frac{1}{\epsilon} \int_0^t |b_k(X_s, \mathcal{L}_{X_s}) - b(X_s, \mathcal{L}_{X_s})|^2 ds$$

$$+ 2 \int_0^t |\sigma_k(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds.$$

Applying [31, Lemma 4.3] we obtain

$$W_2(\mu_k, \mu) \leq \liminf_{t \rightarrow \infty} W_2(\mathcal{L}_{X_{k,t}}, \mathcal{L}_{X_t}) \leq \liminf_{t \rightarrow \infty} (E|X_{k,t} - X_t|^2)^{\frac{1}{2}}.$$

If we can prove  $\lim_{k \rightarrow \infty} \lim_{t \rightarrow \infty} e^{(-2L_1+2L_2+8L^2+\epsilon)t} E\eta_{k,t} = 0$ , (3.22) holds.

By conditions on coefficients  $b, \sigma$  with  $y = 0, \nu = \delta_0$ , we get

$$\begin{aligned} \langle x, b(x, \mu) \rangle &= \langle x, b(x, \mu) - b(0, \delta_0) \rangle + \langle x, b(0, \delta_0) \rangle \\ &\leq -L_1|x|^2 + L_2x \cdot (\mu(|\cdot|^2))^{\frac{1}{2}} + |b(0, \delta_0)| \cdot |x|, \\ |\sigma(x, \mu)|^2 &\leq 2|\sigma(x, \mu) - \sigma(0, \delta_0)|^2 + 2|\sigma(0, \delta_0)|^2 \\ &\leq 4L^2(|x|^2 + \mu(|\cdot|^2)) + 2|\sigma(0, \delta_0)|^2. \end{aligned}$$

Applying Itô's formula and Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} E|X_t|^2 &= E|X_0|^2 + 2E \int_0^t \langle X_s, b(X_s, \mathcal{L}_{X_s}) \rangle ds + E \int_0^t |\sigma(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq E|X_0|^2 + E \int_0^t (-2L_1 + 2L_2 + 8L^2 + \epsilon)|X_s|^2 ds + \left(\frac{1}{\epsilon}|b(0, \delta_0)|^2 + 2|\sigma(0, \delta_0)|^2\right)t. \end{aligned}$$

By Gronwall's inequality, we have

$$E|X_t|^2 \leq (E|X_0|^2 + \left(\frac{1}{\epsilon}|b(0, \delta_0)|^2 + 2|\sigma(0, \delta_0)|^2\right)t) e^{(-2L_1+2L_2+8L^2+\epsilon)t}.$$

Thus, by linear growth condition of coefficients  $b_k$  and  $b$  we get

$$\begin{aligned} &E \int_0^t |b_k(X_s, \mathcal{L}_{X_s}) - b(X(s), \mathcal{L}_{X(s)})|^2 ds \\ &\leq 2E \int_0^t |b_k(X_s, \mathcal{L}_{X_s})|^2 + |b(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq 2E \int_0^t 6K^2(1 + |X_s|^2 + \mathcal{L}_{X_s}(|\cdot|^2)) ds \\ &\leq 12K^2t + 24K^2 \int_0^t (E|X_0|^2 + \left(\frac{1}{\epsilon}|b(0, \delta_0)|^2 + 2|\sigma(0, \delta_0)|^2\right)s) e^{(-2L_1+2L_2+8L^2+\epsilon)s} ds. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} &E \int_0^t |\sigma_k(X_s, \mathcal{L}_{X_s}) - \sigma(X_s, \mathcal{L}_{X_s})|^2 ds \\ &\leq 12K^2t + 24K^2 \int_0^t (E|X_0|^2 + \left(\frac{1}{\epsilon}|b(0, \delta_0)|^2 + 2|\sigma(0, \delta_0)|^2\right)s) e^{(-2L_1+2L_2+8L^2+\epsilon)s} ds. \end{aligned}$$

Thus, we get  $\lim_{k \rightarrow \infty} \lim_{t \rightarrow \infty} e^{(-2L_1+2L_2+8L^2+\epsilon)t} E\eta_{k,t} = 0$ . The proof is complete.  $\square$

#### ACKNOWLEDGEMENT

The first author is supported by National Key R&D Program of China (No. 2023YFA1009200), NSFC (Grants 11871132, 11925102), LiaoNing Revitalization Talents Program (Grant XLYC2202042), and Dalian High-level Talent Innovation Program (Grant 2020RD09). The second author is supported by National Funded Postdoctoral Researcher Program (Grant GZC20230414).



## REFERENCES

- [1] K. Bahlali, M. A. Mezerdi and B. Mezerdi, Stability of McKean-Vlasov stochastic differential equations and applications, *Stoch. Dyn.* **20** (2020), 19 pp.
- [2] P. Billingsley, *Convergence of Probability Measures*, John Wiley & Sons, Inc., New York-London-Sydney, 1968, xii+253 pp.
- [3] V. I. Bogachev, M. Röckner and S. V. Shaposhnikov, Convergence in variation of solutions of nonlinear Fokker-Planck-Kolmogorov equations to stationary measures, *J. Funct. Anal.* **276** (2019), 3681–3713.
- [4] R. Buckdahn, J. Li, S. Peng and C. Rainer, Mean-field stochastic differential equations and associated PDEs, *Ann. Probab.* **45** (2017), 824–878.
- [5] O. A. Butkovsky, On ergodic properties of nonlinear Markov chains and stochastic McKean-Vlasov equations, *Theory Probab. Appl.* **58** (2014), 661–674.
- [6] P. Cannarsa, W. Cheng, C. Cristian and K. Wang, Weak KAM approach to first-order mean field games with state constraints, *J. Dynam. Differential Equations* **35** (2023), 1885–1916.
- [7] P. Cardaliaguet, Notes on mean field games. Notes from P.L. Lions’ lectures at the Collège de France, <https://www.ceremade.dauphine.fr/~cardalia/MFG100629.pdf>, 2012.
- [8] J. F. Chassagneux, D. Crisan and F. Delarue, A probabilistic approach to classical solutions of the master equation for large population equilibria, *Mem. Amer. Math. Soc.* **280** (2022), v+123 pp.
- [9] M. Cheng and Z. Liu, Periodic, almost periodic and almost automorphic solutions for SPDEs with monotone coefficients. *Discrete Contin. Dyn. Syst. Ser. B* **26** (2021), 6425–6462.
- [10] K. L. Chung and R. J. Williams, *Introduction to Stochastic Integration*, Birkhäuser/Springer, New York, 2014, xviii+276 pp.
- [11] G. Da Prato and C. Tudor, Periodic and almost periodic solutions for semilinear stochastic equations. *Stochastic Anal. Appl.* **1** (1995), 13–33.
- [12] A. Eberle, A. Guillin and R. Zimmer, Quantitative Harris-type theorems for diffusions and McKean-Vlasov processes. *Trans. Amer. Math. Soc.* **10** (2019), 7135–7173.
- [13] A. Friedman, *Stochastic Differential Equations and Applications*. Vol. 1. Probability and Mathematical Statistics, Vol. 28. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975. xiii+231 pp.
- [14] I. I. Gihman and A. V. Skorohod, *Stochastic Differential Equations*. Translated from the Russian by Kenneth Wickwire. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 72*. Springer-Verlag, New York-Heidelberg, 1972. viii+354 pp.
- [15] W. R. P. Hammersley, D. Šiška and L. Szpruch, McKean-Vlasov SDEs under measure dependent Lyapunov conditions, *Ann. Inst. H. Poincaré Probab. Statist.* **57** (2021), 1032–1057
- [16] M. Huang, R. P. Malhamé and P. E. Caines, Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle, *Commun. Inf. Syst.* **6** (2006), 221–251.
- [17] M. Huang, R. P. Malhamé and P. E. Caines, Large-population cost-coupled LQG problems with nonuniform agents: individual-mass behavior and decentralized  $\epsilon$ -Nash equilibria, *IEEE Trans. Automat. Control* **52** (2007), 1560–1571.
- [18] R. Iturriaga and K. Wang, A discrete weak KAM method for first-order stationary mean field games, *SIAM J. Appl. Dyn. Syst.* **22**, (2023), 1253–1274.
- [19] M. Kac, Foundations of kinetic theory. In: *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, III, 1954–1955*, 171–197, Berkeley, Los Angeles: University of California Press, (1956).
- [20] J. M. Lasry and P. L. Lions, Jeux à champ moyen. I. Le cas stationnaire. (French) [Mean field games. I. The stationary case], *C. R. Math. Acad. Sci. Paris* **343** (2006), 619–625.
- [21] J. M. Lasry and P. L. Lions, Jeux à champ moyen. II. Horizon fini et contrôle optimal. (French) [Mean field games. II. Finite horizon and optimal control], *C. R. Math. Acad. Sci. Paris* **343** (2006), 679–684.
- [22] J. M. Lasry and P. L. Lions, Mean field games, *Jpn. J. Math.* **2** (2007), 229–260.
- [23] Z. Liu and J. Ma, Existence, uniqueness and exponential ergodicity under Lyapunov conditions for McKean-Vlasov SDEs with Markovian switching. *J. Differential Equations* **337** (2022), 138–167.
- [24] Z. Liu and J. Ma, Existence, uniqueness and ergodicity for McKean-Vlasov SDEs under distribution-dependent Lyapunov conditions. *arXiv preprint* (2023), arXiv:2309.05411.
- [25] H. P. McKean, Propagation of chaos for a class of nonlinear parabolic equations. In: *Lecture Series in Differential Equations* **7** (1967), 41–57.
- [26] H. Qiao, Limit theorems of invariant measures for multivalued McKean-Vlasov stochastic differential equations *J. Math. Anal. Appl.* **528** (2023), 25 pp.

- [27] H. Qiu and Y. Wang, Continuous dependence of recurrent solutions for stochastic differential equations. *Electron. J. Differential Equations* **113** (2020), 9pp.
- [28] P. Ren, H. Tang and F. Y. Wang, Distribution-path dependent nonlinear SPDEs with application to stochastic transport type equation, *arXiv preprint* (2020), arXiv:2007.09188.
- [29] A. S. Sznitman, Nonlinear reflecting diffusion process, and the propagation of chaos and fluctuations associated, *J. Funct. Anal.* **56** (1984), 311–336 .
- [30] A. S. Sznitman, Topics in propagation of chaos. École d'Été de Probabilités de Saint-Flour XIX–1989, 165–251, Lecture Notes in Math. 1464, Springer, Berlin, (1991).
- [31] C. Villani, *Optimal Transport. Old and new*. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 338. Springer-Verlag, Berlin, (2009), xxii+973 pp.
- [32] F. Y. Wang, Distribution dependent SDEs for Landau type equations. *Stochastic Process. Appl.* **128** (2018), 595–621.
- [33] F. Wu, F. Xi and C. Zhu, On a class of McKean-Vlasov stochastic functional differential equations with applications, *J. Differential Equations* bf 371 (2023), 31–49.

Z. LIU: SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN 116024, P. R. CHINA

*Email address:* `zxliu@dlut.edu.cn`

J. MA (CORRESPONDING AUTHOR): SCHOOL OF MATHEMATICS AND STATISTICS, AND CENTER FOR MATHEMATICS AND INTERDISCIPLINARY SCIENCES, NORTHEAST NORMAL UNIVERSITY, CHANGCHUN, 130024, P. R. CHINA

*Email address:* `mathmajun@163.com`