Nucleon Consumption and Mass-Energy Conversion Induced by Dark Matter

Shao-Feng $Ge^{1, 2, *}$ and Xiao-Dong $Ma^{3, 4, \dagger}$

¹Tsung-Dao Lee Institute & School of Physics and Astronomy, Shanghai Jiao Tong University, China

²Key Laboratory for Particle Astrophysics and Cosmology (MOE)

& Shanghai Key Laboratory for Particle Physics and Cosmology,

Shanghai Jiao Tong University, Shanghai 200240, China

³ Key Laboratory of Atomic and Subatomic Structure and Quantum Control (MOE),

Guangdong Basic Research Center of Excellence for Structure and Fundamental Interactions of Matter,

Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China

⁴Guangdong-Hong Kong Joint Laboratory of Quantum Matter,

Guangdong Provincial Key Laboratory of Nuclear Science, Southern Nuclear Science Computing Center,

South China Normal University, Guangzhou 510006, China

We propose the nucleon consumption induced by dark matter as a new scenario to overcome the energy threshold of direct detection. It can be realized with proton $(\chi + p \rightarrow \chi + \ell^+)$ or neutron $(\chi + n \rightarrow \chi + \nu)$ target. Both effective operators and concrete models are provided to illustrate the idea. Since the initial DM and nucleon velocity is only 10^{-3} and 1/4 of the speed of light, respectively, the cross section and hence event rate are determined by the involved particle masses and not affected by the nucleon Fermi motion. Of the two realizations, the proton consumption has richer phenomena with both charged lepton and the daughter nuclei de-excitation to allow double or even triple coincidence to significantly suppress the background. We also illustrate the projected sensitivities at DM and neutrino experiments such as PandaX-4T, DUNE, JUNO, and Hyper-K.

Introduction – The nature of dark matter (DM) is one of the biggest unknowns in our Universe [1–3]. The particle DM is a well-established paradigm that can explain the observed astrophysical and cosmological phenomena with minimal parameter set. Especially, the weakly interacting massive particle (WIMP), whose stability is usually protected by discrete symmetry, is a theoretically well-motivated scenario that can meet the required properties to address the DM conundrum and at the same time have a detectable possibility in terrestrial experiments [4, 5]. However, the DM direct detection experiments have not found signal but constrain the DMnucleus cross section to an unprecedented level for the WIMP mass from GeV to hundreds of GeV [6, 7].

Since the DM-nucleus elastic scattering can only convert the DM kinetic energy to the nucleus recoil energy, it is difficult for a sub-GeV light DM to overcome the energy threshold of a direct detection experiment. To solve this issue, novel low-threshold experiments and theoretical ideas are proposed. On the experimental side [8, 9], for instance, the DM-electron scattering [10, 11], the Migdal effect [12], and the bremsstrahlung processes [13] can enhance the signal strength. On the theoretical side, various DM scenarios have appeared.

The recently proposed fermionic absorption DM is another example of light DM particle that can inject enough energy to the detector to produce a detectable signal with either nuclei [14–19] or electron [16, 20–22] target. The idea is that the DM (χ) is absorbed by nuclear or atomic target (T) while emitting an undetectable neutrino (ν), $\chi + T \rightarrow \nu + T'$, while the target (T') recoils. Such signals have been searched at PandaX-4T [23, 24], Majorana [25], EXO-200 [26], and CDEX [27, 28].

Using the fermion absorption and converting its mass

to recoil energy for detecting an unkown particle has a long history back to the 1940s. Although Pauli also pointed out the possibility of using its electromagnetic dipole interactions to probe the neutrino he proposed in the same letter [29], both its very existence and electromagnetic interactions are hypothetical. With only weak interactions, neutrinos are extremely difficult to detect. So Kan Chang Wang proposed in 1942 a very smart way of using the K-shell electron capture and detecting the mono-energetic daughter nuclei recoil to probe the existence of neutrino [30, 31]. This is essentially absorbing/consuming the known electron and releasing its mass as energy to probe the unknown neutrino.

In this paper, we propose a new absorption mechanism by consuming nucleon rather than the DM particle, to overcome the detection threshold. Unlike the fermionic absorption DM, which is unstable and suffers from stringent indirect detection constraints due to decaying into SM light particles like neutrinos or photons (and possibly electron-positron pair or light mesons were these channels kinematically available) [14, 15, 20–22], the nucleon consuming DM is stable due to a \mathbb{Z}_2 symmetry protecting it from decaying. Since one nucleon is consumed in the process to release almost 1 GeV energy, we anticipate there will be enough energy deposit in the detector and thus can be searched for at the DM direct detection and neutrino experiments.

Nucleon Consumption Operators and Models – For the nucleon consumption scenarios induced by DM, it may happen at the nucleus level if the momentum transfer is not large enough. Then the target nucleus (N)reduces to a lighter one (N'), $\chi + N \rightarrow \chi + N' + \ell/\nu$ where the final-state charged lepton (ℓ) or neutrino (ν) is necessary to balance the fermion flow and angular momentum. However, it is not easy to make the momentum transfer transparent with a three-body final state. On the other hand, a large enough momentum transfer would render the nucleon consumption process to happen at the nucleon level. Since the DM particle remains intact, the consumed nucleon in the initial state will become either a positively charged lepton $(\chi + p \rightarrow \chi + \ell^+)$ or a neutrino $(\chi + n \rightarrow \chi + \nu)$. For both cases, the massive nucleon (either proton p or neutron n) will release its mass around 938 MeV to a light charged lepton (positron e^+ or postively charged muon μ^+) whose mass is much smaller or even a massless neutrino. With a two-body final state, the momentum transfer $q^2 \equiv (p_N - p_\ell)^2 =$ $-m_{\chi}(m_N^2-m_\ell^2)/(m_N+m_{\chi})$ is fixed by the DM (m_{χ}) , nucleon (m_N) , and lepton $(m_\ell$ for charged lepton or neutrino) masses. With both m_{χ} and m_N around the GeV scale, the momentum transfer can probe a length scale much smaller than the nuclei size. In other words, the consumption scenario really happens at the nucleon level rather than with the nucleus as a whole.

For the DM mass below half of the nucleon mass, the same interaction that induces nucleon consumption can also lead to the proton $(p \rightarrow 2\chi + \ell^+)$ or neutron $(n \rightarrow 2\chi + \nu)$ decay. The subsequent dark decay of neutron can address the longstanding discrepancy between the bottleand beam-type measurements of the neutron lifetime [32, 33]. However, the nucleon decay can be forbidden by requiring a heavier DM mass to be on the safe side.

With a nonrelativistic incoming DM particle and the initial proton at rest, the consumed proton mass is released into the final-state DM and charged lepton energies $E_{\chi} = (m_p^2 + 2m_pm_{\chi} + 2m_{\chi}^2 - m_{\ell}^2)/2(m_p + m_{\chi})$ and $E_{\ell} = (m_p^2 + 2m_pm_{\chi} + m_{\ell}^2)/2(m_p + m_{\chi})$, respectively. Both energies are fully determined by the proton (m_p) , DM (m_{χ}) , and lepton (m_{ℓ}) masses. Although the muon mass is not that small, its value is just around 106 MeV and can hence be safely ignored as a good enough approximation. The proton realization can then be detected by the charged lepton and de-exicitation of the daughter nucleus.

While the charged lepton e^+ or μ^+ cannot be replaced by another charged particle, the SM neutrino produced by $\chi + n \rightarrow \chi + \nu$ will escape detection and can be faked by any other neutral particles such as the sterile neutrino whose mass is also small enough or a light dark particle. Of the two realizations, the neutron consumption with only the daughter nucleus de-excitation leaves less signal in the detector. So we would focus on the proton consumption scenario for illustration while most results also apply for the neutron case.

The nucleon consumption can be described in terms of four-fermion effective operators involving a pair of fermion DM particles, proton, and charged lepton are,

$$\mathcal{O}_{\chi\ell p}^{\rm SS(P)} = \overline{\chi}\chi\,\overline{\ell^c}(i\gamma_5)p,\tag{1a}$$




FIG. 1: The Feynman diagram for generating the effective operators responsible for the nucleon consumption induced by DM.

$$\mathcal{O}_{\chi\ell p}^{\mathsf{PS}(\mathsf{P})} = \overline{\chi} i \gamma_5 \chi \,\overline{\ell^c}(i\gamma_5) p,\tag{1b}$$

$$\mathcal{O}_{\chi\ell p}^{\mathsf{VV}(\mathtt{A})} = \overline{\chi}\gamma_{\mu}\chi\,\overline{\ell^{c}}\gamma^{\mu}(\gamma_{5})p,\tag{1c}$$

$$\mathcal{O}_{\chi\ell p}^{\mathtt{AV}(\mathtt{A})} = \overline{\chi}\gamma_{\mu}\gamma_{5}\chi\,\overline{\ell^{c}}\gamma^{\mu}(\gamma_{5})p,\tag{1d}$$

$$\mathcal{O}_{\chi\ell p}^{\mathsf{TT}(\tilde{1})} = \overline{\chi}\sigma_{\mu\nu}\chi\,\overline{\ell^c}\sigma^{\mu\nu}(i\gamma_5)p,\tag{1e}$$

where ℓ^c is the charge conjugation of the lepton field. For the superscript, the first character stands for the Lorentz type of the DM current while the second and the one in parenthesis for the proton-lepton current. The DM current can also take the fermion number violating case, $\overline{\chi^c}\Gamma\chi$. Then only the scalar, pseudo-scalar, and axial-vector operators $\mathcal{O}_{\chi\ell p}^{\mathrm{S,P,A}}$ can survive. The strength of each operator is parametrized by a Wilson coefficient $C_{\chi\ell p}^i$. For the DM-neutron consumption, one may just replace the charged lepton-proton current by the neutrinoneutron current, $(\overline{\nu}\Gamma n)$ or $\overline{\nu^c}\Gamma n$.

The nucleon consumption operators in Eq. (1) can be realized by extending the SM with a Dirac fermion DM candidate $\chi(1,1,0)$ besides three leptoquarks $S_1(3,1,2/3)$, $S_2(3,1,-1/3)$, and $S_3(3,1,-1/3)$. In addition, one needs to impose a discrete \mathbb{Z}_4 symmetry under which the charge assignments are,

$$\chi: i, \quad S_{1,2}: -i, \quad S_3, L, e: -1, \tag{2}$$

while the remaining SM particles are \mathbb{Z}_4 singlets. Here L and e are the SM left- and right-handed leptons. The relevant new physics Lagrangian is

$$\mathcal{L}_{\text{NP}} \ni y_1(\overline{u}\chi_L)S_1 + y_2(d\chi_L)S_2 + y_3(\overline{Q}\epsilon L^c)S_3 + y_4(\overline{u}e^c)S_3 - \mu_S\epsilon_{ijk}S_1^iS_2^jS_3^k + \text{H.c.},$$
(3)

where Q, u, d are the SM quarks while i, j, k are the color indices in the fundamental representation of SU(3)_c. For the Yukawa couples $y_{1,2,3,4}$, the flavor indices have been omitted for simplicity. The dimensional μ_S is the triple scalar coupling.

The Feynman diagram responsible for the nucleon comsuming DM is shown in Fig. 1, which will generate operators with (pseudo-)scalar DM currents in Eq. (1) once quarks condensate into nucleons. While the couplings with S_1 and S_2 already contribute one up and one

down quarks, whether the model induces a proton or neutron consumption is determined by the S_3 Yukawa couplings. A vanishing y_3 prevents neutrino from participating the process and hence only leads to the proton consumption scenario with nonzero y_4 . On the other hand, a nonzero y_3 can induce both proton and neutron consumption. Cancellation between the y_3 and y_4 contributions to the proton decay can also happen by tuning values. However, it is more desirable to adjust the model with a light sterile neutrino $\nu_R(1,1,0)$ that has a \mathbb{Z}_4 charge -1. Furthermore, the SM leptons and quarks all are neutral under the \mathbb{Z}_4 symmetry. In this case, the Yukawa terms associated with $y_{3,4}$ are replaced by $y_3(\vec{d}\nu_R^2)S_3$.

Consumption at Rest and Fermi Motion – Since the halo DM particles travel at only 10^{-3} of the speed of light, it is a good approximation to treat them as at rest. The proton consumption process then happens at rest which leads to mono-energetic DM and charged lepton in the final state. Reparametrizing the DM mass as $x \equiv m_{\chi}/m_p$, the consumption cross section takes the form as,

$$\sigma_{\chi\ell p}^{\rm SS(P)} v_{\chi} \approx \frac{m_p^2}{32\pi \Lambda_{\rm eff}^4} \frac{(1+2x)^4}{(1+x)^4}, \tag{4a}$$

$$\sigma_{\chi\ell p}^{\text{PS}(\text{P})} v_{\chi} \approx \frac{m_p^2}{32\pi \Lambda_{\text{eff}}^4} \frac{(1+2x)^2}{(1+x)^4},$$
(4b)

$$\sigma_{\chi\ell p}^{\rm VV(A)} v_{\chi} \approx \frac{m_p^2}{16\pi \Lambda_{\rm eff}^4} \frac{(1+2x)^2 (3+4x+2x^2)}{(1+x)^4}, \qquad (4c)$$

$$\sigma_{\chi\ell p}^{\text{AV(A)}} v_{\chi} \approx \frac{m_p^2}{16\pi \Lambda_{\text{eff}}^4} \frac{(1+2x)^2 (3+8x+6x^2)}{(1+x)^4}, \qquad (4d)$$

$$\sigma_{\chi\ell p}^{\text{TT}(\tilde{\mathbf{T}})} v_{\chi} \approx \frac{m_p^2}{4\pi \Lambda_{\text{eff}}^4} \frac{(1+2x)^2 (5+10x+6x^2)}{(1+x)^4}, \qquad (4e)$$

where the Wilson coefficients $C_{\chi ep}^i \equiv \Lambda_{\text{eff}}^{-2}$ can be expressed in terms of the effective scale Λ_{eff} . For convenience, one may factorize out the *x*-dependent part, $\sigma_{\chi \ell p}^i v_{\chi} \equiv f^i(x) \times m_p^2/16\pi \Lambda_{\text{eff}}^4$. As shown in Fig. 2, the *x*-dependent f(x)/x that finally appears in the event rate is typically in the range of $10^{-3} \sim 10$. The size of f(x)/x is sizable with the only exception of the pseudo-scalar case. The typical size of the consumption cross section, $\sigma_0 \equiv m_p^2/16\pi \Lambda_{\text{eff}}^4$, is mainly determined by the proton mass m_p and the effective scale Λ_{eff} .

Since the consumption process happens at the nucleon level, the Fermi motion of the bound nucleon should be taken into consideration. The nucleon momentum distribution in nuclei can be described by the Fermi-gas model together with a high-momentum tail accounting for the nucleon-nucleon correlations [34],

$$\hat{\rho}(|\boldsymbol{p}|) = \frac{1}{C} \begin{cases} 1 - 6\left(\frac{k_F a}{\pi}\right)^2, & 0 \le |\boldsymbol{p}| \le k_F, \\ 2R\left(\frac{k_F a}{\pi}\right)^2 \left(\frac{k_F}{|\boldsymbol{p}|}\right)^4, & k_F < |\boldsymbol{p}| \le 4 \,\text{GeV}, \\ 0, & |\boldsymbol{p}| > 4 \,\text{GeV}, \end{cases}$$
(5)




FIG. 2: The dependence of f(x)/x as a function of $x = m_{\chi}/m_p$. Its value is sizable with the only exception of the pseudo-scalar case.

where $a \equiv 2 \,\text{GeV}^{-1}$ is a characteristic size, $C \equiv \frac{4}{3}\pi k_F^3$ the phase space volume, and $R \equiv 1/(1 - k_F/4 \,\text{GeV})$, respectively. The typical Fermi momentum for the Xenon nuclei is $k_F \approx 0.26 \,\text{GeV}$ which corresponds to 1/4 of the speed of light. Nevertheless, the correction induced by the nucleon velocity v_N to the cross section only appears at the quadratic order $\mathcal{O}(v_N^2)$. Being below 10%, it is safe to omit this correction for an estimation of the projected sensitivity at DM and neutrino experiments. Since the dimension-6 four-fermion operators in Eq. (1) scales as $1/\Lambda_{\text{eff}}^2$, the cross section in Eq. (4) has $1/\Lambda_{\text{eff}}^4$ scaling. A 10% deviation in event rate can only lead to a negligible 2.5% uncertainty in the projected sensitivity reach of Λ_{eff} . Being different from the fermionic absorption DM case [18], the nucleon consumption cross section is not affected by the nucleon Fermi motion or Pauli blocking effect since there is no nucleon in the final state.

Search at DM and Neutrino Experiments – As mentioned earlier, the proton consumption case will leave two visible particles, a charged lepton and a daughter nucleus in the excited state. Since both proton and the DM particles are heavy with $\mathcal{O}(\text{GeV})$ masses, the released energy is mainly carried away by the light charged lepton. This also applies for the neutron consumption scenario. Then the charged leptons can directly deposit energy as Cherenkov light in a water detector such as Super-K or Hyper-K as well as scintillation light in others. In addition, positron e^+ can annihilate with a material electron to create two 511 keV gamma photons. For μ^+ , it may also decay inside the detector to produce a Michel positron first. Finally, the daughter nucleus in the excited state will emit a gamma photon. Altogether, the proton consumption can leave rich signal phenomena with double or even triple coincidence that can be used to uniquely select out the signal and at the same time significantly suppress backgrounds. It is then safe to assume background-free search for a first try of sensitivity

estimation.

Probing the proton consumption can be achieved at DM direct detection experiments such as PandaX, XENON, and LZ with low energy thresholds at keV scale. On the other hand, the neutrino experiments like Super-K, Hyper-K, and JUNO have much larger size to compensate their higher thresholds. We take PandaX-4T (4 ton of ¹³¹Xe) [35], DUNE (40 kton of ⁴⁰Ar) [36, 37], JUNO (20 kton of C₁₈H₃₀) [38], and Super-K (22 kton of H₂O) or Hyper-K (190 kton of H₂O) [39] to demonstrate the projected sensitivities at Xenon, Argon, liquid scintillator, and water based experiments, respectively. The event rate is,

$$\frac{d^2N}{dMdt} = \frac{N_A \times N_{T/p}}{\tilde{m}_T} \rho_\chi \frac{\langle \sigma v_\chi \rangle}{m_p x},\tag{6}$$

where the DM number density ρ_{χ}/m_{χ} has been replaced by $\rho_{\chi}/m_p x$ which combines with the f(x) function of σv_{χ} to give f(x)/x as mentioned earlier. Besides the Avogadro number $N_A = 6.022 \times 10^{23}$, $N_{T/p}$ is the proton number in the target T while \tilde{m}_T is the target molar mass,

$$N_{^{131}\text{Xe/p}} = 54, \qquad \tilde{m}_{^{131}\text{Xe}} = 131.292 \,\text{g/mol}, \quad (7a)$$

$$N_{40}_{\rm Ar/p} = 18, \qquad \tilde{m}_{40}_{\rm Ar} = 39.948 \,\mathrm{g/mol}, \quad (7b)$$

$$N_{\rm C_{18}H_{30}/p} = 138, \quad \tilde{m}_{\rm C_{18}H_{30}} = 246.4308 \,\mathrm{g/mol},$$
(7c)

$$N_{\rm H_2O/p} = 10, \qquad \tilde{m}_{\rm H_2O} = 18.02 \,\mathrm{g/mol}, \quad (7d)$$

for the four experiments, respectively. With the DM local density $\rho_{\chi} = 0.4 \,\text{GeV/cm}^3$, the event rates are roughly,

$$\frac{d^2 N}{dMdt} = (9.2, 7.5, 9.3, 6.8) \frac{10^5}{\text{ton} \cdot \text{yr}} \left| \frac{1 \text{ TeV}}{\Lambda_{\text{eff}}} \right|^4 \frac{f^i(x)}{x}, \quad (8)$$

for the different target materials. Considering the huge volume of 2.25 kton and 4 ton for Super-K and PandaX-4T, the total event number predicted with $\Lambda_{eff} = 1 \text{ TeV}$ can reach 10^{10} and 10^6 level within 1 year which is quite significant. This indicates that the projected sensitivity at DM and neutrino experiments can be much higher than 1 TeV.

Fig. 3 shows the projected sensitivity by requiring the signal event number needs to reach at least 100 (solid) or 1 (dashed). For a light sub-GeV DM, the effective scale can be probed up to ~ 10 TeV at the ton-scale PandaX-4T and the impressive $\gtrsim 500 \text{ TeV}$ at Hyper-K for 100 events. In the sub-GeV range, there is no big difference between the scalar $(\mathcal{O}_{\chi\ell p}^{\mathrm{sS}(\mathsf{P})})$ and pseudo-scalar $(\mathcal{O}_{\chi\ell p}^{\mathrm{pS}(\mathsf{P})})$ DM currents since the *x*-dependent function f(x)/x does not differ much as shown in Fig. 2. With increasing DM mass, the probed scale of Λ_{eff} keeps decreasing since the *x*-dependent f(x)/x and hence the event rate keeps decreasing. Between the scalar and pseudo-scalar DM current operators, the sensitivity deceases faster for the pseudo-scalar one, being consistent with the faster decreases of f(x)/x shown in Fig. 2. The sensitivity for the




FIG. 3: The solid (dashed) color curves show the experimental sensitivity on the effective scale Λ_{eff} from different experiments by assuming 1 year exposure with 100 (1) events. The upper (lower) plot is for the interactions with scalar (pseudoscalar) DM current.

vector, axial-vector, and tensor operators is similar as the scalar case with an $\mathcal{O}(1)$ rescaling factor.

For the UV model provided earlier, the effective scale $\Lambda_{\text{eff}}^{-2} \sim y_1 y_2 y_{3,4} \mu_S \Lambda_{\text{QCD}}^3 / (m_{S_1} m_{S_2} m_{S_3})^2$ is determined by not just the new physics scale, such as the S_i masses m_{S_i} , but also the QCD condensation scale Λ_{QCD} . The projected sensitivities then convert to a constraint on the scalar masses, $m_{S_i} \lesssim y_{S_i}^{1/2} \mu_S^{1/6} \Lambda_{\text{QCD}}^{1/2} (\Lambda_{\text{eff}}^{\text{max}})^{1/3}$. Although the QCD scale is just around 200 MeV, the sensivity on m_{S_i} needs not to be small since μ_S can be large.

Discussion and Conclusions – Not just the DM kinetic energy can induce nucleus recoil with elastic scattering, the particle mass can also be absorbed and then converted to the recoil energy. The previous studies mainly focus on the dark sector particles. Our study shows that consuming the nucleons inside a detector target can provide even more efficient conversion from mass to the energy deposit. For comparison, the fermionic absorption DM can only leave a recoil energy $T_r = m_{\chi}^2/2(m_T + m_{\chi})$ where m_T is the target (electron, nucleon, or nucleus) mass. Since m_{χ} can not be too large $(m_{\chi} \ll m_N)$ such

that the fermionic absorption DM can be stable enough to survive until today, the recoil energy is actually much smaller than the DM mass, $T_r \ll m_{\chi}$. Since the nucleon consumption DM mass receives a lower bound, $m_{\chi} \gtrsim m_p/2$, to guarantee the proton stability, our new scenario of proton consumption can give the charged lepton an energy of $E_{\ell} = (m_p^2 + 2m_p m_{\chi} + m_{\ell}^2)/2(m_p + m_{\chi})$ which needs not to be suppressed and the mass conversion to deposit energy is very efficient.

Notes Added – During the finalization of this paper, another paper [40] appeared on arXiv and briefly touched the similar idea of nucleon consumption DM. Although the title of their paper emphasized that baryon destruction is "*catalyzed*" by DM, the scenario of two-component DM has a major difference from ours. Our scheme provides a more economical realization with just a single DM component. The released mass purely comes from the consumed nucleon without resorting to the DM particle.

Acknowledgements

The authors would like to thank Yong Du for useful discussions and clarifying the effect of Fermi motion. SFG is supported by the National Natural Science Foundation of China (12375101, 12090060 and 12090064) and the SJTU Double First Class start-up fund (WF220442604). SFG is also an affiliate member of Kavli IPMU, University of Tokyo. XDM is supported by the Guangdong Major Project of Basic and Applied Basic Research (Grant No.2020B0301030008) and by the NSFC (Grant No. NSFC-12305110).

- * Electronic address: gesf@sjtu.edu.cn
- [†] Electronic address: maxid@scnu.edu.cn
- Bing-Lin Young, "A survey of dark matter and related topics in cosmology," Front. Phys. (Beijing) 12 no. 2, (2017) 121201. [Erratum: Front.Phys.(Beijing) 12, 121202 (2017)].
- [2] A. Arbey and F. Mahmoudi, "Dark matter and the early Universe: a review," Prog. Part. Nucl. Phys. 119 (2021) 103865, [arXiv:2104.11488 [hep-ph]].
- [3] Malcolm Fairbairn, "Galactic Anomalies and Particle Dark Matter," Symmetry 14 no. 4, (2022) 812.
- [4] Leszek Roszkowski, Enrico Maria Sessolo, and Sebastian Trojanowski, "WIMP dark matter candidates and searches—current status and future prospects," Rept. Prog. Phys. 81 no. 6, (2018) 066201, [arXiv:1707.06277 [hep-ph]].
- [5] Marc Schumann, "Direct Detection of WIMP Dark Matter: Concepts and Status," J. Phys. G 46 no. 10, (2019) 103003, [arXiv:1903.03026 [astro-ph.CO]].
- [6] Julien Billard et al., "Direct detection of dark matter—APPEC committee report*," Rept. Prog. Phys.

85 no. 5, (2022) 056201, [arXiv:2104.07634 [hep-ex]].

- [7] Marcin Misiaszek and Nicola Rossi, "Direct Detection of Dark Matter: A Critical Review," Symmetry 16 no. 2, (2024) 201, [arXiv:2310.20472 [hep-ph]].
- [8] Rouven Essig et al., "Snowmass2021 Cosmic Frontier: The landscape of low-threshold dark matter direct detection in the next decade," in Snowmass 2021. 3, 2022. [arXiv:2203.08297 [hep-ph]].
- [9] Andrea Mitridate, Tanner Trickle, Zhengkang Zhang, and Kathryn M. Zurek, "Snowmass white paper: Light dark matter direct detection at the interface with condensed matter physics," Phys. Dark Univ. 40 (2023) 101221, [arXiv:2203.07492 [hep-ph]].
- [10] Rouven Essig, Jeremy Mardon, and Tomer Volansky, *"Direct Detection of Sub-GeV Dark Matter,"* Phys. Rev. D 85 (2012) 076007, [arXiv:1108.5383 [hep-ph]].
- [11] Rouven Essig, Aaron Manalaysay, Jeremy Mardon, Peter Sorensen, and Tomer Volansky, "First Direct Detection Limits on sub-GeV Dark Matter from XENON10," Phys. Rev. Lett. 109 (2012) 021301, [arXiv:1206.2644 [astro-ph.CO]].
- [12] Masahiro Ibe, Wakutaka Nakano, Yutaro Shoji, and Kazumine Suzuki, "Migdal Effect in Dark Matter Direct Detection Experiments," JHEP 03 (2018) 194, [arXiv:1707.07258 [hep-ph]].
- [13] Chris Kouvaris and Josef Pradler, "Probing sub-GeV Dark Matter with conventional detectors," Phys. Rev. Lett. 118 no. 3, (2017) 031803, [arXiv:1607.01789 [hep-ph]].
- [14] Jeff A. Dror, Gilly Elor, and Robert Mcgehee, "Directly Detecting Signals from Absorption of Fermionic Dark Matter," Phys. Rev. Lett. **124** no. 18, (2020) 18, [arXiv:1905.12635 [hep-ph]].
- [15] Jeff A. Dror, Gilly Elor, and Robert Mcgehee,
 "Absorption of Fermionic Dark Matter by Nuclear Targets," JHEP 02 (2020) 134, [arXiv:1908.10861 [hep-ph]].
- [16] Tong Li, Jiajun Liao, and Rui-Jia Zhang, "Dark magnetic dipole property in fermionic absorption by nucleus and electrons," JHEP 05 (2022) 071, [arXiv:2201.11905 [hep-ph]].
- [17] Kai Ma, "Exploring Four Fermion Contact Couplings of a Dark Fermion and an Electron at Hadron Colliders and Direct Detection Experiments," [arXiv:2404.06419 [hep-ph]].
- [18] Shao-Feng Ge and Oleg Titov, "Incoherent Fermionic Dark Matter Absorption with Nucleon Fermi Motion," [arXiv:2405.05728 [hep-ph]].
- [19] Kai Ma, Shao-Feng Ge, Lin-Yun He, and Ning Zhou, "Complementary Search of Fermionic Absorption Operators at Hadron Collider and Direct Detection Experiments," 5, 2024. [arXiv:2405.16878 [hep-ph]].
- [20] Jeff A. Dror, Gilly Elor, Robert McGehee, and Tien-Tien Yu, "Absorption of sub-MeV fermionic dark matter by electron targets," Phys. Rev. D 103 no. 3, (2021) 035001, [arXiv:2011.01940 [hep-ph]]. [Erratum: Phys.Rev.D 105, 119903 (2022)].
- [21] Shao-Feng Ge, Xiao-Gang He, Xiao-Dong Ma, and Jie Sheng, "Revisiting the fermionic dark matter absorption on electron target," JHEP 05 (2022) 191, [arXiv:2201.11497 [hep-ph]].
- [22] Shao-Feng Ge, Kai Ma, Xiao-Dong Ma, and Jie Sheng, "Associated production of neutrino and dark fermion at future lepton colliders," JHEP 11 (2023) 190,

Destruction," [arXiv:2405.18472 [hep-ph]].

[arXiv:2306.00657 [hep-ph]].

- [23] PandaX Collaboration, Linhui Gu et al., "First Search for the Absorption of Fermionic Dark Matter with the PandaX-4T Experiment," Phys. Rev. Lett. 129 no. 16, (2022) 161803, [arXiv:2205.15771 [hep-ex]].
- [24] PandaX Collaboration, Dan Zhang et al., "Search for Light Fermionic Dark Matter Absorption on Electrons in PandaX-47," Phys. Rev. Lett. 129 no. 16, (2022) 161804, [arXiv:2206.02339 [hep-ex]].
- [25] Majorana Collaboration, I. J. Arnquist et al., "Exotic Dark Matter Search with the Majorana Demonstrator," Phys. Rev. Lett. 132 no. 4, (2024) 041001, [arXiv:2206.10638 [hep-ex]].
- [26] EXO-200 Collaboration, S. Al Kharusi et al., "Search for MeV electron recoils from dark matter in EXO-200," Phys. Rev. D 107 no. 1, (2023) 012007, [arXiv:2207.00897 [hep-ex]].
- [27] CDEX Collaboration, W. H. Dai et al., "Exotic Dark Matter Search with the CDEX-10 Experiment at China's Jinping Underground Laboratory," Phys. Rev. Lett. 129 no. 22, (2022) 221802, [arXiv:2209.00861 [hep-ex]].
- [28] CDEX Collaboration, J. X. Liu et al., "First Search for Light Fermionic Dark Matter Absorption on Electrons Using Germanium Detector in CDEX-10 Experiment," [arXiv:2404.09793 [hep-ex]].
- [29] W. Pauli, "On the Earlier and more recent history of the neutrino," Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. 1 (1991) 1–25.
- [30] Kan Chang Wang, "A Suggestion on the Detection of the Neutrino," Phys. Rev. 61 no. 1-2, (1942) 97–97.
- [31] Kan Chang Wang, "Proposed Methods of Detecting the Neutrino," Phys. Rev. 71 no. 9, (1947) 645–646.
- [32] Particle Data Group Collaboration, R. L. Workman et al., "*Review of Particle Physics*," PTEP 2022 (2022) 083C01.
- [33] A. T. Yue, M. S. Dewey, D. M. Gilliam, G. L. Greene, A. B. Laptev, J. S. Nico, W. M. Snow, and F. E. Wietfeldt, "Improved Determination of the Neutron Lifetime," Phys. Rev. Lett. 111 no. 22, (2013) 222501, [arXiv:1309.2623 [nucl-ex]].
- [34] A. Bodek and J. L. Ritchie, "Fermi Motion Effects in Deep Inelastic Lepton Scattering from Nuclear Targets," Phys. Rev. D 23 (1981) 1070.
- [35] PandaX Collaboration, Hongguang Zhang et al., "Dark matter direct search sensitivity of the PandaX-4T experiment," Sci. China Phys. Mech. Astron. 62 no. 3, (2019) 31011, [arXiv:1806.02229 [physics.ins-det]].
- [36] DUNE Collaboration, Babak Abi et al., "Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume I Introduction to DUNE," JINST 15 no. 08, (2020) T08008, [arXiv:2002.02967 [physics.ins-det]].
- [37] DUNE Collaboration, Babak Abi et al., "Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume II: DUNE Physics," [arXiv:2002.03005 [hep-ex]].
- [38] JUNO Collaboration, Fengpeng An et al., "Neutrino Physics with JUNO," J. Phys. G 43 no. 3, (2016) 030401, [arXiv:1507.05613 [physics.ins-det]].
- [39] Hyper-Kamiokande Collaboration, K. Abe et al., "Hyper-Kamiokande Design Report," [arXiv:1805.04163 [physics.ins-det]].
- [40] Yohei Ema, Robert McGehee, Maxim Pospelov, and Anupam Ray, "Dark Matter Catalyzed Baryon