

# CAPPED VERTEX FUNCTIONS FOR $\text{Hilb}^n(\mathbb{C}^2)$

JEFFREY AYERS AND ANDREY SMIRNOV

ABSTRACT. We obtain explicit formulas for capped descendent vertex functions of  $\text{Hilb}^n(\mathbb{C}^2)$  for descendants given by chern classes of tautological bundles. The expression is the result of twisting a well known generating function for normalized Macdonald polynomials. This gives an explicit description of the rational function the capped vertex is an expansion of. Along the way we prove various limit expressions of the capping operator and the bare vertex function.

## CONTENTS

1. Introduction	2
1.1. Content of the Paper	2
2. Acknowledgements	5
3. Background	5
3.1. Hilbert Schemes	5
3.2. Instantons	6
3.3. Quasimaps and Vertex Functions	8
3.4. Capping Operators	9
3.5. Elliptic Hall Algebra	10
3.6. Fock Space	11
3.7. Hopf Structures	11
3.8. Wall crossing operators, and the quantum difference operator	12
3.9. K-theoretic Stable Envelopes	13
4. Results for $r = 2$ splitting	15
4.1. Limit of multiplication by $\mathcal{O}(-1)$ , and of the matrix of tangent weights	15
4.2. Limit of Capping Operator	19
4.3. Limits of Vertex Functions	21
4.4. A relation between capped descendants	25
4.5. Limiting linear equation	26
5. Derivation of Descendants formula	27
References	29

## 1. INTRODUCTION

The K-theoretic capped vertex functions were defined in [6] as partition functions of relative quasimaps to Nakajima varieties. In this paper we consider the capped vertex functions for a variety  $X = \text{Hilb}^n(\mathbb{C}^2)$  given by Hilbert scheme of  $n$ -points in the plane  $\mathbb{C}^2$ . We consider vertex functions with a special type of descendents given by the exterior powers of the tautological bundle. In this case we obtain explicit combinatorial formula for the capped vertex function.

It was conjectured and proved in several special cases [8, 9, 11] that the partition functions corresponding to the capped vertex functions with descendents are Taylor series expansions of rational functions. The explicit formula we obtain in this paper confirms this conjecture for this special choice of descendents. In this section we overview the main definitions and results.

**1.1. Content of the Paper.** In [6], Okounkov introduced the capped vertex function with descendents as the following partition function:

**Definition 1.1.** (Section 7 of [6]) *The capped vertex with descendent  $\tau$  for a Nakajima quiver variety  $X$  is the following generating function:*

$$(1.1) \quad \widehat{V}^{(\tau)}(z) := \sum_d \text{ev}_{p_1, *}\left(\text{QM}_{rel, p_1}^d, \hat{\mathcal{O}}_{vir}^d \otimes \text{ev}_{p_2}^*(\tau)\right) z^d \in K_G(X)[[z]]$$

Here  $\text{QM}_{rel, p_1}^d$  denotes the moduli space of quasimaps from  $\mathbb{P}^1$  to a Nakajima variety  $X$  relative to  $p_1 = 0 \in \mathbb{P}^1$ . By  $\hat{\mathcal{O}}_{vir}^d \in K_G(\text{QM}_{rel, p_1}^d)$  we denote the symmetrized virtual structure sheaf of the moduli space  $\text{QM}_{rel, p_1}^d$ . This moduli space is equipped with a proper map  $\text{ev}_{p_1} : \text{QM}_{rel, p_1}^d \rightarrow X$  and  $\text{ev}_{p_1, *}$  denotes the corresponding push-forward of the  $G$ -equivariant K-theory groups, where  $G$  is a certain symmetry group acting on both the moduli space and  $X$ . For a point  $p_2 = \infty \in \mathbb{P}^2$  we also have a map  $\text{ev}_{p_2} : \text{QM}_{rel, p_1}^d \rightarrow [X]$  where  $[X]$  is the quotient stack corresponding to Nakajima variety  $X$  (we recall that  $X$  is a GIT quotient, while  $[X]$  is a categorical quotient). Using the pull-back map  $\text{ev}_{p_2}^*$  for any class  $\tau \in K_G([X])$  we can construct a K-theory class  $\text{ev}_{p_2}^*(\tau)$  on the moduli space. The resulting vertex function is a generating function of the corresponding push-forwards over all degrees  $d$ . The parameter  $z$  counting the degrees of quasimaps is usually referred to as the *Kähler parameter*.

In this paper we study the capped vertex function for  $X = \text{Hilb}^n(\mathbb{C}^2)$  with descendents

$$(1.2) \quad \tau = \Lambda_u^\bullet(\mathcal{V}) = \sum_{k=1}^n \Lambda^k(\mathcal{V}) u^k$$

where  $\mathcal{V}$  is the tautological bundle on the Hilbert scheme.

Let  $T = (\mathbb{C}^\times)^2$  be a two-dimensional torus acting on  $\mathbb{C}^2$  by  $(x, y) \rightarrow (t_1 x, t_2 y)$ . We consider the induced action of  $T$  on  $\text{Hilb}^n(\mathbb{C}^2)$  and the corresponding  $T$ -equivariant K-theory groups.

Due to a theorem of Haiman, we can identify the direct sum of  $T$ -equivariant  $K$ -theories of  $\text{Hilb}^n(\mathbb{C}^2)$  with an infinite dimensional polynomial space, which we call ‘‘Fock space’’:

**Theorem 1.2.** ([1]) *There is an isomorphism of graded vector spaces*

$$(1.3) \quad \bigoplus_{n=0}^{\infty} K_T(\text{Hilb}^n(\mathbb{C}^2)) = \text{Fock} := \mathbb{Q}[p_1, p_2, \dots] \otimes_{\mathbb{Z}} \mathbb{Q}(t_1, t_2)$$

The main result of this paper is the explicit formula of the following generating function

$$(1.4) \quad F(zq^2/\hbar, t_1, t_2, q, u, y) = \sum_n \widehat{V}_{\text{Hilb}^n(\mathbb{C}^2)}^{(\tau)}(z) y^n \in \text{Fock}[[z, y]]$$

where  $\widehat{V}_{\text{Hilb}^n(\mathbb{C}^2)}^{(\tau)}(z)$  is the capped vertex function for the Hilbert scheme  $\text{Hilb}^n(\mathbb{C}^2)$  with descendent (1.2), and  $\hbar = t_1 t_2$ . In the notations of the Fock space (1.3) we obtain:

**Theorem 1.3.** (Theorem 5.3) *The generating function (1.4) is a Taylor series expansion of the following function:*

$$(1.5) \quad F(zq^2/\hbar, t_1, t_2, q, u, y) = \exp \left( \sum_{k=1}^{\infty} \frac{y^k}{k(1-t_1^{2k})(1-t_2^{2k})} ((1-(-u)^k)\hbar^{2k} p_k + z^k \hbar^k \frac{\hbar^k - \hbar^{-k}}{1-z^k} p_k) \right)$$

When  $z = 0$  and  $u = 0$  this formula describes the generating function of the classes of structure sheaves  $\mathcal{O}_{\text{Hilb}^n(\mathbb{C}^2)} \in K_T(\text{Hilb}^n(\mathbb{C}^2))$ , which can be expressed as a sum of Macdonald polynomials  $H_\lambda$  in the Fock space. Our formula, therefore, may be understood as a  $(z, u)$ -deformation of the well-known Cauchy completeness identity for the Macdonald polynomials:

**Proposition 1.4.** (Proposition 5.1)

$$\sum_{\lambda} \frac{H_{\lambda}}{\Lambda^{\bullet}(T_{\lambda} \text{Hilb}^{|\lambda|}(\mathbb{C}^2))} y^{|\lambda|} = \exp \left( \sum_{k=1}^{\infty} \frac{y^k \hbar^{2k} p_k}{k(1-t_1^{2k})(1-t_2^{2k})} \right)$$

Notice that the  $y$ -coefficients in the expansion of (1.5) are a manifestly rational functions of  $z$ , and thus we get the following corollary

**Corollary 1.5.** (Corollary 5.5) *The capped vertex function  $\widehat{V}_{\text{Hilb}^n(\mathbb{C}^2)}^{(\tau)}(z)$  is a Taylor series expansion of a rational function in  $z$ :*

$$\widehat{V}_{\text{Hilb}^n(\mathbb{C}^2)}^{(\tau)}(z) \in \mathbb{Q}(t_1, t_2, z, q, u)$$

Let us outline the main ideas used to in the proof of (1.5). It is well known that the Hilbert schemes  $\text{Hilb}^n(\mathbb{C}^2)$  are special cases of a more general family of smooth symplectic varieties  $\mathcal{M}(n, r)$  labeled by  $n, r \in \mathbb{N}$ , known as instanton moduli spaces. When the rank

of the instantons  $r = 1$ , we obtain the Hilbert scheme  $\mathcal{M}(n, 1) = \text{Hilb}^n(\mathbb{C}^2)$ . The moduli space  $\mathcal{M}(n, r)$  is an example of the Nakajima quiver varieties and therefore we can study the capped vertex functions associated to  $\mathcal{M}(n, r)$ . Central to us is the result of Okounkov, see Theorem 7.5.23 in [6], on large framing vanishing. This theorem says that when  $r$  is sufficiently large these capped vertex functions are classical, i.e., all terms in the sum (1.1) with  $d > 0$  vanish. For the descendents we study in this paper (1.2) this happens when  $r \geq 2$ . This means that for  $\mathcal{M}(n, r)$  with  $r \geq 2$  we know the corresponding capped vertex explicitly:

$$\widehat{V}_{\mathcal{M}(n,r)}^{(\tau)}(z) = \tau \in K_G(\mathcal{M}(n, r))$$

and the right side is independent of  $z$ .

The moduli space of rank  $r = 2$  instantons is equipped with an action of the torus  $\mathbb{C}_a^* \cong \mathbb{C}^*$  such that the corresponding set of fixed points has the form:

$$(1.6) \quad \mathcal{M}(n, 2)^{\mathbb{C}_a^*} = \coprod_{n_1+n_2=n} \text{Hilb}^{n_1}(\mathbb{C}^2) \times \text{Hilb}^{n_2}(\mathbb{C}^2).$$

Our idea is to extract the information about the capped vertex function for the Hilbert scheme from the limit of  $a \rightarrow 0$  of the  $\mathbb{C}_a^*$ -equivariant capped vertex function of  $\mathcal{M}(n, 2)$ , where  $a$  denotes the equivariant parameter associated with the torus  $\mathbb{C}_a^*$ .

The main technical tool for computing this limit is the following result:

**Theorem 1.6.** (Section 7.4 of [6]) *The capping operator factors into a product*

$$\widehat{V}^{(\tau)}(z) = \Psi(z)V^{(\tau)}(z)$$

where  $V^{(\tau)}(z)$  is the bare vertex function with descendent  $\tau$  and  $\Psi(z)$  is the capping operator.

The bare vertex function  $V^{(\tau)}(z)$  is defined similarly to (1.1), with the moduli space  $\text{QM}_{rel,p_1}^d$  replaced by the moduli space of quasimaps  $\text{QM}_{ns,p_1}^d$  non-singular at  $p_1$ . The bare vertex functions are much simpler to compute, using, for instance, the equivariant localization in K-theory. Using this explicit description of the bare vertex function we obtain the following result.

**Proposition 1.7.** (Proposition 4.10) *We have the following limit of vertex functions:*

$$(1.7) \quad \lim_{a \rightarrow 0} V^{\tau,(2)}(z) = V^{\tau,(1)}(z(-t_1 t_2 q)^{-1}) \otimes V^{\tau,(1)}(z(-t_1 t_2 q))$$

The function  $V^{\tau,(2)}(z)$  in the left side of (1.7) denotes the  $\mathbb{C}_a^*$ -equivariant bare vertex function of for  $\mathcal{M}(n, 2)$ . The right side of (1.7) corresponds to the product of the vertex functions for the Hilbert schemes given by the components of components the  $\mathbb{C}_a^*$ -fixed set (1.6). In words, this proposition says that in the limit  $a \rightarrow 0$ , the bare vertex function of  $\mathcal{M}(n, 2)$  factors, up to some shifts of the Kähler parameter  $z$ , to the product of the vertex functions for the Hilbert schemes.

A similar factorization exists for the capping operator  $\Psi(z)$ . It is known that  $\Psi(z)$  is a fundamental solution matrix of the quantum difference equation associated with the corresponding Nakajima variety [7]. For the instanton moduli  $\mathcal{M}(n, r)$  this equation is the quantum dynamical equation for the quantum toroidal algebra  $\mathcal{U}_\hbar(\widehat{\mathfrak{gl}}_1)$ . Using this algebraic description, following [11] we obtain

**Proposition 1.8.** (Proposition 4.8) *The normalized capping operator  $\widetilde{\Psi}^{(2)}(z)$  for the instanton moduli space  $\mathcal{M}(n, 2)$  has the following limit:*

$$(1.8) \quad \lim_{a \rightarrow 0} \widetilde{\Psi}^{(2)}(z) = Y(z) \widetilde{\Psi}^{(1)}(z\hbar^{-1}) \otimes \widetilde{\Psi}^{(1)}(z\hbar)$$

where  $\widetilde{\Psi}^{(1)}(z\hbar^{-1}) \otimes \widetilde{\Psi}^{(1)}(z\hbar)$  in the right side denotes the tensor product of capping operators for the Hilbert schemes corresponding to the  $\mathbb{C}_a^*$ -fixed components (1.6).

The operator  $Y(z)$  in this proposition is the fusion operator for toroidal algebra  $\mathcal{U}_\hbar(\widehat{\mathfrak{gl}}_1)$ . In the proof of Proposition 4.8 we give an explicit description for its action on the Fock spaces:

$$(1.9) \quad Y(z) = \exp \left( \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right)$$

where  $\alpha_k$  denote the generators of the horizontal Heisenberg subalgebra of  $\mathcal{U}_\hbar(\widehat{\mathfrak{gl}}_1)$ . These generators act on the Fock spaces in a simple way (3.2).

Using the factorization results summarized in the two previous propositions, in the limit  $a \rightarrow 0$  of the Theorem 1.6 we arrive at the following functional relation for the capped vertex functions of the Hilbert scheme:

$$(1.10) \quad \tau \otimes 1 = Y(z) \left( \widehat{V}^{(\tau), (1)}(z) \otimes 1 \right)$$

Finally, since the operator  $Y(z)$  given by (1.9) is invertible and acts on the Fock spaces explicitly, we can solve the linear equation (1.10) for  $\widehat{V}^{(\tau), (1)}(z)$  which gives our main result.

## 2. ACKNOWLEDGEMENTS

We thank H. Dinkins and A. Okounkov for their interest to this work and useful discussions. This work is supported by the NSF grants DMS - 2054527 and DMS-2401380.

## 3. BACKGROUND

**3.1. Hilbert Schemes.** The Hilbert Scheme of  $n$ -points on  $\mathbb{C}^2$  is the scheme parameterizing codimension  $n$ -ideals of  $\mathbb{C}[x, y]$ :

$$\text{Hilb}^n(\mathbb{C}^2) = \{ \mathcal{J} \subset \mathbb{C}[x, y] : \dim(\mathbb{C}[x, y]/\mathcal{J}) = n \}$$

Let  $A \cong (\mathbb{C}^*)^2$  be the 2-dimensional algebraic torus acting on  $\mathbb{C}^2$ . This action scales coordinates:

$$(x, y) \mapsto (t_1x, t_2y)$$

This extends to an action on  $\text{Hilb}^n(\mathbb{C}^2)$ , which scales the symplectic form  $\omega \in H^2(\text{Hilb}^n(\mathbb{C}^2), \mathbb{C})$  with character  $\hbar = t_1t_2$ . Let  $A = \ker(\hbar) \subset T$  be the subtorus that fixes  $\omega$ .

The fixed point set of  $\text{Hilb}^n(\mathbb{C}^2)^T$  is a finite set, labeled by partitions of length  $n$ . For a partition  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ , with each  $\lambda_i$  weakly decreasing,  $\sum_{i=1}^k \lambda_i = n$ , the corresponding fixed ideal is of the following form:

$$\mathcal{J}^T = \{y^{\lambda_1}, xy^{\lambda_2}, \dots, x^k\}$$

See figure 1 for an example:

$y^2$	
$y$	$xy$
$1$	$x$

FIGURE 1. An example of a torus fixed point for the Hilbert Scheme  $\text{Hilb}^5(\mathbb{C}^2)$  corresponding to the partition  $(3, 2)$ . The monomial ideal here is  $\{y^3, xy^2, x^2\}$

**3.2. Instantons.**  $\text{Hilb}^n(\mathbb{C}^2)$  appears as the rank 1 case of a more general moduli space: the moduli space of instantons of rank  $r$ . We describe this moduli space as the moduli of framed torsion free sheaves of rank  $r$ .

Let  $\mathcal{F}$  be a framed torsion free sheaf of rank  $r$  over  $\mathbb{P}^2$  with fixed second chern class  $c_2(\mathcal{F}) = n$ . A framing of a sheaf is a choice of isomorphism

$$\varphi : \mathcal{F}|_{L_\infty} \rightarrow \mathcal{O}_{L_\infty}^{\oplus r}$$

Where  $L_\infty$  is the line at infinity in  $\mathbb{C}^2$ . The moduli space of these sheaves, denoted  $\mathcal{M}(n, r)$  is known to arise as the Nakajima Quiver Variety of the framed double Jordan quiver. We quickly review the construction of this variety:

Let  $Q$  be the framed quiver with one vertex and one edge. Let  $V = \mathbb{C}^n, W = \mathbb{C}^r$ .

$$\text{Rep}(Q) = \text{Hom}(W, V) \oplus \text{Hom}(V, V)$$

This space has a natural  $GL(V)$  action by conjugating the endomorphisms, and multiplying the vertical maps:  $A \in \text{Hom}(V, V), I \in \text{Hom}(W, V)$




FIGURE 2. The quiver giving rise to  $\mathcal{M}(n, r)$

$$\begin{aligned} A &\mapsto gAg^{-1} \\ I &\mapsto gI \end{aligned}$$

This action induces a Hamiltonian action on  $T^*Rep(Q)$ :

$$T^*Rep(Q) = Rep(Q) \oplus Rep(Q)^*$$

with a moment map

$$\mu : T^*Rep(Q) \rightarrow \mathfrak{g}^*$$

Where  $\mathfrak{g} = \text{Lie}(GL(V))$ . With the choice of  $GL(V)$  character  $\theta$ , called the stability parameter, we can define the Nakajima quiver variety:

$$\mathcal{M}(n, r) = \mu^{-1}(0) //_{\theta} GL(V)$$

Here we choose the positive stability condition,  $\theta = 1$

In the special case of  $r = 1$  we recover the Hilbert scheme[5]:

$$\mathcal{M}(n, 1) = \text{Hilb}^n(\mathbb{C}^2)$$

There is again a torus action  $A \cong (\mathbb{C}^*)^r$  on the  $i^{\text{th}}$ -summand in the framing. This scales with coordinate  $a_i$ . Let  $T = A \times \mathbb{C}^2$  with  $\mathbb{C}^2$  scaling the plane coordinates with the same characters  $t_i$  as above. As before we scale the symplectic form with weight  $\hbar = t_1 t_2$ . Let  $\mathbb{C}^* \subset A$  be a subtorus that acts on the rank  $r = r_1 + r_2$  framing space by scaling  $r_2$ :  $r_1 + ar_2$ .

From this we split the framing space from  $r$  to  $r_1 + ar_2$ . The fixed point set of this subtorus decomposes the moduli space into the following product:

$$\mathcal{M}(n, r)^{\mathbb{C}^*} = \coprod_{n_1+n_2=n} \mathcal{M}(n_1, r_1) \times \mathcal{M}(n_2, r_2)$$

Acting by  $T$  produces a fixed point set consisting of tuples of partitions of length  $n$ :

$$(3.1) \quad \mathcal{M}(n, r)^T = \{(\lambda_1, \dots, \lambda_r) : \sum |\lambda_i| = n\}$$

The classes of fixed points  $(\lambda_1, \dots, \lambda_n)$  form a basis in the localized  $K$ -theory of  $\mathcal{M}(n, r)$ . Specifically, the structure sheaves  $\mathcal{O}_{(\lambda_1, \dots, \lambda_n)}$  are mapped to polynomials formed from the

product of Macdonald polynomials in Haiman's normalization. In the rank 1 case, classes of fixed points correspond with Macdonald polynomials.

**3.3. Quasimaps and Vertex Functions.** Our main reference for this section is the notes [6].

Let  $\mathbf{QM}^d$  denote the moduli space of degree  $d$  quasimaps from  $\mathbb{P}^1$  to  $\mathcal{M}(n, r)$ . Let  $\mathbb{C}_q^\times$  be the standard torus acting on  $\mathbb{P}^1$  by scaling coordinates with weight  $q$ :

$$[z_1 : z_2] \mapsto [qz_1 : z_2]$$

There are two fixed points of this action:  $p_1 = 0, p_2 = \infty \in \mathbb{P}^1$ , meaning that the only two nonsingular or regular points of the  $\mathbb{C}_q^\times$ -fixed quasimap may occur at these fixed points. Let  $G = T \times \mathbb{C}_q^\times$  be the group acting on  $\mathbf{QM}^d$ . There is a subset of the moduli space we'll be primarily interested: the open subset of quasimaps of degree  $d$  nonsingular at 0 or  $\infty$ . Denote this open subset  $\mathbf{QM}_p^d \subset \mathbf{QM}^d$ .

The fixed point set  $(\mathbf{QM}_p^d)^G$  consists of pairs  $(\lambda, d_\lambda)$ , where  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$  is a tuple of partitions such that  $|\lambda| = n$  corresponding to a fixed point in  $\mathcal{M}(n, r)$ . The second element in the tuple is the degree  $d_\lambda$ , which assigns a nonnegative integer  $d_\square$  to each box  $\square \in \lambda$  in such a way that the data is organized into  $r$ -plane partitions ([2]). These integers corresponding to the degree data must add up to  $d$ :  $|d_\lambda| = \sum_{\square \in \lambda} d_\square = d$

There is an evaluation map from this open subset to  $\mathcal{M}(n, r)$  by evaluating a map at  $p$ . In general as  $\mathbf{QM}_p^d$  is not proper over  $\mathcal{M}(n, r)$  we don't have the pushforward of this map to K-theory. We can avoid this issue by using localized K-theory instead.

**Definition 3.1.** (Section 7 of [6]) *The bare vertex of  $\mathcal{M}(n, r)$  is the following generating function:*

$$V(z) := \sum_d \text{ev}_{p_1, *}\left(\mathbf{QM}_{p_1}^d, \hat{\mathcal{O}}_{vir}^d\right) z^d \in K_G(\mathcal{M}(n, r))_{loc}[[z]]$$

Where  $\hat{\mathcal{O}}_{vir}^d$  is the symmetrized virtual structure sheaf on  $\mathbf{QM}_{p_1}^d$

In the above definition we use  $z^d = z_1^{d_1} \cdots z_m^{d_m}$  to keep track of degrees in what are usually called *Kähler parameters*.

For the second fixed point at infinity:  $p_2$ , we do not have an evaluation map to the moduli space itself since we don't assume that  $p_2$  is nonsingular. Rather we have an evaluation map to the quotient stack:  $\text{ev}_{p_2} : \mathbf{QM}_{p_1}^d \rightarrow [\mu^{-1}(0)/\prod GL(V_i)]$ . Let  $\tau \in K_G([\mu^{-1}(0)/\prod GL(V_i)])$  be a character of  $G$ , so that we have the following definition:

**Definition 3.2.** (Section 7 of [6]) *The bare vertex with descendent  $\tau$  is the generating function:*

$$V^{(\tau)}(z) := \sum_d \text{ev}_{p_1, *} \left( \text{QM}_{p_1}^d, \hat{\mathcal{O}}_{vir}^d \otimes \tau \left( \mathcal{V}|_{p_2} \right) \right) z^d \in K_G(\mathcal{M}(n, r))_{loc}[[z]]$$

Where  $\tau \left( \mathcal{V}|_{p_2} \right) = \text{ev}_{p_2}^* \left( \tau(\mathcal{V}) \right) \in K_G(\mathcal{M}(n, r))$ , for a Schur functor  $\tau$  of  $\mathcal{V}$ .

Let  $\text{QM}_{rel, p_1}^d$  be the moduli space of degree  $d$  quasimaps from  $\mathbb{P}^1$  to  $\mathcal{M}(n, r)$  relative to  $p_1$ .

**Definition 3.3.** (Section 7 of [6]) *The capped vertex with descendent  $\tau$  is the generating function:*

$$\widehat{V}^{(\tau)}(z) := \sum_d \text{ev}_{p_1, *} \left( \text{QM}_{rel, p_1}^d, \hat{\mathcal{O}}_{vir}^d \otimes \tau \left( \mathcal{V}|_{p_2} \right) \right) z^d \in K_G(\mathcal{M}(n, r))[[z]]$$

Note that,  $\text{ev}_{p_1}$  is proper for relative maps, so we land in non-localized equivariant  $K$ -theory [6].

A remarkable rigidity theorem first proven by Okounkov in [6] is that for sufficiently large rank the capped vertex function obeys a property known as *large frame vanishing*:

**Theorem 3.4.** (Theorem 7.5.23 in [6]) *For every  $\tau$  with  $r$  sufficiently large,*

$$\widehat{V}^{(\tau)}(z) = \tau(\mathcal{V})\mathcal{K}^{1/2} + O(z)$$

Where  $\mathcal{K}$  is the canonical bundle on  $\mathcal{M}(n, r)$

### 3.4. Capping Operators.

**Definition 3.5.** (Section 8 of [6]) *The capping operator is the generating function:*

$$\Psi(z) := \sum_d \text{ev}_{p_1, *} \otimes \text{ev}_{p_2, *} \left( \text{QM}_{rel, p_1, ns, p_2}^d, \hat{\mathcal{O}}_{vir}^d \right) z^d \in K_G(\mathcal{M}(n, r))_{loc}^{\otimes 2}[[z]]$$

The following theorem gives a relation between the three previous definitions:

**Theorem 3.6.** (Section 7.4 of [6]) *The capping operator satisfies the equation*

$$\widehat{V}^{(\tau)}(z) = \Psi(z)V^{(\tau)}(z)$$

In [7], Okounkov and Smirnov showed that the operator  $\Psi(z)$  can be computed explicitly as the fundamental solution of the *quantum difference equation*. This equation is described via objects coming from a certain Hopf algebra acting on the equivariant  $K$ -theory of  $\mathcal{M}(n, r)$

**3.5. Elliptic Hall Algebra.** This section will follow [10], see also section 7.1 of [7].

Let  $\mathbf{Z} = \mathbb{Z}^2$ ,  $\mathbf{Z}^* = \mathbf{Z} \setminus \{(0, 0)\}$ ,

$$\mathbf{Z}^+ = \{(i, j) \in \mathbf{Z}, i > 0, \text{ or } i = 0, j > 0\}$$

and  $\mathbf{Z}^- = -\mathbf{Z}^+$ . Let  $\mathbf{a} = (a_1, a_2) \in \mathbf{Z}^+$ , and define the degree to be the gcd of the integers  $a_1, a_2$ :

$$\deg(\mathbf{a}) = \gcd(a_1, a_2)$$

Set  $\epsilon_{\mathbf{a}} = 1$  if  $\mathbf{a} \in \mathbf{Z}^+$ ,  $\epsilon_{\mathbf{a}} = -1$  if  $\mathbf{a} \in \mathbf{Z}^-$ . If  $\mathbf{a}, \mathbf{b}$  are non-collinear elements, then set  $\epsilon_{\mathbf{a}, \mathbf{b}} = \text{sgn}(\det(\mathbf{a}, \mathbf{b}))$

Let

$$n_k = \frac{(t_1^{k/2} - t_1^{-k/2})(t_2^{k/2} - t_2^{-k/2})(\hbar^{k/2} - \hbar^{-k/2})}{k}$$

**Definition 3.7.** The elliptic hall algebra  $\mathcal{U}_{\hbar}(\widehat{\mathfrak{gl}}_1)$  is an associative unital algebra over  $\mathbb{C}(t_1^{1/2}, t_2^{1/2})$  generated by elements  $e_{\mathbf{a}}, K_{\mathbf{a}}$  where  $\mathbf{a} \in \mathbf{Z}^+$  with the following relations:

- $K_{\mathbf{a}}$  is central for all  $\mathbf{a} \in \mathbf{Z}^+$ , and  $K_{\mathbf{a}+\mathbf{b}} = K_{\mathbf{a}}K_{\mathbf{b}}$
- If  $\mathbf{a}, \mathbf{b}$  are collinear, then

$$[e_{\mathbf{a}}, e_{\mathbf{b}}] = \delta_{\mathbf{a}+\mathbf{b}} \frac{K_{\mathbf{a}}^{-1} - K_{\mathbf{a}}}{n_{\deg(\mathbf{a})}}$$

- If  $\deg(\mathbf{a}) = 1$  and the triangle with vertices  $\{(0, 0), \mathbf{a}, \mathbf{a} + \mathbf{b}\}$  has no interior points then

$$[e_{\mathbf{a}}, e_{\mathbf{b}}] = \epsilon_{\mathbf{b}, \mathbf{a}} K_{\alpha(\mathbf{a}, \mathbf{b})} \frac{\Psi_{\mathbf{a}, \mathbf{b}}}{n_1}$$

Where

$$\alpha(\mathbf{a}, \mathbf{b}) = \begin{cases} \frac{\epsilon_{\mathbf{a}}(\epsilon_{\mathbf{a}}(\mathbf{a}) + \epsilon_{\mathbf{b}}(\mathbf{b}) - \epsilon_{\mathbf{a}+\mathbf{b}}(\mathbf{a}+\mathbf{b}))}{2} & \epsilon_{\mathbf{a}, \mathbf{b}} = 1 \\ \frac{\epsilon_{\mathbf{b}}(\epsilon_{\mathbf{a}}(\mathbf{a}) + \epsilon_{\mathbf{b}}(\mathbf{b}) - \epsilon_{\mathbf{a}+\mathbf{b}}(\mathbf{a}+\mathbf{b}))}{2} & \epsilon_{\mathbf{a}, \mathbf{b}} = -1 \end{cases}$$

And the elements  $\Psi_{\mathbf{a}, \mathbf{b}}$  are defined by

$$\sum_{k=0}^{\infty} \Psi_{k\mathbf{a}} z^k = \exp \left( \sum_{l=1}^{\infty} n_l e_{l\mathbf{a}} z^l \right)$$

**3.5.1. Heisenberg Subalgebras.** For  $w \in \mathbb{Q} \cup \{\infty\}$ , let  $d(w)$  and  $n(w)$  be the denominator, and numerator of  $w$  with  $d(w) \geq 0$ , and  $d(w), n(w)$  coprime. If  $w = \infty$ , set  $d(w) = 0, n(w) = 1$ . Set

$$\alpha_k^w = e_{(d(w)k, n(w)k)}, \quad k \in \mathbb{Z} \setminus \{0\}$$

to be the element that generates a Heisenberg subalgebra. Visually, we can think of the algebra  $\mathcal{U}_{\hbar}(\widehat{\mathfrak{gl}}_1)$  as an integer lattice, with the Heisenberg subalgebras of  $\mathcal{U}_{\hbar}(\widehat{\mathfrak{gl}}_1)$  corresponding

to the lines of slope  $w$ .

The upper triangular  $R$ -matrix for a wall  $w \in \mathbb{Q}$  corresponding to a particular Heisenberg subalgebra is given by

$$R_w^+ = \exp \left( - \sum_{k=1}^{\infty} n_k \alpha_k^w \otimes \alpha_{-k}^w \right)$$

The lower triangular  $R$ -matrix is given by

$$R_w^- = \exp \left( - \sum_{k=1}^{\infty} n_k \alpha_{-k}^w \otimes \alpha_k^w \right)$$

**3.6. Fock Space.** In the rank 1 instanton moduli space, the localized equivariant  $K$ -theory is known to be isomorphic to the Fock space. Define the Fock space to be

$$\mathbf{Fock} := \mathbb{Q}[p_1, p_2, \dots] \otimes_{\mathbb{Z}} \mathbb{Q}(t_1, t_2)$$

Then there is an isomorphism of graded vector spaces

$$\bigoplus_{n=0}^{\infty} K_T(\text{Hilb}^n(\mathbb{C}^2))_{loc} = \mathbf{Fock}$$

Where the grading of  $\mathbf{Fock}$  is defined to be  $\deg(p_k) = k$ . As previously mentioned, the fixed point set of  $\text{Hilb}^n(\mathbb{C}^2)$  consists of partitions of length  $n$ . A basis of  $\mathbf{Fock}$  is the Macdonald polynomials in Haiman's normalization [1], and by the above isomorphism we refer to this as the fixed point basis.

With this identification, by splitting the framing as  $\mathcal{W} = a_1 + \dots + a_r$  we can then get a canonical isomorphism of  $\mathcal{U}_h(\widehat{\mathfrak{gl}}_1)$ -algebras

$$\bigotimes_{i=1}^r K_T(\mathbf{Fock}) = K_T \mathcal{M}(n, r)$$

**3.7. Hopf Structures.** The elliptic hall algebra is a triangular hopf algebra, where different structures arise from a choice of slope. The coproduct was described in [7], and has the following expression

$$\Delta_s : \mathcal{U}_h(\widehat{\mathfrak{gl}}_1) \rightarrow \mathcal{U}_h(\widehat{\mathfrak{gl}}_1) \otimes \mathcal{U}_h(\widehat{\mathfrak{gl}}_1)$$

Explicitly this coproduct acts on the slope 0 Heisenberg subalgebra as follows, where we define  $\Delta = \Delta_0$ :

$$\begin{aligned}\Delta(\alpha_{-k}^0) &= \alpha_{-k}^0 \otimes 1 + K^{-k} \otimes \alpha_{-k}^0 \\ \Delta(\alpha_k^0) &= \alpha_k^0 \otimes K^k + 1 \otimes \alpha_{-k}^0 \\ \Delta(K) &= K \otimes K\end{aligned}$$

This in turn allows us to determine how the Hopf structure interacts with the power sum basis, given the following action of the creation/annihilation operators:

$$(3.2) \quad \alpha_k^0(f) = \begin{cases} -k \frac{\partial}{\partial p_k}(f) & k > 0 \\ \frac{-p_{-n} f}{(t_1^{k/2} - t_1^{-k/2})(t_2^{k/2} - t_2^{-k/2})} & k < 0 \end{cases}$$

Define the evaluation map

$$ev^{(r)} : \mathcal{U}_h(\widehat{\mathfrak{gl}}_1) \rightarrow \text{End}(\text{Fock}^{\otimes a})$$

as

$$ev^{(r)}(\alpha) = ev_{(a_1)} \otimes \cdots \otimes ev_{(a_k)} (\Delta^{(r)}(\alpha))$$

**3.8. Wall crossing operators, and the quantum difference operator.** Consider the following power series defined in section 7 of [7]

$$B_w(z) =: \exp \left( \sum_{k=0}^{\infty} \frac{n_k \hbar^{-krd(w)/2}}{1 - z^{-kd(w)} q^{kn(w)} \hbar^{-krd(w)/2}} \alpha_{-k}^w \alpha_k^w \right) :$$

Called the wall crossing operator for wall  $w \in \mathbb{Q} \cup \{\infty\}$ . Here the symbol  $::$  means to first act with all operators  $\alpha_k^w$  for  $k > 0$ .

**Theorem 3.8** (Theorem 9 in [7]). *The quantum difference operator is defined as the operator given by the following product*

$$\mathbf{M}(z) = \mathcal{O}(1) \prod_{-1 \leq w < 0}^{\leftarrow} B_w(z)$$

Here, the left facing arrow denotes the order of the product. The quantum difference equation (QDE) for the rank  $r$  instanton moduli space is given by

**Theorem 3.9.** (Section 7 of [7])

$$\Psi^{(r)}(zq)\mathcal{O}(1) = \mathbf{M}(z)\Psi^{(r)}(z)$$

Where  $\Psi^{(r)}(z)$  is the capping operator 3.5. We normalize the solution so that

$$\Psi^{(r)}(0) = 1$$

**3.9. K-theoretic Stable Envelopes.** We briefly recall the basic definition and properties of K-theoretic Stable Envelopes. See Section 2.1 of [7], or Sections 9.1-9.2 of [6] for a more detailed exposition.

As before let  $T = A \times \mathbb{C}^2$  be the torus acting on  $\mathcal{M}(n, r)$  with the subtorus  $A$  scaling the symplectic form with weight  $\hbar$ . The K-theoretic stable envelope is a map between equivariant K-theories

$$\text{Stab} : K_T(\mathcal{M}(n, r)^A) \rightarrow K_T(\mathcal{M}(n, r))$$

We require three pieces of data before defining the stable envelope.

The first is a choice of chamber, which is a decomposition of  $\text{Lie}_{\mathbb{R}}(A) = \text{cochar}(A) \otimes_{\mathbb{Z}} \mathbb{R}$  into connected components. We fix a chamber  $\mathfrak{C}$  corresponding to  $a_1 < a_2 < \dots < a_r$ . This determines which directions are attracting and which are repelling. If we have a single parameter  $a$ , then this choice corresponds to sending  $\frac{a_1}{a_2} = a \rightarrow 0$ . Throughout the paper we normalize these parameters as  $(a_1, a_2) = (1, a^{-1})$

This choice of chamber determines a decomposition of the normal bundle to  $\mathcal{M}(n, r)$  at a fixed point  $\lambda$

$$N_{\lambda} = N_{\lambda}^{+} \oplus N_{\lambda}^{-}$$

into attracting and repelling directions with an attracting set

$$\text{Attr}(\lambda) = \{x \in \mathcal{M}(n, r) : \lim_{z \rightarrow 0} \sigma \cdot z \in \lambda\}$$

For  $\sigma$  a cocharacter in  $\mathfrak{C}$ .

The full attracting set  $\text{Attr}^f$  is, by definition, the smallest closed subset of  $X$  containing  $\text{Attr}(\lambda)$  that is closed under  $\text{Attr}(\cdot)$

The second choice is a polarization  $T^{1/2} \in K_G(\mathcal{M}(n, r))$ , which is a half of the decomposition of the tangent bundle. Let  $\mathcal{W}$  be the tautological bundle associated to the framing of  $\mathcal{M}(n, r)$ . As an element of the equivariant K-theory,

$$\mathcal{W} = a_1 + \dots + a_r$$

Let  $\mathcal{V}$  be the rank  $n$  tautological bundle of  $\mathcal{M}(n, r)$ . The weights of  $\mathcal{V}$  at a fixed point  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$  are given by

$$\mathcal{V}_{\lambda} = \sum_{\square \in (\lambda_1, \lambda_2, \dots, \lambda_r)} \varphi_{\lambda_1, \lambda_2, \dots, \lambda_r}(\square)$$

Here

$$(3.3) \quad \varphi_{\lambda_1, \lambda_2, \dots, \lambda_r}(\square) = a_{n(\square)} t_1^{y(\square)} t_2^{x(\square)}$$

Where  $n(\square) = i$  if  $\square \in \lambda_i$ , and  $x(\square), y(\square)$  denote the standard coordinates of a box. The function  $\varphi_{\lambda_1, \lambda_2, \dots, \lambda_r}(\square)$  gives the weights associated to a box in a tuple of partitions. For  $\mathcal{M}(n, r)$  this polarization can be taken to be

$$T^{1/2} = \mathcal{W}^* \otimes \mathcal{V} + \frac{1}{t_2} \mathcal{V}^* \otimes \mathcal{V} - \frac{1}{\hbar} \mathcal{V}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{V}$$

The symbol  $*$  means taking dual in equivariant  $K$ -theory, and inverts all weights:  $*$  :  $w \mapsto w^{-1}$

This choice of polarization is due to section 2.2.7 of [3], and is the natural choice for any Nakajima variety. With this choice of polarization, the tangent bundle to  $\mathcal{M}(n, r)$  can be decomposed as

$$T\mathcal{M}(n, r) = T^{1/2} + \hbar (T^{1/2})^*$$

Finally, we require a choice of slope  $s \in \text{Pic}(\mathcal{M}(n, r)) \otimes_{\mathbb{Z}} \mathbb{Q}$  which should be suitably generic, see section 2.1 of [7] for details.

**Proposition 3.10.** ([6],[7]) *Let  $T, A$  be as above, then for an arbitrary choice of chamber  $\mathfrak{C}$ , polarization  $T^{1/2}$ , and slope  $s$ , there exists a unique map*

$$\text{Stab}_{(r)} : K_T(\mathcal{M}(n, r)^A) \rightarrow K_T(\mathcal{M}(n, r))$$

satisfying the following three axioms:

- *Support condition:*  $\text{supp}(\text{Stab}) \subset \text{Attr}^f$
- *Degree condition:*

$$\deg_A \text{Stab}|_{F_2 \times F_1} \otimes s|_{F_1} \subset \deg_A \text{Stab}|_{F_2 \times F_2} \otimes s|_{F_2}$$

Where  $F_1, F_2$  are fixed components, and  $\deg_A$  is the degree of a Laurent polynomial given by the Newton polygon:

$$\deg_A \sum f_{\mu} z^{\mu} = \text{Convex Hull}(\{\mu, f_{\mu} \neq 0\})$$

Considered up to shift by a slope  $s$ .

- *Normalization condition:*

$$\text{Stab}|_{F \times F} = (-1)^{rkT_{>0}^{1/2}} \left( \frac{\det N_{\lambda}^{-}}{\det T_{\neq 0}^{1/2}} \right)^{1/2} \otimes \Lambda^{\bullet}(N_{\lambda}^{-})$$

4. RESULTS FOR  $r = 2$  SPLITTING

Our vertex function is related to Okounkov's vertex function, denoted  $V_O$  by a  $z$ -shift and multiple of the operator of multiplication by the tautological bundle:

$$V_O(z) = \mathcal{O}(r)V(zq^r)$$

Using localization in rank  $r$  the relation between the capped and bare descendent functions in the basis of torus fixed points for  $\mathcal{M}(n, r)$  is

$$\widehat{V}^{(\tau)}(z) = \Lambda^\bullet(T\mathcal{M}(n, r)^\vee)\Psi(-z)\Lambda^\bullet(T\mathcal{M}(n, r)^\vee)^{-1}V_O^{(\tau)}(z)$$

Where  $\Lambda^\bullet(T\mathcal{M}(n, r)^\vee)$  is the normalized matrix of tangent weights for rank  $r$ . In particular, for  $X = \text{Hilb}^n(\mathbb{C}^2)$  we have

$$(4.1) \quad \widehat{V}^{(\tau)}(z) = \Lambda^\bullet(TX^\vee)\Psi(-z)\Lambda^\bullet(TX^\vee)^{-1}V_O^{(\tau)}(z)$$

The difference equation satisfied by  $\Psi(z)$  is

$$(4.2) \quad \Psi(z/q^2)\mathcal{O}(-1) = M(z)\Psi(z)$$

**4.1. Limit of multiplication by  $\mathcal{O}(-1)$ , and of the matrix of tangent weights.** Let

$$Stab_{(r)} : K_T(\text{Hilb}_n(\mathbb{C}^2) \times \cdots \times \text{Hilb}_n(\mathbb{C}^2)) \rightarrow K_T(\mathcal{M}(n, r))$$

denote the K-theoretic stable envelope map. Splitting the framing as  $r = r_1 + r_2$ , we use the following result known as the *triangle lemma*

**Proposition 4.1.** (Prop 9.2.8 of [6]) *The following diagram commutes:*

$$\begin{array}{ccc} K_T(\mathcal{M}(n, r_1) \times \mathcal{M}(n, r_2)) & \xrightarrow{Stab_{r_1 r_2}} & K_T(\mathcal{M}(n, r)) \\ & \swarrow \underbrace{Stab_{\underbrace{(1, \dots, 1)}_{r_1}} \otimes Stab_{\underbrace{(1, \dots, 1)}_{r_2}}} & \searrow \underbrace{Stab_{\underbrace{(1, \dots, 1)}_r}} \\ & K_T(\text{Hilb}_n(\mathbb{C}^2) \times \cdots \times \text{Hilb}_n(\mathbb{C}^2)) & \end{array}$$

Define  $\Delta_{r_1 r_2}$  to be the diagonal of the stable envelope in the basis of fixed points.

**Example 4.2.** *In the rank 2 case the diagonal has the following explicit form*

$$\Delta_{(1,1)} = \hbar^{-n} \left( \frac{\det N_\lambda^-}{\det T^{1/2}} \right)^{1/2} \otimes \Lambda^\bullet(N_\lambda^-)$$

We're specifically interested in the rank 2 case, and splitting it as a product of Hilbert Schemes. In this case we have the identification of the splitting with the Hilbert Schemes, and thus we know that  $Stab_{(1)} \otimes Stab_{(1)}$  is just the identity.

By  $\mathcal{O}(-1)$  we denote the diagonal matrix in the torus fixed point basis of multiplication by  $\mathcal{O}(-1) \in K_T(\mathcal{M}(n, r))$ . With this notation  $\mathcal{O}(-1) \otimes 1$  denotes the diagonal matrix in the fixed point basis given by

$$(\mathcal{O}(-1) \otimes 1)|_{\lambda} = \prod_{\square \in \lambda_1} \varphi_{\lambda_1, \lambda_2}(\square)$$

**Proposition 4.3.** *We have the following limit of  $\mathcal{O}(-1)$  in rank 2*

$$\lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \mathcal{O}(-1) a^{2n} = \hbar^n \{ \hbar^{n_2} \mathcal{O}(-1) \otimes 1 \}$$

*Proof.* The fiber of the tautological bundle  $\mathcal{O}(-1)$  at a fixed point  $\lambda = (\lambda_1, \lambda_2)$  has weights  $\prod_{(i,j) \in \lambda_k} t_1^{j-1} t_2^{i-1} a_k^{-1}$ . The rank 2 tautological bundle is a tensor product of lower rank bundles that sum to 2, and split as  $\mathcal{O}(-1) = \mathcal{O}(-1) \otimes a^{-n_1} \mathcal{O}(-1)$ . The operator  $\Delta_{(1,1)}$  in rank 2 in the basis of fixed points is expressed as

$$\Delta_{(1,1)} = \hbar^{-n} \left( \frac{\det N_{\lambda}^{-}}{\det T^{1/2}} \right)^{1/2} \otimes \Lambda^{\bullet}(N_{\lambda}^{-})$$

where  $T^{1/2} = \hbar \mathcal{W}^* \otimes \mathcal{V}$ . In particular writing out the expression for the half tangent plane in the basis of fixed points we have

$$\begin{aligned} \hbar \mathcal{W}^* \otimes \mathcal{V} &= \hbar (V_1 + a^{-1} V_2) \left( \frac{1}{u_1} + \frac{a}{u_2} \right) \\ &= \hbar \left( \sum_{\lambda_1} (t_1^{j-1} t_2^{i-1} + a t_1^{j-1} t_2^{i-1}) + \sum_{\lambda_2} (t_1^{j-1} t_2^{i-1} + a^{-1} t_1^{j-1} t_2^{i-1}) \right) \end{aligned}$$

The repelling part of this half tangent space is

$$T_{<0}^{1/2} = \hbar \sum_{\lambda_2} a^{-1} t_1^{j-1} t_2^{i-1}$$

Thus we see that

$$\det(T_{<0}^{1/2}) = \left( \frac{\hbar^{n_2}}{a^{n_2}} 1 \otimes \mathcal{O}(-1) \right)$$

Giving us

$$\frac{1}{\det(T_{<0}^{1/2})} = \left( \frac{\hbar^{n_2}}{a^{n_2}} 1 \otimes \mathcal{O}(1) \right)$$

The repelling weights have the following expression

$$\Lambda^\bullet(N_\lambda^-) = \prod_{w \in N_\lambda^-} (1 - w^{-1})$$

As these weights are proportional to  $a$ :  $w \sim a^{-1}$ , we get that in the limit

$$\lim_{a \rightarrow 0} \Lambda^\bullet(N_\lambda^-) = 1$$

The diagonal of the stable envelope has the following expression in the basis of fixed points

$$\Delta_{(1,1)}^{-1} = \hbar^n \frac{\Lambda^\bullet(N_\lambda^-)}{\det(T_{<0}^{1/2})} = \hbar^{-n} \Lambda^\bullet(N_\lambda^-) \left( \frac{\hbar^{n_2}}{a^{n_2}} 1 \otimes \mathcal{O}(1) \right)$$

Combining these, and simplifying we see

$$\begin{aligned} \Delta_{(1,1)}^{-1} \mathcal{O}(-1) a^n &= \hbar^n \Lambda^\bullet(N_\lambda^-) \left( \frac{\hbar^{n_2}}{a^{n_2}} 1 \otimes \mathcal{O}(1) \right) (\mathcal{O}(-1) \otimes a^{-n_1} \mathcal{O}(-1)) a^n \\ &= \hbar^n \Lambda^\bullet(N_\lambda^-) \left( \frac{\hbar^{n_2}}{a^n} \mathcal{O}(-1) \otimes 1 \right) a^n \\ &= \hbar^n \Lambda^\bullet(N_\lambda^-) (\hbar^{n_2} \mathcal{O}(-1) \otimes 1) \end{aligned}$$

Taking the limit  $a \rightarrow 0$  we arrive at the desired expression

$$\lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \mathcal{O}(-1) a^n = \hbar^n (\hbar^{n_2} \mathcal{O}(-1) \otimes 1)$$

□

**Proposition 4.4.** *We have that that limit of the Stable envelope when canceled with the diagonal terms gives the wall 0  $R$ -matrix.*

$$\lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \cdot \text{Stab}_{(2)} = R_{w_0}^+$$

*Proof.* By section 2.3.3 of [7] we can write the Stable envelope as a product of  $R$ -matrices, as we fixed the positive chamber we have

$$\text{Stab}_{(2)} = \text{Stab}_{+, \infty} \cdots R_{w_2}^+ R_{w_1}^+ R_{w_0}^+$$

By definition  $\text{Stab}_{+, \infty} = \Delta_{(1,1)}$  (section 2.3.3 of [7]). Thus, multiplying by the inverse of  $\Delta_{(1,1)}$  leaves us with a product of wall  $R$ -matrices.

By equation (31) of [7] the wall  $R$ -matrices have 1 on the diagonal, and off diagonal terms proportional to  $a$ . In the limit of  $a \rightarrow 0$  all nonzero wall  $R$ -matrices go to 1, except the zeroth wall  $R_{w_0}^+$ . □

Let  $J(z)$  denote the fusion operator [7], which is expressed as the following sum:

$$J(z) = \exp \left( - \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right)$$

**Proposition 4.5.** *The wall 0 R-matrix is equal to the fusion operator at 0.*

$$R_{w_0}^+ = J(0)$$

*Proof.* See proposition 8 of [7]. □

**Corollary 4.6.** *We have the following limit of the stable envelope:*

$$\lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \cdot S^{(2)} = J(0)$$

*Proof.* Apply the previous two propositions. □

Let  $G$  denote the matrix of tangent weights

$$G = \frac{\mathcal{O}(-r)}{\Lambda^\bullet(T\mathcal{M}(n, r)^\vee)}$$

**Proposition 4.7.** *The matrix of tangent weights has the limit*

$$\lim_{a \rightarrow 0} G = (-1)^n G \mathcal{O}(-1) \otimes G(-\hbar)^d$$

where  $(-\hbar)^d$  denotes the scalar operator which takes value  $(-\hbar)^n$  on the Hilbert scheme of  $n$  points.

*Proof.* By definition of  $G$ , we have that  $G = \frac{\mathcal{O}(-2)}{\Lambda^\bullet(T\mathcal{M}(n, 2)^\vee)}$ . Here the tangent weights of the rank 2 tangent space further decompose as the following product:

$$\Lambda^\bullet(TX_2^\vee) = \Lambda^\bullet(TH_1^\vee) \oplus \Lambda^\bullet(TH_2^\vee) \oplus \Lambda^\bullet(N_2^+) \oplus \Lambda^\bullet(N_2^-)$$

Where  $\Lambda^\bullet(TH_1^\vee) \oplus \Lambda^\bullet(TH_2^\vee)$  are the weights of the decomposed rank 1 moduli spaces. Here  $H_1$  refers to the first Hilbert Scheme in the decomposition, and  $H_2$  refers to the second.  $\Lambda^\bullet(N_2^+), \Lambda^\bullet(N_2^-)$  are the attracting/repelling weights respectively.

The weights of  $\Lambda^\bullet(TH_1^\vee) \oplus \Lambda^\bullet(TH_2^\vee)$  have no  $a$ , terms, and the weights of the attracting/repelling bundle are proportional to  $a$ :

$$\Lambda^\bullet(N_2^\pm) \sim a^\pm$$

Multiplying  $G$  by  $\Lambda^\bullet(N_2^+)$ , and taking the limit  $a \rightarrow 0$  we see the following:

$$\begin{aligned} \Delta_{(1,1)} \Lambda^\bullet(N_2^+) G a^{2n} &= \Delta_{(1,1)} \Lambda^\bullet(N_2^+) \frac{\mathcal{O}(-2)}{\Lambda^\bullet(TX_2^\vee)} a^{2n} \\ &= \Delta_{(1,1)} \Lambda^\bullet(N_2^+) \frac{\mathcal{O}(-2)}{\Lambda^\bullet(TH_1^\vee) \oplus \Lambda^\bullet(TH_2^\vee) \oplus \Lambda^\bullet(N_2^+) \oplus \Lambda^\bullet(N_2^-)} a^{2n} \\ &= \Delta_{(1,1)} \frac{\mathcal{O}(-2)}{\Lambda^\bullet(TH_1^\vee) \oplus \Lambda^\bullet(TH_2^\vee) \oplus \Lambda^\bullet(N_2^-)} a^{2n} \end{aligned}$$

Taking the limit  $a \rightarrow 0$  and using Proposition 4.3 this becomes

$$(-1)^n \frac{1}{(-\hbar^2)^n} \hbar^d \mathcal{O}(-2) \otimes \frac{1}{\Lambda^\bullet(TH_2^\vee)}$$

The rank 1 weights are expressed as the following product of the rank 1 glue matrix and the rank 1 line bundle:

$$\Lambda^\bullet(TH_i^\vee) = \mathcal{O}(-1)G^{-1}$$

Which yields

$$(-1)^n (-\hbar^2)^{-n} \hbar^d \mathcal{O}(-2) G \mathcal{O}(1) \otimes \mathcal{O}(-1) G \mathcal{O}(1) = (-1)^n (-\hbar^2)^{-n} \hbar^d \mathcal{O}(-1) G \otimes G$$

□

**4.2. Limit of Capping Operator.** Normalizing the capping operator to be in the basis of fixed points, we have the following expression for the limit of the rank 2 capping operator

**Proposition 4.8.** *We have the following limit of the fundamental solution matrix:*

$$(4.3) \quad J(z)J(0)^{-1} \times \lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \Lambda^\bullet(N_2) \Psi^{(2)}(z) \Lambda^\bullet(N_2)^{-1} \Delta_{(1,1)}$$

$$(4.4) \quad = \Lambda^\bullet(N_1) \Psi^{(1)}(z\hbar^{-1}) \Lambda^\bullet(N_1)^{-1} \otimes \Lambda^\bullet(N_1) \Psi^{(1)}(z\hbar) \Lambda^\bullet(N_1)^{-1}$$

*Proof.* Consider the following modified capping operator

$$\tilde{\Psi}^{(r)}(z) = \Delta_{\underbrace{(1, \dots, 1)}_r}^{-1} \Lambda^\bullet(N_r) \Psi^{(r)}(z) \Lambda^\bullet(N_r)^{-1} \Delta_{(r_1, r_2)}$$

, gotten by conjugating our capping operator by the normal weights. Our equation becomes

$$J(z)J(0)^{-1} \times \lim_{a \rightarrow 0} \tilde{\Psi}^{(2)}(z) = \tilde{\Psi}^{(1)}(z\hbar^{-1}) \otimes \tilde{\Psi}^{(1)}(z\hbar)$$

Note that in rank 1  $\Delta_{(1)} \Delta_{(1)} = 1$

In [11] it was shown that the capping operator in rank  $r$  obtains the following factorization in the basis of fixed points:

$$\lim_{a \rightarrow 0} \Psi^{(r)}(z) = Y^{(r_1)(r_2)}(z) \Psi^{(r_1)}(z) \otimes \Psi^{(r_2)}(z)$$

Where

$$Y^{(r_1)(r_2)}(z) = ev^{(r_1)} \otimes ev^{(r_2)} \left( \exp \left( \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right) \right)$$

It was proven in [11] that  $Y^{(r_1)(r_2)}(z)$  is the strictly lower triangular solution of the wKZ equation in  $\text{Fock}^{(r_1)} \otimes \text{Fock}^{(r_2)}$ . The fusion operator  $J$  is expressed as the following sum:

$$J(z) = \exp \left( - \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right)$$

Evaluating at  $z = 0$  and inverting yields

$$J(0)^{-1} = \exp \left( \sum_{k=1}^{\infty} n_k K^{-k} \otimes K^k \alpha_{-k} \otimes \alpha_k \right)$$

Multiplying and simplifying:

$$\begin{aligned} J(z)J(0)^{-1} &= \exp \left( - \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right) \exp \left( \sum_{k=1}^{\infty} n_k K^{-k} \otimes K^k \alpha_{-k} \otimes \alpha_k \right) \\ &= \exp \left( \sum_{k=1}^{\infty} \left( n_k K^{-k} \otimes K^k - \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \right) \alpha_{-k} \otimes \alpha_k \right) \\ &= \exp \left( - \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{z^k - 1 K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right) \\ &= \exp \left( - \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right) \\ &= \exp \left( \sum_{k=1}^{\infty} \frac{n_k K^{-k} \otimes K^k}{1 - z^{-k} K^{-k} \otimes K^k} \alpha_{-k} \otimes \alpha_k \right) \\ &= J(z^{-1})^{-1} \end{aligned}$$

Applying the operator  $ev^{(r_1)} \otimes ev^{(r_2)}$  gives that

$$Y^{(r_1)(r_2)}(z) = ev^{(r_1)} \otimes ev^{(r_2)} (J(z)J(0)^{-1})$$

In particular, as  $Y^{(1)(1)}(z) = J(z)J(0)^{-1}$ , so our original limit becomes

$$Y^{(1)(1)}(z) \times \lim_{a \rightarrow 0} \Delta_{(1,1)}^{-1} \tilde{\Psi}^{(2)}(z) \Delta_{(1,1)} = \tilde{\Psi}^{(1)}(z\hbar^{-1}) \otimes \tilde{\Psi}^{(1)}(z\hbar)$$

Which is the same as theorem 3 in [11]

□

**4.3. Limits of Vertex Functions.** Let  $\mathcal{V}$  denote the tautological bundle of rank  $n$  on  $\mathcal{M}(n, r)$ . Let  $c_k(\mathcal{V})$  be the diagonal matrix of multiplication by  $c_k(\mathcal{V})$  in the basis of fixed points.

**Proposition 4.9.** *We have the following limits of classical chern classes*

$$(4.5) \quad \lim_{a \rightarrow 0} c_k(\mathcal{V}) = c_k(\mathcal{V}) \otimes 1$$

where  $c_k(\mathcal{V})$  on the right and left denote the matrices in the fixed point basis of rank 2 and 1 respectively.

*Proof.* The chern classes in rank 2 are given by elementary symmetric polynomials

$$c_k = e_k(x_1, \dots, x_n) = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} x_{j_1} \cdots x_{j_k}$$

Where  $n$  is the same as in  $\mathcal{M}(n, 2)$ , and the  $x_{j_i}$  are the Grothendieck roots. Restricting the Chern classes to fixed points gives an expression in terms of the weights of each box in the fixed point. For  $(\lambda_1 \lambda_2) \in \mathcal{M}(n, 2)^T$

$$c_k|_{(\lambda_1 \lambda_2)} = e_k(x_{\square} = \varphi_{\lambda_1, \lambda_2}(\square))$$

By our rank decomposition  $r = r_1 + ar_2$  we can see that the only factor that carries the  $a$  term are the boxes in the second rank 1 space. This corresponds to a limit of  $\varphi_{\lambda_1, \lambda_2}(\square) = a_{n(\square)} t_1^{y(\square)} t_2^{x(\square)}$  as follows:

$$\lim_{a \rightarrow 0} \varphi_{\lambda_1, \lambda_2}(\square) = \begin{cases} \varphi_{\lambda_1}(\square) & \square \in \lambda_1 \\ 0 & \square \in \lambda_2 \end{cases}$$

In particular for the chern classes in rank 2 means that we have a limit decomposition as  $\lim_{a \rightarrow 0} c_k|_{\lambda_1 \lambda_2} = c_k|_{\lambda_1}$ . This restriction to fixed points is equivalent to the tensor product of classical chern classes

$$\lim_{a \rightarrow 0} c_k = c_k \otimes 1$$

□

**Proposition 4.10.** *We have the following limit of vertex functions:*

$$\lim_{a \rightarrow 0} V^{c_k(\mathcal{V}), (2)}(z) = V^{c_k(\mathcal{V}), (1)}(z(-t_1 t_2 q)^{-1}) \otimes V^{1, (1)}(z(-t_1 t_2 q))$$

*Proof.* We reprove it here, but see [11] for the original proof.

Let  $T_0^{1/2}$  denote the 0th degree part of the half tangent space. Explicitly this means that for the tautological bundle  $\mathcal{V}$  restricted to a fixed point  $\lambda$ , there are no powers of  $q$  that arise

when looking at the weights in each  $\square \in \lambda$ .

Notice that the 0th degree part is exactly the tangent space to the fixed point  $\lambda$ , since the degree 0 quasimaps are the constant maps to  $\mathcal{M}(n, r)$ , meaning  $\mathcal{M}(n, r)$  itself. So we have

$$T_\lambda \mathcal{M}(n, r) = T_0^{1/2}$$

With this notation we now can write the  $G$ -character of the virtual tangent space.

$$T_{(\lambda, d_\lambda)}^{vir} \mathbf{QM}_{p_2}^d(n, r) = \frac{T^{1/2} - T_\lambda \mathcal{M}(n, r)}{q - 1} + T_\lambda \mathcal{M}(n, r)$$

The equivariant localization formula gives the following expression of the bare vertex:

$$(4.6) \quad V^{c_k(\mathcal{V}), (2)}(z) = \sum_{d=0}^{\infty} \sum_{p \in \mathbf{QM}_{p_2}^d} a(p) z^d \in K_G(\mathcal{M}(n, r))_{loc} \otimes \mathbb{Q}[[z]]$$

Where  $a(p)$  represents the contribution of a fixed point  $p$ .

Assume that in the basis (1.1) the vertex function has the expansion  $V^{\tau, (2)}(z) = \sum_{|\lambda|=n} V_\lambda^{c_k(\mathcal{V}), (2)}(z)[\lambda]$ , where  $[\lambda] \in K_G(\mathcal{M}(n, r))_{loc}$  represents the localized  $K$ -theoretic class of a fixed point  $\lambda = (\lambda_1, \lambda_2)$ . By equation 4.6 we know that the only contributing factors for the coefficient in the basis of fixed points are those associated to a fixed point  $p \in \mathbf{QM}_{p_2}^d$ . So we can write

$$V_\lambda^{c_k(\mathcal{V}), (2)}(z) = \sum_{d_\lambda} a_\lambda((\lambda, d_\lambda)) z^{|d|}$$

Explicitly writing out  $a_\lambda((\lambda, d_\lambda))$  via localization in  $K$ -theory we get the following power series

$$V_\lambda^{c_k(\mathcal{V}), (2)}(z) = \sum_{d_\lambda} \hat{a}_\lambda(T_{(\lambda, d_\lambda)}^{vir} \mathbf{QM}_{p_2}^d(n, r) - T_\lambda \mathcal{M}(n, r)) c_k(\mathcal{V}) (-1)^{|d|r} q^{\frac{-r|d|}{2}} z^{|d|}$$

Where

$$\hat{a}(x + y - z) = \frac{\hat{a}(x)\hat{a}(y)}{\hat{a}(z)}, \quad \hat{a}(x) = \frac{1}{x^{1/2} - x^{-1/2}}$$

is Okounkov's roof function, and  $T_{(\lambda, d_\lambda)}^{vir} \mathbf{QM}_{p_2}^d(n, r)$  is the  $G$ -character of the virtual tangent space. The subtraction of the tangent space to a fixed point is a normalization to ensure what our vertex function's power series expansion is of the form

$$V_\lambda^{c_k(\mathcal{V}), (2)} = 1 + O(z)$$

We want to compute  $\lim_{a \rightarrow 0} V^{\tau, (2)}(z)$  under the splitting  $r = r_1 + ar_2$ . By localization this proposition is equivalent to the following factorization of coefficients for the bare vertex

$$\lim_{a \rightarrow 0} V_{\lambda}^{c_k(\mathcal{V}), (2)}(z) = V_{\lambda_1}^{c_k(\mathcal{V}), (1)}(z(-t_1 t_2 q)) V_{\lambda_2}^{1, (1)}(z(-t_1 t_2 q)^{-1})$$

To prove this factorization, we'll investigate what happens to each term in the rank splitting at  $a \rightarrow 0$ , and notice that the factorization falls out of a computation. First, we'll investigate how the tautological bundles split in the rank decomposition: Notice that the tautological bundle  $\mathcal{V}$  for the rank 2 case splits as  $\mathcal{V}_{\lambda_1, d_1} + a\mathcal{V}_{\lambda_2, d_2}$ , and the tautological bundle of the framing splits as  $\mathcal{W} = \mathcal{W}_1 + a\mathcal{W}_2$ . Looking at each term of the half tangent space to  $\mathcal{M}(n, r)$  we can identify what happens in the limit.

First consider the terms  $\mathcal{V} \otimes \mathcal{V}^*$ . A weight  $w \in \mathcal{V} \otimes \mathcal{V}^*$  appears in  $\hat{a}_{\lambda}(T_{(\lambda, d_{\lambda})}^{\text{vir}} \mathbf{QM}_{p_2}^d(n, r))$  as an expression of the form

$$\begin{aligned} \hat{a}(-w(1 - t_1^{-1})(1 - t_2^{-1})) &= \hat{a}(wt_2^{-1} - w(t_1 t_2)^{-1} - w + wt_1^{-1}) \\ &= \frac{\hat{a}(wt_2^{-1})\hat{a}(wt_1^{-1})}{\hat{a}(w(t_1 t_2)^{-1})\hat{a}(w)} \\ &= \frac{1}{(wt_2^{-1})^{1/2} - (wt_2^{-1})^{-1/2}} \left( (w(t_1 t_2)^{-1})^{1/2} - (w(t_1 t_2)^{-1})^{-1/2} \right) \\ &\times \left( (w)^{1/2} - (w)^{-1/2} \right) \frac{1}{(wt_1^{-1})^{1/2} - (wt_1^{-1})^{-1/2}} \\ &= \frac{(w^{1/2} - w^{-1/2})(w^{1/2} t_1^{-1/2} t_2^{-1/2} - w^{-1/2} t_1^{1/2} t_2^{1/2})}{(w^{1/2} t_1^{-1/2} - w^{-1/2} t_1^{1/2})(w^{1/2} t_2^{-1/2} - w^{-1/2} t_2^{1/2})} \end{aligned}$$

If the weight  $w \in \mathcal{V}_{\lambda_i, d_i} \otimes \mathcal{V}_{\lambda_j, d_j}^*$  where  $i \neq j$  then  $w \sim a^{\pm 1}$  and hence the expression is equal to 1 in the limit  $a \rightarrow 0$ . As a result we only need to look at the terms  $\mathcal{V}_{\lambda_i, d_i} \otimes \mathcal{V}_{\lambda_j, d_j}^*$  where  $i = j$ , and by our splitting this amounts to  $\mathcal{V}_{\lambda_1, d_1} \otimes \mathcal{V}_{\lambda_1, d_1}^*$  and  $\mathcal{V}_{\lambda_2, d_2} \otimes \mathcal{V}_{\lambda_2, d_2}^*$ . Therefore the contributions of weights from  $\mathcal{V}_{\lambda_i, d_i} \otimes \mathcal{V}_{\lambda_j, d_j}^*$  in rank 2 factor to a product of rank 1 contributions.

Next, looking at the weights appearing in  $\mathcal{V}_{\lambda_i, d_i} \otimes \mathcal{W}_j$  with  $i \neq j$  first consider the terms where  $i = 1, j = 2$ . The terms which appear in  $T^{1/2}$  are of the form  $\mathcal{V}_{\lambda_1, d_1} \otimes \mathcal{W}_2^* + \hbar^{-1} \mathcal{V}_{\lambda_1, d_1}^* \otimes \mathcal{W}_2$ . These in turn contribute the following to the virtual tangent bundle

$$\frac{\mathcal{V}_{\lambda_1, d_1} \otimes \mathcal{W}_2^* + \hbar^{-1} \mathcal{V}_{\lambda_1, d_1}^* \otimes \mathcal{W}_2 - (\mathcal{V}_{\lambda_1, 0} \otimes \mathcal{W}_2^* + \hbar^{-1} \mathcal{V}_{\lambda_1, 0}^* \otimes \mathcal{W}_2)}{q - 1}$$

Notice that the extra  $T_\lambda \mathcal{M}(n, r)$  is removed due to localization in  $K$ -theory. The weights of  $\mathcal{V}_{\lambda_1, 0}$  are of the form  $\sum_{\square \in \lambda_1} \varphi_{\lambda_1}(\square)$ . Writing out the above expression in terms of weights, and simplifying we get

$$\begin{aligned}
& \frac{\mathcal{V}_{\lambda_1, d_1} \otimes \mathcal{W}_2^* + \hbar^{-1} \mathcal{V}_{\lambda_1, d_1}^* \otimes \mathcal{W}_2 - (\mathcal{V}_{\lambda_1, 0} \otimes \mathcal{W}_2^* + \hbar^{-1} \mathcal{V}_{\lambda_1, 0}^* \otimes \mathcal{W}_2)}{q-1} \\
&= \frac{a^{-1} u_2^{-1} (\sum_{\square \in \lambda_1} \varphi_{\lambda_1}(\square) q^{d_\square} - \varphi_{\lambda_1}(\square))}{q-1} + \frac{a u_2 (\sum_{\square \in \lambda_1} \varphi_{\lambda_1}^{-1}(\square) q^{-d_\square} - \varphi_{\lambda_1}^{-1}(\square))}{(q-1)\hbar} \\
&= a^{-1} u_2^{-1} \left( \frac{(\sum_{\square \in \lambda_1} \varphi_{\lambda_1}(\square) q^{d_\square} - \varphi_{\lambda_1}(\square))}{q-1} \right) + \hbar^{-1} a u_2 \left( \frac{(\sum_{\square \in \lambda_1} \varphi_{\lambda_1}^{-1}(\square) q^{-d_\square} - \varphi_{\lambda_1}^{-1}(\square))}{(q-1)\hbar} \right) \\
&= a^{-1} u_2^{-1} \left( \frac{\sum_{\square \in \lambda_1} \varphi_{\lambda_1}(\square) (q^{d_\square} - 1)}{q-1} \right) \\
&+ \hbar^{-1} a u_2 \left( \frac{\sum_{\square \in \lambda_1} \varphi_{\lambda_1}^{-1}(\square) (q^{-d_\square} - 1)}{(q-1)\hbar} \right) \\
&= a^{-1} u_2^{-1} \left( \sum_{\square \in \lambda_1} \varphi_{\lambda_1}(\square) \sum_{i=0}^{d_\square-1} q^i \right) + \hbar^{-1} a u_2 \left( \hbar^{-1} q^{-1} \sum_{\square \in \lambda_1} \varphi_{\lambda_1}^{-1}(\square) \sum_{i=0}^{d_\square-1} q^{-i} \right) \\
&= \sum_{\square \in \lambda_1} \left( \frac{\varphi_{\lambda_1}(\square)}{a u_2} \left( \sum_{i=0}^{d_\square-1} q^i \right) + \frac{a u_2}{\hbar \varphi_{\lambda_1}(\square)} \left( \frac{1}{\hbar q} \sum_{i=0}^{d_\square-1} q^{-i} \right) \right)
\end{aligned}$$

The contribution of this to the bare vertex is

$$\begin{aligned}
& \prod_{\square \in \lambda_1} \prod_{i=0}^{d_\square-1} \left( \frac{\hat{a} \left( \frac{\varphi_{\lambda_1}(\square)}{a u_2} q^i \right)}{\hat{a} \left( \frac{a u_2}{\hbar^2 q^{i+1} \varphi_{\lambda_1}(\square)} \right)} \right) \\
&= \prod_{\square \in \lambda_1} \prod_{i=0}^{d_\square-1} \frac{\left( \frac{a u_2}{\hbar^2 q^{i+1} \varphi_{\lambda_1}(\square)} \right)^{1/2} - \left( \frac{a u_2}{\hbar^2 q^{i+1} \varphi_{\lambda_1}(\square)} \right)^{-1/2}}{\left( \left( \frac{\varphi_{\lambda_1}(\square)}{a u_2} q^i \right)^{1/2} - \left( \frac{\varphi_{\lambda_1}(\square)}{a u_2} q^i \right)^{-1/2} \right)}
\end{aligned}$$

In the limit  $a \rightarrow 0$  we get the following product which contributes to the vertex in the first component:

$$\prod_{\square \in \lambda_1} \prod_{i=0}^{d_{\square}-1} (-\hbar^{1/2} q^{1/2}) = (-\hbar^{1/2} q^{1/2})^{|\lambda_1|}$$

Similarly computing for  $i = 2, j = 1$  results in a contribution of  $(-\hbar^{1/2} q^{1/2})^{-|d_2|}$ . If  $i = j$ , then the terms that appear have no  $a$  term, and thus in the limit the terms will factor as  $\mathcal{V}_{\lambda_i, \bar{\lambda}_i} \otimes \mathcal{W}_i$  for  $i = 1, 2$ . Using the previous proposition, we know that in the limit the descendent corresponding to  $c_k(\mathcal{V})$  factors as  $c_k \otimes 1$ . Combining all these will yield the following limit:

$$\begin{aligned} \lim_{a \rightarrow 0} V_{(\lambda_1, \lambda_2)}^{\tau, (2)}(z) &= \sum_{d_{\lambda_1}, d_{\lambda_2}} \hat{a}_{\lambda}(T_{(\lambda_1, d_{\lambda_1})}^{\text{vir}} \mathbf{QM}_{p_2}^{d_1}(n_1, r_1)) \hat{a}_{\lambda}(T_{(\lambda_2, d_{\lambda_2})}^{\text{vir}} \mathbf{QM}_{p_2}^{d_2}(n_2, r_2)) \\ &\times (c_k \otimes 1) (-\hbar^{1/2} q^{1/2})^{|\lambda_1|} (-\hbar^{1/2} q^{1/2})^{-|d_2|} (-1)^{|d|r} q^{\frac{-r|d|}{2}} z^{|\lambda_1|+d_2} \\ &= V_{\lambda_1}^{\tau, (1)}(z(-t_1 t_2 q)) V_{\lambda_2}^{(1)}(z(-t_1 t_2 q)^{-1}) \end{aligned}$$

□

**4.4. A relation between capped descendants.** Let us consider the following descendent:

$$(4.7) \quad \tau_n(u) = (u + x_1) \dots (u + x_n) = \sum_{i=0}^n u^{n-i} c_i(\mathcal{V})$$

where  $c_i(\mathcal{V})$  denotes the  $i$ -th Chern class of the tautological bundle  $\mathcal{V}$ , and  $x_1, \dots, x_n$  the Chern roots of  $\mathcal{V}$  over  $\mathcal{M}(n, r)$ . Explicitly, it is  $i$ -th symmetric polynomial in the Chern roots. We will also need the dual generating function

$$(4.8) \quad \bar{\tau}_n(u) = (1 + ux_1^{-1}) \dots (1 + ux_n^{-1}) = \sum_{i=0}^n u^i c_i(\mathcal{V}^*)$$

where  $c_i(\mathcal{V}^*)$  is the  $i$ -th Chern class of the dual bundle  $\mathcal{V}^*$ . We note that  $\mathcal{O}(1) = x_1 \dots x_n$  and therefore:

$$\tau_n(u) = \mathcal{O}(1) \bar{\tau}_n(u)$$

We recall that the operator  $\mathbf{M}(z)$  is the operator of quantum multiplication by  $\mathcal{O}(1)$  and the above identity can be upgraded to capped descendent vertices:

$$(4.9) \quad \hat{\mathbf{V}}^{\tau_n(u)}(z/q^2) = \mathbf{M}(-z/q) \hat{\mathbf{V}}^{\bar{\tau}_n(u)}(z)$$

Our goal is to compute the generating function

$$F(z, t_1, t_2, q, u, y) = \sum_{n=0}^{\infty} \hat{\mathbf{V}}^{\bar{\tau}_n(u)}(z) y^n$$

**4.5. Limiting linear equation.** Now we look at the descendent equation in rank 2. Let  $X_r = \mathcal{M}(n, r)$ , then using theorem 3.4 we have that the capped vertex function with chern class descendents are independent of  $z$  and thus we have:

$$(4.10) \quad c_k(\mathcal{V}^*) = \Lambda^\bullet(TX_2^\vee)\Psi(z)\Lambda^\bullet(TX_2^\vee)^{-1}V_O^{(c_k(\mathcal{V}^*))}(z)$$

From our formula for the matrix of tangent weights, we know that  $\Lambda^\bullet(TX_2^\vee) = \Lambda^\bullet(TH^\vee) \oplus \Lambda^\bullet(N_2^-)$ , where  $\Lambda^\bullet(TH^\vee)$  are the weights of the Hilbert Schemes in the decomposition of rank 2.  $\Lambda^\bullet(N_2^-)$  are the weights of the repelling bundle in rank 2.

Expressing each of these factors as their components we get the following expression of the capped vertex:

$$\widehat{V}^{(c_k(\mathcal{V}^*))}(z) = \Lambda^\bullet(TH^\vee)\Lambda^\bullet(N_2)\Psi^{(2)}(z)\Lambda^\bullet(N_2)^{-1}\Lambda^\bullet(TH^\vee)^{-1}V_O^{(c_k(\mathcal{V}^*))}(z)$$

taking the  $a = 0$  limit of this equation, by Propositions 4.3-4.9 we obtain:

$$\begin{aligned} c_k(\mathcal{V}^*) \otimes 1 &= \Lambda^\bullet(TH^\vee) \times (J(0)J(-z)^{-1})^{-1} \times \Psi^{(1)}(-z\hbar^{-1}q^{-2}) \otimes \Psi^{(1)}(-z\hbar q^{-2}) \\ &\quad \times \Lambda^\bullet(TH^\vee)^{-1} \times V_O^{(c_k(\mathcal{V}^*)),(1)}(-z(\hbar q)^{-1}) \otimes V_O^{1,(1)}(-z(\hbar q)^1) \end{aligned}$$

We'll work to show that the capped vertex in rank 2 decomposes to a tensor product of capped vertices in rank 1

**4.5.1. First component.** In the first component of this limit we see the following version of the qde in rank 1:

$$(J(0)J(-z)^{-1})^{-1}\Psi^{(1)}(-z\hbar^{-1}q^{-2})\frac{\mathcal{O}(1)}{\Lambda^\bullet(TH^\vee)}V_O^{(c_k(\mathcal{V}^*)),(1)}(-z/\hbar)$$

Performing a change of variables with  $Q = -z\hbar^{-1}$ , and noticing that  $\Lambda^\bullet(N_1)^{-1}\Lambda^\bullet(TH^\vee)^{-1} = \Lambda^\bullet(TX_1^\vee)^{-1}$  we see the following expression:

$$\begin{aligned} (J(0)J(-z)^{-1})^{-1}\Psi^{(1)}\left(-\frac{Q}{q^2}\right)\frac{\mathcal{O}(1)}{\Lambda^\bullet(TX_1^\vee)}V_O^{(c_k(\mathcal{V}^*)),(1)}(Q) &= (J(0)J(-z)^{-1})^{-1}\Psi^{(1)}\left(-\frac{Q}{q^2}\right)GV_O^{(c_k(\mathcal{V}^*)),(1)}(Q) \\ &= (J(0)J(-z)^{-1})^{-1}G\widehat{V}_O^{(c_k(\mathcal{V}^*)),(1)}(Q) \end{aligned}$$

**4.5.2. Second component.** In the second component we have the following product:

$$\Psi^{(1)}(-z\hbar q^{-2})\frac{1}{\Lambda^\bullet(TX_1^\vee)}V_O^{1,(1)}(-z\hbar q)$$

By a similar change of variables as in the first component we set  $Q = -z\hbar q$  so that the product becomes

$$\begin{aligned}
 \Psi^{(1)}\left(-\frac{Q}{q^3}\right) \frac{1}{\Lambda^\bullet(TX_1^\vee)} V_O^{1,(1)}(Q) &= M^{(1)}\left(-\frac{Q}{q}\right) \Psi^{(1)}\left(-\frac{Q}{q}\right) G V_O^{1,(1)}(Q) \\
 &= M^{(1)}\left(-\frac{Q}{q}\right) G \widehat{V}_O^{1,(1)}(Q) \\
 &= M^{(1)}\left(-\frac{Q}{q}\right) G
 \end{aligned}$$

Where we've used the qde formula 4.2 in the first line, and that the capped vertex with descendent 1 is trivial in rank 1.

Thus the capped vertex in the limit  $a \rightarrow 0$  appears as the following tensor:

$$c_k(\mathcal{V}^*) \otimes 1 = G \widehat{V}^{(c_k(\mathcal{V}^*)),(1)}(Q) \otimes M^{(1)}\left(-\frac{Q}{q}\right) G$$

Which means that we can solve

$$(J(z)J(0)^{-1})^{-1} \left( \widehat{V}^{(c_k(\mathcal{V}^*))}(z) \otimes 1 \right) = c_k(\mathcal{V}^*) \otimes 1$$

For  $\widehat{V}^{(c_k(\mathcal{V}^*))}(z)$ . In particular, the right hand side is independent of  $z$ .

## 5. DERIVATION OF DESCENDANTS FORMULA

Let  $H_\lambda$  denote the Macdonald polynomial in Haiman's normalization. Our parameters are related to  $q, t$  via the relation  $q = 1/t_1^2, t = 1/t_2^2$ . Then there is a generating function relating normalized  $H_\lambda$  to a sum of the power sums:

**Proposition 5.1.**

$$\sum_{\lambda} \frac{H_{\lambda}}{\Lambda^\bullet(T_{\lambda} \text{Hilb}^{|\lambda|}(\mathbb{C}^2))} y^{|\lambda|} = \exp\left(\sum_{k=1}^{\infty} \frac{y^k \hbar^{2k} p_k}{k(1-t_1^{2k})(1-t_2^{2k})}\right)$$

*Proof.* This is the kernel identity for Macdonald scalar product. See [4], with  $Y = 1$ . □

**Proposition 5.2.** *We have the following generating function given by 4.7:*

$$(5.1) \quad \sum_{n=0}^{\infty} \bar{\tau}_n(u) y^n = \exp\left(\sum_{k=1}^{\infty} \frac{y^k p_k}{k(1-t_1^{2k})(1-t_2^{2k})} \hbar^{2k} (1 - (-u)^k)\right)$$

*Proof.* See corollary 6.4 in [4] □

Rescaling  $p_i \rightarrow p_i/a_i$  and sending  $a \rightarrow \infty$ , from this proposition we obtain the following twisted version of the above normalized Macdonald generating series:

$$(5.2) \quad \sum_{n=0}^{\infty} \mathcal{O}_{\text{Hilb}^n(\mathbb{C}^2)} y^n = \exp \left( \sum_{k=1}^{\infty} \frac{(-1)^k y^k \hbar^{2k} p_k}{k(1-t_1^{2k})(1-t_2^{2k})} \right)$$

From these formulas we have

$$\sum_{n_1, n_2=0}^{\infty} (\tau_{n_1}(u) \otimes \mathcal{O}_{\text{Hilb}^{n_2}(\mathbb{C}^2)}) y^{n_1+n_2} = \exp \left( \sum_{k=1}^{\infty} \frac{y^k \hbar^{2k} ((1 - (-u)^k) p_k^{(1)} + (-1)^k p_k^{(2)})}{k(1-t_1^{2k})(1-t_2^{2k})} \right)$$

Next, recall that the product of the operator  $J(z)$  and normalized part of the transition matrix from the basis of Macdonald polynomials to the basis of power sums acts by (section 7 of [7])

$$J(z)J(0)^{-1} : p_k^{(1)} \rightarrow p_k^{(1)}, \quad p_k^{(2)} \rightarrow p_k^{(2)} + (-1)^k z^k \hbar^k \frac{\hbar^k - \hbar^{-k}}{1 - z^k} p_k^{(1)}$$

Thus, applying this operator to (5) we obtain

$$\begin{aligned} & J(z)J(0)^{-1} \sum_{n_1, n_2=0}^{\infty} (\tau_{n_1}(u) \otimes \mathcal{O}_{\text{Hilb}^{n_2}(\mathbb{C}^2)}) y^{n_1+n_2} = \\ & = \exp \left( \sum_{k=1}^{\infty} \frac{y^k}{k(1-t_1^{2k})(1-t_2^{2k})} \left( (1 - (-u)^k) \hbar^{2k} p_k^{(1)} + (-1)^k \hbar^{2k} p_k^{(2)} + z^k \hbar^k \frac{\hbar^k - \hbar^{-k}}{1 - z^k} p_k^{(1)} \right) \right) \end{aligned}$$

Taking the first component in the tensor product corresponds to  $p_k^{(2)} = 0$ , which gives:

$$\exp \left( \sum_{k=1}^{\infty} \frac{y^k}{k(1-t_1^{2k})(1-t_2^{2k})} \left( (1 - (-u)^k) \hbar^{2k} p_k^{(1)} + \hbar^k z^k \frac{\hbar^k - \hbar^{-k}}{1 - z^k} p_k^{(1)} \right) \right)$$

We thus proved the following theorem:

**Theorem 5.3.** *The generating function for the capped descendent vectors has the following form:*

$$(5.3) \quad \begin{aligned} & F(zq^2/\hbar, t_1, t_2, q, u, y) = \\ & \exp \left( \sum_{k=1}^{\infty} \frac{y^k}{k(1-t_1^{2k})(1-t_2^{2k})} \left( (1 - (-u)^k) \hbar^{2k} p_k^{(1)} + z^k \hbar^k \frac{\hbar^k - \hbar^{-k}}{1 - z^k} p_k^{(1)} \right) \right) \end{aligned}$$

**Corollary 5.4.** *The generating function for the capped vertices  $\hat{V}^{(c_k)}(-zt_1 t_2/q)$  does not depend on  $q$ .*

**Corollary 5.5.**  *$\hat{V}^\tau(z)$  is a rational function in  $\mathbb{Q}(t_1, t_2, z, q, u_1, u_2)$*

## REFERENCES

- [1] Mark Haiman. *Hilbert schemes, polygraphs, and the Macdonald positivity conjecture*. 2000. arXiv: [math/0010246](https://arxiv.org/abs/math/0010246) [[math.AG](#)].
- [2] I.G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford classic texts in the physical sciences. Clarendon Press, 1998. ISBN: 9780198504504. URL: <https://books.google.com/books>
- [3] Davesh Maulik and Andrei Okounkov. *Quantum Groups and Quantum Cohomology*. 2018. arXiv: [1211.1287](https://arxiv.org/abs/1211.1287) [[math.AG](#)].
- [4] Anton Mellit. *Plethystic identities and mixed Hodge structures of character varieties*. 2016. arXiv: [1603.00193](https://arxiv.org/abs/1603.00193) [[math.AG](#)].
- [5] Hiraku Nakajima. “Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras”. In: *Duke Mathematical Journal* 76.2 (1994), pp. 365–416. DOI: [10.1215/S0012-7094-94-07613-8](https://doi.org/10.1215/S0012-7094-94-07613-8). URL: <https://doi.org/10.1215/S0012-7094-94-07613-8>.
- [6] Andrei Okounkov. *Lectures on K-theoretic computations in enumerative geometry*. 2017. arXiv: [1512.07363](https://arxiv.org/abs/1512.07363) [[math.AG](#)].
- [7] Andrei Okounkov and Andrey Smirnov. *Quantum difference equation for Nakajima varieties*. 2016. DOI: [10.48550/ARXIV.1602.09007](https://arxiv.org/abs/10.48550/ARXIV.1602.09007). URL: <https://arxiv.org/abs/1602.09007>.
- [8] R. Pandharipande and A. Pixton. *Descendent theory for stable pairs on toric 3-folds*. 2012. arXiv: [1011.4054](https://arxiv.org/abs/1011.4054) [[math.AG](#)].
- [9] R. Pandharipande and A. Pixton. “Descendents on local curves: rationality”. In: *Compositio Mathematica* 149.1 (Nov. 2012), pp. 81–124. ISSN: 1570-5846. DOI: [10.1112/s0010437x12000498](https://doi.org/10.1112/s0010437x12000498). URL: <http://dx.doi.org/10.1112/S0010437X12000498>.
- [10] Olivier Schiffmann and Eric Vasserot. “The elliptic Hall algebra and the K-theory of the Hilbert scheme of  $A^2$ ”. In: *Duke Mathematical Journal* 162.2 (Feb. 2013). ISSN: 0012-7094. DOI: [10.1215/00127094-1961849](https://doi.org/10.1215/00127094-1961849). URL: <http://dx.doi.org/10.1215/00127094-1961849>.
- [11] Andrey Smirnov. *Rationality of capped descendent vertex in K-theory*. 2016. arXiv: [1612.01048](https://arxiv.org/abs/1612.01048) [[math.AG](#)].