

A CHARACTERIZATION OF THE L^2 -RANGE OF THE GENERALIZED SPECTRAL PROJECTIONS RELATED TO THE HODGE-DE RHAM LAPLACIAN

ABDELHAMID BOUSSEJRA AND KHALID KOUFANY

ABSTRACT. In this paper, we establish a characterization of the L^2 -range of generalized spectral projections on the bundle of differential forms over the real hyperbolic space $H^n(\mathbb{R})$. As an intermediate result, we obtain a characterization of the L^2 -range of the Poisson transform on the bundle of differential forms on the boundary $\partial H^n(\mathbb{R})$. This results confirm a conjecture by Strichartz regarding differential forms.

CONTENTS

1. Introduction	1
2. Notations and Preliminaries	4
2.1. Differential forms on $H^n(\mathbb{R})$ and $\partial H^n(\mathbb{R})$	4
2.2. Fourier-Helgason transform	6
3. The Helgason Fourier restriction theorem	7
4. Poisson transforms on differential forms	9
5. Announcement of the main theorems	13
6. Asymptotic expansion of τ -spherical functions	15
7. Proof of Theorem 5.2 – Strichartz’s conjecture for Poisson transforms	23
8. Proof of Theorem 5.3 – Inversion formula for the Poisson transform	25
9. Proof of Theorem 5.1 – Strichartz’s conjecture for spectral projections	27
10. Appendix	28
References	31

1. INTRODUCTION

Let $T_{\mathbb{C}}^*H^n(\mathbb{R})$ be the complexified cotangent bundle of the real hyperbolic space $H^n(\mathbb{R})$ and let $\wedge^p H^n(\mathbb{R}) := \wedge^p T_{\mathbb{C}}^*H^n(\mathbb{R})$ be the p -th exterior vector bundle ($0 \leq p \leq n$). Its sections are differential p -forms on $H^n(\mathbb{R})$. We write $C^\infty(\wedge^p H^n(\mathbb{R}))$, $C_c^\infty(\wedge^p H^n(\mathbb{R}))$ and $L^2(\wedge^p H^n(\mathbb{R}))$ for smooth sections, smooth sections with compact support and square-integrable sections, respectively. Consider the exterior differentiation operator $d: C^\infty(\wedge^p H^n(\mathbb{R})) \rightarrow C^\infty(\wedge^{p+1} H^n(\mathbb{R}))$ and the co-differentiation operator $\delta: C^\infty(\wedge^{p+1} H^n(\mathbb{R})) \rightarrow C^\infty(\wedge^p H^n(\mathbb{R}))$.

Date: August 22, 2024.

Keywords : Strichartz’s conjecture, differential forms bundle, Poisson transform, Helgason-Fourier transform, generalized spectral projections.

2010 AMS Classification : 43A90, 43A85, 58J50

Let $\Delta = dd + \delta d$ be the Hodge-de Rham Laplacian which is a self-adjoint operator on $L^2(\bigwedge^p H^n(\mathbb{R}))$. According to the Hodge-de Rham decomposition of $L^2(\bigwedge^p H^n(\mathbb{R}))$ any $f \in L^2(\bigwedge^p H^n(\mathbb{R}))$ can be decomposed into a sum

$$f = f_d + f_\delta + f_0, \quad (1.1)$$

where $f_d \in \overline{dC_c^\infty(\bigwedge^{p-1} H^n(\mathbb{R}))}$, $f_\delta \in \overline{\delta C_c^\infty(\bigwedge^{p+1} H^n(\mathbb{R}))}$, and f_0 is an L^2 -harmonic form. The harmonic component f_0 is non zero only when n is even and $p = n/2$. The orthogonal decomposition (1.1) yields to three projectors (de-Rham projectors). Following Strichartz we are interested in the first two. More precisely, in [26], Strichartz initiated the spectral theory of the Hodge-de Rham Laplacian Δ by considering the decomposition of forms into exact differentials (d) and coexact differentials (δ). In the following, we will briefly outline his approach. By using a local presentation of p -forms, Strichartz found explicit expressions for operators \mathcal{Q}_λ^d and $\mathcal{Q}_\lambda^\delta$ which satisfy

$$\begin{aligned} f_d &= \int_0^\infty \mathcal{Q}_\lambda^d f \, d\lambda, \\ f_\delta &= \int_0^\infty \mathcal{Q}_\lambda^\delta f \, d\lambda, \end{aligned}$$

with

$$\begin{aligned} \Delta \mathcal{Q}_\lambda^d f &= -(\lambda^2 + (\rho - p + 1)^2) \mathcal{Q}_\lambda^d f, \\ \Delta \mathcal{Q}_\lambda^\delta f &= -(\lambda^2 + (\rho - p)^2) \mathcal{Q}_\lambda^\delta f. \end{aligned}$$

He noticed that for a dense set of f and for any point z ,

$$\|f_\delta\|_2^2 = \pi \lim_{r \rightarrow \infty} \int_0^\infty \frac{1}{r} \int_{B_r(z)} \langle \mathcal{Q}_\lambda^\delta f, \mathcal{Q}_\lambda^\delta f \rangle_x \, dx \, d\lambda, \quad (1.2)$$

and

$$\|f_d\|_2^2 = \pi \lim_{r \rightarrow \infty} \int_0^\infty \frac{1}{r} \int_{B_r(z)} \langle \mathcal{Q}_\lambda^d f, \mathcal{Q}_\lambda^d f \rangle_x \, dx \, d\lambda. \quad (1.3)$$

Then he made the following conjecture, *loc. cit.* [26, page 146]: “It is natural to conjecture that (1.2) and (1.3) continue to hold for all L^2 , and conversely, the finiteness of (1.2) and (1.3) should characterize the spectral decomposition of L^2 forms”.

In this paper, we focus on proving the aforementioned conjecture. Our result will extend the “classical” Strichartz’s conjecture (refer to [26, Conjecture 4.5, Conjecture 4.6]) from scalar-valued functions on Riemannian symmetric spaces of noncompact type to the differential forms bundle on $H^n(\mathbb{R})$. We should mention that [26, Conjecture 4.5, Conjecture 4.6] has been investigated for rank one symmetric spaces (see e.g. [8–10, 12, 20]) and proved for any symmetric space of noncompact type in [22].

The method employed by Strichartz for (1.2) and (1.3) has the disadvantage of being local: he realized $H^n(\mathbb{R})$ in \mathbb{R}^{n+1} and used local harmonic analysis to define \mathcal{Q}_λ^d and $\mathcal{Q}_\lambda^\delta$, and to prove (1.2) and (1.3) through direct computations.

Our approach for handling this conjecture is based on representation theory. We will view $H^n(\mathbb{R})$ as the rank one symmetric space G/K where $G = \mathrm{SO}_0(n, 1)$ and $K = \mathrm{SO}(n)$ its maximal compact subgroup. As working in the setting of the exterior bundle involves some difficulties, we shall identify $\bigwedge^p H^n(\mathbb{R})$ with the G -homogeneous vector bundle $G \times_K \bigwedge^p \mathbb{C}^n$ and subsequently use representation theory, in particular the Plancherel decomposition of $L^2(G \times_K \bigwedge^p \mathbb{C}^n)$. To be more precise, let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition and let

$\mathfrak{a} \subset \mathfrak{p}$ be a maximal abelian subspace of \mathfrak{p} . Let $\tau = \tau_p = \bigwedge^p Ad^*$ ($0 \leq p \leq n$) be the p -th exterior product of the coadjoint representation of K on $V_\tau = \bigwedge^p \mathfrak{p}_\mathbb{C}^* \simeq \bigwedge^p \mathbb{C}^n$ (Notice that τ_p is irreducible if $p \neq \frac{n}{2}$). Then as G -homogeneous vector bundles we have $\bigwedge^p T_\mathbb{C}^* H^n(\mathbb{R}) = G \times_K V_\tau$. So we shall identify the space $L^2(\bigwedge^p H^n(\mathbb{R}))$ with the space $L^2(G, \tau)$ of square integrable function $f : G \rightarrow V_\tau$, such that $f(gk) = \tau(k)^{-1} f(g)$ for $g \in G$ and $k \in K$. According to the Plancherel Theorem, the space $L^2(G, \tau)$ splits into a direct sum of a continuous part $L^2(G, \tau)_{\text{cont}}$ and possibly of a discrete part $L^2(G, \tau)_{\text{disc}}$. The latter is non-trivial if $p = \frac{n}{2}$.

Due to the Helgason Fourier inversion formula (see e.g. [25]), any $f \in L^2(G, \tau)$ (or in the continuous part for $p = \frac{n}{2}$) can be written as

$$f = \sum_{\sigma \in \widehat{M}(\tau)} \int_0^\infty \mathcal{Q}_{\sigma, \lambda}^\tau f \, d\lambda,$$

where $\widehat{M}(\tau)$ is the set of unitary irreducible representations σ of $M = \text{SO}(n-1)$ that occur in the restriction $\tau|_M$ and $\mathcal{Q}_{\sigma, \lambda}^\tau$ is the generalized partial spectral projection defined for $f \in L^2(G, \tau)$ by

$$\mathcal{Q}_{\sigma, \lambda}^\tau f(g) = \nu_\sigma(\lambda) (\mathcal{P}_{\sigma, \lambda}^\tau (\mathcal{F}_{\sigma, \lambda}^\tau f(\cdot)))(g), \quad \text{for a.e. } \lambda \in \mathfrak{a}^*.$$

Above, $\mathcal{F}_{\sigma, \lambda}^\tau$ is the partial Helgason Fourier transform (see (2.4)), $\mathcal{P}_{\sigma, \lambda}^\tau$ is the Poisson transform (see (4.1)) and $\nu_\sigma(\lambda)$ is the Plancherel density. Moreover, $\mathcal{Q}_{\sigma, \lambda}^\tau f$ is (for a.e. $\lambda \in (0, \infty)$) a joint eigenform of the algebra of G -invariant differential operators acting on smooth sections of $G \times_K V_\tau$. Therefore the family of operators $\mathcal{Q}_\lambda^\tau := (\mathcal{Q}_{\sigma, \lambda}^\tau; \sigma \in \widehat{M}(\tau))$ gives a spectral decomposition of the space of L^2 -forms of degree p .

Our main result is a characterization of the L^2 -range of the generalized spectral projection \mathcal{Q}_λ^τ , see Theorem 5.1. In particular, if $\tau = \tau_p$ with p generic (say $p \neq \frac{n-1}{2}, \frac{n}{2}$) we prove that the operator \mathcal{Q}^{τ_p} is an isomorphism from the space $L^2(G, \tau_p)$ onto $\mathcal{E}_{p, \mathbb{R}_>}^2(G, \tau_p) \oplus \mathcal{E}_{p-1, \mathbb{R}_>}^2(G, \tau_p)$, where $\mathcal{E}_{p, \mathbb{R}_>}^2(G, \tau_p)$ (resp. $\mathcal{E}_{p-1, \mathbb{R}_>}^2(G, \tau_p)$) is a space of V_τ -valued measurable functions f on $\mathbb{R}_> \times G$, such that $f_\lambda = f(\lambda, \cdot)$ is for a.e. λ a co-closed p -eigenforms (resp. closed $(p-1)$ -eigenforms) of Δ and satisfying

$$\sup_R \int_0^\infty d\lambda \frac{1}{R} \int_{B(R)} d(gK) \|f_\lambda(g)\|_\tau^2 < \infty,$$

see (5.1).

The projection on $\mathcal{E}_{p, \mathbb{R}_>}^2(G, \tau_p)$ (resp. $\mathcal{E}_{p-1, \mathbb{R}_>}^2(G, \tau_p)$) being $\mathcal{Q}_{\sigma_p}^{\tau_p}$ (resp. $\mathcal{Q}_{\sigma_{p-1}}^{\tau_p}$). Furthermore, for $\lambda \in \mathbb{R} \setminus \{0\}$, we prove that

$$\lim_{R \rightarrow \infty} \int_0^\infty d\lambda \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{Q}_\lambda^\tau f(g)\|_\tau^2 = \frac{1}{\pi} \|f\|_{L^2(G, \tau)}^2,$$

for any L^2 -differential form $f \in L^2(G, \tau)$. The operator \mathcal{Q}_λ^d (resp. $\mathcal{Q}_\lambda^\delta$) of Strichartz can then be seen as our $\mathcal{Q}_{\sigma_{p-1}, \lambda}^{\tau_p}$ (resp. $\mathcal{Q}_{\sigma_p, \lambda}^{\tau_p}$). This provides a positive answer to the conjecture of Strichartz on differential forms.

To prove Theorem 5.1 we will first study the uniform L^2 -boundedness and an image characterization of the Poisson transform of L^2 -differential forms on the boundary $\partial H^n(\mathbb{R}^n)$. This is the purpose of the second main theorem of this paper, see Theorem 5.2. The proof of this theorem is based on a scattering formula for Poisson integrals, see Theorem 6.8. The key

point of proving this latter theorem is to establish an asymptotic expansion for the τ -spherical functions associated to the unitary principal series, see Proposition 6.4.

One should note that the Poisson transform characterization for symmetric spaces of noncompact type is closely related to Helgason's conjecture (see [17]). This subject has been extensively studied by many authors, in particular for vector bundles (see e.g. [4,6,7,24,27,29]).

Let us outline the content of this paper. After general preliminaries (section 2) we will set up the Helgason Fourier transform on the bundle of differential forms and prove a Helgason Fourier restriction theorem (section 3). We obtain then (section 4) as a consequence a uniform estimate of the Poisson transform. Once all materials are introduced we give the announcement of the main theorems (section 5). The delicate part of the paper (section 6) deals with the proof of the asymptotic expansion of τ -spherical functions from which we get the asymptotic expansion of Poisson integrals. The obtained asymptotic properties are then used in section 7 to prove Theorem 5.2. Additionally, we establish, in Section 8, an inversion formula for the Poisson transform, see Theorem 5.3. Finally, we reserve Section 9 for the proof of Theorem 5.1. An appendix is added. It contains the proof of technical lemmas.

2. NOTATIONS AND PRELIMINARIES

Let $H^n(\mathbb{R})$ be the n -dimensional real hyperbolic space ($n \geq 2$) viewed as the rank one symmetric space of the noncompact type G/K where $G = \mathrm{SO}_0(n, 1)$ and $K = \mathrm{SO}(n)$.

Let $\mathfrak{g} \simeq \mathfrak{so}(n, 1)$ and $\mathfrak{k} \simeq \mathfrak{so}(n)$ be the Lie algebras of G and K respectively and write $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ for the Cartan decomposition of \mathfrak{g} . The tangent space $T_o(G/K) \simeq \mathfrak{p}$ of $G/K = H^n(\mathbb{R})$ at the origin $o = eK$ will be identified with the vector space \mathbb{R}^n .

There exists an element $H_0 \in \mathfrak{p}$ such that $\mathfrak{a} = \mathbb{R}H_0$ is a Cartan subspace in \mathfrak{p} . Let $A = \exp \mathfrak{a} = \{a_t = e^{tH_0}, t \in \mathbb{R}\}$ be the corresponding analytic Lie subgroup of G . We define $\alpha \in \mathfrak{a}^*$ by $\alpha(H_0) = 1$. Then the positive restricted root subsystem is $\Sigma^+(\mathfrak{g}, \mathfrak{a}) = \{\alpha\}$ and the corresponding positive Weyl chamber is $\mathfrak{a}^+ = \{H \in \mathfrak{a}, \alpha(H) > 0\} \simeq (0, \infty)$. We will identify $\mathfrak{a}_\mathbb{C}^*$ and \mathbb{C} via the map $\lambda\alpha \mapsto \lambda$. Then the half-sum of positive roots is $\rho = (n-1)\alpha/2 = (n-1)/2$.

Let $\mathfrak{n} = \mathfrak{g}_\alpha$ be the positive root subspace, $N = \exp(\mathfrak{n})$, $A^+ = \exp(\mathfrak{a}^+)$ and $\overline{A^+} = \exp(\overline{\mathfrak{a}^+})$. The group G has an Iwasawa decomposition $G = KAN$. Thus, each $g \in G$ can be uniquely written as $g = k(g)e^{H(g)}n(g)$, where $k(g) \in K$, $H(g) \in \mathfrak{a}$ and $n(g) \in N$. Furthermore, according to the Cartan decomposition $G = K\overline{A^+}K$, any $g \in G$ decomposes as $g = k_1e^{A^+(g)}k_2$, where $k_1, k_2 \in K$ and where $A^+(g) \in \overline{\mathfrak{a}^+}$ is uniquely determined by g .

We define the invariant measure on G/K by

$$\int_{G/K} d(gK)f(gK) = \int_0^\infty dt(2 \sinh t)^{n-1} \int_K dk f(ka_t). \quad (2.1)$$

for any integrable function f on G/K . Here K is equipped with its normalized Haar measure.

2.1. Differential forms on $H^n(\mathbb{R})$ and $\partial H^n(\mathbb{R})$. A differential p -form on $H^n(\mathbb{R})$ is a section of the p -th exterior power of the complexified cotangent bundle of $H^n(\mathbb{R})$, $\bigwedge^p H^n(\mathbb{R}) := \bigwedge^p T_c^* H^n(\mathbb{R})$.

For $1 \leq p \leq n$, let τ_p be the standard representation of K on $V_{\tau_p} = \bigwedge^p \mathbb{C}^n$. Notice that τ_p is equivalent to the p -th exterior power of the coadjoint representation Ad^* of K on $\mathfrak{p}_\mathbb{C}^*$. Let $G \times_K V_{\tau_p}$ be the the G -homogeneous vector bundle associated to τ_p . Since $T_{eK}H^n(\mathbb{R}) \simeq \mathfrak{p} \simeq$

\mathbb{R}^n , then as G -homogeneous vector bundles we have $\bigwedge^p H^n(\mathbb{R}) = G \times_K V_{\tau_p}$. As usual we shall identify the space $C^\infty(\bigwedge^p H^n(\mathbb{R}))$ of smooth differential p -forms on $H^n(\mathbb{R})$ with the space $C^\infty(G, \tau_p)$ of smooth functions $f : G \rightarrow V_{\tau_p}$ which are right K -covariant, i.e.,

$$f(gk) = \tau(k^{-1})f(g) \quad \text{for all } g \in G, k \in K. \quad (2.2)$$

Similarly, we identify the space $L^2(\bigwedge^p H^n(\mathbb{R}))$ of square integrable p -forms with the space $L^2(G, \tau)$ of V_τ -valued functions on G satisfying the transformation rule (2.2) and

$$\|f\|_{L^2(G, \tau)} := \left(\int_{G/K} d(gK) \|f(g)\|_\tau^2 \right)^{\frac{1}{2}} < \infty.$$

Here $\|\cdot\|_\tau$ denotes a norm on V_τ making τ unitary. Notice also that $\|f(gK)\|_\tau = \|f(g)\|_\tau$ is well defined for f satisfying (2.2).

It is known that the representation τ_p is irreducible unless n is even and $p = \frac{n}{2}$ in which case it decomposes as $\tau_{n/2} = \tau_{n/2}^+ \oplus \tau_{n/2}^-$ with the corresponding decomposition of the representation space $\bigwedge^{\frac{n}{2}} \mathbb{C}^n = \bigwedge_+^{\frac{n}{2}} \mathbb{C}^n \oplus \bigwedge_-^{\frac{n}{2}} \mathbb{C}^n$, where $\bigwedge_\pm^{\frac{n}{2}} \mathbb{C}^n = \{\alpha \in \bigwedge^{\frac{n}{2}} \mathbb{C}^n; \star \alpha = \mu_\pm \alpha\}$. In this case we have $\bigwedge^{\frac{n}{2}} H^n(\mathbb{R}) = \bigwedge_+^{\frac{n}{2}} H^n(\mathbb{R}) \oplus \bigwedge_-^{\frac{n}{2}} H^n(\mathbb{R})$ with $\bigwedge_\pm^p H^n(\mathbb{R}) = G \times_K \bigwedge_\pm^{\frac{n}{2}} \mathbb{C}^n$. Here \star is the Hodge operator and $\mu_\pm = \pm 1$ if $\frac{n}{2}$ is even and $\mu_\pm = \pm i$ if $\frac{n}{2}$ is odd. Notice that the Hodge operator induces the equivalence $\tau_p \sim \tau_{n-p}$, hence, hereafter we shall restrict our discussion to $0 \leq p \leq \frac{n}{2}$.

To distinguish between representations of K and representations of M , we will henceforth use the Greek letter σ to denote those representations of M . This notation will help clearly differentiate the representations associated with these two groups throughout our discussion, ensuring that the analysis remains precise and unambiguous.

For $0 \leq q \leq n-1$, let σ_q be the standard representation of M on $V_{\sigma_q} = \bigwedge^q(\mathbb{C}e_2 \oplus \cdots \oplus \mathbb{C}e_n) = \bigwedge^q \mathbb{C}^{n-1}$, where $(e_j)_{j=1}^n$ is the natural basis of \mathbb{C}^n . Then the branching rules for $(K, M) = (\text{SO}(n), \text{SO}(n-1))$ is given as follows (see e.g. [9, 19, 25]) :

- (1) if $p < \frac{n-1}{2}$, then $\tau_{p|M} = \sigma_{p-1} \oplus \sigma_p$;
- (2) if $p = \frac{n-1}{2}$, then $\tau_{p|M} = \sigma_{p-1} \oplus \sigma_p^+ \oplus \sigma_p^-$;
- (3) if $p = \frac{n}{2}$, then $\tau_{\frac{n}{2}|M}^\pm = \sigma_{\frac{n}{2}}$

We will refer to *generic case* for $1 \leq p \leq \frac{n}{2}$ with $p \neq \frac{n-1}{2}, \frac{n}{2}$ (we will not deal with the well-known case $p = 0$, see e.g. [9], [20]) and *special cases* for the cases where n is odd and $p = \frac{n-1}{2}$ or n even and $p = \frac{n}{2}$.

For $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$, we denote by $\widehat{M}(\tau)$ the set of representations in \widehat{M} that occur (with multiplicity one) in the restriction of τ to M . According to the above branching rules we have,

$$\widehat{M}(\tau) = \begin{cases} \widehat{M}(\tau_p) = \{\sigma_{p-1}, \sigma_p\} & \text{for } p < \frac{n-1}{2}, \\ \widehat{M}(\tau_p) = \{\sigma_{p-1}, \sigma_p^+, \sigma_p^-\} & \text{for } p = \frac{n-1}{2}, \\ \widehat{M}(\tau_p^\pm) = \{\sigma_p\} & \text{for } p = \frac{n}{2}. \end{cases}$$

Let $\sigma \in \widehat{M}(\tau)$. Since $T_{eM}K/M = \mathfrak{k}/\mathfrak{m} \simeq \mathfrak{a}^\perp \simeq \mathbb{R}^{n-1}$ then the space $\bigwedge^q \partial H^n(\mathbb{R})$ of differential q -forms on $\partial H^n(\mathbb{R})$ is canonically isomorphic to the homogenous bundle $K \times_M V_\sigma = K \times_M \bigwedge^q \mathbb{C}^{n-1}$, with $q = p, p-1$. The space $L^2(\bigwedge^q \partial H^n(\mathbb{R}))$ of L^2 differential q -forms on

$\partial H^n(\mathbb{R})$ will be identified with the space $L^2(K, \sigma)$ of vector valued functions $F : K \rightarrow V_\sigma$ satisfying the identity

$$F(km) = \sigma(m)^{-1}F(k), \text{ for all } k \in K, m \in M \quad (2.3)$$

and such that

$$\|F\|_{L^2(K, \sigma)} = \left(\int_K \|F(k)\|^2 dk \right)^{\frac{1}{2}} < \infty.$$

Similarly, we will also identify the space $C^{-\omega}(\wedge^q \partial H^n(\mathbb{R}))$ of q -hyperforms on $\partial H^n(\mathbb{R})$ with the space $C^{-\omega}(K, \sigma)$ of vector-valued hyperfunctions $F : K \rightarrow V_\sigma$ satisfying the identity (2.3).

2.2. Fourier-Helgason transform. Below we review some known facts on the Fourier-Helgason transform on differential forms following mainly the notations in [25].

For a fixed $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$ and a given $\sigma \in \widehat{M}(\tau)$ we define the partial Helgason Fourier transform of $f \in C_c^\infty(G, \tau)$, to be the V_σ -valued function on $\mathfrak{a}_\mathbb{C}^* \times K = \mathbb{C} \times K$ given by (see [13, 25]),

$$\mathcal{F}_\sigma^\tau f(\lambda)(k) := \sqrt{d_{\tau, \sigma}} \int_G dg e^{(i\lambda - \rho)H(g^{-1}k)} P_\sigma \tau(\kappa(g^{-1}k))^{-1} f(g),$$

where $d_{\tau, \sigma} = \frac{d_\tau}{d_\sigma} = \frac{\dim \tau}{\dim \sigma}$ and P_σ the orthogonal projection of V_τ onto its σ -isotypic component $V_\tau(\sigma) = V_\sigma$. One can see that $\mathcal{F}_\sigma^\tau f(\lambda)$ belongs to $C^\infty(K, \sigma)$. In the sequel we will set

$$\mathcal{F}_{\sigma, \lambda}^\tau f(k) = \mathcal{F}_\sigma^\tau f(\lambda)(k). \quad (2.4)$$

The Fourier transform $\mathcal{F}^\tau f$ of $f \in C_c^\infty(G, \tau)$ is defined as the family

$$\mathcal{F}^\tau f = (\lambda \mapsto \mathcal{F}_{\sigma, \lambda}^\tau f)_{\sigma \in \widehat{M}(\tau)}. \quad (2.5)$$

Theorem 2.1 (see [25, Theorems 6.19 – 6.23, 2.27]). (1) *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}$. Then for a differential form $f \in C_c^\infty(G, \tau)$ we have the following inversion and Plancherel formulas*

$$f(g) = \sum_{\sigma \in \widehat{M}(\tau)} \sqrt{d_{\tau, \sigma}} \int_0^\infty \nu_\sigma(\lambda) d\lambda \int_K dk e^{-(i\lambda + \rho)H(g^{-1}k)} \tau(\kappa(g^{-1}k)) \mathcal{F}_{\sigma, \lambda}^\tau f(k), \quad (2.6)$$

$$\|f\|_{L^2(G, \tau)}^2 = \sum_{\sigma \in \widehat{M}(\tau)} \int_0^\infty \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2. \quad (2.7)$$

Moreover, the Fourier transform $\mathcal{F}^\tau = (\mathcal{F}_\sigma^\tau)_{\sigma \in \widehat{M}(\tau)}$ extends to bijective isometry from $L^2(G, \tau)$ onto the space

$$\bigoplus_{\sigma \in \widehat{M}(\tau)} L^2(\mathbb{R}_+; L^2(K, \sigma), \nu_\sigma(\lambda) d\lambda).$$

(2) *Let $\tau = \tau_{\frac{n}{2}}^\pm$. Then for $f^\pm \in C_c^\infty(G, \tau_p^\pm)$ we have the following inversion and Plancherel formulas*

$$\begin{aligned} f^\pm(g) = & \int_0^\infty \nu_{\sigma_{\frac{n}{2}}}(\lambda) d\lambda \int_K dk e^{-(i\lambda + \rho)H(g^{-1}k)} \tau_p^\pm(\kappa(g^{-1}k)) \mathcal{F}_\lambda^\pm f^\pm(k) \\ & + 2^{1-2n} n \int_K dk e^{-(\frac{1}{2} + \rho)H(g^{-1}k)} \tau_p^\pm(\kappa(g^{-1}k)) \mathcal{F}_{-\frac{i}{2}}^\pm f^\pm(k), \end{aligned} \quad (2.8)$$

$$\|f^\pm\|_{L^2(G, \tau_p^\pm)}^2 = \int_0^\infty \nu_{\sigma_{\frac{n}{2}}}(\lambda) d\lambda \|\mathcal{F}_\lambda^\pm f^\pm\|_{L^2(K, \sigma_{\frac{n}{2}})}^2 + 2^{1-2n} n \left\langle \mathcal{F}_{-\frac{i}{2}}^\pm f^\pm, \mathcal{F}_{\frac{i}{2}}^\pm f^\pm \right\rangle_{L^2(K, \nu_{\sigma_{\frac{n}{2}}})}. \quad (2.9)$$

Moreover, the Fourier transform \mathcal{F}^\pm extends to a bijective isometry from $L^2(G, \tau_p^\pm)$ onto

$$L^2(\mathbb{R}_+; L^2(K, \sigma_p), \nu_{\sigma_{\frac{n}{2}}}(\lambda) d\lambda) \oplus \mathcal{H}_{\pi^\pm},$$

where π^\pm is the discrete series of G with trivial infinitesimal character.

For $1 \leq p \leq \frac{n-1}{2}$, $q = p-1$, p , the Plancherel densities $\nu_q(\lambda) := \nu_{\sigma_q}(\lambda)$ in the above theorem (part (1)) are given by

$$\nu_q(\lambda) = d_{pq}^{-1} \frac{1}{2^{2n-3} \Gamma^2(\frac{n}{2}) (\lambda^2 + (\rho - q)^2)} \begin{cases} \lambda^2 \prod_{k=1}^{\frac{n-1}{2}} (\lambda^2 + k^2) & \text{when } n \text{ is odd} \\ \lambda \tanh(\pi\lambda) \prod_{k=1}^{\frac{n}{2}} (\lambda^2 + (k - \frac{1}{2})^2) & \text{when } n \text{ is even} \end{cases} \quad (2.10)$$

For $p = \frac{n}{2}$, the Plancherel density $\nu_{\sigma_{\frac{n}{2}}}$ in the above theorem (part (2)) is given by

$$\nu_{\sigma_{\frac{n}{2}}}(\lambda) = \frac{1}{2^{2n-3} \Gamma^2(\frac{n}{2})} \lambda \tanh(\pi\lambda) \prod_{k=2}^{n/2} \left[\lambda^2 + \left(k - \frac{1}{2}\right)^2 \right]. \quad (2.11)$$

Lemma 2.2. For $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$ and any $\sigma \in \widehat{M}(\tau)$, the following estimate holds,¹

$$\nu_\sigma(\lambda) \asymp \lambda^2 (|\lambda| + 1)^{n-1}, \quad \lambda \in \mathbb{R}. \quad (2.12)$$

Proof. This is a direct consequence of (2.10) and (2.11). \square

3. THE HELGASON FOURIER RESTRICTION THEOREM

Next, we are interested in proving a Helgason Fourier restriction theorem of suitable sections of the bundle $G \times_K V_\tau$ and fixed $\tau \in \{\tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm\}$.

Before stating the main theorem of this section we will first define the Radon transform of $f \in C_c^\infty(G, \tau)$ as the V_τ -valued function (see e.g. [11])

$$\mathcal{R}^\tau f(g) = \mathcal{R}^\tau f(t, k) = e^{\rho t} \int_N f(ka_t n) dn, \quad g = ka_t n \in G.$$

Then, for each $\sigma \in \widehat{M}(\tau)$ we define the partial Radon transform of $f \in C_c^\infty(G, \tau)$ by

$$\mathcal{R}_\sigma^\tau f(t, k) = \sqrt{d_{\tau, \sigma}} P_\sigma \mathcal{R}^\tau f(t, k).$$

For every real number $R > 0$ set $B(R) := \{gK \in G/K, d(o, gK) < R\}$, where $d(\cdot, \cdot)$ is the hyperbolic distance on $G/K = H^n(\mathbb{R})$. In view of the inequality (see [18, page 476]),

$$d(eK, ka_t nK) \geq |t| \quad k \in K, n \in N,$$

we easily see that, if $f \in C_c^\infty(G, \tau)$ with $\text{supp } f \subset B(R)$, then

$$\text{supp } \mathcal{R}_\sigma^\tau f \subset [-R, R] \times K. \quad (3.1)$$

¹Here $\phi \asymp \psi$ for two positive functions means that their quotient $\frac{\phi}{\psi}$ stay uniformly bounded.

Finally, note that the partial Radon transform is related to the partial Helgason Fourier transform through the following formula :

$$\mathcal{F}_{\sigma,\lambda}^{\tau} f(k) = \mathcal{F}_{\sigma}^{\tau} f(\lambda, k) = \mathcal{F}_{\mathbb{R}} [\mathcal{R}_{\sigma}^{\tau} f(\cdot, k)](\lambda), \quad k \in K, \quad (3.2)$$

where $\mathcal{F}_{\mathbb{R}} \phi(\lambda) = \int_{\mathbb{R}} e^{-i\lambda t} \phi(t) dt$ is the Euclidean Fourier transform on $\mathfrak{a} = \mathbb{R}$.

We come now to the statement of the uniform continuity estimate for restriction of the Helgason Fourier transform.

Theorem 3.1 (Helgason Fourier restriction theorem). *Let $\tau \in \{\tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^{\pm}\}$ and $\sigma \in \widehat{M}(\tau)$.*

(1) *Suppose n even. There exists a positive constant C such that for any $\lambda \in \mathbb{R} \setminus \{0\}$ and $R > 1$ we have*

$$\|\mathcal{F}_{\sigma,\lambda}^{\tau} f\|_{L^2(K,\sigma)} \leq CR^{1/2} \nu_{\sigma}(\lambda)^{-1/2} \|f\|_{L^2(G,\tau)}, \quad (3.3)$$

for every $f \in L^2(G,\tau)$ supported in $B(R)$.

In the special case $p = \frac{n}{2}$, the above estimate must be understood for $f = f^{\pm} \in L^2(G, \tau_{\frac{n}{2}}^{\pm})_{\text{cont}}$ supported in $B(R)$.

(2) *Suppose n odd. There exists a positive constant C such that for any $\lambda \in \mathbb{R} \setminus \{0\}$ and $R > 0$ we have*

$$\|\mathcal{F}_{\sigma,\lambda}^{\tau} f\|_{L^2(K,\sigma)} \leq CR^{1/2} \nu_{\sigma}(\lambda)^{-1/2} \|f\|_{L^2(G,\tau)}, \quad (3.4)$$

for every $f \in L^2(G,\tau)$ supported in $B(R)$.

Proof. It is sufficient to show (3.3) (resp. (3.4)) for $f \in C_c^{\infty}(G,\tau)$. By the Plancherel formula (2.7)- (2.9) we have

$$\|f\|_{L^2(G,\tau)}^2 = \sum_{\sigma \in \widehat{M}(\tau)} \int_0^{\infty} \nu_{\sigma}(\lambda) d\lambda \|\mathcal{F}_{\sigma,\lambda}^{\tau} f\|_{L^2(K,\sigma)}^2. \quad (3.5)$$

If $p = \frac{n}{2}$, the discrete term

$$2^{1-2n} n \left\langle \mathcal{F}_{\sigma_p, -\frac{i}{2}}^{\tau_p^{\pm}} f^{\pm}, \mathcal{F}_{\sigma_p, \frac{i}{2}}^{\tau_p^{\pm}} f^{\pm} \right\rangle_{L^2(K,\sigma_p)}$$

must be added to the right-hand side of (3.5). So, we need only to prove that there exists $C > 0$ such that for any $R > 1$ (resp. $R > 0$) and $f \in C_c^{\infty}(G,\tau)$ with $\text{supp } f \subset B(R)$, we have

$$\nu_{\sigma}(\lambda) \|\mathcal{F}_{\sigma,\lambda}^{\tau} f\|_{L^2(K,\sigma)}^2 \leq CR \int_0^{\infty} \nu_{\sigma}(\lambda) d\lambda \|\mathcal{F}_{\sigma,\lambda}^{\tau} f\|_{L^2(K,\sigma)}^2. \quad (3.6)$$

(1) Suppose n even. Use (3.2) to rewrite (3.6) as

$$\|\nu_{\sigma}(\lambda)^{1/2} \mathcal{F}_{\mathbb{R}}[\mathcal{R}_{\sigma}^{\tau} f(t, \cdot)](\lambda)\|_{L^2(K,\sigma)}^2 \leq CR \int_0^{\infty} \nu_{\sigma}(\lambda) d\lambda \|\mathcal{F}_{\mathbb{R}}[\mathcal{R}_{\sigma}^{\tau} f(t, \cdot)](\lambda)\|_{L^2(K,\sigma)}^2. \quad (3.7)$$

Since $\nu_{\sigma}(\lambda) \asymp \lambda^2(1+|\lambda|)^{n-1}$ (see (2.12)), we may replace $\nu_{\sigma}(\lambda)^{1/2}$ by $\lambda(1+|\lambda|)^{\rho}$ and $(1+|\lambda|)^{\rho}$ by a positive smooth function $\eta_{\rho}(\lambda)$ on \mathbb{R} , according to [1, Lemma 4].

Then the inequality (3.7) is equivalent to

$$CR^{-1} \|\mathcal{F}_{\mathbb{R}}[T * \mathcal{R}_{\sigma}^{\tau} f(t, \cdot)](\lambda)\|_{L^2(K,\sigma)}^2 \leq \int_{\mathbb{R}} d\lambda \|\mathcal{F}_{\mathbb{R}}[T * \mathcal{R}_{\sigma}^{\tau} f(t, \cdot)](\lambda)\|_{L^2(K,\sigma)}^2, \quad (3.8)$$

where T is the tempered distribution on \mathbb{R} such that $\mathcal{F}_{\mathbb{R}}T(\lambda) = \lambda\eta_{\rho}(\lambda)$. Now, let us prove (3.8). As T has compact support, let $R_0 > 1$ such that $\text{supp}T \subset [-R_0, R_0]$. By (3.1) we have

$$\text{supp}T * \mathcal{R}_{\sigma}^{\tau}f(\cdot, k) \subset [-(R_0 + R), (R_0 + R)],$$

for all $k \in K$. Therefore by the Cauchy-Schwarz inequality and the Euclidean Plancherel theorem we have

$$\begin{aligned} \|\mathcal{F}_e[T * \mathcal{R}_{\sigma}^{\tau}f(t, \cdot)](\lambda)\|_{L^2(K, \sigma)}^2 &\leq (R + R_0) \int_K dk \int_{\mathbb{R}} dt \|T * \mathcal{R}_{\sigma}^{\tau}f(\cdot, k)(t)\|^2 \\ &= \left(1 + \frac{R_0}{R}\right) R \int_K dk \int_{\mathbb{R}} d\lambda \|\mathcal{F}_{\mathbb{R}}[T * \mathcal{R}_{\sigma}^{\tau}f(\cdot, k)](\lambda)\|^2, \end{aligned}$$

thus, the result follows.

(2) Suppose n odd. To show (3.6), let $f \in C_c^{\infty}(G, \tau)$ with $\text{supp}f \subset B(R)$. Since in this case $\nu(\lambda)^{1/2}$ is polynomial in λ , it follows that

$$\begin{aligned} \|\nu_{\sigma}(\lambda)^{1/2}\mathcal{F}_{\sigma, \lambda}^{\tau}f(k)\| &= \|\nu_{\sigma}(\lambda)^{1/2}\mathcal{F}_{\mathbb{R}}\mathcal{R}_{\sigma}^{\tau}f(\cdot, k)(\lambda)\| \\ &= \|\mathcal{F}_{\mathbb{R}}[\nu_{\sigma}\left(\frac{d}{dt}\right)^{1/2}(\mathcal{R}_{\sigma}^{\tau}f(\cdot, k))](\lambda)\| \\ &\leq \int_{\mathbb{R}} dt \|\nu_{\sigma}\left(\frac{d}{dt}\right)^{1/2}(\mathcal{R}_{\sigma}^{\tau}f(t, k))\|. \end{aligned}$$

As $\nu_{\sigma}\left(\frac{d}{dt}\right)^{1/2}$ is a differential operator, we have $\text{supp}[\nu_{\sigma}\left(\frac{d}{dt}\right)^{1/2}\mathcal{R}_{\sigma}^{\tau}f(\cdot, k)] \subset [-R, R]$. Then the Cauchy-Schwarz inequality and the Euclidean Plancherel formula imply

$$\begin{aligned} \|\nu_{\sigma}(\lambda)^{1/2}\mathcal{F}_{\sigma, \lambda}^{\tau}f(k)\|^2 &\leq c_0 R \int_{\mathbb{R}} d\lambda \left\|\mathcal{F}_{\mathbb{R}}\left[\nu_{\sigma}\left(\frac{d}{dt}\right)^{1/2}\mathcal{R}_{\sigma}^{\tau}f(\cdot, k)\right]\right\|^2 \quad c_0 = 1, 2 \\ &= c_0 R \int_{\mathbb{R}} \nu_{\sigma}(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^{\tau}f(k)\|^2. \end{aligned}$$

Integrating both parts of the above inequality with respect to k , the result follows from (2.7). \square

4. POISSON TRANSFORMS ON DIFFERENTIAL FORMS

In this section we shall introduce the Poisson transform for differential forms on $\partial H^n(\mathbb{R})$. Let $\tau_p = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}$. Recall from section 2 that $\tau_{p|M} = \sigma_{p-1} \oplus \sigma_p$, with σ_p irreducible unless n is odd and $p = \frac{n-1}{2}$.

We set $V_{\sigma_p} = \bigwedge^p \mathbb{C}^{n-1}$ and $V_{\sigma_{p-1}} = e_1 \wedge (\bigwedge^{p-1} \mathbb{C}^{n-1})$ and we denote by $i_q^p : V_{\sigma_q} \rightarrow V_{\tau_p}$ the natural inclusion.

Let $\sigma_{q, \lambda}$ be the representation of $P = MAN$ on $V_{q, \lambda} = V_{\sigma_q}$ given by

$$\sigma_{q, \lambda}(man) = \sigma_q(m)a^{\rho-i\lambda}.$$

This defines a principal series representation on the Hilbert space $L^2(G, \sigma_{q, \lambda})$ consisting of P -covariant functions $F : G \rightarrow V_{\sigma_q}$ whose restriction to K belong to $L^2(K, \sigma_q)$. The group action being the left regular representation. We denote by $C^{-\omega}(G, \sigma_{q, \lambda})$ the space of its hyperfunction vectors. Then as G -homogenous vector bundles we have

$$C^{-\omega}(G, \sigma_{q, i(q-\rho)}) = C^{-\omega}\left(\bigwedge^q \partial H^n(\mathbb{R})\right).$$

By the Iwasawa decomposition, the restriction map from G to K gives a K -isomorphism from $L^2(G, \sigma_{q,\lambda})$ onto $L^2(K, \sigma_q)$. This yields to the compact picture with the group action given by

$$\pi_{\sigma,\lambda}(g)F(k) = e^{(i\lambda-\rho)H(g^{-1}k)}F(\kappa(g^{-1}k)).$$

The space $C^{-\omega}(K, \sigma_q)$ of its hyperfunctions sections is then K -isomorphic to $C^{-\omega}(G, \sigma_{q,\lambda})$.

For $\lambda \in \mathbb{C}$, the Poisson transform is the G -equivariant map

$$\mathcal{P}_{\sigma_q,\lambda}^{\tau_p} : C^{-\omega}(G, \sigma_{q,\lambda}) \rightarrow C^\infty(G, \tau_p)$$

defined by

$$\mathcal{P}_{\sigma_q,\lambda}^{\tau_p} F(g) := \sqrt{d_{\tau_p,\sigma_q}} \int_K dk \tau_p(k) \iota_q^p F(gk),$$

which in the compact picture

$$\mathcal{P}_{\sigma_q,\lambda}^{\tau_p} : C^{-\omega}(K, \sigma_q) \rightarrow C^\infty(G, \tau_p)$$

is given by

$$\mathcal{P}_{\sigma_q,\lambda}^{\tau_p} F(g) = \sqrt{d_{\tau_p,\sigma_q}} \int_K dk e^{-(i\lambda+\rho)H(g^{-1}k)} \tau_p(\kappa(g^{-1}k)) \iota_q^p F(k), \quad (4.1)$$

Notice that the Poisson transform $\mathcal{P}_{\sigma_p,\lambda}^{\tau_p}$ is up to a constant the Poisson transform $\Phi_p^{\rho-i\lambda}$ investigated by Gaillard in [15].

To state Gaillard's result let us review the description the algebra $\mathbb{D}(G, \tau_p)$ of G -invariant differential operators on $C^\infty(G, \tau_p)$.

Let $d : C^\infty(\wedge^k H^n(\mathbb{R})) \rightarrow C^\infty(\wedge^{k+1} H^n(\mathbb{R}))$ be the exterior differentiation operator. Use the Hodge-star operator $\star : \wedge^k H^n(\mathbb{R}) \rightarrow \wedge^{n-k} H^n(\mathbb{R})$ to define the co-differentiation operator $\delta = (-1)^{n(k+1)+1} \star d \star : C^\infty(\wedge^k H^n(\mathbb{R})) \rightarrow C^\infty(\wedge^{k-1} H^n(\mathbb{R}))$. Then the Hodge-de Rham Laplacian is defined on $C^\infty(\wedge H^n(\mathbb{R}))$ by $\Delta = d\delta + \delta d$.

In the sequel, one should work with the equivalent operators of \star , d , δ and Δ according to the identification $C^\infty(\wedge^p H^n(\mathbb{R})) \simeq C^\infty(G, \tau_p)$, but we continue using the notation \star , d , δ and Δ .

As stated in [16, Theorem 3.1], see also [25, Corollary 2.2], we have

$$\mathbb{D}(G, \tau) = \begin{cases} \mathbb{C}[d\delta, \delta d] & \text{if } \tau = \tau_p, p < \frac{n-1}{2} \\ \mathbb{C}[d\delta, \star d] & \text{if } \tau = \tau_p, p = \frac{n-1}{2}, \\ \mathbb{C}[\Delta] & \text{if } \tau = \tau_p^\pm, p = \frac{n}{2}, \\ \mathbb{C}[\star, d \star d] & \text{if } \tau = \tau_p, p = \frac{n}{2}, \end{cases}$$

In particular, $\mathbb{D}(G, \tau)$ is a commutative algebra except for $\tau = \tau_{n/2}$ (last case).

For $1 \leq p \leq \frac{n}{2}$ and $\lambda \in \mathbb{C}$, we consider the following space of coclosed p -eigenforms of the Hodge-de Rham Laplacian:

$$\mathcal{E}_{\sigma_p,\lambda}(G, \tau_p) = \{f \in C^\infty(G, \tau_p) : \Delta f = (\lambda^2 + (\rho - p)^2)f, \text{ and } \delta f = 0\}$$

Then Gaillard's result may be stated as follows:

Theorem 4.1 ([15, Theorem 2']). *Let $\tau_p = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}$ and $\lambda \in \mathbb{C}$. The Poisson transform*

$$\mathcal{P}_{\sigma_p,\lambda}^{\tau_p} : C^{-\omega}(K, \sigma_p) \rightarrow \mathcal{E}_{\sigma_p,\lambda}(G, \tau_p)$$

is a K -isomorphism if and only if

$$i\lambda \notin \mathbb{Z}_{\leq 0} - \rho \cup \{p - \rho\}. \quad (4.2)$$

A similar result was announced in [21, Theorem 4].

Next, to handle $\mathcal{P}_{\sigma, \lambda}^{\tau_p}$ for all σ arising from the restricting τ_p to M , we need a result analogous to Theorem 4.1. To this end we introduce the following common eigenspaces:

For $1 \leq p < \frac{n-1}{2}$,

$$\mathcal{E}_{\sigma_{p-1}, \lambda}(G, \tau_p) = \{f \in C^\infty(G, \tau_p) : \Delta f = (\lambda^2 + (\rho - p + 1)^2)f \text{ and } df = 0\}.$$

For $p = \frac{n-1}{2}$,

$$\mathcal{E}_{\sigma_{p-1}, \lambda}(G, \tau_p) = \{f \in C^\infty(G, \tau_p) : \Delta f = (\lambda^2 + 1)f, \text{ and } df = 0\},$$

and

$$\mathcal{E}_{\sigma_p^\pm, \lambda}(G, \tau_p) = \{f \in C^\infty(G, \tau_p) : \star df = \pm i^{p^2-1} \lambda f \text{ and } \delta f = 0\}.$$

Corollary 4.2. *Let $\lambda \in \mathbb{C}$.*

(1) *For $p < \frac{n-1}{2}$, the Poisson transform*

$$\mathcal{P}_{\sigma_{p-1}, \lambda}^{\tau_p} : C^{-\omega}(K, \sigma_{p-1}) \rightarrow \mathcal{E}_{\sigma_{p-1}, \lambda}(G, \tau_p)$$

is a K -isomorphism if and only if

$$i\lambda \notin \mathbb{Z}_{\leq 0} - \rho \cup \{\rho - p + 1\}. \quad (4.3)$$

(2) *For $p = \frac{n-1}{2}$, the Poisson transform*

$$\mathcal{P}_{\sigma_p^\pm, \lambda}^{\tau_p} : C^{-\omega}(K, \sigma_p^\pm) \rightarrow \mathcal{E}_{\sigma_p^\pm, \lambda}(G, \tau_p)$$

is a K -isomorphism if and only if

$$i\lambda \notin \mathbb{Z}_{\leq 0} - \rho.$$

Proof. Notice that the Hodge operator induces the equivalences $\tau_p \sim \tau_{n-p}$ and $\sigma_q \sim \sigma_{n-1-q}$.

(1) By using the identity (see [25])

$$\star(\mathcal{P}_{p-1, \lambda}^{\tau_p} f) = (-1)^{p(p-1)} \mathcal{P}_{n-p, \lambda}^{\tau_{n-p}}(\star f),$$

as well as the relations

$$\delta \star = (-1)^{n+1-p^2} \star d$$

$$\Delta \star = \star \Delta,$$

we easily see that the following diagram

$$\begin{array}{ccc} C^{-\omega}(K, \sigma_{p-1}) & \xrightarrow{\mathcal{P}_{\sigma_{p-1}, \lambda}^{\tau_p}} & \mathcal{E}_{\sigma_{p-1}, \lambda}(G, \tau_p) \\ \star \downarrow & & \downarrow \star \\ C^{-\omega}(K, \sigma_{n-p}) & \xrightarrow{\mathcal{P}_{\sigma_{n-p}, \lambda}^{\tau_{n-p}}} & \mathcal{E}_{\sigma_{n-p}, \lambda}(G, \tau_{n-p}) \end{array}$$

is commutative and the first part of the corollary follows from Theorem 4.1.

(2) Let $p = \frac{n-1}{2}$. Then $\sigma_p = \sigma_p^+ \oplus \sigma_p^-$. By using [25, (4.24) and (4.38)] we have

$$\text{Im } \mathcal{P}_{\sigma_{\frac{n-1}{2}}^\pm, \lambda}^{\tau_{\frac{n-1}{2}}} \subset \mathcal{E}_{\sigma_{\frac{n-1}{2}}^\pm, \lambda}(G, \tau_{\frac{n-1}{2}}).$$

Since $(\star d)^2 = (-1)^{\frac{n(n-1)}{2}} \delta d$, it follows that $\mathcal{E}_{\sigma_p, \lambda}(G, \tau_p) = \mathcal{E}_{\sigma_p^+, \lambda}(G, \tau_p) \oplus \mathcal{E}_{\sigma_p^-, \lambda}(G, \tau_p)$. This together with Theorem 4.1 give the desired result. \square

It remains to prove the analogue of Theorem 4.1 for $p = \frac{n}{2}$ and the pair (τ_p^\pm, σ_p) . As we were unable to derive this result directly from Gaillard's theorem, we will instead rely on a more general result established in [24] by Olbrich. In fact, we could have applied Olbrich's result to the other cases, but we intentionally chose to remain within the framework of differential forms.

Let

$$\mathcal{E}_{\sigma_p, \lambda}(G, \tau_p^\pm) = \left\{ f^\pm \in C^\infty(G, \tau_p^\pm) : \Delta f^\pm = \left(\lambda^2 + \frac{1}{4} \right) f^\pm \right\}, \quad p = \frac{n}{2}.$$

Then the corollary below follows from [24, Theorem 4.16].

Corollary 4.3. *For $p = \frac{n}{2}$, the Poisson transform*

$$\mathcal{P}_{\sigma_p, \lambda}^{\tau_p^\pm} : C^{-\omega}(K, \sigma_p) \rightarrow \mathcal{E}_{\sigma_p, \lambda}(G, \tau_p^\pm)$$

is a K -isomorphism if and only if

$$i\lambda \notin \mathbb{Z}_{\leq 0} - \rho \cup \left\{ -\frac{1}{2} \right\}.$$

Below, we will begin discussing the precise image of square-integrable forms on $\partial H^n(\mathbb{R})$ under the Poisson transform. This image is expected to constitute a subspace of the eigenspaces we introduced earlier.

As a consequence of Theorem 3.1, we have the following uniform estimate for the Poisson transform.

Proposition 4.4. *Let $\tau \in \Lambda$ and $\sigma \in \widehat{M}(\tau)$.*

(1) *Suppose n even. Then there exists a positive constant C such that for $\lambda \in \mathbb{R} \setminus \{0\}$ we have*

$$\sup_{R>1} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^\tau F(g)\|_\tau^2 \leq C \nu_\sigma(\lambda)^{-1} \|F\|_{L^2(K, \sigma)}^2,$$

for any $F \in L^2(K, \sigma)$.

(2) *Suppose n odd. Then there exists a positive constant C such that for $\lambda \in \mathbb{R} \setminus \{0\}$ we have*

$$\sup_{R>0} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^\tau F(g)\|_\tau^2 \leq C \nu_\sigma(\lambda)^{-1} \|F\|_{L^2(K, \sigma)}^2,$$

for any $F \in L^2(K, \sigma)$.

Proof. Since λ is real and τ is unitary, we have

$$\int_{B(R)} d(gK) \langle \mathcal{P}_{\sigma, \lambda}^\tau F(g), f(g) \rangle_\tau = \int_K dk \langle F(k), \mathcal{F}_{\sigma, \lambda}^\tau f(k) \rangle_\sigma,$$

for any $F \in L^2(K, \sigma)$ and $f \in L^2(G, \tau)$ with $\text{supp } f \subset B(R)$. Hence

$$\left| \int_{B(R)} d(gK) \langle \mathcal{P}_{\sigma, \lambda}^\tau F(g), f(g) \rangle_\tau \right| \leq \|F\|_{L^2(K, \sigma)} \left(\int_K dk \|\mathcal{F}_{\sigma, \lambda}^\tau f(k)\|_\sigma^2 \right)^{1/2},$$

and by the Helgason Fourier restriction theorem (Theorem 3.1) we get

$$\left| \int_{B(R)} d(gK) \langle \mathcal{P}_{\sigma,\lambda}^\tau F(g), f(g)_\tau \rangle \right| \leq C \nu_\sigma(\lambda)^{-1/2} R^{1/2} \|F\|_{L^2(K,\sigma)} \|f\|_{L^2(G,\tau)}.$$

The desired estimate follows by taking the supremum over all f with $\|f\|_{L^2(G,\tau)} = 1$. \square

To obtain more accurate understanding on the L^2 -range of $\mathcal{P}_{\sigma,\lambda}^\tau$, Proposition 4.4 motivates us to introduce the eigenspace

$$\mathcal{E}_{\sigma,\lambda}^2(G, \tau) = \mathcal{E}_{\sigma,\lambda}(G, \tau) \cap B^*(G, \tau),$$

where $B^*(G, \tau)$ is the space of $f \in L_{\text{loc}}^2(G, \tau)$ such that

$$\|f\|_*^2 = \begin{cases} \sup_{R>1} \frac{1}{R} \int_{B(R)} d(gK) \|f(g)\|_\tau^2 < \infty & \text{if } n \text{ is even,} \\ \sup_{R>0} \frac{1}{R} \int_{B(R)} d(gK) \|f(g)\|_\tau^2 < \infty & \text{if } n \text{ is odd.} \end{cases}$$

According to Theorem 4.1, Corollary 4.2 and Corollary 4.3, for $\lambda \in \mathbb{R} \setminus \{0\}$, the Poisson transform

$$\mathcal{P}_{\sigma,\lambda}^\tau : L^2(K, \sigma) \rightarrow \mathcal{E}_{\sigma,\lambda}^2(G, \tau) \quad (4.4)$$

is a continuous injective linear map for any $\tau \in \{\tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm\}$ and $\sigma \in \widehat{M}(\tau)$. We will show below, that (4.4) is in fact a bijective map.

5. ANNOUNCEMENT OF THE MAIN THEOREMS

Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$ and $\sigma \in \widehat{M}(\tau)$. For $f \in L^2(G, \tau)$ we define the function $\mathcal{Q}_{\sigma,\lambda}^\tau f \in C^\infty(G, \tau)$ for a.e., $\lambda \in (0, +\infty)$ by

$$\mathcal{Q}_{\sigma,\lambda}^\tau f(g) = \sqrt{d_{\tau,\sigma} \nu_\sigma(\lambda)} \int_K dk e^{-(i\lambda + \rho)H(g^{-1}k)} \tau(\kappa(g^{-1}k))^{-1} \mathcal{F}_{\sigma,\lambda}^\tau f(k).$$

For $\tau = \tau_{n/2}^\pm$ we would need to consider $\mathcal{Q}_{\sigma,\lambda}^\tau$ defined on $L^2(G, \tau)_{\text{cont}}$.

Notice that by the Plancherel Theorem (Theorem 2.1) the operator $\mathcal{Q}_{\sigma,\lambda}^\tau$ is well defined and we have the following spectral decomposition

$$f = \sum_{\sigma \in \widehat{M}(\tau)} \int_0^\infty d\lambda \mathcal{Q}_{\sigma,\lambda}^\tau f.$$

Next, we define the operator $\mathcal{Q}_\lambda^\tau = \left(\mathcal{Q}_{\sigma,\lambda}^\tau \right)_{\sigma \in \widehat{M}(\tau)} : L^2(G, \tau) \rightarrow \oplus_{\sigma \in \widehat{M}(\tau)} \mathcal{E}_{\sigma,\lambda}(G, \tau)$. The family $(\mathcal{Q}_\lambda^\tau)_{\lambda \in \mathbb{R}_>}$ are called here generalized spectral projections. Write

$$\mathcal{Q}^\tau f(\lambda, g) = \mathcal{Q}_\lambda^\tau f(g) \text{ for a.e. } (\lambda, g) \in (0, +\infty) \times G.$$

To investigate an image characterization of \mathcal{Q}^τ , let us introduce the space

$$\mathcal{E}_{\mathbb{R}_>}^2(G, \tau) := \oplus_{\sigma \in \widehat{M}(\tau)} \mathcal{E}_{\sigma, \mathbb{R}_>}^2(G, \tau) \quad (5.1)$$

where $\mathcal{E}_{\sigma, \mathbb{R}_>}^2(G, \tau)$ is the space of V_τ -valued measurable functions ψ on $\mathbb{R}_> \times G$ such that $\psi(\lambda, \cdot) \in \mathcal{E}_{\sigma, \lambda}(G, \tau)$ for a.e. $\lambda \in (0, +\infty)$ and $\|\psi\|_+ < \infty$, with

$$\|\psi\|_+^2 := \begin{cases} \sup_{R>1} \int_0^\infty d\lambda \frac{1}{R} \int_{B(R)} d(gK) \|\psi(\lambda, g)\|_\tau^2 < \infty & \text{if } n \text{ is even,} \\ \sup_{R>0} \int_0^\infty d\lambda \frac{1}{R} \int_{B(R)} d(gK) \|\psi(\lambda, g)\|_\tau^2 < \infty & \text{if } n \text{ is odd.} \end{cases}$$

We also set

$$\|\phi\|_+^2 = \sum_{\sigma \in \widehat{M}(\tau)} \|\phi_\sigma\|_+^2, \quad (5.2)$$

for $\phi = (\phi_\sigma)_{\sigma \in \widehat{M}(\tau)}$.

Now, we state our main theorems.

Theorem 5.1 (Strichartz's conjecture for the spectral projections). *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$.*

(1) *There exists a positive constant C , such that for any $f \in L^2(G, \tau)$ we have*

$$C^{-1} \|f\|_{L^2(G, \tau)} \leq \|\mathcal{Q}^\tau f\|_+ \leq C \|f\|_{L^2(G, \tau)}, \quad (5.3)$$

Furthermore, we have

$$\lim_{R \rightarrow \infty} \int_0^\infty d\lambda \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{Q}_\lambda^\tau f(g)\|_\tau^2 = \frac{1}{\pi} \|f\|_{L^2(G, \tau)}^2. \quad (5.4)$$

For $\tau = \tau_{\frac{n}{2}}^\pm$, one have to consider f in the continuous part of $L^2(G, \tau)$.

(2) For $\tau \neq \tau_{\frac{n}{2}}^\pm$, the linear map \mathcal{Q}^τ is a topological isomorphism from $L^2(G, \tau)$ onto $\mathcal{E}_{\mathbb{R}_>}^2(G, \tau)$.

(3) For $\tau = \tau_{\frac{n}{2}}^\pm$, the linear map \mathcal{Q}^τ is a topological isomorphism from $L^2(G, \tau)_{\text{cont}}$ onto $\mathcal{E}_{\mathbb{R}_>}^2(G, \tau)$.

Since the spectral projection $\mathcal{Q}_{\sigma, \lambda}^\tau f$ can be written as

$$\mathcal{Q}_{\sigma, \lambda}^\tau f(g) = \nu_\sigma(\lambda) \mathcal{P}_{\sigma, \lambda}^\tau (\mathcal{F}_{\sigma, \lambda}^\tau f)(g) \quad \text{for a.e. } \lambda \in \mathbb{R}_> ,$$

the above theorem will be a consequence of the following theorem.

Theorem 5.2 (Strichartz's conjecture for the Poisson transform). *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$, $\sigma \in \widehat{M}(\tau)$ and $\lambda \in \mathbb{R} \setminus \{0\}$.*

(1) *There exists a positive constant C independent of λ such that for any $F \in L^2(K, \sigma)$,*

$$C^{-1} \nu_\sigma(\lambda)^{-1/2} \|F\|_{L^2(K, \sigma)} \leq \|\mathcal{P}_{\sigma, \lambda}^\tau F\|_* \leq C \nu_\sigma(\lambda)^{-1/2} \|F\|_{L^2(K, \sigma)}. \quad (5.5)$$

Furthermore, we have

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^\tau F(g)\|_\tau^2 = \frac{1}{\pi} \nu_\sigma(\lambda)^{-1} \|F\|_{L^2(K, \sigma)}^2. \quad (5.6)$$

(2) *The Poisson transform $\mathcal{P}_{\sigma, \lambda}^\tau$ is a topological isomorphism from $L^2(K, \sigma)$ onto $\mathcal{E}_{\sigma, \lambda}^2(G, \tau)$.*

Moreover, we establish an L^2 -inversion formula for the Poisson transform.

Theorem 5.3 (Inversion formula). *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^{\pm}$, $\sigma \in \widehat{M}(\tau)$ and $\lambda \in \mathbb{R} \setminus \{0\}$. Let $f \in \mathcal{E}_{\sigma, \lambda}^2(G, \tau)$. Then its boundary value F is given by the following inversion formula*

$$F(k) = \pi \nu_{\sigma}(\lambda) \lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) e_{\sigma, \lambda}^{\tau}(k^{-1}g) f(g) \quad \text{in } L^2(K, \sigma),$$

where $e_{\sigma, \lambda}^{\tau}(g) = \sqrt{d_{\tau, \sigma}} e^{(i\lambda - \rho)H(g^{-1})} P_{\sigma} \tau^{-1}(\kappa(g^{-1}))$.

The rest of the paper will be devoted to the proof of Theorem 5.2, Theorem 5.3 and Theorem 5.1.

6. ASYMPTOTIC EXPANSION OF τ -SPHERICAL FUNCTIONS

Let (τ, V_{τ}) be an irreducible representation of K . A function $\Phi: G \rightarrow \text{End}(V_{\tau})$ with $\Phi(e) = \text{Id}$ is called τ -spherical if

(a) Φ is τ -radial, i.e.,

$$\Phi(k_1 g k_2) = \tau(k_2)^{-1} \Phi(g) \tau(k_1)^{-1}, \quad \forall k_1, k_2 \in K, \forall g \in G;$$

(b) $\Phi(\cdot)v$ is an eigenfunction for the algebra $\mathbb{D}(G, \tau)$ for one nonzero $v \in V_{\tau}$ (hence for all $v \in V_{\tau}$).

Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^{\pm}$ and $\sigma \in \widehat{M}(\tau)$. For $\lambda \in \mathbb{C}$, define the Eisenstein integral

$$\Phi_{\sigma, \lambda}^{\tau}(g) = d_{\tau, \sigma} \int_K dk e^{-(i\lambda + \rho)H(g^{-1}k)} \tau(\kappa(g^{-1}k)) P_{\sigma} \tau(k)^{-1}, \quad (6.1)$$

where P_{σ} is the projection of V_{τ} onto V_{σ} . Then $\Phi_{\sigma, \lambda}^{\tau}$ is the τ -spherical function associated to the principal series representation $\pi_{\sigma, \lambda} = \text{Ind}_{MAN}^G(\sigma \otimes e^{\rho - i\lambda} \otimes 1)$. Moreover, since τ_M is multiplicity free any τ -spherical function is given by (6.1), for some $(\sigma, \lambda) \in \widehat{M}(\tau) \times \mathbb{C}$. For $\Re(i\lambda) > 0$, we have the following asymptotic behavior (see e.g. [5, Proposition 5.2]),

$$\lim_{t \rightarrow \infty} e^{(\rho - i\lambda)t} \Phi_{\sigma, \lambda}^{\tau}(a_t) = d_{\tau, \sigma} \mathbf{c}(\lambda, \tau) P_{\sigma}, \quad (6.2)$$

where $\mathbf{c}(\lambda, \tau)$ is the generalized Harish-Chandra c -function given by

$$\mathbf{c}(\lambda, \tau) = \int_{\overline{N}} d\overline{n} e^{-(i\lambda + \rho)H(\overline{n})} \tau(\kappa(\overline{n})) \in \text{End}_M(V_{\tau}). \quad (6.3)$$

Above $d\overline{n}$ denotes the Haar measure on $\overline{N} = \theta(N)$ with the normalization

$$\int_{\overline{N}} e^{-2\rho(H(\overline{n}))} d\overline{n} = 1.$$

The integral (6.3) converges for $\lambda \in \mathbb{C}$ such that $\Re(i\lambda) > 0$ and it has a meromorphic continuation to \mathbb{C} .

By the Cartan decomposition, the τ -spherical function is completely determined by its restriction to A . Since $\Phi_{\sigma, \lambda}(a_t) \in \text{End}_M(V_{\tau})$ it follows from Schur's lemma that it is scalar on each M -isotypic component of V_{τ} . Thus, there exists $\varphi_{\eta, \lambda}: \mathbb{R} \rightarrow \mathbb{C}$ such that

$$\Phi_{\sigma, \lambda}^{\tau}(a_t) = \sum_{\eta \in \widehat{M}(\tau)} \varphi_{\eta, \lambda}(t) \text{Id}_{V_{\eta}}.$$

The scalar components of $\Phi_{\sigma, \lambda}^{\tau}$ have been given explicitly in [25]. Let us give more details.

Generic case. In the sequel, we shall use the notation $\Phi_{q,\lambda}^p$ for $\Phi_{\sigma_q,\lambda}^{\tau_p}$ and $\varphi_{q,\lambda}$ for $\varphi_{\sigma_q,\lambda}$ where $q = p - 1, p$.

(1) The scalar components $\varphi_{p-1,\lambda}, \varphi_{p,\lambda}$ of $\Phi_{p,\lambda}^p$ are given by

$$\varphi_{p-1,\lambda}(t) = \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t), \quad (6.4)$$

$$\varphi_{p,\lambda}(t) = \frac{n}{n-p} \phi_\lambda^{(\frac{n}{2}-1, -\frac{1}{2})}(t) - \frac{p}{n-p} \cosh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t). \quad (6.5)$$

(2) The scalar components $\varphi_{p-1,\lambda}, \varphi_{p,\lambda}$ of $\Phi_{p-1,\lambda}^p$ are given by

$$\varphi_{p-1,\lambda}(t) = \frac{n}{p} \phi_\lambda^{(\frac{n}{2}-1, -\frac{1}{2})}(t) - \frac{n-p}{p} \cosh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t), \quad (6.6)$$

$$\varphi_{p,\lambda}(t) = \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t). \quad (6.7)$$

Special case $p = \frac{n-1}{2}$.

(1) The scalar components $\varphi_{p-1,\lambda}, \varphi_{p,\lambda}^+, \varphi_{p,\lambda}^-$ of $\Phi_{\sigma_{p-1},\lambda}^{\tau_p}$ are given by

$$\varphi_{p-1,\lambda}(t) = \frac{2n}{n-1} \phi_\lambda^{(\frac{n}{2}-1, -\frac{1}{2})}(t) - \frac{n+1}{n-1} \cosh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t), \quad (6.8)$$

$$\varphi_{p,\lambda}^\pm(t) = \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t). \quad (6.9)$$

(2) The scalar components $\varphi_{p-1,\lambda}, \varphi_{p,\lambda}^+, \varphi_{p,\lambda}^-$ of $\Phi_{\sigma_p^\pm,\lambda}^{\tau_p}$ are given by

$$\varphi_{p-1,\lambda}(t) = \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t), \quad (6.10)$$

$$\varphi_{p,\lambda}^+(t) = \frac{2n}{n+1} \phi_\lambda^{(\frac{n}{2}-1, -\frac{1}{2})}(t) - \frac{n-1}{n+1} \cosh(t) \phi_\lambda^{(n/2, -1/2)}(t) \pm \frac{i2\lambda}{n+1} \sinh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t), \quad (6.11)$$

$$\varphi_{p,\lambda}^-(t) = \frac{2n}{n+1} \phi_\lambda^{(\frac{n}{2}-1, -\frac{1}{2})}(t) - \frac{n-1}{n+1} \cosh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t) \mp \frac{i2\lambda}{n+1} \sinh(t) \phi_\lambda^{(\frac{n}{2}, -\frac{1}{2})}(t). \quad (6.12)$$

Special case $p = \frac{n}{2}$. The scalar component $\varphi_{p,\lambda}^\pm$ of $\Phi_{\sigma_p^\pm}^{\tau_p}$ is given by

$$\varphi_{p,\lambda}^+(t) = \varphi_{p,\lambda}^-(t) = \cosh\left(\frac{t}{2}\right) \phi_{2\lambda}^{(\frac{n}{2}-1, \frac{n}{2}+1)}\left(\frac{t}{2}\right). \quad (6.13)$$

Above $\phi_\lambda^{(\alpha,\beta)}$ denotes the Jacobi function

$$\phi_\lambda^{(\alpha,\beta)}(t) = {}_2F_1\left(\frac{i\lambda + \alpha + \beta + 1}{2}, \frac{-i\lambda + \alpha + \beta + 1}{2}; \alpha + 1; -\sinh^2 t\right),$$

where $\alpha, \beta, \lambda \in \mathbb{C}$ with $\alpha \neq -1, -2, \dots$, (see, e.g. [25] [23]).

We recall below some known results on Jacobi functions that will be needed in the sequel (see [14, 23]).

For $\lambda \notin i\mathbb{Z}$, $\phi_\lambda^{\alpha,\beta}$ has the following development

$$\phi_\lambda^{(\alpha,\beta)}(t) = c_{\alpha,\beta}(\lambda) \Psi_\lambda^{\alpha,\beta}(t) + c_{\alpha,\beta}(-\lambda) \Psi_{-\lambda}^{\alpha,\beta}(t), \quad (6.14)$$

where

$$c_{\alpha,\beta}(\lambda) = \frac{2^{-i\lambda+\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(i\lambda)}{\Gamma\left(\frac{i\lambda+\alpha+\beta+1}{2}\right)\Gamma\left(\frac{i\lambda+\alpha-\beta+1}{2}\right)},$$

and

$$\begin{aligned} \Psi_{\lambda}^{\alpha,\beta}(t) &= (2 \sinh t)^{i\lambda-\alpha-\beta-1} {}_2F_1\left(\frac{\alpha+\beta+1-i\lambda}{2}, \frac{-\alpha+\beta+1-i\lambda}{2}; 1-i\lambda; -\frac{1}{\sinh^2 t}\right) \\ &= e^{(i\lambda-\alpha-\beta-1)t}(1+o(1)) \quad \text{as } t \rightarrow \infty. \end{aligned} \tag{6.15}$$

Furthermore, there exists a constant $C > 0$ such that

$$\Psi_{\lambda}^{\alpha,\beta}(t) = e^{(i\lambda-\alpha-\beta-1)t} (1 + e^{-2t}\Theta(\lambda, t)), \quad \text{with } |\Theta(\lambda, t)| \leq C,$$

for all $\lambda \in \mathbb{R}$ and all $t \geq 1$.

Lemma 6.1. *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^{\pm}$ and $\sigma \in \widehat{M}(\tau)$. For $\lambda \in \mathbb{C}$ such that $\Re(i\lambda) > 0$, there exists a meromorphic \mathbb{C} -valued function $c_{\sigma}(\lambda, \tau)$ such that*

$$d_{\tau,\sigma} \mathbf{c}(\lambda, \tau) P_{\sigma} = c_{\sigma}(\lambda, \tau) P_{\sigma}. \tag{6.16}$$

Moreover,

(1) *in the generic case*

$$c_{\sigma}(\lambda, \tau) = \begin{cases} \frac{i\lambda + \rho - p}{2(n-p)} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda) & \text{for } (\tau, \sigma) = (\tau_p, \sigma_p) \\ \frac{i\lambda - \rho + p - 1}{2p} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda) & \text{for } (\tau, \sigma) = (\tau_p, \sigma_{p-1}) \end{cases} \tag{6.17}$$

(2) *in the special cases $p = \frac{n-1}{2}, \frac{n}{2}$,*

$$c_{\sigma}(\lambda, \tau) = \begin{cases} \frac{i\lambda - 1}{n-1} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda) & \text{for } (\tau, \sigma) = (\tau_{\frac{n-1}{2}}, \sigma_{\frac{n-1}{2}-1}) \\ \frac{2i\lambda}{n+1} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda) & \text{for } (\tau, \sigma) = (\tau_{\frac{n-1}{2}}, \sigma_{\frac{n-1}{2}}^{\pm}) \\ \frac{1}{2} c_{\frac{n}{2}-1, \frac{n}{2}+1}(2\lambda) & \text{for } (\tau, \sigma) = (\tau_{\frac{n}{2}}^{\pm}, \sigma_{\frac{n}{2}}) \end{cases}$$

Proof. (1) Suppose p generic and $q = p-1, p$ and let $\lambda \in \mathbb{C}$ with $\Re(i\lambda) > 0$. A direct computation using (6.14) and (6.15) together with the identity $c_{\frac{n}{2}-1, -\frac{1}{2}}(\lambda) = \frac{i\lambda+\rho}{2n} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda)$ gives

$$\lim_{t \rightarrow \infty} e^{(\rho-i\lambda)t} \Phi_{q,\lambda}^p(a_t) = c_q(\lambda, p) \text{Id}_{\Lambda^q \mathbb{C}^{n-1}},$$

where

$$c_{p-1}(\lambda, p) = \frac{i\lambda - \rho + p - 1}{2p} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda), \quad c_p(\lambda, p) = \frac{i\lambda + \rho - p}{2(n-p)} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda).$$

Then the identity (6.16) follows from (6.2).

(2) The proof for the special cases runs as before. \square

Remark 6.2. *Let $\lambda \in \mathbb{R} \setminus \{0\}$, $\tau \in \Lambda$ and $\sigma \in \widehat{M}(\tau)$. Then as expected the Plancherel density is related to the c -function $c_{\sigma}(\lambda, \tau)$ by the following formula*

$$\nu_{\sigma}(\lambda) = \frac{1}{2\pi} d_{\tau,\sigma} |c_{\sigma}(\lambda, \tau)|^{-2}. \tag{6.18}$$

The main result of this section is to establish, for nonzero real parameter λ , an asymptotic expansion of the translated τ -spherical functions $\Phi_{\sigma,\lambda}^\tau(g^{-1}x)$, ($x, g \in G$). Before stating the result let us say that $f_1, f_2 \in B^*(G, \tau)$ are asymptotically equivalent, and we denote $f_1 \simeq f_2$, if

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|f_1(g) - f_2(g)\|_\tau^2 = 0. \quad (6.19)$$

Proposition 6.3. *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$, $\sigma \in \widehat{M}(\tau)$ and $\lambda \in \mathbb{R} \setminus \{0\}$. Then for any $g \in G$ we have the following asymptotic expansion in $B^*(G, \tau)$,*

$$\Phi_{\sigma,\lambda}^\tau(g)v \simeq \tau^{-1}(k_2(g)) \sum_{s \in W} e^{(is\lambda - \rho)A^+(g)} d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g))v, \quad (6.20)$$

where $v \in V_\tau$ and $g = k_1(g)e^{A^+(g)}k_2(g)$.

Proof. Let us first show that the right hand-side of (6.20) belongs to $B^*(G, \tau)$. Since $d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} = c_\sigma(s\lambda, \tau) P_\sigma$ and $\dim \sigma = \dim s\sigma$ for any $s \in W$, we have

$$\begin{aligned} \frac{1}{R} \int_{B(R)} d(gK) \left\| \left[\tau^{-1}(k_2(g)) \sum_{s \in W} e^{(is\lambda - \rho)A^+(g)} d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g))v \right] \right\|^2 \\ \leq \sum_{s \in W} \frac{1}{R} \int_{B(R)} d(gK) e^{-2\rho A^+(g)} |c_{s\sigma}(s\lambda, \tau)|^2 \|P_{s\sigma} \tau^{-1}(k_1(g))v\|^2 \\ \leq 2 |c_\sigma(\lambda, \tau)|^2 \frac{d_\sigma}{d_\tau} \|v\|^2 \frac{1}{R} \int_0^R dt e^{-2\rho t} (2 \sinh t)^{2\rho} \\ \leq 2 |c_\sigma(\lambda, \tau)|^2 \frac{d_\sigma}{d_\tau} \|v\|^2, \end{aligned}$$

for any $R > 0$, what was required to be shown. Above we have also used the fact $|c_{s\sigma}(s\lambda, \tau)|^2 = |c_\sigma(\lambda, \tau)|^2$.

Now we prove the asymptotic expansion (6.20). We have

$$\begin{aligned} \int_{B(R)} d(gK) \left\| \Phi_{\sigma,\lambda}^\tau(g)v - \tau^{-1}(k_2(g)) \sum_{s \in W} e^{(is\lambda - \rho)A^+(g)} d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g))v \right\|^2 \\ = \int_0^R \int_K \left\| \left[\Phi_{\sigma,\lambda}^\tau(a_t) - \sum_{s \in W} e^{(is\lambda - \rho)t} d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \right] \tau^{-1}(k)v \right\|^2 (2 \sinh t)^{2\rho} dk dt. \end{aligned}$$

We claim that for any $\lambda \in \mathbb{R} \setminus \{0\}$ there exists a positive constant C_λ such that for every $t > 0$ we have

$$\left\| \left[\Phi_{\sigma,\lambda}^\tau(a_t) - \sum_{s \in W} e^{(is\lambda - \rho)t} d_{\tau,s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \right] \tau^{-1}(k)v \right\| \leq C_\lambda e^{-\rho t - t} \|v\|. \quad (6.21)$$

Thus, from the definition (6.19), it is clear that (6.21) implies (6.20). So, we are left to prove the estimate (6.21). By continuity it is sufficient to prove it for $t \geq 1$.

Case : p generic and $q = p$. Let us introduce the temporary notation $\Phi_{p,\lambda}^p$ for $\Phi_{\sigma_p,\lambda}^{\tau_p}$. According to (6.14) we may rewrite the scalar components of $\Phi_{p,\lambda}^p$ (see (6.4), (6.5)) as

$$\begin{aligned}\varphi_{p-1,\lambda}(t) &= \sum_{s \in W} c_{\frac{n}{2}, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho - 1)t} (1 + e^{-2t} \Theta_1(s\lambda, t)) \\ \varphi_{p,\lambda}(t) &= \frac{n}{n-p} \sum_{s \in W} c_{\frac{n}{2}-1, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho)t} (1 + e^{-2t} \Theta_2(s\lambda, t)) \\ &\quad - \frac{p}{n-p} (1 + e^{-2t}) \sum_{s \in W} \frac{1}{2} c_{\frac{n}{2}, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho)t} (1 + e^{-2t} \Theta_3(s\lambda, t))\end{aligned}$$

with $|\Theta_i(\lambda, t)| \leq C_i$ ($i = 1, 2, 3$) for all $t \geq 1$. Now, using the identity

$$c_{\frac{n}{2}-1, -\frac{1}{2}}(\lambda) = \frac{i\lambda + \rho}{2n} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda),$$

a direct calculation yields to

$$\varphi_{p,\lambda}(t) = \frac{n}{(n-p)} \sum_{s \in W} \frac{is\lambda + \rho - p}{is\lambda + \rho} c_{\frac{n}{2}-1, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho)t} + e^{-(\rho+2)t} \Theta(\lambda, p, t),$$

with $|\Theta(\lambda, p, t)| \leq C$ for all $t \geq 1$ and $\lambda \in \mathbb{R}$.

Next, let $v \in V_{\tau_p}$ and write $v = v_{p-1} + v_p$, then $\Phi_{p,\lambda}^p(a_t)v = \varphi_{p-1,\lambda}(t)v_{p-1} + \varphi_{p,\lambda}(t)v_p$. Since $c_p(\lambda, p) = \frac{i\lambda + \rho - p}{i\lambda + \rho} c_{\frac{n}{2}-1, -\frac{1}{2}}(\lambda)$ (see (6.17)), we get

$$\begin{aligned}\Phi_{p,\lambda}^p(a_t)v &= e^{-t} \sum_{s \in W} c_{\frac{n}{2}, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho)t} (1 + e^{-2t} \Theta_1(s\lambda, t)) v_{p-1} \\ &\quad + \frac{n}{(n-p)} \left[\sum_{s \in W} c_p(\lambda, p) e^{(is\lambda - \rho)t} + e^{-(\rho+2)t} \Theta(\lambda, p, t) \right] v_p\end{aligned}$$

from which we deduce that

$$\Phi_{p,\lambda}^p(a_t)v - \frac{n}{(n-p)} \left[\sum_{s \in W} c_p(s\lambda, p) e^{(is\lambda - \rho)t} \right] v_p = e^{-t} e^{-\rho t} K(\lambda, p, t)v,$$

where $K(\lambda, p, t) \in \text{End}(V_{\tau_p})$ with $\|K(\lambda, p, t)\| \leq C_\lambda$ for all $t \geq 1$. As $s\sigma_p = \sigma_p$, for any $s \in W$, $\frac{n}{n-p} = d_{\tau_p, \sigma_p}$ and $d_{\tau_p, \sigma_p} \mathbf{c}(\lambda, \tau_p) = c_p(\lambda, \tau_p) P_{\sigma_p}$, we obtain

$$\Phi_{p,\lambda}^p(a_t)v - \sum_{s \in W} e^{(is\lambda - \rho)t} d_{\tau_p, \sigma_p} \mathbf{c}(s\lambda, \tau_p) P_{\sigma_p} v = e^{-t} e^{-\rho t} K(\lambda, p, t)v,$$

and consequently

$$\|\Phi_{p,\lambda}^p(a_t) - \sum_{s \in W} e^{(is\lambda - \rho)t} d_{\tau_p, \sigma_p} \mathbf{c}(s\lambda, \tau_p) P_{\sigma_p}\| \leq C_\lambda e^{-t} e^{-\rho t}$$

for all $t \geq 1$, as to be shown.

Case p generic and $q = p - 1$. The proof is similar to the previous case, so we omit it.

Special case $p = \frac{n-1}{2}$. For $\sigma = \sigma_{\frac{n-1}{2}-1}$ the proof is similar to the generic case. We shall now deal with the case $\sigma = \sigma_{\frac{n-1}{2}}^+$. Using (6.11)-(6.12) and following the same method as before, we get

$$\varphi_{p,\lambda}^\pm(t) = \frac{1}{n+1} \sum_{s \in W} (is\lambda \pm i\lambda) c_{\frac{n}{2}, -\frac{1}{2}}(s\lambda) e^{(is\lambda - \rho)t} + e^{-\rho t} e^{-t} I^\pm(\lambda, t),$$

with $|I^\pm(\lambda, t)| \leq C_\lambda$, for all $t \geq 1$. Therefore

$$\Phi_{\sigma_{\frac{n-1}{2}}^+}^{\tau_{\frac{n-1}{2}}} (a_t)v = \frac{2i\lambda}{n+1} c_{\frac{n}{2}, -\frac{1}{2}}(\lambda) e^{(i\lambda-\rho)t} v_p^+ - \frac{2i\lambda}{n+1} c_{\frac{n}{2}, -\frac{1}{2}}(-\lambda) e^{(-i\lambda-\rho)t} v_p^- + e^{-t} e^{-\rho t} I^+(\lambda, t),$$

and the estimate (6.21) follows.

The case $\sigma = \sigma_{(n-1)/2}^-$ is obtained in the same way since $\Phi_{\sigma_{n-1/2}^-}^{\tau_p}$ and $\Phi_{\sigma_{n-1/2}^+}^{\tau_p}$ have the same scalar components.

Special case $p = \frac{n}{2}$. It is similar to the generic case using (6.13), so we omit details. \square

Next, we will prove the following asymptotic for the translated τ -spherical functions.

Proposition 6.4 (Key formula). *Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$, and $\sigma \in \widehat{M}(\tau)$. Let $\lambda \in \mathbb{R} \setminus \{0\}$ and let $g \in G$ be fixed. Then for any $x \in G$ we have the following expansion in $B^*(G, \tau)$,*

$$\Phi_{\sigma, \lambda}^\tau(g^{-1}x)v \simeq \tau(k_2(x))^{-1} \sum_{s \in W} e^{(is\lambda-\rho)A^+(x)} e^{(is\lambda-\rho)H(g^{-1}k_1(x))} d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau(\kappa(g^{-1}k_1(x)))^{-1} v,$$

where $v \in V_\tau$ and $x = k_1(x)e^{A^+(x)}k_2(x)$.

Proof. It follows from the estimate (6.20), that

$$\Phi_{\sigma, \lambda}^\tau(g^{-1}x)v \simeq \tau^{-1}(k_2(g^{-1}x)) \sum_{s \in W} e^{(is\lambda-\rho)A^+(g^{-1}x)} d_{\tau, s\sigma} \mathbf{c}(\tau, s\lambda) P_{s\sigma} \tau^{-1}(k_1(g^{-1}x))v. \quad (6.22)$$

We rewrite the right-hand side of (6.22) in the form

$$\tau^{-1}(k_2(x)) \sum_{s \in W} e^{(is\lambda-\rho)A^+(x)+H(g^{-1}k_1(x))} d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k_1(x)))v + R_g(x),$$

where

$$\begin{aligned} R_g(x) &= \tau^{-1}(k_2(g^{-1}x)) \sum_{s \in W} e^{(is\lambda-\rho)A^+(g^{-1}x)} d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g^{-1}x))v \\ &\quad - \tau^{-1}(k_2(x)) \sum_{s \in W} e^{(is\lambda-\rho)(A^+(x)+H(g^{-1}k_1(x)))} d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k_1(x)))v. \end{aligned}$$

So, we have to show that $R_g \simeq 0$. To do so, we need the following lemmas, which will be proved in the appendix (see Lemma A and Lemma B).

Lemma 6.5. *For any $g, x \in G$ we have*

$$A^+(gx) = A^+(x) + H(gk_1(x)) + E(g, x), \quad (6.23)$$

where the function E satisfies the estimate

$$0 < E(g, x) \leq e^{2(A^+(g)-A^+(x))}. \quad (6.24)$$

Lemma 6.6. *For any $g \in G$ we have*

$$\lim_{R \rightarrow \infty} \tau^{-1}(k_2(ge^{RH_0})) P_{\sigma} \tau^{-1}(k_1(ge^{RH_0})) = P_{\sigma} \tau^{-1}(\kappa(g)).$$

Now, use the identity (6.23) to rewrite $R_g(x)$ as $R_g(x) = I_g(x) + J_g(x)$, where

$$\begin{aligned} I_g(x) &= \tau^{-1}(k_2(g^{-1}x)) \sum_{s \in W} e^{(is\lambda - \rho)(A^+(x) + H(g^{-1}k_1(x)))} \times \\ &\quad \times \left[e^{(is\lambda - \rho)E(g^{-1},x)} - 1 \right] d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g^{-1}x))v, \\ J_g(x) &= \sum_{s \in W} e^{(is\lambda - \rho)(A^+(x) + H(g^{-1}k_1(x)))} \times \\ &\quad \times \left[\tau^{-1}(k_2(g^{-1}x)) d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(k_1(g^{-1}x)) - \tau^{-1}(k_2(x)) d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k_1(x))) \right] v. \end{aligned}$$

We shall prove that $I_g \simeq 0$ and $J_g \simeq 0$.

Since

$$d_{\tau, s\sigma} \mathbf{c}(s\lambda, \tau) P_{s\sigma} = c_{s\sigma}(s\lambda, \tau) P_{s\sigma} \quad \text{and} \quad |c_{s\sigma}(s\lambda, \tau)| = |c_\sigma(\lambda, \tau)| \quad (6.25)$$

we have

$$\begin{aligned} &\frac{1}{R} \int_{B(R)} d(xK) \| I_g(x) \|^2 \leq \\ &2 \| v \|^2 |c_\sigma(\lambda, \tau)|^2 \sum_{s \in W} \frac{1}{R} \int_{B(R)} d(xK) e^{-2\rho(A^+(x) + H(g^{-1}k_1(x)))} |e^{(is\lambda - \rho)E(g,x)} - 1|^2. \end{aligned}$$

From Lemma (6.5) we have the estimate $|E(g, x)| \leq c_g e^{-2A^+(x)}$ and therefore

$$|e^{(i\lambda - \rho)E(g,x)} - 1|^2 \leq (\lambda^2 + \rho^2) c_g e^{-2A^+(x)}.$$

Put $\gamma = 2(\lambda^2 + \rho^2) c_g |c_\sigma(\lambda, \tau)|^2 \| v \|^2$. Then we have

$$\begin{aligned} &\frac{1}{R} \int_{B(R)} d(xK) \| I_g(x) \|^2 \leq \gamma \frac{1}{R} \int_{B(R)} d(xK) e^{-2\rho(A^+(x) + H(g^{-1}k_1(x)))} e^{-2A^+(x)} \\ &\leq \gamma \frac{1}{R} \int_0^R e^{-(2\rho+2)t} (2 \sinh t)^{2\rho} dt \left(\int_K e^{-2\rho(Hg^{-1}k)} dk \right) \end{aligned}$$

from which we get $I_g \simeq 0$.

Next we shall prove that $J_g \simeq 0$. Using (6.25) and the Cartan decomposition $x = ka_t h$ we easily see that

$$\begin{aligned} &\frac{1}{R} \int_{B(R)} d(xK) \| J_g(x) \|^2 \\ &\leq 2 |c_\sigma(\lambda, \tau)|^2 \sum_{s \in W} \frac{1}{R} \int_0^R \int_K dk dt e^{-2\rho t} (2 \sinh t)^{2\rho} e^{-2\rho(H(g^{-1}k))} \times \\ &\quad \times \| (\tau^{-1}(k_2(g^{-1}ka_t)) P_{s\sigma} \tau^{-1}(k_1(g^{-1}ka_t)) - P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k))) v \|^2 \\ &\leq 2 |c_\sigma(\lambda, \tau)|^2 \sum_{s \in W} \int_0^1 \int_K dk dt e^{-2\rho(H(g^{-1}k))} \times \\ &\quad \times \| (\tau^{-1}(k_2(g^{-1}ka_{tR})) P_{s\sigma} \tau^{-1}(k_1(g^{-1}ka_{tR})) - P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k))) v \|^2. \end{aligned}$$

By Lemma 6.6 we have

$$\lim_{R \rightarrow \infty} \| (\tau^{-1}(k_2(g^{-1}ka_{tR})) P_{s\sigma} \tau^{-1}(k_1(g^{-1}ka_{tR})) - P_{s\sigma} \tau^{-1}(\kappa(g^{-1}k))) v \|^2 = 0.$$

Since

$$\begin{aligned} e^{-2\rho H(g^{-1}k)} & \left\| \left(\tau^{-1}(k_2(g^{-1}ka_{tR})P_{s\sigma}\tau^{-1}(k_1(g^{-1}ka_{tR})) - P_{s\sigma}\tau^{-1}(\kappa(g^{-1}k)) \right) v \right\|^2 \\ & \leq 2e^{-2\rho H(g^{-1}k)} \|v\|^2 \end{aligned}$$

and noting that $\int_K e^{-2\rho H(g^{-1}k)} dk = 1$, we obtain by Lebesgue's dominated convergence theorem that

$$\lim_{R \rightarrow \infty} \int_K dk e^{-2\rho H(g^{-1}k)} \left\| \left(\tau^{-1}(k_2(g^{-1}ka_{tR})P_{s\sigma}\tau^{-1}(k_1(g^{-1}ka_{tR})) - P_{s\sigma}\tau^{-1}(\kappa(g^{-1}k)) \right) v \right\|^2 = 0.$$

Thus $\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(xK) \|J_g(x)\|^2 = 0$, and the proposition follows. \square

For any $\lambda \in \mathbb{R}$, $g \in G$ and $v \in V_\tau$, we define $p_{\sigma,\lambda}^{g,v} \in L^2(K, \sigma)$ by

$$p_{\sigma,\lambda}^{g,v}(k) = \sqrt{d_{\tau,\sigma}} e^{(i\lambda - \rho)H(g^{-1}k)} P_\sigma \tau(\kappa(g^{-1}k))^{-1} v. \quad (6.26)$$

By the symmetric formula (see e.g. [13], Proposition 3.3), for any $x \in G$ we have

$$\mathcal{P}_{\sigma,\lambda}^\tau(p_{\sigma,\lambda}^{g,v})(x) = \Phi_{\sigma,\lambda}^\tau(g^{-1}x)v. \quad (6.27)$$

Lemma 6.7. (1) For $\lambda \in \mathbb{R} \setminus \{0\}$, the set of finite combinations of $p_{\sigma,\lambda}^{g,v}$ ($v \in V_\tau$ and $g \in G$) is a dense subspace of $L^2(K, \sigma)$.

(2) For any $s \in W = \{Id, w\} \simeq \{1, -1\}$ and $\lambda \in \mathbb{R} \setminus \{0\}$, there exists a unique unitary isomorphism $U_{s,\lambda} : L^2(K, \sigma) \rightarrow L^2(K, s\sigma)$ such that

$$U_{s,\lambda} p_{\sigma,\lambda}^{g,v} = p_{s\sigma,s\lambda}^{g,v}. \quad (6.28)$$

Moreover, for $F_1 \in L^2(K, \sigma)$, $F_2 \in L^2(K, s\sigma)$ we have $\mathcal{P}_{\sigma,\lambda}^\tau F_1 = \mathcal{P}_{s\sigma,s\lambda}^\tau F_2$ if and only if $U_{s,\lambda} F_1 = F_2$ i.e. $U_{s,\lambda} = \left(\mathcal{P}_{s\sigma,s\lambda}^\tau \right)^{-1} \circ \mathcal{P}_{\sigma,\lambda}^\tau$.

Proof. The assertion (1) follows from the injectiveness of the Poisson transform $\mathcal{P}_{\sigma,\lambda}^\tau$ for $\lambda \in \mathbb{R} \setminus \{0\}$, see Theorem 4.1, Corollary 4.2 and Corollary 4.3.

(2) $U_{s,\lambda}$ is well defined in $\text{span}(p_{\sigma,\lambda}^{g,v})$ since $\mathcal{P}_{s\sigma,s\lambda}^\tau$ remains injective for $\lambda \in \mathbb{R} \setminus \{0\}$. The identity (6.28) follows from (6.27) and the identity

$$\Phi_{\sigma,\lambda}^\tau = \Phi_{s\sigma,s\lambda}^\tau. \quad (6.29)$$

To show that $U_{s,\lambda}$ is unitary, let $g_1, g_2 \in G$ and $v_1, v_2 \in V_\tau$. Then by using (6.27) (6.29) and noting that $P_{s\sigma}$ is self-adjoint, we get

$$\begin{aligned} & \langle U_{s,\lambda} p_{\sigma,\lambda}^{g_1,v_1}, U_{s,\lambda} p_{\sigma,\lambda}^{g_2,v_2} \rangle_{L^2(K, s\sigma)} \\ & = d_{\tau,\sigma} \left\langle \int_K dk e^{(i\lambda - \rho)H(g_1^{-1}k)} e^{(-i\lambda - \rho)H(g_2^{-1}k)} \tau(\kappa(g_2^{-1}k)) P_{s\sigma} \tau^{-1}(\kappa(g_1^{-1}k)) v_1, v_2 \right\rangle \\ & = \langle \Phi_{s\sigma,s\lambda}^\tau(g_1^{-1}g_2)v_1, v_2 \rangle \\ & = \langle \Phi_{\sigma,\lambda}^\tau(g_1^{-1}g_2)v_1, v_2 \rangle \\ & = \langle p_{\sigma,\lambda}^{g_1,v_1}, p_{\sigma,\lambda}^{g_2,v_2} \rangle_{L^2(K, \sigma)}, \end{aligned}$$

as to be shown. \square

Theorem 6.8. *Let $\tau = \tau_1, \dots, \tau_{n-1}, \tau_{\frac{n}{2}}^{\pm}$ and let $\sigma \in \widehat{M}(\tau)$. For $\lambda \in \mathbb{R} \setminus \{0\}$ and $F \in L^2(K, \sigma)$, we have the following asymptotic expansion for the Poisson transform in $B^*(G, \tau)$,*

$$\mathcal{P}_{\sigma, \lambda}^{\tau} F(x) \simeq \tau(k_2(x))^{-1} \sum_{s \in W} e^{(is\lambda - \rho)A^+(x)} \sqrt{d_{\tau, s\sigma}} \mathbf{c}(s\lambda, \tau) U_{s, \lambda} F(k_1(x)), \quad (6.30)$$

for any $x = k_1(x)e^{A^+(x)}k_2(x) \in G$.

Proof. Notice that both sides of (6.30) depend continuously on $F \in L^2(K, \sigma)$. Therefore in view of the item (1) in Lemma 6.7, to prove the asymptotic expansion holds in $L^2(K, \sigma)$ it is sufficient to prove it for the functions $F = p_{\sigma, \lambda}^{g, v}$ (see (6.26)). Since

$$\mathcal{P}_{\sigma, \lambda}^{\tau} p_{\sigma, \lambda}^{g, v}(x) = \Phi_{\sigma, \lambda}^{\tau}(g^{-1}x)v,$$

we get by Proposition 6.4

$$\mathcal{P}_{\sigma, \lambda}^{\tau} p_{\sigma, \lambda}^{g, v}(x) \simeq \sqrt{d_{\tau, \sigma}} \tau(k_2(x))^{-1} \sum_{s \in W} e^{(is\lambda - \rho)A^+(x)} \mathbf{c}(s\lambda, \tau) p_{s\sigma, s\lambda}^{g, v}(k_1(x)),$$

then we use the identity (6.28) to conclude. \square

7. PROOF OF THEOREM 5.2 – STRICHARTZ'S CONJECTURE FOR POISSON TRANSFORMS

Proof item (1) of Theorem 5.2. The right hand-side of the estimate (5.5) follows from Proposition 4.4 and the left hand side from the equality (5.6). Now, let us prove (5.6). Let $F \in L^2(K, \sigma)$ and let φ_F be the V_{τ} -valued function defined by

$$\varphi_F(g) = \sqrt{d_{\tau, \sigma}} \tau(k_2(g))^{-1} \left(\mathbf{c}(\lambda, \tau) e^{(i\lambda - \rho)A^+(g)} F(k_1(g)) + \mathbf{c}(-\lambda, \tau) e^{(-i\lambda - \rho)A^+(g)} U_{\omega, \lambda} F(k_1(g)) \right).$$

Here ω is a representative of the non-trivial element of the Weyl group. Using the change of variable $g = k_1 e^{tH_0} k_2$ (see (2.1)), the fact that τ and $U_{\omega, \lambda}$ are unitary, and $d_{\tau, \sigma} \mathbf{c}(\lambda, \tau) P_{\sigma} = c_{\sigma}(\lambda, \tau) P_{\sigma}$, we have

$$\begin{aligned} \frac{1}{R} \int_{B(R)} d(gK) \|\varphi_F(g)\|_{\tau}^2 &= \frac{2}{d_{\tau, \sigma}} |c_{\sigma}(\lambda, \tau)|^2 \|F\|_{L^2(K, \sigma)}^2 \left(\frac{1}{R} \int_0^R dt (2e^{-t} \sinh t)^{n-1} \right) + \\ &+ 2\Re \left(c_{\sigma}(\lambda, \tau) \overline{c_{\omega\sigma}(\omega\lambda, \tau)} \int_K dk \langle F(k), U_{\omega, \lambda} F(k) \rangle \frac{1}{R} \int_0^R dt e^{2(i\lambda - \rho)t} (2 \sinh t)^{n-1} \right). \end{aligned} \quad (7.1)$$

The Riemann-Lebesgue theorem implies

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_0^R dt e^{2(i\lambda - \rho)t} (2 \sinh t)^{n-1} = 0.$$

Taking the limit in (7.1) and noting that $\frac{2}{d_{\tau, \sigma}} |c_{\sigma}(\lambda, \tau)|^2 = \frac{1}{\pi} \nu_{\sigma}(\lambda)^{-1}$, we get

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\varphi_F(g)\|_{\tau}^2 = \frac{1}{\pi} \nu_{\sigma}(\lambda, \tau)^{-1} \|F\|_{L^2(K, \sigma)}^2, \quad (7.2)$$

To conclude, we write

$$\begin{aligned} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^{\tau} F(g)\|_{\tau}^2 &= \frac{1}{R} \int_{B(R)} d(gK) \left(\|\varphi_F(g)\|_{\tau}^2 + \|\mathcal{P}_{\sigma, \lambda}^{\tau} F(g) - \varphi_F(g)\|_{\tau}^2 \right. \\ &\quad \left. + 2\Re \langle \mathcal{P}_{\sigma, \lambda}^{\tau} F(g) - \varphi_F(g), \varphi_F(g) \rangle_{\tau} \right), \end{aligned}$$

hence, (5.6) follows from (7.2), Theorem 6.8 and the Cauchy-Schwarz inequality.

It remains to show that $\mathcal{P}_{\sigma,\lambda}^\tau$ is an isomorphism. Before, we need to establish some intermediate results. Let $\tau = \tau_1, \dots, \tau_{\frac{n-1}{2}}, \tau_{\frac{n}{2}}^\pm$ and $\sigma \in \widehat{M}(\tau)$ of dimension d_σ . Let $\widehat{K}(\sigma) \subset \widehat{K}$ be the subset of unitary equivalence classes of irreducible representations containing σ upon restriction to K . Consider an element (δ, V_δ) in $\widehat{K}(\sigma)$, with $d_\delta = \dim V_\delta$. From [9] or [19] it follows that σ occurs in $\delta|_M$ with multiplicity one, and therefore $\dim \text{Hom}_M(V_\delta, V_\sigma) = 1$. Choose the orthogonal projection $P_\delta : V_\delta \rightarrow V_\sigma$ to be a generator of $\text{Hom}_M(V_\delta, V_\sigma)$. Fix an orthonormal basis $\{v_j : 1 \leq j \leq d_\delta\}$ of V_δ . Then the family $\{\phi_j^\delta : 1 \leq j \leq d_\delta, \delta \in \widehat{K}(\sigma)\}$ defined by

$$k \mapsto \phi_j^\delta(k) = P_\delta(\delta(k^{-1})v_j)$$

is an orthogonal basis of the space $L^2(K, \sigma)$, see, e.g., [28]. Hence, the Fourier series expansion of each F in $L^2(K, \sigma)$ is given by

$$F(k) = \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} a_j^\delta \phi_j^\delta(k),$$

with

$$\|F\|_{L^2(K, \sigma)}^2 = \sum_{\delta \in \widehat{K}(\sigma)} \frac{d_\sigma}{d_\delta} \sum_{j=1}^{d_\delta} |a_j^\delta|^2.$$

For $\lambda \in \mathbb{C}$ and $(\delta, V_\delta) \in \widehat{K}(\sigma)$, define the following Eisenstein integral

$$\Phi_{\lambda, \delta} := \Phi_{\sigma, \lambda}^{\tau, \delta} : G \rightarrow \text{End}(V_\delta, V_\tau)$$

by

$$\Phi_{\lambda, \delta}(g)(v) = \sqrt{d_{\tau, \sigma}} \int_K dk e^{-(i\lambda + \rho)H(g^{-1}k)} \tau(\kappa(g^{-1}k)) P_\delta(\delta(k^{-1})v), \quad (7.3)$$

where $g \in G$ and $v \in V_\delta$.

Proposition 7.1. *We have*

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\Phi_{\lambda, \delta}(g)\|_{HS}^2 = \frac{d_\sigma}{\pi} \nu_\sigma(\lambda)^{-1} \quad (7.4)$$

Here $\|\cdot\|_{HS}$ stands for the Hilbert-Schmidt norm.

Proof. Since $\Phi_{\lambda, \delta}(g)v_j = \mathcal{P}_{\sigma, \lambda}^\tau(P_\delta \delta(\cdot)^{-1}v_j)(g)$ for any $j = 1, \dots, d_\delta$, then by (5.6) we have

$$\begin{aligned} \lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\Phi_{\lambda, \delta}(g)\|_{HS}^2 &= \sum_{j=1}^{d_\delta} \lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^\tau(P_\delta \delta(\cdot)^{-1}v_j)(g)\|^2 \\ &= \frac{\nu_\sigma^{-1}(\lambda)}{\pi} \sum_{j=1}^{d_\delta} \|P_\delta \delta(\cdot)^{-1}v_j\|_{L^2(K, \sigma)}^2 \\ &= \frac{\nu_\sigma^{-1}(\lambda)}{\pi} d_\sigma, \end{aligned}$$

as to be shown. □

Proof of item (2) Theorem 5.2. We will now prove that the Poisson transform is a surjective map from $L^2(K, \sigma)$ onto $\mathcal{E}_{\sigma, \lambda}^2(G, \tau)$. Let $f \in \mathcal{E}_{\sigma, \lambda}^2(G, \tau)$. Since $\lambda \in \mathbb{R} \setminus \{0\}$, then by Theorem 4.1, Corollary 4.2 and Corollary 4.3, there exists $F \in C^{-\omega}(K, \sigma)$ such that $\mathcal{P}_{\sigma, \lambda}^\tau F = f$. From the Fourier expansion $F(k) = \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} a_j^\delta P_\delta(\delta(k^{-1})) v_j$ we get

$$f(g) = \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} a_j^\delta \Phi_{\lambda, \delta}(g) v_j,$$

then Schur orthogonality relations show that

$$\int_{B(R)} d(gK) \|f(g)\|_\tau^2 = \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} \frac{|a_j^\delta|^2}{d_\delta} \int_{B(R)} d(gK) \|\Phi_{\lambda, \delta}(g)\|_{HS}^2.$$

Therefore, for any $R > 0$,

$$\|f\|_*^2 \geq \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} \frac{|a_j^\delta|^2}{d_\delta} \frac{1}{R} \int_{B(R)} d(gK) \|\Phi_{\lambda, \tau}(g)\|_{HS}^2$$

In the above inequality the summation over $\delta \in \widehat{K}(\sigma)$ remains true in particular over $\delta \in \Pi$ for any finite subset $\Pi \subset \widehat{K}(\sigma)$. Thus by (7.4) we get,

$$\|f\|_*^2 \geq \frac{\nu_\sigma(\lambda)^{-1}}{\pi} \sum_{\delta \in \Pi} \sum_{j=0}^{d_\delta} \frac{d_\sigma}{d_\delta} |a_j^\delta|^2.$$

Since Π is arbitrary, we obtain

$$\frac{\nu_\sigma(\lambda)^{-1}}{\pi} \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} \frac{d_\sigma}{d_\delta} |a_j^\delta|^2 \leq \|f\|_*^2 < \infty,$$

which proves that $F \in L^2(K, \sigma)$ and

$$\frac{\nu_\sigma(\lambda)^{-1}}{\pi} \|F\|_{L^2(K, \sigma)}^2 \leq \|f\|_*^2.$$

This completes the proof of Theorem 5.2.

8. PROOF OF THEOREM 5.3 – INVERSION FORMULA FOR THE POISSON TRANSFORM

Let $f \in \mathcal{E}_{\sigma, \lambda}^2(G, \tau)$, then by Theorem 5.2, there exists a unique $F \in L^2(K, \sigma)$ such that $f = \mathcal{P}_{\sigma, \lambda}^\tau F$. Now define the function F_R by

$$F_R(k) = \pi \nu_\sigma(\lambda) \frac{1}{R} \int_{B(R)} e^{(i\lambda - \rho)H(g^{-1}k)} \tau^{-1}(\kappa(g^{-1}k)) f(g) d(gK), \text{ for any } R \geq 0,$$

and consider its Fourier expansion

$$F_R(k) = \sum_{\delta \in \widehat{K}(\sigma)} \sum_{j=1}^{d_\delta} c_j^\delta(R) \sqrt{\frac{d_\delta}{d_\sigma}} P_\delta(\delta(k^{-1})) v_j$$

where the coefficients are given by

$$c_j^\delta(R) = \sqrt{\frac{d_\delta}{d_\sigma}} \langle F_R, \overline{P_\delta \delta^{-1}(\cdot) v_j} \rangle$$

Above we have used the orthonormal basis $k \rightarrow \sqrt{\frac{d_\delta}{d_\sigma}} P_\delta(\delta(k^{-1})v_j)$ of $L^2(K, \sigma)$ instead of the orthogonal basis from Section 7.

Let us show that $F_R \rightarrow F$ as $R \rightarrow \infty$ in $L^2(K, \sigma)$. To do so, it is enough to prove that $\lim_{R \rightarrow \infty} \|F_R\|_{L^2(K, \sigma)} = \|F\|_{L^2(K, \sigma)}$.

From the expansion of F , $F(k) = \sum_{\delta' \in \widehat{K}(\sigma)} \sum_{\ell=1}^{d_{\delta'}} a_\ell^{\delta'} \sqrt{\frac{d_{\delta'}}{d_\sigma}} P_{\delta'}(\delta'(k^{-1})v_j)$ and since $f = \mathcal{P}_{\sigma, \lambda}^\tau F$, we get

$$f(g) = \sum_{\delta' \in \widehat{K}(\sigma)} \sum_{\ell=1}^{d_{\delta'}} a_\ell^{\delta'} \sqrt{\frac{d_{\delta'}}{d_\sigma}} \Phi_{\lambda, \delta'}(g) v_\ell$$

where $\Phi_{\lambda, \delta'}$ is given by (7.3). Thus,

$$F_R(k) = \pi \nu_\sigma(\lambda) \sum_{\delta', \ell} \sqrt{\frac{d_{\delta'}}{d_\sigma}} a_\ell^{\delta'} \frac{1}{R} \int_{B(R)} d(gK) e_{\sigma, \lambda}^\tau(k^{-1}g) \Phi_{\lambda, \delta'}(g) v_\ell.$$

Therefore

$$c_j^\delta(R) = \pi \nu_\sigma(\lambda) \sum_{\delta', \ell} a_\ell^{\delta'} \frac{\sqrt{d_{\delta'} d_\delta}}{d_\sigma} \int_{B(R)} d(gK) \langle \Phi_{\lambda, \delta'}(g) v_\ell, \Phi_{\lambda, \delta}(g) v_j \rangle$$

which, by Schur's orthogonality relation, can be written as

$$c_j^\delta(R) = \pi \nu_\sigma(\lambda) \frac{a_j^\delta}{d_\sigma} \frac{1}{R} \int_{B(R)} d(gK) \|\Phi_{\lambda, \delta}(g)\|_{\text{HS}}^2.$$

Next, using Proposition 7.1, we get $\lim_{R \rightarrow \infty} c_j^\delta(R) = a_j^\delta$, thus

$$\lim_{R \rightarrow \infty} \|F_R\|_{L^2(K, \sigma)}^2 = \lim_{R \rightarrow \infty} \sum_{\delta, j} |c_j^\delta(R)|^2 = \sum_{\delta, j} |a_j^\delta|^2 = \|F\|_{L^2(K, \sigma)}^2. \quad (8.1)$$

To justify interchanging the sum and limit signs in (8.1), we would note that

$$\begin{aligned} \frac{1}{R} \int_{B(R)} d(gK) \|\Phi_{\lambda, \delta}(g)\|_{\text{HS}}^2 &= \sum_j \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{P}_{\sigma, \lambda}^\tau(P_\delta \delta^{-1}(\cdot) v_j)(g)\|^2 \\ &\leq \sum_j \|\mathcal{P}_{\sigma, \lambda}^\tau(P_\delta \delta^{-1}(\cdot) v_j)\|_*^2 \\ &\leq C \nu_\sigma(\lambda)^{-1} \sum_j \|P_\delta \delta^{-1}(\cdot) v_j\|_{L^2(K, \sigma)}^2 \quad \text{by (5.5)} \\ &\leq C \nu_\sigma(\lambda)^{-1} \sum_j \frac{d_\sigma}{d_\delta} = C \nu_\sigma(\lambda)^{-1} d_\sigma. \end{aligned}$$

Hence

$$|c_j^\delta(R)| \leq \pi C |a_j^\delta|,$$

for any δ and any j . This completes the proof of Theorem 5.3.

9. PROOF OF THEOREM 5.1 – STRICHARTZ'S CONJECTURE FOR SPECTRAL PROJECTIONS

Proof of item (1) of Theorem 5.1. Assume that $p \neq \frac{n}{2}$ and $\tau = \tau_p$. Let $f \in L^2(G, \tau)$, and define $\mathcal{Q}_\sigma^\tau f$ on $(0, +\infty) \times G$ by $\mathcal{Q}_\sigma^\tau f(\lambda, g) = \mathcal{Q}_{\sigma, \lambda}^\tau f(g)$. Then by definition

$$\|\mathcal{Q}^\tau f\|_+^2 = \sum_{\sigma \in \widehat{M}(\tau)} \|\mathcal{Q}_\sigma^\tau f\|_+^2.$$

Using the identity $\mathcal{Q}_{\sigma, \lambda}^\tau f = \nu(\lambda) \mathcal{P}_{\sigma, \lambda}^\tau (\mathcal{F}_{\sigma, \lambda}^\tau f)$ and Theorem 5.1 we get

$$\begin{aligned} \|\mathcal{Q}_\sigma^\tau f\|_+^2 &= \sup_{R>0} \int_0^\infty d\lambda \left(\frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{Q}_{\sigma, \lambda}^\tau f(g)\|_\tau^2 \right) \\ &\leq \int_0^\infty d\lambda \|\mathcal{Q}_{\sigma, \lambda}^\tau f\|_*^2 \\ &\leq C \int_0^\infty \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2. \end{aligned}$$

Thus

$$\|\mathcal{Q}^\tau f\|_+^2 \leq C \sum_{\sigma \in \widehat{M}(\tau)} \int_0^\infty \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2 = C \|f\|_{L^2(G, \tau)}^2,$$

by the Plancherel formula. This proves the right hand side of (5.3).

Furthermore, by (5.6) and (5.5) we have

$$\lim_{R \rightarrow \infty} \frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{Q}_\lambda^\tau f(g)\|_\tau^2 = \frac{1}{\pi} \sum_{\sigma \in \widehat{M}(\tau)} \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2,$$

and

$$\frac{1}{R} \int_{B(R)} d(gK) \|\mathcal{Q}_\lambda^\tau f(g)\|_\tau^2 \leq \sum_{\sigma \in \widehat{M}(\tau)} \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2.$$

Then by the Lebesgue dominated convergence and the Placherel formula (2.7) we get

$$\begin{aligned} \lim_{R \rightarrow \infty} \frac{1}{R} \int_0^\infty d\lambda \int_{B(R)} d(gK) \|\mathcal{Q}_\lambda^\tau f(g)\|_\tau^2 &= \frac{1}{\pi} \sum_{\sigma \in \widehat{M}(\tau)} \int_0^\infty \nu_\sigma(\lambda) d\lambda \|\mathcal{F}_{\sigma, \lambda}^\tau f\|_{L^2(K, \sigma)}^2 \\ &= \frac{1}{\pi} \|f\|_{L^2(G, \tau)}^2, \end{aligned}$$

which prove (5.4) and indeed the left hand side of (5.3).

The proof for $p = \frac{n}{2}$ is similar to the above taking in account $L^2(G, \tau_{\frac{n}{2}}^\pm)_{\text{cont}}$.

Proof of item (2) of Theorem 5.1. Assume that $p \neq \frac{n}{2}$ and $\tau = \tau_p$. Linearity together with (5.3) imply the injectivity of \mathcal{Q}^τ . It remains then to prove that \mathcal{Q}^τ is onto. Let $\psi \in \mathcal{E}_{\mathbb{R}^>}^2(G, \tau)$. Then for any λ , $\psi(\lambda, \cdot) = \sum_{\sigma \in \widehat{M}(\tau)} \psi_{\sigma, \lambda}$ with $\psi_{\sigma, \lambda} \in \mathcal{E}_{\sigma, \mathbb{R}^>}^2(G, \tau)$. Since

$$\sup_{R>0} \frac{1}{R} \int_{B(R)} d(gK) \|\psi_{\sigma, \lambda}(g)\|_\tau^2 < \infty \quad \text{a.e. } \lambda \in (0, \infty),$$

then applying Theorem 5.2 we can assert the existence of $F_{\sigma,\lambda} \in L^2(K, \sigma)$ such that for a.e. λ we have $\psi_{\sigma,\lambda} = \nu_\sigma(\lambda) \mathcal{P}_{\sigma,\lambda}^\tau F_{\sigma,\lambda}$. Furthermore, it follows from the estimate (5.5) that

$$\sup_{R>0} \frac{1}{R} \int_{B(R)} d(gK) \|\psi_{\sigma,\lambda}(g)\|_\tau^2 \geq C^{-2} \nu_\sigma(\lambda) \int_K dk \|F_{\sigma,\lambda}(k)\|_\sigma^2,$$

hence

$$C^{-2} \int_0^\infty \int_K dk \|F_{\sigma,\lambda}(k)\|_\sigma^2 \nu_\sigma(\lambda) d\lambda \leq \|\psi_{\sigma,\lambda}\|_+^2 < \infty,$$

which proves that

$$F_\lambda = \sum_{\sigma \in \widehat{M}(\sigma)} F_{\sigma,\lambda} \in \oplus_{\sigma \in \widehat{M}(\sigma)} L^2(\mathbb{R}_+; L^2(K, \sigma), \nu_\sigma(\lambda) d\lambda).$$

Now by the Plancherel theorem (Theorem 2.1), there exists $f \in L^2(G, \tau)$ such that $F_\lambda = \mathcal{F}^\tau f(\lambda) = (\mathcal{F}_{\sigma,\lambda}^\tau f)_\sigma$, thus, for any $g \in G$,

$$\begin{aligned} \psi(\lambda, g) &= \sum_{\sigma \in \widehat{M}(\sigma)} \nu_\sigma(\lambda) \mathcal{P}_{\sigma,\lambda}^\tau (\mathcal{F}_{\sigma,\lambda}^\tau f)(g) \\ &= \mathcal{Q}_\lambda^\tau(f)(g) = \mathcal{Q}^\tau f(\lambda, g), \end{aligned}$$

as required to be shown.

The case $p = \frac{n}{2}$ can be treated similarly. This completes the proof of Theorem 5.1.

10. APPENDIX

In this appendix we will provide the proof lemma 6.6 and Lemma 6.5, we have used to show the key formula in Proposition 6.4. For the convenience of the reader, we will recall their statement.

Elements of $G = \text{SO}_0(n, 1)$ are of the form $g = \begin{pmatrix} A & b \\ c^\top & d \end{pmatrix}$ where $A \in \mathcal{M}(n, \mathbb{R})$, $b, c \in \mathbb{R}^n$ (column vectors) and $d > 0$ a real number, with the relations

$$\begin{aligned} AA^\top &= I_n + cc^\top, \\ A^\top b &= dc, \\ b^\top b &= d^2 - 1. \end{aligned} \tag{10.1}$$

The Cartan decomposition of the Lie algebra $\mathfrak{so}(n, 1)$ is $\mathfrak{so}(n, 1) = \mathfrak{k} \oplus \mathfrak{p}$, where

$$\begin{aligned} \mathfrak{k} &= \left\{ \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix} : x \in \mathcal{M}_n(\mathbb{R}), x^\top = -x \right\}, \\ \mathfrak{p} &= \left\{ \begin{pmatrix} 0_n & y \\ y^\top & 0 \end{pmatrix} : y \in \mathbb{R}^n \text{ (column)} \right\}. \end{aligned}$$

The tangent space of G/K at eK is identified to \mathfrak{p} .

Fix the Cartan subspace $\mathfrak{a} = \mathbb{R}H_0$ of \mathfrak{p} , where

$$H_0 = \begin{pmatrix} 0_n & e_1 \\ e_1^\top & 0 \end{pmatrix} \in \mathfrak{p},$$

and $e_1 = (1, 0, \dots, 0)^\top \in \mathbb{R}^n$. The subgroups K , A and N are given by

$$\begin{aligned} K &= \left\{ k = \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix} : u \in \text{SO}(n) \right\}, \\ A &= \left\{ a_t = e^{tH_0} = \begin{pmatrix} \cosh(t) & 0 & \sinh(t) \\ 0 & I_{n-1} & 0 \\ \sinh(t) & 0 & \cosh(t) \end{pmatrix} : t \in \mathbb{R} \right\}, \\ N &= \left\{ n_y = \begin{pmatrix} 1 - \frac{1}{2}\|y\|^2 & y^\top & \frac{1}{2}\|y\|^2 \\ -y & I_{n-1} & y \\ -\frac{1}{2}\|y\|^2 & y^\top & 1 + \frac{1}{2}\|y\|^2 \end{pmatrix} : y \in \mathbb{R}^{n-1} \text{ (column)} \right\}. \end{aligned}$$

Recall that the action of K on the tangent space corresponds to the adjoint action of K on \mathfrak{p} : if $k = \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix} \in K$ and $Y = \begin{pmatrix} 0 & y \\ y^\top & 0 \end{pmatrix} \in \mathfrak{p}$, then $Ad(k)Y = \begin{pmatrix} 0 & uy \\ (uy)^\top & 0 \end{pmatrix}$. Thus, the adjoint action of K on \mathfrak{p} is identified with the linear action of $\text{SO}(n)$ on \mathbb{R}^n .

For any $g \in G$, we have the Cartan and the Iwasawa decompositions,

$$\begin{aligned} g &= k_1(g)e^{A^+(g)}k_2(g) \in K\overline{A^+}K \\ &= \kappa(g)e^{H(g)}n(g) \in KAN. \end{aligned}$$

Then a direct calculation yields²,

$$H(g) = \log |c^\top e_1 + d| \quad \text{and} \quad A^+(g) = \log \left(d + \sqrt{d^2 - 1} \right), \quad (10.2)$$

where $g = \begin{pmatrix} A & b \\ c^\top & d \end{pmatrix}$.

Lemma A. *For any $g, x \in G$ we have*

$$A^+(g^{-1}x) = A^+(x) + H(g^{-1}k_1(x)) + E(g, x),$$

where the function E satisfies the estimate

$$0 < E(g, x) \leq e^{2(A^+(g) - A^+(x))}.$$

Proof. Put $E(g, x) = A^+(g^{-1}x) - A^+(x) - H(g^{-1}k_1(x))$ and write $x = ke^{tH_0}h$ with respect to the Cartan decomposition. Then we have

$$E(g, ke^{tH_0}h) = A^+(g^{-1}ke^{tH_0}) - H(g^{-1}k) - t.$$

Next write $g^{-1} = \begin{pmatrix} A & b \\ c^\top & d \end{pmatrix}$ and $k = \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix}$. Then according to (10.2),

$$\begin{aligned} &E(g, ke^{tH_0}h) \\ &= \log \left(|\sinh(t)c^\top u e_1 + d \cosh(t)| + \sqrt{|\sinh(t)c^\top u e_1 + d \cosh(t)|^2 - 1} \right) - \log |c^\top u e_1 + d| - t. \end{aligned}$$

Thus

$$\begin{aligned} E(g, ke^{tH_0}h) &\leq \log \left(1 + e^{-2t} \frac{|c^\top u e_1 - d|}{|c^\top u e_1 + d|} \right) \\ &\leq \log \left(1 + e^{-2t} e^{2A^+(g)} \right) \end{aligned}$$

²Notice that from the relation (10.1), $d^2 - 1$ is positive

and the result follows. \square

For $g \in G$, we denote by $\pi_o(g)$ its K -component in the polar decomposition $G = K \exp(\mathfrak{p})$. Then, if $g = \begin{pmatrix} A & b \\ c^\top & d \end{pmatrix}$, one can prove that

$$\pi_0(g) = \begin{pmatrix} A - \frac{1}{1+d}bc^\top & 0 \\ 0 & 1 \end{pmatrix}. \quad (10.3)$$

Recall that, if $g = k_1(g)e^{tH_0}k_2$ is the Cartan decomposition of g , then

$$\pi_0(g) = k_1(g)k_2(g). \quad (10.4)$$

Lemma B. *For any $g \in G$ we have*

$$\lim_{R \rightarrow \infty} \tau^{-1}(k_2(ge^{RH_0})P_{\sigma\tau^{-1}}(k_1(ge^{RH_0})) = P_{\sigma\tau^{-1}}(\kappa(g)). \quad (10.5)$$

Proof. Case $\tau = \tau_p$ with p generic. Set $\mathbb{C}^n = \text{span}(e_1, e_2, \dots, e_n)$ and $\mathbb{C}^{n-1} = \text{span}(e_2, \dots, e_n)$, then $\tau_p|_M = \sigma_{p-1} \oplus \sigma_p$ with de decomposition of the representation space

$$\bigwedge^p \mathbb{C}^n = e_1 \wedge \left(\bigwedge^{p-1} \mathbb{C}^{n-1} \right) \oplus \bigwedge^p \mathbb{C}^{n-1}.$$

The projection on σ_p is given by $P_{\sigma_p}\xi = e_1 \wedge \xi = \varepsilon_{e_1}\xi$ (the left exterior product by e_1) and the projection on σ_{p-1} is given by $P_{\sigma_{p-1}}\xi = \iota_{e_1}\xi = \langle \xi, e_1 \rangle$ (the interior product by e_1).

Since

$$\tau_p(k)\varepsilon_v\tau_p(k^{-1}) = \varepsilon_{kv} \quad \text{and} \quad \tau_p(k)\iota_v\tau_p(k^{-1}) = \iota_{kv},$$

we have

$$P_{\sigma_p}\tau_p(k^{-1})\xi = \tau_p(k^{-1})(\varepsilon_{ke_1}\xi) \quad \text{and} \quad P_{\sigma_{p-1}}\tau_p(k^{-1})\xi = \tau_p(k^{-1})(\iota_{ke_1}\xi).$$

Thus, in the case $q = p$,

$$\begin{aligned} & \tau_p^{-1}(k_2(ge^{RH_0}))P_{\sigma_p}\tau_p^{-1}(k_1(ge^{RH_0}))\xi - P_{\sigma_p}\tau_p^{-1}(\kappa(g))\xi \\ &= \tau_p^{-1}(k_2(ge^{RH_0}))\tau_p^{-1}(k_1(ge^{RH_0}))\varepsilon_{k_1(ge^{RH_0})e_1}\xi - \tau_p^{-1}(\kappa(g))\varepsilon_{\kappa(g)e_1}\xi. \end{aligned}$$

Using (10.4) we have $k_2(ge^{RH_0})k_1(ge^{RH_0}) = \pi_0(ge^{RH_0})$, and from the Iwasawa decomposition $g = \kappa(g)a_t n_y$, we get $\pi_0(ge^{RH_0}) = \kappa(g)\pi_0(e^{tH_0}n_y e^{RH_0})$. But

$$e^{tH_0}n_y e^{RH_0} = \begin{pmatrix} \cosh(t+R) - \frac{e^{t-R}}{2}\|y\|^2 & e^t y^\top & \sinh(t+R) + \frac{e^{t-R}}{2}\|y\|^2 \\ -e^{-R}y & I_{n-1} & e^{-R}y \\ \sinh(t+R) - \frac{e^{t-R}}{2}\|y\|^2 & e^t y^\top & \cosh(t+R) + \frac{e^{t-R}}{2}\|y\|^2 \end{pmatrix}. \quad (10.6)$$

Hence, if $\pi_0(e^{tH_0}n_y e^{RH_0}) = \begin{pmatrix} B & 0 \\ 0 & 1 \end{pmatrix}$, then using (10.3) we obtain

$$\begin{aligned} B &= \begin{pmatrix} \cosh(t+R) - \frac{e^{t-R}}{2}\|y\|^2 & e^t y^\top \\ -e^{-R}y & I_{n-1} \end{pmatrix} - \frac{1}{1 + \cosh(t+R) + \frac{e^{t-R}}{2}\|y\|^2} \times \\ & \quad \begin{pmatrix} \sinh^2(t+R) - \frac{e^{2(t-R)}}{4}\|y\|^4 & e^t \left(\sinh(t+R) + \frac{e^{t-R}}{2}\|y\|^2 \right) y^\top \\ e^{-R} \left(\sinh(t+R) - \frac{e^{t-R}}{2}\|y\|^2 \right) y & e^{t-R} y y^\top \end{pmatrix} \end{aligned}$$

and one can check easily that $B \rightarrow I_{n-1}$ when $R \rightarrow +\infty$.

On the other hand, $k_1(ge^{RH_0})e_1 = \kappa(g)k_1(e^{tH_0}n_y e^{RH_0})e_1$. Then from (10.6) we get

$$k_1(e^{tH_0}n_y e^{RH_0})e_1 = \left(\left(\cosh(t+R) + \frac{e^{r-R}}{2} \|y\|^2 \right)^1 - 1 \right)^{-\frac{1}{2}} \left(\frac{\sinh(t+R) + \frac{e^{t-R}}{2} \|y\|^2}{e^{-R}y} \right)$$

$$\xrightarrow[R \rightarrow \infty]{} e_1$$

and therefore, $k_1(ge^{RH_0})e_1 \xrightarrow[R \rightarrow \infty]{} \kappa(g)e_1$. This proves (10.5) for $q = p$ in the generic case.

The case $q = p - 1$ is similar to the above, and it is left to the reader.

Case $\tau_p = \tau_{\frac{p}{2}}^{\pm}$. In this case $\tau_{\frac{p}{2}|M}^{\pm} = \sigma_{\frac{p}{2}}$ and the projection P_{σ} on σ is the identity. Then (10.5) follows from the proof for p the generic and $q = p$.

Case $\tau_p = \tau_{\frac{n-1}{2}}$. In this case $\tau_{p|M} = \sigma_{p-1} \oplus \sigma_p^+ \oplus \sigma_p^-$, and the representation space is decomposed as

$$\bigwedge^{\frac{n-1}{2}} \mathbb{C}^n = e_1 \wedge \left(\bigwedge^{\frac{n-1}{2}-1} \mathbb{C}^{n-1} \right) \oplus \bigwedge_{+}^{\frac{n-1}{2}} \mathbb{C}^{n-1} \oplus \bigwedge_{-}^{\frac{n-1}{2}} \mathbb{C}^{n-1}.$$

The projection P_{σ} on σ_{p-1} is $\xi \rightarrow \varepsilon_{e_1} \xi = e_1 \wedge \xi$ and (10.5) follows from the proof for p generic. On the other hand, the projection $P_{\sigma_{\pm}^p}$ on σ_{\pm}^p is given by

$$P_{\sigma_{\pm}^p} \xi = \frac{1}{2} e_1 \wedge \xi \pm \frac{1}{2} i^{p(p+2)} \star (e_1 \wedge \xi) = \frac{1}{2} \varepsilon_{e_1} \xi \pm \frac{1}{2} i^{p^2+2} \iota_{e_1} (\star \xi).$$

Again (10.5) follows from the proof for p generic. \square

We hereby announce that the Strichartz conjecture for the bundle of spinors over $H^n(\mathbb{R})$ is discussed in the forthcoming paper [3].

REFERENCES

- [1] Anker J.-Ph. A basic inequality for scattering theory on Riemannian symmetric spaces of the noncompact type. *Amer. J. Math.* **113** (1991), no. 3, 391–398. [8](#)
- [2] Baldoni-Silva M.W. Branching theorems for semisimple Lie groups of real rank one. *Rend. Sem. Mat. Univ. Padova* **61** (1979), 229–250. [2](#), [5](#), [24](#)
- [3] Boussejra A.; Koufany K. An L^2 -range characterization of the generalized spectral projections related to the Dirac operator. Preprint. [31](#)
- [4] Bensaïd S.; Boussejra A.; Koufany K. On Poisson transforms for spinors. *Tunis. J. Math.* **5** (2023) no.4, 771–792. [4](#)
- [5] Bensaïd S.; Boussejra A.; Koufany K. On Poisson transforms for differential forms on real hyperbolic spaces. [arXiv:2112.09994](#). [15](#)
- [6] Boussejra A., Ouald Chaib A. A characterization of the L^2 -range of the Poisson transforms on a class of vector bundles over the quaternionic hyperbolic spaces. *J. Geom. Phys.* **194** (2023), Paper No. 105019, 24 pp. [4](#)
- [7] Boussejra A.; Imesmad N.; Ouald Chaib A. L^2 -Poisson integral representations of eigensections of invariant differential operators on a homogeneous line bundle over the complex Grassmann manifold $SU(r, r + b)/S(U(r) \times U(r + b))$. *Ann. Global Anal. Geom.* **61** (2022), no. 2, 399–426. [4](#)
- [8] Boussejra A.; Ourchane N. Characterization of the L^p -range of the Poisson transform on the octonionic hyperbolic plane *J. Lie Theory* **28** (2018), no. 3, 805–828. [2](#)
- [9] Boussejra A.; Sami H. Characterization of the L^p -Range of the Poisson Transform in Hyperbolic Spaces $\mathbf{B}(\mathbb{F}^n)$. *J. Lie Theory* **12** (2002) no. 1, 1–14. [2](#), [5](#), [24](#)
- [10] Boussejra A.; Intissar A. Caractérisation des intégrales de Poisson-Szegö de $L^2(\partial B^n)$ dans la boule de Bergman B^n , $n \geq 2$. *C. R. Acad. Sci. Paris Sér. I Math.* **315** (1992), no.13, 1353–1357. [2](#)
- [11] Branson Thomas P.; Ólafsson, G.; Schlichtkrull, H. A Bundle valued Radon transform, with applications to invariant wave equations. *Quart. J. Math. Oxford* (2) **45**, (1994), 429–461. [7](#)

- [12] Bray William O. Aspects of harmonic analysis on real hyperbolic space, in: *Fourier Analysis, Orono, ME, 1992*, Lect. Notes Pure Appl. Math., vol. 157, Dekker, New York, 1994, pp. 77–102. [2](#)
- [13] Camporesi. R, The Helgason Fourier transform for homogeneous vector bundles over Riemannian symmetric spaces. *Pac. J. Math.* **179** (2) (1997) 263–300. [6](#), [22](#)
- [14] Flensted-Jensen M. Paley-Wiener type theorems for a differential operator connected with symmetric spaces Ark. Mat. **10**(1972), 143–162. [16](#)
- [15] Gaillard P.-Y. Transformation de Poisson de formes différentielles. Le cas de l'espace hyperbolique. *Comment. Math. Helv.* **61** (1986), no. 4, 581–616. [10](#)
- [16] Gaillard P.-Y. Invariant syzygies and semisimple groups. *Adv. Math.* **92** (1992), no. 1, 27–46. [10](#)
- [17] Helgason S. A duality for symmetric spaces with applications to group representations. *Adv. Math.* **5** (1970) 1–154. [4](#)
- [18] Helgason S. Groups and Geometric Analysis, Integral geometry, Invariant Differential operators and Spherical Functions. Academic Press, New York, 1984. [7](#)
- [19] Ikeda A.; Taniguchi Y. Spectra and eigenforms of the Laplacian on S^n and $P^n(\mathbb{C})$. *Osaka J. Math.* **15** (1978), 515–546. [5](#), [24](#)
- [20] Ionescu Alexandru D. On the Poisson Transform on Symmetric Spaces of Real Rank One. *J. Funct. Anal.* **174** (2000), no.2, 513—523. [2](#), [5](#)
- [21] Juhl, A. On the Poisson transformation for differential forms on hyperbolic spaces. *Seminar Analysis of the Karl-Weierstrass-Institute of Mathematics, 1986/87 (Berlin, 1986/87)*, 224–236, Teubner-Texte Math., **106**, Teubner, Leipzig, 1988. [11](#)
- [22] Kaizuka K. A characterization of the L^2 -range of the Poisson transform related to Strichartz conjecture on symmetric spaces of noncompact type *Adv. Math.* **303** (2016), 464—501. [2](#)
- [23] Koornwinder Tom H. Jacobi functions and analysis on non-compact semisimple Lie groups. *Special functions: group theoretical aspects and applications*, 1–85, Math. Appl., Reidel, Dordrecht, 1984. [16](#)
- [24] Olbrich M. Die Poisson-Transformation für homogene Vektorbiindel, Ph.D. Thesis, Humboldt Universitlt, Berlin, 1994. [4](#), [12](#)
- [25] Pedon E. Analyse harmonique des formes différentielles sur l'espace hyperbolique réel. Thèse Université Henri Poincaré-Nancy (1997). [3](#), [5](#), [6](#), [10](#), [11](#), [15](#), [16](#)
- [26] Strichartz Robert S. Harmonic analysis as spectral theory of Laplacians. *J. Funct. Anal.* **87** (1989) 51–148. [2](#)
- [27] van der Ven H. Vector valued Poisson transforms on Riemannian symmetric spaces of rank one. *J. Funct. Anal.* **119** (1994), no. 2, 358–400. [4](#)
- [28] Wallach N. R. Harmonic analysis on homogenous spaces. *Pure and Applied Mathematics*, No. **19**. Marcel Dekker, Inc., New York, 1973. xv+361 pp. [24](#)
- [29] Yang, A. Poisson transform on vector bundles. *Trans. Amer. Math. Soc.*, **350** (3), 857-887 (1998). [4](#)

ABDELHAMID BOUSSEJRA : DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY IBN TOFAIL, KENITRA, MOROCCO

Email address: boussejra.abdelhamid@uit.ac.ma

KHALID KOUFANY : UNIVERSITÉ DE LORRAINE, CNRS, IECL, F-54000 NANCY, FRANCE

Email address: khalid.koufany@univ-lorraine.fr