

Existence-Uniqueness Theory and Small-Data Decay for a Reaction-Diffusion Model of Wildfire Spread

A. George Morgan*

Abstract

I examine some analytical properties of a nonlinear reaction-diffusion system that has been used to model the propagation of a wildfire. I establish global-in-time existence and uniqueness of bounded mild solutions to the Cauchy problem for this system given bounded initial data. In particular, this shows that the model does not allow for thermal blow-up. If the initial temperature and fuel density also satisfy certain integrability conditions, the L^2 -norms of these global solutions are uniformly bounded in time. Additionally, I use a bootstrap argument to show that small initial temperatures give rise to solutions that decay to zero as time goes to infinity, proving the existence of initial states that do not develop into travelling combustion waves.

Keywords– Wildfire spread models, solid-fuel combustion, reaction-diffusion equations in combustion theory, existence and uniqueness of solutions to reaction-diffusion equations, time decay of solutions to reaction-diffusion equations

MSC classes– 35K57, 80A25

1 Introduction

In this article, I consider a reaction-diffusion model of wildfire propagation in a fuel layer with a very large spatial extent. Fix a spatial dimension $d \geq 1$. Consider a certain amount of solid fuel (trees, shrubberies, and so on) spread throughout \mathbb{R}^d with dimensionless mass density at spatial location x and time t given by $Y(x, t)$. A fire with a dimensionless temperature field $T(x, t)$ is moving through space and consuming this fuel: temperature is normalized so that $T = 0$ is the ambient temperature of the system's environment. Also, we suppose the reaction rate governing combustion is given by the Arrhenius law

$$r(T) = \begin{cases} e^{-\frac{1}{T}} & T > 0 \\ 0 & T \leq 0. \end{cases} \quad (1.1)$$

Under these assumptions, Weber et al. [19] were the first to model the evolution of T and Y using the following Cauchy problem: given two parameters $\lambda, \beta \geq 0$,

$$\begin{cases} T_t = (\Delta - \lambda)T + Yr(T) \\ Y_t = -\beta Yr(T) \\ T|_{t=0} = T_0(x) \\ Y|_{t=0} = Y_0(x). \end{cases} \quad (1.2)$$

*Department of Mathematics, University of Toronto, 40 St. George St., Room 6290, Toronto, Ontario, CA, M5S 2E4. Institutional email: adam.morgan@mail.utoronto.ca

We always assume $Y_0(x) \geq 0$ everywhere. Up to a different choice of reaction rate, these are precisely the equations of first-order solid-fuel combustion at infinite Lewis number, with an added heat loss term [3, §1.3]. Strictly speaking, in [19] and other work $Y(x, t)$ is actually the *mass fraction* of fuel. However, I want to allow for an infinite initial amount of fuel, and in this situation the mass fraction does not make sense and it is more natural to regard $Y(x, t)$ as a mass density. For an instructive derivation of (1.2) and an explanation of why it is useful for data assimilation, see Mandel et al. [11]. The discussion in Mandel et al. highlights that $r(0) = 0$ implies no combustion takes place at the ambient temperature $T = 0$. Consequently, (1.2) admits solutions that take the form of travelling combustion waves; this had already been confirmed using formal asymptotics and numerics in [19].

In light of the existence of travelling wave solutions, (1.2) has attracted attention from researchers in both mathematical modelling and dynamical systems theory. On the more concrete side, (1.2) has been used to develop computationally efficient reduced-order models of wildfire dynamics [5, 9]. Also, Johnston [8] applied (1.2) in a modelling pipeline where remote sensing data was leveraged to estimate realistic model parameter values. Additionally, variants of the model incorporating the effects of wind [2, 11] and radiative heat transfer [1, 12, 15, 18] have also been considered. When it comes to rigorous results on the behaviour of solutions to (1.2), Billingham [4] proved the existence of travelling waves given suitable initial data and $d = 1$. Linear stability of these travelling waves was examined numerically by Varas and Vega [17]. Existence and linear stability of travelling waves for related combustion models in $d = 1$ with *finite* Lewis number has also been considered, see for example [7, 14]. There is certainly no shortage of other interesting work on travelling waves in wildland fire propagation and other solid-fuel systems, but a complete literature review is beyond the scope of this brief article.

While there is plenty of mathematical work on the properties of travelling wave solutions to (1.2), to my knowledge there are no published results on the existence and uniqueness of global-in-time solutions to (1.2) for *generic* bounded initial data (see, however, the existence result for a related model in [3, §4.4]). In particular, with the exception of the lumped-parameter analysis in [15], I have not found any estimates that preclude finite-time blow-up of $|T(x, t)|$. Thermal blow-up phenomena indeed arise in certain solid-fuel combustion models [3, §3.2], so understanding whether or not blow-up occurs in solutions to (1.2) is a meaningful question. Also, in [11], the authors use a heuristic lumped-parameter argument to predict that a fire with a sufficiently low temperature eventually burns itself out: in the jargon of combustion theory, this means the model predicts a nonzero “auto-ignition temperature”. I have not, however, found any rigorous proof of the claim that (1.2) has a nonzero auto-ignition temperature.

In this article, I prove three novel results to address the knowledge gaps discussed in the previous paragraph:

- (i) for any bounded initial state (T_0, Y_0) there exists a unique, bounded mild solution to (1.2) (to be defined precisely below) valid on the entire time interval $[0, \infty)$, provided $\lambda > 0$: in particular, thermal blow-up is impossible (corollary 2.7);
- (ii) further, if Y_0 is also integrable and T_0 is also square-integrable, then the L_x^2 norm of $T(x, t)$ is bounded uniformly in time (corollary 2.7 as well);
- (iii) given an initial state T_0 that is small in a particular physically reasonable norm, the solution to (1.2) satisfies $\lim_{t \rightarrow 0} |T(x, t)| = 0$ uniformly in x for every $\lambda \geq 0$: morally, this confirms that (1.2) indeed has a nonzero auto-ignition temperature, and that not *every* initial state gives rise to a travelling wave solution (theorem 2.10);

I hope these results prove useful to further investigations into the *nonlinear* stability of travelling waves and to the development of rigorous error estimates for numerical discretizations of (1.2).

1.1 Notation

For definitions of any of the objects appearing below, see [6].

- d always denotes a positive integer greater than or equal to 1.
- $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$.
- C always denotes some nonnegative constant that may change from line to line. If α is some real parameter, then $C(\alpha)$ denotes a constant depending on α . If $a, b \in \mathbb{R}$, we say that $a \lesssim b$ if there exists C such that $a \leq bC$. If $a, b \in \mathbb{R}$, we say that $a \simeq b$ if $a \lesssim b$ and $b \lesssim a$.
- For $k \in \mathbb{N}_0$, and a nice function $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$, $D_x^k \varphi$ denotes the set of all order k partial derivatives of φ .
- For $p \in [1, \infty]$, $L_x^p(\mathbb{R}^d)$ denotes the usual Lebesgue space of real-valued functions of $x \in \mathbb{R}^d$.
- For $k \in \mathbb{N}_0$, $H_x^k(\mathbb{R}^d)$ denotes the L_x^2 -based inhomogeneous order k Sobolev space of real-valued functions on \mathbb{R}^d , and $\dot{H}_x^k(\mathbb{R}^d)$ denotes the L_x^2 -based homogeneous order k Sobolev space of real-valued functions on \mathbb{R}^d .
- Given a Banach space A and a time interval $[0, t_*]$, we let $C_t^0 A$ denote the space of continuous curves $[0, t_*] \rightarrow A$. When the value of t_* has a substantial impact on the topic at hand, this value will be made clear.

2 Results

2.1 Preliminaries

First, we recall some decay and smoothing estimates for the linear heat equation.

Lemma 2.1.

(i) Fix any $p \in [1, \infty]$. For any $\varphi \in L_x^p$,

$$\left\| e^{t\Delta} \varphi \right\|_{L_x^p} \leq \|\varphi\|_{L_x^p} \quad \forall t > 0. \quad (2.1)$$

(ii) For any $p, q, \ell \in [1, \infty]$ with $p^{-1} = q^{-1} - \ell^{-1}$, any $\varphi \in L_x^q$, and any $k \in \mathbb{N}_0$, we have

$$\left\| D_x^k e^{t\Delta} \varphi \right\|_{L_x^p} \lesssim t^{-\frac{1}{2}(\frac{d}{\ell} + k)} \|\varphi\|_{L_x^q} \quad \forall t > 0. \quad (2.2)$$

Proof. All these bounds follow from the time-decay properties of the heat kernel on \mathbb{R}^d ; for example, see [10, pp. 41-42] for an outline of how to prove (2.2). \square

Additionally, we need a result about the function $r(T)$ defined in (1.1).

Lemma 2.2. For any $p \in \mathbb{N}_0$, there exists a constant $C(p)$ such that

$$r(T) \leq C(p)T^p \quad \forall T \in [0, 1].$$

Proof. This is obvious since $r(T)$ vanishes to all orders at $T = 0$: it is the canonical example of a smooth, non-analytic function. \square

2.2 Mild Solutions and their Basic Properties

We may use Duhamel's principle to re-formulate (1.2) in a way that lets us introduce solutions with limited regularity:

Definition 2.3. Let A, A_0 , and B be Banach subspaces of L_x^∞ , with $A \subseteq A_0$. Suppose we are given an initial datum $(T_0(x), Y_0(x)) \in A_0 \times B$. We say that $(T(x, t), Y(x, t)) \in C_t^0 A \times C_t^0 B$ is a **mild solution** of (1.2) if this pair satisfies

$$T(x, t) = e^{-\lambda t} e^{t\Delta} T_0 + \int_0^t e^{-\lambda(t-\tau)} e^{(t-\tau)\Delta} Y(x, \tau) r(T(x, \tau)) \, d\tau \quad (2.3a)$$

$$Y(x, t) = Y_0(x) \exp\left(-\beta \int_0^t r(T(x, \sigma)) \, d\sigma\right). \quad (2.3b)$$

Below, the time interval on which a mild solution is valid will be specified when it's vital to know. Immediately, one sees that mild solutions have some important properties.

- (i) First, if $Y_0(x) \geq 0$ everywhere then $Y(x, t) \geq 0$ also holds. This makes perfect sense: an initially nonnegative fuel density should never become negative.
- (ii) Second, owing to the smoothing properties of the heat flow $e^{t\Delta}$, $T(x, t)$ is smooth in x for every $t > 0$. However, $Y(x, t)$ is only as smooth as $Y_0(x)$.

2.3 *A Priori* Estimates and Existence-Uniqueness Theory

In this subsection I prove some basic *a priori* bounds on the Lebesgue and Sobolev norms of mild solutions to (1.2) with $\lambda > 0$. These bounds are then used to help construct global-in-time mild solutions of (1.2). We start with the easiest bound possible:

Proposition 2.4. For any $p \in [1, \infty]$, if (T, Y) is a mild solution of (1.2), then

$$\sup_{t>0} \|Y(x, t)\|_{L_x^p} \leq \|Y_0(x)\|_{L_x^p} \quad (2.4)$$

provided the right-hand side is finite.

Proof. Since $r(T)$ is nonnegative, $Y(x, t) \leq Y_0(x)$ almost everywhere. □

According to Mandel et al. [11, §4.1], $Y(x, t)$ is expected to settle to a nonzero steady value for $t \gg 1$. I therefore do not think one can prove a variant of the above result that includes a time-dependent decay factor on the right-hand side.

Proposition 2.5. Suppose $\lambda > 0$. If we are given an initial datum $(T_0, Y_0) \in L_x^\infty \times L_x^\infty$ and a corresponding mild solution $(T, Y) \in C_t^0 L_x^\infty \times C_t^0 L_x^\infty$ of (1.2), then

$$\|T(x, t)\|_{L_x^\infty} \leq e^{-\lambda t} \|T_0(x)\|_{L_x^\infty} + \lambda^{-1} \|Y_0(x)\|_{L_x^\infty}. \quad (2.5)$$

Proof. Using lemma 2.1 and $|r(T)| \leq 1$, we have

$$\|T\|_{L_x^\infty} \leq e^{-\lambda t} \|T_0\|_{L_x^\infty} + \sup_{t>0} \|Y r(T)\|_{L_x^\infty} \int_0^t e^{-\lambda(t-\tau)} \, d\tau \leq e^{-\lambda t} \|T_0\|_{L_x^\infty} + \lambda^{-1} (1 - e^{-\lambda t}) \|Y_0\|_{L_x^\infty}. \quad \square$$

Proposition 2.6. Suppose $\lambda > 0$. If we are given an initial datum $(T_0, Y_0) \in (L_x^2 \cap L_x^\infty) \times L_x^\infty$ and a corresponding mild solution $(T, Y) \in C_t^0 (L_x^2 \cap L_x^\infty) \times C_t^0 L_x^\infty$ of (1.2), then

$$\|T\|_{L_x^2}^2 \lesssim \min \left\{ \|T_0\|_{L_x^2}^2 + C \left(\|T_0\|_{L_x^\infty}, \|Y_0\|_{L_x^\infty}, \lambda, \beta \right) \|Y_0 - Y\|_{L_x^1}, \|T_0\|_{L_x^2}^2 e^{(\|Y_0\|_{L_x^\infty} - \lambda)t} \right\} \quad (2.6)$$

In particular, if $(T_0, Y_0) \in (L_x^2 \cap L_x^\infty) \times (L_x^1 \cap L_x^\infty)$ then

$$\sup_{t>0} \|T(x, t)\|_{L_x^2} \leq C \left(\|T_0\|_{L_x^2 \cap L_x^\infty}, \|Y_0\|_{L_x^1 \cap L_x^\infty}, \lambda, \beta \right). \quad (2.7)$$

Proof. Recall that $T(x, t)$ is smooth in x , so we can write $T_t = (\Delta - \lambda)T + Yr(T)$ where equality is understood in an L_x^2 sense. We then use a standard energy argument:

$$\begin{aligned}
\frac{d}{dt} \frac{1}{2} \|T\|_{L_x^2}^2 &= -\|T\|_{\dot{H}_x^1}^2 - \frac{\lambda}{2} \|T\|_{L_x^2}^2 + \int T(x, t) Y(x, t) r(T(x, t)) \, dx \\
&\leq \int T(x, t) Y(x, t) r(T(x, t)) \, dx \\
&= \beta^{-1} \int T(x, t) (-Y_t(x, t)) \, dx \\
&\leq \beta^{-1} \sup_{t>0} \|T(x, t)\|_{L_x^\infty} \int (-Y_t(x, t)) \, dx.
\end{aligned} \tag{2.8}$$

Integrating both sides of the above over $[0, t]$ and using proposition 2.5 gives

$$\|T(x, t)\|_{L_x^2}^2 \lesssim \|T_0(x)\|_{L_x^2}^2 + C \left(\|T_0\|_{L_x^\infty}, \|Y_0\|_{L_x^\infty}, \lambda, \beta \right) \int_0^t \int (-Y_\tau(x, \tau)) \, dx \, d\tau.$$

Then, we use Fubini's theorem and the fundamental theorem of calculus to obtain one part of (2.6). The special case (2.7) follows since $Y_0(x) - Y(x, t) \leq Y_0(x)$. For the other part of (2.6), we keep the λ term around, instead use $Tr(T) \lesssim T^2$ in (2.8), place Y in L_x^∞ , and conclude with an application of Grönwall's inequality. \square

Next, I demonstrate how these *a priori* estimates may be applied to obtain global-in-time mild solutions to (1.2).

Corollary 2.7. *Suppose $\lambda > 0$ and we are given an initial datum $(T_0, Y_0) \in L_x^\infty \times L_x^\infty$. Then, (1.2) admits a unique mild solution $(T, Y) \in C_t^0 L_x^\infty \times C_t^0 L_x^\infty$ valid for $t \in [0, \infty)$. Further, if we also have $T_0 \in L_x^2$ and $Y_0 \in L_x^1$, then the mild solution obeys (2.7).*

Proof. Fixed-point iteration (see for example [16]) can be used to establish local-in-time existence and uniqueness. Proposition 2.5 allows us to extend this local solution to a global one. With this solution in hand, we apply proposition 2.6 to conclude. \square

Consequently, solutions to (1.2) do not exhibit thermal blow-up when $\lambda > 0$. In the case $\lambda = 0$ the techniques from proposition 2.5 would yield

$$\|T(x, t)\|_{L_x^\infty} \leq \|T_0(x)\|_{L_x^\infty} + t \|Y_0(x)\|_{L_x^\infty}.$$

While this does not give a nice uniform-in-time bound, it at least says that thermal blow-up does not occur in finite time irrespective of λ .

Even though the mild solutions constructed above are smooth with respect to x , they are not necessarily classical solutions (that is, they are not also continuously differentiable in t). Indeed, the PDE $T_t = \Delta T + Yr(T)$ tells us that T_t only has the spatial regularity of Y , itself controlled by the regularity of Y_0 . However, if Y_0 is continuous, then so is T_t , and we indeed have a classical solution. In particular, the choice $Y_0 \equiv 1$ used in [11] gives rise to a unique global classical solution.

For the sake of completeness, I mention that we can obtain coarse estimates on ∇T without too much trouble:

Proposition 2.8. *Pick $p, q, \ell \in [1, \infty]$ with $p^{-1} = q^{-1} - \ell^{-1}$ and $\ell > d$. If $(T_0, Y_0) \in (L_x^p \cap L_x^\infty) \times (L_x^q \cap L_x^\infty)$ then the corresponding mild solution (T, Y) of (1.2) satisfies*

$$\|\nabla T(x, t)\|_{L_x^p} \lesssim t^{-\frac{1}{2}} \|T_0(x)\|_{L_x^p} + t^{\frac{1}{2}(1-\frac{d}{\ell})} \|Y_0\|_{L_x^q} \quad \forall t > 0. \tag{2.9}$$

In particular,

$$\|\nabla T(x, t)\|_{L_x^\infty} \lesssim t^{-\frac{1}{2}} \|T_0(x)\|_{L_x^\infty} + t^{\frac{1}{2}} \|Y_0\|_{L_x^\infty} \quad \forall t > 0. \tag{2.10}$$

Proof. Using (2.2) and $|r(T)| \leq 1$, we have

$$\|\nabla T(x, t)\|_{L_x^p} \lesssim t^{-\frac{1}{2}} \|T_0\|_{L_x^p} + \|Y_0\|_{L_x^q} \int_0^t (t - \tau)^{-\frac{1}{2}(\frac{d}{\tau} + 1)} d\tau$$

and we're all done once computing the integral. \square

Thus solutions of (1.2) do not develop sharp edges in finite time. I expect that one can get rid of the temporally growing term in (2.9) using a more thorough analysis.

2.4 Decay of Small-Data Solutions

Next I show that, if the initial temperature T_0 is sufficiently small, then $\lim_{t \rightarrow \infty} |T(x, t)| = 0$ uniformly. This confirms that (1.2) has a nonzero auto-ignition temperature up to some mild assumptions on T_0 . From our work in the last section, we know that $\lambda = 0$ gives the worst L_x^∞ temperature bounds possible, so in this section we'll set $\lambda = 0$ for concreteness: if we can establish decay with $\lambda = 0$ then it is even simpler to establish decay with $\lambda > 0$. While this means we don't have access to the existence-uniqueness results from the previous section, we can still build a temporally-decaying global solution using the assumption of small initial data and the bootstrap principle. This strategy for establishing time decay is well-known in the theory of dispersive PDEs, see for instance [13].

Throughout this section, we suppose that $\epsilon_0 \in (0, 1)$ is a small parameter (to be constrained later) and

$$\|T_0\|_{L_x^1 \cap L_x^\infty} < \epsilon_0 \quad \text{and} \quad \|Y_0\|_{L_x^\infty} < \infty. \quad (2.11)$$

The requirement that T_0 is small in L_x^1 is a minor technical one giving us access to a decay estimate. Now, given $t_* \in (0, \infty]$ and a sufficiently nice function $T(x, t)$ defined for $t \in [0, t_*]$, we set

$$\|T(x, t)\|_{X_{t_*}} \doteq \sup_{t \in [0, t_*]} \left[(1+t)^{\frac{d}{2}} \|T(x, t)\|_{L_x^\infty} + \|T(x, t)\|_{L_x^1} \right].$$

This norm naturally gives rise to a Banach space X_{t_*} defined by

$$X_{t_*} \doteq \left\{ T: [0, t_*] \rightarrow L_x^1 \cap L_x^\infty \text{ continuous} \mid \|T(x, t)\|_{X_{t_*}} < \infty \right\}.$$

The first order of business is to prove a new *a priori* estimate for solutions that are small in X_{t_*} .

Proposition 2.9. *Take $\epsilon_1 \in [\epsilon_0, 1)$. Assume that (2.11) holds and that there is a corresponding mild solution $(T, Y) \in X_{t_*} \times C_t^0 L_x^\infty$ to (1.2) valid on $[0, t_*]$. Further, assume that $\|T\|_{X_{t_*}} \leq \epsilon_1$. Then, there exists an absolute constant $C > 0$ such that*

$$\|T(x, t)\|_{X_{t_*}} \leq 2\epsilon_0 + C \|Y_0\|_{L_x^\infty} \epsilon_1^{11}. \quad (2.12)$$

Proof. Let's look at the L_x^∞ part of the X_{t_*} norm first. Together with the hypothesis of our claim, lemma 2.2 (with $p = 11$) implies $\|r(T)\|_{L_x^\infty} \leq C \epsilon_1^{11} (1+t)^{-\frac{11}{2}d}$ and $\|r(T)\|_{L_x^1} \leq C \epsilon_1^{11} (1+t)^{-5d}$. Combining these estimates with (2.1), (2.2), and (2.11), we have for $t > 1$

$$\begin{aligned} \|T(x, t)\|_{L_x^\infty} &\leq (1+t)^{-\frac{d}{2}} \epsilon_0 + \int_0^t \left\| e^{(t-\tau)\Delta} Y(x, \tau) r(T(x, \tau)) \right\|_{L_x^\infty} d\tau \\ &\leq (1+t)^{-\frac{d}{2}} \epsilon_0 + \int_0^{t/2} (t-\tau)^{-\frac{d}{2}} \|Y(x, \tau) r(T(x, \tau))\|_{L_x^1} d\tau + \int_{t/2}^t \|Y(x, \tau) r(T(x, \tau))\|_{L_x^\infty} d\tau \\ &\leq (1+t)^{-\frac{d}{2}} \epsilon_0 + C \|Y_0\|_{L_x^\infty} \epsilon_1^{11} \left[\int_0^{t/2} (1+t-\tau)^{-\frac{d}{2}} (1+\tau)^{-5d} d\tau + \int_{t/2}^t (1+\tau)^{-\frac{11}{2}d} \right] \\ &\leq (1+t)^{-\frac{d}{2}} \left[\epsilon_0 + C \|Y_0\|_{L_x^\infty} \epsilon_1^{11} \right]. \end{aligned}$$

For $t \leq 1$ we may apply an almost identical approach, but we don't split the Duhamel term and we only use (2.1). This takes care of the L_x^∞ piece. The L_x^1 estimate is obtained by a similar but even easier argument using (2.1) to control $\left\| e^{(t-\tau)\Delta} Y r(T) \right\|_{L_x^1}$. \square

Note that there is nothing particularly special about the power of $p = 11$ appearing in the above proof: any large enough natural number will do, and the rapid vanishing of $r(T)$ at $T = 0$ guarantees we can pick such a large p .

Theorem 2.10. *There exists $\epsilon_0 = \epsilon_0(\|Y_0\|_{L_x^\infty}) \in (0, 1)$ such that, under the hypothesis (2.11), (1.2) admits a unique global-in-time mild solution $(T, Y) \in X_\infty \times C_t^0 L_x^\infty$ obeying the decay estimate*

$$\|T(x, t)\|_{L_x^\infty} \lesssim \epsilon_0 (1 + t)^{-\frac{d}{2}}. \quad (2.13)$$

In particular, $\lim_{t \rightarrow \infty} |T(x, t)| = 0$ uniformly in x .

Proof. Using fixed-point iteration, one finds a $t_* > 0$ such that (1.2) has a unique local-in-time mild solution valid on $[0, t_*]$ with $T(x, t) \in X_{t_*}$. Further, the map $\tau \mapsto \|T\|_\tau$ is easily shown to be continuous for $\tau \in [0, t_*]$. Accordingly, for sufficiently small t_* we know $\|T\|_{X_{t_*}} \leq 3\epsilon_0$. Now, apply proposition 2.9 with $\epsilon_1 = 3\epsilon_0$ to discover

$$\|T\|_{X_{t_*}} \leq 2\epsilon_0 \left(1 + C \|Y_0\|_{L_x^\infty} \epsilon_0^{10}\right).$$

By choosing $\epsilon_0 < \frac{1}{2} \left(C \|Y_0\|_{L_x^\infty}\right)^{-\frac{1}{10}}$, the above becomes $\|T\|_{X_{t_*}} \leq \frac{5}{2}\epsilon_0$. The bootstrap principle (see for example [16]) allows us to iterate this argument and produce a global solution satisfying $\|T\|_{X_\infty} \lesssim \epsilon_0$. \square

The ϵ_0 from theorem 2.10 should be regarded as a lower bound on the auto-ignition temperature for the model (1.2), but my methods cannot determine how sharp this bound is.

3 Acknowledgements

I'd like to thank Carrie Clark for helpful comments on the draft manuscript, and Dominic Shillingford for invaluable discussions.

References

- [1] M. I. Asensio and L. Ferragut. On a wildland fire model with radiation. *Internat. J. Numer. Methods Engrg.*, 54(1):137–157, 2002.
- [2] Petro Babak, Anne Bourlioux, and Thomas Hillen. The effect of wind on the propagation of an idealized forest fire. *SIAM J. Appl. Math.*, 70(4):1364–1388, 2009.
- [3] Jerrold Bebernes and David Eberly. *Mathematical problems from combustion theory*, volume 83 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1989.
- [4] John Billingham. Phase plane analysis of one-dimensional reaction diffusion waves with degenerate reaction terms. *Dyn. Stab. Syst.*, 15(1):23–33, 2000.
- [5] Felix Black, Philipp Schulze, and Benjamin Unger. Efficient wildland fire simulation via nonlinear model order reduction. *Fluids*, 6(8), 2021.
- [6] Lawrence Craig Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.

- [7] Anna Ghazaryan and Christopher K. R. T. Jones. On the stability of high Lewis number combustion fronts. *Discrete Contin. Dyn. Syst.*, 24(3):809–826, 2009.
- [8] Andrew Johnston. Wildfire modeling with data assimilation, December 2022. Master’s thesis, available at scholarsarchive.byu.edu/etd/9788/.
- [9] Alan Lattimer, Jeff Borggaard, Serkan Gugercin, Kray Luxbacher, and Brian Lattimer. Computationally efficient wildland fire spread models. 07 2016.
- [10] Felipe Linares and Gustavo Ponce. *Introduction to nonlinear dispersive equations*. Universitext. Springer, New York, second edition, 2015.
- [11] Jan Mandel, Lynn S. Bennethum, Jonathan D. Beezley, Janice L. Coen, Craig C. Douglas, Minjeong Kim, and Anthony Vodacek. A wildland fire model with data assimilation. *Math. Comput. Simulation*, 79(3):584–606, 2008.
- [12] Koondanibha Mitra, Qiyao Peng, and Cordula Reisch. Studying wildfire fronts using advection-diffusion-reaction models. 2024. Arxiv preprint, available at arxiv.org/abs/2401.17468.
- [13] A. George Morgan. Scattering for the quartic generalized Benjamin-Bona-Mahony equation, 2024. Arxiv preprint, available at arxiv.org/abs/2402.09605.
- [14] Abdolrahman Razani. On the existence of premixed laminar flames. *Bull. Austral. Math. Soc.*, 69(3):415–427, 2004.
- [15] Cordula Reisch, Adrián Navas-Montilla, and İlhan Özgen Xian. Analytical and numerical insights into wildfire dynamics: exploring the advection–diffusion–reaction model. *Computers & Mathematics with Applications*, 158:179–198, 2024.
- [16] Terence Tao. *Nonlinear dispersive equations: local and global analysis*, volume 106 of *CBMS Regional Conference Series in Mathematics*. Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2006.
- [17] Fernando Varas and José M. Vega. Linear stability of a plane front in solid combustion at large heat of reaction. *SIAM J. Appl. Math.*, 62(5):1810–1822, 2002.
- [18] R. O. Weber. Toward a comprehensive wildfire spread model. *International Journal of Wildland Fire*, 1(4):245–248, 1991.
- [19] R. O. Weber, G. N. Mercer, H. S. Sidhu, and B. F. Gray. Combustion waves for gases ($Le = 1$) and solids ($Le \rightarrow \infty$). *Proc. Roy. Soc. London Ser. A*, 453(1960):1105–1118, 1997.