

Kerr-MOG-(A)dS black hole and its shadow in scalar-tensor-vector gravity theory

Wentao Liu,¹ Di Wu,² Xiongjun Fang,^{1,*} Jiliang Jing,¹ and Jieci Wang^{1,†}

¹*Department of Physics, Key Laboratory of Low Dimensional Quantum Structures and Quantum Control of Ministry of Education, and Synergetic Innovation Center for Quantum Effects and Applications, Hunan Normal University, Changsha, Hunan 410081, P. R. China*

²*School of Physics and Astronomy, China West Normal University, Nanchong, Sichuan 637002, P. R. China*

The scalar-tensor-vector gravity (STVG) theory has attracted significant interest due to its ability to effectively address the issue of galaxy rotation curves and clusters of galaxies without considering the influence of dark matter. In this paper, we construct rotating black hole solutions with a cosmological constant in the STVG theory (i.e., Kerr-MOG-(A)dS black hole solutions), where the import of a gravitational charge as a source modifies the gravitational constant, determined by $G = G_N(1 + \alpha)$. For Kerr-MOG-dS spacetime, the observer is situated at a specific location within the domain of outer communication, rather than being located infinitely far away. Since black hole shadows are shaped by light propagation in spacetime, the interaction between the MOG parameter and the cosmological constant is expected to produce novel effects on these shadows. As the cosmological constant Λ increases, the apparent size of the black hole shadow on the screen decreases. Additionally, the shadow expands with an increase in the MOG parameter α , reaching a maximum at a certain value, and its shape becomes more rounded under an arbitrary rotation parameter, which leads to degeneracy between different black hole parameters. However, by employing numerical backward ray-tracing techniques, we have found that gravitational lensing and the frame-dragging effect effectively distinguish this degeneracy. Our work contributes to a deeper understanding of black holes in modified gravity, their observational signatures, and constraints.

I. INTRODUCTION

Although the no-hair theorem suggests that black holes are characterized by their mass, spin, and electric charge, however, the electric charge is usually negligible for astrophysical black holes. The reason is that a black hole with mass M and electric charge Q will not gravitationally attract particles with mass m and electric charge e if $eQ > Mm$. Given the m/e ratio for electrons is approximately 10^{-21} , it is highly unlikely for black holes to maintain a significant electric charge [1]. Moreover, processes including vacuum polarization, pair production, and neutralization ensure that stellar-mass black holes cannot sustain significant electric charge. Any accumulated charge dissipates quickly and escapes observation by gravitational wave detectors [1–3]. However, by performing general-relativistic simulations of charged black holes, Bozola et al. have proposed that the gravitational wave event GW150914 could have a charge-to-mass ratio as high as 0.3 [4–6]. Consequently, the term “charge” could be more appropriately interpreted as dark charge or gravitational charge, sparking renewed interest in the scalar-tensor-vector gravity (STVG) theory, also known as modified gravity (MOG) theory, which was first proposed nearly two decades ago [7]. The STVG theory introduces additional massive vector and scalar fields to the metric tensor field, positing that the associated “charge” is a gravitational charge denoted by $Q_5 \equiv \kappa M$, which is determined by a body’s mass M and its interaction with the vector field.

On the observational side, despite numerous attempts to detect dark matter, all experimental efforts up to now have yet to

achieve recognized success [8, 9]. Therefore, on the theoretical side, STVG theory has attracted a great deal of attention because it not only successfully explains the rotation curves of galaxies and the properties of galaxy clusters, but also does so without considering dark matter [10–14]. In addition, observational data from the Event Horizon Telescope (EHT) on the shadows of Sgr A* and M87* [15–17] support the predictions of STVG theory and there are criteria to distinguish this theory from general relativity [18]. The axially symmetric solution in STVG theory provided by Moffat et al. is called the Kerr-MOG black hole [19]. To date, several new black hole solutions related to this theory have been obtained [20–22]. Building on the established solutions in STVG theory, extensive research has been conducted on various aspects of black holes, including their thermodynamic properties [23], geodesic motion [24], shadows [25–33], and quasi-normal modes [34, 35], etc.

Despite much progress in the last few years in constructing black hole solutions in STVG theory and investigating their physical properties, rotating (A)dS black holes in STVG theory remain to be the virgin territory and thus need to be explored deeply. On the other hand, our current universe is in a state of accelerated expansion, and the simplest explanation for such accelerated expansion is a positive cosmological constant [36, 37]. Thus, it is significant to construct rotating dS black hole solutions in STVG theory. Additionally, the study of rotating AdS black holes has already shed light on the nature of gravity through gauge-gravity dualities [38–40], so it is very important and remarkable to seek the rotating AdS black hole solution in STVG theory. These three aspects motivate us to conduct the present work.

In this work, we aim to extend the Kerr-MOG solution [19] to more general cases with a cosmological constant, thereby obtaining Kerr-MOG-(A)dS black holes. Given the recent groundbreaking observations by the EHT, theoretical research

* fangxj@hunnu.edu.cn

† jcwang@hunnu.edu.cn

on the properties of black holes frequently focuses on the analysis of black hole images [41]. Hence, we have also investigated the effects of the MOG parameter and a positive cosmological constant on black hole shadows, as well as the numerical estimation of the angular radius of the supermassive black holes Sgr A* and M87* within this theoretical framework. The remaining part of this paper is organized as follows. In Sec. II, we provide a concise overview of the STVG theory and give the Kerr-like solution of the field equations with a cosmological constant, i.e., Kerr-MOG-(A)dS black holes. In Sec. III, utilizing the geodesic equation, we derive the orbital equations for photons in the context of Kerr-MOG-dS spacetime. In Sec. IV, we examine the shadows of black holes, with a primary focus on the apparent shapes and distortions of the shadow. Also, we numerically calculate these two observables, R_s and δ_s , to characterize the shadow. Sec. V focuses on providing an overview of our conclusions while also contemplating the possible development and exploration of related research topics in the future.

II. STVG THEORY AND BLACK HOLE SOLUTION

In this section, we first briefly review the STVG theory and then proceed to construct new rotating black hole solutions using the Kerr-Schild-(A)dS metric ansatz as a seed solution.

A. STVG theory

The action corresponding to STVG theory is given by [7]

$$\mathcal{S} = \mathcal{S}_M + \mathcal{S}_G + \mathcal{S}_\phi + \mathcal{S}_S, \quad (1)$$

with

$$\begin{aligned} \mathcal{S}_G &= \frac{1}{16\pi} \int d^4x \sqrt{-g} \left[\frac{1}{G} (R - 2\Lambda) \right], \\ \mathcal{S}_\phi &= \int d^4x \sqrt{-g} \left(-\frac{1}{4} B^{ab} B_{ab} + \frac{1}{2} \mu^2 \phi^a \phi_a \right), \\ \mathcal{S}_S &= \int d^4x \sqrt{-g} \left[\frac{1}{G^3} \left(\frac{1}{2} g^{ab} \nabla_a G \nabla_b G - V(G) \right) \right. \\ &\quad \left. + \frac{1}{\mu^2 G} \left(\frac{1}{2} g^{ab} \nabla_a \mu \nabla_b \mu - V(\mu) \right) \right], \end{aligned} \quad (2)$$

where the symbol \mathcal{S}_M represents the matter action, \mathcal{S}_G represents the action for Einstein gravity, and R and Λ are the Ricci scalar and cosmological constant, respectively. Additionally, the action \mathcal{S}_ϕ describes a Proca field ϕ^a with mass μ . The field strength, represented by $B_{ab} = \partial_a \phi_b - \partial_b \phi_a$, satisfies the following equations:

$$\nabla_b B^{ab} = \frac{1}{\sqrt{-g}} \partial_b \left(\sqrt{-g} B^{ab} \right) = 0, \quad (3)$$

$$\partial_c B_{ab} + \partial_a B_{bc} + \partial_b B_{ca} = 0. \quad (4)$$

The potentials $V(G)$ and $V(\mu)$ are linked to the scalar fields $G(x)$ and $\mu(x)$ within the action \mathcal{S}_S , respectively.

Under the assumption of vacuum solution, the action \mathcal{S} can be simplified to:

$$\begin{aligned} \mathcal{S} &= \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G} (R - 2\Lambda) \right. \\ &\quad \left. - \frac{1}{4} B^{ab} B_{ab} + \frac{1}{2} \mu^2 \phi^a \phi_a \right]. \end{aligned} \quad (5)$$

The field equation of motion derived from varying the action in Eq. (5) with respect to the metric is given by

$$\mathcal{G}_{ab} + \Lambda g_{ab} = -8\pi G T_{ab}, \quad (6)$$

where $\mathcal{G}_{ab} = R_{ab} - \frac{1}{2} R g_{ab}$ is Einstein tensor, and the energy-momentum tensor is

$$\begin{aligned} T_{ab} &= -\frac{1}{4\pi} \left(B_a{}^c B_{bc} - \frac{1}{4} g_{ab} B^{cd} B_{cd} \right) \\ &\quad + \frac{\mu^2}{4\pi} \left(\phi_a \phi_b - \frac{1}{2} g_{ab} \phi^c \phi_c \right). \end{aligned} \quad (7)$$

The first and second terms of the energy-momentum tensor are of the order $\sim (\partial\phi)^2$ and $\sim \mu^2 \phi^2$, respectively. To solve the equation (6), one can typically use the weak field approximation and neglect the higher-order terms including g^2 , $g\phi$, and ϕ^2 [10, 14]. Concurrently, according to Refs. [11], the particle mass of the ϕ^a field in the present universe can be fitted as $m_\phi \sim 10^{-28} eV$, making it negligible for a black hole solution. In STVG theory, it is important to note that the fifth force charge Q_5 is tied to a particle's inertial mass as $Q_5 = \kappa M$. This parameter κ affects the gravitational constant, given by $G = G_N + \kappa^2$, with the convention $\kappa^2 = \alpha G_N$. Then, we can obtain the relationship as $G = (1 + \alpha) G_N$, which can also be thought of as a constant unaffected by the spacetime coordinates. By setting α to zero, the theory reduces to General Relativity (GR). Thus, α can be considered as a parameter that measures the deviation of STVG from GR.

In the static case, assuming the vector field ϕ_a takes the form [19]

$$\phi_a = \left(-\sqrt{\alpha G_N} \frac{M}{r}, 0, 0, 0 \right), \quad (8)$$

and setting $G_N = 1$ and $\beta = \alpha/(1 + \alpha)$, the corresponding asymptotic (A)dS metric solution is given by [22]

$$\begin{aligned} ds^2 &= - \left(1 - \frac{2M_D}{r} + \frac{\beta M_D^2}{r^2} - \frac{\Lambda}{3} r^2 \right) dt^2 \\ &\quad + \left(1 - \frac{2M_D}{r} + \frac{\beta M_D^2}{r^2} - \frac{\Lambda}{3} r^2 \right)^{-1} dr^2 \\ &\quad + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\varphi^2. \end{aligned} \quad (9)$$

The symbol M_D stands for the ADM mass, which can be related to the Newtonian mass M through the equation $M_D = (1 + \alpha)M$ [29].

B. Rotating-(A)dS solutions

Rotating black hole solutions play a crucial role in astrophysics. Considering the total spin angular momentum repre-

sented by $J = Ma$, adjustments to the vector potential and metric, equations (8) to (9), are necessary to incorporate the spin parameter. In order to obtain the axisymmetric solutions for asymptotically (A)dS spacetimes, we adopt the Kerr-Schild-(A)dS ansatz [42, 43]

$$g_{ab} = \bar{g}_{ab} - 2\mathcal{H}k_a k_b, \quad (10)$$

where \mathcal{H} is an arbitrary scalar function. The vector \mathbf{k} is null with respect to both the full metric g_{ab} and the Kerr-(A)dS metric \bar{g}_{ab} , with the latter serving as the background. Alternatively, by introducing the coordinate $(\bar{t}, \bar{r}, \bar{\vartheta}, \bar{\varphi})$, the line elements can be expressed as follows:

$$d\bar{s}^2 = - \left(1 - \frac{\Lambda}{3}\bar{r}^2\right) \frac{\Delta_{\bar{\vartheta}}}{\chi} d\bar{t}^2 + \frac{\Sigma}{\Delta_{\bar{r}}} d\bar{r}^2 + \frac{\Sigma}{\Delta_{\bar{\vartheta}}} d\bar{\vartheta}^2 + \frac{(\bar{r}^2 + a^2) \sin^2 \bar{\vartheta}}{\chi} d\bar{\varphi}^2 - 2\mathcal{H} \left(\frac{\Delta_{\bar{\vartheta}}}{\chi} d\bar{t} + \frac{\Sigma}{\Delta_{\bar{r}}} d\bar{r} - \frac{a \sin^2 \bar{\vartheta}}{\chi} d\bar{\varphi} \right)^2, \quad (11)$$

with

$$\begin{aligned} \Sigma &= \bar{r}^2 + a^2 \cos^2 \bar{\vartheta}, & \Delta_{\bar{\vartheta}} &= 1 + \frac{\Lambda}{3} a^2 \cos^2 \bar{\vartheta}, \\ \chi &= 1 + \frac{\Lambda}{3} a^2, & \Delta_{\bar{r}} &= (\bar{r}^2 + a^2) \left(1 - \frac{\Lambda}{3} \bar{r}^2\right). \end{aligned} \quad (12)$$

For the metric ansatz (11), the Ricci scalar is

$$R = g^{ab} R_{ab} = 4\Lambda - 2 \left(\frac{\partial^2}{\partial \bar{r}^2} + \frac{4\bar{r}}{\Sigma} \frac{\partial}{\partial \bar{r}} + \frac{2}{\Sigma} \right) \mathcal{H}, \quad (13)$$

and the vacuum gravitational field equations (with cosmological constant) are $R = 4\Lambda$. Therefore, one can obtain the following partial differential equation for the \mathcal{H} function:

$$\left(\frac{\partial^2}{\partial \bar{r}^2} + \frac{4\bar{r}}{\Sigma} \frac{\partial}{\partial \bar{r}} + \frac{2}{\Sigma} \right) \mathcal{H} = 0. \quad (14)$$

The above equation has the general solution,

$$\mathcal{H} = \frac{GQ_5^2}{2\Sigma} - \frac{GM\bar{r}}{\Sigma}, \quad (15)$$

where GM and GQ_5^2 are constants of integration, as in Ref. [19]. In STVG theory, the charge $Q_5 > 0$ represents a fifth force with a gravitational origin and is proportional to the mass of the source particle, given by $Q_5^2 = \alpha G_N M^2$. Together with the relation $G = G_N(1 + \alpha)$, \mathcal{H} can be rewritten as follows:

$$\mathcal{H} = \frac{G_N^2 M^2 \alpha (1 + \alpha)}{2\Sigma} - \frac{G_N M \bar{r} (1 + \alpha)}{\Sigma}. \quad (16)$$

Then, the seed metric (11) with scalar function (16) can be expressed as

$$ds^2 = - \frac{\Delta_{\bar{\vartheta}}}{\chi} \left(1 - \frac{\Lambda \bar{r}^2}{3} + \frac{h \Delta_{\bar{\vartheta}}}{\chi \Sigma}\right) d\bar{t}^2 + \frac{\Sigma}{\Delta_{\bar{r}}} \left(1 - \frac{h}{\Delta_{\bar{r}}}\right) d\bar{r}^2 + \frac{\Sigma}{\Delta_{\bar{\vartheta}}} d\bar{\vartheta}^2 + \left(\frac{\bar{r}^2 + a^2}{\chi} - \frac{ha^2 \sin^2 \bar{\vartheta}}{\chi^2 \Sigma} \right) \sin^2 \bar{\vartheta} d\bar{\varphi}^2 - \frac{2h \Delta_{\bar{\vartheta}}}{\chi \Delta_{\bar{r}}} d\bar{t} d\bar{r} + \left(\frac{\Delta_{\bar{\vartheta}} d\bar{t}}{\chi \Sigma} + \frac{d\bar{r}}{\Delta_{\bar{r}}} \right) \frac{2ha \sin^2 \bar{\vartheta}}{\chi} d\bar{\varphi}, \quad (17)$$

where

$$h = G_N M (1 + \alpha) (G_N M \alpha - 2\bar{r}). \quad (18)$$

The non-zero components of the effective tensor, defined as $E_{ab} = R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} + 8\pi G T_{ab}$, associated with the metric, are

$$\begin{aligned} E_{00} &= G_\alpha \Delta_{\bar{\vartheta}} \left[(1 - \bar{r}^2 \Lambda / 3) \Sigma_2 + h \Delta_{\bar{\vartheta}} \right] + 8\pi G T_{00}, \\ E_{01} &= G_\alpha h \chi \Delta_{\bar{r}}^{-1} \Delta_{\bar{\vartheta}} + 8\pi G T_{01}, \\ E_{03} &= -G_\alpha \Delta_{\bar{\vartheta}} (h + 2\Delta_{\bar{r}}) a \sin^2 \bar{\vartheta} + 8\pi G T_{03}, \\ E_{11} &= G_\alpha (h - \Delta_{\bar{r}}) \chi^2 \Delta_{\bar{r}}^{-2} \Sigma^2 + 8\pi G T_{11}, \\ E_{13} &= -G_\alpha h \chi \Delta_{\bar{\vartheta}}^{-1} \Sigma a \sin^2 \bar{\vartheta} + 8\pi G T_{13}, \\ E_{22} &= G_\alpha \chi^2 \Delta_{\bar{\vartheta}}^{-1} \Sigma^2 + 8\pi G T_{22}, \\ E_{33} &= G_\alpha \sin^2 \bar{\vartheta} \left[(\bar{r}^2 + a^2) \Sigma_2 + ha^2 \sin^2 \bar{\vartheta} \right] + 8\pi G T_{33}, \end{aligned} \quad (19)$$

where

$$\begin{aligned} G_\alpha &= G_N^2 M^2 \alpha (1 + \alpha) \chi^{-2} \Sigma^{-3}, \\ \Sigma_2 &= \Sigma + a^2 \Delta_{\bar{\vartheta}} - a^2 (1 - \bar{r}^2 \Lambda / 3) (\cos^2 \bar{\vartheta} - \sin^2 \bar{\vartheta}). \end{aligned} \quad (20)$$

The physical significance of these parameters becomes clearer in the Boyer-Lindquist coordinates $(t, r, \vartheta, \varphi)$, which offer a more appropriate asymptotic form for the metric. Applying the Boyer-Lindquist transformation,

$$\begin{aligned} d\bar{r} &= dr, & d\bar{\vartheta} &= d\vartheta, \\ d\bar{t} &= dt - \frac{h}{(1 - \frac{\Lambda}{3} r^2) (\Delta_{\bar{r}} + h)} dr, \\ d\bar{\varphi} &= d\varphi - \frac{\Lambda}{3} a dt - \frac{ah}{(r^2 + a^2) (\Delta_{\bar{r}} + h)} dr, \end{aligned} \quad (21)$$

the metric (17) can be rewritten in the following form,

$$ds^2 = - \frac{\Delta_r}{\Sigma} \left(dt - \frac{a \sin^2 \vartheta}{\chi} d\varphi \right)^2 + \frac{\Sigma}{\Delta_r} dr^2 + \frac{\Sigma}{\Delta_{\vartheta}} d\vartheta^2 + \frac{\Delta_{\vartheta} \sin^2 \vartheta}{\Sigma} \left(a dt - \frac{(r^2 + a^2)}{\chi} d\varphi \right)^2, \quad (22)$$

with

$$\Delta_{\vartheta} = \Delta_{\bar{\vartheta}}, \quad \Delta_r = \Delta_{\bar{r}} - 2G_N M_D r + \frac{\alpha G_N^2 M_D^2}{(1 + \alpha)}.$$

To ensure the effective tensor $E_{ab} = 0$ is satisfied, we can simply obtain the MOG vector potential as:

$$\phi = \phi_a dx^a = \frac{M\sqrt{G_N}\alpha r}{\Sigma} \left(-dt + \frac{a \sin^2 \vartheta}{\chi} d\varphi \right). \quad (23)$$

In Appendix A, we provide the specific forms of all the energy-momentum tensors contained in equations (19), which are sourced from the MOG vector potential (23). Equations (22) and (23) describe the Kerr-MOG-(A)dS black hole solutions. When $\Lambda = 0$, the solutions recovers the usual Kerr-MOG solution [19]; when $a = 0$, it reduces to the static (A)dS cases [22]. This metric (22) has the similar form as the Kerr-Newman-(A)dS metric. However, in this case, the ADM mass M_D is associated with the coupling parameter α , and the source of the vector field is gravitational. This distinction is crucial because astrophysical objects, including black holes, are electrically neutral.

The equations of the horizons are given by:

$$\Delta_r = \frac{\Lambda}{3} (r - r_m)(r - r_h)(r_c - r)(r + r_b) = 0, \quad (24)$$

where r_m , r_h , and r_c correspond to the Cauchy horizon, event horizon, and dS cosmological horizon, respectively [22]. The parameters r_b , r_m , r_h , and r_c satisfy the condition $r_b + r_m + r_h + r_c = 0$. Moreover, $r_c \geq r_h > r_m > 0$ and r_b is negative.

III. PHOTON ORBITS

In this section, we provide a brief derivation of photon trajectories in the gravitational field of Kerr-MOG-(A)dS black holes. When a black hole is located between the observer and the extended background source, not all photons emitted by the source can reach the observer after being deflected by the black hole gravitational field. The apparent shape of the black hole is represented by its shadow contour, which is closely related to the geodesics of photons within the black hole spacetime.

The geodesics corresponding to photons geometry are defined by the Hamilton-Jacobi equation:

$$\frac{\partial \mathcal{S}}{\partial \tau} = -\frac{1}{2} g^{ab} \frac{\partial \mathcal{S}}{\partial x^a} \frac{\partial \mathcal{S}}{\partial x^b}, \quad (25)$$

where τ is an affine parameter along the null geodesics, and \mathcal{S} is the Jacobi action. We can set up the Killing fields $\xi^a = (\partial/\partial t)^a$ and $\psi^a = (\partial/\partial \varphi)^a$, then the corresponding energy \mathcal{E} and the angular momentum in the direction of the axis of symmetry L can be defined as [44–46]:

$$\mathcal{E} = -g_{ab} \xi^a \dot{x}^b = -g_{00} \dot{t} - g_{03} \dot{\varphi}, \quad (26)$$

$$L = g_{ab} \psi^a \dot{x}^b = g_{03} \dot{t} + g_{33} \dot{\varphi}. \quad (27)$$

When \mathcal{S} is separable, the Jacobi action for the photons can be expressed in a straightforward and general form:

$$\mathcal{S} = -\mathcal{E}t + L\varphi + \mathcal{S}_r(r) + \mathcal{S}_\vartheta(\vartheta). \quad (28)$$

Combining Eqs. (25) and (28) and setting $G_N = 1$, we obtain two equations of motion for the propagating photons

$$\dot{t} = \frac{(r^2 + a^2)k}{\Sigma \Delta_r} + \frac{a(L\chi - a\mathcal{E} \sin^2 \vartheta)}{\Sigma \Delta_\vartheta}, \quad (29)$$

$$\dot{\varphi} = \frac{a\chi k}{\Sigma \Delta_r} + \frac{\chi(L\chi \csc^2 \vartheta - a\mathcal{E})}{\Sigma \Delta_\vartheta}, \quad (30)$$

with

$$k = (r^2 + a^2)\mathcal{E} - aL\chi. \quad (31)$$

Substituting Eqs. (29) and (30) into the null geodesics equation $0 = -g_{ab} \dot{x}^a \dot{x}^b$, one can obtain

$$\frac{k^2 - \rho^4 \dot{r}^2}{\Delta_r} - \frac{\rho^4 \dot{\vartheta}^2 + (L\chi \csc \vartheta - a\mathcal{E} \sin \vartheta)^2}{\Delta_\vartheta} = 0. \quad (32)$$

By using the Carter constant \mathcal{K} to separate variables in the above equation, we can obtain two additional equations of motion for photons [47]

$$\dot{r} = \frac{1}{\Sigma} \sqrt{\mathcal{R}(r)}, \quad (33)$$

$$\dot{\vartheta} = \frac{1}{\Sigma} \sqrt{\Theta(\vartheta)}. \quad (34)$$

The explicit forms of the radial and angular equations are as follows:

$$\mathcal{R} = k^2 - [\mathcal{K} + (L\chi - a\mathcal{E})^2] \Delta_r, \quad (35)$$

$$\Theta = [\mathcal{K} + (L\chi - a\mathcal{E})^2] \Delta_\vartheta - (a\mathcal{E} \sin \vartheta - L\chi \csc \vartheta)^2. \quad (36)$$

These equations determine the behavior of light propagation within the spacetime influenced by rotating black holes.

The radial motion equation of the photon is of particular interest, as the silhouette of the black hole shadow can be derived from the orbits with constant $r = r_p$ that fulfill the conditions

$$\mathcal{R}(r)|_{r=r_p} = 0, \quad \frac{d}{dr} \mathcal{R}(r)|_{r=r_p} = 0. \quad (37)$$

To determine the contour of black hole shadow, one can introduce the impact parameters $\xi = L/\mathcal{E}$ and $\eta = \mathcal{K}/\mathcal{E}^2$, and solve the system of equations accordingly [48]. The resulting impact parameters will provide the necessary information:

$$\xi = \frac{M_D (a^2 + 2M\alpha r_p - 3r_p^2) + \chi r_p (a^2 + r_p^2)}{a\chi [M - r_p + M\alpha + \frac{1}{3}r_p\Lambda (a^2 + 2r_p^2)]}, \quad (38)$$

$$\eta = \frac{r_p^2 \mathcal{W} - M_D \Lambda r_p^5 (2a^2 + \frac{4}{3}M\alpha r_p) + r_p^6 - r_p^6 \chi^2}{a^2 [M - r_p + M\alpha + \frac{1}{3}r_p\Lambda (a^2 + 2r_p^2)]^2}, \quad (39)$$

where

$$\mathcal{W} = 4Mr_p(1 + \alpha) [a^2 + M^2\alpha(1 + \alpha)] - 4M^2\alpha(1 + \alpha)\Delta_r - r_p^2 [r_p - 3M(1 + \alpha)]^2. \quad (40)$$

These two impact parameters are equivalent to the results provided in Ref. [26] when $\Lambda = 0$. For simplicity, we set $M = 1$ in subsequent calculations.

IV. SHADOWS OF BLACK HOLES

While AdS solutions play a crucial role in AdS/CFT correspondences, they lack empirical support from specific astronomical observations. Therefore, in this section, we primarily focus on the black hole shadows in Kerr-MOG-dS spacetime. For this, we need to adopt the analysis approach of A. Grenzebach [49] and the numerical backward ray-tracing method [50–54] to investigate the shadow of Kerr-MOG black holes in the presence of a positive cosmological constant.

A. Apparent shape

In the Kerr-MOG-dS spacetime, there exist two unstable light orbits within the domain of outer communication, which extends from the outer horizon r_h to the cosmological horizon r_c : a prograde orbit and a retrograde orbit. Both orbits serve as limit curves for lightlike geodesics that spiral towards them. Similarly to the circular lightlike geodesics in the Schwarzschild spacetime [55], all spherical lightlike geodesics in the domain of outer communication are also unstable and can serve as limit curves for past-oriented light rays from an observer position that spiral towards them. Hence, the photon region determines the boundary curve of the black hole shadow. Assume that the observer is positioned at (r_0, ϑ_0) , where $r_h < r_0 < r_c$. We define an orthonormal tetrad as [56]:

$$\begin{aligned} e_0 &= \frac{r^2 + a^2}{\sqrt{\Sigma\Delta_r}} \partial_t + \frac{a\chi}{\sqrt{\Sigma\Delta_r}} \partial_\varphi \Big|_{(r_0, \vartheta_0)}, \\ e_1 &= \sqrt{\frac{\Delta_\vartheta}{\Sigma}} \partial_\vartheta \Big|_{(r_0, \vartheta_0)}, \\ e_2 &= -\frac{\chi \csc \vartheta}{\sqrt{\Sigma\Delta_\vartheta}} \partial_\varphi - \frac{a \sin \vartheta}{\sqrt{\Sigma\Delta_\vartheta}} \partial_t \Big|_{(r_0, \vartheta_0)}, \\ e_3 &= -\sqrt{\frac{\Delta_r}{\Sigma}} \partial_r \Big|_{(r_0, \vartheta_0)}. \end{aligned} \quad (41)$$

It is easy to verify that $e_0, e_1, e_2,$ and e_3 are orthonormal. In this chosen tetrad, e_0 represents the observer four-velocity, e_3 corresponds to the spatial direction towards the center of the black hole, and the combination $e_0 \pm e_3$ is tangential to the principal null congruences of the metric. For each light ray $\lambda(s)$ with coordinate representation $t(s), r(s), \vartheta(s), \varphi(s)$, the general form of tangent vector is

$$\dot{\lambda} = \dot{t} \partial_t + \dot{r} \partial_r + \dot{\vartheta} \partial_\vartheta + \dot{\varphi} \partial_\varphi. \quad (42)$$

Following the approach in Ref. [49], the tangent vector at the observer can be expressed as

$$\dot{\lambda} = \gamma (\sin \theta \cos \psi e_1 + \sin \theta \sin \psi e_2 + \cos \theta e_3 - e_0), \quad (43)$$

where γ is a scalar factor and can be solved by substituting Eqs. (29), (30), (33), and (34), and Eq. (43) into Eq. (42). The result gives

$$\gamma = -\frac{k}{\sqrt{\Sigma\Delta_r}}, \quad (44)$$

and then yield

$$\sin \theta = \frac{\sqrt{\eta\Delta_r + (a - \chi\xi)^2 \Delta_r}}{r^2 + a^2 - a\chi\xi} \Big|_{r=r_0}, \quad (45)$$

$$\sin \psi = \frac{\csc \vartheta (a \cos^2 \vartheta - a + \chi\xi)}{\sqrt{\eta\Delta_\vartheta + (a - \chi\xi)^2 \Delta_\vartheta}} \Big|_{\vartheta=\vartheta_0}. \quad (46)$$

In order to use analytical parameter to represent the boundary curve of the shadow [49], we utilized the stereographic projection from the celestial sphere onto a plane, where standard Cartesian coordinates are employed,

$$x(r_p) = -2 \tan \left[\frac{\theta(r_p)}{2} \right] \sin [\psi(r_p)], \quad (47)$$

$$y(r_p) = -2 \tan \left[\frac{\theta(r_p)}{2} \right] \cos [\psi(r_p)]. \quad (48)$$

This is a set of parameter equations concerning the radius of the photon sphere r_p [55].

For the non-rotating black hole case with $a = 0$, the expression for the impact parameter has a singularity, necessitating separate consideration of the static case. Using the circular orbit condition (37) and setting $\eta + \xi^2 \chi^2 := 1/X$ [49], the Eqs. (45)-(46) have the form:

$$\sin \theta^{(a=0)} = \frac{\sqrt{\Delta_r|_{a=0}}}{r^2 \sqrt{X}} \Big|_{r=r_0}, \quad (49)$$

$$\sin \psi^{(a=0)} = \sqrt{X} \chi \xi \csc \vartheta \Big|_{\vartheta=\vartheta_0}, \quad (50)$$

where

$$X = \frac{1}{2r_p^2} - \frac{(1 + \alpha)}{2r_p^3} - \frac{\Lambda}{3}, \quad (51)$$

and the radius of the photon sphere is

$$r_p = \frac{3}{2} (1 + \alpha) \left(1 + \sqrt{1 - \frac{8\alpha}{9(1 + \alpha)}} \right). \quad (52)$$

In this case, the system of Eqs. (47)-(48) will become parameter equations only concerning the impact parameter ξ . Note that the expression of the photon sphere radius is independent of Λ . From this, one can deduce that the shape of a static black hole is a standard circle, with its radius dependent on the MOG parameter α .

For simplicity, we consider fixing the position of the observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$. Where the angle ϑ_0 is defined as the inclination angle between the observer's line of sight and the rotation axis of the black hole. Form Eqs. (47) and (48) one can depict the contour of the black hole shadow under varying MOG parameter α , spin a , or cosmological constant Λ . In Fig. 1, the shadows of static black holes and rotating black holes are shown under $\Lambda = 10^{-4}$ or $\Lambda = 10^{-3}$. When the cosmological constant is relatively small, the shadow scale




FIG. 1. Shadows cast by black holes with a cosmological constant, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$. The (a) and (c) are static cases $a/M = 0$, corresponding to $\Lambda = 10^{-4}$ and $\Lambda = 10^{-3}$, respectively. The (b) and (d) are rotating case $\alpha = 0.02$, corresponding to $\Lambda = 10^{-4}$ and $\Lambda = 10^{-3}$, respectively.

exhibits an increase with the growth of the MOG parameter. However, as the cosmological constant increases, this trend changes. This shift can be attributed to the observer's position r_0 approaching the cosmological horizon r_c , causing the observer's position to have a more significant effect on the shadow size than the effects of the MOG modification. In Fig. 2, we present the shadow contour of a near-extreme black hole with different α and Λ , i.e., $a/M = 0.99$. The results show that, as the MOG parameter increases, the range of the photon sphere radius narrows, and its contour increasingly resembles a standard circle. When the cosmological constant is relatively small, the shadow size increases with the growth of the MOG parameter. For a fixed a and α , the apparent shape of the shadow varies with Λ . As Λ decreases, the shadow's size expands. In Fig. 3, the same phenomenon that is clearly visible in Fig. 1-(c) can also be observed in rotating black holes. Furthermore, when the MOG parameter increases significantly, the rotation-induced distortion of the photon orbits becomes less noticeable. It is worth noting that in Figs. 1-3, the purple solid line represents the shadow of the Schwarzschild black hole, serving as a scale reference.¹

The contours of static versus rotating black holes will be extremely similar when the MOG parameter is sufficiently




FIG. 2. Shadows cast by black holes with a spin $a/M = 0.99$, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$. The (a), (b), (c) and (d) corresponding to $\alpha = 0$, $\alpha = 1$, $\alpha = 2$ and $\alpha = 3$, respectively.

large [57], making it difficult to infer black hole parameters solely from the black hole shadow. However, does the frame-dragging effect caused by rotation in black hole spacetimes also affect the MOG parameter? For this, we can use the numerical backward ray-tracing method, which adopts the same orthonormal tetrad as the analytical methods in this paper and sets the number of pixels to $n = 2048$. Figs. 4 and 5 effectively demonstrate the distortion of space by MOG black holes and the gravitational lensing effects these objects produce. Specifically, the gravitational lensing observed can be utilized to examine theories of gravity [58–60]. We can observe that although increasing the MOG parameter makes the contours of rotating black holes approach those of static black holes, it does not affect the frame-dragging effect induced by rotation. Moreover, Figs. 4 and 5 serve as double checks for Figs. 1-(a) and 2-(b), respectively, with each displaying the same black hole contours.

B. Observable measurements

Up to this point, we have determined that parameters such as spin a , MOG parameter α , and cosmological constant Λ will significantly affect the apparent shape of the shadow in Kerr-MOG-dS spacetime. To characterize the shadow, it is essential to have easily measurable and reliable properties of observable measurements. We adopt two such observable measurements: the on-screen radius R_s and distortion parameter δ_s , following an approach analogous to the one taken in Ref. [61]. The parameter R_s characterizes the on-screen size of the shadow, corresponding to the radius of the reference circle

¹ The coordinates x/M and y/M in Figs. 1-3 represent the on-screen scale, which is proportional to the actual celestial coordinate scales \bar{x}/M and \bar{y}/M as described by the geometric relationships in equations (47) and (48), depending on the observer's position r_0 . For example, in the Schwarzschild case, where $r_0 = 50$, the relation $\bar{x} \approx 50.897x$ holds.




FIG. 3. Shadows cast by black holes with a cosmological constant, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$. The (a) and (c) are spin $a/M = 0.5$, corresponding to $\Lambda = 10^{-4}$ and $\Lambda = 10^{-3}$, respectively. The (b) and (d) are spin $a/M = 0.99$, corresponding to $\Lambda = 10^{-4}$ and $\Lambda = 10^{-3}$, respectively.

depicted as a blue dashed line in Fig. 6. The circle passes through three points: the top position $P_A = (x_t, y_t)$, the bottom position $P_B = (x_b, y_b)$, and the point $P_C = (x_r, 0)$, which corresponds to the unstable retrograde circular orbit as observed from the equatorial plane by an observer. Additionally, the point $P_O = (x_o, 0)$ represents the center of the reference circle, and x_o can be obtained from the coordinates of P_A, P_B , and P_C :

$$x_o = \frac{x_r^2 - x_t^2 - y_t^2}{2(x_r - x_t)}. \quad (53)$$

Afterwards, the difference between the shaded left-hand point $P_D = (x_d, 0)$ and the reference circle left-hand point $P_D = (\tilde{x}_r, 0)$ needs to be taken into account, with the size of this difference being evaluated by $D = |x_d - \tilde{x}_r|$. Further, the two observable measurements take the form

$$R_s = \frac{(x_t - x_r)^2 + y_t^2}{2|x_r - x_t|}, \quad (54)$$

$$\delta_s = \frac{D}{R_s}. \quad (55)$$

We numerically calculated these two observable measurements under the condition of fixing the observer position at $r_0 = 50$ with an inclination angle of $\vartheta_0 = \pi/2$. The results are presented in Fig. 7. Note that the static case is not included in the results because the contour of the shadow coincides with the reference circle, resulting in no distortion. Figs. 7-(b) and 7-(d) show that the cosmological constant Λ has little effect on δ_s . But δ_s decreases as the parameter α




FIG. 4. Shadows cast by static black holes ($a/M = 0$) with a cosmological constant, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$.




FIG. 5. Shadows cast by extremely rotating black holes ($a/M = 0.99$) with a cosmological constant, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$.

increases, which suggests that the Kerr-MOG-dS black hole gets less deformed compared to the Kerr-dS black hole. Figs. 7-(a) and 7-(c) clearly show the radius R_s of the shadow. From these results, it becomes evident that when the cosmological constant $\Lambda = 10^{-4}$, R_s increases as the parameter α increases with $\alpha \in [0, 3]$. However, when the cosmological constant $\Lambda = 10^{-3}$, there is a maximum value for R_s , corresponding




FIG. 6. Shadows cast by black holes, as seen by an observer at $r_0 = 50$ and $\vartheta_0 = \pi/2$. The observable measurements for the apparent shape of the black hole are the radius of the reference circle R_s and the distortion parameter $\delta_s = D/R_s$, where D is the difference between the left endpoints of the reference circle and of the shadow.




FIG. 7. observable measurements R_s (left) and δ_s (right) as functions of α are presented, respectively, for $\Lambda = 10^{-4}$ (top) and $\Lambda = 10^{-3}$ (bottom). In all cases, the observer is placed at $r_0 = 50$ and $\vartheta_0 = \pi/2$.

to $\alpha \approx 1.815$. This is consistent with the peculiar phenomena observed in Figs. 1-(c), 3-(c), and 3-(d) describing the shadow cast by MOG-dS black holes in Sec. IV.

For a approximatively estimation, utilizing the Kerr-MOG-dS metric (22), we calculate the angular radius of a black hole shadow, defined as $\theta_{\text{BH}} = \bar{R}_s \frac{\mathcal{M}}{D_O}$, with D_O representing the distance from the observer to the black hole. The radius \bar{R}_s , which is proportional to the on-screen radius R_s as described by the geometric relationships in equations (47) and

$\theta_{\text{BH}}(\mu\text{arcsec})$	$\alpha = 0$	$\alpha = 0.1$	$\alpha = 0.2$	$\alpha = 0.3$	$\alpha = 0.4$
$\Lambda = 0.0000$	24.737	26.747	28.738	30.712	32.670
$\Lambda = 0.0001$	23.644	25.562	27.460	29.342	31.208
$\Lambda = 0.0002$	22.497	24.317	26.118	27.902	29.671
$\Lambda = 0.0003$	21.288	23.004	24.701	26.382	28.047
$\Lambda = 0.0000$	24.751	26.760	28.750	30.722	32.679
$\Lambda = 0.0001$	23.658	25.573	27.470	29.351	31.215
$\Lambda = 0.0002$	22.510	24.328	26.127	27.909	29.677
$\Lambda = 0.0003$	21.298	23.013	24.709	26.388	28.053

TABLE I. The numerical estimation of the angular radius of the supermassive black hole Sgr A* in our galaxy using the metric of a Kerr-MOG-dS black hole. The top four rows correspond to $a/M = 0.1$; the bottom four rows to $a/M = 0.99$.

$\theta_{\text{BH}}(\mu\text{arcsec})$	$\alpha = 0$	$\alpha = 0.1$	$\alpha = 0.2$	$\alpha = 0.3$	$\alpha = 0.4$
$\Lambda = 0.0000$	19.859	21.473	23.081	24.656	26.229
$\Lambda = 0.0001$	18.982	20.522	22.046	23.556	25.055
$\Lambda = 0.0002$	18.062	19.522	20.968	22.401	23.821
$\Lambda = 0.0003$	17.090	18.457	19.816	21.164	22.517
$\Lambda = 0.0000$	19.871	21.484	23.064	24.664	26.236
$\Lambda = 0.0001$	18.993	20.531	22.054	23.563	25.061
$\Lambda = 0.0002$	18.071	19.531	20.975	22.406	23.826
$\Lambda = 0.0003$	17.099	18.475	19.837	21.185	22.521

TABLE II. The numerical estimation of the angular radius of the supermassive black hole M87* using the metric of a Kerr-MOG-dS black hole. The top four rows correspond to $a/M = 0.1$; the bottom four rows to $a/M = 0.99$.

(48) (see detail in footnote 1). Specifically, for a black hole possessing a mass of \mathcal{M} and situated D_O away from the observer, we express the angular radius θ_{BH} quantitatively as $\theta_{\text{BH}} = 9.87098 \times 10^{-6} \bar{R}_s \left(\frac{\mathcal{M}}{M_\odot}\right) \left(\frac{1\text{kpc}}{D_O}\right) \mu\text{as}$ [62].

In Tables I and II, we present the calculated angular radius of the black holes at the Galactic center, Sgr A*, and M87*, respectively, using the Kerr-MOG-dS metric. Here, we utilize the latest observations which indicate that the mass of the black hole Sgr A* is $\mathcal{M} = 4.0 \times 10^6 M_\odot$ with an observer distance of $D_O = 8.3\text{kpc}$ [63], and for the black hole M87*, the mass is $\mathcal{M} = 6.5 \times 10^9 M_\odot$ with an observer distance of $D_O = 16.8\text{Mpc}$ [16]. It should be noted that we have highlighted the data corresponding to the observed range of angular diameters for the Sgr A* and M87* black holes from recent observations.² By combining it with the data in Tables I and

² The angular diameters of the Sgr A* and M87* black holes are measured as $51.8 \pm 2.3 \mu\text{as}$ [63] and $42 \pm 3.0 \mu\text{as}$ [15, 64, 65], respectively.

II, one can find that there is room for the theoretical model of such a Kerr-MOG-dS black hole. Furthermore, the inclusion of the cosmological constant provides a larger parameter space for the MOG parameters.

V. CONCLUSIONS AND OUTLOOKS

In this work, we have extend the Kerr-MOG black hole solution [19] to more general cases with a cosmological constant (namely, the Kerr-MOG-dS and Kerr-MOG-AdS black hole solutions) and investigate the black hole shadows of the Kerr-MOG-dS case. Due to the presence of the positive constant Λ , the observer should be positioned between the black hole's event horizon and the cosmological horizon, rather than at infinity. The apparent shape of the shadow depends on the observer's position.

A summary of three novel consequences of the shadows of the Kerr-MOG-dS black hole obtained in this paper are listed in the following order:

- **Size:** As the cosmological constant increases, the size of the black hole shadow decreases. But the size of the shadow does not appear to be monotonic as α changes. For example, it would reach a maximum at $\alpha \approx 1.815$ with $\Lambda = 10^{-3}$.
- **Shape:** The shape of the shadow becomes more rounded as the MOG parameter α increases, causing the contours of rotating black holes to approach those of static black holes. However, this change does not impact the frame-dragging effect induced by rotation. The distortion of space is consistent with that observed in GR.
- **Observable measurements:** For observable measurements, we consider the radius R_s , which is associated with the apparent size, and the distortions δ_s , which relate to the deformation of the shadow. The results show that, with the position of the observer held constant and the cosmological constant sufficiently large, the radius R_s reaches a maximum value as α increases. This increase in α concurrently reduces the distortion δ_s .

Our results indicate that the interplay between the cosmologi-

cal constant and MOG parameter leads to intriguing phenomena in the black hole shadow contours. If the Earth were situated at the edge of the universe in this black hole spacetime, the black hole spin parameter and MOG parameter would become degenerate. Additionally, we used the latest observational data from M87* and Sgr A* to impose certain parameter constraints on Kerr-MOG-dS black holes, confirming the validity of STVG theory.

There are two promising further topics to be pursued in the future. An intriguing topic is to explore other observable measurements of the shadow of the Kerr-MOG-dS black hole. To prevent certain shadow degeneracy caused by black hole parameters, we can explore other distortion parameters, such as the oblateness parameter K_s , and the "thickness" parameter T of the shadow [66]. In order to visualize the shadow in a real astronomical setting, one should rely on the emissions originating from the accretion disk surrounding the black hole [67, 68]. And the other is to investigate the properties of the Kerr-MOG-AdS black hole, such as black hole thermodynamics [69–73], thermodynamic topological classification [74–77], null hypersurface caustics [78–80], phase transition criticality [81–83], Joule-Thomson expansion [84], etc.

ACKNOWLEDGMENTS

The authors sincerely thank Professor Songbai Chen for his contributions to this subject and Dr. Long Fen for the profound discussions. This work was supported by the National Natural Science Foundation of China under Grants No. 12122504, No. 12375046, No. 12035005, No. 12205243, No. 12375053, by the innovative research group of Hunan Province under Grant No. 2024JJ1006, by the Natural Science Foundation of Hunan Province under grant No. 2023JJ30384, by the Hunan provincial major sci-tech program under grant No.2023zk1010, by the Sichuan Science and Technology Program under Grant No. 2023NSFSC1347, and by the Doctoral Research Initiation Project of China West Normal University under Grant No. 21E028.

Appendix A: Energy-momentum tensor

In this appendix, we provide the energy-momentum tensor corresponding to the metric (17). The MOG vector potential (23) in the coordinates $(\bar{t}, \bar{r}, \bar{\vartheta}, \bar{\varphi})$ is given as follows:

$$\bar{\phi} = -\frac{\sqrt{G_N \alpha M \bar{r}}}{\Sigma} \left(\frac{\sin^2 \bar{\vartheta}}{\chi} + \cos^2 \bar{\vartheta} \right) d\bar{t} - \frac{h\sqrt{G_N \alpha M \bar{r}}}{\Delta_r \Delta_{\bar{r}}} d\bar{r} + \frac{\sqrt{G_N \alpha M \bar{r}}}{\chi \Sigma} a \sin^2 \bar{\vartheta} d\bar{\varphi}, \quad (\text{A1})$$

All non-zero components of T_{ab} are then given as follows:

$$\begin{aligned}
T_{00} &= -\frac{1}{24\pi\chi^2\Sigma^3}G_N M^2\alpha\Delta_{\bar{\vartheta}}\left[(3-\bar{r}^2\Lambda)\Sigma_2+3h\Delta_{\bar{\vartheta}}\right], & T_{01} &= -\frac{1}{8\pi\chi\Delta_{\bar{r}}\Sigma^2}G_N M^2\alpha h\Delta_{\bar{\vartheta}}, \\
T_{11} &= -\frac{1}{8\pi\Delta_{\bar{r}}^2\Sigma}G_N M^2\alpha(\Delta_r-2\Delta_{\bar{r}}), & T_{03} &= \frac{1}{8\pi\chi^2\Sigma^3}G_N M^2\alpha\Delta_{\bar{\vartheta}}(\Delta_r+\Delta_{\bar{r}})a\sin^2\bar{\vartheta}, \\
T_{22} &= -\frac{1}{8\pi\Delta_{\bar{\vartheta}}\Sigma}G_N M^2\alpha, & T_{13} &= \frac{1}{8\pi\chi\Delta_{\bar{r}}\Sigma^2}G_N M^2\alpha h a\sin^2\bar{\vartheta}, \\
T_{33} &= -\frac{\sin^2\bar{\vartheta}}{8\pi\chi^2\Sigma^3}G_N M^2\alpha\left[(\bar{r}^2+a^2)\Sigma_2+ha^2\sin^2\bar{\vartheta}\right].
\end{aligned} \tag{A2}$$

This set of equations corresponds one-to-one with the effective tensor component equations (19). It is clear that this will lead to the effective tensor $E_{ab} = 0$, ensuring that the solutions satisfy the field equations.

-
- [1] G. W. Gibbons, Vacuum Polarization and the Spontaneous Loss of Charge by Black Holes, *Commun. Math. Phys.* **44**, 245 (1975).
- [2] R. D. Blandford and R. L. Znajek, Electromagnetic extractions of energy from Kerr black holes, *Mon. Not. Roy. Astron. Soc.* **179**, 433 (1977).
- [3] V. Cardoso, C. F. B. Macedo, P. Pani, and V. Ferrari, Black holes and gravitational waves in models of minicharged dark matter, *JCAP* **05**, 054, arXiv:1604.07845 [hep-ph].
- [4] G. Bozzola and V. Paschalidis, General Relativistic Simulations of the Quasicircular Inspiral and Merger of Charged Black Holes: GW150914 and Fundamental Physics Implications, *Phys. Rev. Lett.* **126**, 041103 (2021), arXiv:2006.15764 [gr-qc].
- [5] P. K. Gupta, T. F. M. Spieksma, P. T. H. Pang, G. Koekoek, and C. V. D. Broeck, Bounding dark charges on binary black holes using gravitational waves, *Phys. Rev. D* **104**, 063041 (2021), arXiv:2107.12111 [gr-qc].
- [6] G. Carullo, D. Laghi, N. K. Johnson-McDaniel, W. Del Pozzo, O. J. C. Dias, M. Godazgar, and J. E. Santos, Constraints on Kerr-Newman black holes from merger-ringdown gravitational-wave observations, *Phys. Rev. D* **105**, 062009 (2022), arXiv:2109.13961 [gr-qc].
- [7] J. W. Moffat, Scalar-tensor-vector gravity theory, *JCAP* **03**, 004, arXiv:0506021 [gr-qc].
- [8] H. An, S. Ge, W.-Q. Guo, X. Huang, J. Liu, and Z. Lu, Direct Detection of Dark Photon Dark Matter Using Radio Telescopes, *Phys. Rev. Lett.* **130**, 181001 (2023), arXiv:2207.05767 [hep-ph].
- [9] X. Ning *et al.* (PandaX), Limits on the luminance of dark matter from xenon recoil data, *Nature* **618**, 47 (2023).
- [10] J. W. Moffat and S. Rahvar, The MOG weak field approximation and observational test of galaxy rotation curves, *Mon. Not. Roy. Astron. Soc.* **436**, 1439 (2013), arXiv:1306.6383 [astro-ph.GA].
- [11] J. W. Moffat and S. Rahvar, The MOG weak field approximation – II. Observational test of *Chandra* X-ray clusters, *Mon. Not. Roy. Astron. Soc.* **441**, 3724 (2014), arXiv:1309.5077 [astro-ph.CO].
- [12] J. W. Moffat and V. T. Toth, Rotational velocity curves in the Milky Way as a test of modified gravity, *Phys. Rev. D* **91**, 043004 (2015), arXiv:1411.6701 [astro-ph.GA].
- [13] J. W. Moffat, Structure Growth and the CMB in Modified Gravity (MOG), (2014), arXiv:1409.0853 [astro-ph.CO].
- [14] S. Rahvar, Hamiltonian formalism for dynamics of particles in MOG, *Mon. Not. Roy. Astron. Soc.* **514**, 4601 (2022), arXiv:2206.02453 [gr-qc].
- [15] K. Akiyama *et al.* (Event Horizon Telescope), First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole, *Astrophys. J. Lett.* **875**, L1 (2019), arXiv:1906.11238 [astro-ph.GA].
- [16] K. Akiyama *et al.* (Event Horizon Telescope), First M87 Event Horizon Telescope Results. VI. The Shadow and Mass of the Central Black Hole, *Astrophys. J. Lett.* **875**, L6 (2019), arXiv:1906.11243 [astro-ph.GA].
- [17] K. Akiyama *et al.* (Event Horizon Telescope), First Sagittarius A* Event Horizon Telescope Results. VI. Testing the Black Hole Metric, *Astrophys. J. Lett.* **930**, L17 (2022), arXiv:2311.09484 [astro-ph.HE].
- [18] J. W. Moffat and V. T. Toth, Masses and shadows of the black holes Sagittarius A* and M87* in modified gravity, *Phys. Rev. D* **101**, 024014 (2020), arXiv:1904.04142 [gr-qc].
- [19] J. W. Moffat, Black Holes in Modified Gravity (MOG), *Eur. Phys. J. C* **75**, 175 (2015), arXiv:1412.5424 [gr-qc].
- [20] J. W. Moffat, Regular Rotating MOG Dark Compact Object, *Eur. Phys. J. C* **81**, 119 (2021), arXiv:1806.01903 [gr-qc].
- [21] J. W. Moffat and V. T. Toth, The cosmological background and the “external field” in modified gravity (MOG), *Eur. Phys. J. C* **81**, 836 (2021), arXiv:2109.11133 [gr-qc].
- [22] W. Liu, X. Fang, J. Jing, and J. Wang, Gravito-electromagnetic perturbations of MOG black holes with a cosmological constant: quasinormal modes and ringdown waveforms, *JCAP* **11**, 057, arXiv:2306.03599 [gr-qc].
- [23] J. R. Mureika, J. W. Moffat, and M. Faizal, Black hole thermodynamics in MODified Gravity (MOG), *Phys. Lett. B* **757**, 528 (2016), arXiv:1504.08226 [gr-qc].
- [24] H.-C. Lee and Y.-J. Han, Innermost stable circular orbit of Kerr-MOG black hole, *Eur. Phys. J. C* **77**, 655 (2017), arXiv:1704.02740 [gr-qc].
- [25] J. W. Moffat, Modified Gravity Black Holes and their Observable Shadows, *Eur. Phys. J. C* **75**, 130 (2015), arXiv:1502.01677 [gr-qc].
- [26] H.-M. Wang, Y.-M. Xu, and S.-W. Wei, Shadows of Kerr-like black holes in a modified gravity theory, *JCAP* **03**, 046,

- arXiv:1810.12767 [gr-qc].
- [27] S.-W. Wei and Y.-X. Liu, Merger estimates for rotating Kerr black holes in modified gravity, *Phys. Rev. D* **98**, 024042 (2018), arXiv:1803.09530 [gr-qc].
- [28] M. Guo, N. A. Obers, and H. Yan, Observational signatures of near-extremal Kerr-like black holes in a modified gravity theory at the Event Horizon Telescope, *Phys. Rev. D* **98**, 084063 (2018), arXiv:1806.05249 [gr-qc].
- [29] P. Sheoran, A. Herrera-Aguilar, and U. Nucamendi, Mass and spin of a Kerr black hole in modified gravity and a test of the Kerr black hole hypothesis, *Phys. Rev. D* **97**, 124049 (2018), arXiv:1712.03344 [gr-qc].
- [30] Z. Zhang, S. Chen, and J. Jing, Images of Kerr-MOG black holes surrounded by geometrically thick magnetized equilibrium tori, (2024), arXiv:2404.12223 [gr-qc].
- [31] X. Qiao, M. Wang, Q. Pan, and J. Jing, Kerr-MOG black holes with stationary scalar clouds, *Eur. Phys. J. C* **80**, 509 (2020).
- [32] R. A. Konoplya and A. Zhidenko, Shadows of parametrized axially symmetric black holes allowing for separation of variables, *Phys. Rev. D* **103**, 104033 (2021), arXiv:2103.03855 [gr-qc].
- [33] Z. Younsi, A. Zhidenko, L. Rezzolla, R. Konoplya, and Y. Mizuno, New method for shadow calculations: Application to parametrized axisymmetric black holes, *Phys. Rev. D* **94**, 084025 (2016), arXiv:1607.05767 [gr-qc].
- [34] L. Manfredi, J. Mureika, and J. Moffat, Quasinormal Modes of Modified Gravity (MOG) Black Holes, *Phys. Lett. B* **779**, 492 (2018), arXiv:1711.03199 [gr-qc].
- [35] H. Jiang, X. Du, Q. Pan, S. Chen, and J. Jing, Merger estimates for a disformal Kerr black hole in quadratic degenerate higher-order scalar-tensor theories*, *Chin. Phys. C* **48**, 055103 (2024).
- [36] A. G. Riess *et al.* (Supernova Search Team), Observational evidence from supernovae for an accelerating universe and a cosmological constant, *Astron. J.* **116**, 1009 (1998), arXiv:9805201 [astro-ph].
- [37] S. Perlmutter *et al.* (Supernova Cosmology Project), Measurements of Ω and Λ from 42 High Redshift Supernovae, *Astrophys. J.* **517**, 565 (1999), arXiv:9812133 [astro-ph].
- [38] J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2**, 231 (1998), arXiv:hep-th/9711200.
- [39] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, Gauge theory correlators from noncritical string theory, *Phys. Lett. B* **428**, 105 (1998), arXiv:hep-th/9802109.
- [40] E. Witten, Anti-de Sitter space and holography, *Adv. Theor. Math. Phys.* **2**, 253 (1998), arXiv:hep-th/9802150.
- [41] S. Chen, J. Jing, W.-L. Qian, and B. Wang, Black hole images: A review, *Sci. China Phys. Mech. Astron.* **66**, 260401 (2023), arXiv:2301.00113 [astro-ph.HE].
- [42] G. W. Gibbons, H. Lu, D. N. Page, and C. N. Pope, The General Kerr-de Sitter metrics in all dimensions, *J. Geom. Phys.* **53**, 49 (2005), arXiv:0404008 [hep-th].
- [43] T. Malek and V. Pravda, Kerr-Schild spacetimes with (A)dS background, *Class. Quant. Grav.* **28**, 125011 (2011), arXiv:1009.1727 [gr-qc].
- [44] R. M. Wald, *General Relativity* (Chicago Univ. Pr., Chicago, USA, 1984).
- [45] X.-X. Zeng, K.-J. He, and G.-P. Li, Effects of dark matter on shadows and rings of Brane-World black holes illuminated by various accretions, *Sci. China Phys. Mech. Astron.* **65**, 290411 (2022), arXiv:2111.05090 [gr-qc].
- [46] W. Liu, X. Fang, J. Jing, and A. Wang, Gauge invariant perturbations of general spherically symmetric spacetimes, *Sci. China Phys. Mech. Astron.* **66**, 210411 (2023), arXiv:2201.01259 [gr-qc].
- [47] B. Carter, Global structure of the Kerr family of gravitational fields, *Phys. Rev.* **174**, 1559 (1968).
- [48] S. Chandrasekhar, The Mathematical Theory of Black Holes, *Fundam. Theor. Phys.* **9**, 5 (1984).
- [49] A. Grenzebach, V. Perlick, and C. Lämmerzahl, Photon Regions and Shadows of Kerr-Newman-NUT Black Holes with a Cosmological Constant, *Phys. Rev. D* **89**, 124004 (2014), arXiv:1403.5234 [gr-qc].
- [50] M. Wang, S. Chen, and J. Jing, Shadows of Bonnor black dihole by chaotic lensing, *Phys. Rev. D* **97**, 064029 (2018), arXiv:1710.07172 [gr-qc].
- [51] M. Wang, S. Chen, and J. Jing, Kerr black hole shadows in Melvin magnetic field with stable photon orbits, *Phys. Rev. D* **104**, 084021 (2021), arXiv:2104.12304 [gr-qc].
- [52] Z. Hu, Z. Zhong, P.-C. Li, M. Guo, and B. Chen, QED effect on a black hole shadow, *Phys. Rev. D* **103**, 044057 (2021), arXiv:2012.07022 [gr-qc].
- [53] Z. Zhong, Z. Hu, H. Yan, M. Guo, and B. Chen, QED effects on Kerr black hole shadows immersed in uniform magnetic fields, *Phys. Rev. D* **104**, 104028 (2021), arXiv:2108.06140 [gr-qc].
- [54] S. Chen and J. Jing, Kerr black hole shadows cast by extraordinary light rays with Weyl corrections, *Sci. China Phys. Mech. Astron.* **67**, 250411 (2024), arXiv:2308.16479 [gr-qc].
- [55] V. Perlick and O. Y. Tsupko, Calculating black hole shadows: Review of analytical studies, *Phys. Rept.* **947**, 1 (2022), arXiv:2105.07101 [gr-qc].
- [56] E. F. Eiroa and C. M. Sendra, Shadow cast by rotating braneworld black holes with a cosmological constant, *Eur. Phys. J. C* **78**, 91 (2018), arXiv:1711.08380 [gr-qc].
- [57] F. Camilloni, T. Harmark, M. Orselli, and M. J. Rodriguez, Blandford-Znajek jets in MOdified Gravity, *JCAP* **01**, 047, arXiv:2307.06878 [gr-qc].
- [58] T. Liu, M. Biesiada, S. Tian, and K. Liao, Robust test of general relativity at the galactic scales by combining strong lensing systems and gravitational wave standard sirens, *Phys. Rev. D* **109**, 084074 (2024), arXiv:2404.05907 [gr-qc].
- [59] A. Chowdhuri, S. Ghosh, and A. Bhattacharyya, A review on analytical studies in Gravitational Lensing, *Front. Phys.* **11**, 1113909 (2023), arXiv:2303.02069 [gr-qc].
- [60] S. Ghosh and A. Bhattacharyya, Analytical study of gravitational lensing in Kerr-Newman black-bounce spacetime, *JCAP* **11**, 006, arXiv:2206.09954 [gr-qc].
- [61] K. Hioki and K.-i. Maeda, Measurement of the Kerr Spin Parameter by Observation of a Compact Object's Shadow, *Phys. Rev. D* **80**, 024042 (2009), arXiv:0904.3575 [astro-ph.HE].
- [62] L. Amarilla and E. F. Eiroa, Shadow of a rotating braneworld black hole, *Phys. Rev. D* **85**, 064019 (2012), arXiv:1112.6349 [gr-qc].
- [63] K. Akiyama *et al.* (Event Horizon Telescope), First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way, *Astrophys. J. Lett.* **930**, L12 (2022), arXiv:2311.08680 [astro-ph.HE].
- [64] F. Long, J. Wang, S. Chen, and J. Jing, Shadow of a rotating squashed Kaluza-Klein black hole, *JHEP* **10**, 269, arXiv:1906.04456 [gr-qc].
- [65] K. Pal, K. Pal, R. Shaikh, and T. Sarkar, A rotating modified JNW spacetime as a Kerr black hole mimicker, *JCAP* **11**, 060, arXiv:2305.07518 [gr-qc].
- [66] M. Wang, S. Chen, and J. Jing, Chaotic shadow of a non-Kerr rotating compact object with quadrupole mass moment, *Phys. Rev. D* **98**, 104040 (2018), arXiv:1801.02118 [gr-qc].
- [67] Z. Zhang, S. Chen, X. Qin, and J. Jing, Polarized image of a Schwarzschild black hole with a thin accretion disk as pho-

- ton couples to Weyl tensor, *Eur. Phys. J. C* **81**, 991 (2021), [arXiv:2106.07981 \[gr-qc\]](#).
- [68] Z. Zhang, S. Chen, and J. Jing, Image of Bonnor black dihole with a thin accretion disk and its polarization information, *Eur. Phys. J. C* **82**, 835 (2022), [arXiv:2205.13696 \[gr-qc\]](#).
- [69] D. Christodoulou, Reversible and irreversible transformations in black hole physics, *Phys. Rev. Lett.* **25**, 1596 (1970).
- [70] J. M. Bardeen, B. Carter, and S. W. Hawking, The Four laws of black hole mechanics, *Commun. Math. Phys.* **31**, 161 (1973).
- [71] S.-Q. Wu and D. Wu, Thermodynamical hairs of the four-dimensional Taub-Newman-Unti-Tamburino spacetimes, *Phys. Rev. D* **100**, 101501 (2019), [arXiv:1909.07776 \[hep-th\]](#).
- [72] W. Cong, D. Kubiznak, and R. B. Mann, Thermodynamics of AdS Black Holes: Critical Behavior of the Central Charge, *Phys. Rev. Lett.* **127**, 091301 (2021), [arXiv:2105.02223 \[hep-th\]](#).
- [73] M. B. Ahmed, W. Cong, D. Kubizňák, R. B. Mann, and M. R. Visser, Holographic Dual of Extended Black Hole Thermodynamics, *Phys. Rev. Lett.* **130**, 181401 (2023), [arXiv:2302.08163 \[hep-th\]](#).
- [74] S.-W. Wei, Y.-X. Liu, and R. B. Mann, Black Hole Solutions as Topological Thermodynamic Defects, *Phys. Rev. Lett.* **129**, 191101 (2022), [arXiv:2208.01932 \[gr-qc\]](#).
- [75] D. Wu, Topological classes of rotating black holes, *Phys. Rev. D* **107**, 024024 (2023), [arXiv:2211.15151 \[gr-qc\]](#).
- [76] D. Wu and S.-Q. Wu, Topological classes of thermodynamics of rotating AdS black holes, *Phys. Rev. D* **107**, 084002 (2023), [arXiv:2301.03002 \[hep-th\]](#).
- [77] D. Wu, Classifying topology of consistent thermodynamics of the four-dimensional neutral Lorentzian NUT-charged spacetimes, *Eur. Phys. J. C* **83**, 365 (2023), [arXiv:2302.01100 \[gr-qc\]](#).
- [78] A. Al Balushi and R. B. Mann, Null hypersurfaces in Kerr-(A)dS spacetimes, *Class. Quant. Grav.* **36**, 245017 (2019), [arXiv:1909.06419 \[gr-qc\]](#).
- [79] M. T. N. Imseis, A. Al Balushi, and R. B. Mann, Null hypersurfaces in Kerr-Newman-AdS black hole and super-entropic black hole spacetimes, *Class. Quant. Grav.* **38**, 045018 (2021), [arXiv:2007.04354 \[gr-qc\]](#).
- [80] D. Wu, P. Wu, and H. Yu, Shadowless rapidly rotating yet not ultraspinning Kerr-AdS4 and Kerr-Newman-AdS4 black holes, *Phys. Rev. D* **104**, L121501 (2021), [arXiv:2107.13255 \[gr-qc\]](#).
- [81] D. Kubiznak and R. B. Mann, P-V criticality of charged AdS black holes, *JHEP* **07**, 033, [arXiv:1205.0559 \[hep-th\]](#).
- [82] M. B. Ahmed, W. Cong, D. Kubiznak, R. B. Mann, and M. R. Visser, Holographic CFT phase transitions and criticality for rotating AdS black holes, *JHEP* **08**, 142, [arXiv:2305.03161 \[hep-th\]](#).
- [83] D. Wu, S.-Y. Gu, X.-D. Zhu, Q.-Q. Jiang, and S.-Z. Yang, Topological classes of thermodynamics of the static multi-charge AdS black holes in gauged supergravities, (2024), [arXiv:2402.00106 \[hep-th\]](#).
- [84] O. Ökcü and E. Aydiner, Joule-Thomson expansion of Kerr-AdS black holes, *Eur. Phys. J. C* **78**, 123 (2018), [arXiv:1709.06426 \[gr-qc\]](#).