

SMOOTH ORBIT EQUIVALENCE RIGIDITY FOR DISSIPATIVE GEODESIC FLOWS

JAVIER ECHEVARRÍA CUESTA

ABSTRACT. Let M be a smooth closed orientable surface. A Gaussian thermostat on M can be seen as the geodesic flow of a certain metric connection with torsion. These flows may not preserve any smooth volume form. We prove that if two Gaussian thermostats on M with negative thermostat curvature are related by a smooth orbit equivalence isotopic to the identity, then the two background metrics are conformally equivalent via a smooth diffeomorphism of M isotopic to the identity. We also give a relationship between the thermostat forms themselves. Finally, we prove the same result for Anosov magnetic flows.

1. INTRODUCTION

Let (M, g) be a smooth closed oriented Riemannian surface, and let $\lambda \in \mathcal{C}^\infty(SM, \mathbb{R})$ be a smooth function on the unit tangent bundle $\pi : SM \rightarrow M$. We concern ourselves with the dynamical system governed by the equation

$$\nabla_{\dot{\gamma}} \dot{\gamma} = \lambda(\gamma, \dot{\gamma}) J \dot{\gamma},$$

where $J : TM \rightarrow TM$ is the complex structure on M induced by the orientation.

This equation defines a flow $\varphi_t := (\gamma(t), \dot{\gamma}(t))$ on SM which reduces to the geodesic flow when $\lambda = 0$. The flow models the motion of a particle under the influence of a force orthogonal to the velocity and with magnitude λ . Its generating vector field is $F = X + \lambda V$, where X is the geodesic vector field on SM , and V is the vertical vector field. The system (M, g, λ) is called a (*generalized*) *thermostat*.

If λ does not depend on the velocity, i.e., if it corresponds to a function on M , then φ_t is the *magnetic flow* associated with the magnetic field $\lambda \mu_a$, where μ_a is the area form of (M, g) . When λ depends linearly on the velocity, i.e., when it corresponds to a 1-form on M , we instead obtain a *Gaussian thermostat*, which is reversible in the sense that the flip $(x, v) \mapsto (x, -v)$ on SM conjugates φ_t with φ_{-t} (just as in the case of geodesic flows). The resulting flows are interesting from a dynamical point of view because, contrary to geodesic or magnetic flows, they may not preserve any smooth volume form (see [DP07]). Gaussian thermostats also appear in geometry as the geodesic flows of certain metric connections with torsion (see [PW08]). We thus think of them as dissipative geodesic flows.

We are interested in rigidity results for generalized thermostats satisfying the Anosov property. By [Ghy84, Theorem A], these flows are topologically orbit equivalent to the geodesic flow of any metric of constant negative curvature on M via a Hölder homeomorphism which is in fact isotopic to the identity. In particular, this tells us that the flows of generalized thermostats are transitive and topologically mixing, so the idea is that the richness of the chaotic orbits should allow one to recover information about the system.

The set up is as follows. Given two generalized thermostats (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ on the same surface M , we assume there is a smooth orbit equivalence $\phi : \tilde{S}M \rightarrow SM$ which is isotopic to the identity. Here $\tilde{S}M$ is the unit tangent bundle with respect to the metric \tilde{g} on M , and orbit equivalence means that oriented orbits are mapped to oriented orbits, i.e., there exists $c \in \mathcal{C}^\infty(SM, \mathbb{R}_{>0})$ such that $\phi_* \tilde{F} = cF$. In particular, ϕ is a conjugacy if c is identically 1. There is a natural identification of SM with $\tilde{S}M$ by scaling the fibers via the map

$$s : SM \rightarrow \tilde{S}M, \quad (x, v) \mapsto (x, v/\|v\|_{\tilde{g}}). \quad (1.1)$$

By saying that ϕ is isotopic to the identity we mean that $s \circ \phi : \tilde{S}M \rightarrow \tilde{S}M$ is isotopic to the identity in the usual sense.

Question: If both thermostat flows are Anosov, what is the relationship, if any, between (g, λ) and $(\tilde{g}, \tilde{\lambda})$?

The work in [GLP24] gives an answer in the case where $\lambda = \tilde{\lambda} = 0$ and ϕ is a conjugacy. The metrics g and \tilde{g} must be isometric via an isometry isotopic to the identity. Instead of starting with a smooth conjugacy isotopic to the identity, they start with the equivalent assumption that both metrics have the same marked length spectrum. The two assumptions are also equivalent for magnetic flows, but having the same marked length spectra only guarantees a Hölder continuous conjugacy in general.

Still with ϕ as a conjugacy, the paper [Gro99] deals with the mixed case where (M, g, λ) is a magnetic system and $(M, \tilde{g}, 0)$ is geodesic, but at the cost of additional assumptions: \tilde{g} has negative Gaussian curvature, M has the same area with respect to g and \tilde{g} , and neither λ nor its first derivative are too big. The conclusion is then that g and \tilde{g} are isometric via an isometry isotopic to the identity and that $\lambda = 0$.

More recently, progress has been made with [Reb23] to understand a deformative version of our question in the purely magnetic case, framed through the lens of marked length spectrum rigidity.

1.1. Main results. Beyond its physical motivation, the magnetic case represents the first step towards the broader goal of understanding generalized thermostats: it corresponds to the case where $\lambda = \lambda_0$ has Fourier degree 0 (see §2.1.4). Our first main result is the following:

Theorem 1.1. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ be two Anosov magnetic systems on a smooth closed orientable surface M . If there is a smooth orbit equivalence isotopic to the identity between them, then there exists a smooth diffeomorphism $\psi : M \rightarrow M$, isotopic to the identity, such that $\psi^*\tilde{g} = e^{2f}g$ for some $f \in C^\infty(M, \mathbb{R})$. Moreover, if the orbit equivalence is a conjugacy and $f = 0$, then $\lambda = 0$ if and only if $\tilde{\lambda} = 0$.*

We note that finding a relationship between λ and $\tilde{\lambda}$ in the general magnetic case remains an open question. A key similarity between geodesic and magnetic flows is that they preserve the Liouville measure on SM . As we will explain, this allows most of the key arguments from the paper [GLP24] to also go through in the magnetic case.

For this reason, the main emphasis of this paper is instead on Gaussian thermostats. These correspond to the case where $\lambda = \lambda_{-1} + \lambda_1$ or, equivalently,

$$\lambda = \pi_1^*\theta \tag{1.2}$$

for some 1-form θ on M , where $(\pi_1^*\theta)(x, v) := \theta_x(v)$ denotes the restriction to SM of smooth differential forms (so that we may see them as functions on SM). We will denote a Gaussian thermostat (M, g, λ) by (M, g, θ) to highlight its particular form.

One can also study Gaussian thermostats using an *external vector field* E . This is the vector field on M characterized by $\theta_x(v) = g(E(x), Jv)$, that is, the vector field dual to $\star\theta$, where \star is the Hodge star operator of the metric g (given by oriented rotation by $\pi/2$ in the case of 1-forms).

As we allow λ to have Fourier degree 1, we introduce the possibility of new dynamical features absent from the geodesic and magnetic cases. For instance, by [DP07, Theorem A], a Gaussian thermostat preserves an absolutely continuous invariant measure on SM if and only if $\star\theta$ is exact. This means that the Liouville measure may no longer be preserved, and it allows for fractal SRB measures.

The *thermostat curvature* of (M, g, θ) is the quantity

$$\mathbb{K} = \pi^*(K_g + \operatorname{div}_{\mu_a} E), \tag{1.3}$$

where K_g is the Gaussian curvature of (M, g) . If $\mathbb{K} < 0$, then the flow is Anosov by [Woj00, Theorem 5.2], in analogy with the geodesic case. Note that equation (1.3) is a particular case of the more general definition

$$\mathbb{K} := \pi^*K_g - H(\lambda) + \lambda^2 + F(V(\lambda)) \tag{1.4}$$

used for any generalized thermostat (M, g, λ) .

This leads us to our next main result.

Theorem 1.2. *Let (M, g, θ) and $(M, \tilde{g}, \tilde{\theta})$ be two Gaussian thermostats with $\mathbb{K}, \tilde{\mathbb{K}} < 0$ on a smooth closed orientable surface M . If there is a smooth orbit equivalence isotopic to the identity between them, then there exists a smooth diffeomorphism $\psi : M \rightarrow M$,*

isotopic to the identity, such that $\psi^*\tilde{g} = e^{2f}g$ for some $f \in C^\infty(M, \mathbb{R})$. Moreover, if either $\star\theta$ or $\tilde{\star}\tilde{\theta}$ is closed, then $\star(\psi^*\tilde{\theta} - \theta)$ is exact.

As shown in Lemma 4.6, the scaling map defined in (1.1) yields a smooth orbit equivalence isotopic to the identity between the Gaussian thermostats (M, g, θ) and $(M, e^{2f}g, \theta + \star df)$, with a time-change by e^f . This implies that the conformal factor in our main result is optimal and that it is necessary to leave room for an exact difference when relating the 1-forms. However, it is unclear at this stage whether the closedness condition is really necessary to establish this last relationship.

Ideally, one would like to extend this result to the general Anosov case. The only place where we use the negative thermostat curvature is in showing that the Gaussian thermostats satisfy the attenuated tensor tomography problem of order 1 (see §2.3.2). We do not have this issue in the purely magnetic case, which is why we were able to simply assume the more general Anosov property in Theorem 1.1. Removing the negative thermostat curvature assumption should also allow one to mix the magnetic case with Gaussian thermostats, i.e., to take $\lambda = \lambda_{-1} + \lambda_0 + \lambda_1$.

As pointed out above, there are still open questions regarding the rigidity of λ for λ of Fourier degree 1. It is also unclear at this stage how much information is gained from having a genuine conjugacy versus an orbit equivalence, and whether the conjugating diffeomorphism ϕ itself must have some particular form as in the purely geodesic case (see [GLP24, Corollary 1.2]).

After this work, a natural question is whether anything can be said for λ of Fourier degree ≥ 2 . As we show with the no-go Lemma 2.16, the current argument does not work for these thermostats. However, there are interesting examples of such systems. For instance, when λ is the real part of a holomorphic differential of degree ≥ 2 , the corresponding thermostat admits an interpretation as coupled vortex equations (see [MP19]). It was also shown in [MP20] that the geodesic flow of an affine connection on M is, up to a time-change, the flow of a generalized thermostat with λ of the form $\lambda = \lambda_{-3} + \lambda_{-1} + \lambda_1 + \lambda_3$. Just as we have shown that a non-trivial Anosov magnetic system (M, g, λ) cannot be smoothly conjugate to an Anosov geodesic flow $(M, g, 0)$ by a conjugacy isotopic to the identity, it would be interesting to further categorize generalized thermostats.

Finally, we note that Theorem A.5, which applies to generalized thermostats, was placed in the appendix to improve the overall exposition of the paper, but it represents a new result related to the injectivity of the generalized thermostat X-ray transform.

1.2. Strategy. Our main inspiration is the approach in [GLP24]. Indeed, we show that a smooth orbit equivalence isotopic to the identity determines the complex structure of the metric g up to biholomorphisms isotopic to the identity (Proposition 4.2).

This allows us to conclude that the two metrics g and \tilde{g} must be conformally equivalent via a smooth diffeomorphism of M isotopic to the identity.

To show that the orbit equivalence determines the complex structure, we rely on Torelli's theorem (Theorem 2.8), which tells us that it is enough to show that the period matrix of the underlying Riemann surface is preserved. To be able to conclude that the resulting diffeomorphism is isotopic to the identity, we use the fact that the argument can be repeated on any finite cover.

The period matrix is defined in terms of holomorphic 1-forms on M . We show with Theorem 2.15 that these can always be associated to the first Fourier modes of certain distributions $\mathcal{D}_{\text{tr},+}(SM)$ on SM satisfying a transport equation and with non-negative Fourier modes (see §2.2.2). Asking for these distributions to only have non-negative Fourier modes is a critical requirement for the rest of the argument, but it does not carry over to the case of generalized thermostats when λ has Fourier degree ≥ 2 .

We then establish in Lemma 3.6 a pairing formula showing that the integral of any holomorphic 1-form over a thermostat geodesic γ on M (i.e., the periods of the period matrix) is the same as the integral over $\pi^{-1}(\gamma) \subseteq SM$ of an associated 2-current invariant by F and living in a certain subspace $\mathcal{F}(SM)$ (see §2.1.3). This pairing formula then tells us that the smooth orbit equivalence preserves the period matrix.

At a high level, there are two main challenges and departures from [GLP24]: the first is in handling a general orbit equivalence instead of a conjugacy, and the second is in dealing with the fact that Gaussian thermostats may not be volume-preserving.

The presence of a non-zero divergence with respect to the Liouville form manifests itself in a few ways. First, instead of flow-invariant distributions, the right object of study becomes solutions to the dual transport equation. This subspace is no longer preserved by the pullback of the orbit equivalence, so we have to introduce the space $\mathcal{F}(SM)$ of 2-currents mentioned above and establish a one-to-one relationship with the distributions solving the transport equation (Lemma 2.3). We then have to check that the wavefront set analysis is unaffected by factoring the correspondence through this space (Lemma 2.4) and that $\mathcal{D}_{\text{tr},+}(SM)$ is mapped to $\mathcal{D}_{\text{tr},+}(\tilde{S}M)$ (Proposition 3.3).

Another complication due to the dissipation is in showing that any holomorphic 1-form can be seen as the first Fourier mode of an element in $\mathcal{D}_{\text{tr},+}(SM)$, as previously mentioned. The heavy lifting to address this issue is done in Appendix A. Furthermore, again due to the divergence, we have to explain why Gaussian thermostats with negative thermostat curvature satisfy the attenuated tensor tomography problem of order 1 (Theorem 2.12).

For the pairing formula previously described, we have replaced the role of the Liouville form with that of a certain form defined in (2.2). Finally, to relate λ with $\tilde{\lambda}$, we rely on new arguments which at their core involve the smooth Livšic theorem.

1.3. Organization of the paper. In Section 2, we introduce the background tools necessary for the rest of the paper. Specifically, §2.1 provides a short introduction to the geometry and Fourier analysis of the unit tangent bundle. It also introduces the new objects needed to deal with the divergence of the generalized thermostats. In §2.2, we review the complex geometry and harmonic analysis on a surface, while §2.3 delves into hyperbolic dynamics and tensor tomography.

In Section 3, we explain how a smooth orbit equivalence acts on holomorphic differentials, and we establish the pairing formula needed to show that period matrices are preserved. We then present the proofs of our main results in Section 4.

Appendix A delves into the question of finding distributional solutions, with prescribed Fourier modes, of the relevant transport equation for a generalized thermostat.

Acknowledgements. I would like to thank my advisor, Gabriel Paternain, for suggesting this project and guiding me while working on it.

2. PRELIMINARIES

In what follows, (M, g) is a smooth closed oriented Riemannian surface, and we take an arbitrary $\lambda \in \mathcal{C}^\infty(SM, \mathbb{R})$. Whenever we use additional assumptions, it will be clearly stated in the result statements. We will sometimes need a second generalized thermostat $(M, \tilde{g}, \tilde{\lambda})$. All the objects depending on the metric will then be labeled accordingly. Finally, we denote by $\phi : \tilde{S}M \rightarrow SM$ a smooth orbit equivalence between the thermostats (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$. Once again, we will specify when we assume it to be isotopic to the identity.

2.1. Unit tangent bundle of the surface. We review some basics of the unit tangent bundle $\pi : SM \rightarrow M$ defined by

$$SM := \{(x, v) \in TM \mid \|v\|_g = 1\}.$$

2.1.1. Geometry of SM . As previously, let X be the geodesic vector field on SM , and let V be the vertical vector field generating the circle action on the fibers. We define $H := [V, X]$. The vector fields $\{X, H, V\}$ form an orthonormal basis on SM for the Sasaki metric (the natural lift of g to SM). We set $\mathbb{H} := \mathbb{R}H$ and $\mathbb{V} := \mathbb{R}V$. We also note that the geodesic vector field splits into $X = \eta_+ + \eta_-$ where η_\pm are the *raising* and *lowering* Guillemin-Kazhdan operators given by

$$\eta_\pm := \frac{1}{2}(X \mp iH). \tag{2.1}$$

The *Liouville* 1-form $\alpha \in \mathcal{C}^\infty(SM, T^*(SM))$ is defined by $\alpha(X) = 1$ and $\alpha(H) = \alpha(V) = 0$. It is invariant by the geodesic flow in the sense that $\mathcal{L}_X\alpha = 0$. The 2-form

$d\alpha$ is non-degenerate on the *contact plane* $\mathbb{H} \oplus \mathbb{V}$, and it satisfies $\iota_X d\alpha = 0$. Hence

$$\mu := -\alpha \wedge d\alpha$$

is a volume form invariant by the geodesic flow. We call it the *Liouville* volume form. It corresponds to the Riemannian volume form induced by the Sasaki metric on SM . From now on, the L^2 space on SM is defined as $L^2(SM) := L^2(SM, \mu)$.

We also define the 1-forms β, ψ on SM by $\beta(H) = 1 = \psi(V)$ and $\beta(X) = \beta(V) = 0 = \psi(X) = \psi(H)$. It is easy to check that $d\alpha = \psi \wedge \beta$ so that $\mu = \alpha \wedge \beta \wedge \psi$. We set $(\mathbb{R}X)^* := \mathbb{R}\alpha$, $(\mathbb{R}F)^* := \mathbb{R}\lambda$, $\mathbb{H}^* := \mathbb{R}\beta$, and $\mathbb{V}^* := \mathbb{R}\psi$. We refer to [PSU23, Chapter 3] for further details on the geometric structure on SM .

2.1.2. Appearance of the divergence. The key difference between generalized thermostats and geodesic or magnetic flows is that the generating vector field F might not preserve the Liouville volume form μ . Recall that the *divergence* of the vector field F with respect to the volume form μ is the function $\operatorname{div}_\mu F \in \mathcal{C}^\infty(SM, \mathbb{R})$ uniquely defined by

$$\mathcal{L}_F \mu = (\operatorname{div}_\mu F) \mu.$$

The following result is proved in [DP07, Lemma 3.2].

Lemma 2.1. *Let (M, g, λ) be a generalized thermostat. Then, we have:*

$$\mathcal{L}_F \mu = V(\lambda) \mu, \quad \mathcal{L}_H \mu = 0, \quad \mathcal{L}_V \mu = 0.$$

In the geodesic and magnetic cases, we have $V(\lambda) = 0$, so the Liouville volume form is preserved. Another way in which the divergence manifests itself is when calculating the adjoint operators with respect to the L^2 inner product on SM :

$$F^* = -(F + V(\lambda)), \quad H^* = -H, \quad V^* = -V.$$

This is relevant when extending differential operators to act on the space of distributions. Recall that any differential operator P with smooth real-valued coefficients acts on a distribution $u \in \mathcal{D}'(SM)$ by duality, that is $\langle Pu, \varphi \rangle_{\mathcal{D}'(SM)} := \langle u, P^* \varphi \rangle_{\mathcal{D}'(SM)}$ for any $\varphi \in \mathcal{C}^\infty(SM)$. The subspace of distributional solutions to the transport equation

$$\mathcal{D}'_{\operatorname{tr}}(SM) := \{u \in \mathcal{D}'(SM) \mid (F + V(\lambda))u = 0\}$$

thus corresponds to the distributions $u \in \mathcal{D}'(SM)$ such that $\langle u, F\varphi \rangle_{\mathcal{D}'(SM)} = 0$ for all $\varphi \in \mathcal{C}^\infty(SM)$. If $V(\lambda) = 0$, these are simply the distributions invariant by the flow.

2.1.3. Divergence and smooth orbit equivalences. It will prove important to understand how the divergence of a system interacts with smooth orbit equivalences.

The next result, which we have stated in a broader setting than the one we are studying in this paper to highlight its generality, relates the divergences of two flows associated by a smooth orbit equivalence.

Lemma 2.2. *Let N and \tilde{N} be two orientable manifolds endowed with nowhere-vanishing volume forms μ and $\tilde{\mu}$, and smooth vector fields Y and \tilde{Y} . Suppose $\phi : \tilde{N} \rightarrow N$ is a smooth orbit equivalence between the flows generated by \tilde{Y} and Y . If we write $\phi_*\tilde{Y} = cY$ with $c \in \mathcal{C}^\infty(N, \mathbb{R}_{>0})$ and $\phi^*\mu = (\det \phi)\tilde{\mu}$ with $\det \phi \in \mathcal{C}^\infty(\tilde{N}, \mathbb{R})$, then*

$$(\det \phi)\phi^*(\operatorname{div}_\mu Y) = \tilde{Y} \left(\frac{\det \phi}{\phi^*c} \right) + \frac{\det \phi}{\phi^*c} \operatorname{div}_{\tilde{\mu}} \tilde{Y}.$$

In particular, if ϕ preserves the orientation, i.e., $\det \phi > 0$, then

$$(\phi^*c)\phi^*(\operatorname{div}_\mu Y) = \tilde{Y} \left(\ln \left(\frac{\det \phi}{\phi^*c} \right) \right) + \operatorname{div}_{\tilde{\mu}} \tilde{Y}.$$

Proof. We compute

$$\begin{aligned} \phi^*(\mathcal{L}_Y \mu) &= \phi^*(d(\iota_Y \mu)) \\ &= d(\iota_{\phi_*^{-1}Y} \phi^*(\mu)) \\ &= d \left(\frac{\det \phi}{\phi^*c} \iota_{\tilde{Y}} \tilde{\mu} \right) \\ &= d \left(\frac{\det \phi}{\phi^*c} \right) \wedge \iota_{\tilde{Y}} \tilde{\mu} + \frac{\det \phi}{\phi^*c} d(\iota_{\tilde{Y}} \tilde{\mu}) \\ &= \left(\tilde{Y} \left(\frac{\det \phi}{\phi^*c} \right) + \frac{\det \phi}{\phi^*c} \operatorname{div}_{\tilde{\mu}} \tilde{Y} \right) \tilde{\mu}. \end{aligned}$$

On the other hand, we also have

$$\begin{aligned} \phi^*(\mathcal{L}_Y \mu) &= \phi^*(\operatorname{div}_\mu Y \mu) \\ &= \phi^*(\operatorname{div}_\mu Y) \phi^*\mu \\ &= \phi^*(\operatorname{div}_\mu Y) (\det \phi) \tilde{\mu}, \end{aligned}$$

so putting these together yields the desired result since $\tilde{\mu}$ is nowhere-vanishing. \square

In the geodesic and magnetic cases, the pullback ϕ^* of the smooth orbit equivalence $\phi : \tilde{SM} \rightarrow SM$ sends the space $\mathcal{D}'_{\operatorname{tr}}(SM)$ to $\mathcal{D}'_{\operatorname{tr}}(\tilde{SM})$. More generally, however, the divergence term $V(\lambda)$ appearing in the transport equation breaks this down.

Instead, a more useful perspective is to look at the following subspace of 2-currents (or distributional 2-forms) on SM invariant by F :

$$\mathcal{F}(SM) := \{ \sigma \in \mathcal{D}'(SM, \Lambda^2 T^*(SM)) \mid \iota_F \sigma = d\sigma = 0 \}.$$

This set only depends on the foliation corresponding to F , i.e., it is invariant under time-changes, so we get a \mathbb{C} -linear isomorphism $\phi^* : \mathcal{F}(SM) \rightarrow \mathcal{F}(\tilde{SM})$. The 2-form

$$\omega := \iota_F \mu \tag{2.2}$$

then allows us to establish a relationship with solutions to the transport equation.

Lemma 2.3. *The map $L : \mathcal{D}'_{\text{tr}}(SM) \rightarrow \mathcal{F}(SM)$ given by $u \mapsto u\omega$ is a \mathbb{C} -linear isomorphism.*

Proof. Using Cartan's magic formula and Lemma 2.1, note that

$$\begin{aligned} d(u\omega) &= du \wedge \omega + u d\omega \\ &= du \wedge \omega + u \mathcal{L}_F \mu \\ &= du \wedge \omega + u V(\lambda) \mu \\ &= (Fu + V(\lambda)u) \mu. \end{aligned}$$

Therefore, $u\omega$ is closed if and only if $(F + V(\lambda))u = 0$. Since F never vanishes, any 2-current σ on SM satisfying $\iota_F \sigma = 0$ must be of the form $\sigma = u\omega$ for some $u \in \mathcal{D}'(SM)$. \square

Thanks to this identification, we can now define a map $\Phi : \mathcal{D}'_{\text{tr}}(SM) \rightarrow \mathcal{D}'_{\text{tr}}(\tilde{S}M)$ associated to the smooth orbit equivalence $\phi : \tilde{S}M \rightarrow SM$ via the following diagram:

$$\begin{array}{ccc} \mathcal{F}(SM) & \xrightarrow{\phi^*} & \mathcal{F}(\tilde{S}M) \\ L \uparrow & & \downarrow \tilde{L}^{-1} \\ \mathcal{D}'_{\text{tr}}(SM) & \xrightarrow{\Phi} & \mathcal{D}'_{\text{tr}}(\tilde{S}M) \end{array} \quad (2.3)$$

This point of view does not affect the wavefront set analysis.

Lemma 2.4. *If ϕ preserves the orientation, then, for all $u \in \mathcal{D}'_{\text{tr}}(SM)$, we have*

$$\text{WF}(\Phi u) = \text{WF}(\phi^* u).$$

Proof. Let $q \in C^\infty(\tilde{S}M, \mathbb{R}_{>0})$ be the function such that $\phi^* \omega = q \tilde{\omega}$. Then, we get

$$\begin{aligned} \Phi u &= \tilde{L}^{-1} \phi^* L u \\ &= \tilde{L}^{-1} \phi^*(u\omega) \\ &= \tilde{L}^{-1}(q \phi^* u \tilde{\omega}) \\ &= q \phi^* u. \end{aligned}$$

Since multiplication by the nowhere-vanishing function q is elliptic, we get the result by elliptic regularity (see [Hö03, Theorem 8.3.2]). \square

By the properties of wavefront sets under pullback operators (see [Hö03, Theorem 8.2.4] for instance), we thus obtain

$$\text{WF}(\Phi u) = d\phi^T(\text{WF}(u))$$

for all $u \in \mathcal{D}'_{\text{tr}}(SM)$, where $d\phi^T : T^*(SM) \rightarrow T^*(\tilde{S}M)$ is the symplectic lift of ϕ^{-1} to the cotangent bundles given by

$$d\phi^T(y, \eta) := \left(\phi^{-1}(y), d\phi^T_{\phi^{-1}(y)}\eta \right), \quad (y, \eta) \in T^*(SM).$$

2.1.4. *Fourier decomposition.* The space $\mathcal{C}^\infty(SM)$ breaks up as

$$\mathcal{C}^\infty(SM) = \bigoplus_{k \in \mathbb{Z}} \Omega_k, \quad \Omega_k := \{u \in \mathcal{C}^\infty(SM) \mid Vu = iku\}.$$

This decomposition is orthogonal with respect to the L^2 inner product on SM and with \mathcal{C}^∞ being replaced by L^2 . For any $u \in \mathcal{C}^\infty(SM)$, we shall write $u = \sum_{k \in \mathbb{Z}} u_k$, where each $u_k \in \Omega_k$ is given by

$$u_k(x, v) := \frac{1}{2\pi} \int_0^{2\pi} u(\rho_t(x, v)) e^{-ikt} dt, \quad (2.4)$$

with ρ_t being the flow generated by V . More generally, any distribution $u \in \mathcal{D}'(SM)$ can be decomposed as $u = \sum_{k \in \mathbb{Z}} u_k$, where each $u_k \in \mathcal{D}'(SM)$ is defined by

$$\langle u_k, \varphi \rangle := \langle u, \varphi_{-k} \rangle_{\mathcal{D}'(SM)}, \quad \varphi \in \mathcal{C}^\infty(SM),$$

and satisfies $Vu_k = iku_k$.

If a distribution on SM only has finitely many non-trivial Fourier modes, we say that it has *finite Fourier degree*. The smallest $m \in \mathbb{N}$ such that $u_k = 0$ for all $|k| > m$ is then called the *Fourier degree* of u .

It also worth noting that the ladder operators η_\pm in (2.1) take their name from the fact that they act as raising/lowering operators on the Fourier decomposition, that is,

$$\eta_\pm : \Omega_k \rightarrow \Omega_{k \pm 1}$$

for all $k \in \mathbb{Z}$. In particular, we have $(Xu)_k = \eta_- u_{k+1} + \eta_+ u_{k-1}$ for any $u \in \mathcal{D}'(SM)$.

2.2. **Complex geometry.** The conformal class of the Riemannian metric g and the orientation of M induce a complex structure $J : TM \rightarrow TM$ on M , making it into a Riemann surface which we denote by (M, J) .

2.2.1. *Complex structures.* The *Teichmüller space* of M , denoted by $\mathcal{T}(M)$, is the space of complex structures on M modulo the equivalence relation that $J \sim \tilde{J}$ if and only if there exists a diffeomorphism $\psi : M \rightarrow M$, isotopic to the identity, such that $\psi^* \tilde{J} = J$. We will denote such an equivalence class of complex structures by $[J]$.

The *mapping class group* $\text{MCG}(M)$ is defined as the quotient of orientation preserving diffeomorphisms on M modulo isotopy. They act on $\mathcal{T}(M)$ by pullback, and the quotient space $\mathcal{M}(M) := \mathcal{T}(M)/\text{MCG}(M)$ is the *moduli space of complex structures* on M . See [FM11] for a thorough introduction.

Each complex structure J determines a canonical line bundle $\kappa := T_{1,0}^*M$ on M . We will denote by $H_J^0(M, \kappa^{\otimes k})$ the space of J -holomorphic sections of the k -th tensor power of the canonical line bundle κ . Locally, its elements have the form $w(z)dz^k$ for $k \geq 0$ and $w(z)d\bar{z}^{-k}$ for $k < 0$.

2.2.2. Fiberwise holomorphic distributions. Each subspace Ω_k of Fourier modes can be identified with $\mathcal{C}^\infty(M, \kappa^{\otimes k})$, the set of smooth sections of the bundle $\kappa^{\otimes k}$. Indeed, we have a \mathbb{C} -linear isomorphism

$$\pi_k^* : \mathcal{C}^\infty(M, \kappa^{\otimes k}) \rightarrow \Omega_k$$

given by restriction to SM , i.e., in local coordinates (for $k \geq 0$),

$$\pi_k^*(w dz^k)(x, v) = w(x)(dz(v))^k.$$

This is a generalization of the map π_1^* which we have already encountered in (1.2) to identify smooth 1-forms with $\Omega_{-1} \oplus \Omega_1$. Note that the definition of π_k^* depends on the choice of the metric g . We denote by π_{k*} its L^2 -adjoint. In local coordinates, we have

$$(\pi_{k*}u)(x) = \left(\int_{\pi^{-1}(x)} u(x, \cdot) \right) dz^k.$$

Once we extend the operators to distributions by duality, the projection onto the k -th Fourier mode is simply given by $(2\pi)^{-1}\pi_k^*\pi_{k*}$ acting on $\mathcal{D}'(SM)$.

Under this identification, we can essentially think of the raising/lowering operators η_\pm as $\bar{\partial}$ and ∂ operators thanks to the following result (see [PSU14, Lemma 2.1] and the ensuing discussion).

Lemma 2.5. *For $k \geq 0$, the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{C}^\infty(M, \kappa^{\otimes k}) & \xrightarrow{\pi_k^*} & \Omega_k \\ \bar{\partial} \downarrow & & \downarrow \eta_- \\ \mathcal{C}^\infty(M, \kappa^{\otimes k} \otimes \bar{\kappa}) & \xrightarrow{\pi_{k-1}^*} & \Omega_{k-1} \end{array}$$

For $k \leq 0$, the operator π_k^* also intertwines the operators ∂ and η_+ .

As a result, for $k \geq 0$, the operator π_k^* gives us an identification

$$H_J^0(M, \kappa^{\otimes k}) \cong \Omega_k \cap \ker \eta_-.$$

We also introduce the following terminology:

Definition 2.6. A distribution $u \in \mathcal{D}'(SM)$ is said to be *fiberwise holomorphic* if $u_k = 0$ for all $k < 0$.

Equivalently, if we define the *Szegő projectors* $S_{\pm} : \mathcal{D}'(SM) \rightarrow \mathcal{D}'(SM)$ by

$$S_+u = \sum_{k \geq 0} u_k, \quad S_-u = \sum_{k \leq 0} u_k,$$

then a distribution u is fiberwise holomorphic if and only if $S_+u = u$. The projectors satisfy the commutation relations

$$[S_+, X]u = \eta_+u_{-1} - \eta_-u_0, \quad [S_-, X]u = \eta_-u_1 - \eta_+u_0. \quad (2.5)$$

We will be interested in the family of fiberwise holomorphic distributions that satisfy the transport equation:

$$\mathcal{D}'_{\text{tr},+}(SM) := \{u \in \mathcal{D}'(SM) \mid (F + V(\lambda))u = 0, S_+u = u\}. \quad (2.6)$$

2.2.3. Torelli's theorem. The complex vector space $H_J^0(M, \kappa)$ of J -holomorphic 1-forms has the same dimension as the genus of M (see [FK92, Proposition III.2.7]). Given a canonical basis $\{a_j, b_j\}$ of the homology $H_1(M; \mathbb{Z})$ on M , the following result gives us the existence of a useful basis (see [FK92, Proposition, p. 63]).

Proposition 2.7. *There exists a unique basis $\{\zeta_j\}$ for $H_J^0(M, \kappa)$ with the property*

$$\int_{a_j} \zeta_k = \delta_{jk}. \quad (2.7)$$

Furthermore, the matrix $\Pi(J)$ with (j, k) -entry

$$(\Pi(J))_{jk} := \int_{b_j} \zeta_k$$

is symmetric with positive definite imaginary part.

The space of symmetric matrices with positive definite imaginary part and size given by the genus of M is called the *Siegel upper half-space* $\mathcal{H}(M)$. We thus get a well-defined *period matrix* map

$$\Pi : \mathcal{T}(M) \rightarrow \mathcal{H}(M).$$

The following form of Torelli's theorem tells us that period matrices capture a lot of the information about the complex structure.

Theorem 2.8. *Assume that M has genus ≥ 2 . If $\Pi(J) = \Pi(\tilde{J})$, then there exists an orientation-preserving diffeomorphism $\psi : M \rightarrow M$ such that $\psi^*\tilde{J} = J$.*

We refer to [FK92, Theorem III.12.3] for a proof.

2.3. Hyperbolic dynamics. We now further assume that the flow of the generalized thermostat (M, g, λ) is *Anosov* (or *uniformly hyperbolic*).

2.3.1. *Definition.* Recall that the Anosov property means that there exists a flow-invariant continuous splitting

$$T(SM) = \mathbb{R}F \oplus E^s \oplus E^u$$

and uniform constants $C \geq 1$ and $0 < \rho < 1$ such that for all $t \geq 0$ we have

$$\|d\varphi_t|_{E^s}\| \leq C\rho^t, \quad \|d\varphi_{-t}|_{E^u}\| \leq C\rho^t. \quad (2.8)$$

In the geodesic case, the contact form α is preserved, so $\ker \alpha = \mathbb{H} \oplus \mathbb{V} = E^s \oplus E^u$. It is then known that $E^s \cap \mathbb{V} = \{0\} = E^u \cap \mathbb{V}$. For a generalized thermostat, we instead know by [DP07, Lemma 4.1] that

$$(\mathbb{R}F \oplus E^s) \cap \mathbb{V} = \{0\} = (\mathbb{R}F \oplus E^u) \cap \mathbb{V}. \quad (2.9)$$

Here $\mathbb{R}F \oplus E^{s/u}$ are the *weak stable and unstable bundles*. This implies that there exist $r^{s/u} \in \mathcal{C}^0(SM, \mathbb{R})$ such that

$$Y^s := H + r^s V \in \mathbb{R}F \oplus E^s, \quad Y^u := H + r^u V \in \mathbb{R}F \oplus E^u. \quad (2.10)$$

In fact, the weak stable and unstable bundles are \mathcal{C}^1 (see [Has94, Corollary 1.8]), so the functions $r^{s/u}$ are also \mathcal{C}^1 (and smooth along the flow since each bundle $\mathbb{R}F \oplus E^{s/u}$ is φ_t -invariant). The Anosov property implies that $r^s \neq r^u$ everywhere. One may in fact show that $r^s < r^u$, so the basis $\{F, Y^s, Y^u\}$ is positively-oriented.

Lemma 2.9. *Let (M, g, λ) be an Anosov generalized thermostat. Then, the functions $r^{s/u} \in \mathcal{C}^1(SM, \mathbb{R})$ uniquely characterized by (2.10) satisfy $r^s < r^u$.*

Proof. Since $r^s \neq r^u$ everywhere, it suffices to show the inequality at a single point. By compactness, we can pick $(x, v) \in SM$ such that $V(\lambda)(x, v) = 0$. Let us define

$$\xi(t) := d\varphi_{-t}(V_{\varphi_t(x, v)}).$$

Differentiating with respect to t and setting $t = 0$, we obtain

$$\dot{\xi}(0) = [F, V]_{(x, v)}.$$

Using that $[V, F] = H + V(\lambda)V$ yields

$$\dot{\xi}(0) = -H_{(x, v)}.$$

Since $r^s(x, v) \neq r^u(x, v)$, there exists a unique constant $c \in \mathbb{R}$ such that $V_{(x, v)} + cX_{(x, v)}$ belongs to $E^s \oplus E^u$. Therefore, given that E^s and E^u are uniformly attracting and repelling sets on $E^s \oplus E^u$ respectively, we must have $r^s(x, v) < r^u(x, v)$ at this point. \square

Remark 2.10. Note that, when $\lambda = 0$ and $K_g < 0$, i.e., in the geodesic case with negative curvature, we have the stronger statement $r^s < 0 < r^u$ because $[X, H] = K_g V$.




FIGURE 1. The relevant subbundles in tangent and cotangent spaces.

The dual bundles are defined by

$$(\mathbb{R}F)^*(E^s \oplus E^u) = 0 = (E^s)^*(\mathbb{R}F \oplus E^s) = (E^u)^*(\mathbb{R}F \oplus E^u).$$

One can check we have similar estimates to (2.8) for $(E^s)^*$ and $(E^u)^*$, with $d\varphi_t$ replaced by $d\varphi_t^{-T}$ (inverse transpose). Translated to the setting of the cotangent bundle, property (2.9) then becomes

$$(E^s)^* \cap \mathbb{H}^* = \{0\} = (E^u)^* \cap \mathbb{H}^*. \quad (2.11)$$

Further note that

$$(E^s)^* \oplus (E^u)^* = \Sigma,$$

where

$$\Sigma := \{(y, \eta) \in T^*(SM) \mid \eta(F(y)) = 0\}$$

is the characteristic set of the operator F (usually defined without the zero section).

2.3.2. Tensor tomography. The tensor tomography problem is interesting in its own right, particularly as it pertains to the injectivity of the X-ray transform for thermostats. We will need the following property in the case $n = 1$.

Definition 2.11. We say that a thermostat (M, g, λ) satisfies the *attenuated tensor tomography problem of order n* if having $(F + V(\lambda))u = f$ with $f, u \in \mathcal{C}^\infty(SM)$ and f of Fourier degree $n \geq 0$ implies that u is of Fourier degree $\max(n - 1, 0)$.

The term ‘attenuated’ refers to the presence of the divergence $V(\lambda)$ in the transport equation. Note that such a term appears for Gaussian thermostats and generalized thermostats of higher Fourier degree, but not for magnetic or geodesic flows.

The fact that geodesic flows satisfy the (attenuated) tensor tomography problem was first proved in negative curvature in [GK80] for $n \geq 0$ and then generalized to the Anosov case in [DS03] for $n \leq 1$, [PSU14] for $n \leq 2$, and [Gui17] for $n \geq 2$. It

was also shown in [DP05] that Anosov magnetic flows satisfy the (attenuated) tensor tomography problem of order $n \leq 1$.

For generalized thermostats of higher Fourier degree, the non-attenuated and attenuated versions of the tensor tomography problem are different. In [DP07], it was proved that Gaussian thermostats (potentially mixed with a magnetic component) satisfy the non-attenuated tensor tomography problem of order $n \leq 1$. We instead need:

Theorem 2.12. *Any Gaussian thermostat (M, g, θ) with $\mathbb{K} < 0$ satisfies the attenuated tensor tomography problem of order 1.*

This result is a consequence of the work in [AR21]. Their argument heavily relies on the negative thermostat curvature assumption. In particular, most of the heavy lifting is done by [AR21, Theorem 3.1], where the *Carleman estimates* for Gaussian thermostats with negative curvature are established (akin to the work in [PS23]):

Theorem 2.13. *Let (M, g, θ) be a Gaussian thermostat with $\mathbb{K} \leq -\kappa$ for some $\kappa > 0$. For any integer $m \geq 1$ and parameter $s > 0$, we have*

$$\sum_{k \geq m} |k|^{2s+1} \|u_k\|^2 \leq \frac{1}{\kappa s} \sum_{k \geq m+1} |k|^{2s+1} \|(Fu)_k\|^2$$

for all $u \in C^\infty(SM)$.

The rest of the argument is then relatively straightforward for our case, which is less general than the one tackled in [AR21]. We include it here for the sake of completeness, but also to show how it can be simplified.

Proposition 2.14. *Let (M, g, θ) be a Gaussian thermostat with $\mathbb{K} < 0$. Suppose $f \in C^\infty(SM)$ has finite Fourier degree and $u \in C^\infty(SM)$ satisfies $(F + V(\lambda))u = f$. Then u also has finite Fourier degree.*

Proof. We follow the argument from [AR21, Theorem 5.1]. Let $m' \geq 0$ be the Fourier degree of f . Since $(F + V(\lambda))u = f$, we obtain

$$(Fu)_k = -i\lambda_1 u_{k-1} + i\lambda_{-1} u_{k+1} \quad \text{for all } |k| \geq m' + 1.$$

As a result, there exists $C > 0$ such that

$$\|(Fu)_k\|^2 \leq C(\|u_{k-1}\|^2 + \|u_{k+1}\|^2) \quad \text{for all } |k| \geq m' + 1.$$

Pick $\kappa > 0$ such that $\mathbb{K} \leq -\kappa$, fix $s > eC/\kappa$, and let $m \geq \max(2s + 1, m' + 1)$. We can apply Theorem 2.13 to get

$$\begin{aligned} \sum_{|k| \geq m} |k|^{2s+1} \|u_k\|^2 &\leq \frac{C}{\kappa s} \sum_{|k| \geq m+1} |k|^{2s+1} (\|u_{k-1}\|^2 + \|u_{k+1}\|^2) \\ &\leq \frac{C}{\kappa s} \sum_{|k| \geq m} (|k| + 1)^{2s+1} \|u_k\|^2. \end{aligned}$$

Since $m \geq 2s + 1$, we note that

$$(|k|+1)^{2s+1} = \left(1 + \frac{1}{|k|}\right)^{2s+1} |k|^{2s+1} \leq \left(1 + \frac{1}{|k|}\right)^{|k|} |k|^{2s+1} \leq e|k|^{2s+1} \quad \text{for all } |k| \geq m,$$

so that

$$\sum_{|k| \geq m} |k|^{2s+1} \|u_k\|^2 \leq \frac{eC}{\kappa s} \sum_{|k| \geq m} |k|^{2s+1} \|u_k\|^2.$$

It hence follows that

$$\left(1 - \frac{eC}{\kappa s}\right) \sum_{|k| \geq m} |k|^{2s+1} \|u_k\|^2 \leq 0.$$

However, we have $1 - eC/(\kappa s) > 0$ by design, so $u_k = 0$ for all $|k| \geq m$. □

Proof of Theorem 2.12. By Proposition 2.14, we know that u is of finite Fourier degree. Suppose, for the sake of contradiction, that u is of degree $k \geq 1$. Then, using the equation $(F + V(\lambda))u = f$, we have

$$\left(\eta_+ + \left(1 + \frac{1}{k}\right) \lambda_1 V\right) u_k = (\eta_+ + \lambda_1 V + i\lambda_1) u_k = 0$$

and

$$\left(\eta_- + \left(1 + \frac{1}{k}\right) \lambda_{-1} V\right) u_{-k} = (\eta_- + \lambda_{-1} V - i\lambda_{-1}) u_{-k} = 0.$$

By [AZ17, Proposition 6.1], it follows that $u_{\pm k} = 0$, which is a contradiction. □

Finally, the proofs of our theorems rely on the possibility of lifting arbitrary holomorphic 1-forms to solutions of the transport equation. As explained in Appendix A, where we have relegated most of the work on this front, this is again related to the injectivity of the X-ray transform for thermostats.

Theorem 2.15. *Let (M, g, λ) , with λ of Fourier degree ≤ 1 , be an Anosov thermostat. For any holomorphic (resp. anti-holomorphic) 1-form τ on M , there exists $u \in H^{-1}(SM)$ with $u_k = 0$ for all $k \leq 0$ (resp. $k \geq 0$) such that $(F + V(\lambda))u = 0$ and $u_1 = \pi_1^* \tau$ (resp. $u_{-1} = \pi_1^* \tau$).*

Proof. Let us treat the case where the 1-form τ is holomorphic. The anti-holomorphic case is completely analogous. Using Lemma 2.5, we know that $\pi_1^* \tau \in \Omega_1$ is in the kernel of η_- . We can hence apply Theorem A.5, which tells us that there exists $v \in H^{-1}(SM)$ with $(F + V(\lambda))v = 0$ such that $v_1 = \pi_1^* \tau$ and $v_{-1} = v_0 = 0$. We project this distribution onto its positive Fourier components to get $u := S_+ v = \sum_{k \geq 1} v_k$. For all $k \in \mathbb{Z}$, we then have

$$((F + V(\lambda))u)_k = \eta_+ u_{k-1} + \eta_- u_{k+1} + ik(\lambda_1 u_{k-1} + \lambda_0 u_k + \lambda_{-1} u_{k+1}) = 0,$$

which entails that $(F + V(\lambda))u = 0$. □

We note that we cannot hope to get such a result for an arbitrary $\lambda \in \mathcal{C}^\infty(SM, \mathbb{R})$.

Lemma 2.16. *Suppose $\lambda = \lambda_m + \lambda_{-m}$ where $m \geq 2$ and $\eta_- \lambda_{-m} = 0$. Since λ_{-m} has isolated zeroes, there exists $a \in \Omega_1$ with $\eta_- a = 0$ and $\lambda_{-m} a \neq 0$. Then, there is no $u \in H^{-1}(SM)$ with $u_k = 0$ for all $k \leq 0$ such that $(F + V(\lambda))u = 0$ and $u_1 = a$.*

Proof. Suppose such a distribution u exists. For any $k \in \mathbb{Z}$, we must have

$$0 = ((F + V(\lambda))u)_k = \eta_+ u_{k-1} + \eta_- u_{k+1} + ik(\lambda_m u_{k-m} + \lambda_{-m} u_{k+m}).$$

Therefore, applying this to $k = -m+1$, we get $\lambda_{-m} a = \lambda_{-m} u_1 = 0$, a contradiction. \square

3. ACTION ON HOLOMORPHIC DIFFERENTIALS

We have seen that, by passing through a specific type of 2-currents instead of directly using the pullback ϕ^* , the linear map $\Phi : \mathcal{D}'_{\text{tr}}(SM) \rightarrow \mathcal{D}'_{\text{tr}}(\tilde{S}M)$ defined in (2.3) sends distributional solutions to the transport equation of one generalized thermostat to those of the second. In this section, we want to show that Φ can also be seen as acting on holomorphic differentials from one complex surface to another when λ is of Fourier degree ≤ 1 and the attenuated tensor tomography problem of order 1 is satisfied.

3.1. Action on fiberwise holomorphic distributions. We start by studying the action of Φ on the subspace $\mathcal{D}'_{\text{tr},+}(SM)$ defined in (2.6). This will require some microlocal analysis.

We introduce $\mathcal{C} \subseteq \{(y, \eta) \in T^*(SM) \mid \eta(F(y)) = 0\}$, the closed cone enclosed by $(E^s)^*$ and $(E^u)^*$ in the half-space $\{(y, \eta) \in T^*(SM) \mid \eta(V(y)) \geq 0\}$. See Figure 1.

Lemma 3.1. *Let (M, g, λ) be an Anosov generalized thermostat. If $u \in \mathcal{D}'_{\text{tr},+}(SM)$, then $\text{WF}(u) \subseteq \mathcal{C}$ and $u_k \in \Omega_k$ for all $k \in \mathbb{Z}$.*

Proof. The argument is essentially the same as that of [GLP24, Lemma 2.5]. Let us give the details.

By definition, each $u \in \mathcal{D}'_{\text{tr},+}(SM)$ satisfies $S_+ u = u$. Using the wavefront set description of the Schwartz kernel of S_+ (see [Gui17, Lemma 3.10]), we thus get

$$\text{WF}(u) = \text{WF}(S_+ u) \subset \{(y, \eta) \in T^*(SM) \mid \eta(V(y)) \geq 0\}.$$

Given that $(F + V(\lambda))u = 0$, elliptic regularity tells us that

$$\text{WF}(u) \subset \Sigma = \{(y, \eta) \in T^*(SM) \mid \eta(F(y)) = 0\}.$$

By propagation of singularities for real principal type differential operators (see [Hö09, Theorem 26.1.1]), we further know that $\text{WF}(u)$ is invariant by the symplectic lift of the flow φ_t . Given the Anosov property, the maximal flow-invariant subset of $T^*(SM)$ contained in $\{\eta(F(y)) = 0, \eta(V(y)) \geq 0\}$ is \mathcal{C} , so this gives us the first claim.

For the second claim, recall that $u_k = (2\pi)^{-1}\pi_k^*\pi_{k*}u$. The pushforward operator π_{k*} only selects the wavefront set in $(\mathbb{R}F)^* \oplus \mathbb{H}^*$ (see [FT99, Proposition 11.3.3]), which is empty given that $\mathcal{C} \cap ((\mathbb{R}F)^* \oplus \mathbb{H}^*) = \{0\}$ by property (2.11). Therefore $u_k \in \Omega_k$. \square

We will also need the following lemma with the same proof as [GLP24, Lemma 3.3].

Lemma 3.2. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ be Anosov generalized thermostats. Suppose there exists a smooth orbit equivalence $\phi : \tilde{S}M \rightarrow SM$ isotopic to the identity between them. Then ϕ preserves the natural orientation of the weak unstable bundle, namely that given by the basis $\{F, Y^u\}$.*

Armed with this, we can show that Φ maps fiberwise holomorphic distributional solutions to the transport equation of one thermostat to those of the second. For this step of the proof, however, we restrict to thermostats where λ has Fourier degree ≤ 1 and the attenuated tensor tomography problem of order 1 is satisfied.

Proposition 3.3. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$, with λ and $\tilde{\lambda}$ of Fourier degree ≤ 1 , be Anosov thermostats satisfying the attenuated tensor tomography problem of order 1. Suppose there exists a smooth orbit equivalence $\phi : \tilde{S}M \rightarrow SM$ isotopic to the identity between them. Then the map Φ defined in (2.3) yields a \mathbb{C} -linear isomorphism*

$$\Phi : \mathcal{D}'_{\text{tr},+}(SM) \rightarrow \mathcal{D}'_{\text{tr},+}(\tilde{S}M).$$

Proof. Since $d\phi^T$ maps connected sets to connected sets, $(E^u)^*$ to $(\tilde{E}^u)^*$, and $(E^s)^*$ to $(\tilde{E}^s)^*$, $d\phi^T(\mathcal{C})$ must be one of the four cones depicted on the right of Figure 1 (inside the characteristic set $\tilde{\Sigma}$). It follows that $d\phi^T(\mathcal{C}) = \pm\tilde{\mathcal{C}}$ because any other cone would entail that ϕ reverses the orientation, which is impossible since it is assumed to be isotopic to the identity. If $d\phi^T(\mathcal{C}) = -\tilde{\mathcal{C}}$, then ϕ would flip the orientation of the weak unstable leaves, which contradicts Lemma 3.2. Therefore $d\phi^T(\mathcal{C}) = \tilde{\mathcal{C}}$.

Let $u \in \mathcal{D}'_{\text{tr},+}(SM)$ and $\tilde{u} := \Phi u$. By Lemma 3.1, we know that $\text{WF}(u) \subseteq \mathcal{C}$. By Lemma 2.4, we thus know that $\text{WF}(\tilde{u}) = d\phi^T(\text{WF}(u)) \subseteq \tilde{\mathcal{C}}$. Then $\tilde{S}_-\tilde{u} \in \mathcal{C}^\infty(\tilde{S}M)$ and, since $(\tilde{F} + \tilde{V}(\tilde{\lambda}))\tilde{u} = 0$, we also have that

$$\begin{aligned} (\tilde{F} + \tilde{V}(\tilde{\lambda}))\tilde{S}_-\tilde{u} &= [\tilde{F} + \tilde{V}(\tilde{\lambda}), \tilde{S}_-]\tilde{u} \\ &= \left(\tilde{\eta}_+ + \tilde{\lambda}_1\tilde{V} + i\tilde{\lambda}_1\right)\tilde{u}_0 - \left(\tilde{\eta}_- + \tilde{\lambda}_{-1}\tilde{V} - i\tilde{\lambda}_{-1}\right)\tilde{u}_1. \end{aligned}$$

Since $(M, \tilde{g}, \tilde{\lambda})$ satisfies the tensor tomography problem of order 1 by assumption, it follows that $\tilde{S}_-\tilde{u}$ is of Fourier degree 0. Hence \tilde{u} is fiberwise holomorphic. The fact that Φ is an isomorphism is then clear as it admits an inverse, namely the map associated to $(\phi^{-1})^*$ by (2.3). \square

3.2. Extension operator. Next, we show how Φ can be seen as acting on holomorphic differentials from one complex surface to another. Let us start by noting that the map

$$\pi_{1*} : \mathcal{D}'_{\text{tr},+}(SM) \rightarrow H_J^0(M, \kappa). \quad (3.1)$$

is well-defined. Indeed, if $u \in \mathcal{D}'_{\text{tr},+}(SM)$, then $Xu + V(\lambda u) = (F + V(\lambda))u = 0$, which means that $(Xu)_0 = 0$ and hence $\eta_- u_1 = 0$. By Lemma 2.5, this is equivalent to $\bar{\partial}\pi_{1*}u = 0$.

Thanks to Theorem 2.15, we know that the map (3.1) is surjective. We can thus define a right-inverse

$$e_1 : H_J^0(M, \kappa) \rightarrow \mathcal{D}'_{\text{tr},+}(SM)$$

such that $\pi_{1*} \circ e_1 = \text{id}_{H_J^0(M, \kappa)}$. We call it an *extension operator*. We may then define the map

$$\Psi : H_J^0(M, \kappa) \rightarrow H_{\tilde{J}}^0(M, \kappa)$$

by the commutative diagram

$$\begin{array}{ccc} \mathcal{D}'_{\text{tr},+}(SM) & \xrightarrow{\Phi} & \mathcal{D}'_{\text{tr},+}(\tilde{S}M) \\ e_1 \uparrow & & \downarrow \pi_{1*} \\ H_J^0(M, \kappa) & \xrightarrow{\Psi} & H_{\tilde{J}}^0(M, \kappa). \end{array} \quad (3.2)$$

3.3. Period preservation. The following result shows that the induced mapping of holomorphic differentials we have just defined preserves additional structure.

Proposition 3.4. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$, with λ and $\tilde{\lambda}$ of Fourier degree ≤ 1 , be Anosov thermostats satisfying the attenuated tensor tomography problem of order 1. The \mathbb{C} -linear map*

$$\Psi : H_J^0(M, \kappa) \rightarrow H_{\tilde{J}}^0(M, \kappa)$$

is an isomorphism. It preserves periods in the sense that, for all $[\gamma] \in H_1(M; \mathbb{Z})$ and $\tau \in H_J^0(M, \kappa)$, we have

$$\int_{[\gamma]} \tau = \int_{[\gamma]} \Psi \tau.$$

Recall that there is a push-forward map $\pi_* : \mathcal{C}^\infty(SM, \Omega^2(SM)) \rightarrow \mathcal{C}^\infty(M, \Omega^1(M))$ given by integration along fibers. It satisfies $d\pi_* = \pi_* d$ (see [BT82, Proposition 6.14]), and it extends to currents. By [BT82, Proposition 6.15], we have the projection formula

$$\int_{\pi^{-1}(\gamma)} \sigma = \int_{\gamma} \pi_* \sigma \quad (3.3)$$

for any smooth oriented curve γ on M and any 2-form σ on SM .

Lemma 3.5. *Let (M, g, λ) be a generalized thermostat. For any $u \in \mathcal{D}'(SM)$, we have*

$$\pi_*(u\omega) = \frac{1}{2} \star (\pi_{-1*}u + \pi_{1*}u).$$

Proof. It suffices to establish the claim for $u \in \mathcal{C}^\infty(SM)$. Recall that $\omega := \iota_F\mu$. A quick computation then yields

$$\omega = \beta \wedge \psi + \lambda \alpha \wedge \beta. \quad (3.4)$$

Note that $\pi^*\mu_a = \alpha \wedge \beta$, where μ_a is the area form on M .

Pick $x \in M$ and take $w \in S_xM$. Then, by definition, we have

$$\pi_*(u\omega)_x(w) = \int_{S_xM} \iota_{\tilde{w}}(u\omega),$$

where $\tilde{w} \in T(SM)$ is a lift of w under $d\pi$. We take $\tilde{w} = (w, 0)$, i.e., no component in the subbundle $\mathbb{V} = \ker d\pi$. Then, since $\psi(\tilde{w}) = 0$, we get

$$\pi_*(u\omega)_x(w) = \int_{S_xM} \iota_{\tilde{w}}(\beta)u\psi.$$

Given that $\iota_{\tilde{w}}\beta_{(x,v)} = g_x(w, Jv)$, we obtain

$$\begin{aligned} \pi_*(u\omega)_x(Jw) &= \int_{S_xM} g_x(w, v)u(x, v) dv \\ &= \int_0^{2\pi} (\cos t)u(\rho_t(x, w)) dt \\ &= \frac{1}{2} \int_0^{2\pi} (e^{it} + e^{-it})u(\rho_t(x, w)) dt \\ &= \pi(u_{-1} + u_1)(x, w), \end{aligned}$$

where in the last equality we used the formula (2.4). In terms of 1-forms, since $u_k = (2\pi)^{-1}\pi_k^*\pi_{k*}u$, we proved that

$$-\star \pi_*(u\omega) = \frac{1}{2} (\pi_{-1*}u + \pi_{1*}u).$$

We conclude by applying \star to both sides. □

We can then integrate this identity, applied to solutions of the transport equation, over closed thermostat geodesics to obtain the following result.

Lemma 3.6. *Let (M, g, λ) be an Anosov generalized thermostat and γ a closed thermostat geodesic. For any $u \in \mathcal{D}'_{\text{tr}}(SM)$, the pairing $\langle \pi^{-1}(\gamma), u\omega \rangle$ is well-defined and*

$$\int_{\pi^{-1}(\gamma)} u\omega = \frac{1}{2} \int_{[\gamma]} \star(\pi_{-1*}u + \pi_{1*}u).$$

Proof. By the wavefront set calculus, the pairing $\langle \pi^{-1}(\gamma), u\omega \rangle$ is well-defined whenever

$$N^*(\pi^{-1}(\gamma)) \cap \text{WF}(u) = \emptyset \quad (3.5)$$

(see [Hö03, Corollary 8.2.7] for instance). The conormal bundle $N^*(\pi^{-1}(\gamma))$ consists of a line contained in $(\mathbb{R}F)^* \oplus \mathbb{H}^*$, so Lemma 3.1 and property (2.11) tell us that the intersection with $\text{WF}(u)$ is indeed empty. It follows that the pairing $\langle \pi^{-1}(\gamma), u\omega \rangle$ is well-defined and extends the pairing computed for $u \in \mathcal{C}^\infty(SM)$.

We can then apply the projection formula (3.3) and Lemma 3.5. As seen in Lemma 2.3, the 2-current $u\omega$ is closed if $u \in \mathcal{D}'_{\text{tr}}(SM)$, so $\star(\pi_{-1*}u + \pi_{1*}u)$ is also closed given that $\pi_*d = d\pi_*$, which implies that its integral only depends on the homology class $[\gamma]$. \square

As $\pi_1^*\star = -V\pi_1^*$, for any $u \in \mathcal{D}'(SM)$ we may write

$$\star(\pi_{-1*}u + \pi_{1*}u) = i(\pi_{-1*}u - \pi_{1*}u).$$

Therefore, if $\tau \in H^0_j(M, \kappa)$, $[\gamma] \in H_1(M; \mathbb{Z})$, and γ is any thermostat geodesic whose homology class is $[\gamma]$, Lemma 3.6 gives us

$$2i \int_{\pi^{-1}(\gamma)} e_1(\tau)\omega = \int_{[\gamma]} \tau. \quad (3.6)$$

We can now tackle the proof of Proposition 3.4.

Proof of Proposition 3.4. Let $[\gamma] \in H_1(M; \mathbb{Z})$ and let $\gamma, \tilde{\gamma}$ be two thermostat geodesics (with respect to (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$) whose homology class is $[\gamma]$. Since ϕ is isotopic to the identity, we know that $[\phi(\pi^{-1}(\tilde{\gamma}))] = [\pi^{-1}(\gamma)]$ in $H_2(SM; \mathbb{Z})$.

We claim that the pairing $\langle \phi(\pi^{-1}(\tilde{\gamma})), e_1(\tau)\omega \rangle$ is well-defined. The tangent space to $\pi^{-1}(\tilde{\gamma})$ is $(\mathbb{R}\tilde{F}) \oplus \tilde{V}$. By property (2.9), it trivially intersects the closed cone $\tilde{\mathcal{B}} \subseteq \tilde{E}^s \oplus \tilde{E}^u$ enclosed by \tilde{E}^s and \tilde{E}^u , and whose orthogonal projection onto $\tilde{\mathbb{H}} \oplus \tilde{V}$ avoids \tilde{V} . Since ϕ preserves the orientation, the same arguments as in Proposition 3.3 tell us that the tangent space to $\phi(\pi^{-1}(\tilde{\gamma}))$ intersects the closed cone \mathcal{B} trivially. As a result, its conormal $N^*(\phi(\pi^{-1}(\tilde{\gamma})))$ avoids the closed cone \mathcal{C} , which contains $\text{WF}(e_1(\tau))$ by Lemma 3.1. The wavefront set condition (3.5) is hence satisfied.

The 2-current $e_1(\tau)\omega$ is closed by Lemma 2.3. By the Hodge decomposition theorem, we may hence write $e_1(\tau)\omega = \sigma + df$ for some harmonic 2-current σ and 1-current f with $\text{WF}(f) = \text{WF}(e_1(\tau))$. Thanks to the wavefront set condition, the same argument as for $e_1(\tau)\omega$ then shows that both pairings $\langle \pi^{-1}(\gamma), df \rangle$ and $\langle \phi(\pi^{-1}(\tilde{\gamma})), df \rangle$ are well-defined. They must be equal to 0 since df is exact. We thus get

$$\int_{\pi^{-1}(\gamma)} e_1(\tau)\omega = \int_{\pi^{-1}(\gamma)} \sigma = \int_{\phi(\pi^{-1}(\tilde{\gamma}))} \sigma = \int_{\phi(\pi^{-1}(\tilde{\gamma}))} e_1(\tau)\omega,$$

where in the second equality we have used the fact that σ is harmonic and $[\pi^{-1}(\gamma)] = [\phi(\pi^{-1}(\tilde{\gamma}))]$ in $H_2(SM; \mathbb{Z})$.

We can now use (3.6) and unravel the definitions to obtain

$$\begin{aligned} \int_{[\gamma]} \tau &= 2i \int_{\pi^{-1}(\gamma)} e_1(\tau) \omega \\ &= 2i \int_{\phi(\pi^{-1}(\tilde{\gamma}))} e_1(\tau) \omega = 2i \int_{\pi^{-1}(\tilde{\gamma})} \phi^*(e_1(\tau) \omega) \\ &= 2i \int_{\pi^{-1}(\tilde{\gamma})} \Phi(e_1(\tau)) \tilde{\omega} = \int_{[\gamma]} \pi_{1*} \Phi(e_1(\tau)) = \int_{[\gamma]} \Psi \tau. \end{aligned}$$

□

4. END OF THE PROOFS

4.1. Torelli's theorem. The work from the previous section, when combined with Torelli's theorem, tells us that a smooth orbit equivalence isotopic to the identity determines the class $[J]$ in the moduli space $\mathcal{M}(M)$ of complex structures on M .

Proposition 4.1. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$, with λ and $\tilde{\lambda}$ of Fourier degree ≤ 1 , be Anosov thermostats satisfying the attenuated tensor tomography problem of order 1. If there exists a smooth orbit equivalence isotopic to the identity between them, then $[J] = [\tilde{J}]$ in $\mathcal{M}(M)$. Equivalently, there exists a diffeomorphism $\psi : M \rightarrow M$ such that $\psi^* \tilde{J} = J$ and $\psi^* \tilde{g} = e^{2f} g$ for some $f \in C^\infty(M, \mathbb{R})$.*

Proof. By Proposition 3.4, the map $\Psi : H_J^0(M, \kappa) \rightarrow H_{\tilde{J}}^0(M, \kappa)$ is a period-preserving \mathbb{C} -linear isomorphism. This means that (M, J) and (M, \tilde{J}) have the same period matrix. Indeed, given a canonical basis $\{a_j, b_j\}$ of the homology $H_1(M; \mathbb{Z})$ on M , let $\{\zeta_j\}$ be a basis for $H_J^0(M, \kappa)$ such that property (2.7) is satisfied. Then $\{F\zeta_j\}$ is a basis for $H_{\tilde{J}}^0(M, \kappa)$ such that (2.7) is also satisfied, and

$$(\pi(J))_{jk} = \int_{b_j} \zeta_k = \int_{b_j} F\zeta_k = (\pi(\tilde{J}))_{jk}.$$

Since the surface M must be of genus ≥ 2 , Theorem 2.8 tells us that there exists an orientation-preserving diffeomorphism $\psi : M \rightarrow M$ such that $\psi^* J = \tilde{J}$.

□

In this section, we want to show something stronger, namely, that the class of the complex structure J is determined in Teichmüller space $\mathcal{T}(M)$.

Proposition 4.2. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ be either*

(a) *two Anosov magnetic systems, or*

(b) two Gaussian thermostats with $\mathbb{K}, \tilde{\mathbb{K}} < 0$.

If there exists a smooth orbit equivalence isotopic to the identity between them, then $[J] = [\tilde{J}]$ in $\mathcal{T}(M)$. Equivalently, there exists a diffeomorphism $\psi : M \rightarrow M$, isotopic to the identity, such that $\psi^* \tilde{J} = J$ and $\psi^* \tilde{g} = e^{2f} g$ for some $f \in \mathcal{C}^\infty(M, \mathbb{R})$.

The reason we needed to specify the nature of the two thermostats, as opposed to Proposition 4.1 which deals with more general Anosov thermostats of Fourier degree ≤ 1 , is that our proof relies the following technical lemma (see [GLP24, Lemma 3.8]).

Lemma 4.3. *Let J and \tilde{J} be two complex structures on M compatible with orientation such that $[J] \neq [\tilde{J}]$ in $\mathcal{T}(M)$. Then, there exists a finite cover $M' \rightarrow M$ such that the lifts $[J'], [\tilde{J}'] \in \mathcal{T}(M')$ are not in the same $\text{MCG}(M')$ -orbit.*

Indeed, when lifting the thermostats to finite covers, the properties of being Anosov and having negative thermostat curvature are preserved, but, a priori, satisfying the attenuated tensor tomography problem of order 1 is not.

Proof Proposition 4.2. Suppose, for the sake of contradiction, that $[J] \neq [\tilde{J}]$ in $\mathcal{T}(M)$. By Lemma 4.3, there exists a finite cover $M' \rightarrow M$ such that the lifts $[J']$ and $[\tilde{J}']$ are not in the same $\text{MCG}(M')$ -orbit.

Since the smooth orbit equivalence between the flows of (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ is isotopic to the identity, it can be lifted to a smooth orbit equivalence isotopic to the identity between the flows of $(M, p^*g, \lambda \circ dp)$ and $(M, p^*\tilde{g}, \tilde{\lambda} \circ dp)$.

In case (a), we know that the lifted Anosov magnetic flows on M' are again Anosov because the cover is finite. They hence satisfy the (attenuated) tensor tomography problem of order 1. In case (b), we know that the lifted Gaussian thermostats also have negative thermostat curvature since the property is local. By Theorem 2.12, we thus conclude that they satisfy the attenuated tensor tomography problem of order 1.

We can then apply Proposition 4.1 to obtain a contradiction. \square

Proposition 4.2 gives us most of Theorem 1.1 and Theorem 1.2. All that is left is studying the rigidity of the function λ in each case.

4.2. Rigidity of the magnetic field. Since $\lambda \in \mathcal{C}^\infty(SM, \mathbb{R})$ does not depend on the velocity in the magnetic case, we will think of it as living in $\mathcal{C}^\infty(M, \mathbb{R})$.

Lemma 4.4. *Let (M, g, λ) be an Anosov magnetic flow. Then, the 2-form ω on SM defined in (2.2) is exact.*

Proof. By (3.4), we have

$$\omega = -d\alpha + \pi^*(\lambda\mu_a).$$

By the Gauss-Bonnet theorem and the fact M must be of genus ≥ 2 , we know that

$$\int_M K_g \mu_a = 2\pi\chi(M) < 0,$$

so $[K_g \mu_a] \neq 0$ in $H^2(M; \mathbb{R}) \cong \mathbb{R}$. It follows that we may write $\lambda \mu_a = cK_g \mu_a + d\varrho$ for some 1-form ϱ on M and $c \in \mathbb{R}$. The constant c is explicitly given by

$$c = \frac{1}{2\pi\chi(M)} \int_M \lambda \mu_a. \quad (4.1)$$

But then, since $d\psi = -\pi^*(K_g \mu_a)$, we obtain

$$\pi^*(\lambda \mu_a) = c\pi^*(K_g \mu_a) + d\pi^*\varrho = d(-c\psi + \pi^*\varrho),$$

which allows us to write $\omega = d\tau$ for the 1-form

$$\tau := -\alpha - c\psi + \pi^*\varrho. \quad (4.2)$$

□

Knowing how to find primitives of ω in the magnetic case then unlocks the following.

Proposition 4.5. *Let (M, g, λ) and $(M, g, \tilde{\lambda})$ be two Anosov magnetic systems with the same background metric g . Suppose there is a smooth conjugacy $\phi : SM \rightarrow SM$, isotopic to the identity, between them. Then, we have $[\tilde{\lambda} \mu_a] = \pm[\lambda \mu_a]$ in $H^2(M; \mathbb{R})$. Moreover, $\lambda = 0$ if and only if $\tilde{\lambda} = 0$.*

Proof. Define the τ as in (4.2) to be a primitive of ω . Contracting it with F yields

$$\tau(F) = -1 - c\pi^*\lambda + \pi_1^*\varrho. \quad (4.3)$$

Moreover, we know that the Anosov magnetic flows are transitive and ϕ is isotopic to the identity, so $\phi^*\mu = \mu$. Since $\phi_*\tilde{F} = F$, contracting $\phi^*\mu = \mu$ with \tilde{F} yields $\phi^*\omega = \tilde{\omega}$. This can be rewritten as $\phi^*d\tau = d\tilde{\tau}$, which in turn implies

$$d(\phi^*\tau - \tilde{\tau}) = 0.$$

Since $\pi^* : H^1(M; \mathbb{R}) \rightarrow H^1(SM; \mathbb{R})$ is an isomorphism, there exists a closed 1-form φ on M and $f \in C^\infty(SM)$ such that

$$\phi^*\tau - \tilde{\tau} = \pi^*\varphi + df.$$

Contracting with \tilde{F} yields

$$\tau(F) \circ \phi = \tilde{\tau}(\tilde{F}) + \pi_1^*\varphi + \tilde{F}f.$$

By (4.3), we thus get

$$-1 + (-c\pi^*\lambda + \pi_1^*\varrho) \circ \phi = -1 - \tilde{c}\pi^*\tilde{\lambda} + \pi_1^*\tilde{\varrho} + \pi_1^*\varphi + \tilde{F}f,$$

which simplifies to

$$(-c\pi^*\lambda + \pi_1^*\varrho) \circ \phi = -\tilde{c}\pi^*\tilde{\lambda} + \pi_1^*\tilde{\varrho} + \pi_1^*\varphi + \tilde{F}f. \quad (4.4)$$

If we integrate with respect to μ , we obtain (since $\phi^*\mu = \mu$)

$$c \int_{SM} \pi^*(\lambda)\mu = \tilde{c} \int_{SM} \pi^*(\tilde{\lambda})\mu.$$

We thus have $c^2 = \tilde{c}^2$ by (4.1), which shows that the cohomology classes of $\lambda\mu_a$ and $\tilde{\lambda}\mu_a$ in $H^2(M; \mathbb{R})$ are the same up to a sign.

If $\lambda = 0$, we may take $\varrho = 0$, and we know that $c = 0$ thanks to (4.1). It follows that $\tilde{c} = 0$. Let $\tilde{\gamma}$ be a closed magnetic geodesic of $(M, g, \tilde{\lambda})$ and $\tilde{\Gamma} := (\tilde{\gamma}, \dot{\tilde{\gamma}}) \subseteq SM$. Relation (4.4) allows us to write

$$\int_{\tilde{\Gamma}} \pi_1^*(\tilde{\varrho} + \varphi) = 0.$$

By the smooth Livšic theorem [dILMM86, Theorem 2.1] and [DP05, Theorem B], this means that $\tilde{\varrho} + \varphi$ is exact. Since φ is closed, we get $d\tilde{\varrho} = 0$, which in turn implies $\tilde{\lambda} = 0$, as desired. \square

We can now conclude the proof of Theorem 1.1.

Proof of Theorem 1.1. If two Anosov magnetic systems (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$ are related by a smooth conjugacy $\phi : \tilde{SM} \rightarrow SM$, isotopic to the identity, Proposition 4.2 yields a diffeomorphism $\psi : M \rightarrow M$ isotopic to the identity such that $\psi^*\tilde{g} = e^{2f}g$ for some $f \in C^\infty(M, \mathbb{R})$.

If ϕ is a conjugacy and $f = 0$, the map $\phi \circ d\psi : SM \rightarrow SM$ gives us a smooth conjugacy isotopic to the identity between the Anosov magnetic flows $(M, g, \psi^*\tilde{\lambda})$ and (M, g, λ) . Thus Proposition 4.5 tells us that $[\psi^*(\tilde{\lambda})\mu_a] = \pm[\lambda\mu_a]$ in $H^2(M; \mathbb{R})$ and that $\lambda = 0$ if and only if $\tilde{\lambda} = 0$. \square

4.3. Rigidity of the thermostat 1-form. Given the conclusion of Proposition 4.2, it behooves us to understand the behavior of (generalized) thermostat flows under a conformal re-scaling of the metric.

Lemma 4.6. *Let (M, g, λ) be a generalized thermostat, and define a conformal re-scaling $\tilde{g} := e^{-2f}g$ of the metric, for some $f \in C^\infty(M, \mathbb{R})$. Then, the scaling map $s : SM \rightarrow \tilde{SM}$ defined in (1.1), which in this case is simply $(x, v) \mapsto (x, e^f v)$, satisfies*

$$s_*X = e^{-f}(\tilde{X} - \tilde{\pi}_1^*(\star df)\tilde{V}), \quad s_*V = \tilde{V}.$$

In particular, the map s represents a smooth orbit equivalence isotopic to the identity from (M, g, λ) to

$$(M, \tilde{g}, e^f(\lambda \circ s^{-1}) - \tilde{\pi}_1^*(\star df)) \tag{4.5}$$

*with a time-change $s_*F = e^{-f}\tilde{F}$.*

Proof. The first statement is proved in [CP22, Lemma B.1]. The conclusion then follows from the calculation

$$\begin{aligned} s_*F &= s_*X + (\lambda \circ s^{-1})s_*V \\ &= e^{-f}(\tilde{X} - \tilde{\pi}_1^*(\star df)\tilde{V}) + (\lambda \circ s^{-1})\tilde{V}. \end{aligned}$$

□

In what follows, let θ be the 1-form on M satisfying $\pi_1^*\theta = \lambda_{-1} + \lambda_1$. If λ is of Fourier degree ≤ 1 , we may more succinctly write the thermostat (4.5) as

$$(M, e^{-2f}g, e^f\lambda_0 + \tilde{\pi}_1^*(\theta - \star df)).$$

Proposition 4.7. *Let (M, g, λ) and $(M, \tilde{g}, \tilde{\lambda})$, with λ and $\tilde{\lambda}$ of Fourier degree ≤ 1 , be two Anosov thermostats. Suppose there is a smooth orbit equivalence $\phi : \tilde{S}M \rightarrow SM$, isotopic to the identity, between them. If $\star\theta$ or $\tilde{\star}\tilde{\theta}$ is closed, then $\star\theta - \tilde{\star}\tilde{\theta}$ is exact.*

Proof. By Lemma 2.1, we have $\operatorname{div}_\mu F = V(\lambda) = -\pi_1^*(\star\theta)$, so an application of Lemma 2.2 gives us

$$(\phi^*c)\phi^*(\pi_1^*(\star\theta)) = \tilde{\pi}_1^*(\tilde{\star}\tilde{\theta}) - \tilde{F} \left(\ln \left(\frac{\det \phi}{\phi^*c} \right) \right),$$

where $c \in \mathcal{C}^\infty(SM, \mathbb{R}_{>0})$ is such that $\phi_*\tilde{F} = cF$. Therefore, if $\tilde{\gamma}$ is a closed thermostat geodesic on M with respect to $(M, \tilde{g}, \tilde{\theta})$, and $\tilde{\Gamma} := (\tilde{\gamma}, \dot{\tilde{\gamma}}) \subseteq \tilde{S}M$, we have

$$\int_{\phi(\tilde{\Gamma})} c\pi_1^*(\star\theta) = \int_{\tilde{\Gamma}} \phi^*(c\pi_1^*(\star\theta)) = \int_{\tilde{\Gamma}} \tilde{\pi}_1^*(\tilde{\star}\tilde{\theta}).$$

Without loss of generality, suppose that $d(\star\theta) = 0$. Then, since ϕ is isotopic to the identity and integrals of 1-forms over curves are independent of the parametrization, we have

$$\int_{\phi(\tilde{\Gamma})} c\pi_1^*(\star\theta) = \int_{\tilde{\Gamma}} \tilde{\pi}_1^*(\star\theta).$$

Putting these together, we conclude that

$$\int_{\tilde{\Gamma}} \tilde{\pi}_1^*(\star\theta - \tilde{\star}\tilde{\theta}) = 0.$$

An application of the smooth Livšic theorem [dlLMM86, Theorem 2.1] and [DP07, Theorem B] allows us to conclude that $\star\theta - \tilde{\star}\tilde{\theta}$ is exact, as desired.

□

We can now conclude the proof of Theorem 1.2.

Proof of Theorem 1.2. If two Gaussian thermostats (M, g, θ) and $(M, \tilde{g}, \tilde{\theta})$ with negative thermostat curvature are related by a smooth orbit equivalence isotopic to the identity, then Proposition 4.2 tells us that there exists a diffeomorphism $\psi : M \rightarrow M$ isotopic to the identity such that $\psi^*\tilde{g} = e^{2f}g$ for some $f \in \mathcal{C}^\infty(M, \mathbb{R})$.

It remains to show that, if either $\star\theta$ or $\tilde{\star}\tilde{\theta}$ is closed, then $\star(\psi^*\tilde{\theta}-\theta)$ is exact. The map $\phi \circ d\psi$ yields a smooth orbit equivalence isotopic to the identity between the Anosov Gaussian thermostats $(M, e^{2f}g, \psi^*\tilde{\theta})$ and (M, g, θ) . By Lemma 4.6, we may assume without loss of generality that $f = 0$. Proposition 4.7 then allows us to conclude. \square

APPENDIX A. SOLUTIONS TO THE TRANSPORT EQUATION AS EXTENSIONS

A key ingredient that we use in this paper is Theorem 2.15: it allows us to extend any holomorphic 1-form τ on M (seen as a function on SM) to a fiberwise holomorphic distribution $u \in \mathcal{D}'(SM)$ satisfying the transport equation $(F + V(\lambda))u = 0$. We say that the distribution u is an extension of τ in the sense that $u_1 = \pi_1^*\tau$.

This can be seen as part of a larger theme, which is to find distributional solutions of the transport equation $(F + V(\lambda))u = 0$ with some prescribed Fourier modes. The problem is closely related to the study of the surjectivity of the adjoint of the X-ray transform for thermostats, which in turn is key to understanding the injectivity of the X-ray transform operator itself.

As an example, the following extension result generalizes [Ain15, Theorem 1.6] and [AZ17, Theorem 1.7], which cover the cases of magnetic flows and Gaussian thermostats respectively.

Theorem A.1. *Let (M, g, λ) be an Anosov generalized thermostat. For any $f \in \mathcal{C}^\infty(M)$, there exists $u \in H^{-1}(SM)$ such that $(F + V(\lambda))u = 0$ and $u_0 = \pi^*f$.*

The argument crucially relies on the *Pestov identity* (see [DP07, Theorem 3.3]).

Theorem A.2. *Let (M, g, λ) be a generalized thermostat. Then*

$$\|VFu\|^2 = \|FVu\|^2 - \langle \mathbb{K}Vu, Vu \rangle + \|Fu\|^2 \quad (\text{A.1})$$

for all $u \in \mathcal{C}^\infty(SM)$.

Recall that the thermostat curvature \mathbb{K} for a generalized thermostat is defined in (1.4). In both [Ain15] and [AZ17], the proofs introduce the following property:

Definition A.3. Let $\alpha \in [0, 1]$. A generalized thermostat (M, g, λ) is α -controlled if

$$\|Fu\|^2 - \langle \mathbb{K}u, u \rangle \geq \alpha \|Fu\|^2$$

for all $u \in \mathcal{C}^\infty(SM)$.

They then show that magnetic flows and Gaussian thermostats are α -controlled for some $\alpha > 0$ whenever they are Anosov. However, using the Pestov identity (A.1) for generalized thermostats, the same proof as in [AZ17, Theorem 3.1] goes through for the more general case.

Theorem A.4. *Let (M, g, λ) be an Anosov generalized thermostat. Then, there exists $\alpha > 0$ such that*

$$\|Fu\|^2 - \langle \mathbb{K}u, u \rangle \geq \alpha (\|Fu\|^2 + \|u\|^2)$$

for all $u \in C^\infty(SM)$. In particular, (M, g, λ) is α -controlled.

The rest of the argument in [AZ17] can then also be recycled to prove Theorem A.1.

The next theorem is again in the spirit of extending functions with low Fourier degree: it deals with functions induced from 1-forms on M . The result requires a technical condition, which is that the smooth 1-form θ being considered needs to be *solenoidal* (or divergence-free) in the sense that $\delta\theta = 0$. Here δ is the co-differential with respect to the metric g acting on 1-forms, i.e., $\delta = -\star d\star$. If we write $\pi_1^*\theta = a_{-1} + a_1 \in \Omega_{-1} \oplus \Omega_1$, then $\delta\theta = 0$ if and only if $\eta_+a_{-1} + \eta_-a_1 = 0$ (see [PSU14]).

Theorem A.5. *Let (M, g, λ) be an Anosov generalized thermostat. For any solenoidal smooth 1-form θ on M , there exists $u \in H^{-1}(SM)$ with $u_0 = 0$ such that $(F + V(\lambda))u = 0$ and $u_{-1} + u_1 = \pi_1^*\theta$.*

This is a generalization of [Ain15, Theorem 1.7] and [AZ17, Theorem 1.8], which again deal with the magnetic and Gaussian thermostat cases respectively. However, adapting the proofs requires some care. We will need the following lemma.

Lemma A.6. *For any $\lambda \in C^\infty(SM)$, the L^∞ -norms of its Fourier coefficients $\{\lambda_k\}_{k \in \mathbb{Z}}$ are rapidly decaying in the sense that, for all $\alpha \in \mathbb{N}$, we have*

$$\sup_{k \in \mathbb{Z}} \|\lambda_k\|_{L^\infty(SM)} k^\alpha < +\infty.$$

Proof. For any point on SM , let $U \subseteq SM$ be an open neighborhood admitting smooth coordinates $(x, \theta) \in \mathbb{R}^2 \times \mathbb{S}^1$ such that $V \equiv \partial_\theta$. The Sobolev embedding theorem gives us a constant $C > 0$ such that

$$\|\lambda_k\|_{L^\infty(U)} \leq C \sum_{|\beta| \leq 2} \|D^\beta \lambda_k\|_{L^2(U)}$$

for all $k \in \mathbb{Z}$. Here we use the multi-index notation $\beta = (\beta_1, \beta_2, \beta_3) \in \mathbb{N}^3$ and define $D^\beta := \partial_{x_1}^{\beta_1} \partial_{x_2}^{\beta_2} \partial_\theta^{\beta_3}$.

Using the explicit formula (2.4) on U , we may write

$$\lambda_k(x, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \lambda(x, \theta + t) e^{-ikt} dt.$$

Therefore, still on the domain U , we can see that

$$D^\beta \lambda_k = (D^\beta \lambda)_k.$$

By compactness, we can cover SM with a finite number of such open sets $\{U_j\}$. Then, we get

$$\begin{aligned} \|\lambda_k\|_{L^\infty(SM)} &= \max_j \|\lambda_k\|_{L^\infty(U_j)} \\ &\leq \max_j C \sum_{|\beta| \leq 2} \|(D^\beta \lambda)_k\|_{L^2(U_j)} \\ &\leq C \sum_{|\beta| \leq 2} \|(D^\beta \lambda)_k\|_{L^2(SM)}. \end{aligned}$$

Since the L^2 -norms of the Fourier coefficients of a smooth function are rapidly decaying, we obtain the desired result. \square

The following lemma has the same proof as [AZ17, Theorem 1.8]. The argument relies on the Pestov identity and Theorem A.4.

Lemma A.7. *Let (M, g, λ) be an Anosov generalized thermostat. Then there exists a constant $C > 0$ such that*

$$\|u\|_{H^1(SM)} \leq C \|VFu\|$$

for all $u \in \bigoplus_{|k| \geq 1} \Omega_k$

Next, we define the projection operator $T : \mathcal{C}^\infty(SM) \rightarrow \bigoplus_{|k| \geq 2} \Omega_k$ by

$$Tu := \sum_{|k| \geq 2} u_k.$$

We also define $Q : \mathcal{C}^\infty(SM) \rightarrow \bigoplus_{|k| \geq 2} \Omega_k$ as $Q := TVF$.

Lemma A.8. *Let (M, g, λ) be an Anosov generalized thermostat. Then there exists a constant $C > 0$ such that*

$$\|u\|_{H^1(SM)} \leq C \|Qu\|$$

for all $u \in \bigoplus_{|k| \geq 1} \Omega_k$.

Proof. In this proof, we will let $C > 0$ be a constant which can change from line to line to simplify the notation.

Let $u \in \bigoplus_{|k| \geq 1} \Omega_k$. From the definition of Q , we have

$$\|VFu\|^2 = \|(Fu)_1\|^2 + \|(Fu)_{-1}\|^2 + \|Qu\|^2.$$

From Lemma (A.7), we know that

$$\|u\|_{H^1(SM)} \leq C \|VFu\|,$$

so it remains to show that $\|(Fu)_{\pm 1}\| \leq C \|Qu\|$.

By Theorem A.4 and the Pestov identity (A.1), we have

$$\|VFu\|^2 \geq \|Fu\|^2 + \alpha\|FVu\|^2 + \alpha\|Vu\|^2.$$

Therefore, we obtain

$$\|Qu\|^2 \geq \alpha\|Vu\|^2 \geq \alpha \sum_{k \in \mathbb{Z}} k^2 \|u_k\|^2. \quad (\text{A.2})$$

It also gives us

$$\begin{aligned} \|Qu\|^2 &\geq \alpha\|FVu\|^2 \\ &\geq \alpha\|(FVu)_1\|^2 + \alpha\|(FVu)_{-1}\|^2 \\ &= \alpha \left\| 2i\eta_- u_2 - \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k+1} u_k \right\|^2 + \alpha \left\| -2i\eta_+ u_{-2} - \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k-1} u_k \right\|^2. \end{aligned}$$

Therefore

$$\left\| 2i\eta_- u_2 - \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k+1} u_k \right\| \leq C\|Qu\|$$

and

$$\left\| 2i\eta_+ u_{-2} + \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k-1} u_k \right\| \leq C\|Qu\|.$$

By the reverse triangle inequality, we get

$$\|2\eta_- u_2\| - \left\| \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k+1} u_k \right\| \leq C\|Qu\|$$

and

$$\|2\eta_+ u_{-2}\| - \left\| \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k-1} u_k \right\| \leq C\|Qu\|.$$

By the triangle inequality, the Cauchy-Schwarz inequality, Lemma A.6, and property (A.2), we obtain

$$\begin{aligned} \left\| \sum_{k \in \mathbb{Z}} k^2 \lambda_{-k \pm 1} u_k \right\| &\leq \sum_{k \in \mathbb{Z}} k^2 \|\lambda_{-k \pm 1}\|_{L^\infty(SM)} \|u_k\| \\ &\leq \left(\sum_{k \in \mathbb{Z}} k^2 \|\lambda_{-k \pm 1}\|_{L^\infty(SM)}^2 \right)^{1/2} \left(\sum_{k \in \mathbb{Z}} k^2 \|u_k\|^2 \right)^{1/2} \\ &\leq C \left(\sum_{k \in \mathbb{Z}} k^2 \|u_k\|^2 \right)^{1/2} \\ &\leq C\|Qu\|. \end{aligned}$$

This gives us

$$\|\eta_- u_2\| \leq C\|Qu\| \quad \text{and} \quad \|\eta_+ u_{-2}\| \leq C\|Qu\|.$$

The result then follows because

$$\begin{aligned} \|(Fu)_1\| &= \left\| \eta_- u_2 + \sum_{k \in \mathbb{Z}} ik \lambda_{-k+1} u_k \right\| \\ &\leq \|\eta_- u_2\| + \sum_{k \in \mathbb{Z}} k \|\lambda_{-k+1}\|_{L^\infty(SM)} \|u_k\| \\ &\leq C \|Qu\| \end{aligned}$$

and

$$\begin{aligned} \|(Fu)_{-1}\| &= \left\| \eta_+ u_{-2} + \sum_{k \in \mathbb{Z}} ik \lambda_{-k-1} u_k \right\| \\ &\leq \|\eta_+ u_{-2}\| + \sum_{k \in \mathbb{Z}} k \|\lambda_{-k-1}\|_{L^\infty(SM)} \|u_k\| \\ &\leq C \|Qu\|. \end{aligned}$$

□

The rest of the proof of Theorem A.5 then goes exactly as in [AZ17].

REFERENCES

- [Ain15] Gareth Ainsworth. The magnetic ray transform on Anosov surfaces. *Discrete and Continuous Dynamical Systems*, 35(5):1801–1816, 2015.
- [AR21] Yernat M. Assylbekov and Franklin T. Rea. The attenuated ray transforms on Gaussian thermostats with negative curvature. [arXiv:2102.04571](#), 2021.
- [AZ17] Yernat M. Assylbekov and Hanming Zhou. Invariant distributions and tensor tomography for Gaussian thermostats. *Communications in Analysis and Geometry*, 25(5):895–926, 2017.
- [BT82] Raoul Bott and Loring W. Tu. *Differential Forms in Algebraic Topology*. Springer, 1982.
- [CP22] Mihajlo Cekić and Gabriel P. Paternain. Resonant forms at zero for dissipative Anosov flows. [arXiv:2211.06255](#), 2022.
- [dlLMM86] Rafael de la Llave, José Manuel Marco, and Roberto Moriyón. Canonical perturbation theory of Anosov systems and regularity results for the Livsic cohomology equation. *Annals of Mathematics*, 123(3):537–611, 1986.
- [DP05] Nurlan S. Dairbekov and Gabriel P. Paternain. Longitudinal KAM-cocycles and action spectra of magnetic flows. *Mathematical Research Letters*, 12:719–730, 2005.
- [DP07] Nurlan S. Dairbekov and Gabriel P. Paternain. Entropy production in Gaussian thermostats. *Communications in Mathematical Physics*, 269:533–543, 2007.
- [DS03] Nurlan S. Dairbekov and Vladimir A. Sharafutdinov. Some problems of integral geometry on Anosov manifolds. *Ergodic Theory and Dynamical Systems*, 23(1):59–74, 2003.
- [FK92] Hershel M. Farkas and Irwin Kra. *Riemann Surfaces*, volume 71 of *Graduate Texts in Mathematics*. Springer-Verlag, 2 edition, 1992.
- [FM11] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, 2011.

- [FT99] Friedrich Gerard Friedlander and Mark Toshi. *Introduction to the Theory of Distributions*. Cambridge University Press, 2 edition, 1999.
- [Ghy84] Étienne Ghys. Flots d’Anosov sur les 3-variétés fibrées en cercles. *Ergodic Theory and Dynamical Systems*, 4:67–80, 1984.
- [GK80] Victor Guillemin and David Kazhdan. Some inverse spectral results for negatively curved 2-manifolds. *Topology*, 19(3):301–312, 1980.
- [GLP24] Colin Guillarmou, Thibault Lefeuvre, and Gabriel P. Paternain. Marked length spectrum rigidity for Anosov surfaces. *Duke Mathematical Journal*, 2024.
- [Gro99] Stéphane Grogniet. Flots magnétiques en courbure négative . *Ergodic Theory and Dynamical Systems*, 19(2):413–436, 1999.
- [Gui17] Colin Guillarmou. Invariant distributions and X-ray transform for Anosov flows. *Journal of Differential Geometry*, 105(2):177–208, 2017.
- [Has94] Boris Hasselblatt. Regularity of the Anosov splitting and of horospheric foliations. *Ergodic Theory and Dynamical Systems*, 14(4):645–666, 1994.
- [Hö03] Lars Hörmander. *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*. Classics in Mathematics. Springer, 2 edition, 2003.
- [Hö09] Lars Hörmander. *The Analysis of Linear Partial Differential Operators IV. Fourier Integral Operators*. Classics in Mathematics. Springer, 2 edition, 2009.
- [MP19] Thomas Mettler and Gabriel P. Paternain. Holomorphic differentials, thermostats and Anosov flows. *Mathematische Annalen*, 373:553–580, 2019.
- [MP20] Thomas Mettler and Gabriel P. Paternain. Convex projective surfaces with compatible Weyl connection are hyperbolic. *Analysis & PDE*, 13(4):1073–1097, 2020.
- [PS23] Gabriel P. Paternain and Mikko Salo. Carleman estimates for geodesic X-ray transforms. *Annales Scientifiques de l’École Normale Supérieure*, 56(5):1339–1380, 2023.
- [PSU14] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Spectral rigidity and invariant distributions on Anosov surfaces. *Journal of Differential Geometry*, 98:147–181, 2014.
- [PSU23] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. *Geometric Inverse Problems: With Emphasis on Two Dimensions*, volume 204 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2023.
- [PW08] Piotr Przytycki and Maciej P. Wojtkowski. Gaussian thermostats as geodesic flows of nonsymmetric linear connections. *Communications in Mathematical Physics*, 277:759–769, 2008.
- [Reb23] James Marshall Reber. Deformative magnetic marked length spectrum. *Bulletin of the London Mathematical Society*, 55(6):3077–3096, 2023.
- [Woj00] Maciej P. Wojtkowski. W-flows on Weyl manifolds and Gaussian thermostats. *Journal de Mathématiques Pures et Appliquées*, (10):953–974, 2000.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, CAMBRIDGE CB3 0WB, UK

Email address: je396@cam.ac.uk